

THE STABLE COOPERATIONS OF MORAVA K -THEORY AND THE FIBER PRODUCT OF AUTOMORPHISM GROUPS OF FORMAL GROUP LAWS

MASATERU INOUE

ABSTRACT. The Hopf algebra $K(n)_*(K(n))$, the stable cooperations of Morava K -theory at an odd prime, is well-known. Whereas the main part of $K(n)_*(K(n))$ corepresents the group-valued functor consisting of the automorphisms of a formal group law of height n , relations between the whole structure of $K(n)_*(K(n))$ and formal group laws have not been investigated well. Firstly, we constitute the fiber product of two natural transformations between the automorphism groups of formal group laws, and the corepresenting Hopf algebra C_* . Next, we construct a Hopf algebra homomorphism from C_* to $K(n)_*(K(n))$ via Boardman's stable comodule algebras. By Yagita's results and the algebra structure of $K(n)_*(K(n))$, we see that the homomorphism is an isomorphism. The isomorphism clarifies the relationship between the Hopf algebra structure of $K(n)_*(K(n))$ including the exterior algebra part and formal group laws.

1. INTRODUCTION

Throughout this paper, we fix an odd prime p . Let $K(n)_*(K(n))$ be the stable cooperations of n -th Morava K -theory $K(n)$ at p . For a spectrum E and a space X , we assign the degree n to an element of the cohomology $E^n(X)$. Hence an element of the homology $E_n(X)$ is of degree $-n$.

Our aim is to describe the Hopf algebra structure of $K(n)_*(K(n))$ in terms of formal group laws in a similar way to [6, 7]. The Hopf algebra $K(n)_*(K(n))$ is well-known by [14, 15, 18]. The coefficient ring is $K(n)^* \cong \mathbb{F}_p[v_n, v_n^{-1}]$ with $\deg v_n = -2(p^n - 1)$, and there is a $K(n)_*$ -algebra isomorphism

$$K(n)_*(K(n)) \cong \Sigma(n) \otimes E(\tau_0, \tau_1, \dots, \tau_{n-1}), \quad (\Sigma(n) = K(n)_*[t_1, t_2, \dots] / (v_n t_i^{p^n} - v_n^{p^i} t_i)) \quad (1.1)$$

with $\deg \tau_i = -(2p^i - 1)$, $\deg t_i = -2(p^i - 1)$. Here E is an exterior algebra. The coproduct of τ_i is given by

$$\Delta(\tau_i) = 1 \otimes \tau_i + \sum_{j=0}^i \tau_j \otimes t_{i-j}^{p^j}$$

The coproduct on

$$\Sigma(n) \cong K(n)_* \otimes_{BP_*} BP_*(BP) \otimes_{BP_*} K(n)_*$$

is induced by the one on the stable cooperations $BP_*(BP)$ of the Brown-Peterson spectrum BP . See [1, 11] for more details on BP . Though $\Sigma(n)$ corepresents the group-valued functor of the automorphisms of the Honda formal group law H_n with the p -series $[p](x) = v_n x^{p^n}$, little is known about relations between the whole structure of $K(n)_*(K(n))$ including the exterior algebra part and formal group laws.

We construct three group-valued functors $\text{Aut}_{H_n}(-)$, $\text{Aut}_{g_a}(-)$ and $\text{Aut}_{G_a}(-)$ consisting of the automorphism groups of the formal group laws H_n , g_a and G_a in Sections 2,3 and 4. There exist natural transformations

$$\alpha : \text{Aut}_{H_n}(-) \longrightarrow \text{Aut}_{g_a}(-), \quad \beta : \text{Aut}_{G_a}(-) \longrightarrow \text{Aut}_{g_a}(-).$$

2020 *Mathematics Subject Classification*. Primary 55N22; Secondary 55S25, 57T05.

Key words and phrases. Morava K -theory, Steenrod algebra, stable cooperations, Hopf algebras, formal group laws, Boardman's stable comodule algebras.

The fiber product $C(-)$ of α and β is corepresented by a Hopf algebra C_* which is isomorphic to $K(n)_*(K(n))$. To relate $K(n)_*(K(n))$ to C_* , we need stable comodule algebras defined by Boardman [2]. This leads to a Hopf algebra homomorphism $\kappa^* : C_* \rightarrow K(n)_*(K(n))$. By using Yagita's results [18] and the algebra structure of $K(n)_*(K(n))$, we see that κ^* is an isomorphism. We emphasize that κ^* is constructed naturally, although we can not prove that κ^* is an isomorphism in a self-contained manner.

Firstly, we survey [6] before explaining the details of this paper. The author has introduced functors $\text{Aut}_{\hat{g}_a}(-)$ and $\text{Op}_H(-)$ from the category of non-negative graded commutative \mathbb{F}_2 -algebras to the category of groups, and a natural transformation $\text{Op}_H(-) \rightarrow \text{Aut}_{\hat{g}_a}(-)$ as follows. It induces the Hopf algebra structure of the mod 2 dual Steenrod algebra $H_*(H)$ directly. Let H be the mod 2 Eilenberg-MacLane space, and R_* a non-negative graded commutative \mathbb{F}_2 -algebra. For the additive formal group law $\hat{g}_a(x, y) = x + y$ over R_* , a formal power series $f(x) = x + \sum_{i \geq 2} a_i x^i \in R_*[[x]]$ is called a *strict automorphism* of \hat{g}_a if it satisfies $f(x + y) = f(x) + f(y)$. We denote the set of strict automorphisms of \hat{g}_a by $\text{Aut}_{\hat{g}_a}(R_*)$. The product of $\text{Aut}_{\hat{g}_a}(R_*)$ is defined as $f \cdot g = g \circ f(x)$. Then $\text{Aut}_{\hat{g}_a}(-)$ is a group-valued functor. It is easily seen that $\text{Aut}_{\hat{g}_a}(-)$ is corepresented by the Hopf algebra

$$\bar{A}_* = \mathbb{F}_2[\xi_1, \xi_2, \dots, \xi_n, \dots] \quad (\deg \xi_n = -(2^n - 1))$$

with coproduct $\xi_n \mapsto \sum_{0 \leq i \leq n} \xi_{n-i}^{2^i} \otimes \xi_i$. In other words, we have a natural isomorphism

$$\text{Aut}_{\hat{g}_a}(-) \cong \text{Hom}_{\mathbb{F}_2\text{-alg}}(\bar{A}_*, -).$$

Let X be a CW-complex, and $\gamma : H^*(X) \rightarrow H^*(X) \hat{\otimes} R_*$ a natural transformation with degree preserving. Here the degree n part of $H^*(X) \hat{\otimes} R_*$ consists of $\prod_{n \geq 0} H^{k+n}(X) \otimes_{\mathbb{F}_2} R_n$ by abuse of notation.

Definition 1.1. A natural transformation $\gamma_X : H^*(X) \rightarrow H^*(X) \hat{\otimes} R_*$ is said to be *multiplicative* over R_* if γ_X satisfies the conditions:

- (i) The following diagram is commutative:

$$\begin{array}{ccc} H^*(X) \otimes H^*(Y) & \xrightarrow{\gamma_X \otimes \gamma_Y} & (H^*(X) \hat{\otimes} R_*) \otimes (H^*(Y) \hat{\otimes} R_*) \\ \downarrow \times & & \downarrow \\ & & (H^*(X) \hat{\otimes} H^*(Y)) \hat{\otimes} (R_* \otimes R_*) \\ & & \downarrow \times \otimes \mu \\ H^*(X \times Y) & \xrightarrow{\gamma_{X \times Y}} & H^*(X \times Y) \hat{\otimes} R_* \end{array}$$

where \times is the cross product, μ is the product on R_* , and the unlabeled map interchanges the second and third factors.

- (ii) $\gamma_{S^1}(u_1) = u_1 \otimes 1$ for the canonical generator u_1 of $H^1(S^1)$.

We define a functor $\text{Op}_H(-)$ which sends each non-negative graded \mathbb{F}_2 -algebra R_* to the set of multiplicative operations over R_* . For $\gamma_X, \gamma'_X \in \text{Op}_H(R_*)$, the product

$$\gamma_X \cdot \gamma'_X : H^*(X) \xrightarrow{\gamma'_X} H^*(X) \hat{\otimes} R_* \xrightarrow{\gamma_X \otimes 1} H^*(X) \hat{\otimes} R_* \otimes R_* \xrightarrow{1 \otimes \mu} H^*(X) \hat{\otimes} R_*. \quad (1.2)$$

gives $\text{Op}_H(R_*)$ a group structure, and $\text{Op}_H(-)$ is a group-valued functor. There exists a universal multiplicative operation

$$\rho_X : H^*(X) \longrightarrow H^*(X) \hat{\otimes} H_*(H)$$

in the following sense. Given $\gamma_X \in \text{Op}_H(R_*)$, we obtain a unique algebra homomorphism $\bar{\gamma} : H_*(H) \rightarrow R_*$ such that

$$\gamma_X = (1 \otimes \bar{\gamma}) \circ \rho_X : H^*(X) \xrightarrow{\rho_X} H^*(X) \hat{\otimes} H_*(H) \xrightarrow{1 \hat{\otimes} \bar{\gamma}} H^*(X) \hat{\otimes} R_*.$$

Therefore we have a natural isomorphism

$$\mathrm{Op}_H(-) \cong \mathrm{Hom}_{\mathbb{F}_2\text{-alg}}(H_*(H), -).$$

Note that ρ_X coincides with Milnor's coaction, which is studied in [8]. The universal multiplicative operation ρ_X is almost the same as the *universal stable comodule algebra* in [2, Theorem 12.8]. Though the author [6] has constructed ρ_X by using spectra, Boardman [2] has introduced ρ_X by studying various operations in terms of comonads.

To relate $\mathrm{Op}_H(-)$ to $\mathrm{Aut}_{\hat{g}_a}(-)$, we substitute the classifying space $X = B\mathbb{Z}/2$ into multiplicative operations γ_X . For the canonical generator x of $H^1(B\mathbb{Z}/2)$, we have $H^*(B\mathbb{Z}/2) \cong \mathbb{F}_2[x]$. The i -th projection $p_i : B\mathbb{Z}/2 \times B\mathbb{Z}/2 \rightarrow B\mathbb{Z}/2$ ($i = 1, 2$) induces $x_i = p_i^*(x) \in H^1(B\mathbb{Z}/2 \times B\mathbb{Z}/2)$, and we obtain $H^*(B\mathbb{Z}/2 \times B\mathbb{Z}/2) \cong \mathbb{F}_2[x_1, x_2]$. The classifying space $B\mathbb{Z}/2$ is an H -space, and we denote the product by $m : B\mathbb{Z}/2 \times B\mathbb{Z}/2 \rightarrow B\mathbb{Z}/2$. The pullback $m^*(x) = x_1 + x_2$ coincides with the additive formal group law \hat{g}_a .

For $\gamma \in \mathrm{Op}_H(R_*)$, we see that $\gamma_{B\mathbb{Z}/2}(x)$ is an element of the ring $R_*[[x]] \cong H^*(B\mathbb{Z}/2) \hat{\otimes} R_*$. Condition (i) in Definition 1.1 and the naturality of γ with respect to the product m lead to $\gamma(x) + \gamma(y) = \gamma(x + y)$. Condition (ii) implies $\gamma(x) = x + \sum_{i \geq 2} a_i x^i$. These mean that $\gamma(x)$ is a strict automorphism of \hat{g}_a over R_* . Assigning $\gamma(x)$ to γ , we have the natural transformation of group-valued functors

$$F : \mathrm{Op}_H(-) \cong \mathrm{Hom}_{\mathbb{F}_2\text{-alg}}(H_*(H), -) \longrightarrow \mathrm{Aut}_{\hat{g}_a}(-) \cong \mathrm{Hom}_{\mathbb{F}_2\text{-alg}}(\bar{A}, -).$$

The natural transformation F and the universal multiplicative operation ρ induce the Hopf algebra homomorphism

$$\overline{F(\rho)} : \bar{A}_* \longrightarrow H_*(H).$$

Moreover, Mui's work [9, 10] induces other multiplicative operations

$$S_n : H^*(X) \longrightarrow H^*(X) \hat{\otimes} D[n]_*,$$

where $D[n]_* = \mathbb{F}_2[\xi[n]_1, \dots, \xi[n]_n]$ with $\deg \xi[n]_i = -(2^n - 1)$. Since F is a natural transformation, we have the commutative diagram of algebras

$$\begin{array}{ccc} \bar{A}_* & \xrightarrow{\overline{F(\rho)}} & H_*(H) \\ & \searrow \overline{F(S_n)} & \downarrow \bar{S}_n \\ & & D[n]_* \end{array}$$

It is easily seen that $\overline{F(S_n)} : \bar{A}_* \rightarrow D[n]_*$ is given by $\xi_i \mapsto \xi[n]_i$ ($i \leq n$) and $\xi_i \mapsto 0$ ($i > n$). Therefore $\overline{F(S_n)}$ is an isomorphism in some range of degree for sufficiently large n , and we see that $\overline{F(\rho)} : \bar{A}_* \rightarrow H_*(H)$ is injective. From Serre's result [12], we obtain the Poincare series of $H_*(H)$, which see that $\overline{F(\rho)}$ is an isomorphism. Thus, we determine the Hopf algebra structure $H_*(H)$ directly via additive formal group laws.

Next, we recall the mod p dual Steenrod algebra $H_*(H)$ from [7]. Let H be the mod p Eilenberg-MacLane spectrum. To obtain $H_*(H)$, we need to use the classifying space $B\mathbb{Z}/p$ and a 2-dimensional additive formal group law instead of \hat{g}_a . For the standard generators $\epsilon \in H^1(B\mathbb{Z}/p)$ and $x \in H^2(B\mathbb{Z}/p)$, we have $H^*(B\mathbb{Z}/p) \cong E(\epsilon) \otimes \mathbb{F}_p[x]$. The multiplication m on $B\mathbb{Z}/p$ induces

$$m^*(\epsilon) = \epsilon_1 + \epsilon_2, \quad m^*(x) = x_1 + x_2,$$

where $\epsilon_i = p_i^*(\epsilon)$, $x_i = p_i^*(x)$ ($i = 1, 2$) for the i -th projection $p_i : B\mathbb{Z}/p \times B\mathbb{Z}/p \rightarrow B\mathbb{Z}/p$. We regard the pair $(m^*(\epsilon), m^*(x))$ as a 2-dimensional graded additive formal group law, which is denoted by \hat{G}_a . Let

R_* be a non-negative graded \mathbb{F}_p -algebra. If a pair of homogeneous power series

$$f(1)(\epsilon, x) = \epsilon + \sum_{i=0}^{\infty} a_i x^{p^i}, \quad f(2)(\epsilon, x) = x + \sum_{i=1}^{\infty} b_i x^{p^i} \in E(\epsilon) \otimes R_*[[x]]$$

with $\deg f(1) = 1$, $\deg f(2) = 2$ is an automorphism of \hat{G}_a over R_* , then the pair $(f(1), f(2))$ is called a *quasi-strict* automorphism of \hat{G}_a over R_* . If $a_1 = 0$, then it is said to be *strict* in [4]. Let $\text{Aut}_{\hat{G}_a}(R_*)$ be the set of quasi-strict automorphism groups of \hat{G}_a over R_* . The remaining parts proceed in a similar way to [6]. The natural transformation $\text{Op}_H(-) \rightarrow \text{Aut}_{\hat{G}_a}(-)$ induces the Hopf algebra structure of $H_*(H)$.

In [7], the author mainly uses the functor consisting of the automorphisms of a 1-dimensional additive formal group law over the dual numbers $R_*[\epsilon]/(\epsilon^2)$ instead of \hat{G}_a , and describes $\text{Aut}_{\hat{G}_a}(-)$ in Appendix. The functor is naturally isomorphic to $\text{Aut}_{\hat{G}_a}(-)$. Before [7], Yamaguchi [19, Appendix] has constructed the functor by using the dual numbers. Hopkins [5, Section 7] has called \hat{G}_a a supergroup $\mathbb{G}_a^{1,1}$ before [7]. The results have been studied previously though the author [7] has not refer to them.

Lastly, we outline this paper. In Section 2, we recall the Honda formal group law H_n of height n and the automorphism group functor $\text{Aut}_{H_n}(-)$. In Section 3, we survey formal group law chunks in [3], which is a truncated form of 1-dimensional formal group laws. We need only an additive formal group law chunk g_a and the automorphism group functor $\text{Aut}_{g_a}(-)$. In Section 4, we extend g_a and $\text{Aut}_{g_a}(-)$ to a 2-dimensional additive formal group law chunk G_a and the automorphism group functor $\text{Aut}_{G_a}(-)$, respectively. In a similar way to [7], the Hopf algebra B_* which corepresents the functor $\text{Aut}_{G_a}(-)$ yields the elements $\tau_0, \dots, \tau_{n-1}$ of the exterior algebra part of $K(n)_*(K(n))$. In Section 5, we construct natural transformations

$$\alpha : \text{Aut}_{H_n}(-) \longrightarrow \text{Aut}_{g_a}(-), \quad \beta : \text{Aut}_{G_a}(-) \longrightarrow \text{Aut}_{g_a}(-)$$

and the fiber product $C(-)$ of α and β . We define C_* by the Hopf algebra which corepresents the group-valued functor $C(-)$. In Section 6, we survey stable comodule algebras and multiplicative operations for a cohomology E in [2]. We define a multiplicative operation over an E_* -module R_*

$$\theta_X : E^*(X) \longrightarrow E^*(X) \hat{\otimes}_{E_*} R_*$$

in Definition 6.4. We write the set of multiplicative operations over R_* for $\text{Op}_E(R_*)$. According to [2], the multiplicative operations $\text{Op}_E(-)$ is only a set. This is because a multiplicative operation θ_X is not an E_* -module homomorphism in general. Therefore $\text{Op}_E(R_*)$ is not a group. If $E = H\mathbb{Z}/2$, $H\mathbb{Z}/p$ or $K(n)$, then $E_*(E)$ is a Hopf algebra. Hence we regard θ_X as an E_* -module homomorphism. This leads to the product $\theta_X \cdot \theta'_X$ in $\text{Op}_E(R_*)$ in a similar way to (1.2). Therefore $\text{Op}_E(-)$ is a group-valued functor. Moreover there is a universal multiplicative operation $\rho_X : E^*(X) \rightarrow E^*(X) \hat{\otimes}_{E_*} (E)$, which is called the universal comodule algebra in [2]. This induces a natural isomorphism

$$\text{Op}_E(-) \cong \text{Hom}_{E_*\text{-alg}}(E_*(E), -)$$

of group-valued functors. In section 7, we show that C_* is isomorphic to $K(n)_*(K(n))$. Substituting $X = \mathbb{C}P^\infty$, $\mathbb{C}P^{n-1}$, $B(\mathbb{Z}/p)^{[2p^n-1]}$ into multiplicative operations θ_X , we construct natural transformations

$$\kappa_1 : \text{Op}(-) \longrightarrow \text{Aut}_{H_n}(-), \quad \kappa_2 : \text{Op}(-) \longrightarrow \text{Aut}_{g_a}(-), \quad \kappa_3 : \text{Op}(-) \longrightarrow \text{Aut}_{G_a}(-).$$

By the definition of $C(-)$, we obtain

$$\kappa : \text{Op}_{K(n)}(-) \longrightarrow C(-).$$

It induces the Hopf algebra homomorphism $\kappa_* : C_* \rightarrow K(n)_*(K(n))$.

2. FORMAL GROUP LAWS OF HEIGHT n

Let \mathbf{Gp} be the category of groups, and \mathbf{Alg} the category of $K(n)_*$ -algebras.

In this section, we survey the Honda formal group law H_n of height n and the automorphisms of H_n . See [11, 16] for more details.

We express the coefficient ring of the Morava K -theory $K(n)$ as $K(n)^* = \mathbb{F}_p[v_n, v_n^{-1}]$, where $\deg(v_n) = -2(p^n - 1)$. We denote a $K(n)_*$ -algebra by R_* . Note that an element of R_n is of degree $-n$.

- Definition 2.1.**
- (1) The *Honda formal group law* H_n of height n over R_* is the formal group laws whose p -series $[p]_{H_n}$ is $v_n x^{p^n}$.
 - (2) Let x be an indeterminate of degree 2. A homogeneous power series $f(x)$ of degree 2 in $R_*[[x]]$ is a *strict automorphism* of H_n if $f(x)$ is an automorphism of H_n and $f(x) \equiv x \pmod{(x^2)}$.
 - (3) We denote by $\text{Aut}_{H_n}(R_*)$ the set of strict automorphism groups of H_n over R_* .
 - (4) We define a product of $f(x), g(x) \in \text{Aut}_{H_n}(R_*)$ by

$$f(x) \cdot g(x) = g \circ f(x).$$

where \circ is the composition of strict automorphisms. Then

$$\text{Aut}_{H_n}(-) : \mathbf{Alg} \longrightarrow \mathbf{Gp}$$

is a functor.

It is well-known [11] that the Hopf algebra

$$\Sigma(n) = K(n)_* \otimes_{BP_*} BP_*(BP) \otimes_{BP_*} K(n)_* \cong K(n)_*[t_1, t_2, \dots] / (t_i^{p^n} - v_n^{p^i - 1} t_i) \quad (\deg t_i = -(2p^i - 2))$$

corepresents the automorphism of H_n . Here BP is the Brown-Peterson spectrum.

Remark . The product in Definition 2.1 (4) is different from the usual product of automorphisms of H_n . Therefore the group-valued functor $\text{Aut}_{H_n}(-)$ is corepresented by $\bar{\Sigma}(n)$, the conjugation of $\Sigma(n)$. By abuse of notation, we denote the conjugation $c(t_i)$ by simply t_i . Hence we express $\bar{\Sigma}(n)$ as

$$\bar{\Sigma}(n) = K(n)_*[t_1, t_2, \dots] / (t_i^{p^n} - v_n^{p^i - 1} t_i).$$

Lemma 2.2. *Let $\bar{\Sigma}(n)$ be the conjugation of the Hopf algebra $\Sigma(n)$. We denote elements of $\bar{\Sigma}(n)$ as above. Assigning*

$$f(x) = \sum_{i \geq 0}^{H_n} \theta(t_i) x^{p^i} \in \text{Aut}_{H_n}(R_*)$$

to an element θ of $\text{Hom}_{K(n)_*\text{-alg}}(\bar{\Sigma}(n), R_*)$, we obtain the natural isomorphism

$$\text{Hom}_{K(n)_*\text{-alg}}(\bar{\Sigma}(n), -) \cong \text{Aut}_{H_n}(-)$$

Here \sum^{H_n} is the formal sum of H_n .

The Hopf algebra $\bar{\Sigma}(n)$ relates to $K(n)^*(\mathbb{C}P^\infty)$ later.

3. ADDITIVE FORMAL GROUP LAW CHUNKS

In [3, Section 3], a truncated form of a formal group law is called a *formal group law chunk*. We recall some definitions.

Definition 3.1. A *formal group law chunk* over a ring R of size n is a polynomial $F(x, y) \in R[x, y] / (x, y)^{n+1}$ which satisfies the conditions:

- (i) $F(0, x) = F(x, 0) = x$;
- (ii) $F(x, y) = F(y, x)$;
- (iii) $F(x, F(y, z)) = F(F(x, y), z)$.

Definition 3.2. Let F, G be formal group law chunks of size n . A *morphism* from F to G is a $f(x) \in R[x]/(x)^{n+1}$ which satisfies

- (i) The constant term of $f(x)$ is 0;
- (ii) $f(F(x, y)) = G(f(x), f(y))$

If $f(x) \equiv x \pmod{(x)^2}$, then it is called a *strict isomorphism*.

We need the definitions with grading. We denote a $K(n)_*$ -algebra by R_* .

Definition 3.3. (1) Let x and y be indeterminates of degree 2. A *formal group law chunk* over R_* of size n is a homogeneous polynomial $F(x, y) \in R_*[x, y]/(x, y)^{n+1}$ of degree 2 which satisfies the conditions in Definition 3.1.

- (2) A formal group law chunk $F(x, y) = x + y$ over R_* of size n is said to be *additive*. In particular, we denote an additive formal group law chunk of size $p^n - 1$ by $g_a(x, y) = x + y \in R_*[x, y]/(x, y)^{p^n}$.

Definition 3.4. (1) Let F, G be formal group law chunks of size n over R_* . A *morphism* from F to G is a homogeneous polynomial $f(x) \in R_*[x]/(x)^{n+1}$ of degree 2 which satisfies the condition of Definition 3.2. If $f(x) \equiv x \pmod{(x)^2}$, it is called a *strict isomorphism*.

- (2) We denote the strict automorphisms of g_a over R_* by $\text{Aut}_{g_a}(R_*)$. Then $\text{Aut}_{g_a}(-) : \mathbf{Alg} \rightarrow \mathbf{Gp}$ is a functor, and the product of $\text{Aut}_{g_a}(R_*)$ is defined as Definition 2.1 (4).

An automorphism $f(x)$ of g_a over R_* satisfies $f(x) + f(y) = f(x + y)$.

Lemma 3.5. *The group-valued functor $\text{Aut}_{g_a}(-)$ is corepresented by the Hopf algebra*

$$A_* = K(n)_*[\xi_1, \dots, \xi_{n-1}] \quad (\deg \xi_i = -2(p^i - 1))$$

whose coproduct is given by $\xi_k \mapsto \sum_{i=0}^k \xi_{k-i}^{p^i} \otimes \xi_i$. In other words, we have a natural isomorphism

$$\text{Hom}_{K(n)_*\text{-alg}}(A_*, -) \cong \text{Aut}_{g_a}(-) \quad (\theta \mapsto f(x) = x + \sum_{i=1}^{n-1} \theta(\xi_i)x^{p^i}).$$

Proof. We can easily see that a strict automorphism $f(x)$ of g_a is expressed as

$$f(x) = \sum_{k=0}^{n-1} a_k x^{p^k}, \tag{3.1}$$

where $a_0 = 1$ and $a_k \in R_{2p^k-2}$ ($k > 0$). We set the product of $\text{Aut}_{g_a}(R_*)$ as Definition 3.4 (2). The coproduct of A_* follows from calculation of the composition of strict automorphisms. \square

The Hopf algebra A_* relates to $K(n)^*(\mathbb{C}P^{p^n-1})$ later.

4. 2-DIMENSIONAL ADDITIVE FORMAL GROUP LAW CHUNKS

In [4], we define higher dimensional formal group laws and homomorphisms over a commutative ring. The author [7, Appendix] constructs graded version of the definitions. In fact, we need only a 2-dimensional additive formal group law over an \mathbb{F}_p -algebra, which leads to the dual Steenrod algebra for an odd prime p .

In this section, we introduce only 2-dimensional additive formal group law chunks, which corresponds to the partial coproduct of $K(n)^*(B(\mathbb{Z}/p)^{[2p^n-1]})$. We denote a $K(n)_*$ -algebra by R_* .

Definition 4.1. Let $R_*[x_1, x_2]/(x_1, x_2)^{p^n} \otimes E(\epsilon_1, \epsilon_2)$ be a graded R_* -algebra with $\deg \epsilon_i = 1$, $\deg x_i = 2$ ($i = 1, 2$). A pair

$$G_a(\epsilon_1, x_1; \epsilon_2, x_2) = (G_a(1)(\epsilon_1, x_1; \epsilon_2, x_2), G_a(2)(\epsilon_1, x_1; \epsilon_2, x_2))$$

of elements of $R_*[x_1, x_2]/(x_1, x_2)^{p^n} \otimes E(\epsilon_1, \epsilon_2)$ is called a *2-dimensional additive formal group law chunk* over R_* if it satisfies

$$G_a(1) = \epsilon_1 + \epsilon_2, \quad G_a(2) = x_1 + x_2.$$

The formal group law chunk G_a satisfies the definition of 2-dimensional formal group law chunks. See [7, Appendix] for more details on 2-dimensional formal group laws and isomorphisms. Note that $G_a(1)$ and $G_a(2)$ are homogeneous polynomials with $\deg G_a(1) = 1$ and $\deg G_a(2) = 2$, respectively.

Definition 4.2. (1) A pair $f(\epsilon, x) = (f(1)(\epsilon, x), f(2)(\epsilon, x))$ of elements of $R_*[x]/(x^{p^n}) \otimes E(\epsilon)$ with $\deg \epsilon = 1$, $\deg x = 2$ is an *automorphism* of G_a if it satisfies the following conditions:

- (i) The polynomials $f(1)$ and $f(2)$ are homogeneous of degree 1 and 2, respectively.
- (ii) $f(G_a(1), G_a(2)) = G_a(f(\epsilon_1, x_1); f(\epsilon_2, x_2))$, that is

$$\begin{aligned} f(1)(G_a(1), G_a(2)) &= G_a(1)(f(1)(\epsilon_1, x_1), f(2)(\epsilon_1, x_1); f(1)(\epsilon_2, x_2), f(2)(\epsilon_2, x_2)), \\ f(2)(G_a(1), G_a(2)) &= G_a(2)(f(1)(\epsilon_1, x_1), f(2)(\epsilon_1, x_1); f(1)(\epsilon_2, x_2), f(2)(\epsilon_2, x_2)) \end{aligned}$$

- (iii) The constant terms of $f(1), f(2)$ are 0.
- (2) A *quasi-strict* automorphism $f(\epsilon, x)$ of G_a satisfies that

$$f(1)(\epsilon, x) \equiv \epsilon + a_0x, \quad f(2)(\epsilon, x) \equiv x \pmod{(x)^2},$$

where $a_0 \in R_1$.

- (3) We denote the strict automorphisms of G_a over R_* by $\text{Aut}_{G_a}(R_*)$. Then $\text{Aut}_{G_a}(-) : \mathbf{Alg} \rightarrow \mathbf{Gp}$ is a functor, and the product of $\text{Aut}_{G_a}(R_*)$ is defined as

$$f \cdot g(\epsilon, x) = g \circ f(\epsilon, x) = (g(1)(f(1)(\epsilon, x), f(2)(\epsilon, x)), g(2)(f(1)(\epsilon, x), f(2)(\epsilon, x))),$$

where $f = (f(1), f(2))$, $g = (g(1), g(2)) \in \text{Aut}_{G_a}(R_*)$.

The functor is corepresented as follows.

Proposition 4.3. *Let*

$$B_* = K(n)_*[\xi_1, \dots, \xi_{n-1}] \otimes E(\tau_0, \dots, \tau_{n-1}) \quad (\deg \tau_n = -(2p^n - 1), \deg \xi_n = -2(p^n - 1))$$

be a Hopf algebra with coproduct

$$\tau_k \mapsto \tau_k \otimes 1 + \sum_{i=0}^k \xi_{k-i}^{p^i} \otimes \tau_i, \quad \xi_k \mapsto \sum_{i=0}^k \xi_{k-i}^{p^i} \otimes \xi_i.$$

Then B_* corepresents the group-valued functor $\text{Aut}_{G_a}(-)$. In other words, we have a natural isomorphism

$$\text{Hom}_{K(n)_*\text{-alg}}(B_*, -) \cong \text{Aut}_{G_a}(-) \quad (\theta \mapsto (f(1), f(2)) = \left(\epsilon + \sum_{i=0}^{n-1} \theta(\tau_i)x^{p^i}, x + \sum_{i=1}^{n-1} \theta(\xi_i)x^{p^i} \right)).$$

Proof. Condition (ii) in Definition 4.2 (1) is equivalent to

$$\begin{aligned} f(1)(\epsilon_1 + \epsilon_2, x_1 + x_2) &= f(1)(\epsilon_1, x_1) + f(1)(\epsilon_2, x_2) \\ f(2)(\epsilon_1 + \epsilon_2, x_1 + x_2) &= f(2)(\epsilon_1, x_1) + f(2)(\epsilon_2, x_2). \end{aligned} \tag{4.1}$$

In a similar calculation to [7, Appendix], it is easily seen that a quasi-strict automorphism $f = (f(1), f(2))$ is expressed as

$$\begin{aligned} f(1)(\epsilon, x) &= \epsilon + a_0x + a_1x^p + \dots + a_{n-1}x^{p^{n-1}} \in R_*[x]/(x)^{p^n} \otimes E(\epsilon) \\ f(2)(\epsilon, x) &= x + b_1x^p + \dots + b_{n-1}x^{p^{n-1}} \in R_*[x]/(x)^{p^n} \otimes E(\epsilon), \end{aligned} \tag{4.2}$$

where $a_i \in R_{2p^i-1}$, $b_i \in R_{2p^i-2}$. The coproduct is easily checked from Definition 4.2 (3). \square

The Hopf algebra B_* relates to the comodule algebra of $K(n)_*(B(\mathbb{Z}/p)^{[2p^n-1]})$.

5. THE FIBER PRODUCT OF AUTOMORPHISM GROUPS OF FORMAL GROUP LAWS

In this section, we study relations among the functors $\text{Aut}_{H_n}(-)$, $\text{Aut}_{g_a}(-)$ and $\text{Aut}_{G_a}(-)$ in Sections 2, 3 and 4 and constitute a new Hopf algebra C_* which is isomorphic to $K(n)_*(K(n))$.

Firstly, we study relations between H_n and g_a . The p -series of the Honda formal group law $H_n(x, y)$ is $[p]_{H_n}(x) = v_n x^{p^n}$. Via the quotient map

$$R_*[[x, y]] \longrightarrow R_*[x, y]/(x, y)^{p^n},$$

we regard H_n as a formal group law chunk of size $p^n - 1$, which is denoted by \bar{H}_n . The p -series $[p]_{\bar{H}_n}(x) \equiv 0$ implies $\bar{H}_n(x, y) = x + y \in R_*[x, y]/(x, y)^{p^n}$. Therefore, we obtain $\bar{H}_n(x, y) = g_a(x, y)$. It is easily seen that a strict automorphism $f(x)$ of H_n maps to the strict automorphism $\bar{f}(x)$ of g_a through the projection $R_*[[x]] \rightarrow R_*[x]/(x^{p^n})$. Explicitly, it is denoted by

$$f(x) = \sum_{i=0}^{\infty} {}^{H_n} a_i x^{p^i} \mapsto \bar{f}(x) = \sum_{i=0}^{n-1} a_i x^{p^i}, \quad (5.1)$$

where $a_0 = 1$.

Proposition 5.1. *The map (5.1) gives the natural transformation*

$$\alpha : \text{Aut}_{H_n}(R_*) \longrightarrow \text{Aut}_{g_a}(R_*).$$

Moreover α induces the Hopf algebra homomorphism

$$\alpha^* : A_* \cong K(n)_*[\xi_1, \dots, \xi_{n-1}] \longrightarrow \bar{\Sigma}(n) \cong K(n)_*[t_1, t_2, \dots]/(t_i^{p^n} - v_n^{p^i-1} t_i)$$

which is defined by $\alpha^*(\xi_i) = t_i$.

Proof. The latter part follows from Lemmas 2.2 and 3.5. □

Secondly, we discuss relations between G_a and g_a .

Proposition 5.2. *A natural group homomorphism*

$$\beta : \text{Aut}_{G_a}(R_*) \cong \text{Hom}(B_*, R_*) \longrightarrow \text{Aut}_{g_a}(R_*) \cong \text{Hom}(A_*, R_*)$$

is defined by

$$f(\epsilon, x) = (f(1)(\epsilon, x), f(2)(\epsilon, x)) \mapsto f(2)(\epsilon, x).$$

This leads to the Hopf algebra homomorphism

$$\beta^* : A_* = K(n)_*[\xi_1, \dots, \xi_{n-1}] \longrightarrow B_* = K(n)_*[\xi_1, \dots, \xi_{n-1}] \otimes E(\tau_0, \dots, \tau_{n-1}),$$

which is given by $\xi_i \mapsto \xi_i$.

Proof. We can express an element $f(\epsilon, x) = (f(1)(\epsilon, x), f(2)(\epsilon, x))$ of $\text{Aut}_{G_a}(R_*)$ as (4.2). It follows from (3.1) that $f(2)(\epsilon, x) \in R_*[[x]]$ is an automorphism of g_a . By Lemma 3.5 and Proposition 4.3, we obtain the homomorphism β^* . □

Definition 5.3. We denote by a functor $C(-)$ the fiber product of α and β . In other words, the group $C(R_*)$ is the fiber product of groups

$$\begin{array}{ccc} C(R_*) & \longrightarrow & \text{Aut}_{H_n}(R_*) \\ \downarrow & & \downarrow \alpha \\ \text{Aut}_{G_a}(R_*) & \xrightarrow{\beta} & \text{Aut}_{g_a}(R_*) \end{array} \quad (5.2)$$

where R_* is a $K(n)_*$ -algebra.

The functor $C(-)$ is corepresentable as follows.

Theorem 5.4. *The group-valued functor $C(-)$ is corepresented by the Hopf algebra*

$$\bar{\Sigma}(n) \otimes_{K(n)_*} E(\tau_0, \dots, \tau_{n-1}).$$

Here $\bar{\Sigma}(n)$ is the conjugation of $\Sigma(n)$ in Lemma 2.2. The coproduct of τ_i is given by

$$\Delta(\tau_i) = \tau_i \otimes 1 + \sum_{j=0}^i t_{i-j}^{p^j} \otimes \tau_j.$$

Proof. Since α and β are natural transformations of group-valued functors, the fiber product $C(R_*)$ inherits the group structure from α and β . The fiber product (5.2) implies that $C(-)$ is corepresented by the pushout

$$\begin{array}{ccc} C_* & \longleftarrow & \bar{\Sigma}(n)_* \\ \uparrow & & \uparrow \alpha^* \\ B_* & \longleftarrow & A_* \\ & & \downarrow \beta^* \end{array}$$

In other words, there exists a Hopf algebra C_* such that $K(-) \cong \text{Hom}_{K(n)_*\text{-alg}}(C_*, -)$, and C_* is isomorphic to

$$\bar{\Sigma}(n) \otimes_{A_*} B_* \cong \bar{\Sigma}(n) \otimes_{K(n)_*} E(\tau_0, \dots, \tau_{n-1}).$$

The coproducts $\Delta(\tau_i)$ derive from the coproduct of B_* in Proposition 4.3. □

This suggests that $K(n)_*(K(n))$ is expressed in terms of formal group laws.

6. BOARDMAN'S STABLE COMODULE ALGEBRAS

We recall stable comodules and stable comodule algebras from [2]. Let E be $H\mathbb{Z}/2$, $H\mathbb{Z}/p$, $K(n)$, the complex cobordism spectrum MU , the complex K -theory KU , or BP in this section. However we only need the case $E = K(n)$ for our purpose. In this section, we denote tensor products and completed tensor products over E_* by \otimes and $\hat{\otimes}$, respectively.

We introduce some categories. Let **Ho** denotes the homotopy category of CW complexes. A CW complex is called simply a space. **Stab** denotes the stable homotopy category of spectra. We write X for a space, and (X, o) for a space with a basepoint or a spectrum.

We denote the category of graded abelian groups by **Ab**, which consists of graded abelian groups and homomorphisms of degree 0. Let **FMod** be the category of complete Hausdorff filtered E^* -modules and continuous E^* -module homomorphisms of degree 0. Objects of **FMod** are completed Hausdorff E^* -modules with the filtration. In other words, an object M has submodules $F^a M \subset M$, and satisfies $M = \varprojlim_a M/F^a M$.

For a filtered E^* -module M with $F^a M \subset M$, we obtain the completed filtered E^* -module $\hat{M} = \varprojlim_a M/F^a M$. Since a usual tensor product $M \otimes N$ is rarely completed, we define the completed tensor product $M \hat{\otimes} N$ in **FMod** as

$$M \hat{\otimes} N = \varprojlim_{a,b} [(F/F^a M) \otimes (N/F^b N)],$$

where $F^b N \subset N$ is the filtration of N .

A complete Hausdorff commutative filtered E^* -algebra A has the product $\phi : A \otimes A \rightarrow A$, the unit $E^* \rightarrow A$ and the filtration of ideals $F^a A$ of A . Then ϕ is continuous, and we identify ϕ with the completed product $A \hat{\otimes} A \rightarrow A$. We denote by **FAlg** the category of complete Hausdorff commutative filtered E^* -algebras. The full subcategory **DAlg** of **FAlg** consists of objects M with trivial filtration $\{0\} \subset M$. Note that M is a discrete and completed E^* -module.

The cohomology $E^*(X)$ has the profinite filtration

$$F^a E^*(X) = \text{Ker}[E^*(X) \longrightarrow E^*(X_a)],$$

where X_a runs through all finite subcomplexes of X . Then the completion $E^*(X)^\wedge$ is an object of **FAlg**. If (X, o) is a spectrum, then $E^*(X, o)$ has the profinite filtration by using all finite subspectra of X similarly. The completed cohomology $E^*(X, o)^\wedge$ is an object of **FMod**. The homology $E_*(X)$ is a discrete E^* -module with trivial filtration, and therefore $E_*(X)$ is completed.

A ring spectrum E leads to the left or right action on the stable cooperations $E_*(E) = \pi_*(E \wedge E)$. Hence $E_*(E)$ is an E^* -bimodule. We only deal with E such that $E_*(E)$ is free as a left or right E^* -module. Moreover the product of E induces the product on $E_*(E)$

$$\phi : E_*(E) \otimes_{E_*(E)} E_*(E) \xrightarrow{\times} E_*(E \wedge E) \longrightarrow E_*(E),$$

where \times is the cross product. Then $E_*(E)$ is a commutative E^* -algebra with the left or right E^* -action. As usual, we denote the right unit function by $\eta_R : E^* \rightarrow E_*(E)$.

We use the following notations in [2], because of ambiguity of tensor products and bimodules. We rewrite ϕ as

$$\phi : E1_*(E2) \otimes_1 E1_*(E3) \xrightarrow{\times} E1_*(E2 \wedge E3) \longrightarrow E1_*(E),$$

which we call the *action scheme* of ϕ . Here Ei is a copy of E tagged for identification, and \otimes_i denotes \otimes_{Ei_*} .

The stable operations $E^*(E)$ acts on the cohomology $E^*(X)$. We have

$$\lambda_X : E^*(E) \otimes E^*(X) \longrightarrow E^*(X),$$

which is represented as $\lambda_X : E1^*(E2) \otimes_2 E2^*(X) \longrightarrow E1^*(X)$. The partial dualization of λ_X yields the Milnor coaction of E

$$\rho_X : E^*(X) \longrightarrow E^*(X) \hat{\otimes} E_*(E),$$

whose action scheme is $E2^*(X) \rightarrow E1^*(X) \hat{\otimes}_1 E1_*(E2)$. Note that $E_*(E)$ is a discrete E^* -module. Since ρ_X preserves the action of $E2_*$, the completion

$$\rho_X : E^*(X)^\wedge \longrightarrow E^*(X) \hat{\otimes} E_*(E)$$

is a natural transformation of functors from **Ho** to **FMod**. We can define the Milnor coaction for a spectrum (X, o) , which is denoted by

$$\rho_{(X, o)} : E^*(X, o)^\wedge \longrightarrow E^*(X, o) \hat{\otimes} E_*(E).$$

The composition and the unit of $E^*(E)$ have action scheme

$$E1^*(E2) \otimes_2 E2^*(E3) \xrightarrow{\circ} E1^*(E3), \quad \xi : E1^* \longrightarrow E1^*(E2). \quad (6.1)$$

By dualization, we obtain the coproduct and the counit

$$\psi : E1_*(E3) \rightarrow E1_*(E2) \otimes_2 E2_*(E3), \quad \epsilon : E_*(E) = \pi_*(E \wedge E) \longrightarrow E_* = \pi_*(E). \quad (6.2)$$

Since the composition (6.1) is associative, the comultiplication (6.2) is coassociative. Therefore $E_*(E)$ is a Hopf algebroid.

The cohomology $E^*(X)$ is an $E^*(E)$ -module. Therefore $E^*(X)$ is an $E_*(E)$ -comodule with the coaction ρ_X . By [2, Theorem 11.14], $E^*(X)^\wedge$ with the coaction ρ_X is a stable E^* -comodule in the following sense.

Definition 6.1 ([2]Definition 11.11). A *stable (E^* -cohomology) comodule* structure on a Hausdorff filtered E^* -module M , that is an object of **FMod**, consists of a coaction $\rho_M : M \rightarrow M \hat{\otimes} E_*(E)$ that is a

continuous morphism of filtered E^* -modules, that is a morphism in **FMod**, and satisfies the axioms:

$$\begin{array}{ccc}
 M & \xrightarrow{\rho_M} & M \hat{\otimes} E_*(E) & & M & \xrightarrow{\rho_M} & M \hat{\otimes} E_*(E) & & (6.3) \\
 \downarrow \rho_M & & \downarrow M \otimes \psi & & \searrow \cong & & \downarrow M \otimes \epsilon & & \\
 M \hat{\otimes} E_*(E) & \xrightarrow{\rho_M \otimes 1} & M \hat{\otimes} E_*(E) \hat{\otimes} E_*(E), & & & & M \hat{\otimes} E_* \cong M. & &
 \end{array}$$

Here ρ_M has action scheme $M2 \rightarrow M1 \hat{\otimes} E1_*(E2)$.

Then ρ_X has the following properties.

Theorem 6.2 ([2]Theorem 11.14). (1) For any space X , there is a natural coaction

$$\rho_X : E^*(X) \hat{\rhd} \rightarrow E^*(X) \hat{\otimes} E_*(E)$$

in **FMod** that makes $E^*(X) \hat{\rhd}$ a stable comodule.

(2) $\rho_X : E^*(X) \hat{\rhd} \rightarrow E^*(X) \hat{\otimes} E_*(E)$ is universal. In other words, given a discrete E^* -module N_* , any natural transformation of functors from **Ho** to **Ab**

$$\theta_X : E^*(X) \hat{\rhd} \rightarrow E^*(X) \hat{\otimes} N_*,$$

is induced from ρ_X by a unique morphism $f : E_*(E) \rightarrow N_*$ of left E^* -modules, as

$$\theta_X : E^*(X) \hat{\rhd} \xrightarrow{\rho_X} E^*(X) \hat{\otimes} E_*(E) \xrightarrow{1 \otimes f} E^*(X) \hat{\otimes} N_*.$$

Note that θ_X is not a morphism of **FMod** and that Theorem 6.2 is valid for any spectrum (X, o) .

From now on, we work in **FAlg**. Then we think ρ_X as a natural transformation of functors from **Ho** to **FAlg**. That is to say, we have the commutative diagram in [2, (12.3)]:

$$\begin{array}{ccc}
 E^*(X) \otimes E^*(Y) & \xrightarrow{\rho_X \otimes \rho_Y} & (E^*(X) \hat{\otimes} E_*(E)) \otimes (E^*(Y) \hat{\otimes} E_*(E)) & & (6.4) \\
 \downarrow \times & & \downarrow & & \\
 & & (E^*(X) \hat{\otimes} E^*(Y)) \hat{\otimes} (E_*(E) \otimes E_*(E)) & & \\
 & & \downarrow \times \otimes \phi & & \\
 E^*(X \times Y) & \xrightarrow{\rho_{X \times Y}} & E^*(X \times Y) \hat{\otimes} E_*(E), & &
 \end{array}$$

where \times is the cross product and the unlabeled map interchanges the second and third factors. If $X = Y$, then the cohomology $E^*(X)$ has inner product. Therefore ρ_X is a homomorphism of E^* -algebras.

Definition 6.3 ([2] Definition 12.7.). Let M be a E -cohomology stable comodule. We call M a (E -cohomology) stable comodule algebra if M is an object of **FAlg** and its coaction ρ_M is a morphism in **FAlg**. That is to say, the following diagram is commutative:

$$\begin{array}{ccc}
 M \otimes M & \xrightarrow{\rho_M \otimes \rho_M} & (M \hat{\otimes} E_*(E)) \otimes (M \hat{\otimes} E_*(E)) & & (6.5) \\
 \downarrow \mu & & \downarrow & & \\
 & & (M \hat{\otimes} M) \hat{\otimes} (E_*(E) \otimes E_*(E)) & & \\
 & & \downarrow \times \otimes \phi & & \\
 M & \xrightarrow{\rho_M} & M \hat{\otimes} E_*(E), & &
 \end{array}$$

where μ is the product of M and the unlabeled map interchanges the second and third factors.

Boardman [2] has not studied on products of $\theta_X : E^*(X) \rightarrow E^*(X) \hat{\otimes} R_*$ in Theorem 6.2 (2). We generalize stable comodule algebras as follows.

Definition 6.4. Let R_* be a discrete E^* -module and $E^*(-) \hat{\otimes} R_*$ a functor from **Ho** to **Ab**. We call a natural transformation of graded abelian group-valued functors

$$\theta_X : E^*(X) \longrightarrow E^*(X) \hat{\otimes} R_*$$

a *multiplicative operation* over R_* if the following diagram is commutative:

$$\begin{array}{ccc} E^*(X) \otimes_{\mathbb{Z}} E^*(Y) & \xrightarrow{\theta_X \otimes \theta_Y} & (E^*(X) \hat{\otimes} R_*) \otimes_{\mathbb{Z}} (E^*(Y) \hat{\otimes} R_*) \\ \downarrow \times & & \downarrow \\ & & (E^*(X) \hat{\otimes} E^*(Y)) \hat{\otimes} (R_* \otimes R_*) \\ & & \downarrow \times \otimes \phi \\ E^*(X \times Y) & \xrightarrow{\theta_{X \times Y}} & E^*(X \times Y) \hat{\otimes} R_* \end{array}$$

We denote the set of multiplicative operations over R_* by $\text{Op}_E(R_*)$.

The completed cohomology $E^*(X)^\wedge$ with the coaction $\rho_X : E^*(X)^\wedge \rightarrow E^*(X) \hat{\otimes} E_*(E)$ is a stable comodule algebra as follows.

Theorem 6.5 ([2] Theorem 12.8.). (1) *For any space X , the coaction ρ_X makes $E^*(X)^\wedge$ a stable comodule algebra.*

(2) ρ_X is universal. That is to say, for a discrete commutative E^* -algebra R_* , any multiplicative operation over R_*

$$\theta_X : E^*(X)^\wedge \longrightarrow E^*(X) \hat{\otimes} R_*$$

is induced from a unique E^* -algebra homomorphism $\bar{\theta} : E_*(E) \rightarrow R_*$ as follows:

$$\theta_X : E^*(X)^\wedge \xrightarrow{\rho_X} E^*(X) \hat{\otimes} E_*(E) \xrightarrow{1 \otimes \bar{\theta}} E^*(X) \hat{\otimes} R_*.$$

Consequently, there exists a natural isomorphism of functors from **DAlg** to **Set**

$$\text{Op}_E(R_*) \cong \text{Hom}_{E^*\text{-alg}}(E_*(E), R_*) \quad (\theta_X \mapsto \bar{\theta}),$$

where **Set** is the category of sets.

In general, a multiplicative operation $\theta_X : E^*(X)^\wedge \rightarrow E^*(X) \hat{\otimes} R_*$ is not an E^* -module homomorphism. Therefore we can not define iteration of multiplicative operations θ_X, θ'_X over R_*

$$(\theta'_X \otimes 1) \circ \theta_X : E^*(X) \rightarrow E^*(X) \hat{\otimes} R_* \hat{\otimes} R_*. \quad (6.6)$$

Since ρ_X is an E^* -module homomorphism, we can obtain iteration of ρ

$$(\rho_X \otimes 1) \circ \rho_X : E^*(X) \rightarrow E^*(X) \hat{\otimes} E_*(E) \hat{\otimes} E_*(E)$$

as Definition 6.1.

Lemma 6.6. *Let θ_X be an element of $\text{Op}_E(R_*)$, (S^k, o) the k -sphere spectrum and $u_k \in E^k(S^k, o)$ the standard generator. Then we have*

$$\theta_{(S^k, o)}(u_k) = u_k \otimes 1 \in E^*(S^k, o) \otimes R_*.$$

Proof. It follows from [2, (11.24)] that the universal coaction ρ_X on u_k is expressed as

$$\rho_{(S^k, o)}(u_k) = u_k \otimes 1 \in E^*(S^k, o) \otimes E_*(E).$$

By Theorem 6.5, we have

$$\theta_{(S^k, o)}(u_k) = u_k \otimes \bar{\theta}(1) = u_k \otimes 1 \in E^*(S^k, o) \otimes R_*,$$

where $\bar{\theta} : E_*(E) \rightarrow R_*$ is an E^* -algebra homomorphism induced from θ_X . \square

In the case that $E = H\mathbb{Z}/2$ or $H\mathbb{Z}/p$, we see that $E_*(E)$ is a Hopf algebra because of $E_* = \mathbb{Z}/2$ or \mathbb{Z}/p . Then any multiplicative operation

$$\theta_X : E^*(X)^\wedge \longrightarrow E^*(X) \hat{\otimes} R_*$$

is an E^* -module homomorphism. If $E_*(E)$ is a Hopf algebra, then we can define iteration (6.6). This is because we can define iteration of multiplicative operations in [6, 7].

Proposition 6.7. *Let a spectrum E be $H\mathbb{Z}2$, $H\mathbb{Z}p$ or $K(n)$, and $\theta_X \in \text{Op}_E(R_*)$.*

- (1) *The multiplicative operation $\theta_X : E^*(X) \rightarrow E^*(X) \hat{\otimes} R_*$ is an E^* -module homomorphism. Furthermore, we have the commutative diagram*

$$\begin{array}{ccc} E^*(X) \otimes E^*(Y) & \xrightarrow{\rho_X \otimes \rho_Y} & (E^*(X) \hat{\otimes} R_*) \otimes (E^*(Y) \hat{\otimes} R_*) \\ \downarrow \times & & \downarrow \\ & & (E^*(X) \hat{\otimes} E^*(Y)) \hat{\otimes} (R_* \otimes R_*) \\ & & \downarrow \times \otimes \phi \\ E^*(X \times Y) & \xrightarrow{\rho_{X \times Y}} & E^*(X \times Y) \hat{\otimes} R_* \end{array} \quad (6.7)$$

Therefore θ_X is a morphism in **FAlg** as Definition 6.3.

- (2) *We define a product on the multiplicative operations $\text{Op}_E(R_*)$ by the following composition*

$$\theta \cdot \theta' : E^*(-) \xrightarrow{\theta'} E^*(-) \hat{\otimes} R_* \xrightarrow{\theta \otimes 1} E^*(-) \hat{\otimes} R_* \hat{\otimes} R_* \xrightarrow{1 \otimes \phi} E^*(-) \hat{\otimes} R_*, \quad (6.8)$$

where $\theta, \theta' \in \text{Op}_E(R_*)$ and ϕ is the product of R_* . Then $\text{Op}_E(-)$ is a functor from **DAlg** to **Gp**. Furthermore the natural transformation in Theorem 6.5 (2)

$$\text{Op}_E(R_*) \cong \text{Hom}_{E^*\text{-alg}}(E_*(E), R_*)$$

is a group isomorphism equipped with the coproduct ψ on $E_*(E)$.

Proof. (1) Since $E_*(E)$ is a Hopf algebra, any multiplicative operation θ_X is an E_* -homomorphism as above. This leads to the commutative diagram (6.7).

- (2) By Theorem 6.5, the product $\theta \cdot \theta'$ is expressed as

$$E^*(X)^\wedge \xrightarrow{\rho_X} E^*(X) \hat{\otimes} E_*(E) \xrightarrow{\rho_X \otimes 1} E^*(X) \hat{\otimes} E_*(E) \hat{\otimes} E_*(E) \xrightarrow{1 \otimes \bar{\theta} \otimes \bar{\theta}'} E^*(X) \hat{\otimes} R_* \hat{\otimes} R_* \xrightarrow{1 \otimes \phi} E^*(X) \hat{\otimes} R_*.$$

By the commutative diagrams (6.3), we have

$$(\rho_X \otimes 1) \circ \rho_X = (1 \otimes \psi) \circ \rho_X : E^*(X)^\wedge \longrightarrow E^*(X) \hat{\otimes} E_*(E) \hat{\otimes} E_*(E).$$

Therefore we see that the product $\theta \cdot \theta'$ of $\text{Op}_E(R_*)$ corresponds to

$$\theta \cdot \theta' : E_*(E) \xrightarrow{\psi} E_*(E) \otimes E_*(E) \xrightarrow{\bar{\theta} \otimes \bar{\theta}'} R_* \otimes R_* \xrightarrow{\phi} R_* \square$$

From now on, we set $E = K(n)$, and abbreviate $\text{Op}_{K(n)}$ to Op . It is well-known [11] that $K(n)_*(K(n))$ is a Hopf algebra. Therefore we can adapt Proposition 6.7 for $E = K(n)$.

By using stable comodule algebras, we can prove this as follows. The p -th power map $p : \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$ induces the following formal p -series

$$p^*(x) = [p](x) = v_n x^{p^n} \in K(n)^*(\mathbb{C}P^\infty) \cong K(n)^*[[x]],$$

where x is the standard generator of $K(n)^2(\mathbb{C}P^\infty)$. The universal coaction ρ_X for $X = \mathbb{C}P^\infty$ is expressed as

$$\rho_{\mathbb{C}P^\infty} : K(n)^*(\mathbb{C}P^\infty) \cong K(n)^*[[x]] \longrightarrow K(n)^*(\mathbb{C}P^\infty) \hat{\otimes} K(n)_*(K(n)) \cong K(n)_*(K(n))[[x]].$$

We denote the formal power series $\rho_{\mathbb{C}P^\infty}(x) \in K(n)_*(K(n))[[x]]$ by $b(x) = \sum_{i=0}^{\infty} b_i x^i$ ($b_i \in K(n)_{2i-2}(K(n))$). The commutative diagram (6.4) implies

$$\begin{aligned} b([p](x)) &= [p]_R(b(x)) \\ b(v_n x^{p^n}) &= \eta_R(v_n)(b(x))^{p^n}, \end{aligned} \quad (6.9)$$

where $[p]_R(x) = \eta_R(v_n)x^{p^n} \in K(n)_*(K(n))[[x]]$. It follows from Lemma 6.6 in the case $S^2 = \mathbb{C}P^1$ that $b_0 = 0$ and $b_1 = 1$. The coefficient of x^{p^n} in the equality (6.9) yields $v_n = \eta_R(v_n)$.

7. RELATIONS BETWEEN $C(-)$ AND $\text{Op}(-)$

In this section, we construct a natural transformation κ between the functors $\text{Op}(-) = \text{Op}_{K(n)_*}(-)$ and $C(-)$ in Definition 5.3. Any $K(n)_*$ -algebra R_* can be treated as a completed Hausdorff $K(n)_*$ -algebra with trivial filtration. Therefore we identify the category \mathbf{Alg} as \mathbf{DAlg} , and $\text{Op}(-)$ and $C(-)$ are functors from \mathbf{Alg} to \mathbf{Gp} . The transformation κ leads to the Hopf algebra homomorphism $\kappa^* : C_* \rightarrow K(n)_*(K(n))$. By using [18] and the algebra structure of $K(n)_*(K(n))$, we see that κ^* is an isomorphism.

To obtain κ , we need three natural transformations

$$\kappa_1 : \text{Op}(-) \longrightarrow \text{Aut}_{H_n}(-), \quad \kappa_2 : \text{Op}(-) \longrightarrow \text{Aut}_{g_a}(-), \quad \kappa_3 : \text{Op}(-) \longrightarrow \text{Aut}_{G_a}(-)$$

such that the following diagram is commutative:

$$\begin{array}{ccc} \text{Op}(-) & \xrightarrow{\kappa_1} & \text{Aut}_{H_n}(-) \\ \kappa_3 \downarrow & \searrow \kappa_2 & \downarrow \alpha \\ \text{Aut}_{G_a}(-) & \xrightarrow{\beta} & \text{Aut}_{g_a}(-). \end{array} \quad (7.1)$$

Let $\theta_X : K(n)^*(X) \rightarrow K(n)^*(X) \hat{\otimes} R_*$ be a multiplicative operation, $\mathbb{C}P^\infty$ the infinite complex projective space, $\mathbb{C}P^{p^n-1}$ the $(p^n - 1)$ -dimensional complex projective space, and $B(\mathbb{Z}/p)^{[2p^n-1]}$ the $(2p^n - 1)$ -skeleton of the classifying space $B(\mathbb{Z}/p)$. Substituting $X = \mathbb{C}P^\infty, \mathbb{C}P^{p^n-1}, B(\mathbb{Z}/p)^{[2p^n-1]}$ into θ_X , we have the natural transformations $\kappa_1, \kappa_2, \kappa_3$.

Before proving these results, we fix generators of $K(n)^*(\mathbb{C}P^\infty)$, $K(n)^*(\mathbb{C}P^{p^n-1})$, $K(n)^*(B(\mathbb{Z}/p)^{[2p^n-1]})$. Let x be the canonical generator of $K(n)^2(\mathbb{C}P^\infty)$ which satisfies $i^*(x) = u_2 \in K(n)^2(S^2, o)$. Here $i : S^2 \rightarrow \mathbb{C}P^\infty$ is the usual inclusion, and u_k is the standard generator of $K(n)^k(S^k, o)$. Then we have

$$K(n)^*(\mathbb{C}P^\infty) \cong K(n)^*[[x]].$$

The inclusion $j_1 : \mathbb{C}P^{p^n-1} \rightarrow \mathbb{C}P^\infty$ and $j_2 : B(\mathbb{Z}/p)^{[2p^n-1]} \rightarrow \mathbb{C}P^{p^n-1}$ induce the elements $j_1^*(x) \in K(n)^2(\mathbb{C}P^{p^n-1})$ and $j_2^* \circ j_1^*(x) \in K(n)^2(B(\mathbb{Z}/p)^{[2p^n-1]})$, respectively. Using the same notation, we denote the elements by x . Then we obtain

$$K(n)^*(\mathbb{C}P^{p^n-1}) \cong K(n)^*[x]/(x^{p^n}).$$

The usual inclusion $i : S^1 \rightarrow B(\mathbb{Z}/p)^{[2p^n-1]}$ leads to the element u of $K(n)^1(B(\mathbb{Z}/p)^{[2p^n-1]})$ such that $i^*(u) = u_1 \in K(n)^1(S^1, o)$, and we obtain

$$K(n)(B(\mathbb{Z}/p)^{[2p^n-1]}) \cong E(u) \otimes K(n)^*[x]/(x^{p^n}).$$

Firstly, we study the case $X = \mathbb{C}P^\infty$. The product $\mu : \mathbb{C}P^\infty \times \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$ induces the coproduct of $K(n)^*(\mathbb{C}P^\infty) \cong K(n)^*[[x]]$ which is characterized by the p -series $[p](x) = v_n x^{p^n}$.

Proposition 7.1. *For $\theta_X \in \text{Op}(R_*)$ and the above generator $x \in K(n)^*(\mathbb{C}P^\infty)$, the power series*

$$\theta_{\mathbb{C}P^\infty}(x) \in K(n)^*(\mathbb{C}P^\infty) \hat{\otimes} R_* \cong R_*[[x]]$$

is a strict automorphism of H_n over R_ , that is, $\theta_{\mathbb{C}P^\infty}(x) \in \text{Aut}_{H_n}(R_*)$. Moreover, this induces the natural transformation*

$$\kappa_1 : \text{Op}(R_*) \cong \text{Hom}(K(n)_*(K(n)), R_*) \longrightarrow \text{Aut}_{H_n}(R_*) \cong \text{Hom}(\bar{\Sigma}(n), R_*) \quad (\theta \mapsto \theta_{\mathbb{C}P^\infty}(x)). \quad (7.2)$$

*of functors from **Alg** to **Gp**.*

Proof. We consider the diagram:

$$\begin{array}{ccc} K(n)^*(\mathbb{C}P^\infty) & \xrightarrow{\theta_{\mathbb{C}P^\infty}} & K(n)^*(\mathbb{C}P^\infty) \hat{\otimes} R_* \\ \downarrow p^* & & \downarrow p^* \otimes 1 \\ K(n)^*(\times^p \mathbb{C}P^\infty) & \xrightarrow{\theta} & K(n)^*(\times^p \mathbb{C}P^\infty) \hat{\otimes} R_* \\ \uparrow \cong \times & & \uparrow \times \otimes \phi \\ \hat{\otimes}^p K(n)^*(\mathbb{C}P^\infty) & \xrightarrow{\hat{\otimes}^p \theta_{\mathbb{C}P^\infty}} & \hat{\otimes}^p (K(n)^*(\mathbb{C}P^\infty) \hat{\otimes} (\hat{\otimes}^p R_*)) \\ \uparrow & & \uparrow \\ \hat{\otimes}^p K(n)^*(\mathbb{C}P^\infty) & \xrightarrow{\hat{\otimes}^p \theta_{\mathbb{C}P^\infty}} & \hat{\otimes}^p (K(n)^*(\mathbb{C}P^\infty) \hat{\otimes} R_*) \\ \downarrow \Delta^* & & \downarrow \Delta^* \otimes 1 \\ K(n)^*(\mathbb{C}P^\infty) & \xrightarrow{\theta_{\mathbb{C}P^\infty}} & K(n)^*(\mathbb{C}P^\infty) \hat{\otimes} R_* \end{array} \quad (7.3)$$

where $p : \times^p \mathbb{C}P^\infty = \mathbb{C}P^\infty \times \cdots \times \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$ is the p times product, $\Delta : \mathbb{C}P^\infty \rightarrow \times^p \mathbb{C}P^\infty$ is the diagonal, and $\theta_X \in \text{Op}(R_*)$. Since θ_X is natural on X , the upper diagram and the diagram including curved arrows are commutative. It follows from the commutative diagram (6.7) in Theorem 6.5 that the middle diagram is commutative. Since

$$[p](\theta_{\mathbb{C}P^\infty}(x)) = \theta_{\mathbb{C}P^\infty}([p](x)) \in K(n)^*(\mathbb{C}P^\infty) \hat{\otimes} R_* \cong R_*[[x]],$$

it implies that $\theta_{\mathbb{C}P^\infty}(x)$ is an automorphism of H_n over R_* . The standard inclusion $i : S^2 \rightarrow \mathbb{C}P^\infty$ induces

$$i^*(\theta_{\mathbb{C}P^\infty}(x)) = i^* \left(\sum_{i \leq 1} a_i x^i \right) = a_1 u_2.$$

By Lemma 6.6, we have $a_1 = 1$. Therefore $\theta_{\mathbb{C}P^\infty}(x)$ is a strict automorphism of H_n over R_* , and we obtain the natural transformation κ_1 . For elements θ, θ' of $\text{Op}(R_*)$, it follows from Definition 2.1 (4) and Proposition 6.7 (2) that

$$\begin{aligned} \kappa_1(\theta \cdot \theta') &= \phi \circ (\theta'_{\mathbb{C}P^\infty} \otimes 1) \circ \theta_{\mathbb{C}P^\infty}(x) = \theta'_{\mathbb{C}P^\infty}(\theta_{\mathbb{C}P^\infty}(x)) \\ &= \theta'_{\mathbb{C}P^\infty} \circ \theta_{\mathbb{C}P^\infty}(x) = \theta_{\mathbb{C}P^\infty}(x) \cdot \theta'_{\mathbb{C}P^\infty}(x) \in R_*[[x]] \cong K(n)^*[[x]] \hat{\otimes} R_*. \end{aligned}$$

Therefore κ_1 is a group homomorphism. \square

Secondly, we study the case $X = \mathbb{C}P^{p^n-1}$. Then $K(n)^*(\mathbb{C}P^{p^n-1}) \cong K(n)^*[x]/(x^{p^n})$. Though $\mathbb{C}P^{p^n-1}$ is not an H -space, we introduce the formal group structure of $K(n)^*[x]/(x^{p^n})$ by usage of partial multiplications

$$\mu_k : \mathbb{C}P^k \times \mathbb{C}P^{p^n-1-k} \longrightarrow \mathbb{C}P^{p^n-1}.$$

For $Y = \bigcup_{k=0}^{p^n-1} \mathbb{C}P^k \times \mathbb{C}P^{p^n-1-k}$, we can define $\mu : Y \longrightarrow \mathbb{C}P^{p^n-1}$, which induces

$$\mu^* : K(n)^*[x]/(x^{p^n}) \longrightarrow K(n)^*[x \times 1, 1 \times x]/(x \times 1, 1 \times x)^{p^n} \cong K(n)^*(Y).$$

Because of degree requirements, we have $\mu^*(x) = x \times 1 + 1 \times x$, which coincides with the additive group law chunk $g_a(x \times 1, 1 \times x)$ of size $p^n - 1$.

Proposition 7.2. *For $\theta_X \in \text{Op}(R_*)$ and the generator $x \in K(n)^*(\mathbb{C}P^{p^n-1}) \cong K(n)_*[x]/(x^{p^n})$, the polynomial*

$$\theta_{\mathbb{C}P^{p^n-1}}(x) \in K(n)^*(\mathbb{C}P^{p^n-1}) \hat{\otimes} R_* \cong R_*[x]/(x^{p^n})$$

is a strict automorphism of g_n over R_ , that is, $\theta_{\mathbb{C}P^{p^n-1}}(x) \in \text{Aut}_{g_a}(R_*)$. Furthermore, this induces the natural transformation*

$$\kappa_2 : \text{Op}(R_*) \cong \text{Hom}(K(n)_*(K(n)), R_*) \longrightarrow \text{Aut}_{g_a}(R_*) \cong \text{Hom}(A_*, R_*) \quad (\theta \mapsto \theta_{\mathbb{C}P^{p^n-1}}(x))$$

*of functors from **Alg** to **Gp**.*

Proof. Since θ_X is a multiplicative operation over R_* , we have the commutative diagram

$$\begin{array}{ccc} K(n)^*(\mathbb{C}P^k) \otimes K(n)^*(\mathbb{C}P^{p^n-1-k}) & \xrightarrow{\theta_{\mathbb{C}P^k} \otimes \theta_{\mathbb{C}P^{p^n-1-k}}} & (K(n)^*(\mathbb{C}P^k) \hat{\otimes} R_*) \otimes (K(n)^*(\mathbb{C}P^{p^n-1-k}) \hat{\otimes} R_*) \\ \downarrow \times & & \downarrow \\ & & (K(n)^*(\mathbb{C}P^k) \hat{\otimes} K(n)^*(\mathbb{C}P^{p^n-1-k})) \hat{\otimes} (R_* \hat{\otimes} R_*) \\ & & \downarrow \times \otimes \phi \\ K(n)^*(\mathbb{C}P^k \times \mathbb{C}P^{p^n-1-k}) & \xrightarrow{\theta_{\mathbb{C}P^k \times \mathbb{C}P^{p^n-1-k}}} & K(n)^*(\mathbb{C}P^k \times \mathbb{C}P^{p^n-1-k}) \hat{\otimes} R_* \end{array}$$

This induces

$$\theta_{\mathbb{C}P^k \times \mathbb{C}P^{p^n-1-k}}(x \times 1 + 1 \times x) = \theta_{\mathbb{C}P^k}(x) \times 1 + 1 \times \theta_{\mathbb{C}P^{p^n-1-k}}(x).$$

Therefore we see

$$\theta_Y(x \times 1 + 1 \times x) = \theta_Y(x \times 1) + \theta_Y(1 \times x) = \theta_{\mathbb{C}P^{p^n-1}}(x) \times 1 + 1 \times \theta_{\mathbb{C}P^{p^n-1}}(x).$$

By the equality and Lemma 6.6, we have $\theta_{\mathbb{C}P^{p^n-1}}(x) \in \text{Aut}_{g_a}(R_*)$. In a similar way to the proof of Proposition 7.1, we see that κ_2 is a group homomorphism. \square

Thirdly, we investigate $X = B(\mathbb{Z}/p)^{[2p^n-1]}$. In a similar way to $X = \mathbb{C}P^{p^n-1}$, we obtain the partial multiplications

$$m_k : B(\mathbb{Z}/p)^{[2k+1]} \times B(\mathbb{Z}/p)^{[2p^n-2k-2]} \longrightarrow X = B(\mathbb{Z}/p)^{[2p^n-1]}.$$

For $Z = \bigcup_{k=0}^{2p^n-2} B(\mathbb{Z}/p)^{[2k+1]} \times B(\mathbb{Z}/p)^{[2p^n-2k-2]}$, the multiplication $m : Z \rightarrow B(\mathbb{Z}/p)^{[2p^n-1]}$ induces

$$m^* : E(u) \otimes K(n)^*[x]/(x^{p^n-1}) \longrightarrow E(u \times 1, 1 \times u) \otimes K(n)^*[x \times 1, 1 \times x]/(x \times 1, 1 \times x)^{p^n}.$$

Then we can describe $m^*(x) = 1 \times x + x \times 1$, and $m^*(u) = u \times 1 + 1 \times u + \lambda v_n u x^{p^n-1}$ because of degree requirements. Since $B\mathbb{Z}/p$ is a homotopy associative H -space, one can easily check $m^*(u) = u \times 1 + 1 \times u$. Hence the pair $(m^*(u), m^*(x))$ coincides with the 2-dimensional additive formal group law chunk $G_a(u_1, u_2; x_1, x_2) = (G_a(1)(u_1, u_2; x_1, x_2), G_a(2)(u_1, u_2; x_1, x_2))$.

We can prove the following proposition in a similar way to Proposition 7.2.

Proposition 7.3. *Let θ_X be an element of $\text{Op}(R_*)$, and u and x the generators of $K(n)^*(B(\mathbb{Z}/p)^{[2p^n-1]}) \cong E(u) \otimes K(n)^*[x]/(x^{p^n})$. Then the pair*

$$\Theta(u, x) = (\theta_{B(\mathbb{Z}/p)^{[2p^n-1]}}(u), \theta_{B(\mathbb{Z}/p)^{[2p^n-1]}}(x))$$

of elements of $K(n)^(B(\mathbb{Z}/p)^{[2p^n-1]}) \hat{\otimes} R_* \cong E(u) \otimes R_*[x]/(x^{p^n})$ is a strict automorphism of G_a over R_* , that is, $\Theta(u, x) \in \text{Aut}_{G_a}(R_*)$. Assigning Θ to $\theta_X \in \text{Op}(R_*)$, we have the natural transformation*

$$\kappa_3 : \text{Op}(R_*) \cong \text{Hom}(K(n)_*(K(n)), R_*) \longrightarrow \text{Aut}_{G_a}(R_*) \cong \text{Hom}(B_*, R_*) \quad (7.4)$$

*of functors from **Alg** to **Gp**.*

Lastly, we construct a Hopf algebra homomorphism $\kappa^* : C_* \rightarrow K(n)_*(K(n))$, and show that κ^* is an isomorphism.

Theorem 7.4. *Let $C(-)$ be the functor in Definition 5.3. The natural transformation $\kappa_1, \kappa_2, \kappa_3$ induce the natural transformation*

$$\kappa : \text{Op}(R_*) \cong \text{Hom}_{K(n)\text{-alg}}(K(n)_*(K(n)), R_*) \longrightarrow C(R_*) \cong \text{Hom}_{K(n)\text{-alg}}(C_*, R_*),$$

*of functors from **Alg** to **Gp**. Therefore κ yields the Hopf algebra homomorphism*

$$\kappa^* : C_* \longrightarrow K(n)_*(K(n)).$$

Proof. It follows from the surjection

$$j_1^* : K(n)^*(\mathbb{C}P^\infty) \cong K(n)^*[x] \rightarrow K(n)_*(\mathbb{C}P^{p^n-1}) \cong K(n)^*[x]/(x^{p^n}) \quad (x \mapsto x)$$

that we have $\kappa_2 = \alpha \circ \kappa_1$. By the injection

$$j_2^* : K(n)^*(\mathbb{C}P^{p^n-1}) \cong K(n)^*[x]/(x^{p^n}) \rightarrow K(n)^*(B(\mathbb{Z}/p)^{[2p^n-1]}) \cong E(u) \otimes K(n)^*[x]/(x^{p^n}) \quad (x \mapsto x),$$

we have $\theta_{\mathbb{C}P^{p^n-1}}(x) = \theta_{B(\mathbb{Z}/p)^{[2p^n-1]}}(x)$. By Proposition 5.2, we obtain $\kappa_2 = \beta \circ \kappa_3$. The commutative diagram

$$\begin{array}{ccccc} \text{Op}(R_*) & & & & \\ & \searrow^{\kappa_1} & & & \\ & & C(R_*) & \longrightarrow & \text{Aut}_{H_n}(R_*) \\ & \searrow^{\kappa} & \downarrow & & \downarrow \alpha \\ & & \text{Aut}_{G_a}(R_*) & \xrightarrow{\beta} & \text{Aut}_{g_a}(R_*) \\ & \searrow_{\kappa_3} & & & \end{array}$$

induces the natural transformation κ and the corresponding homomorphism κ^* . □

We recall [2, Theorem 14.32 (a)], whose proof is due to Yagita [18].

Theorem 7.5. [2, 18] *Let $\rho_X : K(n)_*(X) \rightarrow K(n)^*(X) \hat{\otimes} K(n)_*(K(n))$ be the universal stable comodule algebra in Theorem 6.5. We express $\rho_{\mathbb{C}P^\infty}(x)$ and $\rho_{B(\mathbb{Z}/p)^{[2p^n-1]}}(u)$ as*

$$\rho_{\mathbb{C}P^\infty}(x) = \sum_{i=1}^{\infty} x^i \otimes b_i, \quad \rho_{B(\mathbb{Z}/p)^{[2p^n-1]}}(u) = \sum_{i=0}^{p^n-1} x^i \otimes a_i + \sum_{i=0}^{p^n-1} ux^i \otimes c_i,$$

respectively. Then $K(n)_(K(n))$ is generated by a_{p^i} ($0 \leq i < n$) and b_{p^i} ($i > 0$) as a $K(n)_*$ -algebra.*

Theorem 7.5 implies that $\kappa : C_* \rightarrow K(n)_*(K(n))$ is surjective. The algebra structure of $K(n)_*(K(n))$ is known by [11, 14, 15, 16, 17, 18], and these induce the following main theorem.

Theorem 7.6. $\kappa^* : C_* \rightarrow K(n)_*(K(n))$ *is an isomorphism as a Hopf algebra.*

Remark . The elements τ_i and t_j of C_* correspond to the conjugations of the elements τ_i and t_j of $K(n)_*(K(n))$ in (1.1).

REFERENCES

- [1] J. F. Adams, *Stable homotopy and generalised homology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, Ill.-London, 1974. x+373 pp.
- [2] J. M. Boardman, *Stable operations in generalized cohomology*. Handbook of algebraic topology, 585-686, North-Holland, Amsterdam, 1995.
- [3] R. Burklund, *A classification of formal group laws over torsion-free rings*, <https://math.mit.edu/research/undergraduate/spur/documents/2015burklund.pdf>
- [4] M. Hazewinkel, *Formal groups and applications*. Pure and Applied Mathematics, 78. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. xxii+573pp.
- [5] M. Hopkins, *Complex oriented cohomology theories and the language of stacks course notes for 18.917, TAUGHT BY MIKE HOPKINS*, <https://www.semanticscholar.org/paper/COMPLEX-ORIENTED-COHOMOLOGY-THEORIES-AND-THE-OF-FOR/d213545f8c1debase8fb642925ca32516276d56f>
- [6] M. Inoue, The Steenrod algebra and the automorphism group of additive formal group law. *J. Math. Kyoto Univ.* 45 (2005), no. 1, 39–55.
- [7] M. Inoue, Odd primary Steenrod algebra, additive formal group laws, and modular invariants. *J. Math. Soc. Japan* 58 (2006), no. 2, pp. 311–332.
- [8] J. Milnor, The Steenrod algebra and its dual. *Ann. of Math. (2)* 67 (1958), 150–171.
- [9] H. Mui, Modular invariant theory and cohomology algebras of symmetric groups. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* 22 (1975), no. 3, 319–369.
- [10] H. Mui, Dickson invariants and the Milnor basis of the Steenrod algebra, *Topology and applications*. *Colloq. Math. Soc. János Bolyai* 41 (1983), 345–355.
- [11] D. C. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, Pure and Applied Mathematics, 121. Academic Press, Inc., Orlando, FL, 1986. xx+413
- [12] J. P. Serre, Cohomologie modulo 2 des complexes d'Eilenberg-MacLane. *Comment. Math. Helv.* 27 (1953), 198–232.
- [13] N. E. Steenrod and D. B. A. Epstein, *Cohomology Operations*, Annals of Math. Studies No. 50, Princeton University Press, 1962
- [14] U. Würgler *On products in a family of cohomology theories associated to the invariant prime ideals of $\pi_*(BP)$* . *Comment. Math. Helv.* 52 (1977), no. 4, 457–481.
- [15] U. Würgler *On the relation of Morava K-theories to Brown-Peterson homology*. *Topology and algebra*, Monograph. Enseign. Math., 26, Univ. Genève, Geneva, 1978, 269–280
- [16] U. Würgler *Morava K-theories: a survey*. Algebraic topology Poznań 1989, 111–138, Lecture Notes in Math., 1474, Springer, Berlin, 1991.
- [17] N. Yagita, The exact functor theorem for BP_*/I_n -theory. *Proc. Japan Acad.* 52 (1976), no. 1, 1–3.
- [18] N. Yagita, A topological note on the Adams spectral sequence based on Morava's K-theory. *Proc. Amer. Math. Soc.* 72 (1978), no. 3, 613–617.
- [19] A. Yamaguchi, *On excess filtration on the Steenrod algebra*. *Proceedings of the Nishida Fest (Kinosaki 2003)*, 423–449, *Geom. Topol. Monogr.*, 10, Geom. Topol. Publ., Coventry, 2007.

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF SCIENCE, OKAYAMA UNIVERSITY OF SCIENCE, 1-1 RIDAICHO, KITA-KU, OKAYAMA-SHI 700-0005, JAPAN
Email address: inoue@xmath.ous.ac.jp