

On recurrence in zero-dimensional locally compact flow with compactly generated phase group

Xiongping Dai

Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China

Abstract

Let X be a zero-dimensional locally compact Hausdorff space not necessarily metric and G a compactly generated topological group not necessarily abelian or countable. We define recurrence at a point for any continuous action of G on X , and then, show that if \overline{Gx} is compact for all $x \in X$, the conditions (i) this dynamics is pointwise recurrent, (ii) X is a union of G -minimal sets, (iii) the G -orbit closure relation is closed in $X \times X$, and (iv) $X \ni x \mapsto \overline{Gx} \in 2^X$ is continuous, are pairwise equivalent. Consequently, if this dynamics is distal, then it is equicontinuous.

Keywords: Recurrence · Distality · Zero dimensional flow · Compactly generated group

2010 MSC: 37B05, 54H15

0 Introduction. Let (G, X) be a flow with phase group G and with phase space X , which is in the following sense unless stated otherwise:

- Ⓐ G is a Hausdorff semi-topological group, namely: G is a group with a Hausdorff topology under which $G \times G \xrightarrow{(s,t) \mapsto st} G$ is jointly continuous but $G \xrightarrow{t \mapsto t^{-1}} G$ need not continuous;
- Ⓑ X is a locally compact Hausdorff space; and moreover,
- Ⓒ there is an action of G on X , denoted $G \times X \xrightarrow{(t,x) \mapsto tx} X$ such that
 - (i) Continuity: $(t, x) \mapsto tx$ is jointly continuous;
 - (ii) Transformation group: $ex = x$ and $(st)x = s(tx)$ for all $x \in X$ and $s, t \in G$, where e is the identity of G ;
 - (iii) Poisson stability: Gx is relatively compact (i.e. \overline{Gx} is compact) in X for all $x \in X$.

By Ⓑ there is a compatible uniformity structure, denoted \mathcal{U} , on X . In the sequel, \mathfrak{N}_x stands for the neighborhood filter at x of X .

0.1. Let Y be a Hausdorff space and 2^Y the collection of non-empty closed subsets of Y . For a net $\{Y_i\}$ in 2^Y and $K \in 2^Y$, we say ' $Y_i \rightarrow K$ in 2^Y ' if

$$K = \{y \in Y \mid \exists \text{ a subnet } \{Y_{i_k}\} \text{ from } \{Y_i\} \text{ and } y_{i_k} \in Y_{i_k} \text{ s.t. } y_{i_k} \rightarrow y\}.$$

Lemma. If $K_i \rightarrow K$ in 2^Y and $\{K_{i_n}\}$ is any subnet of $\{K_i\}$ with $K_{i_n} \rightarrow K'$, then $K' \subseteq K$.

Email address: xpdai@nju.edu.cn (Xiongping Dai)

If Y is discrete and $\{Y_i\}_{i=1}^\infty$ with $Y_i \rightarrow K$, then $K = \bigcap_{n \geq 1} \bigcup_{i \geq n} Y_i = \limsup_{i \rightarrow \infty} Y_i$; in other words, $K = \{Y_i, \text{ i.o.}\}$.

0.2. Associated to (G, X) we define the orbit closure mapping $\mathcal{O}_{G,X}: X \rightarrow 2^X$ by $x \mapsto \overline{Gx}$. Then we say:

- a) $\mathcal{O}_{G,X}$ is ‘continuous’, provided that if $x_i \rightarrow x$ in X and $\overline{Gx_i} \rightarrow \mathfrak{D}$ in 2^X , then $\overline{Gx} = \mathfrak{D}$.
- b) $\mathcal{O}_{G,X}$ is upper semi-continuous, provided that for every $x \in X$ and every neighborhood V of \overline{Gx} there is a $U \in \mathfrak{N}_x$ such that $\overline{Gy} \subseteq V$ for all $y \in U$.

Clearly, if $\mathcal{O}_{G,X}$ is upper semi-continuous, then $\mathcal{O}_{G,X}$ is ‘continuous’. However, no converse is valid in general. Given $U \subset X$, define

- c) $U_G^* = \{x \in X \mid \overline{Gx} \subseteq U\}$.

If U is closed, then $U_G^* = \bigcap_{g \in G} gU$ is closed.

0.3. The ‘orbit closure relation R_o of (G, X) ’ is defined by $(x, y) \in R_o$ if $y \in \overline{Gx}$. It is not difficult to prove that

- a) $\mathcal{O}_{G,X}$ is ‘continuous’ iff R_o is closed in $X \times X$. (Warning: R_o closed $\nRightarrow \mathcal{O}_{G,X}$ upper semi-continuous in general.)

Indeed, suppose $\mathcal{O}_{G,X}$ is ‘continuous’, and let $R_o \ni (x_i, y_i) \rightarrow (x, y)$. Then $\overline{Gx_i} \rightarrow \overline{Gx}$ by 0.2 and $y_i \in \overline{Gx_i}$ with $y_i \rightarrow y$. So $y \in \overline{Gx}$ by 0.1 and $(x, y) \in R_o$. Thus R_o is closed. Conversely, assume R_o is closed, and let $x_i \rightarrow x$ in X and $\overline{Gx_i} \rightarrow \mathfrak{D}$ in 2^X . If $\overline{Gx} \neq \mathfrak{D}$, then $\overline{Gx} \not\subseteq \mathfrak{D}$. For $y \in \mathfrak{D} \setminus \overline{Gx}$ there is a net $y_{i_k} \in \overline{Gx_{i_k}}$ with $y_{i_k} \rightarrow y$. Since $(x_{i_k}, y_{i_k}) \in R_o$, so $(x, y) \in R_o$ and $y \in \overline{Gx}$ contrary to $y \in \mathfrak{D} \setminus \overline{Gx}$. Thus $\overline{Gx} = \mathfrak{D}$.

- b) If X is compact here, then $\mathcal{O}_{G,X}$ is ‘continuous’ iff it is upper semi-continuous iff it is continuous iff R_o is closed.

0.4. We say that a set K in X is ‘minimal’ under (G, X) if $\overline{Gx} = K$ for all $x \in K$. In this case K is a closed invariant subset of X , and moreover, K is compact by (iii).

Lemma. Let R_o be the orbit closure relation of (G, X) . Suppose $y \in X$ such that $R_o[y]$ is closed: if $(x_n, y) \in R_o$ with $x_n \rightarrow x$, then $(x, y) \in R_o$. Then \overline{Gy} is minimal.

Proof. Since \overline{Gy} is compact, there is a point $x \in \overline{Gy}$ such that \overline{Gx} is minimal. There is a net $t_n \in G$ with $t_n y \rightarrow x$. Since $(t_n y, y) \in R_o$, so $(x, y) \in R_o$ and $y \in \overline{Gx}$. Thus $\overline{Gy} = \overline{Gx}$ is minimal. \square

0.5. A subset A of G is ‘syndetic’ in G if there exists a compact subset K of G with $G = KA$. A set B is said to be ‘thick’ in G if for all compact set K in G there corresponds an element $t \in B$ such that $Kt \subseteq B$. It is a well-known fact that A is syndetic in G if and only if it intersects non-voidly with every thick subset of G .

- a) A point $x \in X$ is ‘almost periodic’ (a.p) under (G, X) if $N_G(x, U) = \{t \in G \mid tx \in U\}$ is a syndetic subset of G for all $U \in \mathfrak{N}_x$. Since \overline{Gx} is compact by (iii) here, $x \in X$ is a.p if and only if \overline{Gx} is minimal under (G, X) (cf., e.g. [1, Note 2.5.1]). Thus, $N_G(x, U)$ is discretely syndetic in G for every a.p point x of (G, X) .
- b) Following [5] we say that:
 - 1) (G, X) is ‘weakly a.p’ if given $\alpha \in \mathscr{U}$ there is a finite subset F of G such that $Ftx \cap \alpha[x] \neq \emptyset$ for all $x \in X$ and $t \in G$.

- 2) (G, X) is ‘locally weakly a.p.’ if given $\alpha \in \mathcal{U}$ and $x \in X$ there is a finite subset F of G and some $U \in \mathfrak{R}_x$ such that $Fty \cap \alpha[y] \neq \emptyset$ for all $y \in U$ and $t \in G$.

Clearly, if (G, X) is locally weakly a.p, it is pointwise a.p.

0.6. A Hausdorff space is said to be *zero dimensional* if it has a base consisting of clopen sets. For example, the coset spaces are locally compact zero-dimensional in a totally disconnected locally compact topological group.

Recall that G is *compactly generated*, provided that there exists a compact subset Γ of G , called a *generating set*, such that $G = \bigcup_{r=1}^{\infty} \Gamma^r$. In this case, G is σ -compact and we may assume $e \in \Gamma = \Gamma^{-1}$. In this case, the generating set is not unique; e.g., $t^{-1}\Gamma t$, for $t \in G$, is so.

0.7. In this note we consider the relationships of the following important dynamics of (G, X) :

- (1) The pointwise “recurrence” in certain sense (cf. Def. 2).
- (2) The pointwise almost periodicity.
- (3) X is a union of minimal sets.
- (4) R_o is closed.
- (5) $\mathcal{O}_{G,X}$ is ‘continuous’.
- (6) $\mathcal{O}_{G,X}$ is continuous.
- (7) The local weak almost periodicity.

It is evident that $(6) \Rightarrow (5) \Leftrightarrow (4) \Rightarrow (3) \Leftrightarrow (2) \Rightarrow (1)$. However, even for $G = \mathbb{Z}$ or \mathbb{R} and X a compact metric space, that “(1) \Rightarrow (5)” is not generally true is shown by very simple examples.

Note that if (G, X) is distal and $(G, X \times X)$ satisfies (5), then it is not difficult to prove that the regionally proximal relation of (G, X) is equal to Δ_X , the diagonal of $X \times X$, so that (G, X) is equicontinuous [3] (cf. Theorem 6c below). In McMahon and Wu [7], there is an example of a countable group G acting on a zero-dimensional compact X such that (G, X) is distal non-equicontinuous. So in this example, $(G, X \times X)$, with zero-dimensional compact phase space $X \times X$ and with countable (so σ -compact) phase group G , satisfies that (1) \nRightarrow (5) and (3) \nRightarrow (5). Thus we need impose some more restrictive condition on G for (1) or (3) \Rightarrow (5).

Let X be a zero-dimensional compact space. In [4, Theorems 5 and 6], Gottschalk shows (1) \Rightarrow (5) \Leftrightarrow (7) for \mathbb{Z}_+ -flow on X , and see also [1, Theorem 7.12] for \mathbb{Z} -flow on X .

Recall that G is *generative* provided that G is abelian and generated by some compact neighborhood of e (cf. [5, Def. 6.01]). Using “replete semigroup”, in [5] Gottschalk and Hedlund formulate a definition of recurrence. Then, with this notion, they prove that (1) \Rightarrow (5) \Leftrightarrow (7) for each flow with generative phase group and with zero-dimensional phase space (cf. [5, Theorems 7.07 and 7.08]).

In [2], Auslander et al. introduce a definition of recurrence in terms of “cone”. With their notion of recurrence, they show that (1) \Rightarrow (5) for every flow (G, X) with finitely generated phase group G not necessarily abelian and with zero-dimensional compact metric phase space X (cf. [2, Theorem 1.8]).

The purpose of this note is to formulate a recurrence and extend the theorem of Auslander et al. [2, Theorem 1.8] to any flow (G, X) with compactly generated phase group G not necessarily countable or abelian, and, with zero-dimensional phase space X not necessarily compact metrizable, as follows:

0.8 Theorem (see Thm. 4a). *Let (G, X) be a flow, where G is compactly generated and X is zero dimensional. Then the following conditions are pairwise equivalent:*

- (1) The pointwise “recurrence” in certain sense (cf. Def. 2b).
- (2) The pointwise almost periodicity.
- (3) X is a union of minimal sets.
- (4) R_o is closed.
- (5) $\mathcal{O}_{G,X}$ is “continuous”.
- (6) $\mathcal{O}_{G,X}$ is continuous.
- (7) For every compact open subset U of X , U_G^* is compact open in X .
- (8) (G, X) is locally weakly a.p.

Comparing with the case X compact, we have to face the essential point that a net in X need not have a convergent subnet. In addition, comparing with the case $G = \mathbb{Z}_+$, for any point x and any closed subset U of X we have no the “first visit time” $t \in G$ with $tx \in U$.

0.9 Remark. It should be mentioned that a part of Theorem 0.8 has been proved by Reid [9] under the framework that G is “equicontinuously generated” by a subset S of $\text{Homeo}(X)$; that is, $G = \bigcup_{r \geq 1} S^r$, where $\text{id}_X \in S = S^{-1}$, and S is equicontinuous (cf. Def. 6b). However, our approaches are different completely with and much simpler than Reid [9, Theorems 1.2, 1.3] of using very technical arguments of compact-open invariant sets. It turns out that Theorem 0.8 concisely proved here implies the “equicontinuously generated” case of Reid [9] (see Theorems 8e and 8f for alternative simple proofs of Reid’s results).

1 (Word length and cone). Let G be compactly generated with a generating set Γ such that $e \in \Gamma = \Gamma^{-1}$ as in Def. 0.6.

1a. For any $g \in G$ with $g \neq e$, the Γ -length of g , denoted $|g|$, is the smallest integer r such that $g = \gamma_1 \cdots \gamma_r$ with $\gamma_i \in \Gamma$ for $1 \leq i \leq r$, and, write $\mathbf{K}(g) = \Gamma^{|g|-1} \cdot g$, where $\Gamma^0 = \{e\}$ and $\mathbf{K}(g)$ is a compact subset of G . Clearly, $e \notin \mathbf{K}(g)$ for $g \in G$ with $|g| \geq 1$.

1b. A subset \mathbf{C} of G is called a Γ -cone in G if there exists a net $\{g_i\}$ in G such that $|g_i| \nearrow \infty$ and $\mathbf{K}(g_i) \rightarrow \mathbf{C}$ in 2^G . Here $|g_i| \nearrow \infty$ means that if $i \leq i'$ then $|g_i| \leq |g_{i'}|$. Clearly, every cone in G is a closed set in G . Moreover, if G is discrete (i.e. finitely generated), then $e \notin \mathbf{C}$.

1c Lemma (cf. [2, Prop. 1.2 and Prop. 1.5] for G finitely generated). *Let G be compactly generated with a generating set Γ . Then:*

- (1) *If $\{g_i\}$ is a net in G with $|g_i| \nearrow \infty$, then $\mathbf{K}(g_i) \rightarrow \mathbf{C} \neq \emptyset$.*
- (2) *If \mathbf{C} is a Γ -cone in G , then \mathbf{C} is a discretely thick subset of G .*

Proof. Let $\{g_i\}$ be a net in G with $|g_i| \nearrow \infty$ and $\mathbf{K}(g_i) \rightarrow \mathbf{C}$. Let K be a finite subset of G with $e \in K$. We shall find an element $t \in G$ with $Kt \subseteq \mathbf{C}$. Since G is compactly generated with the generating set Γ , there is an integer $r \geq 1$ such that $K \subseteq \Gamma^r$. For each i , let $\ell_i = |g_i|$ ($\nearrow \infty$) and write

$$g_i = \gamma_1 \cdots \gamma_{\ell_i-r-1} \gamma_{\ell_i-r} \cdots \gamma_{\ell_i} \in \Gamma^{\ell_i}, \quad \alpha_i = \gamma_{\ell_i-r} \cdots \gamma_{\ell_i} \in \Gamma^{r+1}.$$

Then $K\alpha_i \subseteq \mathbf{K}(g_i)$ for i with $\ell_i > r$. Since Γ^{r+1} is compact, we can assume (a subnet of) $\alpha_i \rightarrow t$ in G . Thus $t \in \mathbf{C}$ and $Kt \subseteq \mathbf{C} \neq \emptyset$ by Lemma 0.1. This also proves that \mathbf{C} is a discretely thick subset of G . The proof is completed. \square

From the choice of α_i in the above proof of Lemma 1c we can easily conclude the following lemma:

1d Lemma (cf. [9, Lem. 3.6]). *Let F be any finite subset of G . Then there is an integer $n > 1$ such that for all $g \in G$ with $|g| \geq n$, there is an element $t \in G$ such that $|t| = n$ and $Ft \subseteq K(g)$.*

2 (Recurrence). Let (G, X) be a flow with a compactly generated phase group G and $x \in X$. We will define recurrence at x in two ways.

2a. Let Γ be a generating set of G . We say that x is Γ -recurrent of type I under (G, X) if $Cx \cap U \neq \emptyset$ for all Γ -cone C in G and every $U \in \mathfrak{N}_x$. See [2, Def. 1.6] for G finitely generated and X a compact metric space in terms of sequence.

We say that x is *recurrent of type I* under (G, X) if x is Γ -recurrent of type I for all generating set Γ of G . Clearly, if x is recurrent of type I, then tx is recurrent of type I for every $t \in G$.

2b. Let Γ be a generating set of G . We say that x is Γ -recurrent of type II under (G, X) if for each $U \in \mathfrak{N}_x$ and every net $\{g_i\}$ in G with $|g_i| \rightarrow \infty$ there exists an integer $n \geq 1$ such that there is a subnet $\{g_{i_j}\}$ from $\{g_i\}$ and $c_j \in K(g_{i_j})$ with $c_j x \in U$ and $|c_j| \leq n$. See [9, Def. 1.1].

3 Lemma. *Let (G, X) be a flow, where G is compactly generated with a generating set Γ , and, let $x \in X$. Then:*

- (1) *If x is an a.p point, then x is Γ -recurrent of type II (cf. [9, Prop. 3.7(iv)]).*
- (2) *If x is Γ -recurrent of type II, then x is Γ -recurrent of type I.*

Proof. (1). Let $U \in \mathfrak{N}_x$ and simply write $A = N_G(x, U)$. Then A is a discretely syndetic subset of G by 0.5-a). So there is a finite set F in G such that $F^{-1}A = G$, or equivalently, $Ft \cap A \neq \emptyset$ for all $t \in G$. Let $n_1 = \max\{|f| + 1 \mid f \in F\}$. Let $\{g_i\}$ be a net in G with $|g_i| \rightarrow \infty$. By Lemma 1d, there exists an integer $n > n_1$ such that there is a subnet $\{g_{i_j}\}$ from $\{g_i\}$ such that for each j , there is an element $t_j \in G$ with $Ft_j \subseteq K(g_{i_j})$ and $|t_j| = n$. Take $f_j \in F$ with $c_j = f_j t_j \in A$, then $|c_j| \leq 2n$ and $c_j x \in U$ for all j . Thus x is Γ -recurrent of type II.

(2). Suppose x is Γ -recurrent of type II. Let C be a cone in G . That is, $K(g_i) \rightarrow C$, where $\{g_i\}$ is a net in G with $|g_i| \nearrow \infty$. Let $U, V \in \mathfrak{N}_x$ with $\overline{V} \subseteq U$. By Def. 2b, there is an integer $n \geq 1$ and a subnet $\{g_{i_j}\}$ from $\{g_i\}$ and $c_j \in K(g_{i_j})$ such that $c_j x \in V$ and $|c_j| \leq n$. Since Γ^n is compact and $c_j \in \Gamma^n$, we can assume (a subnet of) $c_j \rightarrow c$ in G and $c_j x \rightarrow cx \in \overline{V}$. Since $c \in C$ by definition and $Cx \cap U \neq \emptyset$, x is Γ -recurrent of type I. The proof is complete. \square

4. We are ready to state and concisely prove our main theorems of this note self-closely. Recall that $\mathcal{O}_{G,X}: X \rightarrow 2^X$ is defined by $x \mapsto \overline{Gx}$ and $U_G^* = \{x \in X \mid \overline{Gx} \subseteq U\}$ for all subset U of X .

4a Theorem. *Let (G, X) be any flow, where G is compactly generated with a generating set Γ and X is zero-dimensional. Then the following conditions are pairwise equivalent:*

- (1) *(G, X) is pointwise Γ -recurrent of type I.*
- (2) *(G, X) is pointwise Γ -recurrent of type II.*
- (3) *(G, X) is pointwise a.p.*
- (4) *X is a union of minimal sets.*
- (5) *The G -orbit closure relation R_o is closed.*
- (6) *$\mathcal{O}_{G,X}$ is \lceil continuous \rceil .*
- (7) *$\mathcal{O}_{G,X}$ is upper semi-continuous.*
- (8) *Given any compact-open subset U of X , U_G^* is compact open.*
- (9) *(G, X) is locally weakly a.p.*

Proof. (2) \Rightarrow (1): By (2) of Lemma 3.

(3) \Rightarrow (2): By (1) of Lemma 3.

(4) \Rightarrow (3): By a) of 0.5.

(5) \Rightarrow (4): By Lemma 0.4.

(6) \Rightarrow (5): By a) of 0.3.

(7) \Rightarrow (6): By Def. 0.2.

(8) \Rightarrow (7): Assume (8). Let $x \in X$ and V an open neighborhood of \overline{Gx} . Because X is locally compact zero-dimensional and \overline{Gx} is compact, there is a compact-open set U with $\overline{Gx} \subseteq U \subseteq V$. Then $x \in \overline{Gx} \subseteq U_G^* \subseteq V$. Since U_G^* is open, $\mathcal{O}_{G,X}$ is upper semi-continuous. Thus (8) \Rightarrow (7) holds.

(1) \Rightarrow (8): Assume (1). Let U be a compact-open subset of X . Then U_G^* is an G -invariant compact set. To prove that U_G^* is open, suppose the contrary that U_G^* is not open. Then $U_G^* \neq \emptyset$ and there is a net $x_i \in U \setminus U_G^*$ and a point $x \in U_G^*$ with $x_i \rightarrow x$. We can obviously take $g_i \in G$ such that $g_i x_i \notin U$ and $|g_i| = \min\{|t| : t \in G, tx \notin U\}$. Since U is clopen and $\overline{Gx} \subseteq U_G^*$, obviously (a subnet of) $|g_i| \rightarrow \infty$. For all i we write $g_i = \gamma_{i,1} \gamma_{i,2} \cdots \gamma_{i,|g_i|}$ and $g'_i = \gamma_{i,2} \cdots \gamma_{i,|g_i|}$. Then $g'_i x_i \in U$ and $\gamma_{i,1} \in \Gamma$ for all i . Since U and Γ are compact, we can assume (a subnet of) $\gamma_{i,1} \rightarrow \gamma$ and $g'_i x_i \rightarrow y' \in U$. Let $y = \gamma y'$. Then $y = \lim_i g_i x_i \notin U$. Let $K(g_i^{-1}) \rightarrow C$, a Γ -cone in G . Given $c \in C$, we can take a subnet $\{g_{i_j}\}$ from $\{g_i\}$ and $c_j = k_j g_{i_j}^{-1}$ with $k_j \in \Gamma^{|g_{i_j}|-1}$ and $c_j \rightarrow c$. Thus by $|k_j| < |g_{i_j}|$, it follows that $c y = \lim_j c_j y = \lim_j (k_j g_{i_j}^{-1})(g_{i_j} x_{i_j}) \in U$. So $Cy \cap (X \setminus U) = \emptyset$, contrary to that y is Γ -recurrent of type I by (1). Thus (1) implies (8).

(9) \Rightarrow (3): By b) of 0.5.

(7) \Rightarrow (9): Let $\alpha \in \mathcal{U}$ and $x \in X$. Let $V \in \mathfrak{N}_x$ be so small that $\alpha[y] \supseteq V$ for all $y \in V$. Since \overline{Gx} is compact minimal, there exists a finite set $\{z_1, \dots, z_k\} \subset X$ with $W_i \in \mathfrak{N}_{z_i}$, $t_i \in G$ such that $t_i W_i \subseteq V$ and $\overline{Gx} \subseteq \bigcup_{i=1}^k W_i$. By (7), there is a $U \in \mathfrak{N}_x$ with $U \subseteq V$ such that $\overline{Gy} \subseteq \bigcup_{i=1}^k W_i$ for all $y \in U$. Then for $F = \{t_1, \dots, t_k\}$, $Fty \cap \alpha[y] \neq \emptyset$ for all $y \in U$ and $t \in T$. Thus (7) implies (9).

The proof of Theorem 4a is thus completed. \square

4b Theorem. Let (G, X) be any flow with G compactly generated with a generating set Γ and with X zero-dimensional, where \overline{Gx} need not be compact for $x \in X$. Let V be an open subset of X and write $V_\infty = \bigcap_{g \in G} gV$. Suppose V_∞ is compact. Then V_∞ is open if and only if $\overline{Gx} \cap V_\infty = \emptyset$ for every $x \in X \setminus V_\infty$.

Proof. Necessity is obvious for $X \setminus V_\infty$ is closed G -invariant. To show sufficiency, suppose the contrary that V_∞ is not open (so $V_\infty \neq \emptyset$). Then there is a net $x_i \in V \setminus V_\infty$ with $x_i \rightarrow x \in V_\infty$. Since X is a locally compact zero-dimensional Hausdorff space, we can find a compact-open neighborhood U of V_∞ with $U \subseteq V$. Clearly, $V_\infty = \bigcap_{g \in G} gU$. We may assume $\{x_i\} \subset U$. Take $g_i \in G$ such that $g_i x_i \notin U$ and $|g_i| = \min\{|t| : t \in G, tx_i \notin U\}$. Since V_∞ is compact G -invariant, $|g_i| \rightarrow \infty$. By compactness of U , we can assume (a subnet of) $g_i x_i \rightarrow y \notin U$ (cf. Proof of (1) \Rightarrow (8) in Theorem 4a). Let $K(g_i^{-1}) \rightarrow C$. Then $Cy \subseteq U$ (cf. Proof of (1) \Rightarrow (8) in Theorem 4a). Next we shall prove that \overline{Cy} contains an G -orbit so $\overline{Cy} \cap V_\infty \neq \emptyset$, a contradiction.

Indeed, let \mathcal{F} be the collection of finite subsets of G with a partial order by inclusion. Since C is discretely thick in G by Lemma 1c, we have for $F \in \mathcal{F}$ that there exists an element $t_F \in C$ with $Ft_F \subseteq C$. Since 2^U is a compact Hausdorff space and $t_F y \in U$ and $Ft_F y \in 2^U$, we can assume (a subnet of) $Ft_F y \rightarrow Q \subseteq \overline{Cy}$ and $t_F y \rightarrow y'$. Given $t \in G$ there is some $F_0 \in \mathcal{F}$ with $t \in F_0$. Then $t \in F$ for all $F \geq F_0$ so that $ty' = \lim_F tt_F y \in \lim_F Ft_F y$. Thus $ty' \in \overline{Cy}$ for all $t \in G$. Then $\overline{Gy'} \subseteq \overline{Cy}$. The proof is completed. \square

5 Remark. It should be noticed that since we do not know whether the Γ -recurrence of type I at a point of X is equivalent to that of type II, so (1) \Leftrightarrow (2) in Theorem 4a is not trivial. In addition, Theorem 4b can be utilized for proving (5) \Rightarrow (8) in Theorem 4a.

6. Let (G, X) be a flow.

6a. We say (G, X) is distal if we have for $x, y \in X$ with $x \neq y$ that $\Delta_X \cap \overline{G(x, y)} = \emptyset$.

6b. We say (G, X) is equicontinuous if given $\varepsilon \in \mathcal{U}$ and $x \in X$ there exists an index $\delta \in \mathcal{U}$ such that $g\delta[x] \subseteq \varepsilon[gx]$ for all $g \in G$. If δ is independent of $x \in X$, then we say (G, X) is uniformly equicontinuous in this case.

Using Theorem 4a we can then conclude the following generalization of an important theorem of R. Ellis:

6c Theorem (cf. [3, Thm. 2] for G generative and [2, Cor. 1.9] for G finitely generated with X compact). *Let (G, X) be a distal flow with X zero-dimensional and G compactly generated. Then (G, X) is equicontinuous.*

Proof. Since (G, X) is distal, $(G, X \times X)$ is pointwise a.p. To show (G, X) equicontinuous, suppose the contrary that (G, X) is not equicontinuous; then there is a point $x \in X$, an index $\varepsilon \in \mathcal{U}$, a net $x_n \rightarrow x$ and a net $t_n \in G$ such that $(t_n x, t_n x_n) \notin \varepsilon$. By (7) of Theorem 4a, we can assume $t_n x \rightarrow x_*$ and $t_n x_n \rightarrow y_*$ with $(x_*, y_*) \notin \varepsilon$. Let $s_n = t_n^{-1}$ and $x'_n = t_n x, y'_n = t_n x_n$. Then $(s_n x'_n, s_n y'_n) \rightarrow (x, x)$ and $(x'_n, y'_n) \rightarrow (x_*, y_*)$. Using (7) of Theorem 4a for $(G, X \times X)$, it follows that $x_* = y_*$, contrary to $(x_*, y_*) \notin \varepsilon$. The proof is complete. \square

7 (The case G to be finitely generated [2]). Let G be finitely generated with a generator set Γ such that $\Gamma = \Gamma^{-1}$, and let (G, X) be a flow. We shall show that our Γ -recurrence of type I (Def. 2a) is consistent with [2] in this case. See Lemma 7d below.

7a. Let $B_r = \{g \in G \mid g \neq e, |g| \leq r\}$, for all $r \geq 1$. For $g \in G$ with $g \neq e$, write $K(g) = B_{|g|-1} \cdot g$.

7b. A subset C of G is an AGW-cone, namely, a cone in the sense of Auslander, Glasner and Weiss [2], if there is a sequence $g_n \in G$ with $|g_n| \rightarrow \infty$ such that for each $r \geq 1$ there exists n_r such that $B_r \cap K(g_n)$ is independent of n for all $n \geq n_r$, and, $C = \lim_{n \rightarrow \infty} K(g_n)$. Since G is discrete here, so $c \in C$ iff $c \in K(g_n)$ as n sufficiently large. Moreover, $e \notin C$.

7c (cf. [2, Def. 1.6]). We say that $x \in X$ is AGW-recurrent, if $Cx \cap U \neq \emptyset$ for every $U \in \mathfrak{R}_x$ and every AGW-cone C in G .

7d Lemma. *Let X be a compact metric space and $x \in X$. Then x is AGW-recurrent under (G, X) if and only if x is Γ -recurrent of type I in the sense of Def. 2a.*

Proof. 1). Let C be a Γ -cone in G with $K(g_i) = \Gamma^{|g_i|-1} \cdot g_i \rightarrow C$, where $\{g_i\}$ is a net in G such that $|g_i| \nearrow \infty$. For all integer $k \geq 1$, there is an $i_k \in \{i\}$ such that $|g_i| \geq k$ for all $i \geq i_k$ and $|g_{i_k}| < |g_{i_{k+1}}|$. Clearly $|g_{i_k}| \rightarrow \infty$ as $k \rightarrow \infty$. Moreover, the sequence $\{g_{i_k}\}_{k=1}^\infty$ is a subnet of $\{g_i\}$. Indeed, for $i' \in \{i\}$, there is an integer $k \geq 1$ with $|g_{i'}| < k \leq |g_{i_{k'}}|$ for all $k' > k$ so $i' \leq i_{k'}$ for all $k' > k$. Therefore, using $\{i_k\}$ in place of $\{i\}$ if necessary, we can assume $\{g_i\}_{i=1}^\infty$ is a sequence in G with $|g_i| \nearrow \infty$ as $i \rightarrow \infty$.

Each B_r is finite, so we may choose a subsequence $\{i_k\}$ from $\{i\}$ so that each $K(g_{i_k}) \cap B_r$ is eventually constant. Then by a diagonal process we can choose a subsequence (and relabel)

$\{i_k\}_{k=1}^\infty$ from $\{i\}$ such that $K(g_{i_k}) \rightarrow C$ as $k \rightarrow \infty$. Then C is an AGW-cone in G such that $C \subseteq \mathcal{C}$ by Lemma 0.1.

2). Let C be a cone in G with $C = \lim_i K(g_i)$ in the sense of Def. 7b, where $\{g_i\}_{i=1}^\infty$ is a sequence in G such that $|g_i| \rightarrow \infty$ and for each $r \geq 1$ there exists an $n_r \geq 1$ such that $B_r \cap K(g_i)$ is independent of i for all $i \geq n_r$. By induction, we may require $n_r < n_{r+1}$ and $|g_{n_r}| < |g_{n_{r+1}}|$ for $r \geq 1$. Then

$$C = \bigcup_{r=1}^\infty (B_r \cap K(g_{n_r})).$$

Let $K(g_{n_r}) \rightarrow C$ in the sense of Def. 1b. Let $c \in C$. Then there exists a subnet $\{n_{r'}\}$ from $\{n_r\}$ and $k_{n_{r'}} \in K(g_{n_{r'}})$ such that $k_{n_{r'}} \rightarrow c$. Since G is discrete and $\{c\}$ is an open neighborhood of c in G , we can assume $k_{n_{r'}} = c$ for all r' . Since we have for all n_r that $n_{r'} \geq n_r$ as r' sufficiently large, so $c \in C$ and $C \subseteq C$.

The above discussion of 1) and 2) implies that x is AGW-recurrent iff x is also Γ -recurrent of type I. The proof is complete. \square

7e Theorem (cf. [2, Thm. 1.8]). *Let (G, X) be a flow, where G is finitely generated and X is a zero-dimensional compact metric space. Then the following conditions are equivalent:*

- (i) (G, X) is pointwise AGW-recurrent.
- (ii) (G, X) is pointwise a.p.
- (iii) The G -orbit closure relation is closed.

Proof. In view of Lemma 7d, this is a special case of Theorem 4a. \square

7f Proposition. *Let $f: X \rightarrow X$ be a homeomorphism of X , which is thought of as a \mathbb{Z} -flow. Then $x \in X$ is recurrent of type I (cf. Def. 2a) iff there exists a net $\{i_n\}$ in \mathbb{Z} with $i_n \rightarrow \infty$ and a net $\{j_n\}$ in \mathbb{Z} with $j_n \rightarrow -\infty$ such that $f^{i_n}x \rightarrow x$ and $f^{j_n}x \rightarrow x$ simultaneously.*

Proof. Note that a Γ -cone C in \mathbb{Z} with $\Gamma = \{-1, 0, 1\}$ is either $C = \mathbb{N}$ or $C = -\mathbb{N}$, where \mathbb{N} is the set of positive integers. Then the statement follows easily from this fact. \square

7g (Regular almost periodicity [5]). A point x is said to be *regularly a.p* under (G, X) if $N_G(x, U)$ contains a syndetic subgroup of G for every $U \in \mathfrak{R}_x$. We say (G, X) is *regularly a.p* if every $\alpha \in \mathcal{U}$ there is a syndetic subgroup A of G such that $Ax \subseteq \alpha[x]$ for all $x \in X$.

7h Proposition (cf. [5, Thm. 7.10] for G abelian finitely generated). *Let (G, X) be a flow with G finitely generated and with X zero-dimensional compact. If (G, X) is pointwise regularly a.p, then (G, X) is regularly a.p.*

Proof. By [5, Lemma 5.04], (G, X) is distal. Further by Corollary 6c, it follows that (G, X) is equicontinuous. Finally by [5, Remark 5.02, Theorem 5.17], we see that (G, X) is regularly a.p. The proof is complete. \square

7i Proposition (cf. [5, Thm. 7.11] for G abelian finitely generated). *Let (G, X) be a flow with G finitely generated and with X a compact metric space. Then (G, X) is regularly a.p if and only if (G, X) is pointwise regularly a.p and weakly a.p.*

Proof. The proof of [5, Thm. 7.11] is still valid for this case using Theorem 4a in place of [5, Theorem 7.08]. We omit the details here. \square

8 (The case G to be equicontinuously generated). Let X be a locally compact Hausdorff space, $C_u(X, X)$ the space of continuous mappings from X to itself equipped with the topology of uniform convergence, $\text{Homeo}(X)$ the group of self-homeomorphisms of X onto itself as a subspace of $C_u(X, X)$. Clearly, $\text{Homeo}(X)$ is a Hausdorff semi-topological group so that $(\text{Homeo}(X), X)$ defined by $\text{Homeo}(X) \times X \xrightarrow{(f, x) \mapsto fx} X$ is a flow.

In the sequel let $S \subset \text{Homeo}(X)$ such that: 1) $e = \text{id}_X \in S = S^{-1}$ and 2) S is equicontinuous on X (cf. Def. 6b).

8a. Set $\langle S \rangle = \bigcup_{r=1}^{\infty} S^r$. Clearly $\langle S \rangle$ is a subgroup of $\text{Homeo}(X)$, which is said to be *equicontinuously generated with the generating set S* .

8b. Write $\Gamma = \text{cls}_u S$, where cls_u denotes the closure relative to $C_u(X, X)$. Then:

Lemma. Γ is a compact subset of $C_u(X, X)$ and $\Gamma \subset \text{Homeo}(X)$ such that $e \in \Gamma = \Gamma^{-1}$.

Proof. Since S is equicontinuous, so is Γ . Then Γ is compact in $C_u(X, X)$ by Ascoli's Theorem (cf. [6, Theorem 7.17]). Moreover, Γx is compact in X . We need prove that $\Gamma \subset \text{Homeo}(X)$. For this, let $S \ni f_n \rightarrow f \in \Gamma$.

Let $x \neq y$ and $\varepsilon \in \mathcal{U}$ with $(x, y) \notin \varepsilon$. If $fx = fy = z$, then for $\delta \in \mathcal{U}$ we have $x, y \in f_n^{-1}\delta[z]$ as n sufficiently large, contrary to equicontinuity of S . Thus f is injective. Let $x \in X$. Since $f_n X = X$, there is a point $x_n \in X$ with $f_n x_n = x$ or $x_n = f_n^{-1}x$. We can assume (a subnet of) $x_n \rightarrow x'$. Then $fx' = x$ so $fX = X$. Thus f is a continuous bijection.

Let (a subnet of) $f_n^{-1} \in S \rightarrow g \in \Gamma$. Since $e = f_n f_n^{-1} \rightarrow fg$, so $fg = e$, $g = f^{-1} \in \Gamma$, and $f \in \text{Homeo}(X)$. The proof is complete. \square

8c. Set $\langle \Gamma \rangle = \bigcup_{r=1}^{\infty} \Gamma^r$. Then $\langle \Gamma \rangle$ is a compactly generated subgroup of $\text{Homeo}(X)$ with a generating set Γ . Moreover, $(\langle \Gamma \rangle, X)$ is a flow. It is evident that S^r is dense in Γ^r for all $r \geq 1$ and $\langle S \rangle$ is dense in $\langle \Gamma \rangle$ under the topology of uniform convergence.

Following Def. 2b, $x \in X$ is S -recurrent of type II under $(\langle S \rangle, X)$ if for every $U \in \mathfrak{R}_x$ and every net $\{g_i\}$ in $\langle S \rangle$ with $|g_i| \rightarrow \infty$, there is an integer n , a subnet $\{g_{i_j}\}$ from $\{g_i\}$ and $c_j \in S^{|g_{i_j}|-1} \cdot g_{i_j}$ such that $c_j x \in U$ and $|c_j| \leq n$. We do not know if a point S -recurrent of type II is Γ -recurrent of type II, yet it is Γ -recurrent of type I by Lemma 8d below.

8d Proposition. *If x is S -recurrent of type II under $(\langle S \rangle, X)$, then x is Γ -recurrent of type I under $(\langle \Gamma \rangle, X)$.*

Proof. Let $\{g_i\}$ be a net in $\langle \Gamma \rangle$ with $|g_i| \nearrow \infty$ and $K(g_i) \rightarrow C$. Let $U, V \in \mathfrak{R}_x$ with $\overline{V} \subseteq U$. We need prove that $Cx \cap U \neq \emptyset$. Since $\langle S \rangle$ is dense in $\langle \Gamma \rangle$, we can assume $g_i \in \langle S \rangle$ for all i . Further by definition, there is an integer $n \geq 1$, a subnet $\{g_{i_j}\}$ from $\{g_i\}$ and $c_j \in S^{|g_{i_j}|-1} \cdot g_{i_j} \subseteq K(g_{i_j})$ such that $c_j x \in V$ and $|c_j| \leq n$. By $c_j \in \Gamma^n$ and Γ^n is compact, it follows that (a subnet of) $c_j \rightarrow c \in C$ and $cx \in U$. The proof is complete. \square

8e Theorem (cf. [8, Thm. 2.2], [9, Thm. 1.2] using compact-open invariant sets). *Let X be zero-dimensional, let G be an equicontinuously generated group. Let V be an open subset of X and $V_{\infty} = \bigcap_{g \in G} gV$. Suppose V_{∞} is compact. Then V_{∞} is open if and only if $\overline{Gx} \cap V_{\infty} = \emptyset$ for every $x \in X \setminus V_{\infty}$.*

Proof. Let $G = \langle S \rangle$, where S is the equicontinuously generating set of G . Then $\overline{Gx} = \overline{\langle \Gamma \rangle x}$ for all $x \in X$. Since V_{∞} is compact closed, so $V_{\infty} = \bigcap_{g \in \langle \Gamma \rangle} gV$. Then Theorem 8e follows from Theorem 4b. \square

8f Theorem (cf. [9, Thm. 1.3] using Thm. 8e). *Let X be zero-dimensional and $G = \langle S \rangle$ such that \overline{Gx} is compact for all $x \in X$. Then the following are pairwise equivalent:*

- i) (G, X) is pointwise S -recurrent of type II.
- ii) (G, X) is pointwise a.p.
- iii) Given any compact open subset V of X , V_G^* is open.
- iv) The orbit closure relation R_o of (G, X) is closed.
- v) There exists an extension $\rho: (G, X) \rightarrow (G, Y)$ such that:
 - 1) Y is locally compact zero-dimensional,
 - 2) $gy = y$ for all $y \in Y$ and $g \in G$, and
 - 3) $\rho^{-1}y$ is a minimal subset of (G, X) for each $y \in Y$.

Proof. By Lemma 8d, $(\langle \Gamma \rangle, X)$ is pointwise Γ -recurrent of type I. Since $\overline{Gx} = \overline{\langle \Gamma \rangle x}$ for all $x \in X$, x is a.p. under (G, X) iff x is a.p. under $(\langle \Gamma \rangle, X)$. Then i) \Leftrightarrow ii) \Leftrightarrow iii) \Leftrightarrow iv) \Leftrightarrow iv') by Theorem 4a, where

iv') $\mathcal{O}_{G,X}: X \rightarrow 2^X$ is upper semi-continuous.

v) \Rightarrow ii): This is evident by 3) of v).

iv') \Rightarrow v): Assume iv'). Define $Y = X/R_o$ and let $\rho: X \rightarrow Y$ be the canonical quotient mapping. Then ρ is closed (cf. [6, Thm. 3.12]), and moreover, by iii) it is easy to see that Y is a Hausdorff space. Let $y \in Y$ and $U \in \mathfrak{N}_y$. Set $V = \rho^{-1}U$ and $x \in \rho^{-1}y$. Then $\overline{Gx} = \rho^{-1}y \subseteq V$. Further, there exists a compact open neighborhood W of \overline{Gx} . By iii), W_G^* is compact open ρ -saturated with $x \in W_G^*$. Thus $\rho W_G^* \subseteq U$ is a clopen compact neighborhood of y in Y . This shows that Y is locally compact zero-dimensional. Thus (G, Y) is a flow having the properties 1), 2) and 3).

The proof of Theorem 8f is thus completed. \square

9 Remark. The Γ -recurrence of type I (Def. 2) is conceptually dependent of the generating set Γ of G . So it should be interested to generalize Proposition 7f from $G = \mathbb{Z}$ to a more general case.

Acknowledgments. This work was supported by National Natural Science Foundation of China (Grant No. 11790274) and PAPD of Jiangsu Higher Education Institutions.

References

- [1] J. Auslander and X. Dai, *Minimality, distality and equicontinuity for semigroup actions on compact Hausdorff spaces*, Discrete Contin. Dyn. Syst., 39 (2019), 4647–4711.
- [2] J. Auslander, E. Glasner and B. Weiss, *On recurrence in zero dimensional flows*, Forum Math., 19 (2007), 107–114.
- [3] R. Ellis, *Distal transformation groups*, Pacific J. Math., 8 (1958), 401–405.
- [4] W. H. Gottschalk, *Almost periodic points with respect to transformation semi-groups*, Annals of Math., 47 (1946), 762–766.
- [5] W. H. Gottschalk and G. A. Hedlund, *Topological Dynamics*, Amer. Math. Soc. Coll. Publ. Vol. 36, Amer. Math. Soc., Providence, R.I., 1955.
- [6] J. L. Kelley, *General Topology*. GTM 27, Springer-Verlag, New York Berlin Heidelberg, 1955.
- [7] D. C. McMahon and T. S. Wu, *On the connectedness of homomorphisms in topological dynamics*, Trans. Amer. Math. Soc., 217 (1976), 257–270.
- [8] C. D. Reid, *Distal actions on coset spaces in totally disconnected locally compact groups*, J. Topology Anal., 12 (2020), 491–532.
- [9] C. D. Reid, *Equicontinuity, orbit closures and invariant compact open sets for group actions on zero-dimensional spaces*, Groups Geom. Dyn., 14 (2020), 413–425.