

# THERMODYNAMIC FORMALISM AND LARGE DEVIATION PRINCIPLE OF MULTIPLICATIVE ISING MODELS

JUNG-CHAO BAN, WEN-GUEI HU, AND GUAN-YU LAI

**ABSTRACT.** The aim of this study is three-fold. First, we investigate the thermodynamics of the Ising models with respect to 2-multiple Hamiltonians. This extends the previous results of [Chazotte and Redig, Electron. J. Probably., 2014] to  $\mathbb{N}^d$ . Second, we establish the large deviation principle (LDP) of the average  $\frac{1}{N}S_N^G$ , where  $S_N^G$  is a 2-multiple sum along a semigroup generated by  $k$  numbers which are  $k$  co-primes. This extends the previous results [Ban et al. Indag. Math., 2021] to a board class of the long-range interactions. Finally, the results described above are generalized to the multidimensional lattice  $\mathbb{N}^d, d \geq 1$ .

## 1. INTRODUCTION

In this article, we investigate the thermodynamic formalism (e.g., Gibbs measures, entropy and pressure functions) and the large deviation principle of the multiplicative Ising models on  $\mathbb{N}^d, d \geq 1$ . Before presenting the main results, we provide the motives behind this study. Consider the lattice spin systems with Ising  $\pm 1$  spins on  $\mathbb{N}$ . The Hamiltonian of the standard Ising model on each configuration  $\sigma \in \{-1, +1\}^{\mathbb{N}}$  is

$$H(\sigma) = -\beta \left( \sum_{i \in \mathbb{N}} J \sigma_i \sigma_{i+1} + h \sum_{i \in \mathbb{N}} \sigma_i \right),$$

where the parameter  $\beta$  is the inverse temperature,  $J$  is the coupling strength and  $h$  stands for the magnetic field. Note that the Hamiltonian  $H(\sigma)$  is a nearest neighbor translation-invariant interaction. The thermodynamics of the Ising models, e.g., the existence of the Gibbs measure, phase transition problem, entropy and pressure functions have been studied in depth. The best reference on this topic is [16].

Motivated from multiple ergodic theory (we refer the reader to [11, 19] for detailed definitions and recent advances in multiple ergodic theory) and the previous works of Kifer [20], Kifer and Varadhan [21], Fan, Liao and Ma [13] on the non-conventional averages, Carinci et al. [7] and Chazottes and Redig [8] investigate the ‘multiplicative Ising models’. That is, they consider the Hamiltonian on each

---

*Key words and phrases.* Gibbs measures, multiplicative shift, large deviation principle, multiple sum, free energy function.

Ban is partially supported by the Ministry of Science and Technology, ROC (Contract MOST 109-2115-M-004-002-MY2 and 108-2115-M-004-003). Hu is partially supported by the National Natural Science Foundation of China (Grant 11601355).

configuration  $\sigma \in \{-1, +1\}^{\mathbb{N}}$  as

$$H^{[m]}(\sigma) = -\beta \left( \sum_{i \in \mathbb{N}} J \sigma_i \sigma_{2i} + h \sum_{i \in \mathbb{N}} \sigma_i \right).$$

It should be noted that the Hamiltonian  $H^{[m]}(\sigma)$  is a long-range non-translation invariant interaction. For  $h = 0$ , the existence and uniqueness of the Gibbs measure with respect to  $H^{[m]}(\sigma)$  is constructed [8], which is multiplication invariant.

**Theorem 1.1** (Theorem 3.2 [8]). *Let*

$$\mu_N(\sigma_{\mathcal{N}_{2N}}) = \frac{e^{-H_N(\sigma_{\mathcal{N}_{2N}})}}{\sum_{\sigma_i = \pm 1, i \in \mathcal{N}_{2N}} e^{-H_N(\sigma_{\mathcal{N}_{2N}})}} 1$$

*be the finite-volume probability measure corresponding to the Hamiltonian*

$$H_N(\sigma_{\mathcal{N}_{2N}}) = -\beta \sum_{i=1}^N \sigma_i \sigma_{2i}.$$

*Then*

1. *Unique limit measure : The measure  $\mu_N$  have a unique weak limit denoted by  $\mu_\infty$  which is the Gibbs measure.*
2. *Independent Ising layers : Under  $\mu_\infty$ , the  $\{\tau_\ell^i = \sigma_{i2^\ell}\}_{\ell=0}^\infty, i \in 2\mathbb{N}_0 + 1$  are independent and distributed according to the standard Ising model measure  $\mu_\infty^{\text{Ising}}$  with a free boundary condition on the left.*
3. *Multiplication invariance : The measure  $\mu_\infty$  is multiplication invariant, i.e., for all  $i \in \mathbb{N}$ ,  $\sigma = (\sigma_j)_{j=1}^\infty$  and  $T_i \sigma = (\sigma_{ij})_{j=1}^\infty$  have the same distribution.*

Carinci et al. [7] established the large deviation principle of the average  $\frac{1}{N} S_N(\sigma) := \frac{1}{N} \sum_{i=1}^N \sigma_i \sigma_{2i}$  by calculating the rigorous formula of the free energy function

$$F_r(\beta) = F_{\mathbb{P}_r}(\beta) = \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{E}_{\mathbb{P}_r} (e^{\beta S_N}) ,$$

where  $\mathbb{P}_r$  is the product of Bernoulli measures with the parameter  $r$  on two symbols  $\{-1, +1\}$ .

**Theorem 1.2** (Theorem 4.1 [7]). *The explicit expression of the free energy function associated to the multiple sum  $S_N$  is*

$$F_r(\beta) = \log \left( (r(1-r))^{\frac{3}{4}} |v^T \cdot e_+| \Lambda_+ \right) + \mathcal{G}(\beta),$$

*where*

$$\mathcal{G}(\beta) = \frac{1}{2} \sum_{\ell=1}^{\infty} \frac{1}{2^\ell} \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^\ell \right)$$

*and  $\Lambda_\pm = e^\beta \left( \cosh(h) \pm \sqrt{\sinh^2(h) + e^{-4\beta}} \right)$ ,  $v^T = (e^{h/2}, e^{-h/2})$ ,  $h = \frac{1}{2} \log(r/(1-r))$ ,  $e_+ = \frac{w_+}{\|w_+\|}$  with  $w_+ = (-e^{-\beta}, e^{h+\beta} - \Lambda_+)^T$ .*

---

<sup>1</sup> $\mathcal{N}_{2N} = \{i \in \mathbb{N} : i \leq 2N\}$

Theorem 1.2 has been recently generalized to multidimensional lattice  $\mathbb{N}^d$ ,  $d \geq 1$  [2].

**Theorem 1.3** (Theorem 3.2 [2]). *For any  $d \geq 1$  and  $\mathbf{p} = (p_1, \dots, p_d) \in \mathbb{N}^d$ , the following statements hold true.*

1. *The explicit expression of the free energy function associated to the multiple sum<sup>2</sup>*

$$S_{N_1 \times \dots \times N_d}^{\mathbf{p}} = \frac{1}{N_1 \dots N_d} \sum_{\mathbf{i} \in \mathcal{N}_{N_1 \times \dots \times N_d}} \sigma_{\mathbf{i}} \sigma_{\mathbf{i} \cdot \mathbf{p}}$$

is

$$F_r(\beta) = \frac{2p_1 \dots p_d - 1}{2p_1 \dots p_d} \log(r(1-r)) + \frac{p_1 \dots p_d - 1}{p_1 \dots p_d} \log |v^T \cdot e_+|^2 + \log \Lambda_+ + \mathcal{G}(\beta),$$

where  $\Lambda_{\pm}$ ,  $v^T$ ,  $h$ ,  $e_+$  are defined as above and

$$\mathcal{G}(\beta) = \sum_{\ell=1}^{\infty} \frac{(p_1 \dots p_d - 1)^2}{(p_1 \dots p_d)^{\ell+1}} \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^{\ell} \right).$$

2. *The function  $F_r(\beta)$  is differentiable with respect to  $\beta \in \mathbb{R}$ .*
3. *The multiple average satisfies a LDP with rate function given by*

$$I_r(x) = \sup_{\beta \in \mathbb{R}} (\beta x - F_r(\beta)).$$

Furthermore, if  $(F_r)'(\eta) = y$ , then  $I_r(y) = \eta y - F_r(\eta)$ .

The aim of this paper is to study the multiplicative Ising models on  $\mathbb{N}^d$  and the objective is three-fold.

1. The thermodynamics of the Ising model with respect to 2-multiple Hamiltonian: We establish the thermodynamics formalism of the multiplicative Ising models on multidimensional lattice  $\mathbb{N}^d$  with respect to the Hamiltonian  $\sum_{i \in \mathbb{N}} \sigma_i \sigma_{2i}$  (call it *2-multiple Hamiltonian*). Theorem 3.3 extends Theorem 1.1 to  $\mathbb{N}^d$  and the formula of the Kolmogorov-Sinai (KS) entropy with respect to the limiting measure constructed in Theorem 3.3 is presented in Theorem 3.9.

2. The large deviation principle of the average  $\frac{1}{N} S_N^G$ : Let  $k \geq 1$  and  $p_1, \dots, p_k$  be co-primes, and  $G = \langle p_1, \dots, p_k \rangle = \{1 = \ell_1 < \ell_2 < \dots\}$  be a semigroup generated by  $p_1, \dots, p_k$ . The rigorous formula for the free energy function and the large deviation principle with respect to the average  $S_N^G(\sigma) := \sum_{i=1}^N \sigma_{\ell_j(i)} \sigma_{\ell_j(i)+1}$  on  $\mathbb{N}$  is given in Theorem 4.1. Clearly, 2-multiple Hamiltonian is a special case of the  $S_N^G$  with  $G = \langle 2 \rangle$ . Therefore, Theorem 4.1 can be seen as a generalization of Theorem 1.2 to a board class of the long-range non-translation invariant interactions.

3. The  $\mathbb{N}^d$  version of the aforementioned results: The  $\mathbb{N}^d$  version of Theorem 4.1 (or the generalization of Theorem 1.3) are presented in Theorems 4.4 and 4.5. We remark that Theorem 4.4 provides a general formula for  $F_r(\beta)$ . Meanwhile, Theorem 4.5 presents a explicit formula for  $F_r(\beta)$  if the sum of  $F_r(\beta)$  is along some specific direction. In this circumstance, we call  $F_r(\beta)$  the **directional free energy**

---

<sup>2</sup> $\mathbf{i} \cdot \mathbf{p} = (i_1 p_1, \dots, i_d p_d)$  and  $\mathcal{N}_{N_1 \times \dots \times N_d} = \{\mathbf{i} = (i_1, \dots, i_d) \in \mathbb{N}^d : i_j \leq N_j, \forall 1 \leq j \leq d\}$

**function** which is defined in Section 4.2. We emphasize that the calculation of the rigorous formula for  $F_r(\beta)$  along other directions is extremely difficult since the independent sublattices according to the constrains  $\mathbf{G}$  are quite hard to analysis. The notation  $\mathbf{G}$  represents the semigroup generated by the vectors  $\mathbf{p}_1, \dots, \mathbf{p}_k \in \mathbb{N}^d$ , the formal definition can be found in Section 2. And the generalizations of Theorems 3.3 and 3.9 are given in Theorems 3.6 and 3.11 respectively.

We remark that Peres et al. [22] computed the Hausdorff and Minkowski dimensions of the set

$$X_\Omega^G = \{(x_k)_{k=1}^\infty \in \Sigma_m : x|_{iG} \in \Omega \text{ for all } i, \gcd(i, G) = 1\},$$

where  $\Omega \subseteq \Sigma_m := \{0, \dots, m-1\}^\mathbb{N}$  and  $x|_{iG} := (x_{i\ell_k})_{k=1}^\infty$ . The work of [22] concerns the decomposition of the lattices  $\mathbb{N}$  into independent sublattices according to the multiple constraints  $G$  and calculates its density among the entire lattice  $\mathbb{N}$ . The study can be regarded as a study of the thermodynamic formalism of  $X_\Omega^G$  with the potential function  $S_N^G$  in  $\mathbb{N}^d, d \geq 1$ . In addition, the challenge of the current problem is to look at the same issue as [22] in  $\mathbb{N}^d, d \geq 1$ . The results can be summarized in the following table.

Thermodynamics and LDP		
Limit measure and KS entropy	2-multiple, $d = 1$	Theorem 1.1
	2-multiple, $d \geq 1$	Theorem 3.1
		Theorem 3.3
		Theorem 3.9
	$S_{N_1 \times \dots \times N_d}^{\mathbf{G}}, d \geq 1$	Theorem 3.4
		Theorem 3.6
		Theorem 3.11
Formula of the free energy function	2-multiple, $d = 1$	Theorem 1.2
	2-multiple, $d \geq 1$	Theorem 1.3
	$S_N^G, d = 1$	Theorem 4.1
	$S_{N_1 \times \dots \times N_d}^{\mathbf{G}}, d \geq 1$	Theorem 4.4 and 4.5

## 2. PRELIMINARIES

In this section, we provide necessary materials and results on the decomposition of the multidimensional lattice  $\mathbb{N}^d$  into independent sublattices and calculate their densities.

Let  $\mathbf{p}_1, \dots, \mathbf{p}_k \in \mathbb{N}^d$  with  $\mathbf{p}_i = (p_{i1}, \dots, p_{id})$  for all  $1 \leq i \leq k$ . Denote by  $\mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k} = \{(i_1, \dots, i_d) \in \mathbb{N}^d : p_{j1} \nmid i_1 \text{ or } \dots \text{ or } p_{jd} \nmid i_d \text{ for all } 1 \leq j \leq k\}$ ,  $\mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k} = \{(p_{11}^{\ell_1} \dots p_{k1}^{\ell_k}, \dots, p_{1d}^{\ell_1} \dots p_{kd}^{\ell_k}) : \ell_j \geq 0 \text{ for all } 1 \leq j \leq k\}$  and

$$\mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{i}) = \{(i_1 p_{11}^{\ell_1} \dots p_{k1}^{\ell_k}, \dots, i_d p_{1d}^{\ell_1} \dots p_{kd}^{\ell_k}) : \ell_j \geq 0 \text{ for all } 1 \leq j \leq k\}$$

is the lattice  $\mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  that starts at  $\mathbf{i}$ . We also need the following definitions.

**Definition 2.1.** For  $N_1, \dots, N_d$  and  $M_1, \dots, M_d \geq 1$ , let

1.  $S^{(j)} = \langle p_{1j}, \dots, p_{kj} \rangle = \{1 = \ell_1^{(j)} < \ell_2^{(j)} < \dots\}$  for all  $1 \leq j \leq d$ .

2.  $\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} = \{\mathbf{i} \in \mathcal{N}_{N_1 \times \dots \times N_d} : i_j \ell_{M_j}^{(j)} \leq N_j < i_j \ell_{M_{j+1}}^{(j)} \text{ for all } 1 \leq j \leq d\}$   
be the subset of  $\mathcal{N}_{N_1 \times \dots \times N_d}$  which satisfies  $i_j \ell_{M_j}^{(j)} \leq N_j < i_j \ell_{M_{j+1}}^{(j)}$  for all  $1 \leq j \leq d$ .
3.  $\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} = \mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \cap \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  be the subset of  $\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}$  which belongs to  $\mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ .

The following lemmas are  $\mathbb{N}^d$  version of Lemmas 2.4 and 2.7 in [1].

**Lemma 2.2.** For  $\mathbf{p}_1, \dots, \mathbf{p}_k \in \mathbb{N}^d$  with  $\gcd(p_{is}, p_{js}) = 1$  for all  $1 \leq i \neq j \leq k$  and  $1 \leq s \leq d$ ,

$$\mathbb{N}^d = \bigsqcup_{\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}} \mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{i}).$$

*Proof.* We first claim that for all  $\mathbf{i} \neq \mathbf{i}' \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ ,  $\mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{i}) \cap \mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{i}') = \emptyset$ . If we suppose not, then there exist  $\mathbf{i} \neq \mathbf{i}' \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  such that  $\mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{i}) \cap \mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{i}') \neq \emptyset$ . Since  $\mathbf{i} \neq \mathbf{i}'$ ,

$$(i_1 p_{11}^{\ell_1} \dots p_{k1}^{\ell_k}, \dots, i_d p_{1d}^{\ell_1} \dots p_{kd}^{\ell_k}) = (i'_1 p_{11}^{\ell'_1} \dots p_{k1}^{\ell'_k}, \dots, i'_d p_{1d}^{\ell'_1} \dots p_{kd}^{\ell'_k})$$

for some  $\ell_i \neq \ell'_i, 1 \leq i \leq k$ . Without loss of generality, we may assume  $\ell_i > \ell'_i$ . Then by the  $i_s p_{is}^{\ell_i - \ell'_i} \prod_{j \neq i} p_{js}^{\ell_j} = i'_s \prod_{j \neq i} p_{js}^{\ell'_j}$  for all  $1 \leq s \leq d$  and  $\gcd(p_{is}, p_{js}) = 1$  for all  $1 \leq i \neq j \leq k$  and  $1 \leq s \leq d$ , we have  $p_{is} | i'_s$  for all  $1 \leq s \leq d$ . This contradicts with  $\mathbf{i}' \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ .

It remains to show that the equality holds. For  $\mathbf{i} \in \mathbb{N}^d$ , there exist  $\ell_{js} \geq 0$  for all  $1 \leq s \leq d$  and  $1 \leq j \leq k$  such that

$$(i_1, \dots, i_d) = \left( i'_1 \prod_{j=1}^k p_{j1}^{\ell_{j1}}, \dots, i'_d \prod_{j=1}^k p_{jd}^{\ell_{jd}} \right),$$

where  $p_{js} \nmid i'_s$  for all  $1 \leq j \leq k$  and for all  $1 \leq s \leq d$ . Take  $\ell_j = \min\{\ell_{js} : 1 \leq s \leq d\}$  for all  $1 \leq j \leq k$ . Then, we have  $(i_1, \dots, i_d) \in \mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\frac{i_1}{\prod_{j=1}^k p_{j1}^{\ell_j}}, \dots, \frac{i_d}{\prod_{j=1}^k p_{jd}^{\ell_j}})$  and  $(\frac{i_1}{\prod_{j=1}^k p_{j1}^{\ell_j}}, \dots, \frac{i_d}{\prod_{j=1}^k p_{jd}^{\ell_j}}) \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ . The converse is then clear.  $\square$

**Lemma 2.3.** For  $M_i, N_i \geq 1$  for all  $1 \leq i \leq d$ , we have the following assertions.

1.  $|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| = \prod_{i=1}^d \left( \left\lfloor \frac{N_i}{\ell_{M_i}^{(i)}} \right\rfloor - \left\lfloor \frac{N_i}{\ell_{M_{i+1}}^{(i)}} \right\rfloor \right).$
2.  $\lim_{N_1, \dots, N_d \rightarrow \infty} \frac{|\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} = \prod_{i=1}^k \left( 1 - \frac{1}{p_{i1} p_{i2} \dots p_{id}} \right).$
3.  $\lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} \sum_{M_1=1}^{N_1} \dots \sum_{M_d=1}^{N_d} |\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| \log a_{M_1, \dots, M_d} =$   
 $\sum_{M_1, \dots, M_d=1}^{\infty} \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} |\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| \log a_{M_1, \dots, M_d}.$

*Proof.*

1. Since  $\mathbf{i} \in \mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}$  if and only if  $i_j \ell_{M_j}^{(j)} \leq N_j < i_j \ell_{M_j+1}^{(j)}$  for all  $1 \leq j \leq d$ .  
 d. It follows that  $\frac{N_j}{\ell_{M_j+1}^{(j)}} < i_j \leq \frac{N_j}{\ell_{M_j}^{(j)}}$  for all  $1 \leq j \leq d$ . Therefore

$$|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| = \prod_{i=1}^d \left( \left\lfloor \frac{N_i}{\ell_{M_i}^{(i)}} \right\rfloor - \left\lfloor \frac{N_i}{\ell_{M_i+1}^{(i)}} \right\rfloor \right).$$

2. Let the complement of  $\mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  be

$$\mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}^c = \bigcup_{j=1}^k \mathcal{S}_j = \bigcup_{j=1}^k \{\mathbf{i} : p_{js} \mid i_s \text{ for all } 1 \leq s \leq d\}.$$

Since  $\gcd(p_{is}, p_{js}) = 1$  for all  $1 \leq i \neq j \leq k$  and  $1 \leq s \leq d$ , we have for any  $1 \leq \ell \leq k$ ,

$$\bigcap_{w=1}^{\ell} \mathcal{S}_{j_w} = \left\{ \mathbf{i} : \prod_{w=1}^{\ell} p_{j_w s} \mid i_s \text{ for all } 1 \leq s \leq d \right\}.$$

Then the inclusion-exclusion principle infers that

$$\begin{aligned} \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{|\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} &= \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \cap \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} \\ &= \lim_{N_1, \dots, N_d \rightarrow \infty} 1 - \frac{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \cap \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}^c|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} \\ &= \lim_{N_1, \dots, N_d \rightarrow \infty} 1 - \left( \sum_{n=1}^k \frac{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \cap \mathcal{S}_n|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} \right. \\ &\quad - \sum_{1 \leq n_1 \neq n_2 \leq k} \frac{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \cap \mathcal{S}_{n_1} \cap \mathcal{S}_{n_2}|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} \\ &\quad + \sum_{1 \leq n_1 \neq n_2 \neq n_3 \leq k} \frac{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \cap \mathcal{S}_{n_1} \cap \mathcal{S}_{n_2} \cap \mathcal{S}_{n_3}|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} \\ &\quad \left. - \dots + (-1)^{k-1} \frac{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \cap \mathcal{S}_1 \cap \dots \cap \mathcal{S}_k|}{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}|} \right) \\ &= \prod_{i=1}^k \left( 1 - \frac{1}{p_{i1} p_{i2} \dots p_{id}} \right). \end{aligned}$$

3. Define

$$\hat{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} = \begin{cases} |\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| & , \text{ if } M_j \leq N_j \text{ for all } 1 \leq j \leq d, \\ 0 & , \text{ otherwise.} \end{cases}$$

Then

$$\begin{aligned} &\lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} \sum_{M_1=1}^{N_1} \dots \sum_{M_d=1}^{N_d} |\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| \log a_{M_1, \dots, M_d} \\ &= \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} \sum_{M_1, \dots, M_d=1}^{\infty} \hat{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \log a_{M_1, \dots, M_d}. \end{aligned}$$

We claim that  $\sum_{M_1, \dots, M_d=1}^{\infty} \frac{\hat{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \log a_{M_1, \dots, M_d}}{N_1 \dots N_d}$  converges uniformly in  $N_1, \dots, N_d$  by Weierstrass M-test with

$$\begin{aligned} \left| \frac{\hat{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \log a_{M_1, \dots, M_d}}{N_1 \dots N_d} \right| &\leq \left| \frac{|\mathcal{J}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| \log a_{M_1, \dots, M_d}}{N_1 \dots N_d} \right| \\ &\leq \prod_{i=1}^d \left( \frac{1}{\ell_{M_i}^{(i)}} - \frac{1}{\ell_{M_i+1}^{(i)}} \right) \log a_{M_1, \dots, M_d} \end{aligned}$$

for all  $N_1, \dots, N_d \geq 1$  and

$$\begin{aligned} &\sum_{M_1, \dots, M_d=1}^{\infty} \prod_{i=1}^d \left( \frac{1}{\ell_{M_i}^{(i)}} - \frac{1}{\ell_{M_i+1}^{(i)}} \right) \log a_{M_1, \dots, M_d} \\ &\leq \sum_{M_1, \dots, M_d=1}^{\infty} \prod_{i=1}^d \left( \frac{1}{\ell_{M_i}^{(i)}} - \frac{1}{\ell_{M_i+1}^{(i)}} \right) \log C^{M_1 \dots M_d} < \infty, \end{aligned}$$

whenever  $a_{M_1, \dots, M_d} \leq C^{M_1 \dots M_d}$  for all  $M_1, \dots, M_d \geq 1$ .

Thus,

$$\begin{aligned} &\lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} \sum_{M_1=1}^{N_1} \dots \sum_{M_d=1}^{N_d} |\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| \log a_{M_1, \dots, M_d} \\ &= \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} \sum_{M_1, \dots, M_d=1}^{\infty} \hat{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \log a_{M_1, \dots, M_d} \\ &= \sum_{M_1, \dots, M_d=1}^{\infty} \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} \hat{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d} \log a_{M_1, \dots, M_d} \\ &= \sum_{M_1, \dots, M_d=1}^{\infty} \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \dots N_d} |\mathcal{K}_{N_1 \times \dots \times N_d; M_1, \dots, M_d}| \log a_{M_1, \dots, M_d}. \end{aligned}$$

The proof is complete.  $\square$

Before state the main results, we need more definitions for semigroup  $\mathbf{G}$  which is generated by vectors  $\mathbf{p}_1, \dots, \mathbf{p}_k \in \mathbb{N}^d$ . For  $d, k \geq 1$ ,  $\mathbf{p}_i = (p_{i1}, \dots, p_{id}) \in \mathbb{N}^d$ ,  $1 \leq i \leq k$  and for each  $1 \leq j \leq d$ ,  $\gcd(p_{ij}, p_{i'j}) = 1$  for all  $1 \leq i \neq i' \leq k$ . Let  $\mathbf{G} = \langle \mathbf{p}_1, \dots, \mathbf{p}_k \rangle = \{ \mathbf{1} = l_1^{(j)} \prec^{(j)} l_2^{(j)} \prec^{(j)} \dots \}$  be the semigroup generated by  $\mathbf{p}_1, \dots, \mathbf{p}_k$  with an order  $\prec^{(j)}$  such that  $\mathbf{G}$  follows  $S^{(j)}$  for each  $1 \leq j \leq d$ . In particular, when  $k = 1$ ,  $\mathbf{G} = \langle \mathbf{p}_1 \rangle = \{ l_n = \mathbf{p}_1^{n-1} \}_{n=1}^{\infty}$ <sup>3</sup>.

### 3. THE INFINITE-VOLUME LIMIT $\mu_{\infty}$ ON $\mathbb{N}^d$ AND KOLMOGOROV-SINAI ENTROPY

In this section, we prove the  $\mathbb{N}^d$  version of Theorem 3.1 in [8] for layer stationarity and multiplication invariance (Theorems 3.1 and 3.4), and then obtain the limit measures (Theorems 3.3 and 3.6). We also give the  $\mathbb{N}^d$  version of Lemma 3.4 in [8] (Lemmas 3.8 and 3.10) to obtain the Kolmogorov-Sinai entropies (Theorems 3.9 and 3.11).

<sup>3</sup> $\mathbf{p}_1^{n-1} = (p_{11}^{n-1}, \dots, p_{1d}^{n-1})$

### 3.1. Existence and invariance of the limit measure on $\mathbb{N}^d$ .

3.1.1. *2-multiple Hamiltonian.* Let  $\mathbf{p}_1 = (p_1, \dots, p_d) \in \mathbb{N}^d$  and  $\tau^{\mathbf{i}} = \{\tau_\ell^{\mathbf{i}} = \sigma_{\mathbf{p}_1^\ell \cdot \mathbf{i}}\}_{\ell=0}^\infty$  for each  $\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1}$ . Then we have following result which is the  $\mathbb{N}^d$  version of Theorem 3.1 in [8].

**Theorem 3.1.** *Suppose that the  $\{\tau^{\mathbf{i}} : \mathbf{i} \in \mathcal{I}_{\mathbf{p}_1}\}$  form an i.i.d. stationary processes, then the distribution of the corresponding  $\sigma$  is multiplication invariant.*

*Proof.* We are going to show that for every finite collection of points  $\mathbf{v}_1, \dots, \mathbf{v}_k \in \mathbb{N}^d$ , and  $\mathbf{m} \in \mathbb{N}^d$  the joint distribution of

$$(\sigma_{\mathbf{m} \cdot \mathbf{v}_1}, \dots, \sigma_{\mathbf{m} \cdot \mathbf{v}_k})$$

coincides with that of

$$(\sigma_{\mathbf{v}_1}, \dots, \sigma_{\mathbf{v}_k}).$$

First, we write  $\mathbf{v}_j = \mathbf{i}_j \cdot \mathbf{p}_1^{\ell_j}$  with  $\mathbf{i}_j \in \mathcal{I}_{\mathbf{p}_1}$  for all  $1 \leq j \leq k$  and  $\mathbf{m} = \mathbf{i} \cdot \mathbf{p}_1^\ell$  with  $\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1}$ . Denote  $\mathbf{i}'_j \cdot \mathbf{p}_1^{\ell'_j} = \mathbf{i} \cdot \mathbf{i}_j$  with  $\mathbf{i}'_j \in \mathcal{I}_{\mathbf{p}_1}$  for all  $1 \leq j \leq k$ .

Then by the definition of  $\tau$ , we have to prove that the joint distribution of

$$\tau_{\ell_j + \ell + \ell'_j}^{\mathbf{i}'_j}, \quad j \in \{1, \dots, k\}$$

coincides with that of

$$\tau_{\ell_j}^{\mathbf{i}_j}, \quad j \in \{1, \dots, k\}.$$

Denote  $\{\mathbf{i}_1, \dots, \mathbf{i}_k\} = \{\mathbf{i}_{n_1}, \dots, \mathbf{i}_{n_M}\}$  where  $\mathbf{i}_{n_i} \neq \mathbf{i}_{n_j}$  for all  $1 \leq i \neq j \leq M$ . For  $1 \leq w \leq M$ , define

$$X^w = \left( \tau_{\ell_j + \ell + \ell'_j}^{\mathbf{i}'_j} : 1 \leq j \leq k, \mathbf{i}_j = \mathbf{i}_{n_w} \right)$$

and

$$Y^w = \left( \tau_{\ell_j}^{\mathbf{i}_j} : 1 \leq j \leq k, \mathbf{i}_j = \mathbf{i}_{n_w} \right).$$

Then, by the independence of the different layers, the joint distribution of  $\tau_{\ell_j + \ell + \ell'_j}^{\mathbf{i}'_j}, j \in \{1, \dots, k\}$  and  $\tau_{\ell_j}^{\mathbf{i}_j}, j \in \{1, \dots, k\}$  coincide with the joint distribution of  $\otimes_{w=1}^M X^w$  and  $\otimes_{w=1}^M Y^w$  respectively, where  $\otimes$  denotes independent joining.

Therefore, it remains to show that for each  $1 \leq w \leq M$ , the distributions of  $X^w$  and  $Y^w$  coincide. Since the layers  $\tau^{\mathbf{i}'_j}$  and  $\tau^{\mathbf{i}_j}$  are i.i.d., we have  $X^w$  and  $Y^w$  have the same distribution. The proof is complete.  $\square$

**Remark 3.2.** We remark that for  $\mathbf{i}, \mathbf{i}_1 \neq \mathbf{i}_2 \in \mathcal{I}_{\mathbf{p}_1}$  if  $\mathbf{i} \cdot \mathbf{i}_1 = \mathbf{i}'_1 \cdot \mathbf{p}_1^{\ell_1}$  and  $\mathbf{i} \cdot \mathbf{i}_2 = \mathbf{i}'_2 \cdot \mathbf{p}_1^{\ell_2}$  with  $\mathbf{i}'_1, \mathbf{i}'_2 \in \mathcal{I}_{\mathbf{p}_1}$ , then  $\mathbf{i}'_1 \neq \mathbf{i}'_2$ . When  $\ell_1 = \ell_2$ , the statement easily true. If we assume  $\ell_1 > \ell_2$  and  $\mathbf{i}'_1 = \mathbf{i}'_2$ , then  $\mathbf{i} \cdot \mathbf{i}_1 = \mathbf{i} \cdot \mathbf{i}_2 \cdot \mathbf{p}_1^{\ell_1 - \ell_2}$ . That gives  $\mathbf{i}_1 = \mathbf{i}_2 \cdot \mathbf{p}_1^{\ell_1 - \ell_2}$  which contradicts with  $\mathbf{i}_1 \in \mathcal{I}_{\mathbf{p}_1}$ .

As a consequence of Theorem 3.1 with the existence of the infinte-volume limit in each layer of  $\tau$  spins we have the following results.



**Theorem 3.3.** *Let*

$$\mu_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}}) = \frac{e^{-H_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}})}}{\sum_{\sigma_{\ell_{\mathbf{j}(\mathbf{i})}} = \pm 1, \mathbf{i} \in \mathcal{N}_{N_1 \times \dots \times N_d}} e^{-H_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}})}}$$

be the finite-volume probability measure corresponding to the Hamiltonian

$$H_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}}) = -\beta \sum_{i_1=1}^{N_1} \dots \sum_{i_d=1}^{N_d} \sigma_{l_{\mathbf{j}(\mathbf{i})}} \sigma_{l_{\mathbf{j}(\mathbf{i})+1}},$$

where  $\mathbf{j}(\mathbf{i})$  is the unique number such that  $\mathbf{i} = (k_1, \dots, k_d) \cdot l_{\mathbf{j}(\mathbf{i})}$  with  $(k_1, \dots, k_d) \in \mathcal{I}_{\mathbf{p}_1}$ .

Then

1. *Unique limit measure :* The measure  $\mu_{N_1 \times \dots \times N_d}$  have a unique weak limit (as  $N_1, \dots, N_d \rightarrow \infty$ ) denoted by  $\mu_\infty$  which is Gibbs.
2. *Independent Ising layers :* Under  $\mu_\infty$ , the  $\tau^{\mathbf{i}}, \mathbf{i} \in \mathcal{I}_{\mathbf{p}_1}$ , are independent and distributed according to the standard Ising model measure  $\mu_\infty^{Ising}$  with a free boundary condition on the left.
3. *Multiplication invariance :* The measure  $\mu_\infty$  is multiplication invariant.

3.1.2. *Generalization of 2-multiple Hamiltonian to  $S_{N_1 \times \dots \times N_d}^{\mathbf{G}}$  on  $\mathbb{N}^d$ .* Let  $\tau^{\mathbf{i}} = \{\tau_k^{\mathbf{i}} = \sigma_{\mathbf{i}, l_k^{(j)}}\}_{k=1}^\infty$  for each  $\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ . For each  $1 \leq i \leq d$ ,  $\gcd(p_{ji}, p_{j'i}) = 1$  for all  $1 \leq j \neq j' \leq k$ .

**Theorem 3.4.** *Suppose that the  $\{\tau^{\mathbf{i}} : \mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}\}$  form an i.i.d. stationary processes, then the distribution of the corresponding  $\sigma$  is multiplication invariant.*

*Proof.* We have to show that for every finite collection of points  $\mathbf{v}_1, \dots, \mathbf{v}_N \in \mathbb{N}^d$  and  $\mathbf{m} \in \mathbb{N}^d$ , the joint distribution of

$$(\sigma_{\mathbf{m} \cdot \mathbf{v}_1}, \dots, \sigma_{\mathbf{m} \cdot \mathbf{v}_N})$$

coincides with that of

$$(\sigma_{\mathbf{v}_1}, \dots, \sigma_{\mathbf{v}_N}).$$

First, we write  $\mathbf{v}_j = \mathbf{i}_j \cdot \mathbf{p}_1^{\ell_{j1}} \dots \mathbf{p}_k^{\ell_{jk}}$  with  $\mathbf{i}_j \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  for all  $1 \leq j \leq N$  and  $\mathbf{m} = \mathbf{i} \cdot \mathbf{p}_1^{m_1} \dots \mathbf{p}_k^{m_k}$  with  $\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ . Note that  $\mathbf{i}'_j \cdot \mathbf{p}_1^{\ell'_{j1}} \dots \mathbf{p}_k^{\ell'_{jk}} = \mathbf{i} \cdot \mathbf{i}_j$  with  $\mathbf{i}'_j \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  for all  $1 \leq j \leq k$ . For convenience, we denote  $\ell_j = (\ell_{j1}, \dots, \ell_{jk})$ ,  $\ell'_j = (\ell'_{j1}, \dots, \ell'_{jk})$  for all  $1 \leq j \leq k$  and  $m = (m_1, \dots, m_k)$ .

Then by the definition of  $\tau$ , we have to prove that the joint distribution of

$$\tau_{\ell_j + m + \ell'_j}^{\mathbf{i}'_j}, \quad j \in \{1, \dots, k\}$$

coincides with that of

$$\tau_{\ell_j}^{\mathbf{i}_j}, \quad j \in \{1, \dots, k\}.$$

Denote  $\{\mathbf{i}_1, \dots, \mathbf{i}_k\} = \{\mathbf{i}_{n_1}, \dots, \mathbf{i}_{n_M}\}$  where  $\mathbf{i}_{n_i} \neq \mathbf{i}_{n_j}$  for all  $1 \leq i \neq j \leq M$ . For  $1 \leq w \leq M$ , we define

$$X^w = \left( \tau_{\ell_j + m + \ell'_j}^{\mathbf{i}'_j} : 1 \leq j \leq k, \mathbf{i}_j = \mathbf{i}_{n_w} \right)$$

and

$$Y^w = \left( \tau_{\ell_j}^{\mathbf{i}_j} : 1 \leq j \leq k, \mathbf{i}_j = \mathbf{i}_{n_w} \right).$$

Then, by the independence of the different layers, the joint distribution of  $\tau_{\ell_j+m+\ell'_j}^{\mathbf{i}'_j}, j \in \{1, \dots, k\}$  and  $\tau_{\ell_j}^{\mathbf{i}_j}, j \in \{1, \dots, k\}$  coincide with the joint distribution of  $\otimes_{w=1}^M X^w$  and  $\otimes_{w=1}^M Y^w$  respectively.

Therefore, it remains to show that for each  $1 \leq w \leq M$ , the distributions of  $X^w$  and  $Y^w$  coincide. Since the layers  $\tau^{\mathbf{i}'_j}$  and  $\tau^{\mathbf{i}_j}$  are i.i.d., we have that  $X^w$  and  $Y^w$  have the same distribution. The proof is complete.  $\square$

**Remark 3.5.** If  $\mathbf{i}, \mathbf{i}_1 \neq \mathbf{i}_2 \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  and  $\mathbf{i} \cdot \mathbf{i}_1 = \mathbf{i}'_1 \cdot \mathbf{p}_1^{\ell'_{11}} \cdots \mathbf{p}_k^{\ell'_{1k}}, \mathbf{i} \cdot \mathbf{i}_2 = \mathbf{i}'_2 \cdot \mathbf{p}_1^{\ell'_{21}} \cdots \mathbf{p}_k^{\ell'_{2k}}$  with  $\mathbf{i}'_1, \mathbf{i}'_2 \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ , then  $\mathbf{i}'_1 \neq \mathbf{i}'_2$ . When  $(\ell'_{11}, \dots, \ell'_{1k}) = (\ell'_{21}, \dots, \ell'_{2k})$ , the statement clearly true. If  $(\ell'_{11}, \dots, \ell'_{1k}) \neq (\ell'_{21}, \dots, \ell'_{2k})$  and  $\mathbf{i}'_1 = \mathbf{i}'_2$ , without loss of generality, we may assume  $\ell'_{11} > \ell'_{21}$ . This implies  $\mathbf{i} \cdot \mathbf{i}_1 \cdot \mathbf{c}_1 = \mathbf{i} \cdot \mathbf{i}_2 \cdot \mathbf{p}_1^{\ell'_{11}-\ell'_{21}} \cdot \mathbf{c}_2$ , then  $\mathbf{i}_1 \cdot \mathbf{c}_1 = \mathbf{i}_2 \cdot \mathbf{p}_1^{\ell'_{11}-\ell'_{21}} \cdot \mathbf{c}_2$  where  $\mathbf{c}_1, \mathbf{c}_2$  are the vectors only multipliable by  $\mathbf{p}_i$ 's for  $2 \leq i \leq k$ . Then the coprime property of  $\mathbf{p}_1$  and  $\mathbf{p}_i, 2 \leq i \leq k$  gives  $\mathbf{p}_1 \mid \mathbf{i}_1$  which contradicts  $\mathbf{i}_1 \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ .

As a consequence of Theorem 3.1 with the existence of the infinte-volume limit in each layer of  $\tau$  spins we have the following theorem.

**Theorem 3.6.** *Let*

$$\mu_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}}) = \frac{e^{-H_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}})}}{\sum_{\sigma_{l_{\mathbf{j}(\mathbf{i})}}^{(j)} = \pm 1, \mathbf{i} \in \mathcal{N}_{N_1 \times \dots \times N_d}} e^{-H_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}})}}$$

be the finite-volume probability measure corresponding to the Hamiltonian

$$H_{N_1 \times \dots \times N_d}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}}) = -\beta \sum_{i_1=1}^{N_1} \cdots \sum_{i_d=1}^{N_d} \sigma_{l_{\mathbf{j}(\mathbf{i})}}^{(j)} \sigma_{l_{\mathbf{j}(\mathbf{i})+1}}^{(j)},$$

where  $\mathbf{j}(\mathbf{i})$  is the unique number such that  $\mathbf{i} = (k_1, \dots, k_d) \cdot l_{\mathbf{j}(\mathbf{i})}^{(j)}$  with  $(k_1, \dots, k_d) \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ . Then

1. *Unique limit measure :* The measure  $\mu_{N_1 \times \dots \times N_d}$  have a unique weak limit with respect to order  $\prec^{(j)}$  denoted by  $\mu_{\infty}^{(j)}$  which is Gibbs.
2. *Independent Ising layers :* Under  $\mu_{\infty}^{(j)}$ , the  $\tau^{\mathbf{i}}, \mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$  are independent and distributed according to the standard Ising model measure  $\mu_{\infty}^{Ising}$  with a free boundary condition on the left.
3. *Multiplication invariance :* The measure  $\mu_{\infty}^{(j)}$  is multiplication invariant.

**Remark 3.7.** We remark that the order  $\prec^{(j)}$  leads  $\mu_{N_1 \times \dots \times N_d}$  to be Markov on each independent layer  $\tau^{\mathbf{i}}, \mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ .

**3.2. Kolmogorov-Sinai entropy with respect to the limit measure on  $\mathbb{N}^d$ .**

**3.2.1. 2-multiple Hamiltonian.** Recall that  $\mathcal{L}_{N_1 \times \dots \times N_d}(\mathbf{i}) = \mathcal{M}_{\mathbf{p}_1}(\mathbf{i}) \cap \mathcal{N}_{N_1 \times \dots \times N_d}$  is the subset of  $\mathcal{M}_{\mathbf{p}_1}(\mathbf{i})$ , which belongs to the  $\mathcal{N}_{N_1 \times \dots \times N_d}$  lattice. Then we have results as follows.

**Lemma 3.8.** *Let  $\phi : \mathbb{N} \rightarrow \mathbb{R}$  be a measurable function such that there exist  $C' > 0$  and  $r > 0$  such that  $|\phi(n)| \leq C'n^r$  for all  $n \in \mathbb{N}$ . Then we have*

$$\lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \cdots N_d} \sum_{\mathbf{i} \in \mathcal{I}_{\mathbf{p}}} \phi(|\mathcal{L}_{N_1 \times \dots \times N_d}(\mathbf{i})|) = \sum_{\ell=1}^{\infty} \frac{(p_1 \cdots p_d - 1)^2}{(p_1 \cdots p_d)^{\ell+1}} \phi(\ell).$$

*Proof.* The proof is a direct consequence of Lemma 2.3. More precisely,

$$\begin{aligned} & \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \cdots N_d} \sum_{\mathbf{i} \in \mathcal{I}_{\mathbf{p}}} \phi(|\mathcal{L}_{N_1 \times \dots \times N_d}(\mathbf{i})|) \\ &= \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \cdots N_d} \sum_{\ell=1}^{K(N_1, \dots, N_d)} \left(1 - \frac{1}{p_1 \cdots p_d}\right) \left(\frac{N_1 \cdots N_d}{(p_1 \cdots p_d)^{\ell-1}} - \frac{N_1 \cdots N_d}{(p_1 \cdots p_d)^{\ell}}\right) \phi(\ell) \\ &= \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \cdots N_d} \sum_{\ell=1}^{K(N_1, \dots, N_d)} \frac{N_1 \cdots N_d (p_1 \cdots p_d - 1)^2}{(p_1 \cdots p_d)^{\ell+1}} \phi(\ell) \\ &= \sum_{\ell=1}^{\infty} \frac{(p_1 \cdots p_d - 1)^2}{(p_1 \cdots p_d)^{\ell+1}} \phi(\ell), \end{aligned}$$

where the last equality holds by Weierstrass M-test with  $|\phi(\ell)| \leq C'\ell^r$  and  $K(N_1, \dots, N_d)$  is the maximum cardinality of  $\mathcal{L}_{N_1 \times \dots \times N_d}(\mathbf{i})$  for all  $\mathbf{i} \in \mathcal{I}_{\mathbf{p}}$ .  $\square$

Theorem 3.9 below is an immediate consequence of Lemma 3.8.

**Theorem 3.9.** *The explicit formula for the KS entropy of  $\mu_{\infty}$  is*

$$\begin{aligned} & - \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \cdots N_d} \mathbb{E}_{\mu_{\infty}} \log \mu_{\infty}(\sigma_{\mathcal{N}_{N_1 \times \dots \times N_d}}) \\ &= - \sum_{k=1}^{\infty} \frac{(p_1 \cdots p_d - 1)^2}{(p_1 \cdots p_d)^{k+1}} \mathbb{E}_{\mu_{\infty}^{I_{\text{sing}}}} \log \mu_{\infty}^{I_{\text{sing}}}(\tau_0, \dots, \tau_{k-1}). \end{aligned}$$

**3.2.2. Generalization of 2-multiple Hamiltonian to  $S_{N_1 \times \dots \times N_d}^{\mathbf{G}}$  on  $\mathbb{N}^d$ .** For each  $1 \leq j \leq d$  and  $\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k} \cap \mathcal{N}_{N_1 \times \dots \times N_d}$ , let  $\mathcal{L}_{N_1 \times \dots \times N_d}^{(j)}(\mathbf{i})$  be the subset of  $\mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{i})$  which satisfies  $i_j p_{1j}^{\ell_1} \cdots p_{kj}^{\ell_k} \leq N_j$ . Then the similar result of Lemma 3.8 is obtained.

**Lemma 3.10.** *Let  $\phi : \mathbb{N} \rightarrow \mathbb{R}$  be a measurable function such that there exist  $C' > 0$  and  $r > 0$  such that  $|\phi(n)| \leq C'n^r$  for all  $n \in \mathbb{N}$ . Then we have*

$$\begin{aligned} & \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \cdots N_d} \sum_{\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}} \phi(|\mathcal{L}_{N_1 \times \dots \times N_d}^{(j)}(\mathbf{i})|) \\ &= \prod_{i=1}^k \left(1 - \frac{1}{p_{i1} \cdots p_{id}}\right) \sum_{M_1, \dots, M_d=1}^{\infty} \left[ \prod_{i=1}^d \left(\frac{1}{\ell_{M_i}^{(i)}} - \frac{1}{\ell_{M_i+1}^{(i)}}\right) \right] \phi(M_j). \end{aligned}$$

*Proof.* Applying the same argument of the Lemma 3.8 with Lemma 2.3, the proof is complete.  $\square$

By Lemma 3.10, we have the following result.

**Theorem 3.11.** *For each  $1 \leq j \leq d$ , the explicit formula for the KS entropy of  $\mu_\infty^{(j)}$  is*

$$\begin{aligned} & - \lim_{N_1, \dots, N_d \rightarrow \infty} \frac{1}{N_1 \cdots N_d} \mathbb{E}_{\mu_\infty} \log \mu_\infty^{(j)}(\sigma_{N_1 \times \dots \times N_d}) \\ & = -C \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k^{(j)}} - \frac{1}{\ell_{k+1}^{(j)}} \right) \mathbb{E}_{\mu_\infty^{Ising}} \log \mu_\infty^{Ising}(\tau_0, \dots, \tau_{k-1}), \end{aligned}$$

where  $C$  is defined in Theorem 4.5.

#### 4. FREE ENERGY FUNCTIONS AND LARGE DEVIATION PRINCIPLE

In this section, we obtain the generalization of Theorem 1.2 (Theorem 4.1), and we consider two types of generalizations of Theorem 1.3 (Theorems 4.4 and 4.5).

**4.1. Generalization of 2-multiple Hamiltonian to  $S_N^G$  on  $\mathbb{N}$ .** Let  $k \geq 1$ ,  $p_1, p_2, \dots, p_k$  be co-primes, and  $G = \langle p_1, p_2, \dots, p_k \rangle = \{1 = \ell_1 < \ell_2 < \dots\}$  be a semigroup generated by  $p_1, p_2, \dots, p_k$ . Denote  $\gamma(G) = \sum_{i=1}^{\infty} \frac{1}{\ell_i}$ . The LDP of  $S_N^G = \sum_{i=1}^N \sigma_{\ell_{j(i)}} \sigma_{\ell_{j(i)+1}}$ ,  $j(i)$  is the unique number such that  $i = i' \ell_j$  and  $p_n \nmid i'$  for all  $1 \leq n \leq k$ , is presented in the following theorem.

**Theorem 4.1.** *The following statements hold true.*

1. *The explicit expression of the free energy function associated to the multiple sum  $S_N^G$  is*

$$F_r(\beta) = \frac{1 + \gamma(G)^{-1}}{2} \log(r(1-r)) + \gamma(G)^{-1} \log |v^T \cdot e_+|^2 + \log \Lambda_+ + \mathcal{G}(\beta),$$

where

$$\mathcal{G}(\beta) = \gamma(G)^{-1} \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k} - \frac{1}{\ell_{k+1}} \right) \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^k \right).$$

2. *The function  $F_r(\beta)$  is differentiable with respect to  $\beta \in \mathbb{R}$ .*
3. *The multiple average  $\frac{S_N^G}{N}$  satisfies a LDP with rate function given by*

$$I_r(x) = \sup_{\beta \in \mathbb{R}} (\beta x - F_r(\beta)).$$

Furthermore, if  $(F_r)'(\eta) = y$ , then  $I_r(y) = \eta y - F_r(\eta)$ .

*Proof.*

1. By the  $\mathbb{N}^d$  version of Lemma 2.7 in [1] and the similar argument of Theorem 3.2 in [2], we obtain

$$\begin{aligned}
F_r(\beta) &= B \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k} - \frac{1}{\ell_{k+1}} \right) \log(r(1-r))^{\frac{k+1}{2}} Z(\beta, h, k+1) \\
&= B \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k} - \frac{1}{\ell_{k+1}} \right) \log(r(1-r))^{\frac{k+1}{2}} \\
&\quad + B \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k} - \frac{1}{\ell_{k+1}} \right) \log(|v^T \cdot e_+|^2 \Lambda_+^k + (2 \cosh(h) - |v^T \cdot e_+|^2) \Lambda_-^k) \\
&= B \frac{1+\gamma(G)}{2} \log(r(1-r)) + B \log |v^T \cdot e_+|^2 + B \gamma(G) \log \Lambda_+ + \mathcal{G}(\beta) \\
&= \frac{1+\gamma(G)^{-1}}{2} \log(r(1-r)) + \gamma(G)^{-1} \log |v^T \cdot e_+|^2 + \log \Lambda_+ + \mathcal{G}(\beta),
\end{aligned}$$

where the constant  $B = \prod_{i=1}^k \left(1 - \frac{1}{p_i}\right) = \gamma(G)^{-1}$ ,

$$Z(\beta, h, k+1) = v^T \begin{bmatrix} e^{\beta+h} & e^{-\beta} \\ e^{-\beta} & e^{\beta-h} \end{bmatrix}^k v$$

and

$$\mathcal{G}(\beta) = \gamma(G)^{-1} \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k} - \frac{1}{\ell_{k+1}} \right) \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^k \right).$$

2. The proof is similar to the (2) of Theorem 3.2 in [2]. More precisely, we are going to show the sum

$$\sum_{k=1}^{\infty} \left( \frac{1}{\ell_k} - \frac{1}{\ell_{k+1}} \right) \left[ \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^k \right) \right]'$$

coverge uniformly with respect to  $\beta \in \mathbb{R}$ , where the notation  $'$  stays for the derivative with respect to  $\beta$ . Then, we apply the same reasoning on

$$\left[ \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^k \right) \right]'$$

and then apply the Weierstrass M-test, and the proof is complete.

3. The proof is a direct consequence of Theorem 4.1 (2) and Gärtner-Ellis Theorem [2].  $\square$

#### Corollary 4.2.

1. For  $k = 1$ ,  $p_1 \geq 1$ ,  $G = \langle p_1 \rangle = \{p_1^{i-1} : i \in \mathbb{N}\}$  and  $\gamma(S) = \frac{p_1}{p_1-1}$ . The explicit expression of the free energy function associated to the multiple sum  $S_N^G$  is

$$F_r(\beta) = \frac{2p_1-1}{2p_1} \log(r(1-r)) + \frac{2(p_1-1)}{p_1} \log |v^T \cdot e_+| + \log \Lambda_+ + \mathcal{G}(\beta),$$

where

$$\mathcal{G}(\beta) = \frac{p_1-1}{p_1} \sum_{k=1}^{\infty} \frac{p_1-1}{p_1^k} \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^k \right).$$




FIGURE 1. Plot of the  $F_r(\beta)$  with  $p_i, 1 \leq i \leq 5$  for different  $r$  values.

2. In addition,  $p_1 = 2$ ,  $G = \langle 2 \rangle = \{\ell_i = 2^{i-1} : i \in \mathbb{N}\}$  and  $\gamma(S) = 2$ . The explicit expression of the free energy function associated to the multiple sum  $S_N^G$  is

$$F_r(\beta) = \frac{3}{4} \log(r(1-r)) + \log |v^T \cdot e_+| + \log \Lambda_+ + \mathcal{G}(\beta),$$

where

$$\mathcal{G}(\beta) = \frac{1}{2} \sum_{k=1}^{\infty} \frac{1}{2^k} \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^k \right).$$

Which coincides with the result in Theorem 1.2.

**Example 4.3.** Figure 1 illustrates the free energy function for different  $r \in (0, 1)$  which is obtained from Theorem 4.1 by truncating the sum to the first 100 terms. The graph is obtained for the case  $d = 1$  with  $p_1 = 2, p_2 = 3, p_3 = 5, p_4 = 7, p_5 = 11$ ,  $G = \langle 2, 3, 5, 7, 11 \rangle$  and  $\gamma(G)^{-1} = \left(1 - \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{5}\right) \left(1 - \frac{1}{7}\right) \left(1 - \frac{1}{11}\right) = \frac{16}{77}$ .

**4.2. Generalization of 2-multiple Hamiltonian to  $S_{N_1 \times \dots \times N_d}^{\mathbf{G}}$  on  $\mathbb{N}^d$ .** Recall that  $S^{(j)} = \langle p_{1j}, \dots, p_{kj} \rangle = \{1 = \ell_1^{(j)} < \ell_2^{(j)} < \dots\}$  is the semigroup generated by  $p_{1j}, \dots, p_{kj}$ . And  $\mathbf{G} = \langle \mathbf{p}_1, \dots, \mathbf{p}_k \rangle = \{\mathbf{1} = l_1^{(j)} \prec^{(j)} l_2^{(j)} \prec^{(j)} \dots\}$  be the semigroup generated by  $\mathbf{p}_1, \dots, \mathbf{p}_k$  with an order  $\prec^{(j)}$  such that  $\mathbf{G}$  follows  $S^{(j)}$  for each  $1 \leq j \leq d$ . The LDP of  $S_{N_1 \times \dots \times N_d}^{\mathbf{G}} = \sum_{i_1=1}^{N_1} \dots \sum_{i_d=1}^{N_d} \sigma_{l_{\mathbf{j}(\mathbf{i})}^{(j)}} \sigma_{l_{\mathbf{j}(\mathbf{i})+1}^{(j)}}$ ,  $\mathbf{j}(\mathbf{i})$  is the unique number such that  $\mathbf{i} = (k_1, \dots, k_d) \cdot l_{\mathbf{j}(\mathbf{i})}^{(j)}$  with  $(k_1, \dots, k_d) \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k}$ , is described by the following energy function.

The *directional free energy function* is defined as following. For  $1 \leq j \leq d$ ,

$$F_r(\beta) := \lim_{N_1, \dots, N_d \rightarrow \infty} \log \mathbb{E}_r \left( e^{\beta \sum_{i_1=1}^{N_1} \dots \sum_{i_d=1}^{N_d} \sigma_{l_{\mathbf{j}(\mathbf{i})}^{(j)}} \sigma_{l_{\mathbf{j}(\mathbf{i})+1}^{(j)}}} \right).$$

The following result takes the sum over all elements  $\mathbf{i}$  if it is less than the maximum element on its layer intersection with the  $N_1 \times \cdots \times N_d$  lattice (in the  $\prec^{(j)}$  sense).

That is, we take the sum over all  $\mathbf{i} \in \mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{j}(\mathbf{i}))$  if

$$\mathbf{i} \prec^{(j)} \max \{\mathbf{i}' : \mathbf{i}' \in \mathcal{M}_{\mathbf{p}_1, \dots, \mathbf{p}_k}(\mathbf{j}(\mathbf{i})) \cap \mathcal{N}_{N_1 \times \cdots \times N_d}\}.$$

**Theorem 4.4.** *For  $1 \leq j \leq d$ , the explicit formula of the directional energy function associated to the multiple sum  $S_{N_1 \times \cdots \times N_d}^{\mathbf{G}}$  converges to*

$$F_r(\beta) = \prod_{i=1}^k \left(1 - \frac{1}{p_{i1} \cdots p_{id}}\right) \sum_{k_1, \dots, k_d=1}^{\infty} \prod_{i=1}^d \left(\frac{1}{\ell_{k_i}^{(i)}} - \frac{1}{\ell_{k_i+1}^{(i)}}\right) \log(r(1-r))^{\frac{b_{k_1, \dots, k_d}+1}{2}} Z(\beta, h, b_{k_1, \dots, k_d}+1),$$

where  $b_{k_1, \dots, k_d}$  is the number of the elements in  $\mathbf{G}$  less than or equal to the maximum element (with the order  $\prec^{(j)}$ ) in the  $\ell_{k_1}^{(1)} \times \cdots \times \ell_{k_d}^{(d)}$  lattice.

*Proof.* The proof is directly by the Lemmas 2.2 and 2.3 with the observation  $|b_{k_1, \dots, k_d}| \leq k_j$ .  $\square$

Due to the explicit expression of Theorem 4.4 is difficult to obtain, we consider the following type directional free energy function which takes the sum over all  $\mathbf{i} \in \mathcal{I}_{\mathbf{p}_1, \dots, \mathbf{p}_k} \cap \mathcal{N}_{N_1 \times \cdots \times N_d}$  layers with the layer members  $\mathbf{i} \cdot \mathbf{p}_1^{\ell_1} \cdots \mathbf{p}_k^{\ell_k}$  satisfy  $i_j p_{1j}^{\ell_1} \cdots p_{kj}^{\ell_k} \leq N_j$ . Then we have following result.

**Theorem 4.5.** *For  $1 \leq j \leq d$ ,*

1. *The explicit expression of the directional energy function associated to the multiple sum  $S_{N_1 \times \cdots \times N_d}^{\mathbf{G}}$  is*

$$F_r(\beta) = \frac{C + C\gamma(S^{(j)})}{2} \log(r(1-r)) + C \log |v^T \cdot e_+|^2 + C\gamma(S^{(j)}) \log \Lambda_+ + \mathcal{G}(\beta),$$

where  $C = \prod_{i=1}^k \left(1 - \frac{1}{p_{i1} \cdots p_{id}}\right) \prod_{1 \leq i \neq j \leq d} \gamma(S^{(i)})$  and

$$\mathcal{G}(\beta) = C \sum_{k=1}^{\infty} \left(\frac{1}{\ell_k^{(j)}} - \frac{1}{\ell_{k+1}^{(j)}}\right) \log \left(1 + \left(\frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1\right) \left(\frac{\Lambda_-}{\Lambda_+}\right)^k\right).$$

2. *The function  $F_r(\beta)$  is differentiable with respect to  $\beta \in \mathbb{R}$ .*
3. *The multiple average  $\frac{S_{N_1 \times \cdots \times N_d}^{\mathbf{G}}}{N_1 \cdots N_d}$  satisfies a LDP with rate function given by*

$$I_r(x) = \sup_{\beta \in \mathbb{R}} (\beta x - F_r(\beta)).$$

Furthermore, if  $(F_r)'(\eta) = y$ , then  $I_r(y) = \eta y - F_r(\eta)$ .

*Proof.*

1. By Lemmas 2.2 and Lemma 2.3 with the similar process of Theorem 3.2 in [2], we have

$$\begin{aligned} F_r(\beta) &= \lim_{N_1, \dots, N_d \rightarrow \infty} \log \mathbb{E}_r \left( e^{\beta \sum_{i=1}^{N_1} \cdots \sum_{i_d=1}^{N_d} \sigma_{\mathbf{j}(\mathbf{i})}^{(j)} \sigma_{\mathbf{j}(\mathbf{i})+1}^{(j)}} \right) \\ &= \prod_{i=1}^k \left(1 - \frac{1}{p_{i1} \cdots p_{id}}\right) \sum_{k_1=1}^{\infty} \cdots \sum_{k_d=1}^{\infty} \prod_{i=1}^d \left(\frac{1}{\ell_{k_i}^{(i)}} - \frac{1}{\ell_{k_i+1}^{(i)}}\right) \log(r(1-r))^{\frac{k_j+1}{2}} Z(\beta, h, k_j+1). \end{aligned}$$




FIGURE 2. Plot of the  $F_r(\beta)$  with  $\mathbf{p}_i, 1 \leq i \leq 5$  for different  $r$  values.

Then replace  $k_j$  by  $k$ , we obtain

$$\begin{aligned} F_r(\beta) &= C \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k^{(j)}} - \frac{1}{\ell_{k+1}^{(j)}} \right) \log(r(1-r))^{\frac{k+1}{2}} Z(\beta, h, k+1) \\ &= \frac{C + C\gamma(S^{(j)})}{2} \log(r(1-r)) + C \log |v^T \cdot e_+|^2 + C\gamma(S^{(j)}) \log \Lambda_+ + \mathcal{G}(\beta), \end{aligned}$$

where  $C = \prod_{i=1}^k \left( 1 - \frac{1}{p_{i1} \cdots p_{id}} \right) \prod_{1 \leq i \neq j \leq d} \gamma(S^{(i)})$  and

$$\mathcal{G}(\beta) = C \sum_{k=1}^{\infty} \left( \frac{1}{\ell_k^{(j)}} - \frac{1}{\ell_{k+1}^{(j)}} \right) \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^k \right).$$

2. By a similar argument of Theorem 4.1 (2), we have  $F_r(\beta)$  is differentiable with respect to  $\beta \in \mathbb{R}$ .
3. The proof is a direct consequence of Theorem 4.5 (2) and Gärtner-Ellis Theorem [2].  $\square$

**Corollary 4.6.** For  $k = 1, \mathbf{p}_1 = (p_1 \cdots p_d, 1, \dots, 1) \in \mathbb{N}^d$  and  $j = 1$ , then  $C = \left( 1 - \frac{1}{p_1 \cdots p_d} \right)$ ,  $S^{(1)} = \langle p_1 \cdots p_d \rangle = \{\ell_n^{(1)} = (p_1 \cdots p_d)^{n-1}\}_{n=1}^{\infty}$  and  $\gamma(S^{(1)}) = \frac{p_1 \cdots p_d}{p_1 \cdots p_d - 1}$ . These give the directional energy function associated to the sum  $S_{N_1 \times \dots \times N_d}^{\mathbf{p}_1}$  is

$$F_r(\beta) = \frac{2p_1 \cdots p_d - 1}{2p_1 \cdots p_d} \log(r(1-r)) + \frac{p_1 \cdots p_d - 1}{p_1 \cdots p_d} \log |v^T \cdot e_+|^2 + \log \Lambda_+ + \mathcal{G}(\beta),$$

where

$$\mathcal{G}(\beta) = \sum_{\ell=1}^{\infty} \frac{(p_1 \cdots p_d - 1)^2}{(p_1 \cdots p_d)^{\ell+1}} \log \left( 1 + \left( \frac{2 \cosh(h)}{|v^T \cdot e_+|^2} - 1 \right) \left( \frac{\Lambda_-}{\Lambda_+} \right)^{\ell} \right).$$

Which is coincides with the formula in Theorem 1.3.



**Example 4.7.** Figure 2 shows the free energy behaviour for the multidimensional case  $d = 2$  with  $j = 1$ ,  $\mathbf{p}_1 = (2, 3)$ ,  $\mathbf{p}_2 = (3, 5)$ ,  $\mathbf{p}_3 = (5, 7)$ ,  $\mathbf{p}_4 = (7, 11)$ ,  $\mathbf{p}_5 = (11, 2)$ ,  $S^{(1)} = S^{(2)} = \langle 2, 3, 5, 7, 11 \rangle$ ,  $\gamma(S^{(1)})^{-1} = \gamma(S^{(2)})^{-1} = \frac{16}{77}$  and  $C = (1 - \frac{1}{6})(1 - \frac{1}{15})(1 - \frac{1}{35})(1 - \frac{1}{77})(1 - \frac{1}{22})\frac{77}{16} = \frac{2261}{660}$ .

## REFERENCES

1. J. C. Ban, W. G. Hu, and G. Y. Lai, *On the entropy of multidimensional multiplicative integer subshifts*, Journal of Statistical Physics **182** (2021), no. 2, 1–20.
2. J. C. Ban, W. G. Hu, and G. Y. Lai, *Large Deviation Principle of Multidimensional Multiple Averages on  $\mathbb{N}^d$* , Indagationes Mathematicae (2021), to appear.
3. J. C. Ban, W. G. Hu, and S. S. Lin, *Pattern generation problems arising in multiplicative integer systems*, Ergodic Theory and Dynamical Systems **39** (2019), no. 5, 1234–1260.
4. R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, 1982.
5. J. Bourgain, *Double recurrence and almost sure convergence*, Journal für die reine und angewandte Mathematik **404** (1990), 140–161.
6. G. Brunet, *Dimensions of “self-affine sponges” invariant under the action of multiplicative integers*, arXiv preprint arXiv:2010.03230 (2020).
7. G. Carinci, J. R. Chazottes, C. Giardinà, and F. Redig, *Nonconventional averages along arithmetic progressions and lattice spin systems*, Indagationes Mathematicae **23** (2012), no. 3, 589–602.
8. J. R. Chazottes and F. Redig, *Thermodynamic formalism and large deviations for multiplication-invariant potentials on lattice spin systems*, Electronic Journal of Probability **19** (2014).
9. A. Dembo and O. Zeitouni, *LDP for finite dimensional spaces*, Large deviations techniques and applications, Springer, 2009, pp. 11–70.
10. R. S. Ellis, *Entropy, large deviations, and statistical mechanics*, vol. 1431, Taylor & Francis, 2006.
11. A. H. Fan, *Some aspects of multifractal analysis*, Geometry and Analysis of Fractals, Springer, 2014, pp. 115–145.
12. ———, *Multifractal analysis of weighted ergodic averages*, Advances in Mathematics **377** (2021), 107488.
13. A. H. Fan, L. M. Liao, and J. H. Ma, *Level sets of multiple ergodic averages*, Monatshefte für Mathematik **168** (2012), no. 1, 17–26.
14. A. H. Fan, J. Schmeling, and M. Wu, *Multifractal analysis of some multiple ergodic averages*, Advances in Mathematics **295** (2016), 271–333.

15. H. Furstenberg, Y. Katznelson, and D. Ornstein, *The ergodic theoretical proof of Szemerédi's theorem*, Bulletin of the American Mathematical Society **7** (1982), no. 3, 527–552.
16. H. O. Georgii, *Gibbs measures and phase transitions*, vol. 9, Walter de Gruyter, 2011.
17. B. Host and B. Kra, *Nonconventional ergodic averages and nilmanifolds*, Annals of Mathematics (2005), 397–488.
18. R. Kenyon, Y. Peres, and B. Solomyak, *Hausdorff dimension for fractals invariant under multiplicative integers*, Ergodic Theory and Dynamical Systems **32** (2012), no. 5, 1567–1584.
19. N. Frantzikinakis, *Some open problems on multiple ergodic averages*, arXiv preprint arXiv:1103.3808 (2011).
20. Y. Kifer, *Nonconventional limit theorems*, Probability theory and related fields **148** (2010), no. 1-2, 71–106.
21. Y. Kifer and S. R. S. Varadhan, *Nonconventional limit theorems in discrete and continuous time via martingales*, The Annals of Probability **42** (2014), no. 2, 649–688.
22. Y. Peres, J. Schmeling, S. Seuret, and B. Solomyak, *Dimensions of some fractals defined via the semigroup generated by 2 and 3*, Israel Journal of Mathematics **199** (2014), no. 2, 687–709.
23. Y. Peres and B. Solomyak, *Dimension spectrum for a nonconventional ergodic average*, Real Analysis Exchange **37** (2012), no. 2, 375–388.
24. M. Pollicott, *A nonlinear transfer operator theorem*, Journal of statistical physics **166** (2017), no. 3-4, 516–524.

(Jung-Chao Ban) DEPARTMENT OF MATHEMATICAL SCIENCES, NATIONAL CHENGCHI UNIVERSITY, TAIPEI 11605, TAIWAN, ROC.

MATH. DIVISION, NATIONAL CENTER FOR THEORETICAL SCIENCE, NATIONAL TAIWAN UNIVERSITY, TAIPEI 10617, TAIWAN. ROC.

*Email address:* jcbn@nccu.edu.tw

(Wen-Guei Hu) COLLEGE OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU, 610064, CHINA

*Email address:* wghu@scu.edu.cn

(Guan-Yu Lai) DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL YANG MING CHIAO TUNG UNIVERSITY, HSINCHU 30010, TAIWAN, ROC.

*Email address:* guanyu.am04g@g2.nctu.edu.tw