ZERO-SUM PARTITIONS OF ABELIAN GROUPS

SYLWIA CICHACZ 1 AND KAROL SUCHAN 2,1

ABSTRACT. The following problem has been known since the 80's. Let Γ be an Abelian group of order m (denoted $|\Gamma| = m$), and let t and $\{m_i\}_{i=1}^t$, be positive integers such that $\sum_{i=1}^t m_i = m - 1$. Determine when $\Gamma^* = \Gamma \setminus \{0\}$, the set of non-zero elements of Γ , can be partitioned into disjoint subsets $\{S_i\}_{i=1}^t$ such that $|S_i| = m_i$ and $\sum_{s \in S_i} s = 0$ for every $i, 1 \le i \le t$. Such a subset partition is called a zero-sum partition.

 $|I(\Gamma)| \neq 1$, where $I(\Gamma)$ is the set of involutions in Γ , is a necessary condition for the existence of zerosum partitions. In this paper we show that the additional condition of $m_i \ge 4$ for every $i, 1 \le i \le t$, is sufficient. Moreover, we present some applications of zero-sum partitions to irregular, magic- and anti-magictype labelings of graphs.

1. INTRODUCTION
1.1. Preliminaries. For standard terms and notation in graph theory, the reader is referred to the textbook by Diestel [9] and the monograph by Brandstädt et al. [3]. An introduction to magic- and anti-magic-type labelings can be found in the monograph by Bača et al. [2], and the fundamental results in abstract algebra that we use can be found in the textbook by Gallian [12].
Let Γ be an Abelian group of order m with the operation denoted by +. For convenience, we will denote

Let Γ be an Abelian group of order m with the operation denoted by +. For convenience, we will denote $\sum_{i=1}^{k} a$ by ka, the inverse of a by -a, and a + (-b) by a - b. Moreover, we will write $\sum_{a \in S} a$ for the sum of all elements in S. The identity element of Γ will be denoted by 0, and the set of non-zero elements of Γ by

$$\Gamma \cong \mathbb{Z}_{p_1^{\alpha_1}} \times \mathbb{Z}_{p_2^{\alpha_2}} \times \ldots \times \mathbb{Z}_{p_k^{\alpha_k}}, \text{with } m = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \ldots \cdot p_k^{\alpha_k},$$

 $\sum_{i=1}^{k} a \text{ by } ka, \text{ the inverse of } a \text{ by } -a, \text{ and } a + (-b) \text{ by } a - b.$ Horeover, we will write $\sum_{a \in S} a \text{ for enclosing of } a \text{ for enclosing of } a \text{ letters of enclosed of } b \text{ for enclosed of$

Recall that, for a prime number p, a group the order of which is a power of p is called a *p*-group. Given a finite Abelian group Γ , Sylow p-subgroup of Γ is the maximal subgroup L of Γ the order of which is a power of p. For example, by the fundamental theorem of Abelian groups, it is easy to see that any Abelian group Γ can be factorized as $\Gamma \cong L \times H$, where L is the Sylow 2-group of Γ and the order of H is odd.

¹AGH University of Science and Technology, al. A. Mickiewicza 30, 30-059 Krakow, Poland

²Universidad Diego Portales, Av. Ejército Libertador 441, 8370191 Santiago, Chile

E-mail addresses: cichacz@agh.edu.pl, karol.suchan@mail.udp.cl.

²⁰¹⁰ Mathematics Subject Classification. 05E16, 20K01, 05C25, 05C78.

Key words and phrases. Abelian group, zero-sum sets, irregular labeling, magic-type labeling, anti-magic-type labeling.

This work was partially supported by the Faculty of Applied Mathematics AGH UST statutory tasks within subsidy of Ministry of Science and Higher Education.

Let us denote the number of involutions in Γ by $|I(\Gamma)|$. Since any cyclic finite group of even order has exactly one involution, if e is the number of cyclic components in the factorization of Γ whose order is even, then $|I(\Gamma)| = 2^e - 1$.

Recall that the sum of all elements of a group Γ is equal to the sum of its involutions and the identity element. The following lemma is well known. The readers can consult [8] for a proof.

Lemma 1.1 ([8]). Let Γ be an Abelian group.

- If $|I(\Gamma)| = 1$, then $\sum_{g \in \Gamma} g = \iota$, where ι is the involution. - If $|I(\Gamma)| \neq 1$, then $\sum_{g \in \Gamma} g = 0$.

1.2. Main Problem. In 1981 Tannenbaum introduced the following problem of partitioning in Abelian groups.

Problem 1.2 ([18]). Let Γ be an Abelian group of order m. Let t be a positive integer and $\{m_i\}_{i=1}^t$ an integer partition of m-1. Let $\{w_i\}_{i=1}^t$ be arbitrary elements of Γ (not necessary distinct). Determine when the elements of Γ^* can be partitioned into subsets $\{S_i\}_{i=1}^t$ (i.e., subsets S_i are pairwise disjoint and their union is Γ^*) such that $|S_i| = m_i$ and $\sum_{s \in S_i} s = w_i$ for every $i, 1 \le i \le t$.

If a respective subset partition of Γ^* exists, then we say that $\{m_i\}_{i=1}^t$ is *realizable* in Γ^* with $\{w_i\}_{i=1}^t$. We will not indicate Γ^* or $\{w_i\}_{i=1}^t$ explicitly when it is clear from the context. Note that realizability of $\{m_i\}_{i=1}^t$ implies that $\sum_{i=1}^t w_i = \sum_{g \in \Gamma} g$.

Note that we use sequences $\{m_i\}_{i=1}^t$ only for ease of presentation, whereas we interpret them as multisets. In other words, only the values present in the sequence and their multiplicities are relevant for a sequence to be realizable, not their order.

Since, throughout the paper, we often consider subsets of fixed cardinalities, let us present an abbreviated notation. Given a set, any of its subsets of cardinality k is called a k-subset.

The case most studied in the literature is the Zero-Sum Partition (ZSP) problem, i.e., when $w_i = 0$ for every $i, 1 \le i \le t$. (A respective subset partition of Γ^* is called a zero-sum partition of Γ .)

In general, in the context of any subset Z of Γ , we will say that $\{S_i\}_{i=1}^t$, a partition of Z into zero-sum subsets with $|S_i| = m_i$ for every $i, 1 \le i \le t$, realizes $\{m_i\}_{i=1}^t$ in Z.

Note that, by Lemma 1.1, an Abelian group Γ with $|I(\Gamma)| = 1$ does not admit zero-sum partitions (for every t and $\{m_i\}_{i=1}^t$). Moreover, if $\{m_i\}_{i=1}^t$ is realizable in Γ^* , then necessarily $m_i \ge 2$ for every $i, 1 \le i \le t$. It was proved that this condition is sufficient only if $|I(\Gamma)| = 0$ [18, 20] or $|I(\Gamma)| = 3$ [20]. I.e., in these cases, for every positive integer t and integer partition $\{m_i\}_{i=1}^t$ of m-1 with $m_i \ge 2$ for every $i, 1 \le i \le t, \{m_i\}_{i=1}^t$ is realizable in Γ^* (with $\{w_i\}_{i=1}^t$ where $w_i = 0$ for every $i, 1 \le i \le t$). In [6], we generalized this condition with the following definition.

Definition 1.3. Let Γ be a finite Abelian group of order m. We say that Γ has x-Zero-Sum Partition Property (x-ZSPP) if, for every positive integer t and integer partition $\{m_i\}_{i=1}^t$ of m-1 with $m_i \geq x$ for every $i, 1 \leq i \leq t$, there exists a subset partition $\{S_i\}_{i=1}^t$ of Γ^* with $|S_i| = m_i$ and $\sum_{s \in S_i} s = 0$ for every i, $1 \leq i \leq t$.

For every finite Abelian group Γ such that $|I(\Gamma)| = 0$ (equivalently, the order of Γ is odd), a stronger version of 2-ZSPP was proved. Let us give a brief explanation.

We call a 6-subset C of an Abelian group Γ good if $C = \{c, d, -c - d, -c, -d, c + d\}$ for some c and d in Γ . Notice that the sum of elements of a good 6-subset is 0. Moreover, it can be partitioned into three zero-sum 2-subsets or two zero-sum 3-subsets.

The following definition was given by Tannenbaum [18].

Definition 1.4. Let Γ be a finite Abelian group of order m = 6k + s for a non-negative integer k and $s \in \{1,3,5\}$. A partition of Γ^* into k good 6-subsets and (s-1)/2 zero-sum 2-subsets is called a Skolem partition of Γ^* .

Note that, if Γ is an Abelian group of order m such that every integer partition $\{m_i\}_{i=1}^t$ of m-1 with $m_i \in \{2,3\}$ for every $i, 1 \leq i \leq t$, is realizable in Γ^* , then also every integer partition $\{m_i'\}_{i=1}^{t'}$ of m-1 with $m'_i \geq 2$ for every $i, 1 \leq i \leq t'$, is realizable. In a similar way, it is easy to see that if a Skolem partition of Γ^*

exists, then every integer partition $\{m_i\}_{i=1}^t$ of m-1 with $m_i \in \{2,3\}$ for every $i, 1 \le i \le t$, is realizable. So the following theorem by Tannenbaun indeed offers a stronger version of 2-ZSPP.

Theorem 1.5 ([18]). Let Γ be a finite Abelian group such that $|I(\Gamma)| = 0$, then Γ^* has a Skolem partition.

The following theorem was first conjectured by Kaplan et al. [14] (they also showed the necessity), and later proved by Zeng [20].

Theorem 1.6 ([20]). Let Γ be a finite Abelian group of order m. Γ has 2-ZSPP if and only if $|I(\Gamma)| \in \{0,3\}$.

The above theorem confirms for the cases of $|I(\Gamma)| \in \{0,3\}$ the following conjecture stated by Tannenbaum.

Conjecture 1.7 ([18]). Let Γ be a finite Abelian group of order m with $|I(\Gamma)| > 1$. Let $R = \Gamma^* \setminus I(\Gamma)$. For every positive integer t and integer partition $\{m_i\}_{i=1}^t$ of m-1 with $m_i \ge 2$ for every i, $1 \le i \le |R|/2$, and $m_i \ge 3$ for every i, $|R|/2 + 1 \le i \le t$, there is a subset partition $\{S_i\}_{i=1}^t$ of Γ^* such that $|S_i| = m_i$ and $\sum_{s \in S_i} s = 0$ for every i, $1 \le i \le t$.

It was shown independently by a few authors that the conjecture is also true for $\Gamma \cong (\mathbb{Z}_2)^n$ with n > 1(notice that in this case $I(\Gamma) = \Gamma^*$, so $R = \emptyset$).

Theorem 1.8 ([4, 10, 19]). Let $\Gamma \cong (\mathbb{Z}_2)^n$ with n > 1, then Γ has 3-ZSPP.

Note that, in general, even the weaker version of Conjecture 1.7 posed in [5] is still open.

Conjecture 1.9 ([5]). Let Γ be a finite Abelian group with $|I(\Gamma)| > 1$. Then Γ has 3-ZSPP.

For every Abelian group Γ with more than one involution and large enough order, first Cichacz and Tuza [7] showed that it has 4-ZSPP, and later Müyesser and Pokrovskiy showed that it indeed has 3-ZSPP [16]. This result was improved by Cichacz and Suchan [6] for every Γ of order 2^n for some integer n > 1 such that $I(\Gamma) \neq 1$ with the following theorem.

Theorem 1.10 ([6]). Let Γ be such that $|I(\Gamma)| > 1$ and $|\Gamma| = 2^n$ for some integer n > 1. Then Γ has 3-ZSPP.

Recall that, for every Abelian group Γ , there exist groups L and H, with $|L| = 2^{\eta}$ for a nonnegative integer η and $|H| = \rho$ for an odd positive integer ρ , such that $\Gamma \cong L \times H$. Moreover, a finite Abelian group Γ has no involutions if and only if its order is odd. So, if $\eta = 0$, then Γ has 2-ZSPP by Theorem 1.6. On the other hand, if $\rho = 1$, then Γ has 3-ZSPP by Theorem 1.10.

Let Γ be an Abelian group such that $\Gamma \cong L \times H$, where $|L| = 2^{\eta}$ and $|H| = \rho$ for a positive integer η and an odd positive integer ρ , with $\rho > 1$. The main contribution of this paper is to show that Γ has 3-ZSPP if $(\rho \mod 6) \in \{1,3\}$. Moreover, for $(\rho \mod 6) = 5$, we show that Γ has 4-ZSPP. Therefore we improve the result from [7] and, towards proving Conjecture 1.9, we leave open only the question if, for $(\rho \mod 6) = 5$, Γ has not only 4-ZSPP, but also 3-ZSPP. We also present a new infinite family of groups for which Conjecture 1.7 holds. As a complement, to illustrate relations of zero-sum group partitioning to graph theory, we present some applications in irregular, magic-type, and anti-magic-type graph labeling.

Our proofs use as base cases zero-sum partitions of $((\mathbb{Z}_2)^3 \times \mathbb{Z}_3)^*$ and $((\mathbb{Z}_2)^3 \times \mathbb{Z}_5)^*$. They were analyzed by a computer program that we created, and sample zero-sum partitions that certify that the corresponding sequences are realizable are given in the annexes.

2. Main Result

Let $\operatorname{Bij}(\Gamma)$ denote the set of all bijections from Γ to itself. A *complete mapping* of a group A is defined as $\varphi \in \operatorname{Bij}(A)$ such that the mapping $\theta \colon g \mapsto g^{-1}\varphi(g)$ is also bijective (some authors refer to θ , rather than φ , as the complete mapping) [15].

In the proof of our main result we will use the following property of complete mappings:

Lemma 2.1 ([5, 20]). Let Γ be a finite Abelian group such that $|I(\Gamma)| \neq 1$. Then there exist $\phi, \varphi \in Bij(\Gamma)$ (not necessarily distinct) such that $g + \phi(g) + \varphi(g) = 0$ for every $g \in \Gamma$. In particular, we may assume that $\phi(0) = \varphi(0) = 0$.

In the following, we will consider integer partitions $\{m_i\}_{i=1}^t$ of m-1 for some positive integer t, with $m_i \geq 3$ for every $i, 1 \leq i \leq t$. Given such a sequence, since $m_i \geq 3$ for every $i, 1 \leq i \leq t$, we can modify the sequence $\{m_i\}_{i=1}^t$ by subdividing each term larger than 5 into a combination of terms 3, 4, and 5. It is easy to check that if the non-zero elements of a group Γ can be partitioned into zero-sum subsets of cardinalities corresponding to the elements of the new sequence, then the same holds for the old sequence. So we can focus only on sequences with $m_i \in \{3, 4, 5\}$ for every $i, 1 \leq i \leq t$.

Let Z be a subset of Γ . Let \mathcal{K} be a zero-sum partition of Z with cardinalities in $\{3, 4, 5\}$, and let $a = |\{S \in \mathcal{K} : |S| = 3\}|, b = |\{S \in \mathcal{K} : |S| = 4\}|$ and $c = |\{S \in \mathcal{K} : |S| = 5\}|$. In this situation, we say that \mathcal{K} realizes (a, b, c) in Z. If there exists a family realizing (a, b, c) in Z, we say that (a, b, c) is realizable in Z. Note that a triple (a, b, c) can be seen as a compact representation of a sequence $\{m_i\}_{i=1}^t$ in which a elements are equal to 3, b elements are equal to 4, and c elements are equal to 5.

Recall that the quotient group Γ modulo H for a subgroup H of Γ is denoted by Γ/H .

The following three theorems present the main results of the paper.

Theorem 2.2. Let Γ be a finite Abelian group such that $\Gamma \cong L \times H$, with $|L| = 2^{\eta}$ for some positive integer η , |I(L)| > 1, and $|H| \equiv 1 \pmod{6}$. Then Γ has 3-ZSPP.

Proof. By Theorem 1.6, we can assume that $I(L) \ge 7$ and, by Theorem 1.10, that |H| > 1.

By Theorem 1.5, there exists a partition of H^* into good 6-subsets:

$$H = \{0\} \cup \left(\bigcup_{i=1}^{|H^*|/6} \{b_i, c_i, -b_i - c_i, -b_i, -c_i, b_i + c_i\}\right).$$

By Theorem 1.10, L has 3-ZSPP and there are $\phi, \varphi \in \text{Bij}(L)$ such that $a + \phi(a) + \varphi(a) = 0$ for every $a \in L$ by Lemma 2.1. Thus

$$\Gamma^* = \bigcup_{a \in L^*} \{(a,0)\} \cup \left(\bigcup_{a \in L} \bigcup_{i=1}^{|H^*|/6} \{(a,b_i), (\phi(a),c_i), (\varphi(a),-b_i-c_i), (-a,-b_i), (-\phi(a),-c_i), (-\varphi(a),b_i+c_i)\} \right),$$

where the 6-subsets are good. Let $M^* = \bigcup_{a \in L^*} \{(a, 0)\}$, note that $M^* \cong L^*$, thus it has 3-ZSPP. Let

$$W = \bigcup_{a \in L} \bigcup_{i=1}^{|H^*|/6} \{(a, b_i), (\phi(a), c_i), (\varphi(a), -b_i - c_i), (-a, -b_i), (-\phi(a), -c_i), (-\varphi(a), b_i + c_i)\}$$

We will prove that any triple (α, β, γ) such that $3\alpha + 4\beta + 5\gamma = |\Gamma| - 1$ is realizable in Γ^* . Assume that $r_1, \ldots, r_{\alpha} = 3$, $r_{\alpha+1}, \ldots, r_{\alpha+\gamma} = 5$, and $r_{\alpha+\gamma+1}, \ldots, r_{\alpha+\beta+\gamma} = 4$. Let l be such that $\sum_{i=1}^{l-1} r_i \leq |L^*|$ and $\sum_{i=1}^{l} r_i > |L^*|$. Let $r'_l = |L^*| - \sum_{i=1}^{l-1} r_i$ and $r''_l = r_l - r'_l$. If $(r'_l = 0 \text{ or } r'_l \geq 3)$ and $(r''_l = 0 \text{ or } r''_l \geq 2)$, then the sequence $r_1, \ldots, r_{l-1}, r'_l$ is realized by a zero-sum partition $A_1, \ldots, A_{l-1}, A'_l$ of M^* by Theorem 1.10, and the sequence $r''_l, r_{l+1}, r_{l+2}, \ldots, r_{\alpha+\beta+\gamma}$ is realized by a zero-sum partition $A''_l, A_{l+1}, A_{l+2}, \ldots, A_{\alpha+\beta+\gamma}$ of W since W is the union of good subsets and $|W| = \sum_{i=l+1}^{\alpha+\beta+\gamma} r_i + r''_l$. So we are done. Hence we have to settle the cases where $r''_l = 1$ or $r'_l = 1$ or $r'_l = 2$.

Case 1 $r_l'' = 1$.

Then l is even and r_{l+1} is odd, since $|L^*|$ is odd and |W| is even (thus $|W| - r''_l$ requires at least one odd term among $r_{l+1}, \ldots, r_{\alpha+\beta+\gamma}$, and all odd terms are put at the beginning of the sequence). Suppose first $r_l = 3$, then $r'_l = 2$ and since $|I(L)| \ge 7$ and |L| is even, there is $|L^*| > 7$. This implies that $r_{l-1} = r_{l-2} = 3$. Note that because $|W| \equiv 0 \pmod{3}$ it has to be $\gamma \ge 1$ or $\beta \ge 2$. If $\gamma \ge 1$, then (l-2,0,1) is realizable in M^* and $(\alpha - l + 2, \beta, \gamma - 1)$ is realizable in W. For $\beta \ge 2$ the triple (l - 3, 2, 0) is realizable in M^* and $(\alpha - l + 3, \beta - 2, \gamma)$ is realizable in W. Suppose that $r_l = 5$. If $\beta > 0$, then the triple $(\alpha, 1, l - \alpha - 1)$ is realizable in M^* and $(0, \beta - 1, \gamma + \alpha - l + 1)$ is realizable in W.

Assume now that $\beta = 0$. If $|L^*| > 7$, then for $\alpha \ge 2$ the triple $(\alpha - 2, 0, l - \alpha + 1)$ is realizable in M^* and $(2, 0, \gamma + \alpha - l - 1)$ is realizable in W. For $\alpha \le 1$ there is $r_{l-1} = 5$ and $(\alpha + 3, 0, l - \alpha - 2)$ is realizable in M^* and $(1, 2, \gamma + \alpha - l)$ is realizable in W. Observe that in this case, by the construction of W, the two zero-sum 4-subset in W can be split into four zero-sum 2-subsets and we obtain a realization

of (α, β, γ) in Γ^* . If $|L^*| = 7$ then $\alpha = 1$ and $L \cong (\mathbb{Z}_2)^3$ what implies $\Gamma \cong (\mathbb{Z}_2)^3 \times H$ for $|H| \ge 13$, $\gamma = (|\Gamma^*| - 4)/5 = (8|H| - 4)/5 \ge 20$. Take the following three zero-sum 5-subsets:

They cover M^* and four inverse pairs of elements from W. W' obtained by the deletion of these 15 elements from W is of the following form (let $i_1 = (1, 0, 0), i_2 = (0, 1, 0)$ and $i_3 = (0, 0, 1)$):

$$W' = \bigcup_{a \in \{i_1, i_2, i_3\}} \{(\phi(a), c_1), (\varphi(a), -b_1 - c_1), (-\phi(a), c_1), (-\varphi(a), b_1 + c_1)\}$$
$$\cup \left(\bigcup_{a \in (\mathbb{Z}_2)^3 \setminus \{i_1, i_2, i_3\}} \{(a, b_1), (\phi(a), c_1), (\varphi(a), -b_1 - c_1), (-a, -b_1), (-\phi(a), -c_1), (-\varphi(a), b_1 + c_1)\}\right)$$
$$\cup \left(\bigcup_{a \in (\mathbb{Z}_2)^3} \bigcup_{i=2}^{|H^*|/6} \{(a, b_i), (\phi(a), c_i), (\varphi(a), -b_i - c_i), (-a, -b_i), (-\phi(a), -c_i), (-\varphi(a), b_i + c_i)\}\right).$$

Note there are $8\frac{|H|-4}{3}$ zero-sum 3-subsets in W', thus the sequence $((\gamma - 2)/2, \gamma - 3, 0)$ is realizable in W'. A zero-sum 4-subset in W' can be split into two zero-sum 2-subsets and we obtain a realization of $(1, 0, \gamma)$ in Γ^* .

Case 2 $r'_l = 2$.

If $r_l = 3$, then we are done as in Case 1. Therefore, we can assume $r_l = 5$ (for $r_l = 4$ we got a contradiction, since |W| is divisible by 4). For $\alpha \ge 1$, we obtain that $(\alpha - 1, 0, l)$ is realizable in M^* and $(1, \beta, \gamma - l)$ is realizable in W. For $\beta \ge 1$, since $|L^*| \ge 7$, there exists a realization of (1, 1, l - 2) in M^* and a realization of $(1, \beta, \gamma - l)$ in W. By the construction of W, one zero-sum 4-subset in W can be split into two zero-sum 2-subset and we obtain a realization of $(0, \beta, \gamma)$ in Γ^* . The only missing case is $\alpha = \beta = 0$. If $|L^*| > 7$, then actually $|L^*| \ge 15$ and $r_{l-1} = r_{l-2} = 5$. Thus (4, 0, l - 3) is realizable in M^* . Since W is a union of good 6-subsets, there exists a partition into four zero-sum 2-subsets and $(\gamma - l - 1)$ zero-sum 5-subsets.

Assume now that $|L^*| = 7 = |I(L)|$. But then, as in Case 1, there is $\Gamma \cong (\mathbb{Z}_2)^3 \times H$ and we proceed the same way.

Case 3. $r'_l = 1$.

Then l > 1, and $r_l' \ge 2$ is even. Assume first that $r_l = 3$. Since $|W| \equiv 0 \pmod{3}$, there is $\beta + \gamma > 0$. If now $\beta > 0$, then there exists a realization of (l - 2, 1, 0) in M^* and a realization of $(\alpha - l + 2, \beta - 1, \gamma)$ in W. Therefore, we can assume that $\beta = 0$. If now $\gamma = 1$, then $|W| \equiv 1 \pmod{3}$, a contradiction. Thus, for $\beta = 0$, there is $\gamma \ge 2$. If $|L^*| > 7$, then $|L^*| \ge 15$ and $r_{l-1} = r_{l-2} = r_{l-3} = 3$ and (l - 4, 0, 2) is realizable in M^* , and $(\alpha - l + 4, 0, \gamma - 2)$ is realizable in W. For $|L^*| = 7 = |I(L)|$, there is $\Gamma \cong (\mathbb{Z}_2)^3 \times H$. The set

$$\bigcup_{a \in ((\mathbb{Z}_2)^3)^*} \{(a,0)\} \cup \left(\bigcup_{a \in (\mathbb{Z}_2)^3} \{(a,b_1), (\phi(a),c_1), (\varphi(a),-b_1-c_1), (-a,-b_1), (-\phi(a),-c_1), (-\varphi(a),b_1+c_1)\} \right)$$

can be partitioned into 15 zero-sum 3-subsets and 2 zero-sum 5-subsets in the following way:

$$\begin{split} &\{(0,0,1,0),(0,0,0,b_1),(0,0,1,-b_1)\}, \quad \{(1,0,0,0),(0,0,1,b_1),(1,0,1,-b_1)\}, \\ &\{(1,1,1,0),(0,1,1,b_1),(1,0,0,-b_1)\}, \quad \{(0,1,0,b_1),(0,0,0,c_1),(0,1,0,-b_1-c_1)\}, \\ &\{(1,0,1,b_1),(0,0,1,c_1),(1,0,0,-b_1-c_1)\}, \quad \{(0,1,0,0),(0,1,0,c_1),(0,0,0,-c_1)\}, \\ &\{(0,0,0,b_1+c_1),(0,1,0,-c_1),(0,1,0,-b_1)\}, \quad \{(0,0,1,b_1+c_1),(0,0,1,-c_1),(0,0,0,-b_1)\}, \\ &\{(0,1,0,b_1+c_1),(1,0,0,-c_1),(1,1,0,-b_1)\}, \quad \{(1,0,0,b_1+c_1),(0,1,1,-c_1),(1,1,1,-b_1)\}, \\ &\{(1,0,0,b_1),(1,1,1,c_1),(0,1,1,-b_1-c_1)\}, \quad \{(1,0,1,b_1+c_1),(1,1,0,-c_1),(0,1,1,-b_1-c_1)\}, \\ &\{(1,0,1,0),(1,0,1,c_1),(0,1,1,b_1+c_1),(1,1,0,-b_1-c_1),(1,0,1,-c_1)\}, \\ &\{(1,0,1,0),(1,0,1,c_1),(0,1,1,b_1+c_1),(1,1,0,-b_1-c_1)\}, \\ &\{(1,0,1,0),(1,0,1,c_1),(0,1,1,b_1+c_1),(1,0,0,-b_1-c_1),(1,0,1,-c_1)\}, \\ &\{(1,0,1,0),(1,0,0,c_1),(1,0,0,-b_1-c_1)\}, \\ &\{(1,0,1,0),(1,0,0,c_1),(1,0,0,-b_1-c_1)\}, \\ &\{(1,0,1,0),(1,0,0,c_1),(1,0,0,-b_1-c_1),(1,0,0,-c_1)\}, \\$$

$$\{(1,1,0,0), (1,1,0,b_1+c_1), (1,1,1,b_1+c_1), (0,0,0,-b_1-c_1), (1,1,1,-b_1-c_1)\}$$

Since the set $\bigcup_{a \in (\mathbb{Z}_2)^3} \bigcup_{i=2}^{|\mathcal{I}|^{-1}} \{(a, b_i), (\phi(a), c_i), (\varphi(a), -b_i - c_i), (-a, -b_i), (-\phi(a), -c_i), (-\varphi(a), b_i + c_i)\}$ is a union

of good 6-subsets, we obtain that $(\alpha, 0, \gamma)$ is realizable in Γ^* .

Suppose now that $r_l = 5$. If now $r_{l-1} = 5$, then let $r'_{l-1} = r_{l-1} - 2 \ge 3$ and $r''_{l-1} = 2$. Let us re-define $r'_l := 3$ and $r''_l := r_l - 3$. The sequence $r_1, \ldots, r_{l-2}, r'_{l-1}, r'_l$ is realized by a zero-sum partition $A_1, \ldots, A_{l-2}, A'_{l-1}, A'_l$ of M^* by Theorem 1.10, and the sequence $r''_{l-1}, r''_l, r_{l+1}, \ldots, r_{\alpha+\beta+\gamma}$ is realized by a zero-sum partition A''_{l-1}, A''_l of M^* by Theorem 1.10, and the sequence $r''_{l-1}, r''_l, r_{l+1}, \ldots, r_{\alpha+\beta+\gamma}$ is realized by a zero-sum partition $A''_{l-1}, A''_l, A_{l+1}, \ldots, A_{\alpha+\beta+\gamma}$ of W. Thus $r_{l-1} = 3$. If now $\beta > 0$, then there exists a realization of $(\alpha - 1, 1, 0)$ in $\bigcup_{a \in L^*} \{(a, 0)\}$ and a realization of $(1, \beta - 1, \gamma)$ in W. If $|L^*| > 7$, then $|L^*| \ge 15$ and $r_{l-1} = r_{l-2} = r_{l-3} = 3$. Since $\beta = 0$ and $\gamma \ge 2$, we obtain that $(\alpha - 3, 0, 2)$ is realizable in M^* , and $(3, 0, \gamma - 2)$ is realizable in W. Let $|L^*| = 7 = |I(L)|$. Then $L \cong (\mathbb{Z}_2)^3$, $\alpha = 2$ and we proceed as above to obtain a realization of the sequence $(2, 0, \gamma)$.

We will need now the following lemma (see Lemma 2.4 in [6]):

Lemma 2.3 ([6]). Let $\Gamma \cong A \times \mathbb{Z}_{2^{n_1}}$ be an Abelian group such that |A| = 4n, |I(A)| > 1, $n_1 \in \{1, 2\}$, and n is a positive integer. Let a, b, c be non-negative integers with $3a + 4b + 5c = |\Gamma| - 1$ and $b \ge (2^{n_1} - 1)n$. If there exists a subset partition \mathcal{K} realizing $(a, b - (2^{n_1} - 1)n, c)$ in A^* , then (a, b, c) is realizable in Γ^* .

Theorem 2.4. Let Γ be a finite Abelian group such that $\Gamma = L \times H$ such that $|L| = 2^{\eta}$ for some natural number η , |I(L)| > 1 and $|H| \equiv 3 \pmod{6}$. Then Γ has the 3-Zero-Sum Partition Property.

Proof. By Theorem 1.6, we can assume that $|I(L)| \ge 7$. By Theorem 1.10, L has 3-ZSPP and there are $\phi, \varphi \in \text{Bij}(L)$ such that $a + \phi(a) + \varphi(a) = 0$ for every $a, a \in L$, and $\phi(0) = \varphi(0) = 0$ by Lemma 2.1.

Case 1. |H| > 3.

By Theorem 1.5 there exists a Skolem partition of H^* :

$$H = \{0, b, -b\} \cup \left(\bigcup_{i=1}^{(|H^*|-2)/6} \{b_i, c_i, -b_i - c_i, -b_i, -c_i, b_i + c_i\}\right)$$

Let

$$W = \bigcup_{a \in L} \bigcup_{i=1}^{(|H^*|-2)/6} \{(a, b_i), (\phi(a), c_i), (\varphi(a), -b_i - c_i), (-a, -b_i), (-\phi(a), -c_i), (-\varphi(a), b_i + c_i)\}$$

and $M^* = \bigcup_{a \in L^*} \{(a, 0)\}.$

Thus

$$\Gamma^* = \{(0,b), (0,-b)\} \cup \left(\bigcup_{a \in L^*} \{(a,0), (\phi(a),b), (\varphi(a),-b)\}\right) \cup W.$$

We will prove that any triple (α, β, γ) such that $3\alpha + 4\beta + 5\gamma = |\Gamma| - 1$ is realizable in Γ^* . If $\alpha + \gamma \ge |L|$, since (|L|, 0, 0) is realizable in $\bigcup_{a \in L^*} \{(a, 0), (\phi(a), b), (\varphi(a), -b)\}$ and W is a union of good 6-subsets, then (α, β, γ) is realizable in Γ^* . Thus we assume that $\alpha + \gamma < |L|$. Observe that

$$\Gamma^* = M^* \cup \left(\bigcup_{a \in L} \{(a, b), (-a, -b)\} \right) \cup W.$$

Since $|L^*| \equiv 3 \pmod{4}$, $M^* \cong L^*$, the sequence $(1, (|L^*| - 3)/4, 0)$ is realizable in M^* and the sequence $(\alpha + \gamma - 1, (|L|(|H^*| - 2) - 3(\alpha + \gamma - 1))/4, 0)$ is realizable in W since $\alpha + \gamma - 1 \leq |L|, |W| = |L|(|H^*| - 2)$, and $\alpha + \gamma - 1$ is even. Because in the set $\bigcup_{a \in L} \{(a, b), (-a, -b)\}$ there are |L| zero-sum 2-subsets, there exists

a realization of (α, β, γ) in Γ^* .

Case 2. |H| = 3.

Thus $H \cong \mathbb{Z}_3$ and

$$\Gamma^* = M^* \cup \left(\bigcup_{a \in L} \{ (a, 1), (-a, 2) \} \right).$$

Assume first that $((\mathbb{Z}_2)^3 \times \mathbb{Z}_3)^*$. By Theorem 1.6, $(\mathbb{Z}_2)^2 \times \mathbb{Z}_3$ has 2-ZSPP. The triples (a, b, c) with b < 3 were analyzed by a computer program we created, sample realizations can be found in the annexes. Therefore, for $A = (\mathbb{Z}_2)^2 \times \mathbb{Z}_3$ and $n_1 = 1$, by Lemma 2.3, we obtain that every triple (a, b, c) is realizable in $((\mathbb{Z}_2)^3 \times \mathbb{Z}_3)^*$.

We will prove now that any triple (α, β, γ) such that $3\alpha + 4\beta + 5\gamma = |\Gamma| - 1$ is realizable in $\Gamma^* \ncong ((\mathbb{Z}_2)^3 \times \mathbb{Z}_3)^*$. If $3(\alpha + \gamma) \le |L^*|$, then for $|L^*| - 3(\alpha + \gamma) \equiv 0 \pmod{4}$ the sequence $(\alpha + \gamma, (|L^*| - 3(\alpha + \gamma)))/(4, 0)$ is realizable in M^* by Theorem 1.10. We will obtain γ zero-sum 5-subsets by taking γ zero-sum 2-subsets from the set $\bigcup_{a \in L} \{(a, 1), (-a, 2)\}$. If $|L^*| - 3(\alpha + \gamma) \equiv 2 \pmod{4}$ and $\gamma > 0$, then the sequence

 $(\alpha + \gamma - 1, (|L^*| - 3(\alpha + \gamma) - 2)/4, 1)$ is realizable in M^* by Theorem 1.10, and we proceed as before. For $\gamma = 0$, there is $\alpha = 1$ or $\alpha \ge 3$ since $|\Gamma^*| \equiv 3 \pmod{4}$. Since also $|L^*| \equiv 3 \pmod{4}$, if $\alpha = 1$, then the sequence $(1, (|L^*| - 3)/4, 0)$ is realizable in M^* , and the remaining zero-sum 4-subsets can be found in the set $\bigcup_{a \in L} \{(a, 1), (-a, 2)\}$. For $\alpha \ge 3$ the sequence $(\alpha - 2, (|L^*| - 3\alpha + 6)/4, 0)$ is realizable in M^* . Take one zero-sum

3-subset from this realization, say $T = \{(t_1, 0), (t_2, 0), (t_3, 0)\}$ then those 3-subsets: $\{(t_1, 0), (t_2, 1), (t_3, 2)\}, \{(t_2, 0), (t_3, 1), (t_1, 2)\}$ and $\{(t_3, 0), (t_1, 1), (t_2, 2)\}$ are zero-sum in Γ and the remaining zero-sum 4-subsets can be found in $\bigcup_{a \in U \setminus \{t_1, t_2, t_3\}} \{(a, 1), (-a, 2)\}.$ Hence we are done.

 $a \in L \setminus \{t_1, t_2, t_3\}$

The only missing case is $3(\alpha + \gamma) > |L^*|$. Since $L \not\cong (\mathbb{Z}_2)^3$, then there exists a subgroup B of Γ such that $|B| = 2^{\eta-2} = |L|/4$, |I(B)| > 1 and $\Gamma/B \cong \mathbb{Z}_2 \times \mathbb{Z}_6$. Since $\mathbb{Z}_2 \times \mathbb{Z}_6 = \{(0,0), (0,2), (0,4)\} \cup \{(0,3), (1,3), (1,0)\} \cup \{\{(0,1), (1,4), (1,1), (0,5), (1,2), (1,5)\}$, we can choose a set of coset representatives for the subgroup B in Γ , say A, such that

$$A = \{0\} \cup \{a, -a\} \cup \{e_1, e_2, e_1 + e_2\} \cup \{c, d, -c - d, -c, -d, c + d\},\$$

where $2e_1, 2e_2 \in A$. Since |I(B)| > 1, there exist $\phi, \varphi \in \text{Bij}(B)$ such that $g + \phi(g) + \varphi(g) = 0$ for every $g \in B$ by Lemma 2.1. Moreover, without loss of generality, we can assume that $\phi(0) = \varphi(0) = 0$. Note that $|B| = |\Gamma|/12 = |L|/4$. If now $\alpha + \gamma < |L|/2 - 1$, then write

$$\Gamma^* = B^* \cup \left(\bigcup_{g \in B} \{a + g, -a - g\} \cup \bigcup_{g \in B} \{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\} \right) \\ \cup \left(\bigcup_{g \in B} \{c + g, d + \phi(g), -c - d + \varphi(g), -c - g, -d - \phi(g), c + d - \varphi(g)\} \right),$$

where the latter 6-subsets are good.

If now $|B^*|$ is divisible by 3, then the sequence $(|B^*|/3, 0, 0)$ is realizable in B^* by Theorem 1.10 and (|B|, 0, 0) is realizable in $\bigcup_{g \in B} \{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\}$. Since $|B^*|/3 + |B| = (|L| - 1)/3 < \alpha + \gamma$, we can find the remaining $\alpha + \gamma - (|L| - 1)/3 \le 2|B|$ zero-sum 3-subsets in

$$\left(\bigcup_{g\in B} \left\{c+g, d+\phi(g), -c-d+\varphi(g), -c-g, -d-\phi(g), c+d-\varphi(g)\right\}\right).$$

Note that we are left now only with zero-sum 2-subsets, thus we are done.

If $|B^*|$ is not divisible by 3, then since $|B^*| \equiv 3 \pmod{4}$ there is $|B^*| \equiv 7 \pmod{12}$ or $|B^*| \equiv 11 \pmod{12}$. Moreover $\beta > 1$ because $|\Gamma| \not\equiv 0 \pmod{5}$ and $\alpha + \gamma < |L|/2 - 1 < (3|L| - 1)/5 = |\Gamma^*|/5$. Thus the sequence $((|B^*| - 4)/3, 1, 0)$ is realizable in B^* for $|B^*| \equiv 7 \pmod{12}$ and $((|B^*| - 8)/3, 2, 0)$ is realizable in B^* for $|B^*| \equiv 11 \pmod{12}$ by Theorem 1.10 and we proceed as above.

For $\alpha + \gamma \geq |L|/2 - 1$ set

$$\begin{split} \Gamma^* = & \{a, -a\} \cup \left(\bigcup_{g \in B^*} \{a + g, \phi(g), -a + \varphi(g)\} \cup \bigcup_{g \in B} (\{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\} \right) \\ & \cup \left(\bigcup_{g \in B} \{c + g, d + \phi(g), -c - d + \varphi(g), -c - g, -d - \phi(g), c + d - \varphi(g)\} \right), \end{split}$$

The sequence (|B| - 1, 0, 0) is realizable in $\bigcup_{g \in B^*} \{a + g, \phi(g), -a + \varphi(g)\}$ and (|B|, 0, 0) is realizable in $\bigcup_{g \in B} \{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\}$. Since $|B| - 1 + |B| = |L|/2 - 1 \le \alpha + \gamma$, we can find the remaining $\alpha + \gamma - |L|/2 + 1$ zero-sum 3-subsets in

$$\left(\bigcup_{g\in B} \left\{c+g, d+\phi(g), -c-d+\varphi(g), -c-g, -d-\phi(g), c+d-\varphi(g)\right\}\right).$$

Note that we are left now only with zero-sum 2-subsets, thus we are done.

Theorem 2.5. Let Γ be a finite Abelian group such that $\Gamma = L \times H$ such that $|L| = 2^{\eta}$ for some natural number η , |I(L)| > 1 and $|H| \equiv 5 \pmod{6}$. Then Γ has the 4-Zero-Sum Partition Property.

Proof. By Theorem 1.6, we can assume that $|I(L)| \ge 7$. By Theorem 1.10, the group L has 3-ZSPP and there are $\phi, \varphi \in \operatorname{Bij}(L)$ such that $a + \phi(a) + \varphi(a) = 0$ for every $a \in L$ and $\phi(0) = \varphi(0) = 0$ by Lemma 2.1. Let $r_1 + \ldots + r_t = |\Gamma| - 1$. Assume that r_1, \ldots, r_s are all odd, and r_{s+1}, \ldots, r_t are all even. Then s is odd because $|\Gamma|$ is even. We may restrict ourselves to sequences where $4 \le r_i \le 7$ for every i. We refine the sequence r_1, \ldots, r_s to $r'_1 = \ldots = r'_s = 3$. Note that $s < |\Gamma|/5$.

Case 1. |H| > 5.

By Theorem 1.5, there exists a Skolem partition of H^* :

$$H = \{0, b, -b, c, -c\} \cup \left(\bigcup_{i=1}^{(|H^*|-4)/6} \{b_i, c_i, -b_i - c_i, -b_i, -c_i, b_i + c_i\} \right).$$

Let

$$W = \bigcup_{a \in L} \bigcup_{i=1}^{(|H^*|-4)/6} \{(a, b_i), (\phi(a), c_i), (\varphi(a), -b_i - c_i), (-a, -b_i), (-\phi(a), -c_i), (-\varphi(a), b_i + c_i)\}$$

and $M^* = \bigcup_{a \in L^*} \{(a, 0)\}$. Thus

$$\Gamma^* = \{(0,b), (0,-b)\} \cup \left(\bigcup_{a \in L^*} \{(a,0), (\phi(a),b), (\varphi(a),-b)\}\right) \cup \left(\bigcup_{a \in L} \{(a,c), (-a,-c)\}\right) \cup W.$$

If $s \ge |L^*|$, then we can realize $|L^*|$ zero-sum 3-subsets in $\bigcup_{a \in L^*} \{(a, 0), (\phi(a), b), (\varphi(a), -b)\}$. Since $s < |\Gamma|/5$, the remaining $s - |L^*|$ zero-sum 3-subsets can be realized in W, the remaining elements in Γ^* form zero-sum

the remaining $s - |L^*|$ zero-sum 3-subsets can be realized in W, the remaining elements in 1^{-*} form zero-sum 2-subsets, so we are done.

Assume now that $s < |L^*|$. Note that we can write Γ^* also in the following way:

$$\Gamma^* = M^* \cup \left(\bigcup_{a \in L} \{ (a, b), (-a, -b) \} \right) \cup \left(\bigcup_{a \in L} \{ (a, c), (-a, -c) \} \right) \cup W.$$

Since $|L^*| \equiv 3 \pmod{4}$, $L^* \cong \bigcup_{a \in L^*} \{(a,0)\}$, the sequence $(1, (|L^*| - 3)/4, 0)$ is realizable in M^* by Theorem 1.10, and the sequence $(s-1, (|L|(|H^*| - 4) - 3(s-1))/4, 0)$ is realizable in W since $s-1 \leq |L|$, $|W| = |L|(|H^*| - 4)$ and s-1 is even. Because there exist 2|L| zero-sum 2-subsets in the set $\left(\bigcup_{a \in L} \{(a,b), (-a,-b)\}\right) \cup \left(\bigcup_{a \in L} \{(a,c), (-a,-c)\}\right)$, there exists the desired subset partition of Γ^* .

Case 2. |H| = 5. Thus $H \cong \mathbb{Z}_5$ and

$$\Gamma^* = M^* \cup \left(\bigcup_{a \in L} \bigcup_{b \in \mathbb{Z}_5^*} \{ (a, b), (-a, -b) \} \right).$$

Assume first that $((\mathbb{Z}_2)^3 \times \mathbb{Z}_5)^*$. By Theorem 1.6, $(\mathbb{Z}_2)^2 \times \mathbb{Z}_5$ has 2-ZSPP. The triples (a, b, c) with b < 5 were analyzed by a computer program we created, sample realizations can be found in the annexes. Therefore, for $A = (\mathbb{Z}_2)^2 \times \mathbb{Z}_5$ and $n_1 = 1$, by Lemma 2.3, we obtain that every triple (a, b, c) is realizable in $((\mathbb{Z}_2)^3 \times \mathbb{Z}_5)^*$.

From now we assume that $L \not\cong (\mathbb{Z}_2)^3$. If $3s \leq |L^*|$, then, for $|L^*| - 3s \equiv 0 \pmod{4}$, the sequence $(s, (|L^*| - 3s)/4, 0)$ is realizable in $M^* \cong L^*$ by Theorem 1.10. We will obtain s zero-sum subsets of odd cardinality by taking zero-sum 2-subsets from the set $\bigcup \bigcup \{(a,b), (-a,-b)\}$. If $|L^*| - 3s \equiv 2 \pmod{4}$, $a \in L \ b \in \mathbb{Z}_5^*$

then the sequence $(s-1, (|L^*|-3s-2)/4, 1)$ is realizable in M^* by Theorem 1.10, and we proceed as before. The missing case is $3s > |L^*|$. Since $L \ncong (\mathbb{Z}_2)^3$, there exists a subgroup B of Γ such that |B| = $2^{\eta-2} = |L|/4, |I(B)| > 3 \text{ and } \Gamma/B \cong \mathbb{Z}_2 \times \mathbb{Z}_{10}. \text{ Since } \mathbb{Z}_2 \times \mathbb{Z}_{10} = \{(0,0), (0,5), (1,5), (1,0)\} \cup \{(0,2), (0,8)\} \cup \{(0,2), (0,2)\} \cup \{(0,2), (0,2)\}$ $\{(1,2),(1,8)\} \cup \{(0,1),(1,3),(1,6),(0,9),(1,7),(1,4)\} \cup \{(1,1),(0,3),(0,6),(1,9),(0,7),(0,4)\} we can choose (1,2),(1,3),(1,4),(1,3),(1,6),(1,7),(1,4)\} \cup \{(1,1),(0,3),(0,6),(1,9),(0,7),(0,4)\}$ a set of coset representatives for the subgroup B in Γ , say A, such that

$$A = \{a, 0, -a\} \cup \{b, -b\} \cup \{e_1, e_2, e_1 + e_2\} \cup \bigcup_{i=1,2} \{c_i, d_i, -c_i - d_i, -c_i, -d_i, c_i + d_i\},\$$

where $2e_1, 2e_2 \in A$. Since |I(B)| > 1, there are $\phi, \varphi \in Bij(B)$ such that $g + \phi(g) + \varphi(g) = 0$ for every $g \in B$ by Lemma 2.1. Moreover without loss of generality we can assume that $\phi(0) = \varphi(0) = 0$. If now s < |L|/2 - 1, then write

$$\begin{split} \Gamma^* = & \left(\bigcup_{g \in B^*} \{g\} \right) \cup \left(\bigcup_{g \in B} \{a + g, -a - g\} \right) \cup \left(\bigcup_{g \in B} \{b + g, -b - g\} \right) \\ & \cup \left(\bigcup_{g \in B} \{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\} \right) \\ & \cup \left(\bigcup_{g \in B} \{c_i + g, d_i + \phi(g), -c_i - d_i + \varphi(g), -c_i - g, -d_i - \phi(g), c_i + d_i - \varphi(g)\} \right), \end{split}$$

where the latter 6-subsets are good.

If now $|B^*|$ is divisible by 3, then the sequence $(|B^*|/3, 0, 0)$ is realizable in $\bigcup_{g \in B^*} \{g\}$ by Theorem 1.10 and (|B|, 0, 0) is realizable in $\bigcup_{g \in B} \{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\}$. Since $|B^*|/3 + |B| = (|L| - 1)/3 < \alpha + \gamma$, we can find the remaining $\alpha + \gamma - (|L| - 1)/3 \le 2|B|$ zero-sum 3-subsets in

$$\left(\bigcup_{g\in B} \left\{c+g, d+\phi(g), -c-d+\varphi(g), -c-g, -d-\phi(g), c+d-\varphi(g)\right\}\right)$$

Note that we are left now only with zero-sum 2-subsets, thus we are done.

If $|B^*|$ is not divisible by 3, then the sequence $((|B^*| - 4)/3, 1, 0)$ is realizable in $\bigcup_{g \in B^*} \{g\}$ for $|B^*| \equiv 7$ (mod 12) and (($|B^*| - 8$)/3, 2, 0) is realizable in $\bigcup_{g \in B^*} \{g\}$ for $|B^*| \equiv 11 \pmod{12}$ by Theorem 1.10 and we proceed as above.

For $s \geq |L|/2 - 1$ set

$$\begin{split} \Gamma^* = & \{a, -a\} \cup \left(\bigcup_{g \in B^*} \{a + g, \phi(g), -a + \varphi(g)\} \right) \cup \left(\bigcup_{g \in B} \{b + g, -b - g\} \right) \\ & \cup \left(\bigcup_{g \in B} \{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\} \right) \\ & \cup \left(\bigcup_{\substack{g \in B \\ i = 1, 2}} \{c_i + g, d_i + \phi(g), -c_i - d_i + \varphi(g), -c_i - g, -d_i - \phi(g), c_i + d_i - \varphi(g)\} \right), \end{split}$$

where the latter 6-subsets are good.

Since $|B| = |\Gamma|/20 = |L|/4$ the sequence (|L|/2 - 1, 0, 0) is realizable in $\left(\bigcup_{g \in B^*} \{a + g, \phi(g), -a + \varphi(g)\}\right)$

 $\cup \left(\bigcup_{g \in B} \{e_1 + g, e_2 + \phi(g), e_1 + e_2 + \varphi(g)\} \right).$ Since s < |L| and $0 \le s - |L|/2 + 1$, we can find the remaining $s - |L|/2 + 1 \le |L|/2 = 2|B|$ zero-sum 3-subsets in

$$\bigcup_{\substack{g \in B \\ i=1,2}} \left\{ c_i + g, d_i + \phi(g), -c_i - d_i + \varphi(g), -c_i - g, -d_i - \phi(g), c_i + d_i - \varphi(g) \right\}.$$

Note that we are left now only with zero-sum 2-subsets, thus we are done.

Putting the main results together, we obtain the following corollary.

Corollary 2.6. Every finite Abelian group Γ has the 4-Zero-Sum Partition Property if and only if $|I(\Gamma)| \neq 1$.

Proof. If $|I(\Gamma)| = 1$, then Γ does not have x-ZSPP for any x because $\sum_{g \in \Gamma} g \neq 0$.

Assume now that $|I(\Gamma)| \neq 1$. By the fundamental theorem of finite Abelian groups, the group $\Gamma \cong L \times H$ such that $|L| = 2^{\eta}$ for some natural number η , $|I(L)| \neq 1$ and |H| is odd. If $\eta = 0$ then we are done by Theorem 1.6 since $|I(\Gamma)| = 0$. Assume now that $\eta > 0$. So |I(L)| > 1, and we can apply Theorems 2.2, 2.4 or 2.5.

Corollary 2.7. Let m be an integer with $m \ge 4$. Then one of the following cases hold:

- (1) $m \mod 4 = 0$ and there exists an Abelian group Γ of order m that has 4-ZSPP.
- (2) $m \mod 4 \in \{1, 3\}$ and every Abelian group Γ of order m has 2-ZSPP.
- (3) $m \mod 4 = 2$ and every Abelian group Γ of order m does not have x-ZSPP for any x.

Proof. Note that, given a positive integer m, if $m \equiv 2 \pmod{4}$, then every group Γ of order m has exactly one involution and so does not have x-ZSPP for any x. But, if $m \ge 4$ and $m \not\equiv 2 \pmod{4}$, then there exists an Abelian group Γ of order m that has 4-ZSPP. Indeed, if $m \mod 4 \in \{1,3\}$, then m is odd. In this case \mathbb{Z}_m has no involutions, thus has 2-ZSPP by Theorem 1.6. If $m \mod 4 = 0$, then $\Gamma = (\mathbb{Z}_2)^2 \times \mathbb{Z}_{m/4}$ has at least 3 involutions, thus has 4-ZSPP by Corollary 2.6.

3. New family of groups

Recall that Theorems 1.6 and 1.8 present cases where Conjecture 1.7 holds. Here we present a new infinite family of groups where it also is the case:

Theorem 3.1. Let $\Gamma \cong (\mathbb{Z}_2)^{2\eta} \times H$ for some natural number η and $|H| \equiv 1 \pmod{6}$. Let $R = \Gamma^* \setminus I(\Gamma)$. For any positive integer t and an integer partition $\{m_i\}_{i=1}^t$ of m-1, with $m_i \ge 2$ for every $i, 1 \le i \le |R|/2$, and $m_i \ge 3$ for every $i, |R|/2 + 1 \le i \le t$, there is a subset partition $\{S_i\}_{i=1}^t$ of Γ^* such that $|S_i| = m_i$ and $\sum_{s \in S_i} s = 0$ for every $i, 1 \le i \le t$.

Proof. By Theorem 1.5 there exists a partition of H^* into good 6-subsets:

$$H^* = \bigcup_{i=1}^{|H^*|/6} \{b_i, c_i, -b_i - c_i, -b_i, -c_i, b_i + c_i\}$$

Let $L = (\mathbb{Z}_2)^{2\eta}$, observe that $I(\Gamma) \cong L^*$. By Theorem 1.6 we can assume that I(L) > 3 and hence $\eta \ge 2$. By Theorem 1.8 the group L has 3-ZSPP and there are $\phi, \varphi \in \text{Bij}(L)$ such that $a + \phi(a) + \varphi(a) = 0$ for every $a \in L$ by Lemma 2.1. Thus

$$\Gamma^* = \bigcup_{a \in L^*} \{(a,0)\} \cup \left(\bigcup_{a \in L} \bigcup_{i=1}^{|H^*|/6} \{(a,b_i), (\phi(a),c_i), (\varphi(a),-b_i-c_i), (-a,-b_i), (-\phi(a),-c_i), (-\varphi(a),b_i+c_i)\} \right),$$

where the latter 6-subsets are good. Let $M^* = \bigcup_{a \in L^*} \{(a, 0)\}$, as before M^* has 3-ZSPP since $L^* \cong M^*$. Let

$$W = \bigcup_{a \in L} \bigcup_{i=1}^{|H^*|/6} \{ (a, b_i), (\phi(a), c_i), (\varphi(a), -b_i - c_i), (-a, -b_i), (-\phi(a), -c_i), (-\varphi(a), b_i + c_i) \}.$$

Observe that $2^{2\eta} - 1 \equiv 0 \pmod{3}$ for every natural number η , W = R and $|W| \equiv 0 \pmod{12}$.

Analogously to realizable triples, we define realizable quadruples. Let X be a subset of Γ . Let \mathcal{K} be a family of pairwise disjoint zero-sum subsets of X with cardinalities in $\{2, 3, 4, 5\}$, and let $\omega = |\{S \in \mathcal{K} : |S| = 2\}|$, $\alpha = |\{S \in \mathcal{K} : |S| = 3\}|, \beta = |\{S \in \mathcal{K} : |S| = 4\}|$ and $\gamma = |\{S \in \mathcal{K} : |S| = 5\}|$, with $2\omega + 3\alpha + 4\beta + 5\gamma = |X|$ (so \mathcal{K} is a subset partition of X). In this situation, we say that \mathcal{K} realizes the quadruple $(\omega, \alpha, \beta, \gamma)$ in X. If there exists a family realizing $(\omega, \alpha, \beta, \gamma)$ in X, we say that $(\omega, \alpha, \beta, \gamma)$ is realizable in X.

We will prove that any quadruple $(\omega, \alpha, \beta, \gamma)$ such that $2\omega + 3\alpha + 4\beta + 5\gamma = |\Gamma| - 1, \omega \le |R|/2$ is realizable in Γ^* . Assume that $r_1, \ldots, r_{\alpha} = 3, r_{\alpha+1}, \ldots, r_{\alpha+\gamma} = 5, r_{\alpha+\gamma+1}, \ldots, r_{\alpha+\beta+\omega} = 2$ and $r_{\alpha+\gamma+\omega+1}, \ldots, r_{\alpha+\beta+\omega+\gamma} = 4$. Let l be such that $\sum_{i=1}^{l-1} r_i \le |L^*|$ and $\sum_{i=1}^{l} r_i > |L^*|$. Let $r'_l = |L^*| - \sum_{i=1}^{l-1} r_i$ and $r''_l = r_l - r'_l$. If $(r'_l = 0 \text{ or } r'_l \ge 3)$ and $(r''_l = 0 \text{ or } r''_l \ge 2)$, then the sequence $r_1, \ldots, r_{l-1}, r'_l$ is realized by a zero-sum partition $A_1, \ldots, A_{l-1}, A'_l$ of $M^* = \bigcup_{\alpha \in L^*} \{(a, 0)\} \cong L^*$ by Theorem 1.10 and the sequence $r''_l, r_{l+1}, r_{l+2}, \ldots, r_{\alpha+\beta+\gamma+\omega}$

is realized by a zero-sum partition $A_l'', A_{l+1}, A_{l+2}, \ldots, A_{\alpha+\beta+\gamma+\omega}$ of W, so we are done. Hence we have to settle the cases where $r_l'' = 1$ or $r_l' = 1$ or $r_l' = 2$.

Case 1 $r_l'' = 1$.

Then l is even and $r_{l-1} = r_l = 5$ since $|L^*| \equiv 3 \pmod{6}$ is odd and |W| is even. The sequence $(0, \alpha + 2, 0, l-2)$ is realizable in M^* and $(\omega + 2, 0, \beta, \gamma - l + 2)$ is realizable in W.

Case 2 $r'_l = 2$.

Since $|L^*| \equiv 0 \pmod{3}$ and $|W| \equiv 0 \pmod{12}$, we can assume $r_{l-1} = r_l = r_{l+1} = 5$. For $\alpha \ge 1$ we obtain that $(0, \alpha - 1, 0, l)$ is realizable in M^* and $(\omega, 1, \beta, \gamma - l)$ is realizable in W. For $\beta \ge 1$, since $|L^*| \ge 7$, there exists a realization of (0, 1, 1, l-2) in M^* and a realization of $(\omega + 2, 1, \beta - 1, \gamma - l)$ in W. The only missing case is $\alpha = \beta = 0$. Then $|L^*| \ge 15$ and $r_{l-1} = r_{l-2} = 5$, thus (0, 4, 0, l-3) is realizable in M^* and $(\omega + 4, 0, 0, \gamma - l - 1)$ in W.

Case 3.
$$r'_{l} = 1$$
.

Then l > 1, $r_l'' = 4$ and $r_{l-1} = 5$. The sequence $(0, \alpha+2, 0, l-2)$ is realizable in M^* , and $(\omega+2, 0, \beta, \gamma-l+2)$ is realizable in W.

4. Some Applications

In this section we present some applications of Corollary 2.6.

4.1. Group irregular labeling for digraphs. Let $\overrightarrow{G} = (V, A)$ be a digraph on the set of vertices V with the set of arcs A. If $\psi: A \to \Gamma$ is a labeling of arcs of \overrightarrow{G} with elements of Γ such that $\varphi_{\psi}: V \to \Gamma$, the labeling of vertices of \overrightarrow{G} with elements of Γ defined by

$$\varphi_{\psi}(x) = \sum_{y \in N^-(x)} \psi((y,x)) - \sum_{y \in N^+(x)} \psi((x,y)),$$

is injective, then we say that ψ is a Γ -irregular labeling of \vec{G} .

Recall that, given a digraph $\overrightarrow{G} = (V, A)$, its underlying graph is the graph G = (V, E) in which $\{x, y\} \in E$ if and only if $(x, y) \in A$ or $(y, x) \in A$. Given a digraph \overrightarrow{G} and its underlying graph G, the subdigraph \overrightarrow{C} of \overrightarrow{G} induced by the vertex set of a connected component C in G is called a *weakly connected component* of \overrightarrow{G} . Thus we have $\overrightarrow{C} = \overrightarrow{G}[V(C)]$.

We have the following result:

Lemma 4.1 ([7]). A digraph $\overrightarrow{G} = (V, A)$ with no isolated vertices has a Γ -irregular labeling if and only if there exists an injective mapping φ from V to Γ such that $\sum_{x \in V(\overrightarrow{C})} \varphi(x) = 0$ for every weakly connected component \overrightarrow{C} of \overrightarrow{G} .

Cichacz and Tuza also showed the following:

Theorem 4.2 ([7]). Any digraph \overrightarrow{G} of order n with no weakly connected components of order less than 3 has a Γ -irregular labeling for every Γ such that $|\Gamma| \ge 2n + 2\sqrt{n-1/2} - 1$.

They also showed the following:

Lemma 4.3 ([7]). Let \overrightarrow{G} be a digraph of order n with no weakly connected component of order less than 3, and let Γ be a finite Abelian group such that $|\Gamma| - |I(\Gamma)| \ge 2n$. Then \overrightarrow{G} has a Γ -irregular labeling.

By Corollary 2.6, we obtain the following:

Corollary 4.4. Any digraph \vec{G} of order n with no weakly connected components of cardinality less than 4 has a Γ -irregular labeling for every Γ such that $|I(\Gamma)| \neq 1$ and $|\Gamma| \geq n+5$.

Proof. Let $\{\overrightarrow{C}_i\}_{i=1}^t$ be the weakly connected components of \overrightarrow{G} . By Lemma 4.1, there exists a Γ -irregular labeling of a digraph \overrightarrow{G} with weakly connected components $\{\overrightarrow{C}_i\}_{i=1}^t$ if and only if there exist in Γ pairwise disjoint subsets $\{S_i\}_{i=1}^t$ such that $|S_i| = |V(\overrightarrow{C}_i)|$ and $\sum_{s \in S_i} s = 0$ for every $i, 1 \le i \le t$. Let $m_i = |V(\overrightarrow{C}_i)|$ for every $i, 1 \le i \le t$, and let $m_{t+1} = |\Gamma^*| - n$. Since $m_{t+1} \ge 4$, using the sequence $\{m_i\}_{i=1}^{t+1}$, by Theorem 1.5 and Corollary 2.6, we get the result.

If we consider all Abelian groups, then by Corollary 4.4 and Lemma 4.3 one can easily see the following:

Corollary 4.5. Any digraph \overrightarrow{G} of order n with no weakly connected components of order less than 4 has a Γ -irregular labeling for every Γ such that $|\Gamma| \geq 2n + 1$.

We think that the results from Corollary 4.4 can be achieved for all Abelian groups and digraphs with no weakly connected components of order less than 3, therefore we finish this section with the following conjecture.

Conjecture 4.6. There exists a constant K such that any digraph \overrightarrow{G} of order n with no weakly connected components of order less than 3 has a Γ -irregular labeling for every Γ such that $|\Gamma| \ge n + K$.

4.2. Magic and anti-magic labeling in graphs with twins. Froncek in [11] defined the notion of a group distance magic labeling of a graph G = (V, E). In such a labeling, the vertices of the graph are labeled through a bijection with the elements of an Abelian group Γ . The weight of each vertex is computed as the sum (in Γ) of the labels assigned to its neighbors. If all weights are the same, then it is a Γ -distance magic labeling. If all weights are different, then it is a Γ -distance anti-magic labeling [1].

For ease of presentation of the results in this subsection we will create labelings using only the non-zero elements of Γ , and speak of Γ^* -distance magic labeling, etc. It is easy to see that these results can be amended to results on labelings with all elements of Γ by dealing with the 0 element accordingly.

Recall that in a graph G = (V, E), a set of vertices $M, M \subset V$, is a module if for every pair of vertices x, y, with $x, y \in M$, there is $N(x) \setminus M = N(y) \setminus M$. In other words, a module M is a set of vertices that share the same neighbors outside of M. Two vertices x and y, with $x, y \in V$, are called *twins* if $\{x, y\}$ is a module. If additionally $\{x, y\} \in E$, then they are *true twins*, and they are *false twins* otherwise. For

ease of presentation, we assume that every vertex x is both a false and true twin of itself (in the literature, the relation of being twins is usually considered only for distinct vertices). In this context, notice that the relation of being false twins is an equivalence relation, and the same holds for the relation of being true twins. So they define equivalence classes corresponding to inclusion maximal sets of (false or true) twins. Any set of (false or true) twins is a module.

Based on the above observations, we can create Γ^* -distance magic and anti-magic labelings for groups and graphs that satisfy certain conditions detailed in the following propositions.

Proposition 4.7. Let G = (V, E) be a graph with a partition of the vertex set into sets of false twins $\{V_i\}_{i=1}^t$ for a positive integer t. Let $m_i = |V_i|$ for every $i, 1 \le i \le t$. Let Γ be an Abelian group of order m such that $\sum_{i=1}^t m_i = m - 1$. If the non-zero elements of Γ can be partitioned into zero-sum sets $\{S_i\}_{i=1}^t$ such that $|S_i| = m_i$ for every $i, 1 \le i \le t$, then G has a Γ^* -distance magic labeling.

Proof. Let us assign to the vertices of every V_i , $1 \le i \le t$, unique elements of the corresponding S_i , as in the statement of the proposition. It can be done, since $|V_i| = |S_i|$. Let us show that, under this labeling, the weight of every vertex in V is 0.

Take any vertex $v, v \in V_a$, for any $a, 1 \leq a \leq t$. Consider the sum of labels of the neighbors of v. Consider any $b, b \neq a$ and $1 \leq b \leq t$. Since V_b is a module, either v is adjacent to no vertices in V_b or to all of them. In both cases, the contribution of $N(v) \cap V_b$ to the weight of v is 0.

The only set of neighbors of v left to consider is $N(v) \cap V_a$. v has no neighbors in V_a , since V_a is a set of false twins. So the contribution of $N(v) \cap V_a$ to the weight of v is also 0.

Proposition 4.8. Let G = (V, E) be a graph with a partition of the vertex set into sets of true twins $\{V_i\}_{i=1}^t$ for a positive integer t. Let $m_i = |V_i|$ for every $i, 1 \le i \le t$. Let Γ be an Abelian group of order m such that $\sum_{i=1}^t m_i = m - 1$. If the non-zero elements of Γ can be partitioned into zero-sum subsets $\{S_i\}_{i=1}^t$ such that $|S_i| = m_i$ for every $i, 1 \le i \le t$, then G has a Γ^* -distance anti-magic labeling.

Proof. Let us do the labeling as in the proof of Proposition 4.7. Like therein, take any vertex $v, v \in V_a$, for any $a, 1 \leq a \leq t$, and consider its weight (the sum of labels of the neighbors of v). Let γ be the element of Γ assigned to v.

Like in the proof of Proposition 4.7, for any $b, b \neq a$ and $1 \leq b \leq t$, the contribution of $N(v) \cap V_b$ to the weight of v is 0. But now V_a is a clique. The sum of the elements of Γ assigned to the vertices in V_a is 0. So the sum of the labels assigned to $N(v) \cap V_a$ is $-\gamma$, the element of Γ inverse to γ .

Since the assignment of non-zero elements of Γ to the elements of V is a bijection, by uniqueness of inverses, the sums of labels of neighbors of the vertices in V are unique too.

The following corollaries are direct consequences of Propositions 4.7 and 4.8, Corollary 2.6, and above mentioned properties of twins.

Corollary 4.9. Let G = (V, E) be a graph in which every vertex has at least 3 false twins. Let Γ be a finite Abelian group of order |V| + 1 such that $|I(\Gamma)| \neq 1$. Then G has a Γ^* -distance magic labeling.

Notice that Corollary 4.9 extends the results on Γ -distance magic labelings of complete k-partite graphs presented in [6].

Corollary 4.10. Let G = (V, E) be a graph in which every vertex has at least 3 true twins. Let Γ be a finite Abelian group of order |V| + 1 such that $|I(\Gamma)| \neq 1$. Then G has a Γ^* -distance anti-magic labeling.

For the last two results of this subsection, let us recall the definition of a G-join introduced by Sabidussi [17].

Definition 4.11 ([17]). Let G = (V, E) be a graph and $\{X_v\}_{v \in V}$ a collection of graphs indexed by V. The G-join of $\{X_v\}_{v \in V}$ is the graph H given by $V(H) = \{(x, v) \mid x \in X_v, v \in V\}$, $E(H) = \{\{(x, v), (x', v')\} \mid \{v, v'\} \in E, \text{ or } v = v' \text{ and } \{x, x'\} \in E(X_v)\}$.

Notice that, given a graph G = (V, E), in the graph H that is a G-join of $\{X_v\}_{v \in V}$, the set of vertices $\{(x, v) \mid x \in X_v\}$ for every $v, v \in V$, is a module in H. Moreover, if X_v is an empty graph, then the vertices of H in $\{(x, v) \mid x \in X_v\}$ are false twins. If X_v is a complete graph, then the vertices of H in $\{(x, v) \mid x \in X_v\}$ are true twins.

Given a vertex $v \in V$, let us use t(v) to denote the set of true twins of v and f(v) for the set of false twins of v. In such a context, the vertex v can be treated as the unique representative of the equivalence class t(v)or f(v), respectively.

Recall that X is an *empty graph* if $E(X) = \emptyset$. In other words, when its vertex set is stable. We use K_k to denote the complete graph on k vertices and $\overline{K_k}$ for the empty graph on k vertices.

Corollary 4.12. Let G = (V, E) be any graph. Let H be the G-join of a sequence of empty graphs of appropriate orders such that every vertex in H has at least 3 false twins. Let Γ be a finite Abelian group of order |V(H)| + 1 such that $|I(\Gamma)| \neq 1$. Then H has a Γ^* -distance magic labeling.

Proof. Let $\{f(v_i)\}_{i=1}^t$ be the equivalence classes of false twins in G with the corresponding unique representatives $\{v_i\}_{i=1}^t$. Notice that, if $|f(v_i)| \ge 4$ for every $i, 1 \le i \le t$, then we can simply take H = G to have H and Γ satisfy the conditions of Corollary 4.9. So, to define the graph H, we use the G-join of $\{X_v\}_{v\in V}$ where, for every $v_i, 1 \le i \le t$, we put $X_{v_i} = \overline{K_{p_i}}$, with $p_i = 4 - |f(v_i)|$. For every other vertex $v, v \in V \setminus \{v_i\}_{i=1}^t$, we put $X_v = \overline{K_1}$. It is easy to check that H and Γ obtained in this way satisfy the conditions of Corollary 4.9.

With a similar reasoning, we obtain the following corollary.

Corollary 4.13. Let G = (V, E) be any graph. Let H be the G-join of a sequence of complete graphs of appropriate orders such that every vertex in H has at least 3 true twins. Let Γ be a finite Abelian group of order |V(H)| + 1 such that $|I(\Gamma)| \neq 1$. Then H has a Γ^* -distance anti-magic labeling.

Note that the equivalence classes of twins in a graph can be found in linear time [13]. So if a respective zerosum partition of Γ^* is given, or it can be found efficiently, then we could efficiently compute the corresponding labeling efficiently.

Note that, by a reasoning similar to the ones used to prove Corollaries 4.9, 4.10, 4.12 and 4.13, and based on Corollary 2.7, the corresponding results can be made stronger: to require only one twin for every vertex if the order n of the corresponding graph satisfies $(n + 1) \mod 4 \in \{1, 3\}$, and using 2-ZSPP. In case $(n + 1) \mod 4 = 2$, such a Γ does not exist, but it can be "fixed" by adding an additional twin for one of the vertices, and thus reducing to the case $(n + 1) \mod 4 = 3$.

5. FINAL REMARKS

Recall that any group Γ can be factorized as $\Gamma \cong L \times H$, where L is the Sylow 2-group of Γ and the order of H is odd. In this context, towards proving Conjecture 1.9, we leave open only the question if Γ has not only 4-ZSPP, but also 3-ZSPP, in the case where $(|H| \mod 6) = 5$. On the other hand, however we presented some new groups for which Conjecture 1.7 holds, for many groups Γ with $|I(\Gamma)| > 3$ the conjecture is still open. In particular, for groups with $|I(\Gamma)| = 2^{2\eta+1} - 1 < |\Gamma^*|$ for any positive integer η , we know nothing.

Finally, let us complement Definition 1.4. Let S be a subset of cardinality m = 6k + s, for a positive integer k and $s \in \{0, 2, 4\}$, of a finite Abelian group Γ . A partition of S into k good 6-subsets and s/2zero-sum 2-subsets is called a *Skolem partition of S*. It is known that \mathbb{Z}_m , for m = 6k + s with any positive integer k, has a Skolem partition when $s \in \{0, 4\}$, but the situation is slightly different for s = 2. For $m \equiv 2$ or 8 (mod 24), there exists a Skolem partition, whereas for $m \equiv 14$ or 20 (mod 24), such a partition does not exist (see Lemma 4 in [18]). The use of good subsets in the proof of Theorem 3.1 motivates the following problem.

Problem 5.1. Characterize every finite Abelian group Γ such that $R = \Gamma^* \setminus I(\Gamma)$ has a Skolem partition.

A complete characterization of existence of Skolem partitions would be a useful tool towards proving the following conjecture. Note that doing it would settle Conjecture 1.7 completely.

Conjecture 5.2. Let Γ be a finite Abelian group of order m with $|I(\Gamma)| > 1$. Let $R = \Gamma^* \setminus I(\Gamma)$. If the set R has a Skolem partition, then for any positive integer t and an integer partition $\{m_i\}_{i=1}^t$ of m-1, with $m_i \ge 2$ for every i, $1 \le i \le |R|/2$, and $m_i \ge 3$ for every i, $|R|/2 + 1 \le i \le t$, there is a subset partition $\{S_i\}_{i=1}^t$ of Γ^* such that $|S_i| = m_i$ and $\sum_{s \in S_i} s = 0$ for every i, $1 \le i \le t$.

On the algorithmic side, given a finite Abelian group Γ of order m and an integer partition $\{m_i\}_{i=1}^t$ of m-1, it would be interesting to know the time complexity (and an efficient algorithm) of finding a zero-sum partition that realizes $\{m_i\}_{i=1}^t$ (if it exists).

References

- M. Anholcer, S. Cichacz, D. Froncek, R. Simanjuntak, and J. Qiu. Group distance magic and antimagic hypercubes. Discrete Mathematics, 344(12):112625, 2021.
- [2] M. Bača, M. Miller, J. Ryan, and A. Semaničová-Feňovčíková. Magic and Antimagic Graphs. Springer, 2019.
- [3] A. Brandstädt, V. B. Le, and J. P. Spinrad. Graph classes: a survey. SIAM, 1999.
- [4] L. Caccetta and R.-Z. Jia. Binary labeling of graphs. Graphs and Combinatorics, 13(2):119–137, 1997.
- [5] S. Cichacz. Zero sum partition of abelian groups into sets of the same order and its applications. The Electronic Journal of Combinatorics, 25(1):1–20, 2018.
- S. Cichacz and K. Suchan. Zero-sum partitions of abelian groups of order 2ⁿ. Discrete Mathematics & Theoretical Computer Science, 25(1):#6, 2023.
- [7] S. Cichacz and Z. Tuza. Realization of digraphs in abelian groups and its consequences. Journal of Graph Theory, 100(2):331–345, 2022.
- [8] D. Combe, A. M. Nelson, and W. D. Palmer. Magic labellings of graphs over finite abelian groups. Australasian Journal of Combinatorics, 29:259–272, 2004.
- [9] R. Diestel. Graph Theory, volume 173 of Graduate Texts in Mathematics. Springer, 2017.
- [10] Y. Egawa. Graph labelings in elementary abelian 2-groups. Tokyo Journal of Mathematics, 20(2):365–379, 1997.
- [11] D. Froncek. Group distance magic labeling of cartesian product of cycles. Australasian Journal of Combinatorics, 55:167– 174, 2013.
- [12] J. Gallian. Contemporary abstract algebra. Cengage Learning, 2016.
- [13] M. Habib and C. Paul. A survey of the algorithmic aspects of modular decomposition. Computer Science Review, 4(1):41–59, 2010.
- [14] G. Kaplan, A. Lev, and Y. Roditty. On zero-sum partitions and anti-magic trees. Discrete Mathematics, 309(8):2010–2014, 2009.
- [15] H. Mann. On orthogonal latin squares. Bull. Amer. Math. Soc., 50:249-257, 1944.
- [16] A. Müyesser and A. Pokrovskiy. A random hall-paige conjecture, 2022.
- [17] G. Sabidussi. Graph derivatives. Mathematische Zeitschrift, 76:385–401, 1961.
- [18] P. Tannenbaum. Partitions of abelian groups into sets with zero sums. Congressus Numerantium, 33:341–348, 1981.
- [19] P. Tannenbaum. Partitions of \mathbb{Z}_2^n . SIAM Journal on Algebraic Discrete Methods, 4(1):22–29, 1983.
- [20] X. Zeng. On zero-sum partitions of abelian groups. Integers, 15:Paper No. A44: 16 pp, 2015.

Throughout the annexes, the notation "a*3 b*4 c*5" refers to *a* subsets of size 3, *b* subsets of size 4, and *c* subsets of size 5. In each subset partition, we present one set per line. Each tuple represents one element of the partitioned set, with the elements of the tuple following the order of the direct product.

```
A. ZERO-SUM PARTITIONS OF ((\mathbb{Z}_2)^3 \times \mathbb{Z}_3)^* with b \leq 2
```

```
[2, 2, 2, 3]
[5, 2, 0]
Γ
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 2]],
[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 2]],
[[0, 1, 0, 1], [1, 0, 0, 1], [1, 1, 0, 1]],
[[0, 0, 1, 2], [0, 1, 0, 2], [0, 1, 1, 2]],
[[1, 0, 1, 1], [1, 1, 1, 1], [1, 0, 1, 2], [1, 1, 1, 2]],
[[1, 0, 1, 0], [1, 1, 0, 0], [0, 1, 1, 1], [0, 0, 0, 2]]
]
A partition for subsets of sizes: 5*3 2*4 0*5
[2, 2, 2, 3]
[6, 0, 1]
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 2]],
[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 2]],
[[0, 1, 0, 1], [1, 0, 0, 1], [1, 1, 0, 1]],
[[0, 0, 1, 2], [0, 1, 0, 2], [0, 1, 1, 2]],
[[1, 1, 0, 0], [0, 1, 1, 1], [1, 0, 1, 2]],
[[1, 0, 1, 0], [1, 0, 1, 1], [1, 1, 1, 1], [0, 0, 0, 2], [1, 1, 1, 2]]
]
A partition for subsets of sizes: 6*3 0*4 1*5
[2, 2, 2, 3]
[3, 1, 2]
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 2]],
[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 2]],
[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 0, 2], [0, 0, 1, 2]],
[[1, 1, 0, 0], [1, 0, 0, 1], [1, 1, 1, 1], [0, 1, 0, 2], [1, 1, 1, 2]],
[[1, 0, 1, 0], [1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 2], [1, 0, 1, 2]]
٦
A partition for subsets of sizes: 3*3 1*4 2*5
[2, 2, 2, 3]
[0, 2, 3]
Γ
[[0, 0, 1, 0], [0, 1, 0, 0], [1, 0, 0, 0], [1, 1, 1, 0]],
[[1, 0, 1, 0], [1, 1, 0, 0], [0, 0, 0, 1], [0, 1, 1, 2]],
[[0, 0, 1, 1], [0, 1, 0, 1], [0, 1, 1, 1], [1, 0, 0, 1], [1, 0, 0, 2]],
[[1, 1, 0, 1], [0, 0, 0, 2], [0, 0, 1, 2], [0, 1, 0, 2], [1, 0, 1, 2]],
[[0, 1, 1, 0], [1, 0, 1, 1], [1, 1, 1, 1], [1, 1, 0, 2], [1, 1, 1, 2]]
A partition for subsets of sizes: 0*3 2*4 3*5
```

```
[2, 2, 2, 3]
[1, 0, 4]
Γ
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [1, 0, 1, 0], [1, 1, 0, 0], [0, 0, 0, 1], [1, 1, 1, 2]],
[[0, 0, 1, 1], [0, 1, 0, 1], [0, 1, 1, 1], [1, 0, 0, 1], [1, 0, 0, 2]],
[[1, 1, 0, 1], [0, 0, 0, 2], [0, 0, 1, 2], [0, 1, 0, 2], [1, 0, 1, 2]],
[[1, 1, 1, 0], [1, 0, 1, 1], [1, 1, 1, 1], [0, 1, 1, 2], [1, 1, 0, 2]]
]
A partition for subsets of sizes: 1*3 0*4 4*5
                         B. ZERO-SUM PARTITIONS OF ((\mathbb{Z}_2)^3 \times \mathbb{Z}_5)^* with b \leq 4
[2, 2, 2, 5]
[13, 0, 0]
Г
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],
[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],
[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],
[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 3]],
[[0, 0, 1, 2], [0, 0, 0, 4], [0, 0, 1, 4]],
[[1, 0, 0, 2], [0, 1, 1, 4], [1, 1, 1, 4]],
[[1, 0, 0, 1], [0, 1, 0, 2], [1, 1, 0, 2]],
[[0, 1, 0, 3], [1, 1, 1, 3], [1, 0, 1, 4]],
[[1, 0, 0, 3], [1, 1, 0, 3], [0, 1, 0, 4]],
[[1, 1, 0, 0], [0, 1, 1, 2], [1, 0, 1, 3]],
[[1, 1, 1, 1], [0, 0, 0, 2], [1, 1, 1, 2]],
[[1, 0, 1, 0], [1, 0, 1, 2], [0, 0, 0, 3]]
]
A partition for subsets of sizes: 13*3 0*4 0*5
[2, 2, 2, 5]
[9, 3, 0]
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],
[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],
[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],
[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 3]],
[[0, 0, 1, 2], [0, 0, 0, 4], [0, 0, 1, 4]],
[[1, 0, 0, 2], [0, 1, 1, 4], [1, 1, 1, 4]],
[[1, 1, 1, 2], [0, 1, 0, 4], [1, 0, 1, 4]],
[[1, 0, 0, 1], [0, 1, 0, 2], [1, 1, 0, 2]],
[[0, 0, 0, 2], [0, 1, 1, 2], [1, 0, 0, 3], [1, 1, 1, 3]],
[[1, 1, 1, 1], [0, 0, 0, 3], [0, 1, 0, 3], [1, 0, 1, 3]],
[[1, 0, 1, 0], [1, 1, 0, 0], [1, 0, 1, 2], [1, 1, 0, 3]]
٦
A partition for subsets of sizes: 9*3 3*4 0*5
```

[2, 2, 2, 5][10, 1, 1] E . [[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 3]],[[0, 0, 1, 2], [0, 0, 0, 4], [0, 0, 1, 4]],[[1, 0, 0, 2], [0, 1, 1, 4], [1, 1, 1, 4]],[[1, 1, 1, 2], [0, 1, 0, 4], [1, 0, 1, 4]],[[1, 0, 0, 1], [0, 1, 0, 2], [1, 1, 0, 2]],[[1, 1, 0, 0], [0, 0, 0, 2], [1, 1, 0, 3]],[[1, 1, 1, 1], [0, 0, 0, 3], [0, 1, 0, 3], [1, 0, 1, 3]],[[1, 0, 1, 0], [0, 1, 1, 2], [1, 0, 1, 2], [1, 0, 0, 3], [1, 1, 1, 3]]٦ A partition for subsets of sizes: 10*3 1*4 1*5 [2, 2, 2, 5] [6, 4, 1]Γ [[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 3]],[[0, 0, 1, 2], [0, 0, 0, 4], [0, 0, 1, 4]],[[0, 1, 1, 2], [1, 0, 0, 2], [1, 0, 1, 2], [0, 1, 0, 4]],[[1, 0, 0, 1], [0, 0, 0, 3], [0, 1, 0, 3], [1, 1, 0, 3]],[[1, 1, 1, 1], [1, 1, 1, 2], [1, 0, 1, 3], [1, 0, 1, 4]],[[1, 1, 0, 0], [0, 1, 0, 2], [0, 1, 1, 4], [1, 1, 1, 4]],[[1, 0, 1, 0], [0, 0, 0, 2], [1, 1, 0, 2], [1, 0, 0, 3], [1, 1, 1, 3]]٦ A partition for subsets of sizes: 6*3 4*4 1*5 [2, 2, 2, 5][7, 2, 2] Е [[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 3]],[[0, 0, 1, 2], [0, 0, 0, 4], [0, 0, 1, 4]],[[1, 0, 0, 2], [0, 1, 1, 4], [1, 1, 1, 4]],[[1, 1, 0, 2], [1, 1, 1, 2], [1, 0, 0, 3], [1, 0, 1, 3]],[[1, 1, 1, 1], [0, 0, 0, 2], [0, 1, 0, 3], [1, 0, 1, 4]],[[0, 1, 1, 2], [0, 0, 0, 3], [1, 1, 0, 3], [1, 1, 1, 3], [0, 1, 0, 4]],[[1, 0, 1, 0], [1, 1, 0, 0], [1, 0, 0, 1], [0, 1, 0, 2], [1, 0, 1, 2]]٦ A partition for subsets of sizes: 7*3 2*4 2*5

[2, 2, 2, 5] [8, 0, 3] Γ [[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 3]],[[0, 0, 1, 2], [0, 0, 0, 4], [0, 0, 1, 4]],[[1, 0, 0, 2], [0, 1, 1, 4], [1, 1, 1, 4]],[[1, 1, 1, 2], [0, 1, 0, 4], [1, 0, 1, 4]],[[1, 0, 0, 1], [1, 1, 1, 1], [0, 0, 0, 2], [1, 0, 0, 3], [1, 1, 1, 3]],[[1, 1, 0, 0], [0, 1, 0, 2], [1, 1, 0, 2], [0, 0, 0, 3], [0, 1, 0, 3]],[[1, 0, 1, 0], [0, 1, 1, 2], [1, 0, 1, 2], [1, 0, 1, 3], [1, 1, 0, 3]]٦ A partition for subsets of sizes: 8*3 0*4 3*5 [2, 2, 2, 5] [4, 3, 3] [[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],[[1, 0, 1, 1], [1, 1, 0, 1], [1, 1, 1, 1], [1, 0, 0, 2]],[[0, 0, 1, 2], [0, 1, 0, 2], [0, 1, 1, 2], [0, 0, 0, 4]],[[1, 0, 1, 2], [1, 1, 0, 2], [0, 0, 0, 3], [0, 1, 1, 3]],[[1, 0, 0, 1], [0, 1, 0, 3], [1, 0, 0, 3], [0, 0, 1, 4], [0, 1, 1, 4]],[[0, 0, 0, 2], [1, 1, 1, 2], [1, 0, 1, 3], [1, 0, 1, 4], [1, 1, 1, 4]],[[1, 0, 1, 0], [1, 1, 0, 0], [1, 1, 0, 3], [1, 1, 1, 3], [0, 1, 0, 4]]٦ A partition for subsets of sizes: 4*3 3*4 3*5 [2, 2, 2, 5] [5, 1, 4] Γ [[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],[[0, 1, 0, 1], [0, 1, 1, 1], [0, 0, 1, 3]],[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 3]],[[0, 0, 0, 2], [0, 0, 1, 2], [0, 1, 0, 2], [0, 1, 1, 4]],[[1, 0, 0, 2], [1, 0, 1, 2], [0, 0, 0, 3], [0, 0, 0, 4], [0, 0, 1, 4]],[[1, 0, 0, 1], [0, 1, 0, 3], [1, 0, 0, 3], [1, 0, 1, 4], [1, 1, 1, 4]],[[1, 1, 0, 2], [1, 0, 1, 3], [1, 1, 0, 3], [1, 1, 1, 3], [0, 1, 0, 4]],[[1, 0, 1, 0], [1, 1, 0, 0], [1, 1, 1, 1], [0, 1, 1, 2], [1, 1, 1, 2]]٦ A partition for subsets of sizes: 5*3 1*4 4*5

```
[2, 2, 2, 5]
[1, 4, 4]
Г
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [1, 0, 1, 0], [1, 1, 0, 0], [1, 1, 1, 0]],
[[0, 0, 0, 1], [0, 0, 1, 1], [0, 1, 0, 1], [0, 1, 1, 2]],
[[1, 0, 0, 1], [1, 0, 1, 1], [1, 1, 0, 1], [1, 1, 1, 2]],
[[0, 0, 0, 2], [0, 0, 1, 2], [0, 1, 0, 2], [0, 1, 1, 4]],
[[1, 0, 0, 2], [1, 0, 1, 2], [0, 0, 0, 3], [0, 0, 0, 4], [0, 0, 1, 4]],
[[1, 1, 1, 1], [0, 1, 0, 3], [0, 1, 1, 3], [0, 1, 0, 4], [1, 0, 0, 4]],
[[1, 1, 0, 2], [0, 0, 1, 3], [1, 1, 0, 3], [1, 1, 1, 3], [1, 1, 0, 4]],
[[0, 1, 1, 1], [1, 0, 0, 3], [1, 0, 1, 3], [1, 0, 1, 4], [1, 1, 1, 4]]
٦
A partition for subsets of sizes: 1*3 4*4 4*5
[2, 2, 2, 5]
[2, 2, 5]
Γ
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],
[[1, 1, 1, 0], [0, 0, 1, 1], [0, 1, 0, 1], [1, 0, 0, 3]],
[[0, 1, 1, 1], [1, 0, 0, 1], [1, 0, 1, 1], [0, 1, 0, 2]],
[[1, 1, 1, 1], [0, 0, 0, 2], [0, 0, 1, 2], [0, 1, 1, 2], [1, 0, 1, 3]],
[[1, 0, 0, 2], [1, 0, 1, 2], [0, 0, 0, 3], [0, 0, 0, 4], [0, 0, 1, 4]],
[[1, 0, 1, 0], [0, 1, 0, 3], [0, 1, 0, 4], [0, 1, 1, 4], [1, 1, 0, 4]],
[[1, 1, 1, 2], [0, 1, 1, 3], [1, 1, 0, 3], [1, 1, 1, 3], [1, 0, 1, 4]],
[[1, 1, 0, 0], [1, 1, 0, 1], [1, 1, 0, 2], [0, 0, 1, 3], [1, 1, 1, 4]]
٦
A partition for subsets of sizes: 2*3 2*4 5*5
[2, 2, 2, 5]
[3, 0, 6]
Γ
[[0, 0, 1, 0], [0, 1, 0, 0], [0, 1, 1, 0]],
[[1, 0, 0, 0], [0, 0, 0, 1], [1, 0, 0, 4]],
[[1, 1, 1, 0], [0, 0, 1, 1], [1, 1, 0, 4]],
[[0, 1, 0, 1], [0, 1, 1, 1], [1, 0, 0, 1], [0, 0, 0, 3], [1, 0, 1, 4]],
[[1, 1, 1, 1], [0, 0, 0, 2], [0, 0, 1, 2], [0, 1, 0, 2], [1, 0, 0, 3]],
[[1, 0, 0, 2], [1, 0, 1, 2], [0, 1, 0, 3], [0, 0, 0, 4], [0, 1, 1, 4]],
[[1, 1, 0, 2], [0, 0, 1, 3], [0, 1, 1, 3], [1, 0, 1, 3], [0, 0, 1, 4]],
[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 2], [1, 1, 1, 2], [1, 1, 1, 4]],
[[1, 0, 1, 0], [1, 1, 0, 0], [1, 1, 0, 3], [1, 1, 1, 3], [0, 1, 0, 4]]
٦
A partition for subsets of sizes: 3*3 0*4 6*5
```

```
[2, 2, 2, 5]
[0, 1, 7]
[
[[0, 0, 1, 0], [0, 1, 0, 0], [1, 0, 0, 0], [1, 1, 1, 0]],
[[1, 0, 1, 0], [1, 1, 0, 0], [0, 0, 0, 1], [0, 0, 1, 1], [0, 1, 0, 3]],
[[0, 1, 0, 1], [0, 1, 1, 1], [1, 0, 0, 1], [0, 0, 0, 3], [1, 0, 1, 4]],
[[1, 1, 1, 1], [0, 0, 0, 2], [0, 0, 1, 2], [0, 1, 0, 2], [1, 0, 0, 3]],
[[1, 0, 0, 2], [1, 0, 1, 2], [0, 1, 1, 3], [0, 0, 0, 4], [0, 1, 0, 4]],
[[0, 1, 1, 0], [0, 0, 1, 3], [0, 0, 1, 4], [1, 0, 0, 4], [1, 1, 1, 4]],
[[1, 1, 1, 2], [1, 0, 1, 3], [1, 1, 0, 3], [1, 1, 1, 3], [0, 1, 1, 4]],
[[1, 0, 1, 1], [1, 1, 0, 1], [0, 1, 1, 2], [1, 1, 0, 2], [1, 1, 0, 4]]
]
A partition for subsets of sizes: 0*3 1*4 7*5
```

C. PROGRAM TO CHECK A SUBSET PARTITION

We offer a simple program in Python 3 that allows to check easily if a subset partition is a zero-sum partition. It can be executed in a terminal (or in an online Python environment like https://www.online-python.com/ or https://trinket.io/python), with the three elements of the input: description of the group, sizes of the sets in the partition, and the partition itself, copied-pasted from the annexes.

import json

```
group = json.loads(input())
sizes = json.loads(input())
sets = json.loads(input())
ok = True
for set in sets:
    sums = [0 for pos in range(len(group))]
    for elem in set:
        for pos in range(len(group)):
            sums[pos] = (sums[pos] + elem[pos]) % group[pos]
    if not sums == [0 for i in range(len(group))]:
        ok = False
        break
if ok:
   print("Zero-sum partition")
else:
   print("Not a zero-sum partition")
```