

EQUIVARIANT INSTANTON FLOER HOMOLOGY AND CALCULATIONS FOR THE BINARY POLYHEDRAL SPACES

GARD OLAV HELLE

ABSTRACT. We calculate the equivariant instanton Floer homology, in the sense of Miller Eismeier, for the trivial $SO(3)$ -bundle over the binary polyhedral spaces with coefficients in a PID R for which $2 \in R$ is invertible. Along the way we modify a part of the algebraic construction needed to define the equivariant instanton Floer groups.

CONTENTS

1. Introduction	2
1.1. Organization of Paper	8
2. Equivariant Instanton Floer Homology	9
2.1. The Framed Instanton Floer Complex	9
2.2. The Donaldson Model	12
3. Binary Polyhedral Spaces and Flat Connections	15
3.1. Binary Polyhedral Spaces	15
3.2. A Few Results from Representation Theory	16
3.3. Arrangement of the Flat Connections	17
4. Determination of Moduli Spaces and Differentials	18
4.1. The ADHM Description of Instantons	20
4.2. Determination of the Low-Dimensional Moduli Spaces	23
4.3. The Structure of $DCI(\bar{Y}_\Gamma)$	29
5. The Algebraic Construction of Equivariant Floer Homology	33
5.1. Conventions	34
5.2. The Bar Construction and the Functors C_A^\pm	35
5.3. Tate Homology	40
5.4. The Equivariant Instanton Floer Groups	49
5.5. The Index Spectral Sequences	57
6. Calculations for Binary Polyhedral Spaces	61
6.1. The Case I^+	61
6.2. The Case I^-	66
6.3. The Case I^∞	73
6.4. Calculations for Y_Γ	75
Appendix A. Character Tables, Dynkin Graphs and Flat Connections	77
A.1. Cyclic Groups	77
A.2. Binary Dihedral Groups	78
A.3. The Binary Tetrahedral Group	79
A.4. The Binary Octahedral Group	80
A.5. The Binary Icosahedral Group	81
Appendix B. Equivariant $SU(2)$ -Bundles over S^4 and the Chern-Simons Invariant	82
B.1. Conventions on Quaternions	82

Date: March 18, 2022.

B.2.	Classification of Γ -equivariant $SU(2)$ -bundles over S^4	83
B.3.	Application of the Equivariant Index Theorem	86
B.4.	The Chern-Simons Invariant	89
B.5.	Connections to Group Cohomology	91
	References	93

The research contained in this paper was conducted during the authors time as a PhD fellow at the University of Oslo and formed a part of the authors thesis.

1. INTRODUCTION

Instanton Floer homology has appeared in many forms in the literature. In this paper we will be concerned with a fairly recent version: the equivariant instanton Floer homology in the sense of Miller Eismeyer [Eis19]. A pair (Y, E) consisting of a closed oriented 3-manifold Y and an $SO(3)$ -bundle $E \rightarrow Y$ is said to be weakly admissible if either Y is a rational homology sphere, $H_*(Y; \mathbb{Q}) \cong H_*(S^3; \mathbb{Q})$, or the second Stiefel-Whitney class $w_2(E)$ admits no torsion lift in $H^2(Y; \mathbb{Z})$. Associated with such a pair and a commutative ring of coefficients R (required to be a principal ideal domain), there are four (relatively) $\mathbb{Z}/8$ graded groups

$$(1.1) \quad I^+(Y, E; R), \quad I^-(Y, E; R), \quad I^\infty(Y, E; R) \quad \text{and} \quad \tilde{I}(Y, E; R),$$

that together constitute the equivariant instanton Floer homology of the pair (Y, E) . The first three groups carry module structure over $H^{-*}(BSO(3); R)$ and fit into an exact triangle

$$(1.2) \quad \begin{array}{ccc} I^+(Y, E; R) & \xrightarrow{[3]} & I^-(Y, E; R) \\ & \nwarrow [-4] \quad \nearrow [0] & \\ & I^\infty(Y, E; R) & \end{array},$$

where the numbers denote the degrees of the maps, while the final group is a module over $H_*(SO(3); R)$. The construction of the equivariant instanton Floer groups is naturally divided in two steps. The first step consists of applying Morse-theoretic techniques to the Chern-Simons functional, defined on a framed configuration space of connections in the bundle, to construct a type of Morse-Bott complex $\widetilde{CI}(Y, E; R)$ equipped with an action of $C_*(SO(3); R)$. The second step, purely algebraic in nature, extracts the equivariant groups from this complex using the bar construction from algebraic topology.

There are two main purposes of this paper. The first is to rework some of the algebra needed to set up the equivariant Floer groups. Below we will elaborate further on why we have found this to be necessary. The second is to calculate the equivariant Floer groups associated with the trivial $SO(3)$ -bundle over the binary polyhedral spaces, that is, the orbit spaces $Y_\Gamma := S^3/\Gamma$ for finite subgroups $\Gamma \subset SU(2)$. For reasons that will become apparent later, our calculations will be restricted to the case where 2 is invertible in the ring of coefficients R . In fact, we will focus primarily on the calculations for \overline{Y}_Γ , the manifold Y_Γ equipped with the opposite orientation, and only briefly treat the calculations for Y_Γ .

To be able to explain our results we will first elaborate further on the brief explanation of the equivariant instanton Floer groups given above. To simplify the exposition we will assume that Y is a rational homology sphere and that $E \rightarrow Y$ is the trivial $SO(3)$ -bundle, both of which are satisfied in our situation of binary polyhedral spaces. In this case one may just as well replace E by the trivial $SU(2)$ -bundle, so we will make this assumption as well. Let \mathcal{A}_E denote the space of connections in E and let \mathcal{G}_E be the group of gauge transformations, that is, the group of bundle automorphisms

$u: E \rightarrow E$. The gauge group acts by pullback on the space of connections, but this action is not free. Indeed, the stabilizer of a connection $A \in \mathcal{A}_E$ may be identified, through evaluation at a point $b \in Y$, with the centralizer of the holonomy group $\text{Hol}_A(b) \subset \text{Aut}(E_b) \cong \text{SU}(2)$. To remedy this one may fix a base point $b \in Y$ and define the space of framed connections by

$$\tilde{\mathcal{A}}_E := \mathcal{A}_E \times E_b.$$

Here the gauge group acts diagonally, through evaluation in the second component. This action is free and in suitable Sobolev completions (see [Eis19, Section 2.2]) the configuration space

$$\tilde{\mathcal{B}}_E := \tilde{\mathcal{A}}_E / \mathcal{G}_E$$

is a smooth Hilbert manifold. The right $\text{SU}(2)$ -action on the fiber descends to a right $\text{SO}(3) = \text{SU}(2)/\{\pm 1\}$ -action on the configuration space.

The Chern-Simons functional $\text{cs}: \tilde{\mathcal{B}}_E \rightarrow \mathbb{R}/\mathbb{Z} \cong S^1$ may be defined for $[A, u] \in \tilde{\mathcal{B}}_E$ represented by $(A, u) \in \tilde{\mathcal{A}}_E$ by

$$(1.3) \quad \text{cs}([A, u]) = \frac{1}{8\pi^2} \int_Y \text{Tr}(a \wedge da + \frac{2}{3} a \wedge a \wedge a) \in \mathbb{R}/\mathbb{Z}$$

where $a \in \Omega^1(Y, \mathfrak{su}(2))$ is a connection form representing A in a global trivialization of the bundle (see [Don02, p. 18]). The fact that this is independent of the gauge equivalence class of A , or equivalently the choice of global trivialization of the bundle, is shown in [Fre02, Proposition 1.27]. Clearly, cs is independent of the framing coordinate and is therefore $\text{SO}(3)$ -invariant. The equivariant framed Floer complex $\widetilde{CI}(Y, E)$ is constructed as a type of Morse-Bott complex for the functional $\text{cs}: \tilde{\mathcal{B}}_E \rightarrow S^1$. This means that the complex is, in a certain sense, generated by the set of critical orbits of cs and the differentials are defined via fiber products with the spaces of negative gradient flow lines between the various critical orbits. First, the set of critical points of cs regarded as a map $\mathcal{A}_E \rightarrow \mathbb{R}/\mathbb{Z}$ is precisely the set of flat connections in E , so that \mathcal{C} , the set of critical orbits in $\tilde{\mathcal{B}}_E$, consists precisely of the $\text{SO}(3)$ -orbits of the gauge equivalence classes of the flat connections. Furthermore, the negative gradient flow equation for cs is, for a path of connections $A_t \in \mathcal{A}_E$, given by, disregarding the normalizing constant,

$$\frac{dA_t}{dt} = - * F_{A_t}$$

where F_{A_t} is the curvature of the connection A_t and $*: \Omega^2(Y, \mathfrak{su}(E)) \rightarrow \Omega^1(Y, \mathfrak{su}(E))$ is the Hodge star operator. Now, a path of connections A_t may in a natural way be identified with a connection \mathbf{A} in the bundle $\mathbb{R} \times E \rightarrow \mathbb{R} \times Y$ over the cylinder. In terms of this relation the negative gradient flow equation takes the form of the familiar anti-self-dual or instanton equation

$$F_{\mathbf{A}}^+ = 0.$$

In general, as in finite dimensional Morse theory, one is typically forced to introduce a perturbation of the functional to ensure that the critical points are non-degenerate and that the spaces of flow lines are cut out transversally. A flat connection α , regarded as a critical point of cs , is non-degenerate if the twisted cohomology group $H_\alpha^1(Y; \mathfrak{su}(E))$ vanishes. Given a pair of non-degenerate flat connections α and β and a relative homotopy class $z \in \pi_1(\tilde{\mathcal{B}}_E, \alpha, \beta)$ one may form a configuration space $\tilde{\mathcal{B}}_{z, \mathbb{R} \times E}(\alpha, \beta)$ of connections framed at $(0, b) \in \mathbb{R} \times Y$ approaching α at $-\infty$ and β at $+\infty$, whose corresponding path in $\tilde{\mathcal{B}}_E$ belongs to the homotopy class z , modulo gauge transformations that approach α -harmonic and β -harmonic gauge transformations at $\pm\infty$. This space also carries a residual action of $\text{SO}(3)$ and there are equivariant end-point maps

$$(1.4) \quad \alpha \xleftarrow{e_+} \tilde{\mathcal{B}}_{z, \mathbb{R} \times E}(\alpha, \beta) \xrightarrow{e_-} \beta$$

defined by parallel transport of the framing to $\pm\infty$ along $\mathbb{R} \times \{b\}$. The moduli space of framed instantons $\widetilde{\mathcal{M}}_z(\alpha, \beta) = \{[\mathbf{A}, u] \in \widetilde{\mathcal{B}}_{\mathbb{R} \times Y, z}(\alpha, \beta) : F_{\mathbf{A}}^+ = 0\}$ is then cut out transversally provided the framed analogue of the anti-self-dual operator

$$D_{\mathbf{A}} = (d_{\mathbf{A}}^*, d_{\mathbf{A}}^+) : \Omega^1(\mathbb{R} \times Y, \mathfrak{su}(E)) \rightarrow \Omega^0(\mathbb{R} \times Y, \mathfrak{su}(E)) \oplus \Omega^+(\mathbb{R} \times Y, \mathfrak{su}(E))$$

is (in appropriate Sobolev completions) surjective for all $[\mathbf{A}, u] \in \widetilde{\mathcal{M}}_z(\alpha, \beta)$. For the trivial $\mathrm{SU}(2)$ -bundle over the binary polyhedral spaces all of the flat connections are non-degenerate and the moduli spaces are cut out transversally, so there is no need to introduce a perturbation (see [Aus90, Section 4.5]).

In [Eis19] it is shown, more generally, that there exist generic perturbations that achieve the above (equivariant) transversality conditions. We will assume that such a perturbation has been chosen, leaving it out from the notation. There is then a finite set \mathcal{C} of critical orbits. Given $\alpha \in \mathcal{C}$ the possible stabilizers in $\mathrm{SO}(3)$ are $\{1\}$, $\mathrm{SO}(2)$ or $\mathrm{SO}(3)$, so that

$$(1.5) \quad \alpha \cong \mathrm{SO}(3), \quad \alpha \cong \mathrm{SO}(3)/\mathrm{SO}(2) \cong S^2 \quad \text{or} \quad \alpha \cong *.$$

We say that α is irreducible, reducible or fully reducible respectively. Given a pair of critical orbits $\alpha, \beta \in \mathcal{C}$ there is then for each homotopy class $z \in \pi_1(\widetilde{\mathcal{B}}_E, \alpha, \beta)$ a finite-dimensional moduli space $\widetilde{\mathcal{M}}_z(\alpha, \beta)$ equipped with an $\mathrm{SO}(3)$ -action and equivariant end-point maps. There is a natural \mathbb{R} -action by translations on the moduli space. Provided $\alpha \neq \beta$ this action is free and the quotient is denoted by $\widetilde{\mathcal{M}}_z^0(\alpha, \beta)$. The situation is summarized in the following diagram,

$$\begin{array}{ccccc} & & \widetilde{\mathcal{M}}_z(\alpha, \beta) & & \\ & \swarrow e_- & \downarrow / \mathbb{R} & \searrow e_+ & \\ \alpha & \xleftarrow{e_-} & \widetilde{\mathcal{M}}_z^0(\alpha, \beta) & \xrightarrow{e_+} & \beta, \end{array}$$

where the upper part is obtained by restriction from diagram (1.4). A relative grading, depending on the relative homotopy class z , may then be defined by

$$\mathrm{gr}_z(\alpha, \beta) = \dim \widetilde{\mathcal{M}}_z(\alpha, \beta) - \dim \alpha \in \mathbb{Z}.$$

This function satisfies a few key properties.

- (i) For $\alpha, \beta, \gamma \in \mathcal{C}$ and relative homotopy classes z, w connecting α to β and β to γ respectively one has

$$\mathrm{gr}_{z*w}(\alpha, \gamma) = \mathrm{gr}_z(\alpha, \beta) + \mathrm{gr}_w(\beta, \gamma).$$

- (ii) For $\alpha, \beta \in \mathcal{C}$ and relative homotopy classes z, z' connecting them one has

$$\mathrm{gr}_z(\alpha, \beta) - \mathrm{gr}_{z'}(\alpha, \beta) \in 8\mathbb{Z}.$$

Moreover, for any integer $n \equiv \mathrm{gr}_z(\alpha, \beta) \pmod{8}$ there exists a homotopy class w such that $\mathrm{gr}_w(\alpha, \beta) = n$.

The relative grading $\mathrm{gr} : \mathcal{C} \times \mathcal{C} \rightarrow \mathbb{Z}/8$ is then defined by $\mathrm{gr}(\alpha, \beta) \equiv \mathrm{gr}_z(\alpha, \beta) \pmod{8}$ for any choice of z . In our case of a trivial bundle, we may lift this to an absolute grading $j : \mathcal{C} \rightarrow \mathbb{Z}/8$ by taking the trivial product connection θ as a reference point, that is, $j(\alpha) := \mathrm{gr}(\alpha, \theta)$.

As a graded module, the framed Floer complex $\widetilde{CI}(Y, E; R)$ may now be defined to be the totalization of the bigraded module,

$$\widetilde{CI}(Y, E; R)_{s,t} = \bigoplus_{j(\alpha) \equiv s} C_t^{gm}(\alpha; R)$$

for all $s, t \in \mathbb{Z}$, where the direct sum is taken over all $\alpha \in \mathcal{C}$ with $j(\alpha) \equiv s \pmod{8}$. Here, for a smooth manifold X , $C_*^{gm}(X; R)$ is a chain complex of free R -modules generated by (equivalence

classes of) maps $\sigma: P \rightarrow X$ from a certain set of stratified topological spaces with quite a lot of additional structure (see [Eis19, Section 6]). Importantly, this version of geometric homology supports a notion of fiber products on the chain level, and the homology $H_*^{gm}(X; R)$ coincides with the usual singular homology of X . The differential in $\widetilde{CI}(Y, E)$ is then defined on a generator $c = [\sigma: P \rightarrow \alpha] \in C_t^{gm}(\alpha) \subset \widetilde{CI}(Y, E)_{s,t}$ by taking the sum of the internal differential $\partial^{gm} c \in C_{t-1}^{gm}(\alpha)$ and various fiber products with (compactified) moduli spaces

$$[P \times_{\alpha} \overline{\mathcal{M}}_z(\alpha, \beta) \rightarrow \beta] \in C_{t+\text{gr}_z(\alpha, \beta)-1}^{gm}(\beta).$$

We will state the key results that make all of this precise in section 2.

From this point we omit the reference to the ring of coefficients in the notation. The framed Floer complex $\widetilde{CI}(Y, E)$ carries two important pieces of additional structure. First, $A := C_*^{gm}(\text{SO}(3))$ is a differential graded algebra and the action on $C_*^{gm}(\alpha)$ for $\alpha \in \mathcal{C}$ gives $\widetilde{CI}(Y, E)$ the structure of a right differential graded A -module. Second, $\widetilde{CI}(Y, E)$ is filtered by index:

$$F_p \widetilde{CI}(Y, E)_n = \bigoplus_{j(\alpha) \equiv s, s \leq p} C_{n-s}^{gm}(\alpha)$$

for $p \in \mathbb{Z}$. This is a filtration by differential graded A -submodules and is naturally referred to as the index filtration.

When 2 is invertible in the ground ring, Miller Eismeier shows in [Eis19, Section 7] that it is possible to replace the pair $(\widetilde{CI}(Y, E), C_*^{gm}(\text{SO}(3)))$ with a simpler pair $(DCI(Y, E), H_*(\text{SO}(3)))$. The complex $DCI(Y, E)$, called the Donaldson model, is defined as a graded module to be the totalization of the bigraded module

$$DCI(Y, E)_{s,t} = \bigoplus_{j(\alpha) \equiv s} H_t(\alpha)$$

and the differential has a concrete description. Furthermore, as $1/2 \in R$, $H_*(\text{SO}(3)) \cong \Lambda_R[u]$ is an exterior algebra on a single generator u in degree 3. Using the results of [Aus95], we are able to describe these complexes explicitly for all the binary polyhedral spaces. This is the key input needed for our calculations.

The fourth group, $\widetilde{I}(Y, E)$, in (1.1) is simply defined to be the homology of $\widetilde{CI}(Y, E)$. To explain the other three groups, we have to delve into some differential graded algebra. Let A be an augmented differential graded (DG) algebra. In [Eis19, Appendix A] four functors $C_A^+, C_A^{+,tw}, C_A^-, C_A^\infty$ from the category of right DG A -modules (degree 0 maps assumed) to the category of left DG $C_A^-(R)$ -modules are constructed. The basic tool employed is the two sided bar construction $B(M, A, N)$ associated with the algebra A , a right A -module M and a left A -module N (see Definition 5.1). Then

$$(1.6) \quad C_A^+(M) := B(M, A, R) \quad \text{and} \quad C_A^-(M) := \text{Hom}_A(B(R, A, A), M).$$

Furthermore, $C_A^{+,tw}(M) := B(M, A, D_A)$ where D_A is a dualizing object associated with A (see Definition 5.11). There is a natural transformation $N: C_A^{+,tw} \rightarrow C_A^-$, called the norm map, and the Tate functor C_A^∞ is defined to be the mapping cone of N . It is this that eventually gives rise to the exact triangle connecting the three Floer groups in (1.2). For a right A -module M the homology of $C_A^+(M)$ is denoted by $H_A^+(M)$ and similarly for the other functors.

Provided the ground ring R is a principal ideal domain and A is degreewise free over R , these functors are all exact and preserve quasi-isomorphisms (isomorphisms upon passage to homology). Aside from the role played in the construction of C_A^∞ the functor $C_A^{+,tw}$ is mostly inessential. Indeed, provided A satisfies a Poincaré duality hypothesis of degree $d \in \mathbb{Z}$, valid in the situations of interest with $d = 3$, there is a degree d isomorphism $H_A^+(M) \cong H_A^{+,tw}(M)$ for all right A -modules M .

To define the equivariant Floer groups in (1.1), Miller Eismeier also constructs what he calls completed versions, \hat{C}_A^\bullet for $\bullet \in \{+, -, (+, tw), \infty\}$, of the above functors. The equivariant Floer groups $I^\bullet(Y, E)$ are then defined by applying \hat{C}_A^\bullet to $\widetilde{CI}(Y, E)$, with $A = C_*^{gm}(\text{SO}(3))$, and then passing to homology. To explain these completed versions it is important to note that the bar construction $B(M, A, N)$ is obtained as the totalization of a double complex $(B_{*,*}(M, A, N), \partial', \partial'')$. It therefore carries two filtrations, one by the first simplicial degree, and one by the second internal degree. In [Eis19, Definition A.6] the completed bar construction $\hat{B}(M, A, N)$ is defined to be the completion of $B(M, A, N)$ with respect to the filtration by internal degree, and the finitely supported cobar construction $c\hat{B}(N, A, M) \subset \text{Hom}_A(B(N, A, A), M)$ is defined to be the submodule of functionals $B(N, A, A) \rightarrow M$ vanishing on $B_{p,q}(N, A, A)$ for all sufficiently large p . The completed versions \hat{C}_A^+ and \hat{C}_A^- are then defined by replacing $B(M, A, R)$ with $\hat{B}(M, A, R)$ and $\text{Hom}_A(B(R, A, A), M)$ with $c\hat{B}(R, A, M)$ in (1.6), while $\hat{C}_A^{+,tw}(M) := \hat{B}(M, A, \hat{D}_A)$ where \hat{D}_A is a completed version of the dualizing object. Finally, the norm map extends to a map $\hat{N} : \hat{C}_A^{+,tw}(M) \rightarrow \hat{C}_A^-(M)$ and $\hat{C}_A^\infty(M)$ is defined to be the cone of this extended map.

Let $M = \widetilde{CI}(Y, E)$ be equipped with the index filtration and let $A = C_*^{gm}(\text{SO}(3))$. The intention of the above completions is to make sure that for $\bullet \in \{+, -, (+, tw), \infty\}$ the induced filtration $F_p \hat{C}_A^\bullet(M) := \hat{C}_A^\bullet(F_p M)$ is well-behaved; by which we mean exhaustive and complete Hausdorff (see [Boa99, Definition 2.1]). This is to ensure that the associated spectral sequence converges conditionally, in the sense of [Boa99], to the correct target. However, the fact that this induced filtration is well-behaved in the above sense is not justified in [Eis19]. In fact, we find this difficult to believe in the case of $\hat{C}_A^{+,tw}$ and \hat{C}_A^∞ . To remedy this potential issue we suggest a different approach to the completed functors. Rather than defining completed functors on the whole category of A -modules, we simply promote the existing functors C_A^\bullet to functors between the corresponding filtered categories and then compose with a full completion functor to ensure that the resulting filtrations are well-behaved. To be precise, given a right DG A -module M equipped with an increasing filtration by DG A -submodules $\{F_p M\}_p$ we define

$$\begin{aligned} \hat{C}_A^\bullet(M) &:= \lim_q \text{colim}_p C_A^\bullet(F_p M / F_q M) \quad \text{filtered by} \\ F_p \hat{C}_A^\bullet(M) &:= \lim_{q < p} C_A^\bullet(F_p M / F_q M) \quad \text{for } p \in \mathbb{Z} \end{aligned}$$

for each $\bullet \in \{+, -, (+, tw), \infty\}$. This approach has the advantage that essentially all the relevant properties of the functors C_A^\bullet extend in a straightforward way to the functors \hat{C}_A^\bullet .

For $A = C_*^{gm}(\text{SO}(3))$ and $M = \widetilde{CI}(Y, E)$ we define filtered complexes of $C_A^-(R)$ -modules

$$CI^\bullet(Y, E) = \hat{C}_A^\bullet(M) \quad \text{for } \bullet \in \{+, -, (+, tw), \infty\}$$

and define $I^\bullet(Y, E)$ to be its homology. When $\frac{1}{2} \in R$ we also define $DCI^\bullet(Y, E)$ as above with $A = \Lambda_R[u]$ and $M = DCI(Y, E)$. Our main results concerning this construction are simply that all of the desired formal properties of the groups mentioned at the beginning of the section are valid. This is established in section 5. Furthermore, by the nature of our definitions we are with certainty able to establish the existence of the index spectral sequences

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_{\text{SO}(3)}^\bullet(\alpha)_t \implies I_{s+t}^\bullet(Y, E)$$

of $H^{-*}(B\text{SO}(3))$ -modules converging conditionally to the desired target. Here, $H_{\text{SO}(3)}^\bullet(\alpha) := H_A^\bullet(C_*^{gm}(\alpha))$. All of these spectral sequences carry a periodicity isomorphism $E_{s,t}^r \cong E_{s+8,t}^r$ for all $s, t \in \mathbb{Z}$, compatible with the module structure, the differentials and the target.

We should note that this modification of the completion procedure has no essential consequences for any of the results in the main body of [Eis19] as far as we can see. Moreover, we explain, see Remark 5.46, that our construction coincides with the original one for $I^\pm(Y, E)$.

For $A = \Lambda_R[u]$ with $|u| = 3$ it holds true that $C_A^-(R) = H_A^-(R) = R[U]$, a polynomial algebra on a single generator U of degree -4 . In this situation we give concrete models for $C_A^+(M)$, $C_A^-(M)$ and $C_A^\infty(M)$. In the first two cases these are easy to extract and are also essentially (up to signs) contained in [Eis19], but the chain level model for C_A^∞ seems to be new. For $M = \widetilde{CI}(Y, E)$ or $M = DCI(Y, E)$ equipped with the index filtration we are able to extend these models to the completed functors $\hat{C}_A^\pm(M)$ and $\hat{C}_A^\infty(M)$. In all cases these models express the relevant complex as a suitable totalization of a double complex $(D_{*,*}, \partial', \partial'')$ concentrated in the right half plane in the $+$ case, the left half-plane in the $-$ case and the whole plane in the ∞ case. Each nonzero column $(D_{s,*}, \partial'')$ is a shifted copy of M and the horizontal differential ∂' is given by the action of $u \in \Lambda_R[u]$ up to a sign. Moreover, the action of $U \in R[U]$ is easy to express in these models. This leads to a new proof of the fact that $U: I^\infty(Y, E) \rightarrow I^\infty(Y, E)$ is an isomorphism ($\frac{1}{2} \in R$ assumed) established in [Eis19]. We find these models very convenient to use at various places in our calculations.

We now move over to the calculational content of this paper. The finite subgroups of $SU(2)$ are naturally divided into two infinite families consisting of cyclic groups C_m and binary dihedral groups D_n^* , $n \geq 2$, as well as three exceptional ones: the binary tetrahedral group T^* , the binary octahedral group O^* and the binary icosahedral group I^* . The quotients S^3/C_m are lens spaces and S^3/I^* is the famous Poincaré homology sphere. Calculations in these cases are contained in [Eis19, Section 7] for one of the two possible orientations.

Calculations of equivariant instanton Floer groups for binary polyhedral spaces have been considered before. Austin and Braam introduced a type of equivariant instanton Floer groups in [AB96]. Their theory, although similar to the one considered here, relies on using differential forms to express the equivariant complexes and is therefore restricted to using real coefficients. Austin calculated these groups for binary polyhedral spaces in [Aus95]. Most of the necessary background material needed to set up the Donaldson model $DCI(Y, E)$ is therefore contained in this paper. The key idea is to exploit a natural bijection between instantons over the cylinder $\mathbb{R} \times S^3/\Gamma$ and Γ -invariant instantons on Γ -equivariant $SU(2)$ -bundles over $S^4 = (\mathbb{R} \times S^3) \cup \{\pm\infty\}$ with the suspended Γ -action, and then use an equivariant version of the well-known ADHM correspondence to determine the relevant moduli spaces. We review and expand on several of the necessary results in section 4. In particular, to state the equivariant ADHM-correspondence clearly we have included a classification of Γ -equivariant $SU(2)$ -bundles over S^4 , whose proof is given in Appendix B.

The conclusion of the above is that we are able to describe the complex $DCI(\overline{Y}_\Gamma)$ for all finite subgroups $\Gamma \subset SU(2)$ explicitly. The description can be summarized as follows. Given a finite subgroup $\Gamma \subset SU(2)$, we construct a graph \mathcal{S}_Γ whose vertices correspond to the flat connections in the trivial $SU(2)$ -bundle over $Y_\Gamma = S^3/\Gamma$. In all cases this graph is a tree. Attached to an edge connecting two vertices α and β there is a symbol encoding the action of the differential between these. If α and β are both irreducible the symbol is $(n_{\alpha\beta}|n_{\beta\alpha})$ and if only, say α , is irreducible the symbol is $n_{\alpha\beta}$, where the $n_{\alpha\beta}$ are certain integers. Then the labeled graph \mathcal{S}_Γ contains all the information needed to set up $DCI(\overline{Y}_\Gamma)$. The grading is determined by $j(\alpha) = 4p(\alpha)$ where $p(\alpha)$ is the length of a minimal path connecting α to the trivial connection θ in \mathcal{S}_Γ . In particular, $DCI(\overline{Y}_\Gamma)_{s,t} = 0$ unless $4|s$. This has the implication that the only possibly nontrivial part of the differential is $\partial: DCI_{4s,0} \rightarrow DCI_{4(s-1),3}$. This differential has components $\partial_{\alpha\beta}: H_0(\beta) \rightarrow H_3(\alpha)$, which is nonzero precisely when α and β are adjacent in \mathcal{S}_Γ and α is irreducible. In that case, the differential is specified by $\partial_{\alpha\beta}(b_\beta) = n_{\alpha\beta}t_\alpha$ where $b_\beta \in H_0(\beta)$ and $t_\alpha \in H_3(\alpha)$ are generators. The

labeled graphs \mathcal{S}_Γ for all the finite subgroups $\Gamma \subset \mathrm{SU}(2)$ are contained in Proposition 4.25 and examples of the resulting Donaldson complexes are given in Example 4.26.

In the final section of the paper we explicitly calculate $I^+(\overline{Y}_\Gamma)$, $I^-(\overline{Y}_\Gamma)$ and $I^\infty(\overline{Y}_\Gamma)$ for each finite subgroup $\Gamma \subset \mathrm{SU}(2)$ and observe that the exact triangle relating them splits into a short exact sequence. The calculations are contained in Theorem 6.7, Theorem 6.13, Theorem 6.14 and Theorem 6.16. The main tools in the calculations are the explicit complexes $DCI(\overline{Y}_\Gamma)$, the index spectral sequences of Theorem 5.58 and the explicit models for DCI^+ , DCI^- mentioned earlier. Interestingly, we obtain essentially the same results as in [Aus95]. Indeed, the fact that $\frac{1}{2} \in R$ ensures that exactly the same type of degeneration pattern seen in Austin's calculations occur for us as well.

In the final part of Appendix B we also include a discussion of the Chern-Simons invariant of the flat connections in the trivial $\mathrm{SU}(2)$ -bundle over the various binary polyhedral spaces. The main result we wish to bring out is that this numerical invariant can easily be calculated with the help of a few results established in section 4. Methods for these calculations are certainly known, see for instance [KK90] or [Auc94]. However, we also show, possibly more interestingly, that this invariant can in a natural way be identified with the second Chern class of the holonomy representation associated with the flat connection.

1.1. Organization of Paper. We start in section 2 by stating the key theorem of [Eis19] and some additional details on the geometric homology functor C_*^{gm} needed to rigorously define the framed Floer complex $\widehat{CI}(Y, E)$. Moreover, we also introduce the Donaldson model $DCI(Y, E)$ under a simplifying assumption valid in the case of binary polyhedral spaces. Finally, we state the precise result that ensures that $DCI(Y, E)$ also may be used to calculate the various flavours of equivariant instanton Floer homology.

In section 3 we recall the classification of finite subgroups of $\mathrm{SU}(2)$ and specify our conventions concerning the binary polyhedral spaces. We also recall some simple representation theory and the McKay correspondence. This is needed to efficiently arrange the set of flat connections in the trivial $\mathrm{SU}(2)$ -bundle over the binary polyhedral spaces. At the end of the section we define the graphs \mathcal{S}_Γ .

In section 4 we review the key results from [Aus95] that explicitly determine the low-dimensional instanton moduli spaces needed to set up the Donaldson model for the binary polyhedral spaces. Here we have, for the purpose of clarity and precision, expanded on Austin's rather terse exposition in a number of places. In the final part of the section we give an explicit description of $DCI(\overline{Y}_\Gamma)$ for each finite subgroup $\Gamma \subset \mathrm{SU}(2)$.

In section 5 we move into the world of differential graded algebra. First, we review the construction of the functors C_A^\bullet ; their invariance and functorial properties for $\bullet \in \{+, -, (+, tw), \infty\}$ mainly following [Eis19, Appendix A.2, A.3, A.5]. Detailed calculations are included in the case $A = \Lambda_R[u]$ for $|u| = 3$, but we note that all of these calculations easily extend to the case of $|u|$ of arbitrary odd degree. After this we explain how to extend the functors to the category of filtered modules and establish all the necessary functoriality and invariance results in detail. We then define the complexes $CI^\bullet(Y, E)$ for $\bullet \in \{+, -, (+, tw), \infty\}$ calculating the various flavors of equivariant Floer homology, and similarly for $DCI^\bullet(Y, E)$. In the final part of the section we recall some theory on spectral sequences and their convergence properties needed for our later calculations, and then go on to set up the index spectral sequences.

In the final section we calculate the various flavors of equivariant instanton Floer homology for the trivial $\mathrm{SU}(2)$ -bundle over the binary polyhedral spaces \overline{Y}_Γ . In the last part of the section we explain how $DCI(Y_\Gamma)$ can be recovered from $DCI(\overline{Y}_\Gamma)$ and then give the calculations for Y_Γ , omitting some details.

In Appendix A we have included various facts concerning the finite subgroups $\Gamma \subset \mathrm{SU}(2)$. This include explicit realizations within $\mathrm{Sp}(1)$, information on their complex representation theory, character tables for T^* , O^* and I^* , extended Dynkin diagrams and complete lists over the 1-dimensional quaternionic representations. This is needed to determine all the flat connections in the trivial $\mathrm{SU}(2)$ -bundle over \overline{Y}_Γ and for the indexing and differentials in the complexes $DCI(\overline{Y}_\Gamma)$.

In Appendix B we have included proofs of two results used in section 4: the classification of Γ -equivariant $\mathrm{SU}(2)$ -bundles over S^4 and an index calculation. As a byproduct, we obtain a simple procedure for the computation of the Chern-Simons invariants for the various flat connections over the binary polyhedral spaces. We also discuss a relation between these invariants and the group cohomology of the binary polyhedral groups.

2. EQUIVARIANT INSTANTON FLOER HOMOLOGY

In this section we will expand on, and make more rigorous, the definition of the framed instanton Floer complex $\widetilde{CI}(Y, E)$. We will assume, as in the introduction, that Y is a rational homology sphere and that $E \rightarrow Y$ is the trivial $\mathrm{SU}(2)$ -bundle as this is sufficient for our purpose. For the more general case of weakly admissible bundles we refer to [Eis19]. We will assume as in the introduction that a perturbation of the Chern-Simons functional has been fixed such that every critical orbit is non-degenerate and all the relevant moduli spaces are cut out transversally. As already noted, there is no need to introduce a perturbation in the case of binary polyhedral spaces (see [Aus95, Section 4.5]). The set of critical orbits will always be denoted by \mathcal{C} .

2.1. The Framed Instanton Floer Complex. Recall that we have fixed a basepoint $b \in Y$ and that $\widetilde{\mathcal{B}}_E$ is a configuration space of framed connections in E modulo gauge equivalence. Moreover, for each pair $\alpha, \beta \in \mathcal{C}$ and a relative homotopy class $z \in \pi_1(\widetilde{\mathcal{B}}_E, \alpha, \beta)$ there is a configuration space $\widetilde{\mathcal{B}}_{z, \mathbb{R} \times E}(\alpha, \beta)$ of connections in $\mathbb{R} \times E \rightarrow \mathbb{R} \times Y$ framed at $(0, b)$ that approaches α and β respectively as t tends to $\pm\infty$, modulo gauge. There is a residual $\mathrm{SO}(3)$ -action and equivariant end-point maps $(e_-, e_+): \widetilde{\mathcal{B}}_{z, \mathbb{R} \times E}(\alpha, \beta) \rightarrow \alpha \times \beta$. In reality, these spaces are completed with respect to certain weighted Sobolev norms (see [Eis19, Section 2.2] for the specifics). We may now state the main theorem required to set up the framed instanton Floer complex. Here we have extracted the relevant part of [Eis19, Theorem 6.8].

Theorem 2.1. *Let $E \rightarrow Y$ be the trivial $\mathrm{SU}(2)$ -bundle over Y and let \mathcal{C} be the finite set of critical orbits of the (perturbed) Chern-Simons functional.*

(a). *For each pair $\alpha, \beta \in \mathcal{C}$ and relative homotopy class z between them in $\widetilde{\mathcal{B}}_E$ there is a number $\mathrm{gr}_z(\alpha, \beta) \in \mathbb{Z}$. If $\gamma \in \mathcal{C}$, w is a relative homotopy class from β to γ and $z * w$ denotes the concatenation of paths then*

$$\mathrm{gr}_{z*w}(\alpha, \gamma) = \mathrm{gr}_z(\alpha, \beta) + \mathrm{gr}_w(\beta, \gamma).$$

Moreover, if z' is a different homotopy class from α to β it holds true that $\mathrm{gr}_z(\alpha, \beta) - \mathrm{gr}_{z'}(\alpha, \beta) \in 8\mathbb{Z}$. Hence, $\mathrm{gr}(\alpha, \beta) \in \mathbb{Z}/8$ is well defined.

(b). *For each pair $\alpha, \beta \in \mathcal{C}$ and relative homotopy class $z \in \pi_1(\widetilde{\mathcal{B}}_E, \alpha, \beta)$ there is a moduli space of framed instantons $\widetilde{\mathcal{M}}_z(\alpha, \beta)$ approaching α as $t \rightarrow -\infty$ and β as $t \rightarrow \infty$. This is a smooth right $\mathrm{SO}(3)$ -manifold of dimension $\mathrm{gr}_z(\alpha, \beta) + \dim(\alpha)$.*

(c). *There is a smooth \mathbb{R} -action by translations and parallel transport of the framing on $\widetilde{\mathcal{M}}_z(\alpha, \beta)$ with smooth quotient denoted by $\widetilde{\mathcal{M}}_z^0(\alpha, \beta)$.*

(d). There are smooth $\mathrm{SO}(3)$ -equivariant end-point maps as in the diagram

$$\begin{array}{ccc} & \widetilde{\mathcal{M}}_z(\alpha, \beta) & \\ e_- \swarrow & \downarrow / \mathbb{R} & \searrow e_+ \\ \alpha \xleftarrow{e_-} & \widetilde{\mathcal{M}}_z^0(\alpha, \beta) & \xrightarrow{e_+} \beta \end{array}$$

obtained by parallel transport of the framing towards $\pm\infty$. Here we identify $\alpha \cong E_b/\Gamma_\alpha$ and $\beta \cong E_b/\Gamma_\beta$, where $\Gamma_\alpha, \Gamma_\beta \subset \mathrm{Aut}(E_b)$ correspond to the stabilizers of α and β .

(e). For each $\alpha \in \mathcal{C}$ there is a two-element set $\Lambda(\alpha)$ corresponding to a choice of orientation. For a pair α, β any choice of orientations in $\Lambda(\alpha)$ and $\Lambda(\beta)$ induces an orientation in the fiber of $e_- : \widetilde{\mathcal{M}}_z^0(\alpha, \beta) \rightarrow \alpha$. If either of the choices are changed the resulting fiber orientation is reversed.

(f). If $\mathrm{gr}_z(\alpha, \beta) - \dim(\alpha) \leq 10$ there is a compactification

$$\widetilde{\mathcal{M}}_z^0(\alpha, \beta) \subset \overline{\mathcal{M}}_z(\alpha, \beta)$$

into a topological $\mathrm{SO}(3)$ -manifold with corners and smooth structure on each stratum. The endpoint maps extend over the compactification. Furthermore, the $\mathrm{SO}(3)$ action is free.

(g). Given any choice of elements from $\Gamma(\alpha)$, $\Gamma(\beta)$ and $\Gamma(\gamma)$ there is a decomposition respecting fiber orientations

$$(-1)^{\dim \alpha} \partial \overline{\mathcal{M}}_z(\alpha, \beta) = \coprod_{w_1, w_2: w_1 * w_2 = z} \overline{\mathcal{M}}_{w_1}(\alpha, \gamma) \times_\gamma \overline{\mathcal{M}}_{w_2}(\gamma, \beta).$$

The function $\mathrm{gr}: \mathcal{C} \times \mathcal{C} \rightarrow \mathbb{Z}/8$ is the relative grading on the set of critical orbits. Since the bundle $E \rightarrow Y$ is trivial we may define an absolute grading $j: \mathcal{C} \rightarrow \mathbb{Z}/8$ by setting $j(\alpha) := \mathrm{gr}(\alpha, \theta)$, where θ denotes the trivial product connection. The relative grading may be recovered from j :

$$j(\alpha) - j(\beta) = \mathrm{gr}(\alpha, \theta) - \mathrm{gr}(\beta, \theta) = \mathrm{gr}(\alpha, \beta) + \mathrm{gr}(\beta, \theta) - \mathrm{gr}(\beta, \theta) = \mathrm{gr}(\alpha, \beta).$$

To construct the framed Floer complex Miller Eismeier uses a type of geometric homology. For each smooth manifold X he constructs a functorial chain complex $C_*^{gm}(X)$ whose generators are equivalence classes of maps from a set of spaces called strong δ -chains. For the precise definition see [Eis19, Def.:6.1]. This class of spaces contains all topological manifolds with corners and smooth structure on each stratum, so in particular contains the compactified instanton moduli spaces in part (f) of the above theorem.

Proposition 2.2. *Given a commutative ring of coefficients R there is a functor $C_*^{gm}(-; R)$ from the category of smooth manifolds to the category of homological chain complexes of R -modules. For a smooth manifold X of dimension d the following holds true.*

- (i) *There is a natural isomorphism $H_*^{sing}(X; R) \cong H_*^{gm}(X; R)$.*
- (ii) *$C_n^{gm}(X; R)$ is a free R -module for each $n \in \mathbb{Z}$.*
- (iii) *$C_n^{gm}(X; R) = 0$ for all $n < 0$ and $n > d + 1$.*

There is one additional property of C_*^{gm} that is needed for the definition of the framed instanton Floer complex. First, for a pair of smooth manifolds X and Y there is a cross product $C_i^{gm}(X; R) \times C_j^{gm}(Y; R) \rightarrow C_{i+j}^{gm}(X \times Y)$. This ensures that $C_*^{gm}(\mathrm{SO}(3); R)$ obtains the structure of a differential graded algebra and for any right $\mathrm{SO}(3)$ -manifold X , $C_*^{gm}(X; R)$ obtains the structure of a right differential graded $C_*^{gm}(\mathrm{SO}(3); R)$ -module.

The framed Floer complex $\widetilde{CI}(Y, E)$ will be defined to be the chain complex associated with a multicomplex, whose definition we first recall.

Definition 2.3. A (homological) multicomplex is a pair $(M_{*,*}, \{\partial^r\}_{r \geq 0})$ consisting of a bigraded module $M_{*,*}$ and differentials $\partial^r: M \rightarrow M$ of bidegree $(-r, r-1)$ for $r \geq 0$ such that

- (i) $\sum_{i+j=k} \partial^i \circ \partial^j = 0 : M_{s,t} \rightarrow M_{s-k,t-k+2}$ for all $k \geq 0$, $s, t \in \mathbb{Z}$, and
- (ii) for each $a \in M_{s,t}$ there exists $r_0 \geq 0$ such that $\partial^r(a) = 0$ for all $r \geq r_0$.

The associated chain complex (M_*, ∂) is defined by $M_n = \bigoplus_{s+t=n} M_{s,t}$ for all $n \in \mathbb{Z}$ and $\partial = \sum_{r \geq 0} \partial^r$.

For a commutative ring of coefficients R define the bigraded R -module $\widetilde{CI}(Y, E; R)_{*,*}$ by

$$\widetilde{CI}(Y, E; R)_{s,t} = \bigoplus_{j(\alpha) \equiv s} C_t^{gm}(\alpha; R)$$

for all $s, t \in \mathbb{Z}$. Observe that since the grading function j takes values in $\mathbb{Z}/8$ there is a periodicity isomorphism $\widetilde{CI}(Y, E; R)_{s,t} \cong \widetilde{CI}(Y, E; R)_{s+8,t}$ for all $s, t \in \mathbb{Z}$. We make a preliminary definition before we define the differentials $\{\partial^r\}_{r \geq 0}$. For each pair $\alpha, \beta \in \mathcal{C}$ fix the unique relative homotopy class z from α to β satisfying $-2 \leq \text{gr}_z(\alpha, \beta) \leq 5$. Then according to part (f) of the above theorem the moduli space $\widetilde{\mathcal{M}}_z^0(\alpha, \beta)$ admits a compactification $\overline{\mathcal{M}}_z(\alpha, \beta)$ and we have a diagram

$$\alpha \xleftarrow{e_-} \overline{\mathcal{M}}_z(\alpha, \beta) \xrightarrow{e_+} \beta.$$

By [Eis19, Lemma 6.6] there is an induced fiber product map

$$f_{\alpha\beta} : C_*^{gm}(\alpha; R) \rightarrow C_{*+\text{gr}_z(\alpha, \beta)-1}^{gm}(\beta; R)$$

defined as follows. Given a basic chain $\sigma : P \rightarrow \alpha$ with $\dim P = t$ representing an element $[\sigma : P \rightarrow \alpha] \in C_t^{gm}(\alpha; R)$ we form the fiber product $P \times_\alpha \overline{\mathcal{M}}_z(\alpha, \beta)$ as in the commutative diagram

$$\begin{array}{ccc} P & \xleftarrow{\quad} & P \times_\alpha \overline{\mathcal{M}}_z(\alpha, \beta) \\ \downarrow \sigma & & \downarrow \tau \\ \alpha & \xleftarrow{e_-} & \overline{\mathcal{M}}_z(\alpha, \beta) \xrightarrow{e_+} \beta, \end{array} \quad \begin{array}{c} \nearrow e_+ \circ \tau \end{array}$$

and define

$$f_{\alpha\beta}([\sigma : P \rightarrow \alpha]) = [e_+ \circ \tau : P \times_\alpha \overline{\mathcal{M}}_z(\alpha, \beta) \rightarrow \beta] \in C_{t+\text{gr}_z(\alpha, \beta)-1}^{gm}(\beta; R).$$

In case $\text{gr}_z(\alpha, \beta) \leq 0$ the compactified moduli space $\overline{\mathcal{M}}_z(\alpha, \beta)$ is empty and the map $f_{\alpha\beta}$ vanishes. Otherwise, we note that the dimension formula

$$(2.1) \quad \dim \overline{\mathcal{M}}_z(\alpha, \beta) = \text{gr}_z(\alpha, \beta) + \dim \alpha - 1$$

ensures that the chain $e_+ \circ \tau : P \times_\alpha \overline{\mathcal{M}}_z(\alpha, \beta) \rightarrow \beta$ has dimension $\dim P + \text{gr}_z(\alpha, \beta) - 1$ as asserted. Finally, we note that the maps $f_{\alpha\beta}$ commute with the action of $C_*^{gm}(\text{SO}(3); R)$.

Definition 2.4. The framed instanton Floer complex $\widetilde{CI}(Y, E; R)$ with coefficients in R is defined to be the chain complex associated with the multicomplex $(\widetilde{CI}(Y, E; R)_{*,*}, \{\partial^r\}_{r \geq 0})$ where

$$\widetilde{CI}(Y, E; R)_{s,t} = \bigoplus_{\alpha \in \mathcal{C} : j(\alpha) \equiv s} C_t^{gm}(\alpha; R)$$

and the differentials $\partial^r : \widetilde{CI}(Y, E; R)_{s,t} \rightarrow \widetilde{CI}(Y, E; R)_{s-r,t+r-1}$ for $r \geq 0$ are defined as follows:

- (i) ∂^0 is the sum of the internal differentials $\partial^{gm} : C_*^{gm}(\alpha; R) \rightarrow C_{*-1}^{gm}(\alpha; R)$.
- (ii) For $1 \leq r \leq 5$ the differential ∂^r is the sum of the maps $f_{\alpha\beta} : C_*^{gm}(\alpha; R) \rightarrow C_{*+\text{gr}_z(\alpha, \beta)-1}^{gm}(\beta; R)$ for pairs α, β with $\text{gr}_z(\alpha, \beta) = r$.
- (iii) $\partial^r = 0$ for $r > 5$.

The homology of $\widetilde{CI}(Y, E; R)$ is denoted by $\widetilde{I}(Y, E)$ and is called the framed instanton Floer homology of the pair (Y, E) . The $C_*^{gm}(\mathrm{SO}(3); R)$ -module structure of $C_*^{gm}(\alpha; R)$ for $\alpha \in \mathcal{C}$ gives $\widetilde{CI}(Y, E; R)$ the structure of a right differential graded $C_*^{gm}(\mathrm{SO}(3); R)$ -module.

The filtration by $C_*^{gm}(\mathrm{SO}(3))$ -modules given degreewise by

$$F_p \widetilde{CI}(Y, E; R)_n := \bigoplus_{s \leq p} \widetilde{CI}(Y, E; R)_{s, n-s}$$

is called the index filtration.

Remark 2.5. In the above we have omitted discussing orientations. To get the complex $\widetilde{CI}(Y, E)$ in the above form one has to fix an orientation $\mathfrak{o}_\alpha \in \Lambda(\alpha)$ for each critical orbit $\alpha \in \mathcal{C}$. The verification that $\partial^2 = 0$ and that this complex has the structure of a right differential graded $C_*^{gm}(\mathrm{SO}(3))$ -module are contained in [Eis19, Lemma 6.11].

From this point we will omit the ring of coefficients from the notation. The complex given in the above definition is the unrolled framed instanton Floer complex. Instead of having a $\mathbb{Z}/8$ -graded complex, we have a \mathbb{Z} -graded complex equipped with an evident periodicity isomorphism

$$\widetilde{CI}(Y, E)_n \cong \widetilde{CI}(Y, E)_{n+8}$$

for all $n \in \mathbb{Z}$, compatible with the $C_*^{gm}(\mathrm{SO}(3))$ -action and the differential. This is more convenient when we apply the algebraic machinery of section 5 to extract the equivariant Floer groups. It should be noted that the above periodicity isomorphisms interact with the index filtration in the following way

$$F_p \widetilde{CI}(Y, E)_n \cong F_{p+8} \widetilde{CI}(Y, E)_{n+8}$$

for all $p, n \in \mathbb{Z}$.

2.2. The Donaldson Model. In the previous definition we introduced the framed Floer complex $\widetilde{CI}(Y, E)$ as a differential graded module over the differential graded algebra $C_*^{gm}(\mathrm{SO}(3))$. Since we are still working on the chain level, the complex and action are difficult to handle. It is shown in [Eis19, Section 7.2] that provided $2 \in R$ is invertible, one may replace the pair $(C_*^{gm}(\mathrm{SO}(3)), \widetilde{CI}(Y, E))$ with a simpler pair $(H_*(\mathrm{SO}(3)), DCI(Y, E))$. The complex $DCI_*(Y, E)$ is called the Donaldson model and is associated with a multicomplex whose bigraded module is given by

$$DCI(Y, E)_{s,t} = \bigoplus_{\alpha \in \mathcal{C} : j(\alpha) \equiv s} H_t(\alpha).$$

Note that $H_*(\mathrm{SO}(3); R) = \Lambda_R[u]$ is an exterior algebra on a single generator u of degree 3 as $\frac{1}{2} \in R$. For $\alpha \in \mathcal{C}$ we have by (1.5)

$$(2.2) \quad H_*(\alpha) \cong \begin{cases} R \oplus R[3] & \alpha \text{ irreducible} \\ R \oplus R[2] & \alpha \text{ reducible} \\ R & \alpha \text{ fully reducible} \end{cases}$$

Here our convention regarding the shift operation is that if C is a graded module then $C[n]_{n+i} = C_i$ for all $i, n \in \mathbb{Z}$. The above isomorphisms are made into identifications by fixing a base point and an orientation for each orbit $\alpha \in \mathcal{C}$. The $\Lambda_R[u] = H_*(\mathrm{SO}(3))$ -action is uniquely determined by the map $\cdot u : DCI_{*,*} \rightarrow DCI_{*,*+3}$, which has components $\mathrm{id} : R = H_0(\alpha) \rightarrow H_3(\alpha) = R$ for α irreducible and vanishes otherwise.

The complete description of the differentials in this complex is given in [Eis19, p.:175-176]. In the following we will only concern ourselves with the components relevant in the situation for binary polyhedral spaces. We therefore make the following assumption (see Lemma 4.17)

- (i) For each pair $\alpha, \beta \in \mathcal{C}$ we have $\mathrm{gr}(\alpha, \beta) \equiv 0 \pmod{4}$.

This implies that $DCI(Y, E)_{s,t}$ can only be nontrivial if $4|s$ and $0 \leq t \leq 3$. The only possibly nontrivial differential is therefore ∂^4 of bidegree $(-4, 3)$, whose components are $R = H_0(\alpha) \rightarrow H_3(\beta) = R$ for $\alpha, \beta \in \mathcal{C}$ with $j(\alpha) - j(\beta) = 4$ and β irreducible.

Recall that for $\alpha, \beta \in \mathcal{C}$ we defined z to be the unique homotopy class between α and β for which $-2 \leq \text{gr}_z(\alpha, \beta) \leq 5$. Combining this with the above assumption and the boundary formula in part (g) of Theorem 2.1 we may conclude that $\mathcal{M}_z^0(\alpha, \beta) = \overline{\mathcal{M}}_z(\alpha, \beta)$ is compact without boundary. If $\text{gr}_z(\alpha, \beta) = 0$ the moduli space is empty and if $\text{gr}_z(\alpha, \beta) = 4$ the dimension is $\dim \alpha + 3$ (provided it is nonempty). For simplicity of notation we write $\overline{\mathcal{M}}_{\alpha, \beta} = \overline{\mathcal{M}}_z(\alpha, \beta)$ with this choice of homotopy class z implicit.

To define the differential ∂^4 we have to make a little technical detour. For each orbit $\alpha \in \mathcal{C}$ let $\tilde{\alpha} \rightarrow \alpha$ denote its universal cover. For irreducible orbits α we have $\tilde{\alpha} = \text{SU}(2)$, while for reducible or fully reducible orbits $\tilde{\alpha} = \alpha$. These coverings carry $\text{SU}(2)$ -actions such that the projections $\tilde{\alpha} \rightarrow \alpha$ are equivariant along $\text{SU}(2) \rightarrow \text{SO}(3)$. For each pair $\alpha, \beta \in \mathcal{C}$ define $\overline{\mathcal{M}}_{\alpha, \beta}^{\text{cov}}$ to be the pullback as in the diagram

$$\begin{array}{ccc} \overline{\mathcal{M}}_{\alpha, \beta}^{\text{cov}} & \xrightarrow{(\tilde{e}_-, \tilde{e}_+)} & \tilde{\alpha} \times \tilde{\beta} \\ \downarrow & & \downarrow \\ \overline{\mathcal{M}}_{\alpha, \beta} & \xrightarrow{(e_-, e_+)} & \alpha \times \beta. \end{array}$$

Thus $\overline{\mathcal{M}}_{\alpha, \beta}^{\text{cov}}$ is a 1, 2 or 4-sheeted covering of $\overline{\mathcal{M}}_{\alpha, \beta}$ depending on whether α and/or β are irreducible. The free $\text{SO}(3)$ -action on $\overline{\mathcal{M}}_{\alpha, \beta}$ lifts to a free $\text{SU}(2)$ -action on $\overline{\mathcal{M}}_{\alpha, \beta}^{\text{cov}}$ such that $(\tilde{e}_-, \tilde{e}_+)$ is equivariant. Define

$$\mathcal{M}_{\alpha, \beta} := \overline{\mathcal{M}}_{\alpha, \beta}^{\text{cov}} / \text{SU}(2) \quad \text{and} \quad X_{\alpha, \beta} := (\tilde{\alpha} \times \tilde{\beta}) / \text{SU}(2)$$

and let $e: \mathcal{M}_{\alpha, \beta} \rightarrow X_{\alpha, \beta}$ be the map induced by $(\tilde{e}_-, \tilde{e}_+)$.

The component $\partial^4: R = H_0(\alpha) \rightarrow H_3(\beta) = R$, for β irreducible, is then defined to be multiplication by the degree of the map

$$e: \mathcal{M}_{\alpha, \beta} \rightarrow X_{\alpha, \beta}.$$

From the dimension formula we have $\dim \mathcal{M}_{\alpha, \beta} = \dim \alpha = \dim X_{\alpha, \beta}$, so as these spaces are compact oriented manifolds this makes sense. Note that $X_{\alpha, \beta} \cong \tilde{\alpha}$ as β is irreducible.

For later purposes it will be important to relate the degree of the map $e: \mathcal{M}_{\alpha, \beta} \rightarrow X_{\alpha, \beta}$ to the degree of a different map determined by Austin in [Aus95]. Recall that if X and Y are closed, oriented manifolds of dimension n and Y is connected then the degree of a map $f: X \rightarrow Y$ is defined by the relation $f_*([X]) = \deg(f)[Y]$ where $f_*: H_n(X) \rightarrow H_n(Y)$ is the induced map and $[X], [Y]$ denote the fundamental classes. If X and Y in addition carry smooth structures and f is a smooth map one may also calculate the degree as

$$\deg(f) = \sum_{x \in f^{-1}(y)} \text{sgn}(df_x: T_x X \rightarrow T_y Y),$$

for a regular value y , where $\text{sgn}(df_x: T_x X \rightarrow T_y Y)$ is ± 1 depending on whether this map preserves or reverses orientation, respectively. The proof of the following proposition is a simple exercise in differential topology, so we leave it out.

Proposition 2.6. *In the statements below assume that X, Y and Z are smooth, closed, oriented manifolds and that Y and Z are connected.*

- (a) *Let G be a compact Lie group acting freely on the manifolds X and Y of the same dimension. Then if $f: X \rightarrow Y$ is a smooth G -equivariant map it holds true that*

$$\deg(f: X \rightarrow Y) = \deg(f/G: X/G \rightarrow Y/G).$$

- (b) Let $f: X \rightarrow Y$ be a smooth map between manifolds of equal dimension and let $p: \tilde{Y} \rightarrow Y$ be a finite, connected covering space with the induced orientation. Let $q: \tilde{X} = f^*\tilde{Y} \rightarrow X$ be the pull-back covering and let $\tilde{f}: \tilde{X} \rightarrow \tilde{Y}$ be the induced map. Then

$$\deg(f: X \rightarrow Y) = \deg(\tilde{f}: \tilde{X} \rightarrow \tilde{Y}).$$

- (c) Let $(f, g): X \rightarrow Y \times Z$ be a smooth map and assume that $\dim X = \dim Y + \dim Z$. If $y \in Y$ is a regular value for f , so that $f^{-1}(y) \subset X$ is a closed, oriented submanifold of dimension $\dim f^{-1}(y) = \dim Z$, then

$$\deg((f, g): X \rightarrow Y \times Z) = \deg(g|_{f^{-1}(y)}: f^{-1}(y) \rightarrow Z).$$

Lemma 2.7. Let $\alpha, \beta \in \mathcal{C}$ be a pair with β irreducible and $\text{gr}(\alpha, \beta) = 4$. Consider the diagram

$$\alpha \xleftarrow{e_-} \overline{\mathcal{M}}_z(\alpha, \beta) \xrightarrow{e_+} \beta,$$

where z is the relative homotopy class with $\text{gr}_z(\alpha, \beta) = 4$. Then if $j(\alpha) \equiv s \pmod{8}$, the component $H_0(\alpha) \rightarrow H_3(\beta)$ of $\partial^4: DCI_{s,0} \rightarrow DCI_{s-4,3}$ is given by multiplication by the degree of the map

$$e_+|_{e_-^{-1}(*): e_-^{-1}(*)} \rightarrow \beta$$

where $* \in \alpha$ is a point.

Proof. Write $\overline{\mathcal{M}}_{\alpha,\beta} = \overline{\mathcal{M}}_z(\alpha, \beta)$ as earlier. By part (c) of the above proposition we have

$$\deg(e_+|_{e_-^{-1}(*): e_-^{-1}(*)} \rightarrow \beta) = \deg((e_-, e_+): \overline{\mathcal{M}}_{\alpha,\beta} \rightarrow \alpha \times \beta),$$

and by part (b)

$$\deg((e_-, e_+): \overline{\mathcal{M}}_{\alpha,\beta} \rightarrow (\alpha \times \beta) = \deg((\tilde{e}_-, \tilde{e}_+): \overline{\mathcal{M}}_{\alpha,\beta}^{\text{cov}} \rightarrow \tilde{\alpha} \times \tilde{\beta}).$$

Finally by part (a) the latter integer coincides with the degree of

$$e: \overline{\mathcal{M}}_{\alpha,\beta}^{\text{cov}}/\text{SU}(2) = \mathcal{M}_{\alpha,\beta} \rightarrow X_{\alpha,\beta} = (\tilde{\alpha} \times \tilde{\beta})/\text{SU}(2),$$

which by definition is the component of ∂^4 from $R = H_0(\alpha) \rightarrow H_3(\beta) = R$. \square

Definition 2.8. Assume that $\text{gr}(\alpha, \beta) \equiv 0 \pmod{4}$ for all $\alpha, \beta \in \mathcal{C}$. Then the Donaldson model $DCI(Y, E)$ for the Floer complex is the chain complex associated with the multicomplex

$$(DCI(Y, E)_{*,*}, \{\partial^r\}_{r \geq 0}) \text{ where } DCI(Y, E)_{s,t} = \bigoplus_{\alpha \in \mathcal{C}: j(\alpha) \equiv s} H_t(\alpha),$$

∂^4 is given as in Lemma 2.7 and $\partial^r = 0$ for $r \neq 4$.

The complex $DCI(Y, E)$ enjoys the same periodicity as $\widetilde{CI}(Y, E)$. Furthermore, it also carries a natural filtration

$$F_p DCI(Y, E)_n := \bigoplus_{s \leq p} DCI(Y, E)_{s, n-s}$$

by $H_*(\text{SO}(3)) \cong \Lambda_R[u]$ -submodules. This filtration is also referred to as the index filtration.

Finally, we state the result that allows the replacement of the pair $(C_*^{gm}(\text{SO}(3)), \widetilde{CI}(Y, E))$ by the pair $(\Lambda_R[u], DCI(Y, E))$.

Proposition 2.9. [Eis19, Corollary 7.6] Assume that $2 \in R$ is invertible. There is a quasi-isomorphism $i: \Lambda_R[u] \rightarrow C_*^{gm}(\text{SO}(3); R)$ of differential graded algebras and there is a zigzag of $\Lambda_R[u]$ -equivariant quasi-isomorphisms

$$DCI(Y, E) \xrightarrow{f} X \xleftarrow{g} \widetilde{CI}(Y, E).$$

Furthermore, X also carries a filtration, f and g are filtration preserving and the induced maps

$$\frac{F_p DCI(Y, E)}{F_{p-1} DCI(Y, E)} \longrightarrow \frac{F_p X}{F_{p-1} X} \longleftarrow \frac{F_p \widetilde{CI}(Y, E)}{F_{p-1} \widetilde{CI}(Y, E)}$$

are quasi-isomorphisms for all p .

Remark 2.10. The statement in the cited result is slightly weaker. The fact that the intermediate objects carry a type of index filtration and that all the quasi-isomorphisms preserve this filtration, inducing quasi-isomorphisms on the minimal filtration quotients, is obtained by a close inspection of the quite involved proof.

3. BINARY POLYHEDRAL SPACES AND FLAT CONNECTIONS

The purpose of this section is to fix our conventions concerning the binary polyhedral spaces and to recall the basic representation theory needed to effectively classify the gauge equivalence classes of flat $SU(2)$ -connections. We also recall the McKay correspondence between the finite subgroups of $SU(2)$ and the simply laced extended Dynkin diagrams, that is, the diagrams of type \tilde{A}_n , \tilde{D}_n , \tilde{E}_6 , \tilde{E}_7 or \tilde{E}_8 . This surprising relation is vital for our work in the next section and is essential in the definition of the graphs \mathcal{S}_Γ , given in the end of the section, mentioned in the introduction.

To avoid making this rather basic section too long we have moved a number of facts concerning the finite subgroups of $SU(2)$ into Appendix A. This includes their complex representation theory, the corresponding McKay graphs used in the construction of \mathcal{S}_Γ and complete lists of the flat $SU(2)$ -connections over the binary polyhedral spaces.

3.1. Binary Polyhedral Spaces. We briefly recall the classification of finite subgroups of $SU(2)$. Let C_l denote the cyclic group of order l for $l \geq 1$, let D_k denote the dihedral group of order $2k$ for $k \geq 2$, and let T , O and I denote the subgroups of $SO(3)$ that leave a regular tetrahedron, octahedron and icosahedron, respectively, in \mathbb{R}^3 invariant. T is called the tetrahedral group, O the octahedral group and I the icosahedral group. Define D_k^* , T^* , O^* and I^* in $SU(2)$ by pulling back D_k , T , O and I along the standard double covering homomorphism $SU(2) \rightarrow SO(3)$. Every cyclic group C_l may also be realized as a subgroup of $SU(2)$ by embedding them in a copy of $U(1) \subset SU(2)$. We have thus constructed two infinite families C_l , D_k^* of subgroups in $SU(2)$, as well as the three exceptional ones T^* , O^* and I^* . It is a classical fact that up to conjugacy these exhaust all the finite subgroups of $SU(2)$. A proof may be found in [Wol11, p. 83].

Theorem 3.1. *Let $\Gamma \subset SU(2)$ be a finite subgroup. Then Γ is isomorphic to precisely one of the groups C_l for $l \geq 1$, D_k^* for $k \geq 2$, T^* , O^* or I^* . Moreover, if two finite subgroups $\Gamma, \Gamma' \subset SU(2)$ are isomorphic, then they are conjugate in $SU(2)$.*

Let \mathbb{C}^2 carry the canonical complex orientation and orient $S^3 \subset \mathbb{C}^2$ by the outward pointing normal first convention; that is, an ordered basis $(v_1, v_2, v_3) \in T_x S^3 \subset T_x \mathbb{C}^2 \cong \mathbb{C}^2$ is positive if and only if (x, v_1, v_2, v_3) is a positive basis for \mathbb{C}^2 . The standard left action of $SU(2)$ on \mathbb{C}^2 restricts to a transitive and free action by orientation preserving isometries on the unit sphere $S^3 \subset \mathbb{C}^2$.

Definition 3.2. For any finite subgroup $\Gamma \subset SU(2)$ let Γ act on $S^3 \subset \mathbb{C}^2$ by restricting the standard free action of $SU(2)$. Define Y_Γ to be the quotient manifold S^3/Γ equipped with the Riemannian metric and orientation induced from the standard round metric and orientation of S^3 . Define \overline{Y}_Γ to be the same Riemannian manifold equipped with the opposite orientation. The spaces Y_Γ and \overline{Y}_Γ are called binary polyhedral spaces.

Note that by the above theorem Y_Γ is determined up to $SU(2)$ -equivariant isometry by the isomorphism class of Γ .

Lemma 3.3. *For $\Gamma \subset \mathrm{SU}(2)$ we have*

$$H_i(Y_\Gamma; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{for } i = 0, 3 \\ \Gamma^{ab} & \text{for } i = 1 \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad H^i(Y_\Gamma; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{for } i = 0, 3 \\ \Gamma^{ab} & \text{for } i = 2 \\ 0 & \text{otherwise,} \end{cases}$$

where $\Gamma^{ab} = \Gamma/[\Gamma, \Gamma]$ is the abelianization of Γ . In particular, $H_*(Y_\Gamma; \mathbb{Q}) \cong H_*(S^3; \mathbb{Q})$, so the Y_Γ are rational homology 3-spheres.

Proof. As Y_Γ is a connected, closed and orientable the (co)homology is concentrated in degrees $0 \leq i \leq 3$ and $H_i(Y_\Gamma) \cong H^i(Y_\Gamma) \cong \mathbb{Z}$ for $i = 0, 3$. By construction, S^3 is a universal cover of Y_Γ so that $\pi_1(Y_\Gamma) \cong \Gamma$ and hence $H_1(Y_\Gamma) \cong \Gamma^{ab}$. Since this group is finite it follows by the universal coefficient theorem that $H^1(Y_\Gamma) = 0$. By Poincaré duality we conclude that $H_2(Y_\Gamma) = 0$ and $H^2(Y_\Gamma) \cong \Gamma^{ab}$. The final statement follows from the universal coefficient theorem as $H_1(Y_\Gamma; \mathbb{Q}) \cong \mathbb{Q} \otimes \Gamma^{ab} = 0$. \square

3.2. A Few Results from Representation Theory. The framed Floer complex $\widetilde{CI}(Y, E)$ associated with the trivial $\mathrm{SU}(2)$ -bundle $E \rightarrow Y_\Gamma$ is generated, in the sense of Definition 2.4, by the gauge equivalence classes of the flat connections in E . By a well-known result (see for instance [Tau11] for a proof) these gauge equivalence classes are in natural bijection with the equivalence classes of representations

$$\Gamma = \pi_1(Y_\Gamma) \rightarrow \mathrm{SU}(2)$$

or in other words, as $\mathrm{SU}(2) \cong \mathrm{Sp}(1)$, the isomorphism classes of 1-dimensional quaternionic representations of Γ . It is convenient to express this set in terms of the complex representation theory of Γ , since in this setting we have the calculational power of character theory at our disposal. The following three results may be extracted from [BD85, p. II.6].

Lemma 3.4. *Let G be a finite group. The set $\mathrm{Irr}(G, \mathbb{C})$ of isomorphism classes of irreducible complex representations admits a decomposition into disjoint subsets*

$$\mathrm{Irr}(G, \mathbb{C}) = \mathrm{Irr}(G, \mathbb{C})_{\mathbb{R}} \cup \mathrm{Irr}(G, \mathbb{C})_{\mathbb{C}} \cup \mathrm{Irr}(G, \mathbb{C})_{\mathbb{H}}.$$

The set $\mathrm{Irr}(G, \mathbb{C})_{\mathbb{C}}$ consists of the classes $[V]$ where $V \not\cong V^$, while $\mathrm{Irr}(G, \mathbb{C})_{\mathbb{R}}$ and $\mathrm{Irr}(G, \mathbb{C})_{\mathbb{H}}$ consists of the classes $[W]$ admitting a conjugate linear equivariant map $s : W \rightarrow W$ satisfying $s^2 = 1$ or $s^2 = -1$, respectively.*

An irreducible representation V is said to be of real, complex or quaternionic type according to whether $[V] \in \mathrm{Irr}(G, \mathbb{C})_K$ for $K = \mathbb{R}, \mathbb{C}, \mathbb{H}$, respectively. There is a simple test for determining the type of an irreducible complex representation.

Lemma 3.5. *Let V be an irreducible complex representation of the finite group G with character χ_V . Then*

$$\frac{1}{|G|} \sum_{g \in G} \chi_V(g^2) = \begin{cases} 1 & \text{if } [V] \in \mathrm{Irr}(G, \mathbb{C})_{\mathbb{R}} \\ 0 & \text{if } [V] \in \mathrm{Irr}(G, \mathbb{C})_{\mathbb{C}} \\ -1 & \text{if } [V] \in \mathrm{Irr}(G, \mathbb{C})_{\mathbb{H}} \end{cases}$$

Finally we recall how the set $\mathrm{Irr}(G, \mathbb{H})$ of isomorphism classes of irreducible quaternionic representations may be recovered from $\mathrm{Irr}(G, \mathbb{C})$. Let $\mathrm{Rep}_K(G)$ denote the category of finite dimensional G -representations over $K = \mathbb{C}, \mathbb{H}$. There is a restriction functor $r : \mathrm{Rep}_{\mathbb{H}}(G) \rightarrow \mathrm{Rep}_{\mathbb{C}}(G)$ given by pullback along the inclusion $\mathbb{C} \hookrightarrow \mathbb{H}$ and an extension functor $e : \mathrm{Rep}_{\mathbb{C}}(G) \rightarrow \mathrm{Rep}_{\mathbb{H}}(G)$ given by $e(V) = \mathbb{H} \otimes_{\mathbb{C}} V$. Then we have the following result.

Proposition 3.6. *There is a decomposition into disjoint subsets*

$$\mathrm{Irr}(G, \mathbb{H}) = \mathrm{Irr}(G, \mathbb{H})_{\mathbb{R}} \cup \mathrm{Irr}(G, \mathbb{H})_{\mathbb{C}} \cup \mathrm{Irr}(G, \mathbb{H})_{\mathbb{H}}.$$

Furthermore, the following maps are bijections

$$\begin{aligned} r : \text{Irr}(G, \mathbb{H})_{\mathbb{H}} &\rightarrow \text{Irr}(G, \mathbb{C})_{\mathbb{H}} \\ e : \frac{1}{2} \text{Irr}(G, \mathbb{C})_{\mathbb{C}} &\rightarrow \text{Irr}(G, \mathbb{H})_{\mathbb{C}} \\ e : \text{Irr}(G, \mathbb{C})_{\mathbb{R}} &\rightarrow \text{Irr}(G, \mathbb{H})_{\mathbb{R}}, \end{aligned}$$

where $\frac{1}{2} \text{Irr}(G, \mathbb{C})_{\mathbb{C}}$ denotes the set of unordered pairs $\{[V], [V^*]\}$ for $[V] \in \text{Irr}(G, \mathbb{C})_{\mathbb{C}}$.

Let $\text{Irr}^n(G, \mathbb{C})_K$ denote the set of irreducible G -representations of dimension n of type $K = \mathbb{R}, \mathbb{C}, \mathbb{H}$.

Corollary 3.7. *The set of isomorphism classes of 1-dimensional quaternionic representations of the finite group G are in bijection with*

$$\text{Irr}^2(G, \mathbb{C})_{\mathbb{H}} \cup \frac{1}{2} \text{Irr}^1(G, \mathbb{C})_{\mathbb{C}} \cup \text{Irr}^1(G, \mathbb{C})_{\mathbb{R}}.$$

In particular, if $G = \Gamma$ is a finite subgroup of $\text{SU}(2)$ then the above set is in bijection with the set of gauge equivalence classes of flat connections in the trivial $\text{SU}(2)$ -bundle over Y_{Γ} .

Note that a flat connection is irreducible, reducible or fully reducible if and only if the associated representation is of quaternionic, complex or real type, respectively.

3.3. Arrangement of the Flat Connections. The McKay correspondence [McK80] sets up a bijection between the isomorphism classes of the nontrivial finite subgroups of $\text{Sp}(1) \cong \text{SU}(2)$ and the simply-laced extended Dynkin diagrams. The extended Dynkin graph is constructed from the group Γ as follows. Let R_0, R_1, \dots, R_n be a complete set of representatives for the elements of $\text{Irr}(\Gamma, \mathbb{C})$, where we take R_0 to be the trivial representation. Let Q denote the two dimensional representation associated with the inclusion $\Gamma \subset \text{SU}(2)$ and define a matrix $A = (a_{ij})_{ij}$ by

$$Q \otimes R_i = \bigoplus_{j=0}^n a_{ij} R_j \quad \text{for } 0 \leq i \leq n.$$

One may show that A is symmetric with $a_{ii} = 0$ and $a_{ij} \in \{0, 1\}$. Define an unoriented graph $\overline{\Delta}_{\Gamma}$ by taking $I = \{0, 1, \dots, n\}$ as the set of vertices and an edge connecting i to j if and only if $a_{ij} = a_{ji} = 1$. This graph will then be an extended Dynkin diagram of type $\tilde{A}_n, \tilde{D}_n, \tilde{E}_6, \tilde{E}_7$ or \tilde{E}_8 . Furthermore, if we let Δ_{Γ} be the graph obtained from $\overline{\Delta}_{\Gamma}$ by deleting the vertex 0 corresponding to the trivial representation we obtain the underlying Dynkin diagram of type A_n, D_n, E_6, E_7 or E_8 . The precise correspondence is given in the following table where $l, k \geq 2$.

Γ	C_l	D_k^*	T^*	O^*	I^*
Δ_{Γ}	A_{l-1}	D_{k+2}	E_6	E_7	E_8

The set of vertices, say $\{1, 2, \dots, n\}$, in a Dynkin diagram corresponds to a set of simple roots $\{r_i : 1 \leq i \leq n\}$ in the associated root system. There is then a unique maximal positive root $r_{\max} = \sum_{i=1}^n d_i r_i$ where $d_i \in \mathbb{N}$ are positive integers. The final fact we wish to bring out is that these integers are determined by $d_i = \dim_{\mathbb{C}} R_i$.

By mapping an irreducible representation R_i to its dual representation R_i^* we obtain an involution $\iota : \{0, 1, \dots, n\} \rightarrow \{0, 1, \dots, n\}$, i.e., $R_{\iota(i)} \cong R_i^*$. This map extends to a graph involution of $\overline{\Delta}_{\Gamma}$ as

$$a_{\iota(i)\iota(j)} = \dim_{\mathbb{C}} \text{Hom}_{\Gamma}(Q \otimes R_i^*, R_j^*) = \dim_{\mathbb{C}} \text{Hom}_{\Gamma}(Q \otimes R_j, R_i) = a_{ji} = a_{ij}$$

for all $0 \leq i, j \leq n$. In view of Lemma 3.4 we see that the fixed points of ι correspond to the vertices of quaternionic or real type, while the nontrivial orbits are pairs $\{i, j\}$ for which $i \neq j$ and $R_i, R_j \cong R_i^*$ are of complex type. The set of vertices in the quotient graph $\overline{\Delta}_{\Gamma}/(\iota)$ may therefore

by Proposition 3.6 be identified with $\text{Irr}(\Gamma, \mathbb{H})$. In particular, we may identify our set \mathcal{C} of flat connections with a subset of the vertices in this quotient graph. The following may be observed from the quotient graphs given in Appendix A.

Lemma 3.8. *For each finite subgroup $\Gamma \subset \text{SU}(2)$ the quotient graph $\overline{\Delta}_\Gamma/(\iota)$ is a connected tree, except for $\Gamma = C_l$ with l odd in which case it has the form*



From the above lemma it follows that for each pair of vertices there is a unique minimal edge path connecting them.

Definition 3.9. [Aus95, p. 297] Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup. Define a graph \mathcal{S}_Γ by letting the vertices be the set of 1-dimensional quaternionic representations of Γ and an (unoriented) edge connecting a pair of distinct vertices α and β if and only if the minimal edge path connecting α to β in $\overline{\Delta}_\Gamma/(\iota)$ does not pass through a vertex corresponding to a 1-dimensional quaternionic representation different from α and β .

Two vertices in \mathcal{S}_Γ are said to be adjacent if there is an edge connecting them.

At the end of the next section we will attach symbols to the edges of the graph \mathcal{S}_Γ such that the resulting labeled graph contains all the necessary information needed to set up the complex $\text{DCI}(\overline{Y}_\Gamma)$. All of the graphs \mathcal{S}_Γ will be given in Proposition 4.25.

4. DETERMINATION OF MODULI SPACES AND DIFFERENTIALS

In this section we review the key results that allow us to determine the Donaldson models $\text{DCI}(\overline{Y}_\Gamma)$, for the finite subgroups $\Gamma \subset \text{SU}(2)$, explicitly. The main result of [Aus95] necessary for the calculations, originally due to Kronheimer, is the following.

Theorem 4.1. *Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup and let α, β be flat connections in the trivial $\text{SU}(2)$ -bundle over \overline{Y}_Γ . Let z be the unique homotopy class from α to β with $-2 \leq \text{gr}_z(\alpha, \beta) \leq 5$. Then if α and β are adjacent in \mathcal{S}_Γ (see Def. 3.9) then in the diagram*

$$\alpha \xleftarrow{e_-} \overline{\mathcal{M}}_z(\alpha, \beta) \xrightarrow{e_+} \beta$$

there is an identification $e_-^{-1}() \cong Y_{\Gamma'}$ for some other finite subgroup $\Gamma' \subset \text{SU}(2)$. If α and β are not adjacent in \mathcal{S}_Γ , then $\overline{\mathcal{M}}_z(\alpha, \beta)$ is empty.*

There is also fairly simple graphical procedure to determine the subgroup Γ' from the positions of α and β in the extended Dynkin diagram $\overline{\Delta}_\Gamma$ associated with Γ under the McKay correspondence (see Proposition 4.12).

The main idea of the proof is to translate the problem of determining instantons over $\mathbb{R} \times Y_\Gamma$ with L^2 curvature modulo gauge transformations into the problem of determining Γ -invariant instantons in Γ -equivariant $\text{SU}(2)$ -bundles over S^4 modulo Γ -invariant gauge transformations, and then to solve the latter problem with the use of the equivariant ADHM correspondence.

We begin by setting up the first correspondence. Recall that the instantons on the trivial $\text{SU}(2)$ -bundle over the cylinder $\mathbb{R} \times Y_\Gamma$ with L^2 -curvature are naturally partitioned into subsets by three pieces of data: the limiting flat connections $\alpha_\pm \in \mathcal{C} \cong \text{Irr}^1(\Gamma, \mathbb{H})$ and the relative homotopy class z traced out by the connection. For our purpose it is more convenient to replace this latter invariant by the relative Chern number (see [Don02, p. 48])

$$\hat{c}_2(A) := \frac{1}{8\pi^2} \int_{\mathbb{R} \times Y_\Gamma} \text{tr } F_A^2 = \frac{1}{8\pi^2} \int_{\mathbb{R} \times Y_\Gamma} |F_A|^2 \in \mathbb{R}.$$

Thus to any instanton A with L^2 -curvature over $\mathbb{R} \times Y_\Gamma$ we may associate a unique triple

$$(\alpha_+, \alpha_-, \hat{c}_2(A)) \in \text{Irr}^1(\Gamma, \mathbb{H}) \times \text{Irr}^1(\Gamma, \mathbb{H}) \times \mathbb{R}.$$

Let Γ act on $\mathbb{R} \times S^3$ by $\gamma \cdot (t, x) \mapsto (t, \gamma x)$ such that $q = (1 \times p): \mathbb{R} \times S^3 \rightarrow \mathbb{R} \times Y_\Gamma$ is an orientation preserving covering map. Regard $S^4 \subset \mathbb{C}^2 \oplus \mathbb{R}$ such that the suspended action $\Gamma \times S^4 \rightarrow S^4$ takes the form $\gamma \cdot (x, t) = (\gamma x, t)$. By composing the conformal equivalence $\mathbb{R} \times S^3 \cong \mathbb{C}^2 - \{0\}$, $(t, x) \mapsto e^t x$, with the inverse of stereographic projection from the north pole we obtain an equivariant conformal equivalence $\tau: \mathbb{R} \times S^3 \cong S^4 - \{N, S\}$, where $N, S = (0, \pm 1) \in S^4$ are the poles. Explicitly,

$$\tau(t, x) = (\cosh(t)x, \tanh(t)).$$

Let S^4 carry the orientation that makes this map orientation preserving, that is, the opposite of the standard orientation.

Recall that a Γ -equivariant vector-bundle over a Γ -space X is a vector bundle $E \rightarrow X$ with a Γ -action on the total space making the projection equivariant and such that $\gamma: E_x \rightarrow E_{\gamma x}$ is linear for each $x \in X$ and $\gamma \in \Gamma$. If E carries an $\text{SU}(2)$ -structure, that is, a Hermitian metric and a fixed unitary trivialization of $\Lambda_{\mathbb{C}}^2 E$, then we call E a Γ -equivariant $\text{SU}(2)$ -bundle provided the action preserves this additional structure. In that case, there is an induced action of Γ on the space of connections \mathcal{A}_E . A connection is said to be invariant if it is fixed under this action. The subspace of invariant connections is denoted by \mathcal{A}_E^Γ . The action of the full gauge group does not preserve this subspace, but the subgroup of equivariant gauge transformations $\mathcal{G}_E^\Gamma \subset \mathcal{G}_E$ does. Therefore, the natural configuration space in this equivariant setting is $\mathcal{B}_{E, \Gamma} := \mathcal{A}_E^\Gamma / \mathcal{G}_E^\Gamma$. We may also include a framing coordinate to obtain the framed configuration space $\tilde{\mathcal{B}}_{E, \Gamma}$. All of the above may, of course, be spelled out in terms of principal bundles as well.

Let $F \rightarrow \mathbb{R} \times Y_\Gamma$ be the trivial $\text{SU}(2)$ -bundle. Then $E := q^*F \rightarrow \mathbb{R} \times S^3$ obtains the structure of a Γ -equivariant vector bundle in a natural way. Moreover, since q is a finite covering map, pullback induces a bijection $\mathcal{A}_F \cong \mathcal{A}_E^\Gamma$ that matches the subsets of anti-self-dual connections with curvature in L^2 . Let B be an ASD connection in F with curvature in L^2 and let $A = q^*(B)$ be the pullback connection in E . According to Uhlenbeck's removable singularities theorem there exists a bundle $\tilde{E} \rightarrow S^4$ with connection \tilde{A} and a bundle map $\rho: E \rightarrow \tilde{E}$ covering $\tau: \mathbb{R} \times S^3 \rightarrow S^4$ such that $\rho^*(\tilde{A}) = A$. The following diagram summarizes the situation

$$(4.1) \quad \begin{array}{ccccc} (F, B) & \longleftarrow & (E, A) & \xrightarrow{\rho} & (\tilde{E}, \tilde{A}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{R} \times Y_\Gamma & \xleftarrow{q} & \mathbb{R} \times S^3 & \xrightarrow{\tau} & S^4. \end{array}$$

According to Austin [Aus95] the Γ -action extends over \tilde{E} making \tilde{A} invariant and the flat limiting connections of the connection B are encoded in the isomorphism classes of the representations \tilde{E}_N and \tilde{E}_S over the fixed points. Furthermore, the following calculation shows that $c_2(\tilde{E})$ is uniquely determined by $\hat{c}_2(B)$

$$(4.2) \quad \begin{aligned} 8\pi^2 c_2(\tilde{E})[S^4] &= \int_{S^4} \text{tr}(F_{\tilde{A}}^2) = \int_{S^4} |F_{\tilde{A}}|^2 = \int_{\mathbb{R} \times S^3} |F_A|^2 \\ &= |\Gamma| \int_{\mathbb{R} \times Y_\Gamma} |F_B|^2 = 8\pi^2 |\Gamma| \hat{c}_2(B). \end{aligned}$$

Here we have used the fact that the Chern-Weil integrand $\text{tr}(F_A^2)$ coincides pointwise with the norm $|F_A|^2$ for ASD connections, the conformal invariance of the integral of 2-forms in dimension four and that $\mathbb{R} \times S^3 \rightarrow \mathbb{R} \times Y_\Gamma$ is a finite Riemannian covering map with fibers of cardinality $|\Gamma|$.

The following theorem classifies the Γ -equivariant $SU(2)$ -bundles over S^4 . The proof is given in Appendix B (see Theorem B.6).

Theorem 4.2. *A Γ -equivariant $SU(2)$ -bundle $\tilde{E} \rightarrow S^4$ is determined up to isomorphism by the ordered triple $([\tilde{E}_N], [\tilde{E}_S], k)$, where $k = c_2(\tilde{E})[S^4]$ is the second Chern number of the underlying $SU(2)$ -bundle and $[\tilde{E}_N], [\tilde{E}_S]$ denote the isomorphism classes of the Γ -representations over the fixed points. Moreover, for each pair $\alpha, \beta \in \text{Irr}^1(\Gamma, \mathbb{H})$ there is a constant $c \in \mathbb{Z}$ such that for each $k \equiv c \pmod{|\Gamma|}$ there exists a Γ -equivariant $SU(2)$ -bundle with invariants (α, β, k) .*

Let $\tilde{E} \rightarrow S^4$ be a Γ -equivariant $SU(2)$ -bundle with invariants (α_+, α_-, k) . Let $E = \tau^*(E) \rightarrow \mathbb{R} \times S^3$ and $F = E/\Gamma \rightarrow \mathbb{R} \times Y_\Gamma$. Then we may reverse the above procedure to produce a map from the invariant instantons in \tilde{E} to the instantons over $\mathbb{R} \times Y_\Gamma$ with limiting flat connection α_\pm and relative Chern number $k/|\Gamma|$. More is true, the Γ -equivariant gauge transformations in \tilde{E} correspond to gauge transformations in F that approach α_\pm -harmonic gauge transformations on the ends. Let $b_0 \in Y_\Gamma$ denote a basepoint and let $\tilde{b} \in S^3$ be a lift. Let $\tilde{\mathcal{M}}_{\Gamma, \infty}(\tilde{E}) \subset \tilde{\mathcal{B}}_{\tilde{E}, \Gamma}$ be the moduli space of invariant instantons in \tilde{E} framed at $(0, \tilde{b})$. Then the above discussion yields a bijective correspondence

$$\tilde{\mathcal{M}}_\Gamma(\tilde{E}) \cong \tilde{\mathcal{M}}_z(\alpha_-, \alpha_+),$$

where z is the relative homotopy class corresponding to the relative Chern number $k/|\Gamma|$ and the latter space is framed at $(0, b) \in \mathbb{R} \times Y_\Gamma$. Recall that the end-point maps $e_\pm : \tilde{\mathcal{M}}_z(\alpha_-, \alpha_+) \rightarrow \alpha_\pm$ were defined by parallel transport of the framing to $\pm\infty$. Under the above bijection this corresponds to parallel transport of the framing along the great circle through the basepoint $(\tilde{b}, 0) \in S^4$ and the poles $N, S = (0, \pm 1)$. The equivariant ADHM correspondence, which we turn to next, describes the instanton moduli space with fixed framing at N and we then see that this corresponds to the fiber $e_+^{-1}(*)$. If we change the orientation in Y_Γ we may precompose with the map $(t, x) \mapsto (-t, x)$ in $\mathbb{R} \times Y_\Gamma$ to see that the ADHM correspondence will describe the fiber $e_-^{-1}(*)$ instead.

4.1. The ADHM Description of Instantons. We first recall the non-equivariant ADHM classification of $SU(2)$ -instantons over S^4 [DK90, Sec. 3.3]. This construction describes the various moduli spaces of instantons over S^4 in terms of linear algebraic data. In the following we regard $\mathbb{C}^2 \cup \{\infty\} = S^4$ where the orientation is determined by the complex orientation on \mathbb{C}^2 . Recall that an $SU(2)$ -bundle over S^4 , or any other closed, oriented 4-manifold, is determined up to isomorphism by the second Chern number $k = c_2(E)[S^4] \in \mathbb{Z}$. The bundle can only support an ASD connection if $k \geq 0$ and in case $k = 0$ every ASD connection is flat.

For a fixed integer $k \geq 1$ let \mathcal{H} be a Hermitian vector space of dimension k and let E_∞ be a 2-dimensional complex vector space with symmetry group $SU(2)$. Define the complex vector space

$$M := \text{Hom}(\mathbb{C}^2 \otimes \mathcal{H}, \mathcal{H}) \oplus \text{Hom}(\mathcal{H}, E_\infty) \oplus \text{Hom}(E_\infty, \mathcal{H}).$$

By identifying $\mathbb{C}^2 \otimes \mathcal{H} = \mathcal{H} \oplus \mathcal{H}$ we may write an element $m \in M$ in terms of its components $m = (\tau_1, \tau_2, \pi, \sigma)$ where $\tau_1, \tau_2 \in \text{End}(\mathcal{H})$, $\pi : \mathcal{H} \rightarrow E_\infty$ and $\sigma : E_\infty \rightarrow \mathcal{H}$. Let $U(\mathcal{H})$, the group of unitary automorphisms of \mathcal{H} , act on M by

$$g \cdot (\tau_1, \tau_2, \pi, \sigma) = (g\tau_1g^{-1}, g\tau_2g^{-1}, \pi g^{-1}, g\sigma).$$

Set $\mathfrak{u}(\mathcal{H}) := \text{Lie}(U(\mathcal{H}))$ and define $\mu = (\mu_\mathbb{C}, \mu_\mathbb{R}) : M \rightarrow \text{End}(\mathcal{H}) \oplus \mathfrak{u}(\mathcal{H})$ by

$$(4.3) \quad \begin{aligned} \mu_\mathbb{C}(\tau_1, \tau_2, \pi, \sigma) &= [\tau_1, \tau_2] + \sigma\pi \\ \mu_\mathbb{R}(\tau_1, \tau_2, \pi, \sigma) &= [\tau_1, \tau_1^*] + [\tau_2, \tau_2^*] + \sigma\sigma^* - \pi^*\pi. \end{aligned}$$

These maps are equivariant when $U(\mathcal{H})$ acts by conjugation on $\text{End}(\mathcal{H})$ and $\mathfrak{u}(\mathcal{H})$.

Given $(\tau_1, \tau_2, \pi, \sigma) \in M$ we may for each $z = (z_1, z_2) \in \mathbb{C}^2$ form the sequence

$$(4.4) \quad \mathcal{H} \xrightarrow{A_z} \mathbb{C}^2 \otimes \mathcal{H} \oplus E_\infty \xrightarrow{B_z} \mathcal{H}$$

where

$$A_z = \begin{pmatrix} \tau_1 - z_1 \\ \tau_2 - z_2 \\ \pi \end{pmatrix} \quad \text{and} \quad B_z = \begin{pmatrix} -(\tau_2 - z_2) & (\tau_1 - z_1) & \sigma \end{pmatrix}.$$

The condition $\mu_{\mathbb{C}}(\tau_1, \tau_2, \pi, \sigma) = 0$ is equivalent to $B_z \circ A_z = 0$ for all $z \in \mathbb{C}^2$. Define $U_{reg} \subset M$ to be the open set consisting of the $(\tau_1, \tau_2, \pi, \sigma)$ for which A_z is injective and B_z is surjective for all $z \in \mathbb{C}^2$. From the sequence (4.4) we may form a bundle homomorphism R between the trivial bundles over \mathbb{C}^2 with fibers $\mathbb{C}^2 \otimes \mathcal{H}$ and $\mathbb{C}^2 \otimes \mathcal{H} \oplus E_\infty$, respectively, by $R_z = (A_z, B_z^*)$. This map extends naturally to a bundle homomorphism

$$\begin{array}{ccc} \gamma \oplus \gamma \oplus \cdots \oplus \gamma & \xrightarrow{R} & \mathbb{C}^2 \otimes \mathcal{H} \oplus E_\infty \\ & \searrow & \swarrow \\ & \mathbb{C}^2 \cup \{\infty\} = S^4, & \end{array}$$

where γ is the $SU(2)$ -bundle with $c_2(\gamma)[S^4] = -1$ and we take k copies of γ on the left hand side. This fact is most easily understood using quaternions and the model $\mathbb{H}P^1 \cong S^4$ (see [Ati79]). If $m = (\tau_1, \tau_2, \pi, \sigma) \in U_{reg} \cap \mu_{\mathbb{C}}^{-1}(0)$ it follows that R is everywhere injective and we obtain an $SU(2)$ -bundle

$$E := \text{Im}(R)^\perp \cong \text{Coker}(R)$$

with $c_2(E)[S^4] = k$, and the fiber over the point at ∞ is the space E_∞ we started with. This bundle carries a natural connection A obtained by orthogonal projection from the product connection in the trivial bundle. This connection is ASD provided $\mu_{\mathbb{R}}(m) = 0$ as well.

The above construction associates a pair (E, A) consisting of an $SU(2)$ -bundle $E \rightarrow S^4$ with $c_2(E) = k$ and an ASD connection A in E to each element $m \in U_{reg} \cap \mu^{-1}(0)$. If m is replaced by $g \cdot m$ for $g \in U(\mathcal{H})$ the resulting pair (E', A') is equivalent to (E, A) by a bundle isomorphism $\rho^g : E \rightarrow E'$ fixing the fiber at ∞ . The content of the ADHM classification is that this construction produces all instantons in the $SU(2)$ -bundle over S^4 with Chern number k .

Theorem 4.3. (*The ADHM-Correspondence*) *The assignment $(\tau_1, \tau_2, \pi, \sigma) \mapsto (E, A)$ descends to an equivalence*

$$(U_{reg} \cap \mu^{-1}(0))/U(\mathcal{H}) \cong \mathcal{M}_\infty(k)$$

where $\mathcal{M}_\infty(k)$ denotes the moduli space of gauge equivalence classes of instantons framed at ∞ in the bundle of Chern number k over $S^4 = \mathbb{C}^2 \cup \{\infty\}$.

For the proof of the theorem and additional details we refer to [DK90, Section 3.3]. A part of the proof involves giving an inverse procedure to the construction sketched above. We will need one fact from this inverse procedure. Given an $SU(2)$ -bundle $E \rightarrow S^4$ with $c_2(E)[S^4] = k$ and ASD connection A , the space \mathcal{H} is recovered by

$$(4.5) \quad \text{Ker}[D_A^* : \Gamma(S^- \otimes E) \rightarrow \Gamma(S^+ \otimes E)],$$

where S^\pm are the complex spinor bundles associated with the unique spin structure on S^4 and D_A^* is the formal adjoint of the Dirac operator twisted by the connection A .

Let $\Gamma \subset SU(2)$ be a finite subgroup. Then according to [Aus95] the ADHM correspondence is natural with respect to the standard linear action $\Gamma \times \mathbb{C}^2 \rightarrow \mathbb{C}^2$ and the corresponding suspended action on $\mathbb{C}^2 \cup \{\infty\} = S^4$. This means that if $E \rightarrow S^4$ is a Γ -equivariant bundle with a Γ -invariant ASD connection A , then the spaces E_∞, \mathcal{H} are complex Γ -representations and $m = (\tau_1, \tau_2, \pi, \sigma) \in$

$U_{reg} \cap \mu^{-1}(0)$ associated with (E, A) belongs to the subspace M^Γ of invariants. Conversely, if \mathcal{H} and E_∞ are Γ -representations and $m \in U_{reg} \cap \mu^{-1}(0) \cap M^\Gamma$ then the associated bundle E naturally obtains the structure of a Γ -equivariant bundle such that the connection A is invariant.

As we have seen the Γ -equivariant $SU(2)$ -bundles over S^4 are determined up to isomorphism by the triple of invariants (α, β, k) as in Theorem 4.2. We need to know which representation \mathcal{H} that produces invariant instantons in the bundle with data (α, β, k) . The relevant formula is obtained by an application of the equivariant index theorem to the twisted Dirac operator in (4.5). Let $R(\Gamma)$ denote the complex representation ring of Γ and let $\epsilon : R(\Gamma) \rightarrow \mathbb{Z}$ denote the augmentation. Furthermore, let as before Q denote the 2-dimensional representation associated with the inclusion $\Gamma \subset SU(2)$.

Lemma 4.4. *Let $E \rightarrow S^4$ be a Γ -equivariant $SU(2)$ -bundle with invariants $([E_N], [E_S], c_2(E)[S^4]) = (\alpha, \beta, k)$ and set*

$$\mathcal{H} = \text{Ind}[D_A^* : \Gamma(S^- \otimes E) \rightarrow \Gamma(S^+ \otimes E)] \in R(\Gamma)$$

where A is a Γ -invariant connection in E . Then we have the following relations in the representation ring $R(\Gamma)$:

$$\begin{aligned} (2 - Q)\mathcal{H} &= \alpha - \beta \\ \epsilon(\mathcal{H}) &= k. \end{aligned}$$

These formulas are given in [Aus95, Sec. 4.3]. We have included a proof in Appendix B.11. The following lemma ensures that the above two equations determine the virtual representation \mathcal{H} uniquely.

Lemma 4.5. *Let $\Gamma \subset SU(2)$ be a finite subgroup and let Q be the canonical 2-dimensional representation of Γ . Then the kernel of the map $R(\Gamma) \rightarrow R(\Gamma)$ given by $V \mapsto (2 - Q)V$ is given by $\{nR : n \in \mathbb{Z}\}$ where $R \cong \mathbb{C}[\Gamma]$ denotes the regular representation.*

Proof. Assume that $(2 - Q)V = 0$ in $R(\Gamma)$. In terms of the associated characters this means that $(2 - \chi_Q(g))\chi_V(g) = 0$ for all $g \in \Gamma$. Since Q is a faithful representation of Γ it follows that $\chi_Q(g) \neq \chi_Q(1) = 2$ for all $g \neq 1$, from which we deduce that $\chi_V(g) = 0$ for all $g \neq 1$. There exist representations U, W of Γ such that $V = U - W$ in $R(\Gamma)$. Therefore,

$$\chi_V(1)/|\Gamma| = (\chi_V, 1_\Gamma) = (\chi_U, 1_\Gamma) - (\chi_W, 1_\Gamma) = \dim_{\mathbb{C}} U^\Gamma - \dim_{\mathbb{C}} W^\Gamma \in \mathbb{Z},$$

where 1_Γ is the character of the trivial representation and (\cdot, \cdot) is the Hermitian inner product on characters. As the character of the regular representation R is given by $\chi_R(1) = |\Gamma|$ and $\chi_R(g) = 0$ for $g \neq 1$, we conclude that for $m = \chi_V(1)/|\Gamma|$ we have $\chi_V = m\chi_R$, and hence $V = mR$ in $R(\Gamma)$. \square

Recall that $R(\Gamma)$ is, as an abelian group, freely generated by the isomorphism classes of the irreducible representations of Γ . Therefore, if we label the irreducible representations by R_0, \dots, R_m , then every element in $R(\Gamma)$ may be expressed uniquely as a linear combination $\sum_{i=0}^m n_i[R_i]$ for $n_i \in \mathbb{Z}$. Such a sum is called a virtual representation and in the case all $n_i \geq 0$ we call it an actual representation since it determines the representation $\bigoplus_i n_i R_i$ up to isomorphism.

We are now in a position to state the equivariant ADHM correspondence. This is a more precise version of [Aus95, Lemma 4.3].

Theorem 4.6. *(The Equivariant ADHM Correspondence) Let E be a Γ -equivariant $SU(2)$ -bundle over S^4 with $([E_N], [E_S], c_2(E)[S^4]) = (\alpha, \beta, k)$. Let $\mathcal{H} \in R(\Gamma)$ be the unique element satisfying $(2 - Q)\mathcal{H} = \alpha - \beta$ and $\epsilon(\mathcal{H}) = k$. Then if \mathcal{H} is an actual representations the moduli space $\widetilde{\mathcal{M}}_{\Gamma, \infty}(E)$ of Γ -invariant instantons modulo Γ -equivariant gauge transformations framed at $\infty = N$ is given by*

$$(U_{reg} \cap \mu^{-1}(0) \cap M^\Gamma) / U(\mathcal{H})^\Gamma,$$

where $U(\mathcal{H})^\Gamma$ is the group of unitary Γ -equivariant automorphisms of \mathcal{H} and

$$M^\Gamma = \text{Hom}_\Gamma(Q \otimes \mathcal{H}, \mathcal{H}) \oplus \text{Hom}_\Gamma(\mathcal{H}, E_\infty) \oplus \text{Hom}_\Gamma(E_\infty, \mathcal{H}).$$

If \mathcal{H} is not an actual representation, the moduli space is empty.

Note that in the above statement $E_\infty = E_N$.

4.2. Determination of the Low-Dimensional Moduli Spaces. The ADHM equations (4.3) are a particularly simple example of the hyper-Kähler moment map equations associated with an action of a Lie group on a hyper-Kähler manifold (see [Hit+87]). We briefly review the set up in the linear case relevant for our purpose.

Let V be a (left) quaternionic vector space equipped with a compatible real inner product $g: V \times V \rightarrow \mathbb{R}$, i.e., multiplication by unit quaternions are orthogonal. Let $\text{Sp}(V)$ denote the group of \mathbb{H} -linear orthogonal automorphisms of V . Define an $\text{Im } \mathbb{H} := \text{Span}_{\mathbb{R}}\{i, j, k\}$ valued 2-form $\omega: V \times V \rightarrow \text{Im } \mathbb{H}$ by the formula

$$\omega(v, w) = ig(iv, w) + jg(jv, w) + kg(kv, w).$$

Note that the unique symplectic inner product $s: V \times V \rightarrow \mathbb{H}$ determined by g is given by $s = g - \omega$.

Suppose that G is a compact Lie group acting on V through a homomorphism $\rho: G \rightarrow \text{Sp}(V)$. The Lie algebra \mathfrak{g} of G acts on V through the derivative $d\rho_1: \mathfrak{g} \rightarrow \text{sp}(V) = \text{Lie}(\text{Sp}(V))$. A hyper-Kähler moment map for the action $G \times V \rightarrow V$ is a G -equivariant map $\mu: V \rightarrow \text{Hom}(\mathfrak{g}, \text{Im } \mathbb{H})$, where G acts on the target through the adjoint representation, satisfying

$$d\mu_x(v) = [\xi \mapsto \omega(\xi \cdot x, v)] \in \text{Hom}(\mathfrak{g}, \text{Im } \mathbb{H})$$

for all $x \in V$ and $v \in V \cong T_x V$. A moment map is clearly unique up to a constant and one may check that the unique moment map vanishing at $0 \in V$ is given by

$$(4.6) \quad \mu(x) = [\xi \in \mathfrak{g} \mapsto \frac{1}{2}\omega(\xi \cdot x, x) \in \text{Im } \mathbb{H}].$$

Definition 4.7. For a pair $(V, \rho: G \rightarrow \text{Sp}(V, g))$ as above we define the hyper-Kähler quotient of V by G to be the space

$$\mu^{-1}(0)/G.$$

In general neither $\mu^{-1}(0)$ nor $\mu^{-1}(0)/G$ will be manifolds due to singularities. However, if $U \subset V$ is an open G -invariant subset on which G acts freely, then $\mu^{-1}(0) \cap U$ and the quotient by G will be manifolds. Moreover, in this case the quotient inherits the structure of a hyper-Kähler manifold.

Example 4.8. (The ADHM equations) Let \mathcal{H} and E_∞ be a pair of Hermitian vector space and define

$$V = \text{Hom}(\mathbb{C}^2 \otimes \mathcal{H}, \mathcal{H}) \oplus \text{Hom}(\mathcal{H}, E_\infty) \oplus \text{Hom}(E_\infty, \mathcal{H})$$

equipped with the Hermitian metric induced from \mathcal{H} and E_∞ . Using the standard basis of \mathbb{C}^2 an element of V is written $(\tau_1, \tau_2, \pi, \sigma)$. Let $G = U(\mathcal{H})$ act on V by $g \cdot (\tau_1, \tau_2, \pi, \sigma) = (g\tau_1 g^{-1}, g\tau_2 g^{-1}, \pi g^{-1}, g\sigma)$. Define a quaternionic structure map $J: V \rightarrow V$ by

$$J(\tau_1, \tau_2, \pi, \sigma) = (-\tau_2^*, \tau_1^*, -\sigma^*, \pi^*).$$

Then the action of G commutes with J and therefore defines a homomorphism $\rho: G \rightarrow \text{Sp}(V)$. If we use the trace inner product in $\mathfrak{u}(\mathcal{H})$ and the decomposition $\text{Im}(\mathbb{H}) \cong \mathbb{R} \oplus \mathbb{C}$ to identify

$$\text{Hom}(\mathfrak{u}(\mathcal{H}), \text{Im } \mathbb{H}) \cong \mathfrak{u}(\mathcal{H}) \otimes \text{Im}(\mathbb{H}) \cong \mathfrak{u}(\mathcal{H}) \oplus \mathfrak{u}(\mathcal{H}) \otimes \mathbb{C} \cong \mathfrak{u}(\mathcal{H}) \oplus \text{End}_{\mathbb{C}}(\mathcal{H}),$$

the resulting moment map $\mu: V \rightarrow \mathfrak{u}(\mathcal{H}) \oplus \text{End}_{\mathbb{C}}(\mathcal{H})$ vanishing at 0 is given by the ADHM equations in (4.3).

Kronheimer [Kro89] used this construction in the following situation. Let $\Gamma \subset \mathrm{SU}(2)$ be a finite subgroup and let Q and R denote the canonical representation and the regular representation, respectively. Assume that Q and R are equipped with Γ -invariant Hermitian metrics and fix a Γ -invariant quaternionic structure $j : Q \rightarrow Q$ compatible with the metric. Let

$$P = Q \otimes \mathrm{End}(R) \cong \mathrm{Hom}(Q \otimes R, R)$$

equipped with the induced Hermitian metric. By combining the quaternionic structure on Q with the real structure $f \mapsto f^*$ on $\mathrm{End}(R)$ one obtains a Γ -equivariant quaternionic structure $J : P \rightarrow P$ compatible with the Hermitian metric. As this structure is preserved by the action of Γ , the subspace $K = P^\Gamma$ carries the same structure.

The group $U(R)$ acts on P by conjugation in the second factor. This action preserves the quaternionic structure and the Hermitian metric. The subgroup $U(R)^\Gamma$ of Γ -equivariant unitary automorphisms preserves the subspace K and therefore defines a homomorphism $U(R)^\Gamma \rightarrow \mathrm{Sp}(K)$. Here, the subgroup $U(1)$ of scalars is contained in the kernel so the homomorphism descends to a homomorphism $F := U(R)^\Gamma / U(1) \rightarrow \mathrm{Sp}(K)$. Let $\mu : K \rightarrow \mathrm{Hom}(\mathfrak{f}, \mathrm{Im} \mathbb{H})$ be the unique hyper-Kähler moment vanishing at $0 \in K$. In this situation Kronheimer identified the hyper-Kähler quotient.

Lemma 4.9. [Kro89] *There is a homeomorphism $\mu^{-1}(0)/F \cong \mathbb{C}^2/\Gamma$ that restricts to an isometry*

$$\frac{\mu^{-1}(0) - \{0\}}{F} \cong \frac{\mathbb{C}^2 - \{0\}}{\Gamma}.$$

Remark 4.10. A consequence of this is that F acts freely on $\mu^{-1}(0) - \{0\}$.

The space K and the group F admit simple descriptions in terms of the McKay graph $\overline{\Delta}_\Gamma$ associated with Γ . Let R_0, \dots, R_m be a complete set of representatives for $\mathrm{Irr}(\Gamma, \mathbb{C})$, where we take $R_0 = \mathbb{C}$ to be the trivial representation. The regular representation $R = \mathbb{C}[\Gamma]$ decomposes as

$$R \cong \bigoplus_{i=0}^m n_i R_i,$$

where $n_i = \dim_{\mathbb{C}} R_i$. Recall from subsection 3.3 that a matrix A was defined by $Q \otimes R_i = \bigoplus_j a_{ij} R_j$, or equivalently $a_{ij} = \dim_{\mathbb{C}} \mathrm{Hom}_\Gamma(Q \otimes R_i, R_j)$. Furthermore, the McKay graph $\overline{\Delta}_\Gamma$ was defined by taking the irreducible representations as vertices and an (unoriented) edge connecting R_i to R_j precisely when $a_{ij} = a_{ji} = 1$. Using this

$$\begin{aligned} K = \mathrm{Hom}_\Gamma(Q \otimes R, R) &= \bigoplus_{i,j} \mathrm{Hom}_{\mathbb{C}}(\mathbb{C}^{n_i}, \mathbb{C}^{n_j}) \otimes \mathrm{Hom}_\Gamma(Q \otimes R_i, R_j) \\ &= \bigoplus_{i \rightarrow j \text{ in } \overline{\Delta}_\Gamma} \mathrm{Hom}(\mathbb{C}^{n_i}, \mathbb{C}^{n_j}) \end{aligned}$$

where each edge is repeated twice, once with each orientation, and

$$F = U(R)^\Gamma / U(1) = U \left(\bigoplus_i n_i R_i \right)^\Gamma / U(1) = \left(\prod_i U(n_i) \right) / U(1).$$

This has a simple graphical interpretation on the Dynkin graph $\overline{\Delta}_\Gamma$ corresponding to Γ . Attach the vector space \mathbb{C}^{n_i} to the vertex corresponding to R_i . Then an element of K is an assignment of a linear map $f_{ji} : \mathbb{C}^{n_i} \rightarrow \mathbb{C}^{n_j}$ whenever $a_{ij} = 1$. An element $g = (g_i)_i \in \prod_i U(n_i)$ acts on $f = (f_{ji})_{i,j} \in K$ by $(g \cdot f)_{ji} = g_j f_{ji} g_i^{-1}$.

Example 4.11. For $\Gamma = O^*$ an element of $K = \text{Hom}_{O^*}(Q \otimes R, R)$ is given by assigning a linear map to each arrow in the following diagram

$$\begin{array}{ccccccc} & & & \mathbb{C}^2 & & & \\ & & & \updownarrow & & & \\ \mathbb{C} & \rightleftarrows & \mathbb{C}^2 & \rightleftarrows & \mathbb{C}^3 & \rightleftarrows & \mathbb{C}^4 & \rightleftarrows & \mathbb{C}^3 & \rightleftarrows & \mathbb{C}^2 & \rightleftarrows & \mathbb{C} \end{array}$$

The next result is the key step in relating Kronheimer's hyper-Kähler quotient to the space occurring in the equivariant ADHM correspondence. Recall that an $\text{SU}(2)$ -representation α of Γ is either

- (1) of quaternionic type and irreducible as a complex representation,
- (2) of complex type and splits $\alpha = \lambda \oplus \lambda^*$ with $\lambda \not\cong \lambda^*$ or
- (3) of real type and splits $\alpha = 2\eta$ for some η with $\eta \cong \eta^*$.

By the vertex or vertices corresponding to α in $\overline{\Delta}_\Gamma$ we will mean the vertex corresponding to α in case (1), the vertices corresponding to λ and λ^* in case (2) and the vertex corresponding to η in case (3). The following is a more refined version of [Aus95, Lemma 4.6]. Austin gives no explicit proof, but states that one may verify it by a case by case analysis.

Proposition 4.12. *Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup and let α and β be $\text{SU}(2)$ -representations that are adjacent in the graph \mathcal{S}_Γ (Definition 3.9). Then the solution \mathcal{H} of the equation $(2-Q)\mathcal{H} = \alpha - \beta$ in $R(\Gamma)$ with minimal $\epsilon(\mathcal{H}) = k > 0$ is given in the following way.*

- (i) Delete the vertex or vertices corresponding to β in $\overline{\Delta}_\Gamma$. The resulting graph has one or two components. Let $\Delta_{\mathcal{H}}$ be the component containing α .
- (ii) Up to isomorphism there is a unique nontrivial finite subgroup $\Gamma' \subset \text{SU}(2)$ and an isomorphism of graphs $\phi: \Delta_{\Gamma'} \rightarrow \Delta_{\mathcal{H}}$.
- (iii) Let S_0, S_1, \dots, S_r be the irreducible representations of Γ' , where S_0 is the trivial representation. Then

$$\mathcal{H} := \bigoplus_{j=1}^r (\dim_{\mathbb{C}} S_j) R_{\phi(j)}.$$

Furthermore, if $Q' = m_1 S_{i_1} \oplus m_2 S_{i_2}$ is the isotypical decomposition of the canonical representation of Γ' , then $\alpha = m_1 R_{\phi(i_1)} \oplus m_2 R_{\phi(i_2)}$ is the isotypical decomposition of α . In particular, ϕ takes the vertex or vertices associated with Q' to the vertex or vertices corresponding to α .

Remark 4.13. The isotypical decomposition of the canonical representation Q of a finite subgroup $\Gamma \subset \text{SU}(2)$ is $Q = Q$ for Γ non-cyclic and $Q = \rho \oplus \rho^*$ for some 1-dimensional representation ρ for Γ cyclic.

Corollary 4.14. *Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup and let $\alpha \neq \beta$ be $\text{SU}(2)$ -representations that are not adjacent in \mathcal{S}_Γ . Let $\alpha = \alpha_0, \alpha_1, \dots, \alpha_s = \beta$ be the vertices of the minimal edge path connecting α to β in \mathcal{S}_Γ and let $\mathcal{H}_i \in R(\Gamma)$ be the minimal positive solution of the equation $(2-Q)\mathcal{H}_i = \alpha_i - \alpha_{i+1}$ for $0 \leq i \leq s-1$. Then $\mathcal{H} := \bigoplus_{i=0}^{s-1} \mathcal{H}_i$ is the minimal positive solution $(2-Q)\mathcal{H} = \alpha - \beta$ in $R(\Gamma)$.*

Proof. The calculation

$$(2-Q)\mathcal{H} = \sum_{i=0}^{s-1} (2-Q)\mathcal{H}_i = \sum_{i=0}^{s-1} \alpha_i - \alpha_{i+1} = \alpha_0 - \alpha_s = \alpha - \beta$$

shows that \mathcal{H} solves the relevant equation. By Lemma 4.5 every other solution is given by $\mathcal{H} + nR$ for some $n \in \mathbb{Z}$, where R is the regular representation. To see that \mathcal{H} is the minimal positive solution, it therefore suffices to show that $\mathcal{H} - R$ is not positive, i.e., not an actual representation.

Let $\Delta_i = \Delta_{\mathcal{H}_i}$ be the subgraph of $\overline{\Delta}_\Gamma$ described in the above proposition. By the construction of these graphs and Lemma 3.8 we conclude that $\Delta_0 \subset \Delta_1 \subset \dots \subset \Delta_{s-1}$. This implies that the representation \mathcal{H} is only supported on the vertices of Δ_{s-1} , which by construction does not contain the vertices corresponding to $\beta = \alpha_s$. Hence, in the isotypical decomposition $\mathcal{H} = \bigoplus_{i=0}^n k_i R_i$ there exists some i with $k_i = 0$. We therefore conclude that $\mathcal{H} - R$ is not an actual representation. \square

Let $\Gamma, \alpha, \beta, \mathcal{H}, \Gamma'$ and $\phi: \Delta_{\Gamma'} \rightarrow \Delta_{\mathcal{H}}$ be defined as in Proposition 4.12 and set

$$N = M^\Gamma = \text{Hom}_\Gamma(\mathcal{H} \otimes Q, \mathcal{H}) \oplus \text{Hom}_\Gamma(\mathcal{H}, E_\infty) \oplus \text{Hom}_\Gamma(E_\infty, \mathcal{H}),$$

as in Theorem 4.6. This space is endowed with a quaternionic structure and compatible metric preserved by the action of $G := U(\mathcal{H})^\Gamma$ as in Example 4.8. Note that $E_\infty = \alpha$. Let R' and Q' denote the regular and canonical representation of Γ' respectively and set

$$K = \text{Hom}_{\Gamma'}(Q' \otimes R', R') \quad \text{and} \quad F = U(R')^{\Gamma'} / U(1)$$

as in Kronheimer's construction. The following is a more precise version of [Aus95, Lemma 4.7].

Proposition 4.15. *In the above situation there is an isometric isomorphism $f: N \rightarrow K$ of quaternionic vector spaces and an isomorphism of groups $\tau: G \rightarrow F$ such that $f(g \cdot n) = \tau(g)f(n)$ for all $n \in N$ and $g \in G$.*

Proof. Let S be the regular representation of Γ' and write $S = S_0 \oplus S'$, where S_0 denotes the trivial representation. Then

$$K = \text{Hom}_{\Gamma'}(Q' \otimes S, S) = \text{Hom}_{\Gamma'}(Q' \otimes S', S') \oplus \text{Hom}_{\Gamma'}(Q', S') \oplus \text{Hom}_{\Gamma'}(S', Q')$$

since $\text{Hom}_{\Gamma'}(Q', S_0) = 0$ and $\text{Hom}_{\Gamma'}(Q' \otimes S', S_0) \cong \text{Hom}_{\Gamma'}(S', Q')$ as Q' is self-dual. We will now match these components with the components of N . Let $n_i = \dim_{\mathbb{C}} S_i$. Recall that the graph $\Delta_{\Gamma'}$ is the graph obtained from $\overline{\Delta}_{\Gamma'}$ by deleting the vertex corresponding to S_0 . Using part (iii) of the above proposition we can write $\mathcal{H} = \bigoplus_{i=1}^r n_i R_{\phi(i)}$ where $\phi: \Delta_{\Gamma'} \rightarrow \Delta_{\mathcal{H}}$ is the isomorphism of graphs. Then

$$\begin{aligned} (4.7) \quad \text{Hom}_\Gamma(Q \otimes \mathcal{H}, \mathcal{H}) &= \bigoplus_{i \rightarrow j \text{ in } \Delta_{\Gamma'}} \text{Hom}_{\mathbb{C}}(\mathbb{C}^{n_i}, \mathbb{C}^{n_j}) \otimes \text{Hom}_\Gamma(Q \otimes R_{\phi(i)}, R_{\phi(j)}) \\ &\cong \bigoplus_{i \rightarrow j \text{ in } \Delta_{\Gamma'}} \text{Hom}_{\mathbb{C}}(\mathbb{C}^{n_i}, \mathbb{C}^{n_j}) \otimes \text{Hom}_{\Gamma'}(Q' \otimes S_i, S_j) = \text{Hom}_\Gamma(Q' \otimes S', S'). \end{aligned}$$

To make this an \mathbb{H} -linear isometric isomorphism we specify the isomorphisms

$$(4.8) \quad \text{Hom}_\Gamma(Q \otimes R_{\phi(i)}, R_{\phi(j)}) \cong \mathbb{C} \cong \text{Hom}_{\Gamma'}(Q' \otimes S_i, S_j)$$

for i adjacent to j in $\Delta_{\Gamma'}$ as follows. Choose an arbitrary orientation on the edges in $\Delta_{\Gamma'}$. Then for each positive arrow $i \rightarrow j$ in $\Delta_{\Gamma'}$ we pick basis vectors $u_{ij} \in \text{Hom}_\Gamma(Q \otimes R_{\phi(i)}, R_{\phi(j)})$ and $v_{ij} \in \text{Hom}_{\Gamma'}(Q' \otimes S_i, S_j)$ of unit length. Then $u_{ji} = Ju_{ij} \in \text{Hom}_\Gamma(Q \otimes R_{\phi(j)}, R_{\phi(i)})$ and $v_{ji} = Jv_{ij} \in \text{Hom}_{\Gamma'}(Q' \otimes S_j, S_i)$ are also unit basis vectors. We then define the isomorphisms in (4.8) by sending u_{ij} to v_{ij} for each $i \rightarrow j$ in $\Delta_{\Gamma'}$.

Next, by the final part of the above proposition we know that if $Q' = m_1 S_{i_1} \oplus m_2 S_{i_2}$ is the isotypical decomposition, then $\alpha = E_\infty = m_1 R_{\phi(i_1)} \oplus m_2 R_{\phi(i_2)}$ is the isotypical decomposition of α . We may therefore identify

$$\begin{aligned} (4.9) \quad \text{Hom}_{\Gamma'}(S', Q') &= \bigoplus_{j=1}^2 \text{Hom}_{\mathbb{C}}(\mathbb{C}^{n_{i_j}}, \mathbb{C}^{m_j}) \otimes \text{Hom}_{\Gamma'}(S_{i_j}, S_{i_j}) \\ &\cong \bigoplus_{j=1}^2 \text{Hom}_{\mathbb{C}}(\mathbb{C}^{n_{i_j}}, \mathbb{C}^{m_j}) \otimes \text{Hom}_\Gamma(R_{\phi(i_j)}, R_{\phi(i_j)}) \cong \text{Hom}_\Gamma(\mathcal{H}, E_\infty) \end{aligned}$$

and similarly $\text{Hom}_{\Gamma'}(Q', S') \cong \text{Hom}_{\Gamma}(E_{\infty}, \mathcal{H})$. Once again one should match unit vectors carefully to ensure that the direct sum of these isomorphisms is an \mathbb{H} -linear isometry. The direct sum of these isomorphisms gives the required \mathbb{H} -linear isometry $N \cong K$.

The (orthogonal) decomposition $\mathcal{H} = \bigoplus_{i=1}^r n_i R_{\phi(i)}$ induces an isomorphism $G = U(\mathcal{H})^{\Gamma} \cong \prod_{i=1}^r U(n_i)$ and this group only acts on the matrix components in the decompositions (4.7) and (4.9). In the latter case it only acts on factors coming from \mathcal{H} . Moreover,

$$F \cong \left(\prod_{i=0}^r U(n_i) \right) / U(1) \cong \prod_{i=1}^r U(n_i)$$

as $n_0 = 1$, and the action can again be seen to be on the matrix component in (4.7) and (4.9), and in the latter case only on the factors coming from S' . Therefore, the isomorphism

$$F \cong \prod_{i=1}^r U(n_i) \cong G$$

does the trick and the proof is complete. \square

Proposition 4.16. *Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup and let α and β be $\text{SU}(2)$ -representations adjacent in the graph \mathcal{S}_{Γ} . Let $E \rightarrow S^4$ be the Γ -equivariant $\text{SU}(2)$ -bundle with $[E_N] = \alpha$, $[E_S] = \beta$ and minimal $k = c_2(E)[S^4] > 0$. Then the moduli space $\widetilde{\mathcal{M}}_{\Gamma, \infty}(E)$ of Γ -invariant instantons in E framed at $N = \infty$ is given by*

$$(\mathbb{C}^2 - \{0\})/\Gamma' \cong \mathbb{R} \times S^3/\Gamma'$$

where the finite subgroup $\Gamma' \subset \text{SU}(2)$ is determined as in Proposition 4.12.

Proof. We continue to use the notation of the two preceding propositions. According to the equivariant ADHM correspondence the moduli space $\widetilde{\mathcal{M}}_{\Gamma, \infty}(E)$ is given by

$$(4.10) \quad \frac{\mu^{-1}(0) \cap U_{\text{reg}} \cap M^{\Gamma}}{U(\mathcal{H})^{\Gamma}},$$

where in the above notation $N = M^{\Gamma}$ and $G = U(\mathcal{H})^{\Gamma}$. Let $\mu_N: N \rightarrow \mathfrak{g} \otimes \text{Im } \mathbb{H}$ and $\mu_K: K \rightarrow \mathfrak{f} \otimes \text{Im } \mathbb{H}$ be the unique hyper-Kähler moment maps vanishing at 0 associated with the action of G on N and F on K . Due to the uniqueness of the moment map it follows that the isomorphisms $f: N \rightarrow K$ and $\psi: G \rightarrow F$ of the previous proposition induce a homeomorphism $\mu_N^{-1}(0) \cong \mu_K^{-1}(0)$ equivariant along the map ψ . The group F acts freely on $\mu_K^{-1}(0) - \{0\}$ (see Remark 4.10), so by Lemma 4.9 we obtain isometries of hyper-Kähler 4-manifolds

$$\frac{\mu_N^{-1}(0) - \{0\}}{G} \cong \frac{\mu_K^{-1}(0) - \{0\}}{F} \cong \frac{\mathbb{C}^2 - \{0\}}{\Gamma'}.$$

To complete the proof we have to show that the left hand side coincides with the space in (4.10). This will be achieved by proving that

- (1) μ_N is the restriction of μ to $N = M^{\Gamma} \subset M$, so that $\mu_N^{-1}(0) = \mu^{-1}(0) \cap N$ and
- (2) $U_{\text{reg}} \cap \mu_N^{-1}(0) = \mu_N^{-1}(0) - \{0\}$.

We may decompose an element $m \in M^{\Gamma}$ as

$$m = (\tau_1, \tau_2, \pi, \sigma) \in \text{Hom}_{\Gamma}(Q \otimes \mathcal{H}, \mathcal{H}) \oplus \text{Hom}_{\Gamma}(\mathcal{H}, E_{\infty}) \oplus \text{Hom}_{\Gamma}(E_{\infty}, \mathcal{H}),$$

where we regard $Q = \mathbb{C}^2$, as earlier. Here π and σ are Γ -equivariant, while τ_1, τ_2 satisfy the properties (see [Kro89, Eq. 2.2])

$$(4.11) \quad \begin{aligned} \gamma^{-1} \tau_1 \gamma &= a \tau_1 + b \tau_2 \\ \gamma^{-1} \tau_2 \gamma &= -\bar{b} \tau_1 + \bar{a} \tau_2 \end{aligned} \quad \text{for all } \gamma = \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \in \Gamma \subset \text{SU}(2),$$

where on the left hand side γ is regarded as an isometry $\mathcal{H} \rightarrow \mathcal{H}$. Using the above description and the moment map equations (4.3)

$$\begin{aligned}\mu_{\mathbb{C}}(\tau_1, \tau_2, \pi, \sigma) &= [\tau_1, \tau_2] + \sigma\pi \\ \mu_{\mathbb{R}}(\tau_1, \tau_2, \pi, \sigma) &= [\tau_1, \tau_1^*] + [\tau_2, \tau_2^*] + \sigma\sigma^* - \pi^*\pi\end{aligned}$$

one easily verifies that $\mu_{\mathbb{R}}$ and $\mu_{\mathbb{C}}$ map $N = M^\Gamma$ into $\mathfrak{u}(\mathcal{H})^\Gamma$ and $\text{End}_\Gamma(\mathcal{H})$, respectively. By the uniqueness of the moment map it follows that μ_N is the restriction of μ to $N = M^\Gamma$. This proves the first point.

Before we proceed to the second point recall that $U_{reg} \subset M$ was defined to be the set of $m = (\tau_1, \tau_2, \pi, \sigma)$ for which

$$\begin{aligned}A_z &= (\tau_1 - z_1, \tau_2 - z_2, \pi)^t: \mathcal{H} \oplus \mathcal{H} \oplus E_\infty \rightarrow \mathcal{H} \\ B_z &= (-\tau_2 - z_2, \tau_1 - z_1, \sigma): \mathcal{H} \oplus \mathcal{H} \oplus E_\infty \rightarrow \mathcal{H}\end{aligned}$$

were injective and surjective, respectively, for all $z = (z_1, z_2) \in \mathbb{C}^2$ (see (4.4)). Observe that the equation $\mu_{\mathbb{R}}(m) = 0$ may be rewritten as

$$\tau_1\tau_1^* + \tau_2\tau_2^* + \sigma\sigma^* = \tau_1^*\tau_1 + \tau_2^*\tau_2 + \pi^*\pi,$$

or equivalently $B_0B_0^* = A_0^*A_0$. From this it follows that

$$\text{Ker}(A_0) = \text{Ker}(A_0^*A_0) = \text{Ker}(B_0B_0^*) = \text{Ker}(B_0^*).$$

Hence, A_0 is injective if and only if B_0 is surjective. In view of the easily verified fact that for any $(z_1, z_2) \in \mathbb{C}^2$ we have $\mu(\tau_1 - z_1, \tau_2 - z_2, \pi, \sigma) = \mu(\tau_1, \tau_2, \pi, \sigma)$, the above argument applies to show that if $m \in \mu_N^{-1}(0)$, then for any $z \in \mathbb{C}^2$, A_z is injective if and only if B_z is surjective.

We may now verify the second claim as follows. As $U_{reg} \cap \mu_N^{-1}(0) \subset \mu_N^{-1}(0) - \{0\}$, it suffices to show that if $m = (\tau_1, \tau_2, \pi, \sigma) \in \mu_N^{-1}(0) - U_{reg}$ then $m = 0$. To this end assume that $m \in \mu_N^{-1}(0) - U_{reg}$. In view of the above considerations this means that there exists $z = (z_1, z_2) \in \mathbb{C}^2$ and $h \neq 0 \in \mathcal{H}$ such that $A_z(h) = 0$. Then h belongs to $\text{Ker } \pi$ and is a common eigenvector for τ_1, τ_2 with eigenvalues z_1, z_2 respectively. Using the equations in (4.11) we deduce that $\gamma \cdot h$ is a common eigenvector for τ_1, τ_2 with eigenvalues

$$\begin{pmatrix} z_1^\gamma \\ z_2^\gamma \end{pmatrix} := \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \quad \text{for all } \gamma = \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \in \Gamma \subset \text{SU}(2).$$

Let $L := \text{Span}_{\mathbb{C}}\{\gamma h : \gamma \in \Gamma\}$. This is a nontrivial Γ -invariant subspace of \mathcal{H} satisfying $\tau_i(L) \subset L$ for $i = 1, 2$ and $\pi(L) = 0$. Moreover, since $\gamma \cdot h \in \text{Ker}(A_{(z_1^\gamma, z_2^\gamma)}) = \text{Ker}(B_{(z_1^\gamma, z_2^\gamma)}^*)$, it follows that $\tau_i^*(\gamma h) = \overline{z_i^\gamma} h$ for $i = 1, 2$ and $\sigma^*(\gamma h) = 0$. Therefore, $\tau_i^*(L) \subset L$ for $i = 1, 2$ and $\sigma^*(L) = 0$ as well. Hence, $\tau_i(L^\perp) \subset L^\perp$ for $i = 1, 2$ and $\sigma(E_\infty) \subset L^\perp$.

To summarize, we have proved that there is an orthogonal splitting of Γ -modules $\mathcal{H} = L \oplus L^\perp$ such that $\tau_i = \tau_i' \oplus \tau_i''$ preserves the splitting for $i = 1, 2$, $\pi(L) = 0$ and $\text{Im}(\sigma) \subset L^\perp$. Pick $\lambda \neq 1 \in U(1)$ and define $g \in U(\mathcal{H})^\Gamma$ by setting $g = \lambda \text{id}$ on L and $g = \text{id}$ on L^\perp . Then $(g\tau_1g^{-1}, g\tau_2g^{-1}, \pi g^{-1}, g\sigma) = (\tau_1, \tau_2, \pi, \sigma)$, so as $g \neq 1$ and $U(\mathcal{H})^\Gamma$ acts freely on $\mu_N^{-1}(0) - \{0\}$ we conclude that $m = 0$ as desired. \square

Proof of Proposition 4.1. Given an adjacent pair $\alpha, \beta \in \mathcal{C}$ let $E \rightarrow S^4$ be the unique Γ -equivariant $\text{SU}(2)$ -bundle with $[E_N] = \alpha$ and $[E_S] = \beta$ and minimal $c_2(E)[S^4] > 0$. Then by the above proposition the moduli space $\widetilde{\mathcal{M}}_{\Gamma, \infty}(E)$ of Γ -invariant instantons framed at $\infty = N$ may be identified with $\mathbb{R} \times Y_{\Gamma'}$ for some other finite subgroup $\Gamma' \subset \text{SU}(2)$. Using the correspondence between invariant instantons in $E \rightarrow S^4$ and instantons over $\mathbb{R} \times \widetilde{Y}_\Gamma$, taking the framing and orientation reversal into account, one obtains

$$\mathbb{R} \times Y_{\Gamma'} \cong e_-^{-1}(*) \subset \widetilde{\mathcal{M}}_z(\alpha, \beta),$$

where $e_-: \widetilde{\mathcal{M}}_z(\alpha, \beta) \rightarrow \alpha$ is the end-point map and z is the homotopy class from α to β corresponding to the relative Chern number $\hat{c}_2 = c_2(E)/|\Gamma|$. According to [Aus95] the natural translation action on the left hand side corresponds to the translation action on the right hand side. Hence,

$$Y_{\Gamma'} \cong e_-^{-1}(*) \subset \widetilde{\mathcal{M}}_z^0(\alpha, \beta).$$

Since the end-point map is $\text{SO}(3)$ -equivariant it follows that $\widetilde{\mathcal{M}}_z^0(\alpha, \beta) = e_-^{-1}(*) \cdot \text{SO}(3)$ is compact. Therefore, $\widetilde{\mathcal{M}}_z^0(\alpha, \beta) = \overline{\mathcal{M}}_z(\alpha, \beta)$, and as $\text{gr}_z(\alpha, \beta) = \dim(e_-^{-1}(*)) + 1 = 4$, we conclude that z is indeed the unique homotopy class with $-2 \leq \text{gr}_z(\alpha, \beta) \leq 5$.

To complete the proof we have to verify that $\widetilde{\mathcal{M}}_z(\alpha, \beta)$ is empty when α and β are not adjacent in \mathcal{S}_{Γ} . Let $\alpha = \alpha_0, \alpha_1, \dots, \alpha_s = \beta$ be the vertices of the minimal edge path connecting α to β in \mathcal{S}_{Γ} . Let z_i denote the unique homotopy class connecting α_i to α_{i+1} with $\text{gr}_{z_i}(\alpha_i, \alpha_{i+1}) = 4$ for each $0 \leq i < s$. Let \mathcal{H}_i be the minimal positive solution of $(2 - Q)\mathcal{H}_i = \alpha_i - \alpha_{i+1}$ for $0 \leq i < s$. By Corollary 4.14 the minimal positive solution of $(2 - Q)\mathcal{H} = \alpha - \beta$ is given by $\mathcal{H} = \bigoplus_{i=0}^{s-1} \mathcal{H}_i$. Put $k = \epsilon(\mathcal{H})$ and let $E^l \rightarrow S^4$ denote the Γ -equivariant $\text{SU}(2)$ -bundle with invariants $([E_N^l], [E_S^l], c_2(E^l)[S^4]) = (\alpha, \beta, k + |\Gamma|l)$ for $l \in \mathbb{Z}$. The representation \mathcal{H}_l used in the description of the moduli space $\widetilde{\mathcal{M}}_{\Gamma, \infty}(E^l)$ in the equivariant ADHM correspondence is then given by $\mathcal{H}_l = \mathcal{H} + lR$, where R is the regular representation. The dimension of the moduli space increases linearly with l so, as $\mathcal{H}_0 = \mathcal{H}$ is the minimal positive solution of $(2 - Q)\mathcal{H} = \alpha - \beta$, we conclude that $\widetilde{\mathcal{M}}_{\Gamma, \infty}(E^l)$ is empty for $l < 0$. This space corresponds to the fiber $e_-^{-1}(*) \subset \widetilde{\mathcal{M}}_w(\alpha, \beta)$ where $w = z_0 * z_1 * \dots * z_{s-1}$. By Theorem 2.1 part (a) we have

$$\text{gr}_w(\alpha, \beta) = \sum_{i=0}^{s-1} \text{gr}_{z_i}(\alpha_i, \alpha_{i+1}) = 4s > 4,$$

since by assumption $s > 1$. Consequently, the fiber $e_-^{-1}(*) \subset \widetilde{\mathcal{M}}_z(\alpha, \beta)$ where z is the unique homotopy class with $\text{gr}_z(\alpha, \beta) = 0, 4$, corresponds to $\widetilde{\mathcal{M}}_{\Gamma, \infty}(E^l)$ for some $l < 0$ and must therefore be empty. This implies that the whole moduli space must be empty and the proof is complete. \square

4.3. The Structure of $DCI(\overline{Y}_{\Gamma})$. In this section we determine the complexes $DCI(\overline{Y}_{\Gamma})$ explicitly. Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup and let \mathcal{C} denote the set of gauge equivalence classes of flat connections in the trivial $\text{SU}(2)$ -bundle over \overline{Y}_{Γ} . In this section we make the convention that the relative and absolute gradings

$$\text{gr}: \mathcal{C} \times \mathcal{C} \rightarrow \mathbb{Z}/8 \quad \text{and} \quad j: \mathcal{C} \rightarrow \mathbb{Z}/8,$$

introduced after Theorem 2.1, are always defined with respect to the orientation \overline{Y}_{Γ} of S^3/Γ . Furthermore, the vertices of the graph \mathcal{S}_{Γ} , introduced in Definition 3.9, i.e., the set of isomorphism classes of $\text{SU}(2)$ -representations of Γ , will be identified freely with the set \mathcal{C} .

Lemma 4.17. *For any pair $\alpha, \beta \in \mathcal{C}$ that are adjacent in \mathcal{S}_{Γ} it holds true that $\text{gr}(\alpha, \beta) = 4$. For each $\alpha \in \mathcal{C}$ let $p(\alpha)$ denote the number of edges in the minimal edge path connecting the trivial representation θ to α . Then $j(\alpha) \equiv 4p(\alpha) \pmod{8}$ for all $\alpha \in \mathcal{C}$.*

Proof. Suppose that α and β are adjacent in \mathcal{S}_{Γ} and let z be the unique homotopy class with $-2 \leq \text{gr}_z(\alpha, \beta) \leq 5$. Then according to Theorem 4.1 the fiber of $e_-: \overline{\mathcal{M}}_z(\alpha, \beta) \rightarrow \alpha$ has dimension 3. Using the dimension formula (2.1) we deduce that

$$\text{gr}(\alpha, \beta) = \dim \overline{\mathcal{M}}_z(\alpha, \beta) - \dim(\alpha) + 1 = \dim e_-^{-1}(*) + 1 = 4$$

proving the first assertion. For the second let $\alpha \in \mathcal{C}$ be arbitrary and let $\theta = \alpha_0, \alpha_1, \dots, \alpha_s = \alpha$ denote the vertices in the minimal edge path connecting θ to α in \mathcal{S}_{Γ} . Then $p(\alpha) = s$ and by the

additivity of the grading

$$j(\alpha) = \text{gr}(\alpha, \theta) = \sum_{i=0}^{s-1} \text{gr}(\alpha_{i+1}, \alpha_i) = 4s \in 4\mathbb{Z}/8.$$

Consequently, $j(\alpha) \equiv 4p(\alpha) \pmod{8}$ as required. \square

To determine the differentials in $DCI(\overline{Y}_\Gamma)$ we need the following result, which is contained in the proof of [Aus95, Lemma 5.1].

Lemma 4.18. *Let $\alpha, \beta \in \mathcal{C}$ be adjacent in \mathcal{S}_Γ with β irreducible, let $\mathcal{H} \in R(\Gamma)$ be the solution of $(2 - Q)\mathcal{H} = \alpha - \beta$ given by Proposition 4.12 and let $\Gamma' \subset \text{SU}(2)$ be the corresponding finite subgroup. Let z be the unique homotopy class with $\text{gr}_z(\alpha, \beta) = 4$. Then the degree of the map*

$$e_-^{-1}(\ast) = Y_{\Gamma'} \rightarrow \beta = \text{SO}(3)$$

is up to a sign given by $2 \dim_{\mathbb{C}} \mathcal{H} / |\Gamma'|$.

Remark 4.19. We are not aware of a procedure to pin down the orientation of $Y_{\Gamma'}$ in the above statement. For this reason we are only able to determine the degree up to a sign. As we will see, this inaccuracy will not affect our calculations in any relevant way.

For $i = 0, 1$ let $\mathcal{C}^i = \{\alpha \in \mathcal{C} : j(\alpha) \equiv 4i \pmod{8}\}$ such that $\mathcal{C} = \mathcal{C}^0 \cup \mathcal{C}^1$ is a disjoint union. Let us fix generators $b_\alpha \in H_0(\alpha)$ for all $\alpha \in \mathcal{C}$, and $t_\alpha \in H_2(\alpha)$ for all reducible α and $t_\alpha := b_\alpha u \in H_3(\alpha)$ for all irreducible α .

Definition 4.20. Let $\mathcal{C}^{irr} \subset \mathcal{C}$ be the subset of irreducibles. For each adjacent pair $\alpha, \beta \in \mathcal{C}$ with $\beta \in \mathcal{C}^{irr}$ define $n_{\beta\alpha}$ to be the integer $2 \dim_{\mathbb{C}} \mathcal{H} / |\Gamma'|$ where $\mathcal{H} \in R(\Gamma)$ is the minimal positive solution of the equation $(2 - Q)\mathcal{H} = \alpha - \beta$. If α and β are not adjacent in \mathcal{S}_Γ we define $n_{\beta\alpha} = 0$.

Theorem 4.21. *Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup. Then the multicomplex $(DCI(\overline{Y}_\Gamma)_{*,*}, \{\partial^r\}_{r \geq 0})$ is given by*

$$DCI_{8s,*} = \bigoplus_{\alpha \in \mathcal{C}^0} H_*(\alpha) \text{ and } DCI_{8s+4,*} = \bigoplus_{\beta \in \mathcal{C}^1} H_*(\beta)$$

for all $s \in \mathbb{Z}$ and $DCI_{s,} = 0$ otherwise. The differentials are given by $\partial^r = 0$ for $r \neq 4$ and $\partial^4 : DCI_{4s,0} \rightarrow DCI_{4(s-1),3}$ is determined on generators by*


$$\partial^4(b_\beta) = \sum_{\alpha \in \mathcal{C}^{irr}} n_{\alpha\beta} t_\alpha$$

where the integers $n_{\alpha\beta}$ are defined above.

Proof. The description of $DCI(Y, E)$ as a graded module is a simple consequence of the definition and Lemma 4.17. By the final part of Theorem 4.1 the moduli space $\overline{\mathcal{M}}_z(\alpha, \beta)$, where z is the homotopy class with $-2 \leq \text{gr}_z(\alpha, \beta) \leq 5$, is empty unless α and β are adjacent in \mathcal{S}_Γ . The statement concerning the differentials therefore follows from Lemma 2.7 and Lemma 4.18. \square


We will end this section by giving a few applications of Proposition 4.12 and show how we can easily extract both the grading and the integers $n_{\alpha\beta}$ provided we know where the one dimensional quaternionic representations are placed in the graph $\overline{\Delta}_\Gamma$. All the necessary information is contained in Appendix A.

Example 4.22. ($\Gamma = O^*$) There are four 1-dimensional quaternionic representations; two of real type θ, η and two of quaternionic type α, β . The graph $\overline{\Delta}_{O^*}$ is shown below.



The integers are the dimensions of the corresponding irreducible complex representations and we have labeled the vertices corresponding to the 1-dimensional quaternionic representation. Here there are no representations of complex type, in particular $\overline{\Delta}_\Gamma = \overline{\Delta}_\Gamma/(\iota)$. One may immediately conclude that the indexing is given by $j(\theta) = j(\beta) = 0$ and $j(\alpha) = j(\eta) = 4$. To determine the integer $n_{\beta\eta}$ we need to solve the equation $(2-Q)\mathcal{H} = \eta - \beta$ in $R(O^*)$. According to the recipe given in Proposition 4.12 we first delete the vertex corresponding to β and let $\Delta_{\mathcal{H}}$ be the component containing η ; that is, $\Delta_{\mathcal{H}}$ is just the single vertex corresponding to η . We then recognize this graph as Δ_{C_2} , from which we deduce that $\Gamma' = C_2$ and $\mathcal{H} = \frac{1}{2}\eta$. Therefore, $n_{\beta\eta} = 2 \dim_{\mathbb{C}} \mathcal{H} / |\Gamma'| = 2/2 = 1$. The situation with θ and α is completely symmetric so $n_{\alpha\theta} = 1$ as well.

The case of $n_{\beta\alpha}$ is more interesting. If we delete α , the component $\Delta_{\mathcal{H}}$ containing β is given by



We recognize this as the Dynkin graph D_6 which corresponds to the group D_4^* of order 16. We find the integral weights of \mathcal{H} , as given in the above graph, by knowledge of the dimensions of the irreducible representations of D_4^* (see Appendix A.2). From this we calculate

$$\dim_{\mathbb{C}} \mathcal{H} = 3 \cdot 1 + 4 \cdot 2 + 2 \cdot 1 + 3 \cdot 2 + 2 \cdot 2 + 1 \cdot 1 = 24$$

and hence $n_{\beta\alpha} = 2 \cdot 24/16 = 3$. By symmetry $n_{\alpha\beta} = 3$ as well.

The calculation of $n_{\alpha\theta}$ and $n_{\beta\eta}$ in the above example generalizes.

Lemma 4.23. *Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup and let Q be the canonical representation. Suppose ρ is a 1-dimensional complex representation of real type and that $\rho \otimes Q$ is irreducible so that $\eta = 2\rho$ and $\beta = \rho \otimes Q$ correspond to a pair of distinct vertices in \mathcal{S}_Γ . Then $n_{\beta\eta} = 1$.*

Proof. We trivially have $(2-Q)\rho = 2\rho - Q \otimes \rho = \eta - \beta$ so that $\mathcal{H} = \rho$. This means that $\Delta_{\mathcal{H}}$ consists of a single vertex and that the associated subgroup is $\Gamma' = C_2$. It follows that $n_{\beta\eta} = 2 \dim_{\mathbb{C}} \mathcal{H} / |\Gamma'| = 2/2 = 1$. \square

We also consider the most involved case $\Gamma = D_n^*$. The 1-dimensional quaternionic representations and the graph $\overline{\Delta}_{D_n^*}$ are given in Appendix A.2. For n even we label the quaternionic representations by

$$\theta, \eta_1, \eta_2, \eta_3, \alpha_1, \alpha_2, \dots, \alpha_{n/2}$$


and for n odd by

$$\theta, \eta, \alpha_1, \alpha_2, \dots, \alpha_{(n-1)/2}, \lambda.$$

Here θ, η and the η_i are fully reducible, the α_i are irreducible and λ is reducible.

Lemma 4.24. *In the above situation we have $n_{\alpha_i, \alpha_{i+1}} = n_{\alpha_{i+1}, \alpha_i} = 2$ for all i .*

Proof. Due to the symmetry in the graph $\overline{\Delta}_{D_n^*}$ it suffices to show that $n_{\alpha_{i+1}, \alpha_i} = 2$ for all i . We therefore have to solve $(2-Q)\mathcal{H} = \alpha_i - \alpha_{i+1}$ for \mathcal{H} . The relevant portion of the graph $\overline{\Delta}_{D_n^*}$ is shown below



Note that α_i is the $2i - 1$ 'th vertex labeled with 2 from left to right. Following the description in Proposition 4.12 we delete the vertex corresponding to α_{i+1} and let $\Delta_{\mathcal{H}}$ be the component containing α_i . This graph has a total of $2(i + 1)$ vertices so we recognize $\Delta_{\mathcal{H}} = D_{2(i+1)} = \Delta_{D_{2i}^*}$. Thus $\Gamma' = D_{2i}^*$ of order $8i$. The weights of \mathcal{H} are therefore given by attaching the integer 2 to each internal vertex (of degree ≥ 2) and 1 to the rest. This yields

$$n_{\alpha_{i+1}\alpha_i} = 2 \dim_{\mathbb{C}} \mathcal{H} / |D_{2i}^*| = 2(1 + 1 + 4(2i - 1) + 2) / 8i = 2$$

as required. \square

We note that for cyclic groups there are no irreducible $\text{SU}(2)$ -representations, from which it follows that $\text{DCI}(\overline{Y}_{C_m})$ has trivial differential for each m . There are now only a finite number of $n_{\alpha\beta}$ left to calculate. We state the calculations of these below without proof and trust that the interested reader will verify these for themselves.

For a finite subgroup $\Gamma \subset \text{SU}(2)$ we attach labels to the edges in the graph \mathcal{S}_{Γ} to express all the information needed to set up $\text{DCI}(\overline{Y}_{\Gamma})$. If α and β are adjacent and both irreducible we attach the symbol $(n_{\alpha\beta}|n_{\beta\alpha})$ to the edge connecting α to β such that $n_{\alpha\beta}$ is closest to α . If α and β are adjacent and only α is irreducible we attach $n_{\alpha\beta}$ to the edge.


Proposition 4.25. *The graphs \mathcal{S}_{Γ} with labels for the finite subgroups $\Gamma \subset \text{SU}(2)$ are given below.*

$$\begin{array}{ccc}
\mathcal{S}_{I^*} & \mathcal{S}_{O^*} & \mathcal{S}_{T^*} \\
\theta \text{ --- } \alpha \text{ --- } \beta & \theta \text{ --- } \alpha \text{ --- } \beta \text{ --- } \eta & \theta \text{ --- } \alpha \text{ --- } \lambda \\
1 \quad (3|4) & 1 \quad (3|3) \quad 1 & 1 \quad 3 \\
\\
\mathcal{S}_{C_{2m}} & \mathcal{S}_{C_{2m+1}} & \\
\theta \text{ --- } \lambda_1 \text{ --- } \lambda_2 \cdots \cdots \lambda_{m-1} \text{ --- } \eta & \theta \text{ --- } \lambda_1 \text{ --- } \lambda_2 \cdots \cdots \lambda_{m-1} \text{ --- } \lambda_m & \\
\\
\mathcal{S}_{D_{2m}^*} & & \\
\begin{array}{c} \theta \searrow 1 \\ \eta_1 \swarrow 1 \end{array} \alpha_1 \text{ --- } \alpha_2 \cdots \cdots \alpha_{m-1} \text{ --- } \alpha_m \begin{array}{c} 1 \swarrow \eta_2 \\ 1 \searrow \eta_3 \end{array} \\
(2|2) \quad \quad \quad (2|2) & & \\
\\
\mathcal{S}_{D_{2m+1}^*} & & \\
\begin{array}{c} \theta \searrow 1 \\ \eta \swarrow 1 \end{array} \alpha_1 \text{ --- } \alpha_2 \cdots \cdots \alpha_{m-1} \text{ --- } \alpha_m \text{ --- } \lambda \\
(2|2) \quad \quad \quad (2|2) \quad 2 & &
\end{array}$$

In all cases θ denotes the trivial representation. The letters α and β are reserved for representation of quaternionic type, λ for representations of complex type and η for representations of real type.

The notation for the 1-dimensional quaternionic representations is compatible with the notation in Appendix A.

Example 4.26. The multicomplexes $(\text{DCI}(\overline{Y}_{\Gamma})_{*,*}, \partial^4)$ for $\Gamma = T^*, O^*, I^*$ are shown below for (s, t) with $0 \leq s \leq 8$ and $0 \leq t \leq 3$.



5. THE ALGEBRAIC CONSTRUCTION OF EQUIVARIANT FLOER HOMOLOGY

The equivariant instanton Floer homology associated with the trivial $SU(2)$ -bundle over a closed oriented 3-manifold Y comes in three flavors: $I^+(Y)$, $I^-(Y)$, $I^\infty(Y)$, the positive, negative and Tate homology, respectively. These are all constructed algebraically from the complex $\widehat{CI}(Y)$, the associated action of $C_*^{gm}(SO(3); R)$ and the index filtration $\{F_p \widehat{CI}(Y)\}_{p \in \mathbb{Z}}$.

The construction proceeds as follows. First, following [Eis19, Appendix A], we review the construction of four functors C_A^+ , C_A^- , $C^{(+, tw)}$ and C_A^∞ from the category of right modules over a differential graded R -algebra A to the category of differential graded R -modules. In fact, $C_A^-(R)$ will carry the structure of a differential graded algebra and the three functors take values in the subcategory of left $C_A^-(R)$ -modules. Under the assumptions that the ground ring R is a principal ideal domain and the algebra A is degreewise free over R , the four functors will be exact and preserve quasi-isomorphisms.

Next, we promote the functors to the category of filtered right A -modules. If M is a right A -module equipped with an increasing filtration

$$\cdots \subset F_p M \subset F_{p+1} M \subset \cdots \subset M$$

of A -submodules, the exactness of the functors ensure that $C_A^\bullet(M)$ is naturally filtered by $F_p C_A^\bullet(M) := C_A^\bullet(F_p M)$ for each $\bullet \in \{+, -, (+, tw), \infty\}$. However, this filtration may not be very well behaved. To remedy this issue we pass to the so-called full completion, thereby obtaining four new functors \hat{C}_A^\bullet , $\bullet \in \{+, -, (+, tw), \infty\}$ from the category of filtered right A -modules to the category of filtered left $C_A^-(R)$ -modules; to be specific,

$$\hat{C}_A^\bullet(M) = \lim_q \operatorname{colim}_p C_A^\bullet(F_p M / F_q M).$$

For $A = C_*^{gm}(\mathrm{SO}(3); R)$ the groups $I^\bullet(Y)$ for $\bullet \in \{+, -, \infty\}$ are obtained by applying \hat{C}_A^\bullet to $M = \widehat{CI}(Y)$ equipped with the index filtration and then passing to homology. These groups will then be modules over $H(C_A^-(R))$, which, provided $\frac{1}{2} \in R$, is isomorphic to the polynomial algebra $R[U]$ with a single generator U of degree -4 . The mod 8 periodicity of M will be preserved so that $I_n^\bullet \cong I_{n+8}^\bullet$ for all $n \in \mathbb{Z}$.

The above constructions are also functorial in a suitable sense depending on $\bullet \in \{+, -, \infty\}$, in the differential graded algebra A . Moreover, this functoriality will preserve quasi-isomorphisms in the situation of interest; that is, we may equally well calculate $I^\bullet(Y)$ using $A = \Lambda_R[u]$ and $M = DCI(Y)$. When we proceed to the calculations for binary polyhedral spaces in the next section, we will indeed work in this context.

5.1. Conventions. Throughout R will be a fixed principal ideal domain. A module or algebra will always mean an R -module or R -algebra, and the same applies in the graded or differential graded setting. We will work with homological gradings and differentials ∂ of degree -1 . We follow the conventions on graded and differential graded (DG) modules given in [Mac95, Chapter VI]. We briefly recall the most central concepts. If M and N are DG modules then $M \otimes N$ and $\operatorname{Hom}(M, N)$ are the DG modules given by

$$(M \otimes N)_n = \bigoplus_{p+q=n} M_p \otimes N_q \quad \text{and} \quad \operatorname{Hom}(M, N)_n = \prod_{q-p=n} \operatorname{Hom}(M_p, N_q)$$

with differentials $\partial_{M \otimes N} = \partial_M \otimes 1 + 1 \otimes \partial_N$ and $\partial_{\operatorname{Hom}(M, N)} = \operatorname{Hom}(1, \partial_N) - \operatorname{Hom}(\partial_M, 1)$. This means that $\partial_{M \otimes N}(a \otimes b) = \partial_M a \otimes b + (-1)^{|a|} a \otimes \partial_N b$ and $\partial_{\operatorname{Hom}(M, N)}(f) = \partial_N \circ f - (-1)^{|f|} f \circ \partial_M$, where we write $|x|$ for the degree of an element in a graded module. There is a natural interchange isomorphism $\tau : M \otimes N \cong N \otimes M$ given by $m \otimes n \mapsto (-1)^{|m||n|} n \otimes m$.

For each integer $p \in \mathbb{Z}$ let $R[p]$ denote the complex with a single copy of R concentrated in degree p . For a DG module M the shifted complex $M[p]$ is defined to be $R[p] \otimes M$. Therefore, $M[p]_{n+p} = M_n$ for all $n \in \mathbb{Z}$ with differential $\partial_{M[p]} = (-1)^p \partial_M$. It is also possible to define $M[p] = M \otimes R[p]$, but this will result in different sign conventions.

A differential graded algebra is a graded algebra A equipped with a differential satisfying the Leibniz rule $\partial_A(ab) = \partial_A(a)b + (-1)^{|a|} a \partial_A b$ for all $a, b \in A$. We will always assume that A is unital, i.e., there is a unit $1 \in A_0$ for the multiplication. A left or right A -module is a graded left or right A -module equipped with a differential satisfying a corresponding Leibniz rule. This ensures that $H(M)$ is a graded $H(A)$ -module. If M is a left or right A -module then so is $M[p]$. For $a \in A$ and $x \in M[p]_n = M_{n-p}$ the induced product is given by $a \cdot x = (-1)^{|a|p} ax$ in the left module context and by $x \cdot a = xa$ in the right module context. This is a consequence of our left shifting (suspension) convention.

We also mention that there is a natural adjunction isomorphism of DG modules

$$\operatorname{Hom}(M \otimes N, P) \cong \operatorname{Hom}(M, \operatorname{Hom}(N, P))$$

given by $(f: M \otimes N \rightarrow P) \mapsto (g: M \rightarrow \text{Hom}(N, P))$ where $f(m \otimes n) = g(m)(n)$. This isomorphism admits various natural generalizations to the case where M , N and P carry additional module structures over various DG algebras, see [Mac95, Chapter VI.8]. All the signs occurring in the upcoming theory can be deduced from these conventions: the interchange morphism, the natural adjunction and the left shift convention.

Finally, a homomorphism $f: M \rightarrow N$ of DG modules will mean a degree 0 chain map unless otherwise stated. We say that f is a quasi-isomorphism if the induced map in homology, $H(f): H(M) \rightarrow H(N)$, is an isomorphism.

5.2. The Bar Construction and the Functors C_A^\pm . Let A be a unital DG algebra and assume that A is equipped with an augmentation $\epsilon: A \rightarrow R$, that is, ϵ is a map of DG algebras where R is regarded as a DG algebra concentrated in degree 0. The augmentation gives R the structure of an A -bimodule. We define $\bar{A} = A/R \cdot 1$ and note that there is a canonical identification of \bar{A} with the augmentation ideal $\text{Ker}(\epsilon)$. Given a right A -module M and a left A -module N define the bigraded module $B_{*,*}(M, A, N)$ by

$$B_{p,q}(M, A, N) = (M \otimes \bar{A}^{\otimes p} \otimes N)_q.$$

We adopt the standard notation and write

$$m[a_1|a_2|\cdots|a_p]n := m \otimes a_1 \otimes a_2 \otimes \cdots \otimes a_p \otimes n \in B_{p,q}(M, A, N).$$

Here, p is the simplicial degree, $q = |m| + (\sum_{i=1}^p |a_i|) + |n|$ is the internal degree and $p + q$ is the total degree. Let $\partial^s: B_{p,q}(M, A, N) \rightarrow B_{p-1,q}(M, A, N)$ and $\partial^i: B_{p,q}(M, A, N) \rightarrow B_{p,q-1}(M, A, N)$ act on $x = m[a_1|\cdots|a_p]n$ by

$$\begin{aligned} \partial^s(x) &= ma_1[a_2|\cdots|a_p]n + \sum_{i=1}^{p-1} (-1)^i m[a_1|\cdots|a_i a_{i+1}|\cdots|a_p]n \\ &\quad + (-1)^p m[a_1|\cdots|a_{p-1}]a_p n \\ \partial^i(x) &= (-1)^p (\partial_M m[a_1|\cdots|a_p]n + \sum_{i=1}^p (-1)^{\epsilon_i-1} m[a_1|\cdots|\partial a_i|\cdots|a_p]n \\ &\quad + (-1)^{\epsilon_p} m[a_1|\cdots|a_p]\partial_N n), \end{aligned}$$

where $\epsilon_i = |m| + |a_1| + \cdots + |a_i|$. Note that ∂^i is the usual tensor product differential in $M \otimes \bar{A}^{\otimes p} \otimes N$ twisted by a sign depending on the simplicial degree. We call ∂^s the simplicial differential and ∂^i the internal differential. It holds true that $(\partial^i)^2 = 0$, $(\partial^s)^2 = 0$ and $\partial^i \partial^s + \partial^s \partial^i = 0$. This ensures that the triple $(B_{*,*}(M, A, N), \partial^s, \partial^i)$ is a double complex.

Definition 5.1. [GM74, Appendix A] The bar construction $B(M, A, N)$ is the total complex associated with the double complex $(B_{*,*}(M, A, N), \partial^s, \partial^i)$ introduced above, that is,

$$B_k(M, A, N) = \bigoplus_{p+q=k} B_{p,q}(M, A, N) \quad \text{and} \quad \partial = \partial^s + \partial^i.$$

The bar construction is functorial in the triple (M, A, N) . For a triple $(f, g, h): (M, A, N) \rightarrow (M', A', N')$ where g is a map of augmented DG algebras and f, h are chain maps linear along g , there is an induced map $B(f, g, h): B(M, A, N) \rightarrow B(M', A', N')$ of DG modules. There is also a natural augmentation $B(M, A, N) \rightarrow M \otimes_A N$ given by $m[a_1|\cdots|a_p]n \mapsto 0$ for $p > 0$ and $m[]n \mapsto m \otimes n$.

Remark 5.2. The DG module $B(M, A, A)$ inherits a right A -module structure from the last factor. The augmentation $B(M, A, A) \rightarrow M$ is a map of right A -modules and should be regarded as an analogue of a projective resolution of M . Moreover, $B(M, A, N) \cong B(M, A, A) \otimes_A N$, so that

$B(M, A, N)$ is a derived tensor product $M \otimes_A^L N$ in a certain sense. These statements are made precise in the language of model categories in [BMR14, Section 10.2].

Following [Eis19] we introduce the notation

$$BA := B(R, A, R) \quad \text{and} \quad EA := B(R, A, A).$$

Furthermore, we write $[a_1|a_2|\cdots|a_p] := 1_R[a_1|a_2|\cdots|a_p]1_R \in BA$ and similar conventions apply for $B(R, A, N)$ and $B(M, A, R)$.

Lemma 5.3. *For any right A -module M and left A -module N there is a natural map of DG modules $\psi_{M,N}: B(M, A, N) \rightarrow B(M, A, R) \otimes B(R, A, N)$ given by*

$$m[a_1|\cdots|a_p]n \mapsto \sum_{i=0}^p (-1)^{(p-i)\epsilon_i} m[a_1|\cdots|a_i] \otimes [a_{i+1}|\cdots|a_p]n,$$

where $\epsilon_i = |m| + |a_1| + \cdots + |a_i|$. The map $\psi_{R,R}: BA \rightarrow BA \otimes BA$ along with the augmentation $BA \rightarrow R$ give BA the structure of a DG coalgebra. The map $\psi_{M,R}$ gives $B(M, A, R)$ the structure of a right DG BA -comodule and the map $\psi_{R,N}$ gives $B(R, A, N)$ the structure of a left DG BA -comodule.

Definition 5.4. [Eis19, Definition A.5] Let A be an augmented DG algebra and let M be a right A -module. Define the positive and negative A -chains of M to be

$$C_A^+(M) := B(M, A, R) \quad \text{and} \quad C_A^-(M) := \text{Hom}_A(EA, M).$$

The homology of $C_A^\pm(M)$ is denoted by $H_A^\pm(M)$ and is called the positive and negative A -homology of M , respectively.

The above constructions are functorial in the following sense. A homomorphism $f: A \rightarrow B$ of augmented DG algebras induces maps

$$f_*: C_A^+(f^{-1}M) \rightarrow C_B^+(M) \quad \text{and} \quad f^*: C_B^-(M) \rightarrow C_A^-(f^{-1}M)$$

where M is a B -module and $f^{-1}M$ is the A -module obtained from M by restriction along f . Furthermore, a homomorphism $g: M \rightarrow N$ of A -modules induces maps

$$g^+: C_A^+(M) \rightarrow C_A^+(N) \quad \text{and} \quad g^-: C_A^-(M) \rightarrow C_A^-(N).$$

The following invariance result is proven in [Eis19, Theorem A.1, A.2]; it relies on the standing assumption that R is a PID.

Proposition 5.5. *Let $f: A' \rightarrow A$ is a homomorphism of augmented DG algebras and let $g: M \rightarrow M'$ be a homomorphism of A -modules.*

- (i) *If A and A' are degreewise free and f is a quasi-isomorphism, then f_* and f^* are quasi-isomorphisms.*
- (ii) *If A is degreewise free and g is a quasi-isomorphism, then g^+ and g^- are quasi-isomorphisms.*

The DG modules $C_A^\pm(M)$ carry additional structure. First, we have the following sequence of isomorphisms of DG modules

$$C_A^-(R) = \text{Hom}_A(EA, \epsilon^{-1}(R)) \cong \text{Hom}_R(EA \otimes_A R, R) \cong \text{Hom}(BA, R),$$

which identifies $C_A^-(R)$ with the R -dual of BA . The DG coalgebra structure of BA dualizes to give $C_A^-(R)$ the structure of a DG algebra. Next, for any right A -module M the DG modules $C_A^\pm(M)$

obtain left $C_A^-(R)$ -module structures in the following way. Write $BA^\vee = \text{Hom}_R(BA, R) = C_A^-(R)$, then the structure map $C_A^-(R) \otimes C_A^+(M) \rightarrow C_A^+(M)$ is given by the following composition

$$(5.1) \quad \begin{aligned} & BA^\vee \otimes C_A^+(M) \xrightarrow{1 \otimes \psi_M} BA^\vee \otimes C_A^+(M) \otimes BA \\ & \xrightarrow{\tau \otimes 1} C_A^+(M) \otimes BA^\vee \otimes BA \xrightarrow{1 \otimes \text{ev}} C_A^+(M) \end{aligned}$$

where $\psi_M = \psi_{M,R}$ is the comodule structure map of $C_A^+(M) = B(M, A, R)$ in Lemma 5.3, τ is the interchange isomorphism $a \otimes b \mapsto (-1)^{|a||b|} b \otimes a$ and $\text{ev}: BA^\vee \otimes BA \rightarrow R$ is the evaluation map. In the other case the structure map $C_A^-(R) \otimes C_A^-(M) \rightarrow C_A^-(M)$ is given by the following composition

$$(5.2) \quad BA^\vee \otimes \text{Hom}_A(EA, M) \xrightarrow{\zeta} \text{Hom}_A(BA \otimes EA, M) \xrightarrow{\psi_{EA}^*} \text{Hom}_A(EA, M)$$

where $\zeta(f \otimes g)(a \otimes b) = (-1)^{|a||g|} f(a) \cdot g(b)$ and $\psi_{EA}^* = \text{Hom}_A(\psi_{R,A}, 1)$ is the dual of the comodule structure map $\psi_{R,A}: EA \rightarrow BA \otimes EA$.

The properties of the functors C_A^\pm that will be important for us are summarized in the following theorem. Here the morphisms in the category of right or left modules over a DG algebra A are taken to be the A -linear chain maps of degree 0.

Theorem 5.6. *Let A be a degreewise R -free augmented DG algebra. Then the assignments*

$$M \mapsto C_A^\pm(M) \quad \text{and} \quad (g: M \rightarrow M') \mapsto (g^\pm: C_A^\pm(M) \rightarrow C_A^\pm(M'))$$

define functors C_A^\pm from the category of right A -modules to the category of left $C_A^-(R)$ -modules. Moreover, these functors preserve quasi-isomorphisms and short exact sequences.

We will now treat the case $A = \Lambda_R[u]$ with $|u| = 3$ in detail. These calculations will be important for later purposes and will serve as an excellent example of how the above constructions work. We should note that all of these calculations easily generalize to the case where the degree of u is any odd number.

We will write $R[U]$ for the graded polynomial algebra on a single generator U of even non-zero degree. This is also a coalgebra with counit given by the identity in degree 0 and diagonal determined by

$$U^p \mapsto \sum_{i=0}^p U^i \otimes U^{p-i}$$

where $U^0 = 1$. This structure is compatible with the algebra structure and turns $R[U]$ into a Hopf algebra, but this will not be important for us.

Lemma 5.7. *Let $A = \Lambda_R[u]$ with $|u| = 3$. Then there is an isomorphism of DG coalgebras $BA \cong R[V]$ where $|V| = 4$, and an isomorphism of DG algebras $C_A^-(R) = BA^\vee \cong R[U]$ where $|U| = -4$. In particular, these complexes have trivial differentials so that $H_A^+(R) = R[V]$ and $H_A^-(R) = R[U]$. The first isomorphism is given by*

$$V^p \mapsto (-1)^{p(p+1)/2} [u] \cdots [u] \quad (p \text{ times})$$

for $p \geq 0$. The second is obtained by dualization of the first where U^p is determined by the evaluation $U^p(V^p) = 1$ for all $p \geq 0$. Furthermore, the left BA^\vee -module structure of BA is given by $U^s \cdot V^p = V^{p-s}$, where we interpret $V^i = 0$ for $i < 0$.

Proof. We have $A = \Lambda_R[u]$ so that $A_0 = R$, $A_3 = Ru$ and $A_n = 0$ otherwise. The augmentation is necessarily given by the identity in degree 0. Hence, $\overline{A} = Ru$ concentrated in degree 3. From the definition of $B(R, A, R)$ we find that $BA_{p,3p} = R \cdot Z^p$ for $p \geq 0$, where $Z^p = [u] \cdots [u]$ with

u repeated p times, and $B_{p,q} = 0$ otherwise. It follows immediately that the differential vanishes identically. Using the formula for the coalgebra structure map in Lemma 5.3 we find that

$$\psi_{R,R}(Z^p) = \sum_{i=0}^p (-1)^{(p-i)3i} Z^i \otimes Z^{p-i}.$$

Define $V^p = (-1)^{p(p+1)/2} Z^p$ for $p \geq 0$. Then using the identity

$$\frac{p(p+1)}{2} = (p-i)i + \frac{i(i+1)}{2} + \frac{(p-i)(p-i+1)}{2}$$

we deduce that the above formula transforms into $\psi_{R,R}(V^p) = \sum_{i=0}^p V^i \otimes V^{p-i}$. Hence, $BA \cong R[V]$ as DG coalgebras with trivial differentials.

Define $U^p \in BA^\vee$ to be the dual of V^p , i.e., $U^p: BA \rightarrow R$ has degree $-4p$ and acts by $U^p(V^p) = 1$ and vanishes otherwise. The differential is still of course trivial. The product $U^s \cdot U^t$ has degree $-4(s+t)$ and must therefore be an R -multiple of U^{s+t} . The calculation

$$(U^s \cdot U^t)(V^{s+t}) = (U^s \otimes U^t) \left(\sum_{i=0}^{s+t} V^i \otimes V^{s+t-i} \right) = U^s(V^s) U^t(V^t) = 1$$

shows that $U^s \cdot U^t = U^{s+t}$. Hence, $C_A^-(R) \cong BA^\vee \cong R[U]$ as DG algebras with trivial differentials. Finally, we need to check that the left module structure $BA^\vee \otimes BA \rightarrow BA$ is given by $U^s \cdot V^p = V^{p-s}$. Using the definition we find

$$U^s \cdot V^p = \sum_{i=0}^p (-1)^{|U^s||V^i|} V^i \cdot U^s(V^{p-i}) = \begin{cases} V^{p-s} & \text{if } p \geq s \\ 0 & \text{otherwise.} \end{cases}$$

as required. \square

We will also need explicit calculations of $H_A^\pm(M)$ when $M = H_*(\alpha; R)$ for $\alpha = \text{SO}(3)/\text{SO}(2)$ or $\alpha = \text{SO}(3)$, under the assumption $\frac{1}{2} \in R$. In other words, $M = R \oplus R[2]$ with the trivial A -module structure or $M = A = \Lambda_R[u]$ (see (2.2)). For this purpose we give concrete models for the complexes $C_A^\pm(M)$. Note that a right $\Lambda_R[u]$ -module M is simply a DG module equipped with a degree 3 map $u: M \rightarrow M$ satisfying $u^2 = 0$ and $u\partial_M = \partial_M u$ (because A acts from the right).

Proposition 5.8. *Let M be a right $A = \Lambda_R[u]$ -module. Define bigraded modules $D_{*,*}^\pm$ by*

$$D_{p,q}^+ = \begin{cases} M_{q-3p} & p \geq 0 \\ 0 & p < 0 \end{cases} \quad \text{and} \quad D_{p,q}^- = \begin{cases} M_{q-3p} & p \leq 0 \\ 0 & p > 0 \end{cases}.$$

Define $U: D_{p,q}^\pm \rightarrow D_{p-1,q-3}^\pm$ to be the identity $M_{q-3p} \rightarrow M_{q-3p}$ when $p \geq 1$ and $p \leq 0$, respectively, and otherwise to be zero. Define

$$\begin{aligned} \partial' &= (-1)^{p+q+1} u: D_{p,q}^\pm \rightarrow M_{q-3p} \rightarrow M_{q-3p+3} = D_{p-1,q}^\pm \\ \partial'' &= \partial_M: D_{p,q}^\pm \rightarrow M_{q-3p} \rightarrow M_{q-3p-1} = D_{p,q-1}^\pm, \end{aligned}$$

in the range it makes sense. Then $(D_{*,*}^+, \partial', \partial'')$ and $(D_{*,*}^-, \partial', \partial'')$ are double complexes and there are natural isomorphisms of DG $R[U]$ -modules

$$\text{Tot}^\oplus(D_{*,*}^+, \partial', \partial'') \cong C_A^+(M) \quad \text{and} \quad \text{Tot}^\Pi(D_{*,*}^-, \partial', \partial'') \cong C_A^-(M).$$

Proof. We have $\bar{A} = R \cdot u$ with u in degree 3. Therefore, $B_{p,q} = (M \otimes \bar{A}^{\otimes p})_q \cong M_{q-3p} \otimes \bar{A}_{3p}^{\otimes p} \cong M_{q-3p}$. The isomorphism is given by $m \mapsto m[u|u] \cdots [u]$ with u repeated p times. For simplicity of notation write this as $m \otimes Z^p$. Then, using the formulas for the differentials, we find $\partial^s(m \otimes Z^p) = mu \otimes Z^{p-1}$ and $\partial^i(m \otimes Z^p) = (-1)^p \partial_M m \otimes Z^p$. The action of U is given by first applying the comodule structure

map to $m \otimes Z^p$, and then evaluating U on the right factors in the resulting sum (U has even degree so the interchange introduces no sign). We have

$$\psi_{M,R}(m \otimes Z^p) = \sum_{i=0}^p (-1)^{(p-i)(|m|+3i)} (m \otimes Z^i) \otimes Z^{p-i}$$

so that $U \cdot (m \otimes Z^p) = (-1)^{|m|+p-1} (m \otimes Z^{p-1}) \cdot U(Z^1) = (-1)^{|m|+p} m \otimes Z^{p-1}$ as U was defined to be the dual of $V^1 = -Z^1$. It is now easy to check that when we introduce the sign change $m \in D_{p,q}^+ = M_{q-3p} \mapsto (-1)^{p(p+1)/2+p|m|} m \otimes Z^p$ the U action and the differentials are given as in the proposition.

In the second case, $EA = B(R, A, A)$ has generators over R of the form $Z^p = [u|u| \cdots |u]$ and $Z^p \otimes u = [u|u| \cdots |u]u$ in degrees $4p$ and $4p+3$ respectively. It is convenient as earlier to introduce the sign twist $V^p = (-1)^{p(p+1)/2} Z^p$, since then the differential is given by $\partial(V^p) = V^{p-1} \otimes u$ and $\partial(V^p \otimes u) = 0$ and the comodule structure map $\psi_{R,A}: EA \rightarrow BA \otimes EA$ is given by

$$\psi_{R,A}(V^p) = \sum_{i=0}^p V^i \otimes V^{p-i}.$$

An element $f \in \text{Hom}_A(EA, M)_n$ is by A -linearity uniquely determined by the elements $f(V^p) \in M_{n+4p}$ for $p \geq 0$. The differential is given by

$$(\partial f)(V^p) = \partial_M(f(V^p)) - (-1)^{|f|} f(\partial V^p) = \partial_M(f(V^p)) - (-1)^{|f|} f(V^{p-1})u$$

and the action of U is given by

$$(U \cdot f)(V^p) = (U \otimes f)(\psi_{R,A}(V^p)) = (U \otimes f)(V \otimes V^{p-1}) = f(V^{p-1}).$$

It is now easy to check that the complex $C_A^-(M) = \text{Hom}_A(EA, M)$ can be described as the product totalization of the double complex given in the proposition. \square

For later reference we record the following facts from the above proof. The isomorphism $\text{Tot}^\oplus(D^+, \partial', \partial'') \cong C_A^+(M)$ is given by sending an element $m \in M_{q-3p} = D_{p,q}^+$ to

$$(5.3) \quad (-1)^{p(q-p)+p(p+1)/2} m \overbrace{[u|u| \cdots |u]}^p \in B(M, A, R)_{p,q}$$

and the isomorphism $C_A^-(M) \cong \text{Tot}^\Pi(D^-, \partial', \partial'')$ is given by sending an element $f \in \text{Hom}_A(EA, M)_n$ to

$$(5.4) \quad ((-1)^{p(p+1)/2} f(\overbrace{[u|u| \cdots |u]}^p))_{p \geq 0} \in \prod_{p \geq 0} M_{n+4p} = \prod_{p \geq 0} D_{-p,n+p}^-.$$

It should be noted that there are other choices of signs possible. We have chosen the signs that ensure that the action of U corresponds to the identity map. A part of the double complex

$(D_{*,*}^+, \partial', \partial'')$ corresponding to a right A -module M is shown below

$$\begin{array}{ccccccc}
& \vdots & & \vdots & & \vdots & \\
& \downarrow & & \downarrow & & \downarrow & \\
0 & \longleftarrow & M_2 & \xleftarrow{u} & M_{-1} & \xleftarrow{-u} & M_{-4} \longleftarrow \cdots \\
& \searrow \partial_M & & \searrow \partial_M & & \searrow \partial_M & \\
0 & \longleftarrow & M_1 & \xleftarrow{-u} & M_{-2} & \xleftarrow{u} & M_{-5} \longleftarrow \cdots \\
& \searrow \partial_M & & \searrow \partial_M & & \searrow \partial_M & \\
0 & \longleftarrow & M_0 & \xleftarrow{u} & M_{-3} & \xleftarrow{-u} & M_{-6} \longleftarrow \cdots \\
& \searrow \partial_M & & \searrow \partial_M & & \searrow \partial_M & \\
0 & \longleftarrow & M_{-1} & \xleftarrow{-u} & M_{-4} & \xleftarrow{u} & M_{-7} \longleftarrow \cdots \\
& \downarrow & & \downarrow & & \downarrow & \\
& \vdots & & \vdots & & \vdots &
\end{array}$$

The dotted arrows represent the action of U . An element x of $C_A^+(M)_n$ can be represented by a sequence $(m_{n-4p})_{p \geq 0}$, where almost all terms vanish. This element x is a cycle if and only if $\partial_M m_{n-4p} = (-1)^n m_{n-4(p-1)} u$ for each $p \geq 0$. Each column is a shifted copy of the original complex M . The double complex $(D_{*,*}^-, \partial', \partial'')$ is obtained by shifting the whole diagram from the right half plane over to the left half plane. A more concise representation of this double complex is given in the following diagram

$$\cdots \longleftarrow M[-9] \xleftarrow{\pm u} M[-6] \xleftarrow{\mp u} M[-3] \xleftarrow{\pm u} M \longleftarrow 0.$$

Note that in this case the internal differential in $M[-3n]$, $n \geq 0$, is not twisted by a sign, and that the $\pm u$ notation means that the sign depends on the internal degree.

With these concrete models in hand it is easy to calculate the following.

Corollary 5.9. *We have $H_A^+(A) \cong R$, $H_A^-(A) \cong R[3]$, with necessarily trivial $R[U]$ -module structure, and $H_A^+(R \oplus R[2]) \cong R[W]$ with $|W| = 2$, $H_A^-(R \oplus R[2]) \cong R[Z][2]$ with $|Z| = -2$. The $R[U]$ -module structures are determined by $U \cdot W^p = W^{p-2}$ and $U \cdot Z^p = Z^{p+2}$.*

Remark 5.10. The notation $R[W]$ and $R[Z][2]$ in the above result is only used to effectively express the underlying graded modules and the action of $R[U]$, it should not be interpreted to mean that $H_A^\pm(R \oplus R[2])$ carry any (co)algebra structure.

5.3. Tate Homology. We will now proceed to the construction of $C_A^\infty(M)$.

Definition 5.11. For an augmented DG algebra A define the dualizing object by

$$D_A := C_A^-(A) = \text{Hom}_A(EA, A).$$

The dualizing object is a (left-left) $A - C_A^-(R)$ -bimodule. The left $C_A^-(R)$ -module structure is the one already given in (5.2), while the left A -module structure is obtained from the second factor in $D_A = \text{Hom}_A(EA, A)$, that is, $(a \cdot f)(x) := af(x)$ for all $a \in A$, $f \in \text{Hom}_A(EA, A)$ and $x \in EA$. This justifies the following definition.

Definition 5.12. For a right A -module M define the twisted positive A -chains of M to be the DG module

$$C_A^{+,tw}(M) := B(M, A, D_A).$$

The homology of this complex is denoted by $H_A^{+,tw}(M)$ and is called the positive twisted A -homology of M .

The complex $C_A^{+,tw}(M)$ carries a left $C_A^-(R)$ -module structure induced from the left $C_A^-(R)$ -module structure of D_A .

Definition 5.13. The norm map $N_M: C_A^{+,tw}(M) \rightarrow C_A^-(M)$ is for each right A -module M defined to be the composite

$$B(M, A, D_A) \xrightarrow{\epsilon} M \otimes_A D_A \xrightarrow{\xi} \text{Hom}_A(EA, M),$$

where ϵ is the augmentation and ξ is the adjoint of

$$M \otimes_A (D_A \otimes EA) \xrightarrow{1 \otimes \text{ev}} M \otimes_A A = M.$$

Explicitly, for $m[a_1 | \cdots | a_p]v \in B(M, A, D_A)$ and $x \in EA$,

$$(5.5) \quad N_M(m[a_1 | \cdots | a_p]v)(x) = \begin{cases} m \cdot v(x) & \text{for } p = 0 \\ 0 & \text{for } p > 0. \end{cases}$$

Lemma 5.14. The norm map $N_M: C_A^{+,tw}(M) \rightarrow C_A^-(M)$ is $C_A^-(R)$ -linear and natural in M .

Proof. The naturality in M is clear from the definition. To verify the $C_A^-(R)$ -linearity let $f \in C_A^-(R)$ and $w = m[a_1 | \cdots | a_p]u \in C_A^{+,tw}(M) = B(M, A, D_A)$. Then $f \cdot w = (-1)^\eta m[a_1 | \cdots | a_p](f \cdot w)$ for $\eta = p + |m| + |a_1| + \cdots + |a_p|$ and therefore $N_M(f \cdot w) = 0 = f \cdot N_M(w)$ provided $p > 0$. We may therefore assume that $p = 0$ such that $w = m[u]$. Let $v = N_M(w)$ and note that $|v| = |m| + |u|$. Let $x \in EA$ and write $\psi_{R,A}(x) = \sum_i y_i \otimes z_i \in BA \otimes EA$ where $\psi_{R,A}$ is the map of Lemma 5.3. Then by definition $(f \cdot v)(x) = (f \otimes v)(\psi_{R,A}(x))$, which equals

$$(5.6) \quad \sum_i (-1)^{|v||y_i|} f(y_i) v(z_i) = \sum_i (-1)^{|m||y_i| + |u||y_i|} m \cdot f(y_i) u(z_i).$$

On the other hand, $N_M(f \cdot m[u]) = (-1)^{|f||m|} N_M(m[(f \cdot u)])$, and by (5.5) this coincides with

$$(-1)^{|f||m|} \sum_i m \cdot (f \otimes u)(y_i \otimes z_i) = \sum_i (-1)^{|f||m| + |u||y_i|} m \cdot f(y_i) u(z_i).$$

The final expression is equal to the final expression in (5.6) because $f: BA \rightarrow R$ so that $f(y_i)$ can only be nonzero if $|f| = -|y_i|$. Hence, $f \cdot N_M(w) = N_M(f \cdot w)$ and the proof is complete. \square

The Tate complex $C_A^\infty(M)$ will be defined to be the mapping cone complex of the norm map. At a later stage we will also have to consider the cone of a map of nonzero degree, so we fix our conventions in the following definition.

Definition 5.15. Let B be a DG algebra and suppose that $f: M \rightarrow N$ is a degree d chain map of left B -modules. This means that

- (a) $f(M_n) \subset N_{n+d}$ for all $n \in \mathbb{Z}$,
- (b) $f(bm) = (-1)^{|b|d} b f(m)$ for all $b \in B$ and $m \in M$ and
- (c) $f \circ \partial_M = (-1)^d \partial_N \circ f$.

The left B -module $\text{Cone}(f)$ is defined by $\text{Cone}(f)_k = N_k \oplus M_{k-d-1}$ for all $k \in \mathbb{Z}$, with differential

$$\partial_{\text{Cone}(f)} \cdot \begin{pmatrix} n \\ m \end{pmatrix} = \begin{pmatrix} \partial_N & (-1)^d f \\ 0 & (-1)^{d+1} \partial_M \end{pmatrix} \begin{pmatrix} n \\ m \end{pmatrix} = \begin{pmatrix} \partial_N n + (-1)^d f m \\ (-1)^{d+1} \partial_M m \end{pmatrix}$$

and B -module structure $b \cdot (n, m) = (bn, (-1)^{|b|(d+1)} bm)$.

Remark 5.16. If $B = R$ is concentrated in degree 0 and f has degree 0, then this definition agrees with [Mac95, p. 46]. The above definition is what we get when we regard f as a degree 0 map $f: M[d] \rightarrow N$ and take into account the sign twist in the B -action upon shifting M .

The cone complex sits in a natural short exact sequence

$$0 \longrightarrow N \longrightarrow \text{Cone}(f) \longrightarrow M[d+1] \longrightarrow 0$$

of B -modules, whose connecting homomorphism $\delta: H(M)_n \rightarrow H_{n+d}(N)$ recovers $H(f)$ up to a sign. In particular, there is a natural exact triangle in homology.

Definition 5.17. Let M be a right A -module. Then the Tate complex of M is defined to be the $C_A^-(R)$ -module

$$C_A^\infty(M) := \text{Cone}(N_M: C_A^{+,tw}(M) \rightarrow C_A^-(M)).$$

The homology of this complex is denoted by $H_A^\infty(M)$ and is called the Tate homology of M .

Given a map $g: M \rightarrow M'$ of right A -modules, there are induced maps

$$g^{+,tw}: C_A^{+,tw}(M) \rightarrow C_A^{+,tw}(M') \quad \text{and} \quad g^\infty: C_A^\infty(M) \rightarrow C_A^\infty(M')$$

of $C_A^-(R)$ -modules. The key properties of the functors $C_A^{+,tw}$ and C_A^∞ are summarized in the following theorem.

Theorem 5.18. Let A be a degreewise R -free augmented DG algebra. Then the assignments

$$(M \mapsto C_A^\bullet(M)) \quad \text{and} \quad (g: M \rightarrow M') \mapsto (g^\bullet: C_A^\bullet(M) \rightarrow C_A^\bullet(M'))$$

for $\bullet \in \{(+, tw), \infty\}$, define functors $C_A^{+,tw}, C_A^\infty$ from the category of right A -modules to the category of left $C_A^-(R)$ -modules. They are both exact and preserve quasi-isomorphisms. Moreover, there is a natural exact triangle of $H_A^-(R)$ -modules

$$\begin{array}{ccc} H_A^{+,tw}(M) & \xrightarrow{H(N_M)} & H_A^-(M) \\ & \swarrow \scriptstyle [-1] & \searrow \\ & H_A^\infty(M) & \end{array}$$

Suppose $f: A \rightarrow B$ is a quasi-isomorphism of augmented DG algebras. For a given right B -module M there is no induced map between $C_A^\bullet(f^{-1}M)$ and $C_B^\bullet(M)$ for $\bullet \in \{(+, tw), \infty\}$. Rather one has to introduce an intermediate object to compare them.

Definition 5.19. Let $f: A \rightarrow B$ be a homomorphism of augmented DG algebras. Define the relative dualizing object to be the (left-left) $C_A^-(R) - B$ -bimodule

$$D_f := C_A^-(f^{-1}B) = \text{Hom}_A(EA, f^{-1}B),$$

where $f^{-1}B$ is regarded as a (right-left) $A - B$ -bimodule.

Given a right B -module M define the relative positive twisted complex to be $C_f^{+,tw}(M) := B(M, B, D_f)$ equipped with the left $C_A^-(R)$ -module structure induced from D_f .

The relative positive twisted complex is a functor from the category of right B -modules to the category of left $C_A^-(R)$ -modules. Given a map $g: M \rightarrow N$ of right B -modules the induced map is given by $B(g, 1, 1): B(M, B, D_f) \rightarrow B(N, B, D_f)$.

Lemma 5.20. In the situation of the above definition, if A and B are degreewise free, the relative functor $C_f^{+,tw}$ preserves quasi-isomorphisms and short exact sequences.

Proof. The assumptions that A and B are degreewise free along with the fact that R is PID ensure that the relative dualizing object D_f is degreewise flat. This is verified as in [Eis19, Lemma A.10]. It then follows from the invariance theorem [Eis19, Theorem A.1] that $C_f^{+,tw} = B(-, B, D_f)$ is an exact functor preserving quasi-isomorphisms. \square

Proposition 5.21. *Let $f: A \rightarrow B$ be a homomorphism of augmented DG algebras and let M be a right B -module. There is a relative norm map $N_M^f: C_f^{+,tw}(M) \rightarrow C_A^-(f^{-1}M)$ fitting into the following commutative diagram, natural in M ,*

$$\begin{array}{ccccc} C_B^{+,tw}(M) & \xrightarrow{\beta_M} & C_f^{+,tw}(M) & \xleftarrow{\alpha_M} & C_A^{+,tw}(f^{-1}M) \\ \downarrow N_M^B & & \downarrow N_M^f & & \downarrow N_M^A \\ C_B^-(M) & \xrightarrow{f^*} & C_A^-(f^{-1}M) & \xleftarrow{=} & C_A^-(f^{-1}M). \end{array}$$

All of the maps in the diagram are $C_A^-(R)$ -linear, $C_B^-(R)$ -linear or linear along $f^*: C_B^-(R) \rightarrow C_A^-(R)$, as appropriate.

Moreover, if f is a quasi-isomorphism and A and B are degreewise R -free, then all the horizontal maps in the diagram are quasi-isomorphisms.

Proof. Define $g: D_B \rightarrow D_f$ and $h: D_A \rightarrow D_f$ to be the maps

$$g = f^*: C_B^-(B) \rightarrow C_A^-(f^{-1}B) \quad \text{and} \quad h = f^-: C_A^-(A) \rightarrow C_A^-(f^{-1}B).$$

Note that D_A is a $C_A^-(R) - A$ -bimodule, D_f is a $C_A^-(R) - B$ -bimodule and D_B is a $C_B^-(R) - B$ -bimodule. Moreover, observe that g is a homomorphism along $(f^*, 1): (C_B^-(R), B) \rightarrow (C_A^-(R), B)$ and that h is a homomorphism along $(1, f): (C_A^-(R), A) \rightarrow (C_A^-(R), B)$. We may therefore define the upper horizontal maps in the diagram by

$$\begin{aligned} \beta_M &:= B(1, 1, g): B(M, B, D_B) \rightarrow B(M, B, D_f) \\ \alpha_M &:= B(1, f, h): B(f^{-1}M, A, D_A) \rightarrow B(M, B, D_f). \end{aligned}$$

It is clear by construction that these maps are natural in M . Furthermore, by the above observations β_M is linear along $f^*: C_B^-(R) \rightarrow C_A^-(R)$ and α_M is $C_A^-(R)$ -linear.

We define the relative norm map $N_M^f: B(M, B, D_f) \rightarrow C_A^-(f^{-1}M)$ to be the composition of the augmentation $B(M, B, D_f) \rightarrow M \otimes_B D_f$ and the map $\kappa: M \otimes_B D_f \rightarrow C_A^-(M)$ which is given as the adjoint of

$$M \otimes_B (\text{Hom}_A(EA, f^{-1}B) \otimes EA) \xrightarrow{1 \otimes \text{ev}} M \otimes_B f^{-1}B \cong f^{-1}M.$$

Then N_M^B is $C_B^-(R)$ -linear while N_M^f and N_M^A are $C_A^-(R)$ -linear. Consider the diagram

$$\begin{array}{ccccc} B(M, B, D_B) & \xrightarrow{B(1,1,g)} & B(M, B, D_f) & \xleftarrow{B(1,f,h)} & B(f^{-1}M, A, D_A) \\ \downarrow \epsilon & & \downarrow \epsilon & & \downarrow \epsilon \\ M \otimes_B D_B & \xrightarrow{1 \otimes_B g} & M \otimes_B D_f & \xleftarrow{1 \otimes_f h} & f^{-1}M \otimes_A D_A \\ \downarrow \xi & & \downarrow \kappa & & \downarrow \xi \\ \text{Hom}_B(EB, B) & \xrightarrow{Ef^*} & \text{Hom}_A(EA, f^{-1}M) & \xleftarrow{=} & \text{Hom}_A(EA, f^{-1}M), \end{array}$$

where ϵ denotes the augmentations and ξ are the natural maps in the definition of the norm map (see 5.13). In particular, the outer compositions $\xi \circ \epsilon$ are the respective norm maps and by definition $\kappa \circ \epsilon = N_M^f$. We wish to verify that the diagram commutes. The two upper squares commute by the

naturality of the augmentation in the bar construction. To verify that the lower squares commute, consider the diagram obtained by passing to the adjoints of the vertical maps ξ , κ and ξ in the bottom half of the diagram. The resulting diagram is then seen to commute using the fact that each of the maps ξ , κ and ξ was defined as the adjoint of $1 \otimes \text{ev}$ for a suitable evaluation. By omitting the middle row of the above diagram, we obtain the commutative diagram in the statement of the proposition.

Finally, we need to verify that all the horizontal maps in the diagram are quasi-isomorphisms provided that $f: A \rightarrow B$ is a quasi-isomorphism of degreewise R -free algebras. We already know that this is true for $f^*: C_B^-(M) \rightarrow C_A^-(f^{-1}M)$ by Proposition 5.5. For the upper horizontal maps, note first that $g = f^*$ and $h = f^-$ are quasi-isomorphisms by the same result. Then, since D_A , D_B and D_f are degreewise flat, one concludes by [Eis19, Theorem A.1] that $\beta_M = B(1, 1, g)$ and $\alpha_M = B(1, f, g)$ are quasi-isomorphisms as well. This completes the proof. \square

From this result one quickly obtains the required invariance result. Nevertheless, we will need the more refined statement of the above proposition in the next section.

Corollary 5.22. *Let $f: A \rightarrow B$ be a quasi-isomorphism of degreewise R -free augmented DG algebras. Then for every right B -module M there are natural isomorphisms of $H_A^-(R) \cong H_B^-(R)$ modules*

$$H_A^{+,tw}(f^{-1}M) \cong H_B^{+,tw}(M) \quad \text{and} \quad H_A^\infty(f^{-1}M) \cong H_B^\infty(M).$$

Moreover, these isomorphisms are compatible with the exact triangles of Theorem 5.18.

A DG algebra A is said to satisfy Poincaré duality of degree $d \in \mathbb{Z}$ provided there is a weak equivalence $A \simeq \text{Hom}(A, R[d])$ of A - A -bimodules, where the right and left actions on the target are induced by the left action on A and the trivial left action on $R[d]$, respectively. Here, weak equivalence means that there exist a finite number $A = X_0, X_1, \dots, X_r = \text{Hom}(A, R[d])$ of A -bimodules and for each $0 \leq i < r$ there is a quasi-isomorphism $X_i \rightarrow X_{i+1}$ or $X_{i+1} \rightarrow X_i$ of A -bimodules. According to [Eis19, Theorem A.19] there is under this hypothesis, an isomorphism $H_A^{+,tw}(M) \cong H_A^+(M)[n]$ of $H_A^-(R)$ -modules. We will give a different proof of this result for $A = \Lambda_R[u]$ with $|u| = 3$. This case is sufficient for the application to equivariant instanton Floer homology when $\frac{1}{2} \in R$.

We will now consider these constructions for $A = \Lambda_R[u]$ with $|u| = 3$ as we did for C_A^\pm . The Tate complex $C_A^\infty(M)$, even in this simple case, is big and difficult to handle explicitly. To remedy this we will construct a much simpler and very explicit complex that also computes the Tate homology $H_A^\infty(M)$. Indeed, we will show that if we take one of the double complexes $(D_{*,*}^\pm, \partial', \partial'')$ of Proposition 5.8 and extend them in the natural way to the whole plane, then a suitable totalization of this complex will do the trick.

Lemma 5.23. *Let $A = \Lambda_R[u]$ with $|u| = 3$. Then there is an isomorphism of left A -modules $A \otimes R[\alpha] \cong D_A$ where $|\alpha| = -4$. Under this isomorphism the differential on $A \otimes R[\alpha]$ is given by $\partial(1 \otimes \alpha^p) = -u \otimes \alpha^{p+1}$ and $\partial(u \otimes \alpha^p) = 0$ for $p \geq 0$, and the action of $R[U] = C_A^-(R)$ is given by $U \cdot (a \otimes \alpha^p) = a \otimes \alpha^{p+1}$ for $a \in A$ and $p \geq 0$.*

Furthermore, the map $\rho: R \rightarrow A \otimes R[\alpha]$ of degree 3 given by $1 \mapsto u \otimes \alpha^0$ is an A -linear quasi-isomorphism.

Proof. Using the explicit model of Theorem 5.8 we see that

$$(D_A)_n = C_A^-(A)_n \cong \begin{cases} A_0 & \text{for } n = 4m, m \leq 0 \\ A_3 & \text{for } n = 4m + 3, m \leq 0 \\ 0 & \text{otherwise} \end{cases}$$

and a quick check shows that these identifications are compatible with the left A -module structure. The required isomorphism is therefore obtained by letting $1 \otimes \alpha^p$ and $u \otimes \alpha^p$ correspond to the

generators $1 \in A_0 \cong C_A^-(A)_{-4p}$ and $u \in A_3 = C_A^-(A)_{-4p+3}$, respectively, for $p \geq 0$. The same result shows that the differential and $R[U]$ -action are determined by the formulas $\partial(u \otimes \alpha^p) = 0$, $\partial(1 \otimes \alpha^p) = -u \otimes \alpha^{p+1}$ and $U \cdot (a \otimes \alpha^p) = a \otimes \alpha^{p+1}$ for $a \in A$ and $p \geq 0$.

From this description we find that $\partial: (D_A)_{-4p} \rightarrow (D_A)_{-4(p+1)-3}$ is an isomorphism for each $p \geq 0$. Therefore, the map $\rho: R \rightarrow A \otimes R[\alpha]$ of degree 3 given by $1 \mapsto u \otimes \alpha^0$ is an A -linear quasi-isomorphism. \square

In the following result we establish the isomorphism $H_A^{+,tw}(M) \cong H_A^+(M)[3]$ of $R[U]$ -modules for each right A -module M . It also contains the key ingredients needed to establish the simplified model for $C_A^\infty(M)$.

Lemma 5.24. *Let $A = \Lambda_R[u]$ with $|u| = 3$ and let M be a right A -module. Then there is a quasi-isomorphism $f_M: C_A^+(M) \rightarrow C_A^{+,tw}(M)$ of degree 3 and a chain homotopy $s_M: C_A^+(M) \rightarrow C_A^{+,tw}(M)$ of degree 0 such that*

$$f_M \circ U - U \circ f_M = \partial \circ s_M - s_M \circ \partial.$$

In particular, $H(f_M): H_A^+(M) \rightarrow H_A^{+,tw}(M)$ is a degree 3 isomorphism of $R[U]$ -modules. Furthermore, f_M and s_M are natural in the module M , and in terms of the explicit models $C_A^+(M) \cong \text{Tot}^\oplus(D_{,*}^+, \partial', \partial'')$ and $C_A^-(M) \cong \text{Tot}^\Pi(D_{*,*}^-, \partial', \partial'')$ of Proposition 5.8 the compositions $N_M \circ f_M$ and $N_M \circ s_M$ are given in degree n by*

$$\bigoplus_{p \geq 0} M_{n-4p} \xrightarrow{\pi} M_n \xrightarrow{(-1)^n u} M_{n+3} \xrightarrow{\iota} \prod_{q \geq 0} M_{n+3+4q}$$

$$\bigoplus_{p \geq 0} M_{n-4p} \xrightarrow{\pi} M_n \xrightarrow{\iota} \prod_{q \geq 0} M_{n+4q},$$

respectively.

Proof. Recall that $C_A^+(M) = B(M, A, R)$ and $C_A^{+,tw}(M) = B(M, A, D_A)$. It is convenient to introduce the notation

$$\begin{aligned} m \otimes V^p &:= (-1)^{p(p+1)/2} m[u|u| \cdots |u] \in B(M, A, R) \\ m \otimes V^p \otimes \psi &:= (-1)^{p(p+1)/2} m[u|u| \cdots |u]\psi \in B(M, A, D_A) \end{aligned}$$

for $m \in M$ and $\psi \in D_A$, where u is repeated p times in both formulas. In terms of this notation the differentials and actions of $U \in R[U]$ of $B(M, A, R)$ and $B(M, A, D_A)$ are given by the formulas

$$\begin{aligned} U(m \otimes V^p) &= (-1)^{|m|} m \otimes V^{p-1} \\ U(m \otimes V^p \otimes \psi) &= m \otimes V^p \otimes (U\psi) \\ \partial(m \otimes V^p) &= (-1)^p (\partial_M m + mu \otimes V^{p-1}) \\ \partial(m \otimes V^p \otimes \psi) &= (-1)^p (\partial_M m \otimes V^p + mu \otimes V^{p-1}) \otimes \psi \\ &\quad + m \otimes (V^{p-1} \otimes u\psi + (-1)^{|m|} V^p \otimes \partial_{D_A} \psi). \end{aligned}$$

By the previous lemma there is an isomorphism $D_A \cong A \otimes R[\alpha]$ with $|\alpha| = -4$. To simplify the notation write $a\alpha^p := a \otimes \alpha^p$ for $a \in A$. There is also an A -linear quasi-isomorphism $\rho: R \rightarrow D_A$ of degree 3 given by $\rho(1) = u\alpha^0$. This map induces a quasi-isomorphism

$$f_M := B(1, 1, \rho): B(M, A, R) \rightarrow B(M, A, D_A)$$

of degree 3, which in terms of the above notation is given by $f_M(m \otimes V^p) = (-1)^{|m|} m \otimes V^p \otimes u\alpha^0$ (the sign is forced upon us by the fact that we regard ρ as a map of degree 3). Using the above

formulas we find that

$$(5.7) \quad (f_M \circ U - U \circ f_M)(m \otimes V^p) = m \otimes V^{p-1} \otimes u\alpha^0 - (-1)^{|m|} m \otimes V^p \otimes u\alpha^1.$$

Define $s_M: B(M, A, R) \rightarrow B(M, A, D_A)$ of degree 0 by $s_M(m \otimes V^p) = m \otimes V^p \otimes \alpha^0$. A straightforward calculation using the above formulas and the formula for the differential in $D_A \cong A \otimes R[\alpha]$ given in the previous lemma shows that

$$\begin{aligned} \partial s_M(m \otimes V^p) &= (-1)^p (\partial_M m \otimes V^p \otimes \alpha^0 + mu \otimes V^{p-1} \otimes \alpha^0) \\ &\quad + m \otimes V^{p-1} \otimes u\alpha^0 - (-1)^{|m|} m \otimes V^p \otimes u\alpha^1 \\ s_M \partial(m \otimes V^p) &= (-1)^p (\partial_M m \otimes V^p \otimes \alpha^0 + mu \otimes V^{p-1} \otimes \alpha^0). \end{aligned}$$

Combining this with (5.7) we obtain $f_M \circ U - U \circ f_M = \partial s_M - s_M \partial$ as required.

Finally, we need to consider the compositions of f_M and s_M with the norm map $N_M: C_A^{+,tw}(M) \rightarrow C_A^-(M)$ in terms of the models given in Proposition 5.8. By (5.3) $m \in M_{q-3p} = D_{p,q}^+$ corresponds to $(-1)^{|m|p} m \otimes V^p$, which is mapped by f_M to $(-1)^{|m|(p+1)} m \otimes V^p \otimes u\alpha^0$. The norm map kills off all such elements with $p > 0$ and sends $m \otimes V^0 \otimes u\alpha^0 \mapsto mu \cdot \alpha^0$. This is the functional that sends V^0 to $mu \in M_{q+3}$ and vanishes on V^q for $q > 0$. By (5.4) the identification $C_A^-(M)_n \cong \prod_{q \geq 0} M_{n+4q}$ is given by $\psi \mapsto (\psi(V^q))_{q \geq 0}$. We conclude that $N_M \circ f_M$ is given by the composition in the statement of the lemma. The verification for $N_M \circ s_M$ is analogous. \square

We need two simple but slightly technical results before we can reach our goal.

Lemma 5.25. *Let X and Y be DG $R[U]$ -modules and suppose that $f: X \rightarrow Y$ is a degree d map of $R[U]$ -modules up to homotopy; that is, f is a chain map of degree d and there is a chain homotopy $s: X \rightarrow Y$ such that $fU - Uf = \partial_Y s + (-1)^d s \partial_X$. Let $\text{Cone}(f)$ be given as in Definition 5.15, except that we redefine the $R[U]$ -module structure by setting*

$$U \cdot \begin{pmatrix} y \\ x \end{pmatrix} = \begin{pmatrix} U & (-1)^{|s|} s \\ 0 & U \end{pmatrix} \begin{pmatrix} y \\ x \end{pmatrix} = \begin{pmatrix} Uy - (-1)^d sx \\ Ux \end{pmatrix}$$

for all $(y, x) \in Y_n \oplus X_{n-d-1} = \text{Cone}(f)_n$. Then $\text{Cone}(f)$ is a DG $R[U]$ -module and the maps in the natural short exact sequence

$$0 \longrightarrow Y \longrightarrow \text{Cone}(f) \longrightarrow X[d+1] \longrightarrow 0$$

are (strict) maps of $R[U]$ -modules.

Proof. It suffices to check that $\partial_{\text{Cone}(f)} U = U \partial_{\text{Cone}(f)}$. This follows easily from the relations $\partial_X U = U \partial_X$, $\partial_Y U = U \partial_Y$ and $fU - Uf = \partial_Y s + (-1)^d s \partial_X$. Next, the maps in the natural exact sequence are given degreewise by the inclusion $Y_n \rightarrow Y_n \oplus X_{n-d-1}$ and the projection $Y_n \oplus X_{n-d-1} \rightarrow X_{n-d-1}$ and are seen to commute with the action of U . \square

Lemma 5.26. *Let $f: X \rightarrow Y$ be a degree 3 map of DG $R[U]$ -modules up to homotopy, and let $g: Y \rightarrow Z$ be a degree 0 map of DG $R[U]$ -modules. Then $g \circ f: X \rightarrow Z$ is a degree 3 map of $R[U]$ -modules up to homotopy and there is a commutative diagram*

$$\begin{array}{ccccccc} 0 & \longrightarrow & Z & \longrightarrow & \text{Cone}(g \circ f) & \longrightarrow & X[4] \longrightarrow 0 \\ & & \downarrow = & & \downarrow \text{Cone}(f,1) & & \downarrow f[1] \\ 0 & \longrightarrow & Z & \longrightarrow & \text{Cone}(g) & \longrightarrow & Y[1] \longrightarrow 0 \end{array}$$

with short exact rows. If $\text{Cone}(g \circ f)$ is given the DG $R[U]$ -module structure of the previous lemma, then the maps $\text{Cone}(f, 1)$ and $f[1]$ are maps of DG $R[U]$ -modules up to homotopy and all the other maps in the diagram are strict maps of DG $R[U]$ -modules.

Proof. By assumption there is a chain homotopy $s: X \rightarrow Y$ of degree 0 such that $fU - Uf = \partial s - s\partial$. Then for $h := g \circ f$ and $t := g \circ s$ we have $hU - Uh = \partial t - t\partial$ as well. This shows that h is a map of $R[U]$ -modules up to homotopy. By functoriality of the cone construction we obtain a commutative diagram of DG modules with short exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & Z & \longrightarrow & \text{Cone}(g \circ f) & \longrightarrow & X[4] \longrightarrow 0 \\ & & \downarrow = & & \downarrow \text{Cone}(f, 1) & & \downarrow f[1] \\ 0 & \longrightarrow & Z & \longrightarrow & \text{Cone}(g) & \longrightarrow & Y[1] \longrightarrow 0. \end{array}$$

Here, $\text{Cone}(f, 1): \text{Cone}(g \circ f)_n = Z_n \oplus X_{n-4} \rightarrow Z_n \oplus Y_{n-1} = \text{Cone}(g)_n$ is given by $(z, x) \mapsto (z, -fx)$, and $f[1]: X[4]_n = X_{n-4} \rightarrow Y_{n-1} = Y[1]_n$ is given by $x \mapsto -fx$. Equip $\text{Cone}(g \circ f)$ with the adjusted $R[U]$ -module structure given in the previous lemma. Then all the horizontal maps in the diagram are strict maps of $R[U]$ -modules. The right hand vertical map is a map of $R[U]$ -modules up to homotopy, so the only thing we have to verify is that $\text{Cone}(f, 1)$ is a map of $R[U]$ -modules up to homotopy.

We claim that $v: \text{Cone}(h) \rightarrow \text{Cone}(g)$ given by $v(z, x) = (0, sx)$ is the required homotopy. Indeed, using matrix notation for the various maps,

$$\begin{aligned} C(f, 1)U - UC(f, 1) &= \begin{pmatrix} 1 & 0 \\ 0 & -f \end{pmatrix} \begin{pmatrix} U & t \\ 0 & U \end{pmatrix} - \begin{pmatrix} U & 0 \\ 0 & U \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -f \end{pmatrix} \\ &= \begin{pmatrix} 0 & t \\ 0 & Uf - fU \end{pmatrix} \\ \partial_{\text{Cone}(g)}v + v\partial_{\text{Cone}(h)} &= \begin{pmatrix} \partial_Z & g \\ 0 & -\partial_Y \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & s \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & s \end{pmatrix} \begin{pmatrix} \partial_Z & -h \\ 0 & \partial_X \end{pmatrix} \\ &= \begin{pmatrix} 0 & g \circ s \\ 0 & s\partial_X - \partial_Y s \end{pmatrix}, \end{aligned}$$

where $C(f, 1)$ is shorthand for $\text{Cone}(f, 1)$. Therefore, as $t = g \circ s$ and $\partial_Y s - s\partial_X s = fU - Uf$, we deduce that $\text{Cone}(f, 1)$ is $R[U]$ -linear up to homotopy. \square

Given a sequence of R -modules $\{D_s\}_{s \in \mathbb{Z}}$ we write $\prod_{s \rightarrow -\infty} D_s$ for the submodule of $\prod_s D_s$ consisting of the sequences $(x_s)_{s \in \mathbb{Z}}$ for which there exists s_0 such that $x_s = 0$ for all $s \geq s_0$. Similarly, $\prod_{s \rightarrow \infty} D_s$ is the submodule of $(x_s)_{s \in \mathbb{Z}}$ with $x_s = 0$ for all sufficiently small s .

Definition 5.27. Given a double complex $E = (D_{*,*}, \partial', \partial'')$, define $\text{Tot}^{\Pi, -\infty}(E)$ and $\text{Tot}^{\Pi, +\infty}(E)$ to be the complexes that in degree n are given by

$$\prod_{s \rightarrow -\infty} D_{s, n-s} \quad \text{and} \quad \prod_{s \rightarrow \infty} D_{s, n-s}$$

respectively, with differential $\partial = \partial' + \partial''$.

Theorem 5.28. Let $A = \Lambda_R[u]$ with $|u| = 3$ and let M be a right DG A -module. Define $D_{p,q}^\infty = M_{q-3p}$ for all $p, q \in \mathbb{Z}$. Set

$$\begin{aligned} \partial' &= (-1)^{p+q+1}u: D_{p,q}^\infty = M_{q-3p} \rightarrow M_{q-3p+3} = D_{p-1,q}^\infty \\ \partial'' &= \partial_M: D_{p,q}^\infty = M_{q-3p} \rightarrow M_{q-3p-1} = D_{p,q-1}^\infty \end{aligned}$$

and let $U: D_{p,q}^\infty = M_{q-3p} \rightarrow M_{q-3p} = D_{p-1,q-3}^\infty$ be the identity for all p, q . Then there is a quasi-isomorphism, natural in M ,

$$\text{Tot}^{\Pi, -\infty}(D_{*,*}, \partial', \partial'') \rightarrow C_A^\infty(M),$$

which is a homomorphism of $R[U]$ -modules up to homotopy.

Proof. By Lemma 5.24 there is a quasi-isomorphism $f = f_M: C_A^+(M) \rightarrow C_A^{+,tw}(M)$ and a homotopy $s = s_M: C_A^+(M) \rightarrow C_A^{+,tw}(M)$ such that $fU - Uf = \partial s - s\partial$. We apply Lemma 5.26 for $f: C_A^+(M) \rightarrow C_A^{+,tw}(M)$ and the norm map $N_M: C_A^{+,tw}(M) \rightarrow C_A^-(M)$ to obtain the commutative diagram with exact rows

$$(5.8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & C_A^-(M) & \longrightarrow & \text{Cone}(N_M \circ f) & \longrightarrow & C_A^+(M)[4] \longrightarrow 0 \\ & & \downarrow = & & \downarrow \text{Cone}(f,1) & & \downarrow f[1] \\ 0 & \longrightarrow & C_A^-(M) & \longrightarrow & C_A^\infty(M) & \longrightarrow & C_A^{+,tw}(M)[1] \longrightarrow 0, \end{array}$$

where we recall that $C_A^\infty(M) = \text{Cone}(N_M)$. By the same result $\text{Cone}(f, 1)$ is a map of $R[U]$ -modules up to homotopy. As both the outer vertical maps are quasi-isomorphisms, it follows by passage to the long exact sequences in homology and the 5-lemma that $\text{Cone}(f, 1)$ is a quasi-isomorphism. Thus, $\text{Cone}(f, 1)$ is the required map. This quasi-isomorphism is natural in M , since both f and N_M are natural transformations in M and the cone construction is functorial.

The remaining task is to establish an isomorphism $\text{Cone}(N_M \circ f) \cong \text{Tot}^{\Pi, -\infty}(D_{*,*}^\infty, \partial', \partial'')$. Using the identifications of Proposition 5.8 we obtain

$$\text{Cone}(N_M \circ f)_n = C_A^-(M)_n \oplus C_A^+(M)_{n-4} \cong \left(\prod_{q \geq 0} M_{n+4q} \right) \oplus \left(\bigoplus_{p \geq 0} M_{n-4(p+1)} \right)$$

for each $n \in \mathbb{Z}$. Moreover, by Lemma 5.24 $h := N_M \circ f$ and $t := N_M \circ s$ are given on $(x_p)_{p \geq 0} \in C_A^+(M)_n$ by $h((x_p)_p) = ((-1)^{|x_0|} x_0 u, 0, \dots)$ and $t((x_p)_{p \geq 0}) = (x_0, 0, 0, \dots)$. Therefore, using the formulas

$$\partial_{\text{Cone}(h)} = \begin{pmatrix} \partial_{C_A^-(M)} & -h \\ 0 & \partial_{C_A^+(M)} \end{pmatrix} \quad \text{and} \quad U = \begin{pmatrix} U & t \\ 0 & U \end{pmatrix},$$

as well as the description of the differentials and the U -action in terms of the models for $C_A^\pm(M)$, we deduce that $\text{Cone}(h) \cong \text{Tot}^{\Pi, -\infty}(D_{*,*}^\infty, \partial', \partial'')$. \square

A part of the double complex $(D_{*,*}^\infty, \partial', \partial'')$ is shown below.

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \vdots & & \vdots \\ & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longleftarrow & M_8 & \xleftarrow{u} & M_5 & \xleftarrow{-u} & M_2 & \xleftarrow{u} & M_{-1} & \longleftarrow \cdots \\ & & \downarrow \partial_M & & \downarrow \partial_M & & \downarrow \partial_M & & \downarrow \partial_M & \\ \cdots & \longleftarrow & M_7 & \xleftarrow{-u} & M_4 & \xleftarrow{u} & M_1 & \xleftarrow{-u} & M_{-2} & \longleftarrow \cdots \\ & & \downarrow \partial_M & & \downarrow \partial_M & & \downarrow \partial_M & & \downarrow \partial_M & \\ \cdots & \longleftarrow & M_6 & \xleftarrow{u} & M_3 & \xleftarrow{-u} & M_0 & \xleftarrow{u} & M_{-3} & \longleftarrow \cdots \\ & & \downarrow \partial_M & & \downarrow \partial_M & & \downarrow \partial_M & & \downarrow \partial_M & \\ \cdots & \longleftarrow & M_5 & \xleftarrow{-u} & M_2 & \xleftarrow{u} & M_{-1} & \xleftarrow{-u} & M_{-4} & \longleftarrow \cdots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\ & & \vdots & & \vdots & & \vdots & & \vdots & \end{array}$$

This is indeed the natural concatenation of the two double complexes $D_{*,*}^+$ and $D_{*,*}^-$.

Corollary 5.29. *Let M be a right $A = \Lambda_R[u]$ -module. Then $U: H_A^\infty(M) \rightarrow H_A^\infty(M)$ is an isomorphism.*

Proof. $U: \text{Tot}^{\Pi, -\infty}(D_{*,*}^\infty, \partial', \partial'') \rightarrow \text{Tot}^{\Pi, -\infty}(D_{*,*}^\infty, \partial', \partial'')$ is an isomorphism. \square

As in the cases C_A^\pm we will need calculations for $M = R$, $M = R \oplus R[2]$ and $M = \Lambda_R[u]$. This is easily achieved with the help of the above explicit model.

Corollary 5.30. *We have*

$$H_A^\infty(A) = 0, \quad H_A^\infty(R) = R[T, T^{-1}] \quad \text{and} \quad H_A^\infty(R \oplus R[2]) = R[S, S^{-1}],$$

where $|T| = -4$, $|S| = -2$ and the $R[U]$ -action is given by $U \cdot T^i = T^{i+1}$ and $U \cdot S^i = S^{i+2}$ for all $i \in \mathbb{Z}$.

We note once again that our notation should not be interpreted to mean that there is any product structure in $H_A^\infty(R \oplus R[2])$ or $H_A^\infty(R)$.

5.4. The Equivariant Instanton Floer Groups. In this section we will give the definition of the various flavors of equivariant instanton Floer homology. Before we can get to this we have to recall a few definitions concerning filtered DG modules. Let M be a DG module equipped with an increasing filtration

$$\cdots \subset F_{p-1}M \subset F_pM \subset F_{p+1}M \subset \cdots \subset M$$

by DG submodules. The minimal filtration quotients $F_pM/F_{p-1}M$ enter many times in the following discussion, so we introduce the simpler notation $\overline{F}_pM := F_pM/F_{p-1}M$. The filtration gives rise to a homological spectral sequence with

$$(5.9) \quad E_{p,q}^1 = H_{p+q}(\overline{F}_pM) \implies H_{p+q}(M)$$

(we will elaborate on this in the next section). However, without any additional information, this spectral sequence may fail to converge in any sense to the desired target $H(M)$.

The following properties of a filtration are highly desirable in order to achieve any reasonable convergence properties of the above spectral sequence. The filtration $\{F_pM\}_p$ of M is said to be [Boa99, Definition 2.1]

- (i) exhaustive if $\text{colim}_p F_pM = \bigcup_p F_pM = M$,
- (ii) Hausdorff if $\lim_p F_pM = \bigcap_p F_pM = 0$ and
- (iii) complete if $\text{Rlim}_p F_pM = 0$.

Here Rlim is the first (and only nonzero) right derived functor of \lim (see [Boa99, Section 1] for additional details). It is important to note that these limits are calculated degreewise. In particular, if the filtration is degreewise bounded below, that is, for each degree n there exists $p_0(n)$ such that $F_pM_n = 0$ for all $p \leq p_0$, then the filtration is automatically complete Hausdorff. Similarly, if the filtration is degreewise bounded above then the filtration is automatically exhaustive.

Definition 5.31. Let M be a DG module equipped with an increasing filtration $\{F_pM\}_p$ by DG submodules. Then the full completion of $(M, \{F_pM\}_p)$ is defined to be the complex

$$\hat{M} := \lim_q \text{colim}_p F_pM/F_qM$$

equipped with the filtration $F_p\hat{M} := \lim_q F_pM/F_qM$.

Remark 5.32. The standard completion of a filtered module M is typically defined by $\lim_q M/F_qM$ (see for instance [Boa99, p. 2.7]) and is naturally filtered by $\{\lim_q F_pM/F_qM\}_p$. We have therefore chosen to use the terminology full completion to distinguish it from this notion. The lemma below demonstrates that the two notions coincide if the filtration of M is exhaustive.

A map $f: M \rightarrow M'$ between filtered DG modules is filtered if $f(F_p M) \subset F_p M'$ for all $p \in \mathbb{Z}$. There is therefore a category of filtered DG modules and filtered maps. A sequence $M' \rightarrow M \rightarrow M''$ of filtered DG modules and filtered maps is short exact if $F_p M' \rightarrow F_p M \rightarrow F_p M''$ is short exact for each $p \in \mathbb{Z}$. The relevant properties of the full completion are summarized in the following lemma.

Lemma 5.33. *Let M be a filtered DG module. Then the following holds true.*

- (i) *The filtration $F_p \hat{M} = \lim_{q < p} F_p M / F_q M$ of \hat{M} is exhaustive and complete Hausdorff.*
- (ii) *For each $q < p$ there is a natural isomorphism $F_p M / F_q M \rightarrow F_p \hat{M} / F_q \hat{M}$.*
- (iii) *If the filtration of M is exhaustive, then $\hat{M} = \lim_q M / F_q M$.*
- (iv) *If the filtration of M is complete Hausdorff, then $\hat{M} = \operatorname{colim}_p F_p M = \bigcup_p F_p M$.*

The full completion $M \mapsto \hat{M}$ defines a functor from the category of filtered modules to itself, which preserves short exact sequences.

Proof. Let M be a filtered DG module and set $N = \operatorname{colim}_p F_p M = \bigcup_p F_p M$ filtered by $F_p N = F_p M$. Then the full completion of M coincides with the standard completion of N , i.e., $\hat{M} = \lim_q N / F_q N$. Therefore part (i) and (ii) follow from the corresponding properties of the standard completion (see [Boa99, Proposition 2.8]). If the filtration of M is exhaustive, then $\operatorname{colim}_p F_p M / F_q M = M / F_q M$ for each q and part (iii) easily follows. For part (iv)

$$\hat{M} = \lim_q \operatorname{colim}_p F_p M / F_q M = \lim_q N / F_q N = N = \operatorname{colim}_p F_p M.$$

It is clear that $M \mapsto \hat{M}$ is functorial with respect to filtered maps. The exactness of \hat{M} follows from the exactness of sequential colimits and exactness of the standard completion in the category of R -modules (see for instance [AM69, Corollary 10.3]). \square

Remark 5.34. If B is a DG algebra and M is a left or right B -module equipped with a filtration by B -submodules the definition of the full completion and the above lemma apply without change. The reason for this is that quotients, limits and colimits are formed in the category of graded modules, with the B -action and differentials carried along.

Recall from the previous section that for a degreewise R -free augmented DG algebra A , four functors C_A^\bullet , $\bullet \in \{+, -, (+, tw), \infty\}$, from the category of right A -modules to the category of left $C_A^-(R)$ -modules were defined. By Theorem 5.6 and Theorem 5.18 these are all exact and preserve quasi-isomorphisms. Therefore, if M is a right A -module equipped with a filtration $F_p M$ by A -submodules, $C_A^\bullet(M)$ carries a natural filtration $F_p C_A^\bullet(M) := C_A^\bullet(F_p M)$ by $C_A^-(R)$ -submodules. Now, even if the initial filtration of M is exhaustive and complete Hausdorff, the resulting filtration of $C_A^\bullet(M)$ may fail to satisfy these properties. This motivates the following definition.

Definition 5.35. Let A be a degreewise R -free augmented DG algebra. For $\bullet \in \{+, -, (+, tw), \infty\}$ define a functor \hat{C}_A^\bullet from the category of filtered right A -modules to the category of filtered left $C_A^-(R)$ -modules by

$$\begin{aligned} \hat{C}_A^\bullet(M) &:= \lim_q \operatorname{colim}_p C_A^\bullet(F_p M / F_q M) \quad \text{filtered by} \\ F_p \hat{C}_A^\bullet(M) &:= \lim_{q < p} C_A^\bullet(F_p M / F_q M) \quad \text{for all } p \in \mathbb{Z}. \end{aligned}$$

In other words, \hat{C}_A^\bullet is the composition of the natural extension of C_A^\bullet to filtered A -modules followed by the full completion functor. We denote the homology of $\hat{C}_A^\bullet(M)$ by $\hat{H}_A^\bullet(M)$.

The various flavors of equivariant instanton Floer homology are obtained by applying this construction to the framed Floer complex.

Definition 5.36. Let $A = C_*^{gm}(\mathrm{SO}(3))$ and let $M = \widetilde{CI}(Y, E)$ be the framed Floer complex associated with an $\mathrm{SO}(3)$ -bundle E over a closed oriented three-manifold Y , equipped with the index filtration. For $\bullet \in \{+, -, (+, tw), \infty\}$ define

$$CI^\bullet(Y, E) := \hat{C}_A^\bullet(M) \quad \text{and} \quad I^\bullet(Y, E) := H(CI^\bullet(Y, E)).$$

The groups $I^\bullet(Y, E)$, for $\bullet \in \{+, -, (+, tw), \infty\}$, are called the positive, negative, positive twisted and Tate equivariant instanton Floer homologies of (Y, E) , respectively. These homology groups are modules over $H_A^-(R)$.

For $A = \Lambda_R[u]$ and $M = DCI(Y, E)$ we similarly define the complexes

$$DCI^\bullet(Y, E) = \hat{C}_A^\bullet(M).$$

Remark 5.37. For $A = C_*(\mathrm{SO}(3); R)$ and $M = R$ it holds true that $H_A^-(R) \cong H^{-*}(\mathrm{BSO}(3); R)$ (see [Eis19, Lemma A.21]).

The norm map $N_M: C_A^{+, tw}(M) \rightarrow C_A^-(M)$ is a natural in M and therefore extends to a natural map $\hat{N}_M: \hat{C}_A^{+, tw}(M) \rightarrow \hat{C}_A^-(M)$ for any filtered module M . The following simple lemma ensures that $\hat{C}_A^\infty(M)$ is naturally isomorphic to $\mathrm{Cone}(\hat{N}_M)$.

Lemma 5.38. *Let A be a degree-wise R -free augmented DG algebra and let M be a filtered right A -module. Then there is a natural (filtered) isomorphism of $C_A^-(R)$ -modules*

$$\mathrm{Cone}(\hat{N}_M) \cong \hat{C}_A^\infty(M).$$

In particular, we have the following exact triangle of $H_A^-(R)$ -modules.

$$\begin{array}{ccc} \hat{H}_A^{+, tw}(M) & \xrightarrow{H(\hat{N}_M)} & \hat{H}_A^-(M) \\ & \swarrow [-1] \quad \searrow [0] & \\ & \hat{H}_A^\infty(M) & \end{array}$$

Proof. For $q < p$ and n there is a natural identification

$$\frac{F_p C_A^\infty(M)}{F_q C_A^\infty(M)}_n \cong C_A^\infty \left(\frac{F_p M}{F_q M} \right)_n = C_A^- \left(\frac{F_p M}{F_q M} \right)_n \oplus C_A^{+, tw} \left(\frac{F_p M}{F_q M} \right)_{n-1}.$$

In terms of the right hand side the differential is given by

$$\begin{pmatrix} \partial_{p,q}^- & N_{p,q} \\ 0 & -\partial_{p,q}^{+, tw} \end{pmatrix},$$

where $\partial_{p,q}^-$ and $\partial_{p,q}^{+, tw}$ denote the differential in the first and second summand respectively and $N_{p,q}$ denotes the evident norm map. Applying $\lim_q \mathrm{colim}_p$ to this identification, taking into account that this is calculated degree-wise with differentials and A -module structure carried along, we obtain $\hat{C}_A^\infty(M)_n \cong \hat{C}_A^-(M)_n \oplus \hat{C}_A^{+, tw}(M)_{n-1}$ in each degree n . The differential on the right hand side is given by

$$\begin{pmatrix} \lim_q \mathrm{colim}_p \partial_{p,q}^- & \lim_q \mathrm{colim}_p N_{p,q} \\ 0 & \lim_q \mathrm{colim}_p \partial_{p,q}^{+, tw} \end{pmatrix}.$$

Therefore, the right hand side is precisely $\mathrm{Cone}(\hat{N}_M)$, since by definition $\hat{N}_M = \lim_q \mathrm{colim}_p N_{p,q}$. \square

To verify that $DCI^\bullet(Y, E)$ also calculates $I^\bullet(Y, E)$ when $\frac{1}{2} \in R$ we will establish the necessary invariance results for the functors \hat{C}_A^\bullet . The appropriate tool is the following lemma, which is a simple consequence of the Eilenberg-Moore comparison theorem of [Wei94, Theorem 5.5.11]. For a filtered map $f: M \rightarrow N$ between filtered DG modules we use the notation $\bar{f}_p: \bar{F}_p M \rightarrow \bar{F}_p N$ for the induced maps between the the minimal filtration quotients.

Lemma 5.39. *Let M and N be DG modules equipped with exhaustive and complete Hausdorff filtrations. Then if $f: M \rightarrow N$ is a filtered map such that the induced map $\bar{f}_p: \bar{F}_p M \rightarrow \bar{F}_p N$ is a quasi-isomorphism for each $p \in \mathbb{Z}$, then f is a quasi-isomorphism.*

Proposition 5.40. *Let A be a degreewise R -free augmented DG algebra. Suppose $f: M \rightarrow N$ is a filtered map of filtered right A -modules such that $\bar{f}_p: \bar{F}_p M \rightarrow \bar{F}_p N$ is a quasi-isomorphism for each $p \in \mathbb{Z}$. Then for $\bullet \in \{+, -, (+, tw), \infty\}$ the induced map*

$$\hat{f}^\bullet: \hat{C}_A^\bullet(M) \rightarrow \hat{C}_A^\bullet(N)$$

is a (filtered) quasi-isomorphism of $C_A^-(R)$ -modules. Furthermore, upon passage to homology, these isomorphisms are compatible with the exact triangle of Lemma 5.38.

Proof. Let $\bullet \in \{+, -, (+, tw), \infty\}$ and note that the functors C_A^\bullet and \hat{C}_A^\bullet are covariant in M . For simplicity of notation let $g = \hat{f}^\bullet: \hat{C}_A^\bullet(M) \rightarrow \hat{C}_A^\bullet(N)$ denote the induced filtered map of $C_A^-(R)$ -modules. Then by Lemma 5.33 part (ii) and the exactness of C_A^\bullet , there is for each p a commutative diagram

$$\begin{array}{ccc} C_A^\bullet(\bar{F}_p M) & \xrightarrow{(\bar{f}_p)^\bullet} & C_A^\bullet(\bar{F}_p N) \\ \downarrow \cong & & \downarrow \cong \\ \bar{F}_p \hat{C}_A^\bullet(M) & \xrightarrow{\bar{g}_p} & \bar{F}_p \hat{C}_A^\bullet(N). \end{array}$$

By assumption \bar{f}_p is a quasi-isomorphism, so since the functor C_A^\bullet preserves quasi-isomorphisms it follows that $(\bar{f}_p)^\bullet$ is a quasi-isomorphism. As the vertical maps are isomorphisms, we conclude that \bar{g}_p is a quasi-isomorphism. By Lemma 5.33 part (i) the filtrations of $\hat{C}_A^\bullet(M)$ and $\hat{C}_A^\bullet(N)$ are exhaustive and complete Hausdorff. We conclude that $g = \hat{f}^\bullet$ is a quasi-isomorphism using the comparison result; Lemma 5.39. The lemma applies equally well to the filtered subcomplexes. Consequently, the restrictions $g_p: F_p \hat{C}_A^\bullet(M) \rightarrow F_p \hat{C}_A^\bullet(N)$ are also quasi-isomorphisms for all p . This shows that g is in fact a filtered quasi-isomorphism. \square

Next we have to establish a similar invariance result for variations in the DG algebra.

Proposition 5.41. *Let $f: A \rightarrow B$ be a quasi-isomorphism of degreewise R -free augmented algebras and let M be a filtered B -module. Then there is for each $\bullet \in \{+, -, (+, tw), \infty\}$ a natural isomorphism of $H_A^-(R) \cong H_B^-(R)$ -modules*

$$\hat{H}_A^\bullet(f^{-1}M) \cong \hat{H}_B^\bullet(M),$$

compatible with the exact triangles of Lemma 5.38.

Proof. We first consider the case $\bullet = +$. For each pair $q < p$ there is by Proposition 5.5 a natural quasi-isomorphism

$$f_*: C_A^+(F_p f^{-1}M / F_q f^{-1}M) \rightarrow C_B^+(F_p M / F_q M)$$

linear along $f^*: C_B^-(R) \rightarrow C_A^-(R)$. By applying $\lim_q \operatorname{colim}_p$ we obtain a filtered map $g = \hat{f}_*: \hat{C}_A^+(f^{-1}M) \rightarrow \hat{C}_B^+(M)$ linear along $f^*: C_B^-(R) \rightarrow C_A^-(R)$. By construction, the following diagram commutes for each p .

$$\begin{array}{ccc} C_A^+(\bar{F}_p f^{-1}M) & \xrightarrow{(\bar{f}_*)^p} & C_B^+(\bar{F}_p M) \\ \downarrow \cong & & \downarrow \cong \\ \bar{F}_p \hat{C}_A^+(f^{-1}M) & \xrightarrow{\bar{g}_p} & \bar{F}_p \hat{C}_B^+(M) \end{array}$$

As the upper horizontal map is a quasi-isomorphism, so is the lower horizontal map and we conclude that $g = \hat{f}_*$ is a filtered quasi-isomorphism by Lemma 5.39. For $\bullet = -$ there is for each pair $q < p$ a natural quasi-isomorphism

$$f^*: C_B^-(F_p M / F_q M) \rightarrow C_A^-(F_p f^{-1} M / F_q f^{-1} M).$$

By passing to the colimit and limit we obtain a filtered map $\hat{f}^*: \hat{C}_B^-(M) \rightarrow \hat{C}_A^-(f^{-1} M)$, linear along $f^*: C_B^-(R) \rightarrow C_A^-(R)$. The same argument as above shows that this is a filtered quasi-isomorphism.

For the two remaining cases recall that there is a relative functor $C_f^{+,tw}$, see Definition 5.19, from the category of right B -modules to the category of left $C_A^-(R)$ -modules. By Lemma 5.20 this functor is exact and preserves quasi-isomorphism, so we may promote it to a functor $\hat{C}_f^{+,tw}$ between the corresponding filtered categories as we did with the others. By passing to the colimit and then the limit in the commutative diagram of Proposition 5.21 applied to $F_p M / F_q M$ for all $q < p$, we obtain a commutative diagram

$$\begin{array}{ccccc} \hat{C}_B^{+,tw}(M) & \xrightarrow{\hat{\beta}_M} & \hat{C}_f^{+,tw}(M) & \xleftarrow{\hat{\alpha}_M} & \hat{C}_A^{+,tw}(f^{-1} M) \\ \downarrow \hat{N}_M^B & & \downarrow \hat{N}_M^f & & \downarrow \hat{N}_M^A \\ \hat{C}_B^-(M) & \xrightarrow{\hat{f}^*} & \hat{C}_A^-(f^{-1} M) & \xleftarrow{=} & \hat{C}_A^-(f^{-1} M) \end{array}$$

of filtered DG modules over $C_A^-(R)$ or $C_B^-(R)$ as appropriate. Furthermore, all the maps are filtered and $C_A^-(R)$ -linear, $C_B^-(R)$ -linear or linear along $f^*: C_B^-(R) \rightarrow C_A^-(R)$ as appropriate. By the same proposition, as $f: A \rightarrow B$ is a quasi-isomorphism of degreewise free DG algebras, all the horizontal arrows in the diagram applied to $F_p M / F_q M$ are quasi-isomorphisms. This implies, by the now familiar argument, that all the horizontal arrows in the above diagram are filtered quasi-isomorphisms. In particular, there is a natural isomorphism $\hat{H}_A^{+,tw}(f^{-1} M) \cong \hat{H}_B^{+,tw}(M)$. Moreover, we obtain a zigzag of quasi-isomorphisms between the cones of the vertical maps

$$\text{Cone}(\hat{N}_M^B) \longrightarrow \text{Cone}(\hat{N}_M^f) \longleftarrow \text{Cone}(\hat{N}_{f^{-1}M}^A).$$

In view of Lemma 5.38, we obtain an isomorphism $\hat{H}_A^\infty(f^{-1} M) \cong \hat{H}_B^\infty(M)$ linear along $H_A^-(R) \cong H_B^-(R)$ as well. The above diagram also ensures that these isomorphisms are compatible with the corresponding exact triangles. \square

The key application of these invariance results is the following.

Corollary 5.42. *Suppose that $\frac{1}{2} \in R$. Then for each $\bullet \in \{+, -, (+, tw), \infty\}$ there is an isomorphism $H(DCI^\bullet(Y, E)) \cong I^\bullet(Y, E)$ of $H_A^-(R) \cong R[U]$ -modules. These isomorphisms are compatible with the corresponding exact triangles.*

Proof. As $\frac{1}{2} \in R$ there are, according to Proposition 2.9, a quasi-isomorphism $i: \Lambda_R[u] \rightarrow C_*^{gm}(\text{SO}(3))$ of DG algebras and a zigzag

$$\widetilde{CI}(Y, E) \xrightarrow{f} X \xleftarrow{g} DCI(Y, E)$$

of filtered objects and filtered $\Lambda_R[u]$ -homomorphisms such that \bar{f}_p and \bar{g}_p are quasi-isomorphism for each $p \in \mathbb{Z}$. The required result now follows from Propositions 5.40 and 5.41. \square

For $A = \Lambda_R[u]$ we can extend a number of the results from the previous section.

Proposition 5.43. *Let $A = \Lambda_R[u]$ and let M be a filtered A -module. Then there is a filtered quasi-isomorphism $\hat{f}_M: \hat{C}_A^+(M) \rightarrow \hat{C}_A^{+,tw}(M)$ of degree 3 and a (filtered) homotopy $\hat{s}_M: \hat{C}_A^+(M) \rightarrow$*

$\hat{C}_A^{+,tw}(M)$ such that $\hat{f}_M U - U \hat{f}_M = \partial \hat{s}_M - \hat{s}_M \partial$ for $U \in R[U] \cong C_A^-(R)$. Moreover, there is a (filtered) quasi-isomorphism

$$\text{Cone}(\hat{N}_M \circ \hat{f}) \rightarrow \text{Cone}(\hat{N}_M) \cong \hat{C}_A^\infty(M),$$

which is $R[U]$ -linear up to homotopy, where the source is given the $R[U]$ -module structure determined by the homotopy $\hat{t}_M = \hat{N}_M \circ \hat{s}_M$ as in Lemma 5.25. Furthermore, the resulting isomorphism in homology is compatible with the associated exact triangles.

Proof. Recall from Lemma 5.24 that there is for each right A -module N a quasi-isomorphism $f_N: C_A^+(N) \rightarrow C_A^{+,tw}(N)$ of degree 3 and a homotopy $s_N: C_A^+(N) \rightarrow C_A^{+,tw}(N)$ such that $f_N U - U f_N = \partial s_N - s_N \partial$. Let $f_{p,q}$ and $s_{p,q}$ denote the maps obtained for $N = F_p M / F_q M$ for $q < p$. By naturality we may pass to the colimit over p and then to the limit over q to obtain maps $\hat{f}_M: \hat{C}_A^+(M) \rightarrow \hat{C}_A^{+,tw}(M)$ and $\hat{s}_M: \hat{C}_A^+(M) \rightarrow \hat{C}_A^{+,tw}(M)$. As $f_{p,q} U - U f_{p,q} = s_{p,q} \partial - \partial s_{p,q}$ is valid for each pair $q < p$, it follows that $\hat{f}_M U - U \hat{f}_M = \partial \hat{s}_M - \hat{s}_M \partial$. The fact that \hat{f}_M is a quasi-isomorphism is proved as before using Lemma 5.39.

The final statement follows by applying $\lim_q \text{colim}_p$ to the diagram (5.8) in the proof of Theorem 5.28 for $F_p M / F_q M$ and another application of Lemma 5.39. It is clear from this construction that one obtains an isomorphism between the corresponding exact triangles in homology. \square

Combining the above proposition with Corollary 5.42 we obtain the following.

Corollary 5.44. *Suppose that $\frac{1}{2} \in R$. Then there is a degree 3 isomorphism $I^+(Y, E) \cong I^{+,tw}(Y, E)$ of $R[U]$ -modules and there is an exact triangle of $R[U]$ -modules*

$$\begin{array}{ccc} I^+(Y, E) & \xrightarrow{[3]} & I^-(Y, E) \\ & \swarrow [-4] \quad \searrow [0] & \\ & I^\infty(Y, E) & \end{array},$$

where the numbers specify the degrees of the maps.

We will now restrict ourselves to the cases of interest, namely,

$$(A, M) = (C_*^{gm}(\text{SO}(3)), \widetilde{CI}(Y, E)) \quad \text{or} \quad (A, M) = (\Lambda_R[u], DCI(Y, E)),$$

where M is equipped with the index filtration. Recall from Definition 2.8 and Definition 2.4 that M was obtained from a multicomplex $(M_{*,*}, \{\partial^r\}_{r=0}^5)$, and that the index filtration was defined to be the column filtration, i.e., $F_p M_n = \bigoplus_{s \leq p} M_{s,n-s}$. In both cases it holds true that $M_{s,t} = 0$ for all $t > 4$ and $t < 0$ (for $M = DCI(Y, E)$ this is true for $t \geq 4$). This implies that the filtration is degreewise finite and therefore exhaustive and complete Hausdorff. Explicitly,

$$(5.10) \quad F_p M_n = \begin{cases} M_n & \text{for all } p \geq n \\ 0 & \text{for all } p \leq n - 5 \end{cases}$$

The following lemma indicates why it is desirable to introduce the full completions of the various functors.

Lemma 5.45. *The filtration of $C_A^+(M)$, $F_p C_A^+(M) = C_A^+(F_p M)$, is degreewise bounded above, exhaustive and Hausdorff. The filtration of $C_A^-(M)$ is degreewise bounded below and complete Hausdorff. If $M \neq 0$ the filtration of $C_A^+(M)$ fails to be complete and the filtration of $C_A^-(M)$ fails to be exhaustive. Therefore,*

$$\hat{C}_A^+(M) = \lim_q C_A^+(M/F_q M) \quad \text{and} \quad \hat{C}_A^-(M) = \text{colim}_p C_A^-(F_p M).$$

Proof. We will use the underlying bigrading of the multicomplex $M_{*,*}$. As a graded module (not DG) we have $C_A^+(M) = B(M, A, R) \cong M \otimes BA$. Therefore, in degree n

$$(5.11) \quad C_A^+(M)_n = \bigoplus_{u+r=n} M_u \otimes BA_r = \bigoplus_{s+t+r=n} M_{s,t} \otimes BA_r,$$

and $F_p C_A^+(M)$ is obtained by imposing the condition $s \leq p$ in the above right hand direct sum. First, notice that A and hence BA are supported in nonnegative degrees. Then, as $M_{s,t} = 0$ for $t < 0$, it follows $F_p C_A^+(M)_n = C_A^+(M)_n$ for $p \geq n$. This shows that the filtration is degreewise bounded above and hence exhaustive. It also follows from the above description that the filtration is Hausdorff. On the other hand

$$(5.12) \quad \lim_p C_A^+(M/F_p M)_n \cong \lim_p \bigoplus_{\substack{s+t+r=n \\ s > p}} M_{s,t} \otimes BA_r \cong \prod_{s+t+r=n} M_{s,t} \otimes BA_r.$$

This does not coincide with the direct sum in equation (5.11) if $M \neq 0$, due to the periodicity $M_{s,t} \cong M_{s+8,t}$ for all s, t . Thus, $C_A^+(M)$ fails to be complete if $M \neq 0$.

Next, there is an isomorphism $C_A^-(M) = \text{Hom}_A(EA, M) \cong \text{Hom}(BA, M)$ of graded modules. In a fixed degree n , we find

$$(5.13) \quad C_A^-(M)_n = \text{Hom}(BA, M)_n = \prod_{s+t-r=n} \text{Hom}(BA_r, M_{s,t})$$

and as above $F_p C_A^-(M)$ is obtained by imposing the condition $s \leq p$ in the right hand product. For the factor $\text{Hom}(BA_r, M_{s,t})$ to be nonzero it is necessary that $r \geq 0$ and $0 \leq t \leq 4$, and therefore $s = n + r - t \geq n - 4$. Hence, $F_p C_A^-(M)_n = 0$ for $p < n - 4$ showing that the filtration is degreewise bounded below, and thus complete Hausdorff. Moreover, the above also implies that the right hand product in (5.13) subject to the condition $s \leq p$ is finite and therefore coincides with the direct sum. Consequently, $\text{colim}_p F_p C_A^-(M)$ is given by

$$(5.14) \quad \text{colim}_p \bigoplus_{\substack{s+t-r=n \\ s \leq p}} \text{Hom}(BA_r, M_{s,t}) = \bigoplus_{s+t-r=n} \text{Hom}(BA_r, M_{s,t}),$$

which does not coincide with the product in (5.13) as long as $M \neq 0$. This shows that the filtration fails to be exhaustive if $M \neq 0$.

The description of $\hat{C}_A^+(M)$ and $\hat{C}_A^-(M)$ given in the statement is now a consequence of Lemma 5.33 part (iii) and (iv). \square

Remark 5.46. Miller [Eis19, Appendix A] defines completed versions of C_A^\pm using what he names the completed bar construction and the finitely supported cobar construction. He defines the completed bar construction, $\hat{B}(M, A, N)$, to be the completion of $B(M, A, N)$ with respect to the filtration by internal degree, while the finitely supported cobar construction $c\hat{B}(N, A, M) \subset \text{Hom}_A(B(N, A, A), M)$ is the subcomplex consisting of those functionals that vanish on $B_{p,q}(N, A, A)$ for all sufficiently large p . The above lemma, in particular equations (5.12) and (5.14), verifies that our $\hat{C}_A^+(M)$ and $\hat{C}_A^-(M)$ coincide with Miller's $\hat{B}(M, A, R)$ and $c\hat{B}(R, A, M)$ for $M = \widetilde{CI}(Y, E)$ and $M = DCI(Y, E)$ with A as appropriate.

The final part of this section is devoted to extending the concrete models of Proposition 5.8 and Theorem 5.28 to the complex $DCI^\bullet(Y, E)$ for $\bullet \in \{+, -, \infty\}$. We will therefore assume that $\frac{1}{2} \in R$ throughout.

Theorem 5.47. *Let $A = \Lambda_R[u]$ and let $(D_{*,*}^\pm, \partial', \partial'')$ be the double complexes associated with $M = DCI(Y, E)$ in Proposition 5.8. Then there are isomorphisms of DG $R[U]$ -modules*

$$DCI^+(Y, E) \cong \text{Tot}^\Pi(D_{*,*}^+, \partial', \partial'') \quad \text{and} \quad DCI^-(Y, E) \cong \text{Tot}^\oplus(D_{*,*}^-, \partial', \partial'').$$

In particular, there are identifications

$$DCI^+(Y)_n \cong \prod_{p \geq 0} M_{n-4p} \quad \text{and} \quad DCI^-(Y)_n \cong \bigoplus_{p \geq 0} M_{n+4p}$$

for each $n \in \mathbb{Z}$. An element $x \in DCI^\pm(Y)_n$ can therefore be expressed by a sequence $x = (m_{n+4p})_p$ where we require $p \leq 0$ in the $+$ case and $p \geq 0$ and almost all $m_{n+4p} = 0$ in the $-$ case. The differential is in both cases given by

$$\partial x = (\partial_M m_{n+4p} - (-1)^n m_{n+4(p-1)} u)_p$$

and the action of $R[U]$ is determined by $U \cdot x = (m_{n+4(p+1)})_p$.

Proof. The isomorphism $C_A^+(M) \cong \text{Tot}^\oplus(D_{*,*}^+, \partial', \partial'')$ of Proposition 5.8 yields in degree n

$$C_A^+(M)_n \cong \bigoplus_{s \geq 0} M_{n-4s} = \bigoplus_{s \geq 0, t} M_{t, n-4s-t}$$

and $F_p C_A^+(M)$ is obtained by imposing the condition $t \leq p$ in the right hand sum. Now the calculation of the limit $\lim_p C_A^+(M)_n / F_p C_A^+(M)_n$ goes through as in equation (5.12) of the previous proof to show that $\hat{C}_A^+(M)_n \cong \prod_{s \geq 0} M_{n-4s}$.

The other case is analogous and the calculation of the relevant colimit proceeds as in equation (5.14) of the previous proof. \square

Note that the only difference between the above theorem and Proposition 5.8 is that we have interchanged the type of totalization we apply to the double complexes.

We also obtain the following explicit model computing $I^\infty(Y, E)$.

Theorem 5.48. *Let $A = \Lambda_R[u]$, let $(D_{*,*}^\pm, \partial', \partial'')$ and $(D_{*,*}^\infty, \partial', \partial'')$ be the double complexes associated with $M = DCI(Y, E)$ in the above theorem and Theorem 5.28, respectively. Let $\nu, \psi: \text{Tot}^\Pi(D^+, \partial', \partial'') \rightarrow \text{Tot}^\oplus(D^-, \partial', \partial'')$ of degree 3 and 0 respectively, be given degreewise by the compositions*

$$\prod_{s \geq 0} M_{n-4s} \xrightarrow{\pi} M_n \xrightarrow{(-1)^n u} M_{n+3} \xrightarrow{\iota} \bigoplus_{t \geq 0} M_{n+3+4t}$$

$$\prod_{s \geq 0} M_{n-4s} \xrightarrow{\pi} M_n \xrightarrow{\iota} \bigoplus_{t \geq 0} M_{n+4t},$$

where π and ι denote the projection and inclusion, respectively. Then ν is a chain map satisfying $\nu U - U \nu = \partial \psi - \psi \partial$. Define $\text{Cone}(\nu)$ with the $R[U]$ -module structure adjusted by the homotopy ψ as in Lemma 5.25. Then there is an isomorphism $R[U]$ -modules $\text{Tot}^{\Pi, \infty}(D_{*,*}^\infty, \partial', \partial'') \cong \text{Cone}(\nu)$ and a filtered quasi-isomorphism between this complex and $DCI^\infty(Y, E)$, which is $R[U]$ -linear up to homotopy. Moreover, this quasi-isomorphism is compatible with the corresponding exact triangles in homology.

Proof. By Proposition 5.43 there is a filtered quasi-isomorphism $\hat{f}_M: \hat{C}_A^+(M) \rightarrow \hat{C}_A^{+, tw}(M)$ of degree 3 and a filtered homotopy $\hat{s}_M: \hat{C}_A^+(M) \rightarrow \hat{C}_A^{+, tw}(M)$ such that $\hat{f}_M U - U \hat{f}_M = \partial \hat{s}_M - \hat{s}_M \partial$. Furthermore, there is an induced filtered quasi-isomorphism $\text{Cone}(\hat{N}_M \circ \hat{f}_M) \rightarrow \hat{C}_A^\infty(M)$, which is $R[U]$ -linear up to homotopy when the cone is given the adjusted $R[U]$ -module structure determined by the homotopy $\hat{t}_M = \hat{N}_M \circ \hat{s}_M$. In terms of the explicit models of the above theorem $\hat{N}_M \circ \hat{f}_M$ and $\hat{N}_M \circ \hat{s}_M$ are precisely ν and ψ as defined in the statement. The verification of this is done just as in Lemma 5.24. The fact that $\text{Cone}(\nu) \cong \text{Tot}^{\Pi, \infty}(D_{*,*}^\infty, \partial', \partial'')$ is proved just as in Theorem 5.28. \square

Remark 5.49. The proofs of Theorem 5.47 and Theorem 5.48 go through without any essential change with $M = \widetilde{CI}(Y, E)$, regarded as a $\Lambda_R[u]$ -module, in place of $M = DCI(Y, E)$.

We may extract the following noteworthy consequences, showing in particular that the framed homology groups $\tilde{I}(Y, E) := H(DCI(Y, E)) \cong H(\widetilde{CI}(Y, E))$ may be recovered from the $R[U]$ -action on $I^\pm(Y, E)$ up to extension.

Corollary 5.50. *There are long exact sequences*

$$\tilde{I}_n(Y, E) \longrightarrow I^+(Y, E)_n \xrightarrow{U} I^+(Y, E)_{n-4} \longrightarrow \tilde{I}(Y, E)_{n-1}$$

$$\tilde{I}(Y, E)_{n-3} \longrightarrow I^-(Y, E) \xrightarrow{U} I^-(Y, E)_{n-4} \longrightarrow \tilde{I}(Y, E)_{n-4}.$$

Moreover, $U : I^\infty(Y, E) \rightarrow I^\infty(Y, E)$ is an isomorphism.

Proof. Let $M = DCI(Y, E)$ and $A = \Lambda_R[u]$. In terms of the explicit models for $DCI^\pm(Y, E) = \hat{C}_A^\pm(M)$ the action of $U \in R[U]$ are in degree n given by the natural shifts

$$U : \prod_{s \geq 0} M_{n-4s} \rightarrow \prod_{s \geq 0} M_{n-4-4s} \quad \text{and} \quad U : \bigoplus_{s \geq 0} M_{n+4s} \rightarrow \bigoplus_{s \geq 0} M_{n-4+4s},$$

respectively. Explicitly, $(x_s)_{s \geq 0} \mapsto (x_{s+1})_{s \geq 0}$ in the first case and $(y_s)_{s \geq 0} \mapsto (0, y_0, y_1, \dots)$ in the second. From this we deduce that the first map is surjective with kernel $D_{0,*}^+ \cong M$ and the second is injective with cokernel isomorphic to $D_{0,*}^- \cong M$. There are therefore short exact sequences of chain complexes

$$0 \longrightarrow M \longrightarrow D_*^+ \xrightarrow{U} D_*^+[4] \longrightarrow 0$$

$$0 \longrightarrow D_*^-[-4] \xrightarrow{U} D_*^- \longrightarrow M \longrightarrow 0$$

giving rise to the stated long exact sequences in homology. The final statement is a consequence of the fact that the shift

$$U : D_n^\infty = \prod_{s \rightarrow \infty} M_{n-4s} \rightarrow \prod_{s \rightarrow \infty} M_{n-4-4s} = D_{n-4}^\infty$$

is an isomorphism in each degree n and therefore also an isomorphism upon passage to homology. \square

Remark 5.51. The fact that $U : I^\infty(Y, E) \rightarrow I^\infty(Y, E)$ is an isomorphism when $\frac{1}{2} \in R$ was first proved using a spectral sequence argument in [Eis19].

5.5. The Index Spectral Sequences. For $\bullet \in \{+, -, (+, tw), \infty\}$ the complexes $CI^\bullet(Y, E)$ and $DCI^\bullet(Y, E)$ introduced in Definition 5.36 come equipped with filtrations and therefore give rise to spectral sequences. These will be called index spectral sequences as the filtrations are induced by the index filtrations of $\widetilde{CI}(Y, E)$ and $DCI(Y, E)$. The purpose of this section is to recall some theory on spectral sequences and then to establish the basic properties of the index spectral sequences. We will mainly follow Boardman's paper [Boa99], but we will stick to homological notation. All of this will be very useful for our calculations in the next section.

Recall that an (unrolled) exact couple (A, E, i, j, k) consists of two sequences of graded modules $(A_s)_s$ and $(E_s)_s$, indexed over $s \in \mathbb{Z}$, and homomorphisms $i = (i_s : A_{s-1} \rightarrow A_s)_s$, $j = (j_s : A_s \rightarrow E_s)_s$ and $k = (k_s : E_s \rightarrow A_{s-1})_s$ such that the sequence

$$A_{s-1} \xrightarrow{i_s} A_s \xrightarrow{j_s} E_s \xrightarrow{k_s} A_{s-1} \xrightarrow{i_s} A_s$$

is exact for each s . We will generally omit the lower index on the maps, letting the source and target specify the map in question. In our situation the degrees of (i, j, k) will be $(0, 0, -1)$. For each $r \geq 1$ and s set

$$\begin{aligned} Z_s^r &:= k^{-1} \operatorname{Im}[i^{(r-1)} : A_{s-r} \rightarrow A_{s-1}] \\ B_s^r &:= j \operatorname{Ker}[i^{(r-1)} : A_s \rightarrow A_{s+r-1}] \\ E_s^r &:= Z_s^r / B_s^r, \end{aligned}$$

where $i^{(r)}$ denotes the r -fold composition $i \circ \dots \circ i$ for $r \geq 1$ and $i^{(0)}$ is the identity. Furthermore, put $Z_s^\infty := \bigcap_r Z_s^r$, $B_s^\infty := \bigcup_r B_s^r$ and $E_s^\infty := Z_s^\infty / B_s^\infty$. For the later convergence theory it is also important to introduce the derived limit $RE_s^\infty := \operatorname{Rlim}_s Z_s^r$. There are differentials $d^r : E_s^r \rightarrow E_{s-r}^r$ for each s and $r \geq 1$, defined in the following way. Let $x \in Z_s^r$ represent the class $[x] \in E_s^r$. Choose $y \in A_{s-r}$ such that $i^{(r-1)}(y) = k(x)$. Then $d^r([x]) = [j(y)] \in E_{s-r}^r$. The situation is illustrated in the following diagram

$$(5.15) \quad \begin{array}{ccc} & y \in A_{s-r} & \xrightarrow{i^{(r-1)}} & k(x) \in A_{s-1} \\ & \swarrow j & & \nwarrow k \\ j(y) \in E_{s-r}^r & & & x \in E_s^r. \end{array}$$

One may then show that d^r is well-defined, that $d^r \circ d^r = 0$ and that

$$E_s^{r+1} \cong \operatorname{Ker}(d^r : E_s^r \rightarrow E_{s-r}^r) / \operatorname{Im}(d^r : E_{s+r}^r \rightarrow E_s^r)$$

for each $r \geq 1$ and s . In other words, $(E^r, d^r)_{r \geq 1}$ is a spectral sequence. We call it the spectral sequence associated with the exact couple (A, E, i, j, k) . Every exact couple we will meet is covered by the following example (see [Boa99, Section 9]).

Example 5.52. Let C be a DG module equipped with an increasing filtration $\{F_s C\}_s$ by DG submodules. For each $s \in \mathbb{Z}$ let $A_s = H(F_s C)$ and $E_s = H(F_s C / F_{s-1} C)$. Define $i : A_{s-1} \rightarrow A_s$, $j : A_s \rightarrow E_s$ and $k : E_s \rightarrow A_{s-1}$ to be the maps induced by the short exact sequence

$$0 \longrightarrow F_{s-1} C \longrightarrow F_s C \longrightarrow F_s C / F_{s-1} C \longrightarrow 0$$

upon passage to homology. In particular, k is the connecting homomorphism of degree -1 . Then (A, E, i, j, k) is an exact couple. The spectral sequence associated with this exact couple is the spectral sequence mentioned in (5.9). If B is a DG algebra and C is a B -module filtered by B -submodules, then A_s, E_s carry the structure of graded $H(B)$ -modules, and the maps i, j, k are $H(B)$ -linear. Moreover, this structure carries in a natural way over to the whole spectral sequence.

Returning to the situation of an exact couple (A, E, i, j, k) put

$$(5.16) \quad A_\infty := \operatorname{colim}_s A_s, \quad A_{-\infty} := \lim_s A_s \quad \text{and} \quad RA_{-\infty} := \operatorname{Rlim}_s A_s.$$

For us the relevant target for the spectral sequence is the colimit A_∞ , filtered by $F_s A_\infty := \operatorname{Im}(A_s \rightarrow A_\infty)$. For each $s \in \mathbb{Z}$ there is a natural inclusion $\phi_s : F_s A_\infty / F_{s-1} A_\infty \rightarrow E_s^\infty$ given by the composition

$$(5.17) \quad F_s A_\infty / F_{s-1} A_\infty \xrightarrow{\cong} \operatorname{Im} j / B_s^\infty \longrightarrow Z_s^\infty / B_s^\infty = E_s^\infty.$$

Here, the first isomorphism is given by sending $[x] \in F_s A_\infty / F_{s-1} A_\infty$, represented by $x \in F_s A_\infty$, to $[j(y)] \in \operatorname{Im} j / B_s^\infty$, where $y \in A_s$ is a lift of x along the surjection $A_s \rightarrow F_s A_\infty$, and the second map is induced by the inclusion $\operatorname{Im} j = \operatorname{Ker} k \subset Z_s^\infty$ (see [Boa99, Lemma 5.6] and its proof).

Definition 5.53. The spectral sequence $(E^r, d^r)_{r \geq 1}$ associated with the exact couple (A, E, i, j, k) converges strongly to the colimit A_∞ if the filtration $F_s A_\infty = \text{Im}(A_s \rightarrow A_\infty)$ of A_∞ is exhaustive and complete Hausdorff and the map $\phi_s: F_s A_\infty / F_{s-1} A_\infty \rightarrow E_s^\infty$ is an isomorphism for each $s \in \mathbb{Z}$.

Strong convergence is the ideal form of convergence. In theory, if we can calculate E^∞ we would know the subquotients $F_s A_\infty / F_{s-1} A_\infty$ and provided we can solve the extension problems to determine $F_s A_\infty / F_t A_\infty$ for each pair $s > t$, we can recover the target group as

$$A_\infty \cong \lim_s \text{colim}_t F_t A_\infty / F_s A_\infty.$$

Definition 5.54. [Boa99, Definition 5.10] Let (A, E, i, j, k) be an exact couple with associated spectral sequence $(E^r, d^r)_{r \geq 1}$. If $RA_{-\infty} = 0$ and $A_{-\infty} = 0$ (see (5.16)), then the spectral sequence is said to converge conditionally to the colimit A_∞ .

In our situation of a filtered DG module and the corresponding exact couple described in Example 5.52 we have the following result.

Theorem 5.55. [Boa99, Theorem 9.2] Let C be a DG module equipped with an increasing filtration $F_s C$ by DG submodules. Let (A_s, E_s, i, j, k) be the associated exact couple with corresponding spectral sequence $(E^r, d^r)_{r \geq 1}$. If the filtration of C is exhaustive and complete Hausdorff, then the spectral sequence $(E^r, d^r)_{r \geq 1}$ converges conditionally to the colimit $A_\infty = H(C)$.

In the presence of conditional convergence there are two remaining obstructions to achieving strong convergence; namely, the vanishing of the group RE^∞ and another group W . We refer to [Boa99, Lemma 8.5] for the general definition of W , but we note that if the exact couple is constructed from a filtered complex, then W is isomorphic to the kernel of the canonical interchange map (see [HR19])

$$\kappa: \text{colim}_p \lim_q H(F_p C / F_q C) \rightarrow \lim_q \text{colim}_p H(F_p C / F_q C).$$

We then have the following convergence theorem.

Theorem 5.56. [Boa99, Theorem 8.2] Let (A, E, i, j, k) be an exact couple and assume that the associated spectral sequence $(E^r, d^r)_{r \geq 1}$ converges conditionally to the colimit A_∞ . Then if $W = 0$ and $RE^\infty = 0$ the spectral converges strongly to the colimit A_∞ .

The following result gives a few standard criteria for the vanishing of RE^∞ and W in the situation of a filtered complex.

Proposition 5.57. Let C be a filtered complex with associated exact couple (A, E, i, j, k) and spectral sequence $(E^r, d^r)_{r \geq 1}$.

- (i) If the filtration of C is degreewise bounded below, then $RE^\infty = 0$ and $W = 0$.
- (ii) If the filtration of C is degreewise bounded above then $W = 0$.
- (iii) If the spectral sequence degenerates at some finite stage; that is, $E^{r_0} = E^{r_0+1} = \dots = E^\infty$ for some finite $r_0 \geq 1$, then $RE^\infty = 0$ and $W = 0$.

We may now apply this theory to the filtered complexes $CI^\bullet(Y, E)$ for $\bullet \in \{+, -, (+, tw), \infty\}$. This is a good point to explain that the mod 8 periodicity of $\widetilde{CI}(Y, E)$ carries over to the complexes $CI^\bullet(Y, E)$. Let $M = \widetilde{CI}(Y, E)$ and let $A = C_*^{gm}(\text{SO}(3))$ or $A = \Lambda_R[u]$. Then the periodicity may be expressed as an isomorphism $M \cong M[8]$ of A -modules. Moreover, this isomorphism is compatible with the filtration in the sense that $F_{p+8} M \cong (F_p M)[8]$ for all $p \in \mathbb{Z}$. There is therefore an induced isomorphism

$$\begin{aligned} CI^\bullet(Y, E) &= \lim_q \text{colim}_p C_A^\bullet(F_{p+8} M / F_q M) \\ &\cong \lim_q \text{colim}_p C_A^\bullet(F_p M / F_q M)[8] = CI^\bullet(Y, E)[8]. \end{aligned}$$

This isomorphism is seen to be filtered in the same sense: $F_p CI^\bullet(Y, E)_n \cong F_{p+8} CI^\bullet(Y, E)_{n+8}$ for all p and n .

Theorem 5.58. *Let $A = C_*^{gm}(\mathrm{SO}(3))$. There is a conditionally convergent spectral sequence of $H_A^-(R)$ -modules*

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_{\mathrm{SO}(3)}^\bullet(\alpha)_t \implies I^\bullet(Y, E)_{s+t}$$

for each $\bullet \in \{+, -, (+, tw), \infty\}$, where

$$H_{\mathrm{SO}(3)}^\bullet(\alpha) := H_A^\bullet(C_*^{gm}(\alpha))$$

for each critical orbit $\alpha \in \mathcal{C}$. Moreover, the spectral sequences are periodic in the sense that there are isomorphisms $E_{s,t}^r \cong E_{s+8,t}^r$ for all s, t, r . These isomorphisms commute with the differentials and the $H_A^-(R)$ -action and they are compatible with the target in the sense that

$$\begin{array}{ccc} \overline{F}_s I_{s+t}^\bullet & \xrightarrow{\cong} & \overline{F}_{s+8} I_{s+t+8}^\bullet \\ \downarrow & & \downarrow \\ E_{s,t}^\infty & \xrightarrow{\cong} & E_{s+8,t}^\infty \end{array}$$

commutes for all s, t .

The spectral sequence for $I^-(Y, E)$ converges strongly, while the spectral sequence for $I^+(Y, E)$ converges strongly provided $RE^\infty = 0$.

Proof. Let $M = \widetilde{CI}(Y, E)$ equipped with the index filtration. For each $\bullet \in \{+, -, (+, tw), \infty\}$ the complex $CI^\bullet(Y, E) = \hat{C}_A^\bullet(M)$ carries by construction an exhaustive and complete Hausdorff filtration with minimal filtration quotients

$$\overline{F}_p \hat{C}_A^\bullet(M) \cong C_A^\bullet(\overline{F}_p M).$$

There is therefore by Theorem 5.55 a conditionally convergent spectral sequence

$$E_{s,t}^1 = H_A^\bullet(\overline{F}_s M)_{s+t} \implies \hat{H}_A^\bullet(M)_{s+t} = I^\bullet(Y, E)_{s+t}.$$

From the definition of $\widetilde{CI}(Y, E)$ (Definition 2.4) we may identify

$$\overline{F}_s M_{s+t} = \bigoplus_{j(\alpha) \equiv s} C_t^{gm}(\alpha)$$

so that $E_{s,t}^1 \cong \bigoplus_{j(\alpha) \equiv s} H_{\mathrm{SO}(3)}^\bullet(\alpha)_t$ in the notation of the statement.

Next, the periodicity isomorphisms $F_p \hat{C}_A^\bullet(M)_n \cong F_{p+8} \hat{C}_A^\bullet(M)_{n+8}$, compatible with the differentials and the $C_A^-(R)$ -structure, induce a morphism of the associated exact couple. That is, there are isomorphisms $A_s \cong A_{s+8}$, $E_s \cong E_{s+8}$ each s , compatible with the structure maps. It is then a straightforward exercise to verify that this gives rise to the periodicity in the spectral sequence, compatible with the target, as stated.

Finally, for the convergence we know by Lemma 5.45 that the filtration of $\hat{C}_A^+(M)$ is degree-wise bounded above and that the filtration of $\hat{C}_A^-(M)$ is degree-wise bounded below. This implies by Proposition 5.57 and Theorem 5.56 that the spectral sequence associated with $I^-(Y, E)$ converges strongly and that the spectral sequence associated with $I^+(Y, E)$ converges strongly provided $RE^\infty = 0$. \square

Provided $\frac{1}{2} \in R$, these spectral sequences can equally well be constructed from $M = DCI(Y, E)$ equipped with the index filtration and $A = \Lambda_R[u]$. This is what we will assume when we calculate

these groups for binary polyhedral spaces in the next section. Furthermore, in this setting the calculation of the groups

$$H_{\mathrm{SO}(3)}^\bullet(C_*^{gm}(\alpha)) \cong H_{\Lambda_R[u]}^\bullet(H_*(\alpha))$$

for each $\bullet \in \{+, -, (+, tw), \infty\}$, are contained in Lemma 5.7, Corollary 5.9 and Corollary 5.30. From these calculations it becomes clear that the spectral sequences for I^+ and I^- are contained in the upper and lower half-planes respectively, while the spectral sequence for I^∞ is a whole-plane spectral sequence.

6. CALCULATIONS FOR BINARY POLYHEDRAL SPACES

We will from this point require that $2 \in R$ is invertible. The universal example to have in mind is $R = \mathbb{Z}[\frac{1}{2}]$. Our main aim is to give explicit calculations of the $H_{\Lambda_R[u]}^-(R) = R[U]$ -modules

$$I^+(\overline{Y}_\Gamma, E), \quad I^-(\overline{Y}_\Gamma, E) \quad \text{and} \quad I^\infty(\overline{Y}_\Gamma, E)$$

for each finite subgroup $\Gamma \subset \mathrm{SU}(2)$, where $E \rightarrow \overline{Y}_\Gamma$ is the trivial $\mathrm{SU}(2)$ -bundle. To simplify the notation slightly we will omit the reference to the bundle in the notation from this point. In the final part we also include the calculations of $I^\bullet(Y_\Gamma)$ for $\bullet \in \{+, -, \infty\}$, omitting some details.

The two results Theorem 4.21 and Proposition 4.25 give a concrete description of the Donaldson model $\mathrm{DCI}(\overline{Y}_\Gamma)$ in all cases. The key tools needed in the calculations are the index spectral sequence in combination with the explicit models for DCI^\pm of Theorem 5.8. In the I^+ case, we will see that the index spectral sequence immediately degenerates, and we are left with solving an extension problem in the category of $R[U]$ -modules. In the I^- case there are a number of differentials in the index spectral sequence, but we will in each case verify that the spectral sequence degenerates on some finite page E^{r_0} . The extension problems in this case are neatly solved by observing that E_s^∞ is a free $R[U]$ -module for all s .

In [Eis19, Corollary 8.7] Miller calculates $I^\infty(Y)$ for all rational homology spheres. This, of course, includes \overline{Y}_Γ , so our calculation of this group should only be regarded as a verification of his result. In addition, we observe that the norm map vanishes in homology so that the exact triangle relating the three groups splits up into short exact sequences.

Throughout this section we let $A = \Lambda_R[u]$ with $|u| = 3$. Given a finite subgroup $\Gamma \subset \mathrm{SU}(2)$ we will always write \mathcal{C} for the set of critical orbits, or equivalently, the set of 1-dimensional quaternionic representations of Γ . Recall that for each $\alpha \in \mathcal{C}$ we have fixed generators $b_\alpha \in H_0(\alpha)$, $t_\alpha = b_\alpha \cdot u \in H_3(\alpha)$ if α is irreducible and $t_\alpha \in H_2(\alpha)$ if α is reducible.

6.1. The Case I^+ . In this section we will calculate $I^+(\overline{Y}_\Gamma)$ as an $R[U]$ -module for all finite subgroups $\Gamma \subset \mathrm{SU}(2)$. Even though our arguments do not explicitly rely on the index spectral sequence, we note that the motivation for the upcoming arguments is the following observation.

Lemma 6.1. *Let $M = \mathrm{DCI}(\overline{Y}_\Gamma)$ for a finite subgroup $\Gamma \subset \mathrm{SU}(2)$. Then the index spectral sequence*

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_A^+(\alpha)_t \implies I^+(\overline{Y}_\Gamma)_{s+t}$$

degenerates at the E^1 page, that is, all differentials are trivial and $E^1 = E^\infty$.

Proof. From the calculations in Lemma 5.7 and Corollary 5.9 we see that $H_A^+(\alpha)$ vanishes in odd degrees for all types of orbits α . Then, as $j(\alpha) \equiv 0 \pmod{4}$ for all $\alpha \in \mathcal{C}$, it follows that $E_{s,t}^1 = 0$ if s or t is odd. Since the differential d^r has bidegree $(-r, r-1)$ we conclude that $d^r = 0$ for all $r \geq 1$ as required. \square

From this one may determine $I^+(\overline{Y}_\Gamma)$ as an R -module. However, the structure as an $R[U]$ -module is more subtle and this is the reason we spell out our argument more directly in terms of the filtration of the chain complex $DCI^+(\overline{Y}_\Gamma)$.

Let $\Gamma \subset \mathrm{SU}(2)$ be a finite subgroup and let $M = DCI(\overline{Y}_\Gamma)$. Let $\mathcal{C}^{irr} \subset \mathcal{C}$ denote the subset of free critical orbits or equivalently irreducible flat connections. Define $M^{irr} \subset M$ to be the submodule generated by \mathcal{C}^{irr} , i.e.,

$$M_{s,t}^{irr} = \bigoplus_{\substack{\alpha \in \mathcal{C}^{irr} \\ j(\alpha) \equiv s}} H_t(\alpha) \quad \text{and} \quad M_n^{irr} = \bigoplus_{s+t=n} M_{s,t}^{irr}.$$

The following lemma is the key tool in resolving our extension problems in the category of $R[U]$ -modules.

Lemma 6.2. *There is an R -linear map $\psi: M \rightarrow M$ of degree -4 satisfying*

- (i) $\psi(x) \cdot u = (-1)^{|x|} \partial_M x$ for all $x \in M$,
- (ii) $\mathrm{Im}(\psi) \subset M^{irr}$,
- (iii) $\psi(\mathrm{Im}(u)) = 0$ and
- (iv) $\psi(\mathrm{Im} \partial_M) = 0$.

Proof. By Theorem 4.21 the only possibly nontrivial differentials in M are $\partial_M: M_{4s} \rightarrow M_{4s-1}$ for $s \in \mathbb{Z}$. The same result also shows that $u: M_{4s-4}^{irr} \rightarrow M_{4s-1}^{irr} = M_{4s-1}$ is an isomorphism for each s . We may therefore define $\psi: M_{4s} \rightarrow M_{4s-4}^{irr} \subset M_{4s-4}$ by $u^{-1} \circ \partial_M$ and set $\psi = 0: M_t \rightarrow M_{t-4}$ in all other degrees. The relation $\psi(x) \cdot u = (-1)^{|x|} \partial_M x$ is then satisfied for all $x \in M$ and by construction $\mathrm{Im}(\psi) \subset M^{irr}$. For (iii) and (iv) it is then sufficient to note that $\mathrm{Im}(u)$ and $\mathrm{Im}(\partial_M) \subset M^{irr}$ are supported in the degrees M_{4s-1} for $s \in \mathbb{Z}$. \square

Remark 6.3. It is not difficult to show that the conditions (i) and (ii) determine the map ψ uniquely, but we will not use this explicitly.

Recall from Corollary 5.47 that for $M = DCI(\overline{Y}_\Gamma)$ there is an identification $DCI^+(\overline{Y}_\Gamma)_n \cong \prod_{p \geq 0} M_{n-4p}$ for each n . Moreover, the differential is given by

$$\partial(m_{n-4p})_{p \geq 0} = (\partial_M m_{n-4p} - (-1)^n m_{n-4(p+1)} u)_{p \geq 0}.$$

The point of the map ψ should now be clear; given $m \in M_n$ the element $(\psi^p(x))_{p \geq 0} \in \prod_{p \geq 0} M_{n-4p}$ gives an explicit extension of m to a cycle in $DCI^+(Y)$.

Using the description of the differentials in Proposition 4.21 we note that ψ is given on a generator $b_\alpha \in H_0(\alpha) \subset M_{4p}$ by

$$(6.1) \quad \psi(b_\alpha) = u^{-1} \left(\sum_{\beta \in \mathcal{C}^{irr}} n_{\beta\alpha} t_\beta \right) = \sum_{\beta \in \mathcal{C}^{irr}} n_{\beta\alpha} b_\beta$$

(the integers $n_{\beta\alpha}$ were defined in Definition 4.20).

Recall from Lemma 5.45 that

$$DCI^+(\overline{Y}_\Gamma) = \hat{C}_A^+(M) = \lim_p C_A^+(M/F_p M),$$

filtered by $F_p \hat{C}_A^+(M) = \hat{C}_A^+(F_p M) = \lim_{q < p} C_A^+(F_p M/F_q M)$. The concrete model given in Corollary 5.47 applies equally well to these subcomplexes. Recall that we use the notation $\overline{F}_p M = F_p M/F_{p-1} M$.

Lemma 6.4. *Let $M = DCI(\overline{Y}_\Gamma)$ be equipped with the index filtration. Then $F_{4p}M = F_{4p+t}M$ for all $p \in \mathbb{Z}$ and $0 \leq t \leq 3$. Moreover, there are identifications for $s \in \mathbb{Z}$ and $0 \leq t \leq 3$*

$$(6.2) \quad C_A^+(\overline{F}_{4p}M)_{4s+t} \cong \begin{cases} M_{4p+t} & \text{for } s \geq p \\ 0 & \text{for } s < p \end{cases}$$

$$\hat{C}_A^+(F_{4p}M)_{4s+t} \cong \begin{cases} \prod_{q \geq 0} M_{4(p-q)+t} & \text{for } s \geq p \\ \prod_{q \geq 0} M_{4(s-q)+t} & \text{for } s < p. \end{cases}$$

Define $\zeta: C_A^+(\overline{F}_{4p}M) \rightarrow \hat{C}_A^+(F_{4p}M)$ for $x \in M_{4p+t} \cong C^+(\overline{F}_{4p}M)_{4s+t}$, $s \geq p$, by

$$\zeta(x) = (\psi^q(x))_{q \geq 0} \in \prod_{q \geq 0} M_{4(p-q)+t} = \hat{C}_A^+(F_{4p}M),$$

where ψ is the map of Lemma 6.2, and otherwise to be zero. Then ζ is a chain map splitting the exact sequence of DG R -modules (i.e., $\pi \circ \zeta = 1$)

$$0 \longrightarrow \hat{C}_A^+(F_{4p-1}M) \xrightarrow{i} \hat{C}_A^+(F_{4p}M) \xleftarrow{\zeta} C_A^+(\overline{F}_{4p}M) \xrightarrow{\pi} 0.$$

Proof. The fact that $F_{4p+t}M = F_{4p}M$ for $0 \leq t \leq 3$ follows from the structure theorem 4.21. From the same result one concludes that $F_{4p}M_n = M_n$ for $n \leq 4p + 3$ and $F_{4p}M_n = 0$ for $n \geq 4(p + 1)$. Therefore, $\overline{F}_{4p}M_n$ coincides with M_n for $4p \leq n \leq 4p + 3$ and vanishes otherwise. Using this, the given degreewise formulas for $C_A^+(\overline{F}_{4p}M)$ and $\hat{C}_A^+(F_{4p}M)$ follow from Proposition 5.8 and Corollary 5.47, respectively.

Next we have to verify that ζ is a chain map. The differential in $C_A^+(\overline{F}_{4p}M)$ is given by $u: M_{4p} \rightarrow M_{4p+3}$ in degree $4s$ for $s > p$ and is otherwise 0. The differential in $\hat{C}_A^+(F_{4p}M)$ is given in degree $4s + t$ by

$$\partial((m_{4(s-q)+t})_{q \geq 0}) = (\partial_M m_{4(s-q)+t} - (-1)^t m_{4(s-q-1)+t} u)_{q \geq 0},$$

where if $s > p$ it is assumed that $m_{4(s-q)+t} = 0$ for $q \leq s - p$. In view of this formula and the fact that $\psi(x) \cdot u = (-1)^{|x|} \partial_M x$ for all $x \in M$ by Lemma 6.2, we see that for $x \in M_{4p+t} = C_A^+(\overline{F}_{4p}M)_{4s+t}$, $s \geq p$,

$$\partial \circ \zeta(x) = \partial(\psi^q x)_{q \geq 0} = \begin{cases} 0 & \text{if } (s, t) = (p, 0) \text{ or } 1 \leq t \leq 3 \\ (x \cdot u, 0, 0, \dots) & \text{if } s > p, t = 0 \end{cases}$$

In the first case we also have $\partial x = 0$ and hence $\zeta \circ \partial(x) = 0$, while in the second case we have $\partial x = x \cdot u$ so that $\zeta \circ \partial(x) = (\psi^q(x \cdot u))_{q \geq 0} = (x \cdot u, 0, 0, \dots)$ as ψ vanishes on $\text{Im } u$ by Lemma 6.2. In all other degrees ζ vanishes, so we may conclude from the above that ζ is a chain map.

Finally, the projection $\pi: \hat{C}_A^+(F_{4p}M) \rightarrow C_A^+(\overline{F}_{4p}M)$ is in terms of the formulas of (6.2) given by the projection $\prod_{q \geq 0} M_{4(p-q)+t} \rightarrow M_{4p+t}$ in degree $4s + t$ for $s \geq p$ and otherwise 0. From this description it is clear that $\pi \circ \zeta = 1$ and the proof is complete. \square

We need one final lemma describing the interaction between the map ζ and the action of $R[U]$.

Lemma 6.5. *In the situation of the above lemma the map $\zeta: C_A^+(\overline{F}_{4p}M) \rightarrow \hat{C}_A^+(F_{4p}M)$ satisfies $\zeta(Ux) = U\zeta(x)$ for each x with $|x| \neq 4p$. Moreover, in degree $4p$ we have the following commutative diagram*

$$\begin{array}{ccccc} C_A^+(\overline{F}_{4p}M)_{4p} & \xrightarrow{\zeta} & \hat{C}_A^+(F_{4p}M)_{4p} & \xrightarrow{U} & \hat{C}_A^+(F_{4p}M)_{4(p-1)} \\ \downarrow \psi & & & & \downarrow = \\ C_A^+(\overline{F}_{4(p-1)}M)_{4(p-1)} & \xrightarrow{\zeta} & \hat{C}_A^+(F_{4(p-1)}M)_{4(p-1)} & & \end{array}$$

Proof. In degree n for $n < 4p$ the map ζ vanishes, so the first assertion is clear in this case. In degree $4s + t$ with $s > p$ and $0 \leq t \leq 3$ we have, using the formulas of Lemma 6.4,

$$C_A^+(\overline{F}_{4p}M)_{4s+t} = M_{4p+t} \quad \text{and} \quad \hat{C}_A^+(F_{4p}M)_{4s+t} = \prod_{q \geq 0} M_{4(p-q)+t}.$$

It then follows from Proposition 5.8 and Corollary 5.47 that the U action is given by the identity in these degrees. Therefore, $U \circ \zeta = \zeta \circ U$ in this case as well.

For $x \in M_{4p+t} = C_A^+(\overline{F}_{4p}M)_{4p+t}$, where $0 \leq t \leq 3$, we have $Ux = 0$, while

$$U(\zeta(x)) = U((\psi^q x)_{q \geq 0}) = (\psi^{q+1} x)_{q \geq 0} = (\psi^q(\psi x))_{q \geq 0} = \zeta(\psi(x)),$$

where we regard $\psi x \in M_{4(p-1)+t} = C_A^+(\overline{F}_{4(p-1)}M)_{4(p-1)+t}$. This shows that the given diagram commutes in degree $4p + t$ for $0 \leq t \leq 3$. However, in the proof of Lemma 6.2 it was shown that $\psi: M_{4p+t} \rightarrow M_{4(p-1)+t}$ vanishes for $1 \leq t \leq 3$. Therefore $\zeta \circ U = U \circ \zeta$ in degree $4p + t$ for $1 \leq t \leq 3$ as well. This completes the proof. \square

To describe $I^+(\overline{Y}_\Gamma)$ and later $I^-(\overline{Y}_\Gamma)$, it is convenient to introduce the following definition.

Definition 6.6. For an $R[U]$ -module X define the mod 8 periodic $R[U]$ -modules $X^{\Pi,8}$ and $X^{\oplus,8}$ degree-wise by

$$X_n^{\Pi,8} = \prod_{s \in \mathbb{Z}} X_{n+8s} \quad \text{and} \quad X_n^{\oplus,8} = \bigoplus_{s \in \mathbb{Z}} X_{n+8s}.$$

The maps $U: X_n^{\Pi,8} \rightarrow X_{n-4}^{\Pi,8}$ and $U: X_n^{\oplus,8} \rightarrow X_{n-4}^{\oplus,8}$ are defined to be the product and direct sum over the maps $U: X_{n+8s} \rightarrow X_{n-4+8s}$ for $s \in \mathbb{Z}$, respectively.

Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup. Given $\alpha \in \mathcal{C}$ we will use the shorthand notation $H_A^+(\alpha) := H_A^+(H_*(\alpha))$. Write $\mathcal{C} = \mathcal{C}^{irr} \cup \mathcal{C}^{red} \cup \mathcal{C}^{f.red}$ for the decomposition of the critical orbits into irreducible, reducibles and fully reducibles. For each $\eta \in \mathcal{C}^{f.red}$, $\lambda \in \mathcal{C}^{red}$ and $\alpha \in \mathcal{C}^{irr}$ introduce variables V_η , W_λ and a generator g_α so that

$$H_A^+(\eta) = R[V_\eta], \quad H_A^+(\lambda) = R[W_\lambda] \quad \text{and} \quad H_A^+(\alpha) = R \cdot g_\alpha,$$

where $|V_\eta| = 4$, $|W_\lambda| = 2$ and $|g_\alpha| = 0$. This is justified by the calculations of Lemma 5.7 and Corollary 5.9. In the following we regard the relative grading $j: \mathcal{C} \rightarrow \mathbb{Z}/8$ as taking values in $\{0, 4\}$.

Theorem 6.7. *The positive equivariant instanton Floer homology $I^+(\overline{Y}_\Gamma)$ associated with the trivial $\text{SU}(2)$ -bundle over \overline{Y}_Γ is given by*

$$\left[\left(\bigoplus_{\alpha \in \mathcal{C}^{irr}} R \cdot g_\alpha[j(\alpha)] \right) \oplus \left(\bigoplus_{\lambda \in \mathcal{C}^{red}} R[W_\lambda][j(\lambda)] \right) \oplus \left(\bigoplus_{\eta \in \mathcal{C}^{f.red}} R[V_\eta][j(\eta)] \right) \right]^{\Pi,8}.$$

The $R[U]$ -module structure is determined by

$$\begin{aligned} U \cdot V_\eta^p &= \begin{cases} V_\eta^{p-1} & \text{for } p > 0 \\ \sum_{\rho \in \mathcal{C}^{irr}} n_{\rho\eta} g_\rho & \text{for } p = 0 \end{cases} \\ U \cdot W_\lambda^p &= \begin{cases} W_\lambda^{p-2} & \text{for } p > 0 \\ \sum_{\rho \in \mathcal{C}^{irr}} n_{\rho\lambda} g_\rho & \text{for } p = 0 \end{cases} \\ U \cdot g_\alpha &= \sum_{\rho \in \mathcal{C}^{irr}} n_{\rho\alpha} g_\rho, \end{aligned}$$

where we interpret $W_\beta^{-1} = 0$ and the integers $n_{\rho_1\rho_2}$ are given in Proposition 4.25 for each pair $(\rho_1, \rho_2) \in \mathcal{C}$.

Proof. Let as usual $M = DCI(\overline{Y}_\Gamma)$ be equipped with the index filtration. By Lemma 6.4 we have $F_{4p}M = F_{4p+t}M$ for $0 \leq t \leq 3$, and the map $\zeta : C_A^+(\overline{F}_{4p}M) \rightarrow \hat{C}_A^+(F_{4p}M)$ along with the inclusion $i : \hat{C}_A^+(F_{4(p-1)}M) \rightarrow \hat{C}_A^+(F_{4p}M)$ define a chain isomorphism

$$i + \zeta : \hat{C}_A^+(F_{4(p-1)}M) \oplus C_A^+(\overline{F}_{4p}M) \cong \hat{C}_A^+(F_{4p}M)$$

of DG R -modules for each p . By induction we obtain isomorphisms

$$\hat{C}_A^+(F_{4p}M) \cong \hat{C}_A^+(F_{4(p-r-1)}M) \oplus \left(\bigoplus_{q=0}^r C_A^+(\overline{F}_{4(p-q)}M) \right)$$

for each p and $r \geq 1$. Since the filtration $F_p \hat{C}_A^+(M) = \hat{C}_A^+(F_pM)$ is degreewise bounded above and complete Hausdorff, it follows by first passing to the limit over r and then to the colimit over p that we obtain an isomorphism of DG R -modules

$$\hat{C}_A^+(M) \cong \prod_{p \in \mathbb{Z}} C_A^+(\overline{F}_{4p}M).$$

Hence, as products commutes with homology,

$$(6.3) \quad I^+(\overline{Y}_\Gamma) = \hat{H}_A^+(M) \cong \prod_{p \in \mathbb{Z}} H_A^+(\overline{F}_{4p}M).$$

Write $\mathcal{C} = \mathcal{C}^0 \cup \mathcal{C}^1$ where $\mathcal{C}^i = \{\alpha \in \mathcal{C} : j(\alpha) \equiv 4i \pmod{8}\}$ for $i = 0, 1$. Then

$$H_A^+(\overline{F}_{4p}M) = \begin{cases} \bigoplus_{\alpha \in \mathcal{C}^0} H_A^+(\alpha)[4p] & \text{if } p \equiv 0 \pmod{2} \\ \bigoplus_{\alpha \in \mathcal{C}^1} H_A^+(\alpha)[4p] & \text{if } p \equiv 1 \pmod{2}, \end{cases}$$

and we may simplify the expression in (6.3)

$$(6.4) \quad \prod_{p \in \mathbb{Z}} H_A^+(\overline{F}_{4p}M) = \prod_{p \in \mathbb{Z}} \bigoplus_{\alpha \in \mathcal{C}} H_A^+(\alpha)[8p + j(\alpha)] = \left(\bigoplus_{\alpha \in \mathcal{C}} H_A^+(\alpha)[j(\alpha)] \right)^{\Pi, 8}.$$

By replacing each $H_A^+(\alpha)$ with $R[V_\alpha]$, $R[W_\alpha]$ or $R \cdot g_\alpha$ according to whether α is fully reducible, reducible or irreducible we obtain the additive statement of the theorem.

To finish the proof we need to determine the action of U . Returning to the notation of equation (6.3), for a fixed degree n , we have $I^+(\overline{Y}_\Gamma)_n = \hat{H}_A^+(F_{4p}M)_n$ for each p with $4p > n$. This is a consequence of the formula (6.2) given in Lemma 6.4. Fix the minimal p with $4p > n$ and consider the commutative diagram

$$\begin{array}{ccccc} I^+(\overline{Y}_\Gamma)_n & \xleftarrow{\cong} & \hat{H}_A^+(F_{4p}M)_n & \xleftarrow{\phi} & \prod_{q:4q \leq n} H_A^+(\overline{F}_{4q}M)_n \\ \downarrow U & & \downarrow U & & \downarrow U' \\ I^+(\overline{Y}_\Gamma)_{n-4} & \xleftarrow{\cong} & \hat{H}_A^+(F_{4p}M)_{n-4} & \xleftarrow{\phi} & \prod_{q:4q \leq n} H_A^+(\overline{F}_{4(q-1)}M)_{n-4}. \end{array}$$

Here, ϕ is the isomorphism obtained by taking the limit, as $j \rightarrow \infty$, over the maps

$$([x_q])_{j \leq q \leq n/4} \mapsto \sum_{j \leq q \leq n/4} [\zeta(x_q)],$$

where $x_q \in C_A^+(\overline{F}_{4q}M)$ represents $[x_q] \in H_A^+(\overline{F}_{4q}M)$ and U' is the map that forces the right rectangle to commute. Our task is to determine U' . By Lemma 6.5 we have for $y \in H_A^+(\overline{F}_{4q}M)_n$ that $H(\zeta) \circ U(y) = U \circ H(\zeta)(y)$ provided $n > 4q$, while if $n = 4q$, and $y = [x]$ then $U \circ H(\zeta)(y) = [\psi x]$.

It follows that $U' = \prod U$ if n is not divisible by 4. Otherwise, if $4|n$ so that $n = 4(p-1)$, then $U' = \tau + \prod U$, where $\tau = ([x_q])_{q \leq p-1} = [\psi(x_{p-1})]$. Therefore, in the description

$$I^+(\overline{Y}_\Gamma) = \prod_{p \in \mathbb{Z}} \bigoplus_{\alpha \in \mathcal{C}} H_A^+(\alpha)[8p + j(\alpha)]$$

of (6.4), the action of U is given by taking the product over the internal $R[U]$ -module structure of each factor $H_A^+(\alpha)$ and adding the correction term τ . For each p , τ only affects the terms $H_A^+(\alpha)[8p + j(\alpha)]_{8p+j(\alpha)} = H_A^+(\alpha)_0$. In terms of our generators this correspond to V_α^0 if α is fully reducible, W_α^0 if α is reducible and g_α if α is irreducible. Now, we may require that these generators correspond to $b_\alpha \in H_0(\alpha) \subset M_{4q} \cong C_A^+(\overline{F}_{4q}M) = H_A^+(\overline{F}_{4q}M)_{4q}$ under the identifications of Lemma 6.4. Therefore, using the explicit formula (6.1) for ψ , we obtain

$$U \cdot V_\eta^0 = \sum_{\rho \in \mathcal{C}^{irr}} n_{\rho\eta} g_\rho$$

and similarly for the other types of generators. This completes the proof. \square

Example 6.8. For $Y = \overline{Y}_{O^*}$ there are two irreducibles α, β , the trivial connection θ and a fully reducible η . The grading is given by $j(\theta) = j(\beta) = 0$ and $j(\alpha) = j(\eta) = 4$. Furthermore, $n_{\alpha\beta} = n_{\beta\alpha} = 3$ and $n_{\alpha\theta} = n_{\beta\eta} = 1$. This is obtained from Proposition 4.25. The above theorem gives

$$I^+(Y) = (R[V_\theta] \oplus R \cdot g_\beta \oplus R[V_\eta][4] \oplus R \cdot g_\alpha[4])^{\Pi, 8}.$$

This means that

$$I^+(Y)_n = \begin{cases} R\{g_\beta\} \oplus \prod_{i \geq 0} R\{V_\theta^{2i}, V_\eta^{2i-1}\} & n \equiv 0 \pmod{8} \\ R\{g_\alpha\} \oplus \prod_{i \geq 0} R\{V_\theta^{2i+1}, V_\eta^{2i}\} & n \equiv 4 \pmod{8} \\ 0 & \text{otherwise.} \end{cases}$$

For $x = (r_{2i}V_\eta^{2i})_{i \geq 0} \in I^+(Y)_4$, $r_{2i} \in R$, we have $U \cdot x = r_0g_\alpha + (r_{2i}V_\eta^{2i-1})_{i \geq 1} \in I^+(Y)_0$. For $x = rg_\alpha$, $r \in R$, we have $U \cdot x = 3rg_\beta$.

6.2. The Case I^- . In this section we will calculate $I^-(\overline{Y}_\Gamma)$ for all finite subgroups $\Gamma \subset \text{SU}(2)$. In contrast to the calculations in the previous section, there will be a number of nontrivial differentials in the index spectral sequence. Nevertheless, the spectral sequence will still stabilize after a finite number of steps in each case.

Lemma 6.9. *Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup. Then the only possibly nontrivial differentials in the index spectral sequence*

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_A^-(\alpha)_t \implies I^-(\overline{Y}_\Gamma)_{s+t}$$

are $d^{4r} : E_{4(s+r), -4(r-1)}^{4r} \rightarrow E_{4s, 3}^{4r}$ for s and $r \geq 1$.

Proof. As $j(\alpha) \equiv 0 \pmod{4}$ for each $\alpha \in \mathcal{C}$, we have $E_{s,*}^1 = 0$ unless $4|s$. By Lemma 5.7 and Corollary 5.9 the $R[U]$ -module $H_A^-(\alpha)$ is given by $R[U]$ with $|U| = -4$, $R[Z][2]$ with $|Z| = -2$ or $R[3]$ depending on whether α is fully reducible, reducible or irreducible, respectively. This means that $E_{4s,t}^1 = 0$ for $t \geq 4$ and odd $t \leq 2$. Since the differential d^r has bidegree $(-r, r-1)$ we deduce that it can only be nonzero if it lands in the bidegree $(4s, 3)$ for some s . The only such differentials that also begin in a possibly nonzero group are $d^{4r} : E_{4(s-r), -4(r-1)}^{4r} \rightarrow E_{4s, 3}^{4r}$. \square

Our first task will be to give an explicit formula for the nontrivial differentials. Fix a finite subgroup $\Gamma \subset \mathrm{SU}(2)$ and let $M = DCI(\overline{Y}_\Gamma)$ be equipped with the index filtration. Recall that by Lemma 5.45

$$DCI^-(\overline{Y}_\Gamma) = \mathrm{colim}_p C_A^-(F_p M) = \bigcup_p C_A^-(F_p M).$$

In the following we will use the fact from Lemma 6.4 that $F_{4p+t}M = F_{4p}M$ for all $p \in \mathbb{Z}$ and $0 \leq t \leq 3$ without further mention.

Lemma 6.10. *For all $p, s \in \mathbb{Z}$ and $0 \leq t \leq 3$ we have*

$$C_A^-(F_{4p}M)_{4s+t} \cong \begin{cases} \bigoplus_{0 \leq q \leq p-s} M_{4(s+q)+t} & \text{if } s \leq p \\ 0 & \text{if } s > p \end{cases}$$

$$C_A^-(\overline{F}_{4p}M)_{4s+t} \cong \begin{cases} M_{4p+t} & \text{if } s \leq p \\ 0 & \text{if } s > p \end{cases}$$

The differential in $C_A^-(\overline{F}_{4p}M)$ is given by $u: M_{4p} \rightarrow M_{4p+3}$ in degree $4s$ with $s \leq p$, and vanishes otherwise. The differential in $C_A^-(F_{4p}M)$ is given by

$$\partial((x_q)_{0 \leq q \leq p-s}) = (\partial M x_q - (-1)^t x_{q-1} u)_{0 \leq q \leq p-s+1}$$

for $(x_q)_{0 \leq q \leq p-s} \in \bigoplus_{0 \leq q \leq p-s} M_{4(s+q)+t} = C_A^-(F_{4p}M)_{4s+t}$ where $s \leq p$, and we interpret $x_{-1} = 0 = x_{p-s+1}$.

Proof. This is a simple consequence of Proposition 5.8 using the fact that $F_{4p}M_n = M_n$ for $n \leq 4p+3$ and $F_{4p}M_n = 0$ otherwise. \square

For each $\eta \in \mathcal{C}^{f.red}$, $\lambda \in \mathcal{C}^{red}$ and $\alpha \in \mathcal{C}^{irr}$ we introduce variables U_η, Z_λ and a generator h_α . Then, according to Lemma 5.7 and Corollary 5.9,

$$H_A^-(\eta) = R[U_\eta], \quad H_A^-(\lambda) = R[Z_\lambda][2] \quad \text{and} \quad H_A^-(\alpha) = R \cdot h_\alpha$$

where $|U_\eta| = -4$, $|Z_\lambda| = -2$ and $|h_\alpha| = 3$.

Lemma 6.11. *Let $\{E^r, d^r\}_{r \geq 1}$ be the index spectral sequence with*

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_A^-(\alpha)_t \implies I_{s+t}^-(\overline{Y}_\Gamma).$$

Then, for each $r \geq 1$ and s , $E_{4s, -4(r-1)}^{4r} = E_{4s, -4(r-1)}^1$ is a free R -module with generators

$$\{U_\eta^{r-1} : \eta \in \mathcal{C}^{f.red} \ni j(\eta) \equiv 4s\} \cup \{Z_\lambda^{2r-1} : \lambda \in \mathcal{C}^{red} \ni j(\lambda) \equiv 4s\}$$

and $E_{4s,3}^1$ is a free R -module with generators $\{h_\alpha : \alpha \in \mathcal{C}^{irr} \ni j(\alpha) \equiv 4s\}$. Let $p_r : E_{4s,3}^1 \rightarrow E_{4s,3}^{4r}$ denote the natural surjection. Then in terms of these generators the differential $d^r : E_{4(s+r), -4(r-1)}^{4r} \rightarrow E_{4s,3}^{4r}$ is determined by

$$d^{4r}(U_\eta^{r-1}) = p_r \left(\sum_{(\alpha_1, \dots, \alpha_r)} n_{\alpha_1 \eta} \left(\prod_{i=1}^{r-1} n_{\alpha_{i+1} \alpha_i} \right) \cdot h_{\alpha_r} \right)$$

$$d^{4r}(W_\lambda^{2r-1}) = p_r \left(\sum_{(\alpha_1, \dots, \alpha_r)} n_{\alpha_1 \lambda} \left(\prod_{i=1}^{r-1} n_{\alpha_{i+1} \alpha_i} \right) \cdot h_{\alpha_r} \right),$$

where the sums are taken over tuples $(\alpha_1, \dots, \alpha_r) \in (\mathcal{C}^{irr})^r$ for which $(\eta, \alpha_1, \dots, \alpha_r)$, respectively $(\lambda, \alpha_1, \dots, \alpha_r)$, form a path in the graph S_Γ (see Definition 3.9).

Proof. Before we introduce the explicit generators we will work out a formula for the differentials in terms of the complex M . By Lemma 6.9 we have $E_{4(s+r),-4(r-1)}^{4r} = E_{4(s+r),-4(r-1)}^1 = H_A^-(\overline{F}_{4(s+r)}M)_{4(s+1)}$ and by Lemma 6.10 the latter group may be identified with $\text{Ker}(u) \subset M_{4(s+r)}$. Recall from diagram (5.15) that the differential d^{4r} is defined by first applying the connecting homomorphism $\delta: H_A^-(\overline{F}_{4(s+r)}M) \rightarrow H_A^-(F_{4(s+r)-1}M)$, then lifting along the map $H_A^-(F_{4s}M) \rightarrow H_A^-(F_{4(s+r)-1}M)$ and then pushing down along $H_A^-(F_{4s}M) \rightarrow H_A^-(\overline{F}_{4s}M)$. On the chain level we have the following diagram

$$\begin{array}{ccccc} C_A^-(\overline{F}_{4(s+r)}M)_{4(s+1)} & \xrightarrow{\delta} & C_A^-(F_{4(s+r)-1}M)_{4s+3} & \xleftarrow{\iota} & C_A^-(F_{4s}M)_{4s+3} \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ M_{4(s+r)} & \xrightarrow{\delta} & \bigoplus_{q=0}^{r-1} M_{4(s+q)+3} & \xleftarrow{\iota} & M_{4s+3}. \end{array}$$

The arrows correspond to the connecting homomorphism δ are dashed as they are not well-defined before passing to homology. Given $x \in \text{Ker } u \subset M_{4(s+r)}$ the element $\delta(x)$ is represented on the chain level by $(0, 0, \dots, 0, \partial_M x)$. As ι is the inclusion of the first summand, this element does not lift along ι . Let $\psi: M \rightarrow M$ be the map of Lemma 6.2 and let

$$y := (\psi^{r-q}x)_{q=1}^{r-1} \in \bigoplus_{q=1}^{r-1} M_{4(s+q)} \cong C_A^-(F_{4(s+r-1)}M)_{4(s+1)}.$$

Then $\partial(y) = (\partial_M \psi^{r-1}x, 0, 0, \dots, 0, -\partial_M x)$, which implies that $\iota[\partial_M \psi^{r-1}x] = [(0, \dots, 0, \partial_M x)]$ in homology. By Lemma 6.10 we have the following sequence of isomorphisms

$$M_{4s+3} \cong C_A^-(F_{4s}M)_{4s+3} \cong C_A^-(\overline{F}_{4s}M)_{4s+3} \cong H_A^-(\overline{F}_{4s}M)_{4s+3} \cong E_{4s,3}^1.$$

Moreover, according to Lemma 6.9, there are no outgoing differentials from $E_{4s,3}^r$ for $r \geq 1$. We conclude that $d^{4r}(x)$ is given by the image of $\partial_M \psi^{r-1}x$ along the surjection $E_{4s,3}^1 \rightarrow E_{4s,3}^{4r}$ for each $x \in E_{4(s+r),-4(r-1)}^{4r} \subset M_{4(s+r)}$. In other words, we have the following commutative diagram

$$(6.5) \quad \begin{array}{ccccc} E_{4(s+r),-4(r-1)}^{4r} & \xrightarrow{=} & E_{4(s+r),-4(r-1)}^1 & \xrightarrow{\cong} & \text{Ker}(u) \subset M_{4(s+r)} \\ \downarrow d^{4r} & & & & \downarrow \partial_M \circ \psi^{(r-1)} \\ E_{4s,3}^{4r} & \xleftarrow{p_r} & E_{4s,3}^1 & \xrightarrow{\cong} & M_{4s+3}. \end{array}$$

Now, $\text{Ker } u \subset M_{4(s+r)}$ is freely generated by $b_\eta \in H_0(\eta)$, $b_\lambda \in H_0(\lambda)$ for $\eta \in \mathcal{C}^{f,red}$ and $\lambda \in \mathcal{C}^{red}$ with $j(\eta) \equiv j(\lambda) \equiv 4(s+r)$. It is important to note that under the top horizontal identification in (6.5) we may, and will, require that these correspond to the generators U_η^{r-1} and W_λ^{2r-1} in $E_{4(s+r),-4(r-1)}^1$. Similarly, M_{4s+3} is freely generated by $t_\alpha = b_\alpha \cdot u \in H_3(\alpha)$ for $\alpha \in \mathcal{C}^{irr}$ with $j(\alpha) \equiv 4s$, and under the lower horizontal isomorphism we require that these correspond to the generators $h_\alpha \in E_{4s,3}^1$. Recall from equation (6.1) that

$$\psi(b_\eta) = u^{-1} \partial_M(b_\eta) = \sum_{\alpha \in \mathcal{C}^{irr}} u^{-1}(n_{\alpha\eta} t_\alpha) = \sum_{\alpha \in \mathcal{C}^{irr}} n_{\alpha\eta} b_\alpha.$$

and similarly for $\psi(b_\lambda)$. Taking into account the fact that $n_{\alpha\beta} = 0$ whenever α and β are not adjacent in the graph \mathcal{S}_Γ , we obtain

$$\partial_M \psi^{r-1}(b_\eta) = \sum_{(\alpha_1, \dots, \alpha_r)} (n_{\alpha_1 \eta} n_{\alpha_2 \alpha_1} \cdots n_{\alpha_r \alpha_{r-1}}) \cdot t_{\alpha_r},$$

where the sum is taken over $(\alpha_1, \dots, \alpha_r) \in (C^{irr})^r$ for which $(\eta, \alpha_1, \dots, \alpha_r)$ forms an edge path in S_Γ . An analogous formula holds for b_λ , so in view of diagram (6.5) the proof is complete. \square

By combining the above lemma with Theorem 4.25 we obtain complete control over all the differentials in the spectral sequences. The following result takes care of all the extension problems we will meet.

Lemma 6.12. *Let $\Gamma \subset \mathrm{SU}(2)$ be a finite subgroup and let $(E^r, d^r)_{r \geq 1}$ be the index spectral sequence with*

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_A^-(\alpha)_t \implies I^-(\overline{Y}_\Gamma)_{s+t}.$$

Assume that E_s^∞ is a free $R[U]$ -module for $s = 0, 4$. Then there is an isomorphism of $R[U]$ -modules

$$I^-(\overline{Y}_\Gamma) \cong (E_0^\infty \oplus E_4^\infty)^{\oplus, 8}.$$

Proof. Recall that the index spectral sequence is periodic in the sense that $E_{s,t}^r \cong E_{s+8,t}^r$, or equivalently $E_s^r[8] \cong E_{s+8}^r$, for all $s, t \in \mathbb{Z}$ and $r \in \mathbb{N} \cup \{\infty\}$. Therefore, the assumption that E_0^∞ and E_4^∞ are free $R[U]$ -modules implies that E_s^∞ is a free $R[U]$ -module for each $s \in \mathbb{Z}$, as $E_s^1 = 0$ for all s not divisible by 4.

Write $I = I^-(\overline{Y}_\Gamma)$. By Theorem 5.58 $(E^r, d^r)_{r \geq 1}$ is a spectral sequence of $R[U]$ -modules that converges strongly to I . In other words, I carries an exhaustive and complete Hausdorff filtration $\{F_s I\}_s$ of $R[U]$ -submodules and there is an isomorphism $E_s^\infty \cong \overline{F}_s I$ of $R[U]$ -modules for each $s \in \mathbb{Z}$. In view of the fact that E_s^∞ is free over $R[U]$, the short exact sequence

$$0 \longrightarrow F_{s-1} I \xrightarrow{\iota_s} F_s I \xrightarrow{\pi_s} \overline{F}_s I \cong E_s^\infty \longrightarrow 0$$

splits in the category of $R[U]$ -modules for each $s \in \mathbb{Z}$. Hence, $F_{s-1} I \oplus \overline{F}_s I \cong F_s I$ and inductively

$$F_{s-1} I \oplus \left(\bigoplus_{t=0}^r \overline{F}_{s+t} I \right) \cong F_{s+r} I$$

for $s \in \mathbb{Z}$ and $r \geq 1$. These are all isomorphisms of $R[U]$ -modules. By Lemma 6.10 we have $E_{s,t}^1 = 0$ for all s and $t \geq 4$. Therefore, for fixed $n \in \mathbb{Z}$, $(E_s^1)_n = E_{s,n-s}^1 = 0$ for all $s \leq n - 4 =: s_0$. Hence, $(\overline{F}_s I)_n = 0$ for all $s \leq s_0$. Since the filtration of I is complete Hausdorff it follows that $(F_s I)_n = 0$ for all $s \leq s_0$. Consequently, using the above isomorphisms, we obtain

$$I_n = \mathrm{colim}_{s \geq s_0} (F_s I)_n \cong \mathrm{colim}_{s \geq s_0} \bigoplus_{i=s_0}^s (\overline{F}_i I)_n \cong \bigoplus_s (E_s^\infty)_n$$

for each $n \in \mathbb{Z}$. These isomorphisms piece together to an isomorphism $I \cong \bigoplus_s E_s^\infty$ of $R[U]$ -modules. Finally, we may exploit the periodicity of E^∞ and the fact that $E_s^\infty = 0$ unless $4|s$ to simplify:

$$I \cong \bigoplus_s E_s^\infty \cong \bigoplus_s (E_{8s}^\infty \oplus E_{8s+4}^\infty) \cong \bigoplus_s (E_0^\infty[8s] \oplus E_4^\infty[8s]) = (E_0^\infty \oplus E_4^\infty)^{\oplus, 8}$$

and the proof is complete. \square

It is not possible to give a uniform result for the calculation of $I^-(\overline{Y}_\Gamma)$ for all the subgroups $\Gamma \subset \mathrm{SU}(2)$ simultaneously. We will therefore treat the simplest cases $\Gamma = I^*, O^*, T^*, C_m$ first and treat the more involved case of binary dihedral groups afterwards.

We have the following table over the flat connections extracted from Appendix A. The notation is compatible with Proposition 4.25.

Γ	$\mathcal{C}^{f.red}$	\mathcal{C}^{irr}	\mathcal{C}^{red}
C_{2m}	θ, η		$\lambda_1, \dots, \lambda_{m-1}$
C_{2m+1}	θ		$\lambda_1, \dots, \lambda_m$
I^*	θ	α, β	
O^*	θ, η	α, β	
T^*	θ	α	λ

Theorem 6.13. *The negative equivariant instanton Floer homology associated with the trivial $SU(2)$ -bundle over \bar{Y}_Γ for $\Gamma = C_{2m}, C_{2m+1}$ is given by*

$$I^-(\bar{Y}_{C_{2m}}) = \left((R[U_\theta] \oplus R[U_\eta][4m] \oplus \bigoplus_{i=1}^{m-1} R[Z_{\lambda_i}][4i+2]) \right)^{\oplus, 8}$$

$$I^-(\bar{Y}_{C_{2m+1}}) = \left(R[U_\theta] \oplus \bigoplus_{i=1}^m R[Z_{\lambda_i}][4i+2] \right)^{\oplus, 8},$$

while for $\Gamma = I^*, O^*, T^*$ we have $I^-(\bar{Y}_\Gamma) = (X_\Gamma)^{\oplus, 8}$ where X_Γ is the $R[U]$ -submodule of P_Γ generated by G_Γ specified in the following table.

Γ	P_Γ	G_Γ
I^*	$R[U_\theta]$	$\{U_\theta^2\}$
O^*	$R[U_\theta] \oplus R[U_\eta][4]$	$\{U_\theta^1, U_\eta^1\}$
T^*	$R[U_\theta] \oplus R[Z_\lambda][2]$	$\{U_\theta^1, Z_\lambda^0, 3U_\theta^0 - Z_\lambda^1\}$

Proof. For $\Gamma = I^*, O^*, T^*, C_n$ we will write $\mathcal{C} = \mathcal{C}_\Gamma$ for the set of flat connections and $(E^r, d^r)_{r \geq 1}$ will denote the corresponding index spectral sequence with

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_A^-(\alpha)_t \implies I^-(\bar{Y}_\Gamma)_{s+t}.$$

In each case we will calculate E^∞ and observe that E_0^∞ and E_4^∞ are free $R[U]$ -modules. By Lemma 6.12 this is enough to conclude that $I^-(\bar{Y}_\Gamma) = (E_0^\infty \oplus E_4^\infty)^{\oplus, 8}$. In the following we will make consistent use of Lemma 6.9, Theorem 4.25 and Lemma 6.11 that together determine all the differentials explicitly.

C_n. As $\mathcal{C}^{irr} = \emptyset$ there are no nontrivial differentials in the spectral sequence so that $E^1 = E^\infty$. For $n = 2m$ we have $\mathcal{C} = \{\theta, \eta, \lambda_1, \dots, \lambda_{m-1}\}$ with θ, η fully reducible and the λ_i reducible. The grading is given by $j(\lambda_i) \equiv 4i$ and $j(\eta) \equiv 4m$. This gives

$$E_0^\infty \oplus E_4^\infty = R[U_\theta] \oplus R[U_\eta][j(\eta)] \oplus \left(\bigoplus_{i=1}^{m-1} R[Z_{\lambda_i}][j(\lambda_i) + 2] \right).$$

For $n = 2m + 1$ we have $\mathcal{C} = \{\theta, \lambda_1, \dots, \lambda_m\}$ with θ fully reducible and the λ_i reducible. The grading is given by $j(\lambda_i) \equiv 4i$. Hence,

$$E_0^\infty \oplus E_4^\infty = R[U_\theta] \oplus \left(\bigoplus_{i=1}^m R[Z_{\lambda_i}][j(\lambda_i) + 2] \right).$$

In both cases these are free $R[U]$ -modules ($R[Z_\lambda]$ is freely generated by $\{Z_\lambda^0, Z_\lambda^1\}$). The stated results are obtained by applying $X \mapsto X^{\oplus, 8}$ to the above formulas for $E_0^\infty \oplus E_4^\infty$.

I*. In this case we have $\mathcal{C} = \{\theta, \alpha, \beta\}$ with α and β irreducible. The grading is given by $j(\theta) = j(\beta) = 0$ and $j(\alpha) = 4$. The first nontrivial differential in the index spectral sequence is $d^4: E_{8s,0}^4 = R\{U_\theta^0\} \rightarrow R\{h_\alpha\} = E_{8s-4,3}^4$ and is given by $d^4(U_\theta^0) = n_{\alpha\eta}h_\alpha = h_\alpha$. Therefore, $E_{8s+4,3}^\infty = 0 = E_{8s,0}^\infty$. The next differential is $d^8: E_{8s,-4}^8 = R\{U_\theta^1\} \rightarrow R\{h_\beta\} = E_{8(s-1),3}^8$ and is given by $d^8(U_\theta^1) = n_{\beta\alpha}n_{\alpha\eta}h_\beta = 4h_\beta$. As $2 \in R$ is invertible, this is an isomorphism and we conclude that $E_{8s,3}^\infty = 0$ and $E_{8s,-4}^\infty = 0$. There are no more nontrivial differentials in the spectral sequence. Therefore, $E_4^\infty = 0$ and $E_0^\infty \subset R[U_\theta]$ is the free $R[U]$ -submodule generated by U_θ^2 as required.

O*. Here we have $\mathcal{C} = \{\theta, \eta, \alpha, \beta\}$ with θ, η fully reducible and α, β irreducible. The grading is $j(\theta) = j(\beta) = 0$ and $j(\alpha) = j(\eta) = 4$. The first nontrivial differentials are

$$\begin{aligned} d^4: E_{8s+4,0}^4 &= R\{U_\eta^0\} \rightarrow R\{h_\beta\} = E_{8s,3}^4 \\ d^4: E_{8s,0}^4 &= R\{U_\theta^0\} \rightarrow R\{h_\alpha\} = E_{8s-4,3}^4. \end{aligned}$$

These are both isomorphisms as $n_{\beta\eta} = 1 = n_{\alpha\theta}$. This implies that $E_{8s,3}^\infty = E_{8s,3}^5 = 0$ and $E_{8s-4,3}^\infty = E_{8s-4,3}^5 = 0$. There can therefore be no more nontrivial differentials. Hence, $E_0^\infty \subset R[U_\theta]$ and $E_4^\infty \subset R[U_\eta][4]$ are the free $R[U]$ -submodules generated by U_θ^1 and U_η^1 , respectively.

T*. In this case $\mathcal{C} = \{\theta, \alpha, \lambda\}$ with θ fully reducible, α irreducible and λ reducible. The grading is $j(\theta) = j(\lambda) = 0$ and $j(\alpha) = 4$. The first nontrivial differential is $d^4: E_{8s,0}^4 = R\{Z_\lambda^1, U_\theta^0\} \rightarrow R\{h_\alpha\} = E_{8s-4,3}^4$ and is determined by $d^4(U_\theta^0) = n_{\alpha\theta}h_\alpha = h_\alpha$ and $d^4(Z_\lambda^1) = n_{\alpha\lambda}h_\alpha = 3h_\alpha$. This map is surjective with kernel $R\{3U_\theta^0 - Z_\lambda^1\}$. Therefore, $E_{8s-4,3}^\infty = E_{8s-4,3}^4 = 0$ and there are no more nontrivial differentials. We find $E_4^\infty = 0$, $(E_0^\infty)_0 = R\{3U_\theta^0 - Z_\lambda^1\}$ and $(E_0^\infty)_n = (R[U_\theta] \oplus R[Z_\lambda][2])_n$ in all other degrees n . It now suffices to observe that E_0^∞ is indeed the free submodule of $R[U_\theta] \oplus R[Z_\lambda][2]$ generated by $\{Z_\lambda^0, 3U_\theta^0 - Z_\lambda^1, U_\theta^1\}$. This completes the final case and the proof. \square

We will now consider the binary dihedral groups D_m^* . It is necessary to partition the calculations into cases depending on the residue of $m \bmod 4$. We have the following table over the flat connections

	$\mathcal{C}^{f.red}$	\mathcal{C}^{irr}	\mathcal{C}^{red}
D_{4n}^*	$\theta, \eta_1, \eta_2, \eta_3$	$\alpha_1, \dots, \alpha_{2n}$	
D_{4n+1}^*	θ, η	$\alpha_1, \dots, \alpha_{2n}$	λ
D_{4n+2}^*	$\theta, \eta_1, \eta_2, \eta_3$	$\alpha_1, \dots, \alpha_{2n+1}$	
D_{4n+3}^*	θ, η	$\alpha_1, \dots, \alpha_{2n+1}$	λ

in agreement with Appendix A and Proposition 4.25. In all cases the grading of the irreducibles are given by $j(\alpha_i) \equiv 4i \bmod 8$. For D_{4n+2}^* and D_{4n+3}^* all the fully reducibles and reducibles satisfy $j(\rho) = 0$. For D_{4n}^* we have $j(\theta) = j(\eta_1) = 0$, $j(\eta_2) = j(\eta_3) = 4$, while for D_{4n+1}^* we have $j(\theta) = j(\eta) = 0$ and $j(\lambda) = 4$. For the convenience of the reader we include the relevant diagrams for $S_{D_m^*}$ from Proposition 4.25.

$$\begin{array}{c} \theta \searrow 1 \quad \quad \quad S_{D_{2m}^*} \quad \quad \quad 1 \nearrow \eta_2 \\ \eta_1 \swarrow 1 \quad \alpha_1 - \alpha_2 \cdots \cdots \alpha_{m-1} - \alpha_m \quad (2|2) \quad \quad \quad (2|2) \quad \quad \quad 1 \nearrow \eta_3 \end{array}$$

$$\begin{array}{c} \theta \searrow 1 \quad \quad \quad S_{D_{2m+1}^*} \quad \quad \quad \alpha_m - \lambda \\ \eta \swarrow 1 \quad \alpha_1 - \alpha_2 \cdots \cdots \alpha_{m-1} - \alpha_m \quad (2|2) \quad \quad \quad (2|2) \quad \quad \quad 2 \end{array}$$

Theorem 6.14. *The negative instanton Floer homology associated with the trivial $SU(2)$ -bundle over \overline{Y}_Γ for $\Gamma = D_m^*$ is given by $I^-(\overline{Y}_{D_m^*}) = (X_m)^{\oplus, 8}$, where X_m is the $R[U]$ -submodule of P_m*

generated by G_m specified in the following tables.

m	P_m
$4n$	$R[U_\theta] \oplus R[U_{\eta_1}] \oplus R[U_{\eta_2}][4] \oplus R[U_{\eta_3}][4]$
$4n+1$	$R[U_\theta] \oplus R[U_\eta] \oplus R[Z_\lambda][6]$
$4n+2$	$R[U_\theta] \oplus R[U_{\eta_1}] \oplus R[U_{\eta_2}] \oplus R[U_{\eta_3}]$
$4n+3$	$R[U_\theta] \oplus R[U_\eta] \oplus R[Z_\lambda][2]$

m	G_m
$4n$	$\{U_\theta^0 - U_{\eta_1}^0, U_{\eta_2}^0 - U_{\eta_3}^0, U_\theta^n, U_{\eta_2}^n\}$
$4n+1$	$\{U_\theta^0 - U_\eta^0, Z_\lambda^0, U_\theta^n, Z_\lambda^{2n+1}\}$
$4n+2$	$\{U_\theta^0 - U_{\eta_1}^0, U_{\eta_2}^0 - U_{\eta_3}^0, U_\theta^n - U_{\eta_2}^n, U_\theta^{n+1}\}$
$4n+3$	$\{U_\theta^0 - U_\eta^0, Z_\lambda^0, 2U_\theta^n - Z_\lambda^{2n+1}, U_\theta^{n+1}\}$

Proof. Observe first that the modules P_m in the above table coincides with the $R[U]$ -submodule of $E_0^1 \oplus E_4^1$ in the index spectral sequence generated by the reducibles and fully reducibles. Furthermore, the submodule generated by G_m is free in each case. Therefore, following the same procedure as in the proof of Theorem 6.13, it will be sufficient to show that $E_0^\infty \oplus E_4^\infty$ is the $R[U]$ -submodule of P_m generated by G_m in each case.

For $m = 4n, 4n+1$ we have

$$E_{8s,3}^1 = R\{h_{\alpha_2}, h_{\alpha_4}, \dots, h_{\alpha_{2n}}\} \text{ and } E_{8s+4,3}^1 = R\{h_{\alpha_1}, h_{\alpha_3}, \dots, h_{\alpha_{2n-1}}\}$$

for each $s \in \mathbb{Z}$. For $m = 4n+2, 4n+3$ we have the same formulas except that one generator $h_{\alpha_{2n+1}}$ is adjoined to the latter group. To simplify the notation put $h_i := h_{\alpha_i}$ for $1 \leq i \leq 2n+1$. For each $r \geq 1$ let $p_r: E_{4s,3}^1 \rightarrow E_{4s,3}^{4r}$ denote the natural surjection.

Consider first the cases $m = 4n, 4n+1$. To unify the notation slightly we write $(U_1, U_2, U_3, U_4) = (U_\theta, U_{\eta_1}, U_{\eta_2}, U_{\eta_3})$ for $m = 4n$ and $(U_1, U_2, Z) = (U_\theta, U_\eta, Z_\lambda)$ for $m = 4n+1$. We claim that for $1 \leq r \leq n$ the differentials

$$f^r := d^{4r}: E_{8s, -4(r-1)}^{4r} \rightarrow E_{8s-4r, 3}^{4r}$$

$$g^r := d^{4r}: E_{8s+4, -4(r-1)}^{4r} \rightarrow E_{8s-4(r-1), 3}^{4r}$$

are given by the formulas

$$f^r(U_1^{r-1}) = f^r(U_2^{r-1}) = 2^{r-1}p_r(h_r)$$

$$g^r(U_3^{r-1}) = g^r(U_4^{r-1}) = 2^{r-1}p_r(h_{2n-r+1})$$

$$g^r(Z^{2r-1}) = 2^r p_r(h_{2n-r+1}).$$

We verify this by induction on r . Let $1 \leq r \leq n-1$ and assume that the statement is true for all i with $1 \leq i < r$. This implies in particular that $p_r(h_i) = 0$ for $1 \leq i < r$ and $2n-r+1 < i \leq 2n$. By Lemma 6.11 we have

$$f^r(U_1^{r-1}) = p_r \left(\sum_{(\beta_1, \dots, \beta_r)} (n_{\beta_1 \theta} n_{\beta_2 \beta_1} \cdots n_{\beta_r \beta_{r-1}}) \cdot h_{\beta_r} \right),$$

where the sum runs over all $(\beta_1, \dots, \beta_r) \in (\mathcal{C}^{irr})^r$ for which $(\theta, \beta_1, \dots, \beta_r)$ forms an edge path in \mathcal{S}_Γ . As such a path has length r , it must, in view of the graphs $\mathcal{S}_{D_m^*}$ shown above, terminate at some vertex α_i with $i \leq r$. By the inductive hypothesis $p_r(h_{\alpha_i}) = 0$ for $i < r$, so the only nonzero term in the formula corresponds to the path $(\alpha_1, \alpha_2, \dots, \alpha_r)$. As $n_{\alpha_{i+1}\alpha_i} = 2$ for each i and $n_{\alpha_1\theta} = 1$ we

conclude that $f^r(U_1^{r-1}) = 2^{r-1}p_r(h_r)$. The other formulas follow by essentially identical arguments, we only note that the additional factor 2 picked up in the last formula follows from the fact that $n_{\alpha_{2n}\lambda} = 2$. This completes the inductive step and the claim is verified.

The above formulas for the differentials imply, as $2 \in R$ is invertible, that the generators h_i , $1 \leq i \leq 2n$, of $E_{8s,3}^1$ and $E_{8s+4,3}^1$ are killed off two by two until we reach $E_{8s,3}^{4n+1} = 0$, $E_{8s+4,3}^{4n+1} = 0$. Therefore, by Lemma 6.9, $E^{4n+1} = E^\infty$. Moreover, we find $E_{8s,-4(r-1)}^\infty = \text{Ker}(f^r) = R\{(U_1^{r-1} - U_2^{r-1})\}$ for $1 \leq r \leq n$, and similarly

$$E_{8s+4,-4(r-1)}^\infty = \begin{cases} R\{(U_3^{r-1} - U_4^{r-1})\} & \text{for } m = 4n \\ 0 & \text{for } m = 4n + 1 \end{cases}$$

for $1 \leq r \leq n$. In all other degrees we have $E_{s,t}^1 = E_{s,t}^\infty$. We therefore see that E_0^∞ is freely generated as an $R[U]$ -module by $\{(U_1^0 - U_2^0), U_1^n\}$ for both $m = 4n$ and $m = 4n + 1$, while E_4^0 is freely generated by $\{(U_3^0 - U_4^0), U_3^n\}$ for $m = 4n$ and by $\{Z_\lambda^0, Z_\lambda^{2n+1}\}$ for $m = 4n + 1$. We have thus completed the cases $m = 4n, 4n + 1$.

Consider now the cases $m = 4n+2, 4n+3$. As earlier we write $(U_1, U_2, U_3, U_4) = (U_\theta, U_{\eta_1}, U_{\eta_2}, U_{\eta_3})$ for $m = 4n+2$ and $(U_1, U_2, Z) = (U_\theta, U_\eta, Z_\lambda)$. Here $U_i^{r-1}, Z^{2r-1} \in E_{8s,-4(r-1)}^1$ for each i and $r \geq 1$. For $1 \leq r \leq n+1$ the differentials

$$d^{4r} : E_{8s,-4(r-1)}^{4r} \rightarrow E_{8s-4r,3}^{4r}$$

are given by the formulas

$$\begin{aligned} d^{4r}(U_1^{r-1}) &= d^{4r}(U_2^{r-1}) = 2^{r-1}p_r(h_r) \\ d^{4r}(U_3^{r-1}) &= d^{4r}(U_4^{r-1}) = 2^{r-1}p_r(h_{2n+2-r}) \\ d^{4r}(Z^{2r-1}) &= 2^r p_r(h_{2n+2-r}). \end{aligned}$$

This may be verified by induction in exactly the same way as in the cases $n = 4m, 4m + 1$. We conclude that $E_{8s,3}^{4n+5} = 0 = E_{8s+4,3}^{4n+5}$ and hence $E^{4n+5} = E^\infty$. In addition, for $0 \leq r \leq n-1$ we obtain

$$E_{8s,-4r}^\infty = \text{Ker}(d^{4(r+1)}) = \begin{cases} R\{(U_1^r - U_2^r), (U_3^r - U_4^r)\} & \text{if } m = 4n + 2 \\ R\{(U_1^r - U_2^r)\} & \text{if } m = 4n + 3 \end{cases}$$

Furthermore, for $m = 4n + 2$ we have $d^{4(n+1)}(U_i^n) = 2^n p_{n+1}(h_{n+1})$ for each $1 \leq i \leq 4$, and for $m = 4n + 3$ we have

$$2d^{4(n+1)}(U_i^n) = d^{4(n+1)}(Z^{2n+1}) = 2^{n+1}p_{n+1}(h_{2n+1})$$

for $i = 1, 2$. From this we deduce that

$$E_{8s,-4n}^\infty = \begin{cases} R\{(U_1^n - U_2^n), (U_3^n - U_4^n), (U_1^n - U_3^n)\} & \text{if } m = 4n + 2 \\ R\{(U_1^n - U_2^n), (2U_1^{n+1} - Z^{2n+1})\} & \text{if } m = 4n + 3 \end{cases}$$

In all other degrees we have $E_{s,t}^1 = E_{s,t}^\infty$. For $m = 4n + 2$ we conclude that E_0^∞ is freely generated by $\{U_1 - U_2, U_3 - U_4, U_1^n - U_3^n, U_1^{n+1}\}$ and for $m = 4n + 3$ we conclude that E_0^∞ is freely generated by $\{(U_1^0 - U_2^0), Z_\lambda^0, 2U_1^n - Z^{2n+1}, U_1^{n+1}\}$. In both cases $E_4^\infty = 0$. This completes the final two cases and hence the proof. \square

6.3. The Case I^∞ . This is the simplest calculation. Let $\Gamma \subset \text{SU}(2)$ be a finite subgroup and write as usual $M = \text{DCI}(\overline{Y}_\Gamma)$ and \mathcal{C} for the set of flat connections. Introduce a variable T_η for each $\eta \in \mathcal{C}^{f.red}$ and a variable S_λ for each $\lambda \in \mathcal{C}^{red}$, where $|T_\eta| = -4$, $|S_\lambda| = -2$. Then according to Corollary 5.30 we have $H_A^\infty(\alpha) = 0$ for α irreducible, $H_A^\infty(\lambda) = R[S_\lambda, S_\lambda^{-1}]$ for λ reducible and $H_A^\infty(\eta) = R[T_\eta, T_\eta^{-1}]$ for η fully reducible. The $R[U]$ -module structure is given by $U \cdot T_\eta^i = T_\eta^{i+1}$ and $U \cdot S_\lambda^i = S_\lambda^{i-2}$ for each $i \in \mathbb{Z}$.

Definition 6.15. For an $R[U]$ -module X we define the mod 8 periodic $R[U]$ -module $X^{\Pi_\infty, 8}$ degreewise by

$$X_n^{\Pi_\infty, 8} = \prod_{s \rightarrow \infty} X_{n+8s}$$

and define $U: X_n^{\Pi_\infty, 8} \rightarrow X_{n-4}^{\Pi_\infty, 8}$ to be the product over $U: X_{n+8s} \rightarrow X_{n-4+8s}$ for $s \in \mathbb{Z}$.

Theorem 6.16. *The Tate equivariant instanton Floer homology associated with the trivial $SU(2)$ -bundle over \overline{Y}_Γ is given by*

$$I^\infty(\overline{Y}_\Gamma) = \left[\left(\bigoplus_{\eta \in \mathcal{C}^{f, red}} R[T_\eta, T_\eta^{-1}][j(\eta)] \right) \oplus \left(\bigoplus_{\lambda \in \mathcal{C}^{red}} R[S_\lambda, S_\lambda^{-1}][j(\lambda)] \right) \right]^{\Pi_\infty, 8}.$$

Proof. Let $(E^r, d^r)_{r \geq 1}$ be the index spectral sequence with

$$E_{s,t}^1 = \bigoplus_{j(\alpha) \equiv s} H_A^\infty(\alpha)_t \implies I^\infty(\overline{Y}_\Gamma)_{s+t}.$$

From the fact that $j(\alpha) \equiv 0 \pmod{4}$ for each $\alpha \in \mathcal{C}$ and that $H_A^\infty(\alpha)$ is concentrated in even degrees for all types of orbits, we conclude that $E_{s,t}^1 = 0$ for all (s, t) with s odd or t odd. It follows that there are no nontrivial differentials in the spectral sequence so that $E^1 = E^\infty$. This implies by Proposition 5.57 part (iii) and Theorem 5.55 that the spectral sequence converges strongly.

Write $I = I^\infty(\overline{Y}_\Gamma)$. To resolve the extension problems observe first that E_{4s}^1 is a (shifted) direct sum of $R[U]$ -modules $R[T_\eta, T_\eta^{-1}]$ and $R[S_\lambda, S_\lambda^{-1}]$. These contain free $R[U]$ -submodules $R[T_\eta]$ and $R[S_\lambda]$ (freely generated by $\{S_\lambda^0, S_\lambda^1\}$), respectively, and the whole module is the localization of these free submodules in the multiplicatively closed subset $\{U^i\}_{i \geq 0} \subset R[U]$. By Corollary 5.50 $U: I \rightarrow I$ is an isomorphism, and the same is valid for the $R[U]$ -submodule $F_s I$ for each $s \in \mathbb{Z}$. We may therefore for each s construct a splitting of the short exact sequence

$$0 \longrightarrow F_{s-1}I \longrightarrow F_s I \xrightarrow{\xi_s} \overline{F}_s I \cong E_s^\infty \longrightarrow 0$$

by defining ξ to be a section on the free submodule and then extending to the whole module using the universal property of localization. By induction we obtain an isomorphism

$$F_s I / F_t I \cong \bigoplus_{q=t+1}^s E_q^\infty$$

of $R[U]$ -modules for each pair $s > t$. Since the filtration of I is exhaustive and complete Hausdorff we obtain, exploiting the periodicity,

$$I \cong \lim_t \operatorname{colim}_s F_s I / F_t I \cong \prod_{s \rightarrow -\infty} E_s^\infty \cong \prod_{s \rightarrow -\infty} (E_0^\infty[-8s] \oplus E_4^\infty[-8s])$$

and the final term is precisely $(E_0^\infty \oplus E_4^\infty)^{\Pi_\infty, 8} = (E_0^1 \oplus E_4^1)^{\Pi_\infty, 8}$. The proof is completed by observing that

$$E_0^1 \oplus E_4^1 = \left(\bigoplus_{\eta \in \mathcal{C}^{f, red}} R[T_\eta, T_\eta^{-1}][j(\eta)] \right) \oplus \left(\bigoplus_{\lambda \in \mathcal{C}^{red}} R[S_\lambda, S_\lambda^{-1}][j(\lambda)] \right).$$

□

Proposition 6.17. *For each finite subgroup $\Gamma \subset \mathrm{SU}(2)$ the homology norm map*

$$H(N): I^+(\overline{Y}_\Gamma)_n \rightarrow I^-(\overline{Y}_\Gamma)_{n+3}$$

vanishes. In particular, the exact triangle of $R[U]$ -modules of Corollary 5.44 splits into a short exact sequence

$$0 \longrightarrow I^-(\overline{Y}_\Gamma) \longrightarrow I^\infty(\overline{Y}_\Gamma) \longrightarrow I^+(\overline{Y}_\Gamma)[4] \longrightarrow 0 .$$

Proof. The calculations of Theorem 6.7, Theorem 6.13 and Theorem 6.14 show that $I^+(\overline{Y}_\Gamma)$ and $I^-(\overline{Y}_\Gamma)$ are concentrated in even degrees. Therefore, as the norm map has degree 3, it must vanish. \square

6.4. Calculations for Y_Γ . In this section we explain, omitting some details, the necessary modifications needed to calculate the equivariant instanton Floer groups for Y_Γ , that is, S^3/Γ equipped with the standard orientation inherited from S^3 . The tools needed to handle this are contained in [Eis19, Theorem 7.10]. First, the given result states that there is an isomorphism $I^\infty(Y_\Gamma) \cong I^\infty(\overline{Y}_\Gamma)$. Second, in the proof it is verified that $DCI(Y_\Gamma) \cong DCI(\overline{Y}_\Gamma)^\vee$ and from this it is not hard to express $DCI(Y_\Gamma)$ as the totalization of a multicomplex.

Lemma 6.18. *Let $i: \mathcal{C} \rightarrow \mathbb{Z}/8$ denote the grading for Y_Γ and let $j: \mathcal{C} \rightarrow \mathbb{Z}/8$ denote the grading for \overline{Y}_Γ . Then*

$$i(\alpha) = \begin{cases} j(\alpha) - 3 & \text{if } \alpha \in \mathcal{C}^{irr} \\ j(\alpha) - 2 & \text{if } \alpha \in \mathcal{C}^{red} \\ j(\alpha) & \text{if } \alpha \in \mathcal{C}^{f.red} \end{cases}$$

Moreover, $DCI(Y_\Gamma) = \mathrm{Tot}^{oplus}(DCI(Y_\Gamma)_{,*}, \{\partial^r\}_{r=1}^4)$ where*

$$DCI(Y_\Gamma)_{s,t} = \bigoplus_{i(\alpha) \equiv s} H_t(\alpha),$$

$\partial^2 = 0$ and $\partial^1, \partial^3, \partial^4$ are given for $b_\alpha \in H_0(\alpha) \subset DCI(Y_\Gamma)_{4s+1,0}$, $\alpha \in \mathcal{C}^{irr}$, by

$$\partial^1(b_\alpha) = \sum_{\eta \in \mathcal{C}^{f.red}} n_{\alpha\eta} b_\eta$$

$$\partial^3(b_\alpha) = \sum_{\lambda \in \mathcal{C}^{red}} n_{\alpha\lambda} t_\lambda$$

$$\partial^4(b_\alpha) = \sum_{\beta \in \mathcal{C}^{irr}} n_{\alpha\beta} t_\beta.$$

and vanish otherwise.

The patterns seen in the calculations of $I^\pm(\overline{Y}_\Gamma)$ are reversed for $I^\pm(Y_\Gamma)$. Compare the following with Lemma 6.1 and Lemma 6.9.

Lemma 6.19. *Let $(E^r, d^r)_{r \geq 1}$ be the index spectral sequence with*

$$E_{s,t}^1 = \bigoplus_{i(\alpha) \equiv s} H_A^-(\alpha)_t \implies I^-(Y_\Gamma)_{s+t}.$$

Then the spectral sequence immediately degenerates, i.e., $E^1 = E^\infty$.

Let $(E^r, d^r)_{r \geq 1}$ be the index spectral sequence with

$$E_{s,t}^1 = \bigoplus_{i(\alpha) \equiv s} H_A^+(\alpha)_t \implies I^+(Y_\Gamma)_{s+t}.$$

Then the only possibly nontrivial differentials are $d^{2r+1}: E_{4s+1,0}^{2r+1} \rightarrow E_{4s-2r,2r}^{2r+1}$ for $r \geq 0$ and $s \in \mathbb{Z}$.

Proof. In the first case recall that $H_A^-(\alpha) = R \cdot h_\alpha$ with $|h_\alpha| = 3$ for $\alpha \in \mathcal{C}^{irr}$, $H_A^-(\lambda) = R[Z_\lambda][2]$, $|Z_\lambda| = -2$ for $\lambda \in \mathcal{C}^{red}$ and $H_A^-(\eta) = R[U_\eta]$, $|U_\eta| = -4$ for $\eta \in \mathcal{C}^{f.red}$. By the above lemma $i(\rho) \equiv 0 \pmod{2}$ for each $\rho \in \mathcal{C}^{ref} \cup \mathcal{C}^{f.red}$ and $i(\alpha) \equiv 1 \pmod{4}$ for each $\alpha \in \mathcal{C}^{irr}$. Therefore $E_{s,t}^1$ can only be nonzero if the total degree $s+t$ is even. As each differential d^r reduces the total degree by one, they must all vanish and the first assertion follows.

In the second case, $H_A^+(\eta) = R[V_\eta]$, $|V_\eta| = 4$, $H_A^+(\lambda) = R[W_\lambda]$, $|W_\lambda| = 2$ and $H_A^+(\alpha) = R \cdot g_\alpha$, $|g_\alpha| = 0$, where $\eta \in \mathcal{C}^{f.red}$, $\lambda \in \mathcal{C}^{red}$ and $\alpha \in \mathcal{C}^{irr}$. Due to the indexing mentioned above the only possibly nontrivial module $E_{s,t}^1$ with $s+t$ odd are $E_{4s+1,0}^1$. This module is freely generated by g_α for $\alpha \in \mathcal{C}^{irr}$ satisfying $i(\alpha) \equiv 4s+1 \pmod{8}$. Therefore, a nontrivial differential must start in $E_{4s+1,0}^r$ for some r and s . As $E_{s,t}^r = 0$ for each $t > 0$ when s is odd, we conclude that the only possibly nontrivial differentials are $d^{2r+1}: E_{4s+1,0}^{2r+1} \rightarrow E_{4s-2r,2r}^{2r+1}$. \square

For the calculation of $I^-(Y_\Gamma)$ one is therefore left with solving an extension problem. In this case the extension problem is slightly simpler. We only state the conclusion. Note that in the statement below we are using the grading function $j: \mathcal{C} \rightarrow \mathbb{Z}/8$ for \bar{Y}_Γ , regarded as taking values in $\{0, 4\}$.

Theorem 6.20. *The negative equivariant instanton Floer homology $I^-(Y_\Gamma)$ associated with the trivial $SU(2)$ -bundle over Y_Γ is given by*

$$\left[\left(\bigoplus_{\alpha \in \mathcal{C}^{irr}} R \cdot h'_\alpha[j(\alpha)] \right) \oplus \left(\bigoplus_{\lambda \in \mathcal{C}^{red}} R[Z_\lambda][j(\lambda)] \right) \oplus \left(\bigoplus_{\eta \in \mathcal{C}^{f.red}} R[U_\eta][j(\eta)] \right) \right]^{\oplus, 8},$$

where $|h'_\alpha| = 0$, $|Z_\lambda| = -2$ and $|U_\eta| = -4$. The $R[U]$ -module structure is determined by $U \cdot U_\eta^p = U_\eta^{p+1}$, $U \cdot Z_\lambda^p = Z_\lambda^{p+2}$ for $p \geq 0$ and

$$U \cdot h'_\alpha = \left(\sum_{\beta \in \mathcal{C}^{irr}} n_{\alpha\beta} h'_\beta \right) + \left(\sum_{\lambda \in \mathcal{C}^{red}} n_{\alpha\lambda} Z_\lambda^0 \right) + \left(\sum_{\eta \in \mathcal{C}^{f.red}} n_{\alpha\eta} V_\eta^0 \right).$$

To calculate $I^+(Y_\Gamma)$ one may proceed as in Lemma 6.11 to determine the differentials in the spectral sequence explicitly and then resolve the resulting extension problem. However, one can avoid this rather technical endeavor by using the following result. If M is a DG $R[U]$ -module we write $M^\vee = \text{Hom}(M, R)$ for the dual complex. Our convention is that $(\partial_{M^\vee} f)(x) = (-1)^{|f|+1} f(\partial_M x)$ and $(U \cdot f)(x) = f(Ux)$ for $f \in M^\vee = \text{Hom}(M_n, R)$. The content of the following lemma, which applies more generally, is stated in the proof of [Eis19, Theorem 7.10].

Lemma 6.21. *There is an isomorphism of DG $R[U]$ -modules*

$$DCI^+(Y_\Gamma) \cong DCI^-(\bar{Y}_\Gamma)^\vee.$$

Proof. Let $M = DCI(\bar{Y}_\Gamma)$ and recall that $M^\vee \cong DCI(Y_\Gamma)$. Using the explicit models for DCI^\pm given in Theorem 5.47 we find

$$\begin{aligned} DCI^-(\bar{Y}_\Gamma)_n^\vee &= \text{Hom} \left(\bigoplus_{p \geq 0} M_{-n+4p}, R \right) \cong \prod_{p \geq 0} \text{Hom}(M_{-n+4p}, R) \\ &= \prod_{p \geq 0} M_{n-4p}^\vee \cong DCI^+(Y_\Gamma)_n. \end{aligned}$$

It is a straightforward calculation to check that these isomorphisms piece together to give an isomorphism of DG $R[U]$ -modules taking into account that the $\Lambda_R[u]$ -module structure on M^\vee is given by $(f \cdot u)(x) = (-1)^{|u||x|} f(xu) = (-1)^{|x|} f(xu)$. \square

Theorem 6.22. *There is an isomorphisms of $R[U]$ -modules*

$$I^+(Y_\Gamma) \cong I^-(\bar{Y}_\Gamma)^\vee$$

for each finite subgroup $\Gamma \subset \mathrm{SU}(2)$.

Proof. Since R is a principal ideal domain and the complex $DCI^-(\bar{Y}_\Gamma)$ is degreewise free over R (use the explicit model and the fact that $DCI(\bar{Y}_\Gamma)$ is degreewise free), there is a universal coefficient theorem. In view of the above lemma, this takes the form of a short exact sequence

$$\mathrm{Ext}_R^1(I_{n-1}^-(\bar{Y}_\Gamma), R) \longrightarrow I_{-n}^+(Y_\Gamma) \longrightarrow \mathrm{Hom}(I^-(\bar{Y}_\Gamma)_n, R)$$

for each $n \in \mathbb{Z}$. By Theorem 6.13 and Theorem 6.14 we know that $I^-(\bar{Y}_\Gamma)$ is degreewise free over R . Consequently, the first term in the above sequence vanishes and we are left with an isomorphism $I^+(Y_\Gamma) \cong (I^-(\bar{Y}_\Gamma)_\Gamma)^\vee$ of $R[U]$ -modules as required. \square


Note that the calculations of Theorem 6.20 and Theorem 6.7 are compatible with the duality $I^+(\bar{Y}_\Gamma) \cong (I^-(Y_\Gamma))^\vee$ as well.

APPENDIX A. CHARACTER TABLES, DYNKIN GRAPHS AND FLAT CONNECTIONS

This appendix contains various useful facts concerning the binary polyhedral groups. Of particular importance are their complex representation theory, the corresponding McKay graphs and complete lists of the 1-dimensional quaternionic representations.

A.1. Cyclic Groups. The complex representation theory of cyclic groups are of course very well known, so we only include a brief summary for completeness. Let $C_l = \langle g : g^l = 1 \rangle$ be the cyclic group of order l . The irreducible complex representations are $\rho_k : C_l \rightarrow U(1)$ for $0 \leq k < l$, where $\rho_k(g) = e^{\frac{2\pi i k}{l}}$. If we interpret the indices modulo l we have $\rho_k \otimes \rho_{k'} \cong \rho_{k+k'}$ and $\rho_k^* \cong \rho_{-k}$. We embed $C_l \subset \mathrm{SU}(2)$ through the representation $\rho_1 \oplus \rho_{-1}$.


Suppose that l is odd. Then all of the ρ_k , $1 \leq k < l$, are of complex type, so the 1-dimensional quaternionic representations are given by $\theta = 2\rho_0$ and $\lambda_k = \rho_k \oplus \rho_{-k}$ for $1 \leq k < l/2$. The McKay graph $\bar{\Delta}_{C_l} \cong \tilde{A}_{l-1}$ takes the form



where the line indicates the involution symmetry. The quotient graph $\bar{\Delta}_{C_l}/(\iota)$ is given by



Suppose that l is even. Then $\rho_{l/2}$ is of real type, so the flat connections are given by $\theta = 2\rho_0$, $\eta = 2\rho_{k/2}$ and $\lambda_k = \rho_k \oplus \rho_{-k}$ for $1 \leq k < l/2$. The McKay graph takes the form



with quotient graph



A.2. Binary Dihedral Groups. A presentation for the binary dihedral group D_n^* , $n \geq 2$, is given by

$$D_n^* = \langle a, x \mid a^n = x^2, axa = x \rangle.$$

Note that $a^{2n} = a^n x^2 = xa^{-n}x = xx^{-2}x = 1$ follows from the two relations. Each element may be expressed uniquely in the form $a^k x^\epsilon$ for $0 \leq k < 2n$ and $0 \leq \epsilon \leq 1$. The group may be realized within $\text{Sp}(1) \subset \mathbb{H}$ by taking $a = e^{\pi i/n}$ and $x = j$. In this realization $a^n = x^2 = -1$ so this element is central.

Lemma A.1. *We have $[D_n^*, D_n^*] = \langle a^2 \rangle$ so that*

$$D_n^*/[D_n^*, D_n^*] \cong \begin{cases} \mathbb{Z}/4 & \text{if } n \equiv 1 \pmod{2} \\ \mathbb{Z}/2 \times \mathbb{Z}/2 & \text{if } n \equiv 0 \pmod{2} \end{cases}$$

Proof. Using the fact that $xa^k = a^{-k}x$ one sees that every commutator is of the form a^k . Moreover, $axa^{-1}x^{-1} = a^2$ from which it follows that $\langle a^2 \rangle \subset [D_n^*, D_n^*]$. As $\langle a^2 \rangle$ is normal with quotient of order 4 and thus abelian, we must have equality. If n is odd x has order 4 modulo $\langle a^2 \rangle$, so the quotient is cyclic. If n is even every element has order 2 mod $\langle a^2 \rangle$, so the quotient is isomorphic to $\mathbb{Z}/2 \times \mathbb{Z}/2$. \square

Lemma A.2. *The conjugacy classes of D_n^* are given by $\{1\}, \{-1\}, \{a^j, a^{-j}\}$ for $1 \leq j < n$,*

$$\{x, a^2x, \dots, xa^{2n-2}x\} \quad \text{and} \quad \{ax, a^3x, \dots, a^{2n-1}x\}.$$

Proof. The elements $1, -1$ are central so $\{1\}, \{-1\}$ are conjugacy classes. From the calculation $(a^s x)a^j(a^s x)^{-1} = a^{-j}$, valid for any s and j , it follows that $\{a^j, a^{-j}\}$, for $1 \leq j < n$, are conjugacy classes. Next, $a^j x a^{-j} = a^{2j}x$ from which we deduce that $\{x, a^2x, \dots, a^{2n-2}x\}$ is a conjugacy class. Finally, from the relation $a^j(ax)a^{-j} = a^{2j+1}x$ we deduce that $\{ax, a^3x, \dots, a^{2n-1}x\}$ is at least a portion of a conjugacy class. However, since we have already exhausted every element of the group, we conclude that this must in fact be the whole conjugacy class and that we have found all the conjugacy classes. \square

Proposition A.3. *The irreducible representations of D_n^* consist of the four 1-dimensional representations associated with the four distinct homomorphisms $D_n^*/[D_n^*, D_n^*] \rightarrow \{\pm 1, \pm i\}$ and the 2-dimensional representations $\tau_k: D_n^* \rightarrow \text{GL}_2(\mathbb{C})$, for $1 \leq k < n$, given by*

$$\tau_k(a) = \begin{pmatrix} \xi^k & 0 \\ 0 & \xi^{-k} \end{pmatrix} \quad \text{and} \quad \tau_k(x) = \begin{pmatrix} 0 & (-1)^k \\ 1 & 0 \end{pmatrix}$$

where $\xi = e^{\pi i/n}$.


Proof. The statement about the 1-dimensional representations is clear. Let χ_k denote the character of τ_k . Then $\chi_k(a^j) = 2 \cos(\pi j k/n)$, $\chi_k(1) = 2$, $\chi_k(-1) = 2(-1)^k$ and χ_k vanishes on the remaining conjugacy classes. Hence,

$$\begin{aligned} \|\chi_k\|^2 &= \frac{1}{4n} \left(8 + 2 \sum_{j=1}^{n-1} 4 \cos^2(\pi j k/n) \right) = \frac{1}{4n} \left(8 + 4 \sum_{j=1}^{n-1} (\cos(2\pi j k/n) + 1) \right) \\ &= \frac{1}{4n} (8 + 4(n-2)) = 1, \end{aligned}$$

which shows that each τ_k is irreducible. The representations τ_k are inequivalent for $1 \leq k < n$ because the values $\chi_k(a) = 2 \cos(\pi k/n)$ are distinct for $1 \leq k < n$. The sum of the squares of the dimensions of the irreducible representations found is $4 + (n-1)4 = 4n = |D_n^*|$, so conclude that we have found all of them. \square

The representation corresponding to the inclusion $D_n^* \subset \text{Sp}(1) = \text{SU}(2)$ is τ_1 . We denote the four representations corresponding to the homomorphisms $(D_n^*)^{ab} \rightarrow \{\pm 1, \pm i\}$ by ρ_j for $0 \leq j \leq 3$, where we take ρ_0 to be the trivial representation and fix ρ_1 by the requirement $\tau_1 \otimes \tau_1 = \tau_2 \oplus \rho_1 \oplus \rho_0$. Then ρ_0, ρ_1 are of real type, while if n is odd $\rho_2 \cong \rho_3^*, \rho_3$ are of complex type and if n is even ρ_2 and ρ_3 are of real type.

The McKay graph $\overline{\Delta}_{D_n^*} \cong \tilde{D}_{n+2}$ takes the form



where we have included the dimensions of the representations in the graph. For D_n^* there are $n-1$ vertices in the middle corresponding to $\tau_1, \dots, \tau_{n-1}$ from left to right. On the left hand side we have ρ_0, ρ_1 and on the right hand side we have ρ_2, ρ_3 .


Lemma A.4. *The representation τ_k is quaternionic for k odd and real for k even.*

Proof. We use the criterion of Lemma 3.5. Since $\sum_{j=0}^{2n-1} \chi_k(a^{2j}) = 0$ and $(xa^j)^2 = x^2 = -1$ for each j , we obtain $\sum_{g \in D_n^*} \chi_k(g^2) = 4n(-1)^k$ and the result follows. \square

Let $\theta = 2\rho_0$ and $\alpha_k = \tau_{2k-1}$ for $1 \leq k \leq n/2$. For n even the 1-dimensional quaternionic representations are given by

$$\theta, \eta_1 = 2\rho_1, \eta_2 = 2\rho_2, \eta_3 = 2\rho_3, \alpha_1, \dots, \alpha_{n/2}$$


and the quotient graph $\overline{\Delta}_{D_n^*}/(\iota)$ is given by



For n odd the 1-dimensional quaternionic representations are given by

$$\theta, \eta = 2\rho_1, \alpha_1, \dots, \alpha_{(n-1)/2}, \lambda = \rho_2 \oplus \rho_3$$

and the corresponding quotient graph is given by



A.3. The Binary Tetrahedral Group. Let $Q = \{\pm 1, \pm i, \pm j, \pm k\} \subset \text{Sp}(1)$ be the quaternion group (note that $Q \cong D_2^*$). Then T^* may be realized in $\text{Sp}(1) \subset \mathbb{H}$ as

$$T^* = Q \cup \left\{ \frac{1}{2}(\epsilon_0 1 + \epsilon_1 i + \epsilon_2 j + \epsilon_3 k) : \epsilon_r = \pm 1, 0 \leq r \leq 3 \right\}.$$

We have $[T^*, T^*] = Q$ so as $|T^*| = 24$, it follows that $(T^*)^{ab} = T^*/[T^*, T^*] \cong \mathbb{Z}/3$.


The character table of T^* is given below. Here $\xi = e^{2\pi i/3}$ is a primitive third root of unity.

	1	2	3a	3b	4	6a	6b	type
ρ_1	1	1	1	1	1	1	1	\mathbb{R}
ρ_2	1	1	ξ	ξ^2	1	ξ	ξ^2	\mathbb{C}
ρ_2^*	1	1	ξ^2	ξ	1	ξ^2	ξ	\mathbb{C}
ρ_3	2	-2	$-\xi$	$-\xi^2$	0	ξ	ξ^2	\mathbb{C}
ρ_3^*	2	-2	$-\xi^2$	$-\xi$	0	ξ^2	ξ	\mathbb{C}
ρ_4	2	-2	-1	-1	0	1	1	\mathbb{H}
ρ_5	3	3	0	0	-1	0	0	\mathbb{R}

The type of the representation in the right hand column is calculated using Lemma 3.5 and the following table. The first row gives a representative for each conjugacy class and the second row gives the conjugacy class in which the square of this element belongs. Here $x = (1+i+j+k)/2 \in T^*$.

	1	2	3a	3b	4	6a	6b
rep	1	-1	$-x$	$-x^*$	i	x	x^*
sm	1	1	$3b$	$3a$	2	$3b$	$3a$

The canonical representation given by $T^* \hookrightarrow \text{SU}(2)$ corresponds to the irreducible character ρ_4 . From this one may calculate the McKay graph $\overline{\Delta}_{T^*} \cong \tilde{E}_6$ to be



From the type calculation given in the character table and Proposition 3.6, we see that the 1-dimensional quaternionic representations are $\theta = 2\rho_1$, $\alpha = \rho_4$ and $\lambda = \rho_2 \oplus \rho_2^*$. The quotient graph $\overline{\Delta}_{T^*}/(\iota)$ then takes the form



A.4. The Binary Octahedral Group. Let $Q \subset T^* \subset \text{Sp}(1)$ be as in the previous section. Then O^* may be realized as

$$O^* = T^* \cup \{(u+v)/\sqrt{2} : u, v \in Q, u \neq \pm v\} \subset \text{Sp}(1).$$

We have $[O^*, O^*] = T^*$, so as $|O^*| = 48$ it follows that $(O^*)^{ab} = O^*/[O^*, O^*] \cong \mathbb{Z}/2$.


The character table of O^* is given below

	1	2	3	4a	4b	6	8a	8b	type
ρ_1	1	1	1	1	1	1	1	1	\mathbb{R}
ρ_2	1	1	1	1	-1	1	-1	-1	\mathbb{R}
ρ_3	2	2	-1	2	0	-1	0	0	\mathbb{R}
ρ_4	2	-2	-1	0	0	1	$\sqrt{2}$	$-\sqrt{2}$	\mathbb{H}
ρ_5	2	-2	-1	0	0	1	$-\sqrt{2}$	$\sqrt{2}$	\mathbb{H}
ρ_6	3	3	0	-1	-1	0	1	1	\mathbb{R}
ρ_7	3	3	0	-1	1	0	-1	-1	\mathbb{R}
ρ_8	4	-4	1	0	0	-1	0	0	\mathbb{H}


Let $x = (1 + i + j + k)/2$, as in the previous section, let $y = (1 + i)/\sqrt{2}$ and let $z = (i + j)/\sqrt{2}$. The following table contains the information needed for the type calculation given in the above right hand column.

	1	2	3	4a	4b	6	8a	8b
rep	1	-1	-x	i	z	x	y	-y
sm	1	1	3	2	2	3	6	6

The canonical representation given by the inclusion $O^* \hookrightarrow \text{SU}(2)$ corresponds to ρ_4 . From this one obtains the McKay graph $\bar{\Delta}_{O^*} \cong \tilde{E}_7$.



Using the by now standard method one finds the 1-dimensional quaternionic representation to be $\theta = 2\rho_1$, $\alpha = \rho_4$, $\beta = \rho_5$ and $\eta = 2\rho_2$. In this case there are no irreducible representations of complex type, so the quotient graph is simply



A.5. The Binary Icosahedral Group. Let $\phi = (1 + \sqrt{5})/2 = 2\cos(\pi/5)$ be the golden ratio (thus $\phi^2 = \phi + 1$) and let S be the set of quaternions obtained by even coordinate permutations of

$$(\epsilon_0 i + \epsilon_1 \phi^{-1} j + \epsilon_2 \phi k)/2$$

where $\epsilon_r \in \{\pm 1\}$ for $0 \leq r \leq 2$. Note that there are $12 = |A_4|$ even permutations and $8 = 2^3$ sign combinations so $|S| = 12 \cdot 8 = 96$. The binary icosahedral group may be realized in $\text{Sp}(1)$ as $I^* = T^* \cup S$. It is well-known that I^* is a perfect group, that is, $[I^*, I^*] = I^*$ or equivalently $(I^*)^{ab} = 0$. The binary polyhedral space $Y_{I^*} = S^3/I^*$ is also called the Poincaré sphere and is an integral homology sphere.

The character table of I^* is given below.

	1	2	3	4	5a	5b	6	10a	10b	type
ρ_1	1	1	1	1	1	1	1	1	1	\mathbb{R}
ρ_2	2	-2	-1	0	ϕ^{-1}	$-\phi$	1	ϕ	$-\phi^{-1}$	\mathbb{H}
ρ_3	2	-2	-1	0	$-\phi$	ϕ^{-1}	1	$-\phi^{-1}$	ϕ	\mathbb{H}
ρ_4	3	3	0	-1	$-\phi^{-1}$	ϕ	0	ϕ	$-\phi^{-1}$	\mathbb{R}
ρ_5	3	3	0	-1	ϕ	$-\phi^{-1}$	0	$-\phi^{-1}$	ϕ	\mathbb{R}
ρ_6	4	4	1	0	-1	-1	1	-1	-1	\mathbb{R}
ρ_7	4	-4	1	0	-1	-1	-1	1	1	\mathbb{H}
ρ_8	5	5	-1	1	0	0	-1	0	0	\mathbb{R}
ρ_9	6	-6	0	0	1	1	0	-1	-1	\mathbb{H}


Put

$$u = (\phi \cdot 1 + \phi^{-1} \cdot i + j)/2 \in S$$


and let $x \in T^*$ be defined as earlier. We then have the following table for the type calculation in the above right hand column.

	1	2	3	4	5a	5b	6	10a	10b
rep	1	-1	-x	i	u ²	-u	x	u	-u ²
sm	1	1	3	2	5b	5a	3	5a	5b

The character corresponding to the canonical representation $I^* \hookrightarrow \mathrm{Sp}(1) = \mathrm{SU}(2)$ is ρ_2 . From this one may show that the McKay graph $\overline{\Delta}_{I^*} \cong \widetilde{E}_8$ is given by



The 1-dimensional quaternionic representations are $\theta = 2\rho_1$, $\alpha = \rho_2$ and $\beta = \rho_3$. Once again there are no irreducible representations of complex type, so the quotient graph is simply



APPENDIX B. EQUIVARIANT $\mathrm{SU}(2)$ -BUNDLES OVER S^4 AND THE CHERN-SIMONS INVARIANT

In this appendix we give a proof of the classification of Γ -equivariant $\mathrm{SU}(2)$ -bundles over S^4 and apply the equivariant index theorem to the twisted Dirac operators $D_A: \Gamma(S^+ \otimes E) \rightarrow \Gamma(S^- \otimes E)$ to obtain a proof of the equations in Lemma 4.4. As a byproduct of this we obtain a simple way to calculate the Chern-Simons invariants of the flat connections in the trivial $\mathrm{SU}(2)$ -bundle over the binary polyhedral spaces. In the final part we also show that this invariant can in a natural way be related to the algebraic Chern class of the holonomy representation associated with the flat connection.

B.1. Conventions on Quaternions. It will be convenient to employ quaternions in the the upcoming theory, so in this section we fix our conventions. We will regard \mathbb{H} as the 4-dimensional real algebra with standard basis $(1, i, j, k)$ and multiplication rules

$$i^2 = j^2 = k^2 = -1 \text{ and } ij = -ji = k.$$

The involution determined by $1, i, j, k \mapsto 1, -i, -j, -k$ is written as $q \mapsto q^*$. Our convention is that a quaternionic vector space is a left \mathbb{H} -module and that the standard quaternionic structure on \mathbb{H}^n is given by left multiplication. We write $M_n(F)$ for the algebra of $n \times n$ matrices with entries in $F = \mathbb{R}, \mathbb{C}, \mathbb{H}$. The standard identification

$$M_n(\mathbb{H}) \cong \mathrm{End}_{\mathbb{H}}(\mathbb{H}^n)$$

is then given by $A \mapsto (x \mapsto xA^*)$, that is, we multiply the conjugate transpose of A with x , regarded as a row vector, from the right. To avoid confusion we will always write $q \cdot x := xq^*$ for the left action of $q \in \mathrm{Sp}(1)$ on $x \in \mathbb{H}$ and always omit the dot when we mean actual multiplication.

We define the standard inclusion by $\mathbb{C} = \mathrm{Span}_{\mathbb{R}}(1, i) \subset \mathbb{H}$ and the standard identification $\mathbb{C}^2 \cong \mathbb{H}$ by $(z, w) \mapsto z + wj$. Note that the complex orientation on \mathbb{C}^2 corresponds to the standard orientation on $\mathbb{H} \cong \mathbb{R}^4$ in which $(1, i, j, k)$ is a positive basis.

With these conventions in place we fix the isomorphism $\mathrm{Sp}(1) \cong \mathrm{SU}(2)$ by requiring the following diagram to commute

$$\begin{array}{ccccc} \mathrm{Sp}(1) & \xrightarrow{\subset} & \mathbb{H} & \xrightarrow{\cong} & \mathrm{End}_{\mathbb{H}}(\mathbb{H}) \\ \downarrow \cong & & \downarrow & & \downarrow \subset \\ \mathrm{SU}(2) & \xrightarrow{\subset} & M_2(\mathbb{C}) & \xrightarrow{\cong} & \mathrm{End}_{\mathbb{C}}(\mathbb{C}^2). \end{array}$$

Explicitly,

$$q = z + wj \in \mathrm{Sp}(1) \mapsto \begin{pmatrix} z^* & w^* \\ -w & z \end{pmatrix}.$$

These conventions ensure that the standard left action $\mathrm{SU}(2) \times \mathbb{C}^2 \rightarrow \mathbb{C}^2$ corresponds to the above defined left action $\mathrm{Sp}(1) \times \mathbb{H} \rightarrow \mathbb{H}$. We define the orientation of $\mathrm{Sp}(1) = \mathrm{SU}(2)$ by requiring that $(i, j, k) \in \mathrm{Im} \mathbb{H} = \mathfrak{sp}(1)$ is a positive basis. The orbit map $\mathrm{Sp}(1) \rightarrow S^3 \subset \mathbb{H}$ given by $q \mapsto q \cdot 1 = 1q^* = q^*$ is orientation reversing. Since our work in this appendix is very sensitive to orientations, we will for this reason distinguish between the group $\mathrm{Sp}(1) = \mathrm{SU}(2)$ and the unit sphere in the representation space $S^3 \subset \mathbb{H} = \mathbb{C}^2$.

B.2. Classification of Γ -equivariant $\mathrm{SU}(2)$ -bundles over S^4 . Let $\Gamma \subset \mathrm{Sp}(1) = \mathrm{SU}(2)$ be a finite subgroup. Recall that Γ acts on S^3 by restriction of the linear action on $\mathbb{H} = \mathbb{C}^2$. By regarding $S^4 \subset \mathbb{H} \oplus \mathbb{R}$ the suspended action $\Gamma \times S^4 \rightarrow S^4$ takes the form $\gamma \cdot (x, t) = (\gamma \cdot x, t)$. The fixed points of this action are $N := (0, 1)$ and $S := (0, -1)$.

Let $\mathrm{Rep}^1(\Gamma, \mathbb{H})$ denote the set of isomorphism classes of 1-dimensional quaternionic representations of Γ and let $\mathrm{Vec}_\Gamma^1(S^4, \mathbb{H})$ denote the set of isomorphism classes of Γ -equivariant quaternionic line bundles over S^4 . If V is a 1-dimensional quaternionic representation of Γ we write $[V]$ for its isomorphism class in $\mathrm{Rep}^1(\Gamma, \mathbb{H})$. The same applies to Γ -equivariant bundles and $\mathrm{Vec}_\Gamma^1(S^4, \mathbb{H})$. Define

$$\phi: \mathrm{Vec}_\Gamma^1(S^4, \mathbb{H}) \rightarrow \mathrm{Rep}^1(\Gamma, \mathbb{H}) \times \mathrm{Rep}^1(\Gamma, \mathbb{H}) \times \mathbb{Z}$$

by $[E] \mapsto ([E_N], [E_S], c_2(E)[S^4])$. Our aim is to show that this map is injective and identify the image. We will need the following two well-known results for our work.

Lemma B.1. *Let Y be a closed 3-manifold. Then every $\mathrm{Sp}(1) = \mathrm{SU}(2)$ -bundle over Y is trivial.*

Proof. Every closed 3-manifold admits the structure of a finite CW complex with cells of dimension ≤ 3 . We have $B\mathrm{Sp}(1) = \mathbb{H}P^\infty$ and this space has a CW structure with a single cell in each dimension $n = 4k$, $k \geq 0$. By cellular approximation every map $f: Y \rightarrow \mathbb{H}P^\infty$ is homotopic to a map with image in the 3-skeleton $(\mathbb{H}P^\infty)^3 = *$. This implies that every $\mathrm{Sp}(1)$ -bundle must be trivial. \square

Lemma B.2. *Let Y be a closed oriented 3-manifold and let $f, g: Y \rightarrow \mathrm{Sp}(1)$ be a pair of continuous maps. Define $h: Y \rightarrow \mathrm{Sp}(1)$ by $h(y) = f(y)g(y)$ for $y \in Y$. Then*

$$\deg(h) = \deg(f) + \deg(g).$$

Proof. The map h is given by the following composition

$$Y \xrightarrow{\Delta} Y \times Y \xrightarrow{f \times g} \mathrm{Sp}(1) \times \mathrm{Sp}(1) \xrightarrow{\mu} \mathrm{Sp}(1),$$

where Δ is the diagonal map and μ is the multiplication map. Let $\pi_i: \mathrm{Sp}(1) \times \mathrm{Sp}(1) \rightarrow \mathrm{Sp}(1)$ for $i = 1, 2$ be the two projections and let $\iota_i: \mathrm{Sp}(1) \rightarrow \mathrm{Sp}(1) \times \mathrm{Sp}(1)$ for $i = 1, 2$ the two inclusions given by $\iota_1(a) = (a, 1)$ and $\iota_2(a) = (1, a)$ for $a \in \mathrm{Sp}(1)$. Then $\pi_i \circ \iota_i = \mathrm{id}$ for $i = 1, 2$ and $\pi_i \circ \iota_j$ is constant for $i \neq j$. Clearly, $H_3(\mathrm{Sp}(1) \times \mathrm{Sp}(1)) \cong \mathbb{Z} \oplus \mathbb{Z}$ and from this we deduce that

$$q := ((\pi_1)_*, (\pi_2)_*): H_3(\mathrm{Sp}(1) \times \mathrm{Sp}(1)) \rightarrow H_3(\mathrm{Sp}(1)) \oplus H_3(\mathrm{Sp}(1))$$

is an isomorphism with inverse $(\iota_1)_* + (\iota_2)_*$. Furthermore,

$$\mu_* \circ ((\iota_1)_* + (\iota_2)_*) = (\mu \circ \iota_1)_* + (\mu \circ \iota_2)_* = \mathrm{id}_* + \mathrm{id}_*.$$

These considerations imply that the following diagram commutes

$$\begin{array}{ccccc}
H_3(Y) & \xrightarrow{(f,g)_* \circ \Delta_*} & H_3(\mathrm{Sp}(1) \times \mathrm{Sp}(1)) & \xrightarrow{\mu_*} & H_3(\mathrm{Sp}(1)) \\
& \searrow (f_*, g_*) & \downarrow \cong & \nearrow + & \\
& & H_3(\mathrm{Sp}(1)) \oplus H_3(\mathrm{Sp}(1)) & & .
\end{array}$$

Since the upper composition is $h_*: H_3(Y) \rightarrow H_3(\mathrm{Sp}(1))$ by definition, it follows that

$$\deg(h)[\mathrm{Sp}(1)] = h_*([Y]) = f_*([Y]) + g_*([Y]) = (\deg(f) + \deg(g))[\mathrm{Sp}(1)],$$

where $[Y] \in H_3(Y)$ and $[\mathrm{Sp}(1)] \in H_3(\mathrm{Sp}(1))$ are the fundamental classes. Hence, $\deg(h) = \deg(f) + \deg(g)$ as required. \square

Let $\alpha \in \mathrm{Rep}^1(\Gamma, \mathbb{H})$ and choose a homomorphism $\rho_\alpha: \Gamma \rightarrow \mathrm{Sp}(1)$ representing α . Write $\mathrm{Sp}(1)^\alpha$ for the group $\mathrm{Sp}(1)$ equipped with the Γ -action given by $\gamma \cdot q = \rho_\alpha(\gamma)q$. If X and Y are Γ -manifolds we write $[X, Y]_\Gamma$ for the set of Γ -equivariant homotopy classes of equivariant maps.

Lemma B.3. *There exist Γ -equivariant maps $S^3 \rightarrow \mathrm{Sp}(1)^\alpha$. Moreover, for any pair g, g' of such maps it holds true that $\deg g \equiv \deg g' \pmod{|\Gamma|}$.*

Proof. Since Γ acts freely on S^3 the set of equivariant maps $S^3 \rightarrow \mathrm{Sp}(1)^\alpha$ is in natural bijection with the set of sections of the associated principal $\mathrm{Sp}(1)$ -bundle

$$S^3 \times_\Gamma \mathrm{Sp}(1)^\alpha \rightarrow S^3/\Gamma.$$

According to Lemma B.1 this bundle must be trivial, and hence admits global sections. This proves the first assertion.

For the second assertion assume that $f, g: S^3 \rightarrow \mathrm{Sp}(1)^\alpha$ is a pair of equivariant maps. We may then form the map $h: S^3 \rightarrow \mathrm{Sp}(1)$ given by $h(x) = f(x)^*g(x)$. This map is equivariant when Γ acts trivially on $\mathrm{Sp}(1)$ and therefore descends to a map $\bar{h}: S^3/\Gamma \rightarrow \mathrm{Sp}(1)$. Since the covering map $S^3 \rightarrow S^3/\Gamma$ has degree $|\Gamma|$, we deduce that $\deg h = |\Gamma| \deg(\bar{h}) \equiv 0 \pmod{|\Gamma|}$. Finally, by Lemma B.2 and the fact that $\deg f^* = -\deg f$ we obtain $\deg h = \deg(g) - \deg f$ and the proof is complete. \square

Let $\alpha, \beta \in \mathrm{Rep}^1(\Gamma, \mathbb{H})$ have representatives $\rho_\alpha, \rho_\beta: \Gamma \rightarrow \mathrm{Sp}(1)$. Let $\mathbb{H}(\alpha)$ and $\mathbb{H}(\beta)$ denote \mathbb{H} equipped with the Γ -action determined by ρ_α and ρ_β , respectively. Furthermore, write $\mathrm{Sp}(1)^{(\alpha; \beta)}$ for the group $\mathrm{Sp}(1)$ equipped with the Γ -action $\gamma \cdot q = \rho_\alpha(\gamma)q\rho_\beta(\gamma)^*$.

Proposition B.4. *The set of isomorphism classes of Γ -equivariant $\mathrm{Sp}(1)$ -bundles $E \rightarrow S^4$ with $[E_N] = \alpha$ and $[E_S] = \beta$ is in natural bijection with*

$$[S^3, \mathrm{Sp}(1)^{(\alpha; \beta)}]_\Gamma.$$

Proof. This is well-known in the non-equivariant setting, so we will only include the details necessary to adapt the usual proof to the equivariant setting.

Let E be a Γ -equivariant $\mathrm{Sp}(1)$ -bundle over S^4 (regarded as a vector bundle) with $[E_N] = \alpha$ and $[E_S] = \beta$. Let $U_N = S^4 - \{S\}$ and $U_S = S^4 - \{N\}$. Let A be a Γ -invariant $\mathrm{Sp}(1)$ -connection in E . Then, using parallel transport along radial geodesic from S and N , one obtains equivariant trivializations $U_S \times E_S \cong E|_{U_S}$ and $U_N \times E_N \cong E|_{U_N}$. By making suitable choices of unit basis vectors in E_S and E_N , we may identify $E_S \cong \mathbb{H}(\beta)$ and $E_N \cong \mathbb{H}(\alpha)$. The transition function between these trivializations is an equivariant map $U_N \cap U_S \rightarrow \mathrm{Hom}_{\mathbb{H}}(E_S, E_N)$ taking values in the subset of isometries. This subset may be identified with $\mathrm{Sp}(1)^{(\alpha; \beta)}$ through

$$\mathrm{Sp}(1)^{(\alpha; \beta)} \hookrightarrow \mathrm{Hom}_{\mathbb{H}}(\mathbb{H}(\beta), \mathbb{H}(\alpha)) \cong \mathrm{Hom}_{\mathbb{H}}(E_S, E_N),$$

where we recall that by convention the first map is given by $q \mapsto r_q^*$. Let $t: S^3 \rightarrow S^3(\alpha, \beta)$ denote the restriction of this map to the middle sphere. One may verify that the Γ -equivariant homotopy

class of this map is independent of the trivializations chosen. We have therefore constructed one direction of the equivalence.

The inverse may be defined as follows. Let $f: S^3 \rightarrow \mathrm{Sp}(1)^{(\alpha;\beta)}$ represent a homotopy class in $[S^3, \mathrm{Sp}(1)^{(\alpha;\beta)}]_\Gamma$. Let $p: U_N \cap U_S \rightarrow S^3$ be the equivariant projection onto the middle sphere. Using f we may therefore construct an equivariant bundle E by gluing $U_S \times \mathbb{H}(\beta)$ and $U_N \times \mathbb{H}(\alpha)$ along $\psi: U_N \cap U_S \times \mathbb{H}(\beta) \rightarrow U_S \cap U_N \times \mathbb{H}(\alpha)$ given by $\psi(x, v) = (x, f(p(x)) \cdot v) = (x, v f(p(x))^*)$. The equivariance of f and p ensure that the actions of Γ on $U_N \times \mathbb{H}(\alpha)$ and $U_S \times \mathbb{H}(\beta)$ match over the intersection $U_N \cap U_S$. The resulting bundle is therefore a Γ -equivariant quaternionic line bundle. One may verify that the isomorphism class of this bundle is independent of the representative f chosen using the equivariant bundle homotopy theorem of [Seg68, Prop. 1.3].

The verification of the fact that these two constructions are mutual inverses proceeds just as in the non-equivariant case. \square

Recall that we regard $S^4 \subset \mathbb{H} \oplus \mathbb{R}$. Let $u: S^4 - \{0\} \cong \mathbb{H}$ be stereographic projection from the north pole. We give \mathbb{H} the standard orientation and orient S^4 by requiring u to be orientation preserving. Moreover, we orient the middle sphere $S^3 \subset S^4$ by requiring that the restriction $u|_{S^3}: S^3 \cong S^3 \subset \mathbb{H}$ preserves orientation.

Proposition B.5. *Let $\alpha, \beta \in \mathrm{Rep}^1(\Gamma, \mathbb{H})$ have representatives $\rho_\alpha, \rho_\beta: \Gamma \rightarrow \mathrm{Sp}(1)$. Then the natural map*

$$[S^3, \mathrm{Sp}(1)^{(\alpha;\beta)}]_\Gamma \rightarrow [S^3, \mathrm{Sp}(1)^{(\alpha;\beta)}]$$

is injective. In other words, two equivariant maps $f, g: S^3 \rightarrow \mathrm{Sp}(1)^{(\alpha;\beta)}$ are homotopic if and only if they are equivariantly homotopic. Moreover, if we identify $[S^3, \mathrm{Sp}(1)^{(\alpha;\beta)}] \cong \mathbb{Z}$ through $[f] \mapsto \deg(f)$, there is a constant $c \in \mathbb{Z}$ so that the image is $\{n \in \mathbb{Z} : n \equiv c \pmod{|\Gamma|}\}$.

Proof. By Lemma B.3 we can find equivariant maps $g: S^3 \rightarrow \mathrm{Sp}(1)^\alpha$ and $h: S^3 \rightarrow \mathrm{Sp}(1)^\beta$. Given an equivariant map $f: S^3 \rightarrow \mathrm{Sp}(1)^{(\alpha;\beta)}$ define $\tau(f) = \tau_{g,h}(f): S^3 \rightarrow \mathrm{Sp}(1)^\theta$ to be the composition

$$S^3 \xrightarrow{(h,f,g)} \mathrm{Sp}(1)^\alpha \times \mathrm{Sp}(1)^{(\alpha;\beta)} \times \mathrm{Sp}(1)^\beta \xrightarrow{\mu} \mathrm{Sp}(1)^\theta,$$

where $\mathrm{Sp}(1)^\theta$ denotes $\mathrm{Sp}(1)$ equipped with the trivial Γ -action and μ is the equivariant map $(x, y, z) \mapsto x^* y z$. Observe that if $f': S^3 \rightarrow \mathrm{Sp}(1)^{(\alpha;\beta)}$ is another equivariant map, then f and f' are equivariantly homotopic if and only if $\tau(f)$ and $\tau(f')$ are equivariantly homotopic. Indeed, if $f_t: S^3 \rightarrow \mathrm{Sp}(1)^{(\alpha;\beta)}$ is an equivariant homotopy between f and f' , then $\tau(f_t)$ is an equivariant homotopy between $\tau(f)$ and $\tau(f')$ and the same argument applies for the converse using $\tau_{g^{-1}, h^{-1}}$ in place of $\tau = \tau_{g,h}$.

Since the Γ -action on $\mathrm{Sp}(1)^\theta$ is trivial, we obtain a bijection

$$[S^3, \mathrm{Sp}(1)^\theta]_\Gamma \cong [S^3/\Gamma, \mathrm{Sp}(1)] \cong \mathbb{Z},$$

where we give S^3/Γ the orientation induced from S^3 and the second isomorphism is given by $[u] \mapsto \deg(u)$. Given an equivariant map $f: S^3 \rightarrow \mathrm{Sp}(1)^{(\alpha;\beta)}$ let $\overline{\tau(f)}: S^3/\Gamma \rightarrow \mathrm{Sp}(1)$ denote the map induced by $\tau(f)$. The above work then amounts to the fact that the map $[S^3, \mathrm{Sp}(1)^{(\alpha;\beta)}]_\Gamma \rightarrow \mathbb{Z}$ given by $f \mapsto \deg(\overline{\tau(f)})$ is a bijection. To complete the proof we will relate this degree to the degree of f . First, since $S^3 \rightarrow S^3/\Gamma$ has degree $|\Gamma|$, we deduce that $|\Gamma| \deg(\overline{\tau(f)}) = \deg \tau(f)$. Using Lemma B.2 and the definition $\tau(f) = \mu \circ (g, f, h)$ one finds that

$$\deg(\tau(f)) = \deg h - \deg g + \deg(f).$$

Hence, $\deg(f) = (\deg g - \deg h) + |\Gamma| \deg(\overline{\tau(f)})$. We therefore have the following commutative diagram

$$(B.1) \quad \begin{array}{ccc} [S^3, \mathrm{Sp}(1)^{(\alpha; \beta)}]_{\Gamma} & \longrightarrow & [S^3, \mathrm{Sp}(1)^{(\alpha; \beta)}] \\ \downarrow \cong & & \downarrow \cong \\ \mathbb{Z} & \xrightarrow{m} & \mathbb{Z}, \end{array}$$

where the vertical maps are given by $[f] \mapsto \deg(\overline{\tau(f)})$ and $[f] \mapsto \deg f$, from left to right, and $m(n) = |\Gamma|n + (\deg g - \deg h)$. Since m is injective we conclude that the upper horizontal map is injective. The final assertion follows from the fact that $\deg g - \deg h$ is independent of g and h by Lemma B.3. \square

Theorem B.6. *Let $\Gamma \subset \mathrm{Sp}(1)$ be a finite subgroup acting on $S^4 \subset \mathbb{H} \oplus R$ by $\gamma \cdot (x, t) = (\gamma \cdot x, t) = (x\gamma^*, t)$. Then the map*

$$\phi: \mathrm{Vec}_{\Gamma}^1(S^4, \mathbb{H}) \rightarrow \mathrm{Rep}^1(\Gamma, \mathbb{H}) \times \mathrm{Rep}^1(\Gamma, \mathbb{H}) \times \mathbb{Z}$$

given by $[E] \mapsto ([E_N], [E_S], c_2(E)[S^4])$ is injective. Furthermore, for each pair $\alpha, \beta \in \mathrm{Rep}^1(\Gamma, \mathbb{H})$ there is a constant $c \in \mathbb{Z}$ such that $(\alpha, \beta, k) \in \mathrm{Im} \phi$ if and only if $k \equiv c \pmod{|\Gamma|}$.

Proof. The follows from Proposition B.4 and Proposition B.5 in view of the fact that $c_2(E)[S^4]$ coincides with the degree of the transition function $t: S^3 \rightarrow \mathrm{Sp}(1) \subset \mathrm{Hom}_{\mathbb{H}}(E_S, E_N)$. \square

The proof of Proposition B.5 gives a formula for the constant c that will be useful later.

Corollary B.7. *The constant c in the above theorem may be taken to be $c = \deg g - \deg h$ for any choices of equivariant maps $g: S^3 \rightarrow \mathrm{Sp}(1)^{\alpha}$ and $h: S^3 \rightarrow \mathrm{Sp}(1)^{\beta}$.*

B.3. Application of the Equivariant Index Theorem. Let $E \rightarrow S^4$ be a Γ -equivariant $\mathrm{SU}(2)$ bundle and let S^+ and S^- be the complex spinor bundles associated with the unique spin structure on S^4 . It turns out that this spin structure is naturally Γ -equivariant. This means that the spinor bundles carry the structure of Γ -equivariant bundles and that the Clifford multiplication map $TS^4 \rightarrow \mathrm{Hom}_{\mathbb{C}}(S^+, S^-)$ is a map of equivariant bundles. Therefore, if A is a Γ -invariant connection in E the twisted Dirac operator

$$D_A: \Gamma(S^+ \otimes E) \rightarrow \Gamma(S^- \otimes E)$$

is a Γ -equivariant elliptic operator. The non-equivariant index of such an operator takes values in the K -theory of a point $K(*) = \mathbb{Z}$. In the Γ -equivariant setting it takes values in the Γ -equivariant K -theory of a point, namely, $K_{\Gamma}(*) = R(\Gamma)$; the complex representation ring of Γ . It is this index we will calculate in this section.

For this purpose it is more convenient to use the model $\mathbb{H}P^1$ for S^4 . This is the identification space $\mathbb{H}^2 - \{0\} / \sim$ where $(x, y) \sim (qx, qy)$ for $q \in \mathbb{H} - \{0\}$. The equivalence class of (x, y) in $\mathbb{H}P^1$ is denoted by $[x : y]$. There are two canonical charts

$$\mathbb{H} \cong U = \{[x : y] \in \mathbb{H}P^1 : x \neq 0\} \quad \text{and} \quad \mathbb{H} \cong V = \{[x : y] \in \mathbb{H}P^1 : y \neq 0\}$$

given by $z \mapsto [1 : z]$ and $w \mapsto [w : 1]$, respectively. The transition function $\mathbb{H} - \{0\} \rightarrow \mathbb{H} - \{0\}$ (in either direction) is given by $q \mapsto q^{-1}$. This map is orientation preserving so we define the orientation of $\mathbb{H}P^1$ by requiring that both of the canonical charts are positive, where $\mathbb{H} = \mathbb{C}^2$ has the standard orientation.

The tautological quaternionic line bundle $\gamma \rightarrow \mathbb{H}P^1$ is defined to be

$$\gamma := \{([x : y], (z, w)) \in \mathbb{H}P^1 \times \mathbb{H}^2 : \exists q \in \mathbb{H} - \{0\} \ni (z, w) = (qx, qy)\}.$$

Let the trivial bundle $\underline{\mathbb{H}}^2 = \mathbb{H}P^1 \times \mathbb{H}^2$ carry the standard symplectic inner product, i.e., $(z_1, z_2) \cdot (w_1, w_2) = z_1 w_1^* + z_2 w_2^*$. Then $\tilde{\gamma} := \gamma^{\perp}$ is another quaternionic line bundle. We give γ and $\tilde{\gamma}$ the

connections obtained from the product connection in \mathbb{H}^2 by orthogonal projection and equip them with the symplectic inner products inherited from the trivial bundle \mathbb{H}^2 .

The standard action $\mathrm{Sp}(2) \times \mathbb{H}^2 \rightarrow \mathbb{H}^2$ descends to a transitive action $\mathrm{Sp}(2) \times \mathbb{H}P^1 \rightarrow \mathbb{H}P^1$. The diagonal action on $\mathbb{H}P^1 \times \mathbb{H}^2$ preserves the subbundles $\gamma, \tilde{\gamma} \subset \mathbb{H}^2$ and therefore give these the structure of $\mathrm{Sp}(2)$ -equivariant bundles over $\mathbb{H}P^1$. The connections are invariant under this action.

The unique spin structure on $\mathbb{H}P^1$ may now be realized explicitly with $S^+ = \gamma$, $S^- = \tilde{\gamma}$ and Clifford multiplication $\chi: T\mathbb{H}P^1 \cong \mathrm{Hom}_{\mathbb{H}}(\gamma, \tilde{\gamma})$ defined in the following way. Let $\iota: \gamma \rightarrow \mathbb{H}^2$ denote the inclusion and $\pi: \mathbb{H}^2 \rightarrow \tilde{\gamma}$ the projection. Given $p \in \mathbb{H}P^1$, $v \in T_p\mathbb{H}P^1$ and $w \in \gamma_p$ choose a local smooth section s of γ with $s(p) = w$. Then $\chi(v)(w) = \pi(d(\iota(s)))_p$, where d is the product connection in \mathbb{H}^2 . In suitable trivializations of γ and $\tilde{\gamma}$ over $\mathbb{H} \cong U \subset \mathbb{H}P^1$, the map $\chi_q: T_q\mathbb{H} = \mathbb{H} \rightarrow \mathrm{Hom}_{\mathbb{H}}(\mathbb{H}, \mathbb{H})$ for $q \in \mathbb{H}$ takes the form $v \mapsto r_v: \mathbb{H} \rightarrow \mathbb{H}$, where $r_v(w) = vw$ is right multiplication. Finally, we note that this spin structure is $\mathrm{Sp}(2)$ -equivariant.

Remark B.8. With our orientation conventions it holds true that $c_2(\gamma)[\mathbb{H}P^1] = -1$. This can be verified, as suggested in [FU91, p. 85], as follows. First check that the transition function restricted to $S^3 \subset \mathbb{H} \cong U = \mathbb{H}P^1 - \{[0 : 1]\}$ is a diffeomorphism and hence has degree ± 1 . Then explicitly calculate the connection and curvature forms in the chart $\mathbb{H} \cong U$ and conclude that the connection of γ is self-dual. From these two pieces of information the assertion follows. As $\gamma \oplus \tilde{\gamma}$ is trivial it follows that $c_2(\tilde{\gamma})[\mathbb{H}P^1] = 1$ as well.

Let $\Gamma \subset \mathrm{Sp}(1)$ be a finite subgroup. We define a Γ -action on $\mathbb{H}P^1$ by restricting the $\mathrm{Sp}(2)$ -action along $\Gamma \subset \mathrm{Sp}(1) \subset \mathrm{Sp}(2)$, where the latter inclusion is specified by

$$(B.2) \quad q \in \mathrm{Sp}(1) \mapsto \begin{pmatrix} 1 & 0 \\ 0 & q \end{pmatrix}.$$

From the above discussion it follows that the spinor bundles S^{\pm} obtain Γ -equivariant structures and that the spinor connections, are Γ -invariant. By convention the action of $\mathrm{Sp}(2)$ on $\mathbb{H}P^1$ is given by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot [x : y] = [x : y] \begin{pmatrix} a^* & c^* \\ b^* & d^* \end{pmatrix} = [xa^* + yb^* : xc^* + yd^*].$$

From this we see that in the chart $\mathbb{H} \cong U \subset \mathbb{H}P^1$, $z \mapsto [1 : z]$, this action corresponds to the standard linear action $\Gamma \times \mathbb{H} \rightarrow \mathbb{H}$ given by $\gamma \cdot x = x\gamma^*$.

Remark B.9. By regarding $S^4 \subset \mathbb{H} \oplus \mathbb{R}$ as before the map $\psi: \mathbb{H}P^1 \rightarrow S^4$ given by

$$\psi([x : y]) = \frac{1}{\|x\|^2 + \|y\|^2} (2x^*y, \|y\|^2 - \|x\|^2)$$

is a Γ -equivariant, orientation preserving diffeomorphism, when S^4 is given the suspended action and orientation described earlier (note that this is the opposite of the standard orientation on S^4).

Let G be a compact Lie group, X a G -manifold and $P: \Gamma(V) \rightarrow \Gamma(W)$ an elliptic G -operator, that is, V and W are G -equivariant complex vector bundles and P is equivariant with respect the induced actions. Then the G -index of P is defined by

$$\mathrm{Ind}_G(P) = [\mathrm{Ker} P] - [\mathrm{Coker}(P)] \in R(G).$$

The character of this virtual representation is denote by $\mathrm{Ind}_g(P) = \mathrm{tr}(g|_{\mathrm{Ker}(P)}) - \mathrm{tr}(g|_{\mathrm{Coker}(P)}) \in \mathbb{C}$ for each $g \in G$. The fixed-point formula due to Atiyah, Segal and Singer expresses the fact that $\mathrm{Ind}_g(P)$ only depends on information above the fixed point set $F_g := \{x \in X : gx = x\}$. In the case where F_g is a finite set the formula (see [LM89, Theorem 14.3]) simplifies to

$$(B.3) \quad \mathrm{Ind}_g P = \sum_{x \in F_g} \frac{\mathrm{ch}_g(V_x) - \mathrm{ch}_g(W_x)}{\mathrm{ch}_g(\lambda_{-1}(T_x X)_{\mathbb{C}})}.$$

Here, $\text{ch}_g(V_x) = \text{tr}(g: V_x \rightarrow V_x)$, $(T_x X)_{\mathbb{C}}$ is the complexification of $T_x X$ and λ_{-1} is defined for a G representation U by

$$\lambda_{-1}(U) = \sum_{k=0}^{\dim_{\mathbb{C}} U} (-1)^k \Lambda_{\mathbb{C}}^k(U) \in R(G).$$

Let $\Gamma \subset \text{SU}(2) = \text{Sp}(1)$ be a finite subgroup acting on $\mathbb{H}P^1$ in the way described above. This action is free away from the two fixed points $S = [1 : 0]$ and $N = [0 : 1]$. Let $E \rightarrow \mathbb{H}P^1 \cong S^4$ be a Γ -equivariant $\text{Sp}(1)$ -bundle with $[E_S] = \alpha$, $[E_N] = \beta$ and $c_2(E)[\mathbb{H}P^1] = k$. Let A be a Γ -invariant connection in E . Then the twisted Dirac operator $D_A: \Gamma(S^+ \otimes E) \rightarrow \Gamma(S^- \otimes E)$ is an elliptic Γ -operator as described above. For our purpose the formal adjoint $D_A^*: \Gamma(S^- \otimes E) \rightarrow \Gamma(S^+ \otimes E)$ is more relevant.

Remark B.10. Here we regard S^{\pm} and E as complex vector bundles by restriction along $\mathbb{C} \subset \mathbb{H}$. In particular, the tensor products $S^{\pm} \otimes E = S^{\pm} \otimes_{\mathbb{C}} E$ are formed over the complex numbers.

Proposition B.11. *Let $E \rightarrow \mathbb{H}P^1 \cong S^4$ be a Γ -equivariant $\text{SU}(2)$ -bundle with $[E_N], [E_S] = \alpha, \beta \in R(\Gamma)$ and $c_2(E)[\mathbb{H}P^1] = k \in \mathbb{Z}$. Then $\mathcal{H} := \text{Ind}_{\Gamma}(D_A^*) \in R(\Gamma)$ satisfies*

$$(2 - Q)\mathcal{H} = \alpha - \beta,$$

where Q is the 2-dimensional complex representation associated with the inclusion $\Gamma \subset \text{SU}(2)$. Furthermore, if $\epsilon: R(\Gamma) \rightarrow \mathbb{Z}$ denotes the augmentation, then $\epsilon(\mathcal{H}) = k$.

Proof. To apply the formula (B.3) we have to determine the action of Γ on the fibers of $S^{\pm} \otimes E$ over the fixed points $S = [1 : 0]$ and $N = [0 : 1]$ as well as the action on the tangent spaces $T_N \mathbb{H}P^1, T_S \mathbb{H}P^1$. From the definition of $S^+ = \gamma$ and $S^- = \tilde{\gamma}$ we observe that in terms of the inclusions $\gamma, \tilde{\gamma} \subset \underline{\mathbb{H}}^2$

$$\begin{aligned} \gamma_S &= \mathbb{H} \oplus 0 & \gamma_N &= 0 \oplus \mathbb{H} \\ \tilde{\gamma}_S &= 0 \oplus \mathbb{H} & \tilde{\gamma}_N &= \mathbb{H} \oplus 0 \end{aligned}.$$

From the description of the action in (B.2) it follows that $\gamma_S \cong \tilde{\gamma}_N \cong \mathbb{C}^2$, the trivial representation, and $\gamma_N \cong \tilde{\gamma}_S \cong Q$.

Next we need to investigate the action on the tangent spaces. We have already seen that in the chart $\mathbb{H} \cong U \subset \mathbb{H}P^1$ the action corresponds to the standard linear action $\gamma \cdot x = x\gamma^*$. Here $0 \in \mathbb{H}$ corresponds to $S \in \mathbb{H}P^1$. It follows that $T_S \mathbb{H}P^1 \cong Q_{\mathbb{R}}$, that is, the canonical representation regarded as a real representation. In the other chart $\mathbb{H} \cong V \subset \mathbb{H}P^1$ given by $z \mapsto [z : 1]$ one may verify that the action is given by $\gamma \cdot z = \gamma z$. The differentiated action on $\mathbb{H} \cong T_0 \mathbb{H}$ is given by the same formula. The conjugation map $*$: $\mathbb{H} \rightarrow \mathbb{H}$ is real linear and satisfies $(\gamma x)^* = x^* \gamma^*$. Hence, $T \mathbb{H}P_N^1 \cong Q_{\mathbb{R}}$ as well. We need to calculate the term $\lambda_{-1}((Q_{\mathbb{R}})_{\mathbb{C}})$. First, note that $(Q_{\mathbb{R}})_{\mathbb{C}} \cong Q \oplus Q^* \cong 2Q$, since Q is an $\text{SU}(2)$ -representation and therefore self-dual. Thus,

$$\lambda_{-1}((Q_{\mathbb{R}})_{\mathbb{C}}) = \lambda_{-1}(2Q) = (\lambda_{-1}Q)^2 = (\mathbb{C} - Q + \Lambda^2 Q)^2 = (2 - Q)^2,$$

where we have used that $\lambda_{-1}(V \oplus W) = \lambda_{-1}(V)\lambda_{-1}(W)$ and the fact that $\Lambda^2 Q \cong \mathbb{C}$, the trivial representation.

Given an element $V \in R(\Gamma)$ we let $\chi_V: \Gamma \rightarrow \mathbb{C}$ denote the associated character. For any $g \neq 1 \in \Gamma$ the index formula (B.3) now gives $\chi_{\mathcal{H}}(g)$ as

$$\left(\frac{2\chi_{\alpha}(g) - \chi_Q(g)\chi_{\alpha}(g)}{(2 - \chi_Q(g))^2} \right) + \left(\frac{\chi_Q(g)\chi_{\beta}(g) - 2\chi_{\beta}(g)}{(2 - \chi_Q(g))^2} \right) = \frac{\chi_{\alpha}(g) - \chi_{\beta}(g)}{2 - \chi_Q(g)},$$

where the first term corresponds to the fixed point $S = [1 : 0]$ and the second term to $N = [0 : 1]$. This yields the equality of characters $(2 - \chi_Q)\chi_{\mathcal{H}} = \chi_{\alpha} - \chi_{\beta}$, since by the above it holds for all

$g \neq 0$, while for $g = 1$ it is trivially satisfied as $\chi_Q(1) = \chi_\alpha(1) = \chi_\beta(1) = 2$. We therefore obtain the relation $(2 - Q)\mathcal{H} = \alpha - \beta$ in $R(\Gamma)$. The final assertion follows from the regular index theorem

$$\epsilon(\mathcal{H}) = \text{Ind}(D_A^*) = c_2(E)[S^4].$$

□

B.4. The Chern-Simons Invariant. Let Y be a closed oriented 3-manifold and let $E \rightarrow Y$ be the necessarily trivial $\text{Sp}(1) = \text{SU}(2)$ -bundle over Y . The Chern-Simons invariant of a flat connection A in E is the value $\text{cs}(A) \in \mathbb{R}/\mathbb{Z}$, where cs is the Chern-Simons functional defined in equation (1.3). In this section we will explain that the theory of section (4) lead to a simple procedure for the calculation of this invariant when $Y = Y_\Gamma$ for some finite subgroup $\Gamma \subset \text{SU}(2)$.

Equip $\mathfrak{sp}(1) = \text{Im } \mathbb{H}$ with the standard invariant inner product for which i, j, k is an orthonormal basis and write $g(x, y) = -(x, y)$ for its negative. The identification $\mathfrak{sp}(1) \cong \mathfrak{su}(2)$ matches g with the symmetric bilinear form $(x, y) \mapsto \frac{1}{2} \text{tr}(xy)$. In this section we will prefer to work with the group $\text{Sp}(1)$ so we make the following definition (compare equation (1.3)).

Definition B.12. Let X be a manifold. For any $a \in \Omega^1(X, \mathfrak{sp}(1))$ define the Chern-Simons form $\zeta(a) \in \Omega^3(X, \mathbb{R})$ by

$$\zeta(a) = \frac{1}{4\pi^2} [a \wedge_g da + \frac{1}{3} a \wedge_g (a \wedge_{ad} a)].$$

Here \wedge_g and \wedge_{ad} denote the combination of the wedge product and the bilinear maps g and $\text{ad} = [\cdot, \cdot]$, respectively. Note that $f^*(\zeta(a)) = \zeta(f^*(a))$ for any smooth map $f: X \rightarrow Y$ and $a \in \Omega^1(Y, \mathfrak{sp}(1))$.

Let Y be a closed oriented 3-manifold and let $E \rightarrow Y$ be an $\text{Sp}(1)$ -bundle. Fix a global trivialization $E \cong Y \times \text{Sp}(1)$ so that the space of connections \mathcal{A}_E is identified with $\Omega^1(Y, \mathfrak{sp}(1))$. Then for $a \in \Omega^1(Y, \mathfrak{sp}(1))$ we have $\text{cs}(a) = \int_Y \zeta(a) \pmod{\mathbb{Z}}$. Observe that if a is flat, that is, $da + \frac{1}{2} a \wedge_{ad} a = 0$, then

$$(B.4) \quad \zeta(a) = \frac{-1}{24\pi^2} a \wedge_g (a \wedge_{ad} a).$$

Lemma B.13. Let $\theta \in \Omega^1(\text{Sp}(1), \mathfrak{sp}(1))$ be the left Maurer-Cartan form, that is, $\theta_g(v) = dl_{g^{-1}}(v)$ for all $g \in \text{Sp}(1)$ and $v \in T_g \text{Sp}(1)$. Then

$$\int_{\text{Sp}(1)} \zeta(\theta) = 1,$$

when $\text{Sp}(1)$ is oriented by requiring that (i, j, k) is a positive basis for $T_1 \text{Sp}(1) = \mathfrak{sp}(1) = \text{Im } \mathbb{H}$.

Proof. As θ is left invariant and $l_g^* \zeta(\theta) = \zeta(l_g^*(\theta)) = \zeta(\theta)$ for each $g \in \text{Sp}(1)$, it follows that $\zeta(\theta) \in \Omega^3(\text{Sp}(1), \mathbb{R})$ is left invariant. Every left invariant form of top dimension in a compact, connected Lie group is necessarily right invariant as well. Therefore, if we regard $\text{Sp}(1) = S^3 \subset \mathbb{H} = \mathbb{R}^4$, we see that $\zeta(\theta)$ is invariant under the action $\text{Sp}(1) \times \text{Sp}(1) \times S^3 \rightarrow S^3$ given by $(a, b) \cdot x = axb^{-1}$. This action factors through the double covering $\text{Sp}(1) \times \text{Sp}(1) \rightarrow \text{SO}(4)$ where $\text{SO}(4)$ acts in the standard way on $S^3 \subset \mathbb{R}^4$. We conclude that $\zeta(\theta)$ is $\text{SO}(4)$ -invariant.

The Riemannian volume form of S^n equipped with the standard round metric is the restriction of

$$\omega_n = \sum_{i=1}^{n+1} (-1)^{i+1} x_i dx_1 \wedge \cdots \wedge \hat{dx}_i \wedge \cdots \wedge dx_{n+1} \in \Omega^n(\mathbb{R}^{n+1}, \mathbb{R}),$$

where the hat denotes omission. This form is invariant under the standard transitive action of $\text{SO}(n+1)$. Specializing to the case $n = 3$, it follows that there is a constant $c \in \mathbb{R}$ such such that $\zeta(\theta) = c\omega_3$.

We determine the constant c by evaluating at $1 = (1, 0, 0, 0) \in \mathbb{H} = \mathbb{R}^4$. By the formula above we have $(\omega_3)_1 = dx_2 \wedge dx_3 \wedge dx_4$, so that $c = \zeta(\theta)(i, j, k)$. As $\theta_1: \mathfrak{sp}(1) \rightarrow \mathfrak{sp}(1)$ is the identity, we have for $a, b, c \in \mathfrak{sp}(1)$ that

$$\theta \wedge_g (\theta \wedge_{ad} \theta)(a, b, c) = g(a, 2[b, c]) - g(b, 2[a, c]) + g(c, 2[a, b]) = 6g(a, [b, c]),$$

since g is symmetric and $g(b, [a, c]) = g([b, a], c) = -g([a, b], c) = -g(a, [b, c])$. Inserting $(a, b, c) = (i, j, k)$ we find

$$\theta \wedge_g (\theta \wedge_{ad} \theta)(i, j, k) = 6g(i, [j, k]) = 6g(i, 2i) = -12(i, i) = -12.$$

Consequently, using formula (B.4) we obtain $\zeta(\theta)_1(i, j, k) = \frac{1}{2\pi^2}$ and therefore $\zeta(\theta) = \frac{1}{2\pi^2}\omega_3$. The volume of S^3 is given by $\text{Vol}(S^3) = \int_{S^3} \omega = 2\pi^2$, so it follows that $\int_{\text{Sp}(1)} \zeta(\theta) = 1$ as required. \square

In the following we give $Y_\Gamma = S^3/\Gamma$ the orientation induced from the standard orientation on $S^3 \subset \mathbb{H}$.

Lemma B.14. *Let $\alpha \in \text{Rep}^1(\Gamma, \mathbb{H})$ correspond to a flat connection in the $\text{Sp}(1)$ -bundle over Y_Γ . Choose a representative $\rho_\alpha: \Gamma \rightarrow \text{Sp}(1)$ for α and let $\text{Sp}(1)^\alpha$ denote the group $\text{Sp}(1)$ equipped with the Γ -action $\gamma \cdot q = \rho_\alpha(\gamma)q$. Then for any Γ -equivariant map $g: S^3 \rightarrow \text{Sp}(1)^\alpha$ it holds true that*

$$\text{cs}(\alpha) = \deg(g)/|\Gamma| \pmod{\mathbb{Z}}.$$

Proof. Choose a representative $a \in \Omega^1(Y_\Gamma, \mathfrak{sp}(1))$ for the flat connection in a global trivialization of the bundle. Then by definition

$$\text{cs}(\alpha) = \int_{Y_\Gamma} \zeta(a) \pmod{\mathbb{Z}}.$$

Let $p: S^3 \rightarrow Y_\Gamma$ be the quotient map. Since Y_Γ carries the orientation induced from S^3 this map has degree $|\Gamma|$. Let $b = p^*(a) \in \Omega^1(S^3, \mathfrak{sp}(1))$. Since a and b are flat and S^3 is simply connected, there exists a smooth map $g: S^3 \rightarrow \text{Sp}(1)$ such that $b = g^*\theta$, where $\theta \in \Omega^1(\text{Sp}(1), \mathfrak{sp}(1))$ is the left Maurer-Cartan form. Moreover, the map g is unique up to left translation in the sense that if $h: S^3 \rightarrow \text{Sp}(1)$ is another map with $h^*\theta = b$, then $h = l_u \circ g$ for some $u \in \text{Sp}(1)$, where $l_u(v) = uv$ denotes left translation in $\text{Sp}(1)$. For each $\gamma \in \Gamma$ we have

$$\gamma^*(b) = \gamma^*p^*(a) = (p \circ \gamma)^*a = p^*(a) = b.$$

Then as $(g \circ \gamma)^*(\theta) = \gamma^*b = b$ we deduce from the uniqueness of g that there is a unique $\rho(\gamma) \in \text{Sp}(1)$ such that $g \circ \gamma = l_{\rho(\gamma)} \circ g$. One may easily verify that $\rho: \Gamma \rightarrow \text{Sp}(1)$ is a homomorphism, in fact, it encodes the holonomy representation of the flat connection $a \in \Omega^1(Y_\Gamma, \mathfrak{sp}(1))$ we started with. If we replace $g: S^3 \rightarrow \text{Sp}(1)$ with $l_u \circ g$, then $\rho: \Gamma \rightarrow \text{Sp}(1)$ is replaced by $c_u \circ \rho$, where c_u is conjugation by u in $\text{Sp}(1)$. Therefore, for a suitable choice of $u \in \text{Sp}(1)$ we can arrange that $\rho = \rho_\alpha$. From the defining formula $g \circ \gamma = l_{\rho_\alpha(\gamma)} \circ g$ it follows that $g: S^3 \rightarrow \text{Sp}(1)^\alpha$ is equivariant.

Our setup is summarized in the following diagram

$$\theta \xrightarrow{g^*} b \xleftarrow{p^*} a$$

$$\text{Sp}(1) \xleftarrow{g} S^3 \xrightarrow{p} Y_\Gamma.$$

We may now calculate

$$\int_{Y_\Gamma} \zeta(a) = \frac{1}{|\Gamma|} \int_{S^3} p^*(\zeta(a)) = \frac{1}{|\Gamma|} \int_{S^3} g^*(\zeta(\theta)) = \frac{\deg(g)}{|\Gamma|} \int_{\text{Sp}(1)} \zeta(\theta).$$

By the previous lemma $\int_{\text{Sp}(1)} \zeta(\theta) = 1$, so by reducing this modulo integers we obtain $\text{cs}(\alpha) = \text{cs}(a) = \frac{\deg(g)}{|\Gamma|} \pmod{\mathbb{Z}}$. To complete the proof we must show that $\deg(h)/|\Gamma| \pmod{\mathbb{Z}}$ is independent

of the equivariant map $h: S^3 \rightarrow \mathrm{Sp}(1)^\alpha$. This is a consequence of Lemma B.3, which shows that $\deg h \bmod |\Gamma|$ is independent of h . \square

Example B.15. Let $\Gamma \subset \mathrm{Sp}(1) = \mathrm{SU}(2)$ be any finite subgroup. Let $\alpha = Q$ denote the canonical representation associated with the inclusion above. Then the map $g: S^3 \rightarrow \mathrm{Sp}(1)^\alpha$ given by $g(x) = x^*$ is equivariant. Indeed,

$$g(\gamma \cdot x) = g(x\gamma^*) = (x\gamma^*)^* = \gamma x^* = \gamma \cdot g(x)$$

for each $\gamma \in \Gamma$ and $x \in S^3$. Since $S^3 \subset \mathbb{H}$ and $\mathrm{Sp}(1) \subset \mathbb{H}$ both carry the standard orientation for which $(i, j, k) \in T_1 S^3 = T_1 \mathrm{Sp}(1)$ is a positive basis, it follows that $\deg g = -1$ and therefore

$$\mathrm{cs}(\alpha) = \frac{-1}{|\Gamma|} \bmod \mathbb{Z}.$$

If we reverse the orientation of Y_Γ then the value of cs changes by a sign.

Proposition B.16. *Let $\Gamma \subset \mathrm{SU}(2)$ be a finite subgroup and let α, β be a pair of $\mathrm{SU}(2)$ -representations of $\Gamma = \pi_1(Y_\Gamma)$ corresponding to a pair of flat connections in the $\mathrm{SU}(2)$ -bundle over Y_Γ . Let \mathcal{H} be a solution of the equation*

$$(2 - Q)\mathcal{H} = \alpha - \beta$$

in $R(\Gamma)$. Then $\mathrm{cs}(\alpha) - \mathrm{cs}(\beta) = \epsilon(\mathcal{H})/|\Gamma| \in \mathbb{R}/\mathbb{Z}$.

Proof. Let $E \rightarrow S^4$ be a Γ -equivariant $\mathrm{SU}(2)$ -bundle with $[E_N] = \alpha$ and $[E_S] = \beta$. Then by Proposition B.11 the element $\mathcal{H} := \mathrm{Ind}(D_A^*) \in R(\Gamma)$, where A is any Γ -invariant connection in E , satisfies $(2 - Q)\mathcal{H} = \alpha - \beta$ and $\epsilon(\mathcal{H}) = c_2(E)[S^4]$. Moreover, by the classification theorem B.6 and Corollary B.7 the integer $c_2(E)[S^4]$ satisfies a congruence

$$c_2(E)[S^4] \equiv \deg g - \deg h \bmod |\Gamma|,$$

where $g: S^3 \rightarrow \mathrm{Sp}(1)^\alpha$ and $h: S^3 \rightarrow \mathrm{Sp}(1)^\beta$ are any choices of equivariant maps. From Lemma B.14 we obtain

$$\mathrm{cs}(\alpha) - \mathrm{cs}(\beta) \equiv (\deg g - \deg h)/|\Gamma| \equiv c_2(E)[S^4]/|\Gamma| \equiv \epsilon(\mathcal{H})/|\Gamma| \bmod \mathbb{Z}.$$

This shows that the result is valid for a specific solution \mathcal{H} . However, by Lemma 4.5 any other solution is given by $\mathcal{H} + mR$, where $m \in \mathbb{Z}$ and $R = \mathbb{C}[\Gamma]$ is the regular representation. Since $\dim_{\mathbb{C}} R = |\Gamma|$ we conclude that $\epsilon(\mathcal{H})/|\Gamma|$ is independent of the solution \mathcal{H} modulo \mathbb{Z} and the proof is complete. \square

This theorem reduces the calculation of Chern-Simons invariants to solving an equation in $R(\Gamma)$. Employing the graphical solution procedure of Proposition 4.12 one may easily solve for all the flat connections recursively starting at the trivial connection. We will give an example in the next section.

B.5. Connections to Group Cohomology. Given a finite group G its group cohomology may be defined by $H^*(G; \mathbb{Z}) := H^*(BG; \mathbb{Z})$, where BG is its classifying space (see for instance [AM04, Chapter II]). A representation $\rho: G \rightarrow U(n)$ gives rise to characteristic classes $c_i(\rho) \in H^{2i}(\Gamma; \mathbb{Z})$ by pulling back the universal Chern classes $c_i \in H^{2i}(BU(n); \mathbb{Z})$ along the induced map $B\rho: B\Gamma \rightarrow BU(n)$. The purpose of this section is to show that for a finite group $\Gamma \subset \mathrm{SU}(2)$ there is a natural way to identify the Chern-Simons invariant $\mathrm{cs}(\alpha) \in \mathbb{R}/\mathbb{Z}$ with the second Chern class $c_2(\rho_\alpha) \in H^4(\Gamma; \mathbb{Z})$, where $\rho_\alpha: \Gamma \rightarrow \mathrm{SU}(2)$ is the holonomy representation associated with α .

The following calculation is certainly known. We include a quick proof for completeness.

Lemma B.17. *Let $\Gamma \subset \mathrm{SU}(2)$ be a finite subgroup. Then $H^0(\Gamma; \mathbb{Z}) = \mathbb{Z}$, while for $i > 0$*

$$H^i(\Gamma; \mathbb{Z}) \cong \begin{cases} \Gamma^{ab} & \text{for } i \equiv 2 \pmod{4} \\ \mathbb{Z}/|\Gamma| & \text{for } i \equiv 0 \pmod{4} \\ 0 & \text{for } i \equiv 1, 3 \pmod{4} \end{cases}$$

Moreover, for any generator $e \in H^4(\Gamma; \mathbb{Z})$ the cup product $e \cup : H^i(\Gamma; \mathbb{Z}) \cong H^{i+4}(\Gamma; \mathbb{Z})$ is an isomorphism for each $i > 0$.

Proof. Let Γ act freely on S^3 in the standard way. The action induces a fibration $S^3 \rightarrow S^3/\Gamma \rightarrow B\Gamma$ (see for instance [AM04, Lemma 6.2]). The associated Gysin sequence breaks up into the following pieces of information:

- (1) There are isomorphisms $H^i(\Gamma) \cong H^i(S^3/\Gamma)$ for $0 \leq i \leq 2$.
- (2) There is an exact sequence

$$0 \longrightarrow H^3(\Gamma) \longrightarrow H^3(S^3/\Gamma) \xrightarrow{p^*} H^3(S^3) \longrightarrow H^4(\Gamma) \longrightarrow 0,$$

where $p : S^3 \rightarrow S^3/\Gamma$ is the quotient map.

- (3) The map $e \cup : H^i(\Gamma) \rightarrow H^{i+4}(\Gamma)$ is surjective for $i = 0$ and an isomorphism for $i \geq 1$, where $e \in H^4(\Gamma)$ is the Euler class of the fibration.

By Lemma 3.3 we have $H^1(S^3/\Gamma) = 0$ and $H^2(S^3/\Gamma) = \Gamma^{ab}$ so that $H^1(\Gamma) = 0$ and $H^2(\Gamma) \cong \Gamma^{ab}$ by the first point. Since $p : S^3 \rightarrow S^3/\Gamma$ is a covering map of degree $|\Gamma|$, it follows from the second point that $H^3(\Gamma) = 0$ and $H^4(\Gamma) \cong \mathbb{Z}/|\Gamma|$. The rest of the statement now follows from the third point. \square

The above lemma does not give complete information about the product structure in $H^*(\Gamma)$. The underlying Serre spectral sequence giving rise to the Gysin sequence simply does not contain information about the cup product $H^2(\Gamma; \mathbb{Z}) \times H^2(\Gamma; \mathbb{Z}) \rightarrow H^4(\Gamma; \mathbb{Z})$. It is of course known that $H^*(\mathbb{Z}/m, \mathbb{Z}) \cong \mathbb{Z}[\beta]/(m\beta)$ with $|\beta| = 2$ for cyclic groups [Wei94, Exercice 6.7.4]. For the binary icosahedral group I^* one has $(I^*)^{ab} = 0$, so the cup product is necessarily trivial. However, for the remaining groups more refined techniques seem to be required.

Definition B.18. Define $e_\Gamma \in H^4(\Gamma; \mathbb{Z})$ to be $c_2(\iota)$ where $\iota : \Gamma \subset \mathrm{SU}(2)$ denotes the inclusion.

Proposition B.19. *Let $\Gamma \subset \mathrm{SU}(2)$ be a finite subgroup. Then $e_\Gamma \in H^4(\Gamma; \mathbb{Z})$ is a generator. Define $\tau : H^4(\Gamma; \mathbb{Z}) \rightarrow \mathbb{R}/\mathbb{Z}$ to be the group homomorphism given by $\tau(ke_\Gamma) = k/|\Gamma| \pmod{\mathbb{Z}}$. Let α be a flat connection in the trivial $\mathrm{SU}(2)$ -bundle over Y_Γ with associated representation $\rho_\alpha : \Gamma \rightarrow \mathrm{SU}(2)$. Then*

$$\tau(c_2(\rho_\alpha)) = -\mathrm{cs}(\alpha).$$

Proof. According to Lemma B.3 there exists an equivariant map $g : S^3 \rightarrow \mathrm{Sp}(1) \cong \mathrm{SU}(2)$ in the sense that $g(\gamma \cdot q) = \rho_\alpha(\gamma) \cdot g(q)$ for each $\gamma \in \Gamma$ and $q \in S^3$. Composing this map with the orientation reversing orbit map $\mathrm{SU}(2) \rightarrow S^3$ given by $x \mapsto x \cdot 1$, we obtain a map $h : S^3 \rightarrow S^3$, which is equivariant along $\rho_\alpha : \Gamma \rightarrow \mathrm{SU}(2)$. Here $\mathrm{SU}(2)$ acts on $S^3 \subset \mathbb{C}^2$ in the standard way. Note that $\deg h = -\deg g$. There is an induced morphism of fibrations

$$\begin{array}{ccccc} S^3 & \longrightarrow & E\Gamma \times_\Gamma S^3 & \longrightarrow & B\Gamma \\ \downarrow h & & \downarrow & & \downarrow B\rho_\alpha \\ S^3 & \longrightarrow & E\mathrm{SU}(2) \times_{\mathrm{SU}(2)} S^3 & \longrightarrow & B\mathrm{SU}(2), \end{array}$$

where $E\Gamma \rightarrow B\Gamma$ and $E\mathrm{SU}(2) \rightarrow B\mathrm{SU}(2)$ are the universal bundles. As both Γ and $\mathrm{SU}(2)$ act freely on S^3 there are homotopy equivalences

$$E\Gamma \times_\Gamma S^3 \simeq S^3/\Gamma \quad \text{and} \quad E\mathrm{SU}(2) \times_{\mathrm{SU}(2)} S^3 \simeq S^3/\mathrm{SU}(2) = *.$$

The above morphism of fibrations induces a morphism of the associated (cohomology) Serre spectral sequences. As both fibrations have S^3 as fiber the only nontrivial differentials are $d_4: E_4^{p,3} \rightarrow E_4^{p+4,0}$ for $p \geq 0$. In particular, we obtain a commutative diagram

$$\begin{array}{ccc} H^3(S^3; \mathbb{Z}) = E_4^{0,3} & \xrightarrow{d_4} & E_4^{4,0} = H^4(B \mathrm{SU}(2); \mathbb{Z}) \\ \downarrow h^* & & \downarrow B\rho_\alpha^* \\ H^3(S^3; \mathbb{Z}) = E_4^{0,3} & \xrightarrow{d_4} & E_4^{4,0} = H^4(B\Gamma; \mathbb{Z}) \end{array}$$

The upper differential must be an isomorphism since the spectral sequence converges to $H^*(*) = \mathbb{Z}$. Let $u \in H^3(S^3; \mathbb{Z})$ be the unique generator with $d_4(u) = c_2 \in H^4(B \mathrm{SU}(2); \mathbb{Z})$. Then from the commutativity of the diagram we obtain

$$c_2(\alpha) = (B\rho_\alpha)^*(c_2) = d_4(h^*u) = (\deg h)d_4(u) = (\deg h)e_\Gamma.$$

For the final equality, consider the same commutative diagram with ρ_α replaced by the inclusion $\iota: \Gamma \subset \mathrm{SU}(2)$. Then we can take $h: S^3 \rightarrow S^3$ to be the identity, and it follows from the same calculation as above that $e_\Gamma := B\iota^*(c_2) = d^4(u)$. This also shows that e_Γ is a generator for $H^4(B\Gamma; \mathbb{Z})$ as the lower differential must be surjective because $H^4(S^3/\Gamma; \mathbb{Z}) = 0$.

To conclude, recall from Lemma B.14 that for the equivariant map $g: S^3 \rightarrow \mathrm{SU}(2)$ we started with, it holds true that $\mathrm{cs}(\alpha) = \deg(g)/|\Gamma| \pmod{\mathbb{Z}}$. Hence,

$$\tau(c_2(\alpha)) = \tau(\deg(h)e_\Gamma) = -\tau(\deg(g)e_\Gamma) = -\deg(g)/|\Gamma| \pmod{\mathbb{Z}}$$

and the proof is complete. \square

Example B.20. Let $\Gamma = T^*$. There are three flat connections θ , α and λ . Here θ corresponds to the trivial representation, α corresponds to the canonical representation Q , while λ is reducible and corresponds to a representation $\rho \oplus \rho^*$ where $\rho: T^* \rightarrow U(1)$ is one of the 1-dimensional representations (see the character table in A.3). By example B.15 we know that $\mathrm{cs}(\alpha) = -1/|T^*| = -1/24$, and, of course, $\mathrm{cs}(\theta) = 0$. To determine $\mathrm{cs}(\lambda)$ we solve the equation $(2 - Q)\mathcal{H} = \alpha - \lambda$ in $R(\Gamma)$ using the technique of Proposition 4.12. This yields a solution \mathcal{H} with $\epsilon(\mathcal{H}) = \dim_{\mathbb{C}} \mathcal{H} = 15$. Then by Proposition B.16 we obtain

$$\mathrm{cs}(\alpha) - \mathrm{cs}(\lambda) = \epsilon(\mathcal{H})/|T^*| = 15/24 \pmod{\mathbb{Z}}.$$

Hence, $\mathrm{cs}(\lambda) = -2/3 = 1/3 \pmod{\mathbb{Z}}$.

This calculation can be used to determine the cup product $H^2(T^*; \mathbb{Z}) \times H^2(T^*; \mathbb{Z}) \rightarrow H^4(T^*; \mathbb{Z})$ using the above proposition. As $c_1: \mathrm{Hom}(T^*, U(1)) \rightarrow H^2(T^*; \mathbb{Z})$ is an isomorphism, we can take $c_1(\rho) \in H^2(T^*; \mathbb{Z}) \cong \mathbb{Z}/3$ as a generator. Then

$$-c_1(\rho)^2 = c_1(\rho) \cup c_1(\rho^*) = c_2(\rho \oplus \rho^*) = c_2(\lambda) = 8e_\Gamma.$$

In particular, the cup product is an isomorphism from $\mathbb{Z}/3 \otimes \mathbb{Z}/3 \cong \mathbb{Z}/3$ onto the unique cyclic subgroup $8\mathbb{Z}/(24)$ of order 3.

REFERENCES

- [AB96] David M. Austin and Peter J. Braam. “Equivariant Floer theory and gluing Donaldson polynomials”. In: *Topology* 35.1 (1996), pp. 167–200. ISSN: 0040-9383. DOI: 10.1016/0040-9383(95)00000-0.
- [AM04] Alejandro Adem and R. James Milgram. *Cohomology of finite groups*. Second. Vol. 309. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2004, pp. viii+324. ISBN: 3-540-20283-8. DOI: 10.1007/978-3-662-06280-7.
- [AM69] M. F. Atiyah and I. G. Macdonald. *Introduction to commutative algebra*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969, pp. ix+128.

- [Ati79] M. F. Atiyah. *Geometry of Yang-Mills fields*. Scuola Normale Superiore Pisa, Pisa, 1979, p. 99.
- [Auc94] David R. Auckly. “Topological methods to compute Chern-Simons invariants”. In: *Math. Proc. Cambridge Philos. Soc.* 115.2 (1994), pp. 229–251. ISSN: 0305-0041. DOI: 10.1017/S0305004100072
- [Aus90] David M. Austin. “SO(3)-instantons on $L(p, q) \times \mathbf{R}$ ”. In: *J. Differential Geom.* 32.2 (1990), pp. 383–413. ISSN: 0022-040X.
- [Aus95] David M. Austin. “Equivariant Floer groups for binary polyhedral spaces”. In: *Math. Ann.* 302.2 (1995), pp. 295–322. ISSN: 0025-5831. DOI: 10.1007/BF01444497.
- [BD85] Theodor Bröcker and Tammo tom Dieck. *Representations of compact Lie groups*. Vol. 98. Graduate Texts in Mathematics. Springer-Verlag, New York, 1985, pp. x+313. ISBN: 0-387-13678-9. DOI: 10.1007/978-3-662-12918-0.
- [BMR14] Tobias Barthel, J. P. May, and Emily Riehl. “Six model structures for DG-modules over DGAs: model category theory in homological action”. In: *New York J. Math.* 20 (2014), pp. 1077–1159.
- [Boa99] J. Michael Boardman. “Conditionally convergent spectral sequences”. In: *Homotopy invariant algebraic structures (Baltimore, MD, 1998)*. Vol. 239. Contemp. Math. Amer. Math. Soc., Providence, RI, 1999, pp. 49–84. DOI: 10.1090/conm/239/03597.
- [DK90] S. K. Donaldson and P. B. Kronheimer. *The geometry of four-manifolds*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1990, pp. x+440. ISBN: 0-19-853553-8.
- [Don02] S. K. Donaldson. *Floer homology groups in Yang-Mills theory*. Vol. 147. Cambridge Tracts in Mathematics. With the assistance of M. Furuta and D. Kotschick. Cambridge University Press, Cambridge, 2002, pp. viii+236. ISBN: 0-521-80803-0. DOI: 10.1017/CB09780511543098
- [Eis19] S. M. Miller Eismeier. “Equivariant instanton homology”. In: *arXiv: Geometric Topology* (2019).
- [Fre02] Daniel S. Freed. “Classical Chern-Simons theory. II”. In: vol. 28. 2. Special issue for S. S. Chern. 2002, pp. 293–310.
- [FU91] Daniel S. Freed and Karen K. Uhlenbeck. *Instantons and four-manifolds*. Second. Vol. 1. Mathematical Sciences Research Institute Publications. Springer-Verlag, New York, 1991, pp. xxii+194. ISBN: 0-387-97377-X. DOI: 10.1007/978-1-4613-9703-8.
- [GM74] V. K. A. M. Guillemin and J. Peter May. *On the theory and applications of differential torsion products*. Memoirs of the American Mathematical Society, No. 142. American Mathematical Society, Providence, R.I., 1974, pp. ix+94.
- [Hit+87] N. J. Hitchin et al. “Hyper-Kähler metrics and supersymmetry”. In: *Comm. Math. Phys.* 108.4 (1987), pp. 535–589. ISSN: 0010-3616.
- [HR19] Gard Helle and John Rognes. “Boardman’s Whole-Plane Obstruction Group for Cartan-Eilenberg Systems”. In: *Documenta Mathematica* (2019), pp. 1855–1878. ISSN: 1431-0635.
- [KK90] Paul A. Kirk and Eric P. Klassen. “Chern-Simons invariants of 3-manifolds and representation spaces of knot groups”. In: *Math. Ann.* 287.2 (1990), pp. 343–367. ISSN: 0025-5831. DOI: 10.1007/BF01446898.
- [Kro89] P. B. Kronheimer. “The construction of ALE spaces as hyper-Kähler quotients”. In: *J. Differential Geom.* 29.3 (1989), pp. 665–683. ISSN: 0022-040X.
- [LM89] H. Blaine Lawson Jr. and Marie-Louise Michelsohn. *Spin geometry*. Vol. 38. Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 1989, pp. xii+427. ISBN: 0-691-08542-0.
- [Mac95] Saunders Mac Lane. *Homology*. Classics in Mathematics. Reprint of the 1975 edition. Springer-Verlag, Berlin, 1995, pp. x+422. ISBN: 3-540-58662-8.

- [McK80] John McKay. “Graphs, singularities, and finite groups”. In: *The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979)*. Vol. 37. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, R.I., 1980, pp. 183–186.
- [Seg68] Graeme Segal. “Equivariant K -theory”. In: *Inst. Hautes Études Sci. Publ. Math.* 34 (1968), pp. 129–151. ISSN: 0073-8301.
- [Tau11] Clifford Henry Taubes. *Differential geometry*. Vol. 23. Oxford Graduate Texts in Mathematics. Bundles, connections, metrics and curvature. Oxford University Press, Oxford, 2011, pp. xiv+298. ISBN: 978-0-19-960588-0. DOI: 10.1093/acprof:oso/9780199605880.001.0001.
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*. Vol. 38. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1994, pp. xiv+450. ISBN: 0-521-43500-5; 0-521-55987-1. DOI: 10.1017/CB09781139644136.
- [Wol11] Joseph A. Wolf. *Spaces of constant curvature*. Sixth. AMS Chelsea Publishing, Providence, RI, 2011, pp. xviii+424. ISBN: 978-0-8218-5282-8.