arXiv:2203.10371v2 [math.GT] 8 Aug 2022

A GENERALIZATION OF THE HOPF DEGREE THEOREM

MATTHEW D. KVALHEIM

Abstract. The Hopf theorem states that homotopy classes of continuous maps from a closed connected oriented smooth n-manifold M to the n-sphere are classified by their degree. Such a map is equivalent to a section of the trivial n-sphere bundle over M. A generalization of the Hopf theorem is obtained for sections of nontrivial oriented n-sphere bundles over M.

Throughout, let M be a compact connected oriented smooth n-dimensional manifold without boundary. The Hopf degree theorem is as follows [Mil97, p. 51].

Theorem (Hopf)**.** A pair of continuous maps $f, g\colon M \to \mathbb{S}^n$ to the n-sphere are homotopic if and only if $\deg(f) = \deg(g)$.

Homotoping f and g to smooth maps, denoted by the same symbols, does not affect their degrees. Let $c \in \mathbb{S}^n$ be a regular value for both. Then $\deg(f) = I(f, c)$ and $\deg(g) = I(g, c)$ coincide with oriented intersection numbers. If instead c is viewed as the constant map $M \to \{c\} \subset \mathbb{S}^n$, rather than as a point in $\mathbb{S}^n$, these can be viewed as oriented intersection numbers of maps $M \to \mathbb{S}^n$ [Mil97, GP10, Hir94]. Such a map is equivalent to a section of the trivial sphere bundle $M \times \mathbb{S}^n \to M$. This motivates the following question.

Question. What can be said when the sphere bundle is nontrivial and $M \to \{c\}$ is replaced by a general continuous section?

A partial answer is given by the following generalization of the Hopf theorem, which is the special case that π is a trivial bundle and Z is constant.

Theorem 1. Let X, Y, Z be continuous sections of a smooth n-sphere bundle $\pi\colon E \to M$, where E is oriented. Then X, Y are homotopic through sections if and only if $I(X, Z) = I(Y, Z)$.

Remark 1. The proof works by reducing to the situation of the Hopf theorem.

Proof. Homotopy invariance of the oriented intersection number implies that $I(X, Z) = I(Y, Z)$ if X and Y are homotopic.

Conversely, assume that $I(X, Z) = I(Y, Z)$. It suffices to show that X and Y are homotopic through sections to some common other section. By approximation techniques we may assume (after preliminary homotopies through sections) that X, Y, Z are smooth and transverse (cf. [Hir94, p. 56, Ex. 3]). In particular, X, Y intersect Z only at finitely many points in M.

Finiteness and connectedness of M imply that these intersection points are contained in the interior of a compact set $B \subset M$ diffeomorphic to a ball in $\mathbb{R}^n$ [MV94].

University of Pennsylvania, Philadelphia, PA, USA
E-mail address: kvalheim@seas.upenn.edu.
2020 *Mathematics Subject Classification.* Primary 57R19; Secondary 55R25, 55N45.

Since B is smoothly contractible there is a fiber-preserving diffeomorphism

$$\pi^{-1}(B) \approx B \times \mathbb{S}^n$$

with respect to which sections over B may be viewed as $\mathbb{S}^n$-valued and $Z|_B$ is constant. After homotopies through sections we may assume that

(1) $X|_B$ and $Y|_B$ coincide with $-Z|_B$ on a neighborhood of ∂B in B.

It follows that X, Y induce, by collapsing the boundary of B, self-maps

$$\mathbb{S}^n \approx B/\partial B \to \mathbb{S}^n$$

whose degrees coincide with $I(X,Z) = I(Y,Z)$. The Hopf theorem and its proof imply that the X-induced self-map is homotopic to the Y-induced self-map through maps sending $[\partial B] \in \mathbb{S}^n$ to the constant $-Z|_B \in \mathbb{S}^n$ [Mil97, pp. 50–51]. By (1) this homotopy extends by the constant one to yield a global homotopy of X, Y through sections. □

Remark 2 (cf. [Mil97, p. 50, Remarks])**.** Theorem 1 easily generalizes to let M have a nonempty boundary ∂M: continuous sections X, Y disjoint from $Z|_{\partial M}$ are homotopic through sections disjoint from $Z|_{\partial M}$ if and only if $I(X,Z) = I(Y,Z)$.

References

[GP10] V Guillemin and A Pollack, *Differential topology*, AMS Chelsea Publishing, Providence, RI, 2010, Reprint of the 1974 original. MR 2680546

[Hir94] M W Hirsch, *Differential topology*, Graduate Texts in Mathematics, vol. 33, Springer-Verlag, New York, 1994, Corrected reprint of the 1976 original. MR 1336822

[Mil97] J W Milnor, *Topology from the differentiable viewpoint*, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 1997, Based on notes by David W. Weaver, Revised reprint of the 1965 original. MR 1487640

[MV94] P W Michor and C Vizman, *n-transitivity of certain diffeomorphism groups*, Acta Math. Univ. Comenianae **63** (1994), no. 2, 221–225.

Acknowledgments

This work was funded by ONR N00014-16-1-2817, a Vannevar Bush Faculty Fellowship sponsored by the Basic Research Office of the Assistant Secretary of Defense for Research and Engineering.