ON THE STABLE EQUIVALENCES BETWEEN FINITE TENSOR CATEGORIES

YUYING XU AND GONGXIANG LIU

ABSTRACT. We aim to study Morita theory for tensor triangulated categories. For two finite tensor categories having no projective simple objects, we prove that their stable equivalence induced by an exact k-linear monoidal functor can be lifted to a tensor equivalence under some certain conditions.

1. Introduction

Our aim is to study tensor triangulated categories from Hopf algebraic way. In recent years, there has been tremendous interest in developing tensor triangulated categories (see [Bal05, Bal10, NVY1, NVY2] and references therein). However, so far, limited work has been done in Hopf algebraic fields.

Since all finite tensor categories are Frobenius categories [EO04], it follows that their stable categories are in fact tensor triangulated categories. In particular, take two finite dimensional Hopf algebras H and H' and consider their representation categories H-mod and H'-mod. A natural question is: If their stable categories are equivalent as tensor triangulated categories, then what can we say about the relations between H and H'?

An important relation in Hopf algebras is gauge equivalence. Ng and Schauenburg showed in [NS08] that H and H' are gauge equivalent if and only if H-mod and H'-mod are k-linear monoidally equivalent. Nevertheless, it should be pointed out that even if the stable categories of two finite-dimensional algebras are equivalent, the corresponding algebraic structures may be quite different. That is, Morita theory does not work. For example, it is easy to check that some block algebras are not Morita equivalent, although they are stably equivalent [Bro94].

Our main results states that, under some mild conditions, for Hopf algebras if their stable equivalence is induced from an exact k-linear monoidal functor then they are gauge equivalent. We describe this observation in the categorical language as follows:

Proposition 1.1. Let C and C' be two non-semisimple finite tensor categories. Suppose $F: C \to C'$ is an exact k-linear monoidal functor inducing a stable equivalence $\underline{F}: \underline{C} \to \underline{C'}$. If every simple object in C is invertible, then F is a tensor equivalence.

Theorem 1.2. Let C and C' be two non-semisimple finite tensor categories having no projective simple objects such that $\operatorname{FPdim}(C) = \operatorname{FPdim}(C')$. Suppose $F : C \to C'$ is an exact \mathbb{k} -linear monoidal functor inducing a stable equivalence $\underline{F} : \underline{C} \to \underline{C'}$, then F is a tensor equivalence.

1

²⁰²⁰ Mathematics Subject Classification. 16T05, 16G65, 16G10, 18M05.

 $Key\ words\ and\ phrases.$ Hopf algebra, Stable categories, Tensor categories, Frobenius-Perron dimensions. † Supported by NSFC 12171230.

The present paper is built up as follows. Some definitions, notations and results related to stable categories, tensor categories and Hopf algebras are presented in Section 2. We devote Section 3 to give a proof to our main results: Proposition 1.1 and Theorem 1.2.

2. Preliminaries

Let k be an algebraically closed field throughout this paper. For any k-algebra A, the category of finitely generated modules over A is denoted by A-mod. About general background knowledge, the reader is referred to [ARS95] for stable categories, [Mon93] for Hopf algebras and [EGNO15] for tensor categories.

2.1. **Stable categories.** Let \mathcal{C} be a \mathbb{k} -linear abelian category. The *stable category* of \mathcal{C} written as $\underline{\mathcal{C}}$ is defined as follows: The objects of $\underline{\mathcal{C}}$ are the same as those of \mathcal{C} ; For any objects $X, Y \in \underline{\mathcal{C}}$, the morphisms from X to Y are given by the quotient space

$$\underline{\operatorname{Hom}}_{\mathcal{C}}(X,Y) = \operatorname{Hom}_{\mathcal{C}}(X,Y)/\mathcal{P}(X,Y),$$

where $\mathcal{P}(X,Y)$ is the subspace of $\operatorname{Hom}_{\mathcal{C}}(X,Y)$ consisting of homomorphisms which factor through a projective object. We say two \mathbb{k} -linear abelian categories \mathcal{C} and \mathcal{C}' are stably equivalent, if $\underline{\mathcal{C}}$ and $\underline{\mathcal{C}}'$ are \mathbb{k} -linear equivalent.

For simplicity of presentations, we stipulate the following notations.

Notation 2.1. We use A-mod to denote the stable category of A-mod. Talking about any stable categories \underline{C} , the following notations are always used:

• For $X, Y \in \underline{\mathcal{C}}$, let \underline{f} denote the morphism in the quotient space $\underline{\operatorname{Hom}}_{\mathcal{C}}(X, Y)$ represented by $f \in \operatorname{Hom}_{\mathcal{C}}(X, Y)$. We use the diagram below to indicate $\underline{f} = 0$:

$$f: X \xrightarrow{i} P \xrightarrow{j} Y$$
,

where $f = j \circ i$ in Hom(X, Y) and P is a projective object in C.

• Given a k-linear functor $F: \mathcal{C} \to \mathcal{C}'$, if F transforms projective objects to projective objects, then it induces a functor from $\underline{\mathcal{C}}$ to $\underline{\mathcal{C}'}$:

$$\underline{F}: \underline{\mathcal{C}} \to \underline{\mathcal{C}'}, \ X \mapsto F(X), \ f \mapsto F(f),$$

where $X \in \underline{\mathcal{C}}$ and f is a morphism in $\underline{\mathcal{C}}$.

Recall that an artin algebra A is said to be *self-injective* if it is injective as an A-module. A great deal of mathematical effort in the representation theory of algebras has been devoted to the study of self-injective algebras. The following proposition tells us when a stable equivalence can be lifted to a Morita equivalence.

Lemma 2.2. ([Lin96, Proposition 2.5]) Let A and A' be self-injective k-algebras having no projective simple modules and F: A-mod $\to A'$ -mod be an exact functor. Suppose F induces a stable equivalence $\underline{F}: A$ -mod $\to A'$ -mod. Then F is an equivalence if and only if F maps any simple A-module to a simple A'-module.

2.2. **Tensor categories.** We have the following basic properties about tensor categories.

Lemma 2.3. ([EO04, Propositon 2.3]) Any projective object in a tensor category is also injective.

Lemma 2.4. ([KL94, Corollary 2, p.441]) Let P be a projective object in a tensor category C. Then $P \otimes X$ and $X \otimes P$ are both projective for any object $X \in C$.

It is known that a *tensor equivalence* is a k-linear monoidal equivalence. Here we state the relations between tensor equivalences and *gauge equivalences* in the case of Hopf algebras:

Lemma 2.5. ([NS08, Theorem 2.2]) Let H and H' be finite-dimensional Hopf algebras over \mathbb{R} . If H-mod and H'-mod are tensor equivalent, then H is gauge equivalent to H' as Hopf algebras.

An important technical tool in the study of tensor categories is Frobenius-Perron dimensions. Due to [EGNO15, Proposition 4.5.4], one can define the Frobenius-Perron dimensions of objects in a tensor category \mathcal{C} . To be specific, for each object $X \in \mathcal{C}$, FPdim(X) is the largest positive eigenvalue of the matrix of left or right multiplication by X. Furthermore, FPdim is the unique additive and multiplicative map which takes positive values on all simple objects of \mathcal{C} . Here is a lemma which we will need later.

Lemma 2.6. ([EGNO15, Proposition 4.5.7]) Let C and C' be finite tensor categories. If $F: C \to C'$ is an exact k-linear monoidal functor, then $\mathrm{FPdim}(F(X)) = \mathrm{FPdim}(X)$ for any $X \in C$.

Let $\{L_i\}_{i\in I}$ be the set of isomorphic classes of simple objects of C, and P_i denotes the projective cover of L_i for each i.

Definition 2.7. ([EGNO15, Definition 6.1.6]) Let C be a finite tensor category. Then the Frobenius-Perron dimension of C is defined by

$$\operatorname{FPdim}(\mathcal{C}) := \sum_{i \in I} \operatorname{FPdim}(L_i) \operatorname{FPdim}(P_i)$$

For a finite dimensional Hopf algebra H, it is easy to see $FPdim(H-mod) = dim_k(H)$, which can be found in [EGNO15, Example 6.1.9].

2.3. **Tensor triangulated categories.** In retrospect, all finite tensor categories are Frobenius categories by Lemma 2.3. Meanwhile the stable categories of Frobenius categories are triangulated categories [Hap88, Theorem 2.6].

According to [Bal05, Definition 1.1], a tensor triangulated category is a triangulated category having a monoidal structure [Mac98, Chapter VII]

$$\otimes: \mathcal{C} \times \mathcal{C} \to \mathcal{C}$$

and a unit object $\mathbf{1} \in \mathcal{C}$, such that the bifunctor $-\otimes$ – is exact in each variable. (Note that we are not dealing with symmetric structure here.) Then the stable categories of finite tensor categories are naturally tensor triangulated categories.

Two tensor triangulated categories $\underline{\mathcal{C}}$ and $\underline{\mathcal{C}'}$ are said to be tensor triangulated equivalent if there is a monoidal functor making $\underline{\mathcal{C}}$ and $\underline{\mathcal{C}'}$ be triangulated equivalent. Our aim in this paper is to show that a tensor equivalence can be recovered by a stable equivalence as a special form of tensor triangulated equivalences. Note that a stable equivalence induced by an exact $\underline{\mathbb{k}}$ -linear monoidal functor is clearly a tensor triangulated equivalence.

3. Main results

In the begining, we turn to mention the relation between the Chevalley property and the existence of simple projective objects. A Hopf algebra is said to have the Chevalley property, if the tensor product of two simple modules is semisimple. Generally, let us say a

tensor category has the Chevalley property if the category of semisimple objects is a tensor subcategory [AEG01, definition 4.1].

The following lemma is contributed to simplify the assumptions of our results.

Lemma 3.1. Let C be a non-semisimple finite tensor category with the Chevalley property. Then C has no simple projective objects.

Proof. Otherwise, let L be a simple projective object in \mathcal{C} . Since $L \otimes L^*$ is semisimple, $\mathbf{1}$ is a direct summand of it. Moreover, Lemma 2.4 tells us that $L \otimes L^*$ is projective as L is projective. This implies $\mathbf{1}$ is also projective, then \mathcal{C} is semisimple by [EGNO15, Corollary 4.2.13], a contradiction.

A direct consequence of this lemma is:

Corollary 3.2. Let H be a finite-dimensional non-semisimple Hopf algebra with the Chevalley property. Then H-mod has no simple projective modules.

It is easy to see that a tensor category in which every simple object is invertible (in the sense of [EGNO15, Definition 2.11.1]) has the Chevalley property by [EGNO15, Proposition 4.12.4]. With this observation, we are in a position to show our first main conclusion now:

Proof of Proposition 1.1. We claim F maps simple objects to simple objects. Actually, for any simple object $L \in \mathcal{C}$, we have:

$$F(L^*) \otimes F(L) \cong F(L^* \otimes L) \cong F(\mathbb{k}) \cong \mathbb{k}$$

then

$$\operatorname{length}(F(L^*))\operatorname{length}(F(L)) \leq \operatorname{length}(F(L^*) \otimes F(L)) = \operatorname{length}(\mathbb{k}) = 1,$$

where length(-) denotes the length of the Jordan-Hölder series. Hence length(F(L)) = 1 that is F(L) is a simple object.

Since \mathcal{C} and \mathcal{C}' are finite, we may assume $\mathcal{C} \cong A\text{-mod}$, $\mathcal{C}' \cong A'\text{-mod}$ as \mathbb{k} -linear abelian categories, where A and A' are finite-dimensional \mathbb{k} -algebras. In addition, as \mathcal{C} and \mathcal{C}' are tensor categories, A and A' also can be self-injective according to Lemma 2.3. Moreover, \mathcal{C} and \mathcal{C}' have no projective simple objects by Lemma 3.1. As a result, F is a \mathbb{k} -linear equivalence by Lemma 2.2. Consequently it is a tensor equivalence.

Note that a Hopf algebra H is basic if and only if every simple object in the tensor category of finite-dimensional H-modules is invertible. So the following conclusion is directly obtained.

Corollary 3.3. Let H and H' be finite-dimensional non-semisimple Hopf algebras. Suppose $F: H\text{-mod} \to H'\text{-mod}$ is an exact \mathbbm{k} -linear monoidal functor inducing a stable equivalence $F: H\text{-mod} \to H'\text{-mod}$. If F: H is also basic, then F: H and F: H are gauge equivalent.

To prove our second main results, we need several lemmas. First, let us make some basic observations.

Lemma 3.4. Let $f: X \to Y$ be an isomorphism in a non-semisimple finite k-linear abelian category C. If $\underline{f} = 0$ in \underline{C} , then f has the following form:

$$f: X \stackrel{i}{\rightarrowtail} P(Y) \stackrel{p}{\twoheadrightarrow} Y,$$

where (P(Y), p) is a projective cover of Y and $f = p \circ i$.

Proof. According to f = 0 in $\underline{\mathcal{C}}$, we can find a projective object P such that $f = \beta \circ \alpha$, where

$$f: X \stackrel{\alpha}{\rightarrowtail} P \stackrel{\beta}{\twoheadrightarrow} Y.$$

Moreover, since f is an isomorphism, α and β are in fact a monomorphism and an epimorphism respectively. By the universal property of projective cover, there exists an epimorphism $h: P \twoheadrightarrow P(Y)$ such that $p \circ h = \beta$.

As a result, we have:

$$f: X \stackrel{h\alpha}{\rightarrowtail} P(Y) \stackrel{p}{\twoheadrightarrow} Y,$$

and $h \circ \alpha$ is a monomorphism for the reason that f is an isomorphism.

Using the above lemma, we have the following result.

Lemma 3.5. Let C and C' be non-semisimple finite k-linear abelian categories having no projective simple objects. Suppose a k-linear functor $F: C \to C'$ induces a stable equivalence between C and C'. If $F(L_1)$ is simple and $F(L_1) \cong F(L_2)$ in C' for two simple objects L_1 and L_2 , then we have $L_1 \cong L_2$.

Proof. Assume on the contrary that $L_1 \ncong L_2$ which follows $\operatorname{Hom}_{\mathcal{C}}(L_1, L_2) = 0$ by Schur Lemma. Hence, $\operatorname{\underline{Hom}}_{\mathcal{C}}(L_1, L_2) = 0$ in $\operatorname{\underline{\mathcal{C}}}$. However, since F induces a stable equivalence, we know

$$\underline{\operatorname{Hom}}_{\mathcal{C}}(L_1, L_2) \cong \underline{\operatorname{Hom}}_{\mathcal{C}'}(F(L_1), F(L_2)).$$

Next, we claim each isomorphism $f \in \operatorname{Hom}_{\mathcal{C}'}(F(L_1), F(L_2))$ satisfies $\underline{f} \neq 0$, which would follow that $\underline{\operatorname{Hom}_{\mathcal{C}'}}(F(L_1), F(L_2)) \neq 0$. Assume on the contrary that f has the following form:

$$f: F(L_1) \stackrel{i}{\rightarrowtail} P(F(L_2)) \stackrel{j}{\twoheadrightarrow} F(L_2)$$

where $P(F(L_2))$ can be chosen as a projective cover of $F(L_2)$ by Lemma 3.4. Let us consider the following commuting diagram:

$$F(L_1) \xrightarrow{i} P(F(L_2)) \xrightarrow{t} Coker(i)$$

$$\downarrow j \qquad \qquad \downarrow \beta$$

$$F(L_2) \xrightarrow{\alpha} N$$

where $(\operatorname{Coker}(i), t)$ is the cokernel of i and (β, α) is the pushout of (j, t).

There are two cases which may happen:

(1) If N = 0, then we have an epimorphism:

$$P(F(L_2)) \rightarrow \operatorname{Coker}(i) \oplus F(L_2),$$

which follows another epimorphism:

$$P(F(L_2)) = P(P(F(L_2))) \twoheadrightarrow P(\operatorname{Coker}(i)) \oplus P(F(L_2))$$

where $P(F(L_2))$ and $P(\operatorname{Coker}(i))$ denote projective covers of $P(F(L_2))$ and $\operatorname{Coker}(i)$ respectively. Thus, $\operatorname{Coker}(i) = 0$ and consequently $F(L_1) \cong P(F(L_2))$, which contradicts to the assumption that the simple module $F(L_1)$ is not projective.

(2) If $N \neq 0$, since $\alpha \circ f = \alpha \circ j \circ i = \beta \circ t \circ i = 0$, we find $\alpha = 0$. This leads to a contradiction that $N = \text{Im}(\alpha)$.

In conclusion, $\underline{f} \neq 0$ and thus $\underline{\operatorname{Hom}}_{\mathcal{C}'}(F(L_1), F(L_2)) \neq 0$. It follows $\operatorname{Hom}_{\mathcal{C}}(L_1, L_2) \neq 0$, or equivalently $L_1 \cong L_2$.

Next result is a categorical version of a result in representation theory of artin algebras.

Lemma 3.6. ([ARS95, cf. Proposition 1.1, p.336]) Let C and C' be two non-semisimple finite k-linear abelian categories and $F: C \to C'$ be a k-linear functor inducing a stable equivalence $\underline{F}: \underline{C} \to \underline{C'}$. Then for any object $X \in C$, X is an indecomposable non-projective object if and only if F(X) is an indecomposable non-projective object.

Proof. For the reason that \mathcal{C} and \mathcal{C}' are finite \mathbb{k} -linear abelian categories, we can assume $\mathcal{C} \cong A$ -mod, $\mathcal{C}' \cong A'$ -mod as \mathbb{k} -linear abelian categories, where A and A' are finite-dimensional \mathbb{k} -algebras. Since for any indecomposable non-projective A-module X, we have the result that $\mathcal{P}(X,X) \subseteq \operatorname{rad} \operatorname{End}_A(X)$, it follows that $\operatorname{End}_A(X)$ is local if and only if $\operatorname{End}_{A'}(F(X))$ is local. That is X is indecomposable if only if F(X) is indecomposable. As F induces a stable equivalence, F(X) is non-projective. The proof is completed.

According to [EGNO15, Remark 4.3.10], an exact k-linear monoidal functor between two tensor categories is also faithful. Using this fact, we have lemma below.

Lemma 3.7. Let C and C' be two finite tensor categories and $F: C \to C'$ be an exact k-linear monoidal functor. For any simple object $L \in C$, if there is an indecomposable object $X \in C'$ such that $F(X) \cong L$, then X is a simple object.

Proof. If otherwise, then we assume length(X) = $n \ge 2$. We will show length(X) $\ge n$. Indeed, if

$$0 \subseteq X_1 \subseteq X_2 \subseteq \cdots \subseteq X_{n-1} \subseteq X$$

is a Jordan-Hölder series of X, then by F is exact, there is a filtration of F(X) as follows:

$$0 \subseteq F(X_1) \subseteq F(X_2) \subseteq \cdots \subseteq F(X_{n-1}) \subseteq F(X)$$
.

Moreover, F is faithful by reason that F is k-linear monoidal functor. It follows that

$$0 \subsetneq F(X_1) \subsetneq F(X_2) \subsetneq \cdots \subsetneq F(X_{n-1}) \subsetneq F(X).$$

Hence $\operatorname{length}(F(X)) \geq n$, which is a contradiction to $\operatorname{length}(F(X)) = \operatorname{length}(L) = 1$. Thus X must be simple.

With the help of the preceding two lemmas, we can now prove the following result.

Proof of Theorem 1.2. Let $\{L_i\}_{i\in I}$ and $\{L'_j\}_{j\in J}$ be the set of isoclasses of simple objects in \mathcal{C} and \mathcal{C}' respectively, where I and J are finite sets. Moreover, we use P_i (resp. P'_j) to represent a projective cover of each simple object L_i (resp. L'_i).

The trick of the proof is to show F maps simple objects to simple objects. As \underline{F} is a stable equivalence, for any $j \in J$, there is an $X \in \mathcal{C}$ such that $F(X) \cong L'_j$ in $\underline{\mathcal{C}}'$, that is $F(X) \oplus Q' \cong L'_j \oplus Q''$ where Q', Q'' are projective objects in \mathcal{C}' . Note that we can choose X to have no projective summands. Let $X \cong X_1 \oplus X_2 \oplus \cdots \oplus X_n$ be the indecomposable decomposition in \mathcal{C} , then by Lemma 3.6, we have $F(X) \cong F(X_1) \oplus F(X_2) \oplus \cdots \oplus F(X_n)$ which is the indecomposable decomposition of F(X). Thus it happens

$$F(X_1) \oplus F(X_2) \oplus \cdots \oplus F(X_n) \oplus Q' \cong L'_i \oplus Q''.$$

By Krull-Schmdit theorem and that L'_j is not projective, there is some k such that $F(X_k) \cong L'_j$ where $1 \leq k \leq n$. Moreover by Lemma 3.7, every X_k is a simple object in \mathcal{C} . Therefore by

Lemma 3.5 there is an injection $\phi: J \to I$, such that $F(L_{\phi(j)}) \cong L'_j$ holds for each $j \in J$. In addition, as F is an exact functor, there is an epimorphism $F(P_i) \twoheadrightarrow P(F(L_i))$ for any $i \in I$, where $P(F(L_i))$ denotes a projective cover of $F(L_i)$.

Consequently, we can get the following formula:

$$\begin{aligned} \operatorname{FPdim}(\mathcal{C}) &= \sum_{i \in I} \operatorname{FPdim}(L_i) \operatorname{FPdim}(P_i) = \sum_{i \in I} \operatorname{FPdim}(F(L_i)) \operatorname{FPdim}(F(P_i)) \\ &= \sum_{j \in J} \operatorname{FPdim}(F(L_{\phi(j)})) \operatorname{FPdim}(F(P_{\phi(j)})) + \sum_{i \in I \setminus \phi(J)} \operatorname{FPdim}(F(L_i)) \operatorname{FPdim}(F(P_i)) \\ &\geq \sum_{j \in J} \operatorname{FPdim}(F(L_{\phi(j)})) \operatorname{FPdim}(F(P_{\phi(j)})) \\ &\geq \sum_{j \in J} \operatorname{FPdim}(F(L_{\phi(j)})) \operatorname{FPdim}(P(F(L_{\phi(j)}))) \\ &= \sum_{j \in J} \operatorname{FPdim}(L'_j) \operatorname{FPdim}(P'_j) = \operatorname{FPdim}(\mathcal{C}'). \end{aligned}$$

By the condition that FPdim(C) = FPdim(C'), we have:

$$\sum_{i \in I \setminus \phi(J)} \operatorname{FPdim}(F(L_i)) \operatorname{FPdim}(F(P_i)) = 0.$$

However, because for any $i \in I$

$$\operatorname{FPdim}(F(P_i)) \ge \operatorname{FPdim}(F(L_i)) = \operatorname{FPdim}(L_i) \ge 1$$

we get the conclusion that $\operatorname{FPdim}(F(L_i)) = 0$ for any $i \in I \setminus \phi(J)$, that is $L_i = 0$ for any $i \in I \setminus \phi(J)$. As a result, $\phi(J) = I$ and hence $F(L_i) = L'_{\phi^{-1}(i)}$ for any $i \in I$.

Since \mathcal{C} and \mathcal{C}' are finite, using the same method used in proof of Proposition 1.1, we can assume $\mathcal{C} \cong A\operatorname{-mod}$, $\mathcal{C}' \cong A'\operatorname{-mod}$ as $\mathbb{k}\operatorname{-linear}$ abelian categories, where A and A' are finite-dimensional self-injective $\mathbb{k}\operatorname{-algebras}$. By Lemma 2.2, F is a $\mathbb{k}\operatorname{-linear}$ equivalence. Consequently it is a tensor equivalence.

It is direct to see the following corollary.

Corollary 3.8. Let H and H' be finite-dimensional non-semisimple Hopf algebras having no simple projective modules such that $\dim_{\mathbb{K}}(H) = \dim_{\mathbb{K}}(H')$. If an exact \mathbb{K} -linear monoidal functor $F: H\operatorname{-mod} \to H'\operatorname{-mod}$ induces a stable equivalence $\underline{F}: H\operatorname{-mod} \to H'\operatorname{-mod}$, then H and H' are gauge equivalent.

Proof. By $FPdim(H-mod) = dim_k(H)$ we can get the conclusion.

Remark 3.9. We end this section by pointing out that: The condition "the functor F is monoidal" can not be removed in Theorem 1.2. Let us illustrate it with an example. Consider the n^2 -dimensional Taft algebras $T_{n^2}(\omega_1)$ and $T_{n^2}(\omega_2)$, where ω_1 and ω_2 are primitive n-th roots of unity. [KMN12, Corollary 3.3] tells us that $T_{n^2}(\omega_1)$ and $T_{n^2}(\omega_2)$ are gauge equivalent if and only if $\omega_1 = \omega_2$. As the fact that $T_{n^2}(\omega_1)$ and $T_{n^2}(\omega_2)$ are isomorphic as algebras, they are Morita equivalent inducing a functor from $T_{n^2}(\omega_1)$ -mod to $T_{n^2}(\omega_2)$ -mod. This functor satisfies the assumptions of Theorem 1.2, except that F is a monoidal functor when $\omega_1 \neq \omega_2$.

References

- [AEG01] N. Andruskiewitsch; P. Etingof; S. Gelaki, Triangular Hopf algebras with the Chevalley property. Michigan Math. J. 49 (2001), no. 2, 277-298.
- [ARS95] M. Auslander; I. Reiten; S.O. Smalø, Representation theory of Artin algebras. Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1995. xiv+423 pp.
- [Bal10] P. Balmer, Tensor triangular geometry. Proceedings of the International Congress of Mathematicians. Volume II, 85-112, Hindustan Book Agency, New Delhi, 2010.
- [Bal05] P. Balmer, The spectrum of prime ideals in tensor triangulated categories. J. Reine Angew. Math. 588 (2005), 149-168.
- [Bro94] M. Broué, Equivalences of blocks of group algebras. Finite-dimensional algebras and related topics (Ottawa, ON, 1992), 1–26, NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci., 424, Kluwer Acad. Publ., Dordrecht, 1994.
- [EO04] P. Etingof; V. Ostrik, Finite tensor categories. Mosc. Math. J. 4 (2004), no. 3, 627-654, 782-783.
- [EGNO15] P. Etingof; S. Gelaki; D. Nikshych; V. Ostrik, Tensor categories. Mathematical Surveys and Monographs, 205. American Mathematical Society, Providence, RI, 2015. xvi+343 pp.
- [Hap88] D. Happel, Triangulated categories in the representation theory of finite-dimensional algebras. London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988. x+208
- [KMN12] Y. Kashina; S. Montgomery; S.-H. Ng, On the trace of the antipode and higher indicators. Israel J. Math. 188 (2012), 57-89.
- [KL94] D. Kazhdan; G. Lusztig, Tensor structures arising from affine Lie algebras. IV, J. Amer. Math. Soc. 7 (1994), no. 2, 383-453.
- [Lin96] M. Linckelmann, Stable equivalences of Morita type for self-injective algebras and p-groups. Math. Z. 223 (1996), no. 1, 87-100.
- [Mac98] S. Mac Lane, Categories for the working mathematician. Second edition. Graduate Texts in Mathematics, 5. Springer-Verlag, New York, 1998. xii+314 pp.
- [Mon93] S. Montgomery, *Hopf algebras and their actions on rings*. CBMS Regional Conference Series in Mathematics, 82. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1993. xiv+238 pp.
- $[\mathrm{NVY1}]$ D.K. Nakano, K.B. Vashaw, M.T. Yakimov: Noncommutative tensor triangular geometry. preprint arXiv:1909.04304v2
- [NVY2] D.K. Nakano, K.B. Vashaw, M.T. Yakimov: On the spectrum and support theory of a finite tensor category. preprint arXiv:2112.11170
- [NS08] S.-H. Ng; P. Schauenburg, Central invariants and higher indicators for semisimple quasi-Hopf algebras. Trans. Amer. Math. Soc. 360 (2008), no. 4, 1839-1860.

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, CHINA

 $Email\ address: {\tt yuyingxu@smail.nju.edu.cn}$

Email address: gxliu@nju.edu.cn