ON TOPOLOGICAL COMPLEXITY OF GORENSTEIN SPACES

SMAIL BENZAKI AND YOUSSEF RAMI

ABSTRACT. In this paper, using Sullivan's approach to rational homotopy theory of simply-connected finite type CW complexes, we endow the \mathbb{Q} -vector space $\mathcal{E}xt_{C^*(X;\mathbb{Q})}(\mathbb{Q},C^*(X;\mathbb{Q}))$ with a graded commutative algebra structure. This leads us to introduce the $\mathcal{E}xt$ -version of higher (resp. module, homology) topological complexity of X_0 , the rationalization of X (resp. of X over \mathbb{Q}). We then make comparisons between these invariants and their respective ordinary ones for Gorenstein spaces. We also highlight, in this context, the benefit of Adams-Hilton models over a field of odd characteristics especially through two cases, the first one when the space is a 2-cell CW-complex and the second one when it is a suspension.

1 Introduction

Throughout this work we assume that X is a 1-connected CW-complex of finite type and \mathbb{K} a field of arbitrary characteristic. In his famous paper, M. Farber introduced the concept of topological complexity of a space X, TC(X), which is indeed seen as the sectional category, $secat(\Delta_X)$, of the diagonal map $\Delta_X:X\to X\times X$. Recall that if $f:X\to Y$ is a continuous map, secat(f) is the smallest integer m for which there are m+1 local homotopy sections $s_i:U_i\to Y$ for f whose sources form an open covering of X. TC(X) is actually used to assess the complexity of motion planning algorithms of a mechanical system along its configuration space X. Later, Y. Rudyak in [22] generalizes Farber's concept and introduces that of higher topological complexity, $TC_n(X)$ $(n \ge 2)$, which turns out to be the sectional category $secat(\Delta_X^n)$ of the n-diagonal $\Delta_X^n: X \to X^n$. Determining $TC_n(X)$ is as difficult as it is exactly for the Lusternik-Scnirelmann category $cat_{LS}(X) = secta(* \hookrightarrow X)$ introduced in [LS]. Spectacular advances using rational homotopy theory methods have however made it possible to establish better approximation of this latter, namely $cat_{LS}(X_0) \leq cat_{LS}(X)$ with X_0 denoting the rationalization X_0 of X. In that sense, J. Carrasquel used in [5] the characterization à la Félix-Halperin to give an explicit definition of the higher topological complexity $TC_n(X_0)$ which in turn lowers $TC_n(X)$ (cf. Definition 4.1 below). This is achieved thanks to the Sullivan model (called also Sullivan algebra) $(\Lambda V, d)$ associated to X (cf. §2 for more details). He also introduced higher (rational) homology (resp. module) topological complexity $HTC_n(X)$, (resp. $MTC_n(X)$) and showed that they interpolate $zcl_n(X) := nilkerH^*(\Delta_X^n)$ and $TC_n(X_0)$.

In this paper we are interested in the study of the invariant $HTC_n(X)$ when X satisfies the *Gorenstein property* which we introduce right after. Recall first that to every local commutative ring R with residue field \mathbb{K} it is associated the invariant $Ext_R(\mathbb{K}, R)$ and it is said a *Gorenstein ring* if dim $Ext_R(\mathbb{K}, R) = 1$.

Motivated by the deep study of this invariant in the context of local algebra, the authors of [13] introduced the notion of a *Gorenstein space* and thereby obtain new characterizations of Poincaré duality complexes. Recall from [23] that a finite n-dimensional subcomplex X of \mathbb{R}^{n+k} is a Poincaré complex if and only if its Spivak fiber F_X (see §4) is an homotopy sphere. They give, in the particular case where X has finite Lusternik-Schnirelmann category $cat_{LS}(X)$ and $\mathbb{K} = \mathbb{Z}_p$ (p prime or zero), a local analogue of the Spivak characterization in terms of the Eilenberg-Moore Ext functor denoted here by $\mathcal{E}xt$ [13, Theorem 3.1]:

(1)
$$H^*(X, \mathbb{Z}_p) \text{ is a Poincar\'e duality algebra} \Leftrightarrow (F_X)_{(p)} \simeq (\mathbb{S}^k)_{(p)} \\ \Leftrightarrow \dim \mathcal{E}xt_{C^*(X; \mathbb{Z}_p)}(\mathbb{Z}_p, C^*(X; \mathbb{Z}_p)) = 1$$

Address: Moulay Ismaïl University, Department of Mathematics B. P. 11 201 Zitoune, Meknès, Morocco

Emails: smail.benzaki@edu.umi.ac.ma and y.rami@umi.ac.ma

2020 Mathematics Subject Classification. Primary 55P62; Secondary 55M30.

Key words and phrases. Higher topological complexity, Eilenberg-Moore functor, Sullivan algebra, Gorenstein spaces.

(cf. §2 for the definition of Poincaré duality algebras and the functor $\mathcal{E}xt$). Recall that, $C^*(X;\mathbb{Z}_p)$ is endowed with the left module structure over itself. The equivalences in (1) make sense to the following definition [13].

Definition 1.1. X is a Gorenstein space over \mathbb{K} if dim $\mathcal{E}xt_{C^*(X;\mathbb{K})}(\mathbb{K}, C^*(X;\mathbb{K})) = 1$.

For instance, if X is finite dimensional then [13, Proposition 4.5, Theorem 3.1]:

$$X$$
 is Gorenstein over $\mathbb{K} \Leftrightarrow H^*(X,\mathbb{K})$ is a Gorenstein algebra $\Leftrightarrow H^*(X,\mathbb{K})$ is a Poincaré duality algebra.

E.g. every closed manifold is Gorenstein over any field \mathbb{K} . In the rational case, if $\dim \pi_*(X) \otimes \mathbb{Q}$ is finite dimensional then X is Gorenstein over \mathbb{Q} [13, Proposition 3.4]. E.g. every rationally elliptic space X (i.e. both if its rational homology and rational homotopy are finite dimensional) is Gorenstein over \mathbb{Q} . The authors of [13] also introduced a morphism of \mathbb{K} -graded vector spaces:

$$ev_{C^*(X,\mathbb{K})}: \mathcal{E}xt_{C^*(X,\mathbb{K})}(\mathbb{K},C^*(X,\mathbb{K})\to H^*(X,\mathbb{K})$$

(cf. §2) called the *evaluation map* of X over \mathbb{K} . This is a homotopy invariant which allows to better (algebraically) characterize Poincaré duality spaces [15, Corollary 2]:

$$X$$
 is a Poincaré duality space $\Leftrightarrow \left\{ egin{array}{l} (i) \ X \ \mbox{is a Gorenstein space}, \ (ii) \ ev_{C^*(X,\mathbb{K})}
eq 0 \ (iii) \ H^*(X,\mathbb{Q}) \ \mbox{is a Gorenstein algebra} \end{array}
ight.$

Also, using the evaluation map we know, for instance, that every space $Y = X \cup e^n$ with e^n representing a non-zero class in $H_n(X, \mathbb{Q})$ is Gorenstein over \mathbb{Q} [13, Corollary 2.3 and Proposition 3.4].

In fact, fewer space types for which the invariants $HTC_n(X)$, $mTC_n(X)$ and $TC_n(X_0)$ are determined. However, in [17] a new invariant $L(X_0)$ is introduced and it is shown that $cat_{LS}(X_0) + L(X_0) \leq TC_n(X_0)$ for any pure elliptic coformal space. It is also established that $TC_n(X_0) = \dim \pi_*(X) \otimes \mathbb{Q}$ for certain particular families of such spaces.

In this work, inspired by Carrasquel's characterizations we will introduce Ext-versions of the aforementioned topological complexities. We start by endowing the graded \mathbb{Q} -vector space

$$A =: Hom_{(\Lambda V,d)}(\Lambda V \otimes \Lambda(sV), (\Lambda V,d))$$

(cf. §2 for more details) with a homotopy multiplicative structure denoted in all that follows

$$\mu_{A}: A \otimes A \to A.$$

The cohomology of A is neither than $\mathcal{A} = \mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d))$ and our first main result reads:

Theorem 1.1. The graded vector space A endowed with the product $\mu_A := H^*(\mu_A)$ is a graded commutative algebra with unit. Moreover, the evaluation map

$$ev_{(\Lambda V,d)}: \mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d)) \to H(\Lambda V,d)$$

is a morphism of graded algebras.

As a consequence, using [13, Remark 1.3] we have the following:

Corollary 1. The graded vector space $\mathcal{E}xt_{C^*(X,\mathbb{Q})}(\mathbb{Q},C^*(X,\mathbb{Q}))$ is a graded commutative algebra and the evaluation map

$$ev_{C^*(X,\mathbb{Q})}: \mathcal{E}xt_{C^*(X,\mathbb{Q})}(\mathbb{Q}, C^*(X,\mathbb{Q})) \to H^*(X,\mathbb{Q})$$

is a morphism of graded algebras.

Next, recall from [6] that a morphism of graded commutative differential algebra (cdga for short) $\varpi:(C,d)\to (D,d)$ admits a homotopy retraction if there exists a map $r:(\Lambda V\otimes C,d))\to (C,d)$ such that $r\circ \iota=Id_C$, where $\iota:(C,d)\to (\Lambda V\otimes C,d)$ stands for the relative model of ϖ . As a particular

case, consider any Sullivan algebra $(\Lambda V, d)$ and denote by $\mu_{A,n}: A^{\otimes n} \to A$ the n-fold product of $\mu_A: A \otimes A \to A$. Using [5, Definition 9], we state in terms of the cdga projection:

$$\Gamma_m: (\mathbf{A}^{\otimes n}, d) \to \left(\frac{\mathbf{A}^{\otimes n}}{(\ker(\mu_{\mathbf{A},n}))^{m+1}}, \overline{d}\right)$$

the following

Definition 1.2. (a): $sc(\mu_{A,n})$ is the least m such that Γ_m admits a homotopy retractioe

- (b): $msc(\mu_{A,n})$ is the least m such that Γ_m admits a homotopy retraction as $(A^{\otimes n}, d)$ -modules.
- (c): $Hsc(\mu_{A,n})$ is the least m such that $H(\Gamma_m)$ is injective.
- (d): $nil \ker (\mu_{A,n}, \mathbb{Q})$ is the longest non trivial product of elements of $\ker (\mu_{A,n})$.

If $(\Lambda V, d)$ is a Sullivan model of X, these are denoted respectively $\mathrm{TC}_n^{\mathcal{E}xt}(X,\mathbb{Q})$, $\mathrm{mTC}_n^{\mathcal{E}xt}(X,\mathbb{Q})$ HTC $_n^{\mathcal{E}xt}(X,\mathbb{Q})$ and $zcl_n^{\mathcal{E}xt}(X,\mathbb{Q})$. We call theme the rational (resp. module, homology, zero cup length) Ext-version topological complexity.

Our second main result reads as follows

Theorem 1.2. Let X be a 1-connected finite type CW-complex. If X is a Gorenstein space over $\mathbb Q$ and $ev_{C^*(X,\mathbb Q)} \neq 0$, then $HTC_n^{\mathcal Ext}(X,\mathbb Q) \leq HTC_n(X,\mathbb Q)$ for any integer $n \geq 2$. Furthermore, if $(\Lambda V,d)$ is a Sullivan minimal model of X, [f] is the generating class of A and $m = HTC_n^{\mathcal Ext}(X,\mathbb Q)$, then:

$$HTC_n(X, \mathbb{Q}) = HTC_n^{\mathcal{E}xt}(X, \mathbb{Q}) =: m \Leftrightarrow f(1)^{\otimes n} \in (\ker \mu_n)^m \setminus (\ker \mu_n)^{m+1}.$$

Notice, referring to [15] that hypothesis on X imply that it is either a Poincaré duality space on $\mathbb Q$ or else $H^*(X,\mathbb Q)$ is not noetherian and not a Gorenstein graded algebra. It also should be pointed out here that, when X satisfies Poincaré duality property then $HTC_n(X,\mathbb Q)=mTC_n(X,\mathbb Q)$ [6, Corollary 4.8]. As Particular cases, we have:

Corollary 2. Hypothesis of Theorem 1.2 and hence its conclusions are satisfied in the following cases:

- (a) X is rationally elliptic,
- (b) $H_{>N}(X,\mathbb{Z})=0$, for some N, and $H^*(X,\mathbb{Q})$ is a Poincaré duality algebra,
- (c) X is a finite 1-connected CW-complex and its Spivak fiber F_X has finite dimensional cohomology.

Next, let R be a principal ideal domain containing $\frac{1}{2}$. Denote by $\rho(R)$ the least non invertible prime (or ∞) in R and by $CW_r(R)$ the sub-category of finite r-connected CW-complexes X ($r \ge 1$) satisfying $\dim(X) \le r\rho(R)$.

In [16] (see also [2]), in his attempt to extend Sullivan's theory to arbitrary rings, S. Halperin associated to every $X \in CW_r(R)$ an appropriate differential graded Lie algebra (L, ∂) and showed that its Cartan-Eilenberg-Chevally complex $C^*(L, \partial)$ is, on one hand, linked with the cochains algebra $C^*(X, R)$ by a series of quasi-isomorphisms ([16, p. 274]) and, on the other hand, it is quasi-isomorphic to a free commutative differential graded algebra $(\Lambda W, d)$ [16, §7]. $(\Lambda W, d)$ is then called a free commutative model of X or a Sullivan minimal model of X.

In Section 5 we will extend a part of Theorem 1.1 and its corollary to the sub-category $CW_r(R)$ when $R=\mathbb{K}$ is a field of odd characteristic and make use of Adams-Hilton models introduced in [1] to obtain an explicit calculation of the homotopy invariant $\mathcal{A}=\mathcal{E}xt_{C^*(X,\mathbb{K})}(\mathbb{K},C^*(X,\mathbb{K}))$ when X is a suspension or a two-cell CW-complex.

The rest of the paper is organized as follows: In Section 2 we recall notions used to state and to establish our results while in Section 3 we introduce the multiplicative structure μ_A and the proof of Theorem 1.1. Section 4 is devoted to the proof of Theorem 1.2.

2 PRELIMINARIES

To make this work constructive enough, we summarize in this section most of the results established on Gorenstein algebras and Gorenstein spaces. \mathbb{K} will denote an arbitrary ground field unless otherwise stated.

2.1 Eilenberg-Moore Ext: A graded module is a family $A = (A^i)_{i \in \mathbb{Z}}$ of \mathbb{K} -modules denoted also $A = \bigoplus_{i \in \mathbb{Z}} A^i$. Every $a \in A^i$ is of degree i denoted thereafter |a|.

A linear map of graded modules $f:A\to B$ of degree |f| is a \mathbb{K} -linear map sending each A^i to $B^{i+|f|}$. In |f|=0 we call it a morphism of graded modules.

In all that follows, unless otherwise stated, modules are over \mathbb{K} and we will assume that $A^i=0$ if i<0.

A grade algebra A is a graded module together with an associative multiplication $\mu_A:A\otimes A\to A$ that has un identity element $1_A=:1\in A^0$. We will put $\mu_A(x\otimes y)=:xy$. Notice that $|\mu_A|=0$. Moreover, if we have $ab=(-1)^{|a||b|}ba$ for all $a,b\in A$, then A is said to be commutative.

A differential graded algebra (A,d) (dga for short) is a graded algebra A together with a linear map $d:A\to A$ of degree |d|=+1 that is a derivation : $d(ab)=d(a)b+(-1)^{|a|}ad(b)$, and satisfying $d\circ d=0$.

A morphism of dga $f:(A,d)\to (B,d)$ is a linear map of degree zero satisfying f(aa')=f(a)f(a'), and the compatibility with the differential d:f(da)=d(f(a)).

A dga algebra A is said to be augmented if it is endowed with a morphism $\varepsilon:A\to\mathbb{K}$ of graded algebras.

A (left) graded (A,d) module is a graded module M equipped with a linear map $A\otimes M\to M$, $a\otimes m\mapsto am$ of degree zero such that a(bm)=(ab)m and 1m=m, and a differential d satisfying $d(am)=(da)m+(-1)^{|a|}a(dm),\ m\in M,\ a\in A.$

A morphism of (left) graded modules over a dga (A,d) is a morphism $f:(M,d)\to (N,d)$ compatible with the differential: $d\circ f=f\circ d$.

A left (A,d)-module (M,d) is said semi-free if it is the union of an increasing sequence $M(0) \subset M(1) \subset M(2) \cdots \subset M(n) \subset \cdots$ of sub (A,d)-modules such that M(0) and each M(i)/M(i-1) is A-free on a basis of cycles. Such an increasing sequence is called a semi-free filtration of (M,d).

A semi-free resolution of an (A,d)-module (M,d) is an (A,d)-semi-free module (P,d) together with a quasi-isomorphism (i.e. a morphism inducing an isomorphism in homology) $m:(P,d) \stackrel{\simeq}{\longrightarrow} (M,d)$ of (A,d)-modules. Each of M(0) and M(i)/M(i-1) has the form $(A,d)\otimes (V(i),0)$ where V(i) is a free \mathbb{K} -module. Thus the surjections $M(n)\to A\otimes V(n)$ and the differential d satisfy:

$$M(n) = M(n-1) \oplus (A \otimes V(n)), \quad and \quad d: V(n) \to M(n-1).$$

By [14, Prop. 6.6], every (A,d)-module (M,d) has a semi-free resolution $m:(P,d) \xrightarrow{\cong} (M,d)$ and if $m:(P',d) \xrightarrow{\cong} (M,d)$ is a second semi-free resolution, then, there is an equivalence

$$\alpha:(P',d)\longrightarrow (P,d)$$

of (A,d)-modules such that $m \circ \alpha$ and m' are homotopic morphisms; we denote for short $m \circ \alpha \simeq_A m'$. Particularly, let (A,d) be a differential graded algebra and $(P,d) \stackrel{\simeq}{\longrightarrow} (\mathbb{Q},0)$ an (A,d)-semi-free resolution of $(\mathbb{Q},0)$. This defines the graded (A,d)-module

$$Hom_A((P,d),(A,d)) = \bigoplus_{p \geq 0} Hom_A^{p,*}((P,d),(A,d)) = \bigoplus_{p \geq 0} \bigoplus_{i \geq 0} Hom_A(P^i,A^{i+p}),$$

which, endowed with the differential

$$D(f) = d \circ f - (-1)^p f \circ d; \quad f \in Hom_A^{p,*}((P,d),(A,d)),$$

yields the Eilenberg-Moore Ext functor:

$$\mathcal{E}xt_{(A,d)}(\mathbb{K}, (A,d)) = H^*(Hom_A((P,d), (A,d)), D).$$

This is an invariant up to homotopy of differential graded algebras (see [13, Appendix] or [8, Appendix]). More usefully [13, Remark 1.3] if $(A, d) \xrightarrow{\simeq} (B, d)$ is a quasi-isomorphism of differential graded algebras, then $\mathcal{E}xt_{(A,d)}(\mathbb{K},(A,d))$ is identified with $\mathcal{E}xt_{(B,d)}(\mathbb{K},(B,d))$ via natural (induced) isomorphisms

(3)
$$\mathcal{E}xt_{(A,d)}(\mathbb{K},(A,d)) \xrightarrow{\cong} \mathcal{E}xt_{(A,d)}(\mathbb{K},(B,d)) \xleftarrow{\cong} \mathcal{E}xt_{(B,d)}(\mathbb{K},(B,d)).$$

Particularly, $\mathcal{E}xt_{C^*(X;\mathbb{K})}(\mathbb{K},C^*(X;\mathbb{K}))$ and $\mathcal{E}xt_{C_*(\Omega X;\mathbb{K})}(\mathbb{K},C_*(\Omega X;\mathbb{K}))$ depend only on the homotopy class of X.

The highest N such that $[\mathcal{E}xt_{C^*(X;\mathbb{K})}(\mathbb{K},C^*(X;\mathbb{K}))]^N \neq 0$ is called the formal dimension of X. It is denoted $fd(X, \mathbb{K})$.

2.2 Evaluation map and Gorenstein spaces. Let $\rho:(P,d)\stackrel{\sim}{\to}(\mathbb{K},0)$ be a minimal (A,d)-semi-free resolution of $(\mathbb{K}, 0)$. Consider the chain map

$$cev_{(A,d)}: \operatorname{Hom}_{(A,d)}((P,d),(A,d)) \longrightarrow (A,d)$$

given by $f \mapsto f(z)$, where $z \in P$ is a cocycle representing 1 in \mathbb{K} . We call it the *chain evaluation map* of (A, d). Passing to homology, we obtain the natural map

$$ev_{(A,d)}: \mathcal{E}xt_{(A,d)}(\mathbb{K},(A,d)) \longrightarrow H^*(A,d),$$

called the evaluation map of (A,d). The definition of $ev_{(A,d)}$ is independent of the choice of (P,d) and z. The evaluation map of X over \mathbb{K} is by definition the evaluation map of $C^*(X,\mathbb{K})$.

A Poincaré duality algebra over \mathbb{K} is a graded algebra $H = \{H^k\}_{0 \le k \le N}$ such that $H^N = \mathbb{K}\alpha$ and the pairing $<\beta,\gamma>\alpha=\beta\gamma,\;\beta\in H^k,\;\gamma\in H^{N-k}$ defines an isomorphism $H^k\stackrel{\cong}{\to} Hom_{\mathbb{K}}(H^{N-k},\mathbb{K}),$ $0 \le k \le N$. In particular, $H = Hom_{\mathbb{K}}(Hom_{\mathbb{K}}(H,\mathbb{K}),\mathbb{K})$ is necessarily finite dimensional.

A Poincaré space at K is a space whose cohomology with coefficients in K is a Poincaré duality algebra. In this case, the cohomology class α such that $H^N(X,\mathbb{K})=\mathbb{K}\alpha$ has degree $N=fd(X,\mathbb{K})$ [13, Proposition 5.1]. It is called the *fundamental class* X.

A Gorenstein algebra over \mathbb{K} is a differential graded algebra (A,d) whose associated graded vector space $\mathcal{E}xt_{(A,d)}(\mathbb{K},(A,d))$ is one dimensional.

A space X is Gorenstein over \mathbb{K} if the cochain algebra $C^*(X;\mathbb{K})$ is a Gorenstein algebra.

For instance, let X be a simply connected CW complex. If in addition X it finite dimensional then: $C^*(X;\mathbb{K})$ is Gorenstein if and only if $H^*(X;\mathbb{K})$ is a Poincaé duality algebra [13, Theorem 3.1]. But, if X is not finite dimensional, it is Gorenstein provided that $\dim \pi_*(X) \otimes \mathbb{Q} < \infty$ [13, Proposition 3.4]. In this case, dim $H^*(X,\mathbb{Q}) < \infty$ if and only if $ev_{C^*(X;\mathbb{Q})} \neq 0$ [19].

3 THE Q-ALGEBRA
$$\mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d))$$

Along this section, the ground field is Q. Recall from [14] that to every finite-type simply-connected space X it is associated a quasi-isomorphism $(\Lambda V, d) \stackrel{\simeq}{\to} A_{PL}(X)$ from a free commutative differential graded algebra (cdga for short) ($\Lambda V, d$) to the commutative graded algebra $A_{PL}(X)$ of polynomial forms with rational coefficients. This latter is connected to $C^*(X,\mathbb{Q})$ by a sequence of quasi-isomorphisms. More explicitly, $\Lambda V = TV/I$ where I is the graded ideal spanned by $\{v \otimes w - (-1)^{deg(u)deg(v)}w \otimes (-1)^{deg(u)deg(v)}\}$ $v, v, w \in V$, $V = \bigoplus_{n>2} V^n$ is a finite-type graded vector space and the differential d is a derivation defined on V satisfying $d \circ d = 0$. $(\Lambda V, d)$ is called a Sullivan model of X. This model is said minimal if moreover d is decomposable, i.e. $d: V \to \Lambda^{\geq 2}V$ where $\Lambda^{\geq 2}V$ denotes the graded vector space spanned by all the monomials $v_1 \dots v_r$ $(r \ge 2)$. Notice that such a model is unique up to isomorphisms [14].

Let (X, x_0) be a based simply-connected finite type CW-complex and denote by

$$m: (\Lambda V, d) \xrightarrow{\simeq} A_{PL}(X)$$
₅

its minimal Sullivan model. In fact, the multiplicative structure of $(\Lambda V, d)$, $\mu_{\Lambda V}: \Lambda V \otimes_{\mathbb{Q}} \Lambda V \to \Lambda V$ is compatible with the one induced on $C^*(X,\mathbb{Q})$ by the diagonal map $\Delta_X: X \to X \times X$ and the same holds for the augmentation $\varepsilon_{\Lambda V}: (\Lambda V, d) \to (\mathbb{Q}, 0)$ and the inclusion $\iota: \{x_0\} \hookrightarrow X$.

Referring for instance to [10] a $(\Lambda V, d)$ -semi-free resolution of $(\mathbb{Q}, 0)$ has the form $(P, d) = (\Lambda V \otimes \Lambda s V, d) \stackrel{\simeq}{\longrightarrow} (\mathbb{Q}, 0)$ where sV is the suspension of V defined by $(sV)^k = V^{k+1}$ and d(sv) = -s(dv) for all $v \in V$.

We are mow ready to define on $A =: Hom_{(\Lambda V,d)}(\mathbb{Q}, (\Lambda V,d))$ an homotopy multiplication:

$$\mu_{A}: A \otimes A \to A$$

which induces in cohomology a multiplicative structure:

on $\mathcal{A}=:\mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d))$. The proof of Theorem 1.1 and Corollary 1 follows from the following theorem.

Theorem 3.1. The \mathbb{Q} -vector space \mathcal{A} , endowed with $\mu_{\mathcal{A}}$, is a graded commutative algebra with unit. Moreover, the evaluation map is a morphism of graded algebras.

Proof. Let $f,g:P\to\Lambda V$ be elements in A representing two classes in \mathcal{A} . As ΛV is commutative, the left ΛV -module P is also a right ΛV -module by setting $x\cdot a=(-1)^{|x||a|}a\cdot x,\quad x\in P$ and $a\in\Lambda V$.

Multiplicative structure:

First, we consider

$$f \otimes g : P \otimes_{\mathbb{Q}} P \longrightarrow \Lambda V \otimes_{\mathbb{Q}} \Lambda V$$
$$x \otimes y \longmapsto (-1)^{|g||x|} f(x) \otimes g(y),$$

and I the ideal generated by $x \cdot a \otimes y - x \otimes a \cdot y$; $x,y \in P$ and $a \in \Lambda V$. It is straightforward that the map $\mu_{\Lambda V} \circ (f \otimes g)$ sends I to zero, which then induces on the quotient $P \otimes_{\Lambda V} P = P \otimes_{\mathbb{Q}} P/I$, the dashed map

$$P \otimes_{\mathbb{Q}} P \xrightarrow{f \otimes g} \Lambda V \otimes_{\mathbb{Q}} \Lambda V \xrightarrow{\mu_{\Lambda V}} \Lambda V$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \qquad \qquad \downarrow \qquad \qquad \qquad$$

where

$$\mu_{\mathcal{A}}(f \otimes g) =: f.g : P \otimes_{\Lambda V} P \longrightarrow \Lambda V$$
$$x \otimes y \longmapsto (-1)^{|g||x|} f(x)g(y),$$

Now, for $a \in \Lambda V$ and $x, y \in P$, we have $(f \cdot g)((a \cdot x) \otimes y) = (-1)^{|f \cdot g||a|}a(f \cdot g)(x \otimes y)$. Therefore $f \cdot g$ is an ΛV -morphism.

Next, we show that $Q = P \otimes_{\Lambda V} P$ is an ΛV -semi-free resolution. Recall that a semi-free resolution (P,d) of $\mathbb Q$ has the form $W \otimes_{\mathbb Q} \Lambda V$ with $W = \bigoplus_{i=0}^{+\infty} W(i)$ and each W(i) is a free graded $\mathbb Q$ -module and $d:W(k) \to P(k-1)$, with the semi-free filtration given by $P(k) = \bigoplus_{i=0}^k W(i) \otimes_{\mathbb Q} \Lambda V$ [14]. Therefore,

$$Q = (W \otimes_{\mathbb{Q}} \Lambda V) \otimes_{\Lambda V} (W \otimes_{\mathbb{Q}} \Lambda V) = (W \otimes_{\mathbb{Q}} W) \otimes_{\mathbb{Q}} \Lambda V.$$

Let $Z = W \otimes_{\mathbb{Q}} W$ and put $Q(k) = \bigotimes_{i=0}^k Z(i) \otimes_{\mathbb{Q}} \Lambda V$ where $Z(l) = \bigoplus_{i+j=l} W(i) \otimes_{\mathbb{Q}} W(j)$ is obviously a free graded \mathbb{Q} -module since each W(i) is. For any $x \otimes y \in W(i) \otimes W(j)$, we verify easily that

$$dx \otimes y \in P(i-1) \otimes W(j) \subseteq Q(k-1)$$
 and $x \otimes dy \in W(i) \otimes P(j-1) \subseteq Q(k-1)$,

whence $D: Z(k) \to Q(k-1)$. It results that $(Q, D) \stackrel{\simeq}{\longrightarrow} (\mathbb{Q}, 0)$ is an ΛV -semi-free resolution of $(\mathbb{Q}, 0)$. This defines a multiplication

$$\mu_{A}: A \otimes A \to A$$

on $A = Hom_{(\Lambda V,d)}(\Lambda V \otimes \Lambda sV, \Lambda V)$. Therefore, passing to cohomology we acquire a well-defined map of vector spaces:

(6)
$$\mu_{\mathcal{A}} : \mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q}, (\Lambda V,d)) \otimes_{\mathbb{Q}} \mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q}, (\Lambda V,d)) \longrightarrow \mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q}, (\Lambda V,d))$$
(7)
$$[f] \otimes [g] \longmapsto [f \cdot g]$$

which gives rise to a multiplication in $\mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d))$ and we can readily check that it is associative since the product in ΛV is.

Uniqueness:

Let $m': (\Lambda V', d') \to A_{PL}(X)$ be another minimal Sullivan model of X. We know, by [14, Proposition 12. 10], that $(\Lambda V, d)$ and $(\Lambda V', d')$ are isomorphic. The same goes clearly for Q and Q'. It results from the above commutative triangle that μ_A is independent of the choice of minimal model of X.

Unit element:

Let $\varepsilon: \Lambda V \to \mathbb{Q}$ be the augmentation. Recall that $P = \Lambda V \otimes \Lambda s V$ is a $(\Lambda V, d)$ -semi-free resolution of $(\mathbb{Q}, 0)$.

We extend ε to $\varepsilon' = \varepsilon \otimes \varepsilon_{\Lambda sV} : \Lambda V \otimes \Lambda sV \to \mathbb{Q}$, then we compose it with the injection $i : \mathbb{Q} \hookrightarrow \Lambda V$ and obtain $\tilde{\varepsilon} : \Lambda V \otimes \Lambda sV \to \Lambda V$ a representative of a class in $\mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d))$. Now, for $f : \Lambda V \otimes \Lambda sV \to \Lambda V$, a representative of an arbitrary class in $\mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d))$, we have

$$f \cdot \tilde{\varepsilon} : (\Lambda V \otimes \Lambda sV) \otimes_{\Lambda V} (\Lambda V \otimes \Lambda sV) = \Lambda V \otimes \Lambda sV \otimes \Lambda sV \to \Lambda V$$

and the map

makes the following diagram commutative:

$$\mathbb{Q} \xleftarrow{\simeq} \Lambda V \otimes \Lambda s V$$


$$\cong \uparrow \qquad \qquad \downarrow f$$

$$\Lambda V \otimes \Lambda s V \otimes \Lambda s V \xrightarrow{f \cdot \tilde{\varepsilon}} \Lambda V.$$

Thus, it defines a homotopy unit element for $A = Hom_{(\Lambda V,d)}((P,d),(\Lambda V,d))$. Passing to cohomology, we get $[f] \cdot [\tilde{\varepsilon}] = [f]$ and similarly $[\tilde{\varepsilon}] \cdot [f] = [f]$. Henceforth, the class $[\tilde{\varepsilon}]$ defines a unit element for $\mu_{\mathcal{A}}$.

Commutativity:

Let τ be the flip map $\tau:P\otimes_{\Lambda V}P\to P\otimes_{\Lambda V}P; x\otimes y\mapsto (-1)^{|x||y|}y\otimes x.$ The diagram



is commutative.

clearly τ being a quasi-isomorphism, $f \cdot g \sim g \cdot f$ and $[f \cdot g] = (-1)^{|f||g|}[g \cdot f]$ so that, the multiplication on A is homotopy commutative and that on \mathcal{A} is indeed commutative.

We respectively conclude that A is a homotopy commutative differential graded algebra with unit and $\mathcal{E}xt_{(\Lambda V,d)}(\mathbb{Q},(\Lambda V,d))$ is a graded commutative \mathbb{Q} -algebra with unit.

Finally, it is clear that the following diagram, where cev is the chain evaluation map of $(\Lambda V, d)$, is commutative:

$$egin{aligned} \mathrm{A} \otimes \mathrm{A} & \xrightarrow{\mu_{\mathrm{A}}} & \mathrm{A} \\ & \downarrow_{cev \otimes cev} & \downarrow_{cev} \\ (\Lambda V, d) \otimes (\Lambda V, d) & \xrightarrow{\mu_{\Lambda V}} & (\Lambda V, d). \end{aligned}$$

Thus, passing to cohomology

we deduce that the evaluation map is a morphism of graded algebra.

4 $\mathcal{E}xt$ -versions approximations and the main theorem

Let (A,d) be any commutative differential graded algebra model for a space X, $(\Lambda V,d)$ its minimal Sullivan model given by the quasi-isomorphism $\theta:(\Lambda V,d)\stackrel{\simeq}{\to}(A,d)$ [14]. Referring to [6], the cdga morphism

$$\mu_n^{\theta} := (\mathrm{Id}_A, \theta, \dots, \theta) : (A, d) \otimes (\Lambda V, d)^{\otimes n-1} \to (A, d)$$

is a special model, called an *s-model*, for the path fibration $\pi_n: X^I \to X^n$ which is indeed the substitute of the *n*-fold diagonal $\Delta_X^n: X \to X^n$. This allows the following:

Definition 4.1. (a): $TC_n(X_0)$ is the least m such that the projection

$$\rho_m: \left(A \otimes (\Lambda V)^{\otimes n-1}, d\right) \to \left(\frac{A \otimes (\Lambda V)^{\otimes n-1}}{\left(\ker \mu_n^{\theta}\right)^{m+1}}, \overline{d}\right)$$

admits an algebra retraction.

- (b): $\mathrm{mTC}_n(X,\mathbb{Q})$ is the least m such that ρ_m admits a retraction as $(A \otimes (\Lambda V)^{\otimes n-1}, d)$ -module.
- (c): $\mathrm{HTC}_n(X,\mathbb{Q})$ is the least m such that $H(\rho_m)$ is injective.
- (d): $nil \ker H^*(\Delta_X^n, \mathbb{Q})$ is the longest non trivial product of elements of $\ker H^*(\Delta_X^n), \mathbb{Q}$.

It is also well known (cf. for instance [6]) that:

(8)
$$nil \ker H^*(\Delta_X^n, \mathbb{Q}) \le \operatorname{HTC}_n(X, \mathbb{Q}) \le \operatorname{mTC}_n(X, \mathbb{Q}) \le \operatorname{TC}_n(X_0) \le \operatorname{TC}_n(X)$$
.

Now, if we take θ to be the identity of ΛV , $\mu_n^\theta = \mu_n^{Id_{\Lambda V}}$ becomes the n-fold multiplication in ΛV denoted by

(9)
$$\mu_n: (\Lambda V)^{\otimes n} \to \Lambda V.$$

Therefore, $nil \ker H^*(\Delta_X^n, \mathbb{Q}) = nil \ker H^*(\mu_n)$.

In a similar way, we put

(10)
$$\mu_{A,n}: A^{\otimes n} \to A \quad \text{and} \quad \mu_{A,n}: A^{\otimes n} \to A$$

where $A := Hom_{\Lambda V}((P,d),(\Lambda V,d))$ and $\mathcal{A} := H(A) = \mathcal{E}xt_{(\Lambda V,d)}((P,d),(\Lambda V,d))$. Then (cf. Definition 1.2) the following definition actually reads as the $\mathcal{E}xt$ -version of the previous one:

Definition 4.2. (a): $TC_n^{\mathcal{E}xt}(X,\mathbb{Q})$ is the least m such that the projection

$$\Gamma_m: (\mathbf{A}^{\otimes n}, d) \to \left(\frac{\mathbf{A}^{\otimes n}}{(\ker(\mu_{\mathbf{A},n}))^{m+1}}, \overline{d}\right)$$

admits a homotopy retraction.

- (b): $\text{mTC}^{\mathcal{E}xt}_n(X)$ is the least m such that Γ_m admits a homotopy retraction as $(A^{\otimes n}, d)$ -module.
- (c): $\mathrm{HTC}_n^{\mathcal{E}xt}(X)$ is the least m such that $H(\Gamma_m)$ is injective.
- (d): $nil \ker (\mu_{A,n}, \mathbb{Q})$ is the longest non trivial product of elements of $\ker (\mu_{A,n})$.

The same arguments used to establish the inequalities in (8) allow the following:

(11)
$$nil \ker (\mu_{\mathcal{A},n}, \mathbb{Q}) \leq HTC_n^{\mathcal{E}xt}(X, \mathbb{Q}) \leq mTC_n^{\mathcal{E}xt}(X, \mathbb{Q}) \leq TC_n^{\mathcal{E}xt}(X_0).$$

The projections

$$\Gamma_m: (\mathbf{A}^{\otimes n}, d) \to \left(\frac{\mathbf{A}^{\otimes n}}{(\ker(\mu_{\mathbf{A},n}))^{m+1}}, \overline{d}\right) \quad \text{and} \quad \rho_m: ((\Lambda V)^{\otimes n}, d) \to \left(\frac{(\Lambda V)^{\otimes n}}{(\ker\mu_n)^{m+1}}, \overline{d}\right)$$

induce two short exact sequences linked by chain evaluation maps:

$$(12) \qquad 0 \longrightarrow (\ker(\mu_{A,n}))^{m+1} \longrightarrow A^{\otimes n} \xrightarrow{\Gamma_m} \frac{A^{\otimes n}}{(\ker(\mu_{A,n}))^{m+1}} \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \theta$$

$$0 \longrightarrow (\ker\mu_n)^{m+1} \longrightarrow (\Lambda V)^{\otimes n} \xrightarrow{\rho_m} \frac{(\Lambda V)^{\otimes n}}{(\ker\mu_n)^{m+1}} \longrightarrow 0$$

As a consequence, we have:

Theorem 4.1. Let X be a 1-connected finite type CW-complex. If X is a Gorenstein space over $\mathbb Q$ and $ev_{C^*(X,\mathbb Q)} \neq 0$, then $HTC_n^{\mathcal Ext}(X,\mathbb Q) \leq HTC_n(X,\mathbb Q)$ for any integer $n \geq 2$. Furthermore, if $(\Lambda V,d)$ is a Sullivan minimal model of X, [f] is the generating class of A and $m = HTC_n^{\mathcal Ext}(X,\mathbb Q)$, then:

$$HTC_n(X,\mathbb{Q}) = HTC_n^{\mathcal{E}xt}(X,\mathbb{Q}) =: m \Leftrightarrow f(1)^{\otimes n} \in (\ker \mu_n)^m \setminus (\ker \mu_n)^{m+1}.$$

Before giving the proof of the theorem, let us first recall that if X is a finite n-dimensional subcomplex of \mathbb{R}^{n+k} , k>n+1 and M its regular neighborhood, the homotopy fiber F_X of the inclusion $\partial M \hookrightarrow M$ is called the *Spivak fiber* for X and it is a homotopy invariant of X. It is introduced in [23] where it is shown that X is a Poincaré complex if and only if F_X is a homotopy sphere.

The following corollary presents some essential classes of spaces satisfying the previous theorem:

Corollary 3. Hypothesis of 1.2 and hence its conclusions are satisfied in the following cases:

- (a) X is rationally elliptic,
- (b) $H_{>N}(X,\mathbb{Z})=0$, for some N, and $H^*(X,\mathbb{Q})$ is a Poincaré duality algebra,
- (c) X is a finite 1-connected CW-complex and its Spivak fiber F_X has finite dimensional cohomology.

For the sake of completeness, we present below a sketch of the proof of the corollary:

- (a) If X is rationally elliptic, then it is Gorenstein and $ev_{C^*(X,\mathbb{Q})} \neq 0$ thanks respectively to [13, Proposition 3.4] and [19, Theorem A].
- (b) Referring to [8] (cf. also [13, Theorem 3.6]), under the hypothesis, $H^*(X,\mathbb{Q})$ is a Poincaré duality algebra if and only if X is a Gorenstein space over \mathbb{Q} . Thus, by [13, Theorem 2.2] we have $\dim \pi_*(X) \otimes \mathbb{Q} < \infty$ and by [19, Theorem A] we obtain $ev_{C^*(X,\mathbb{Q})} \neq 0$.

(c) Here using [13, Corollary 4.5] we have $H^*(X,\mathbb{Q})$ is a Poincaré duality algebra, hence it is a Gorenstein algebra. It results that X is Gorenstein over \mathbb{Q} [13, Proposition 3.2] and $ev_{C^*(X,\mathbb{Q})} \neq$ 0 as in the previous case.

Proof. (of Theorem 4.1): Let $(\Lambda V, d)$ be a Sullivan minimal model of X. Since X is Gorenstein, $\mathcal{A} \cong \mathbb{Q}\Omega$ where Ω is the generating class represented by a cocycle $f \in A^N$ of degree N = fd(X), the formal dimension of X (cf. §5, [13]). Therefore, the diagram (12) induces in cohomology the following one

$$0 \longrightarrow H^{nN}(\ker(\mu_{\mathbf{A},n})^{m+1}) \longrightarrow (\mathcal{A}^{N})^{\otimes n} \xrightarrow{H^{nN}(\Gamma_{m})} H^{nN}(\frac{\mathbf{A}^{\otimes n}}{\ker(\mu_{\mathbf{A},n})^{m+1}}) \longrightarrow 0$$

$$\downarrow \qquad \qquad \qquad \downarrow ev_{(\Lambda V,d)}^{\otimes n} \qquad \downarrow H^{nN}(\theta)$$

$$H^{nN-1}(\frac{(\Lambda V)^{\otimes n}}{(\ker\mu_{n})^{m+1}}) \longrightarrow H^{nN}((\ker\mu_{n})^{m+1}) \longrightarrow (H^{N}(\Lambda V))^{\otimes n} \xrightarrow{H^{nN}(\rho_{m})} H^{nN}(\frac{(\Lambda V)^{\otimes n}}{(\ker\mu_{n})^{m+1}}) \longrightarrow 0$$

Now, since $ev_{(\Lambda V,d)} = ev_{C^*(X,\mathbb{Q})} \neq 0$, this is also the case for the horizontal arrow $ev_{(\Lambda V,d)}^{\otimes n}$. Thus, if $H^{nN}(\rho_m)$ is injective then $H^{nN}(\Gamma_m)$ is also injective. It results that: $HTC_n^{\mathcal{E}xt}(X,\mathbb{Q}) \leq HTC_n(X,\mathbb{Q})$. Next, let m denote the smallest integer such that $H^{nN}(\Gamma_m)$ is injective or equivalently $f^{\otimes n}$ is a cocycle in $A^{\otimes n}$ and $f^{\otimes n} \in \ker(\mu_{A,n})^{m+1} \setminus \ker(\mu_{A,n})^{m+1}$ (see Remarque below). Moreover, since $(\mathcal{A}^N)^{\otimes n}$ is one dimensional, $H^{nN}(\Gamma_m)$ is indeed a bijection. Hence, $H^{nN}(\rho_m)$ is injective if and only if $H^{nN}(\theta)$ is injective. But, $ev_{(\Lambda V,d)}^{\otimes n}$ being non-zero, the commutativity of the right diagram implies that this is equivalent to $f(1)^{\otimes n} \in (\ker \mu_n)^m \setminus (\ker \mu_n)^{m+1}$. Notice that $\mu_n(f(1)^{\otimes n})$ is a cocycle in $(\Lambda V)^{\otimes n}$ and $ev_{(\Lambda V,d)}^{\otimes n}[f^{\otimes n}] = [f(1)]^{\otimes n} \neq 0$. It results that $HTC_n(X,\mathbb{Q}) = HTC_n^{\mathcal{E}xt}(X,\mathbb{Q}) =: m \Leftrightarrow f(1)^{\otimes n} \in \mathbb{Q}$ $(\ker \mu_n)^m \setminus (\ker \mu_n)^{m+1}$.

Remark 1. An equivalent definition of $HTC_n(X,\mathbb{Q})$ when X is a Poincaré duality space reads as follows: It is the smallest integer $m \geq 0$ such that some cocycle ω representing the fundamental class of $(\Lambda V, d)^{\otimes n}$, can be written as a product of m elements of $ker(\mu_n)$ (not necessarily cocycles). Similarly, for any Gorenstein space X, $HTC_n^{\mathcal{E}xt}(X,\mathbb{Q})$ is the smallest integer m such that some cocycle representing the fundamental class of $A^{\otimes n}$, namely $\Omega = [f]^{\otimes n}$ where [f] designates the generating element of A^N , can be written as a product of length m of elements in $ker(\mu_{A,n})$. Therefore, in order to determine $HTC_n(X,\mathbb{Q})$ we may, using the precedent theorem, calculate $m=HTC_n^{\mathcal{E}xt}(X,\mathbb{Q})$ which is quite simpler since A^* is one dimensional, and afterwards deal with the obstruction to have the equality.

Now, if dim $V < \infty$ and dim $H(\Lambda V, d) = \infty$ then by [19, Theorem A] we have $(\Lambda V, d)$ is a Gorenstein but not a Poincaré duality algebra. Moreover $ev_{(\Lambda V,d)}=0$. Hence, in this case, to compare the invariants $HTC_n^{\mathcal{E}xt}(X,\mathbb{Q})$ and $HTC_n(X,\mathbb{Q})$ we should determine them separately.

USE OF THE ADAMS-HILTON MODEL.

The Adams-Hilton model [1] of an arbitrary CW-complex X over an arbitrary field \mathbb{K} is a chain algebra quasi-morphism $\theta_X: (TV, d) \xrightarrow{\simeq} C_*(\Omega X; \mathbb{K})$ i.e. $H_*(\theta_X)$ is an isomorphism of graded algebras. Here V satisfies $H_{i-1}(V,d_1) \cong H_i(X;\mathbb{K})$ and $d_1:V\to V$ is the linear part of d. (TV,d) is called a *free model* of X.

Now, if K is a field with odd characteristic (thus containing $\frac{1}{2}$) and X is a q-connected $(q \ge 1)$ finite CW-complex such that dim $X \leq q \cdot char(\mathbb{K})$ i.e. $X \in CW_q(\mathbb{K})$ (cf. Introduction) it has a minimal Sullivan model (ΛW , d) [16, Theorem 7.1]. Therefore, using the isomorphism (3) we have successively:

(13)
$$\mathcal{E}xt_{C^*(X;\mathbb{K})}(\mathbb{K}, C^*(X;\mathbb{K})) \stackrel{\cong}{\to} \mathcal{E}xt_{(\Lambda W, d)}(\mathbb{K}, (\Lambda W, d)).$$

and

(14)
$$\mathcal{E}xt_{(TV,d)}(\mathbb{K}, (TV,d)) \stackrel{\cong}{\to} \mathcal{E}xt_{C_*(\Omega X:\mathbb{K})}(\mathbb{K}, C_*(\Omega X;\mathbb{K})).$$

These, combined with the isomorphism of graded K-vector spaces [13, Theorem 2.1]:

(15)
$$\mathcal{E}xt_{C^*(X;\mathbb{K})}(\mathbb{K}, C^*(X;\mathbb{K})) \stackrel{\cong}{\to} \mathcal{E}xt_{C_*(\Omega X;\mathbb{K})}(\mathbb{K}, C_*(\Omega X;\mathbb{K})).$$

gives the isomorphism of graded K-vector spaces:

(16)
$$\mathcal{E}xt_{(\Lambda W,d)}(\mathbb{K},(\Lambda W,d)) \cong \mathcal{E}xt_{(TV,d)}(\mathbb{K},(TV,d)).$$

Argument used in the rational case allows us to conclude that $\mathcal{E}xt_{(\Lambda W,d)}(\mathbb{K},(\Lambda W,d))$ has the structure of a graded commutative algebra with unit. The latter isomorphism serves to endow $\mathcal{E}xt_{(TV,d)}(\mathbb{K},(TV,d))$ with the same structure. It results the following

Proposition 5.1. Let \mathbb{K} be a field with odd characteristic and $X \in CW_q(\mathbb{K})$. Then, the graded vector spaces $\mathcal{E}xt_{C^*(X;\mathbb{K})}(\mathbb{K},C^*(X;\mathbb{K}))$ and $\mathcal{E}xt_{C_*(\Omega X;\mathbb{K})}(\mathbb{K},C_*(\Omega X;\mathbb{K}))$ have isomorphic graded commutative algebra structures with unit. In particular, the Adams-Hilton model can be used to make this structure explicit.

Recall that $\mathcal{E}xt_{(TV,d)}(\mathbb{K},(TV,d))$ is, as in the rational case, obtained in terms of the acyclic closure of \mathbb{K} of the form $(TV \otimes (\mathbb{K} \oplus sV), \delta)$, where the differential δ satisfies $\delta s + sd = id$, d being the differential of TV. That is, for any element $z \otimes sv$ of $TV \otimes (\mathbb{K} \oplus sV)$, we have

$$\delta(z \otimes sv) = dz \otimes sv + (-1)^{|z|} zv \otimes 1 - (-1)^{|z|} z \otimes sdv.$$

Notice that any element f in $Hom^p_{(TV,d)}((TV\otimes (\mathbb{K}\oplus sV),\delta),(TV,d))$ is entirely determined by its image of $1\otimes (\mathbb{K}\oplus sV)$ since $TV\otimes (\mathbb{K}\oplus sV)$ is a left (TV,d)-module acting on the first factor. Thus we have

$$(D(g))(1 \otimes sv) = d \circ f(1 \otimes sv) - (-1)^p f \circ \delta(1 \otimes sv)$$

= $df(1 \otimes sv) - (-1)^{p(|v|+1)} v f(1) + (-1)^p f(1 \otimes sdv).$

Therefore,

(a) An element g in $Hom^{p-1}_{(TV,d)}((TV\otimes (\mathbb{K}\oplus sV),\delta),(TV,d))$ is in Im(D) if and only if g=D(f) for some f in $Hom^p_{(TV,d)}((TV\otimes (\mathbb{K}\oplus sV),\delta),(TV,d))$, i.e.

$$g(1 \otimes sv) = df(1 \otimes sv) - (-1)^{p(|v|+1)}vf(1) + (-1)^{p}f(1 \otimes sdv).$$

Consequently:

- $(17) \quad g \in Im(D) \Leftrightarrow g(1 \otimes sv) = df(1 \otimes sv) (-1)^{p(|v|+1)}vf(1) + (-1)^pf(1 \otimes sdv) \text{ for some } f.$
 - (b) An element $f \in Hom^p_{(TV,d)}((TV \otimes (\mathbb{K} \oplus sV), \delta), (TV,d))$ is in Ker(D) if and only if D(f) = 0, that is, $df(1 \otimes sv) = (-1)^{p(|v|+1)}vf(1) (-1)^pf(1 \otimes sdv)$. Consequently:

(18)
$$f \in Ker(D) \Leftrightarrow df(1 \otimes sv) = (-1)^{p(|v|+1)} vf(1) - (-1)^p f(1 \otimes sdv).$$

Now, since $\deg(d)=-1$, $\mathcal{A}_*=\Big(Hom_{(TV,d)}((TV\otimes(\mathbb{K}\oplus sV),\delta),(TV,d)),D\Big)$ is a DGA_* in the sense of [13]. Using the standard convention $\mathcal{A}^{-q}=\mathcal{A}_q$, for all $q\in\mathbb{Z}$, we obtain a DGA^* whose cohomology at -p is the \mathbb{K} -module:

$$\mathcal{E}xt^{-p}_{(TV,d)}(\mathbb{Q},(TV,d)) = H_p\left(Hom_{(TV,d)}((TV\otimes(\mathbb{K}\oplus sV),\delta),(TV,d)),D\right)$$

More explicitly, if $f \in Hom_{(TV,d)}(TV \otimes (\mathbb{K} \oplus sV), \Lambda V)$ is a cycle of (homological) degree p, it define a cohomological class $[f] \in \mathcal{E}xt^{-p}_{(TV,d)}(\mathbb{Q},(TV,d))$ of degree -p.

5.1 Case of a suspension. Assume that $\mathbb{K} = \mathbb{Q}$ and let X be a simply connected space, and $Y = \Sigma X$ its suspension. The morphism of graded modules $\sigma_* : H_*(X;\mathbb{Q}) \longrightarrow H_*(\Omega\Sigma X;\mathbb{Q})$ induced by the adjoint $\sigma : X \to \Omega\Sigma X$ of $id_{\Sigma X}$ extends to a morphism of graded algebras $T(\sigma_*) : TH_*(X;\mathbb{Q}) \longrightarrow H_*(\Omega\Sigma X;\mathbb{Q})$. In fact, this is an isomorphism of graded algebras since $H_*(X;\mathbb{Q})$ is a free graded \mathbb{Q} -module. Therefore ([4])

$$(TH_*(X;\mathbb{Q}),0) \stackrel{\simeq}{\longrightarrow} C_*(\Omega\Sigma X;\mathbb{Q})$$

is an Adams-Hilton model of $Y = \Sigma X$. It results that

$$\mathcal{E}xt_{C_*(\Omega\Sigma X;\mathbb{Q})}(\mathbb{Q},C_*(\Omega\Sigma X;\mathbb{Q}))\cong \operatorname{Ext}_{TV}(\mathbb{Q},TV)$$

where $V = H_*(X; \mathbb{Q})$. Now of $(S, d) = (\Lambda W, d)$ is a Sullivan model of $Y = \Sigma X$, using (15) we obtain a commutative diagram

$$\operatorname{Ext}_{TV}(\mathbb{Q}, TV) \otimes_{\mathbb{Q}} \operatorname{Ext}_{TV}(\mathbb{Q}, TV) \xrightarrow{\mu_{\mathcal{A}}} \operatorname{Ext}_{TV}(\mathbb{Q}, TV)$$

$$\cong \bigcup \qquad \qquad \bigcup \cong$$

$$\operatorname{\mathcal{E}xt}_{(S,d)}(\mathbb{Q}, (S,d)) \otimes_{\mathbb{Q}} \operatorname{\mathcal{E}xt}_{(S,d)}(\mathbb{Q}, (S,d)) \xrightarrow{\mu_{\mathcal{A}}} \operatorname{\mathcal{E}xt}_{(S,d)}(\mathbb{Q}, (S,d)).$$

This permits the use of Adams-Hilton models to explicitly describe the algebra structure on $\mathcal A$ since it restricts to ordinary Ext which loosen the calculations. Notice that $\Omega(\Sigma X)$ is weakly equivalent to the James space J(X) and referring to [14, Example 7] that there exists a minimal Sullivan model for ΣX of the form $(\Lambda W, d)$ with quadratic differential i.e. such that $d(W) \subset \Lambda^2 W$.

5.2 When X is a 2-cell CW complex. Let \mathbb{K} any field containing $\frac{1}{2}$. In this subsection, we showcase another use of the Adams-Hiton models to help picture the algebra structure on \mathcal{A} . Let then $X=S^q\cup_\varphi e^{q+1}, q\geq 2$, be the space where the cell e^{q+1} is attached by a map of degree r. The Adams-Hilton model of X has the form (TV,d) where V is a \mathbb{K} -vector space generated by a and a' with deg(a)=q-1, $deg(a')=q, \quad da=0$ and $\quad da'=-ra$.

Let us go back to where we left off at the beginning of this section and apply, in this case, the obtained formulas (17) and (18). This respectively yields:

$$(19) g \in Im(D) \Leftrightarrow \begin{cases} g(1) = df(1), \\ g(1 \otimes sa) = df(1 \otimes sa) - (-1)^{pq} af(1), \\ g(1 \otimes sa') = df(1 \otimes sa') - (-1)^{p} rf(1 \otimes sa) - (-1)^{p(q+1)} a'f(1), \end{cases}$$

and

(20)
$$f \in Ker(D) \Leftrightarrow \begin{cases} df(1) = 0, \\ df(1 \otimes sa) = (-1)^{pq} a f(1), \\ df(1 \otimes sa') = (-1)^{p} r f(1 \otimes sa) + (-1)^{p(q+1)} a' f(1). \end{cases}$$

Recall that to any pointed topological space X, it is associated in [13] an invariant called the *formal dimension* of X (with respect to a field \mathbb{K}) defined as follows:

$$fd(X,\mathbb{K}) = \sup\{r \in \mathbb{Z} \mid [\mathcal{E}xt^p_{C^*(X;\mathbb{K})}(\mathbb{K},C^*(X;\mathbb{K}))]^r \neq 0\}.$$

or $fd(X, \mathbb{K}) = -\infty$ if such integer does not exist. In particular [13, Proposition 5.1], if $H^*(X; \mathbb{K})$ is finite dimensional,

$$fd(X, \mathbb{K}) = \sup\{r \in \mathbb{Z} \mid H^r(X; \mathbb{K}) \neq 0\}.$$

Notice that, using cellular homology, we see that $H_*(X,\mathbb{Z}) = H_0(X,\mathbb{Z}) \oplus H_q(X,\mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}/r\mathbb{Z}$. We should then discuss two cases:

(i) If $char(\mathbb{K}) = 0$ or co-prime with r, we have $H^*(X, \mathbb{K}) = H^0(X, \mathbb{K}) \cong \mathbb{K}$. In this case, $H^*(X, \mathbb{K})$ has formal dimension fd(X) = 0, thus, it is a Poincaré duality space. Moreover, since it has finite dimensional cohomology, it is also a Gorenstein space [13, Theorem 3.1].

(ii) If $char(\mathbb{K})$ divides r then, $H^*(X,\mathbb{K}) = H^0(X,\mathbb{K}) \oplus H^q(X,\mathbb{K}) \oplus H^{q+1}(X,\mathbb{K}) \cong \mathbb{K} \oplus \mathbb{K} \oplus \mathbb{K}$. Thus, since $q \geq 2$, X is neither a Poincaré duality space nor a Gorenstein space [8, Theorem 1]. In this case, fd(X) = q + 1, so that $\mathcal{E}xt^k_{(TV,d)}(\mathbb{K},(TV,d)) = 0, \forall k > q + 1$.

Example: In this example, we specify the case where q=2, i.e. $X=S^2 \cup_{\varphi} e^3$. Thus $V=\mathbb{K} a \oplus \mathbb{K} a'$ with |a|=1 and |a'|=2. We give below an explicit computation of it to illustrate the use of Adams-Hilton models.

i. Assume that $char(\mathbb{K}) = 0$ or co-prime with r (we specialize in the case where $\mathbb{K} = \mathbb{Q}$).

Let f be a cycle of degree 0, we have df(1)=0, then f(1) is necessarily a scalar $f(1)=\gamma$. The second equation in (20) implies that $df(sa)=\gamma a$, therefore $f(sa)=-\frac{\gamma}{r}a'+\gamma'a^2$. The last equation in (20) gives, after a simple simplification, $df(sa')=r\gamma'a^2$, then $f(sa')=-\gamma'_1a'\cdot a+\gamma'_2a\cdot a'+\gamma''a^3$,

$$Ker(D) = \frac{\mathbb{Q} \oplus \mathbb{Q}a' \oplus \mathbb{Q}a^2 \oplus \mathbb{Q}a' \cdot a \oplus \mathbb{Q}a \cdot a' \oplus \mathbb{Q}a^3}{\langle x_1 = -rx_2; x_3 = -x_4 + x_5 \rangle}.$$

Now let g be an arbitrary element of degree 1

$$\begin{cases} g(1) = \alpha_1 a \\ g(sa) = \alpha_2 a' \cdot a + \alpha_3 a \cdot a' + \alpha_4 a^3 \\ g(sa') = \alpha_5 a'^2 + \alpha_6 a' \cdot a^2 + \alpha_7 a \cdot a' \cdot a + \alpha_8 a^2 \cdot a' + \alpha_9 a^4, \end{cases}$$

by [19] we have

$$\begin{cases} D(g)(1) = 0 \\ D(g)(sa) = (-\alpha_1 - r\alpha_2 + r\alpha_3)a^2 \\ D(g)(sa') = (\alpha_1 + r\alpha_2 - r\alpha_5)a' \cdot a + (r\alpha_3 - r\alpha_5)a \cdot a' + (r\alpha_4 - r\alpha_6 + r\alpha_7 - r\alpha_8)a^3, \end{cases}$$

therefore

$$Im(D) = \frac{\mathbb{Q}a^2 \oplus \mathbb{Q}a' \cdot a \oplus \mathbb{Q}a \cdot a' \oplus \mathbb{Q}a^3}{\langle x_1 = -x_2 + x_3 \rangle}.$$

We consequently obtain $\mathcal{E}xt^0_{(TV,d)}(\mathbb{Q},(TV,d))=\mathbb{Q}$.

Applying the same process for $i \neq 0$, we recover the previously stated fact $\mathcal{E}xt^i_{(TV,d)}(\mathbb{Q},(TV,d))=0$.

ii. Assume $char(\mathbb{K})$ divides r so that r=0 (mod $char(\mathbb{K})$) and da=da'=0. Recall that in general, we have:

$$(D(f))(1 \otimes sa) = df(1 \otimes sa) - (-1)^{pq}af(1) = 0,$$

and

$$(D(f))(1 \otimes sa') = df(1 \otimes sa') - (-1)^p rf(1 \otimes sa) - (-1)^{p(q+1)}a'f(1) = 0.$$

These become respectively in this case:

$$(21) \qquad (D(f))(1\otimes sa)=-(-1)^{pq}af(1)=0, \text{ and } (D(f))(1\otimes sa')=-(-1)^{p(q+1)}a'f(1)=0.$$

Notice that in this case, for an element f to be in ker(D) it is necessarily that f(1) = 0

Let f be a cycle of degree 0, then we have f(1) = 0, consequently df(sa) = df(sa') = 0, which implies $f(sa) = \gamma_1 a' + \gamma_2 a^2$ and $f(sa') = \gamma_3 a' \cdot a + \gamma_4 a \cdot a' + \gamma_5 a^3$. Therefore

$$Ker(D) = \mathbb{K}a^2 \oplus \mathbb{K}a' \oplus \mathbb{K}a' \cdot a \oplus \mathbb{K}a \cdot a' \oplus \mathbb{K}a^3.$$

Now let g be an arbitrary element of degree 1

$$\begin{cases} g(1) = \alpha_1 a \\ g(sa) = \alpha_2 a' \cdot a + \alpha_3 a \cdot a' + \alpha_4 a^3 \\ g(sa') = \alpha_5 a'^2 + \alpha_6 a' \cdot a^2 + \alpha_7 a \cdot a' \cdot a + \alpha_8 a^2 \cdot a' + \alpha_9 a^4, \end{cases}$$

hence

$$\begin{cases} D(g)(1) = 0 \\ D(g)(sa) = -ag(1) = -\alpha_1 a^2 \\ D(g)(sa') = a'g(1) = \alpha_1 a' \cdot a, \end{cases}$$

therefore

$$Im(D) = \frac{\mathbb{K}a^2 \oplus \mathbb{K}a' \cdot a}{\langle x_1 = -x_2 \rangle} \cong \mathbb{K}(a' \cdot a - a^2).$$

we obtain $\mathcal{E}xt^0_{(TV,d)}(\mathbb{K},(TV,d))\cong \mathbb{K}a^2\oplus \mathbb{K}a'\oplus \mathbb{K}a\cdot a'\oplus \mathbb{K}a^3\cong \mathbb{K}^4.$

An application of the same argument yields

$$\mathcal{E}xt_{(TV,d)}^{-1}(\mathbb{K},(TV,d)) \cong \mathbb{K}a^3 \oplus \mathbb{K}a \cdot a' \oplus \mathbb{K}a' \cdot a \oplus \mathbb{K}a \cdot a' \cdot a \oplus \mathbb{K}a^2 \cdot a' \oplus \mathbb{K}a^4 \cong \mathbb{K}^6,$$

$$\mathcal{E}xt^{-2}_{(TV,d)}(\mathbb{K},(TV,d)) \ \cong \ \mathbb{K}a^4 \oplus \mathbb{K}a^2 \cdot a' \oplus \mathbb{K}a \cdot a' \cdot a \oplus \mathbb{K}a' \cdot a^2 \oplus \mathbb{K}a'^2 \oplus \\ \ \mathbb{K}a \cdot a'^2 \oplus \mathbb{K}a^3 \cdot a' \oplus \mathbb{K}a^2 \cdot a' \cdot a \oplus \mathbb{K}a \cdot a' \cdot a^2 \oplus \mathbb{K}a^5 \\ \cong \ \mathbb{K}^{10}.$$

From the previous cases, it is obvious that $\mathcal{E}xt^{-i}_{(TV,d)}(\mathbb{K},(TV,d))\neq 0,\ \forall i\geq 0.$ Since $fd(X,\mathbb{K})=3$, we have $\mathcal{E}xt^{i}_{(TV,d)}(\mathbb{K},(TV,d))=0,\ \forall i\geq 4.$ It remains then to calculate the cases cases i=-1,-2,-3 which are given successively by applying the same process as follows

$$\mathcal{E}xt^1_{(TV,d)}(\mathbb{K},(TV,d))\cong \mathbb{K}a'\oplus \mathbb{K}a^2, \mathcal{E}xt^2_{(TV,d)}(\mathbb{K},(TV,d))\cong \mathbb{K}\oplus \mathbb{K}a \text{ and } \mathcal{E}xt^3_{(TV,d)}(\mathbb{K},(TV,d))\cong \mathbb{K}.$$

Now notice that for q>2, Adams-Hilton model for $X=S^q\cup_{\varphi}e^{q+1}$ has generators a and a' of degrees respectively q-1 and q, thus the degree of a' does not double that of a, so the previous example is somewhat a special case. However the computational process still holds, and we have, for the case \mathbf{i} , $\mathcal{E}xt^0_{(TV,d)}(\mathbb{K},(TV,d))=\mathbb{K}$ and $\mathcal{E}xt^i_{(TV,d)}(\mathbb{K},(TV,d))=0$ for $i\neq 0$. Whereas for the case $\mathbf{i}\mathbf{i}$, computation process holds but the results differ, since we might have $\mathcal{E}xt^{-i}_{(TV,d)}(\mathbb{K},(TV,d))=0$ for finitely many $i\geq -(q+1)$ (e.g. for q=7 we have $\mathcal{E}xt^7_{(TV,d)}(\mathbb{K},(TV,d))=0$), on the other hand, we always have $\mathcal{E}xt^{-i}_{(TV,d)}(\mathbb{K},(TV,d))=0$, $\forall i< -(q+1)$ since fd(X)=q+1.

REFERENCES

- [1] Adams J. F., Hilton P. J., On the chain algebra of a loop space, Comment. Math. Helv. 30 (1955),305-330.
- [2] D. Anick, Hopf algebras up to homotopy, J. Amer. Math. Soc. 2 (1989) 417-453.
- [3] I. Basabe, J. González, Y. B. Rudyak and D. Tamaki, *Higher topological complexity and its symmetrization*, Algebraic and Geometric Topology 14 (2014) 2103-2124.
- [4] R. Bott and H. Samelson, *On the Pontryagin product in spaces of paths*, Commentarii Mathematici Helvetici, Vol.27 (1), 320-337, (1953).
- [5] Carrasquel-Vera, J. G. Computations in rational sectional category. Bull. Belg. Math. Soc. Simon Stevin 22 (2015), no. 3, 455-469.
- [6] J. Carrasquel, Rational methods applied to sectional category and topological complexity, Contemporary Mathematics, volume 702, 2018.
- [7] M. Farber, Topological Complexity of Motion Planning, Discrete Comput Geom 29:211-221 (2003).
- [8] Y. Félix and S. Halperin, A note on Gorenstein spaces, Journal of Pure and Applied Algebra 223 (2019) 4937-4953.
- [9] Y. Félix and S. Halperin, *Rational LS-category and its applications*, Transactions of the American mathematical society Volume 273, Number I, September 1982.
- [10] Y. F'elix, S. Halperin, C. Jacobson, C. Löfwall and J. C. Thomas, *The radical of the homotopy Lie algebra*, Amer. J. Math. 110. (1988), 301-322.
- [11] Y. Félix, S. Halperin and Jean-Michel Lemaire, *The rational LS-category of products and of Poincaré duality complexes*, Topology vol. 37, No. 4, pp. 749-756, 1998.
- [12] Y. Félix, S Halperin and Jean-Claude Thomas, *LS-catégorie et suite spectrale de Milnor-Moore*, Bulletin de la S. M. F., tome 111 (1983), p. 89-96.
- [13] Y. Félix, S. Halperin and Jean-Claude Thomas, Gorenstein spaces. Adv. Math. 71, 92-112 (1988).
- [14] Y. Félix, Steve Halperin and Jean-Claude Thomas, Rational Homotopy Theory, Graduate Texts in Mathematics 215, Springer Verlag, 2000.
- [15] H. Gammelin, Gorenstein spaces with nonzero evaluation map, Trans. Amer. Math. Soc. 351 (1999), no. 8, pp. 3433-3440. 16E99 (55P62).
- [16] S. Halperin, *Universal enveloping algebras and loop space homology*, Journal of Pure and Applied Algebra 83 (1992) 237-282, North-Holland.

- [17] S. Hamoun, Y. Rami and L. Vandembroucq On the rational topological complexity of coformal elliptic spaces, Journal of Pure and Applied Algebra Volume 227, Issue 7, July 2023, 107318
- [18] L. Lechuga and A. Murillo, *Topological complexity of formal spaces*, Topology and robotics, 105-114, Contemp. Math., 438, Amer. Math. Soc., Providence, RI, 2007.
- [19] A. Murillo, The evaluation map of some Gorenstein algebras, J. Pure. Appl. Algebra 91 (1994) 209-218.
- [20] Y. Rami, A new lower bound for LS-category, Arab. J. Math. (2018) 7:39-47.
- [21] Y. Rami, *Dimension globale et classe fondamentale d'un espace*, Annales de l'institut Fourier, tome 49, no° 1 (1999), p. 333-350.
- [22] Y. B. Rudyak, On higher analogs of topological complexity, Topology and its Applications 157 (2010) 916-920.
- [23] M. Spivak, Spaces satisfying Poincaré duality, Topology 6 (1967), 77-102.