

WATKINS' CONJECTURE FOR ELLIPTIC CURVES OVER FUNCTION FIELDS

JERSON CARO

ABSTRACT. In 2002 Watkins conjectured that given an elliptic curve defined over \mathbb{Q} , its Mordell-Weil rank is at most the 2-adic valuation of its modular degree. We consider the analogous problem over function fields of positive characteristic, and we prove it in several cases. More precisely, every modular semi-stable elliptic curve over $\mathbb{F}_q(T)$ after extending constant scalars, and every quadratic twist of a modular elliptic curve over $\mathbb{F}_q(T)$ by a polynomial with sufficiently many prime factors satisfy the analogue of Watkins' conjecture. Furthermore, for a well-known family of elliptic curves with unbounded rank due to Ulmer, we prove the analogue of Watkins' conjecture.

1. INTRODUCTION

Let \mathcal{E} be an elliptic curve over \mathbb{Q} of conductor N . The modular degree $m_{\mathcal{E}}$ of \mathcal{E} is the minimum degree of all modular parametrizations $\phi : X_0(N) \rightarrow \mathcal{E}$ over \mathbb{Q} . The modularity Theorem [24, 20, 2] implies that it is well-defined. In 2002 Watkins [23] conjectured that for every elliptic curve \mathcal{E} over \mathbb{Q} we have $r \leq \nu_2(m_{\mathcal{E}})$, where ν_2 denotes the 2-adic valuation and $r := \text{rank}_{\mathbb{Z}}(\mathcal{E}(\mathbb{Q}))$.

Let k be a finite field of characteristic $p > 3$, write $A = k[T]$ for the polynomial ring, and let $K = k(T)$ be its fraction field. Let ∞ denote the place of K associated with $1/T$. Let E be a non-isotrivial (see Section 2.3 for the definition) elliptic curve defined over K . Under the assumption that E has split multiplicative reduction at ∞ , there is an analogue to the modularity Theorem cf. Theorem 2.1. Namely, if E is non-isotrivial and has split multiplicative reduction at ∞ and conductor ideal \mathfrak{n} , then there is a non-constant map $\phi_E : X_0(\mathfrak{n}) \rightarrow E$, where $X_0(\mathfrak{n})$ is the corresponding Drinfeld modular curve. Thus, from now on we say that E is modular if it is non-isotrivial and has split multiplicative reduction at ∞ . Given a modular elliptic curve E over K , we say that it satisfies Watkins' conjecture if $\text{rank}_{\mathbb{Z}}(E(K)) \leq \nu_2(m_E)$, where m_E is the minimal degree of a modular parametrization ϕ_E .

Using Atkin-Lehner involutions we prove a potential version of Watkins' conjecture for semi-stable elliptic curves over K (see [7] and [3] for other applications of Atkin-Lehner involutions in the context of Watkins' conjecture).

Theorem 1.1. *Let E be a modular semi-stable elliptic curve defined over K with conductor $\mathfrak{n}_E = (n)\infty$. Let k' be a finite field containing the splitting field of n over k , then Watkins' conjecture holds for $E' = E \times_{\text{Spec } K} \text{Spec } K'$, where $K' := k'(T)$.*

It is not known whether the Mordell-Weil rank of elliptic curves over \mathbb{Q} is unbounded or not. Over K we know that the rank is unbounded thanks to the work of Shafarevitch and Tate [19] in the isotrivial case and Ulmer [21] in the non-isotrivial case. The next result proves Watkins' conjecture for one of the families given by Ulmer, thus, we obtain Watkins' conjecture for elliptic curves over K with arbitrarily large rank.

Theorem 1.2. *Let p be a prime and n be a positive integer, such that $6 \mid p^n + 1$. The elliptic curve*

$$E : y^2 + T^d xy = x^3 - 1$$

where $d = (p^n + 1)/6$ defined over $\mathbb{F}_q(T)$, satisfies Watkins' conjecture.

On the other hand, Esparza-Lozano and Pasten [8] prove that, over \mathbb{Q} , the quadratic twist $\mathcal{E}^{(D)}$ of \mathcal{E} by D satisfies Watkins' conjecture whenever the number of distinct prime divisor of D is big enough. Using results of Papikian [13] on $L(\text{Sym}^2 f, 2)$ over function fields, when f is a Drinfeld modular form, we can prove an analogue over function fields. In the following we write $\omega_K(g)$ for the number of distinct irreducible factors of a polynomial g in A .

Theorem 1.3. *Let E be an elliptic curve over K with minimal conductor among its quadratic twists. Let its conductor be $\mathfrak{n}_\infty = (n_1^2 n_2)_\infty$, where n_1, n_2 are square-free coprime polynomials. Assume that E has a non-trivial K -rational 2-torsion. Let g be a monic square-free polynomial of even degree such that $\gcd(n_1, g) = 1$, and $\omega_K(g) \geq 2\omega_K(\mathfrak{n}) - \nu_2(m_E)$, then Watkins' conjecture holds for $E^{(g)}$.*

The condition that g has even degree is necessary to guarantee that $E^{(g)}$ is modular (cf. Section 4). The previous Theorem will be used to deduce the following:

Corollary 1.4. *Assume that E is a semi-stable modular elliptic curve over K . Then we have that $E^{(g)}$ satisfies Watkins' conjecture whenever $\omega_K(g) \geq 3$. Furthermore, if every prime dividing \mathfrak{n} has non-split multiplicative reduction and $E(K)[2] \cong \mathbb{Z}/2\mathbb{Z}$ then $E^{(g)}$ satisfies Watkins' conjecture for every square-free polynomial $g \in A$ of even degree.*

2. PRELIMINARIES

The idea of this section is to define the associated invariants to Watkins' conjecture over function fields. Write K_∞ for the completion of K at T^{-1} , and let \mathcal{O}_∞ be its ring of integers. Let \mathbb{C}_∞ denote the completion of an algebraic closure of K_∞ .

2.1. Drinfeld Modular Curves. We denote by Ω the Drinfeld upper half plane $\mathbb{C}_\infty - K_\infty$. Notice that $GL(2, K_\infty)$ acts on Ω by fractional linear transformations, in particular, so does the Hecke congruence subgroup associated with an ideal \mathfrak{n} of A

$$\Gamma_0(\mathfrak{n}) = \left\{ g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G : a, b, c, d \in \mathbb{F}_q[T], c \equiv 0 \pmod{\mathfrak{n}}, \det(g) \in \mathcal{O}_\infty \right\}.$$

The compactification of the quotient space $\Gamma_0(\mathfrak{n}) \backslash \Omega$ by the finitely many cusps $\Gamma_0(\mathfrak{n}) \backslash \mathbb{P}^1(K)$ is the Drinfeld modular curve. We denoted it by $X_0(\mathfrak{n})$.

2.2. Drinfeld Modular Forms and Hecke Operators. In this section, we define an analogue of the cuspidal Hecke newforms over \mathbb{C} . Another way to understand Ω is the Bruhat-Tits tree \mathcal{T} of $PGL(2, K_\infty)$, whose oriented edges are in correspondence with the cosets of $GL(2, K_\infty)/K_\infty^\times \cdot \mathcal{J}$ (see Section 4.2 [9]), where

$$\mathcal{J} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL(2, \mathcal{O}_\infty) : c \equiv 0 \pmod{T^{-1}} \right\}.$$

This correspondence gives an action of $GL(2, K_\infty)$ on the real-valued functions on the oriented edges of \mathcal{T} by left-multiplying the argument. Let $\underline{H}_1(\Gamma_0(\mathfrak{n}), \mathbb{R})$ be the finite-dimensional \mathbb{R} -space of real-valued, alternating, harmonic and $\Gamma_0(\mathfrak{n})$ -invariant functions on the oriented edges of \mathcal{T} having finite support modulo $\Gamma_0(\mathfrak{n})$.

For each divisor $\mathfrak{d} = (d)$ of \mathfrak{n} , let $i_{\mathfrak{d}}$ be the map

$$i_{\mathfrak{d}} : (\underline{H}_1(\Gamma_0(\mathfrak{n}/\mathfrak{d}), \mathbb{R}))^2 \longrightarrow \underline{H}_1(\Gamma_0(\mathfrak{n}), \mathbb{R})$$

given by

$$i_{\mathfrak{d}}(f, g)(e) = f(e) + g\left(\begin{pmatrix} d & 0 \\ 0 & 1 \end{pmatrix} \cdot e\right),$$

for every oriented edge e . The subspace of oldforms at level \mathfrak{n} is

$$\underline{H}_!^{old}(\Gamma_0(\mathfrak{n}), \mathbb{R}) = \sum_{\mathfrak{p}|\mathfrak{n}} i_{\mathfrak{p}}((\underline{H}_!(\Gamma_0(\mathfrak{n}/\mathfrak{p}), \mathbb{R}))^2).$$

Denote by $\underline{H}_!^{new}(\Gamma_0(\mathfrak{n}), \mathbb{R})$ to the orthogonal complement of the oldforms with respect to the Petersson-norm (see Section 4.8 Gekeler *op. cit.*) defined over $\underline{H}_!(\Gamma_0(\mathfrak{n}), \mathbb{R})$.

For any nonzero ideal \mathfrak{m} there is a Hecke operator $T_{\mathfrak{m}}$, for example, for \mathfrak{m} relatively prime to \mathfrak{n} is defined by

$$T_{\mathfrak{m}}f(e) = \sum f\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \cdot e\right),$$

where the sum runs over $a, b, d \in A$ such that a, d are monic, $\mathfrak{m} = (ad)$, and $\deg(b) < \deg(d)$, see Section 4.9 Gekeler *op. cit.* for a general definition. Finally, a newform is a normalized Drinfeld modular form $f \in \underline{H}_!^{new}(\Gamma_0(\mathfrak{n}), \mathbb{R})$, and an eigenform for all Hecke operators.

2.3. Elliptic curves. Let E be an elliptic curve defined over K . Assume that E has an affine model

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6. \quad (1)$$

where $a_i \in K$. For this cubic equation, define the usual Weierstrass invariants:

$$\begin{aligned} b_2 &= a_1^2 + 4a_2, & b_4 &= a_1a_3 + 2a_4, & b_6 &= a_3^2 + 4a_6, \\ b_8 &= a_1^2a_6 - a_1a_3a_4 + 4a_2a_6 + a_2a_3^2 - a_4^2, \\ c_4 &= b_2^2 - 24b_4, & c_6 &= -b_2^3 + 36b_2b_4 - 216b_6, \\ \Delta &= -b_2^2b_8 - 8b_4^3 - 27b_6^2 + 9b_2b_4b_6, \\ j_E &= c_4^3\Delta^{-1}. \end{aligned}$$

We say that E is *non-isotrivial* when $j_E \notin k$. Since we assume that $\text{char}(k) > 3$ the conductor of E is cubefree. Denote it by \mathfrak{n}_E and by \mathfrak{n} its finite part, in particular, $\mathfrak{n}_E = \mathfrak{n} \cdot \infty^i$, where $i \in \{0, 1, 2\}$. When E has split multiplicative reduction at ∞ , due to Drinfeld's reciprocity law (Proposition 10.3 [6]) and the fact that E is automorphic (Theorem 9.8 in [5]), there is an analogue of the modularity Theorem over \mathbb{Q} :

Theorem 2.1 (Modularity Theorem). *Let E be an elliptic curve over K of conductor $\mathfrak{n}_E = \mathfrak{n}_0 \cdot \infty$ having split multiplicative reduction at ∞ . There is a non-constant morphism $X_0(\mathfrak{n}) \rightarrow E$ defined over K .*

Remark 2.2. This Theorem gives a bijection between primitive newforms f (i.e., f is a newform such that $f \notin n\underline{H}_!^{new}(\Gamma_0(\mathfrak{n}), \mathbb{Z})$ for $n > 1$) with integer eigenvalues and isogeny classes of modular elliptic curves over K with conductor $\mathfrak{n} \cdot \infty$.

2.3.1. L -functions. There is an attached L -function to an elliptic curve with conductor \mathfrak{n}_E , which has an Euler product expansion

$$L(E, s) = \sum_{n \text{ pos. div.}} \frac{a_n}{|n|^s} = \prod_{\mathfrak{p}} \left(1 - \frac{\alpha_{\mathfrak{p}}}{|\mathfrak{p}|^s}\right)^{-1} \left(1 - \frac{\beta_{\mathfrak{p}}}{|\mathfrak{p}|^s}\right)^{-1},$$

where $\alpha_{\mathfrak{p}}, \beta_{\mathfrak{p}}$ are defined as follows: (1) if $\mathfrak{p} \nmid \mathfrak{n}_E$, $\alpha_{\mathfrak{p}} + \beta_{\mathfrak{p}} = a_{\mathfrak{p}} := |\mathfrak{p}| + 1 - \#E(\mathbb{F}_{\mathfrak{p}})$ and $\alpha_{\mathfrak{p}}\beta_{\mathfrak{p}} = |\mathfrak{p}|$, (2) if $\mathfrak{p} \parallel \mathfrak{n}_E$, $\alpha_{\mathfrak{p}} = 0$ and $\beta_{\mathfrak{p}} = \pm 1$, and (3) if $\mathfrak{p}^2 \mid \mathfrak{n}_E$, $\alpha_{\mathfrak{p}} = \beta_{\mathfrak{p}} = 0$.

Due to results of Grothendieck [10] and Deligne [5] $L(E, s) = L(f_E, s)$, where f_E is the newform associated to E , and $L(E, s)$ is a polynomial in the variable q^{-s} of degree $\deg(\mathfrak{n}) - 4$.

Over this newform f_E we define the L -function attached to its symmetric square $L(\text{Sym}^2 f_E, s)$ with the following local factors

$$L_{\mathfrak{p}}(\text{Sym}^2 f_E, s) = \begin{cases} 1, & \text{if } \mathfrak{p}^2 \mid \mathfrak{n}_E, \\ \left(1 - \frac{1}{|\mathfrak{p}|^s}\right)^{-1}, & \text{if } \mathfrak{p} \parallel \mathfrak{n}_E, \\ \left(1 - \frac{\alpha_{\mathfrak{p}}^2}{|\mathfrak{p}|^s}\right)^{-1} \left(1 - \frac{\alpha_{\mathfrak{p}} \overline{\alpha_{\mathfrak{p}}}}{|\mathfrak{p}|^s}\right)^{-1} \left(1 - \frac{\overline{\alpha_{\mathfrak{p}}}^2}{|\mathfrak{p}|^s}\right)^{-1} & \text{if } \mathfrak{p} \nmid \mathfrak{n}_E. \end{cases}$$

When E is semi-stable Proposition 5.4 from [12] implies that $L(\text{Sym}^2 f_E, s)$ is a polynomial in the variable q^{-s} of degree $2 \deg(\mathfrak{n}_E) - 4$.

2.3.2. Upper Bounds for the Rank of the Mordell-Weil Group. The following is a geometric bound for the Mordell-Weil rank due to Tate [18]

$$\text{rank}_{\mathbb{Z}}(E(K)) \leq \text{ord}_{s=1} L(E, s) \leq \deg(\mathfrak{n}_E) - 4. \quad (2)$$

See [22] for detailed proof. In addition, if the elliptic curve E has a non-trivial K -rational 2-torsion, we can give an upper bound for its Mordell-Weil rank in terms of $\omega_K(\mathfrak{n})$, the number of distinct primes that divide \mathfrak{n} in A .

First of all, notice that the change of variables $X = z/4$, $Y = y/8 - a_1 z/8 - a_3/2$ transforms (1) into

$$y^2 = z^3 + b_2 z^2 + 8b_4 z + 16b_6. \quad (3)$$

Let $\gamma \in K$ be a root of the previous cubic, associated to a non-trivial K -rational 2-torsion point. Then $\gamma \in A$ and the change of variables $z = x + \gamma$ turns (3) into

$$y^2 = x^3 + Ax^2 + Bx \quad (4)$$

where

$$A = 3\gamma + b_2 \quad \text{and} \quad B = 3\gamma^2 + 2b_2\gamma + 8b_4.$$

Let Δ^{\min} be the discriminant of the minimal model (1) and let Δ be the discriminant of (4). Notice that $\Delta = 2^{12} \Delta^{\min}$ by the standard transformation formulas, thus, (4) is a minimal model of E . Now, recall the usual exact sequence related to a 2-descent,

$$0 \longrightarrow \frac{E(K)}{2E(K)} \longrightarrow \text{Sel}_2(E/K) \longrightarrow \text{III}(E/K)[2] \longrightarrow 0. \quad (5)$$

Furthermore, consider the exact sequence from Lemma 6.1 of [15]

$$0 \longrightarrow \frac{E'(K)[\theta']}{\phi(E(K)[2])} \longrightarrow \text{Sel}^{\theta}(E/K) \longrightarrow \text{Sel}_2(E/K) \longrightarrow \text{Sel}^{\theta'}(E'/K). \quad (6)$$

These two exact sequences imply that $\text{rank}_{\mathbb{Z}}(E(K)) + 2 \leq s(E, \theta) + s'(E, \theta)$, where $s(E, \theta) = \dim_{\mathbb{F}_2}(\text{Sel}^{\theta}(E/K))$ and $s'(E, \theta) = \dim_{\mathbb{F}_2}(\text{Sel}^{\theta'}(E'/K))$. In addition, there is a correspondence between Selmer groups and homogeneous spaces (see Chapter 4 from [14]), which shows that $s(E, \theta) \leq \omega_K(A^2 - 4B) + 1$ and $s'(E, \theta) \leq \omega_K(B) + 1$. Thus, we have the following proposition:

Proposition 2.3. *Let E be an elliptic curve with K -rational 2-torsion and Weierstrass minimal model $y^2 = x^3 + Ax^2 + Bx$, then:*

$$\text{rank}_{\mathbb{Z}}(E(K)) \leq \omega_K(A^2 - 4B) + \omega_K(B),$$

consequently, if α (resp. μ) is the number of primes of additive (resp. multiplicative) bad reduction of E/K . Then:

$$\text{rank}_{\mathbb{Z}}(E(K)) \leq \mu + 2\alpha.$$

2.3.3. *Modular Degree.* Let E be a modular elliptic curve defined over K . Let $X_0(\mathfrak{n})$ be the Drinfeld modular curve parametrizing $\phi_E: X_0(\mathfrak{n}) \rightarrow E$ where ϕ_E is non-trivial and of minimal possible degree. The modular degree m_E is the degree of ϕ_E . The following Lemma relates the 2-adic valuations of m_E and $L(\text{Sym}^2 f, 2)$.

Lemma 2.4. *Let E be a modular elliptic curve with conductor \mathfrak{n}_∞ . Then we have that*

$$\nu_2(m_E) = \nu_2(L(\text{Sym}^2 f, 2)) - \nu_2(\text{val}_\infty(j_E)).$$

Proof. Proposition 1.3 in [13] states that

$$m_E = \frac{q^{\deg \mathfrak{n} - 2} (\widetilde{c_E})^2}{-\text{val}_\infty(j_E)} L(\text{Sym}^2 f, 2),$$

where $\widetilde{c_E}$ is the Manin constant and $q = \#k$. By taking 2-adic valuations we obtain

$$\nu_2(m_E) = \nu_2(q^{\deg \mathfrak{n} - 2} (\widetilde{c_E})^2) + \nu_2(L(\text{Sym}^2 f, 2)) - \nu_2(\text{val}_\infty(j_E)),$$

and by Proposition 1.2 from [11] $\widetilde{c_E}$ is a power of q which yields the desired result. \square

3. WATKINS' CONJECTURE FOR SEMI-STABLE ELLIPTIC CURVES

For any ideal $\mathfrak{m} = (m)$, such that $\mathfrak{m} \mid \mathfrak{n} = (n)$, and \mathfrak{m} and $\mathfrak{n}/\mathfrak{m}$ are relatively primes, there is an Atkin-Lehner involution $W_{\mathfrak{m}}$. This involution acts on $\underline{H}_1(\Gamma_0(\mathfrak{n}), \mathbb{R})$ as follows

$$W_{\mathfrak{m}}f(e) = f\left(\begin{pmatrix} ma & b \\ nc & md \end{pmatrix} \cdot e\right),$$

where $a, b, c, d \in A$ and $m^2ab - nbc = \gamma m$ for some $\gamma \in k^\times$. We denote by $\mathcal{W}(\mathfrak{n})$ the 2-elementary abelian group of all Atkin-Lehner involutions. Let f be a primitive newform; since f is primitive, it is determined by its eigenvalues up to sign. By Lemma 11 from [1] the Hecke operators commute with the Atkin-Lehner involutions, hence $W_{\mathfrak{p}}^{(\mathfrak{n})}f$ and f have the same Hecke eigenvalues. By Lemma 1.2 from [16] $\underline{H}_1^{\text{new}}(\Gamma_0(\mathfrak{n}), \mathbb{R})$ is stable under the Atkin-Lehner involutions, and consequently, we have that $W_{\mathfrak{p}}f = \pm f$.

Remark 3.1. Let E be a modular elliptic curve, and f_E be its attached primitive newform, then f_E is an eigenform of every Atkin-Lehner involution.

The following Proposition gives a lower bound of $\nu_2(m_E)$ in terms of $\omega_K(\mathfrak{n})$.

Proposition 3.2. *Let E be an elliptic curve with conductor $\mathfrak{n}_E = \mathfrak{n}_\infty$. Let f_E be the primitive newform associated to E . Over this newform, we define $\mathcal{W}' = \{W \in \mathcal{W} : W(f_E) = f_E\}$, and $\kappa := \dim_{\mathbb{F}_2}([\mathcal{W}(\mathfrak{n}) : \mathcal{W}']) + \dim_{\mathbb{F}_2}(E(K)[2])$. Then $\omega_K(\mathfrak{n}) - \kappa \leq \nu_2(m_E)$.*

Proof. Proposition 10.3 from [6] gives the following isomorphism

$$H^1(X_0(\mathfrak{n}) \otimes K_\infty^{\text{sep}}, \mathbb{Q}_\ell) \cong \underline{H}_1(\Gamma_0(\mathfrak{n}), \mathbb{Q}_\ell) \otimes \text{sp},$$

where sp is the two-dimensional special ℓ -adic representation of $\text{Gal}(K_\infty^{\text{sep}}/K_\infty)$. Furthermore, this isomorphism is compatible with the action of the Atkin-Lehner involutions.

Since $H^1(X_0(\mathfrak{n}) \otimes K_\infty^{\text{sep}}, \mathbb{Q}_\ell)$ is the dual of $V_\ell(J_0(\mathfrak{n}))$, we have that if $\pi : J_0(\mathfrak{n}) \rightarrow E$ is the projection, then $\pi([W(D)]) = \pi([D])$ for every divisor D of degree 0 over $X_0(\mathfrak{n})$ whenever $W \in \mathcal{W}'$. By Remark 3.1 \mathcal{W}' has at most index 2 in $\mathcal{W}(\mathfrak{n})$. Now, as in Proposition 2.1 in [7] we construct a homomorphism $\theta : \mathcal{W}' \rightarrow E(K)[2]$. First of all, we fix a K -rational point $x_0 \in X_0(\mathfrak{n})$, then for $W \in \mathcal{W}'$ we define $\theta(W) = \pi([W(x_0) - (x_0)])$. Notice that $\theta(W) \in E(K)[2]$, since $x_0 \in X_0(\mathfrak{n})(K)$ and

$$\theta(W) = \pi([W(x_0) - (x_0)]) = \pi([W(W(x_0) - (x_0))]) = -\pi([W(x_0) - (x_0)]) = -\theta(W).$$

Now, define $\mathcal{W}'' = \ker \theta$. Let $\mathcal{X} = X_0(\mathfrak{n})/\mathcal{W}''$, and denote by $\psi : X_0(\mathfrak{n}) \rightarrow \mathcal{X}$ that is also defined over K and by \mathcal{J} the Jacobian of \mathcal{X} . We can define $\iota : X_0(\mathfrak{n}) \rightarrow J_0(\mathfrak{n})$ based on x_0 , and $\iota' : \mathcal{X} \rightarrow \mathcal{J}$ based on $\psi(x_0)$, so we obtain a commutative diagram

$$\begin{array}{ccc} X_0(\mathfrak{n}) & \xrightarrow{\iota} & J_0(\mathfrak{n}) \\ \psi \downarrow & & \downarrow \psi_* \\ \mathcal{X} & \xrightarrow{\iota'} & \mathcal{J}. \end{array}$$

Since $\pi([W(x_0) - x_0]) = 0$ for $W \in \mathcal{W}''$, we have that $\pi \circ \iota(w(x)) = \pi \circ \iota(x)$ for all $x \in X_0(\mathfrak{n})$, in particular, $\pi \circ \iota$ factors through \mathcal{X} . Since the image of ι generates to $J_0(\mathfrak{n})$ as a group, there exists $\pi' : \mathcal{J} \rightarrow E$ such that $\pi = \pi' \circ \psi_*$, then

$$[m_E] = \pi \circ \pi^\vee = (\pi' \circ \psi_*) \circ (\psi^* \circ \pi'^\vee) = \pi' \circ [\deg(\psi)] \circ \pi'^\vee = [\#\mathcal{W}''] \circ (\pi' \circ \pi'^\vee).$$

Since the degree of $[i]$ (multiplication by i) is $i \cdot i^*$ or $(i^*)^2$, where i^* denotes the p -free part of i , then $\#\mathcal{W}'' \mid m_E$, since $p \neq 2$. \square

The previous Proposition and Tate's geometric bound (2) allow us to prove Theorem 1.1.

Proof of Theorem 1.1. Recall that $E' = E \times_{\text{Spec } K} \text{Spec } K'$. Since the conductor of E' is also $\mathfrak{n}_E = (n)\infty$, then by Tate's geometric bound (2) $\text{rank}(E'(K')) \leq \deg(n) - 4$. On the other hand, we know that $\omega_{K'}((n)) = \deg(n)$ because k' contains the splitting field of n . Furthermore, since $\dim_{\mathbb{F}_2}([\mathcal{W}(\mathfrak{n}) : \mathcal{W}']) \leq 1$, by Remark 3.1, we have $\kappa \leq 3$, then by Proposition 3.2 we have that

$$\nu_2(m_{E'}) \geq \omega_K((n)) - 3 = \deg(n) - 3 = \deg(\mathfrak{n}_E) - 4 \geq \text{rank}(E'(k'(T))),$$

which yields the desired result. \square

Ulmer [21] exhibits a closed formula for the rank of a family of elliptic curves. Proposition 3.2 together with this formula allow us to show Watkins' conjecture for this family.

Proof of Theorem 1.2. First of all, we notice that $E(\overline{\mathbb{F}_p}(T))[2] = (0)$, since the polynomial $4x^3 + T^{2d}x - 4$ does not have solution over $\overline{\mathbb{F}_p}(T)$. Notice that E is the change of base point of \mathbb{P}^1 given by $[0 : 1] \mapsto \infty$ of

$$E' : y^2 + xy = x^3 - T^m,$$

where $m = p^n + 1$. Theorem 1.5 in [21] shows that $\mathfrak{n}_{E'} = T(1 - 2^4 3^3 T^m)$, then in particular $\mathfrak{n}_E = (T^m - 2^4 3^3)\infty$. We claim that $f(T) = T^m - 2^4 3^3$ always has a root in \mathbb{F}_{p^2} . Let $\alpha \in \mathbb{F}_{p^2}$ such that $\alpha^2 = 3$, and notice that if $\alpha \in \mathbb{F}_p$, $2^2 3\alpha$ is a root of f . If $\alpha \notin \mathbb{F}_p$, since $6 \mid p^n - 1$ we have that $p \equiv -1 \pmod{3}$, then $p \equiv 1 \pmod{4}$ by the law of quadratic reciprocity. This implies that $2^2 3\alpha$ or $2^2 3\alpha\beta$ is a root of f , where $\beta^2 = -1$. Consequently, there is a bijection between the prime divisors of even degree of $T^m - 1$ and $f(T)$.

By definition, $T^m - 1$ factors over $\mathbb{F}_p[T]$ as follows

$$T^m - 1 = \prod_{e \mid m} \Phi_e(T),$$

where $\Phi_n(T)$ is the n^{th} -cyclotomic polynomial. Thus, the number of prime divisors over $\mathbb{F}_q[T]$ of $f(T)$ is

$$\omega_{\mathbb{F}_q(T)}(\mathfrak{n}_E) = \sum_{e \mid m} \frac{\phi(e)}{o_e(q)} - \begin{cases} 0 & \text{if } T^m - 2^4 3^3 \text{ has solution in } \mathbb{F}_q, \\ 1 & \text{otherwise} \end{cases},$$

where $\phi(e)$ is the cardinality of $(\mathbb{Z}/e\mathbb{Z})^\times$ and $o_e(q)$ is the order of q in $(\mathbb{Z}/e\mathbb{Z})^\times$. On the other hand, we know that $\text{rank}(E(\mathbb{F}_p(T))) = \text{rank}(E'(\mathbb{F}_p(T)))$. Theorem 1.5 in [21] states a closed expression for $\text{rank}(E'(\mathbb{F}_q(T)))$

$$\sum_{\substack{e|m \\ e \nmid 6}} \frac{\phi(e)}{o_e(q)} + \begin{cases} 2 & \text{if } 3 \mid q-1 \\ 1 & \text{otherwise} \end{cases} + \begin{cases} 1 & \text{if } 4 \mid q-1 \\ 0 & \text{otherwise} \end{cases}.$$

Since there are 4 divisors of 6 we obtain

$$\sum_{e|m} \frac{\phi(e)}{o_e(q)} \geq \sum_{\substack{e|m \\ e \nmid 6}} \frac{\phi(e)}{o_e(q)} + 4$$

Furthermore, if $3 \mid q-1$ then q is a square since $p \equiv -1 \pmod{3}$; which implies that $T^m - 2^4 3^3$ has solution in \mathbb{F}_q . Hence, Proposition 3.2 implies that

$$\nu_2(m_E) \geq \omega_{\mathbb{F}_q(T)}(\mathbf{n}_E) - 1 = \sum_{e|m} \frac{\phi(e)}{o_e(q)} - 1 \geq \sum_{\substack{e|m \\ e \nmid 6}} \frac{\phi(e)}{o_e(q)} + 3 \geq \text{rank}(E(\mathbb{F}_q(T))).$$

Finally, if $3 \nmid q-1$ we obtain

$$\nu_2(m_E) \geq \omega_{\mathbb{F}_q(T)}(\mathbf{n}_E) - 1 \geq \sum_{e|m} \frac{\phi(e)}{o_e(q)} - 2 \geq \sum_{\substack{e|m \\ e \nmid 6}} \frac{\phi(e)}{o_e(q)} + 2 \geq \text{rank}(E(\mathbb{F}_q(T))),$$

which gives the desired result. \square

4. WATKINS' CONJECTURE FOR QUADRATIC TWISTS

Let E be a modular elliptic curve with conductor \mathbf{n}_E , since $\text{char}(k) > 3$ there exist square-free coprime polynomials $n_1, n_2 \in A$ such that $\mathbf{n}_E = (n_1^2 n_2) \infty$. For $g \in A$ be a monic square-free polynomial, with $(n_1, g) = 1$, we define the quadratic twist $E^{(g)}$ of E by g as follows

$$E^{(g)}: y^2 = x^3 + Agx^2 + Bg^2x.$$

We assume that $\deg(g)$ is even to ensure that $E^{(g)}$ is modular. To see that, notice that if the change of variables $x \mapsto T^{2n}x$ and $y \mapsto T^{3n}y$ makes E a minimal T^{-1} -integral model, then the change $x \mapsto T^{2(n+m)}x$ and $y \mapsto T^{3(n+m)}y$ makes $E^{(g)}$ a minimal T^{-1} -integral model, where $\deg(g) = 2m$; since g is a monic polynomial, both reductions modulo T^{-1} are the same. Note that the conductor $\mathbf{n}_E^{(g)}$ of $E^{(g)}$ is equal to $\mathbf{n}_E(g^2/d)$, where $d = \gcd(n_2, g)$. We denote by $f^{(g)}$ to the associated Drinfeld newform to $E^{(g)}$.

The following lemma gives an upper bound for the Mordell-Weil rank of $E^{(g)}$.

Lemma 4.1. *With the notation above, we have that*

$$\text{rank}_{\mathbb{Z}}(E^{(g)}(K)) \leq \omega_K(n_2) + 2(\omega_K(n_1) + \omega_K(g)).$$

Proof. First of all, we notice that $E^{(g)}$ has multiplicative reduction at \mathfrak{p} if $\mathfrak{p} \mid n_2/d$, $E^{(g)}$ has additive reduction at \mathfrak{p} if $\mathfrak{p} \mid n_1g$, and otherwise $E^{(g)}$ has good reduction at \mathfrak{p} . Then by Proposition 2.3 we obtain that

$$\text{rank}_{\mathbb{Z}}(E^{(g)}(K)) \leq \omega_K(n_2/d) + 2(\omega_K(n_1) + \omega_K(g)),$$

since $\omega_K(n_2/d) \geq \omega_K(n_2)$ we obtain the desired result. \square

To find a lower bound for $\nu_2(m_{E^{(g)}})$, we need to relate $L(\text{Sym}^2 f^{(g)}, 2)$ and $L(\text{Sym}^2 f, 2)$, so, we can use Lemma 2.4 and the fact that $j_E = j_{E^{(g)}}$ (since this two elliptic curves are isomorphic in a quadratic extension of K), but before, we need the following lemma

Lemma 4.2. *Let \mathfrak{p} be a prime ideal of A and let $\left(\frac{\cdot}{\mathfrak{p}}\right) : \mathbb{F}_{\mathfrak{p}} \rightarrow \{-1, 0, 1\}$ be the extended Legendre symbol. Then*

$$a_{\mathfrak{p}}(E^{(g)}) = \left(\frac{g}{\mathfrak{p}}\right) a_{\mathfrak{p}}(E).$$

Proof. If $E^{(g)}$ has additive reduction at \mathfrak{p} , we have that $\mathfrak{p} \mid n_1$ or $\mathfrak{p} \mid g$, then $a_{\mathfrak{p}}(E^{(g)}) = 0$ and there is nothing to prove. On the other hand, assume that $E^{(g)}$ has multiplicative reduction at \mathfrak{p} . By Lemma 2.2 in [4] E has split multiplicative reduction at \mathfrak{p} if and only if $\left(\frac{-c_6(E)}{\mathfrak{p}}\right) = 1$, as a consequence, this quantity is equal to $a_{\mathfrak{p}}(E)$. Furthermore, since $c_6(E^{(g)}) = g^3 c_6(E)$, we have

$$a_{\mathfrak{p}}(E^{(g)}) = \left(\frac{-c_6(E^{(g)})}{\mathfrak{p}}\right) = \left(\frac{-g^3 c_6(E)}{\mathfrak{p}}\right) = \left(\frac{g}{\mathfrak{p}}\right) a_{\mathfrak{p}}(E).$$

Finally, assume that $\mathfrak{p} \nmid n^{(g)}$, Define $M = \{x \in \mathbb{F}_{\mathfrak{p}} : x^3 + Ax^2 + B \neq 0\}$. Consequently, we obtain

$$\begin{aligned} \#E_{\mathfrak{p}}^{(g)}(\mathbb{F}_{\mathfrak{p}}) &= |\mathfrak{p}| + 1 + \sum_{x \in M} \left(\frac{x^3 + Agx^2 + Bg^2x}{\mathfrak{p}}\right) \\ &= |\mathfrak{p}| + 1 + \sum_{x \in M} \left(\frac{g^3(x^3 + Ax^2 + Bx)}{\mathfrak{p}}\right) \\ &= |\mathfrak{p}| + 1 + \left(\frac{g}{\mathfrak{p}}\right) \sum_{x \in M} \left(\frac{x^3 + Ax^2 + Bx}{\mathfrak{p}}\right) \\ &= |\mathfrak{p}| + 1 - \left(\frac{g}{\mathfrak{p}}\right) a_{\mathfrak{p}}(E^{(g)}), \end{aligned}$$

by recalling the definition of $a_{\mathfrak{p}}(E)$ we get the desired result. \square

Proposition 4.3. *Let E be a modular elliptic curve with conductor \mathbf{n}_E and associated primitive newform f . Assume that E' is a quadratic twist of E , with conductor \mathbf{n}'_E and associated primitive newform f' , such that $\text{ord}_{\mathfrak{p}}(\mathbf{n}_E) \leq \text{ord}_{\mathfrak{p}}(\mathbf{n}'_E)$ for all \mathfrak{p} . Thus, there exist n_1, n_2, d, g square-free monic polynomials with $1 = \gcd(n_1, g)$, and $d = \gcd(n_2, g)$ such that $\mathbf{n}_E = (n_1^2 n_2) \infty$ and $\mathbf{n}'_E = \mathbf{n}_E g^2 / d$. Then one has*

$$L(\text{Sym}^2 f', 2) = L(\text{Sym}^2 f, 2) \frac{|d|}{|g|^3} \prod_{\mathfrak{p} \mid d} (|\mathfrak{p}|^2 - 1) \prod_{\mathfrak{p} \nmid g/d} ((|\mathfrak{p}| + 1)^2 - a_{\mathfrak{p}}(E)^2) (|\mathfrak{p}| - 1).$$

Proof. By Lemma 4.2 we have that when $\text{ord}_{\mathfrak{p}}(\mathbf{n}) = \text{ord}_{\mathfrak{p}}(\mathbf{n}')$ the local factors are equal, i.e. $L_{\mathfrak{p}}(\text{Sym}^2 f', 2) = L_{\mathfrak{p}}(\text{Sym}^2 f, 2)$. If $\mathfrak{p} \mid d$, we have that

$$L_{\mathfrak{p}}(\text{Sym}^2 f', s) = L_{\mathfrak{p}}(\text{Sym}^2 f, s)(1 - |\mathfrak{p}|^{-s}),$$

thus, at $s = 2$ we obtain

$$L_{\mathfrak{p}}(\text{Sym}^2 f', 2) = L_{\mathfrak{p}}(\text{Sym}^2 f, 2) \frac{1}{|\mathfrak{p}|^2} (|\mathfrak{p}|^2 - 1).$$

Finally, assume that $\mathfrak{p} \mid (g/d)$. The local factors are related as follows

$$L_{\mathfrak{p}}(\text{Sym}^2 f', s) = L_{\mathfrak{p}}(\text{Sym}^2 f, s) (1 - \alpha_{\mathfrak{p}}^2 |\mathfrak{p}|^{-s}) (1 - \overline{\alpha_{\mathfrak{p}}}^2 |\mathfrak{p}|^{-s}) (1 - |\mathfrak{p}|^{1-s}),$$

therefore at $s = 2$ we obtain

$$L_{\mathfrak{p}}(\text{Sym}^2 f', 2) = L_{\mathfrak{p}}(\text{Sym}^2 f, 2) \frac{1}{|\mathfrak{p}|^3} ((|\mathfrak{p}| + 1)^2 - a_{\mathfrak{p}}(E)^2) (|\mathfrak{p}| - 1),$$

putting all together, we achieve the desired result. \square

Proof of Theorem 1.3. Since E and $E^{(g)}$ are isomorphic over \mathbb{C}_{∞} , we have that $j_E = j_{E^{(g)}}$, thus by Lemma 2.4 we obtain

$$\nu_2(m_{E^{(g)}}) = \nu_2(m_E) + \nu_2(L(\text{Sym}^2 f^{(g)}, 2)) - \nu_2(L(\text{Sym}^2 f, 2)).$$

On the other hand, Proposition 4.3 implies that

$$\nu_2(L(\text{Sym}^2 f^{(g)}, 2)/L(\text{Sym}^2 f, 2)) = \sum_{\mathfrak{p}|d} \nu_2(|\mathfrak{p}|^2 - 1) + \sum_{\mathfrak{p}|g/d} \nu_2(((|\mathfrak{p}| + 1)^2 - a_{\mathfrak{p}}(E)^2)(|\mathfrak{p}| - 1)).$$

We know that $|\mathfrak{p}|^2 - 1 \equiv 0 \pmod{8}$, meanwhile $|\mathfrak{p}| - 1 \equiv 0 \pmod{2}$. As $E(K)[2]$ is non-trivial and it maps injectively into $E_{\mathfrak{p}}(\mathbb{F}_{\mathfrak{p}})$ for every prime $\mathfrak{p} \nmid n\infty$, then $|\mathfrak{p}| + 1 - a_{\mathfrak{p}}(E) \equiv 0 \pmod{2}$, which implies $(|\mathfrak{p}| + 1)^2 - a_{\mathfrak{p}}(E)^2 \equiv 0 \pmod{4}$. As a consequence

$$\nu_2(L(\text{Sym}^2 f^{(g)}, 2)) - \nu_2(L(\text{Sym}^2 f, 2)) \geq 3\omega_K(g).$$

Putting all together, we achieve the result.

$$\nu_2(m_{E^{(g)}}) \geq \nu_2(m_E) + 3\omega_K(g). \quad (7)$$

By Proposition 2.3 we know that $\text{rank}(E^{(g)}) \leq 2(\omega_K(\mathfrak{n}) + \omega_K(g))$. By our assumptions on g we obtain that

$$\nu_2(m_E) + 3\omega_K(g) \geq 2(\omega_K(\mathfrak{n}) + \omega_K(g)),$$

consequently, $\text{rank}(E^{(g)}) \leq \nu_2(m_{E^{(g)}})$. \square

Proof of Corollary 1.4. By Proposition 3.2 we have that $\nu_2(m_E) \geq \omega_K(\mathfrak{n}) - 3$. Since E is semi-stable, \mathfrak{n} is square-free, consequently, Lemma 4.1 implies that $\text{rank}(E^{(g)}) \leq \omega_K(\mathfrak{n}) + 2\omega_K(g)$. Using the equation (7), we have

$$\nu_2(m_{E^{(g)}}) \geq \nu_2(m_E) + 3\omega_K(g) \geq \omega_K(\mathfrak{n}) - 3 + 3\omega_K(g) \geq \omega_K(g) - 3 + \text{rank}(E^{(g)}), \quad (8)$$

hence Watkins' conjecture holds for $E^{(g)}$, whenever $\omega_K(d) \geq 3$. Furthermore, if a prime ideal \mathfrak{p} divides \mathfrak{n} and has non-split multiplicative reduction, by Theorem 3 in [1] $W_{\mathfrak{p}}f = f$, consequently, $\mathcal{W} = \mathcal{W}'$. Therefore, if every prime \mathfrak{p} which divides \mathfrak{n} has non-split multiplicative reduction and $E(K)[2] \cong \mathbb{Z}/2\mathbb{Z}$ Proposition 3.2 implies that $\nu_2(m_E) \geq \omega_K(\mathfrak{n}) - 1$, thus, equation (8) turns into

$$\nu_2(m_{E^{(g)}}) \geq \omega_K(g) - 1 + \text{rank}(E^{(g)}),$$

accordingly, Watkins' Conjecture holds for every square-free polynomial g of even degree. \square

ACKNOWLEDGEMENTS

I want to thank Professor Hector Pasten for suggesting this problem to me and for numerous helpful remarks. I was supported by ANID Doctorado Nacional 21190304.

REFERENCES

- [1] Atkin, A. O., & Lehner, J. (1970). *Hecke operators on $\Gamma_0(m)$* . Mathematische Annalen, 185(2), 134-160.
- [2] Breuil, C. Conrad, B. Diamond, F. and Taylor, R. *On the modularity of elliptic curves over \mathbb{Q} : wild 3-adic exercises*. Journal of the American Mathematical Society, pages 843–939, 2001.
- [3] Jerson Caro and Héctor Pastén. *Watkins’s conjecture for elliptic curves with non-split multiplicative reduction*. Preprint, 2021.
- [4] Conrad, B., Conrad, K. and Helfgott, H. (2005) *Root numbers and ranks in positive characteristic*. Advances in Mathematics, 198(2): 684–731.
- [5] Deligne, P. (1973) *Les constantes des équations fonctionnelles des fonctions L* , 501–597. Lecture Notes in Math., Vol. 349.
- [6] Drinfel’d, V. G. (1974). *Elliptic modules*. Mathematics of the USSR-Sbornik, 23(4), 561.
- [7] Dummigan, N. and Krishnamoorthy, S. (2013) *Powers of 2 in modular degrees of modular abelian varieties*. Journal of Number Theory, 133(2):501–522.
- [8] Esparza-Lozano, J. and Pastén, H. *A conjecture of Watkins for quadratic twists*. Proceedings of the American Mathematical Society, 149(6): 2381–2385, 2021.
- [9] Gekeler, E. and Reversat, A. (1996) *Jacobians of Drinfeld Modular Curves* J.Reine Angew. Math. 476, 27-93.
- [10] Grothendieck, A. (1964) *Formule de Lefschetz et rationalité des fonctions L* , Sémin. Bourbaki 279.
- [11] Pál, A. (2010) *The Manin constant of elliptic curves over function fields*. Algebra & Number Theory, 4(5), 509-545.
- [12] Papikian, M. (2002) *On the degree of modular parametrizations over function fields*. Journal of Number Theory, 97(2): 317–349.
- [13] Papikian, M. (2007) *Analogue of the degree conjecture over function fields*. Transactions of the American Mathematical Society, 359(7): 3483–3503.
- [14] Roberts, D. (2007) *Explicit descent on elliptic curves over function fields*. PhD thesis, University of Nottingham.
- [15] Schaefer, E. and Stoll, M. (2004) *How to do p -descent on an elliptic curve*. Transactions of the American Mathematical Society, 356(3): 1209–1231.
- [16] Schweizer, A. (1998). *Involutory elliptic curves over $\mathbb{F}_q(T)$* . Journal de théorie des nombres de Bordeaux, 10(1), 107-123
- [17] Shioda, T. (1992) *Some remarks on elliptic curves over function fields*. Astérisque, 209(12):99–114.
- [18] Tate, J. (1965). *On the conjectures of Birch and Swinnerton-Dyer and a geometric analog*. Séminaire Bourbaki, 9(306), 415-440.
- [19] Tate, J., & Shafarevich, I. R. (1967). *The rank of elliptic curves*. In Doklady Akademii Nauk (Vol. 175, No. 4, pp. 770-773). Russian Academy of Sciences.
- [20] Taylor, R. and Wiles, A. *Ring-theoretic properties of certain hecke algebras*. Annals of Mathematics, pages 553–572, 1995.
- [21] Ulmer, D. (2002) *Elliptic curves with large rank over function fields*. Annals of Mathematics, pages 295–315.
- [22] Ulmer, D. (2011). *Park City lectures on elliptic curves over function fields*. arXiv preprint arXiv:1101.1939.
- [23] Watkins, M. *Computing the modular degree of an elliptic curve*. Experimental Mathematics, 11(4): 487–502, 2002.
- [24] Wiles, A. *Modular elliptic curves and Fermat’s last theorem*. Annals of Mathematics, pages 443–551, 1995.
- [25] Yazdani, S. *Modular abelian varieties of odd modular degree*. Algebra and Number Theory, 5(1): 37–62, 2011.

Email address: jocaroc@uc.cl

FACULTAD DE MATEMÁTICAS, PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE, CAMPUS SAN JOAQUÍN, AVENIDA VICUÑA MACKENNA 4860, SANTIAGO, CHILE.