

TORUS LORENZ LINKS OBTAINED BY FULL TWISTS ALONG TORUS LINKS

THIAGO DE PAIVA

ABSTRACT. All knots are known to be hyperbolic, satellite, or torus knots, and one important family is Lorenz links, or T-links, which arise from dynamics. However, it remains difficult to determine the geometric type of a Lorenz link from a description via dynamics or as a T-link. In this paper, we consider those T-links that are torus links. We show that T-links obtained by full twists along torus links can never be torus links, aside from a family of cases. This addresses a question of Birman and Kofman.

1. INTRODUCTION

Lorenz links are knotted closed periodic orbits in the flow of the Lorenz system, which is a system of three ordinary differential equations discovered by meteorologist E. N. Lorenz when he was trying to find equations that model weather patterns [12].

Birman and Kofman found that *T-links* exactly coincide with Lorenz links [1, Theorem 1]. For $2 \leq r_1 < \dots < r_k$, and all $s_i > 0$, the T-link $T((r_1, s_1), \dots, (r_k, s_k))$ is defined to be the closure of the following braid

$$(\sigma_1 \sigma_2 \dots \sigma_{r_1-1})^{s_1} (\sigma_1 \sigma_2 \dots \sigma_{r_2-1})^{s_2} \dots (\sigma_1 \sigma_2 \dots \sigma_{r_k-1})^{s_k}.$$

Here σ_i is a standard generator of the braid group B_{r_k} .

Thurston showed that any non-trivial knot in the 3-sphere is either a *torus knot*, a *satellite knot*, or has *hyperbolic complement* [14]. These are called the *geometric types* of the knot.

Birman and Kofman asked the following question [1, Question 6].

Question 1.1. When are Lorenz links torus links?

Recall that a *torus link* is a link that can be embedded on the surface of an unknotted torus T in S^3 . A torus link is denoted by $T(p, q)$ with p, q integers. Here, p, q denote the number of times that $T(p, q)$ wraps around the meridian, longitude, respectively, of T . The torus link $T(p, q)$ has $\gcd(p, q)$ link components. Thus, if $\gcd(p, q) = 1$, then $T(p, q)$ is a knot.

The purpose of this paper is to answer Question 1.1 for T-links obtained by full twists along (p, q) -torus links. The answer to this question was not previously considered in general for this family of T-links. However, as a consequence of Lee's work on the geometric classification of twisted torus knots, it is known when a T-knot of the form $T((r, sr), (p, q))$ is a torus knot

for p, q coprime [8, Theorem 1.1], [10, Theorem 1.1]. Therefore, when the T-link $T((a_1, a_1s_1), \dots, (a_n, a_ns_n), (p, q))$ is a knot we assume that $n > 1$.

Our main theorem is the following.

Theorem 1.2. *Let $p, q, a_1, \dots, a_n, s_1, \dots, s_n$ be positive integers such that $1 < q < p$ and $1 < a_1 < \dots < a_n < p$ with $a_i \neq q$.*

- *If $\gcd(p, q) > 1$, then the T-link*

$$T((a_1, s_1a_1), (a_2, s_2a_2), \dots, (a_n, s_na_n), (p, q))$$

is never a torus link;

- *Otherwise, if $q < a_n$, or $p \neq bq + 1$, or $s_1 > 1$, or $a_2 \neq a_1 + 1$ for $b > 0$, then the T-knot*

$$T((a_1, s_1a_1), (a_2, s_2a_2), \dots, (a_n, s_na_n), (p, q))$$

is never a torus knot for $n > 1$.

Therefore, Theorem 1.2 answers Question 1.1 for T-links obtained by full twists along (p, q) -torus links, aside from a family of cases.

The answer to Question 1.1 is also related to the geometric classification of T-links. The geometric classification of T-links started with the study of the geometric classification of twisted torus knots. The twisted torus knots that are T-knots have the form $T((r, sr), (p, q))$ for p, q coprime. See [2], [8], [10], [9], and [11] for more information about the geometric classification of twisted torus knots. Recently Thiago de Paiva and Jessica Purcell extended these results to general T-knots obtained by full twists along (p, q) -torus knots [5], [6]. More precisely, they proved that there exists $B \gg 0$ such that if each $s_i > B$, the T-knot $T((a_1, a_1s_1), \dots, (a_n, a_ns_n), (p, q))$ is hyperbolic if and only if there is at least one $a_i > q$ that is not a multiple of q [5]. But, they didn't make B explicit. However, Theorem 1.2 will help to find this B as they are either hyperbolic or satellite knots if they are not torus knots.

1.1. Acknowledgment. I thank Jessica Purcell, my supervisor, and Sangyop Lee for some helpful tips.

2. MAIN SECTION

In this section we prove Theorem 1.2. We start by proving some lemmas that will be used later to prove this theorem.

Due to [6, Lemma 2.7], we assume that a_1, \dots, a_n are different from q for the T-link $T((a_1, s_1a_1), \dots, (a_n, s_na_n), (p, q))$.

Lemma 2.1. *Consider $q < a_n$. Then, there is an ambient isotopy that takes the T-link*

$$K = T((a_1, s_1a_1), \dots, (a_n, s_na_n), (p, q))$$

to an equivalent link given by the closure of the braid

$$\begin{aligned} & (\sigma_{a_n-1} \dots \sigma_{a_n-q+1})^{p-a_n} (\sigma_1 \dots \sigma_{a_1-1})^{s_1a_1} \dots (\sigma_1 \dots \sigma_{a_{n-1}-1})^{s_{n-1}a_{n-1}} \\ & (\sigma_1 \dots \sigma_{a_n-1})^{s_na_n+q}, \end{aligned}$$




FIGURE 1. The red strand in the second drawing can be replaced by a strand that goes below the third strand, as illustrated in the third drawing. This isotopy generalizes the isotopy in Figure 6 of [10].

where this isotopy happens in the complement of the braid axis of the last braid. Furthermore, K has braid index equal to a_n .

Proof. We start with the standard braid of K :

$$(\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1} \dots (\sigma_1 \dots \sigma_{a_n-1})^{s_n a_n} (\sigma_1 \dots \sigma_{p-1})^q.$$

If we pull the $(a_n - q + 1)$ -st strand around the braid closure anticlockwise, we see that it first reaches the bottom of the braid then passes through the sub-braid $(\sigma_1 \dots \sigma_{p-1})^q$ and finally ends in the $(a_n + 1)$ -st strand as $q < a_n$. So the $(a_n - q + 1)$ -st strand is connected to the $(a_n + 1)$ -st strand by a strand that goes one time around the braid closure. See the red strand in Figure 1. As this strand is an under strand, we can push it down and shrink it to reduce one strand from the last braid, as shown in Figure 1. After that, this strand becomes an under strand between the $(a_n - q + 1)$ and the $(a_n + 1)$ -st strand. Thus, we obtain the braid

$$B = (\sigma_{a_n-1} \dots \sigma_{a_n-q+1})(\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1} \dots (\sigma_1 \dots \sigma_{a_n-1})^{s_n a_n} (\sigma_1 \dots \sigma_{p-2})^q$$

after this isotopy.

As the sub-braid $(\sigma_{a_n-1} \dots \sigma_{a_n-q+1})$ of B is a braid on the leftmost a_n strands, the $(a_n - q + 1)$ -st strand is still connected to the $(a_n + 1)$ -st strand of B by an under strand that goes one time around the braid closure. Then we can apply the last isotopy again. After that, we obtain the braid

$$(\sigma_{a_n-1} \dots \sigma_{a_n-q+1})^2 (\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1} \dots (\sigma_1 \dots \sigma_{a_n-1})^{s_n a_n} (\sigma_1 \dots \sigma_{p-3})^q.$$

Therefore, after repeating this procedure $p - a_n - 2$ more times, we finally obtain the braid

$$(\sigma_{a_n-1} \dots \sigma_{a_n-q+1})^{p-a_n} (\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1} \dots (\sigma_1 \dots \sigma_{a_{n-1}-1})^{s_{n-1} a_{n-1}} (\sigma_1 \dots \sigma_{a_n-1})^{s_n a_n + q}.$$

As $s_n a_n + q > a_n$, the last braid has at least one full twist on the a_n strands. So, it follows from [7, Corollary 2.4] that the link K has braid index equal to a_n . \square

Lemma 2.2. *Consider $a_n < q < p$. Then, the T-link*

$$K = T((a_1, s_1 a_1), \dots, (a_n, s_n a_n), (p, q))$$

has braid index equal to q and is equivalent to the T-link

$$K' = T((a_1, s_1 a_1), \dots, (a_n, s_n a_n), (q, p)).$$

Proof. As $a_n < q < p$, the T-link K is equivalent to K' [1, Corollary 3]. Since $p > q$, K' has braid index equal to q by [7, Corollary 2.4]. \square

The next lemma was proved by Los [13, Corollary 1.2]. See also [3] for a quick proof sketch using the Markov theorem.

Lemma 2.3. *Consider β_1, β_2 two braids which have minimal braid index and are representations of the same torus link. Suppose that these two closed braids travel around the same braid axis C . Then, there is an isotopy in the complement of C that takes β_1 to β_2 .*

Lemma 2.4. *Consider $q < a_n < p$. Then, the T-link*

$$K = T((a_1, s_1 a_1), \dots, (a_n, s_n a_n), (p, q))$$

is not a torus link.

Proof. We prove this lemma by induction.

For when $n = 1$, consider that $K = T((a_1, s_1 a_1), (p, q))$ is a torus link. By Lemma 2.1, K is equivalent to the link given by the closure of the braid

$$B_1 = (\sigma_{a_1-1} \dots \sigma_{a_1-q+1})^{p-a_1} (\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1 + q}.$$

and has braid index equal to a_1 . So, K is the (a_1, k) -torus link with $k > 0$.

By Lemma 2.3, the link K_2 given by the closure of the braid

$$B_2 = B_1(\sigma_{a_1-1}^{-1} \dots \sigma_1^{-1})^{s_1 a_1} = (\sigma_{a_1-1} \dots \sigma_{a_1-q+1})^{p-a_1} (\sigma_1 \dots \sigma_{a_1-1})^q$$

is the $(a_1, k - s_1 a_1)$ -torus link. The braid B_2 represents the (p, q) -torus link. Thus, a_1 is equal to p or q , which is a contradiction.

Now we can consider that all T-links $T((a_1, s_1 a_1), \dots, (a_i, s_i a_i), (p, q))$ with $q < a_i$ are not torus links for $i < n$.

Suppose that $K = T((a_1, s_1 a_1), \dots, (a_n, s_n a_n), (p, q))$ is a torus link. By Lemma 2.1, K is equivalent to the link given by the closure of the braid

$$(\sigma_{a_n-1} \dots \sigma_{a_n-q+1})^{p-a_n} (\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1} \dots (\sigma_1 \dots \sigma_{a_{n-1}-1})^{s_{n-1} a_{n-1}} (\sigma_1 \dots \sigma_{a_n-1})^{s_n a_n + q}$$

and has braid index equal to a_n . So, K is the (a_n, k) -torus link with $k > 0$. By Lemma 2.3, the link K_1 given by the closure of the braid

$$B_1 = (\sigma_{a_n-1} \dots \sigma_{a_n-q+1})^{p-a_n} (\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1} \dots (\sigma_1 \dots \sigma_{a_{n-1}-1})^{s_{n-1} a_{n-1}} (\sigma_1 \dots \sigma_{a_n-1})^q$$

is the $(a_n, k - s_n a_n)$ -torus link. Since the last isotopy happened in the complement of the braid axis of the last braid, we can apply the isotopy of Lemma 2.1 in the backward direction to see that K_1 is the T-link $T((a_1, s_1 a_1), \dots, (a_{n-1}, s_{n-1} a_{n-1}), (p, q))$.

Suppose that $q > a_{n-1}$. Then, by Lemma 2.2, K_1 has braid index equal to q . So, $k - s_n a_n = q$. This implies that the sub-braid

$$(\sigma_{a_n-1} \dots \sigma_{a_n-q+1})^{p-a_n} (\sigma_1 \dots \sigma_{a_1-1})^{s_1 a_1} \dots (\sigma_1 \dots \sigma_{a_{n-1}-1})^{s_{n-1} a_{n-1}}$$

of B_1 is the trivial braid as K_1 is the torus link $T(a_n, q)$, but this is impossible as $a_{n-1}, \dots, a_1 > 1$. Thus, $q < a_{n-1}$. However, this contradicts the induction hypothesis.

Therefore, K is never a torus link. \square

Lemma 2.5. *Consider $n > 1$. If $s_1 > 1$ or $a_2 \neq a_1 + 1$, then the T-knot*

$$K = T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n), (p, bp + 1))$$

is not a torus knot for $b > 0$.

Proof. Since $p, bp + 1$ are coprime, K is a knot.

Consider that K is a torus knot. By [7, Corollary 2.4], K has braid index equal to p . By Lemma 2.3, the T-knot

$$T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n), (p, 1))$$

is also a torus knot. By a Markov move, the last T-knot is equivalent to the T-knot

$$T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n + 1)).$$

By [7, Corollary 2.4], this T-knot has braid index equal to a_n . Then, the T-knot

$$T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_{n-1}, s_{n-1} a_{n-1}), (a_n, 1)),$$

which is equivalent to the T-knot, by a Markov move,

$$T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_{n-1}, s_{n-1} a_{n-1} + 1)),$$

is also a torus knot. Continuing with this argument, we obtain that the T-knot

$$T((a_1, s_1 a_1), (a_2, s_2 a_2 + 1)),$$

which is equivalent to the T-knot $T((a_1, s_1 a_1), (s_2 a_2 + 1, a_2))$ by Lemma 2.2, is also a torus knot, but this is a contradiction by [8, Theorem 1.1] and [10, Theorem 1.1]. \square

Lemma 2.6. *Let $a_n < q < p$. Assume $n > 1$ when p and q are coprime. If $p \neq bq + 1$, or $s_1 > 1$, or $a_2 \neq a_1 + 1$ for $b > 0$, then the T-link*

$$K = T((a_1, s_1 a_1), \dots, (a_n, s_n a_n), (p, q))$$

is not a torus link.

Proof. As $a_n < q < p$, the T-link

$$K' = T((a_1, s_1 a_1), \dots, (a_n, s_n a_n), (q, p))$$

is equivalent to K and has braid index equal to q by Lemma 2.2.

Suppose that K is a torus link. Then, K is the (q, k) -torus link with $k > 0$. Let r be a positive integer such that $0 \leq p - rq < q$. The T-link

$$K'_{-rq} = T((a_1, s_1 a_1), \dots, (a_n, s_n a_n), (q, p - rq))$$

is the torus link $T(q, k - rq)$ by Lemma 2.3.

If $p - rq = 0$, then K'_{-rq} would be a splittable link. But, by Lemma 3.2 of [4], torus links are irreducible. So, K'_{-rq} can't be a torus link, which is a contradiction.

If $p - rq = 1$, then K' is a knot. However, by Lemma 2.5, K' can't be a torus knot.

Thus, $p - rq > 1$. It is not possible that $a_n > p - rq$ by Lemma 2.4. So, $a_n \leq p - rq$. If $a_n = p - rq$, then, by [6, Lemma 2.7], K'_{-rq} is equivalent to the T-link

$$T((a_1, s_1 a_1), \dots, (a_{n-1}, s_{n-1} a_{n-1}), (a_n, s_n a_n + q)),$$

which has braid index equal to a_n by Lemma 2.2. Thus, $a_n = k - rq$ as $a_n \neq q$. This implies that $k = a_n + rq = p$. Hence, $a_1, \dots, a_n = 0$ as K is the (p, q) -torus knot. This is a contradiction. If $a_n < p - rq$, then K'_{-rq} has braid index equal to $p - rq$ by Lemma 2.2. So, $p - rq = q$ or $k - rq$. It is not possible that $p - rq = q$ since $0 \leq p - rq < q$. If $p - rq = k - rq$, then $k = p$, which we have already ruled out.

Therefore, K is not a torus link. \square

Proposition 2.7. *Let $p, q, a_1, \dots, a_n, s_1, \dots, s_n$ be positive integers such that $1 < q < p$, $\gcd(p, q) > 1$, and $1 < a_1 < \dots < a_n < p$ with $a_i \neq q$. Then, the T-link*

$$K = T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n), (p, q))$$

is never a torus link.

Proof. If $q < a_n$, then K is not a torus link by Lemma 2.4.

Consider now that $a_n < q$. It is not possible that $p = bq + 1$ for $b > 0$ otherwise p and q would be coprime. Thus, from Lemma 2.6, K is not a torus link. \square

Proposition 2.8. *Let $p, q, a_1, \dots, a_n, s_1, \dots, s_n$ be positive integers such that $1 < q < p$, $\gcd(p, q) = 1$, and $1 < a_1 < \dots < a_n < p$ with $a_i \neq q$ and $n > 1$. If $q < a_n$, or $p \neq bq + 1$, or $s_1 > 1$, or $a_2 \neq a_1 + 1$ for $b > 0$, then the T-knot*

$$T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n), (p, q))$$

is never a torus knot.

Proof. If $q < a_n$, then it follows from Lemma 2.4 that K is not a torus knot. If $a_n < q$, under the hypotheses, K is not a torus knot from Lemma 2.6 either. \square

Theorem 1.2. *Let $p, q, a_1, \dots, a_n, s_1, \dots, s_n$ be positive integers such that $1 < q < p$ and $1 < a_1 < \dots < a_n < p$ with $a_i \neq q$.*

- *If $\gcd(p, q) > 1$, then the T-link*

$$T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n), (p, q))$$

is never a torus link;

- *Otherwise, if $q < a_n$, or $p \neq bq + 1$, or $s_1 > 1$, or $a_2 \neq a_1 + 1$ for $b > 0$, then the T-knot*

$$T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n), (p, q))$$

is never a torus knot for $n > 1$.

Proof. It follows from Propositions 2.7 and 2.8. \square

Corollary 2.9. *Let $p, q, a_1, \dots, a_n, s_1, \dots, s_n$ be positive integers such that $1 < q < p$, $\gcd(p, q) = 1$, and $1 < a_1 < \dots < a_n < p$ with $a_i \neq q$. If each s_i is greater than one, then the T-knot*

$$K = T((a_1, s_1 a_1), (a_2, s_2 a_2), \dots, (a_n, s_n a_n), (p, q))$$

is not a torus knot.

Proof. If $n = 1$, then the T-knot K is not a torus knot by [[8], Theorem 1.1]. Otherwise, the T-knot K is not a torus knot by Theorem 1.2. \square

REFERENCES

1. Joan Birman and Ilya Kofman, *A new twist on Lorenz links*, J. Topol. **2** (2009), no. 2, 227–248. MR 2529294 1, 4
2. Thiago de Paiva, *Unexpected essential surfaces among exteriors of twisted torus knots*, arXiv:2110.09873, to appear in Algebr. Geom. Topol., 2021. 2
3. ———, *Hyperbolic knots given by positive braids with at least two full twists*, Proceedings of the American Mathematical Society (2022). 4
4. ———, *Hyperbolic twisted torus links*, arXiv:2202.10975 (2022). 6
5. Thiago de Paiva and Jessica S. Purcell, *Lorenz links obtained by twisting*, in preparation. 2
6. ———, *Satellites and Lorenz knots*, arxiv:2103.09500, 2021. 2, 6
7. John Franks and R. F. Williams, *Braids and the Jones polynomial*, Trans. Amer. Math. Soc. **303** (1987), no. 1, 97–108. MR 896009 4, 5
8. Sangyop Lee, *Torus knots obtained by twisting torus knots*, Algebr. Geom. Topol. **15** (2015), no. 5, 2819–2838. MR 3426694 2, 5, 7
9. ———, *Satellite knots obtained by twisting torus knots: hyperbolicity of twisted torus knots*, Int. Math. Res. Not. IMRN (2018), no. 3, 785–815. MR 3801447 2
10. ———, *Positively twisted torus knots which are torus knots*, J. Knot Theory Ramifications **28** (2019), no. 3, 1950023, 13. MR 3938086 2, 3, 5
11. Sangyop Lee and Thiago de Paiva, *Torus knots obtained by negatively twisting torus knots*, arXiv:2108.10641, 2021. 2

12. E.Ñ. Lorenz, *Deterministic non-periodic flow*, J. Atmos. Sci. **20** (1963), 130–141. 1
13. Jérôme E. Los, *Knots, braid index and dynamical type*, Topology **33** (1994), no. 2, 257–270. MR 1273785 4
14. William P. Thurston, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S.) **6** (1982), no. 3, 357–381. MR 648524 1

SCHOOL OF MATHEMATICS, MONASH UNIVERSITY, VIC 3800, AUSTRALIA

Email address: `thiago.depaivasouza@monash.edu`