F-ALGEBROIDS AND DEFORMATION QUANTIZATION VIA PRE-LIE ALGEBROIDS

JOHN ALEXANDER CRUZ MORALES, JIEFENG LIU, AND YUNHE SHENG

ABSTRACT. In this paper, first we introduce a new approach to the notion of F-algebroids, which is a generalization of F-manifold algebras and F-manifolds, and show that F-algebroids are the corresponding semi-classical limits of pre-Lie formal deformations of commutative associative algebroids. Then we use the deformation cohomology of pre-Lie algebroids to study pre-Lie infinitesimal deformations and extension of pre-Lie *n*-deformations to pre-Lie (n+1)-deformations of a commutative associative algebroid. Next we develop the theory of Dubrovin's dualities of F-algebroids with eventual identities and use Nijenhuis operators on F-algebroids to construct new F-algebroids. Finally we introduce the notion of pre-F-algebroids, which is a generalization of F-manifolds with compatible flat connections. Dubrovin's dualities of pre-F-algebroids with eventual identities, Nijenhuis operators on pre-F-algebroids and their applications to integral systems are discussed.

CONTENTS

1. Introduction	1
2. <i>F</i> -algebroids	3
3. Pre-Lie deformation quantization of commutative associative algebroids	7
4. Some constructions of <i>F</i> -algebroids	12
4.1. (Pseudo-)Eventual identities and Dubrovin's dual of <i>F</i> -algebroids	12
4.2. Nijenhuis operators and deformed <i>F</i> -algebroids	15
5. Pre- <i>F</i> -algebroids and eventual identities	17
5.1. Some Properties of pre- <i>F</i> -algebroids	17
5.2. Eventual identities of pre- <i>F</i> -algebroids	21
5.3. Nijenhuis operators and deformed pre- <i>F</i> -algebroids	26
5.4. Applications to integral systems	28
References	29

1. INTRODUCTION

The concept of Frobenius manifolds was introduced by Dubrovin in [11] as a geometrical manifestation of the Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) associativity equations in the 2-dimensional topological field theories. Hertling and Manin weakened the conditions of a Frobenius manifold and introduced the notion of an *F*-manifold in [13]. Any Frobenius manifold has an underlying *F*-manifold structure. *F*-manifolds appear in many fields of mathematics

Key words and phrases. F-algebroid, pre-F-algebroid, eventual identity, Nijenhuis operator.

such as singularity theory [12], integrable systems [1, 3, 4, 8, 9, 21, 23], quantum K-theory [17], information geometry [29], operad [25] and so on.

The notion of a Lie algebroid was introduced by Pradines in 1967, which is a generalization of Lie algebras and tangent bundles. Just as Lie algebras are the infinitesimal objects of Lie groups, Lie algebroids are the infinitesimal objects of Lie groupoids. See [24] for general theory about Lie algebroids. Lie algebroids are now an active domain of research, with applications in various parts of mathematics, such as geometric mechanics, foliation theory, Poisson geometry, differential equations, singularity theory, operad and so on. The notion of a pre-Lie algebroid (also called a left-symmetric algebroid or a Koszul-Vinberg algebroid) is a geometric generalization of a pre-Lie algebra. Pre-Lie algebras arose from the study of convex homogeneous cones, affine manifolds and affine structures on Lie groups, deformation and cohomology theory of associative algebras and then appear in many fields in mathematics and mathematical physics. See the survey article [5] for more details on pre-Lie algebras and [18, 19, 27, 28] for more details on cohomology and applications of pre-Lie algebroids. In [10], Dotsenko showed that the graded object of the filtration of the operad encoding pre-Lie algebras is the operad encoding *F*-manifold algebras, where the notion of an *F*-manifold algebra is the underlying algebraic structure of an *F*-manifold. In [20], the notion of pre-Lie formal deformations of commutative associative algebras was introduced and it was shown that *F*-manifold algebras are the corresponding semi-classical limits. This result is parallel to that the semi-classical limit of an associative formal deformation of a commutative associative algebra is a Poisson algebra. In this paper, we introduce the notion of *F*-algebroids, which is a generalization of *F*-

In this paper, we introduce the notion of *F*-algebroids, which is a generalization of *F*-manifold algebras and *F*-manifolds. There is a slight difference between this *F*-algebroid and the one introduced in [7]. We introduce the notion of pre-Lie formal deformations of commutative associative algebroids and show that *F*-algebroids are the corresponding semi-classical limits. Viewing a commutative associative algebroid as a pre-Lie algebroid, we show that pre-Lie infinitesimal deformation and extension of pre-Lie *n*-deformations to pre-Lie (n + 1)-deformations of a commutative associative algebroid are classified by the second and the third cohomology groups of the pre-Lie algebroid respectively.

cohomology groups of the pre-Lie algebroid tare enablined by the second that the third cohomology groups of the pre-Lie algebroid respectively. *F*-manifolds with eventual identities were introduced by Manin in [26] and then were studied systematically by David and Strachan in [9]. In this paper, we generalize Dubrovin's dualities of *F*-manifolds with eventual identities to the case of *F*-algebroids. We introduce the notion of (pseudo-)eventual identities on *F*-algebroids and develop the theory of Dubrovin's dualities of *F*-algebroids with eventual identities. We introduce the notion of Nijenhuis operators on *F*algebroids and use them to construct new *F*-algebroids. In particular, a pseudo-eventual identity naturally gives a Nijenhuis operator on an *F*-algebroid.

naturally gives a Nijenhuis operator on an *F*-algebroid. The notion of an *F*-manifold with a compatible flat connection was introduced by Manin in [26]. Applications of *F*-manifolds with compatible flat connections also appeared in Painlevé equations [2, 4, 14, 21] and integral systems [1, 3, 15, 22, 23]. In this paper, we introduce the notion of pre-*F*-algebroids, which is a generalization of *F*-manifolds with compatible flat connections. A pre-*F*-algebroid gives rise to an *F*-algebroid. We also study pre-*F*-algebroids with eventual identities and give a characterization of such eventual identities. Furthermore, The theory of Dubrovin's dualities of pre-*F*-algebroids with eventual identities were developed. We introduce the notion of a Nijenhuis operator on a pre-*F*-algebroid, and show that a Nijenhuis operator gives rise to a deformed pre-*F*-algebroid. Mirror symmetry, roughly speaking, is a duality between symplectic and complex geometry. The theory of Frobenius and F-manifolds plays an important role in this duality. We expect that the notion of F-algebroids might also be relevant in understanding the mirror phenomenon. In particular, the Dubrovin's dual of F-algebroids constructed in this paper should be related to the mirror construction along the way the Dubrovin's dual of Frobenius manifolds is related, at least in some situations, with mirror symmetry. More precisely the question is: Could we consider the construction of Dubrovin's dual of F-algebroids as a kind of mirror construction? In order to answer the question above, we might need to add some extra structures to F-algebroids and include those structures in the construction of the Dubrovin's dual. This would allow us to give a comprehensible interpretation of our construction as a manifestation of a mirror phenomenon. We want to follow this line of thought in future works.

The paper is organized as follows. In Section 2, we introduce the notion of F-algebroids and give some constructions of F-algebroids including the action F-algebroids and direct product F-algebroids. In particular, we show that Poisson manifolds give rise to action F-algebroids naturally. In Section 3, we study pre-Lie formal deformations of a commutative associative algebroid, whose semi-classical limits are F-algebroids. We show that the equivalence classes of pre-Lie infinitesimal deformations of a commutative associative algebroid A are classified by the second cohomology group in the deformation cohomology of A. Furthermore, we study extensions of pre-Lie n-deformations to pre-Lie (n+1)-deformations of a commutative associative algebroid A and show that a pre-Lie *n*-deformation can be extendable if and only if its obstruction class in the third cohomology group of the commutative associative algebroid A is trivial. In Section 4, we first study Dubrovin's duality of F-algebroids with eventual identities. Then we use Nijenhuis operators on F-algebroids to construct deformed F-algebroids. In Section 5, first we introduce the notion of a pre-F-algebroid, and show that a pre-F-algebroid gives rise to an F-algebroid. Then we study Dubrovin's duality of pre-F-algebroids with eventual identities. Finally, we introduce the notion of a Nijenhuis operator on a pre-F-algebroid, and show that a Nijenhuis operator on a pre-F-algebroid gives rise to a deformed pre-F-algebroid. Finally we give some applications to integrable systems.

Acknowledgements. This research was supported by NSFC (11922110,11901501). The second author also supported by the National Key Research and Development Program of China (2021YFA1002000). The first author wants to thank Max Planck Institute for Mathematics where the last part of this work was written for its hospitality and financial support.

2. F-Algebroids

In this section, we introduce the notion of F-algebroids, which are generalizations of F-manifolds and F-manifold algebras. We give some constructions of F-algebroids including the action F-algebroids and direct product F-algebroids.

Definition 2.1. An *F*-manifold algebra is a triple $(g, [-, -], \cdot)$, where (g, \cdot) is a commutative associative algebra and (g, [-, -]) is a Lie algebra, such that for all $x, y, z, w \in g$, the Hertling-Manin relation holds:

(1)
$$P_{x \cdot y}(z, w) = x \cdot P_y(z, w) + y \cdot P_x(z, w),$$

where $P_x(y, z)$ is defined by

(2)
$$P_{x}(y,z) = [x, y \cdot z] - [x, y] \cdot z - y \cdot [x, z].$$

Example 2.2. Any Poisson algebra is an *F*-manifold algebra.

Definition 2.3. An *F*-manifold is a pair (M, \bullet) , where *M* is a smooth manifold and \bullet is a $C^{\infty}(M)$ -bilinear, commutative, associative multiplication on the tangent bundle *TM* such that $(\mathfrak{X}(M), [-, -]_{\mathfrak{X}(M)}, \bullet)$ is an *F*-manifold algebra, where $[-, -]_{\mathfrak{X}(M)}$ is the Lie bracket of vector fields.

The notion of Lie algebroids was introduced by Pradines in 1967, as a generalization of Lie algebras and tangent bundles. See [24] for the general theory about Lie algebroids.

Definition 2.4. A Lie algebroid structure on a vector bundle $A \rightarrow M$ is a pair that consists of a Lie algebra structure $[-, -]_A$ on the section space $\Gamma(A)$ and a vector bundle morphism $a_A : A \rightarrow TM$, called the anchor, such that the following relation is satisfied:

 $[X, fY]_A = f[X, Y]_A + a_A(X)(f)Y, \quad \forall X, Y \in \Gamma(A), \ f \in C^{\infty}(M).$

We usually denote a Lie algebroid by $(A, [-, -]_A, a_A)$, or A if there is no confusion.

Definition 2.5. A commutative associative algebroid is a vector bundle A over M equipped with a $C^{\infty}(M)$ -bilinear, commutative, associative multiplication \cdot_A on the section space $\Gamma(A)$.

We denote a commutative associative algebroid by (A, \cdot_A) .

In the following, we give the notion of *F*-algebroids, which are generalizations of *F*-manifold algebras and *F*-manifolds.

Definition 2.6. An *F*-algebroid is a vector bundle *A* over *M* equipped with a bilinear operation $\cdot_A : \Gamma(A) \times \Gamma(A) \to \Gamma(A)$, a skew-symmetric bilinear bracket $[-, -]_A : \Gamma(A) \times \Gamma(A) \to \Gamma(A)$, and a bundle map $a_A : A \to TM$, called the anchor, such that $(A, [-, -]_A, a_A)$ is a Lie algebroid, (A, \cdot_A) is a commutative associative algebroid and $(\Gamma(A), [-, -]_A, \cdot_A)$ is an *F*-manifold algebra.

We denote an *F*-algebroid by $(A, [-, -]_A, \cdot_A, a_A)$.

Remark 2.7. In [7], the authors had already defined an *F*-algebroid. There is a slight difference between the above definition of an *F*-algebroid and that one. In [7], it is assumed that the base manifold has an *F*-manifold structure (M, \bullet) . An *F*-algebroid defined in [7] is a vector bundle *A* over *M* equipped with a bilinear operation $\cdot_A : \Gamma(A) \times \Gamma(A) \to \Gamma(A)$, a skew-symmetric bilinear bracket $[-, -]_A : \Gamma(A) \times \Gamma(A) \to \Gamma(A)$, and a bundle map $a_A : A \to TM$, such that $(A, [-, -]_A, a_A)$ is a Lie algebroid, (A, \cdot_A) is a commutative associative algebroid, $(\Gamma(A), [-, -]_A, \cdot_A)$ is an *F*manifold algebra and

(3) $a_A(X \cdot_A Y) = a_A(X) \bullet a_A(Y), \quad \forall X, Y \in \Gamma(A).$

Example 2.8. Any *F*-manifold algebra is an *F*-algebroid over a point. Let (M, \bullet) be an *F*-manifold. Then $(TM, [-, -]_{\mathfrak{X}(M)}, \bullet, \mathrm{Id})$ is an *F*-algebroid.

Definition 2.9. Let $(A, [-, -]_A, \cdot_A, a_A)$ and $(B, [-, -]_B, \cdot_B, a_B)$ be *F*-algebroids on *M*. A bundle map $\varphi : A \longrightarrow B$ is called a **homomorphism** of *F*-algebroids, if for all $X, Y \in \Gamma(A)$, the following conditions are satisfied:

 $\varphi(X \cdot_A Y) = \varphi(X) \cdot_B \varphi(Y), \quad \varphi([X, Y]_A) = [\varphi(X), \varphi(Y)]_B, \quad a_B \circ \varphi = a_A.$

Definition 2.10. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid. A section $e \in \Gamma(A)$ is called the **identity** if $e \cdot_A X = X$ for all $X \in \Gamma(A)$. We denote an *F*-algebroid $(A, [-, -]_A, \cdot_A, a_A)$ with an identity e by $(A, [-, -]_A, \cdot_A, e, a_A)$.

Proposition 2.11. Let $(A, [-, -]_A, a_A)$ be a Lie algebroid equipped with a $C^{\infty}(M)$ -bilinear, commutative, associative multiplication $\cdot_A : \Gamma(A) \times \Gamma(A) \to \Gamma(A)$. Define

(4)
$$\Phi(X, Y, Z, W) := P_{X \cdot_A Y}(Z, W) - X \cdot_A P_Y(Z, W) - Y \cdot_A P_X(Z, W), \quad \forall X, Y, Z, W \in \Gamma(A),$$

where P is given by (2). Then Φ is a tensor field of type (4, 1) and

(5)
$$\Phi(X, Y, Z, W) = \Phi(Y, X, Z, W) = \Phi(X, Y, W, Z).$$

Proof. By the commutativity of the associative multiplication \cdot_A , we have

 $\Phi(X,Y,Z,W)=\Phi(Y,X,Z,W)=\Phi(X,Y,W,Z).$

To prove that Φ is a tensor field of type (4, 1), we only need to show

$$\Phi(fX, Y, Z, W) = \Phi(X, Y, fZ, W) = f\Phi(X, Y, Z, W).$$

By a direct calculation, we have

$$\begin{split} & \Phi(fX, Y, Z, W) \\ = & [f(X \cdot_A Y), Z \cdot_A W]_A - Z \cdot_A [f(X \cdot_A Y), W]_A - W \cdot_A [f(X \cdot_A Y), Z]_A \\ & -f(X \cdot_A P_Y(Z, W)) - Y \cdot_A ([fX, Z \cdot_A W]_A - Z \cdot_A [fX, W]_A - W \cdot_A [fX, Z]_A) \\ = & fP_{X \cdot_A Y}(Z, W) - a_A(Z \cdot_A W)(f)(X \cdot_A Y) + a_A(W)(f)(X \cdot_A Y \cdot_A Z) \\ & +a_A(Z)(f)(X \cdot_A Y \cdot_A W) - f(X \cdot_A P_Y(Z, W)) - f(Y \cdot_A P_X(Z, W)) \\ & +a_A(Z \cdot_A W)(f)(X \cdot_A Y) - a_A(W)(f)(X \cdot_A Y \cdot_A Z) - a_A(Z)(f)(X \cdot_A Y \cdot_A W) \\ = & f\Phi(X, Y, Z, W). \end{split}$$

Similarly, we also have $\Phi(X, Y, fZ, W) = f\Phi(X, Y, Z, W)$.

Proposition 2.12. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid with an identity *e*. Then

$$P_e(X,Y) = 0.$$

Proof. It follows from (1) directly.

Definition 2.13. Let $(\mathfrak{g}, [-, -], \cdot)$ be an *F*-manifold algebra. An **action** of \mathfrak{g} on a manifold *M* is a linear map $\rho : \mathfrak{g} \longrightarrow \mathfrak{X}(M)$ from \mathfrak{g} to the space of vector fields on *M*, such that for all $x, y \in \mathfrak{g}$, we have

$$\rho([x, y]) = [\rho(x), \rho(y)]_{\mathfrak{X}(M)}.$$

Given an action of g on M, let $A = M \times g$ be the trivial bundle. Define an anchor map $a_{\rho} : A \longrightarrow TM$, a multiplication $\cdot_{\rho} : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ and a bracket $[-, -]_{\rho} : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ by

(6)
$$a_{\rho}(m,u) = \rho(u)_m, \quad \forall m \in M, u \in \mathfrak{g}.$$

(7)
$$X \cdot_{\rho} Y = X \cdot Y,$$

(8)
$$[X,Y]_{\rho} = \mathcal{L}_{\rho(X)}Y - \mathcal{L}_{\rho(Y)}X + [X,Y], \quad \forall X,Y \in \Gamma(A).$$

where $X \cdot Y$ and [X, Y] are the pointwise $C^{\infty}(M)$ -bilinear multiplication and bracket, respectively.

Proposition 2.14. With the above notations, $(A = M \times g, [-, -]_{\rho}, \cdot_{\rho}, a_{\rho})$ is an *F*-algebroid, which is called an **action** *F*-algebroid, where $[-, -]_{\rho}, \cdot_{\rho}$ and a_{ρ} are given by (8), (7) and (6), respectively.

Proof. Note that the multiplication \cdot_{ρ} is a $C^{\infty}(M)$ -bilinear, commutative and associative multiplication and $(A, [-, -]_{\rho}, a_{\rho})$ is a Lie algebroid. By Proposition 2.11 and the fact that g is an *F*-manifold algebra, for all $u_1, u_2, u_3, u_4 \in \mathfrak{g}$ and $f_1, f_2, f_3, f_4 \in C^{\infty}(M)$, we have

$$\Phi(f_1u_1, f_2u_2, f_3u_3, f_4u_4) = f_1f_2f_3f_4\Phi(u_1, u_2, u_3, u_4) = 0,$$

which implies that $(\Gamma(A), [-, -]_{\rho}, \cdot_{\rho})$ is an *F*-manifold algebra. Thus $(A, [-, -]_{\rho}, \cdot_{\rho}, a_{\rho})$ is an *F*-algebroid.

Example 2.15. Let g be a 2-dimensional vector space with basis $\{e_1, e_2\}$. Then $(g, [-, -], \cdot)$ with the non-zero multiplication \cdot and the bracket [-, -]

$$e_1 \cdot e_1 = e_1, \quad e_1 \cdot e_2 = e_2 \cdot e_1 = e_2, \quad [e_1, e_2] = e_2$$

is an *F*-manifold algebra with the identity e_1 . Let (t_1, t_2) be the canonical coordinate systems on \mathbb{R}^2 . It is straightforward to check that the map $\rho : \mathfrak{g} \longrightarrow \mathfrak{X}(\mathbb{R}^2)$ defined by

$$\rho(e_1) = t_2 \frac{\partial}{\partial t_2}, \quad \rho(e_2) = t_2 \frac{\partial}{\partial t_1} + t_2^2 \frac{\partial}{\partial t_2}$$

is an action of the *F*-manifold algebra g on \mathbb{R}^2 . Then $(A = \mathbb{R}^2 \times g, [-, -]_{\rho}, \cdot_{\rho}, a_{\rho})$ is an *F*-algebroid with an identity $1 \otimes e_1$, where $[-, -]_{\rho}, \cdot_{\rho}$ and a_{ρ} are given by

$$\begin{aligned} a_{\rho}(m,c_{1}e_{1}+c_{2}e_{2}) &= (c_{1}t_{2}\frac{\partial}{\partial t_{2}}+c_{2}t_{2}\frac{\partial}{\partial t_{1}}+c_{2}t_{2}^{2}\frac{\partial}{\partial t_{2}})\mid_{m}, \quad \forall \ m \in \mathbb{R}^{2}, \\ f \otimes (c_{1}e_{1}) \cdot_{\rho} g \otimes (c_{2}e_{i}) &= (fg) \otimes (c_{1}c_{2}e_{i}), \quad f \otimes (c_{1}e_{2}) \cdot_{\rho} g \otimes (c_{2}e_{2}) = 0, \\ [f \otimes (c_{1}e_{1}), g \otimes (c_{2}e_{2})]_{\rho} &= fc_{1}t_{2}\frac{\partial g}{\partial t_{2}} \otimes (c_{2}e_{2}) - gc_{2}(t_{2}\frac{\partial f}{\partial t_{1}}+t_{2}^{2}\frac{\partial f}{\partial t_{2}}) \otimes (c_{1}e_{1}) + fg \otimes (c_{1}c_{2}[e_{1},e_{2}]), \\ \text{where } f, g \in C^{\infty}(\mathbb{R}^{2}), \ c_{1}, c_{2} \in \mathbb{R}, \ i \in \{1,2\}. \end{aligned}$$

Let (M, π) be a Poisson manifold and $(C^{\infty}(M), \cdot, \{-, -\})$ be the corresponding Poisson algebra. Then for a given function f on M, there is a unique vector field H_f on M, called the Hamiltonian vector field of f, such that for any $g \in C^{\infty}(M)$, we have $H_f(g) = \{f, g\}$. Furthermore, the map $H : C^{\infty}(M) \to \mathfrak{X}(M)$ defined by $f \mapsto H_f$ is a homomorphism from the Lie algebra $C^{\infty}(M)$ of smooth functions to the Lie algebra of smooth vector fields, i.e.

$$H_{\{f,g\}} = [H_f, H_g]_{\mathfrak{X}(M)}, \quad \forall f, g \in C^{\infty}(M).$$

Proposition 2.16. Let (M, π) be a Poisson manifold and $(C^{\infty}(M), \cdot, \{-, -\})$ be the corresponding Poisson algebra. Then $(A = M \times C^{\infty}(M), [-, -]_H, \cdot_H, a_H)$ is an *F*-algebroid, where the anchor map a_H , multiplication \cdot_H and bracket $[-, -]_H$ are given by

$$\begin{aligned} a_H(m, f) &= H_f(m), \quad \forall \ m \in M, \ f \in C^{\infty}(M), \\ (f_1 \otimes g_1) \cdot_H (f_2 \otimes g_2) &= (f_1 f_2) \otimes (g_1 g_2), \\ [f_1 \otimes g_1, f_2 \otimes g_2]_H &= f_1 \{g_1, f_2\} \otimes g_2 - f_2 \{g_2, f_1\} \otimes g_1 + f_1 f_2 \otimes \{g_1, g_2\} \end{aligned}$$

where f_1, f_2, g_1, g_2 are smooth functions on M.

Let A_1 and A_2 be vector bundles over M_1 and M_2 respectively. Denote the projections from $M_1 \times M_2$ to M_1 and M_2 by pr₁ and pr₂ respectively. The product vector bundle $A_1 \times A_2 \rightarrow M_1 \times M_2$ can be regarded as the Whitney sum over $M_1 \times M_2$ of the pullback vector bundles pr¹₁ A_1 and pr¹₂ A_2 . Sections of pr¹₁ A_1 are of the form $\sum u_i \otimes X_i^1$, where $u_i \in C^{\infty}(M_1 \times M_2)$ and $X_i^1 \in \Gamma(A_1)$.

Similarly, sections of $\operatorname{pr}_2^! A_2$ are of the form $\sum u_i' \otimes X_i^2$, where $u_i' \in C^{\infty}(M_1 \times M_2)$ and $X_i^2 \in \Gamma(A_2)$. The tangent bundle $T(M_1 \times M_2)$ may in the same way be regarded as the Whitney sum $\operatorname{pr}_1^!(TM_1) \oplus \operatorname{pr}_2^!(TM_2)$. Let $(A_1, [-, -]_{A_1}, a_{A_1})$ and $(A_2, [-, -]_{A_2}, a_{A_2})$ be two Lie algebroids over the base manifolds M_1 and M_2 respectively. We define the anchor $\mathfrak{a} : A_1 \times A_2 \longrightarrow T(M_1 \times M_2)$ by

$$\mathfrak{a}(\sum (u_i \otimes X_i^1) \oplus \sum (u_j' \otimes X_j^2)) = \sum (u_i \otimes a_{A_1}(X_i^1)) \oplus \sum (u_j' \otimes a_{A_2}(X_j^2)).$$

And the Lie bracket on $A_1 \times A_2$ is determined by the following relations with the Leibniz rule

$$\begin{bmatrix} 1 \otimes X^1, 1 \otimes Y^1 \end{bmatrix} = 1 \otimes [X^1, Y^1]_{A_1}, \quad \begin{bmatrix} 1 \otimes X^1, 1 \otimes Y^2 \end{bmatrix} = 0,$$
$$\begin{bmatrix} 1 \otimes X^2, 1 \otimes Y^2 \end{bmatrix} = 1 \otimes [X^2, Y^2]_{A_2}, \quad \begin{bmatrix} 1 \otimes X^2, 1 \otimes Y^1 \end{bmatrix} = 0,$$

for $X^1, Y^1 \in \Gamma(A_1), X^2, Y^2 \in \Gamma(A_2)$. See [24] for more details of the direct product Lie algebroids.

Proposition 2.17. Let $(A_1, [-, -]_{A_1}, \cdot_{A_1}, a_{A_1})$ and $(A_2, [-, -]_{A_2}, \cdot_{A_2}, a_{A_2})$ be two *F*-algebroids over the base manifolds M_1 and M_2 respectively. Then $(A_1 \times A_2, [-, -]], \diamond, \mathfrak{a})$ is an *F*-algebroid over $M_1 \times M_2$, where for

$$X = \sum (u_i \otimes X_i^1) \oplus \sum (u'_j \otimes X_j^2), \quad Y = \sum (v_k \otimes Y_k^1) \oplus \sum (v'_l \otimes Y_l^2),$$

the associative multiplication \diamond is defined by

$$X \diamond Y = \sum (u_i v_k \otimes (X_i^1 \cdot_{A_1} Y_k^1)) \oplus \sum (u'_j v'_l \otimes (X_j^2 \cdot_{A_2} Y_l^2)).$$

Proof. It follows from straightforward verifications.

The *F*-algebroid $(A_1 \times A_2, [-, -]], \diamond, \mathfrak{a})$ is called the direct product *F*-algebroid.

3. PRE-LIE DEFORMATION QUANTIZATION OF COMMUTATIVE ASSOCIATIVE ALGEBROIDS

In this section, we study pre-Lie formal deformations of a commutative associative algebroid, whose semi-classical limits are F-algebroids. Viewing the commutative associative algebroid A as a pre-Lie algebroid, we show that the equivalence classes of pre-Lie infinitesimal deformations of a commutative associative algebroid A are classified by the second cohomology group in the deformation cohomology of A and a pre-Lie n-deformation can be extended to a pre-Lie (n + 1)-deformation if and only if its obstruction class in the third cohomology group of A is trivial.

Definition 3.1. A pre-Lie algebra is a pair (g, *), where g is a vector space and $* : g \otimes g \longrightarrow g$ is a bilinear multiplication satisfying that for all $x, y, z \in g$, the associator

(9)
$$(x, y, z) \triangleq x * (y * z) - (x * y) * z$$

is symmetric in x, y, i.e.

$$(x, y, z) = (y, x, z)$$
, or equivalently, $x * (y * z) - (x * y) * z = y * (x * z) - (y * x) * z$.

Definition 3.2. ([19, 27]) A **pre-Lie algebroid** structure on a vector bundle $A \rightarrow M$ is a pair that consists of a pre-Lie algebra structure $*_A$ on the section space $\Gamma(A)$ and a vector bundle morphism $a_A : A \rightarrow TM$, called the anchor, such that for all $f \in C^{\infty}(M)$ and $X, Y \in \Gamma(A)$, the following conditions are satisfied:

(i)
$$X *_A (fY) = f(X *_A Y) + a_A(X)(f)Y$$
,

(ii) $(fX) *_A Y = f(X *_A Y)$.

We usually denote a pre-Lie algebroid by $(A, *_A, a_A)$. Any pre-Lie algebra is a pre-Lie algebroid over a point.

A connection ∇ on a manifold M is said to be **flat** if the torsion and the curvature of the connection ∇ vanish identically. A manifold M endowed with a flat connection ∇ is called a **flat manifold**.

Example 3.3. Let *M* be a manifold with a flat connection ∇ . Then (TM, ∇, Id) is a pre-Lie algebroid whose sub-adjacent Lie algebroid is exactly the tangent Lie algebroid. We denote this pre-Lie algebroid by $T_{\nabla}M$.

Proposition 3.4. ([19]) *Let* $(A, *_A, a_A)$ *be a pre-Lie algebroid. Define a skew-symmetric bilinear bracket operation* $[-, -]_A$ *on* $\Gamma(A)$ *by*

(10)
$$[X,Y]_A = X *_A Y - Y *_A X, \quad \forall X,Y \in \Gamma(A).$$

Then $(A, [-, -]_A, a_A)$ *is a Lie algebroid, and denoted by* A^c *, called the* **sub-adjacent Lie algebroid** *of* $(A, *_A, a_A)$.

Definition 3.5. Let *E* be a vector bundle over *M*. A **multiderivation** of degree *n* on *E* is a pair (D, σ_D) , where $D \in \text{Hom}(\Lambda^{n-1}\Gamma(E) \otimes \Gamma(E), \Gamma(E))$ and $\sigma_D \in \Gamma(\text{Hom}(\Lambda^{n-1}E, TM))$, such that for all $f \in C^{\infty}(M)$ and sections $X_i \in \Gamma(E)$, the following conditions are satisfied:

$$D(X_1, \dots, fX_i, \dots, X_{n-1}, X_n) = fD(X_1, \dots, X_i, \dots, X_{n-1}, X_n), \quad i = 1, \dots, n-1;$$

$$D(X_1, \dots, X_{n-1}, fX_n) = fD(X_1, \dots, X_{n-1}, X_n) + \sigma_D(X_1, \dots, X_{n-1})(f)X_n.$$

We will denote by $\text{Der}^{n}(E)$ the space of multiderivations of degree $n, n \ge 1$.

Let $(A, *_A, a_A)$ be a pre-Lie algebroid. Recall that the deformation complex of A is a cochain complex $(C^*_{def}(A, A) = \bigoplus_{n \ge 0} \text{Der}^n(A), d_{def})$, where for all $X_i \in \Gamma(A), i = 1, 2 \cdots, n + 1$, the coboundary operator d_{def} : $\text{Der}^n(A) \longrightarrow \text{Der}^{n+1}(A)$ is given by

$$d_{def}\omega(X_{1},\cdots,X_{n+1}) = \sum_{i=1}^{n} (-1)^{i+1} X_{i} *_{A} \omega(X_{1},\cdots,\hat{X}_{i},\cdots,X_{n+1}) + \sum_{i=1}^{n} (-1)^{i+1} \omega(X_{1},\cdots,\hat{X}_{i},\cdots,X_{n},X_{i}) *_{A} X_{n+1} - \sum_{i=1}^{n} (-1)^{i+1} \omega(X_{1},\cdots,\hat{X}_{i},\cdots,X_{n},X_{i} *_{A} X_{n+1}) + \sum_{1 \le i < j \le n} (-1)^{i+j} \omega([X_{i},X_{j}]_{A},X_{1},\cdots,\hat{X}_{i},\cdots,\hat{X}_{j},\cdots,X_{n+1}),$$

in which $\sigma_{d_{def}\omega}$ is given by

$$\sigma_{d_{def}\omega}(X_1, \cdots, X_n) = \sum_{i=1}^n (-1)^{i+1} [a_A(X_i), \sigma_\omega(X_1, \cdots, \hat{X}_i, \cdots, X_n)]_{\mathfrak{X}(M)} + \sum_{1 \le i < j \le n} (-1)^{i+j} \sigma_\omega([X_i, X_j]_A, X_1, \cdots, \hat{X}_i, \cdots, \hat{X}_j, \cdots, X_n)$$

(11)
$$+ \sum_{i=1}^{n} (-1)^{i+1} a_A(\omega(X_1, \cdots, \hat{X}_i, \cdots, X_n, X_i)).$$

The corresponding cohomology, which we denote by $\mathcal{H}^{\bullet}_{def}(A, A)$, is called the **deformation** cohomology of the pre-Lie algebroid.

Since any commutative pre-Lie algebra is a commutative associative algebra, we have the following conclusion obviously.

Lemma 3.6. Any commutative pre-Lie algebroid is a commutative associative algebroid.

Note that in a commutative pre-Lie algebroid, the anchor must be zero.

Definition 3.7. Let (A, \cdot_A) be a commutative associative algebroid. A **pre-Lie formal deforma**tion of A is a sequence of pairs $(\mu_k, \sigma_{\mu_k}) \in \text{Der}^2(A)$ with μ_0 being the commutative associative algebroid multiplication \cdot_A on $\Gamma(A)$ and $\sigma_{\mu_0} = 0$ such that the $\mathbb{R}[[\hbar]]$ -bilinear product \cdot_{\hbar} on $\Gamma(A)[[\hbar]]$ and $\mathbb{R}[[\hbar]]$ -linear map $\mathfrak{a}_{\hbar} : A \otimes \mathbb{R}[[\hbar]] \to TM \otimes \mathbb{R}[[\hbar]]$ determined by

(12)
$$X \cdot_{\hbar} Y = \sum_{k=0}^{\infty} \hbar^{k} \mu_{k}(X, Y),$$

(13)
$$\mathfrak{a}_{\hbar}(X) = \sum_{k=0}^{\infty} \hbar^{k} \sigma_{\mu_{k}}(X), \quad \forall X, Y \in \Gamma(A)$$

is a pre-Lie algebroid.

It is straightforward to check that the rule of pre-Lie algebra product \cdot_{\hbar} on $\Gamma(A)[[\hbar]]$ is equivalent to

(14)
$$\sum_{i+j=k} \left(\mu_i(\mu_j(X,Y),Z) - \mu_i(X,\mu_j(Y,Z)) \right) = \sum_{i+j=k} \left(\mu_i(\mu_j(Y,X),Z) - \mu_i(Y,\mu_j(X,Z)) \right)$$

for $k \ge 0$.

Theorem 3.8. Let (A, \cdot_A) be a commutative associative algebroid and $(A \otimes \mathbb{R}[[\hbar]], \cdot_{\hbar}, \mathfrak{a}_{\hbar})$ a pre-Lie formal deformation of A. Define a bracket $[-, -]_A : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ by

$$[X, Y]_A = \mu_1(X, Y) - \mu_1(Y, X), \quad \forall X, Y \in \Gamma(A).$$

Then $(A, [-, -]_A, \cdot_A, \sigma_{\mu_1})$ is an *F*-algebroid. The *F*-algebroid $(A, [-, -]_A, \cdot_A, \sigma_{\mu_1})$ is called the semi-classical limit of $(A \otimes \mathbb{R}[[\hbar]], \cdot_\hbar, \mathfrak{a}_\hbar)$. The pre-Lie algebroid $(A \otimes \mathbb{R}[[\hbar]], \cdot_\hbar, \mathfrak{a}_\hbar)$ is called a **pre-Lie deformation quantization** of (A, \cdot_A) .

Proof. Define the bracket $[-, -]_{\hbar}$ on $\Gamma(A)[[\hbar]]$ by

$$[X,Y]_{\hbar} = X \cdot_{\hbar} Y - Y \cdot_{\hbar} X = \hbar[X,Y]_A + \hbar^2(\mu_2(X,Y) - \mu_2(Y,X)) + \cdots, \quad \forall X,Y \in \Gamma(A).$$

By the fact that $(A \otimes \mathbb{R}[[\hbar]], \cdot_{\hbar}, \mathfrak{a}_{\hbar})$ is a pre-Lie algebroid, $(A[[\hbar]], [-, -]_{\hbar}, \mathfrak{a}_{\hbar})$ is a Lie algebroid. The \hbar^2 -terms of the Jacobi identity for $[-, -]_{\hbar}$ gives the Jacobi identity for $[-, -]_A$ and \hbar -terms of $[X, fY]_{\hbar} = f[X, Y]_{\hbar} + \mathfrak{a}_{\hbar}(X)(f)Y$ gives

$$[X, fY]_A = f[X, Y]_A + \sigma_{\mu_1}(X)(f)Y.$$

Thus $(A, [-, -]_A, \sigma_{\mu_1})$ is a Lie algebroid.

For k = 1 in (14), by the commutativity of μ_0 , we have

$$\mu_0(\mu_1(X, Y), Z) - \mu_0(X, \mu_1(Y, Z)) - \mu_1(X, \mu_0(Y, Z))$$

= $\mu_0(\mu_1(Y, X), Z) - \mu_0(Y, \mu_1(X, Z)) - \mu_1(Y, \mu_0(X, Z)).$

By a similar proof given by Hertling in [12], we can show that the Hertling-Manin relation holds with $X \cdot_A Y = \mu_0(X, Y)$ and $[X, Y]_A = \mu_1(X, Y) - \mu_1(Y, X)$ for $X, Y \in \Gamma(A)$. Thus $(A, [-, -]_A, \cdot_A, \sigma_{\mu_1})$ is an *F*-algebroid.

In the sequel, we study pre-Lie *n*-deformations and pre-Lie infinitesimal deformations of commutative associative algebroids.

Definition 3.9. Let (A, \cdot_A) be a commutative associative algebroid. A **pre-Lie** *n*-**deformation** of A is a sequence of pairs $(\mu_k, \sigma_{\mu_k}) \in \text{Der}^2(A)$ for $0 \le k \le n$ with μ_0 being the commutative associative algebroid multiplication \cdot_A on $\Gamma(A)$ and $\sigma_{\mu_0} = 0$, such that the $\mathbb{R}[[\hbar]]/(\hbar^{n+1})$ -bilinear product \cdot_\hbar on $\Gamma(A)[[\hbar]]/(\hbar^{n+1})$ and $\mathbb{R}[[\hbar]]/(\hbar^{n+1})$ -linear map $\mathfrak{a}_\hbar : A \otimes \mathbb{R}[[\hbar]] \to TM \otimes \mathbb{R}[[\hbar]]$ determined by

(15)
$$X \cdot_{\hbar} Y = \sum_{k=0}^{n} \hbar^{k} \mu_{k}(X, Y),$$

(16)
$$\mathfrak{a}_{\hbar}(X) = \sum_{k=0}^{n} \hbar^{k} \sigma_{\mu_{k}}(X), \quad \forall X, Y \in \Gamma(A)$$

is a pre-Lie algebroid.

We call a pre-Lie 1-deformation of a commutative associative algebroid (A, \cdot_A) a **pre-Lie** infinitesimal deformation and denote it by $(A, \mu_1, a_A = \sigma_{\mu_1})$.

By direct calculations, $(A, \mu_1, \sigma_{\mu_1})$ is a pre-Lie infinitesimal deformation of a commutative associative algebroid (A, \cdot_A) if and only if for all $X, Y, Z \in \Gamma(A)$

(17)
$$\mu_1(X, Y) \cdot_A Z - X \cdot_A \mu_1(Y, Z) - \mu_1(X, Y \cdot_A Z) = \mu_1(Y, X) \cdot_A Z - Y \cdot_A \mu_1(X, Z) - \mu_1(Y, X \cdot_A Z).$$

Equation (17) means that μ_1 is a 2-cocycle, i.e. $d_{def}\mu_1 = 0.$

Two pre-Lie infinitesimal deformations $A_{\hbar} = (A, \mu_1, \sigma_{\mu_1})$ and $A'_{\hbar} = (A, \mu'_1, \sigma_{\mu'_1})$ of a commutative associative algebroid (A, \cdot_A) are said to be **equivalent** if there exist a family of pre-Lie algebroid homomorphisms Id $+\hbar\varphi : A_{\hbar} \longrightarrow A'_{\hbar}$ modulo \hbar^2 for $\varphi \in \text{Der}^1(A)$. A pre-Lie infinitesimal deformation is said to be **trivial** if there exist a family of pre-Lie algebroid homomorphisms Id $+\hbar\varphi : A_{\hbar} \longrightarrow (A, \cdot_A, a_A = 0)$ modulo \hbar^2 .

By direct calculations, A_{\hbar} and A'_{\hbar} are equivalent pre-Lie infinitesimal deformations if and only if

(18)
$$\sigma_{\mu_1} = \sigma_{\mu'_1},$$

(19)
$$\mu_1(X,Y) - \mu'_1(X,Y) = X \cdot_A \varphi(Y) + \varphi(X) \cdot_A Y - \varphi(X \cdot_A Y).$$

Equation (19) means that $\mu_1 - \mu'_1 = d_{def}\varphi$ and (18) can be obtained by (19). Thus we have

Theorem 3.10. Let (A, \cdot_A) be a commutative associative algebroid. There is a one-to-one correspondence between the space of equivalence classes of pre-Lie infinitesimal deformations of A and the second cohomology group $\mathcal{H}^2_{def}(A, A)$.

It is routine to check that

Proposition 3.11. Let (A, \cdot_A) be a commutative associative algebroid such that

$$\mathcal{H}^2_{\rm def}(A,A)=0.$$

Then all pre-Lie infinitesimal deformations of A are trivial.

Definition 3.12. Let $\{(\mu_1, \sigma_{\mu_1}), \dots, (\mu_n, \sigma_{\mu_n})\}$ a pre-Lie n-deformation of a commutative associative algebroid (A, \cdot_A) . If there exists $(\mu_{n+1}, \sigma_{\mu_{n+1}}) \in \text{Der}^2(A)$ such that

$$\{(\mu_1, \sigma_{\mu_1}), \cdots, (\mu_n, \sigma_{\mu_n}), (\mu_{n+1}, \sigma_{\mu_{n+1}})\}$$

is a pre-Lie (n + 1)-deformation of (A, \cdot_A) , then $\{(\mu_1, \sigma_{\mu_1}), \dots, (\mu_n, \sigma_{\mu_n}), (\mu_{n+1}, \sigma_{\mu_{n+1}})\}$ is called an **extension** of the pre-Lie n-deformation $\{(\mu_1, \sigma_{\mu_1}), \dots, (\mu_n, \sigma_{\mu_n})\}$.

Theorem 3.13. For any pre-Lie n-deformation of a commutative associative algebroid (A, \cdot_A) , the pair $(\Theta_n, \sigma_{\Theta_n}) \in \text{Der}^3(A)$ defined by

$$(20) \ \Theta_n(X, Y, Z) = \sum_{\substack{i+j=n+1, \\ i,j\geq 1}} (\mu_i(\mu_j(X, Y), Z) - \mu_i(X, \mu_j(Y, Z)) - \mu_i(\mu_j(Y, X), Z) + \mu_i(Y, \mu_j(X, Z)))),$$

$$(21) \ \sigma_{\Theta_n}(X, Y) = \sum_{\substack{i+j=n+1, \\ i,j\geq 1}} (\sigma_{\mu_i}(\mu_j(X, Y) - \mu_j(Y, X)) - [\sigma_{\mu_i}(X), \sigma_{\mu_j}(Y)]_{\mathfrak{X}(M)})$$

is a cocycle, i.e. $d_{def}\Theta_n = 0$.

Moreover, the pre-Lie n-deformation $\{(\mu_1, \sigma_{\mu_1}), \cdots, (\mu_n, \sigma_{\mu_n})\}$ extends to some pre-Lie (n+1)-deformation if and only if $[\Theta_n] = 0$ in $\mathcal{H}^3_{def}(A, A)$.

Proof. It is obvious that $\Theta_n(X, Y, Z) = -\Theta_n(Y, Z, X)$ for all $X, Y, Z \in \Gamma(A)$. It is straightforward to check that

$$\begin{split} \Theta_n(X, fY, Z) &= f\Theta_n(X, Y, Z), \\ \Theta_n(X, Y, fZ) &= f\Theta_n(X, Y, Z) + \sigma_{\Theta_n}(X, Y)(f)Z. \end{split}$$

Thus Θ_n is an element of $\text{Der}^3(A)$. By a direct calculation, the cochain $\Theta_n \in \text{Der}^3(A)$ is closed.

Assume that the pre-Lie (n+1)-deformation $\{(\mu_1, \sigma_{\mu_1}), \cdots, (\mu_{n+1}, \sigma_{\mu_{n+1}})\}$ of a commutative associative algebroid (A, \cdot_A) is an extension of the pre-Lie *n*-deformation $\{(\mu_1, \sigma_{\mu_1}), \cdots, (\mu_n, \sigma_{\mu_n})\}$, then we have

$$\sum_{i+j=n+1,i,j\geq 1} \left(\mu_i(\mu_j(X,Y),Z) - \mu_i(X,\mu_j(Y,Z)) - \mu_i(\mu_j(Y,X),Z) + \mu_i(Y,\mu_j(X,Z)) \right)$$

= $X \cdot_A \mu_{n+1}(Y,Z) - Y \cdot_A \mu_{n+1}(X,Z) + \mu_{n+1}(Y,X) \cdot_A Z - \mu_{n+1}(X,Y) \cdot_A Z$
+ $\mu_{n+1}(Y,X) \cdot_A Z - \mu_{n+1}(X,Y) \cdot_A Z.$

Note that the left-hand side of the above equality is just $\Theta_n(X, Y, Z)$. We can rewrite the above equality as

$$\Theta_n(X, Y, Z) = d_{def}\mu_{n+1}(X, Y, Z)$$

We conclude that, if a pre-Lie *n*-deformation of a commutative associative algebroid (A, \cdot_A) extends to a pre-Lie (n + 1)-deformation, then Θ_n is a coboundary.

Conversely, if Θ_n is a coboundary, then there exists an element $(\psi, \sigma_{\psi}) \in \text{Der}^2(A)$ such that

$$\Theta_n(X, Y, Z) = d_{def}\psi(X, Y, Z).$$

It is not hard to check that $\{(\mu_1, \sigma_{\mu_1}), \cdots, (\mu_{n+1}, \sigma_{\mu_{n+1}})\}$ with $\mu_{n+1} = \psi$ is a pre-Lie (n + 1)-deformation of (A, \cdot_A) and thus this pre-Lie (n + 1)-deformation is an extension of the pre-Lie *n*-deformation $\{(\mu_1, \sigma_{\mu_1}), \cdots, (\mu_n, \sigma_{\mu_n})\}$.

4. Some constructions of F-algebroids

In this section, we use eventual identities and Nijenhuis operators to construct *F*-algebroids. In particular, a pseudo-eventual identity naturally gives a Nijenhuis operator on an *F*-algebroid.

4.1. (Pseudo-)Eventual identities and Dubrovin's dual of *F*-algebroids.

Definition 4.1. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid with an identity *e*. A section $\mathcal{E} \in \Gamma(A)$ is called a **pseudo-eventual identity** on the *F*-algebroid if the following equality holds:

(22)
$$P_{\mathcal{E}}(X,Y) = [e,\mathcal{E}]_A \cdot_A X \cdot_A Y, \quad \forall X,Y \in \Gamma(A).$$

A pseudo-eventual identity \mathcal{E} on the *F*-algebroid *A* is called an **eventual identity** if it is invertible, i.e. there is a section $\mathcal{E}^{-1} \in \Gamma(A)$ such that $\mathcal{E}^{-1} \cdot_A \mathcal{E} = \mathcal{E} \cdot_A \mathcal{E}^{-1} = e$.

Denote the set of all pseudo-eventual identities on an F-algebroid A by E(A), i.e.

$$E(A) = \{ \mathcal{E} \in \Gamma(A) \mid P_{\mathcal{E}}(X, Y) = [e, \mathcal{E}]_A \cdot_A X \cdot_A Y, \quad \forall X, Y \in \Gamma(A) \}.$$

Proposition 4.2. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid with an identity *e*. Then E(A) is an *F*-manifold subalgebra of $\Gamma(A)$. Moreover, if $\mathcal{E} \in \Gamma(A)$ is an eventual identity on the *F*-algebroid *A*, then \mathcal{E}^{-1} is also an eventual identity on *A*.

Proof. For the first claim, by a straightforward calculation, E(A) is a subspace of the vector space $\Gamma(A)$.

Then we show that the multiplication \cdot_A of two pseudo-eventual identities is still a pseudo-eventual identity. In fact, for any two pseudo-eventual identities \mathcal{E}_1 and \mathcal{E}_2 , by (1),

$$P_{\mathcal{E}_{1}\cdot_{A}\mathcal{E}_{2}}(X,Y) = \mathcal{E}_{1}\cdot_{A}P_{\mathcal{E}_{2}}(X,Y) + \mathcal{E}_{2}\cdot_{A}P_{\mathcal{E}_{1}}(X,Y)$$

$$= (\mathcal{E}_{1}\cdot_{A}[e,\mathcal{E}_{2}]_{A} + \mathcal{E}_{2}\cdot_{A}[e,\mathcal{E}_{1}]_{A})\cdot_{A}X\cdot_{A}Y$$

$$= [e,\mathcal{E}_{1}\cdot_{A}\mathcal{E}_{2}]_{A}\cdot_{A}X\cdot_{A}Y,$$

where in the last equality we used $P_e(\mathcal{E}_1, \mathcal{E}_2) = 0$. Thus $\mathcal{E}_1 \cdot_A \mathcal{E}_2$ is a pseudo-eventual identity.

Finally, we show that the Lie bracket of two pseudo-eventual identities is also a pseudoeventual identity. By (1), we can show that for any $X, Y, Z, W \in \Gamma(A)$, we have

$$\begin{aligned} P_{[X,Y]_A}(Z,W) &= [X,P_Y(Z,W)]_A - P_Y([X,Z]_A,W) - P_Y(Z,[X,W]_A) \\ &- [Y,P_X(Z,W)]_A + P_X([Y,Z]_A,W) + P_X(Z,[Y,W]_A). \end{aligned}$$

Let \mathcal{E}_1 and \mathcal{E}_2 be eventual identities. Taking $X = \mathcal{E}_1$ and $Y = \mathcal{E}_2$, then by (22),

$$P_{[\mathcal{E}_{2},\mathcal{E}_{2}]_{A}}(Z,W) = [\mathcal{E}_{1}, [e,\mathcal{E}_{2}]_{A} \cdot_{A} Z \cdot_{A} W]_{A} - [e,\mathcal{E}_{2}]_{A} \cdot_{A} [\mathcal{E}_{1},Z]_{A} \cdot_{A} W$$

-[e,\mathcal{E}_{2}]_{A} \cdot_{A} Z \cdot_{A} [\mathcal{E}_{1},W]_{A} - [\mathcal{E}_{2}, [e,\mathcal{E}_{1}]_{A} \cdot_{A} Z \cdot_{A} W]_{A}
+[e,\mathcal{E}_{1}]_{A} \cdot_{A} [\mathcal{E}_{2},Z]_{A} \cdot_{A} W + [e,\mathcal{E}_{1}]_{A} \cdot_{A} Z \cdot_{A} [\mathcal{E}_{2},W]_{A}

On the other hand, by (22), we have

$$[\mathcal{E}_1, [e, \mathcal{E}_2]_A \cdot_A Z \cdot_A W]_A = 2[e, \mathcal{E}_1]_A \cdot [e, \mathcal{E}_2]_A \cdot_A Z \cdot_A W + [\mathcal{E}_1, [e, \mathcal{E}_2]_A]_A \cdot_A Z \cdot_A W + [e, \mathcal{E}_2]_A \cdot_A [\mathcal{E}_1, Z]_A \cdot_A W + [e, \mathcal{E}_2]_A \cdot_A Z \cdot_A [\mathcal{E}_1, W]_A;$$

$$[\mathcal{E}_2, [e, \mathcal{E}_1]_A \cdot_A Z \cdot_A W]_A = 2[e, \mathcal{E}_2]_A \cdot [e, \mathcal{E}_1]_A \cdot_A Z \cdot_A W + [\mathcal{E}_2, [e, \mathcal{E}_1]_A]_A \cdot_A Z \cdot_A W + [e, \mathcal{E}_1]_A \cdot_A [\mathcal{E}_2, Z]_A \cdot_A W + [e, \mathcal{E}_1]_A \cdot_A Z \cdot_A [\mathcal{E}_2, W]_A.$$

Thus

$$P_{[\mathcal{E}_2,\mathcal{E}_2]_A}(Z,W) = [\mathcal{E}_1, [e, \mathcal{E}_2]_A]_A \cdot_A Z \cdot_A W - [\mathcal{E}_2, [e, \mathcal{E}_1]_A]_A \cdot_A Z \cdot_A W$$
$$= [e, [\mathcal{E}_1, \mathcal{E}_2]_A]_A \cdot_A Z \cdot_A W,$$

which implies that $[\mathcal{E}_1, \mathcal{E}_2]_A$ is a pseudo-eventual identity. Therefore, E(A) is an *F*-manifold subalgebra of $\Gamma(A)$.

Assume that \mathcal{E} is an eventual identity on the *F*-algebroid *A*. By Proposition 2.12, we have $P_e(X, Y) = 0$. Applying the Hertling-Manin relation with $X = \mathcal{E}$ and $Y = \mathcal{E}^{-1}$, we obtain

$$0 = P_{\mathcal{E}_A \mathcal{E}^{-1}}(X, Y) = \mathcal{E} \cdot_A P_{\mathcal{E}^{-1}}(X, Y) + \mathcal{E}^{-1} \cdot_A P_{\mathcal{E}}(X, Y).$$

Combining this relation with (22), we have

$$P_{\mathcal{E}^{-1}}(X,Y) = -\mathcal{E}^{-2} \cdot_A [e,\mathcal{E}]_A \cdot_A X \cdot_A Y$$

On the other hand, by $P_e(X, Y) = 0$, we have

$$0 = P_e(\mathcal{E}, \mathcal{E}^{-1}) = -[e, \mathcal{E}]_A \cdot_A \mathcal{E}^{-1} - \mathcal{E} \cdot_A [e, \mathcal{E}^{-1}]_A$$

and then

$$[e, \mathcal{E}]_A \cdot_A \mathcal{E}^{-2} = ([e, \mathcal{E}]_A \cdot_A \mathcal{E}^{-1}) \cdot_A \mathcal{E}^{-1} = (-\mathcal{E} \cdot_A [e, \mathcal{E}^{-1}]_A) \cdot_A \mathcal{E}^{-1} = -[e, \mathcal{E}^{-1}]_A.$$

Thus we have

$$P_{\mathcal{E}^{-1}}(X,Y) = [e,\mathcal{E}^{-1}]_A \cdot_A X \cdot_A Y$$

which implies that \mathcal{E}^{-1} is also an eventual identity on *A*.

A pseudo-eventual identity on an F-algebroid gives a new F-algebroid.

Theorem 4.3. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid with an identity *e*. Then \mathcal{E} is a pseudoeventual identity on *A* if and only if $(A, [-, -]_A, \cdot_{\mathcal{E}}, a_A)$ is an *F*-algebroid, where $\cdot_{\mathcal{E}} : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ is defined by

(23)
$$X \cdot_{\mathcal{E}} Y = X \cdot_{A} Y \cdot_{A} \mathcal{E}, \quad \forall X, Y \in \Gamma(A).$$

Proof. The proof of this theorem is similar to the proof of Theorem 3 in [9]. We give a sketchy proof here for completeness. Assume that \mathcal{E} is a pseudo-eventual identity on A. It is straightforward to check that the multiplication $\cdot_{\mathcal{E}}$ defined by (23) is $C^{\infty}(M)$ -bilinear, commutative and associative.

For $X, Y, Z \in \Gamma(A)$, we set

$$P_X^{\mathcal{E}}(Y,Z) := [X, Y \cdot_{\mathcal{E}} Z]_A - [X, Y]_A \cdot_{\mathcal{E}} Z - Y \cdot_{\mathcal{E}} [X, Z]_A.$$

By a direct calculation, we have

(24)
$$P_X^{\mathcal{E}}(Y,Z) = P_X(\mathcal{E} \cdot_A Y,Z) + P_X(\mathcal{E},Y) \cdot_A Z + [Z,\mathcal{E}]_A \cdot_A X \cdot_A Y.$$

Using (24) and the Hertling-Manin relation, we have

$$P_{X \cdot \varepsilon Y}^{\mathcal{E}}(Z, W) = \mathcal{E} \cdot_A X \cdot_A P_Y(\mathcal{E} \cdot_A Z, W) + \mathcal{E} \cdot_A Y \cdot_A P_X(\mathcal{E} \cdot_A Z, W) + X \cdot_A Y \cdot_A P_{\mathcal{E}}(\mathcal{E} \cdot_A Z, W) + \mathcal{E} \cdot_A X \cdot_A W \cdot_A P_Y(\mathcal{E}, Z) + \mathcal{E} \cdot_A Y \cdot_A W \cdot_A P_X(\mathcal{E}, Z) + X \cdot_A Y \cdot_A W \cdot_A P_{\mathcal{E}}(\mathcal{E}, Z)$$

$$+[X \cdot_A Y \cdot_A \mathcal{E}, \mathcal{E}]_A \cdot_A Z \cdot_A W$$

Since \mathcal{E} is a pseudo-eventual identity on A, by (24), we have

$$P_{X_{\cdot \mathcal{E}}Y}^{\mathcal{E}}(Z, W) - X \cdot_{\mathcal{E}} P_{Y}^{\mathcal{E}}(Z, W) - Y \cdot_{\mathcal{E}} P_{X}^{\mathcal{E}}(Z, W)$$

$$= X \cdot_{A} Y \cdot_{A} \left(P_{\mathcal{E}}(\mathcal{E} \cdot_{A} Z, W) + W \cdot_{A} P_{\mathcal{E}}(\mathcal{E}, Z) \right)$$

$$-Z \cdot_{A} W \cdot_{A} \left([X \cdot_{A} Y \cdot_{A} \mathcal{E}, \mathcal{E}]_{A} + \mathcal{E} \cdot_{A} X \cdot_{A} [Y, \mathcal{E}]_{A} + \mathcal{E} \cdot_{A} Y \cdot_{A} [X, \mathcal{E}]_{A} \right)$$

$$= X \cdot_{A} Y \cdot_{A} \left(P_{\mathcal{E}}(\mathcal{E} \cdot_{A} Z, W) + W \cdot_{A} P_{\mathcal{E}}(\mathcal{E}, Z) \right) - Z \cdot_{A} W \cdot_{A} \left(P_{\mathcal{E}}(\mathcal{E}, X) \cdot_{A} Y + P_{\mathcal{E}}(\mathcal{E} \cdot_{A} X, Y) \right).$$

$$= X \cdot_{A} Y \cdot_{A} \left([e, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} Z \cdot_{A} W + [e, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} Z \cdot_{A} W \right)$$

$$-Z \cdot_{A} W \cdot_{A} \left([e, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} X \cdot_{A} Y + [e, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} X \cdot_{A} Y \right)$$

$$= 2[e, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} X \cdot_{A} Y \cdot_{A} Z \cdot_{A} W - 2[e, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} X \cdot_{A} Y \cdot_{A} Z \cdot_{A} W$$

$$= 0,$$

which implies that $(A, [-, -]_A, \cdot_{\mathcal{E}}, a_A)$ is an *F*-algebroid.

The converse can be proved similarly. We omit the details.

Theorem 4.4. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid with an identity *e*. Then \mathcal{E} is an eventual identity on *A* if and only if $(A, [-, -]_A, \cdot_{\mathcal{E}}, a_A)$ is also an *F*-algebroid with the identity \mathcal{E}^{-1} , which is called the Dubrovin's dual of $(A, [-, -]_A, \cdot_A, a_A)$, where $\cdot_{\mathcal{E}}$ is given by (23). Moreover, *e* is an eventual identity on the *F*-algebroid $(A, [-, -]_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A)$ and the map

(25)
$$(A, [-, -]_A, \cdot_A, e, a_A, \mathcal{E}) \longrightarrow (A, [-, -]_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A, e^{\dagger})$$

is an involution of the set of *F*-algebroids with eventual identities, where $e^{\dagger} := \mathcal{E}^{-2}$ is the inverse of *e* with respect to the multiplication $\cdot_{\mathcal{E}}$.

Proof. By Theorem 4.3, $(A, [-, -]_A, \cdot_{\mathcal{E}}, a_A)$ is an *F*-algebroid. It is obvious that \mathcal{E}^{-1} is the identity with respect to the multiplication $\cdot_{\mathcal{E}}$ defined by (23).

Next, we show that *e* is an eventual identity on $(A, [-, -]_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A)$. Since the identity with respective to the multiplication $\cdot_{\mathcal{E}}$ is \mathcal{E}^{-1} , we need to show that

$$[e, X \cdot_{\mathcal{E}} Y]_A - [e, X]_A \cdot_{\mathcal{E}} Y - X \cdot_{\mathcal{E}} [e, Y]_A = [\mathcal{E}^{-1}, e]_A \cdot_{\mathcal{E}} X \cdot_{\mathcal{E}} Y, \quad \forall X, Y \in \Gamma(A).$$

By a straightforward computation, for any $Z \in \Gamma(A)$, we have

(26)
$$[Z, X \cdot_{\mathcal{E}} Y]_A - [Z, X]_A \cdot_{\mathcal{E}} Y - X \cdot_{\mathcal{E}} [Z, Y]_A = P_Z(\mathcal{E} \cdot_A X, Y) + P_Z(\mathcal{E}, X) \cdot_A Y + [Z, \mathcal{E}]_A \cdot_A X \cdot_A Y.$$

Letting Z = e in (26) and using $P_e(X, Y) = 0$, we have

$$[e, X \cdot_{\mathcal{E}} Y]_A - [e, X]_A \cdot_{\mathcal{E}} Y - X \cdot_{\mathcal{E}} [e, Y]_A = [e, \mathcal{E}]_A \cdot_A X \cdot_A Y = ([e, \mathcal{E}]_A \cdot_A \mathcal{E}^{-2}) \cdot_{\mathcal{E}} X \cdot_{\mathcal{E}} Y.$$

Recall now from the proof of Proposition 4.2 that $[e, \mathcal{E}]_A \cdot_A \mathcal{E}^{-2} = [\mathcal{E}^{-1}, e]_A$. Thus *e* is an eventual identity on the *F*-algebroid $(A, [-, -]_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A)$.

Now we show that the map (25) is an involution. Note that $e^{\dagger} := \mathcal{E}^{-2}$ is the inverse of e with respect to the multiplication $\cdot_{\mathcal{E}}$. By Proposition 4.2, e^{\dagger} is also an eventual identity on the *F*-algebroid $(A, [-, -]_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A)$. Furthermore, for $X, Y \in \Gamma(A)$, we have

$$X \cdot_A Y = X \cdot_{\mathcal{E}} Y \cdot_{\mathcal{E}} \mathcal{E}^{-2} = X \cdot_{\mathcal{E}} Y \cdot_{\mathcal{E}} e^{\dagger},$$

which implies that the map defined by (25) is an involution of the set of *F*-algebroids with eventual identities. \Box

Definition 4.5. An *F*-manifold (M, \bullet, e) is called **semi-simple** if there exists canonical local coordinates (u^1, \dots, u^n) on *M* such that $e = \frac{\partial}{\partial u^1} + \dots + \frac{\partial}{\partial u^n}$ and

$$\frac{\partial}{\partial u^i} \bullet \frac{\partial}{\partial u^j} = \delta_{ij} \frac{\partial}{\partial u^j}, \quad i, j \in \{1, 2, \cdots, n\}$$

Example 4.6. Let (M, \bullet, e) be a semi-simple *F*-manifold. Then $(TM, [-, -]_{\mathfrak{X}(M)}, \bullet, \mathrm{Id})$ is an *F*-algebroid with an identity *e*. It is straightforward to check that any pseudo-eventual identity on $(TM, [-, -]_{\mathfrak{X}(M)}, \bullet, \mathrm{Id})$ is of the form

$$\mathcal{E} = f_1(u^1) \frac{\partial}{\partial u^1} + \dots + f_n(u^n) \frac{\partial}{\partial x_n},$$

where $f_i(u^i) \in C^{\infty}(M)$ depends only on u^i for $i = 1, 2, \dots, n$. Furthermore, it was shown in [9] that if all $f_i(u^i)$ are non-vanishing everywhere, then $\mathcal{E} \in \mathfrak{X}(M)$ is an eventual identity.

4.2. Nijenhuis operators and deformed *F*-algebroids. Recall from [6] that a Nijenhuis operator on a commutative associative algebra (A, \cdot_A) is a linear map $N : A \to A$ such that

(27)
$$N(x) \cdot_A N(y) = N(N(x) \cdot_A y + x \cdot_A N(y) - N(x \cdot_A y)), \quad \forall x, y \in A$$

Recall that a Nijenhuis operator on a Lie algebroid $(A, [-, -]_A, a_A)$ is a bundle map $N : A \to A$ such that

(28)
$$[N(X), N(Y)]_A = N([N(X), Y]_A + [X, N(Y)]_A - N([X, Y]_A)), \quad \forall X, Y \in \Gamma(A).$$

Based on these structures, we introduce the notion of a Nijenhuis operator on an *F*-algebroid as follows.

Definition 4.7. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid. A bundle map $N : A \to A$ is called a **Nijenhuis operator** on the *F*-algebroid A if N is both a Nijenhuis operator on the commutative associative algebra ($\Gamma(A), \cdot_A$) and a Nijenhuis operator on the Lie algebroid ($A, [-, -]_A, a_A$).

Define the deformed operation $\cdot_N : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ and the deformed bracket $[-, -]_N : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ by

(29) $X \cdot_N Y = N(X) \cdot_A Y + X \cdot_A N(Y) - N(X \cdot_A Y),$

(30)
$$[X, Y]_N = [N(X), Y]_A + [X, N(Y)]_A - N([X, Y]_A), \quad \forall X, Y \in \Gamma(A).$$

Theorem 4.8. Let $N : A \longrightarrow A$ be a Nijenhuis operator on an F-algebroid $(A, [-, -]_A, \cdot_A, a_A)$. Then $(A, [-, -]_N, \cdot_N, a_N = a_A \circ N)$ is an F-algebroid and N is an F-algebroid homomorphism from the F-algebroid $(A, [-, -]_N, \cdot_N, a_N = a_A \circ N)$ to $(A, [-, -]_A, \cdot_A, a_A)$.

Proof. Since *N* is a Nijenhuis operator on the commutative associative algebra ($\Gamma(A)$, \cdot_A), it follows that ($\Gamma(A)$, \cdot_N) is a commutative associative algebra ([6]). Since *N* is a Nijenhuis operator on the Lie algebroid (A, $[-, -]_A$, a_A), (A, $[-, -]_N$, a_N) is a Lie algebroid ([16]).

Define

(31)
$$\Phi_N(X, Y, Z, W) := P^N_{X \cup Y}(Z, W) - X \cdot_N P^N_Y(Z, W) - Y \cdot_N P^N_X(Z, W),$$

where $X, Y, Z, W \in \Gamma(A)$ and

$$P_X^N(Y,Z) := [X, Y \cdot_N Z]_N - [X, Y]_N \cdot_N Z - Y \cdot_N [X, Z]_N$$

Since A is an F-algebroid and N is a Nijenhuis operator on A, by a direct calculation, we have

$$\begin{split} \Phi_N(X,Y,Z,W) &= & \Phi(N(X),N(Y),N(Z),W) + \Phi(N(X),N(Y),Z,N(W)) \\ &+ \Phi(N(X),Y,N(Z),N(W)) + \Phi(X,N(Y),N(Z),N(W)) \\ &- N \Big(\Phi(N(X),N(Y),Z,W) + \Phi(N(X),Y,N(Z),W) + \Phi(N(X),Y,Z,N(W)) \Big) \\ &+ \Phi(X,N(Y),N(Z),W) + \Phi(X,Y,N(Z),N(W)) + \Phi(X,N(Y),Z,N(W)) \Big) \\ &+ N^2 \Big(\Phi(N(X),Y,Z,W) + \Phi(X,N(Y),Z,W) \\ &+ \Phi(X,Y,N(Z),W) + \Phi(X,Y,Z,N(W)) \Big) \\ &- N^3 (\Phi(X,Y,Z,W)) \\ &= & 0, \end{split}$$

which implies that

- /-- -- ---

$$P_{X \to V}^{N}(W, Z) - X \cdot_{N} P_{Y}^{N}(W, Z) - Y \cdot_{N} P_{X}^{N}(W, Z) = 0.$$

Thus $(A, [-, -]_N, \cdot_N, a_N = a_A \circ N)$ is an *F*-algebroid. It is obvious that N is an *F*-algebroid homomorphism from the *F*-algebroid $(A, [-, -]_N, \cdot_N, a_N = a_A \circ N)$ to $(A, [-, -]_A, \cdot_A, a_A)$.

Lemma 4.9. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid and *N* a Nijenhuis operator on *A*. For all $k, l \in \mathbb{N},$

- (i) $(A, [-, -]_{N^k}, \cdot_{N^k}, a_{N^k})$ is an *F*-algebroid;
- (ii) N^l is also a Nijenhuis operator on the *F*-algebroid $(A, [-, -]_{N^k}, \cdot_{N^k}, a_{N^k})$;
- (iii) The *F*-algebroids $(A, ([-, -]_{N^k})_{N^l}, (\cdot_{N^k})_{N^l}, a_{N^{k+l}})$ and $(A, [-, -]_{N^{k+l}}, \cdot_{N^{k+l}}, a_{N^{k+l}})$ are the same;
- (iv) N^l is an *F*-algebroid homomorphism between the *F*-algebroid $(A, [-, -]_{N^{k+l}}, \cdot_{N^{k+l}}, a_{N^{k+l}})$ and $(A, [-, -]_{N^k}, \cdot_{N^k}, a_{N^k})$.

Proof. Since the above conclusions with respect to Nijenhuis operators on commutative associative algebras ([6]) and Lie algebroids ([16]) simultaneously hold, by Theorem 4.8, the conclusions follow immediately.

At the end of this section, we show that a pseudo-eventual identity naturally gives a Nijenhuis operator on an F-algebroid.

Proposition 4.10. Let $(A, [-, -]_A, \cdot_A, a_A)$ be an *F*-algebroid with an identity *e* and \mathcal{E} a pseudoeventual identity on A. Then the endomorphism $N = \mathcal{E}_A$ is a Nijenhuis operator on the Falgebroid A. Consequently, $(A, [-, -]_{\mathcal{E}}, \cdot_{\mathcal{E}}, a_{\mathcal{E}})$ is an F-algebroid, where the bracket $[-, -]_{\mathcal{E}}$ is given by

(32) $[X, Y]_{\mathcal{E}} = [\mathcal{E} \cdot_A X, Y]_A + [X, \mathcal{E} \cdot_A Y]_A - \mathcal{E} \cdot_A [X, Y]_A, \quad \forall X, Y \in \Gamma(A),$

the multiplication $\cdot_{\mathcal{E}}$ is given by (23) and $a_{\mathcal{E}}(X) = a_{\mathcal{A}}(\mathcal{E} \cdot_{\mathcal{A}} X)$.

Proof. First, we show that $N = \mathcal{E}_A$ is a Nijenhuis operator on the associative algebra ($\Gamma(A), \cdot_A$). For any $X, Y \in \Gamma(A)$, we have

$$N(X) \cdot_{A} N(Y) - N(N(X) \cdot_{A} Y + X \cdot_{A} N(Y) - N(X \cdot_{A} Y))$$

= $X \cdot_{A} Y \cdot_{A} \mathcal{E}^{2} - \mathcal{E} \cdot_{A} (X \cdot_{A} Y \cdot_{A} \mathcal{E} + X \cdot_{A} Y \cdot_{A} \mathcal{E} - X \cdot_{A} Y \cdot_{A} \mathcal{E})$
= $X \cdot_{A} Y \cdot_{A} \mathcal{E}^{2} - X \cdot_{A} Y \cdot_{A} \mathcal{E}^{2}$

Thus $N = \mathcal{E}_A$ is a Nijenhuis operator on the associative algebra ($\Gamma(A), \cdot_A$).

Then we show that $N = \mathcal{E}_A$ is a Nijenhuis operator on the Lie algebroid $(A, [-, -]_A, a_A)$. It is obvious that N is a bundle map. Since \mathcal{E} is a pseudo-eventual identity on the *F*-algebroid A, taking $Y = \mathcal{E}$ in (22), we have

$$[X \cdot_A \mathcal{E}, \mathcal{E}]_A - [X, \mathcal{E}]_A \cdot_A \mathcal{E} = [\mathcal{E}, e]_A \cdot_A X \cdot_A \mathcal{E}.$$

For any $X, Y \in \Gamma(A)$, expanding $[\mathcal{E} \cdot_A X, \mathcal{E} \cdot_A Y]_A$ using the Hertling-Manin relation and by (33), we have

$$[N(X), N(Y)]_{A} - N([N(X), Y]_{A} + [X, N(Y)]_{A} - N([X, Y]_{A}))$$

$$= [\mathcal{E} \cdot_{A} X, \mathcal{E} \cdot_{A} Y]_{A} - \mathcal{E} \cdot_{A} ([\mathcal{E} \cdot_{A} X, Y]_{A} + [X, \mathcal{E} \cdot_{A} Y]_{A} - \mathcal{E} \cdot_{A} [X, Y]_{A})$$

$$= [\mathcal{E} \cdot_{A} X, \mathcal{E}]_{A} \cdot_{A} Y - [X, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} Y - [\mathcal{E} \cdot_{A} Y, \mathcal{E}]_{A} \cdot_{A} X + [Y, \mathcal{E}]_{A} \cdot_{A} \mathcal{E} \cdot_{A} X$$

$$= ([\mathcal{E} \cdot_{A} X, \mathcal{E}]_{A} - [X, \mathcal{E}]_{A} \cdot_{A} \mathcal{E}) \cdot_{A} Y - ([\mathcal{E} \cdot_{A} Y, \mathcal{E}]_{A} - [Y, \mathcal{E}]_{A} \cdot_{A} \mathcal{E}) \cdot_{A} X$$

$$= [\mathcal{E}, e]_{A} \cdot_{A} X \cdot_{A} \mathcal{E} \cdot_{A} Y - [\mathcal{E}, e]_{A} \cdot_{A} Y \cdot_{A} \mathcal{E} \cdot_{A} X$$

$$= 0.$$

Thus $N = \mathcal{E}_A$ is a Nijenhuis operator on the Lie algebroid $(A, [-, -]_A, a_A)$. Therefore, $N = \mathcal{E}_A$ is a Nijenhuis operator on the *F*-algebroid *A*.

The second claim follows from Theorem 4.8.

Corollary 4.11. Let (M, \bullet) be an *F*-manifold with an identity *e* and \mathcal{E} a pseudo-eventual identity on *M*. Then there is a new *F*-algebroid structure on *TM* given by

$$\begin{split} X \bullet_{\mathcal{E}} Y &= X \bullet Y \bullet \mathcal{E}, \\ [X, Y]_{\mathcal{E}} &= [\mathcal{E} \bullet X, Y]_{\mathfrak{X}(M)} + [X, \mathcal{E} \bullet Y]_{\mathfrak{X}(M)} - \mathcal{E} \bullet [X, Y]_{\mathfrak{X}(M)}, \\ a_{\mathcal{E}}(X) &= \mathcal{E} \bullet X, \quad \forall X, Y \in \mathfrak{X}(M). \end{split}$$

5. Pre-F-algebroids and eventual identities

In this section, first we introduce the notion of a pre-F-algebroid, and show that a pre-Falgebroid gives rise to an F-algebroid. Then we study eventual identities on a pre-F-algebroid, which give new pre-F-algebroids. Finally, we introduce the notion of a Nijenhuis operator on a pre-F-algebroid, and show that a Nijenhuis operator gives rise to a deformed pre-F-algebroid. Finally we give some applications to integrable systems.

5.1. Some Properties of pre-*F*-algebroids.

Definition 5.1. Let (g, \cdot) is a commutative associative algebra and (g, *) is a pre-Lie algebra. Define $\Psi : \otimes^3 g \longrightarrow g$ by

(34)
$$\Psi(x, y, z) := x * (y \cdot z) - (x * y) \cdot z - y \cdot (x * z).$$

(i) The triple $(g, *, \cdot)$ is called a **pre-***F***-manifold algebra** if

(35)
$$\Psi(x, y, z) = \Psi(y, x, z), \quad \forall x, y, z \in \mathfrak{g},$$

 (ii) The triple (g, *, ·) is called a pre-Lie commutative algebra (or PreLie-Com algebra) if

(36)
$$\Psi(x, y, z) = 0, \quad \forall x, y, z \in \mathfrak{g}.$$

It is obvious that a PreLie-Com algebra is a pre-*F*-manifold algebra.

Example 5.2. ([20]) Let (g, \cdot) be a commutative associative algebra with a derivation *D*. Then *the new product*

$$x * y = x \cdot D(y), \quad \forall x, y \in \mathfrak{g}$$

makes $(g, *, \cdot)$ being a PreLie-Com algebra. Furthermore, $(g, [-, -], \cdot)$ is an *F*-manifold algebra, where the bracket is given by

$$[x, y] = x * y - y * x = x \cdot D(y) - y \cdot D(x), \quad \forall x, y \in g.$$

Let $g = \mathbb{R}[u^1, x_2, \dots, x_n]$ be the algebra of polynomials in *n* variables. Denote by $\mathfrak{D}_n = \{\sum_{i=1}^n p_i \partial_{u^i} \mid p_i \in \mathfrak{g}\}$ the space of derivations.

Example 5.3. ([20]) Let g be the algebra of polynomials in *n* variables. Define $\cdot : \mathfrak{D}_n \times \mathfrak{D}_n \longrightarrow \mathfrak{D}_n$ and $* : \mathfrak{D}_n \times \mathfrak{D}_n \longrightarrow \mathfrak{D}_n$ by

$$\begin{array}{lll} (p\partial_{u^i}) \cdot (q\partial_{u^j}) &=& (pq)\delta_{ij}\partial_{u^i}, \\ (p\partial_{u^i}) * (q\partial_{u^j}) &=& p\partial_{u^i}(q)\partial_{u^j}, \quad \forall \ p,q \in \mathfrak{g}. \end{array}$$

Then $(\mathfrak{D}_n, *, \cdot)$ is a PreLie-Com algebra with the identity $e = \partial_{u^1} + \cdots + \partial_{x_n}$. Furthermore, it follows that $(\mathfrak{D}_n, [-, -], \cdot)$ is an *F*-manifold algebra with the identity *e*, where the bracket is given by

$$[p\partial_{u^i}, q\partial_{u^j}] = p\partial_{u^i}(q)\partial_{u^j} - q\partial_{u^j}(p)\partial_{u^i}, \quad \forall \ p, q \in \mathfrak{g}.$$

Definition 5.4. A pre-*F*-algebroid is a vector bundle *A* over *M* equipped with bilinear operations $\cdot_A : \Gamma(A) \times \Gamma(A) \to \Gamma(A), *_A : \Gamma(A) \times \Gamma(A) \to \Gamma(A)$, and a bundle map $a_A : A \to TM$, called the anchor, such that $(A, *_A, a_A)$ is a pre-Lie algebroid, (A, \cdot_A) is a commutative associative algebroid and $(\Gamma(A), [-, -]_A, \cdot_A)$ is a pre-*F*-manifold algebra. In particular, if $(\Gamma(A), *_A, \cdot_A)$ is a PreLie-Com algebra, we call this pre-*F*-algebroid a **PreLie-Com algebroid**.

We denote a pre-*F*-algebroid (or PreLie-Com algebroid) by $(A, *_A, \cdot_A, a_A)$.

Definition 5.5. Let $(A, *_A, \cdot_A, a_A)$ and $(B, *_B, \cdot_B, a_B)$ be pre-*F*-algebroids on *M*. A bundle map $\varphi : A \longrightarrow B$ is called a **homomorphism** of pre-*F*-algebroids, if the following conditions are satisfied:

$$\varphi(X \cdot_A Y) = \varphi(X) \cdot_B \varphi(Y), \quad \varphi(X *_A Y) = \varphi(X) *_B \varphi(Y), \quad a_B \circ \varphi = a_A, \quad \forall X, Y \in \Gamma(A).$$

Proposition 5.6. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid. Then $(A, [-, -]_A, \cdot_A, a_A)$ is an *F*-algebroid, and denoted by A^c , called the **sub-adjacent** *F*-algebroid of the pre-*F*-algebroid, where the bracket $[-, -]_A$ is given by

$$[X, Y]_A = X *_A Y - Y *_A X, \quad \forall X, Y \in \Gamma(A).$$

Proof. Since $(A, *_A, a_A)$ is a pre-Lie algebroid, $(A, [-, -]_A, a_A)$ is a Lie algebroid ([19]). Since $(\Gamma(A), *_A, \cdot_A)$ is a pre-*F*-manifold algebra, $(\Gamma(A), [-, -]_A, \cdot_A)$ is an *F*-manifold algebra ([10]). Thus $(A, [-, -]_A, \cdot_A, a_A)$ is an *F*-algebroid.

The notion of an *F*-manifold with a compatible flat connection was introduced by Manin in [26]. Recall that an *F*-manifold with a compatible flat connection (PreLie-Com manifold) is a triple (M, ∇, \bullet) , where *M* is a manifold, ∇ is a flat connection and \bullet is a $C^{\infty}(M)$ -bilinear, commutative, associative multiplication on the tangent bundle *TM* such that $(TM, \nabla, \bullet, Id)$ is a pre-*F*-algebroid (PreLie-Com algebroid). It is obvious that an *F*-manifold with a compatible flat connection is a special case of pre-*F*-algebroids. An *F*-manifold with a compatible flat connection (resp. PreLie-Com manifold) is called **semi-simple** if its sub-adjacent *F*-manifold is semi-simple.

Proposition 5.7. Let (M, ∇, \bullet, e) be a semi-simple PreLie-Com manifold with the canonical local coordinate systems (u^1, \dots, u^n) . Then we have

$$abla_{rac{\partial}{\partial u^i}}rac{\partial}{\partial u^j}=0, \quad i,j\in\{1,2,\cdots,n\}.$$

Proof. Set $\nabla_{\frac{\partial}{\partial u^j}} \frac{\partial}{\partial u^j} = \sum_k \Gamma_{ij}^k \frac{\partial}{\partial x_k}$. By (36), for any $i, j, k \in \{1, 2, \dots, n\}$, we have

(38)
$$0 = \nabla_{\frac{\partial}{\partial u^{i}}} (\frac{\partial}{\partial u^{j}} \bullet \frac{\partial}{\partial u^{k}}) - (\nabla_{\frac{\partial}{\partial u^{i}}} \frac{\partial}{\partial u^{j}}) \bullet \frac{\partial}{\partial u^{k}} - \frac{\partial}{\partial u^{j}} \bullet (\nabla_{\frac{\partial}{\partial u^{i}}} \frac{\partial}{\partial u^{k}})$$
$$= \sum_{l} \delta_{jk} \Gamma_{lk}^{l} \frac{\partial}{\partial x_{l}} - \Gamma_{lj}^{k} \frac{\partial}{\partial u^{k}} - \Gamma_{lk}^{j} \frac{\partial}{\partial u^{j}}.$$

For $j \neq k$ in (38), we have $\Gamma_{ij}^k = 0$ $(j \neq k)$. For j = k in (38), we have $\Gamma_{ij}^j = 0$. Thus for any $i, j, k \in \{1, 2, \dots, n\}$, we have $\Gamma_{ij}^k = 0$.

We give some useful formulas that will be frequently used in the sequel.

Lemma 5.8. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid. Then $\Psi(X, Y, Z)$ defined by (34) is a tensor field of type (3, 1) and symmetric in all arguments. Furthermore, Ψ satisfies

(39) $\Psi(X \cdot_A Y, Z, W) - \Psi(X, Z, W) \cdot_A Y = \Psi(X \cdot_A Z, Y, W) - \Psi(X, Y, W) \cdot_A Z,$

(40) $\Psi(X \cdot_A Y, Z, W) - \Psi(X \cdot_A Z, Y, W) = \Psi(W \cdot_A Y, X, Z) - \Psi(W \cdot_A Z, X, Y)$

for all $X, Y, Z, W \in \Gamma(A)$.

Proof. It is straightforward to check that $\Psi(X, Y, Z)$ is a tensor field of type (3, 1). The symmetry of $\Psi(X, Y, Z)$ in the first two arguments is the consequence of (35) and in the last two arguments is the consequence of the commutativity of \cdot_A .

By the symmetry of Ψ , we have

$$\begin{split} \Psi(X \cdot_A Y, Z, W) &= \Psi(Z, X \cdot_A Y, W) \\ &= Z *_A ((X \cdot_A Y) \cdot_A W) - (Z *_A (X \cdot_A Y)) \cdot_A W - (X \cdot_A Y) \cdot_A (Z *_A W) \\ &= \Psi(Z, X \cdot_A W, Y) + (Z *_A (X \cdot_A W)) \cdot_A Y + (X \cdot_A W) \cdot_A (Z * Y) \\ &- \Psi(Z, X, Y) \cdot_A W - (Z *_A X) \cdot_A Y \cdot W - X \cdot_A (Z *_A Y) \cdot_A W \\ &- (X \cdot_A Y) \cdot_A (Z *_A W) \\ &= (Z *_A (X \cdot_A W) - (Z *_A X) \cdot_A W - X \cdot_A (Z *_A W)) \cdot_A Y \\ &+ \Psi(Z, X \cdot_A W, Y) - \Psi(Z, X, Y) \cdot_A W \\ &= \Psi(X, Z, W) \cdot_A Y + \Psi(X \cdot_A W, Y, Z) - \Psi(X, Y, Z) \cdot_A W. \end{split}$$

Thus we have

(41)
$$\Psi(X \cdot_A Y, Z, W) - \Psi(X, Z, W) \cdot_A Y = \Psi(X \cdot_A W, Y, Z) - \Psi(X, Y, Z) \cdot_A W.$$

Interchanging Z and W in (41), we have

$$\Psi(X \cdot_A Y, W, Z) - \Psi(X, W, Z) \cdot_A Y = \Psi(X \cdot_A Z, Y, W) - \Psi(X, Y, W) \cdot_A Z$$

By the symmetry of Ψ , (39) follows.

By (39), we have

$$\begin{split} \Psi(X \cdot_A Y, Z, W) &- \Psi(X \cdot_A Z, Y, W) &= \Psi(X, Z, W) \cdot_A Y - \Psi(X, Y, W) \cdot_A Z, \\ \Psi(W \cdot_A Y, X, Z) &- \Psi(W \cdot_A Z, X, Y) &= \Psi(W, X, Z) \cdot_A Y - \Psi(W, X, Y) \cdot_A Z. \end{split}$$

By the symmetry of Ψ , we have

$$\Psi(X, Z, W) \cdot_A Y - \Psi(X, Y, W) \cdot_A Z = \Psi(W, X, Z) \cdot_A Y - \Psi(W, X, Y) \cdot_A Z.$$

Thus (40) holds.

Lemma 5.9. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid with an identity *e*. Then we have

(42)
$$\Psi(e, X, Y) = -(X *_A e) \cdot_A Y,$$

(43)
$$(X *_A e) \cdot_A Y = (Y *_A e) \cdot_A X, \quad \forall X, Y \in \Gamma(A).$$

Proof. (42) follows by a direct calculation. By the symmetry of Ψ and (42), (43) follows.

Lemma 5.10. Let $(A, *_A, \cdot_A, a_A)$ be a PreLie-Com algebroid with an identity e. Then we have (44) $X *_A e = 0, \quad \forall X \in \Gamma(A).$

Proof. The conclusion follows from the following relation

$$X *_A (e \cdot_A e) - (X *_A e) \cdot_A e - (X *_A e) \cdot_A e = 0. \quad \Box$$

Example 5.11. Let $\{u\}$ be a coordinate system of \mathbb{R} . Define an anchor map $a : T\mathbb{R} \longrightarrow T\mathbb{R}$, a multiplication $\cdot : \mathfrak{X}(\mathbb{R}) \times \mathfrak{X}(\mathbb{R}) \longrightarrow \mathfrak{X}(\mathbb{R})$ and a multiplication $* : \mathfrak{X}(\mathbb{R}) \times \mathfrak{X}(\mathbb{R}) \longrightarrow \mathfrak{X}(\mathbb{R})$ by

$$a(f\frac{\partial}{\partial u}) = uf\frac{\partial}{\partial u}, \quad f\frac{\partial}{\partial u} \cdot g\frac{\partial}{\partial u} = fg\frac{\partial}{\partial u}, \quad f\frac{\partial}{\partial u} * g\frac{\partial}{\partial u} = uf\frac{\partial g}{\partial u}\frac{\partial}{\partial u}$$

for all $f, g \in C^{\infty}(\mathbb{R})$. Then $(T\mathbb{R}, *, \cdot, a)$ is a PreLie-Com algebroid with the identity $\frac{\partial}{\partial u}$. Furthermore, $(T\mathbb{R}, [-, -], \cdot, a)$ is an *F*-algebroid with the identity $\frac{\partial}{\partial u}$, where [-, -] is given by

$$[f\frac{\partial}{\partial u},g\frac{\partial}{\partial u}] = u(f\frac{\partial g}{\partial u} - g\frac{\partial f}{\partial u})\frac{\partial}{\partial u}$$

Definition 5.12. Let $(g, *, \cdot)$ be a pre-*F*-manifold algebra (PreLie-Com algebra). An **action** of g on a manifold *M* is a linear map $\rho : g \longrightarrow \mathfrak{X}(M)$ from g to the space of vector fields on *M*, such that for all $x, y \in g$, we have

$$\rho(x \ast y - y \ast x) = [\rho(x), \rho(y)]_{\mathfrak{X}(M)}.$$

Given an action of a pre-*F*-manifold algebra (PreLie-Com algebra) g on *M*, let $A = M \times g$ be the trivial bundle. Define an anchor map $a_{\rho} : A \longrightarrow TM$, a multiplication $\cdot_{\rho} : \Gamma(A) \times \Gamma(A) \longrightarrow$ $\Gamma(A)$ and a bracket $*_{\rho} : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ by

(45)
$$a_{\rho}(m,u) = \rho(u)_m, \quad \forall \ m \in M, u \in g,$$

$$(46) X \cdot_{\rho} Y = X \cdot Y,$$

(47)
$$X *_{\rho} Y = \mathcal{L}_{\rho(X)} Y + X * Y, \quad \forall X, Y \in \Gamma(A),$$

where $X \cdot Y$ and X * Y are the pointwise $C^{\infty}(M)$ -bilinear multiplication and bracket, respectively.

Proposition 5.13. With the above notations, $(A = M \times g, *_{\rho}, \cdot_{\rho}, a_{\rho})$ is a pre-*F*-algebroid (PreLie-Com algebroid), which we call an **action pre-***F***-algebroid** (action PreLie-Com algebroid), where $*_{\rho}, \cdot_{\rho}$ and a_{ρ} are given by (47), (46) and (45), respectively.

Proof. It follows by a similar proof of Proposition 2.14.

It is obvious that the sub-adjacent F-algebroid of the action pre-F-algebroid is an action F-algebroid.

Example 5.14. Consider the PreLie-Com algebra $(\mathfrak{D}_n, \cdot, *)$ given by Example 5.3. Let (t_1, \dots, t_n) be the canonical coordinate systems on \mathbb{R}^n . Define a map $\rho : \mathfrak{D}_n \longrightarrow \mathfrak{X}(\mathbb{R}^n)$ by

$$\rho(p(u^1,\cdots,u^n)\partial_{u^i})=p(t_1,\cdots,t_n)\frac{\partial}{\partial t_i},\quad i\in\{1,2,\cdots,n\}.$$

It is straightforward to check that ρ is an action of the PreLie-Com algebra \mathfrak{D}_n on \mathbb{R}^n . Thus $(A = \mathbb{R}^n \times \mathfrak{D}_n, *_\rho, \cdot_\rho, a_\rho)$ is a PreLie-Com algebroid, where $*_\rho, \cdot_\rho$ and a_ρ are given by

$$\begin{aligned} a_{\rho}(m, p(u^{1}, u^{2}, \cdots, u^{n})\partial_{u^{i}}) &= p(m)\frac{\partial}{\partial t_{i}} \mid_{m}, \quad \forall \ m \in \mathbb{R}^{n}, \\ (f \otimes (p\partial_{u^{i}})) \cdot_{\rho} (g \otimes (q\partial_{u^{j}})) &= (fg) \otimes (pq\delta_{ij}\partial_{u^{i}}), \\ (f \otimes (p\partial_{u^{i}})) *_{\rho} (g \otimes (q\partial_{u^{j}})) &= fp\frac{\partial g}{\partial t_{i}} \otimes (q\partial_{u^{j}}) + (fg) \otimes p\partial_{u^{i}}(q)\partial_{u^{j}}. \end{aligned}$$

where $f, g \in C^{\infty}(\mathbb{R}^n), p, q \in \mathbb{R}[u^1, \cdots, u^n].$

5.2. Eventual identities of pre-*F*-algebroids.

Definition 5.15. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid with an identity *e*. A section $\mathcal{E} \in \Gamma(A)$ is called a **pseudo-eventual identity** on *A* if the following equalities hold:

(48)
$$\Psi(\mathcal{E}, X, Y) = -(\mathcal{E} *_A e) \cdot_A X \cdot_A Y,$$

(49)
$$(X *_A \mathcal{E}) \cdot_A Y = (Y *_A \mathcal{E}) \cdot_A X, \quad \forall X, Y \in \Gamma(A).$$

A pseudo-eventual identity \mathcal{E} on the pre-*F*-algebroid with an identity *e* is called an **eventual** identity if it is invertible.

Proposition 5.16. Let $(A, *_A, \cdot_A, e, a_A)$ be a pre-*F*-algebroid with an identity *e*. If $\mathcal{E} \in \Gamma(A)$ is a pseudo-eventual identity on *A*, then $\mathcal{E} \in \Gamma(A)$ is a pseudo-eventual identity on its sub-adjacent *F*-algebroid A^c .

Proof. By a direct calculation, for $X, Y \in \Gamma(A)$, we have

$$P_{\mathcal{E}}(X, Y) - [e, \mathcal{E}]_A \cdot_A X \cdot_A Y$$

$$= \mathcal{E} *_A (X \cdot_A Y) - (X \cdot_A Y) *_A \mathcal{E} - (\mathcal{E} *_A X) \cdot_A Y + (X *_A \mathcal{E}) \cdot_A Y$$

$$-(\mathcal{E} *_A Y) \cdot_A X + (Y *_A \mathcal{E}) \cdot_A X - (e *_A \mathcal{E}) \cdot_A X \cdot_A Y + (\mathcal{E} *_A e) \cdot_A X \cdot_A Y$$

$$= \Psi(\mathcal{E}, X, Y) + (\mathcal{E} *_A e) \cdot_A X \cdot_A Y - (X \cdot_A Y) *_A \mathcal{E} + (X *_A \mathcal{E}) \cdot_A Y$$

$$+(Y *_A \mathcal{E}) \cdot_A X - (e *_A \mathcal{E}) \cdot_A X \cdot_A Y.$$

By (48) and (49), we have

$$P_{\mathcal{E}}(X, Y) - [e, \mathcal{E}]_A \cdot_A X \cdot_A Y$$

$$= -(X \cdot_A Y) *_A \mathcal{E} + (X *_A \mathcal{E}) \cdot_A Y + (Y *_A \mathcal{E}) \cdot_A X - (e *_A \mathcal{E}) \cdot_A X \cdot_A Y$$

$$= -(e *_A \mathcal{E}) \cdot_A X \cdot_A Y + (X *_A \mathcal{E}) \cdot_A Y + (X *_A \mathcal{E}) \cdot_A Y - (e *_A \mathcal{E}) \cdot_A X \cdot_A Y$$

$$= 2(X *_A \mathcal{E}) \cdot_A Y - 2(e *_A \mathcal{E}) \cdot_A X \cdot_A Y$$

$$= 2(X *_A \mathcal{E}) \cdot_A Y - 2(X *_A \mathcal{E}) \cdot_A Y$$

$$= 0.$$

Thus $\mathcal{E} \in \Gamma(A)$ is a pseudo-eventual identity on its sub-adjacent *F*-algebroid A^c .

By Lemma 5.10, we have

Proposition 5.17. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid with an identity *e* and \mathcal{E} an invertible element in $\Gamma(A)$. If $(A, *_A, \cdot_A, a_A)$ is a PreLie-Com algebroid, then \mathcal{E} is an eventual identity on *A* if and only if (49) holds.

Lemma 5.18. Let $(A, *_A, \cdot_A, e, a_A)$ be a pre-*F*-algebroid. Then for $\mathcal{E} \in \Gamma(A)$, (48) holds if and only if

(50) $\Psi(X, \mathcal{E}_{A}, Y, Z) = \Psi(Y, \mathcal{E}_{A}, X, Z), \quad \forall X, Y, Z \in \Gamma(A).$

Proof. Assume that (50) holds. By (39), we have

(51) $\Psi(\mathcal{E}, X, Z) \cdot_A Y - \Psi(\mathcal{E}, Y, Z) \cdot_A X = \Psi(X, \mathcal{E} \cdot_A Y, Z) - \Psi(Y, \mathcal{E} \cdot_A X, Z) = 0.$

Taking Y = e in (51), we have

$$\Psi(\mathcal{E}, X, Z) = -(\mathcal{E} *_A e) \cdot_A X \cdot_A Z.$$

This implies that (48) holds.

Conversely, if (48) holds, then we have

 $\Psi(\mathcal{E}, X, Z) \cdot_A Y - \Psi(\mathcal{E}, Y, Z) \cdot_A X = -(\mathcal{E} *_A e) \cdot_A X \cdot_A Z \cdot_A Y + (\mathcal{E} *_A e) \cdot_A Y \cdot_A Z \cdot_A X = 0.$

By (39), we have

$$\Psi(X, \mathcal{E} \cdot_A Y, Z) = \Psi(Y, \mathcal{E} \cdot_A X, Z).$$

This implies that (50) holds.

Denote the set of all pseudo-eventual identities on a pre-*F*-algebroid $(A, *_A, \cdot_A, a_A)$ with an identity *e* by $\mathfrak{E}(A)$.

Proposition 5.19. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid with an identity *e*. Then for any $\mathcal{E}_1, \mathcal{E}_2 \in \mathfrak{E}(A), \mathcal{E}_1 \cdot_A \mathcal{E}_2 \in \mathfrak{E}(A)$. Furthermore, if \mathcal{E} is an eventual identity on *A*, then \mathcal{E}^{-1} is also an eventual identity on *A*.

Proof. For the first claim, let $\mathcal{E}_1, \mathcal{E}_2$ be two pseudo-eventual identities on the pre-*F*-algebroid *A*. For all $X, Y, Z \in \Gamma(A)$, since \mathcal{E}_1 and \mathcal{E}_2 are pseudo-eventual identities, by (50), we have

$$\Psi(X, \mathcal{E}_1 \cdot_A \mathcal{E}_2 \cdot_A Y, Z) = \Psi(\mathcal{E}_2 \cdot_A Y, \mathcal{E}_1 \cdot_A X, Z);$$

$$\Psi(Y, \mathcal{E}_2 \cdot_A \mathcal{E}_1 \cdot_A X, Z) = \Psi(\mathcal{E}_1 \cdot_A X, \mathcal{E}_2 \cdot_A Y, Z).$$

Thus by the symmetry of Ψ , we have

$$\Psi(X, \mathcal{E}_1 \cdot_A \mathcal{E}_2 \cdot_A Y, Z) = \Psi(Y, \mathcal{E}_1 \cdot_A \mathcal{E}_2 \cdot_A X, Z).$$

By Lemma 5.18, we have

$$\Psi(\mathcal{E}_1 \cdot_A \mathcal{E}_2, X, Y) = -((\mathcal{E}_1 \cdot_A \mathcal{E}_2) *_A e) \cdot_A X \cdot_A Y.$$

For all $X, Y \in \Gamma(A)$, by (35), we have

$$(X *_A (\mathcal{E}_1 \cdot_A \mathcal{E}_2)) \cdot_A Y - (Y *_A (\mathcal{E}_1 \cdot_A \mathcal{E}_2)) \cdot_A X$$

= $\Psi(\mathcal{E}_1, X, \mathcal{E}_2) \cdot_A Y + (X *_A \mathcal{E}_1) \cdot_A \mathcal{E}_2 \cdot_A Y + (X *_A \mathcal{E}_2) \cdot_A \mathcal{E}_1 \cdot_A Y$
 $-\Psi(\mathcal{E}_1, Y, \mathcal{E}_2) \cdot_A X - (Y *_A \mathcal{E}_1) \cdot_A \mathcal{E}_2 \cdot_A X - (Y *_A \mathcal{E}_2) \cdot_A \mathcal{E}_1 \cdot_A X.$

By (39) and (50), we have

$$\Psi(\mathcal{E}_1, X, \mathcal{E}_2) \cdot_A Y - \Psi(\mathcal{E}_1, Y, \mathcal{E}_2) \cdot_A X = \Psi(\mathcal{E}_1 \cdot_A Y, X, \mathcal{E}_2) - \Psi(\mathcal{E}_1 \cdot_A X, Y, \mathcal{E}_2) = 0.$$

Using the above relation and by (49), we have

$$(X *_A (\mathcal{E}_1 \cdot_A \mathcal{E}_2)) \cdot_A Y - (Y *_A (\mathcal{E}_1 \cdot_A \mathcal{E}_2)) \cdot_A X$$

= $(X *_A \mathcal{E}_1) \cdot_A \mathcal{E}_2 \cdot_A Y + (X *_A \mathcal{E}_2) \cdot_A \mathcal{E}_1 \cdot_A Y - (Y *_A \mathcal{E}_1) \cdot_A \mathcal{E}_2 \cdot_A X - (Y *_A \mathcal{E}_2) \cdot_A \mathcal{E}_1 \cdot_A X$
= $(Y *_A \mathcal{E}_1) \cdot_A \mathcal{E}_2 \cdot_A X + (Y *_A \mathcal{E}_2) \cdot_A \mathcal{E}_1 \cdot_A X - (Y *_A \mathcal{E}_1) \cdot_A \mathcal{E}_2 \cdot_A X - (Y *_A \mathcal{E}_2) \cdot_A \mathcal{E}_1 \cdot_A X$
= $0,$

which implies that

$$(X *_A (\mathcal{E}_1 \cdot_A \mathcal{E}_2)) \cdot_A Y = (Y *_A (\mathcal{E}_1 \cdot_A \mathcal{E}_2)) \cdot_A X.$$

Thus $\mathcal{E}_1 \cdot_A \mathcal{E}_2 \in \mathfrak{E}(A)$.

For the second claim, using relation (50) with X and Y replaced by $\mathcal{E}^{-1} \cdot_A X$ and $\mathcal{E}^{-1} \cdot_A Y$ respectively, we have

$$0 = \Psi(\mathcal{E}^{-1} \cdot_A X, \mathcal{E} \cdot_A \mathcal{E}^{-1} \cdot_A Y, Z) - \Psi(\mathcal{E}^{-1} \cdot_A Y, \mathcal{E} \cdot_A \mathcal{E}^{-1} \cdot_A X, Z)$$

= $\Psi(\mathcal{E}^{-1} \cdot_A X, Y, Z) - \Psi(\mathcal{E}^{-1} \cdot_A Y, X, Z).$

By the symmetry of Ψ and Lemma 5.18, we have

$$\Psi(\mathcal{E}^{-1}, X, Y) = -(\mathcal{E}^{-1} *_A e) \cdot_A X \cdot_A Y$$

By (39) and (50), we have

(52)
$$\Psi(X, \mathcal{E}, \mathcal{E}^{-1}) \cdot_A Y = \Psi(Y, \mathcal{E}, \mathcal{E}^{-1}) \cdot_A X.$$

Furthermore, by a direct calculation, we have

$$(X *_A \mathcal{E}^{-1}) \cdot_A Y \cdot_A \mathcal{E} = \Psi(X, \mathcal{E}, \mathcal{E}^{-1}) \cdot_A Y - (X *_A e) \cdot_A Y + (X *_A \mathcal{E}) \cdot_A Y \cdot_A \mathcal{E}^{-1},$$

$$(Y *_A \mathcal{E}^{-1}) \cdot_A X \cdot_A \mathcal{E} = \Psi(Y, \mathcal{E}, \mathcal{E}^{-1}) \cdot_A X - (Y *_A e) \cdot_A X + (Y *_A \mathcal{E}) \cdot_A X \cdot_A \mathcal{E}^{-1}.$$

By (43), (49) and (52), we have

$$(X *_A \mathcal{E}^{-1}) \cdot_A Y \cdot_A \mathcal{E} = (Y *_A \mathcal{E}^{-1}) \cdot_A X \cdot_A \mathcal{E}.$$

Because \mathcal{E} is invertible, we have

$$(X *_A \mathcal{E}^{-1}) \cdot_A Y = (Y *_A \mathcal{E}^{-1}) \cdot_A X.$$

Thus \mathcal{E}^{-1} is an eventual identity on *A*.

Proposition 5.20. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid with an identity *e*. Then \mathcal{E} is a pseudoeventual identity on *A* if and only if $(A, *_A, \cdot_{\mathcal{E}}, a_A)$ is a pre-*F*-algebroid, where $\cdot_{\mathcal{E}} : \Gamma(A) \times \Gamma(A) \longrightarrow$ $\Gamma(A)$ is given by (23).

Proof. Define

$$\Psi(X, Y, Z) = X *_A (Y \cdot_{\mathcal{E}} Z) - (X *_A Y) \cdot_{\mathcal{E}} Z - Y \cdot_{\mathcal{E}} (X *_A Z), \quad \forall X, Y, Z \in \Gamma(A).$$

By a straightforward computation, we have

(53)
$$\tilde{\Psi}(X, Y, Z) = \Psi(X, \mathcal{E} \cdot_A Y, Z) + \Psi(X, \mathcal{E}, Y) \cdot_A Z + (X *_A \mathcal{E}) \cdot_A Y \cdot_A Z,$$

(54)
$$\Psi(Y, X, Z) = \Psi(Y, \mathcal{E} \cdot_A X, Z) + \Psi(Y, \mathcal{E}, X) \cdot_A Z + (Y *_A \mathcal{E}) \cdot_A X \cdot_A Z.$$

By the symmetry of Ψ , $(A, *_A, \cdot_{\mathcal{E}}, a_A)$ is a pre-*F*-algebroid if and only if

(55)
$$\Psi(X, \mathcal{E} \cdot_A Y, Z) - \Psi(Y, \mathcal{E} \cdot_A X, Z) = (Y *_A \mathcal{E}) \cdot_A X \cdot_A Z - (X *_A \mathcal{E}) \cdot_A Y \cdot_A Z.$$

By the symmetry of Ψ and (40), we have

$$\Psi(X, \mathcal{E} \cdot_A Y, e) - \Psi(Y, \mathcal{E} \cdot_A X, e) = \Psi(e \cdot_A Y, \mathcal{E}, X) - \Psi(e \cdot_A X, \mathcal{E}, Y) = 0.$$

Taking Z = e in (55), we have

$$(X *_A \mathcal{E}) \cdot_A Y = (Y *_A \mathcal{E}) \cdot_A X.$$

This implies that (49) holds. Furthermore, by (49), (55) implies that (50) holds. By Lemma 5.18, (50) is equivalent to (48). Thus \mathcal{E} is a pseudo-eventual identity on $(A, *_A, \cdot_A, e, a_A)$.

Conversely, if \mathcal{E} is a pseudo-eventual identity on $(A, *_A, \cdot_A, e, a_A)$, by Lemma 5.18, we have

$$\Psi(X, \mathcal{E} \cdot_A Y, Z) = \Psi(Y, \mathcal{E} \cdot_A X, Z).$$

Furthermore, by (49), (55) follows. Thus $(A, *_A, \cdot_{\mathcal{E}}, a_A)$ is a pre-*F*-algebroid.

Corollary 5.21. Let (M, ∇, \bullet) be an *F*-manifold with a compatible flat connection and \mathcal{E} a pseudo-eventual identity on *M*. Then $(M, \nabla, \bullet_{\mathcal{E}})$ is also an *F*-manifold with a compatible flat connection, where $\bullet_{\mathcal{E}}$ is given by

(56)
$$X \bullet_{\mathcal{E}} Y = X \bullet Y \bullet \mathcal{E}, \quad \forall X, Y \in \mathfrak{X}(M).$$

Theorem 5.22. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid with an identity *e*. Then \mathcal{E} is an eventual identity on *A* if and only if $(A, *_A, \cdot_{\mathcal{E}}, a_A)$ is a pre-*F*-algebroid with the identity \mathcal{E}^{-1} , which is called the Dubrovin's dual of $(A, *_A, \cdot_A, a_A)$, where $\cdot_{\mathcal{E}}$ is given by (23). Moreover, *e* is an eventual identity on the pre-*F*-algebroid $(A, *_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A)$ and the map

(57)
$$(A, *_A, \cdot_A, e, a_A, \mathcal{E}) \longrightarrow (A, *_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A, e^{\dagger})$$

is an involution of the set of pre-*F*-algebroids with eventual identities, where $e^{\dagger} = \mathcal{E}^{-2}$ is the inverse of *e* with respect to the multiplication $\cdot_{\mathcal{E}}$.

Proof. By Proposition 5.20, the first claim follows immediately. For the second claim, assume that \mathcal{E} is an eventual identity on $(A, *_A, \cdot_A, e, a_A)$. We need to show that *e* is an eventual identity on the pre-*F*-algebroid $(A, *_A, \cdot_E, \mathcal{E}^{-1}, a_A)$, i.e.

(58)
$$\tilde{\Psi}(e, X, Y) = -(e *_A \mathcal{E}^{-1}) \cdot_{\mathcal{E}} X \cdot_{\mathcal{E}} Y,$$

(59)
$$(X *_A e) \cdot_{\mathcal{E}} Y = (Y *_A e) \cdot_{\mathcal{E}} X.$$

By (43), we have

$$(X *_A e) \cdot_{\mathcal{E}} Y - (Y *_A e) \cdot_{\mathcal{E}} X = ((X *_A e) \cdot_A Y - (Y *_A e) \cdot_A X) \cdot_A \mathcal{E} = 0,$$

which implies that (59) holds.

On the one hand, by (48) and (50), we have

$$\begin{split} \Psi(e, X, Y) &= \Psi(e, \mathcal{E} \cdot_A X, Y) + \Psi(e, \mathcal{E}, X) \cdot_A Y + (e *_A \mathcal{E}) \cdot_A X \cdot_A Y \\ &= \Psi(\mathcal{E}, X, Y) + \Psi(\mathcal{E}, e, X) \cdot_A Y + (e *_A \mathcal{E}) \cdot_A X \cdot_A Y \\ &= -(\mathcal{E} *_A e) \cdot_A X \cdot_A Y - (\mathcal{E} *_A e) \cdot_A X \cdot_A Y + (e *_A \mathcal{E}) \cdot_A X \cdot_A Y \\ &= -2(\mathcal{E} *_A e) \cdot_A X \cdot_A Y + (e *_A \mathcal{E}) \cdot_A X \cdot_A Y. \end{split}$$

On the other hand, taking $X = \mathcal{E}$ and $Y = \mathcal{E}^{-1}$ in (48), by the symmetry of Ψ , we have

$$\Psi(e,\mathcal{E},\mathcal{E}^{-1})=\Psi(\mathcal{E},e,\mathcal{E}^{-1})=-(\mathcal{E}*_{A}e)\cdot_{A}\mathcal{E}^{-1},$$

which implies that

$$e *_A e - (e *_A \mathcal{E}) \cdot_A \mathcal{E}^{-1} - (e *_A \mathcal{E}^{-1}) \cdot_A \mathcal{E} = -(\mathcal{E} *_A e) \cdot_A \mathcal{E}^{-1}.$$

Furthermore, by (43), we have

$$(e *_A \mathcal{E}^{-1}) \cdot_A \mathcal{E}^2 = (e *_A e) \cdot_A \mathcal{E} - e *_A \mathcal{E} + \mathcal{E} *_A e = 2\mathcal{E} *_A e - e *_A \mathcal{E}.$$

Thus we have

$$\tilde{\Psi}(e, X, Y) = -(e *_A \mathcal{E}^{-1}) \cdot_A \mathcal{E}^2 \cdot_A X \cdot_A Y = -(e *_A \mathcal{E}^{-1}) \cdot_{\mathcal{E}} X \cdot_{\mathcal{E}} Y.$$

Therefore, *e* is an eventual identity on the *F*-algebroid with a compatible pre-Lie algebroid $(A, *_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A)$.

By Proposition 5.19, $e^{\dagger} = \mathcal{E}^{-2}$ is an eventual identity on the pre-*F*-algebroid $(A, *_A, \cdot_{\mathcal{E}}, \mathcal{E}^{-1}, a_A)$. Then similar to the proof of Theorem 4.4, the map given by (57) is an involution of the set of pre-*F*-algebroids with eventual identities.

Example 5.23. Consider the PreLie-Com algebra $(g, *, \cdot)$ with an identity *e* given by Example 5.2. By a direct calculation, for any $\mathcal{E} \in g$, we have

$$(x * \mathcal{E}) \cdot y - (y * \mathcal{E}) \cdot x = x \cdot D(\mathcal{E}) \cdot y - y \cdot D(\mathcal{E}) \cdot x = 0, \quad \forall x, y \in \mathfrak{g}$$

By Proposition 5.17, \mathcal{E} is a pseudo-eventual identity on g. Thus any element of g is a pseudoeventual identity on g. Furthermore, for any $\mathcal{E} \in g$, there is a new pre-F-manifold algebra structure on g given by

$$x \cdot_{\mathcal{E}} y = x \cdot y \cdot \mathcal{E}, \quad x * y = x \cdot D(y), \quad \forall x, y \in \mathfrak{g}.$$

Example 5.24. Let (M, ∇, \bullet, e) be a semi-simple PreLie-Com manifold with local coordinate systems (u^1, \dots, u^n) . Then any pseudo-eventual identity on *TM* is of the form

$$\mathcal{E} = f_1(u^1)\frac{\partial}{\partial u^1} + \dots + f_n(u^n)\frac{\partial}{\partial u^n}.$$

where $f_i(u^i) \in C^{\infty}(M)$ depends only on u^i for $i = 1, 2, \dots, n$. Furthermore, if all $f_i(u^i)$ are non-vanishing everywhere, then $\mathcal{E} \in \mathfrak{X}(M)$ is an eventual identity.

Example 5.25. Let (u^1, u^2) be a local coordinate systems on \mathbb{R}^2 . Define two multiplications by

$$\frac{\partial}{\partial u^1} \bullet \frac{\partial}{\partial u^i} = \frac{\partial}{\partial u^i}, \quad \frac{\partial}{\partial u^2} \bullet \frac{\partial}{\partial u^2} = 0, \quad \frac{\partial}{\partial u^i} * \frac{\partial}{\partial u^j} = 0, \quad i, j \in \{1, 2\}.$$

Then $(T\mathbb{R}^2, *, \bullet, \mathrm{Id})$ is a PreLie-Com algebroid with the identity $\frac{\partial}{\partial u^1}$ and thus $(T\mathbb{R}^2, *, \bullet, \mathrm{Id})$ is a pre-*F*-algebroid with the identity $\frac{\partial}{\partial u^1}$.

Furthermore, any pseudo-eventual identity on $(T\mathbb{R}^2, *, \bullet, Id)$ is of the form

$$\mathcal{E} = f_1(u^1)\frac{\partial}{\partial u^1} + f_2(u^1, u^2)\frac{\partial}{\partial u^2}$$

with $\frac{\partial f_1}{\partial u^1} = \frac{\partial f_2}{\partial u^2}$, where $f_1 \in C^{\infty}(\mathbb{R}^2)$ depends only on u^1 and f_2 is any smooth function. Furthermore, any pseudo-eventual identity on the sub-adjacent F-algebroid of $(T\mathbb{R}^2, *, \bullet, \mathrm{Id})$ is of the form

$$\mathcal{E} = f_1(u^1)\frac{\partial}{\partial u^1} + f_2(u^1, u^2)\frac{\partial}{\partial u^2}.$$

In particular, if $f_1(u^1)$ is non-vanishing everywhere, then \mathcal{E} is an eventual identity on the subadjacent *F*-algebroid of $(T\mathbb{R}^2, *, \bullet, Id)$.

5.3. Nijenhuis operators and deformed pre-*F*-algebroids. Recall from [19] that a Nijenhuis operator on a pre-Lie algebroid $(A, *_A, a_A)$ is a bundle map $N : A \to A$ such that

(60)
$$N(X) *_A N(Y) = N(N(X) *_A Y + X *_A N(Y) - N(X *_A Y)), \quad \forall X, Y \in \Gamma(A).$$

Definition 5.26. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid. A bundle map $N : A \rightarrow A$ is called a **Nijenhuis operator** on $(A, *_A, \cdot_A, a_A)$ if N is both a Nijenhuis operator on the commutative associative algebra ($\Gamma(A), \cdot_A$) and a Nijenhuis operator on the pre-Lie algebroid $(A, *_A, a_A)$.

Theorem 5.27. Let $N : A \longrightarrow A$ be a Nijenhuis operator on a pre-*F*-algebroid $(A, *_A, \cdot_A, a_A)$. Then $(A, *_N, \cdot_N, a_N = a_A \circ N)$ is a pre-*F*-algebroid and N is a homomorphism from the pre-*F*-algebroid $(A, *_N, \cdot_N, a_N = a_A \circ N)$ to $(A, *_A, \cdot_A, a_A)$, where the operation \cdot_N is given by (29) and the operation $*_N : \Gamma(A) \times \Gamma(A) \longrightarrow \Gamma(A)$ is given by

(61)
$$X *_N Y = N(X) *_A Y + X *_A N(Y) - N(X *_A Y), \quad \forall X, Y \in \Gamma(A).$$

Proof. Since *N* is a Nijenhuis operator on the commutative associative algebra ($\Gamma(A)$, \cdot_A), it follows that ($\Gamma(A)$, \cdot_N) is a commutative associative algebra. Since *N* is a Nijenhuis operator on the pre-Lie algebroid (A, $*_A$, a_A), (A, $*_N$, a_N) is a pre-Lie algebroid ([19]).

Define

(62)
$$\Psi_N(X, Y, Z) := X *_N (Y \cdot_N Z) - (X *_N Y) \cdot_N Z - (X *_N Z) \cdot_N Y, \quad \forall X, Y, Z \in \Gamma(A).$$

By a direct calculation, we have

$$\begin{split} \Psi_N(X,Y,Z) &= \Psi(NX,NY,Z) + \Psi(NX,Y,NZ) + \Psi(X,NY,NZ) \\ &- N\big(\Psi(NX,Y,Z) + \Psi(X,NY,Z) + \Psi(X,Y,NZ)\big) + N^2(\Psi(X,Y,Z)). \end{split}$$

Thus by (35), we have

$$\Psi_N(X, Y, Z) = \Psi_N(Y, X, Z).$$

This implies that $(A, *_N, \cdot_N, a_N = a_A \circ N)$ is a pre-*F*-algebroid. It is not hard to see that *N* is a homomorphism from the pre-*F*-algebroid $(A, *_N, \cdot_N, a_N = a_A \circ N)$ to $(A, *_A, \cdot_A, a_A)$.

Proposition 5.28. Let $(A, *_A, \cdot_A, a_A)$ be a pre-*F*-algebroid with an identity *e* and \mathcal{E} a pseudoeventual identity on *A*. Then the endomorphism $N = \mathcal{E}_A$ is a Nijenhuis operator on the pre-*F*algebroid $(A, *_A, \cdot_A, a_A)$. Furthermore, $(A, *_{\mathcal{E}}, \cdot_{\mathcal{E}}, a_{\mathcal{E}})$ is a pre-*F*-algebroid, where the multiplication $*_{\mathcal{E}}$ is given by

(63)
$$X *_{\mathcal{E}} Y = (\mathcal{E} \cdot_A X) *_A Y + X *_A (\mathcal{E} \cdot_A Y) - \mathcal{E} \cdot_A (X *_A Y), \quad \forall X, Y \in \Gamma(A),$$

the multiplication $\cdot_{\mathcal{E}}$ is given by (23) and $a_{\mathcal{E}}(X) = a_A(\mathcal{E} \cdot_A X)$.

Proof. First, we show that N is a Nijenhuis operator on the pre-Lie algebroid $(A, *_A, a_A)$. By (35), we have

$$\Psi(\mathcal{E} \cdot_A X, \mathcal{E}, Y) = \Psi(Y, \mathcal{E} \cdot_A X, \mathcal{E}), \quad \forall X, Y \in \Gamma(A),$$

which implies that

$$(64) \qquad (\mathcal{E} \cdot_A X) *_A (\mathcal{E} \cdot_A Y) = Y *_A (X \cdot_A \mathcal{E} \cdot_A \mathcal{E}) - (Y *_A (\mathcal{E} \cdot_A X)) \cdot_A \mathcal{E} + ((\mathcal{E} \cdot X) *_A Y) \cdot_A \mathcal{E}.$$

Since \mathcal{E} is a pseudo-eventual identity on A, by (48) and the symmetry of Ψ , we have

$$\Psi(X, \mathcal{E}, Y) = -(\mathcal{E} *_A e) \cdot_A X \cdot_A Y.$$

which implies that

(65)
$$X *_A (\mathcal{E} \cdot_A Y) = -(\mathcal{E} *_A e) \cdot_A X \cdot_A Y - (X *_A \mathcal{E}) \cdot_A Y - (X *_A Y) \cdot_A \mathcal{E}.$$

By (48), (49), (64), (65) and the symmetry of Ψ , we have

$$N(X) *_{A} N(Y) - N(N(X) *_{A} Y + X *_{A} N(Y) - N(X *_{A} Y))$$

$$= (\mathcal{E} \cdot_{A} X) *_{A} (\mathcal{E} \cdot_{A} Y) - \mathcal{E} \cdot_{A} ((\mathcal{E} \cdot_{A} X) *_{A} Y + X *_{A} (\mathcal{E} \cdot_{A} Y) - \mathcal{E} \cdot_{A} (X *_{A} Y))$$

$$= Y *_{A} (X \cdot_{A} \mathcal{E} \cdot_{A} \mathcal{E}) - (Y *_{A} (X \cdot_{A} \mathcal{E})) \cdot_{A} \mathcal{E} - (X *_{A} \mathcal{E}) \cdot_{A} Y \cdot_{A} \mathcal{E} + (\mathcal{E} *_{A} e) \cdot_{A} X \cdot_{A} Y \cdot_{A} \mathcal{E}$$

$$= Y *_{A} (X \cdot_{A} \mathcal{E} \cdot_{A} \mathcal{E}) - (Y *_{A} (X \cdot_{A} \mathcal{E})) \cdot_{A} \mathcal{E} - (Y *_{A} \mathcal{E}) \cdot_{A} X \cdot_{A} \mathcal{E} + (\mathcal{E} *_{A} e) \cdot_{A} X \cdot_{A} Y \cdot_{A} \mathcal{E}$$

$$= \Psi(Y, X \cdot_{A} \mathcal{E}, \mathcal{E}) + (\mathcal{E} *_{A} e) \cdot_{A} X \cdot_{A} Y \cdot_{A} \mathcal{E}$$

$$= -(\mathcal{E} *_{A} e) \cdot_{A} X \cdot_{A} Y \cdot_{A} \mathcal{E} + (\mathcal{E} *_{A} e) \cdot_{A} X \cdot_{A} Y \cdot_{A} \mathcal{E}$$

$$= 0.$$

Thus $N = \mathcal{E}_A$ is a Nijenhuis operator on the pre-Lie algebroid $(A, *_A, a_A)$.

Also, $N = \mathcal{E}_A$ is a Nijenhuis operator on the commutative associative algebra ($\Gamma(A), \cdot_A$). Therefore, $N = \mathcal{E}_A$ is a Nijenhuis operator on the pre-*F*-algebroid ($A, *_A, \cdot_A, a_A$). The second claim follows.

Corollary 5.29. Let (M, ∇, \bullet) be an *F*-manifold with a compatible flat connection and \mathcal{E} a pseudo-eventual identity on *M*. Then there is a new pre-*F*-algebroid structure on *TM* given by

$$\begin{aligned} X \bullet_{\mathcal{E}} Y &= X \bullet Y \bullet \mathcal{E}, \\ X *_{\mathcal{E}} Y &= \nabla_{\mathcal{E} \bullet X} Y + \nabla_{\mathcal{E} \bullet Y} X - \mathcal{E} \bullet (\nabla_X Y), \\ a_{\mathcal{E}}(X) &= \mathcal{E} \bullet X, \quad \forall X, Y \in \mathfrak{X}(M). \end{aligned}$$

5.4. Applications to integral systems.

Theorem 5.30. ([23]) Let (M, ∇, \bullet) be an *F*-manifold with a compatible flat connection. Let (u^1, u^2, \dots, u^n) be the canonical coordinate systems on *M*. If *X* and *Y* in $\mathfrak{X}(M)$ satisfy

$$\nabla_Z X) \bullet W = (\nabla_W X) \bullet Z, \quad (\nabla_Z Y) \bullet W = (\nabla_W Y) \bullet Z, \quad \forall \ W, Z \in \mathfrak{X}(M),$$

then the associated flows

(66)
$$u_t^i = c_{jk}^i X^k u_x^i \quad and \quad u_\tau^i = c_{jk}^i Y^k u_x^j$$

commute, where $\frac{\partial}{\partial u^i} \bullet \frac{\partial}{\partial u^j} = c_{ij}^k \frac{\partial}{\partial u^k}$, $X = X^i \frac{\partial}{\partial u^i}$ and $Y = Y^i \frac{\partial}{\partial u^i}$.

Proposition 5.31. Let (M, ∇, \bullet) be an *F*-manifold with a compatible flat connection and an identity *e*. Assume that $\mathcal{E}_1, \mathcal{E}_2 \in \mathfrak{X}(M)$ are pseudo-eventual identities. Then the flows

(67)
$$u_t^i = c_{jk}^i X^k u_x^i, \quad u_\tau^i = c_{jk}^i Y^k u_x^j, \quad u_s^i = X^p Y^q c_{jk}^i c_{pq}^k u_x^j$$

commute, where $\frac{\partial}{\partial u^i} \bullet \frac{\partial}{\partial u^j} = c_{ij}^k \frac{\partial}{\partial u^k}$, $\mathcal{E}_1 = X^i \frac{\partial}{\partial u^i}$ and $\mathcal{E}_2 = Y^i \frac{\partial}{\partial u^i}$.

Proof. Since $\mathcal{E}_1 \in \mathfrak{X}(M)$ and $\mathcal{E}_2 \in \mathfrak{X}(M)$ are pseudo-eventual identities on (M, ∇, \bullet) , by Proposition 5.19, $\mathcal{E}_1 \bullet \mathcal{E}_2$ is also a pseudo-eventual identity. Thus $\mathcal{E}_1, \mathcal{E}_2$ and $\mathcal{E}_1 \bullet \mathcal{E}_2$ satisfy (49). Furthermore, we have

$$\mathcal{E}_1 \bullet \mathcal{E}_2 = X^p Y^q c_{pq}^k \frac{\partial}{\partial u^k}.$$

By Theorem 5.30, the claim follows.

Theorem 5.32. ([23]) Let (M, ∇, \bullet) be an *F*-manifold with a compatible flat connection. Let (u^1, u^2, \dots, u^n) be the canonical coordinate systems on *M* and $(X_{(1,0)}, \dots, X_{(n,0)})$ a basis of flat vector fields. Define the primary flows by

(68)
$$u_{t_{(p,0)}}^i = c_{jk}^i X_{(p,0)}^k u_x^j.$$

Then there is a well-defined higher flows of the hierarchy defined as

(69)
$$u_{t_{(p,\alpha)}}^i = c_{jk}^i X_{(p,\alpha)}^k u_x^j.$$

by means of the following recursive relations:

(70)
$$\nabla_{\frac{\partial}{\partial u^j}} X^i_{(p,\alpha)} = c^i_{jk} X^k_{(p,\alpha-1)} u^k_x.$$

Furthermore, the flows of the principal hierarchy (69) *commute.*

Proposition 5.33. Let (M, ∇, \bullet) be an *F*-manifold with a compatible flat connection and an identity e. Let $(X_{(1,0)}, \dots, X_{(n,0)})$ be a basis of flat vector fields. Assume that $\mathcal{E} \in \mathfrak{X}(M)$ is a pseudo-eventual identity. Define the primary flows by

(71)
$$u_{t_{(p,0)}}^{i} = c_{jk}^{m} c_{ml}^{i} \mathcal{E}^{l} X_{(p,0)}^{k} u_{x}^{j},$$

where $\mathcal{E} = \mathcal{E}^{i} \frac{\partial}{\partial u^{i}}$. Then there is a well-defined higher flows of the hierarchy defined as

(72)
$$u_{t_{(p,\alpha)}}^i = c_{jk}^m c_{ml}^i \mathcal{E}^l X_{(p,\alpha)}^k u_x^j.$$

by means of the following recursive relations:

(73)
$$\nabla_{\frac{\partial}{\partial u^{j}}} X^{i}_{(p,\alpha)} = c^{m}_{jk} c^{i}_{ml} \mathcal{E}^{l} X^{k}_{(p,\alpha-1)} u^{k}_{x}.$$

Furthermore, the flows of the principal hierarchy (72) commute.

Proof. Since $\mathcal{E} \in \mathfrak{X}(M)$ is a pseudo-eventual identity on (M, ∇, \bullet) , by Proposition 5.20, $(M, \nabla, \bullet_{\mathcal{E}})$ is also an *F*-manifold with a compatible flat connection, where

$$X \bullet_{\mathcal{E}} Y = X \bullet Y \bullet \mathcal{E}, \quad \forall X, Y \in \mathfrak{X}(M).$$

Furthermore, we have

$$\frac{\partial}{\partial u^i} \bullet_{\mathcal{E}} \frac{\partial}{\partial u^j} = c^m_{ij} c^k_{ml} \mathcal{E}^l \frac{\partial}{\partial u^k}.$$

By Theorem 5.32, the claim follows.

References

- [1] A. Arsie, A. Buryak, P. Lorenzoni and P. Rossi, Flat *F*-manifolds, *F*-CohFTs, and integrable hierarchies. *Comm. Math. Phys.* 388 (2021), 291-328. 2
- [2] A. Arsie and P. Lorenzoni, From the Darboux-Egorov system to bi-flat F-manifolds. J. Geom. Phys. 70 (2013), 98-116.2
- [3] A. Arsie and P. Lorenzoni, F-manifolds with eventual identities, bidifferential calculus and twisted Lenard-Magri chains. Int. Math. Res. Not. (2013), 3931-3976.2
- [4] A. Arsie and P. Lorenzoni, F-manifolds, multi-flat structures and Painlevé transcendents. Asian J. Math. 23 (2019), 877-904.2
- [5] D. Burde, Left-symmetric algebras, or pre-Lie algebras in geometry and physics. *Cent. Eur. J. Math.* 4 (2006), 323-357.2
- [6] J. F. Cariñena, J. Grabowski and G. Marmo, Quantum Bi-Hamiltonian Systems. Int. J. Mod. Phys. A 15 (2000), 4797-4810. 15, 16
- [7] J. A. Cruz Morales and A. Torres-Gomez, On *F*-algebroids and Dubrovin's duality. *Arch. Math. (Brno)* 55 (2019), 109-122. 2, 4
- [8] L. David and I. A. B. Strachan, Compatible metrics on a manifold and nonlocal bi-Hamiltonian structures. *Int. Math. Res. Not.* (2004), 3533-3557. 2
- [9] L. David and I. A. B. Strachan, Dubrovin's duality for *F*-manifolds with eventual identities. *Adv. Math.* 226 (2011), 4031-4060. 2, 13, 15
- [10] V. Dotsenko, Algebraic structures of *F*-manifolds via pre-Lie algebras. *Ann. Mat. Pura Appl.* 198 (2019), no. 2, 517-527. 2, 18
- [11] B. Dubrovin, Geometry of 2D topological field theories. Lecture Notes in Math, 1620 (1995). 1
- [12] C. Hertling, Frobenius Manifolds and Moduli Spaces for Singularities. Cambridge Tracts in Math. Cambridge University Press, 2002. 2, 10
- [13] C. Hertling and Y. I. Manin, Weak Frobenius manifolds. Int. Math. Res. Not. (1999), no. 6, 277-286. 1
- [14] M. Kato, T. Mano and J. Sekiguchi, Flat structure on the space of isomonodromic deformations. SIGMA Symmetry Integrability Geom. Methods Appl. 16 (2020), No. 110, 36 pp. 2
- [15] H. Kawakami and T. Mano, Regular flat structure and generalized Okubo system. Comm. Math. Phys. 369 (2019), 403-431.2
- [16] Y. Kosmann-Schwarzbach and F. Magri, Poisson-Nijenhuis structures. Ann. Inst. Henri Poincaré A 53 (1990), 35-81. 15, 16
- [17] Y. P. Lee, Quantum K-theory, I: Foundations. Duke Math. J. 121 (2004), 389-424. 2
- [18] J. Liu, Y. Sheng and C. Bai, Pre-symplectic algebroids and their applications. *Lett. Math. Phys.* 108 (2018), 779-804.2
- [19] J. Liu, Y. Sheng, C. Bai and Z. Chen, Left-symmetric algebroids. *Math. Nachr.* 289 (2016), 1893-1908. 2, 7, 8, 18, 26
- [20] J. Liu, Y. Sheng and C. Bai, F-manifold algebras and deformation quantization via pre-Lie algebras. J. Algebra 559 (2020), 467-495. 2, 18
- [21] P. Lorenzoni, Darboux-Egorov system, bi-flat F-manifolds and Painlevé VI. Int. Math. Res. Not. (2014), 3279-3302.2
- [22] P. Lorenzoni and M. Pedroni, Natural connections for semi-Hamiltonian systems: the case of the ϵ -system. Lett. Math. Phys. 97 (2011), 85-108. 2

 \Box

- [23] P. Lorenzoni, M. Pedroni and A. Raimondo, F-manifolds and integrable systems of hydrodynamic type. Arch. Math. (Brno) 47 (2011), 163-180. 2, 28
- [24] K. C. H. Mackenzie, General theory of Lie groupoids and Lie algebroids. *Lecture Note Series*, 213. *London Mathematical Society.*, Cambridge University Press, Cambridge, 2005. 2, 4, 7
- [25] S. A. Merkulov, Operads, deformation theory and F-manifolds. Asp. Math. 36 (2004), 213-251.2
- [26] Y. I. Manin, F-manifolds with flat structure and Dubrovin's duality. Adv. Math. 198 (2005), 5-26. 2, 19
- [27] M. Nguiffo Boyom, Cohomology of Koszul-Vinberg algebroids and Poisson manifolds. I. Banach Center Publ. 54 (2001), 99-110. 2, 7
- [28] M. Nguiffo Boyom, KV-cohomology of Koszul-Vinberg algebroids and Poisson manifolds. Internat. J. Math. 16 (2005), 1033-1061.2
- [29] C. Noémie and Y. I. Manin, F-manifolds and geometry of information. Bull. Lond. Math. Soc. 52 (2020), 777-792. 2

Departmento de matemáticas, Universidad Nacional de Colombia, Bogotá, Max Planck Institute for Mathematics, Vivatsgasse 7, Bonn, Germany

Email address: jacruzmo@unal.edu.co, alexcruz@mpim-bonn.mpg.de

School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, Jilin, China *Email address*: liujf534@nenu.edu.cn

DEPARTMENT OF MATHEMATICS, JILIN UNIVERSITY, CHANGCHUN 130012, JILIN, CHINA *Email address*: shengyh@jlu.edu.cn