

Connecting Dark Gauge Symmetry to the Standard Model

Ernest Ma

*Department of Physics and Astronomy,
University of California, Riverside, California 92521, USA*

Abstract

Dark matter is postulated to be a neutral Dirac fermion, charged under a dark $U(1)_D$ gauge symmetry. Scalar partners of the quarks and leptons are also charged under $U(1)_D$. The dark gauge boson Z_D and the dark Higgs boson h_D enable either freeze-out or freeze-in mechanisms to account for the correct dark matter relic abundance. Dark number D is connected to baryon number B and lepton number L through $D = 3B + L - (2j)_{[mod\ 2]}$ where j is the intrinsic spin of the particle.

Introduction : The nature of dark matter [1] is unknown, but it is likely to be stabilized by an unbroken symmetry, the simplest of which is dark parity, under which it is odd and all particles of the Standard Model (SM) are even. In many proposed models involving Majorana neutrinos, it is derivable [2] from lepton parity, i.e. $\pi_D = \pi_L(-1)^{2j}$, where $\pi_L = (-1)^L$ and j is the intrinsic spin of the particle. For Dirac neutrinos with the conservation of lepton number L , the analog connection [3] is $D = L - (2j)_{[mod\ 2]}$. If baryon number B is also considered, it may become $D = 3B + L - (2j)_{[mod\ 2]}$. In this paper, it is shown how it comes about from a dark gauged $U(1)_D$ symmetry, which is broken spontaneously by three units [4], instead of the customary two.

To support an anomaly-free $U(1)_D$ gauge symmetry, a neutral Dirac fermion N is proposed with dark charge $D = -1$. It connects to quarks and leptons through their scalar counterparts with $D = 1$. The particle content is therefore very similar to that of the supersymmetric standard model. There are however very important differences. Instead of two Higgs superfields, there is only the one SM Higgs doublet, and there are no gauginos. In contrast, there is a dark gauge boson Z_D and a dark Higgs boson h_D . Together with N as dark matter, they participate in either freeze-out or freeze-in scenarios. Whereas Z_D decays immediately to $N\bar{N}$ if kinematically allowed, h_D decays mainly through its mixing with the SM Higgs boson. If $m_{Z_D} < 2m_N$, then Z_D decays to an SM fermion-antifermion pair, through the latter's dark magnetic moment. The model and its details are described below.

Model : The idea of this model and its implementation are both very simple. All particles of the SM do not transform under $U(1)_D$. To each quark and lepton, a scalar counterpart is added which is charged +1 under $U(1)_D$, thereby connecting to the neutral Dirac fermion N which is charged -1 . The neutral scalar ζ^0 breaks $U(1)_D$ by three units. It cannot couple to $N_L N_L$ or $N_R N_R$, hence the global symmetry D remains, with N having $D = -1$

fermion/scalar	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$U(1)_D$	B	L
$(u, d)_L$	3	2	1/6	0	1/3	0
u_R	3	1	2/3	0	1/3	0
d_R	3	1	-1/3	0	1/3	0
$(\nu, l)_L$	1	2	-1/2	0	0	1
ν_R	1	1	0	0	0	1
l_R	1	1	-1	0	0	1
(ϕ^+, ϕ^0)	1	2	1/2	0	0	0
$N_{L,R}$	1	1	0	-1	0	0
$(\tilde{u}_L, \tilde{d}_L)$	3	2	1/6	1	1/3	0
\tilde{u}_R	3	1	2/3	1	1/3	0
\tilde{d}_R	3	1	-1/3	1	1/3	0
$(\tilde{\nu}_L, \tilde{l}_L)$	1	2	-1/2	1	0	1
$\tilde{\nu}_R$	1	1	0	1	0	1
\tilde{l}_R	1	1	-1	1	0	1
ζ^0	1	1	0	3	0	0

Table 1: Fermions and scalars in the dark $U(1)_D$ model.

and the scalar quarks and leptons having $D = 1$, thereby ensuring the connection $D = 3B + L - (2j)_{[mod\ 2]}$.

Dark $U(1)_D$ Gauge Symmetry: Whereas the SM $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge symmetry is broken by the one Higgs doublet $\Phi = (\phi^+, \phi^0)$, the dark $U(1)_D$ gauge symmetry is broken by the one Higgs singlet ζ^0 . The Higgs potential consisting of Φ and ζ is thus very simple, i.e.

$$V = m_1^2 \Phi^\dagger \Phi + m_2^2 \zeta^* \zeta + \frac{1}{2} \lambda_1 (\Phi^\dagger \Phi)^2 + \frac{1}{2} (\zeta^* \zeta)^2 + \lambda_3 (\Phi^\dagger \Phi) (\zeta^* \zeta). \quad (1)$$

Let $\langle \phi^0 \rangle = v$ and $\langle \zeta \rangle = u$, then

$$v^2 = \frac{-\lambda_2 m_1^2 + \lambda_3 m_2^2}{\lambda_1 \lambda_2 - \lambda_3^2}, \quad u^2 = \frac{-\lambda_1 m_2^2 + \lambda_3 m_1^2}{\lambda_1 \lambda_2 - \lambda_3^2}. \quad (2)$$

The only physical scalars are the SM $h = \sqrt{2} Re(\phi^0)$ and the new dark Higgs boson $h_D =$

$\sqrt{2}Re(\zeta^0)$, and

$$\begin{aligned}
V = & \lambda_1 v^2 h^2 + \lambda_2 u^2 h_D^2 + 2\lambda_3 v u h h_D + \frac{1}{\sqrt{2}} \lambda_1 v h^3 + \frac{1}{\sqrt{2}} \lambda_2 u h_D^3 \\
& + \frac{1}{\sqrt{2}} \lambda_3 v h h_D^2 + \frac{1}{\sqrt{2}} \lambda_3 u h_D h^2 + \frac{1}{8} \lambda_1 h^4 + \frac{1}{8} \lambda_2 h_D^4 + \frac{1}{4} \lambda_3 h^2 h_D^2.
\end{aligned} \tag{3}$$

Hence h_D mixes with h according to

$$M_{hh_D}^2 = \begin{pmatrix} 2\lambda_1 v^2 & 2\lambda_3 v u \\ 2\lambda_3 v u & 2\lambda_2 u^2 \end{pmatrix}, \tag{4}$$

and decays to SM fermions through the h Yukawa couplings. Note that ζ^0 has $D = 3$, hence h_D does not couple to $N_L N_L$ or $N_R N_R$.

As for Z_D , the gauge invariant term $|(\partial^\mu - 3ig_D Z_D^\mu)\zeta|^2$ yields $m_{Z_D} = 3\sqrt{2}g_D u$, together with the interaction

$$\mathcal{L}_{int} = 9\sqrt{2}g_D^2 u h_D Z_D^\mu Z_{D\mu} + \frac{9}{2}g_D^2 h_D^2 Z_D^\mu Z_{D\mu}. \tag{5}$$

This means that h_D may decay into $Z_D Z_D$ if kinematically allowed. It also couples to $N\bar{N}$ indirectly through Z_D in one loop.

Since N is charged under $U(1)_D$, as are the scalar quarks and leptons, they are decay products of Z_D if kinematically allowed. If not, then Z_D may decay into an SM fermion-antifermion pair. The γ^μ coupling is zero from dark gauge invariance. The $\sigma^{\mu\nu}$ coupling is




Figure 1: Dark magnetic moment of SM fermion.

obtained as shown in Fig. 1, where $\tilde{f}_{1,2}$ are the mass eigenstates from $\tilde{f}_L - \tilde{f}_R$ mixing. Let

$\tilde{f}_L = \cos \theta \tilde{f}_1 - \sin \theta \tilde{f}_2$ and $\tilde{f}_R = \sin \theta \tilde{f}_1 + \cos \theta \tilde{f}_2$, with $m_{1,2}^2 \gg m_N^2 \gg m_f^2$, then the dark magnetic moment of f is [5]

$$a_f = \frac{g_f^2 \sin \theta \cos \theta}{4\pi^2} m_f m_N \left[\frac{1}{m_1^2} \left(\ln \frac{m_1^2}{m_N^2} - \frac{7}{4} \right) - \frac{1}{m_2^2} \left(\ln \frac{m_2^2}{m_N^2} - \frac{7}{4} \right) \right], \quad (6)$$

where $g_f^L = g_f^R$ has been assumed. The corresponding interaction is

$$\mathcal{L}_{int} = \frac{ig_D a_f}{2m_f} Z_D^\mu q^\nu \bar{f} \sigma_{\mu\nu} f. \quad (7)$$

Scotogenic Fermion Masses : According to Table 1, quarks and leptons obtain masses as in the SM. However, it is a simple matter to allow some light fermions, such as the Dirac neutrinos, to acquire radiative masses through dark matter [6], by the implementation of a softly broken Z_2 symmetry. For example, let ν_R and $\tilde{\nu}_R$ be odd under this Z_2 , then the dimension-four Yukawa coupling $\bar{\nu}_R(\nu_L \phi^0 - l_L \phi^+)$ is forbidden, but $\bar{\nu}_R N_L \tilde{\nu}_R$ is allowed. Hence the one-loop generation of Dirac neutrino mass is possible as shown in Fig. 2, where the soft




Figure 2: Scotogenic Dirac neutrino mass.

breaking of Z_2 occurs with the scalar trilinear $\tilde{\nu}_R^*(\phi^0 \tilde{\nu}_L - \phi^+ \tilde{l}_L)$ term. Analogous constructions are possible for the electron and muon masses if desired. Note that $U(1)_D$ is also applicable to the dark sector of the recently proposed [7] scotogenic $A_5 \rightarrow A_4$ model of Dirac neutrino masses.

Freeze-Out Scenario : The neutral dark fermion N is a candidate for the dark matter of the Universe from thermal freeze-out. Assuming that $m_N > m_{Z_D}$, the annihilation of




Figure 3: Annihilation of $N\bar{N} \rightarrow Z_D Z_D$.

$N\bar{N} \rightarrow Z_D Z_D$ is shown in Fig. 3. The cross section \times relative velocity is given by [8]

$$\sigma v_{rel} = \frac{g_D^4}{16\pi m_N^2} \left(1 - \frac{m_{Z_D}^2}{m_N^2}\right)^{\frac{3}{2}} \left(1 - \frac{m_{Z_D}^2}{2m_N^2}\right)^{-2}. \quad (8)$$

Setting this value to the canonical $6 \times 10^{-26} \text{ cm}^3/\text{s}$ for a Dirac fermion, and assuming $m_N = 1 \text{ TeV}$ and $m_{Z_D} = 800 \text{ GeV}$, it is satisfied for $g_D = 0.86$. Once produced, Z_D thermalizes with the SM particles through h_D and its interactions listed in Eqs. (3) and (5).

As for direct detection, N interacts with quarks through their scalar counterparts. For simplicity, let $g_u^L = g_u^R = g_d^L = g_d^R = g_0$ and $m_{\tilde{u}_L} = m_{\tilde{u}_R} = m_{\tilde{d}_L} = m_{\tilde{d}_R} = \tilde{m}_0$, then the elastic scattering of N off a Xenon nucleus per nucleon is given by

$$\sigma_0 = \frac{g_0^4 \mu^2}{64\pi(\tilde{m}_0^2 - m_N^2)^2}, \quad (9)$$

where $\mu = m_N m_{Xe}/(m_N + m_{Xe})$ is the reduced mass of N . For $m_N = 1 \text{ TeV}$, and using $m_{Xe} = 122.3 \text{ GeV}$ with $\sigma_0 < 10^{-45} \text{ cm}^2$ [9], the lower limit on \tilde{m}_0/g_0 is about 70 TeV. Note that this severe constraint rules out the possibility that the relic abundance of N comes from this interaction. Without the dark gauge $U(1)_D$ symmetry, N would not be a dark matter candidate in this case.

Freeze-In Scenario : An alternative for N to be dark matter is the freeze-in mechanism [10]. Here N is assumed to be light, and its only production is through Higgs decay [11] as shown in Fig. 4. This scenario works if the reheat temperature of the Universe is 1 to 10 TeV, and




Figure 4: Higgs decay to $N\bar{N}$.

the thermal production of N is very much suppressed, so that its relic abundance comes only from Higgs decay until the latter goes out of thermal equilibrium with the other SM particles. The effective h coupling to $\bar{N}N$ is

$$g_h = \frac{g_f^2 \mu_f m_f}{8\pi^2} F(\tilde{m}_f^2, m_f^2), \quad (10)$$

where $F(x, y) = (x - y)^{-1} - y \ln(x/y)(x - y)^{-2}$, and μ_f is the $h\tilde{f}_L^* \tilde{f}_R$ coupling. The decay rate of $h \rightarrow N\bar{N}$ is then

$$\Gamma_h = \frac{g_h^2 m_h}{8\pi} \sqrt{1 - 4r^2} (1 - 2r^2), \quad (11)$$

where $r = m_N/m_h$. The correct relic abundance is possible if g_h is very small. Hence N could be FIMP (Feebly Interacting Massive Particle) dark matter [10], and for $r \ll 1$, the right number density is obtained for [12]

$$g_h \sim 10^{-12} r^{-1/2}. \quad (12)$$

Assuming $m_N = 2$ GeV and $\tilde{m}_f/g_f = 10^5$ GeV [13], this would require $\mu_f m_f \sim 6$ GeV². In this scenario, both Z_D and h_D should also be much heavier than h .

Possible Light Z_D and h_D : In the freeze-out scenario, it is possible to have light Z_D and h_D , such that h decays to $h_D h_D$, then h_D decays to $Z_D Z_D$, and Z_D decays to an SM fermion-antifermion pair. This would result in multi-lepton final states of the SM Higgs boson decay,

as pointed out previously [14]. The respective decay rates are

$$\Gamma(h \rightarrow h_D h_D) = \frac{\lambda_3^2 v^2}{16\pi m_h} \sqrt{1 - \frac{4m_{h_D}^2}{m_h^2}}, \quad (13)$$

$$\Gamma(h_D \rightarrow Z_D Z_D) = \frac{9g_D^2}{16\pi m_{h_D}} (m_{h_D}^2 + 2m_{Z_D}^2) \sqrt{1 - \frac{4m_{Z_D}^2}{m_{h_D}^2}}, \quad (14)$$

$$\Gamma(Z_D \rightarrow f \bar{f}) = \frac{g_D^2 a_f^2}{96\pi m_{Z_D} m_f^2} (m_{Z_D}^2 + 2m_f^2)(m_{Z_D}^2 + 8m_f^2) \sqrt{1 - \frac{4m_f^2}{m_{Z_D}^2}}. \quad (15)$$

Concluding Remarks : In the SM, fermions dominate in the form of quarks and leptons. Bosons appear as the necessary vector gauge particles of the SM $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge symmetry, and only one physical scalar Higgs boson remains. It is postulated here that in the dark sector, the opposite occurs, with one neutral Dirac fermion N and many scalars, i.e. the scalar partners of the SM quarks and leptons. They are charged under a dark $U(1)_D$ gauge symmetry, which is broken spontaneously through a neutral scalar by 3 units, resulting in one dark gauge boson Z_D and one dark Higgs boson h_D . Consequently, a residual global dark number is identified as $D = 3B + L - (2j)_{[mod\ 2]}$, where j is the intrinsic spin of the particle. The combined SM and dark sectors are also very suitable for obtaining scotogenic Dirac neutrino masses with the imposition of a softly broken Z_2 symmetry.

The stable dark matter is N . It may be thermally produced and annihilates to Z_D in the freeze-out mechanism, or through the rare decay of the SM Higgs boson h in the freeze-in mechanism. In the former case, h may decay to $h_D h_D$, then h_D to $Z_D Z_D$, then Z_D to an SM fermion-antifermion pair, resulting in multilepton final states.

Acknowledgement : This work was supported in part by the U. S. Department of Energy Grant No. DE-SC0008541.

References

- [1] See for example B.-L. Young, *Front. Phys.* **12**, 121201 (2017). DOI: 10.1007/s11467-016-0583-4.
- [2] E. Ma, *Phys. Rev. Lett.* **115**, 011801 (2015).
- [3] E. Ma, *Phys. Lett.* **B809**, 135736 (2020).
- [4] E. Ma, I. Picek, and B. Radovicic, *Phys. Lett.* **B726**, 744 (2013).
- [5] F. S. Queiroz and W. Shepherd, *Phys. Rev.* **D89**, 095024 (2014).
- [6] E. Ma, *Phys. Rev.* **D73**, 077301 (2006).
- [7] E. Ma, arXiv:2202.13031 [hep-ph].
- [8] E. Ma, *Phys. Lett.* **B817**, 136290 (2021).
- [9] E. Aprile *et al.*, XENON Collaboration, *Phys. Rev. Lett.* **121**, 111302 (2018).
- [10] L. J. Hall, K. Jedamzik, J. March-Russell, and S. M. West, *JHEP* **1003**, 080 (2010).
- [11] E. Ma, *LHEP* **2**, 103 (2019) [DOI:10.31526/LHEP.1.2019.103].
- [12] G. Arcadi and L. Covi, *JCAP* **1308**, 005 (2013).
- [13] E. Ma and V. De Romeri, *Phys. Rev.* **D104**, 055004 (2021).
- [14] C.-F. Chang, E. Ma, and T.-C. Yuan, *JHEP* **1403**, 054 (2014).