

ON THE NON-ARCHIMEDEAN MONGE–AMPÈRE EQUATION IN MIXED CHARACTERISTIC

YANBO FANG, WALTER GUBLER, AND KLAUS KÜNNEMANN

ABSTRACT. Let X be a smooth projective variety over a complete discretely valued field of mixed characteristic. We solve non-archimedean Monge–Ampère equations on X assuming resolution and embedded resolution of singularities and subadditivity for $+$ -test ideals on models of X . We follow the variational approach of Boucksom, Favre, and Jonsson proving the continuity of the plurisubharmonic envelope of a continuous metric on an ample line bundle on X . We replace the use of multiplier ideals in equicharacteristic zero by the use of $+$ -test ideals introduced in mixed characteristic by Hacon, Lamarche, and Schwede.

MSC: Primary 14G22, Secondary 14F18, 32P05, 32U15

CONTENTS

1. Introduction	1
2. Model metrics, curvature forms and psh envelopes	3
3. Test ideals in mixed characteristic	5
4. The subadditivity conjecture	7
5. Continuity of the envelope of the zero function	7
6. Resolution of singularities	8
7. Continuity of the envelope	10
8. The Monge–Ampère equation	10
References	11

1. INTRODUCTION

Let L be an ample line bundle on an n -dimensional projective variety X over a non-archimedean field K . The non-archimedean analogue of the famous *Calabi–Yau problem* asks for a given Radon measure μ on the Berkovich analytification X^{an} with $\mu(X^{\text{an}}) = \deg_L(X)$ whether there exists a continuous semipositive metric $\|\cdot\|$ of L , unique up to scaling, such that

$$(1.1) \quad c_1(L, \|\cdot\|)^{\wedge n} = \mu$$

Date: March 24, 2022.

Y. Fang, W. Gubler and K. Künnemann were supported by the collaborative research center SFB 1085 *Higher Invariants - Interactions between Arithmetic Geometry and Global Analysis* funded by the Deutsche Forschungsgemeinschaft.

using the non-archimedean Monge–Ampère measure introduced by Chambert-Loir on the left. We call (1.1) the *non-archimedean Monge–Ampère equation*. The analogous problem over the complex numbers was solved by Yau for Radon measures given by smooth volume forms within the class of smooth metrics. Uniqueness was shown before by Calabi. Later Kołodziej used pluripotential theory to treat more singular measures and solutions [Kol98].

Yuan and Zhang proved uniqueness up to scaling for solutions of the non-archimedean Monge–Ampère equation [YZ17]. In a groundbreaking work, Boucksom, Favre and Jonsson solved the non-archimedean Monge–Ampère equation over a complete discretely valued field K of *residue characteristic zero* assuming that X is smooth and that the support of the Radon measure μ is contained in the skeleton of an SNC-model (a projective regular model with special fiber having simple normal crossing support) [BFJ15]. They assumed also that K is a completion of the function field of a curve at a closed point. This geometric condition was later removed by Burgos, Jell, Martin and the last two authors of this paper [BGJ⁺20]. Boucksom, Favre, and Jonsson used a variational approach to solve (1.1) which relies crucially on the continuity of the semipositive envelope of a continuous metric of L . They show continuity of this envelope by using multiplier ideals on SNC-models [BFJ16]. In their arguments, Hironaka’s resolution of singularities plays an important role in order to have sufficiently many SNC-models at hand. Apart from the multiplier ideals, it is precisely here where residue characteristic zero is used again.

In equicharacteristic $p > 0$, the following existence result was shown by Jell, Martin and the last two authors of this paper [GJKM19]. Similarly as above, it is assumed that K is a completion of the function field of a curve over a perfect field k of characteristic $p > 0$. It is also assumed that resolution of singularities and embedded resolution of singularities hold in dimension $n + 1$, see §6.1 for precise definitions. Then existence of a solution of (1.1) is shown in [GJKM19] if the support of μ is contained in the skeleton of an SNC-model. The proof is along the same lines as in [BFJ16, BFJ15] replacing multiplier ideals by test ideals. Note that this result is unconditional for $n = 2$ by using resolution of singularities for three-folds proved by Cossart and Piltant.

If K has mixed characteristic, then there are results about the non-archimedean Monge–Ampère problem for varieties like curves, abelian varieties and toric varieties based on their special geometry [Thu05, Liu11, BGJK21].

In this paper, we deal with the Monge–Ampère problem for arbitrary smooth projective varieties over a complete discretely valued field K of mixed characteristic $(0, p)$. For a complete local noetherian domain R of mixed characteristic, a theory of test ideals was introduced by Ma and Schwede based on perfectoid ideas [MS18, MS21]. Using perfectoid methods, prismatic techniques, and a p -adic Riemann–Hilbert correspondence, Bhatt was able in 2020 to show a variant of Kodaira vanishing ‘up to finite covers’ in mixed characteristic [Bha20]. Applications to the minimal model program were given by Bhatt, Ma, Patakfalvi, Schwede, Tucker, Waldron, and Witaszek [BMP⁺21] and independently by Takamatsu and Yoshikawa [TY21]. For projective normal schemes over R , these ideas were extended by Hacon, Lamarche and Schwede. They introduced *+test ideals* and showed global generation results [HLS21]. Their construction,

which replaces spaces of global sections by so-called spaces of $+$ -stable sections from [BMP⁺21, TY21], is recalled in §3. Hacon, Lamarche and Schwede conjecture that subadditivity holds for their $+$ -test ideals [HLS21, Conjecture 8.3]. Observe that subadditivity is well known for multiplier resp. test ideals in the equicharacteristic case. We refer to §4 for details.

The contribution of this paper is to show that $+$ -test ideals allow applications to the non-archimedean Monge–Ampère problem in mixed characteristic. More precisely, we prove the following results.

Theorem 1.1. *Let L be an ample line bundle on a smooth projective variety X over K . If resolution of singularities holds for projective K° -models of X and if the subadditivity conjecture holds on regular projective models of X , then the semipositive envelope of a continuous metric $\|\cdot\|$ of L is continuous.*

This will be shown in Theorem 7.1. This result is the key to apply the variational method of Boucksom–Favre–Jonsson as we will see in Theorem 8.1.

Theorem 1.2. *Let L be an ample line bundle on an n -dimensional smooth projective variety X over K . We assume that the following conjectures hold:*

- (i) *resolution of singularities for projective models of X ;*
- (ii) *embedded resolution of singularities for regular projective models of X ;*
- (iii) *subadditivity for $+$ -test ideals on regular projective models of X .*

Then the non-archimedean Monge–Ampère equation (1.1) is solved by a continuous semipositive metric $\|\cdot\|$ of L , unique up to scaling, if the positive Radon measure μ has support in the skeleton of an SNC-model of X .

2. MODEL METRICS, CURVATURE FORMS AND PSH ENVELOPES

Non-archimedean pluripotential theory in higher dimensions was introduced by Boucksom, Favre and Jonsson [BFJ16]. We recall basic notions of non-archimedean pluripotential theory following [GJKM19, 2.1–2.8].

Let X be a proper variety over a non-archimedean field K . A *model* of X is a proper flat scheme \mathcal{X} over $S := \text{Spec } K^\circ$ together with an isomorphism h from X to the generic fiber \mathcal{X}_η of the S -scheme \mathcal{X} . The special fiber of \mathcal{X} is denoted by \mathcal{X}_s . Let L be a line bundle over X . A *model* of (X, L) is given by a model \mathcal{X} of X and a line bundle \mathcal{L} over \mathcal{X} with an isomorphism of L to $h^*(\mathcal{L}|_{\mathcal{X}_\eta})$.

Let $(\mathcal{X}, \mathcal{L})$ be a model of $(X, L^{\otimes m})$ for some $m \in \mathbb{N}_{>0}$. The *model metric* $\|\cdot\|_{\mathcal{L}}$ on L^{an} over X^{an} is determined by $\|s\|_{\mathcal{L}} := \sqrt[m]{|g|}$ on $U^{\text{an}} \cap \text{red}^{-1}(\mathcal{U}_s)$, where \mathcal{U} is open in \mathcal{X} , s is a section of \mathcal{L} over $U := X \cap \mathcal{U}$, b is a frame of \mathcal{L} over \mathcal{U} , $g \in \mathcal{O}_X(U)$ with $s^{\otimes m} = gb$ over U and $\text{red}_{\mathcal{X}}: X^{\text{an}} \rightarrow \mathcal{X}_s$ is the reduction map. A model metric $\|\cdot\|$ on $\mathcal{O}_X^{\text{an}}$ induces a so-called *model function* $-\log \|1\|: X^{\text{an}} \rightarrow \mathbb{R}$. The \mathbb{Q} -vector space of all model functions on X is denoted by $\mathcal{D}(X)$.

Let \mathfrak{a} be an ideal sheaf on a model \mathcal{X} of X which is supported in the special fibre. The exceptional divisor E of the blowup \mathcal{Y} of \mathcal{X} in \mathfrak{a} defines a model $(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}(E))$ of (X, \mathcal{O}_X) . The associated model function in $\mathcal{D}(X)$ is denoted by $\log |\mathfrak{a}|$.

Given a model \mathcal{X} of X one defines $N^1(\mathcal{X}/S)_{\mathbb{Q}}$ as the quotient of $\text{Pic}(\mathcal{X})_{\mathbb{Q}}$ by the subspace generated by line bundles whose restriction to every closed curve C in the special fiber \mathcal{X}_s has degree zero. Call $\alpha \in N^1(\mathcal{X}/S) := N^1(\mathcal{X}/S)_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{R}$ *nef* if $\alpha \cdot C \geq 0$ for all such curves.

The *space of closed (1,1)-forms on X* is defined as

$$\mathcal{Z}^{1,1}(X) := \mathbb{R} \otimes_{\mathbb{Q}} \varinjlim N^1(\mathcal{X}/S)_{\mathbb{Q}}$$

where the direct limit is taken over all isomorphism classes of models of X . For the model metric induced by a model \mathcal{L} of $L^{\otimes m}$, its *curvature form* $c_1(L, \|\cdot\|_{\mathcal{L}})$ is the image of $\mathcal{L}^{\otimes \frac{1}{m}} \in N^1(\mathcal{X}/S)_{\mathbb{Q}}$ in $\mathcal{Z}^{1,1}(X)$. By construction we have a map $dd^c: \mathcal{D}(X) \rightarrow \mathcal{Z}^{1,1}(X)$ given by $g \mapsto c_1(\mathcal{O}_X, \|\cdot\| \cdot e^{-g})$. Call a closed (1,1)-form $\theta \in \mathcal{Z}^{1,1}(X)$ *semipositive* if it can be represented by a nef element in $N^1(\mathcal{X}/S)$ for some model \mathcal{X} . Denote by $N^1(X)$ the quotient of $\text{Pic}(X)_{\mathbb{R}}$ by numerical equivalence. The map $\{\cdot\}: \mathcal{Z}^{1,1}(X) \rightarrow N^1(X)$ induced by the restriction maps $N^1(\mathcal{X}/S) \rightarrow N^1(X)$ sends a closed (1,1)-form θ to its *de Rham class* $\{\theta\}$. A class in $N^1(X)$ is called *ample* if it is an $\mathbb{R}_{>0}$ -linear combination of classes induced by ample line bundles on X .

We fix $\theta \in \mathcal{Z}^{1,1}(X)$. A model function $\varphi \in \mathcal{D}(X)$ is called *θ -plurisubharmonic* (*θ -psh* for short) if the class $\theta + dd^c\varphi \in \mathcal{Z}^{1,1}(X)$ is semipositive. The space of all psh model functions is denoted by $\text{PSH}_{\mathcal{D}}(X, \theta)$. For a continuous function $u \in \mathcal{C}^0(X^{\text{an}})$, its *θ -psh envelope* $P_{\theta}(u): X^{\text{an}} \rightarrow \mathbb{R} \cup \{-\infty\}$ is the function defined by

$$P_{\theta}(u)(x) := \sup\{\varphi(x) \mid \varphi \in \text{PSH}_{\mathcal{D}}(X, \theta), \varphi \leq u\}, \quad (x \in X^{\text{an}}).$$

For $u \in C^0(X^{\text{an}})$, $v \in \mathcal{D}(X)$ and $t \in \mathbb{R}_{>0}$ we have

$$(2.1) \quad P_{\theta}(u) - v = P_{\theta + dd^c v}(u - v),$$

$$(2.2) \quad P_{t\theta}(t\theta) = tP_{\theta}(u).$$

If the de Rham class $\{\theta\} \in N^1(X)$ is *ample*, then $\text{PSH}_{\mathcal{D}}(X, \theta)$ is non-empty, $P_{\theta}(u)$ takes value in \mathbb{R} , and we have

$$(2.3) \quad \sup_{X^{\text{an}}} |P_{\theta}(u) - P_{\theta}(u')| \leq \sup_{X^{\text{an}}} |u - u'|.$$

for all $u, u' \in C^0(X^{\text{an}})$. For further properties of θ -psh model functions and the θ -psh envelope we refer to [BFJ16] and [GJKM19, Section 2]

For $\theta \in \mathcal{Z}^{1,1}(X)$ that has an ample de Rham class $\{\theta\}$, we consider $\varphi \in \text{PSH}_{\mathcal{D}}(X, \theta)$ and we assume that there exists a normal model \mathcal{X} of X such that θ and φ are induced by line bundles \mathcal{M} and \mathcal{L} on \mathcal{X} . The *Monge–Ampère measure* $\text{MA}_{\theta}(\varphi)$ is the discrete measure

$$\text{MA}_{\theta}(\varphi) := \sum_V \text{length}_{\mathcal{O}_{\mathcal{X}_s, \xi_V}}(\mathcal{O}_{\mathcal{X}_s, \xi_V}) \deg_{\mathcal{M} \otimes \mathcal{L}}(V) \delta_{x_V}$$

on X^{an} where V runs through the irreducible components of \mathcal{X}_s , $x_V \in X^{\text{an}}$ is the unique preimage of the generic point ξ_V of V under $\text{red}_{\mathcal{X}}$, and δ_{x_V} denotes the Dirac probability measure supported in x_V . For generalization of the Monge–Ampère measure to all $\theta \in \mathcal{Z}^{1,1}(X)$ with ample de Rham class $\{\theta\}$ and to more general classes of θ -psh functions, we refer to [BFJ15], [GJKM19] and Section 8.

3. TEST IDEALS IN MIXED CHARACTERISTIC

We recall definitions and some properties from the theory of global and asymptotic $+$ -test ideals developed in [HLS21]. The $+$ -test ideals form a mixed characteristic analogue of the theories of multiplier ideals in characteristic zero and test ideals in positive characteristic. The theory is based on the notion of $+$ -stable sections introduced in [BMP⁺21, TY21].

Let (R, \mathfrak{m}, k) be a complete noetherian local ring of mixed characteristic. Let $p > 0$ denote the characteristic of the residue field k . Let \mathcal{X} be a normal integral scheme which is proper over $S := \text{Spec } R$. The canonical sheaf $\omega_{\mathcal{X}}$ is reflexive ([Sta21, Tag 0AWK] and [Har94, Theorem 1.9]) and we fix a canonical divisor $K_{\mathcal{X}}$ with $\omega_{\mathcal{X}} = \mathcal{O}_{\mathcal{X}}(K_{\mathcal{X}})$ [Sta21, Tag 0EBM].

Definition 3.1. For a reflexive sheaf $\mathcal{M} = \mathcal{O}_{\mathcal{X}}(M)$ associated with a divisor M and an effective \mathbb{Q} -divisor B on \mathcal{X} , the *subspace of $+$ -stable sections* of the adjoint line bundle $\omega_{\mathcal{X}} \otimes \mathcal{M}$ relative to B

$$\mathbf{B}^0(\mathcal{X}, B, \mathcal{O}_{\mathcal{X}}(K_{\mathcal{X}} + M)) \subset H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(K_{\mathcal{X}} + M))$$

is defined by

$$\bigcap_{f: \mathcal{Y} \rightarrow \mathcal{X}} \text{Im} \left(H^0(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}(K_{\mathcal{Y}} + [f^*(M - B)])) \rightarrow H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(K_{\mathcal{X}} + M)) \right),$$

where an algebraic closure $\overline{\kappa(\mathcal{X})}$ of the function field $\kappa(\mathcal{X})$ is fixed, and f runs through the finite surjective morphisms $f: \mathcal{Y} \rightarrow \mathcal{X}$ from a normal integral scheme \mathcal{Y} together with an embedding $\kappa(\mathcal{Y}) \hookrightarrow \overline{\kappa(\mathcal{X})}$ [HLS21, Definition 3.2 and Lemma 3.8].

Remark 3.2. If $M - B$ is \mathbb{Q} -Cartier, then Definition 3.1 holds as well if f runs through the alterations $f: \mathcal{Y} \rightarrow \mathcal{X}$ from a normal integral scheme \mathcal{Y} together with an embedding $\kappa(\mathcal{Y}) \hookrightarrow \overline{\kappa(\mathcal{X})}$ [HLS21, Lemma 3.8].

For the rest of this section, we assume that the scheme \mathcal{X} is regular and projective over S .

Definition 3.3. Take a very ample line bundle \mathcal{H} on \mathcal{X} . For an effective \mathbb{Q} -divisor B on \mathcal{X} and for each $i \in \mathbb{N}$ the subspace

$$(3.1) \quad \mathbf{B}^0(\mathcal{X}, B, \omega_{\mathcal{X}} \otimes \mathcal{H}^i) \subset H^0(\mathcal{X}, \omega_{\mathcal{X}} \otimes \mathcal{H}^i)$$

generates the subsheaf $\mathcal{N}_i \subset \omega_{\mathcal{X}} \otimes \mathcal{H}^i$ which defines

$$(3.2) \quad \mathcal{J}_i := \mathcal{N}_i \otimes \mathcal{H}^{-i} \subset \omega_{\mathcal{X}}.$$

The sequence $(\mathcal{J}_i)_{i \in \mathbb{N}}$ is increasing and becomes stationary. Define the *$+$ -test submodule* $\tau_+(\omega_{\mathcal{X}}, B)$ to be \mathcal{J}_i for $i \gg 0$ [HLS21, Definition 4.3].

Remark 3.4. (i) Definition 3.3 does not depend on the choice of \mathcal{H} [HLS21, Proposition 4.5].

(ii) For $i \gg 0$ we have $H^0(\mathcal{X}, \tau_+(\omega_{\mathcal{X}}, B) \otimes \mathcal{H}^i) = \mathbf{B}^0(\mathcal{X}, B, \omega_{\mathcal{X}} \otimes \mathcal{H}^i)$ [HLS21, Proposition 4.7]. If $B' \geq B$, then $\tau_+(\omega_{\mathcal{X}}, B') \subset \tau_+(\omega_{\mathcal{X}}, B)$ and if F is an effective Cartier divisor, then [HLS21, Lemma 4.8]

$$(3.3) \quad \tau_+(\omega_{\mathcal{X}}, B + F) = \tau_+(\omega_{\mathcal{X}}, B) \otimes \mathcal{O}_{\mathcal{X}}(-F).$$

(iii) Equality (3.3) allows one to define $\tau_+(\omega_{\mathcal{X}}, B)$ for a not necessarily effective \mathbb{Q} -divisor B as $\tau_+(\omega_{\mathcal{X}}, B + F) \otimes \mathcal{O}_{\mathcal{X}}(F)$ where F is a Cartier divisor such that $B + F$ is effective [HLS21, Definition 4.14].

Definition 3.5. For an effective \mathbb{Q} -divisor B on \mathcal{X} , the ideal sheaf

$$(3.4) \quad \tau_+(\mathcal{O}_{\mathcal{X}}, B) := \tau_+(\omega_{\mathcal{X}}, K_{\mathcal{X}} + B)$$

is called the *+test ideal associated with B* [HLS21, Definition 4.15, Lemma 4.18]. Using $F = -K_{\mathcal{X}}$ in Remark 3.4(iii), this is indeed an ideal.

Definition 3.6. Let \mathcal{M} be a line bundle. Let $W \subseteq H^0(\mathcal{X}, \mathcal{M})$ be a linear subspace and $\lambda \in \mathbb{Q}_{>0}$. Define the *+test module of the linear series $|W|$* to be

$$(3.5) \quad \tau_+(\omega_{\mathcal{X}}, \lambda \cdot |W|) := \sum_{D \in |W|} \tau_+(\omega_{\mathcal{X}}, \lambda \cdot D).$$

Thanks to the Noetherian assumption, one can pick a finite number of elements D_1, \dots, D_r in $|W|$ such that $\sum_{i=1}^r \tau_+(\omega_{\mathcal{X}}, \lambda \cdot D_i)$ agrees with (3.5) [HLS21, Definition 7.2].

Remark 3.7. If D has \mathbb{Z} -coefficients and $|D| \neq \emptyset$, then $\tau_+(\omega_{\mathcal{X}}, |D|) = \mathfrak{b}_{|D|} \otimes \omega_{\mathcal{X}}$ [HLS21, Lemma 7.5 (b)] where $\mathfrak{b}_{|D|}$ denotes the base ideal of the linear system $|D|$ [Laz04, Definition 1.1.8].

Definition 3.8. Let D be a \mathbb{Q} -divisor with Itaka dimension $\kappa(D) \geq 0$, so mD is a divisor such that $|mD| \neq \emptyset$ for all large and sufficiently divisible $m \in \mathbb{N}$. Define the *asymptotic +test module* of $|D|$ by

$$(3.6) \quad \tau_+(\omega_{\mathcal{X}}, \lambda \cdot ||D||) := \bigcup_{m>0} \tau_+(\omega_{\mathcal{X}}, \frac{\lambda}{m} \cdot |mD|).$$

and the *asymptotic +test ideal* of $|D|$ by

$$(3.7) \quad \tau_+(\mathcal{O}_{\mathcal{X}}, \lambda \cdot ||D||) := \tau_+(\omega_{\mathcal{X}}, \lambda \cdot ||D||) \otimes \omega_{\mathcal{X}}^{-1}.$$

Thanks to the Noetherian assumption, one has

$$\tau_+(\omega_{\mathcal{X}}, \lambda \cdot ||D||) = \tau_+(\omega_{\mathcal{X}}, \frac{\lambda}{m} \cdot |mD|)$$

for sufficiently divisible $m \in \mathbb{N}$ [HLS21, Lemma 7.4].

Remark 3.9. The asymptotic test module satisfies the following properties [HLS21, Lemma 7.5]:

- (i) If $|D| \neq \emptyset$, then $\tau_+(\omega_{\mathcal{X}}, \lambda \cdot |D|) \subset \tau_+(\omega_{\mathcal{X}}, \lambda \cdot ||D||)$;
- (ii) If $\lambda < \mu$, then $\tau_+(\omega_{\mathcal{X}}, \mu \cdot ||D||) \subset \tau_+(\omega_{\mathcal{X}}, \lambda \cdot ||D||)$; if $D - G$ is semiample, then $\tau_+(\mathcal{O}_{\mathcal{X}}, \lambda \cdot ||G||) \subset \tau_+(\mathcal{O}_{\mathcal{X}}, \lambda \cdot ||D||)$;
- (iii) If $k \in \mathbb{N}_{>0}$, then $\tau_+(\omega_{\mathcal{X}}, \lambda \cdot ||D||) = \tau_+(\omega_{\mathcal{X}}, \frac{\lambda}{k} \cdot ||kD||)$ (homogeneity).

One can use asymptotic +test modules to formulate effective global generation properties.

Theorem 3.10. Let $\mathcal{H} = \mathcal{O}_{\mathcal{X}}(H)$ be a globally generated ample line bundle, D be an effective \mathbb{Q} -divisor, E be a divisor, and $\lambda \in \mathbb{Q}_{>0}$. Let $n = \dim \mathcal{X}_s$. If $E - \lambda D$ is big and semi-ample, then

$$(3.8) \quad \tau_+(\omega_{\mathcal{X}}, \lambda \cdot ||D||) \otimes \mathcal{O}_{\mathcal{X}}(nH + E)$$

is globally generated by a sub linear series of $H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(K_{\mathcal{X}} + nH + E))$ for all $n \geq \dim(\mathcal{X} \otimes_R k)$.

Proof. [HLS21, Lemma 7.5 (e)]. \square

4. THE SUBADDITIVITY CONJECTURE

We recall the subadditivity conjecture [HLS21, Conjecture 8.3] and draw a useful consequence for asymptotic test ideals. Let (R, \mathfrak{m}) be a complete noetherian local ring of mixed characteristic $(0, p)$. Let \mathcal{X} be a regular integral scheme of dimension d which is projective over $S = \text{Spec } R$.

Conjecture 4.1. *Given effective \mathbb{Q} -divisors D and E on \mathcal{X} , we have*

$$\tau_+(\mathcal{O}_{\mathcal{X}}, D + E) \subset \tau_+(\mathcal{O}_{\mathcal{X}}, D) \cdot \tau_+(\mathcal{O}_{\mathcal{X}}, E).$$

Corollary 4.2. *If the subadditivity conjecture holds for \mathcal{X} , then for a divisor D on \mathcal{X} and $q, r \in \mathbb{N}_{>0}$, we have*

$$\tau_+(\mathcal{O}_{\mathcal{X}}, qr \cdot \|D\|) \subset \tau_+(\mathcal{O}_{\mathcal{X}}, r \cdot \|D\|)^q.$$

Proof. By homogeneity of asymptotic $+$ -test ideals [HLS21, Lemma 7.5(d)] (Remark 3.9 (iii)), we may assume $r = 1$. For sufficiently divisible m , we deduce from (3.6) that

$$\tau_+(\mathcal{O}_{\mathcal{X}}, q \cdot \|D\|) = \tau_+\left(\mathcal{O}_{\mathcal{X}}, \frac{q}{m} \cdot |mD|\right).$$

For any effective divisor $D_m \sim mD$, using subadditivity in Conjecture 4.1 for the \mathbb{Q} -divisor $\frac{1}{m}D_m$, we get

$$\tau_+\left(\mathcal{O}_{\mathcal{X}}, \frac{q}{m} \cdot D_m\right) = \tau_+\left(\mathcal{O}_{\mathcal{X}}, q \cdot \frac{1}{m}D_m\right) \subset \tau_+\left(\mathcal{O}_{\mathcal{X}}, \frac{1}{m}D_m\right)^q$$

and hence

$$\tau_+(\mathcal{O}_{\mathcal{X}}, q \cdot \|D\|) \subset \tau_+\left(\mathcal{O}_{\mathcal{X}}, \frac{1}{m} \cdot |mD|\right)^q \subset \tau_+(\mathcal{O}_{\mathcal{X}}, \|D\|)^q$$

proving the claim. \square

5. CONTINUITY OF THE ENVELOPE OF THE ZERO FUNCTION

Let K be a complete discretely valued field of mixed characteristic $(0, p)$. Let X be a n -dimensional smooth projective variety over K and L an ample line bundle on X . Let $(\mathcal{X}, \mathcal{A})$ be a model of (X, L) with \mathcal{X} projective and regular and \mathcal{A} ample. For $m \in \mathbb{N}_{>0}$ let

$$\mathfrak{a}_m := H^0(\mathcal{X}, \mathcal{L}^m) \otimes \mathcal{L}^{-m} \subset \mathcal{O}_{\mathcal{X}}$$

denote the m -th base ideal of \mathcal{L} on \mathcal{X} .

Remark 5.1. Write $\omega_{\mathcal{X}} = \mathcal{O}_{\mathcal{X}}(K_{\mathcal{X}})$ for some divisor $K_{\mathcal{X}}$ on \mathcal{X} . If D, E and H are divisors on \mathcal{X} and $m > 0$ such that $\mathcal{O}_{\mathcal{X}}(H)$ is ample and globally generated and $E - mD$ is big and semiample, then

$$\tau_+(\mathcal{O}_{\mathcal{X}}, m \cdot \|D\|) \otimes \mathcal{O}_{\mathcal{X}}(E + K_{\mathcal{X}} + nH)$$

is globally generated. This is just a reformulation of Theorem 3.10.

In the approach of Boucksom–Favre–Jonsson [BFJ16, BFJ15] to pluripotential theory on X^{an} , a model \mathcal{L} of L on the model \mathcal{X} induces a curvature form θ on X^{an} and we denote by $P_\theta(f)$ the θ -psh envelope of a continuous real function f on X^{an} , see [GJKM19, 2.4–2.6] for details. We follow the strategy from [BFJ16] and [GJKM19] to show continuity of the envelope of the zero function $P_\theta(0)$.

Theorem 5.2. *Let θ be the curvature form on X^{an} induced by a model \mathcal{L} on \mathcal{X} of the ample line bundle L on X . If the Subadditivity Conjecture 4.1 holds on \mathcal{X} , then $(m^{-1} \log |\mathfrak{a}_m|)_{m \in \mathbb{N}_{>0}}$ is a sequence of θ -psh model functions which converges uniformly on X^{an} to the envelope $P_\theta(0)$ of the zero function.*

Proof. Consider the graded sequence $(\mathfrak{a}_m)_{m>0}$ of base ideals

$$\mathfrak{a}_m = H^0(\mathcal{X}, \mathcal{L}^m) \otimes \mathcal{L}^{-m} \subset \mathcal{O}_{\mathcal{X}}$$

associated with \mathcal{L} . Write $\mathcal{L} = \mathcal{O}_{\mathcal{X}}(D)$ for some divisor D on \mathcal{X} . Let

$$\mathfrak{b}_m := \tau_+(\mathcal{O}_{\mathcal{X}}, m \cdot \|D\|) \subset \mathcal{O}_{\mathcal{X}}$$

denote the associated asymptotic $+$ -test ideal of exponent m in mixed characteristic. These ideals have the following properties:

- (a) We have $\mathfrak{a}_m \subset \mathfrak{b}_m$ for all $m \in \mathbb{N}_{>0}$.
- (b) We have $\mathfrak{b}_{ml} \subset \mathfrak{b}_m^l$ for all $l, m \in \mathbb{N}_{>0}$.
- (c) There exists $m_0 \geq 0$ such that $\mathfrak{b}_m \otimes \mathcal{A}^{\otimes m_0} \otimes \mathcal{L}^{\otimes m}$ is globally generated for all $m > 0$.

Indeed these properties can be seen as follows: Property (a) follows from Remarks 3.7 and 3.9(i), and Property (b) follows from Corollary 4.2. Property (c) is shown as follows. Choose a divisor H on \mathcal{X} such that $\mathcal{O}_{\mathcal{X}}(H)$ is ample and globally generated. We have $n = \dim \mathcal{X}_s$. As \mathcal{A} is ample, we can choose $m_0 \in \mathbb{N}$ such that the line bundle

$$(5.1) \quad \mathcal{A}^{\otimes m_0} \otimes \mathcal{O}_{\mathcal{X}}(-K_{\mathcal{X}} - (n+1)H)$$

is globally generated. Given $m \in \mathbb{N}$ we apply Remark 5.1 to $E := mD + H$. Then $E - mD = H$ is big and semiample and

$$(5.2) \quad \mathfrak{b}_m \otimes \mathcal{O}_{\mathcal{X}}(mD + H + K_{\mathcal{X}} + nH)$$

is globally generated. Taking the tensor product of (5.1) and (5.2) we see that $\mathfrak{b}_m \otimes \mathcal{A}^{\otimes m_0} \otimes \mathcal{L}^{\otimes m}$ is globally generated.

Finally (a), (b) and (c) imply our claim by the strategy of the proof of [BFJ16, Thm. 8.5]. Step 1 of *loc. cit.* holds not only on quasi-monomial points of X^{an} , but pointwise on the whole X^{an} using [GJKM19, Proposition 2.10]. Then Step 2 of *loc. cit.* works again in our setting using (a) – (c) above. The only difference is that all inequalities hold immediately on X^{an} and not only on the quasi-monomial points of X^{an} . \square

6. RESOLUTION OF SINGULARITIES

Let K° be a complete discrete valuation ring with field of fractions K and $S := \text{Spec } K^\circ$.

Definition 6.1. We say that *resolution of singularities holds over S in dimension n* if for every projective flat S -scheme \mathcal{X} of dimension n with regular generic fibre there exists a regular S -scheme \mathcal{X}' and a projective S -morphism $\mathcal{X}' \rightarrow \mathcal{X}$ which induces an isomorphism on generic fibres over S .

Remark 6.2. Let X be a smooth projective variety over K of dimension n . If one chooses an immersion $X \rightarrow \mathbb{P}_K^n$, then the scheme theoretic image \mathcal{X} of X in $\mathbb{P}_{K^\circ}^n$ defines a projective model of X over S . If resolution of singularities holds over S in dimension $n + 1$, then there is a regular projective model \mathcal{X}' of X over S .

Resolution of singularities in dimension 2 holds [Art86, Theorem (1.1)].

Cossart and Piltant have shown that resolution of singularities holds over S in dimension 3 up to the projectivity of the morphism $\mathcal{X}' \rightarrow \mathcal{X}$ [CP19, Theorem 1.1]. It is only shown in *loc. cit.* that this morphism is locally projective.

It is essential to show that projective models are dominated by SNC-models. In order to prove this we are going to use the following assumption.

Definition 6.3. We say that *embedded resolution of singularities holds over S in dimension n* if for every regular projective flat S -scheme \mathcal{X} of dimension n and every proper closed subset Z of \mathcal{X} , there is a projective morphism of S -schemes $\pi: \mathcal{X}' \rightarrow \mathcal{X}$ of such that the set $\pi^{-1}(Z)$ is the support of a normal crossing divisor and such that π is an isomorphism over $\mathcal{X} \setminus Z$.

Lemma 6.4. *Let \mathcal{X} be an integral projective S -scheme with n -dimensional regular generic fiber X . We assume that resolution of singularities holds over S in dimension $n + 1$. Then for any ample line bundle L on X , there exists $m \in \mathbb{N}_{>0}$ and an ample extension \mathcal{L}' of $L^{\otimes m}$ to a regular S -model \mathcal{X}' of X with a projective morphism $\mathcal{X}' \rightarrow \mathcal{X}$ over S extending the identity on X .*

Proof. The arguments are the same as for [GJKM19, Lemma 7.5]. In a first step, we start with a projective model \mathcal{Y} of X such that L extends to an ample line bundle \mathcal{H} on \mathcal{Y} , possibly replacing L by a positive tensor power. By a result of Lütkebohmert [Lüt93, Lemma 2.2], there is a blowing up $\pi: \mathcal{Z} \rightarrow \mathcal{Y}$ in an ideal sheaf supported in the special fiber of \mathcal{Y} such that the identity on X extends to a morphism $\mathcal{Z} \rightarrow \mathcal{X}$. A property of blowing ups [Har77, Proposition II.7.13] shows that $\pi^*(\mathcal{H}^{\otimes \ell}) \otimes \mathcal{O}_{\mathcal{Z}/\mathcal{Y}}(1)$ is an ample line bundle on \mathcal{Z} with generic fiber $L^{\otimes \ell}$ for sufficiently large ℓ . Replacing \mathcal{X} by \mathcal{Z} and L by $L^{\otimes \ell}$, we conclude that we may assume that L extends to an ample line bundle \mathcal{H} on \mathcal{X} .

By a result of Pépin [Pép13, Thm. 3.1], there is a blowing-up morphism $\pi': \mathcal{Z}' \rightarrow \mathcal{X}$ centered in the special fiber of \mathcal{X} such that \mathcal{Z}' is semi-factorial. So similarly as in the first step, replacing \mathcal{X} by \mathcal{Z}' and L by a positive tensor power, we may assume that L extends to an ample line bundle \mathcal{H} on a semi-factorial projective model \mathcal{X} . This is the conclusion of the second step.

Then we apply resolution of singularities to \mathcal{X} to get a projective morphism $\pi: \mathcal{X}' \rightarrow \mathcal{X}$ which is an isomorphism over the regular locus of \mathcal{X}' . As the generic fiber X of \mathcal{X}' is in this regular locus, we may identify the generic fiber of \mathcal{X}' with X . Since π is projective, there is an $\ell > 0$ such that $\mathcal{L}' :=$

$\pi^*(\mathcal{H}^{\otimes \ell}) \otimes \mathcal{O}_{\mathcal{X}'/\mathcal{X}}(1)$ is an ample line bundle on \mathcal{X}' . However, we do not know if π is a blow up in an ideal supported in the special fiber and hence the restriction F of $\mathcal{O}_{\mathcal{X}'/\mathcal{X}}(1)$ to the generic fiber X of \mathcal{X}' might be non-trivial. Since \mathcal{X} is semi-factorial, F extends to a line bundle \mathcal{F} on \mathcal{X} and we can replace $\mathcal{O}_{\mathcal{X}'/\mathcal{X}}(1)$ by $\mathcal{O}_{\mathcal{X}'/\mathcal{X}}(1) \otimes \varphi^*(\mathcal{F}^{-1})$. Then \mathcal{L}' is a model of $L^{\otimes \ell}$ proving the claim. \square

7. CONTINUITY OF THE ENVELOPE

Let K be a complete discretely valued field of mixed characteristic $(0, p)$ and $S := \text{Spec } K^\circ$. Let L be an ample line bundle on a regular projective variety X of dimension n over K . Let θ be a closed $(1, 1)$ -form on X with ample de Rham class $\{\theta\}$ induced by a model \mathcal{L} of L on a proper model \mathcal{X} of X .

Theorem 7.1. *Assume that Conjecture 6.1 holds over S in dimension $n + 1$ and that the Subadditivity Conjecture 4.1 holds on all regular projective models of X . If $u \in C^0(X^{\text{an}})$, then $P_\theta(u)$ is a uniform limit of θ -psh model functions and thus $P_\theta(u)$ is continuous on X^{an} .*

Proof. Using that u is a uniform limit of model functions on X , we may assume that u is itself a model function by [GJKM19, Proposition 2.9(v)]. The model function u is defined by a vertical \mathbb{Q} -divisor on a proper model \mathcal{X} of X over S . By [GJKM19, Proposition 2.9(7)], we may replace (θ, u) for the proof by $(m\theta, mu)$ for some $m \in \mathbb{N}$. Hence we may assume without loss of generality that u is actually defined by a vertical divisor on \mathcal{X} . If we apply [Lüt93, Lemma 2.2] to \mathcal{X} and a projective model of X as in Remark 6.2, then \mathcal{X} is dominated by a projective model of X . Hence we can assume without loss of generality that \mathcal{X} is a projective model. Then we choose a resolution of singularities $\mathcal{X}' \rightarrow \mathcal{X}$ as in Lemma 6.4 such that some power $L^{\otimes m}$ of L extends to an ample line bundle \mathcal{A} on the regular projective model \mathcal{X}' of X . As before we may assume that $m = 1$. By [GJKM19, Proposition 2.9(4)], we get

$$P_\theta(u) = P_{\theta+dd^c u}(0) + u.$$

By construction the class $\theta + dd^c u$ is induced by a line bundle \mathcal{L} on \mathcal{X}' whose restriction to X is isomorphic to L and Theorem 5.2 shows that $P_{\theta+dd^c u}(0)$ is a uniform limit of $(\theta + dd^c u)$ -psh model functions. This finishes our proof. \square

8. THE MONGE–AMPÈRE EQUATION

Let K be a complete discretely valued field with valuation ring R of mixed characteristic $(0, p)$. In this section, we consider a projective regular variety X over K of dimension n . We assume that resolution of singularities and embedded resolution of singularities hold in dimension $n + 1$ over R . It follows that every projective K° -model of X is dominated by a projective SNC-model. By Lemma 6.4, any ample line bundle on X extends to an ample line bundle on this SNC model after possibly passing to a positive tensor power. As explained in [GJKM19, Section 9], these assumptions are enough to set up a pluripotential theory for θ -psh functions with respect to a closed $(1, 1)$ -form θ on X^{an} with ample de Rham class $\{\theta\}$. All the results of [BFJ16, Sections 1–7] hold. If we assume continuity of the envelope for ample line bundles on X , then monotone

regularization [BFJ16, Theorem 8.7] holds as well in our setting which is crucial to extend the Monge–Ampère measure $\text{MA}_\theta(\varphi)$ from θ -psh model functions to bounded θ -psh functions on X^{an} . Then the results from [BFJ16, Sections 4–6] hold in our setting by the same arguments.

Theorem 8.1. *Let $n \in \mathbb{N}$ such that resolution of singularities and embedded resolution of singularities hold in dimension $n + 1$ over R . Let X be a regular projective variety over K of dimension n and let θ be a closed $(1, 1)$ -form on X^{an} with ample de Rham class $\{\theta\}$. We assume that the subadditivity conjecture 4.1 holds for all projective regular models of X . We consider a positive Radon measure μ on X^{an} of total mass $\{\theta\}^n$ which is supported on the skeleton of a projective SNC-model of X . Then there is a continuous θ -psh function φ on X^{an} such that $\text{MA}_\theta(\varphi) = \mu$ and φ is unique up to additive constants.*

Proof. Uniqueness follows from a result of Yuan and Zhang, see [BFJ15, §8.1]. To prove existence of a θ -psh solution φ , the variational method of Boucksom, Favre, and Jonsson is used. Continuity of the envelope for ample line bundles on X holds by Theorem 7.1. By [BGJ⁺20, Theorems 6.3.2, 6.3.3], we conclude that θ satisfies the orthogonality property. Then existence of a continuous solutions follows from [BFJ15, Theorem 8.1]. \square

Remark 8.2. If L is an ample line bundle on X and if θ is induced by a model $(\mathcal{X}, \mathcal{L})$ of (X, L) , then there is a bijective correspondence between θ -psh functions and semipositive metrics of L , see [BFJ15, §2.6]. It follows that Theorem 7.1 implies Theorem 1.1 and that Theorem 8.1 implies Theorem 1.2.

ACKNOWLEDGEMENTS

We are grateful to Bhargav Bhatt, Huayi Chen and Mattias Jonsson for remarks about previous versions and to Karl Schwede for discussions about the subadditivity conjecture for $+$ -test ideals.

REFERENCES

- [Art86] Michael Artin. Lipman’s proof of resolution of singularities for surfaces. In *Arithmetic geometry (Storrs, Conn., 1984)*, pages 267–287. Springer, New York, 1986.
- [Bha20] Bhargav Bhatt. Cohen–Macaulayness of absolute integral closures. [arXiv:2008.08070](https://arxiv.org/abs/2008.08070), 2020.
- [BMP⁺21] Bhargav Bhatt, Linqun Ma, Zsolt Patakfalvi, Karl Schwede, Kevin Tucker, Joe Waldron, and Jakub Witaszek. Globally $+$ -regular varieties and the minimal model program for threefolds in mixed characteristic. [arXiv:2012.15801v2](https://arxiv.org/abs/2012.15801v2), 2021.
- [BFJ15] Sébastien Boucksom, Charles Favre, and Mattias Jonsson. Solution to a non-Archimedean Monge–Ampère equation. *J. Amer. Math. Soc.*, 28(3):617–667, 2015.
- [BFJ16] Sébastien Boucksom, Charles Favre, and Mattias Jonsson. Singular semipositive metrics in non-Archimedean geometry. *J. Algebraic Geom.*, 25(1):77–139, 2016.
- [BGJK21] José Ignacio Burgos Gil, Walter Gubler, Philipp Jell, and Klaus Künnemann. Pluripotential theory for tropical toric varieties and non-archimedean Monge–Ampère equations. [arXiv:2102.07392](https://arxiv.org/abs/2102.07392), 2021.

- [BGJ⁺20] José Ignacio Burgos Gil, Walter Gubler, Philipp Jell, Klaus Künnemann, and Florent Martin. Differentiability of non-archimedean volumes and non-archimedean Monge–Ampère equations (with an appendix by Robert Lazarsfeld). *Algebraic Geometry*, 7(2):113–152, 2020.
- [CP19] Vincent Cossart and Olivier Piltant. Resolution of singularities of arithmetical threefolds. *J. Algebra*, 529:268–535, 2019.
- [GJKM19] Walter Gubler, Philipp Jell, Klaus Künnemann, and Florent Martin. Continuity of plurisubharmonic envelopes in non-archimedean geometry and test ideals. *Ann. Inst. Fourier (Grenoble)*, 69(5):2331–2376, 2019. With an appendix by José Ignacio Burgos Gil and Martín Sombra.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [Har94] Robin Hartshorne. Generalized divisors on Gorenstein schemes. In *Proceedings of Conference on Algebraic Geometry and Ring Theory in honor of Michael Artin, Part III (Antwerp, 1992)*, volume 8, pages 287–339, 1994.
- [HLS21] Christopher Hacon, Alicia Lamarche, and Karl Schwede. Global generation of test ideals in mixed characteristic and applications. [arXiv:2106.14329](https://arxiv.org/abs/2106.14329), 2021.
- [Kol98] Sławomir Kołodziej. The complex Monge–Ampère equation. *Acta Math.*, 180:69–117, 1998.
- [Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. I. Classical setting: line bundles and linear series*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer-Verlag, Berlin, 2004.
- [Liu11] Yifeng Liu. A non-Archimedean analogue of the Calabi-Yau theorem for totally degenerate abelian varieties. *J. Differential Geom.*, 89(1):87–110, 2011.
- [Lüt93] Werner Lütkebohmert. On compactification of schemes. *Manuscripta Math.*, 80(1):95–111, 1993.
- [MS18] Linquan Ma and Karl Schwede. Perfectoid multiplier/test ideals in regular rings and bounds on symbolic powers. *Invent. Math.*, 214(2):913–955, 2018.
- [MS21] Linquan Ma and Karl Schwede. Singularities in mixed characteristic via perfectoid big Cohen-Macaulay algebras. *Duke Math. J.*, 170(13):2815–2890, 2021.
- [Pép13] Cédric Pépin. Modèles semi-factoriels et modèles de Néron. *Math. Ann.*, 355(1):147–185, 2013.
- [Sta21] The Stacks Project Authors. *Stacks Project*. <http://stacks.math.columbia.edu>, 2021.
- [TY21] Teppei Takamatsu and Shou Yoshikawa. Minimal model program for semi-stable threefolds in mixed characteristic. [arXiv:2012.07324](https://arxiv.org/abs/2012.07324), 2021.
- [Thu05] Amaury Thuillier. *Théorie du potentiel sur les courbes en géométrie analytique non archimédienne. Applications à la théorie d’Arakelov*. PhD thesis, Université de Rennes, 2005. <https://tel.archives-ouvertes.fr/tel-00010990>.
- [YZ17] Xinyi Yuan and Shou-Wu Zhang. The arithmetic Hodge index theorem for adelic line bundles. *Math. Ann.*, 367(3-4):1123–1171, 2017.

Y. FANG, MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY
Email address: yanbo.fang@mathematik.uni-regensburg.de

W. GUBLER, MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY
Email address: walter.gubler@mathematik.uni-regensburg.de

K. KÜNNEMANN, MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY
Email address: klaus.kuennemann@mathematik.uni-regensburg.de