The orthogonal unit group of the trivial source ring^{*}

Robert Boltje

Department of Mathematics University of California Santa Cruz, CA 95064 USA boltje@ucsc.edu Department of Mathematics William & Mary Williamsburg, VA 23185 USA

Rob Carman

wrcarman@wm.edu

October 24, 2022

Abstract

Let G be a finite group, p a prime, and (K, \mathcal{O}, F) a p-modular system. We prove that the trivial source ring of $\mathcal{O}G$ is isomorphic to the ring of coherent G-stable tuples (χ_P) , where χ_P is a virtual character of $K[N_G(P)/P]$, P runs through all p-subgroups of G, and the coherence condition is the equality of certain character values. We use this result to describe the group of orthogonal units of the trivial source ring as the product of the unit group of the Burnside ring of the fusion system of G with the group of coherent G-stable tuples (φ_P) of homomorphisms $N_G(P)/P \to F^{\times}$. The orthogonal unit group of the trivial source ring of $\mathcal{O}G$ is of interest, since it embeds into the group of p-permutation autoequivalences of $\mathcal{O}G$.

1 Introduction

Throughout this paper we fix a finite group G, a prime p, and a p-modular system (K, \mathcal{O}, F) . In other words, \mathcal{O} is a complete discrete valuation ring with residue field F of characteristic p and field of fractions K of characteristic zero.

The class of *p*-permutation $\mathcal{O}G$ -modules and its Grothendieck ring $T(\mathcal{O}G)$, the trivial source ring of $\mathcal{O}G$, play an important role in the modular representation theory of finite groups. For instance, several equivalence relations between block algebras (as for instance splendid Rickard equivalences, see [Ri96], and *p*-permutation equivalences, see [BoXu08] and [BoPe20]) are defined in terms of *p*-permutation bimodules or their chain complexes. They give satisfactory explanations for the phenomenon of *isotypies*, introduced by Broué,

^{*}MR Subject Classification: 20C20, 20C15, 19A22. Keywords: Trivial source ring, *p*-permutation modules, orthogonal units, *p*-permutation equivalences, species

see [Bro90]. Finding new ways to describe *p*-permutation modules or the ring $T(\mathcal{O}G)$ is therefore of interest. In this paper we give a description of $T(\mathcal{O}G)$ in terms of tuples of *coherent characters*. More precisely, we consider a ring homomorphism

$$\beta_G \colon T(\mathcal{O}G) \to \left(\prod_{P \in \mathscr{S}_p(G)} R(K[N_G(P)/P])\right)^G,$$

where $S_p(G)$ denotes the set of *p*-subgroups of *G*, $R(K[N_G(P)/P])$ denotes the ring of characters of $K[N_G(P)/P]$ -modules, and the exponent *G* stands for taking *G*-fixed points under the natural *G*-conjugation action on the product. See Section 3 for a precise definition of the homomorphism β_G , using the Brauer construction for each $P \in S_p(G)$.

Theorem A The ring homomorphism β_G is injective and its image consists of those tuples $(\chi_P) \in (\prod_{P \in S_p(G)} R(K[N_G(P)/P]))^G$ which satisfy the following coherence condition:

(C) For each $P \in S_p(G)$ and each $x \in N_G(P)$ one has $\chi_P(xP) = \chi_{P\langle x_p \rangle}(xP\langle x_p \rangle)$.

Here, x_p denotes the *p*-part of the element $x \in G$ and $\langle x_p \rangle$ the subgroup generated by x_p . We call *G*-fixed tuples (χ_P) of virtual characters satisfying the condition (C) coherent tuples. Theorem A is proved in Section 3. It has also a block-wise version, see Corollary 3.3 and a version related to limiting the set of vertices, see Corollary 3.5. These versions will be crucial when analyzing *p*-permutation equivalences between blocks in terms of the associated coherent character tuples and studying the precise relationship between isotypies and *p*-permutation equivalences, as anticipated in a future paper.

We will present several applications of the above theorem. A crucial tool in these applications is Theorem B. Although it follows immediately from [BaCa20, Theorem 7.1] (see Theorem 4.4), we think it is worth stating it in the introduction. Let S denote a Sylow p-subgroup of G and let $\mathcal{F} := \mathcal{F}_S(G)$ denote the resulting fusion system on S. The Burnside ring $B(\mathcal{F})$ of \mathcal{F} is a subring of the Burnside ring B(S), see 4.3 for a precise definition. One has a natural ring homomorphism $\lambda_G \colon B(G) \to T(\mathcal{O}G)$ induced by mapping a finite G-set X to its \mathcal{O} -linearization, the permutation $\mathcal{O}G$ -module $\mathcal{O}X$. Note that Green's theory of vertices and sources and Green's indecomposability theorem imply that $\lambda_S \colon B(S) \to T(\mathcal{O}S)$ is a ring isomorphism, since S is a p-group. It is an easy verification that the images of the restriction maps

 $\operatorname{res}_{S}^{G} : B(G) \to B(S) \text{ and } \lambda_{S}^{-1} \circ \operatorname{res}_{S}^{G} : T(\mathcal{O}G) \to T(\mathcal{O}S) \xrightarrow{\sim} B(S)$

are contained in $B(\mathcal{F})$. Theorem B states that the resulting restriction maps $B(G) \to B(\mathcal{F})$ and $T(\mathcal{O}G) \to B(\mathcal{F})$ are split surjective.

Theorem B Let S be a Sylow p-subgroup of G and let $\mathcal{F} := \mathcal{F}_S(G)$ denote the resulting fusion system on S. There exists a ring homomorphism $t_S^G \colon B(\mathcal{F}) \to B(G)$ such that the compositions

$$\operatorname{res}_{S}^{G} \circ t_{S}^{G} \colon B(\mathcal{F}) \to B(G) \to B(\mathcal{F}) \tag{1}$$

and

$$\lambda_S^{-1} \circ \operatorname{res}_S^G \circ \lambda_G \circ t_S^G \colon B(\mathcal{F}) \to B(G) \to T(\mathcal{O}G) \to T(\mathcal{O}S) \xrightarrow{\sim} B(S)$$
(2)

are equal to the identity map on $B(\mathcal{F})$. In particular, one obtains decompositions

$$B(G) = \ker(\operatorname{res}_S^G) \oplus t_S^G(B(\mathcal{F})) \quad and \quad T(\mathcal{O}G) = \ker(\lambda_S^{-1} \circ \operatorname{res}_S^G) \oplus (\lambda_G \circ t_S^G)(B(\mathcal{F}))$$

into ideals ker(res^G_S) of B(G) and ker($\lambda_S^{-1} \circ \text{res}_S^G$) of $T(\mathcal{O}G)$ and subrings isomorphic to $B(\mathcal{F})$.

Taking the \mathcal{O} -dual $M^{\circ} := \operatorname{Hom}_{\mathcal{O}}(M, \mathcal{O})$ of a *p*-permutation $\mathcal{O}G$ -module induces a ring automorphism $-^{\circ}: T(\mathcal{O}G) \to T(\mathcal{O}G), [M] \mapsto [M^{\circ}]$, of order 2. We define the *orthogonal* unit group of $T(\mathcal{O}G)$ as

$$\mathcal{O}(T(\mathcal{O}G)) := \{ u \in T(\mathcal{O}G) \mid u \cdot u^{\circ} = 1 \}.$$

Thus, $O(T(\mathcal{O}G))$ is a subgroup of the unit group of the commutative ring $T(\mathcal{O}G)$. It turns out that $O(T(\mathcal{O}G))$ consists precisely of the units of $T(\mathcal{O}G)$ of finite order, see Remark 4.1(b). Orthogonal units of $T(\mathcal{O}G)$ yield *p*-permutation autoequivalences of $\mathcal{O}G$, see [BoPe20] and see Remark 4.1(a), and are therefore of particular interest.

Theorem C Let S be a Sylow p-subgroup of G and let $\mathcal{F} := \mathcal{F}_S(G)$ be the associated fusion system on S. One has a direct product decomposition

$$O(T(\mathcal{O}G)) \cong B(\mathcal{F})^{\times} \times \left(\prod_{P \in \mathscr{S}_p(G)} \operatorname{Hom}(N_G(P)/P, F^{\times})\right)'$$
(3)

where the second factor is defined as the set of all tuples

$$(\varphi_P) \in \left(\prod_{P \in \mathcal{S}_p(G)} \operatorname{Hom}(N_G(P)/P, F^{\times})\right)^G$$

satisfying

$$\varphi_P(xP) = \varphi_{P\langle x_p \rangle}(xP\langle x_p \rangle) \tag{4}$$

for all $P \in S_p(G)$ and $x \in N_G(P)$.

Theorem C is proved in Section 4. It uses Theorems A and B. In Proposition 4.6 we prove a further direct product decomposition of the second factor in (3). It is well-known that if pis odd then $B(S)^{\times} = \{\pm 1\}$ and consequently $B(\mathcal{F})^{\times} = \{\pm 1\}$. This follows for instance from [Yo90, Proposition 6.5]. See also [Bc10a, Lemma 11.2.5] for a direct elementary proof. If $p = 2, B(S)^{\times}$ has been described explicitly by Bouc (see [Bc10a, Section 11.2]) and $B(\mathcal{F})^{\times}$ has been described in some situations by Barsotti and Carman in [BaCa20].

In Section 5 we prove a criterion for a potential tuple of species values to come from an orthogonal unit of $T(\mathcal{O}G)$. A similar result was proved by Yoshida in [Yo90, Proposition 6.5]

for units of the Burnside ring. In Section 6 we give more explicit descriptions of $O(T(\mathcal{O}G))$ for some special cases of G.

Notation Apart from the notations already introduced, we will use the following standard notations:

For any ring A, we denote by A^{\times} the unit group of A, by Z(A) the center of A, and by ${}_{A}$ mod the category of finitely generated left A-modules. For $M, N \in {}_{A}$ mod we write $M \mid N$ if M is isomorphic to a direct summand of N.

For any positive integer n we denote by n_p the largest power of p dividing n and set $n_{p'} = n/n_p$. For any group element x we denote by x_p its p-part and by $x_{p'}$ its p'-part.

By $a \mapsto \bar{a}$ we denote the natural epimorphisms $\mathcal{O} \to F$ and $\mathcal{O}G \to FG$.

We set ${}^{g_a} := gag^{-1}$ for any $g \in G$ and $a \in G$ or $a \in kG$ for a commutative ring k. For subgroups H_1 and H_2 of G we write $H_1 \leq_G H_2$ if $H_1 \leq {}^{g}H_2$ for some $g \in G$.

For any group H, any commutative ring k and any $\varphi \colon H \to k^{\times}$, we denote by k_{φ} , the kH-module whose underlying k-module is equal to k and on which H acts via multiplication by $\varphi(h)$. If φ is the trivial homomorphism we also write k_H for the trivial kH-module.

2 Preliminaries

2.1 (a) Recall that a *p*-permutation $\mathcal{O}G$ -module is a finitely generated $\mathcal{O}G$ -module M with the property that $\operatorname{Res}_{P}^{G}(M)$ is a permutation $\mathcal{O}P$ -module for every *p*-subgroup P of G. Equivalently, every indecomposable direct summand of M has trivial source. Another equivalent condition is that M is isomorphic to a direct summand of a finitely generated permutation $\mathcal{O}G$ -module. The class of *p*-permutation modules over \mathcal{O} is closed under restriction, induction, \mathcal{O} -duals, tensor products $-\otimes_{\mathcal{O}} -$, direct sums, and taking direct summands and Green correspondents.

(b) Let $e \in Z(\mathcal{O}G)$ be an idempotent. We denote the category of *p*-permutation $\mathcal{O}Ge$ modules by $_{\mathcal{O}Ge}$ triv and denote by $T(\mathcal{O}Ge)$ the Grothendieck group of the monoid of isomorphism classes of *p*-permutation $\mathcal{O}Ge$ -modules with respect to the direct sum operation. By the Krull-Schmidt Theorem, $T(\mathcal{O}Ge)$ has a standard \mathbb{Z} -basis consisting of elements [M], where M runs through a set of representatives of the isomorphism classes of indecomposable $\mathcal{O}Ge$ -modules with trivial source. When e = 1, $T(\mathcal{O}G)$ is a commutative ring with multiplication induced by $-\otimes_{\mathcal{O}} -$ and it is also known as the *trivial source ring* of G over \mathcal{O} . We always consider $T(\mathcal{O}Ge)$ as a subgroup of $T(\mathcal{O}G)$.

(c) Similarly, one defines p-permutation $FG\bar{e}$ -modules and the associated trivial source group $T(FG\bar{e})$. The above statements hold as well over F. The functor $F \otimes_{\mathcal{O}} -$ induces a ring isomorphism $T(\mathcal{O}G) \to T(FG)$ which sends the standard basis of $T(\mathcal{O}Ge)$ bijectively onto the standard basis of $T(FG\bar{e})$ and preserves vertices (see [Bro85]). By $Pr(\mathcal{O}Ge)$ we denote the subgroup of $T(\mathcal{O}Ge)$ generated by the standard basis elements [M], where M is a projective indecomposable $\mathcal{O}Ge$ -module. Similarly we define $Pr(FG\bar{e})$.

(d) Note that one has a commutative diagram

$$\begin{array}{rcl} Pr(\mathcal{O}Ge) &\subseteq & T(\mathcal{O}Ge) \stackrel{\kappa_G}{\longrightarrow} R(KGe) \\ & & & & \downarrow \wr & & \downarrow d_G \\ Pr(FG\bar{e}) &\subseteq & T(FG\bar{e}) \stackrel{\eta_G}{\longrightarrow} R(FG\bar{e}) \,, \end{array}$$

where R(KGe) denotes the group of virtual characters of KGe-modules, R(FGe) denotes the group of virtual Brauer characters of FGe-modules, κ_G is induced by the functor $K \otimes_{\mathcal{O}}$ $-: {}_{\mathcal{O}Ge}\mathsf{triv} \to {}_{KGe}\mathsf{mod}, \eta_G$ is the canonical map, the left two vertical maps are induced by the functor $F \otimes_{\mathcal{O}} -$, and the right vertical map is the decomposition map. By [Se78, Théorèmes 34 et 36], the map $\kappa_G: Pr(\mathcal{O}Ge) \to R(KGe)$ is injective and its image consists precisely of those virtual characters in R(KGe) which vanish on *p*-singular elements.

2.2 Lemma Suppose that P is a normal p-subgroup of a finite group H. Let M be an indecomposable p-permutation $\mathcal{O}H$ -module with vertex Q. The following are equivalent:

- (i) P acts trivially on M, i.e., $\operatorname{Res}_P^H(M) \cong \mathcal{O}_P \oplus \cdots \oplus \mathcal{O}_P$.
- (ii) $\mathcal{O}_P \mid \operatorname{Res}_P^H(M)$.
- (iii) $P \leq Q$.

Proof Obviously, (i) implies (ii). If (ii) holds, then $\mathcal{O}_P | \operatorname{Res}_P^H(\operatorname{Ind}_Q^H(\mathcal{O}_Q))$. The Mackey formula and the fact that \mathcal{O}_P has vertex P now implies that (iii) holds. Finally, if (iii) holds then $\operatorname{Res}_P^H(M) | \operatorname{Res}_P^H(\operatorname{Ind}_Q^H(\mathcal{O}_Q))$ and the Mackey formula imply (i).

An analogous version of Lemma 2.2 with identical proof holds over F instead of \mathcal{O} .

2.3 Let $P \in S_p(G)$. Recall from [Bro85] that the *Brauer construction* with respect to P is a functor

 $-(P)\colon {}_{FG}\mathsf{triv}\to {}_{F[N_G(P)/P]}\mathsf{triv}\,,\quad M\mapsto M(P)\,.$

If $M \in {}_{FG}$ triv is indecomoposable and P is a vertex of M then the inflated module $\inf_{N_G(P)/P}^{N_G(P)}(M(P)) \in {}_{N_G(P)}$ triv is the Green correspondent of M. Moreover, for $Q \in S_p(G)$, one has $M(Q) \neq \{0\}$ if and only if $Q \leq_G P$.

The commutative diagram

defines a unique ring homomorphism $T(\mathcal{O}G) \to T(\mathcal{O}[N_G(P)/P])$ which we abusively denote again by -(P). Note however, that this map does not come from a functor $\mathcal{O}_G \mathsf{triv} \to \mathcal{O}_{[N_G(P)/P]}\mathsf{triv}$. For $M \in \mathcal{O}_G \mathsf{triv}$ the image [M](P) of $[M] \in T(\mathcal{O}G)$, can be constructed as follows: Decompose

$$\operatorname{Res}_{N_G(P)}^G(M) = V_1 \oplus \ldots \oplus V_n$$

into indecomposable $\mathcal{O}N_G(P)$ -modules V_1, \ldots, V_n and let I denote the set of all $i \in \{1, \ldots, n\}$ such that V_i satisfies the equivalent conditions of Lemma 2.2 (with $H = N_G(P)$), i.e, Pacts trivially on V_i . For each $i \in I$ one has $V_i \cong \text{Inf}_{N_G(P)/P}^{N_G(P)}(\tilde{V}_i)$ for an indecomposable $\tilde{V}_i \in \mathcal{O}[N_G(P)/P]$ triv, which is uniquely determined up to isomorphism by V_i . Then $[M](P) = \sum_{i \in I} [\tilde{V}_i]$. By abuse of notation we also set

$$M(P) := \bigoplus_{i \in I} \tilde{V}_i \in {}_{\mathcal{O}[N_G(P)/P]}\mathsf{triv}\,,$$

which, by the Krull-Schmidt Theorem is uniquely determined up to isomorphism.

2.4 Lemma Let M be a p-permutation $\mathcal{O}G$ -module and let $P \in S_p(G)$. For any subgroup H satisfying $P \leq H \leq N_G(P)$ one has

$$\operatorname{Res}_{H/P}^{N_G(P)/P}(M(P)) \cong \left(\operatorname{Res}_H^G(M)\right)(P)$$

as $\mathcal{O}[H/P]$ -modules.

Proof Let V be an indecomposable direct summand of $\operatorname{Res}_{N_G(P)}^G(M)$ and let W be an indecomposable direct summand of $\operatorname{Res}_{H}^{N_G(P)}(V)$. By Lemma 2.2, the $\mathcal{O}[N_G(P)]$ -module V satisfies the conditions in Lemma 2.2 if and only if the $\mathcal{O}H$ -module W does. In fact, if V satisfies (i) then W does, and if W satisfies (ii) then V does. The result follows.

3 Proof of Theorem A

In this section we will prove Theorem A and generalize it to statements that involve sums of blocks and restricted vertices.

Using the construction from 2.3, we first define the homomorphism β_G . Note that G acts on the ring $\prod_{P \in \mathcal{S}_p(G)} R(K[N_G(P)/P])$ via ring automorphisms by

$${}^{g}((\chi_P)_{P\in\mathfrak{S}_p(G)}):=\left({}^{g}(\chi_{g^{-1}P})\right)_{P\in\mathfrak{S}_p(G)}.$$

Here, ${}^{g}(\chi_{g^{-1}P}) \in R(K[N_{G}(P)/P])$ denotes the character arising from $\chi_{g^{-1}P}$ via restriction along the isomorphism $N_{G}(P)/P \to N_{G}({}^{g^{-1}}P)/{}^{g^{-1}}P$ induced by $x \mapsto g^{-1}xg$ for $x \in N_{G}(P)$. Thus, a tuple (χ_{P}) is a *G*-fixed point if and only if ${}^{g}\chi_{P} = \chi_{{}^{g}P}$ for all $g \in G$ and $P \in \mathcal{S}_{p}(G)$. The homomorphism β_{G} is now defined as

$$\beta_G \colon T(\mathcal{O}G) \to \left(\prod_{P \in \mathscr{S}_p(G)} R(K[N_G(P)/P])\right)^G, \quad \omega \mapsto \left(\kappa_{N_G(P)/P}(\omega(P))\right)_{P \in \mathscr{S}_p(G)}$$

Since both the maps $-(P): T(\mathcal{O}G) \to T(\mathcal{O}[N_G(P)/P])$ and $\kappa_{N_G(P)/P}$ commute with *G*-conjugations, the image of β_G is contained in the *G*-fixed points, and since both maps are ring homomorphisms, β_G is a ring homomorphism.

Proof of Theorem A. First note that β_G is injective, since, by the commutative diagram in 2.1(d), the diagram

is commutative and the composition of the two bottom maps is injective by a theorem of Conlon, see [Be98, Proposition 5.5.4].

Next we show that every element in the image of β_G satisfies the condition (C) in Theorem A. This can be derived from [Ri96, Lemma 6.2], but we give a detailed proof for the reader's convenience. Let $M \in {}_{\mathcal{O}G}$ triv and $(\chi_P) = \beta_G([M])$. Let $P \in \mathcal{S}_p(G)$, and $x \in N_G(P)$. We need to show that $\chi_P(xP) = \chi_Q(xQ)$ with $Q := P\langle x_p \rangle$. By Lemma 2.4 with $H := P\langle x \rangle$, we may assume that $G = P\langle x \rangle$. We may also assume that M is indecomposable and P < Q. Let R be a vertex of M. If R = Q then M(P) = M = M(Q) and therefore $\chi_P(xP) = \chi_Q(xQ)$. If $R \neq Q$ then R < Q and therefore $M(Q) = \{0\}$ and $\chi_Q(xQ) = 0$. In order to show that $\chi_P(xP) = 0$, we distinguish the two cases $P \leq R$ and $P \leq R$. In the latter case, $M(P) = \{0\}$ and we are done. So assume that $P \leq R < Q$. Then, with $\tilde{\mathcal{O}}$ denoting a finite extension of \mathcal{O} containing a root of unity of order $|\langle x_{p'} \rangle|$, we have

$$\tilde{\mathcal{O}} \otimes_{\mathcal{O}} M \mid \mathrm{Ind}_{R}^{G}(\tilde{\mathcal{O}}) \cong \mathrm{Ind}_{R\langle x_{p'} \rangle}^{G} \mathrm{Ind}_{R}^{R\langle x_{p'} \rangle}(\tilde{\mathcal{O}}) \cong \bigoplus_{\varphi} \mathrm{Ind}_{R\langle x_{p'} \rangle}^{G}(\tilde{\mathcal{O}}_{\varphi}) \,,$$

where φ runs through all homomorphisms $R\langle x_{p'}\rangle \to \tilde{\mathcal{O}}^{\times}$ with $\varphi|_R = 1$. Green's indecomposability theorem implies that $\tilde{\mathcal{O}} \otimes_{\mathcal{O}} M$ is isomorphic to a direct sum of modules of the form $\mathrm{Ind}_{R\langle x_{p'}\rangle}^G(\tilde{\mathcal{O}}_{\varphi})$, for homomorphisms φ as above. Therefore, M(P) = M and $\chi_P(xP)$ equals a sum of characters of $\tilde{\mathcal{O}}G$ -modules of the form $\mathrm{Ind}_{R\langle x_{p'}\rangle}^G(\tilde{\mathcal{O}}_{\varphi})$ evaluated at x. The latter evaluation equals 0, since x is not G-conjugate to an element of $R\langle x_{p'}\rangle$ noting that R < Q.

Conversely, assume that we have a family of virtual characters $\chi_P \in R(K[N_G(P)/P])$ satisfying

$${}^{g}\chi_{P} = \chi_{{}^{g}P} \quad \text{and} \quad \chi_{P}(xP) = \chi_{P\langle x_{p} \rangle}(xP\langle x_{p} \rangle)$$

$$\tag{5}$$

for all $P \in S_p(G)$, $g \in G$, and $x \in N_G(P)$. We will construct an element $\omega \in T(\mathcal{O}G)$ with $\beta_G(\omega) = (\chi_P)_{P \in S_p(G)}$ inductively. Let $P_1, \ldots, P_n \in S_p(G)$ be representatives of the *G*-conjugacy classes and assume that $|P_1| \ge |P_2| \ge \cdots \ge |P_n|$. We will show by induction on $i \in \{1, \ldots, n\}$ that there exists an element $\omega_i \in T(\mathcal{O}G)$ with the following two properties:

(i) ω_i is a Z-linear combination of the standard basis elements [M] with vertex P_i .

(ii) The P_i -component of $\beta_G(\omega_1 + \cdots + \omega_i)$ is equal to χ_{P_i} .

i = 1: Note that $P := P_1$ is a Sylow *p*-subgroup of *G*. Since $N_G(P)/P$ is a *p'*-group, we have $Pr(\mathcal{O}[N_G(P)/P]) = T(\mathcal{O}[N_G(P)/P])$ and $\kappa_{N_G(P)/P} \colon Pr(\mathcal{O}[N_G(P)/P]) \to R_K(N_G(P)/P)$ is an isomorphism. Thus, there exists a \mathbb{Z} -linear combination $\gamma := a_1[V_1] + \cdots + a_k[V_k]$ with indecomposable projective $\mathcal{O}[N_G(P)/P]$ -modules V_1, \ldots, V_k such that

 $\kappa_{N_G(P)/P}(\gamma) = \chi_P$. Then each $W_r := \operatorname{Inf}_{N_G(P)/P}^{N_G(P)}(V_r)$, $r = 1, \ldots, k$, is an indecomposable p-permutation $\mathcal{O}[N_G(P)]$ -module with vertex P. For $r = 1, \ldots, k$, let $M_r \in {}_{OG}$ triv be the Green correspondent of W_r and set $\omega_1 := a_1[M_1] + \cdots + a_k[M_k] \in T(\mathcal{O}G)$. Then ω_1 satisfies (i) and (ii).

 $i \to i+1$: Set $P := P_{i+1}$, let $x \in N_G(P)$ with $x_p \notin P$, and set $Q := P\langle x_p \rangle$. Then Q > P. Let $\beta_G(\omega_1 + \cdots + \omega_i)_P \in R_K(N_G(P)/P)$ denote the *P*-component of $\beta_G(\omega_1 + \cdots + \omega_i)$.

Claim: $\beta_G(\omega_1 + \cdots + \omega_i)_P(xP) = \chi_P(xP)$. To prove the claim, let $j \in \{1, \ldots, i\}$ with $Q =_G P_j$. We first show that

$$\beta_G(\omega_{j+1} + \dots + \omega_i)_P(xP) = 0.$$
(6)

If $\omega_{j+1} + \cdots + \omega_i = 0$ then this is obvious. So assume that M is an indecomposable ppermutation $\mathcal{O}G$ -module such that [M] appears in $\omega_{j+1} + \cdots + \omega_i$ with nonzero coefficient
and let W be an indecomposable $\mathcal{O}N_G(P)$ -module that appears in M(P). It suffices to show
that the $\chi_W(x) = 0$, where χ_W denotes the character of W. So assume that $\chi_W(x) \neq 0$,
which implies $x_p \in R$ for a vertex R of W. Since $W \mid M(P) \mid \operatorname{Res}_{N_G(P)}^G(M)$, we have $R \geq P$.
Thus $Q = P\langle x_p \rangle \leq R$. But by (i), M has a vertex R' with $|R'| \leq |P_j| = |Q|$ and since $W \mid M(P) \mid \operatorname{Res}_{N_G(P)}^G(M)$, we also have $R \leq_G R'$. Thus, $|R| \leq |Q|$ so that $Q \leq R$ implies R = Q. Thus, M has vertex $P_j =_G Q$, in contradiction to M appearing in $\omega_{j+1} + \cdots + \omega_i$.
Hence, (6) holds. Next note that since $\beta_G(\omega_1 + \cdots + \omega_j)$ satisfies the coherence condition
(C), we obtain

$$\beta_G(\omega_1 + \dots + \omega_j)_P(xP) = \beta_G(\omega_1 + \dots + \omega_j)_Q(xQ).$$
(7)

Further, by the induction hypothesis for $j \leq i$, and since the tuple (χ_P) satisfies the coherence condition we have

$$\beta_G(\omega_1 + \dots + \omega_j)_Q(xQ) = \chi_Q(xQ) = \chi_P(xP).$$
(8)

Combining the equations (6), (7) and (8) now proves the claim.

Next we set

$$\psi := \chi_P - \beta_G(\omega_1 + \dots + \omega_i)_P \in R_K(N_G(P)/P),$$

which is a virtual character that vanishes on *p*-singular elements by the claim. By [Se78, Théorèmes 34 et 36], see also 2.1(d), we can write $\psi = \kappa_{N_G(P)/P}(\gamma)$ for a unique Z-linear combination $\gamma = a_1[V_1] + \cdots + a_k[V_k]$ with indecomposable projective $\mathcal{O}[N_G(P)/P]$ -modules V_1, \ldots, V_k . For $r = 1, \ldots, k$, set $W_r := \text{Inf}_{N_G(P)/P}^{N_G(P)}(V_r)$. Then each W_r has vertex *P*. For each $r = 1, \ldots, k$, let M_r denote the Green correspondent of W_r . Then each M_r is an indecomposable *p*-permutation $\mathcal{O}G$ -module with vertex *P*. We set

$$\omega_{i+1} := a_1[M_1] + \dots + a_k[M_k].$$

Thus, ω_{i+1} satisfies (i). Moreover, it satisfies (ii), since

$$\beta_G(\omega_{i+1})_P = \kappa_{N_G(P)/P}(a_1[V_1] + \cdots + a_k[V_k]) = \psi$$

and therefore

$$\beta_G(\omega_1 + \cdots + \omega_{i+1})_P = \beta_G(\omega_1 + \cdots + \omega_i)_P + \psi = \chi_P.$$

This completes the induction proof.

Finally, set $\omega := \omega_1 + \cdots + \omega_n$ and let $i \in \{1, \ldots, n\}$. Then we have

$$\beta_G(\omega)_{P_i} = \beta_G(\omega_1 + \dots + \omega_i)_{P_i} = \chi_{P_i}$$

In fact, the first equation follows from $\omega_j(P_i) = 0$ for all $j \in \{i + 1, ..., n\}$, since $P_i \not\leq_G P_j$, see 2.3, and the second equation follows from (ii). Finally, since the family (χ_P) is *G*-fixed we also have $\beta_G(\omega)_P = \chi_P$ for any $P \in \mathcal{S}_p(G)$.

3.1 Notation We will denote from now on by

$$\left(\prod_{P\in\mathcal{S}_p(G)} R\left(K[N_G(P)/P]\right)\right)' \tag{9}$$

the set of all tuples (χ_P) with $\chi_P \in R(K[N_G(P)/P])$, for $P \in S_p(G)$, which satisfy the two conditions in (5).

3.2 Next we derive a block-wise version of Theorem A. Let $e \in Z(\mathcal{O}G)$ be an idempotent. For $P \in \mathcal{S}_p(G)$ denote by e(P) the unique idempotent in $Z(\mathcal{O}[N_G(P)])$ with $\operatorname{br}_P(e) = \overline{e(P)}$, where $\operatorname{br}_P: Z(\mathcal{O}G) \to Z(F[N_G(P)])$ denotes the Brauer homomorphism. It is a well-known property of the Brauer construction that

$$e(P)M(P) = M(P)$$
 for every $M \in \mathcal{O}_{Ge}$ triv. (10)

Note that, since ${}^{g}e(P) = e({}^{g}P)$ for all $g \in G$ and $P \in S_{p}(G)$, the subgroup $\prod_{P \in S_{p}(G)} R(K[N_{G}(P)/P]e(P))$ of $\prod_{P \in S_{p}(G)} R(K[N_{G}(P)/P])$ is G-stable. Therefore, if we denote by

$$\left(\prod_{P\in\mathcal{S}_p(G)} R\left(K[N_G(P)/P]e(P)\right)\right)'$$

the set of all tuples (χ_P) with $\chi_P \in R(K[N_G(P)/P]e(P))$, for $P \in S_p(G)$, which satisfy the two conditions in (5) then β_G restricts to an injective group homomorphism

$$\beta_G \colon T(\mathcal{O}Ge) \to \left(\prod_{P \in \mathcal{S}_P(G)} R\left(K[N_G(P)/P]e(P)\right)\right)',\tag{11}$$

where we consider e(P) also as a central idempotent of $K[N_G(P)/P]$ via the canonical Kalgebra epimorphism $K[N_G(P)] \to K[N_G(P)/P]$. Note that $T(\mathcal{O}Ge)$ is in general not a subring of $T(\mathcal{O}G)$.

3.3 Corollary For every idempotent $e \in Z(\mathcal{O}G)$, the map in (11) is an isomorphism.

Proof It suffices to prove the surjectivity. So let (χ_P) be a tuple in the codomain of (11). Then, by Theorem A there exists $\omega \in T(\mathcal{O}G)$ with $\beta_G(\omega) = (\chi_P)$. The decomposition $T(\mathcal{O}G) = T(\mathcal{O}Ge) \oplus T(\mathcal{O}G(1-e))$ allows us to write $\omega = \omega e + \omega(1-e)$ with $\omega e \in T(\mathcal{O}Ge)$ and $\omega(1-e) \in T(\mathcal{O}G(1-e))$. The additivity of β_G and the property (10) for the idempotents e and 1-e now imply that $\beta_G(\omega(1-e)) = 0$ and $\beta(\omega e) = (\chi_P)$. **3.4 Notation** Let \mathfrak{X} be a subset of $S_p(G)$ that is closed under *G*-conjugation and under taking subgroups and let $e \in Z(\mathcal{O}G)$ be an idempotent. We denote by $T(\mathcal{O}Ge | \mathfrak{X})$ the \mathbb{Z} -span of the standard basis elements [M], where $M \in _{\mathcal{O}Ge}$ triv is indecomposable and has a vertex belonging to \mathfrak{X} . Moreover, we denote by

$$\left(\prod_{P\in\mathfrak{X}} R\left(K[N_G(P)/P]e(P)\right)\right)'$$

the set of all tuples (χ_P) with $\chi_P \in R(K[N_G(P)/P]e(P))$, for $P \in \mathfrak{X}$, satisfying

$${}^{g}\chi_{P} = \chi_{{}^{g}P}$$
 and $\chi_{P}(xP) = \chi_{Q}(xQ)$

for all $g \in G$, $P \in \mathfrak{X}$ and $x \in N_G(P)$, with $Q := P\langle x_p \rangle$, where we interpret χ_Q as 0 if $Q \notin \mathfrak{X}$. Note that for $M \in T(\mathcal{O}Ge | \mathfrak{X})$, one has $M(P) = \{0\}$ whenever $P \in \mathfrak{S}_p(G) \setminus \mathfrak{X}$. Thus, the map β_G restricts to an injective group homomorphism

$$\beta_G \colon T(\mathcal{O}Ge \mid \mathfrak{X}) \to \left(\prod_{P \in \mathfrak{X}} R\left(K[N_G(P)/P]e(P)\right)\right)',\tag{12}$$

whose codomain we consider as a subset of $\prod_{P \in S_p(G)} R(K[N_G(P)/P]e(P))$ by adding 0's in the missing components.

3.5 Corollary Let $\mathfrak{X} \subseteq \mathfrak{S}_p(G)$ be a subset that is closed under taking *G*-conjugates and under taking subgroups and let $e \in Z(\mathcal{O}G)$ be an idempotent. Then the map in (12) is an isomorphism.

Proof It suffices to show the surjectivity. Suppose that (χ_P) belongs to the codomain of the map (12), and view this tuple as element in $\prod_{P \in \mathcal{S}_p(G)} R(K[N_G(P)/P]e(P))$ by adding 0's in the components indexed by $P \in \mathcal{S}_p(G) \setminus \mathcal{X}$. Then (χ_P) belongs to the codomain of the map in (11). By Corollary 3.3 there exists $\omega \in T(\mathcal{O}Ge)$ with $\beta_G(\omega) = (\chi_P)$ for all $P \in \mathfrak{S}_p(G)$. We'll show that $\omega \in T(\mathcal{O}G \mid \mathfrak{X})$. Let \mathfrak{Y} denote the set of all $P \in \mathfrak{S}_p(G)$ such that there exists an indecomposable p-permutation $\mathcal{O}G$ -module M having a vertex that contains P and occurring with nonzero coefficient in ω . Then \mathcal{Y} is closed under G-conjugation and taking subgroups. It suffices to show that $\mathcal{Y} \subseteq \mathcal{X}$. Let P be a maximal element in \mathcal{Y} . It suffices to show that $P \in \mathfrak{X}$. Suppose this is not the case. Then $0 = \chi_P = \kappa_{N_G(P)/P}(\omega(P))$. Moreover, $\omega(P) \in Pr(\mathcal{O}[N_G(P)/P])$, since P was chosen maximal in \mathcal{Y} . Since $\kappa_{N_G(P)/P}$ is injective on $Pr(\mathcal{O}[N_G(P)/P])$, this implies that $\omega(P) = 0$. We can write $\omega = \omega_1 + \omega_2$, such that P is a vertex of the indecomposable modules appearing in ω_1 and P is not a vertex of the indecomposable modules appearing in ω_2 . By the maximality of P in \mathcal{Y} , we obtain $\omega_1 \neq 0$ but $\omega_1(P) = \omega(P) = 0$. We write $\omega_1 = a_1[M_1] + \cdots + a_k[M_k]$ with representatives M_1, \ldots, M_k of the isomorphism classes of indecomposable *p*-permutation $\mathcal{O}Ge$ -modules with vertex P and $a_1, \ldots, a_k \in \mathbb{Z}$, we obtain $\omega_1(P) = a_1[V_1] + \cdots + a_k[V_k]$, where each V_r is an indecomposable projective $\mathcal{O}[N_G(P)/P]e(P)$ -module whose inflation W_r to $N_G(P)$ is the Green correspondent of M_r , for $r = 1, \ldots, k$. Since the Green correspondence is bijective and the inflation map $T(\mathcal{O}[N_G(P)/P]e(P)) \to T(\mathcal{O}[N_G(P)]e(P))$ is injective, V_1, \ldots, V_r are pairwise non-isomorphic. Thus, $a_1 = \cdots = a_k = 0$ and therefore $\omega_1 = 0$, a contradiction. **3.6 Remark** (a) It is worth mentioning that in Theorem A, the field K can be any finite extension of \mathbb{Q}_p , including \mathbb{Q}_p itself.

(b) Suppose that K contains a root of unity of order $\exp(G)$, the exponent of G, so that R(KG) is isomorphic to the character ring of G. The image of β_G followed by the projection map π_1 onto R(KG) is equal to $\kappa_G(T(\mathcal{O}G))$. The coherence condition (C) in Theorem A implies the well-known fact that characters of p-permutation $\mathcal{O}G$ -modules are p-rational, i.e., with values contained in the group $\mathbb{Q}_p(\zeta)$, where ζ is a root of unity of G of order $\exp(G)_{p'}$. The latter is also the group $R(\mathbb{Q}_p(\zeta)G)$ of characters of $\mathbb{Q}_p(\zeta)$ -modules by a Theorem of Brauer, see [Bra41, Lemma 1^{*}]. It is shown in [McH21] that the factor group $R(\mathbb{Q}_p(\zeta)G)/\kappa_G(T(\mathcal{O}G))$ is an elementary abelian 2-group (trivial if p is odd) and that these groups (for varying G) form a uniserial fibered biset functor over the field \mathbb{F}_2 whose composition factors are indexed by the quaternion groups.

4 The orthogonal unit group of $T(\mathcal{O}G)$

Recall from Section 1 that the orthogonal unit group of $T(\mathcal{O}G)$ is defined by $O(T(\mathcal{O}G)) = \{u \in T(\mathcal{O}G) \mid u \cdot u^\circ = 1\}.$

4.1 Remark (a) Our interest in the orthogonal unit group $O(T(\mathcal{O}G))$ stems from the fact that the functor $\operatorname{Ind}_{\Delta(G)}^{G\times G} : {}_{\mathcal{O}G}\mathsf{triv} \to {}_{\mathcal{O}[G\times G]}\mathsf{triv}$, where we identify G and the diagonal subgroup $\Delta(G) := \{(g,g) \mid g \in G\}$ of $G \times G$ via $g \mapsto (g,g)$, yields an injective ring homomorphism $\Delta : T(\mathcal{O}G) \to T^{\Delta}(\mathcal{O}G, \mathcal{O}G)$, when we identify left $\mathcal{O}[G \times G]$ -modules with $(\mathcal{O}G, \mathcal{O}G)$ bimodules. Here $T^{\Delta}(\mathcal{O}G, \mathcal{O}G)$ denotes the free abelian group on isomorphism classes of indecomposable $(\mathcal{O}G, \mathcal{O}G)$ -bimodules which are p-permutation modules with twisted diagonal vertices when considered as $\mathcal{O}[G \times G]$ -modules. The product on the ring $T^{\Delta}(\mathcal{O}G, \mathcal{O}G)$ is induced by taking the tensor product over $\mathcal{O}G$. If u is an orthogonal unit of $T(\mathcal{O}G)$ then its image $\Delta(u) \in T^{\Delta}(\mathcal{O}G, \mathcal{O}G)$ is a p-permutation autoequivalence of $\mathcal{O}G$, see [BoPe20]. Thus, $O(T(\mathcal{O}G))$ can be considered as a subgroup of the group of p-permutation autoequivalences of $\mathcal{O}G$, a group we are interested in studying in the future.

(b) The group $O(T(\mathcal{O}G))$ is actually equal to the torsion subgroup of the unit group $T(\mathcal{O}G)^{\times}$, see also [Ca18, Section 9]. To see this, consider the map

$$\sigma_G \colon T(\mathcal{O}G) \to \prod_{(P,sP) \in \mathfrak{T}_p(G)} \mathbb{Z}[\zeta] \,,$$

where $\mathfrak{T}_p(G)$ is the set of all pairs (P, sP) with $P \in \mathfrak{S}_p(G)$ and sP a p'-element of $N_G(P)/P$, and where the (P, sP)-component of $\sigma_G(\omega)$ is given by the evaluation of the virtual character $\kappa_{N_G(P)/P}(\omega(P))$ at the element sP. Here ζ can be taken as a root of unity of order $\exp(G)_{p'}$ in some extension of K. The map σ_G is an injective ring homomorphism (see [Be98, Corollary 5.5.5]) whose components are also called the *species* of $T(\mathcal{O}G)$. It commutes with duals: if the (P, sP)-component of $\sigma_G(\omega)$ is equal to $a \in \mathbb{Z}[\zeta]$ then the (P, sP)-component of $\sigma(\omega^\circ)$ is equal to the 'complex conjugate' \bar{a} of a, i.e., the image of a under the unique Galois automorphism of $\mathbb{Q}(\zeta)$ sending ζ to ζ^{-1} . By [Wa82, Theorem 4.12], every element $a \in \mathbb{Z}[\zeta]$ with $a\bar{a} = 1$ has finite order. This implies that the image of every element $u \in O(T(\mathcal{O}G))$ has finite order, and that also u has finite order by the injectivity of σ_G . Conversely, if $u \in T(\mathcal{O}G)^{\times}$ has finite order then also $\sigma_G(u)$ has finite order. Thus, every component $a_{(P,sP)}$ of $\sigma_G(u)$ is a root of unity and satisfies $a_{(P,sP)}\bar{a}_{(P,sP)} = 1$. Since σ_G transforms duals into complex conjugates and σ_G is injective, u satisfies $uu^\circ = 1$.

(c) Taking K-duals of KG-modules induces a ring automorphism $-\circ: R(KG) \to R(KG)$. Again, we call a unit $u \in R(KG)^\circ$ an orthogonal unit if $uu^\circ = 1$. The orthogonal units of R(KG) form a subgroup O(R(KG)) of the unit group $R(KG)^{\times}$. Using the species of R(KG), i.e., the injective ring homomorphism $R(KG) \to \prod_{g \in G} \mathbb{Z}[\zeta]$ with ζ a primitive root of unity of order exp G in some extension field of K, the same arguments as in (b) show that O(R(KG)) equals the torsion subgroup of $R(KG)^{\times}$. Moreover, it follows from [Ya91] that every torsion unit of R(KG) is of the form $\epsilon \cdot \varphi$, where $\epsilon \in \{\pm 1\}$ and $\varphi \in \operatorname{Hom}(G, K^{\times})$ is a linear character with values in K. Therefore, $O(R(KG)) = \{\pm 1\} \times \operatorname{Hom}(G, K^{\times})$, see also [Ca18, Section 9].

4.2 Proposition For each $u \in O(T(\mathcal{O}G))$ there exist unique signs $\epsilon_P \in \{\pm 1\}$ and homomorphisms $\varphi_P \in \operatorname{Hom}(N_G(P)/P, K^{\times})$, for $P \in \mathcal{S}_P(G)$, such that

$$\beta_G(u) = (\epsilon_P \cdot \varphi_P)_{P \in \mathcal{S}_p(G)}.$$


Proof First note that $\beta_G(\omega^\circ) = \beta_G(\omega)^\circ$, for all $\omega \in T(\mathcal{O}G)$, where we use the notation $(\chi_P)^\circ := (\chi_P^\circ)$ for any tuple $(\chi_P) \in \prod_{P \in \mathcal{S}_p(G)} R(K[N_G(P)/P])$. Therefore, if $u \in O(T(\mathcal{O}G))$ then the *P*-component of $\beta_G(u)$ is an orthogonal unit in $R(K[N_G(P)/P])$. The result follows now from Remark 4.1(c).

We will continue the study of $O(T(\mathcal{O}G))$ after the proof of Theorem B.

4.3 Definition ([Re15]) Let S be a Sylow p-subgroup of p and let $\mathcal{F} := \mathcal{F}_S(G)$ denote the associated fusion system on S. The Burnside ring of \mathcal{F} , denoted by $B(\mathcal{F})$ is defined as the subring of the Burnside ring B(S) consisting of all elements $a \in B(S)$ with the property that $\phi_P(a) = \phi_Q(a)$, whenever P and Q are G-conjugate subgroups of S. Here, for $P \leq S$, $\phi_P : B(S) \to \mathbb{Z}$ is the ring homomorphism defined by $\phi_P([X]) := |X^P|$, whenever X is a finite S-set, where X^P denotes the set of P-fixed points of X.

4.4 Theorem ([BaCa20, Theorem 7.1]) Let S be a Sylow p-subgroup of G and let $\mathcal{F} := \mathcal{F}_S(G)$ denote the associated fusion system on S. Then the ring homomorphism $\operatorname{res}_S^G : B(G) \to B(\mathcal{F})$ induced by restricting a finite G-set to S, is split surjective. More precisely, there exists a ring homomorphism $t_S^G : B(\mathcal{F}) \to B(G)$ such that $\operatorname{res}_S^G \circ t_S^G = \operatorname{id}_{B(\mathcal{F})}$. For $a \in B(\mathcal{F})$, the element $t_S^G(a)$ is determined by the equations $\phi_H(t_S^G(a)) = \phi_{P_H}(a)$, for all subgroups $H \leq G$, where P_H is a subgroup of S which is G-conjugate to a Sylow p-subgroup of H.

Proof of Theorem B. That (1) is the identity on $B(\mathcal{F})$ is already stated in Theorem 4.4. And that (2) is the identity on $B(\mathcal{F})$ follows from the obvious commutativity of the diagram



and from (1) being the identity.

For the proof of Theorem C we need the following lemma which follows immediately from Lemma 2.4.

4.5 Lemma Let $H \leq G$. One has a commutative diagram

where the right vertical map $\operatorname{res}_{H}^{G}$ is the ring homomorphism defined by sending $(\chi_{P})_{P \in \mathbb{S}_{p}(G)}$ to $\left(\operatorname{res}_{N_{H}(P)/P}^{N_{G}(P)/P}(\chi_{P})\right)_{P \in \mathbb{S}_{p}(H)}$.

Proof of Theorem C. First note that if we define $-\circ: B(G) \to B(G)$ as the identity, then all four maps in (2) commute with $-\circ$. Since they are also ring homomorphisms, they restrict to a sequence of group homomorphisms

$$\lambda_S^{-1} \circ \operatorname{res}_S^G \circ \lambda_G \circ t_S^G \colon B(\mathcal{F})^{\times} \to B(G)^{\times} \to \mathcal{O}(T(\mathcal{O}G)) \to \mathcal{O}(T(\mathcal{O}S)) = T(\mathcal{O}S)^{\times} \xrightarrow{\sim} B(S)^{\times}$$
(13)

whose composition is the identity of $B(\mathcal{F})^{\times}$. It follows immediately that

$$O(T(\mathcal{O}G)) = (\lambda_G \circ t_S^G)(B(\mathcal{F})) \times \ker\left(\operatorname{res}_S^G \colon O(T(\mathcal{O}G)) \to O(T(\mathcal{O}S))\right)$$
(14)

and that the first factor is isomorphic to $B(\mathcal{F})^{\times}$, since $\lambda_G \circ t_S^G$ is injective.

Next we show that $\beta_G \left(\ker \left(\operatorname{res}_S^G \colon \mathcal{O}(T(\mathcal{O}G)) \to \mathcal{O}(T(\mathcal{O}S)) \right) \right)$ is equal to the group

$$\left(\prod_{P\in\mathcal{S}_p(G)}\operatorname{Hom}(N_G(P)/P, K^{\times})\right)'$$
(15)

consisting of all tuples $(\varphi_P) \in \left(\prod_{P \in \mathbb{S}_p(G)} \operatorname{Hom}(N_G(P)/P, K^{\times})\right)^G$ satisfying

$$\varphi_P(xP) = \varphi_{P\langle x_P \rangle}(xP\langle x_P \rangle), \qquad (16)$$

for all $P \in \mathcal{S}_p(G)$ and $x \in N_G(P)$. First suppose that $u \in O(T(\mathcal{O}G))$ with $\operatorname{res}_S^G(u) = 1 = [\mathcal{O}_S]$. Write $\beta_G(u) = (\epsilon_P \cdot \varphi_P)_{P \in \mathcal{S}_p(G)}$ as in Proposition 4.2. Lemma 4.5 applied to H = S

13

implies that $\epsilon_P = 1$ for all $P \leq S$. Since the tuple (ϵ_P) is a *G*-fixed point, we obtain $\epsilon_P = \epsilon_{\binom{g_P}{P}}$ for all $P \in \mathcal{S}_p(G)$ and $g \in G$. This implies that $\epsilon_P = 1$ for all $P \in \mathcal{S}_p(G)$ and that $\beta_G(u)$ belongs to the group in (15). Conversely, if a tuple (φ_P) belongs to the group in (15), then by Theorem A there exists an element $\omega \in T(\mathcal{O}G)$ with $\beta(\omega) = (\varphi_P)$. Since β_G commutes with duals, we obtain $\beta_G(\omega^\circ) = (\varphi_P)^\circ = (\varphi_P^{-1})$ and therefore $\beta_G(\omega \cdot \omega^\circ) = \beta_G(\omega)\beta_G(\omega)^\circ = 1$. Since β_G is injective, we obtain $\omega \in O(T(\mathcal{O}G))$.

Since β_G is injective we have now proved that the second factor in (14) is isomorphic to the group in (15). Finally, if (φ_P) belongs to the group in (15) then φ_P vanishes on *p*-elements by the condition in (16) for all $P \in S_p(G)$. Thus each φ_P takes values in the roots of unity of K of *p'*-order of K which correspond isomorphically to those of F^{\times} . This concludes the proof of Theorem C.

The next result is a further decomposition of the second factor occurring in the decomposition of $O(T(\mathcal{O}G))$ in Theorem C.

4.6 Proposition One has a decomposition

$$\left(\prod_{P\in\mathfrak{S}_p(G)}\operatorname{Hom}(N_G(P)/P, F^{\times})\right)'\cong\operatorname{Hom}(G, F^{\times})\times\left(\prod_{P\in\mathfrak{S}_p(G)}\operatorname{Hom}(N_G(P)/PC_G(P), F^{\times})\right)',$$

where the second factor denotes the set of all tuples

$$(\varphi_P) \in \left(\prod_{P \in \mathcal{S}_p(G)} \operatorname{Hom}(N_G(P)/PC_G(P), F^{\times})\right)^G$$

satisfying

$$\varphi_P(xPC_G(P)) = \varphi_Q(xQC_G(Q)),$$

for all $P \in \mathfrak{S}_p(G)$ and all $x \in N_G(P)$, where $Q := P\langle x_p \rangle$.

Proof Set

$$H := \left(\prod_{P \in \mathcal{S}_p(G)} \operatorname{Hom}(N_G(P), F^{\times})\right)^G \tag{17}$$

and note that the component-wise inflations map the group $\left(\prod_{P \in S_p(G)} \operatorname{Hom}(N_G(P)/P, F^{\times})\right)'$ isomorphically onto the subgroup H_1 of H, consisting of those tuples $(\varphi_P) \in H$ that satisfy

$$P \leq \ker(\varphi_P) \quad \text{and} \quad \varphi_P(x) = \varphi_{P\langle x_p \rangle}(x)$$
(18)

for all $P \in \mathcal{S}_p(G)$ and all $x \in N_G(P)$. Similarly, the component-wise inflations map the group $\left(\prod_{P \in \mathcal{S}_p(G)} \operatorname{Hom}(N_G(P)/PC_G(P), F^{\times})\right)'$ isomorphically onto the group H_2 of tuples $(\varphi_P) \in H$ satisfying

$$PC_G(P) \leq \ker(\varphi_P) \quad \text{and} \quad \varphi_P(x) = \varphi_{P\langle x_p \rangle}(x)$$
(19)

for all $P \in \mathcal{S}_p(G)$ and all $x \in N_G(P)$.

Note that the homomorphisms ι : Hom $(G, F^{\times}) \to H_1$, $\varphi \mapsto (\varphi|_{N_G(P)})_{P \in \mathfrak{S}_p(G)}$, and $\pi \colon H_1 \to \operatorname{Hom}(G, F^{\times}), \ (\varphi_P) \mapsto \varphi_1$, where $1 \in \mathfrak{S}_p(G)$ denotes the trivial subgroup of G, satisfy $\pi \circ \iota = \operatorname{id}$. Therefore, one has a decomposition

$$\left(\prod_{P \in \mathcal{S}_p(G)} \operatorname{Hom}(N_G(P)/P, F^{\times})\right)' \cong H_1 \cong \iota(\operatorname{Hom}(G, F^{\times})) \times \ker(\pi).$$
(20)

Since ι is injective, the first factor in (20) is isomorphic to $\operatorname{Hom}(G, F^{\times})$. Thus, it suffices to show that $\ker(\pi) = H_2$. Clearly, if $(\varphi_P) \in H_2$ then $\varphi_1 = 1$. Conversely, assume now that $(\varphi_P) \in H$ with $\varphi_1 = 1$. It suffices to show that $PC_G(P) \leq \ker(\varphi_P)$ for all $P \in S_p(G)$. Since every torsion element of F^{\times} has p'-order, φ_P is trivial on any p-element of $N_G(P)$. Thus, it suffices to show that $\varphi_P(c) = 1$ for all p'-elements $c \in C_G(P)$. We show this by induction on |P|. The case P = 1 is clear, since $\varphi_1 = 1$. Let $1 < P \in S_p(G)$ be arbitrary. Choose a normal subgroup Q of P of index p and an element $x \in P$ with $P = Q\langle x \rangle$. Then, since φ_P and φ_Q vanish on p-elements and by the second property in (18) applied to Q and xc, we obtain $\varphi_P(c) = \varphi_P(xc) = \varphi_Q(cc) = \varphi_Q(c)$ which equals 1 by induction.

5 A criterion à la Yoshida

In this section we prove a result analogous to Yoshida's characterization of the mark values of units of the Burnside ring, see [Yo90, Proposition 6.5]. To set the stage, recall from Remark 4.1(b) that one has an injective ring homomorphism

$$\sigma_G \colon T(\mathcal{O}G) \to \left(\prod_{(P,sP)\in\mathfrak{T}_p(G)} K\right)^G, \quad \omega \mapsto \left(\kappa_{N_G(P)/P}(\omega(P))(sP)\right)_{(P,sP)\in\mathfrak{T}_p(G)}$$

where we view the element $\kappa_{N_G(P)/P}(\omega(P)) \in R(K[N_G(P)/P])$ as virtual character that can be evaluated at sP, and where the exponent G stands for taking fixed points under the natural conjugation action ${}^{g}(a_{(P,sP)})_{(P,sP)\in\mathfrak{T}_{p}(G)} := ({}^{g}a_{g^{-1}}_{(P,sP)})_{(P,sP)\in\mathfrak{T}_{p}(G)}$. Here we use the action of G on $\mathfrak{T}_{p}(G)$ given by ${}^{g}(P,sP) = ({}^{g}P, {}^{g}(sP))$. Since every orthogonal unit of $T(\mathcal{O}G)$ has finite order (see Remark 4.1(b) or Theorem C), the ring homomorphism σ_{G} restricts to an injective group homomorphism

$$\sigma_G \colon \mathcal{O}(T(\mathcal{O}G)) \to \left(\prod_{(P,sP) \in \mathfrak{T}_p(G)} \langle \zeta \rangle\right)^G,$$

where $\zeta \in K^{\times}$ is a root of unity that generates the group of roots of unity in K of order dividing $2 \exp(G)_{p'}$ (see again Theorem C).

5.1 Theorem Let $\zeta \in K^{\times}$ be as above and let $(\alpha_{(P,sP)}) \in (\prod_{(P,sP) \in \mathfrak{T}_p(G)} \langle \zeta \rangle)^G$. The following are equivalent:

(i) The tuple $(\alpha_{(P,sP)})$ belongs to $\sigma_G(O(T(\mathcal{O}G)))$.

(ii) For every $P \in S_p(G)$, the function $\psi_P \colon N_G(P)/P \to \langle \zeta \rangle$, $xP \mapsto \alpha_{(Q,xQ)} \cdot \alpha_{(P,1P)}$, where $Q = P \langle x_p \rangle$, is a group homomorphism. **Proof** Note that the functions ψ_P in (ii) are well-defined: If xP = yP for $x, y \in N_G(P)$, then $P\langle x_p \rangle = P\langle y_p \rangle$ and $xP\langle x_p \rangle = yP\langle y_p \rangle$.

Suppose first that $u \in O(T(\mathcal{O}G))$ and $(\alpha_{(P,sP)}) = \sigma_G(u)$. By Proposition 4.2 we have $\beta_G(u) = (\epsilon_P \cdot \varphi_P)$ with $\epsilon_P \in \{\pm 1\}$ and $\varphi_P \in \operatorname{Hom}(N_G(P)/P, K^{\times})$. Thus, $\alpha_{(P,sP)} = \epsilon_P \cdot \varphi_P(sP)$ for all $(P, sP) \in \mathcal{T}_p(G)$. As a consequence, for all $P \in \mathcal{S}_p(G)$ and $xP \in N_G(P)/P$, we have $\psi_P(xP) = \epsilon_Q \cdot \varphi_Q(xQ) \cdot \epsilon_P$, where $Q = P\langle x_p \rangle$. However, by the coherence property of the components of $\beta_G(u)$, we have $\epsilon_Q \cdot \varphi_Q(xQ) = \epsilon_P \cdot \varphi_P(xP)$ so that $\psi_P(xP) = \varphi_P(xP)$ and $\psi_P = \varphi_P$ is a homomorphism.

Conversely, suppose that ψ_P is a homomorphism for every $P \in S_p(G)$. This implies $1 = \psi_P(1P) = \alpha_{(P,1P)} \cdot \alpha_{(P,1P)}$ so that $\epsilon_P := \alpha_{(P,1P)} \in \{\pm 1\}$. For $P \in S_p(G)$, we define $\chi_P : N_G(P)/P \to K$, $xP \mapsto \epsilon_P \cdot \psi_P(xP)$. Since $\psi_P \in \operatorname{Hom}(N_G(P)/P, K^{\times})$, $\chi_P = \epsilon_P \cdot \psi_P \in R(K[N_G(P)/P])$. Since the tuple $(\alpha_{(P,sP)})$ is *G*-fixed, the tuple (χ_P) is *G*-fixed. We check that the tuple (χ_P) satisfies the coherence condition. Let $P \in S_p(G)$, $x \in N_G(P)$ and set $Q := P\langle x_p \rangle$. Then $\chi_P(xP) = \epsilon_P \cdot \psi_P(xP) = \epsilon_P^2 \cdot \alpha_{(Q,xQ)} = \alpha_{(Q,xQ)}$ and also $\chi_Q(xQ) = \epsilon_Q \cdot \psi_Q(xQ) = \epsilon_Q^2 \cdot \alpha_{(Q,xQ)} = \alpha_{(Q,xQ)}$. By Theorem A, there exists $u \in T(\mathcal{O}G)$ with $\beta_G(u) = (\chi_P)$, which implies that the (P, sP)-component of $\sigma_G(u)$ is $\chi_P(sP) = \alpha_{(P,sP)}$, so that $\sigma_G(u) = (\alpha_{(P,sP)})$. Moreover, $\beta_G(u \cdot u^\circ) = \beta_G(u) \cdot \beta_G(u)^\circ = 1$, since $\beta_G(u)^\circ = (\chi_P)^\circ = (\epsilon_P \cdot \psi_P^{-1})$. The injectivity of β_G implies $u \cdot u^\circ = 1$ so that $u \in O(T(\mathcal{O}G))$.

6 Examples

6.1 Proposition Suppose that every p-singular element of G is a p-element. Let S be a Sylow p-subgroup of G and let $\mathcal{F} := \mathcal{F}_S(G)$ be the associated fusion system on S. Moreover, let $\tilde{S}_p(G)$ be a set of representatives of the G-conjugacy classes of p-subgroups of G. Then

$$O(T(\mathcal{O}G)) \cong B(\mathcal{F})^{\times} \times \prod_{P \in \tilde{S}_p(G)} \operatorname{Hom}(N_G(P)/P, F^{\times}).$$

Proof We use the decomposition of $O(T(\mathcal{O}G))$ from Theorem C. Since every *p*-singular element of *G* is a *p*-element, the coherence condition in (4) is satisfied for every tuple $(\varphi_P) \in (\prod_{P \in S_p(G)} \operatorname{Hom}(N_G(P)/P, F^{\times}))^G$. Thus,

$$\left(\prod_{P\in\mathfrak{S}_p(G)}\operatorname{Hom}(N_G(P)/P, F^{\times})\right)' = \left(\prod_{P\in\mathfrak{S}_p(G)}\operatorname{Hom}(N_G(P)/P, F^{\times})\right)^G.$$

Moreover, the natural projection map

$$\left(\prod_{P\in\mathfrak{S}_p(G)}\operatorname{Hom}(N_G(P)/P,F^{\times})\right)^G\to\prod_{P\in\tilde{\mathfrak{S}}_p(G)}\operatorname{Hom}(N_G(P)/P,F^{\times})$$

is an isomorphism.

Note that the hypothesis of the above proposition is satisfied if G is a Frobenius group of the form $S \rtimes E$, where the Frobenius kernel S is a p-group and $E \leq \operatorname{Aut}(S)$ is a p'-group. In particular, if S is cyclic, we have the following corollary.

6.2 Corollary Suppose that $G = S \rtimes E$, where S is a cyclic p-group of order p^n and $E \leq \operatorname{Aut}(S)$ is a p'-group. Then

$$O(T(\mathcal{O}G)) \cong B(S)^{\times} \times \prod_{P \in \tilde{S}_p(G)} \operatorname{Hom}(E, F^{\times})^{n+1}$$

Proof This follows immediately from the previous proposition noting that for every $P \in S_p(G)$ one has $N_G(P) = G$ and $\operatorname{Hom}(G/P, F^{\times}) \cong \operatorname{Hom}(E, F^{\times})$. Moreover, $B(\mathcal{F}) = B(S)$, since S is cyclic.

Recall that G is called *p*-nilpotent if there exists a normal p'-subgroup N of G such that G/N is a *p*-group.

6.3 Proposition Suppose that G is a p-nilpotent group with Sylow p-subgroup S and let $\mathcal{F} := \mathcal{F}_S(G)$ be the associated fusion system on S. Then

$$O(T(\mathcal{O}G)) \cong B(S)^{\times} \times Hom(G, F^{\times}).$$

Proof We use again Theorem C. Since G is p-nilpotent, one has $\mathcal{F}_S(G) = \mathcal{F}_S(S)$ by a theorem of Frobenius. Therefore, two subgroups P and Q of S are G-conjugate if and only if they are S-conjugate. It follows that $B(\mathcal{F}) = B(S)$. Moreover, since G is p-nilpotent, the group $N_G(P)/C_G(P)$ is a p-group for every $P \in \mathcal{S}_p(G)$. The result now follows from Proposition 4.6 and noting that $\operatorname{Hom}(N_G(P)/PC_G(P), F^{\times})$ is trivial for all $P \in \mathcal{S}_p(G)$.

References

- [BaCa20] J. BARSOTTI, R. CARMAN: Burnside rings of fusion systems and their unit groups. J. Group Theory 23 (2020), 709–729.
- [Be98] D. J. Benson: Representations and cohomology. I. Basic representation theory of finite groups and associative algebras. Second edition. Cambridge Studies in Advanced Mathematics, 30. Cambridge University Press, Cambridge, 1998.
- [BoPe20] R. BOLTJE, P. PEREPELITSKY: *p*-permutation equivalences between blocks of group algebras. arXiv:2007.09253.
- [BoXu08] R. BOLTJE, B. XU: On *p*-permutation equivalences: Between Rickard equivalences and isotypies. Transactions of the American Mathematical Society **360** (2008), 5067–5087.
- [Bc10a] S. BOUC: Biset functors for finite groups, Lecture Notes in Mathematics, 1990. Springer-Verlag, Berlin, 2010.
- [Bra41] R. BRAUER: Investigations on group characters. Ann. of Math. 42 (1941), 936–958.

- [Bro85] M. BROUÉ: On Scott modules and p-permutation modules: an approach through the Brauer morphism. Proc. Amer. Math. Soc. 93 (1985), 401–408.
- [Bro90] M. BROUÉ: Isométries parfaites, types de blocs, catégories dérivées. Astérisque 181-182 (1990), 61–92.
- [Ca18] R. CARMAN: Unit groups of representation rings and their ghost rings as inflation functors. J. Algebra 498 (2018), 263–293.
- [McH21] J. R. MCHUGH: On the image of the trivial source ring in the ring of virtual characters of a finite group. *Pacific J. Math.* **315** (2021), 449–476.
- [Re15] S. P. REEH: The abelian monoid of fusion-stable finite sets is free. Algebra Number Theory 9 (2015), 2303–2324.
- [Ri96] J. RICKARD: Splendid equivalences: derived categories and permutation modules. Proceedings of the London Mathematical Society 72 (1996), 331–358.
- [Se78] J. P. SERRE: Représentations linéaires des groupes finis. Third revised edition. Hermann, Paris, 1978.
- [Wa82] L. C. WASHINGTON: Introduction to cyclotomic fields. Graduate Texts in Mathematics, 83. Springer-Verlag, New York, 1982.
- [Ya91] K. YAMAUCHI: On the units in the character ring. Hokkaido Math. J. 20 (1991), 477–479.
- [Yo90] T. YOSHIDA: On the unit groups of Burnside rings. J. Math. Soc. Japan 42 (1990), 31–64.