

Relation between the two geometric Satake equivalence via nearby cycle

Katsuyuki Bando*

May 24, 2022

Abstract

Fargues and Scholze proved the geometric Satake equivalence over the Fargues–Fontaine curve. On the other hand, Zhu proved the geometric Satake equivalence using a Witt vector affine Grassmannian. In this paper, we explain the relation between the two version of the geometric Satake equivalence via nearby cycle.

1 Introduction

Let F be a p -adic local field with a residue field \mathbb{F}_q . Write \mathcal{O}_F for its ring of integers. Put $k := \overline{\mathbb{F}}_q$. Let G be a reductive group scheme over \mathcal{O}_F . Let Λ be either of $\overline{\mathbb{Q}}_\ell$, a finite extension L of \mathbb{Q}_ℓ , its ring of integers \mathcal{O}_L , or its quotient ring. By [FS21, Theorem I.6.3], we have an equivalence of symmetric monoidal categories

$$\mathcal{S}_{\text{FS}}: \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Div}^1}, \Lambda) \xrightarrow{\sim} \text{Rep}(\widehat{G}^{\text{tw}} \rtimes W_F, \Lambda)$$

where $\text{Sat}(\mathcal{H}\text{ck}_{G, \text{Div}^1}, \Lambda)$ is the Satake category, which is the monoidal category of perverse sheaves with coefficients in Λ on the Hecke stack $\mathcal{H}\text{ck}_{G, \text{Div}^1}$ over the moduli space $\text{Div}^1 = \text{Spd } \check{F}/\varphi^{\mathbb{Z}}$ parametrizing degree 1 Cartier divisors on the Fargues–Fontaine curve for F . Let \widehat{G} be the Langlands dual group over Λ and W_F the Weil group of F . The group scheme $\widehat{G}^{\text{tw}} \rtimes W_F$ is a semidirect product of \widehat{G} and W_F with respect to the W_F -action on \widehat{G} obtained by twisting the usual action by an explicit cyclotomic

*Katsuyuki Bando, the University of Tokyo, Graduate School of Mathematical Sciences, Japan, kbando@ms.u-tokyo.ac.jp

action. Let $\text{Rep}(\widehat{G}^{\text{tw}} \rtimes W_F, \Lambda)$ be the monoidal category of $\widehat{G}^{\text{tw}} \rtimes W_F$ -representations on finite projective Λ -modules.

Consider a diagram

$$\begin{array}{ccc} \mathcal{H}\text{ck}_{G, \text{Spd } k} & \xrightarrow{i} & \mathcal{H}\text{ck}_{G, \text{Spd } \mathcal{O}_{\overline{F}}} \xleftarrow{j} \mathcal{H}\text{ck}_{G, \text{Spd } \widehat{F}} \\ & & \downarrow p_F \\ & & \mathcal{H}\text{ck}_{G, \text{Div}^1}. \end{array}$$

We have a nearby cycle functor

$$\Psi := i^* \circ Rj_* \circ (p_F)^* : \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Div}^1}, \Lambda) \rightarrow \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Spd } k}, \Lambda).$$

The main theorem is the following:

Theorem 1.1. *There exists a symmetric monoidal structure on $(\text{Sat}(\mathcal{H}\text{ck}_{G, \text{Spd } k}, \Lambda), \star)$ and a symmetric monoidal equivalence*

$$\mathcal{S}_{YZ} : \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Spd } k}, \Lambda) \xrightarrow{\sim} \text{Rep}(\widehat{G}, \Lambda)$$

such that for the diagram of functors

$$\begin{array}{ccc} \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Div}^1}, \Lambda) & \xrightarrow{\mathcal{S}_{\text{FS}}} & \text{Rep}(\widehat{G}^{\text{tw}} \rtimes W_F, \Lambda) \\ \Psi \downarrow & & \downarrow \text{For} \\ \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Spd } k}, \Lambda) & \xrightarrow{\mathcal{S}_{YZ}} & \text{Rep}(\widehat{G}, \Lambda), \end{array}$$

there is a natural isomorphism

$$\text{For} \circ \mathcal{S}_{\text{FS}} \cong \mathcal{S}_{YZ} \circ \Psi.$$

The equivalence \mathcal{S}_{YZ} with coefficients in $\overline{\mathbb{Q}}_\ell, \mathbb{F}_\ell$ or \mathbb{Z}_ℓ has the same form as the geometric Satake equivalence in [Zhu17] or [Yu19]. We do not prove that the equivalence \mathcal{S}_{YZ} is the same functor as the equivalence in [Zhu17] or [Yu19], which is too complicated to prove since the way of constructing a symmetric monoidal structure on Satake categories is quite different between [FS21] and [Zhu17]. However, we prove that the several properties of [Zhu17] also hold for \mathcal{S}_{YZ} .

The similar results to the above theorem are mentioned in [FS21, Remark I.2.14]. However, we show further that the nearby cycle functor, involving Rj_* , can be used as the vertical functor.

While we are completing this work, a related paper [AGLR22] appeared. The results of Lemma 3.6, Theorem 3.10 and Corollary 3.11 overlap with [AGLR22, Lemma 6.11, Proposition 6.12, Corollary 6.14], respectively. Our proofs of these results are also similar to that of [AGLR22], but slightly different. We prove Lemma 3.6 by reducing to the scheme settings, showing some results on the comparison to schemes. Also, our proofs are simpler than [AGLR22] since the result of [AGLR22] is more general.

Acknowledgement

The author is really grateful to his advisor Naoki Imai for useful discussions and comments.

2 Notations

Throughout this paper, F is a p -adic local field with a residue field \mathbb{F}_q . Write \mathcal{O}_F for its ring of integers. Put $k := \overline{\mathbb{F}_q}$.

Let ℓ be a prime number not equal to p . Let Λ be either of $\overline{\mathbb{Q}_\ell}$, a finite extension L of \mathbb{Q}_ℓ , its ring of integers \mathcal{O}_L , or its quotient ring. The category of finite projective Λ -module is denoted by $\text{Proj}(\Lambda)$. If Λ is a field, then we write $\text{Vect}(\Lambda)$ for $\text{Proj}(\Lambda)$.

Put

$$\text{Div}^1 := \text{Spd } \check{F} / \varphi^{\mathbb{Z}}$$

where φ is the Frobenius. Let G be a reductive group over $\text{Spec } \mathcal{O}_F$. For a small v-stack S with a map $S \rightarrow \text{Div}^1$, or $S \rightarrow \text{Spd } \mathcal{O}_{\check{F}}$, let $\text{Gr}_{G,S}$ be the Beilinson–Drinfeld affine Grassmannian of G over S and $\mathcal{H}ck_{G,S}$ the local Hecke stack defined in [FS21, VI.1].

The Langlands dual group of G over Λ is denoted by \widehat{G}_Λ or simply \widehat{G} . The scheme \widehat{G} has a natural W_F -action. We define the category $\text{Rep}(\widehat{G}, \Lambda)$ as the monoidal category of algebraic \widehat{G} -representations on finite projective Λ -modules. We write $\text{Rep}(\widehat{G} \rtimes W_F, \Lambda)$ for the category of $\widehat{G} \rtimes W_F$ -representations on finite projective Λ -modules such that the \widehat{G} -action is algebraic, and that the W_F -action is smooth. The category $\text{Rep}(\widehat{G}^{\text{tw}} \rtimes W_F, \Lambda)$ is similarly defined with respect to the twisted W_F -action on \widehat{G} appearing in [FS21, Theorem I.6.3].

If Λ is a torsion ring, then the derived category $D_{\text{ét}}(X, \Lambda)$ for a small v-stack X and the six functors with respect to this formalism are defined as in [Sch17, Definition 1.7].

3 Torsion coefficient case

Let Λ be a proper quotient of \mathcal{O}_L where L is a finite extension over \mathbb{Q}_ℓ .

3.1 Generalities on étale cohomology of diamonds

Put $C := \widehat{F}$.

3.1.1 Excision distinguished triangle

Lemma 3.1. *Let X be a small v -stack. For $A \in D_{\text{ét}}(X, \Lambda)$, the functor*

$$R\mathcal{H}\text{om}_{D_{\text{ét}}(X, \Lambda)}(-, A): D_{\text{ét}}(X, \Lambda)^{\text{op}} \rightarrow D_{\text{ét}}(X, \Lambda)$$

preserves distinguished triangles.

Proof. Following [Sch17, Corollary 17.2], write

$$R_{X_{\text{ét}}}: D(X_v, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda)$$

for the right adjoint of the inclusion $D_{\text{ét}}(X, \Lambda) \hookrightarrow D(X_v, \Lambda)$. Then as in the remark just after Lemma 17.8 of [Sch17], we have a natural isomorphism

$$R\mathcal{H}\text{om}_{D_{\text{ét}}(X, \Lambda)}(-, A) \cong R_{X_{\text{ét}}} R\mathcal{H}\text{om}_{D(X_v, \Lambda)}(-, A): D_{\text{ét}}(X, \Lambda)^{\text{op}} \rightarrow D_{\text{ét}}(X, \Lambda).$$

The functor $R\mathcal{H}\text{om}_{D(X_v, \Lambda)}(-, A)$ preserves distinguished triangles since it is the derived functor of the inner hom of v -sheaves. The functor $R_{X_{\text{ét}}}$ preserves distinguished triangles since its left adjoint $D_{\text{ét}}(X, \Lambda) \hookrightarrow D(X_v, \Lambda)$ preserves distinguished triangles. \square

Let $j: \text{Spd } C \rightarrow \text{Spd } \mathcal{O}_C$ and $i: \text{Spd } k \rightarrow \text{Spd } \mathcal{O}_C$ be inclusions.

Lemma 3.2. *For $A \in D_{\text{ét}}(\text{Spd } \mathcal{O}_C, \Lambda)^{bd}$,*

$$j_! Rj^* A \rightarrow A \rightarrow Ri_* i^* A \rightarrow, \quad (3.1)$$

$$\text{and } Ri_* Ri^! A \rightarrow A \rightarrow Rj_* j^* A \rightarrow \quad (3.2)$$

are distinguished triangles.

Proof. The proof of (3.1) is reduced to the case that $A \in D_{\text{ét}}(\text{Spd } \mathcal{O}_C, \Lambda)$ is concentrated in degree 0. The functors j^*, i^* are exact (that is, commute with canonical truncations) by definition. The functor $Rj_!$ is also exact (see [Sch17, Definition/Proposition 19.1]). Moreover, the functor Ri_* is exact. Hence it suffices to show that

$$0 \rightarrow j_! Rj^* A \rightarrow A \rightarrow Ri_* i^* A \rightarrow 0$$

is an exact sequence. This can be checked after taking i^* and j^* , and that is easy.

For (3.2), by (3.1) and Lemma 3.1, we have a distinguished triangle

$$R\mathcal{H}\text{om}(Ri_*i^*\Lambda, A) \rightarrow R\mathcal{H}\text{om}(\Lambda, A) \rightarrow R\mathcal{H}\text{om}(Rj_!j^*\Lambda, A) \rightarrow .$$

By [Sch17, Theorem 1.8(iv), (v)], we have

$$\begin{aligned} R\mathcal{H}\text{om}(Ri_*i^*\Lambda, A) &\cong Ri_*R\mathcal{H}\text{om}(i^*\Lambda, Ri^!A) \cong Ri_*Ri^!R\mathcal{H}\text{om}(\Lambda, A) \cong Ri_*Ri^!A \\ R\mathcal{H}\text{om}(\Lambda, A) &\cong A \\ R\mathcal{H}\text{om}(j_!Rj^*\Lambda, A) &\cong Rj_*R\mathcal{H}\text{om}(j^*\Lambda, Rj^!A) \cong Rj_*Rj^!R\mathcal{H}\text{om}(\Lambda, A) \cong Rj_*j^*A, \end{aligned}$$

hence the lemma follows. \square

3.1.2 Comparison to scheme

Let X be a scheme locally of finite type over $\text{Spec } \mathcal{O}_C$. Then we can define a diamond X^\diamond over $\text{Spd } \mathcal{O}_C$ as follows:

$$X^\diamond : \text{Perf} \rightarrow \text{Sets}$$

$$S \mapsto \left\{ (S^\#, f: S^\# \rightarrow X) \left| \begin{array}{l} S^\# \text{ is an untilt over } \text{Spa } \mathcal{O}_C = \text{Spa}(\mathcal{O}_C, \mathcal{O}_C). \\ f: S^\# \rightarrow X \text{ is a morphism of locally ringed} \\ \text{space over } \text{Spec } \mathcal{O}_C. \end{array} \right. \right\}.$$

The functor $X \mapsto X^\diamond$ defines a morphism of site

$$c_X : (X^\diamond)_v \rightarrow X_{\text{ét}}.$$

It defines a functor

$$c_X^* : D(X_{\text{ét}}, \Lambda) \rightarrow D_{\text{ét}}(X^\diamond, \Lambda).$$

It commute with colimits, and so admits a right adjoint Rc_{X^*} . We have the following proposition:

Proposition 3.3. *Let $f: Y \rightarrow X$ be a separated map between schemes of finite type over \mathcal{O}_C . Then we have*

$$\begin{aligned} (f^\diamond)^*c_X^* &\cong c_Y^*f^*, \\ Rf_*Rc_{Y^*} &\cong Rc_{X^*}Rf_*^\diamond. \end{aligned}$$

Proof. The first isomorphism follows from the definition. The second isomorphism is the adjunction of the first one. \square

Proposition 3.4. *Let $f: Y \rightarrow X$ be a separated map between schemes of finite type over \mathcal{O}_C . Then $f^\diamond: Y^\diamond \rightarrow X^\diamond$ is compactifiable and representable in locally spatial diamonds with $\dim.\text{trg}f < \infty$, and*

$$\begin{aligned} Rf_!^\diamond c_Y^* &\cong c_X^* Rf_!, \\ Rf^! R c_{X*} &\cong R c_{Y*} R(f^\diamond)!. \end{aligned}$$

Proof. The proof is the same as [Sch17, Proposition 27.5].

By Nagata's compactification, we can assume f is proper. Then f^\diamond is also proper and representable in spatial diamonds with $\dim.\text{trg}f < \infty$. We want to see that the natural transformation

$$c_X^* Rf_* \rightarrow Rf_*^\diamond c_Y^*$$

is an equivalence. Let $T \rightarrow \text{Spa } \mathcal{O}_C$ be a smooth surjective map of adic spaces such that T is analytic. Then $X \times_{\text{Spec } \mathcal{O}_C} T$ is an analytic adic space, and it holds that $(X \times_{\text{Spec } \mathcal{O}_C} T)^\diamond = X^\diamond \times_{\text{Spd } \mathcal{O}_C} T^\diamond$. Since $T^\diamond \rightarrow \text{Spd } \mathcal{O}_C$ is surjective and ℓ -cohomologically smooth for all $\ell \neq p$ by [Sch17, Proposition 24.4], the pullback functor $D_{\text{ét}}(X^\diamond, \Lambda) \rightarrow D_{\text{ét}}(X^\diamond \times_{\text{Spd } \mathcal{O}_C} T^\diamond, \Lambda)$ is conservative. Thus, we only have to show the result after taking the fiber product with T . Then the proposition follows from [Hub96, Theorem 3.7.2] and some base change results. \square

Proposition 3.5. *The three functors*

$$\begin{aligned} c_k &:= c_{\text{Spec } k}^*: D((\text{Spec } k)_{\text{ét}}, \Lambda) \rightarrow D_{\text{ét}}(\text{Spd } k, \Lambda), \\ c_{\mathcal{O}_C} &:= c_{\text{Spec } \mathcal{O}_C}^*: D((\text{Spec } \mathcal{O}_C)_{\text{ét}}, \Lambda) \rightarrow D_{\text{ét}}(\text{Spd } \mathcal{O}_C, \Lambda), \\ c_C &:= c_{\text{Spec } C}^*: D((\text{Spec } C)_{\text{ét}}, \Lambda) \rightarrow D_{\text{ét}}(\text{Spd } C, \Lambda) \end{aligned}$$

are isomorphisms.

Proof. For $\text{Spec } C$, the claim follows from a canonical equivalence

$$D_{\text{ét}}(\text{Spd } C, \Lambda) \cong D(|\text{Spd } C|, \Lambda) = D(\Lambda\text{-mod})$$

since $\text{Spd } C$ is strictly totally disconnected. For $\text{Spec } k$, it follows from the fact that the pullback functor

$$\begin{aligned} D_{\text{ét}}(\text{Spd } k, \Lambda) &= D_{\text{ét}}(\text{Spa } k, \Lambda) \\ &\rightarrow D_{\text{ét}}(\text{Spa } C^b, \Lambda) \\ &\simeq D(|\text{Spa } C^b|, \Lambda) \\ &\simeq D(\Lambda\text{-mod}) \end{aligned}$$

is fully faithful by [FS21, Theorem 19.5], and essentially surjective considering constant complexes. Let us consider the case of $\mathrm{Spec} \mathcal{O}_C$. First, the fully faithfulness is reduced to the fully faithfulness of

$$c_{\mathcal{O}_C}^* : D((\mathrm{Spec} \mathcal{O}_C)_{\acute{e}t}, \Lambda) \rightarrow D((\mathrm{Spd} \mathcal{O}_C)_v, \Lambda),$$

that is, the fully faithfulness of the exact functor

$$c_{\mathcal{O}_C}^* : \mathrm{Sh}((\mathrm{Spec} \mathcal{O}_C)_{\acute{e}t}, \Lambda) \rightarrow \mathrm{Sh}((\mathrm{Spd} \mathcal{O}_C)_v, \Lambda).$$

where $\mathrm{Sh}(-, \Lambda)$ means the category of sheaves of Λ -modules. The site $(\mathrm{Spec} \mathcal{O}_C)_{\acute{e}t}$ can be written explicitly, so one can check this directly.

Let $j : \mathrm{Spd} C \rightarrow \mathrm{Spd} \mathcal{O}_C$ and $i : \mathrm{Spd} k \rightarrow \mathrm{Spd} \mathcal{O}_C$ be inclusions. For $A \in D_{\acute{e}t}(\mathrm{Spd} \mathcal{O}_C, \Lambda)$, there exists a distinguished triangle

$$j_! j^* A \rightarrow A \rightarrow i_* i^* A \rightarrow j_! j^* A[1].$$

Hence the object of $D_{\acute{e}t}(\mathrm{Spd} \mathcal{O}_C, \Lambda)$ is determined by $B \in D_{\acute{e}t}(\mathrm{Spd} k, \Lambda)$, $C \in D_{\acute{e}t}(\mathrm{Spd} C, \Lambda)$ and a morphism $i_* B \rightarrow j_! C[1]$. Similarly, the object of $D(\mathrm{Spec} \mathcal{O}_C, \Lambda)$ is determined by $B' \in D((\mathrm{Spec} k)_{\acute{e}t}, \Lambda)$, $C' \in D((\mathrm{Spec} C)_{\acute{e}t}, \Lambda)$ and a morphism $i_* B' \rightarrow j_! C'[1]$.

Since i_* and $j_!$ are compatible with c_X^* by Proposition 3.4, the equivalence of $c_{\mathcal{O}_C}^*$ follows from the equivalence of c_k^* , c_C^* and the fully faithfulness of $c_{\mathcal{O}_C}^*$. \square

3.1.3 Pushforward of constant sheaves

Let $j : \mathrm{Spd} C \rightarrow \mathrm{Spd} \mathcal{O}_C$ and $i : \mathrm{Spd} k \rightarrow \mathrm{Spd} \mathcal{O}_C$ be inclusions.

Lemma 3.6. *If $K \in D_{\acute{e}t}(\mathrm{Spd} \mathcal{O}_C, \Lambda)$ is constant with perfect fiber, then the canonical morphism*

$$K \rightarrow Rj_* j^* K$$

is an isomorphism.

First, we show the scheme version of this lemma.

Lemma 3.7. *Let $j_C : \mathrm{Spec} C \rightarrow \mathrm{Spec} \mathcal{O}_C$ be an inclusion. The canonical morphism*

$$\Lambda \rightarrow Rj_* \Lambda$$

is an isomorphism.

Proof. Let $j_{\overline{F}}: \text{Spec } \overline{F} \rightarrow \text{Spec } \mathcal{O}_{\overline{F}}$ and $i_{\overline{F}}: \text{Spec } k \rightarrow \text{Spec } \mathcal{O}_{\overline{F}}$ be inclusions. Apply [Sta18, Lemma 0EYM] to the inverse system of morphisms of schemes $(j_E: \text{Spec } E \rightarrow \text{Spec } \mathcal{O}_E)_E$ and system of sheaves $(\Lambda_{\text{Spec } E})_E$ where E 's are finite extensions of F . Then we have

$$R^p j_{\overline{F}*} \Lambda \cong \text{colim } h_E^* R^p j_{E*} \Lambda$$

where $h_E: \text{Spec } \mathcal{O}_{\overline{F}} \rightarrow \text{Spec } \mathcal{O}_E$. By the calculation of Galois cohomologies, one has

$$i_{\overline{F}}^* h_E^* R^p j_{E*} \Lambda \cong \begin{cases} \Lambda & (p = 0) \\ \text{Hom}(I_E, \Lambda) & (p = 1) \\ 0 & (\text{otherwise}) \end{cases}$$

where I_E is the inertia group of E . Moreover, for $E_1 \subset E_2$, the map $i_{\overline{F}}^* h_{E_1}^* R^p j_{E_1*} \rightarrow i_{\overline{F}}^* h_{E_2}^* R^p j_{E_2*}$ coincides with a canonical homomorphism induced by the inclusion $I_{E_2} \subset I_{E_1}$. Therefore we have

$$i_{\overline{F}}^* R^p j_{\overline{F}*} \Lambda \cong \begin{cases} \Lambda & (p = 0) \\ 0 & (\text{otherwise}) \end{cases}.$$

That is, $i_{\overline{F}}^* R j_{\overline{F}*} \Lambda \cong \Lambda$. This isomorphism is the inverse of the canonical morphism $\Lambda = i_{\overline{F}}^* \Lambda \rightarrow i_{\overline{F}}^* R j_{\overline{F}*} j_{\overline{F}}^* \Lambda = i_{\overline{F}}^* R j_{\overline{F}*} \Lambda$. It follows that the canonical morphism

$$\Lambda \rightarrow R j_{\overline{F}*} \Lambda \tag{3.3}$$

is an isomorphism.

Finally, let $\alpha: \text{Spec } \widehat{\overline{F}} \rightarrow \text{Spec } \overline{F}$, and $\beta: \text{Spec } \mathcal{O}_{\widehat{\overline{F}}} \rightarrow \text{Spec } \mathcal{O}_{\overline{F}}$ be the natural morphisms. One can show that the induced morphism of sites

$$\begin{aligned} (\text{Spec } \widehat{\overline{F}})_{\text{ét}} &\rightarrow (\text{Spec } \overline{F})_{\text{ét}}, \\ (\text{Spec } \mathcal{O}_{\widehat{\overline{F}}})_{\text{ét}} &\rightarrow (\text{Spec } \mathcal{O}_{\overline{F}})_{\text{ét}} \end{aligned}$$

are isomorphism since these four site can be written explicitly. Therefore, $\alpha^*: D((\text{Spec } \overline{F})_{\text{ét}}, \Lambda) \rightarrow D((\text{Spec } \widehat{\overline{F}})_{\text{ét}}, \Lambda)$ and $\beta^*: D((\text{Spec } \mathcal{O}_{\overline{F}})_{\text{ét}}, \Lambda) \rightarrow D((\text{Spec } \mathcal{O}_{\widehat{\overline{F}}})_{\text{ét}}, \Lambda)$ are equivalences. It follows that the diagram

$$\begin{array}{ccc} D((\text{Spec } \overline{F})_{\text{ét}}, \Lambda) & \xrightarrow{j_{\overline{F}*}} & D((\text{Spec } \mathcal{O}_{\overline{F}})_{\text{ét}}, \Lambda) \\ \alpha^* \downarrow & & \downarrow \beta^* \\ D((\text{Spec } \widehat{\overline{F}})_{\text{ét}}, \Lambda) & \xrightarrow{j_{\widehat{\overline{F}}*}} & D((\text{Spec } \mathcal{O}_{\widehat{\overline{F}}})_{\text{ét}}, \Lambda) \end{array}$$

is commutative up to natural isomorphism since $\alpha^* \cong (\alpha_*)^{-1}$, $\beta^* \cong (\beta_*)^{-1}$. By this diagram and the isomorphism (3.3), we get the lemma. \square

Corollary 3.8. *If $K \in D((\mathrm{Spec} \mathcal{O}_C)_{\acute{\mathrm{e}}\mathrm{t}}, \Lambda)$ is constant with perfect fiber, then the canonical morphism*

$$K \rightarrow Rj_*j^*K$$

is an isomorphism.

Proof. Since the value of the constant complex K is perfect, it can be represented by a bounded complex (M^\bullet) whose terms are finite projective over Λ . Since Λ is a self-injective ring, each M^i is injective, Rj_*j^*K is represented by a complex (Rj_*M^\bullet) , which is isomorphic to a constant complex (M^\bullet) by Lemma 3.7. This proves the claim. \square

Proof of Lemma 3.6. By Corollary 3.8, it suffices to show that the diagram

$$\begin{array}{ccc} D((\mathrm{Spec} C)_{\acute{\mathrm{e}}\mathrm{t}}, \Lambda) & \xrightarrow{j_*} & D((\mathrm{Spec} \mathcal{O}_C)_{\acute{\mathrm{e}}\mathrm{t}}, \Lambda) \\ c_C^* \downarrow & & \downarrow c_{\mathcal{O}_C}^* \\ D_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spd} C, \Lambda) & \xrightarrow{j_*} & D_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spd} \mathcal{O}_C, \Lambda) \end{array}$$

is commutative up to a natural isomorphism. This follows from Proposition 3.5 since $c_C^* \cong (Rc_{C^*})^{-1}$ and $c_{\mathcal{O}_C}^* \cong (Rc_{\mathcal{O}_C^*})^{-1}$ holds. \square

3.2 Nearby cycle functor

Proposition 3.9. *$K \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} C}, \Lambda)^{\mathrm{bd}}$ is universally locally acyclic (ULA) over $\mathrm{Spd} C$, then $Rj_*K \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} \mathcal{O}_C}, \Lambda)^{\mathrm{bd}}$ is ULA over $\mathrm{Spd} \mathcal{O}_C$.*

Proof. Since we are working over $\mathrm{Spd} \mathcal{O}_C$, we may assume that G is split. Fix a maximal split torus $T \subset G$ and a Borel subgroup $B \subset G$ containing T . By [FS21, VI.6.4], for a small v-stack S over Div^1 and $A \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathcal{H}\mathrm{ck}_{G, S}, \Lambda)^{\mathrm{bd}}$, the complex A is ULA over S if and only if

$$\mathrm{CT}_B(A) \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathcal{H}\mathrm{ck}_{T, S}, \Lambda)^{\mathrm{bd}}$$

is ULA over S , where the functor CT_B is the hyperbolic localization. Since CT_B is compatible with Rj_* , we may assume $G = T$.

In this case, since $\mathcal{H}\mathrm{ck}_{T, S} = \mathrm{Gr}_{T, S} \cong X_*(T) \times S$ holds, we need to show that if $K \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spd} C, \Lambda)^{\mathrm{bd}}$ is ULA over $\mathrm{Spd} C$, then $Rj_*K \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spd} \mathcal{O}_C, \Lambda)^{\mathrm{bd}}$ is ULA over $\mathrm{Spd} \mathcal{O}_C$. By [FS21, IV.2.9], we have to show that if $K \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spd} C, \Lambda)^{\mathrm{bd}}$ is locally constant with perfect fibers, then $Rj_*K \in D_{\acute{\mathrm{e}}\mathrm{t}}(\mathrm{Spd} \mathcal{O}_C, \Lambda)^{\mathrm{bd}}$ is locally constant with perfect fibers. This follows from Lemma 3.6. \square

Theorem 3.10. *The pullback functor*

$$j^*: D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G, \text{Spd } \mathcal{O}_C}, \Lambda)^{\text{bd}} \rightarrow D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G, \text{Spd } C}, \Lambda)^{\text{bd}}$$

is an equivalence of categories, and its quasi-inverse is Rj_* .

Proof. The fact that j^* is an equivalence is proved in [FS21, Corollary VI.6.7]. By Proposition 3.9 and the identity $j^*Rj_* = \text{id}$, its quasi-inverse is Rj_* . \square

Corollary 3.11. *Consider the diagram*

$$\begin{array}{ccc} \mathcal{H}\text{ck}_{G, \text{Spd } k} & \xrightarrow{i} & \mathcal{H}\text{ck}_{G, \text{Spd } \mathcal{O}_{\widehat{F}}} \xleftarrow{j} \mathcal{H}\text{ck}_{G, \text{Spd } \widehat{F}} \\ & & \downarrow p_F \\ & & \mathcal{H}\text{ck}_{G, \text{Div}^1}. \end{array}$$

The functors $(p_F)^*$, Rj_* , i^* preserves the object of the Satake categories. Moreover, Rj_* and i^* induce equivalences

$$\begin{aligned} Rj_*: \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } C}, \Lambda) &\xrightarrow{\sim} \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } \mathcal{O}_C}, \Lambda), \\ i^*: \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } \mathcal{O}_C}, \Lambda) &\xrightarrow{\sim} \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } k}, \Lambda). \end{aligned}$$

Proof. Since we are considering the relative locally acyclicity and the relative perversity, $i^*, j^*, (p_F)^*$ preserves ULA objects and flat perverse (i.e. for all Λ -modules M , $A \otimes_{\Lambda}^{\mathbb{L}} M$ is perverse) objects.

Let us prove that Rj_* preserves the object of the Satake categories. Since we are working over $\text{Spd } \mathcal{O}_C$, we may assume that G is split. Fix a maximal split torus $T \subset G$ and a Borel subgroup $B \subset G$ containing T . The pushforward Rj_* commutes with $R\pi_{T, S*} \text{CT}_B(-)[\text{deg}]$ where $\pi_{T, S}: \text{Gr}_{T, S} \rightarrow S$. Hence by [FS21, Proposition VI.6.4, Proposition VI.7.7], we need to show that if $A \in D_{\text{ét}}(\text{Spd } C, \Lambda)$ is étale locally a finite projective Λ -module in degree 0, then so is $Rj_*A \in D_{\text{ét}}(\text{Spd } \mathcal{O}_C, \Lambda)$. This follows from Lemma 3.6.

Moreover, Rj_* induces an equivalence since j^* gives the quasi-inverse. It remains to show that i^* induces an equivalence. As in [FS21, Corollary VI.6.7],

$$i^*: D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{\text{Spd } \mathcal{O}_C}, \Lambda) \xrightarrow{\sim} D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{\text{Spd } k}, \Lambda)$$

is an equivalence. Thus, as above, using the hyperbolic localization, the result follows from the fact that for $A \in D_{\text{ét}}^{\text{ULA}}(\text{Spd } \mathcal{O}_C, \Lambda)$, the object $i^*A \in D_{\text{ét}}(\text{Spd } k, \Lambda)$ is étale locally a finite projective Λ -module in degree 0 if and only if so is $A \in D_{\text{ét}}(\text{Spd } \mathcal{O}_C, \Lambda)$. \square

Therefore, we have a functor

$$\Psi := i^* Rj_*(p_F)^* : \text{Sat}(\mathcal{H}\text{ck}_{\text{Div}^1}, \Lambda) \rightarrow \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } k}, \Lambda) \quad (3.4)$$

called a nearby cycle functor.

3.3 Nearby cycle functor and convolution

In this subsection, we prove that

Proposition 3.12. *Consider the nearby cycle functor*

$$\Psi := i^* Rj_*(p_F)^* : \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Div}^1}, \Lambda) \rightarrow \text{Sat}(\mathcal{H}\text{ck}_{G, \text{Spd } k}, \Lambda).$$

There is a natural isomorphism

$$\Psi(A \star B) \cong \Psi(A) \star \Psi(B)$$

for $A, B \in \text{Sat}(\mathcal{H}\text{ck}_{\text{Div}^1}, \Lambda)$ where $- \star -$ denotes the convolution product defined in [FS21, VI.8].

Apply Theorem 3.10 to a reductive group $G \times_{\mathcal{O}_F} G$, noting that there is an canonical isomorphism

$$\mathcal{H}\text{ck}_{G, S} \times_S \mathcal{H}\text{ck}_{G, S} \cong \mathcal{H}\text{ck}_{G \times_{\mathcal{O}_F} G, S}.$$

We have the following proposition:

Proposition 3.13. *Let $j_{G^2} : \mathcal{H}\text{ck}_{G, \text{Spd } C} \times_{\text{Spd } C} \mathcal{H}\text{ck}_{G, \text{Spd } C} \rightarrow \mathcal{H}\text{ck}_{G, \text{Spd } \mathcal{O}_C} \times_{\text{Spd } \mathcal{O}_C} \mathcal{H}\text{ck}_{G, \text{Spd } \mathcal{O}_C}$ be the natural inclusion. Then*

$$\begin{aligned} j_{G^2}^* : D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G, \text{Spd } \mathcal{O}_C} \times_{\text{Spd } \mathcal{O}_C} \mathcal{H}\text{ck}_{G, \text{Spd } \mathcal{O}_C}, \Lambda)^{\text{bd}} \\ \rightarrow D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G, \text{Spd } C} \times_{\text{Spd } C} \mathcal{H}\text{ck}_{G, \text{Spd } C}, \Lambda)^{\text{bd}} \end{aligned}$$

is an equivalence of categories, and its quasi-inverse is $Rj_{G^2 \star}$.

Proof of Proposition 3.12. Put

$$\widetilde{\mathcal{H}\text{ck}}_{G, \text{Div}^1} := L_{\text{Div}^1}^+ G \backslash L_{\text{Div}^1} G \times^{L_{\text{Div}^1}^+ G} L_{\text{Div}^1} G / L_{\text{Div}^1}^+ G,$$

and put $\widetilde{\mathcal{H}ck}_{G,S} := \widetilde{\mathcal{H}ck}_{G,\text{Div}^1} \times_{\text{Div}^1} S$ for a small v-stack $S \rightarrow \text{Div}^1$. Consider the diagram

$$\begin{array}{ccccc}
\mathcal{H}ck_{G,\text{Div}^1} \times_{\text{Div}^1} \mathcal{H}ck_{G,\text{Div}^1} & \xleftarrow{a_{\text{Div}^1}} & \widetilde{\mathcal{H}ck}_{G,\text{Div}^1} & \xrightarrow{b_{\text{Div}^1}} & \mathcal{H}ck_{G,\text{Div}^1} \\
\uparrow p_{F,G^2} & & \uparrow & & \uparrow p_F \\
\mathcal{H}ck_{G,\text{Spd} C} \times_{\text{Spd} C} \mathcal{H}ck_{G,\text{Spd} C} & \xleftarrow{a_C} & \widetilde{\mathcal{H}ck}_{G,\text{Spd} C} & \xrightarrow{b_C} & \mathcal{H}ck_{G,\text{Spd} C} \\
\downarrow j_{G^2} & & \downarrow & & \downarrow j \\
\mathcal{H}ck_{G,\text{Spd} \mathcal{O}_C} \times_{\text{Spd} \mathcal{O}_C} \mathcal{H}ck_{G,\text{Spd} \mathcal{O}_C} & \xleftarrow{a_{\mathcal{O}_C}} & \widetilde{\mathcal{H}ck}_{G,\text{Spd} \mathcal{O}_C} & \xrightarrow{b_{\mathcal{O}_C}} & \mathcal{H}ck_{G,\text{Spd} \mathcal{O}_C} \\
\uparrow i_{G^2} & & \uparrow & & \uparrow i \\
\mathcal{H}ck_{G,\text{Spd} k} \times_{\text{Spd} k} \mathcal{H}ck_{G,\text{Spd} k} & \xleftarrow{a_k} & \widetilde{\mathcal{H}ck}_{G,\text{Spd} k} & \xrightarrow{b_k} & \mathcal{H}ck_{G,\text{Spd} k}
\end{array}$$

Since $a_{\mathcal{O}_C}$ is an L^+G -torsor and $b_{\text{Div}^1}, b_{\mathcal{O}_C}$ is ind-proper, by using base changes, we have

$$R(b_k)_*(a_k)^*(i_{G^2})^*R(j_{G^2})_*(p_{F,G^2})^*(A \boxtimes_{\Lambda,\text{Div}^1} B) \cong \Psi(A \star B)$$

for $A, B \in D_{\text{ét}}(\mathcal{H}ck_{G,\text{Div}^1}, \Lambda)^{\text{bd}}$. It suffices to show that if $A, B \in D_{\text{ét}}(\mathcal{H}ck_{G,\text{Spd} C}, \Lambda)$ are ULA sheaves, then

$$R(j_{G^2})_*(A \boxtimes_{\Lambda,\text{Spd} C} B) \cong Rj_* A \boxtimes_{\Lambda,\text{Spd} \mathcal{O}_C} Rj_* B. \quad (3.5)$$

First, if $A, B \in D_{\text{ét}}(\mathcal{H}ck_{G,\text{Spd} C}, \Lambda)$ are ULA over $\text{Spd} C$, then

$$A \boxtimes_{\Lambda,\text{Spd} C} B \in D_{\text{ét}}(\mathcal{H}ck_{G,\text{Spd} C} \times_{\text{Spd} C} \mathcal{H}ck_{G,\text{Spd} C}, \Lambda)^{\text{bd}}$$

is ULA over $\text{Spd} C$. In fact, by [FS21, Proposition VI.6.5], it suffices to show that if $A_1, A_2 \in D_{\text{ét}}(\text{Spd} C, \Lambda)$ is locally constant with perfect fibers, then so is $A_1 \boxtimes_{\Lambda,\text{Spd} C} A_2 = A_1 \otimes_{\Lambda}^{\mathbb{L}} A_2 \in D_{\text{ét}}(\text{Spd} C, \Lambda)$. This is clear as $D_{\text{ét}}(\text{Spd} C, \Lambda) \simeq D(\Lambda\text{-mod})$.

Therefore, by Proposition 3.13, we only have to show (3.5) after taking $j_{G^2}^*$. Since pullbacks are compatible with exterior tensor products, we have

$$\begin{aligned}
j_{G^2}^* R(j_{G^2})_*(A \boxtimes_{\Lambda,\text{Spd} C} B) &\cong A \boxtimes_{\Lambda,\text{Spd} C} B \\
&\cong j^* Rj_* A \boxtimes_{\Lambda,\text{Spd} C} j^* Rj_* B \\
&\cong j_{G^2}^*(Rj_* A \boxtimes_{\Lambda,\text{Spd} \mathcal{O}_C} Rj_* B),
\end{aligned}$$

and the proposition follows. \square

3.4 Relation between the two geometric Satake via nearby cycle

Theorem 3.14. *Consider the diagram*

$$\begin{array}{ccc}
 \text{Sat}(\mathcal{H}\text{ck}_{\text{Div}^1}, \Lambda) & \xrightarrow{F_{\text{Div}^1}} & \text{Rep}(W_F, \Lambda) \\
 (p_F)^* \downarrow & & \downarrow \text{For} \\
 \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } C}, \Lambda) & \xrightarrow{F_{\text{Spd } C}} & \text{Proj}(\Lambda) \\
 Rj_* \downarrow & & \parallel \\
 \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } \mathcal{O}_C}, \Lambda) & \xrightarrow{F_{\text{Spd } \mathcal{O}_C}} & \text{Proj}(\Lambda) \\
 i^* \downarrow & & \parallel \\
 \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } k}, \Lambda) & \xrightarrow{F_{\text{Spd } k}} & \text{Proj}(\Lambda),
 \end{array} \tag{3.6}$$

where

$$\begin{aligned}
 F_S &:= F_{G,S}: D_{\text{ét}}(\mathcal{H}\text{ck}_{G,S}, \Lambda) \rightarrow \text{LocSys}(S, \Lambda), \\
 A &\mapsto \bigoplus_i \mathcal{H}^i(R\pi_{G,S*} A)
 \end{aligned}$$

for a small v -stack S over Div^1 or $\text{Spd } \mathcal{O}_E$. Here $\text{LocSys}(S, \Lambda)$ is the category of Λ -local systems on S . Then there are isomorphisms $\text{For} \circ F_{\text{Div}^1} \cong F_{\text{Spd } C} \circ (p_F)^*$ and $F_{\text{Spd } C} \cong F_{\text{Spd } k} \circ \Psi$.

Proof. We can prove this by applying the base change results to the cartesian squares

$$\begin{array}{ccccccc}
 \mathcal{H}\text{ck}_{\text{Div}^1} & \longleftarrow & \mathcal{H}\text{ck}_{\text{Spd } C} & \longrightarrow & \mathcal{H}\text{ck}_{\text{Spd } \mathcal{O}_C} & \longleftarrow & \mathcal{H}\text{ck}_{\text{Spd } k} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \text{Div}^1 & \longleftarrow & \text{Spd } C & \longrightarrow & \text{Spd } \mathcal{O}_C & \longleftarrow & \text{Spd } k.
 \end{array}$$

□

Corollary 3.15. *There exists a symmetric monoidal structure on the category*

$$(\text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } k}, \Lambda), \star)$$

and on the functors

$$\begin{aligned}
 i^* Rj_* &: \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } C}, \Lambda) \rightarrow \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } k}, \Lambda), \\
 F_{\text{Spd } k} &: \text{Sat}(\mathcal{H}\text{ck}_{\text{Spd } k}, \Lambda) \rightarrow \text{Proj}(\Lambda)
 \end{aligned}$$

such that the isomorphism

$$F_{\mathrm{Spd} k} \circ i^* Rj_* \cong F_{\mathrm{Spd} C}$$

in Theorem 3.14 is monoidal, with respect to the monoidal structure of $F_{\mathrm{Spd} C}$ using fusion products (see [FS21, VI.9]).

Proof. This follows from Theorem 3.14 and the fact that $i^* Rj_*: \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{\mathrm{Spd} C}, \Lambda) \rightarrow \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{\mathrm{Spd} k}, \Lambda)$ is an equivalence. \square

The main theorem follows from this corollary.

Theorem 3.16. *There exists a symmetric monoidal structure on the category*

$$(\mathrm{Sat}(\mathcal{H}\mathrm{ck}_{\mathrm{Spd} k}, \Lambda), \star)$$

and on the functor

$$F_{\mathrm{Spd} k}: \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{\mathrm{Spd} k}, \Lambda) \rightarrow \mathrm{Proj}(\Lambda)$$

which induce a symmetric monoidal equivalence

$$\mathcal{S}_{\mathrm{YZ}}: \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} k}, \Lambda) \xrightarrow{\sim} \mathrm{Rep}(\widehat{G}, \Lambda)$$

such that the squares

$$\begin{array}{ccc} \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Div}^1}, \Lambda) & \xrightarrow{\mathcal{S}_{\mathrm{FS}}} & \mathrm{Rep}(\widehat{G}^{\mathrm{tw}} \times W_F, \Lambda) \\ (p_F)^* \downarrow & & \downarrow \mathrm{For} \\ \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} C}, \Lambda) & \xrightarrow{\mathcal{S}_{\mathrm{FS}}} & \mathrm{Rep}(\widehat{G}, \Lambda) \\ i^* Rj_* \downarrow & & \parallel \\ \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} k}, \Lambda) & \xrightarrow{\mathcal{S}_{\mathrm{YZ}}} & \mathrm{Rep}(\widehat{G}, \Lambda) \end{array}$$

naturally commute.

Proof. Apply Tannakian theory to the monoidal natural isomorphism in Corollary 3.15. \square

3.5 Monoidal structure on hyperbolic localization functor

Let P be a parabolic subgroup of G . Let M denote its Levi quotient, and \overline{M} its maximal torus quotient. Define a locally constant function $\mathrm{deg}_P: \mathrm{Gr}_{M, \mathrm{Spd} k} \rightarrow \mathbb{Z}$ as

the composition

$$\begin{aligned} \mathrm{Gr}_{M, \mathrm{Spd} k} &\longrightarrow \mathrm{Gr}_{\overline{M}, \mathrm{Spd} k} \\ &\xrightarrow{\sim} X_*(\overline{M}) \\ &\xrightarrow{1 < 2\rho_G - 2\rho_M, -} \mathbb{Z} \end{aligned}$$

where ρ_G, ρ_M is a half sum of positive roots of G, M , respectively.

Theorem 3.17. *Consider the shifted hyperbolic localization functor*

$$\mathrm{CT}_P[\mathrm{deg}_P] := R(p_k^+)!(q_k^+)^*[\mathrm{deg}_P]: D_{\mathrm{ét}}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} k}, \Lambda)^{\mathrm{bd}} \rightarrow D_{\mathrm{ét}}(\mathcal{H}\mathrm{ck}_{M, \mathrm{Spd} k}, \Lambda)^{\mathrm{bd}}$$

where $p_k^+: \mathrm{Gr}_{P, \mathrm{Spd} k} \rightarrow \mathrm{Gr}_{M, \mathrm{Spd} k}$, $q_k^+: \mathrm{Gr}_{P, \mathrm{Spd} k} \rightarrow \mathrm{Gr}_{G, \mathrm{Spd} k}$ are natural maps. Then $\mathrm{CT}_P[\mathrm{deg}_P]$ induces the functor

$$\mathrm{CT}_P[\mathrm{deg}_P]: \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} k}, \Lambda) \rightarrow \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{P, \mathrm{Spd} k}, \Lambda)$$

and there is a unique symmetric monoidal structure on $\mathrm{CT}_P[\mathrm{deg}_P]$ such that the natural isomorphism

$$F_{G, \mathrm{Spd} k} \cong F_{M, \mathrm{Spd} k} \circ \mathrm{CT}_P[\mathrm{deg}_P]: \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} k}, \Lambda) \rightarrow \mathrm{Proj}(\Lambda) \quad (3.7)$$

is monoidal. Here the symmetric monoidal structures on the categories $\mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} k}, \Lambda)$, $\mathrm{Sat}(\mathcal{H}\mathrm{ck}_{M, \mathrm{Spd} k}, \Lambda)$ and the functors $F_{G, \mathrm{Spd} k}, F_{M, \mathrm{Spd} k}$ are as in Corollary 3.15.

Proof. By the same argument as [FS21, Proposition VI.9.6], there is a natural symmetric monoidal structure on the functor

$$\mathrm{CT}_P[\mathrm{deg}_P]: \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} C}, \Lambda) \rightarrow \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{P, \mathrm{Spd} C}, \Lambda)$$

such that such that the natural isomorphism

$$F_{G, \mathrm{Spd} C} \cong F_{M, \mathrm{Spd} C} \circ \mathrm{CT}_P[\mathrm{deg}_P]: \mathrm{Sat}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} C}, \Lambda) \rightarrow \mathrm{Proj}(\Lambda)$$

is monoidal. The following square is naturally commutative

$$\begin{array}{ccc} D_{\mathrm{ét}}^{\mathrm{ULA}}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} C}, \Lambda) & \xrightarrow{\mathrm{CT}_P[\mathrm{deg}_P]} & D_{\mathrm{ét}}^{\mathrm{ULA}}(\mathcal{H}\mathrm{ck}_{M, \mathrm{Spd} C}, \Lambda) \\ i^*Rj_* \downarrow & & i^*Rj_* \downarrow \\ D_{\mathrm{ét}}^{\mathrm{ULA}}(\mathcal{H}\mathrm{ck}_{G, \mathrm{Spd} k}, \Lambda) & \xrightarrow{\mathrm{CT}_P[\mathrm{deg}_P]} & D_{\mathrm{ét}}^{\mathrm{ULA}}(\mathcal{H}\mathrm{ck}_{M, \mathrm{Spd} k}, \Lambda), \end{array}$$

and the vertical arrows i^*Rj_* induce equivalences of the Satake categories. By this diagram, we can endow the functor $\mathrm{CT}_P[\mathrm{deg}_P]$ with a symmetric monoidal structure. The isomorphism (3.7) is monoidal by definition. The uniqueness of such monoidal structure follows from the fact that $F_{G, \mathrm{Spd} k}$ is faithful. \square

4 Integral coefficient case

In this subsection, Λ is the ring of integers of a finite extension of \mathbb{Q}_ℓ . Let $\lambda \in \Lambda$ be a uniformizer. First, recall the definition of $D_{\text{ét}}(-, \Lambda)$.

Definition 4.1. ([Sch17, Definition 26.1]) *For any small v -stack Y , define*

$$D_{\text{ét}}(Y, \Lambda) \subset D(Y_v, \Lambda)$$

as the full subcategory of all $A \in D(Y_v, \Lambda)$ such that A is derived (λ) -complete, and $A \otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda$ lies in $D_{\text{ét}}(Y, \Lambda/\lambda)$.

Lemma 4.2. ([Sch17, Remark 26.3])

(i) *For any map $f: X \rightarrow Y$ of small v -stacks, the following squares are commutative:*

$$\begin{array}{ccc} D_{\text{ét}}(Y, \Lambda) & \xrightarrow{f^*} & D_{\text{ét}}(X, \Lambda) \\ -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow & & -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow \\ D_{\text{ét}}(Y, \Lambda/\lambda^n) & \xrightarrow{f^*} & D_{\text{ét}}(X, \Lambda/\lambda^n), \end{array} \quad \begin{array}{ccc} D_{\text{ét}}(X, \Lambda) & \xrightarrow{Rf_*} & D_{\text{ét}}(Y, \Lambda) \\ -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow & & -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow \\ D_{\text{ét}}(X, \Lambda/\lambda^n) & \xrightarrow{Rf_*} & D_{\text{ét}}(Y, \Lambda/\lambda^n). \end{array}$$

(ii) *Let $f: X \rightarrow Y$ be a map of small v -stacks which is compactifiable, representable in locally spatial diamonds and with $\dim.\text{trg}f < \infty$. The following squares are commutative:*

$$\begin{array}{ccc} D_{\text{ét}}(X, \Lambda) & \xrightarrow{Rf_!} & D_{\text{ét}}(Y, \Lambda) \\ -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow & & -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow \\ D_{\text{ét}}(X, \Lambda/\lambda^n) & \xrightarrow{Rf_!} & D_{\text{ét}}(Y, \Lambda/\lambda^n), \end{array} \quad \begin{array}{ccc} D_{\text{ét}}(Y, \Lambda) & \xrightarrow{Rf^!} & D_{\text{ét}}(X, \Lambda) \\ -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow & & -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow \\ D_{\text{ét}}(Y, \Lambda/\lambda^n) & \xrightarrow{Rf^!} & D_{\text{ét}}(X, \Lambda/\lambda^n). \end{array}$$

Proof. See [Sch17, Remark 26.3]. □

Also, the definition of universal locally acyclicity is as follows:

Definition 4.3. ([FS21, VII.5]) *Let $f: X \rightarrow S$ be a compactifiable map of small v -stacks representable by locally spatial diamonds with locally $\dim.\text{trg}f < \infty$. Then we define the category of ULA complexes $D_{\text{ét}}^{\text{ULA}}(X/S, \Lambda)$ by $\varprojlim_n D_{\text{ét}}^{\text{ULA}}(X/S, \Lambda/\lambda^n)$.*

Definition 4.4. *We say that $A \in D_{\text{ét}}(\mathcal{H}ck_{G,S}, \Lambda)$ is ULA if its pullback to $\text{Gr}_{G,S}$ is ULA. Write $D_{\text{ét}}^{\text{ULA}}(\mathcal{H}ck_{G,S}, \Lambda)$ for the full subcategory of $D_{\text{ét}}(\mathcal{H}ck_{G,S}, \Lambda)$ consisting of ULA objects.*

Lemma 4.5. *Let S be a small v -stack over Div^1 . For an object $A \in D_{\acute{e}t}(\mathcal{H}ck_{G,S}, \Lambda)$, the following conditions are equivalent:*

- (i) $A \in D_{\acute{e}t}^{\mathrm{ULA}}(\mathcal{H}ck_{G,S}, \Lambda)$.
- (ii) $A \otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \in D_{\acute{e}t}^{\mathrm{ULA}}(\mathcal{H}ck_{G,S}, \Lambda/\lambda^n)$ for all n .

Proof. The square

$$\begin{array}{ccc} D_{\acute{e}t}(\mathcal{H}ck_{G,S}, \Lambda) & \xrightarrow{(-)^*} & D_{\acute{e}t}(\mathrm{Gr}_{G,S}, \Lambda) \\ -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \downarrow & & \downarrow -\otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \\ D_{\acute{e}t}(\mathcal{H}ck_{G,S}, \Lambda/\lambda^n) & \xrightarrow{(-)^*} & D_{\acute{e}t}(\mathrm{Gr}_{G,S}, \Lambda/\lambda^n) \end{array}$$

is commutative. For $A \in D_{\acute{e}t}(\mathcal{H}ck_{G,S}, \Lambda)$, the object π^*A is ULA if and only if $(\pi^*A) \otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n \cong \pi^*(A \otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n)$ is ULA for all n . Hence A is ULA if and only if $A \otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n$ is ULA for all n . \square

This equivalence implies that the essential image of $D_{\acute{e}t}^{\mathrm{ULA}}(\mathcal{H}ck_{G,S}, \Lambda)$ under the categorical equivalence

$$D_{\acute{e}t}(\mathcal{H}ck_{G,S}, \Lambda) \cong \varprojlim_n D_{\acute{e}t}(\mathcal{H}ck_{G,S}, \Lambda/\lambda^n)$$

in [Sch17, Proposition 26.2] is $\varprojlim_n D^{\mathrm{ULA}/S}(\mathcal{H}ck_{G,S}, \Lambda/\lambda^n)$. Also, we define the flat perversity as follows.

Definition 4.6. *For $A \in D_{\acute{e}t}^{\mathrm{ULA}}(\mathcal{H}ck_{G,S}, \Lambda)$, we say that A is flat perverse over Λ if and only if $A \otimes_{\Lambda}^{\mathbb{L}} \Lambda/\lambda^n$ is flat perverse over Λ/λ^n for all n .*

From this definition, we have

$$\mathrm{Sat}(\mathcal{H}ck_{G,S}, \Lambda) = \varprojlim_n \mathrm{Sat}(\mathcal{H}ck_{G,S}, \Lambda/\lambda^n).$$

By passing to the limit of the results in §3.2, 3.3, 3.4, 3.5 and using Lemma 4.2, we get the following theorem:

Theorem 4.7. *Theorem 3.10, Corollary 3.11, Proposition 3.12, Theorem 3.16 and Theorem 3.17 hold even if Λ is the ring of integers in a finite extension over \mathbb{Q}_{ℓ} .*

5 Fractional coefficient case

Moreover, we can prove a similar result for $\Lambda[\ell^{-1}]$ -coefficient.

Definition 5.1. *Let X be a small v-stack. The category $D_{\text{ét}}(X, \Lambda[\ell^{-1}])$ is defined as the category*

$$D_{\text{ét}}(X, \Lambda)[\ell^{-1}]$$

obtained by inverting ℓ in the Hom-sets of $D_{\text{ét}}(X, \Lambda)$. There is a natural functor

$$- \otimes_{\Lambda} \Lambda[\ell^{-1}]: D_{\text{ét}}(X, \Lambda) \rightarrow D_{\text{ét}}(X, \Lambda[\ell^{-1}]).$$

For any map $f: X \rightarrow Y$, the adjoint pair (f^*, Rf_*) with coefficients in Λ induces the adjoint pair

$$\begin{aligned} f^*: D_{\text{ét}}(Y, \Lambda[\ell^{-1}]) &\rightarrow D_{\text{ét}}(X, \Lambda[\ell^{-1}]), \\ Rf_*: D_{\text{ét}}(X, \Lambda[\ell^{-1}]) &\rightarrow D_{\text{ét}}(Y, \Lambda[\ell^{-1}]). \end{aligned}$$

Let $f: X \rightarrow Y$ be a map of small v-stacks which is compactifiable, representable in locally spatial diamonds and with $\dim.\text{trg}f < \infty$. The the adjoint pair $(Rf_!, Rf^!)$ with coefficients in Λ induces the adjoint pair

$$\begin{aligned} Rf_!: D_{\text{ét}}(X, \Lambda[\ell^{-1}]) &\rightarrow D_{\text{ét}}(Y, \Lambda[\ell^{-1}]), \\ Rf^!: D_{\text{ét}}(Y, \Lambda[\ell^{-1}]) &\rightarrow D_{\text{ét}}(X, \Lambda[\ell^{-1}]). \end{aligned}$$

Definition 5.2. *We say that $A \in D_{\text{ét}}(\mathcal{H}\text{ck}_{G,S}, \Lambda[\ell^{-1}])$ is universal locally acyclic (ULA) if A is in the essential image of $D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G,S}, \Lambda)$ under the functor*

$$- \otimes_{\Lambda} \Lambda[\ell^{-1}]: D_{\text{ét}}(\mathcal{H}\text{ck}_{G,S}, \Lambda) \rightarrow D_{\text{ét}}(\mathcal{H}\text{ck}_{G,S}, \Lambda[\ell^{-1}]).$$

Write $D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G,S}, \Lambda[\ell^{-1}])$ for this essential image.

By definition, $D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G,\text{Spd}C}, \Lambda[\ell^{-1}])$ is equivalent to the category

$$D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G,S}, \Lambda)[\ell^{-1}]$$

Also, we define the (flat) perversity as follows.

Definition 5.3. *For $A \in D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G,S}, \Lambda[\ell^{-1}])$, we say that A is perverse over $\Lambda[\ell^{-1}]$ if and only if A is isomorphic to an image of a flat perverse sheaf in $D_{\text{ét}}^{\text{ULA}}(\mathcal{H}\text{ck}_{G,S}, \Lambda)$ under the functor $- \otimes_{\Lambda}^{\mathbb{L}} \Lambda[\ell^{-1}]$.*

From this definition, we have

$$\text{Sat}(\mathcal{H}\text{ck}_{G,S}, \Lambda[\ell^{-1}]) \simeq \text{Sat}(\mathcal{H}\text{ck}_{G,S}, \Lambda)[\ell^{-1}].$$

By inverting ℓ in Theorem 4.7, we get the following theorem:

Theorem 5.4. *Theorem 3.10, Corollary 3.11, Proposition 3.12, Theorem 3.16 and Theorem 3.17 hold even if Λ is a finite extension over \mathbb{Q}_ℓ .*

6 $\overline{\mathbb{Q}}_\ell$ -coefficient case

We can prove the result for the $\overline{\mathbb{Q}}_\ell$ -coefficient case.

Definition 6.1. *Let X be a small v -stack. $D_{\text{ét}}(X, \overline{\mathbb{Q}}_\ell)$ is defined by*

$$D_{\text{ét}}(X, \overline{\mathbb{Q}}_\ell) = \varinjlim_L D_{\text{ét}}(X, L)$$

where L is a finite extension over \mathbb{Q}_ℓ , and if $L/L'/\mathbb{Q}_\ell$ is a tower of finite extension, the transition functor is the functor

$$- \otimes_{L'} L: D_{\text{ét}}(X, L') \rightarrow D_{\text{ét}}(X, L)$$

induced by

$$- \otimes_{\mathcal{O}_{L'}} \mathcal{O}_L: D_{\text{ét}}(X, \mathcal{O}_{L'}) \rightarrow D_{\text{ét}}(X, \mathcal{O}_L).$$

There is a natural functor

$$- \otimes_L \overline{\mathbb{Q}}_\ell: D_{\text{ét}}(X, L) \rightarrow D_{\text{ét}}(X, \overline{\mathbb{Q}}_\ell).$$

For any map $X \rightarrow Y$ of small v -stacks, the adjoint pairs (f^*, Rf_*) with coefficient in a finite extension L over \mathbb{Q}_ℓ induce the adjoint pair

$$\begin{aligned} f^*: D_{\text{ét}}(Y, \overline{\mathbb{Q}}_\ell) &\rightarrow D_{\text{ét}}(X, \overline{\mathbb{Q}}_\ell), \\ Rf_*: D_{\text{ét}}(X, \overline{\mathbb{Q}}_\ell) &\rightarrow D_{\text{ét}}(Y, \overline{\mathbb{Q}}_\ell). \end{aligned}$$

In fact, the following lemma holds:

Lemma 6.2. *Let $L/L'/\mathbb{Q}_\ell$ be a tower of finite extension.*

(i) *For any map $f: X \rightarrow Y$ of small v -stacks, the following squares are commutative:*

$$\begin{array}{ccc} D_{\text{ét}}(Y, L') & \xrightarrow{f^*} & D_{\text{ét}}(X, L') & \xrightarrow{Rf_*} & D_{\text{ét}}(Y, L') \\ - \otimes_{L'} L \downarrow & & - \otimes_{L'} L \downarrow & & - \otimes_{L'} L \downarrow \\ D_{\text{ét}}(Y, L) & \xrightarrow{f^*} & D_{\text{ét}}(X, L) & \xrightarrow{Rf_*} & D_{\text{ét}}(Y, L). \end{array}$$

(ii) Let $f: X \rightarrow Y$ be a map of small v -stacks which is compactifiable, representable in locally spatial diamonds and with $\dim.\mathrm{tr}gf < \infty$. The following squares are commutative:

$$\begin{array}{ccc} D_{\acute{e}t}(X, L') & \xrightarrow{Rf_!} & D_{\acute{e}t}(Y, L') & & D_{\acute{e}t}(Y, L') & \xrightarrow{Rf^!} & D_{\acute{e}t}(X, L') \\ -\otimes_{L'} L \downarrow & & -\otimes_{L'} L \downarrow & & -\otimes_{L'} L \downarrow & & -\otimes_{L'} L \downarrow \\ D_{\acute{e}t}(X, L) & \xrightarrow{Rf_!} & D_{\acute{e}t}(Y, L) & & D_{\acute{e}t}(Y, L) & \xrightarrow{Rf^!} & D_{\acute{e}t}(X, L). \end{array}$$

Proof. For (i), we need to show that the squares

$$\begin{array}{ccc} D_{\acute{e}t}(Y, \mathcal{O}_{L'}) & \xrightarrow{j^*} & D_{\acute{e}t}(X, \mathcal{O}_{L'}) & & D_{\acute{e}t}(X, \mathcal{O}_{L'}) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_{L'}) \\ -\otimes_{\mathcal{O}_{L'} \mathcal{O}_L} \downarrow & & -\otimes_{\mathcal{O}_{L'} \mathcal{O}_L} \downarrow & & -\otimes_{\mathcal{O}_{L'} \mathcal{O}_L} \downarrow & & -\otimes_{\mathcal{O}_{L'} \mathcal{O}_L} \downarrow \\ D_{\acute{e}t}(Y, \mathcal{O}_L) & \xrightarrow{j^*} & D_{\acute{e}t}(X, \mathcal{O}_L) & & D_{\acute{e}t}(X, \mathcal{O}_L) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_L) \end{array}$$

are naturally commutative. By Lemma 4.2, it is enough to prove that the squares

$$\begin{array}{ccc} D_{\acute{e}t}(Y, \mathcal{O}_{L'}/\ell^n) & \xrightarrow{j^*} & D_{\acute{e}t}(X, \mathcal{O}_{L'}/\ell^n) & & D_{\acute{e}t}(X, \mathcal{O}_{L'}/\ell^n) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_{L'}/\ell^n) \\ -\otimes_{\mathcal{O}_{L'}/\ell^n \mathcal{O}_L/\ell^n} \downarrow & & -\otimes_{\mathcal{O}_{L'}/\ell^n \mathcal{O}_L/\ell^n} \downarrow & & -\otimes_{\mathcal{O}_{L'}/\ell^n \mathcal{O}_L/\ell^n} \downarrow & & -\otimes_{\mathcal{O}_{L'}/\ell^n \mathcal{O}_L/\ell^n} \downarrow \\ D_{\acute{e}t}(Y, \mathcal{O}_L/\ell^n) & \xrightarrow{j^*} & D_{\acute{e}t}(X, \mathcal{O}_L/\ell^n) & & D_{\acute{e}t}(X, \mathcal{O}_L/\ell^n) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_L/\ell^n) \end{array}$$

are naturally commutative. The commutativity of the first square follows from the definition. For the second square, since \mathcal{O}_L/ℓ^n is a finite free module over $\mathcal{O}_{L'}/\ell^n$, the outer square in the diagram

$$\begin{array}{ccc} D_{\acute{e}t}(X, \mathcal{O}_{L'}/\ell^n) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_{L'}/\ell^n) \\ -\otimes_{\mathcal{O}_{L'}/\ell^n \mathcal{O}_L/\ell^n} \downarrow & & -\otimes_{\mathcal{O}_{L'}/\ell^n \mathcal{O}_L/\ell^n} \downarrow \\ D_{\acute{e}t}(X, \mathcal{O}_L/\ell^n) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_L/\ell^n) \\ \text{For} \downarrow & & \downarrow \text{For} \\ D_{\acute{e}t}(X, \mathcal{O}_{L'}/\ell^n) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_{L'}/\ell^n) \end{array}$$

is commutative, where the vertical functors For are forgetful functors. As the functor

$$\text{For}: D_{\acute{e}t}(Y, \mathcal{O}_L/\ell^n) \rightarrow D_{\acute{e}t}(Y, \mathcal{O}_{L'}/\ell^n)$$

is conservative, it suffices to show that the following square is naturally commutative:

$$\begin{array}{ccc} D_{\acute{e}t}(X, \mathcal{O}_L/\ell^n) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_L/\ell^n) \\ \text{For} \downarrow & & \downarrow \text{For} \\ D_{\acute{e}t}(X, \mathcal{O}_{L'}/\ell^n) & \xrightarrow{Rj_*} & D_{\acute{e}t}(Y, \mathcal{O}_{L'}/\ell^n). \end{array}$$

This square is the right adjoint of the first square. The proof of (ii) is similar. \square

Definition 6.3. We say that $A \in D_{\text{ét}}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell)$ is universal locally acyclic (ULA) if there exists a finite extension L over \mathbb{Q}_ℓ such that A is in the essential image of $D_{\text{ét}}^{\text{ULA}}(\mathcal{Hck}_{G,S}, L)$ under the functor

$$- \otimes_L \overline{\mathbb{Q}}_\ell: D_{\text{ét}}(\mathcal{Hck}_{G,S}, L) \rightarrow D_{\text{ét}}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell).$$

Write $D_{\text{ét}}^{\text{ULA}}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell)$ for the full subcategory of ULA sheaves in $D_{\text{ét}}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell)$.

By definition, it holds that

$$D_{\text{ét}}^{\text{ULA}}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell) = \varinjlim_L D_{\text{ét}}^{\text{ULA}}(\mathcal{Hck}_{G,S}, L).$$

We define the perversity as follows.

Definition 6.4. We say that $A \in D_{\text{ét}}^{\text{ULA}}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell)$ is perverse if there exists a finite extension L over \mathbb{Q}_ℓ such that A is in the essential image of $\text{Sat}(\mathcal{Hck}_{G,S}, L)$ under the functor

$$- \otimes_L \overline{\mathbb{Q}}_\ell: D_{\text{ét}}^{\text{ULA}}(\mathcal{Hck}_{G,S}, L) \rightarrow D_{\text{ét}}^{\text{ULA}}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell).$$

By definition, it holds that

$$\text{Sat}(\mathcal{Hck}_{G,S}, \overline{\mathbb{Q}}_\ell) = \varinjlim_L \text{Sat}(\mathcal{Hck}_{G,S}, L).$$

By passing the limit in Theorem 5.4 and using Lemma 6.2, we get the following theorem:

Theorem 6.5. *Theorem 3.10, Corollary 3.11, Proposition 3.12, Theorem 3.16 and Theorem 3.17 hold even if $\Lambda = \overline{\mathbb{Q}}_\ell$.*

References

- [AGLR22] J. Anschütz, I. Gleason, J. Lourenço and T. Richarz, On the p-adic theory of local models, 2022, 2201.01234.
- [FS21] L. Fargues and P. Scholze, Geometrization of the local Langlands correspondence, 2021, 2102.13459.
- [Hub96] R. Huber, Étale cohomology of rigid analytic varieties and adic spaces, Aspects of Mathematics, E30, Friedr. Vieweg & Sohn, Braunschweig, 1996.

- [Sch17] P. Scholze, Etale cohomology of diamonds, <http://www.math.uni-bonn.de/people/scholze/EtCohDiamonds.pdf>, 2017.
- [Sta18] T. Stacks Project Authors, *Stacks Project*, <https://stacks.math.columbia.edu>, 2018.
- [Yu19] J. Yu, The integral geometric Satake equivalence in mixed characteristic, 2019, 1903.11132.
- [Zhu17] X. Zhu, Affine Grassmannians and the geometric Satake in mixed characteristic, *Ann. of Math. (2)* 185 (2017), no. 2, 403–492.