

OBSTRUCTION THEORY AND THE LEVEL n ELLIPTIC GENUS

ANDREW SENGER

ABSTRACT. Given a height ≤ 2 Landweber exact \mathbb{E}_∞ -ring E whose homotopy is concentrated in even degrees, we show that any complex orientation of E which satisfies the Ando criterion admits a unique lift to an \mathbb{E}_∞ -complex orientation $\mathrm{MU} \rightarrow E$. As a consequence, we give a short proof that the level n elliptic genus lifts uniquely to an \mathbb{E}_∞ -complex orientation $\mathrm{MU} \rightarrow \mathrm{tmf}_1(n)$ for all $n \geq 2$.

CONTENTS

1. Introduction	1
2. Existence of \mathbb{E}_∞ -Orientations	4
3. Uniqueness of \mathbb{E}_∞ -Orientations	8
4. The level n elliptic genus	9
Appendix A. $K(n)$ -localizations of Landweber exact ring spectra	17
References	17

1. INTRODUCTION

Complex-oriented ring spectra play a central role in the chromatic approach to homotopy theory. Given a homotopy associative ring spectrum E , recall that a complex orientation is a choice of class $u \in \tilde{E}^2(\mathbb{C}\mathbb{P}^\infty)$ with the property that its restriction along $S^2 \cong \mathbb{C}\mathbb{P}^1 \hookrightarrow \mathbb{C}\mathbb{P}^\infty$ is the unit $1 \in E^0(*) \cong \tilde{E}^2(S^2)$. A complex orientation determines an isomorphism of graded rings

$$E^*(\mathbb{C}\mathbb{P}^\infty) \cong E^*[[u]].$$

Complex orientations may also be described in terms of the complex cobordism spectrum MU : complex orientations of E are in natural bijection with maps of homotopy associative ring spectra $\mathrm{MU} \rightarrow E$.

The complex cobordism spectrum MU admits much more structure than that of a homotopy associative ring spectrum: it is an \mathbb{E}_∞ -ring spectrum. When E also admits the structure of an \mathbb{E}_∞ -ring spectrum, it is natural to ask whether a given complex orientation is induced by a map of \mathbb{E}_∞ -ring spectra

$$\mathrm{MU} \rightarrow E.$$

We will call such maps \mathbb{E}_∞ -complex orientations. The \mathbb{E}_∞ -complex orientations of an \mathbb{E}_∞ -ring may be viewed as particularly canonical complex orientations.

Indeed, many of the most familiar (not necessarily complex) orientations admit lifts to \mathbb{E}_∞ -orientations. For example, the Atiyah–Bott–Shapiro orientations [ABS64]

$$\mathrm{MSpin} \rightarrow \mathrm{ko} \quad \text{and} \quad \mathrm{MSpin}^{\mathbb{C}} \rightarrow \mathrm{ku}.$$

were refined to \mathbb{E}_∞ -orientations by Joachim [Joa04], who gave an explicit geometric construction of such an \mathbb{E}_∞ -orientation. Indeed, one expects that any orientation of

geometric origin may, with enough care, be refined to an \mathbb{E}_∞ -orientation. A more sophisticated example is the Ando–Hopkins–Rezk \mathbb{E}_∞ -String orientation of the connective \mathbb{E}_∞ -ring of topological modular forms tmf [AHR10]

$$\mathrm{MString} \rightarrow \mathrm{tmf},$$

which refines the Witten genus [Wit87, Wit88]. While it is expected that this \mathbb{E}_∞ -orientation has a geometric origin, and much work has gone into developing such a viewpoint (for a small sampling, see [ST04, ST11, Cos10, Cos11, DH11, BE21]), such a description has so far remained elusive. In Theorem 1.7 below, we will prove that the Hirzebruch level n elliptic genera [Hir88][Wit88, §5] for $n \geq 2$ may be lifted to \mathbb{E}_∞ -complex orientations

$$\mathrm{MU} \rightarrow \mathrm{tmf}_1(n).$$

1.1. The Ando criterion. An algebraic approximation of what it means for a complex orientation $\mathrm{MU} \rightarrow E$ to be \mathbb{E}_∞ is given by the *Ando criterion*. It asks that the complex orientation be compatible in a suitable sense with the power operations in E . In many cases, the Ando criterion is equivalent to the property that the complex orientation $\mathrm{MU} \rightarrow E$ be a map of \mathbb{H}_∞ -ring spectra.

Now let E denote an \mathbb{E}_∞ , or more generally an \mathbb{H}_∞ -ring spectrum, with a fixed complex orientation u .

Notation 1.1. Given a complex vector bundle $V \rightarrow X$ of dimension d , we let $t_u(V) \in E^{2d}(\mathrm{Th}(V))$ denote the Thom class of V .

Fix a prime p , let ρ denote the vector bundle over BC_p corresponding to the complex regular representation of C_p , and let γ_1 denote the tautological bundle over $\mathbb{C}\mathbb{P}^\infty$. Let $I_{tr} \subset E^*(\mathrm{BC}_p)$ denote the transfer ideal. Recall from [HL18, §7] that there are additive power operations

$$\Psi_u : E^{2*}(\mathrm{Th}(\gamma_1)) \rightarrow E^{2p*}(\mathrm{Th}(\rho \boxtimes \gamma_1))/I_{tr}.$$

Definition 1.2. We say that a complex orientation of E satisfies the Ando criterion at the prime p if

$$\Psi_u(t_u(\gamma_1)) = t_u(\rho \boxtimes \gamma_1)$$

in $E^{2p}(\mathrm{Th}(\rho \boxtimes \gamma_1))/I_{tr}$. We say that a complex orientation of E satisfies the Ando criterion if it satisfies the Ando criterion for all primes p .

Remark 1.3. If E is p -local, then $E^*(\mathrm{BC}_\ell)/I_{tr} = 0$ for all primes $\ell \neq p$. Therefore a complex orientation of a p -local \mathbb{E}_∞ -ring satisfies the Ando criterion if and only if it satisfies the Ando criterion at p .

The complex cobordism spectrum MU , equipped with the canonical complex orientation, satisfies the Ando criterion. It follows that any \mathbb{E}_∞ -complex orientation, or more generally any \mathbb{H}_∞ -complex orientation, satisfies the Ando criterion.

1.2. Results. The first main theorem of this paper states that for many \mathbb{E}_∞ -rings of height ≤ 2 , this condition is sufficient, and that the resulting \mathbb{E}_∞ -complex orientations are determined up to homotopy by their underlying complex orientations:

Theorem 1.4. *Let E denote a height ≤ 2 Landweber exact \mathbb{E}_∞ -ring whose homotopy is concentrated in even degrees. Then any complex orientation $\mathrm{MU} \rightarrow E$ which satisfies the Ando criterion lifts uniquely up to homotopy to an \mathbb{E}_∞ -ring map $\mathrm{MU} \rightarrow E$:*

In the above theorem, we say that a Landweber exact ring spectrum E is of height $\leq n$ if $v_n \in \pi_*(E)/(p, v_1, \dots, v_{n-1})$ is a unit for all primes p .¹ As a corollary, we obtain the following result for height ≤ 2 Lubin–Tate theories:

¹It may happen that $\pi_*(E)/(p, v_1, \dots, v_{n-1})$ is the zero ring. In this case, our convention is to regard the unique element of the zero ring as a unit.

Corollary 1.5. *Let $k \subseteq \overline{\mathbb{F}}_p$ denote a field of characteristic $p > 0$ which is algebraic over the prime field \mathbb{F}_p , and let \mathbb{G} denote a formal group of height ≤ 2 over k . Then any complex orientation of the associated 2-periodic Morava K -theory $K(k, \mathbb{G})$ lifts uniquely up to homotopy to an \mathbb{E}_∞ -complex orientation of $E(k, \mathbb{G})$*

$$\mathrm{MU} \rightarrow E(k, \mathbb{G}).$$

Proof. This follows immediately from Theorem 1.4 and [Zhu20, Theorem 1.2], which implies that every complex orientation of $K(k, \mathbb{G})$ admits a unique lift to a complex orientation of $E(k, \mathbb{G})$ which satisfies the Ando criterion whenever k is algebraic over \mathbb{F}_p . \square

Remark 1.6. Theorem 1.4 was inspired by recent work of Balderrama [Bal21, Theorem 6.5.3]. Balderrama shows that every periodic complex orientation of a Lubin–Tate theory of height ≤ 2 satisfying an analogue of the Ando criterion lifts to an \mathbb{E}_∞ -orientation. He also shows that \mathbb{E}_∞ -refinements of periodic complex orientations of even periodic $K(1)$ -local \mathbb{E}_∞ -rings exist whenever the Ando criterion is satisfied and are unique up to homotopy.

The key observation he makes is the presence of evenness in the Goerss–Hopkins obstruction theory for (periodic) \mathbb{E}_∞ -complex orientations, which implies that the obstructions to existence and uniqueness appearing in his theorem vanish for formal reasons. Our results about existence of \mathbb{E}_∞ -complex orientations will be obtained by observing a similar evenness in the Hopkins–Lawson obstruction theory [HL18].

In contrast to Theorem 1.4, Balderrama does not obtain any uniqueness results at height 2 [Bal21, Remark 6.5.4]. In Remark 3.6, we will prove that \mathbb{E}_∞ -refinements of periodic complex orientations of height 2 Lubin–Tate theories are unique.

Our second main theorem uses Theorem 1.4 to give a simple proof of the following theorem:

Theorem 1.7. *For $n \geq 2$, the Hirzebruch level n elliptic genus lifts uniquely up to homotopy to a map of \mathbb{E}_∞ -rings*

$$\mathrm{MU} \rightarrow \mathrm{tmf}_1(n).$$

While Theorem 1.4 does not apply directly to $\mathrm{tmf}_1(n)$, it does apply to $\mathrm{TMF}_1(n)$, and it is not hard to upgrade the resulting \mathbb{E}_∞ -complex orientation to one of $\mathrm{tmf}_1(n)$. This reduces us to verifying the Ando criterion, which may be done following the strategy of Ando–Hopkins–Strickland [AHS04].

Remark 1.8. During the writing process, [Abs21] has also appeared, which follows the strategy of Ando–Hopkins–Rezk [AHR10] to prove a similar result to Theorem 1.7 for $\mathrm{Tmf}_1(n)$. Our method is rather different from that of Absmeier and completely avoids the consideration of p -adic Eisenstein measures.

1.3. Further questions. One of key inputs in our proof of Theorem 1.4 is Theorem 2.11, which states that the Morava K -theory of certain finite groups is concentrated in even degrees. This is closely related to these groups being *good* in the sense of Hopkins–Kuhn–Ravenel [HKR00, §7].

These groups come in a family, and to see that the Hopkins–Lawson obstruction theory is concentrated in even degrees one would like to show that the entire family has Morava K -theory concentrated in even degrees, cf. Remark 2.13:

Question 1.9. Is the Morava K -theory of the groups $\Gamma_k^{(n)}$ of Definition 2.7 concentrated in even degrees for $k > 2$, at least for $n \gg 0$? The groups $\Gamma_k^{(1)}$ are the extraspecial p -groups of type p_+^{1+2k} .

One could also ask about C_2 -equivariant refinement of our results.

Question 1.10. The complex cobordism spectrum may be refined to a C_2 -equivariant \mathbb{E}_∞ -ring $\mathrm{MU}_\mathbb{R}$, and $\mathrm{tmf}_1(n)$ admits the natural structure of a C_2 -equivariant \mathbb{E}_∞ -ring [Mei21, Theorem 2.20]. Moreover, the Hirzebruch level n elliptic genus admits a refinement to a map of homotopy C_2 -ring spectra $\mathrm{MU}_\mathbb{R} \rightarrow \mathrm{tmf}_1(n)$ [Mei21, Theorem 3.5]. Is there a suitable C_2 -equivariant analogue of Theorem 1.4 which may be used to prove a C_2 -equivariant refinement of Theorem 1.7? See [HL18, Remark 13] for a comment on a C_2 -equivariant version of the Hopkins–Lawson obstruction theory.

On the other hand, it would be very interesting to study \mathbb{E}_∞ -complex orientations at heights ≥ 3 . A natural choice of spectra to study would be Lubin–Tate spectra. Since the obstruction groups don’t vanish for formal reasons at these heights, it seems likely that this will require an explicit analysis of the Goerss–Hopkins obstruction theory for \mathbb{E}_∞ -maps $\mathrm{MU} \rightarrow E(k, \mathbb{G})$. In particular, one would have to compute the E_2 -page.

Problem 1.11. Compute the Goerss–Hopkins obstruction groups for \mathbb{E}_∞ -maps $\mathrm{MU} \rightarrow E(k, \mathbb{G})$ for height ≥ 3 Lubin–Tate theories $E(k, \mathbb{G})$.

1.4. Acknowledgements. The author would like to thank Robert Burklund and Jeremy Hahn for useful comments on a draft. During the course of this work, the author was supported by NSF Grants DGE-1745302 and DMS-2103236.

2. EXISTENCE OF \mathbb{E}_∞ -ORIENTATIONS

In this section, we will prove the half of Theorem 1.4 concerning the existence of \mathbb{E}_∞ -complex orientations. The main tool that we will utilize is an obstruction theory for \mathbb{E}_∞ -complex orientations studied by Hopkins–Lawson [HL18]. Given this obstruction theory, the existence half of Theorem 1.4 reduces to the statement that certain obstruction groups vanish. Using work of Arone–Lesh [AL07], this can be further reduced to the evenness of the Morava K -theory of certain extraspecial p -groups and related groups, which we are able to extract from the literature.

2.1. Hopkins–Lawson obstruction theory. We begin by summarizing the main results of the Hopkins–Lawson obstruction theory [HL18]. First, a definition:

Definition 2.1. Let E denote a homotopy commutative ring. We let $\mathrm{Or}(E)$ denote the space of complex orientations of E , i.e. the fiber

$$\mathrm{Or}(E) \rightarrow \mathrm{Map}(\Sigma^{\infty-2} \mathbb{C}\mathbb{P}^\infty, E) \rightarrow \mathrm{Map}(\Sigma^{\infty-2} \mathbb{C}\mathbb{P}^2, E) \simeq \mathrm{Map}(\mathbb{S}, E)$$

above the unit map $\mathbb{S} \rightarrow E$.

Theorem 2.2 ([HL18, Theorems 1 & 32]). *There exists a diagram of \mathbb{E}_∞ -ring spectra*

$$\mathbb{S} \rightarrow \mathrm{MX}_1 \rightarrow \mathrm{MX}_2 \rightarrow \mathrm{MX}_3 \rightarrow \cdots \rightarrow \mathrm{MU}$$

such that the following hold:

- (1) *The natural map $\varinjlim \mathrm{MX}_n \rightarrow \mathrm{MU}$ is an equivalence.*
- (2) *The \mathbb{E}_∞ -ring MX_1 is equipped with a natural complex orientation inducing an equivalence $\mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MX}_1, E) \xrightarrow{\sim} \mathrm{Or}(E)$ for each \mathbb{E}_∞ -ring E .*
- (3) *Given $m > 0$ and an \mathbb{E}_∞ -ring E , there is a pullback square*

$$\begin{array}{ccc} \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MX}_m, E) & \longrightarrow & \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MX}_{m-1}, E) \\ \downarrow & & \downarrow \\ \{*\} & \longrightarrow & \mathrm{Map}_*(F_m, \mathrm{Pic}(E)), \end{array}$$

where F_m is a certain pointed space described below.

- (4) *The map $\mathrm{MX}_{m-1} \rightarrow \mathrm{MX}_m$ is a rational equivalence if $m > 1$, a p -local equivalence if m is not a power of p , and a $K(n)$ -local equivalence if $m > p^n$.*

- (5) Let E denote an \mathbb{E}_∞ -ring such that π_*E is p -local and p -torsion free. Then an \mathbb{E}_∞ -map $\mathrm{MX}_1 \rightarrow E$ extends to an \mathbb{E}_∞ -map $\mathrm{MX}_p \rightarrow E$ if and only if the corresponding complex orientation of E satisfies the Ando criterion.

Using this theorem, we will reduce the proof of Theorem 1.4 to the following:

Lemma 2.3. *Let E denote a p -complete Landweber exact ring spectrum with homotopy concentrated in even degrees. Then $E^{2n}(F_p) \cong E^{2n+1}(F_{p^2}) \cong 0$ for all $n \in \mathbb{Z}$.*

Remark 2.4. In fact, we only need that $E^{2n+1}(F_{p^2}) \cong 0$. However, we include the statement $E^{2n}(F_p) \cong 0$ since it is no harder for us to prove. This extra evenness implies uniqueness up to homotopy for \mathbb{E}_∞ -refinements of complex orientations of height ≤ 1 . However, we will prove uniqueness in a different way in Section 3.

Question 2.5. Given a p -complete Landweber exact ring spectrum E with homotopy concentrated in even degrees, is $E^{2*+k-1}(F_{p^k}) \cong 0$ for $k \geq 3$?

Proof of existence in Theorem 1.4 assuming Lemma 2.3. We begin by reducing to the case where E is p -complete for some prime p . We make use of the fracture square:

$$\begin{array}{ccc} E & \longrightarrow & \prod_p E_p^\wedge \\ \downarrow & & \downarrow \\ E_{\mathbb{Q}} & \longrightarrow & \left(\prod_p E_p^\wedge \right)_{\mathbb{Q}} \end{array}$$

Note that E_p^\wedge is again even and Landweber exact, cf. Appendix A.

By Theorem 2.2, we have $\mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MU}, R) \simeq \mathrm{Or}(R)$ for a rational \mathbb{E}_∞ -ring R ; it follows further that $\pi_1 \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MU}, R) \cong \pi_1 \mathrm{Or}(R) \cong 0$ if R has homotopy concentrated in even degrees. As a consequence, there are pullback squares of sets:

$$\begin{array}{ccc} \pi_0 \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MU}, E) & \longrightarrow & \pi_0 \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MU}, \prod_p E_p^\wedge) \\ \downarrow & & \downarrow \\ \pi_0 \mathrm{Or}(E_{\mathbb{Q}}) & \longrightarrow & \pi_0 \mathrm{Or}\left(\left(\prod_p E_p^\wedge\right)_{\mathbb{Q}}\right) \end{array} \quad (1)$$

and

$$\begin{array}{ccc} \pi_0 \mathrm{Or}(E) & \longrightarrow & \pi_0 \mathrm{Or}\left(\prod_p E_p^\wedge\right) \\ \downarrow & & \downarrow \\ \pi_0 \mathrm{Or}(E_{\mathbb{Q}}) & \longrightarrow & \pi_0 \mathrm{Or}\left(\left(\prod_p E_p^\wedge\right)_{\mathbb{Q}}\right). \end{array}$$

To lift a complex orientation of E to an \mathbb{E}_∞ -complex orientation, it therefore suffices to lift the induced complex orientation of E_p^\wedge . We may therefore assume that E is p -complete.

Let E now denote an p -complete Landweber exact \mathbb{E}_∞ -ring with homotopy concentrated in even degrees. Using Theorem 2.2, we see that it suffices to show that

$$\pi_0 \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MX}_{p^2}, E) \rightarrow \pi_0 \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MX}_p, E)$$

is surjective.

By Theorem 2.2, there is a fiber sequence

$$\mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MX}_{p^2}, E) \rightarrow \mathrm{Map}_{\mathbb{E}_\infty}(\mathrm{MX}_p, E) \rightarrow \mathrm{Map}_*(F_{p^2}, \mathrm{Pic}(E))$$

Now, there are equivalences

$$\mathrm{Map}_*(F_m, \mathrm{Pic}(E)) \simeq \mathrm{Hom}(\Sigma^\infty F_m, \mathrm{pic}(E)) \simeq \mathrm{Hom}(\Sigma^\infty F_m, \Sigma E),$$

where in the second equivalence we have used the fact that $\Sigma^\infty F_m$ is $(2m-1)$ -connected by [HL18, Corollary 4(5)]. It therefore follows from the above fiber sequence that it suffices to show that

$$E^1(\Sigma^\infty F_{p^2}) \cong 0.$$

Since E is p -complete, this follows from Lemma 2.3. \square

2.2. Proof of Lemma 2.3. In the remainder of this section, we will prove Lemma 2.3. First, we must recall the definition of the spaces F_m .

Recollection 2.6. Let L_m denote the nerve of the topologized poset of proper direct-sum decompositions of \mathbb{C}^m , and let $(L_m)^\diamond$ denote its unreduced suspension. The natural action of $U(m)$ on \mathbb{C}^m endows L_m and $(L_m)^\diamond$ with the structure of $U(m)$ -spaces.

Furthermore, view S^{2m} as a $U(m)$ -space via its identification with the one-point compactification of \mathbb{C}^m , viewed as the fundamental representation of $U(m)$. Then F_m is given by

$$F_m \simeq ((L_m)^\diamond \wedge S^{2m})_{hU(m)}.$$

To prove Lemma 2.3, we will use a result of Arone–Lesh to reduce the study of the E -cohomology of F_m to the E -cohomology of certain groups Γ_k , whose definition we now recall.

Definition 2.7 ([Oli94, Definition 1]). Let $\sigma_0, \dots, \sigma_{k-1} \in \Sigma_{p^k}$ denote the permutations

$$\sigma_r(i) = \begin{cases} i + p^r & \text{if } i \equiv 1, \dots, (p-1)p^r \pmod{p^{r+1}} \\ i - (p-1)p^r & \text{if } i \equiv (p-1)p^r + 1, \dots, p^{r+1} \pmod{p^{r+1}}. \end{cases}$$

We let $\Gamma_k \subset U(p^k)$ denote the subgroup generated by the permutation matrices corresponding to $\sigma_0, \dots, \sigma_{k-1}$, the central S^1 , and the diagonal matrices A_0, \dots, A_{k-1} given by

$$(A_r)_{ii} = \zeta_p^{\lfloor (i-1)/p^r \rfloor},$$

where ζ_p is a primitive p th root of unity. Then Γ_k lies in a central extension

$$1 \rightarrow S^1 \rightarrow \Gamma_k \rightarrow \mathbb{F}_p^{2k} \rightarrow 1.$$

For each $n \geq 1$, there is a normal subgroup $\Gamma_k^{(n)} \subset \Gamma_k$ which only contains the central p^n th roots of unity instead of the full S^1 . Then there are central extensions

$$1 \rightarrow C_{p^n} \rightarrow \Gamma_k^{(n)} \rightarrow \mathbb{F}_p^{2k} \rightarrow 1$$

and exact sequences

$$1 \rightarrow \Gamma_k^{(n)} \rightarrow \Gamma_k \rightarrow S^1 \rightarrow 1.$$

Remark 2.8. The groups $\Gamma_k^{(1)}$ are examples of *extraspecial p -groups*, and in this language are commonly denoted by p_+^{1+2k} .

Proposition 2.9 ([AL07, Propositions 9.6 & 10.3]). *The p -completion of the spectrum*

$$\Sigma^\infty F_m \simeq \Sigma^\infty((L_m)^\diamond \wedge S^{2m})_{hU(m)}$$

is null unless $m = p^k$, in which case it is a summand of the p -completion of

$$\Sigma^k(S^{2p^k})_{h\Gamma_k},$$

where Γ_k acts on S^{2p^k} via the inclusion $\Gamma_k \subset U(p^k)$.

Now, $(S^{2p^k})_{h\Gamma_k}$ may also be described as the Thom spectrum associated to the composition

$$B\Gamma_k \rightarrow \mathrm{BU}(p^k) \rightarrow \mathbb{Z} \times \mathrm{BU},$$

from which it follows that

$$E^*((S^{2p^k})_{h\Gamma_k}) \cong \tilde{E}^{*-2p^k}(B\Gamma_k).$$

Lemma 2.3 therefore reduces to the following lemma:

Lemma 2.10. *Let E denote a p -local Landweber exact ring spectrum whose homotopy is concentrated in even degrees. Then, for $k \leq 2$, $E^*(B\Gamma_k)$ is concentrated in even degrees.*

We will deduce Lemma 2.10 from the following Morava K -theory computations:

Theorem 2.11. *For all $n \geq 0$, the following groups are concentrated in even degrees:*

- (1) (Tezuka–Yagita, [TY89, Theorem 4.2]) $K(n)^*(B\Gamma_1^{(1)})$ at all primes p .
- (2) (Schuster–Yagita, [SY04, Theorem 5.4]) $K(n)^*(B\Gamma_2^{(1)})$ at the prime 2.
- (3) (Yagita, [Yag05, Theorem 1.2]) $K(n)^*(B\Gamma_2^{(2)})$ at odd primes p .

A lemma of Strickland which builds on work of Ravenel–Wilson–Yagita [RWY98] allows us to transport this evenness from Morava K -theory to E -cohomology:

Corollary 2.12. *Given a p -local Landweber exact ring spectrum E whose homotopy is concentrated in even degrees, the following groups are concentrated in even degrees:*

- (1) $E^*(B\Gamma_1^{(1)})$ at all primes p .
- (2) $E^*(B\Gamma_2^{(1)})$ at the prime 2.
- (3) $E^*(B\Gamma_2^{(2)})$ at odd primes p .

Proof. Combine Theorem 2.11 with [Str99, Lemma 8.25].^{2,3} □

Proof of Lemma 2.10. The short exact sequence

$$\Gamma_k^{(n)} \rightarrow \Gamma_k \rightarrow S^1$$

induces a fiber sequence

$$B\Gamma_k^{(n)} \rightarrow B\Gamma_k \rightarrow BS^1.$$

Using the associated Atiyah–Hirzebruch spectral sequence

$$H^*(BS^1; E^*(B\Gamma_k^{(n)})) \Rightarrow E^*(B\Gamma_k),$$

we find that if $E^*(B\Gamma_k^{(n)})$ is even for some n , then $E^*(B\Gamma_k)$ must be as well. Therefore Lemma 2.10 follows from Corollary 2.12. □

Remark 2.13. By the same arguments, to give a positive answer to Question 2.5 it suffices to show that $K(n)^*(B\Gamma_k^{(i_k)})$ is concentrated in even degrees for a fixed i_k not depending on n . This is closely related to the question of whether $\Gamma_k^{(i_k)}$ is a *good group* in the sense of Hopkins–Kuhn–Ravenel [HKR00, §7].

²Note that we cannot apply [Str99, Lemma 8.25] directly to $B\Gamma_k$, since it only applies to spaces with bounded above \mathbb{Q} -cohomology.

³While it is assumed in [Str99] that E is even-periodic, this is not used in the proof of [Str99, Lemma 8.25]. We also note that E may be viewed as a summand of an even-periodic ring spectrum $E[x_2^{\pm 1}]$, so that we may conclude from [Str99, Lemma 8.25] as stated.

3. UNIQUENESS OF \mathbb{E}_∞ -ORIENTATIONS

Our goal in this section is to prove the following result:

Theorem 3.1. *Let E denote an L_2 -local complex orientable \mathbb{E}_∞ -ring with the property that $K(1)_*E$ and $K(1)_*L_{K(2)}E$ are concentrated in even degrees at all primes p . Then each complex orientation of E admits at most one refinement to an \mathbb{E}_∞ -complex orientation up to homotopy.*

Example 3.2. Any Landweber exact ring spectrum E of height ≤ 2 whose homotopy is concentrated in even degrees satisfies the hypotheses of Theorem 3.1. By Lemma A.1, $L_{K(2)}E$ is again Landweber exact and has homotopy concentrated in even degrees. It therefore suffices to show that $K(1)_*E$ is concentrated in even degrees. This is true because $K(1)_*E \cong K(1)_*\mathrm{MU} \otimes_{\pi_*\mathrm{MU}} \pi_*E$, and $K(1)_*\mathrm{MU}$ is concentrated in even degrees.

Combining Theorem 3.1 with Example 3.2, we obtain the uniqueness half of Theorem 1.4.

Example 3.3. The ring spectra $\mathrm{Tmf}_1(n)$ satisfy the hypotheses of Theorem 3.1. This follows from [Wil15, Propositions 2.4 and 2.6], which imply that the p -complete complex K -theory of these spectra and their $K(2)$ -localizations is p -torsionfree and concentrated in even degrees.

Our proof of Theorem 3.1 will be based on the orientation theory of [ABG⁺14] and the following lemma:

Lemma 3.4. *Let E denote an MU -module with the property that $K(1)_*E$ is concentrated in even degrees. Then $[\mathrm{KU}_p, L_{K(1)}E]$ is torsionfree and $[\Sigma\mathrm{KU}_p, L_{K(1)}E] = 0$.*

Definition 3.5. We say that a $(\mathrm{KU}_p)_*$ -module is *pro-free* if it is the p -completion of a free module. Moreover, given a spectrum X , we write $\mathrm{KU}_*^\vee(X)$ for $\pi_*L_{K(1)}(\mathrm{KU} \otimes X)$.

Proof. Since $K(1)_*(\mathrm{KU}_p)$ is even, $\mathrm{KU}_*^\vee(\mathrm{KU}_p)$ is pro-free by [HS99b, Proposition 8.4(f)]. Therefore, by [BH16, Proposition 1.14], there is an isomorphism

$$\pi_* \mathrm{Hom}(\mathrm{KU}_p, L_{K(1)}(\mathrm{KU}_p \otimes E)) \cong \mathrm{Hom}_{(\mathrm{KU}_p)_*}(\mathrm{KU}_*^\vee(\mathrm{KU}_p), \mathrm{KU}_*^\vee(E)).$$

By assumption, $K(1)_*(E)$ is even and hence $\mathrm{KU}_*^\vee(E)$ is even and pro-free by [HS99b, Proposition 8.4(f)]. In particular, it is torsionfree, so that $\mathrm{Hom}_{(\mathrm{KU}_p)_*}(\mathrm{KU}_*^\vee(\mathrm{KU}_p), \mathrm{KU}_*^\vee(E))$ is even and torsionfree.

The result then follows from the following facts:

- (1) Since E is an MU -module, the unit map $L_{K(1)}E \rightarrow L_{K(1)}(\mathrm{MU} \otimes E)$ admits a splitting given by the module structure map.
- (2) There is a splitting of the map $L_{K(1)}\mathrm{MU} \rightarrow L_{K(1)}E(1)$ which is compatible with the unit map [HS99a, Theorem 4.1].
- (3) There is an equivalence of spectra $\mathrm{KU}_p \simeq \bigoplus_{i=0}^{p-2} \Sigma^{2i} L_{K(1)}E(1)$. □

Proof of Theorem 3.1. Using the pullback square of sets (1), we may assume that E is p -complete.

Recall the map $\Sigma^\infty \mathbb{C}\mathbb{P}^\infty \rightarrow \mathrm{bu}$ which is adjoint to the canonical map $\mathbb{C}\mathbb{P}^\infty \rightarrow \mathrm{BU}$. By orientation theory [ABG⁺14], we must show that

$$[\mathrm{bu}, \mathrm{gl}_1(E)] \rightarrow [\Sigma^\infty \mathbb{C}\mathbb{P}^\infty, \mathrm{gl}_1(E)]$$

is injective. Since E is p -complete, $\mathrm{gl}_1(E)$ agrees with $\mathrm{gl}_1(E)_p^\wedge$ in degrees ≥ 2 , so we may as well replace the former by the latter. Letting F denote the fiber of the map $\mathrm{gl}_1(E) \rightarrow L_2\mathrm{gl}_1(E)$, we find that there is a fiber sequence

$$F_p^\wedge \rightarrow \mathrm{gl}_1(E)_p^\wedge \rightarrow L_{K(1) \oplus K(2)}\mathrm{gl}_1(E).$$

It therefore suffices to show that

$$[\mathrm{bu}, F_p^\wedge] \rightarrow [\Sigma^\infty \mathbb{C}\mathbb{P}^\infty, F_p^\wedge]$$

and

$$[\mathrm{bu}, L_{K(1) \oplus K(2)} \mathrm{gl}_1(E)] \rightarrow [\Sigma^\infty \mathbb{C}\mathbb{P}^\infty, L_{K(1) \oplus K(2)} \mathrm{gl}_1(E)]$$

are injective. The first is injective because F_p^\wedge is 3-coconnective by [AHR10, Theorem 4.11] and the cofiber of $\Sigma^\infty \mathbb{C}\mathbb{P}^\infty \rightarrow \mathrm{bu}$ is 4-connective.

To prove that the second is injective, we first note that the Bousfield–Kuhn functor [Bou87, Kuh89] and the chromatic fracture square for $L_{K(1) \oplus K(2)} \mathrm{gl}_1(E)$ imply that there is an exact sequence

$$[\Sigma \mathrm{KU}_p, L_{K(1)} L_{K(2)} E] \rightarrow [\mathrm{bu}, L_{K(1) \oplus K(2)} E] \rightarrow [\mathrm{KU}_p, L_{K(1)} E].$$

Applying Lemma 3.4, we learn that $[\mathrm{bu}, L_{K(2) \oplus K(1)} E]$ is torsionfree. It therefore injects into its rationalization, so that the result follows from the fact that $\Sigma^\infty \mathbb{C}\mathbb{P}^\infty \rightarrow \mathrm{bu}$ is a rational equivalence. \square

Remark 3.6. Balderrama has shown that *periodic* complex orientations of height 2 Lubin–Tate theories $E(k, \mathbb{G})$ which satisfy a version of the Ando criterion admit lifts to periodic \mathbb{E}_∞ -complex orientations $\mathrm{MUP} \rightarrow E(k, \mathbb{G})$ [Bal21, Theorem 6.5.3(3)]. In this remark, we prove that such lifts are unique up to homotopy.

We begin with a result which has been proven by Rezk at height 2 (which is the case that we use) [Rez13] and in general is an unpublished theorem of Hopkins and Lurie. Let \bar{k} denote the algebraic closure of k . Then we have:

$$\pi_* \mathbb{G}_m(E(\bar{k}, \mathbb{G})) := \pi_* \mathrm{Map}(\mathbb{Z}, \mathrm{gl}_1(E(\bar{k}, \mathbb{G}))) \cong \begin{cases} \bar{k}^\times & * = 0 \\ \mathbb{Z}_p & * = 3 \\ 0 & \text{otherwise.} \end{cases}$$

On π_0 , the map $\mathbb{Z} \rightarrow \mathrm{gl}_1(E(\bar{k}, \mathbb{G}))$ corresponding to $a \in \bar{k}^\times$ picks out the Teichmüller lift $[a]$.

Since $\mathrm{Gal}(k)$ has p -cohomological dimension at most 1 [GS06, Proposition 6.1.9], it follows that

$$\pi_0 \mathbb{G}_m(E(k, \mathbb{G})) \cong k^\times.$$

In particular, the map $\pi_0 \mathbb{G}_m(E(k, \mathbb{G})) \rightarrow \pi_0 \mathrm{gl}_1(E(k, \mathbb{G}))$ is injective. Now, by orientation theory it suffices to show that

$$[\mathrm{ku}, \mathrm{gl}_1(E(k, \mathbb{G}))] \rightarrow [\Sigma_+^\infty \mathbb{C}\mathbb{P}^\infty, \mathrm{gl}_1(E(k, \mathbb{G}))]$$

is injective. By what we have proven above about uniqueness of \mathbb{E}_∞ -complex orientations, it suffices to show that

$$[\mathbb{Z}, \mathrm{gl}_1(E(k, \mathbb{G}))] \rightarrow [\mathbb{S}^0, \mathrm{gl}_1(E(k, \mathbb{G}))]$$

is injective, which is what we showed above.

As noted in [Bal21, Remark 6.5.4], this implies that the Goerss–Hopkins obstruction group $\mathrm{Ext}_\Delta^2(\hat{Q}(E(k, \mathbb{G}))_0^\wedge \mathrm{MUP}, \omega)$ is equal to 0.

4. THE LEVEL n ELLIPTIC GENUS

Convention 4.1. In this section n will denote an integer greater than or equal to 2.

In this section, we will prove Theorem 1.7, which states that the Hirzebruch level n elliptic genus lifts uniquely up to homotopy to an \mathbb{E}_∞ -complex orientation $\mathrm{MU} \rightarrow \mathrm{tmf}_1(n)$.

We will begin by recalling some background material about \mathbb{E}_∞ -rings of topological modular forms with level- $\Gamma_1(n)$ structures in Section 4.1. In Section 4.2, we recall from [AHS01, AHS04] how complex orientations may be described in terms of Θ^1 -structures.

We then describe the level n elliptic genus in this language, following Meier [Mei21, §3]. In Section 4.3, we show that a complex orientation for $\mathrm{TMF}_1(n)$ satisfies the Ando criterion at p if and only if its composition along a map $\mathrm{TMF}_1(n) \rightarrow E(k, \mathbb{G})$ to a Lubin–Tate theory does.

In Section 4.4, we recall from [AHS04] how the Ando criterion for Lubin–Tate theories may be rephrased in terms of Θ^1 -structures. We then prove that the Hirzebruch level n genus satisfies the Ando criterion. As a consequence of Theorem 1.4, it lifts uniquely up to homotopy to an \mathbb{E}_∞ -ring map $\mathrm{MU} \rightarrow \mathrm{TMF}_1(n)$. Finally, in Section 4.5, we prove that this \mathbb{E}_∞ -ring map admits a unique up to homotopy lift to an \mathbb{E}_∞ -ring map $\mathrm{MU} \rightarrow \mathrm{tmf}_1(n)$, completing the proof of Theorem 1.7.

4.1. Topological modular forms with level structures. In this section, we will recall some basic facts about the \mathbb{E}_∞ -rings $\mathrm{tmf}_1(n)$, $\mathrm{Tmf}_1(n)$ and $\mathrm{TMF}_1(n)$ of topological modular forms with level- $\Gamma_1(n)$ structure. We begin by recalling the algebraic background.

Definition 4.2. We let $\mathcal{M}_1(n)$ denote the Deligne–Mumford moduli stack of elliptic curves with level- $\Gamma_1(n)$ structure over $\mathbb{Z}[\frac{1}{n}]$. Concretely, given a scheme S on which n is invertible, we have

$$\mathcal{M}_1(n)(S) = \text{elliptic curves } E \text{ over } S \text{ with a point } P \in E[n](S) \text{ of exact order } n.$$

Moreover, we write $\overline{\mathcal{M}}_1(n)$ for the Deligne–Rapoport moduli stack of generalized elliptic curves with level- $\Gamma_1(n)$ structure [DR73]. This is again a Deligne–Mumford stack over $\mathbb{Z}[\frac{1}{n}]$, and $\mathcal{M}_1(n) \subset \overline{\mathcal{M}}_1(n)$ is an open substack.

We denote the universal family of curves by $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_1(n)$, and write $\pi^{\mathrm{sm}} : \mathcal{C}^{\mathrm{sm}} \rightarrow \overline{\mathcal{M}}_1(n)$ for the smooth locus. Then $\mathcal{C}^{\mathrm{sm}}$ admits the natural structure of a group scheme, and we write ω for the line bundle of invariant differentials.

Finally, we write $\mathcal{M} = \mathcal{M}_1(1)$ and $\overline{\mathcal{M}} = \overline{\mathcal{M}}_1(1)$ for the moduli stacks of (generalized) elliptic curves without level structure.

Recollection 4.3 (Goerss–Hopkins–Miller [DFHH14, Chapter 12], Lurie [Lur18b], Hill–Lawson [HL16]). There is a sheaf $\mathcal{O}_{\overline{\mathcal{M}}}^{\mathrm{top}}$ of \mathbb{E}_∞ -ring spectra on the Kummer log-étale site of $\overline{\mathcal{M}}$ with the following properties:

- (1) There is a natural isomorphism of sheaves $\pi_0 \mathcal{O}_{\overline{\mathcal{M}}}^{\mathrm{top}} \cong \mathcal{O}_{\overline{\mathcal{M}}}$.
- (2) There are natural isomorphisms of quasicoherent sheaves $\pi_{2i} \mathcal{O}_{\overline{\mathcal{M}}}^{\mathrm{top}} \cong \omega^i$ and $\pi_{2i+1} \mathcal{O}_{\overline{\mathcal{M}}}^{\mathrm{top}} \cong 0$.
- (3) Write $\widehat{\pi^{\mathrm{sm}}} : \widehat{\mathcal{C}^{\mathrm{sm}}} \rightarrow \overline{\mathcal{M}}$ for the completion of $\pi^{\mathrm{sm}} : \mathcal{C}^{\mathrm{sm}} \rightarrow \overline{\mathcal{M}}$ along the zero section. There is a natural isomorphism of sheaves of rings $\pi_0 \mathrm{Hom}(\Sigma_+^\infty \mathbb{C}\mathbb{P}^\infty, \mathcal{O}^{\mathrm{top}}) \cong (\widehat{\pi^{\mathrm{sm}}})_* \mathcal{O}_{\widehat{\mathcal{C}^{\mathrm{sm}}}}$.

Since the natural morphisms $\overline{\mathcal{M}}_1(n) \rightarrow \overline{\mathcal{M}}$ are Kummer log-étale (we refer the reader to [HL16] for more details), we may define \mathbb{E}_∞ -rings:

$$\mathrm{TMF}_1(n) := \Gamma(\mathcal{M}_1(n), \mathcal{O}^{\mathrm{top}})$$

and

$$\mathrm{Tmf}_1(n) := \Gamma(\overline{\mathcal{M}}_1(n), \mathcal{O}^{\mathrm{top}}).$$

By definition, there are spectral sequences

$$\mathrm{H}^s(\mathcal{M}_1(n), \omega^i) \Rightarrow \pi_{2i-s} \mathrm{TMF}_1(n)$$

and

$$\mathrm{H}^s(\overline{\mathcal{M}}_1(n), \omega^i) \Rightarrow \pi_{2i-s} \mathrm{Tmf}_1(n).$$

Remark 4.4. Let $n \geq 2$. It follows from [Mei20, Proposition 2.4(4)] that $H^s(\mathcal{M}_1(n), \omega^i) = 0$ for all $s > 0$, so that $\mathrm{TMF}_1(n)$ has homotopy groups concentrated in even degrees and that there are natural isomorphisms

$$\pi_{2i}(\mathrm{TMF}_1(n)) \cong \Gamma(\mathcal{M}_1(n), \omega^i).$$

Moreover, $\mathrm{TMF}_1(n)$ is Landweber exact. Indeed, this is a consequence of flatness of $\mathcal{M}_1(n)$ over $\mathbb{Z}[\frac{1}{n}]$, the integrality of $\mathcal{M}_1(n)_{\mathbb{F}_p}$ and the fact that the formal group of an elliptic curve is of height ≤ 2 .

However, because the groups $H^1(\overline{\mathcal{M}}_1, \omega^i)$ do not in general vanish, we do not have a similar theorem for $\mathrm{Tmf}_1(n)$. Instead, we have the \mathbb{E}_∞ -ring spectrum $\mathrm{tmf}_1(n)$ from [Mei21].

Recollection 4.5 ([Mei21, Theorem 1.1]). There is an essentially unique connective \mathbb{E}_∞ -ring spectrum $\mathrm{tmf}_1(n)$ whose homotopy groups are concentrated in even degrees and which is equipped with an \mathbb{E}_∞ -ring map

$$\mathrm{tmf}_1(n) \rightarrow \mathrm{Tmf}_1(n)$$

such that the induced maps

$$\pi_{2i}\mathrm{tmf}_1(n) \rightarrow \pi_{2i}\mathrm{Tmf}_1(n) \rightarrow \Gamma(\overline{\mathcal{M}}_1(n), \omega^i)$$

are isomorphisms.

Remark 4.6. There is a sequence of natural maps

$$\mathrm{tmf}_1(n) \rightarrow \mathrm{Tmf}_1(n) \rightarrow \mathrm{TMF}_1(n).$$

4.2. Θ^1 -structures. In this section, we describe complex orientations in terms of Θ^1 -structures and give a description of the Hirzebruch level n genus in this language.

Suppose that we are given a base Deligne–Mumford stack S and a formal group or generalized elliptic curve G over S . We denote the structure map by $p : G \rightarrow S$ and the zero section by $0 : S \rightarrow G$. Given a line bundle \mathcal{L} on G , we set

$$\Theta^1(\mathcal{L}) := p^*0^*\mathcal{L} \otimes \mathcal{L}^{-1}.$$

There is a canonical trivialization $0^*\Theta^1(\mathcal{L}) \cong \mathcal{O}_S$.

Definition 4.7. A Θ^1 -structure on a line bundle \mathcal{L} over G is a trivialization of $\Theta^1(\mathcal{L})$ which pulls back to the canonical trivialization of $0^*\Theta^1(\mathcal{L})$.

Equivalently, a Θ^1 -structure on \mathcal{L} is an isomorphism $p^*0^*\mathcal{L} \cong \mathcal{L}$ which pulls back to the canonical isomorphism $0^*p^*0^*\mathcal{L} \cong 0^*\mathcal{L}$.

Remark 4.8. It is clear from the definition that there is a natural bijection between Θ^1 -structures on \mathcal{L} and \mathcal{L}^{-1} .

Remark 4.9. Note that a line bundle \mathcal{L} on G admits a Θ^1 -structure if and only if it is pulled back from a line bundle on S via the structure map p .

When G is a generalized elliptic curve, Θ^1 -structures are unique when they exist:

Lemma 4.10. *Suppose that G is a generalized elliptic curve. Then the set of Θ^1 -structures on a line bundle \mathcal{L} over G is either empty or consists of a single element.*

Proof. The set of isomorphisms $p^*0^*\mathcal{L} \cong \mathcal{L}$ is a torsor for $\mathcal{O}_G(G)^\times$, whereas the set of isomorphisms $0^*p^*0^*\mathcal{L} \cong 0^*\mathcal{L}$ is a torsor for $\mathcal{O}_S(S)^\times$. It therefore suffices to show that the map $\mathcal{O}_G(G) \rightarrow \mathcal{O}_S(S)$ induced by pulling back along the zero section is an isomorphism. This follows from [Sta21, Tag 0E0L]. \square

Now let E denote an even weakly periodic homotopy commutative ring spectrum, and let $\mathbb{G}_E = \mathrm{Spf} E^0(\mathbb{C}\mathbb{P}^\infty)$ denote the associated formal group over $\mathrm{Spec} \pi_0 E$. We denote by γ_1 the canonical line bundle over $\mathbb{C}\mathbb{P}^\infty$ and let $\mathcal{O}_{\mathbb{G}_E}(1)$ denote the line bundle over \mathbb{G}_E corresponding to

$$E^0(\mathrm{Th}(\gamma_1)) \cong \widetilde{E}^0(\mathbb{C}\mathbb{P}^\infty) \cong \ker(\mathcal{O}_{\mathbb{G}_E} \xrightarrow{0^*} \pi_0 E).$$

Then we have the following theorem:

Proposition 4.11 ([AHS01, Theorem 2.48]). *There is a natural bijection between complex orientations of E and Θ^1 -structures on $\mathcal{O}_{\mathbb{G}_E}(1)$.*

Proof. Recall that a complex orientation of E consists of an element of $\widetilde{E}^2(\mathbb{C}\mathbb{P}^\infty)$ which restricts to the unit along the map $\widetilde{E}^2(\mathbb{C}\mathbb{P}^\infty) \rightarrow \widetilde{E}^2(\mathbb{C}\mathbb{P}^1) \cong \pi_0 E$. Since the map $E^0(\mathbb{C}\mathbb{P}^\infty) \rightarrow \pi_0 E$ is an infinitesimal thickening, complex orientations of E may be identified with $E^0(\mathbb{C}\mathbb{P}^\infty)$ -module isomorphisms $E^0(\mathbb{C}\mathbb{P}^\infty) \cong \widetilde{E}^2(\mathbb{C}\mathbb{P}^\infty)$ that become equal to the canonical isomorphism $\widetilde{E}^2(\mathbb{C}\mathbb{P}^1) \cong \pi_0 E$ after tensoring down along $E^0(\mathbb{C}\mathbb{P}^\infty) \rightarrow \pi_0 E$.

By Remark 4.8, we may replace $\mathcal{O}_{\mathbb{G}_E}(1)$ with $\mathcal{O}_{\mathbb{G}_E}(1)^{-1}$ in the statement of the proposition. To prove the proposition, it therefore suffices to identify the global sections of $\Theta^1(\mathcal{O}_{\mathbb{G}_E}(1)^{-1})$ with $\widetilde{E}^2(\mathbb{C}\mathbb{P}^\infty)$ and pullback along the zero section with $\widetilde{E}^2(\mathbb{C}\mathbb{P}^\infty) \rightarrow \widetilde{E}^2(\mathbb{C}\mathbb{P}^1) \cong \pi_0 E$. This is an immediate consequence of the definitions. \square

Even though $\mathrm{tmf}_1(n)$ is not weakly even periodic, Meier has shown that its complex orientations may still be described in terms of Θ^1 -structures.

Proposition 4.12 ([Mei21, Lemma 3.2]). *There is a natural bijection between complex orientations of $\mathrm{tmf}_1(n)$ and Θ^1 -structures on $\mathcal{O}_{\widehat{\mathcal{C}}}(1)$ over $\widehat{\mathcal{C}} \rightarrow \overline{\mathcal{M}}_1(n)$.*

We now recall the treatment of Hirzebruch's level n elliptic genus from [Mei21, §3]. The following proposition is a mild rephrasing of [Mei21, Lemma 3.3].

Proposition 4.13. *Let $P : \overline{\mathcal{M}}_1(n) \rightarrow \mathcal{C}^{\mathrm{sm}}$ denote the universal level- $\Gamma_1(n)$ structure.*

- (1) *The pullback of the line bundle $\mathcal{O}_{\mathcal{C}}([0] - [P])$ on \mathcal{C} to $\widehat{\mathcal{C}}$ is naturally isomorphic to $\mathcal{O}_{\widehat{\mathcal{C}}}(1)$.*
- (2) *There is a degree n étale cover $q : \mathcal{C}' \rightarrow \mathcal{C}$ of generalized elliptic curves so that $q^* \mathcal{O}_{\mathcal{C}}([0] - [P])$ admits a (necessarily unique) Θ^1 -structure.*

Since the induced map $\widehat{q} : \widehat{\mathcal{C}}' \xrightarrow{\sim} \widehat{\mathcal{C}}$ is an isomorphism, we obtain a Θ^1 -structure on $\mathcal{O}_{\widehat{\mathcal{C}}}(1)$, and hence by Proposition 4.12 a complex orientation of $\mathrm{tmf}_1(n)$. This is the complex orientation corresponding to the *Hirzebruch level n elliptic genus*.

4.3. Reduction to Lubin–Tate theory. Our goal in this section is to prove Proposition 4.16 below. This proposition implies that to verify the Ando criterion for the Hirzebruch level n elliptic genus, it suffices to verify the Ando criterion after composition with a map $\mathrm{TMF}_1(n) \rightarrow E(k, \mathbb{G})$ to a Lubin–Tate theory. This will be useful to us because work of Ando–Hopkins–Strickland [AHS04] rephrases the Ando criterion for Lubin–Tate theories in terms of Θ^1 -structures.

We begin by recalling some basic facts about about Lubin–Tate theories.

Recollection 4.14 (Goerss–Hopkins [GH04], Lurie [Lur18b]). Let k denote a perfect field of characteristic $p > 0$ and let \mathbb{G} denote a formal group of finite height h over k . To the pair (k, \mathbb{G}) one may associate an \mathbb{E}_∞ -ring spectrum $E(k, \mathbb{G})$, known as the *Lubin–Tate spectrum* of (k, \mathbb{G}) . The ring $\pi_0 E(k, \mathbb{G})$ is naturally isomorphic to the universal deformation ring of (k, \mathbb{G}) , which is noncanonically isomorphic to $\mathbb{W}(k)[[u_1, \dots, u_{h-1}]]$. We let $\mathbb{G}^{\mathrm{univ}}$ denote the universal deformation of \mathbb{G} over $\pi_0 E(k, \mathbb{G})$, and denote by $\omega_{\mathbb{G}^{\mathrm{univ}}}$

its module of invariant differentials. There are natural isomorphisms of $\pi_{2i}E(k, \mathbb{G})$ with $\omega_{\mathbb{G}^{\text{univ}}}^i$.

Construction 4.15. Let k denote a perfect field of characteristic $p > 0$. Associated to a supersingular k -point $(E, \alpha \in E[n](k))$ of $\mathcal{M}_1(n)$, there is a map of \mathbb{E}_∞ -ring spectra $\text{TMF}_1(n) \rightarrow E(k, \widehat{E})$. Indeed, this follows from the description of $E(k, \widehat{E})$ as an oriented deformation ring [Lur18b, §6], Lurie’s Serre–Tate theorem for strict abelian varieties [Lur18a, §7], and the universal property of $\text{TMF}_1(n)$.

Proposition 4.16. *Let k denote a perfect field of characteristic p and let $(E, \alpha \in E[n](k))$ denote an object of $\mathcal{M}_1(n)(k)$ with E a supersingular elliptic curve. Then a complex orientation of $\text{TMF}_1(n)$ satisfies the Ando criterion at p if and only if its composite with the canonical map $\text{TMF}_1(n) \rightarrow E(k, \widehat{E})$ does.*

Our proof of Proposition 4.16 rests on the following lemma:

Lemma 4.17. *Let $E \rightarrow F$ denote a map of complex orientable homotopy commutative ring spectra. Suppose that π_*E and π_*F are p -torsionfree and that the induced map $\pi_*E/p \rightarrow \pi_*F/p$ is an injection. Then the map $E^*(\text{BC}_p)/I_{tr} \rightarrow F^*(\text{BC}_p)/I_{tr}$ is an injection.*

Proof. Since π_*E is p -torsionfree, a choice of complex orientation gives rise to an isomorphism (see [Qui71, Proposition 4.2]):

$$E^*(\text{BC}_p)/I_{tr} \cong E^*[[t]]/\langle p \rangle(t)$$

where $|t| = 2$, $\langle p \rangle(t) = \frac{[p](t)}{t}$ and $[p](t)$ is the p -series of the formal group on E^* . Moreover, since π_*E is p -torsionfree, we may identify $t^{n-1}E^*[[t]]/\langle p \rangle(t), t^n$ with a shift of E^*/p . The analogous statements for F also hold.

Taking the induced complex orientation of F , we identify $E^*(\text{BC}_p)/I_{tr} \rightarrow F^*(\text{BC}_p)/I_{tr}$ with the natural map $E^*[[t]]/\langle p \rangle(t) \rightarrow F^*[[t]]/\langle p \rangle(t)$. To show that this map is injective, it suffices to show that it is injection on the associated graded for the t -adic filtration. This identifies with a shift of the natural map $E^*/p \rightarrow F^*/p$ in each degree, which is an injection by hypothesis. \square

Proof of Proposition 4.16. It is clear that if a complex orientation of $\text{TMF}_1(n)$ satisfies the Ando criterion at p , so does the induced complex orientation of $E(k, \widehat{E})$. To prove the converse, it suffices to show that the induced map

$$\text{TMF}_1(n)^*(\text{BC}_p)/I_{tr} \rightarrow E(k, \widehat{E})^*(\text{BC}_p)/I_{tr}$$

is an injection. We begin by reducing to the case $n \geq 5$. Any map $\text{Spec } k \rightarrow \mathcal{M}_1(n)$ fits into a diagram

$$\begin{array}{ccc} \text{Spec } k' & \longrightarrow & \mathcal{M}_1(n^2) \\ \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & \mathcal{M}_1(n) \end{array}$$

for k' a finite separable extension of k . It follows that there is a diagram

$$\begin{array}{ccc} \text{TMF}_1(n) & \longrightarrow & E(k, \widehat{E}) \\ \downarrow & & \downarrow \\ \text{TMF}_1(n^2) & \longrightarrow & E(k', \widehat{E}_{k'}), \end{array}$$

so that it suffices to show that

$$\mathrm{TMF}_1(n)^*(\mathrm{BC}_p)/I_{tr} \rightarrow \mathrm{TMF}_1(n^2)^*(\mathrm{BC}_p)/I_{tr}$$

and

$$\mathrm{TMF}_1(n^2)^*(\mathrm{BC}_p)/I_{tr} \rightarrow E(k', \widehat{E}_{k'})^*(\mathrm{BC}_p)/I_{tr}$$

are injective.

We may therefore assume that $n \geq 5$ if we can show that the first map is an injection. Both $\pi_*\mathrm{TMF}_1(n)$ and $\pi_*\mathrm{TMF}_1(n^2)$ are p -torsionfree, and

$$\pi_{2i}\mathrm{TMF}_1(n)/p \cong \Gamma(\mathcal{M}_1(n)_{\mathbb{F}_p}, \omega^i) \rightarrow \Gamma(\mathcal{M}_1(n^2)_{\mathbb{F}_p}, \omega^i) \cong \pi_{2i}\mathrm{TMF}_1(n)/p$$

is an injection since $\mathcal{M}_1(n^2)_{\mathbb{F}_p} \rightarrow \mathcal{M}_1(n)_{\mathbb{F}_p}$ is a finite étale cover. Therefore we may apply Lemma 4.17 to show that the first map above is an injection.

We may now assume that $n \geq 5$, so that $\mathcal{M}_1(n)$ is represented by an affine scheme $\mathrm{Spec} R_n$ [Mei20, Proposition 2.4(2)]. Let $\mathfrak{m} \subset R_n$ denote the kernel of the map $R_n \rightarrow k$. Then R_n/\mathfrak{m} is a finite field (hence perfect), and the pair $(E, P \in E[n](k))$ descends to R_n/\mathfrak{m} . As a consequence, there is a factorization $\mathrm{TMF}_1(n) \rightarrow E(R_n/\mathfrak{m}, \widehat{E}) \rightarrow E(k, \widehat{E})$. It follows immediately from Lemma 4.17 that the induced map

$$E(R_n/\mathfrak{m}, \widehat{E})^*(\mathrm{BC}_p)/I_{tr} \rightarrow E(k, \widehat{E})^*(\mathrm{BC}_p)/I_{tr}$$

is an injection, so that it suffices to show that

$$\mathrm{TMF}_1(n)^*(\mathrm{BC}_p)/I_{tr} \rightarrow E(R_n/\mathfrak{m}, \widehat{E})^*(\mathrm{BC}_p)/I_{tr}$$

is an injection. To apply Lemma 4.17, we need to show that $\pi_*\mathrm{TMF}_1(n)/p \rightarrow \pi_*E(R_n/\mathfrak{m}, \widehat{E})/p$ is an injection. By abuse of notation, we let ω denote the invertible R_n -module corresponding to the line bundle ω on $\mathcal{M}_1(n)$. Then the above map can be identified with the \mathfrak{m} -adic completion map $(\omega^{*/2}/p) \rightarrow (\omega^{*/2}/p)_{\mathfrak{m}}^{\wedge}$. This is an injection by the Krull intersection theorem, since $\mathcal{M}_1(n)_{\mathbb{F}_p}$ is an integral scheme. \square

4.4. The Ando criterion and Θ^1 -structures. In this section, we recall from the work of Ando–Hopkins–Strickland how the Ando criterion for Lubin–Tate theories may be rephrased in terms of Θ^1 -structures. We refer the reader to [AHS04] for proofs and further details. We then combine this rephrasing with Proposition 4.16 to prove that the Hirzebruch level n elliptic genus satisfies the Ando criterion. Finally, we deduce from Theorem 1.4 that the Hirzebruch level n elliptic genus lifts uniquely up to homotopy to an \mathbb{E}_{∞} -ring map

$$\mathrm{MU} \rightarrow \mathrm{TMF}_1(n).$$

We begin by recalling from [AHS04, §14] how Θ^1 -structures may be normed along isogenies.

Recollection 4.18 ([AHS04, §14]). Suppose we are given an isogeny $G \rightarrow G'$ of formal groups or elliptic curves. Given a line bundle \mathcal{L} over G , there is a line bundle $N(\mathcal{L})$ over G' , called the *norm* of \mathcal{L} . Moreover, given a Θ^1 -structure s on \mathcal{L} , there is an associated Θ^1 -structure $N(s)$ on $N(\mathcal{L})$, called the *norm* of s .

Remark 4.19. Given an isogeny of formal groups $\mathbb{G} \rightarrow \widetilde{\mathbb{G}}$, there is a natural isomorphism of line bundles $N(\mathcal{O}_{\mathbb{G}}(1)) \cong \mathcal{O}_{\widetilde{\mathbb{G}}}(1)$.

Remark 4.20. Suppose that we are given an elliptic curve E with a point P of exact order n , i.e. a level- $\Gamma_1(n)$ structure, and an isogeny $E \rightarrow \widetilde{E}$ of degree p coprime to n . Then the image \widetilde{P} of P in \widetilde{E} is again a point of exact order n , and there is a natural isomorphism of line bundles $N(\mathcal{O}_E([0] - [P])) \cong \mathcal{O}_{\widetilde{E}}([0] - [\widetilde{P}])$.

Following [AHS04], we may now use this language to rephrase the Ando criterion for Lubin–Tate theory. We begin with some setup.

Recollection 4.21. Let k denote a perfect field of characteristic $p > 0$ and let \mathbb{G} denote a formal group of finite height over k , so that we have an associated Lubin–Tate theory $E(k, \mathbb{G})$. Then there are two ring maps $i, \psi : \pi_0 E(k, \mathbb{G}) \rightarrow E(k, \mathbb{G})^0(\mathrm{BC}_p)/I_{tr}$. The first, i , is induced by the projection $\mathrm{BC}_p \rightarrow *$. The second ψ , is the total power operation. Over the ring $E(k, \mathbb{G})^0(\mathrm{BC}_p)/I_{tr}$, there is a degree p isogeny

$$i^* \mathbb{G}^{\mathrm{univ}} \rightarrow \psi^* \mathbb{G}^{\mathrm{univ}}$$

induced by the total power operation on $E(k, \mathbb{G})^0(\mathbb{C}\mathbb{P}^\infty)$.

Given a Θ^1 -structure s on $\mathcal{O}_{\mathbb{G}^{\mathrm{univ}}}(1)$, there are therefore two naturally induced Θ^1 -structures on $\mathcal{O}_{\psi^* \mathbb{G}^{\mathrm{univ}}}(1) \cong N(\mathcal{O}_{i^* \mathbb{G}^{\mathrm{univ}}}(1))$: the pullback $\psi^*(s)$ and the norm $N(i^*(s))$.

Definition 4.22. We say that a Θ^1 -structure s on $\mathbb{G}^{\mathrm{univ}}$ satisfies the Ando criterion if $\psi^*(s) = N(i^*(s))$.

It follows from [AHS04, §5] that this is compatible with our previous definition of the Ando criterion:

Proposition 4.23. *A complex orientation of $E(k, \mathbb{G})$ satisfies the Ando criterion if and only if the associated Θ^1 -structure on $\mathcal{O}_{\widehat{\mathbb{G}}}(1)$ satisfies the Ando criterion.*

We are now able to prove the main theorem of this section:

Theorem 4.24. *The Hirzebruch level n elliptic genus $\mathrm{MU} \rightarrow \mathrm{tmf}_1(n) \rightarrow \mathrm{TMF}_1(n)$ satisfies the Ando criterion. As a consequence of Theorem 1.4 and Remark 4.4, it lifts uniquely up to homotopy to an \mathbb{E}_∞ -complex orientation*

$$\mathrm{MU} \rightarrow \mathrm{TMF}_1(n).$$

Proof. Choose, for each p not dividing n , $(E, P \in E[n](k)) \in \mathcal{M}_1(n)(k)$ with E supersingular and k a perfect field of characteristic p . By Proposition 4.16, it suffices to show that the induced complex orientation of $E(k, \widehat{E})$ satisfies the Ando criterion.

By the Serre–Tate theorem [Kat81, §1], $\pi_0 E(k, \widehat{E})$ is the universal deformation ring of E . We let E^{univ} denote the universal deformation of E over $\pi_0 E(k, \widehat{E})$, and let $P^{\mathrm{univ}} \in E^{\mathrm{univ}}[n](\pi_0 E(k, \mathbb{G}))$ denote the unique lift of P . The associated formal group $\widehat{E^{\mathrm{univ}}}$ is a universal deformation of \widehat{E} .

Applying Proposition 4.23, we must show that the Θ^1 -structure s on $\mathcal{O}_{\widehat{E^{\mathrm{univ}}}}(1)$ corresponding to the level n elliptic genus satisfies the Ando criterion, i.e. that $\psi^*(s) = N(i^*(s))$. By definition of the level n elliptic genus, there is a degree n étale isogeny $q : (E^{\mathrm{univ}})' \rightarrow E^{\mathrm{univ}}$ and a Θ^1 -structure \bar{s} on $q^* \mathcal{O}_{E^{\mathrm{univ}}}([0] - [P^{\mathrm{univ}}])$ which induces s . By the Serre–Tate theorem, the isogeny of formal groups

$$i^* \widehat{E^{\mathrm{univ}}} \rightarrow \psi^* \widehat{E^{\mathrm{univ}}}$$

over $E(k, \widehat{E})^0(\mathrm{BC}_p)/I_{tr}$ lifts to a diagram of isogenies of elliptic curves

$$\begin{array}{ccc} i^*(E^{\mathrm{univ}})' & \longrightarrow & \psi^*(E^{\mathrm{univ}})' \\ \downarrow q_i & & \downarrow q_\psi \\ i^* E^{\mathrm{univ}} & \longrightarrow & \psi^* E^{\mathrm{univ}}. \end{array}$$

Let $\overline{P^{\mathrm{univ}}}$ denote the image of $i^*(P^{\mathrm{univ}})$ in $\psi^* E^{\mathrm{univ}}$. From the above diagram, we obtain Θ^1 -structures $\psi^*(\bar{s})$ and $N(i^*(\bar{s}))$ on

$$(q_\psi)^* \mathcal{O}_{\psi^* E^{\mathrm{univ}}}([0] - [\overline{P^{\mathrm{univ}}})) \cong N((q_i)^* \mathcal{O}_{i^* E^{\mathrm{univ}}}([0] - [i^*(P^{\mathrm{univ}})])).$$

As these induce $\psi^*(s)$ and $N(i^*(s))$, it suffices to show that $\psi^*(\bar{s}) = N(i^*(\bar{s}))$. But this follows immediately from the uniqueness of Θ^1 -structures over elliptic curves proven in Lemma 4.10. \square

4.5. **Lift to $\mathrm{tmf}_1(n)$.** In this section, we will complete the proof of Theorem 1.7 by proving the following two lemmas:

Lemma 4.25. *Suppose that we are given a complex orientation $\mathrm{MU} \rightarrow \mathrm{Tmf}_1(n)$ with the property that the composite*

$$\mathrm{MU} \rightarrow \mathrm{Tmf}_1(n) \rightarrow \mathrm{TMF}_1(n)$$

lifts to an \mathbb{E}_∞ -ring map. Then this complex orientation lifts uniquely to an \mathbb{E}_∞ -complex orientation $\mathrm{MU} \rightarrow \mathrm{Tmf}_1(n)$.

Lemma 4.26. *Any \mathbb{E}_∞ -complex orientation of $\mathrm{Tmf}_1(n)$ lifts uniquely to an \mathbb{E}_∞ -complex orientation of $\mathrm{tmf}_1(n)$.*

Proof of Lemma 4.25. The uniqueness will follow from Theorem 3.1 and Example 3.3 once we know that a complex orientation of $\mathrm{Tmf}_1(n)$ is determined by the induced complex orientation of $\mathrm{TMF}_1(n)$. This follows from the fact that the map $\pi_* \mathrm{Tmf}_1(n) \rightarrow \pi_* \mathrm{TMF}_1(n)$ is injective in even degrees by [Mei19, Proposition 2.5] and the descent spectral sequence.

It therefore suffices to show that the \mathbb{E}_∞ -map $\mathrm{MU} \rightarrow \mathrm{TMF}_1(n)$ lifts to an \mathbb{E}_∞ -map $\mathrm{MU} \rightarrow \mathrm{Tmf}_1(n)$. We begin with the pullback square of \mathbb{E}_∞ -rings coming from [HL16]:

$$\begin{array}{ccc} \mathrm{Tmf}_1(n) & \longrightarrow & \mathrm{TMF}_1(n) \\ \downarrow & & \downarrow \\ K_1^{\mathrm{cusp}}(n) & \longrightarrow & \Delta^{-1} K_1^{\mathrm{cusp}}(n) \end{array}$$

This square satisfies the following properties:

- (1) The \mathbb{E}_∞ -rings $K_1^{\mathrm{cusp}}(n)$ and $\Delta^{-1} K_1^{\mathrm{cusp}}(n)$ are Landweber exact and of height ≤ 1 .
- (2) The induced map $\pi_* K_1^{\mathrm{cusp}}(n)/p \rightarrow \pi_* \Delta^{-1} K_1^{\mathrm{cusp}}(n)/p$ is injective for all p .

It follows that to construct an \mathbb{E}_∞ -lifting $\mathrm{MU} \rightarrow \mathrm{Tmf}_1(n)$, it suffices to lift the composite

$$\mathrm{MU} \rightarrow \mathrm{TMF}_1(n) \rightarrow \Delta^{-1} K_1^{\mathrm{cusp}}(n)$$

to an \mathbb{E}_∞ -map

$$\mathrm{MU} \rightarrow K_1^{\mathrm{cusp}}(n).$$

By the uniqueness in Theorem 1.4 and (1) above, it suffices to lift the complex orientation and verify that it satisfies the Ando criterion. But we are given a lift of the complex orientation by assumption, and it follows from Lemma 4.17 and (2) above that it satisfies the Ando criterion. \square

Proof of Lemma 4.26. For this, we use orientation theory [ABG⁺14]. We have the sequence of maps

$$\mathrm{bu} \xrightarrow{J} \mathrm{bgl}_1(\mathbb{S}) \rightarrow \mathrm{bgl}_1(\mathrm{tmf}_1(n)) \rightarrow \mathrm{bgl}_1(\mathrm{Tmf}_1(n)),$$

and \mathbb{E}_∞ -complex orientations of tmf_1 and $\mathrm{Tmf}_1(n)$ correspond to nullhomotopies of the respective composites. Since bu is 2-connective, we may as well replace all occurrences of bgl_1 with $\mathrm{bsl}_1 := \tau_{\geq 2} \mathrm{bgl}_1$.

Now, it follows from the definition of $\mathrm{tmf}_1(n)$ [Mei21] that the composite

$$\mathrm{bsl}_1(\mathrm{tmf}_1(n)) \rightarrow \mathrm{bsl}_1(\mathrm{Tmf}_1(n)) \rightarrow \tau_{\geq 3} \mathrm{bsl}_1(\mathrm{Tmf}_1(n))$$

is an equivalence, so that there is a splitting

$$\mathrm{bsl}_1(\mathrm{Tmf}_1(n)) \simeq \mathrm{bsl}_1(\mathrm{tmf}_1(n)) \oplus \Sigma^2 \pi_1 \mathrm{Tmf}_1(n),$$

from which the lemma follows. \square

Proof of Theorem 1.7. Combine Theorem 4.24 with Lemma 4.25 and Lemma 4.26. \square

APPENDIX A. $K(n)$ -LOCALIZATIONS OF LANDWEBER EXACT RING SPECTRA

In this appendix, we prove the following lemma, which the author was unable to find a reference for in the literature:

Lemma A.1. *Let E denote a Landweber exact ring spectrum whose homotopy is concentrated in even degrees. Then $L_{K(n)}E$ is also a Landweber exact ring spectrum whose homotopy is concentrated in even degrees.*

Before we proceed to the proof of Lemma A.1, we need a lemma from commutative algebra.

Lemma A.2. *Let R_* denote a graded commutative ring and let $x_1, \dots, x_n \in R_*$ denote a regular sequence of homogeneous elements. Then the sequence x_1, \dots, x_n remains regular in the completion $(R_*)_{(x_1, \dots, x_n)}^\wedge$.*

Proof. Let $I = (x_1, \dots, x_n)$. We claim that there are short exact sequences

$$0 \rightarrow R_*/I^{k-1} \xrightarrow{x_1} R_*/I^k \rightarrow R_*/(I^k + (x_1)) \rightarrow 0 \quad (2)$$

for all $k \geq 1$. Supposing this for the moment, we find by taking the limit that

$$0 \rightarrow (R_*)_I^\wedge \xrightarrow{x_1} (R_*)_I^\wedge \rightarrow (R_*/x_1)_I^\wedge \rightarrow 0$$

is short exact. In particular, we find that x_1 is a regular element in $(R_*)_I^\wedge$ and that $(R_*)_I^\wedge/x_1 \cong (R_*/x_1)_I^\wedge$. Taking R_*/x_1 as our new ring and the image of x_2, \dots, x_n in R_*/x_1 as our exact sequence, we may conclude by induction on the length of our regular sequence.

It remains to establish (2). It is sufficient to prove that (2) is exact on the associated graded of the I -adic filtration where x_1 is considered as a map of I -adic filtration degree 1. Since x_1, \dots, x_n is regular, this associated graded may be identified with the sequence

$$\begin{array}{c} 0 \rightarrow (R_*/I)[\bar{x}_1, \dots, \bar{x}_n]/(\bar{x}_1, \dots, \bar{x}_n)^{k-1} \xrightarrow{\bar{x}_1} (R_*/I)[\bar{x}_1, \dots, \bar{x}_n]/(\bar{x}_1, \dots, \bar{x}_n)^k \\ \left. \begin{array}{c} \xrightarrow{\hspace{10em}} \\ \xrightarrow{\hspace{10em}} \end{array} \right\} \\ (R_*/I)[\bar{x}_2, \dots, \bar{x}_n]/(\bar{x}_2, \dots, \bar{x}_n)^k \longrightarrow 0, \end{array}$$

which is easily verified to be short exact. \square

Proof of Lemma A.1. By abuse of notation, we let $v_i \in \pi_*E$ inductively denote an arbitrary lift of the class $v_i \in \pi_*E/(p, \dots, v_{i-1})$. Given a positive integer k , let $I_k = (p, v_1, \dots, v_{k-1})$ and let $I_0 = (0)$. It follows from [HS99b, Proposition 7.10] that $\pi_*L_{K(n)}E \cong (v_n^{-1}\pi)_I^\wedge$. In particular, $\pi_{2*-1}L_{K(n)}E = 0$. It is clear that the operation of inverting v_n^{-1} preserves Landweber exactness, so that it suffices to prove that completion with respect to I_n does as well. But this follows immediately from Lemma A.2. \square

REFERENCES

- [ABG⁺14] Matthew Ando, Andrew J. Blumberg, David Gepner, Michael J. Hopkins, and Charles Rezk. Units of ring spectra, orientations and Thom spectra via rigid infinite loop space theory. *J. Topol.*, 7(4):1077–1117, 2014.
- [ABS64] M. F. Atiyah, R. Bott, and A. Shapiro. Clifford modules. *Topology*, 3(suppl. 1):3–38, 1964.
- [Abs21] Dominik Absmeier. Strictly commutative complex orientations of $Tmf_1(N)$. 2021. arXiv:2108.03079.
- [AHR10] Matthew Ando, Michael J. Hopkins, and Charles Rezk. Multiplicative orientations of ko -theory and of the spectrum of topological modular forms. 2010. Unpublished preprint.
- [AHS01] M. Ando, M. J. Hopkins, and N. P. Strickland. Elliptic spectra, the Witten genus and the theorem of the cube. *Invent. Math.*, 146(3):595–687, 2001.
- [AHS04] Matthew Ando, Michael J. Hopkins, and Neil P. Strickland. The sigma orientation is an H_∞ map. *Amer. J. Math.*, 126(2):247–334, 2004.

- [AL07] Gregory Arone and Kathryn Lesh. Filtered spectra arising from permutative categories. *J. Reine Angew. Math.*, 604:73–136, 2007.
- [Bal21] William Balderrama. Algebraic theories of power operations. 2021. arXiv:2108.06802v2.
- [BE21] Daniel Berwick-Evans. Supersymmetric field theories and the elliptic index theorem with complex coefficients. *Geom. Topol.*, 25(5):2287–2384, 2021.
- [BH16] Tobias Barthel and Drew Heard. The E_2 -term of the $K(n)$ -local E_n -Adams spectral sequence. *Topology Appl.*, 206:190–214, 2016.
- [Bou87] A. K. Bousfield. Uniqueness of infinite deloopings for K -theoretic spaces. *Pacific J. Math.*, 129(1):1–31, 1987.
- [Cos10] Kevin Costello. A geometric construction of the Witten genus, I. In *Proceedings of the International Congress of Mathematicians. Volume II*, pages 942–959. Hindustan Book Agency, New Delhi, 2010.
- [Cos11] Kevin Costello. A geometric construction of the Witten genus, II. 2011. arXiv:1112.0816.
- [DFHH14] Christopher L. Douglas, John Francis, André G. Henriques, and Michael A. Hill, editors. *Topological modular forms*, volume 201 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2014.
- [DH11] Christopher L. Douglas and André G. Henriques. Topological modular forms and conformal nets. In *Mathematical foundations of quantum field theory and perturbative string theory*, volume 83 of *Proc. Sympos. Pure Math.*, pages 341–354. Amer. Math. Soc., Providence, RI, 2011.
- [DR73] P. Deligne and M. Rapoport. Les schémas de modules de courbes elliptiques. In *Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972)*, pages 143–316. Lecture Notes in Math., Vol. 349, 1973.
- [GH04] P. G. Goerss and M. J. Hopkins. Moduli spaces of commutative ring spectra. In *Structured ring spectra*, volume 315 of *London Math. Soc. Lecture Note Ser.*, pages 151–200. Cambridge Univ. Press, Cambridge, 2004.
- [GS06] Philippe Gille and Tamás Szamuely. *Central simple algebras and Galois cohomology*, volume 101 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2006.
- [Hir88] Friedrich Hirzebruch. Elliptic genera of level N for complex manifolds. In *Differential geometrical methods in theoretical physics (Como, 1987)*, volume 250 of *NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci.*, pages 37–63. Kluwer Acad. Publ., Dordrecht, 1988.
- [HKR00] Michael J. Hopkins, Nicholas J. Kuhn, and Douglas C. Ravenel. Generalized group characters and complex oriented cohomology theories. *J. Amer. Math. Soc.*, 13(3):553–594, 2000.
- [HL16] Michael Hill and Tyler Lawson. Topological modular forms with level structure. *Invent. Math.*, 203(2):359–416, 2016.
- [HL18] Michael J. Hopkins and Tyler Lawson. Strictly commutative complex orientation theory. *Math. Z.*, 290(1-2):83–101, 2018.
- [HS99a] Mark Hovey and Hal Sadofsky. Invertible spectra in the $E(n)$ -local stable homotopy category. *J. London Math. Soc. (2)*, 60(1):284–302, 1999.
- [HS99b] Mark Hovey and Neil P. Strickland. Morava K -theories and localisation. *Mem. Amer. Math. Soc.*, 139(666):viii+100, 1999.
- [Joa04] Michael Joachim. Higher coherences for equivariant K -theory. In *Structured ring spectra*, volume 315 of *London Math. Soc. Lecture Note Ser.*, pages 87–114. Cambridge Univ. Press, Cambridge, 2004.
- [Kat81] N. Katz. Serre-Tate local moduli. In *Algebraic surfaces (Orsay, 1976–78)*, volume 868 of *Lecture Notes in Math.*, pages 138–202. Springer, Berlin-New York, 1981.
- [Kuh89] Nicholas J. Kuhn. Morava K -theories and infinite loop spaces. In *Algebraic topology (Arcata, CA, 1986)*, volume 1370 of *Lecture Notes in Math.*, pages 243–257. Springer, Berlin, 1989.
- [Lur18a] Jacob Lurie. Elliptic Cohomology I. 2018. Available at <http://www.math.ias.edu/lurie/>.
- [Lur18b] Jacob Lurie. Elliptic Cohomology II: Orientations. 2018. Available at <http://www.math.ias.edu/lurie/>.
- [Mei19] Lennart Meier. Topological modular forms with level structure: decompositions and duality. 2019. arXiv:1806.06709v2.
- [Mei20] Lennart Meier. Additive decompositions for rings of modular forms. 2020. arXiv:1710.03461v3.
- [Mei21] Lennart Meier. Connective models for topological modular forms of level n . 2021. arXiv:2104.12649v1.
- [Oli94] Bob Oliver. p -stubborn subgroups of classical compact Lie groups. *J. Pure Appl. Algebra*, 92(1):55–78, 1994.

- [Qui71] Daniel Quillen. Elementary proofs of some results of cobordism theory using Steenrod operations. *Advances in Math.*, 7:29–56 (1971), 1971.
- [Rez13] Charles Rezk. Power operations in Morava e -theory: structure and calculations (draft). 2013. Available at <https://faculty.math.illinois.edu/rezk/power-ops-ht-2.pdf>.
- [RWY98] Douglas C. Ravenel, W. Stephen Wilson, and Nobuaki Yagita. Brown-Peterson cohomology from Morava K -theory. *K-Theory*, 15(2):147–199, 1998.
- [ST04] Stephan Stolz and Peter Teichner. What is an elliptic object? In *Topology, geometry and quantum field theory*, volume 308 of *London Math. Soc. Lecture Note Ser.*, pages 247–343. Cambridge Univ. Press, Cambridge, 2004.
- [ST11] Stephan Stolz and Peter Teichner. Supersymmetric field theories and generalized cohomology. In *Mathematical foundations of quantum field theory and perturbative string theory*, volume 83 of *Proc. Sympos. Pure Math.*, pages 279–340. Amer. Math. Soc., Providence, RI, 2011.
- [Sta21] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2021.
- [Str99] Neil P. Strickland. Formal schemes and formal groups. In *Homotopy invariant algebraic structures (Baltimore, MD, 1998)*, volume 239 of *Contemp. Math.*, pages 263–352. Amer. Math. Soc., Providence, RI, 1999.
- [SY04] Björn Schuster and Nobuaki Yagita. Morava K -theory of extraspecial 2-groups. *Proc. Amer. Math. Soc.*, 132(4):1229–1239, 2004.
- [TY89] M. Tezuka and N. Yagita. Cohomology of finite groups and Brown-Peterson cohomology. In *Algebraic topology (Arcata, CA, 1986)*, volume 1370 of *Lecture Notes in Math.*, pages 396–408. Springer, Berlin, 1989.
- [Wil15] Dylan Wilson. Orientations and Topological Modular Forms with Level Structure. 2015. arXiv:1507.05116v1.
- [Wit87] Edward Witten. Elliptic genera and quantum field theory. *Comm. Math. Phys.*, 109(4):525–536, 1987.
- [Wit88] Edward Witten. The index of the Dirac operator in loop space. In *Elliptic curves and modular forms in algebraic topology (Princeton, NJ, 1986)*, volume 1326 of *Lecture Notes in Math.*, pages 161–181. Springer, Berlin, 1988.
- [Yag05] Nobuaki Yagita. Integral and BP cohomologies of extraspecial p -groups for odd primes. *Kodai Math. J.*, 28(1):1–30, 2005.
- [Zhu20] Yifei Zhu. Norm coherence for descent of level structures on formal deformations. *J. Pure Appl. Algebra*, 224(10):106382, 35, 2020.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA, USA
Email address: senger@math.harvard.edu