arXiv:2203.14030v1 [math.NT] 26 Mar 2022

WEIGHTED SUM FORMULAS FROM SHUFFLE PRODUCTS OF MULTIPLE ZETA-STAR VALUES

KWANG-WU CHEN AND MINKING EIE

ABSTRACT. In this paper, we are going to perform the shuffle products of $Z_-(n)=\sum_{a+b=m}(-1)^b\zeta(\{1\}^a,b+2)$ and $Z_+^\star(n)=\sum_{c+d=n}\zeta^\star(\{1\}^c,d+2)$ with $m+n=p$. The resulted shuffle relation is a weighted sum formula given by

$$\frac{(p+1)(p+2)}{2}\zeta(p+4)=\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)$$
$$\times\sum_{a+b+c=m}\Big(W_{\boldsymbol{\alpha}}(a,b,c)+W_{\boldsymbol{\alpha}}(a,b,c=0)+W_{\boldsymbol{\alpha}}(a=0,b,c)+W_{\boldsymbol{\alpha}}(a=0,b=m,c=0)\Big),$$

where $W_{\boldsymbol{\alpha}}(a,b,c)=2^{\sigma(a+b+1)-\sigma(a)-(b+1)}(1-2^{1-\alpha_{a+b+1}})$, with $\sigma(r)=\sum_{j=0}^{r}\alpha_j$.

1. Introduction

For an r-tuple $\boldsymbol{\alpha}=(\alpha_1,\alpha_2,\ldots,\alpha_r)$ of positive integers with $\alpha_r\geq 2$, a multiple zeta value $\zeta(\boldsymbol{\alpha})$ and a multiple zeta-star value $\zeta^\star(\boldsymbol{\alpha})$ are defined to be [9,10]

$$\zeta(\boldsymbol{\alpha})=\sum_{1\leq k_1<k_2<\cdots<k_r}k_1^{-\alpha_1}k_2^{-\alpha_2}\cdots k_r^{-\alpha_r},$$

and

$$\zeta^\star(\boldsymbol{\alpha})=\sum_{1\leq k_1\leq k_2\leq\cdots\leq k_r}k_1^{-\alpha_1}k_2^{-\alpha_2}\cdots k_r^{-\alpha_r}.$$

We denote the parameters $w(\boldsymbol{\alpha})=|\,\boldsymbol{\alpha}\,|=\alpha_1+\alpha_2+\cdots+\alpha_r$, $d(\boldsymbol{\alpha})=r$, and $h(\boldsymbol{\alpha})=\#\{i\mid\alpha_i>1,1\leq i\leq r\}$, called respectively the weight, the depth, and the height of $\boldsymbol{\alpha}$ (or of $\zeta(\boldsymbol{\alpha})$, or of $\zeta^\star(\boldsymbol{\alpha})$). For any multiple zeta value $\zeta(\boldsymbol{\alpha})$, we put a bar on top of α_j $(j=1,2,\ldots,r)$ if there is a sign $(-1)^{k_j}$ appearing in the numerator of the summand [18,19]. We call it an alternating multiple zeta value. For example, an alternating multiple zeta value

$$\zeta(\overline{1},\{1\}^m,\overline{2})=\sum_{1\leq k_1<k_2<\cdots<k_{m+2}}\frac{(-1)^{k_1+k_{m+2}}}{k_1\cdots k_{m+1}k_{m+2}^2},$$

where $\{a\}^k$ is k repetitions of a. Its weight is $m+3$, the height is one, and the depth is $m+2$.

Due to Kontsevich, multiple zeta values can be represented by iterated integrals over a simplex of dimension weight:

$$\zeta(\alpha_1,\alpha_2,\ldots,\alpha_r)=\int_{E_{|\boldsymbol{\alpha}|}}\Omega_1\Omega_2\cdots\Omega_{|\boldsymbol{\alpha}|}\quad\text{or}\quad\int_0^1\Omega_1\Omega_2\cdots\Omega_{|\boldsymbol{\alpha}|}$$

Date: 2022-03-26; Version 7.

2020 *Mathematics Subject Classification.* Primary: 11M32; Secondary: 05A15, 33B15.

Key words and phrases. Multiple zeta value, Shuffle product.

The first author (corresponding author) was funded by the Ministry of Science and Technology, Taiwan, R.O.C., under Grant MOST 110-2115-M-845-001.

with $E_{|\boldsymbol{\alpha}|}: 0 < t_1 < t_2 < \cdots < t_{|\boldsymbol{\alpha}|} < 1$ and

$$\Omega_j = \begin{cases} \dfrac{dt_j}{1-t_j} & \text{if } j = 1, \alpha_1+1, \alpha_1+\alpha_2+1, \ldots, \alpha_1+\alpha_2+\cdots+\alpha_{r-1}+1, \\ \dfrac{dt_j}{t_j} & \text{otherwise.} \end{cases}$$

Sometime, we express them as words with just two alphabets:

$$a\{b\}^{\alpha_1-1}a\{b\}^{\alpha_2-1}\cdots a\{b\}^{\alpha_r-1}.$$

The dual then is obtained by exchanging a, b and reversing the order. For example, we express $\zeta(5)$ as ab^4, its dual is a^4b which is $\zeta(1,1,1,2)$.

Once multiple zeta values are expressed as iterated integrals, the shuffle product of two multiple zeta values can be defined as

$$\int_0^1 \Omega_1\Omega_2\cdots\Omega_m \int_0^1 \Omega_{m+1}\Omega_{m+2}\cdots\Omega_{m+n} = \sum_{\sigma}\int_0^1 \Omega_{\sigma(1)}\Omega_{\sigma(2)}\cdots\Omega_{\sigma(m+n)},$$

where σ ranges over all permutations on the set $\{1,2,\ldots,m+n\}$ that preserve the orderings of $\Omega_1\Omega_2\cdots\Omega_m$ and $\Omega_{m+1}\Omega_{m+2}\cdots\Omega_{m+n}$. This is equivalent to saying that for all $1 \le i < j \le m$ or $m+1 \le i < j \le m+n$, we have $\sigma^{-1}(i) < \sigma^{-1}(j)$. It is clear that the shuffle product of two multiple zeta values of weight m and n, respectively, will produce $\binom{m+n}{n}$ multiple zeta values of weight $m+n$.

In this paper, we will investigate the following integral

$$\frac{1}{p!}\int_{\substack{0<t_1<t_2<1\\0<u_1<u_2<1}} \left(\log\frac{1-t_1}{1-u_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p \frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2}.$$

We derive our main theorem by finding its different representations.

Main Theorem For integer $p \ge 0$, we have

$$\frac{(p+1)(p+2)}{2}\zeta(p+4) = \sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)$$
$$\times\sum_{a+b+c=m}\Big(W_{\boldsymbol{\alpha}}(a,b,c)+W_{\boldsymbol{\alpha}}(a,b,c=0)+W_{\boldsymbol{\alpha}}(a=0,b,c)+W_{\boldsymbol{\alpha}}(a=0,b=m,c=0)\Big),$$

where $W_{\boldsymbol{\alpha}}(a,b,c) = 2^{\sigma(a+b+1)-\sigma(a)-(b+1)}(1-2^{1-\alpha_{a+b+1}})$, with $\sigma(r) = \sum_{j=0}^{r}\alpha_j$.

We note that the second, third, and fourth weight in the sum formula are

$$\sum_{a+b+c=m} W_{\boldsymbol{\alpha}}(a,b,c=0) = \sum_{a+b=m} W_{\boldsymbol{\alpha}}(a,b,0) = \sum_{a=0}^{m} 2^{\sigma(m+1)-\sigma(a)-(m-a+1)}(1-2^{1-\alpha_{m+1}}),$$
$$\sum_{a+b+c=m} W_{\boldsymbol{\alpha}}(a=0,b,c) = \sum_{b+c=m} W_{\boldsymbol{\alpha}}(0,b,c) = \sum_{b=0}^{m} 2^{\sigma(b+1)-\alpha_0-(b+1)}(1-2^{1-\alpha_{b+1}}),$$
$$\sum_{a+b+c=m} W_{\boldsymbol{\alpha}}(a=0,b=m,c=0) = W_{\boldsymbol{\alpha}}(0,m,0).$$

For brevity, we hereafter use the lowercase English letters and lowercase Greek letters in the summation, with or without subscripts, to denote the nonnegative and positive integers, unless otherwise specified.

In fact, our main theorem is from the integral

$$J_p = \frac{1}{p!}\int_{\substack{0<t_1<t_2<1\\0<u_1<u_2<1}} \left(\log\frac{1-t_1}{1-u_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p \frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2}.$$

As a replacement of shuffle product, we decompose the domain $\{(t_1,t_2,u_1,u_2)\in\mathbb{R}^4 \mid 0<t_1<t_2<1, 0<u_1<u_2<1\}$ into 6 disjoint simplices of dimension 4:

$$\begin{aligned} &D_1: 0<t_1<t_2<u_1<u_2<1, \quad D_2: 0<u_1<u_2<t_1<t_2<1,\\ &D_3: 0<t_1<u_1<t_2<u_2<1, \quad D_4: 0<t_1<u_1<u_2<t_2<1,\\ &D_5: 0<u_1<t_1<u_2<t_2<1, \quad D_6: 0<u_1<t_1<t_2<u_2<1. \end{aligned}$$

So

$$J_p = \sum_{j=1}^{6}\mathbb{J}(j)$$

with

$$\mathbb{J}(j) = \frac{1}{p!}\int_{D_j} \left(\log\frac{1-t_1}{1-u_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p \frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2}.$$

The main task now is to evaluate $\mathbb{J}(j)$ one by one in terms of multiple zeta values. The weighted sum in our main theorem is $\sum_{j=3}^{6}\mathbb{J}(j)$. The crucial step is to prove

$$\mathbb{J}(1)+\mathbb{J}(2) = J_p - \frac{(p+1)(p+2)}{2}\zeta(p+4) \tag{1.1}$$

and the resulted shuffle relation is

$$\sum_{j=3}^{6}\mathbb{J}(j) = \frac{(p+1)(p+2)}{2}\zeta(p+4). \tag{1.2}$$

We organize this paper as follows: We give some preliminaries and auxiliary tools in Section 2. In Section 3, we break J_p into six parts and evaluate $\mathbb{J}(1)$ and $\mathbb{J}(2)$ in terms of multiple zeta values. In the next section we express $\mathbb{J}(j)$ for $3\le j\le 6$ as the part of the weighted sum of our main theorem. We prove the crucial step Eq. (1.1) in Section 5. In Section 6, we give another application of the identity Eq. (1.2) to another sum formula wihich is also proved by a duality theorem due to Ohno [15]. In Section 7, we relate $\mathbb{J}(3)$ with the weighted sum

$$\sum_{|\boldsymbol{\alpha}|=n+3} 2^{\alpha_2}\zeta^\star(\alpha_1,\{1\}^m,\alpha_2+1).$$

In the final section, we will give some formulas related to weighted alternating Euler sums, e.g.,

$$\sum_{m+n=p} n\zeta(m+1,\overline{n+1}), \quad \text{and} \quad \sum_{m+n=p} n(n-1)\zeta(m+1,\overline{n+1}).$$

2. Some Preliminaries and Auxiliary Tools

Here we list a general integral representation which we use frequently.

Proposition 2.1 [11] For nonnegative integers $b_1,b_2,\dots,b_r,b_{r+1}$, we have

$$\zeta(b_1+1,b_2+1,\dots,b_r+1,b_{r+1}+2) = \frac{1}{b_1!b_2!\cdots b_{r+1}!}\int_{E_{r+2}}\prod_{j=1}^{r+1}\frac{\mathrm{d}t_j}{1-t_j}\left(\log\frac{t_{j+1}}{t_j}\right)^{b_j}\frac{\mathrm{d}t_{r+2}}{t_{r+2}},$$

where E_{r+2} is a simplex defined as $0<t_1<t_2<\cdots<t_{r+2}$.

Proposition 2.2 [11, Proposition 2.1] For integers $p, q, r, \ell \geq 0$,

$$\sum_{|\boldsymbol{\alpha}|=q+r+1} \zeta(\{1\}^p, \alpha_0, \alpha_1, \ldots, \alpha_q + \ell + 1)$$
$$= \frac{1}{p!q!r!\ell!} \int_{0<t_1<t_2<1} \left(\log \frac{1}{1-t_1}\right)^p \left(\log \frac{1-t_1}{1-t_2}\right)^q \left(\log \frac{t_2}{t_1}\right)^r \left(\log \frac{1}{t_2}\right)^\ell \frac{dt_1 dt_2}{(1-t_1)t_2}.$$

The well-known sum formula due to Granville [12] asserted that

$$\sum_{|\boldsymbol{\alpha}|=q+r+1} \zeta(\alpha_0, \alpha_1, \ldots, \alpha_q + 1) = \zeta(q+r+2).$$

If we let $p = \ell = 0$ in the formula of Proposition 2.2, then we have

$$\sum_{|\boldsymbol{\alpha}|=q+r+1} \zeta(\alpha_0, \alpha_1, \ldots, \alpha_q + 1) = \frac{1}{q!r!} \int_{0<t_1<t_2<1} \left(\log \frac{1-t_1}{1-t_2}\right)^q \left(\log \frac{t_2}{t_1}\right)^r \frac{dt_1 dt_2}{(1-t_1)t_2}. \tag{2.1}$$

Therefore the integral

$$\frac{1}{q!r!} \int_{0<t_1<t_2<1} \left(\log \frac{1-t_1}{1-t_2}\right)^q \left(\log \frac{t_2}{t_1}\right)^r \frac{dt_1 dt_2}{(1-t_1)t_2} = \zeta(q+r+2).$$

Our main results are obtained by the convolution of the following sums:

$$Z_-(m) = \sum_{a+b=m} (-1)^b \zeta(\{1\}^a, b+2) \quad \text{and} \quad Z_+^\star(n) = \sum_{c+d=n} \zeta^\star(\{1\}^c, d+2). \tag{2.2}$$

A special case of results proved by Le and Murakami [14] is stated as follows.

$$\sum_{a+b=2w-2} (-1)^{a+1} \zeta(\{1\}^a, b+2) = 2\zeta(\overline{2w}). \tag{2.3}$$

Both $Z_-(n)$, and $Z_+^\star(n)$ are sums of multiple zeta(-star) values of height one and can be expressed as double integrals (see [6])

$$Z_-(n) = \frac{1}{n!} \int_{E_2} \left(\log \frac{1}{1-t_1} - \log \frac{t_2}{t_1}\right)^n \frac{\mathrm{d}t_1 \mathrm{d}t_2}{(1-t_1)t_2},$$
$$Z_+^\star(n) = \frac{1}{n!} \int_{E_2} \left(\log \frac{1}{1-t_2} + \log \frac{t_2}{t_1}\right)^n \frac{\mathrm{d}t_1 \mathrm{d}t_2}{(1-t_1)t_2}.$$

Through the duality theorem $\zeta(\{1\}^a, b+2) = \zeta(\{1\}^b, a+2)$, we see that $Z_-(2m+1) = 0$. Therefore, we combine these with Eq. (2.3), then we have

Proposition 2.3 [6, Proposition 3.1] For any nonnegative integer m, we have $Z_-(2m+1) = 0$ and

$$Z_-(2m) = \zeta^\star(\{2\}^{m+1}) = 2\left(1 - \frac{1}{2^{2m+1}}\right)\zeta(2m+2) = -2\zeta(\overline{2m+2}).$$

The sum of multiple zeta-star values $Z_+^\star(n)$ appeared as the principal term of the evaluation of $\zeta^\star(3, \{2\}^n)$ (see [5]):

$$\zeta^\star(3, \{2\}^n) = Z_+^\star(2n+1) - 2 \sum_{a+b=n} \zeta(\{2\}^a)\zeta(2b+3).$$

Arakawa and Kaneko [1] defined the function

$$\xi_k(s) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{t^{s-1}}{e^t - 1} \mathrm{Li}_k(1 - e^{-t})\, dt,$$

where $\mathrm{Li}_k(s)$ denotes the k-th polylogarithm $\mathrm{Li}_k(s)=\sum_{n=1}^{\infty}\frac{s^n}{n^k}$. It is exactly the multiple zeta-star values of height one

$$\xi_k(s)=\zeta^\star(\{1\}^{s-1},k+1).$$

Many properties of the generalized Arakawa-Kaneko zeta functions have been discovered recently (ref. [3,8, 13]).

Indeed, $Z_+^\star(n)$ has the generating function

$$\sum_{n=0}^{\infty}Z_+^\star(n)(-x)^n=\int_{0<t_1<t_2<1}(1-t_2)^x\left(\frac{t_1}{t_2}\right)^x\frac{dt_1dt_2}{(1-t_1)t_2}.$$

The dual of the above is

$$\int_{0<u_1<u_2<1}\left(\frac{1-u_2}{1-u_1}\right)^x u_1^x\frac{du_1du_2}{(1-u_1)u_2},$$

which can be evaluated as

$$\sum_{k=1}^{\infty}\frac{1}{(k+x)(k+2x)}\frac{\Gamma(k+x)^2}{\Gamma(k)\Gamma(k+2x)}.$$

Fortunately, we have the identity

$$\sum_{k=1}^{\infty}\frac{1}{(k+x)(k+2x)}\frac{\Gamma(k+x)^2}{\Gamma(k)\Gamma(k+2x)}=\sum_{\ell=1}^{\infty}\frac{2(-1)^{\ell-1}}{(\ell+x)^2}$$

by investigating possible poles of both sides of the meromorphic functions. This leads to the evaluation of $Z_+^\star(n)$:

$$\sum_{a+b=w-2}\zeta^\star(\{1\}^a,b+2)=2(w-1)(1-2^{1-w})\zeta(w)=-2(w-1)\zeta(\overline{w}),\tag{2.4}$$

where $w>1$ is an integer. This formula was first proved by Ohno [16, Theorem 8] in 2005.

3. The Main Integral J_p and its $\mathbb{J}(1),\mathbb{J}(2)$ Parts

We begin with the integral

$$J_p=\frac{1}{p!}\int_{E_2\times E_2}\left(\log\frac{1-t_1}{1-u_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p\frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2},$$

where $E_2\times E_2=\{(t_1,t_2,u_1,u_2)\in\mathbb{R}^4\mid 0<t_1<t_2<1,0<u_1<u_2<1\}$. As

$$\begin{aligned}&\frac{1}{p!}\left(\log\frac{1-t_1}{1-u_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p\\&=\frac{1}{p!}\left\{\left(-\log\frac{1}{1-t_1}+\log\frac{t_2}{t_1}\right)+\left(\log\frac{1}{1-u_2}+\log\frac{u_2}{u_1}\right)\right\}^p\\&=\sum_{m+n=p}\frac{1}{m!n!}\left(-\log\frac{1}{1-t_1}+\log\frac{t_2}{t_1}\right)^m\left(\log\frac{1}{1-u_2}+\log\frac{u_2}{u_1}\right)^n,\end{aligned}$$

so we have the evaluation

$$J_p=\sum_{m+n=p}(-1)^mZ_-(m)Z_+^\star(n)$$

where $Z_-(m)$ and $Z_+^\star(n)$ are defined in Eq (2.2). By Proposition 2.3 $Z_-(2m+1)=0$, thus

$$J_p=\sum_{m=0}^{[p/2]}Z_-(2m)Z_+^\star(p-2m)=\sum_{m+n=p}Z_-(m)Z_+^\star(n).$$

We got an expression of the finite convolution of $Z_-(m)$ and $Z_+^\star(n)$ in [6, Corollary 5.4]:

$$J_p = 2((-1)^p - 1)\zeta(p+2,\overline{2}) + (p+2)(p+1+2^{-p-2})\zeta(p+4).$$

In this paper, we consider another expression of J_p. When we decompose $E_2 \times E_2$ into 6 simplices of dimension 4:

$$D_1 : 0 < t_1 < t_2 < u_1 < u_2 < 1, \quad D_2 : 0 < u_1 < u_2 < t_1 < t_2 < 1,$$
$$D_3 : 0 < t_1 < u_1 < t_2 < u_2 < 1, \quad D_4 : 0 < t_1 < u_1 < u_2 < t_2 < 1,$$
$$D_5 : 0 < u_1 < t_1 < u_2 < t_2 < 1, \quad D_6 : 0 < u_1 < t_1 < t_2 < u_2 < 1$$

and let

$$\mathbb{J}(j) = \frac{1}{p!}\int_{D_j}\left(\log\frac{1-t_1}{1-u_2} + \log\frac{t_2}{t_1} + \log\frac{u_2}{u_1}\right)^p \frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2}.$$

Then J_p is breaked into 6 parts:

$$J_p = \sum_{j=1}^{6}\mathbb{J}(j).$$

In the following, we will evaluate $\mathbb{J}(j)$, for $1 \le j \le 6$, in terms of multiple zeta values.

Proposition 3.1 Given any nonnegative integer p, we have

$$\mathbb{J}(1) = \sum_{m+n=p}\sum_{a+b+c=m}\sum_{u+v=n}\sum_{\substack{|\boldsymbol{\alpha}|=a+u+1\\|\boldsymbol{\beta}|=c+v+1}} \zeta(\alpha_0,\alpha_1,\ldots,\alpha_a+1,\{1\}^b,\beta_0,\ldots,\beta_c+1).$$

Proof. We first expand the integrand of $\mathbb{J}(1)$ as

$$\sum_{m+n=p}\frac{1}{m!n!}\left(\log\frac{1-t_1}{1-u_2}\right)^m\left(\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^n.$$

Since $\mathbb{J}(1)$ is an integral on $D_1 : 0 < t_1 < t_2 < u_1 < u_2 < 1$, we replace the factor

$$\log\frac{1-t_1}{1-u_2}$$

by its equal

$$\log\frac{1-t_1}{1-t_2} + \log\frac{1-t_2}{1-u_1} + \log\frac{1-u_1}{1-u_2}.$$

Then we expand the integrand of $\mathbb{J}(1)$ as

$$\sum_{m+n=p}\sum_{a+b+c=m}\sum_{u+v=n}\frac{1}{a!b!c!u!v!}$$
$$\times\left(\log\frac{1-t_1}{1-t_2}\right)^a\left(\log\frac{1-t_2}{1-u_1}\right)^b\left(\log\frac{1-u_1}{1-u_2}\right)^c\left(\log\frac{t_2}{t_1}\right)^u\left(\log\frac{u_2}{u_1}\right)^v.$$

The factor

$$\frac{1}{a!u!}\left(\log\frac{1-t_1}{1-t_2}\right)^a\left(\log\frac{t_2}{t_1}\right)^u$$

forms a sum and the other factor

$$\frac{1}{c!v!}\left(\log\frac{1-u_1}{1-u_2}\right)^c\left(\log\frac{u_2}{u_1}\right)^v$$

forms another sum, so

$$\mathbb{J}(1)=\sum_{m+n=p}\sum_{a+b+c=m}\sum_{u+v=n}\sum_{\substack{|\boldsymbol{\alpha}|=a+u+1\\|\boldsymbol{\beta}|=c+v+1}}\zeta(\alpha_0,\alpha_1,\dots,\alpha_a+1,\{1\}^b,\beta_0,\dots,\beta_c+1). \qquad \square$$

Proposition 3.2 For any nonnegative integer p, we have

$$\mathbb{J}(2)=\sum_{c+d+m=p}(-1)^m\zeta(c+2,\{1\}^m,d+2).$$

Proof. On $D_2:0<u_1<u_2<t_1<t_2<1$, the factor

$$\log\frac{1-t_1}{1-u_2}$$

is replaced by

$$-\log\frac{1-u_2}{1-t_1}$$

and the integrand

$$\frac{1}{p!}\left(-\log\frac{1-u_2}{1-t_1}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p$$

is expanded into

$$\sum_{c+d+m=p}\frac{(-1)^m}{c!\,d!\,m!}\left(\log\frac{1-u_2}{1-t_1}\right)^m\left(\log\frac{u_2}{u_1}\right)^c\left(\log\frac{t_2}{t_1}\right)^d.$$

So that

$$\mathbb{J}(2)=\sum_{c+d+m=p}(-1)^m\zeta(c+2,\{1\}^m,d+2). \qquad \square$$

It is noting that we have another expression of $\mathbb{J}(2)$ in [7, Theorem 3]:

$$\sum_{a+b+m=p}(-1)^m\zeta(a+2,\{1\}^m,b+2)=2[(-1)^p-1]\zeta(p+2,\overline{2})+\left(1-2^{-p-2}\right)\zeta(p+4).$$

4. The Integrals $\mathbb{J}(3),\mathbb{J}(4),\mathbb{J}(5)$, and $\mathbb{J}(6)$ Parts

In this section, we will give the evaluations of $\mathbb{J}(j)$ for $3\le j\le 6$. These expressions give us the part of the weighted sum of our main theorem. Our method of proving the following proposition is inspired by the approach used in [17, Theorem 5.3].

Proposition 4.1 Let p be a nonnegative integer. We have

$$\mathbb{J}(3)=\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\dots,\alpha_m,\alpha_{m+1}+1)\sum_{a+b+c=m}W(a,b,c)$$

with $\sigma(r)=\alpha_0+\alpha_1+\cdots+\alpha_r$ and

$$W(a,b,c)=2^{\sigma(a+b+1)-\sigma(a)-(b+1)}(1-2^{1-\alpha_{a+b+1}}). \tag{4.1}$$

Proof. The integral $\mathbb{J}(3)$ is on $D_3:0<t_1<u_1<t_2<u_2<1$. Therefore we replace the integrand of $\mathbb{J}(3)$ by

$$\frac{1}{p!}\left(\log\frac{1-t_1}{1-u_1}+\log\frac{1-u_1}{1-t_2}+\log\frac{1-t_2}{1-u_2}+\log\frac{u_1}{t_1}+2\log\frac{t_2}{u_1}+\log\frac{u_2}{t_2}\right)^p.$$

Then we expand it as

$$\sum_{\substack{m+n=p\\ a+b+c=m\\ u+v+w=n}} \frac{2^v}{a!b!c!u!v!w!}\left(\log\frac{1-t_1}{1-u_1}\right)^a\left(\log\frac{1-u_1}{1-t_2}\right)^b\left(\log\frac{1-t_2}{1-u_2}\right)^c\left(\log\frac{u_1}{t_1}\right)^u\left(\log\frac{t_2}{u_1}\right)^v\left(\log\frac{u_2}{t_2}\right)^w.$$

This yields

$$\mathbb{J}(3)=\sum_{\substack{m+n=p\\ a+b+c=m\\ u+v+w=n}} 2^v \sum_{\substack{|\boldsymbol{\alpha}|=a+u+1\\ |\boldsymbol{\beta}|=b+v+1\\ |\boldsymbol{\gamma}|=c+w+1}} \zeta(\alpha_0,\alpha_1,\ldots,\alpha_a,\beta_0,\beta_1,\ldots,\beta_b+\gamma_0,\gamma_1,\ldots,\gamma_c+1).$$

We change the variables $\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma}$ to a nonnegative vector variable $\boldsymbol{e}=(e_0,e_1,\ldots,e_{m+1})$. Then

$$\mathbb{J}(3)=\sum_{\substack{m+n=p\\ a+b+c=m\\ |\boldsymbol{e}|=n}}\left(\sum_{i=0}^{e_{a+b+1}} 2^{e_{a+b+1}-i}\right)2^{e_{a+1}+e_{a+2}+\cdots+e_{a+b}}$$
$$\times\zeta(e_0+1,\ldots,e_{a+b}+1,e_{a+b+1}+2,e_{a+b+2}+1,\ldots,e_m+1,e_{m+1}+2).$$

Since

$$\sum_{i=0}^{e_{a+b+1}} 2^{e_{a+b+1}-i}=2^{e_{a+b+1}+1}-1,$$

we can change the nonnegative vector variable $\boldsymbol{e}$ to a new positive variable $\boldsymbol{\alpha}$ and $\mathbb{J}(3)$ becomes

$$\sum_{\substack{m+n=p\\ a+b+c=m\\ |\boldsymbol{\alpha}|=m+n+3}} 2^{\alpha_{a+1}+\cdots+\alpha_{a+b}-b}\left(2^{\alpha_{a+b+1}-1}-1\right)\zeta(\alpha_0,\ldots,\alpha_m,\alpha_{m+1}+1).$$

We rewrite $\mathbb{J}(3)$ as

$$\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)\sum_{a+b+c=m}W_{\boldsymbol{\alpha}}(a,b,c)$$

with $\sigma(r)=\alpha_0+\alpha_1+\cdots+\alpha_r$ and

$$W_{\boldsymbol{\alpha}}(a,b,c)=2^{\sigma(a+b+1)-\sigma(a)-(b+1)}(1-2^{1-\alpha_{a+b+1}}).$$ □

In the same manner as the previous proposition, we obtain the following.

Proposition 4.2 Notation as introduced above, we have

$$\mathbb{J}(4)=\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)\sum_{a+b=m}W_{\boldsymbol{\alpha}}(a,b,0),$$

where $W_{\boldsymbol{\alpha}}(a,b,c)$ is defined in Eq. (4.1) and

$$W_{\boldsymbol{\alpha}}(a,b,0)=2^{\sigma(a+b+1)-\sigma(a)-(b+1)}(1-2^{1-\alpha_{a+b+1}}).$$

Proposition 4.3 Notation as introduced above, then we have

$$\mathbb{J}(5)=\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)W_{\boldsymbol{\alpha}}(0,m,0),$$

where

$$W_{\boldsymbol{\alpha}}(0,m,0)=2^{\sigma(m+1)-\sigma(0)-(m+1)}(1-2^{1-\alpha_{m+1}})=2^{n+2-\sigma(0)}(1-2^{1-\alpha_{m+1}}).$$

Proposition 4.4 Notation as introduced above, then we have

$$\mathbb{J}(6)=\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)\sum_{b+c=m}W_{\boldsymbol{\alpha}}(0,b,c)$$

with $\sigma(r)=\alpha_0+\alpha_1+\cdots+\alpha_r$ and

$$W_{\boldsymbol{\alpha}}(0,b,c)=2^{\sigma(b+1)-\sigma(0)-(b+1)}(1-2^{1-\alpha_{b+1}}).$$

So up to now, our shuffle relation appeared to be the form

$$\begin{aligned}J_p&=\mathbb{J}(1)+\mathbb{J}(2)\\&\quad+\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)\\&\quad\times\left(\sum_{a+b+c=m}W_{\boldsymbol{\alpha}}(a,b,c)+\sum_{a+b=m}W_{\boldsymbol{\alpha}}(a,b,0)+W_{\boldsymbol{\alpha}}(0,m,0)+\sum_{b+c=m}W_{\boldsymbol{\alpha}}(0,b,c)\right).\end{aligned}$$

We write it in a compact form:

$$\begin{aligned}J_p&=\mathbb{J}(1)+\mathbb{J}(2)\\&\quad+\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\ldots,\alpha_m,\alpha_{m+1}+1)\\&\quad\times\sum_{a+b+c=m}\Big\{W_{\boldsymbol{\alpha}}(a,b,c)+W_{\boldsymbol{\alpha}}(a,b,c=0)+W_{\boldsymbol{\alpha}}(a=0,b=m,c=0)+W_{\boldsymbol{\alpha}}(a=0,b,c)\Big\}.\end{aligned}\tag{4.2}$$

5. Another Way to Evaluate $\mathbb{J}(1)$ and $\mathbb{J}(2)$ Parts

The modified Bell polynomials are defined by [2, 8]

$$\exp\left\{\sum_{k=1}^{\infty}\frac{x_kz^k}{k}\right\}=\sum_{m=0}^{\infty}P_m(x_1,x_2,\ldots,x_m)z^m.$$

So that

$$P_m(x_1,x_2,\ldots,x_m)=\sum_{k_1+2k_2+\cdots+mk_m=m}\frac{1}{k_1!k_2!\cdots k_m!}\left(\frac{x_1}{1}\right)^{k_1}\left(\frac{x_2}{2}\right)^{k_2}\cdots\left(\frac{x_m}{m}\right)^{k_m}.$$

In particular, for $m=0,1,2,3$,

$$P_0=1,\quad P_1(x_1)=x_1,\quad P_2(x_1,x_2)=\frac{1}{2}(x_1^2+x_2),\quad P_3(x_1,x_2,x_3)=\frac{1}{3}(x_1^3+3x_1x_2+2x_3).$$

It is well-known that

$$P_n(\zeta(2),-\zeta(4),\ldots,(-1)^{n+1}\zeta(2n))=\zeta(\{2\}^n)\quad\text{and}\quad P_n(\zeta(2),\zeta(4),\ldots,\zeta(2n))=\zeta^\star(\{2\}^n).$$

In order to give another expression of $\mathbb{J}(1)$ we need the following lemma.

Lemma 5.1 [4, Proposition 4.4] For a pair of positive integers k_1, k_2 with $k_1\le k_2$, let

$$Q(y)=\frac{\Gamma(k_1-y)}{\Gamma(k_1)}\frac{\Gamma(k_2+1)}{\Gamma(k_2+1-y)}.$$

Then for any nonnegative integer n,

$$\frac{Q^{(n)}(0)}{n!}=P_n(h_1,h_2,\ldots,h_n)=\sum_{k_1\le\ell_1\le\ell_2\le\cdots\le\ell_n\le k_2}\frac{1}{\ell_1\ell_2\cdots\ell_n}$$

with $h_n = \sum_{j=k_1}^{k_2} \frac{1}{j^n}$.

Proposition 5.2 Let p be a nonnegative integer and

$$\mathbb{J}(1) = \sum_{m+n=p} \frac{1}{m!n!} \int_{0<t_1<t_2<u_1<u_2<1} \left(\log\frac{1-t_1}{1-u_2}\right)^m \left(\log\frac{t_2}{t_1} + \log\frac{u_2}{u_1}\right)^n \frac{dt_1dt_2}{(1-t_1)t_2} \frac{du_1du_2}{(1-u_1)u_2}.$$

Then

$$\begin{aligned}\mathbb{J}(1) &= \sum_{m+n=p} \sum_{c+d=n} \{\zeta^\star(c+2, \{1\}^m, d+2) - \zeta(m+n+4)\} \\ &= \left\{\frac{p(p+3)}{2} + \frac{p+3}{2^{p+2}}\right\} \zeta(p+4).\end{aligned}$$

Proof. Let $S(m,n)$ be the general term inside the summation of $\mathbb{J}(1)$. Then

$$\begin{aligned}\sum_{m=0}^{\infty}\sum_{n=0}^{\infty} S(m,n)x^m y^n &= \int_{0<t_1<t_2<u_1<u_2<1} \left(\frac{1-t_1}{1-u_2}\right)^x \left(\frac{t_2}{t_1}\right)^y \left(\frac{u_2}{u_1}\right)^y \frac{dt_1dt_2}{(1-t_1)t_2} \frac{du_1du_2}{(1-u_1)u_2} \\ &\equiv G(x,y)\end{aligned}$$

is the generating function for the double sequence $\{S(m,n)\}$ and

$$S(m,n) = \frac{1}{m!n!} \left(\frac{\partial}{\partial x}\right)^m \left(\frac{\partial}{\partial y}\right)^n G(x,y)\Big|_{x=y=0}.$$

Beginning with

$$\frac{1}{(1-t_1)^{1-x}} = \sum_{k=1}^{\infty} \frac{\Gamma(k-x)}{\Gamma(k)\Gamma(1-x)} t_1^{k-1},$$

and then integrating with respect to t_1, t_2 and u_1, we obtain that

$$G(x,y) = \sum_{k=1}^{\infty}\sum_{\ell=1}^{\infty} \frac{1}{(k-y)k(k+\ell-y)} \frac{\Gamma(k-x)}{\Gamma(k)\Gamma(1-x)} \int_0^1 (1-u_2)^{-x} u_2^{k+\ell-1}\, du_2.$$

The value of integral is

$$\frac{\Gamma(1-x)\Gamma(k+\ell)}{\Gamma(k+\ell+1-x)}, \quad \text{or} \quad \frac{1}{k+\ell}\frac{\Gamma(1-x)\Gamma(k+\ell+1)}{\Gamma(k+\ell+1-x)}.$$

So

$$G(x,y) = \sum_{k=1}^{\infty}\sum_{\ell=1}^{\infty} \frac{1}{k(k-y)(k+\ell)(k+\ell-y)} \frac{\Gamma(k-x)\Gamma(k+\ell+1)}{\Gamma(k)\Gamma(k+\ell+1-x)}$$

after differentiations with respect to x, y for m, n times, and we use Lemma 5.1,

$$S(m,n) = \sum_{c+d=n} \left(\sum_{k=1}^{\infty}\sum_{\ell=1}^{\infty} \frac{1}{k^{c+2}(k+\ell)^{d+2}} \sum_{k\le\ell_1\le\cdots\le\ell_m\le k+\ell} \frac{1}{\ell_1\ell_2\cdots\ell_m}\right).$$

The general term of $S(m,n)$ is

$$\zeta^\star(c+2, \{1\}^m, d+2) - \zeta(m+n+4)$$

and hence

$$\mathbb{J}(1) = \sum_{m+n=p} \sum_{c+d=n} \{\zeta^\star(c+2, \{1\}^m, d+2) - \zeta(m+n+4)\}.$$

We use a result in [7, Theorem 2]:

$$\sum_{a+b+m=p} \zeta^\star(a+2,\{1\}^m,b+2) = \left(p^2+3p+1+\frac{p+3}{2^{p+2}}\right)\zeta(p+4).$$

Therefore we have

$$\mathbb{J}(1) = \left(\frac{p(p+3)}{2}+\frac{p+3}{2^{p+2}}\right)\zeta(p+4).$$

□

To transform

$$\mathbb{J}(2) = \sum_{c+d+m=p} (-1)^m \zeta(c+2,\{1\}^m,d+2)$$

into multiple zeta values related to $\mathbb{J}(1)$, we need the following reflection formula.

Proposition 5.3 [7, Proposition 4] For an r-tuple $\boldsymbol{\alpha} = (\alpha_1,\alpha_2,\dots,\alpha_r)$ of positive integers with $\alpha_1 \geq 2$, $\alpha_r \geq 2$, we have

$$\zeta(\alpha_1,\dots,\alpha_r) + (-1)^r \zeta^\star(\alpha_r,\dots,\alpha_1) = \sum_{k=1}^{r-1} (-1)^{k+1} \zeta^\star(\alpha_k,\dots,\alpha_1)\zeta(\alpha_{k+1},\dots,\alpha_r). \tag{5.1}$$

Proposition 5.4 Let p be a nonnegative integer and

$$\mathbb{J}(2) = \sum_{c+d+m=p} (-1)^m \zeta(c+2,\{1\}^m,d+2).$$

Then we have

$$\mathbb{J}(2) = J_p - \sum_{c+d+m=p} \zeta^\star(d+2,\{1\}^m,c+2).$$

Proof. By the reflection formula (see Proposition 5.3), we have

$$\zeta(c+2,\{1\}^m,d+2) = \sum_{a+b=m} (-1)^a \zeta^\star(\{1\}^a,c+2)\zeta(\{1\}^b,d+2) + (-1)^{m+1}\zeta^\star(d+2,\{1\}^m,c+2),$$

so that

$$\begin{aligned}\mathbb{J}(2) &= \sum_{m+n=p}\left((-1)^m \sum_{a+b=m}(-1)^a \sum_{c+d=n} \zeta^\star(\{1\}^a,c+2)\zeta(\{1\}^b,d+2) - \sum_{c+d=n}\zeta^\star(d+2,\{1\}^m,c+2)\right)\\ &= \sum_{a+b+c+d=p}(-1)^b\zeta^\star(\{1\}^a,c+2)\zeta(\{1\}^b,d+2) - \sum_{c+d+m=n}\zeta^\star(d+2,\{1\}^m,c+2).\end{aligned}$$

The first sum can be rewritten as

$$\sum_{m+n=p}\sum_{b+d=m}(-1)^b\zeta(\{1\}^b,d+2)\sum_{a+c=n}\zeta^\star(\{1\}^a,c+2),$$

which is precisely equal to J_p expressed as

$$\sum_{m+n=p}\frac{1}{m!n!}\int_{E_2}\left(-\log\frac{1}{1-t_1}+\log\frac{t_2}{t_1}\right)^m\frac{dt_1dt_2}{(1-t_1)t_2}\int_{E_2}\left(\log\frac{1}{1-u_2}+\log\frac{u_2}{u_1}\right)^n\frac{du_1du_2}{(1-u_1)u_2}.$$

□

According to previous propositions concerning $\mathbb{J}(1)$ and $\mathbb{J}(2)$, we have the following

Corollary 5.5 Notation as above, then we have

$$\mathbb{J}(1)+\mathbb{J}(2)=J_p-\frac{(p+1)(p+2)}{2}\zeta(p+4).$$

Now we combine Eq. (4.2) and we conclude our main theorem.

6. An Application of $\mathbb{J}(3)+\mathbb{J}(4)+\mathbb{J}(5)+\mathbb{J}(6)$

Let us begin with another integral

$$\mathbb{I}(p)=\frac{1}{p!}\int_{E_2\times E_2}\left(\log\frac{1-t_1}{1-t_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p\frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2},$$

where p is a nonnegative integer. Since

$$\frac{1}{m!}\int_{0<t_1<t_2<1}\left(\log\frac{1-t_1}{1-t_2}+\log\frac{t_2}{t_1}\right)^m\frac{dt_1}{1-t_1}\frac{dt_2}{t_2}=\zeta^\star(\{1\}^m,2)=(m+1)\zeta(m+2),$$

The integral $\mathbb{I}(p)$ can be written as

$$\sum_{m+n=p}(m+1)\zeta(m+2)\zeta(n+2)\quad\text{or}\quad\frac{p+2}{2}\sum_{m+n=p}\zeta(m+2)\zeta(n+2).$$

On the other hand, we decomposed the integral $\mathbb{I}(p)$ into six parts in a similar way to $\mathbb{J}_p$:

$$\mathbb{I}(p)=\sum_{j=1}^{6}\mathbb{I}(p;j),$$

where

$$\mathbb{I}(p;j)=\frac{1}{p!}\int_{D_j}\left(\log\frac{1-t_1}{1-t_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p\frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2},$$

The first integral $\mathbb{I}(p;1)$ is on $D_1: 0<t_1<t_2<u_1<u_2<1$, and we have

$$\mathbb{I}(p;1)=\sum_{a+b+c=p}\sum_{|\boldsymbol{\alpha}|=a+b+1}\zeta(\alpha_0,\ldots,\alpha_{a-1},\alpha_a+1,c+2).$$

The integral $\mathbb{I}(p;2)$ is

$$\mathbb{I}(p;2)=\sum_{a+b+c=p}\sum_{|\boldsymbol{\beta}|=b+c+1}\zeta(a+2,\beta_0,\ldots,\beta_{b-1},\beta_b+1).$$

The next four parts have corresponding equality with the parts $\mathbb{J}(i)$, for $3\le i\le 6$, as

$$\mathbb{I}(p;3)=\mathbb{J}(4),\quad \mathbb{J}(p;4)=\mathbb{J}(3),\quad \mathbb{J}(p;5)=\mathbb{J}(6),\quad \mathbb{J}(p;6)=\mathbb{J}(5).$$

Therefore, we get

$$\sum_{j=3}^{6}\mathbb{I}(p;j)=\sum_{j=3}^{6}\mathbb{J}(j)=\frac{(p+1)(p+2)}{2}\zeta(p+4).$$

This gives us the following theorem.

Theorem 6.1 Let p be an nonnegative integer. Then

$$\sum_{a+b+c=p}\left(\sum_{|\boldsymbol{\alpha}|=a+b+1}\zeta(\alpha_0,\ldots,\alpha_{a-1},\alpha_a+1,c+2)+\sum_{|\boldsymbol{\beta}|=b+c+1}\zeta(a+2,\beta_0,\ldots,\beta_{b-1},\beta_b+1)\right)$$
$$=\frac{p+2}{2}\sum_{m+n=p}\zeta(m+2)\zeta(n+2)-\frac{(p+1)(p+2)}{2}\zeta(p+4).$$

Here we also give another proof using a duality theorem due to Ohno [15]. Given any nonnegative integer a, we know that the dual of $\zeta(\{1\}^a,2,2)$ is $\zeta(2,a+2)$ and the dual of $\zeta(2,\{1\}^a,2)$ is $\zeta(a+2,2)$, respectively. Then for a nonnegative integer m, we have

$$\sum_{|\boldsymbol{c}|=m}\zeta(c_1+1,\dots,c_a+1,c_{a+1}+2,c_{a+2}+2)=\sum_{|\boldsymbol{d}|=m}\zeta(d_1+2,d_2+a+2),$$
$$\sum_{|\boldsymbol{c}|=m}\zeta(c_1+2,c_2+1,\dots,c_{a+1}+1,c_{a+2}+2)=\sum_{|\boldsymbol{d}|=m}\zeta(d_1+a+2,d_2+2).$$

We substitue the above formulas into $\mathbb{I}(p;1)+\mathbb{I}(p;2)$, then the desired result will be obtained.

7. Another Expression of $\mathbb{J}(3)$

The sum of multiple zeta values

$$\mathbb{J}(3)=\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=p+3}\zeta(\alpha_0,\alpha_1,\dots,\alpha_m,\alpha_{m+1}+1)\sum_{a+b+c=m}W(a,b,c)$$

came from a joint of three sums

$$\sum_{m+n=p}\sum_{a+b+c=m}\sum_{u+v+w=n}2^v\sum_{\substack{|\boldsymbol{\alpha}|=a+u+1\\|\boldsymbol{\beta}|=b+v+1\\|\boldsymbol{\gamma}|=c+w+1}}\zeta(\alpha_0,\alpha_1,\dots,\alpha_a,\beta_0,\beta_1,\dots,\beta_b+\gamma_0,\dots,\gamma_c+1)$$

and it has the integral representation

$$\frac{1}{p!}\int_{D_3}\left(\log\frac{1-t_1}{1-u_2}+\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^p\frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2},$$

which is equal to

$$\sum_{m+n=p}\frac{1}{m!n!}\int_{D_3}\left(\log\frac{1-t_1}{1-u_2}\right)^m\left(\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^n\frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2}.$$

Let $T(m,n)$ be the general term in the above sum. Here we are going to find the value of $T(m,n)$ through its generating function.

Proposition 7.1 For integers $m,n\geq 0$, let

$$T(m,n)=\frac{1}{m!n!}\int_{D_3}\left(\log\frac{1-t_1}{1-u_2}\right)^m\left(\log\frac{t_2}{t_1}+\log\frac{u_2}{u_1}\right)^n\frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2}.$$

Then

$$T(m,n)=\sum_{|\boldsymbol{\alpha}|=n+3}\{\zeta^\star(\alpha_1,\{1\}^m,\alpha_2+1)-\zeta(m+n+4)\}(2^{\alpha_2-1}-1).$$

Proof. The generating function for the double sequence $T(m,n)$ is

$$\sum_{m=0}^\infty\sum_{n=0}^\infty T(m,n)x^my^n\equiv G(x,y)$$
$$=\int_{D_3}\left(\frac{1-t_1}{1-u_2}\right)^x\left(\frac{t_2}{t_1}\right)^y\left(\frac{u_2}{u_1}\right)^y\frac{dt_1dt_2}{(1-t_1)t_2}\frac{du_1du_2}{(1-u_1)u_2}.$$

Like the case of $\mathbb{J}(1)$, $G(x,y)$ can be evaluated as

$$\sum_{k=1}^{\infty}\sum_{\ell=1}^{\infty}\frac{1}{(k-y)(k+\ell-2y)(k+\ell-y)(k+\ell)}\frac{\Gamma(k-x)}{\Gamma(k)}\frac{\Gamma(k+\ell+1)}{\Gamma(k+\ell+1-x)}$$

Hence

$$\begin{aligned}T(m,n)&=\frac{1}{m!n!}\left(\frac{\partial}{\partial x}\right)^m\left(\frac{\partial}{\partial y}\right)^n G(x,y)\Big|_{x=y=0}\\&=\sum_{a+b+c=n}\sum_{k=1}^{\infty}\sum_{\ell=1}^{\infty}\frac{2^b}{k^{a+1}(k+\ell)^{b+c+3}}\sum_{k\le k_1\le k_2\le\cdots\le k_m\le k+\ell}\frac{1}{k_1k_2\cdots k_m}.\end{aligned}$$

It is equal to

$$\sum_{|\boldsymbol{\alpha}|=n+3}\{\zeta^\star(\alpha_1,\{1\}^m,\alpha_2+1)-\zeta(m+n+4)\}(2^{\alpha_2-1}-1),$$

as asserted. □

Corollary 7.2 For any nonnegaitve integer p, we have

$$\mathbb{J}(3)=\sum_{m+n=p}\sum_{|\boldsymbol{\alpha}|=n+3}\Big\{\zeta^\star(\alpha_1,\{1\}^m,\alpha_2+1)-\zeta(p+4)\Big\}(2^{\alpha_2-1}-1).$$

Remark 7.3. We have [7, Theorem 1] a duality theorem:

$$\sum_{|\boldsymbol{\alpha}|=n+3}\zeta^\star(\alpha_1,\{1\}^m,\alpha_2+1)=(m+n+3)\zeta(m+n+4).$$

However, it is still unknown for the weighted sum

$$\sum_{|\boldsymbol{\alpha}|=n+3}\zeta^\star(\alpha_1,\{1\}^m,\alpha_2+1)2^{\alpha_2}.$$

However, our weighted sum formula did provide an approximate evaluation of weighted sum of zeta-star values of such kind.

In the next section, we will try give some weighted alternating Euler sums.

8. Weighted Alternating Euler Sum Formulas

The weighted alternating Euler sum formula

$$\sum_{m+n=p}2^n\zeta(m+1,\overline{n+1})=\frac{p+1}{2}\zeta(p+2)+\zeta(p+1,\overline{1})+\zeta(\overline{p+2})$$

was obtained in [6, Eq. (5.3)]. Now we produce a more general formula which it covers the above.

Theorem 8.1 For any nonnegative integers p,q and a real number λ, we have

$$\begin{aligned}(8.1)\qquad&\sum_{m+n=p}\binom{n}{q}\lambda^{n-q}\zeta(\overline{m+1})\zeta(\overline{n+1})\\&=\sum_{m+n=p}\binom{n}{q}(\lambda+1)^{n-q}\zeta(m+1,\overline{n+1})+\sum_{\substack{m+n=p\\a+b=q}}\binom{m}{a}\binom{n}{b}\lambda^{m-a}(\lambda+1)^{n-b}\zeta(m+1,\overline{n+1}).\end{aligned}$$

Proof. It is known that [6, Eq. (2.3)]:

$$\zeta(\overline{p+1}) = \frac{-1}{p!}\int_0^1 \left(\log\frac{1}{t}\right)^p \frac{dt}{1+t}.$$

We evaluate the following sum

$$S(p) = \sum_{m+n=p} \lambda^n \zeta(\overline{m+1})\zeta(\overline{n+1})$$

as an integral form

$$\frac{1}{p!}\int_{\substack{0<t_1<t_2<1\\0<t_2<t_1<1}} \left(\log\frac{1}{t_1} + \lambda\log\frac{1}{t_2}\right)^p \frac{dt_1dt_2}{(1+t_1)(1+t_2)}.$$

This integral can be decomposed into two parts:

$$\mathbb{K}_1 = \frac{1}{p!}\int_{0<t_1<t_2<1} \left(\log\frac{1}{t_1} + \lambda\log\frac{1}{t_2}\right)^p \frac{dt_1dt_2}{(1+t_1)(1+t_2)}, \text{ and}$$
$$\mathbb{K}_2 = \frac{1}{p!}\int_{0<t_2<t_1<1} \left(\log\frac{1}{t_1} + \lambda\log\frac{1}{t_2}\right)^p \frac{dt_1dt_2}{(1+t_1)(1+t_2)}.$$

It can be seen that

$$\mathbb{K}_1 = \sum_{a+b=p} (\lambda+1)^b \zeta(a+1,\overline{b+1}) \text{ and } \mathbb{K}_2 = \sum_{a+b=p} \lambda^a(\lambda+1)^b\zeta(a+1,\overline{b+1}).$$

Therefore we have

$$\sum_{m+n=p} \lambda^n \zeta(\overline{m+1})\zeta(\overline{n+1}) = \sum_{m+n=p} \zeta(m+1,\overline{n+1})\Big[(\lambda+1)^n(\lambda^m+1)\Big]. \tag{8.2}$$

We differential q times with λ in the above equation, we obtain the final identity. □

We list some examples. Let $\lambda = 0$ in Eq. (8.1). The following identity is obtained.

$$\sum_{m+n=p}\binom{n}{q}\zeta(m+1,\overline{n+1}) = \zeta(\overline{q+1})\zeta(\overline{p+1-q}) - \sum_{a+b=q}\binom{p-a}{b}\zeta(a+1,\overline{p+1-a}).$$

Then

$$\sum_{m+n=p} \zeta(m+1,\overline{n+1}) = \zeta(\overline{1})\zeta(\overline{p+1}) - \zeta(1,\overline{p+1}). \quad \text{(for } q=0\text{, [18, Eq. (2.15)])}$$
$$\sum_{m+n=p} n\zeta(m+1,\overline{n+1}) = \zeta(\overline{2})\zeta(\overline{p}) - p\zeta(1,\overline{p+1}) - \zeta(2,\overline{p}). \quad \text{(for } q=1\text{)}$$
$$\sum_{m+n=p}\binom{n}{2}\zeta(m+1,\overline{n+1}) = \zeta(\overline{3})\zeta(\overline{p-1}) - \binom{p}{2}\zeta(1,\overline{p+1}) - (p-1)\zeta(2,\overline{p}) - \zeta(3,\overline{p-1}). \quad \text{(for } q=2\text{)}$$

Let $\lambda = -1$ in Eq. (8.1). Then

$$\sum_{m+n=p}\binom{n}{q}(-1)^m\zeta(\overline{m+1})\zeta(\overline{n+1}) = (-1)^{p+q}\zeta(p+1-q,\overline{q+1}) + \sum_{a+b=q}\binom{p-b}{a}\zeta(p+1-b,\overline{b+1}).$$

The $q=0$ and $q=1$ cases are

$$\sum_{m+n=p}(-1)^m\zeta(\overline{m+1})\zeta(\overline{n+1}) = \left[1+(-1)^p\right]\zeta(p+1,\overline{1}),$$

$$\sum_{m+n=p}(-1)^m n\zeta(\overline{m+1})\zeta(\overline{n+1})=\left[1+(-1)^{p+1}\right]\zeta(p,\overline{2})+p\zeta(p+1,\overline{1}).$$

Let $\lambda=1$ in Eq. (8.1), we have

$$\sum_{m+n=p}\binom{n}{q}\zeta(\overline{m+1})\zeta(\overline{n+1})=\sum_{m+n=p}\binom{n}{q}2^{n-q}\zeta(m+1,\overline{n+1})+\sum_{\substack{m+n=p\\a+b=q}}\binom{m}{a}\binom{n}{b}2^{n-b}\zeta(m+1,\overline{n+1}).$$

We set $q=0$ in the above identity, we will get [6, Eq. (2.14)]

$$\sum_{m+n=p}\zeta(\overline{m+1})\zeta(\overline{n+1})=\sum_{m+n=p}2^{n+1}\zeta(m+1,\overline{n+1}).$$

The well-known double-stuffle relation [18, Eq. (2.2)]

$$\zeta(\overline{r})\zeta(\overline{s})=\zeta(\overline{r},\overline{s})+\zeta(\overline{s},\overline{r})+\zeta(r+s), \tag{8.3}$$

where $r\geq 1$, $s\geq 1$. We use the above relation and the sum formula [18, Eq. (2.14)]

$$\sum_{m+n=p}\zeta(\overline{m+1},\overline{n+1})=\zeta(\overline{1})\zeta(p+1)-\zeta(\overline{1},p+1),$$

We derive the following

$$\sum_{m+n=p}2^n\zeta(m+1,\overline{n+1})=\frac{p+1}{2}\zeta(p+2)+\zeta(p+1,\overline{1})+\zeta(\overline{p+2}).$$

This identity was obtained in [6, Eq. (5.3)]. We indicate this formula in the first paragraph of this section.

Let $\lambda=-2$ in Eq. (8.1), we have

$$\begin{aligned}\sum_{m+n=p}\binom{n}{q}(-1)^m 2^{n-q}\zeta(\overline{m+1})\zeta(\overline{n+1})=&\sum_{m+n=p}\binom{n}{q}(-1)^m\zeta(m+1,\overline{n+1})\\&+\sum_{\substack{m+n=p\\a+b=q}}\binom{m}{a}\binom{n}{b}2^{m-a}\zeta(m+1,\overline{n+1}).\end{aligned}$$

We set $q=0$ in the above identity, we have

$$\sum_{m+n=p}(-1)^m 2^n\zeta(\overline{m+1})\zeta(\overline{n+1})=\sum_{m+n=p}\left[2^m+(-1)^m\right]\zeta(m+1,\overline{n+1}).$$

Using the double-stuffle relation Eq (8.3) we have

$$\begin{aligned}&\frac{(-2)^{p+1}-1}{3}\zeta(p+2)\\&=\sum_{m+n=p}[(-2)^n+(-2)^m]\zeta(\overline{m+1},\overline{n+1})-\sum_{m+n=p}(-1)^n[1+(-2)^m]\zeta(m+1,\overline{n+1}).\end{aligned}$$

References

[1] T. Arakawa, M. Kanekno, *Multiple zeta values, poly-Bernoulli numbers, and related zeta functions*, Nagoya Math. J. **153**, 189–209 (1999).

[2] K.-W. Chen, *Generalized harmonic numbers and Euler sums*, Int. J. Number Theory **13**, no. 2, 513–528 (2017). https://doi.org/10.1142/S1793042116500883.

[3] K.-W. Chen, *Generalized Arakawa-Kaneko zeta functions*, Integral Transforms Spec. Funct. **30**, no. 4, 282–300 (2019). https://doi.org/10.1080/10652469.2018.1562450.

[4] K.-W. Chen, C.-L. Chung, M. Eie, *Sum formulas and duality theorems of multiple zeta values*, J. Number Theory **158**, 33–53 (2016). https://doi.org/10.1016/j.jnt.2015.06.014.

[5] K.-W. Chen, M. Eie, *Some special Euler sums and* $\zeta^\star(r+2,\{2\}^n)$, arXiv (2018), arXiv:1810.11795v1 [math.NT]

[6] K.-W. Chen, M. Eie, *On the convolutions of sums of multiple zeta(-star) values of height one*, arXiv (2021), arXiv: 2110.00231v1 [math.NT]

[7] K.-W. Chen, M. Eie, *On three general forms of multiple zeta(-star) values*, arXiv (2022), arXiv: 2202.03839v1 [math.NT]

[8] M.-A. Coppo, B. Candelpergher, *The Arakawa-Kaneko zeta function*, Ramanujan J. **22**, no. 2, 153–162 (2010). https://doi.org/10.1007/s11139-009-9205-x.

[9] M. Eie, *Topics in Number Theory*, Monographs in Number Theory **2**, World Scientific, Singapore, 2009. https://doi.org/10.1142/7036.

[10] M. Eie, *The Theory of Multiple Zeta Values with Applications in Combinatorics*, Monographs in Number Theory **7**, World Scientific, Singapore, 2013. https://doi.org/10.1142/8769.

[11] M. Eie, C.-S. Wei, *A short proof for the sum formula and its generalization*, Arch. Math. **91**, 330–338 (2008). https://doi.org/10.1007/s00013-008-2547-6.

[12] A. Granville, *A decomposition of Riemann's zeta-function*, in: *Analytic Number Theory* (Kyoto, 1996), 95–102, London Math. Soc. Lecture Notes Ser. **247**, Cambridge Univ. Press, Cambridge, 1997. https://doi.org/10.1017/CBO9780511666179.009.

[13] L. Kargın, *Poly-p-Bernoulli polynomials and generalized Arakawa-Kaneko zeta function*, Lith. Math. J. **60**, 29–50 (2020). https://doi.org/10.1007/s10986-019-09448-7.

[14] T. Q. T. Le, J. Murakami, *Kontsevich's integrals for the Homfly polynomial and relations between values of multiple zeta functions*, Topology Appl. **62**, no. 2, 193–206 (1995). https://doi.org/10.1016/0166-8641(94)00054-7.

[15] Y. Ohno, *A generalization of the duality and sum formulas on the multiple zeta values*, J. Number Theory **74**, no. 1, 39–43 (1999). https://doi.org/10.1006/jnth.1998.2314.

[16] Y. Ohno, *Sum relations for multiple zeta values*, In: Aoki, T., Kanemitsu S., Nakahara M., Ohno Y. (eds) Zeta Functions, Topology and Quantum Physics, Developments in Mathematics, **14**, Springer, Boston, MA (2005). https://doi.org/10.1007/0-387-24981-8$_8$.

[17] Y. L. Ong, M. Eie, W.-C. Liaw, *On generalizations of weighted sum formulas of multiple zeta values*, Int. J. Number Theory **9**, no. 5, 1185–1198 (2013). https://doi.org/10.1142/S179304211350019X.

[18] L.-P. Teo, *Alternating double Euler sums, hypergeometric identities and a theorem of Zagier*, J. Math. Anal. Appl. **462**, 777–800 (2018). https://doi.org/10.1016/j.jmaaa.2018.02.037.

[19] C. Xu, *Integrals of logarithmic functions and alternating multiple zeta values*, Math. Slovaca **69**, no. 2, 339–356 (2019). https://doi.org/10.1515/ms-2017-0227.

Department of Mathematics, University of Taipei, 10048 Taipei, Taiwan

Email address: kwchen@utaipei.edu.tw

Department of Mathematics, National Chung Cheng University, 168 University Road, Min-Hsiung, Chia-Yi 62145, Taiwan

Email address: minkingeie@gmail.com