

SUMS OF SQUARES IN FUNCTION FIELDS OVER HENSELIAN DISCRETELY VALUED FIELDS

GONZALO MANZANO-FLORES

ABSTRACT. Let $n \in \mathbb{N}$ and let K be a field with a henselian discrete valuation of rank n with hereditarily euclidean residue field. Let F/K be a function field in one variable in one variable. It is known that every sum of squares is a sum of 3 squares. We determine the order of the group of nonzero sums of 3 squares modulo sums of 2 squares in F in terms of equivalence classes of certain discrete valuations of F of rank at most n . In the case of function fields of hyperelliptic curves of genus g , K.J. Becher and J. Van Geel showed that the order of this quotient group is bounded by $2^{n(g+1)}$. We show that this bound is optimal. Moreover, in the case where $n = 1$, we show that if F/K is a hyperelliptic function field such that the order of this quotient group is 2^{g+1} , then F is nonreal.

KEYWORDS: Sums of squares, valuation, function field in one variable, quadratic form, local-global principle, hereditarily pythagorean field.

CLASSIFICATION (MSC 2020): 11E81, 12D15, 12J10

1. INTRODUCTION

Let K be a field. For $d \in \mathbb{N}$, let $S_d(K)$ denote the set of nonzero sums of d squares in K . Let $S(K) = \bigcup_{n \in \mathbb{N}} S_n(K)$, the multiplicative group of nonzero sums of squares in K . The Pythagoras number $p(K)$ of a field K is the smallest integer d such that $S(K) = S_d(K)$ if such an integer d exists, otherwise it is infinite. We call a finitely generated field extension of transcendence degree one a *function field in one variable*, for short. In [20] it was shown that $p(F) = 2$ for every function field in one variable F over \mathbb{R} . However, the same does not hold for function field in one variables over $\mathbb{R}((t_1)) \dots ((t_n))$ when $n \geq 1$. For example, S. Tikhonov in [17, Example 3.8] showed that tX is a sum of three squares but not a sum of two squares in the function field F of the curve $Y^2 = (tX - 1)(X^2 + 1)$ over $\mathbb{R}((t))$, which shows that $p(F) > 2$. However, in that example it was not known whether tX was the only nontrivial element of the quotient group of nonzero sums of squares in F modulo the subgroup of nonzero sums of two squares in F .

Let $n \in \mathbb{N}$. Consider a function field in one variable F over the base field $\mathbb{R}((t_1)) \dots ((t_n))$. It was shown in [3, Theorem 6.12] that $p(F) \leq 3$ and that the quotient $S(F)/S_2(F)$, which measures the failure of every sum of squares to be

Date: 22.02.2023.

a sum of two squares, is a finite group. In this article, we determine the order of $\mathcal{S}(F)/\mathcal{S}_2(F)$ in terms of the cardinality of a certain finite set of valuations on F . In fact, we consider situations with slightly more general base fields (than $\mathbb{R}((t_1)) \dots ((t_n))$) namely fields carrying a nontrivial henselian valuation with value group \mathbb{Z}^n endowed with the lexicographic order, and such that the Pythagoras number of the rational function field over its residue field is bounded by 2^r , for some $r \in \mathbb{N}^+$. Concretely, for a function field in one variable $F/\mathbb{R}((t_1)) \dots ((t_n))$, we show that $|\mathcal{S}(F)/\mathcal{S}_2(F)| = 2^m$, where m is the number of equivalence classes of discrete valuations of rank at most n and with nonreal residue field not containing $\sqrt{-1}$. In this case, this generalizes [3, Theorem 6.12]. It is natural to wonder whether we can bound the order of $\mathcal{S}(F)/\mathcal{S}_2(F)$ in terms of the genus of F/K . In [2], reduction theory of arithmetic curves over $\mathbb{R}[[t]]$ is applied to obtain the bound $|\mathcal{S}(F)/\mathcal{S}_2(F)| \leq 2^{g+1}$ for any function field in one variable F of genus g over $\mathbb{R}((t))$. In the case of a function field F of a hyperelliptic curve of genus g over $\mathbb{R}((t_1)) \dots ((t_n))$, it is a consequence of [5, Theorem 3.10] that

$$|\mathcal{S}(F)/\mathcal{S}_2(F)| \leq 2^{n(g+1)}.$$

It is known that in the case where $n = 1$ and $g = 1$, the bound is optimal. In fact, for the nonreal function field F of the curve $Y^2 = -(X^2 + 1)(X^2 + t^2)$ over $\mathbb{R}((t))$, it was shown in [5, Example 5.12] that $|\mathcal{S}(F)/\mathcal{S}_2(F)| = 2^2$. In this article, for any $g \in \mathbb{N}$, we will construct a hyperelliptic curve of genus g over K such that $|\mathcal{S}(F)/\mathcal{S}_2(F)| = 2^{n(g+1)}$. This shows the optimality of the previously mentioned bounds for function field in one variables over $\mathbb{R}((t))$ or for hyperelliptic function fields over $\mathbb{R}((t_1)) \dots ((t_n))$. We ask in Section 4 whether our bounds in terms of valuations may equally serve to find a bound in terms of g and n for function field in one variables F/K of genus g .

2. PRELIMINARIES

For a ring R , let R^\times denote its group of invertible elements. Let K be a field. For a valuation v on K , we denote by \mathcal{O}_v the valuation ring associated to v , by Γ_v the value group $v(K^\times)$, which is an ordered abelian group, by \mathfrak{m}_v the maximal ideal of \mathcal{O}_v and by κ_v the residue field $\mathcal{O}_v/\mathfrak{m}_v$. The residue in κ_v of an element $a \in \mathcal{O}_v$ will be denoted by \bar{a} . For a valuation ring \mathcal{O} of K , the corresponding valuation on K with valuation ring \mathcal{O} and value group $K^\times/\mathcal{O}^\times$ is denoted by $v_{\mathcal{O}}$. We call two valuations v and w on K *equivalent* if $\mathcal{O}_v = \mathcal{O}_w$, equivalently, if and only if there exists an order-isomorphism $\gamma : \Gamma_w \rightarrow \Gamma_v$ such that $v = \gamma \circ w$. See [8, Proposition 2.1.3].

The *rank* of an ordered abelian group Γ is defined as the number of its proper convex subgroups. For a valuation v on K , we denote by $\text{rk}(v) \in \mathbb{N}$ the rank of Γ_v , and we say that $\text{rk}(v)$ is the rank of v . Let $n \in \mathbb{N}$. We call a valuation on K having value group $(\mathbb{Z}^n, \leq_{\text{leq}})$, where \leq_{leq} is the lexicographic order, a \mathbb{Z}^n -*valuation* for short. Note that $(\mathbb{Z}^0, \leq_{\text{leq}})$ is the trivial additive group and that for a \mathbb{Z}^n -valuation v on K we have that $\text{rk}(v) = n$.

For $i \in \mathbb{N}$, let $V_i(K)$ be the set of \mathbb{Z}^i -valuations on K , where $V_0(K)$ is the trivial valuation on K . We set

$$V(K) = \bigcup_{i \in \mathbb{N}} V_i(K).$$

Let $\Omega_n(K) = \{\mathcal{O}_v \mid v \in V_n(K)\}$, where $\Omega_0(K) = K$. We set

$$\Omega(K) = \bigcup_{i \in \mathbb{N}} \Omega_i(K).$$

We observe that every $\mathcal{O} \in \Omega_n(K)$ is a valuation ring of Krull dimension n by [8, Lemma 2.3.1]. For valuation rings $\mathcal{O}, \mathcal{O}'$ of a field K and valuations w, w' on K corresponding to \mathcal{O} and \mathcal{O}' , respectively, we say that \mathcal{O}' is a *coarsening* of \mathcal{O} or that \mathcal{O} is a *refinement* of \mathcal{O}' , respectively, w' is a coarsening of w or w is a refinement of w' , if $\mathcal{O} \subseteq \mathcal{O}'$. Note that a field K is trivially a coarsening of every valuation ring of itself. We denote by $\mathcal{O}\mathcal{O}'$ the smallest subring of K containing both \mathcal{O} and \mathcal{O}' .

Let $n, d \in \mathbb{N}$ with $d \leq n$. We denote by $\pi_d : \mathbb{Z}^n \rightarrow \mathbb{Z}^d$ the projection on the first d components. Note that π_d is a homomorphism of ordered abelian groups with respect to the lexicographic orders on \mathbb{Z}^n and \mathbb{Z}^d . Dually, we denote by $\pi^d : \mathbb{Z}^n \rightarrow \mathbb{Z}^d$ the projection on the last d components of \mathbb{Z}^n . Note that π^d is a group homomorphism, but it is not order-preserving when $d < n$.

2.1. Proposition. *Let $n \in \mathbb{N}$. Let v be a \mathbb{Z}^n -valuation on K . Let \mathcal{O}' be a coarsening of \mathcal{O}_v . Then $\mathcal{O}' = \mathcal{O}_{\pi_r \circ v}$, for a unique $r \leq n$.*

Proof. First, note that $K^\times / \mathcal{O}_v^\times$ is an abelian group, which has an order \leq , given by $x\mathcal{O}_v^\times \leq y\mathcal{O}_v^\times$ if and only if $yx^{-1} \in \mathcal{O}_v$. Thus Γ_v is order-isomorphic to $K^\times / \mathcal{O}_v^\times$. Let $\mathcal{O} = \mathcal{O}_v$. Let \mathfrak{p} be the maximal ideal of \mathcal{O}' . Then \mathfrak{p} is a prime ideal of \mathcal{O} and localizing \mathcal{O} at \mathfrak{p} we obtain $\mathcal{O}' = \mathcal{O}_{\mathfrak{p}}$. Thus, we have a surjective order-preserving group homomorphism of value groups $\varphi : K^\times / \mathcal{O}^\times \rightarrow K^\times / \mathcal{O}_{\mathfrak{p}}^\times$ sending $x\mathcal{O}^\times$ to $x\mathcal{O}_{\mathfrak{p}}^\times$. In fact, if $a\mathcal{O}^\times \leq b\mathcal{O}^\times$, then $ab^{-1} \in \mathcal{O} \subseteq \mathcal{O}_{\mathfrak{p}}$ and hence $a\mathcal{O}_{\mathfrak{p}} \leq b\mathcal{O}_{\mathfrak{p}}$, for $a, b \in K^\times$. Since $K^\times / \mathcal{O}^\times \simeq \mathbb{Z}^n$, then the quotient group $(K^\times / \mathcal{O}^\times) / \ker \varphi$ is order-isomorphic to \mathbb{Z}^r , for some $r \in \mathbb{N}$ such that $r \leq n$. Hence $K^\times / \mathcal{O}_{\mathfrak{p}}^\times \simeq \mathbb{Z}^r$ by the first isomorphism theorem. Let $v' = \pi_r \circ v$. Since π_r is order-preserving, we have that $\mathcal{O}_{v'} \in \Omega_r(K)$, which is a coarsening of \mathcal{O}_v . By [8, Lemma 2.3.1] and since \mathbb{Z}^n has only n proper convex subgroups, we have that $\mathcal{O}' = \mathcal{O}_{v'}$. \square

Let $v \in V(K)$ and set $n = \text{rk}(v)$. For $1 \leq i \leq n$, we denote by e_i^n the n -tuple (e_1, \dots, e_n) such that

$$e_j = \begin{cases} 1 & \text{if } j = (n+1) - i, \\ 0 & \text{if } j \neq (n+1) - i. \end{cases}$$

Note that e_1^n is the minimal positive element of \mathbb{Z}^n . An n -tuple $(t_1, \dots, t_n) \in K^n$ is called a *parametrical system* of v if $v(t_i) = e_i^n$ for all $1 \leq i \leq n$. For a \mathbb{Z} -valuation v we call $t \in K$ a *uniformizer* of v if $v(t) = 1$.

Definition. Let $n \in \mathbb{N}$. Let v be a \mathbb{Z}^n -valuation on K and let \mathcal{O}' be a coarsening of \mathcal{O}_v . By Proposition 2.1 there exists a unique $r \leq n$, such that $\mathcal{O}' = \mathcal{O}_{\pi_r \circ v}$. Let $v' = \pi_r \circ v$. Let $x, y \in \mathcal{O}_{v'}$ be such that $x - y \in \mathfrak{m}_{v'}$ and $x, y \notin \mathfrak{m}_{v'}$. Since $y \in \mathcal{O}_{v'}^\times$ and $\mathfrak{m}_{v'} \subseteq \mathfrak{m}_v$, we have that $(x - y)y^{-1} = x/y - 1 \in \mathfrak{m}_v$, and hence $v(x) = v(y)$. We denote by $\bar{v} : \kappa_{v'} \rightarrow \mathbb{Z}^{n-r} \cup \{\infty\}$, $\bar{x} \mapsto (\pi^{n-r} \circ v)(x)$, where x is any lifting of \bar{x} , the *residual valuation of v modulo v'* , which is well-defined by the above.

Let $r, d \in \mathbb{N}$. On the other hand, let v' be a \mathbb{Z}^r -valuation on K and let \bar{v} be a \mathbb{Z}^d -valuation on $\kappa_{v'}$. Let $(t_1, \dots, t_r) \in K^r$ be a parametrical system of v' . For $a \in K^\times$, there exists $a_1, \dots, a_r \in \mathbb{Z}$ such that $a = t_1^{a_1} \dots t_r^{a_r} u$, for some $u \in \mathcal{O}_{v'}^\times$. We can define a valuation $v : K \rightarrow \mathbb{Z}^r \times \mathbb{Z}^d \cup \{\infty\}$, by $v(a) = (v'(a), \bar{v}(\bar{u}))$. The valuation v is called *the composition of v' and \bar{v} with respect to (t_1, \dots, t_r)* . Note that \mathcal{O}_v is a refinement of $\mathcal{O}_{v'}$.

2.2. Example. Let $n \in \mathbb{N}^+$. We consider $K_n = k((t_1)) \dots ((t_n))$ the field of iterated Laurent series over a field k . By induction on n , we show that there exists a \mathbb{Z}^n -valuation v on K_n such that $(t_1, \dots, t_n) \in K_n^n$ is a parametrical system of v . For $n = 1$, the valuation v_{t_1} on K_1 , given by $v_{t_1}(\sum_{i=m}^\infty a_i t_1^i) = m$, when $a_m \in k^\times$, is a \mathbb{Z} -valuation on K_1 such that t_1 is a parametrical system v_{t_1} . Let $n > 1$. Let v_{t_n} be the t_n -adic valuation on K_n , that is, the \mathbb{Z} -valuation corresponding to the valuation ring $K_{n-1}[[t_n]]$. By the induction hypothesis, since $\kappa_{v_{t_n}} = K_{n-1}$, we may consider a \mathbb{Z}^{n-1} -valuation w on K_{n-1} such that $(t_1, \dots, t_{n-1}) \in \kappa_{v_{t_n}}^{n-1}$ is a parametrical system of w . Let v be the composition of v_{t_n} and w with respect to t_n . Hence v is a \mathbb{Z}^n -valuation on K_n , and since $v(t_n) = (v_{t_n}(t_n), w(\bar{1})) = (1, 0, \dots, 0) = e_n^n$, we have that $(t_1, \dots, t_n) \in K_n^n$ is a parametrical system of v . Note that different choices of uniformizers of v_{t_n} can lead to distinct \mathbb{Z}^n -valuations on K_n (e.g. $t_n t_1$ instead of t_n).

2.3. Remark. Let $r \in \mathbb{N}$. Fixing a \mathbb{Z}^r -valuation v' on K , a parametrical system $(t_1, \dots, t_r) \in K^r$ for v' , and $d \in \mathbb{N}$, the above defined assignments define mutually inverse bijections between equivalence classes of \mathbb{Z}^{r+d} -valuations v , where \mathcal{O}_v are refinements of $\mathcal{O}_{v'}$ and equivalence classes of \mathbb{Z}^d -valuations \bar{v} on $\kappa_{v'}$, such that $\kappa_v = \kappa_{\bar{v}}$. See [8, Section 2.3].

We call $W \subseteq \Omega(K) \setminus \Omega_0(K)$ *saturated* if for all $\mathcal{O} \in W$ we have $\mathcal{O}' \in W$ for every nontrivial coarsening \mathcal{O}' of \mathcal{O} . Let $S, S' \subseteq V(K)$. We say that S and S' are *equivalent* if $\{\mathcal{O}_v \mid v \in S\} = \{\mathcal{O}_v \mid v \in S'\}$. We say that $S \subseteq V(K) \setminus V_0(K)$ is *coherent* if

- S is a set of pairwise non-equivalent valuations,
- and, if $v \in S$ is a \mathbb{Z}^n -valuation, then $\pi_i \circ v \in S$, for all $1 \leq i \leq n$.

2.4. Proposition. *Let $S \subseteq V(K) \setminus V_0(K)$ be such that $W = \{\mathcal{O}_v \mid v \in S\}$ is finite and saturated. Then there exists a coherent set $S' \subseteq V(K)$ equivalent to S .*

Proof. Let $n = \max\{\dim \mathcal{O}_v \mid v \in S\}$. We prove the statement by induction on n . If $n = 1$, there is nothing to show, since every subset of $V_1(K)$ is a set of pairwise of

non-equivalent \mathbb{Z} -valuations, which is trivially coherent. Assume now that $n > 1$. By the induction hypothesis, we have that for any $S \subseteq \bigcup_{1 \leq i \leq n-1} V_i(K)$ such that $W = \{\mathcal{O}_v \mid v \in S\}$ is finite and saturated, there exists a coherent set $S' \subseteq V(K)$ with $W = \{\mathcal{O}_v \mid v \in S'\}$. Let $S \subseteq \bigcup_{1 \leq i \leq n} V_i(K)$ be such that $W = \{\mathcal{O}_v \mid v \in S\}$ is finite and saturated. Without loss of generality, we may assume that S consist of a set of non-equivalent valuations. Let $S_{n-1} = S \setminus V_n(K)$. Then, by the induction hypothesis, S_{n-1} is equivalent to a coherent subset S'_{n-1} of $\bigcup_{1 \leq i \leq n-1} V_i(K)$. Let $v \in S'_{n-1}$. Let $V_v = \{w \in S \mid \mathcal{O}_w \subsetneq \mathcal{O}_v\}$. Let $(t_1, \dots, t_{n-1}) \in K^{n-1}$ be a parametrical system of v . For $w \in V_v$, let \bar{w} be the residual valuation of w modulo v . Let w' be the composition of v and \bar{w} , with respect to (t_1, \dots, t_{n-1}) . Thus, for every $w \in V_v$, we can define a \mathbb{Z}^n -valuation w' on K , which is equivalent to w , by Remark 2.3. Let $S(v) = \{w' \mid w \in V_v\}$. Then $S(v) \cup S'_{n-1}$ is a coherent set equivalent to $V_v \cup S'_{n-1}$. We obtain that

$$S' = \bigcup_{v \in S'_{n-1} \cap V_{n-1}(K)} S(v) \cup S'_{n-1},$$

is a coherent finite set equivalent to S . \square

Let $v, w \in V(K)$. We call $(\gamma_v, \gamma_w) \in \Gamma_v \times \Gamma_w$ *compatible with respect to (v, w)* if $\pi_r(\gamma_v) = \pi_r(\gamma_w)$, where $r \in \mathbb{N}$ is such that $\mathcal{O}_v \mathcal{O}_w \in \Omega_r(K)$.

Let $n \in \mathbb{N}$. We recall that π^1 is the projection on the last component of \mathbb{Z}^n . Let $S \subseteq V(K) \setminus V_0(K)$ be a finite coherent set. We define the group homomorphism

$$(2.5) \quad \Phi_S : K^\times \rightarrow \prod_{v \in S} \mathbb{Z}, a \mapsto (\pi^1(v(a)))_{v \in S}.$$

2.6. Proposition. *Let K be a field. Let $S \subseteq V(K) \setminus V_0(K)$ be a finite coherent set. Then Φ_S is surjective.*

Proof. Let $(e_v)_{v \in S}$ be the canonical basis of $\prod_{v \in S} \mathbb{Z}$ as a \mathbb{Z} -module. For every $v \in S$, we show that there exists $x_v \in K^\times$ such that $\Phi_S(x_v) = e_v$. Consider $v \in S$ and let $\gamma_v = e_1^{\text{rk}(v)}$. For $w \in S$ a refinement of v , let $\gamma_w = e_{\text{rk}(w) - \text{rk}(v) + 1}^{\text{rk}(w)}$ in Γ_w , otherwise $\gamma_w = 0$. We claim that for every $w, w' \in S$, the pair $(\gamma_w, \gamma_{w'})$ in $\Gamma_w \times \Gamma_{w'}$, is compatible with respect to (w, w') . Let $w, w' \in S$. If w, w' are both not refinements of v , then $(\gamma_w, \gamma_{w'})$ are trivially compatible. Assume w is a refinement of v and w' is not a refinement of v . It follows from Proposition 2.1 that $\mathcal{O}_w \subseteq \mathcal{O}_v \subsetneq \mathcal{O}_w \mathcal{O}_{w'} \subseteq K$, because every valuation ring has a unique coarsening of a fixed rank. Let $d \in \mathbb{N}$ be such that $\mathcal{O}_w \mathcal{O}_{w'} \in \Omega_d(K)$. Then $d < \text{rk}(v) \leq \text{rk}(w)$ and hence $\pi_d(\gamma_w) = 0 = \pi_d(\gamma_{w'})$ in \mathbb{Z}^d . Finally, we assume that w and w' are refinements of v . Let $d \in \mathbb{N}$ be such that $\mathcal{O}_w \mathcal{O}_{w'} \in \Omega_d(K)$. Then $\mathcal{O}_w \mathcal{O}_{w'} \subseteq \mathcal{O}_v$ and hence we have that $\text{rk}(v) \leq d < \text{rk}(w), \text{rk}(v) \leq d < \text{rk}(w')$. Thus, we have $\pi_d(\gamma_w) = \pi_d(\gamma_{w'})$, that is, $(\gamma_w, \gamma_{w'})$ are compatible. Therefore, for every $w, w' \in S$, the pair $(\gamma_w, \gamma_{w'}) \in \Gamma_w \times \Gamma_{w'}$ is compatible. By [16, Theorem 5] there exists $x_v \in K^\times$ such that $w(x_v) = \gamma_w$ for all $w \in W$. Hence $\pi^1(v(x_v)) = \pi^1(\gamma_v) = 1$,

and $\pi^1(w(x_v)) = 0$ for all $w \in S \setminus \{v\}$. Therefore $\Phi_S(x_v) = e_v$. Since $v \in S$ was arbitrarily chosen, we conclude that Φ_S is surjective. \square

2.7. Proposition. *Let $n \in \mathbb{N}$. Assume that K carries a henselian \mathbb{Z}^n -valuation. Then $|\Omega_i(K)| = 1$ for all $1 \leq i \leq n$.*

Proof. We prove this by induction on n . For $n = 1$, follows from [3, Proposition 2.2]. We assume that $n > 1$. Let $v_{n-1} = \pi_{n-1} \circ v$. Then v_{n-1} is a henselian \mathbb{Z}^{n-1} -valuation on K , which is a coarsening of v , by [8, Corollary 4.1.4] and Proposition 2.1. By the induction hypothesis $|\Omega_i(K)| = 1$, for all $1 \leq i \leq n-1$. Let \bar{v} be the residual valuation of v modulo v_{n-1} . Then \bar{v} is a henselian \mathbb{Z} -valuation on $\kappa_{v_{n-1}}$, by [8, Corollary 4.1.4], and it is the unique \mathbb{Z} -valuation on $\kappa_{v_{n-1}}$, by [3, Proposition 2.2]. Therefore, by Remark 2.3, we have that $\Omega_n(K) = \{\mathcal{O}_v\}$ and hence $|\Omega_n(K)| = 1$. \square

By a function field in one variable F/K we mean a finitely generated field extension of transcendence degree one.

2.8. Lemma. *Let F/K be a function field in one variable and $r \in \mathbb{N}$. Let w be a valuation on F such that $\kappa_w/\kappa_{w|_K}$ is transcendental. Then $\text{rk}(w) = \text{rk}(w|_K)$ and $\kappa_w/\kappa_{w|_K}$ is a function field in one variable.*

Proof. Let $v = w|_K$. It follows from [8, Proposition 3.4.1, Theorem 3.4.3] that $\text{rk}(w) = \text{rk}(v)$. We claim that κ_w/κ_v is a function field in one variable. Let $\alpha \in \kappa_w$ be a transcendental element over κ_v and let $\theta \in \mathcal{O}^\times$ be such that $\bar{\theta} = \alpha$. It follows by [8, Theorem 3.2.4] that $K(\theta)/K$ is transcendental. Let $w' := w|_{K(\theta)}$. It follows from [8, Corollary 2.2.2] that w' is the Gauss extension of v with respect to θ . In particular $\kappa_{w'} = \kappa_v(\alpha)$. Since F/K is a function field in one variable, we have that $F/K(\theta)$ is finite, then $\kappa_w/\kappa_v(\alpha)$ is a finite field extension and hence κ_w/κ_v is a function field in one variable. \square

We fix a valued field (K, v) and F/K a function field in one variable. An extension w of v to F is called *residually transcendental* if κ_w/κ_v is transcendental. We say that F/K is *ruled* if F is a rational function field over some finite extension of K .

2.9. Proposition. *Let v be a valuation on K . Let F/K be a ruled extension. Let w be a residually transcendental extension of v . Then κ_w/κ_v is ruled.*

Proof. See [15, Theorem 3.3]. \square

We define a set of equivalence classes of valuation extensions of coarsenings of v to F in the following way. For $i \in \mathbb{N}$, let $\Omega_i(F/v)$ denote the set of valuation rings $\mathcal{O} \in \Omega_i(F)$ such that $v_{\mathcal{O}}$ is a residually transcendental extension of a coarsening of v . We set

$$\Omega(F/v) = \bigcup_{i \in \mathbb{N}} \Omega_i(F/v).$$

2.10. Proposition. *Let $n \in \mathbb{N}$ and let $v \in V_n(K)$. Let F/K be a function field in one variable. Let $\mathcal{O} \in \Omega(F/v)$. Then $v_{\mathcal{O}}$ is a \mathbb{Z}^r -valuation, for some $r \leq n$.*

Proof. This follows directly from Proposition 2.1 and Lemma 2.8. \square

For $i \in \mathbb{N}^+$, we set

$$\Omega_i^*(F/v) = \{\mathcal{O} \in \Omega_i(F/v) \mid \kappa_{\mathcal{O}}/\kappa_{\mathcal{O} \cap K} \text{ is nonruled}\}.$$

We set

$$\Omega(F/v) = \bigcup_{i \in \mathbb{N}} \Omega_i^*(F/v).$$

2.11. Lemma. *Let F/K be a function field in one variable. Then $\Omega^*(F/v)$ is saturated.*

Proof. Let w be a valuation on F such that $\mathcal{O}_w \in \Omega^*(F/v)$. Let w' be a coarsening of w . We claim that $\mathcal{O}_{w'} \in \Omega^*(F/v)$. It follows by [8, Theorem 3.2.4] that $\kappa_{w'}/\kappa_{w'|_K}$ is transcendental, and hence is a function field in one variable, by Lemma 2.8. Let \bar{w} be the residual valuation of w modulo w' . Let $\nu = w|_K$. Note that $w'|_K$ is a coarsening of ν . Let $\bar{\nu}$ be the residual valuation of ν modulo $w'|_K$. If $\kappa_{w'}/\kappa_{w'|_K}$ were ruled, then $\kappa_{\bar{w}}/\kappa_{\bar{\nu}}$ would be ruled, by Proposition 2.9, and hence κ_w/κ_{ν} would be ruled because $\kappa_w = \kappa_{\bar{w}}, \kappa_{\nu} = \kappa_{\bar{\nu}}$, which is a contradiction. Therefore $\mathcal{O}_{w'} \in \Omega^*(F/v)$. \square

Let F/K be a function field in one variable. It is natural to wonder whether the set $\Omega^*(F/v)$ is finite. We shall give a positive answer in Theorem 2.12 under the assumption that $v \in V(K)$. Assuming that v is a \mathbb{Z} -valuation on K , it was shown in [2, Theorem 5.3] by K. Becher and D. Grimm $|\Omega^*(F/v)| \leq g + 1$, where g is the genus of F/K . We will show in Example 4.5 that the above bound is optimal.

2.12. Theorem. *Let n be a positive integer. Assume that K carries a \mathbb{Z}^n -valuation v such that κ_v is perfect. Let F/K be a function field in one variable. Then $\bigcup_{1 \leq i \leq n} \Omega_i^*(F/v)$ is finite.*

Proof. We prove the statement by induction on n . If $n = 1$, then v is a \mathbb{Z} -valuation on K , and it follows by [2, Theorem 5.3] that $\Omega_1^*(F/v)$ is finite. Assume now that $n > 1$. By the induction hypothesis, for any positive integer $s < n$, for any field L carrying a \mathbb{Z}^s -valuation v' and for any function field in one variable E/L , the set $\bigcup_{1 \leq i \leq s} \Omega_i^*(E/v')$ is finite. Let $v_1 = \pi_1 \circ v$ and let \bar{v} be the residual valuation of v modulo v_1 . Then \bar{v} is a \mathbb{Z}^{n-1} -valuation on κ_{v_1} , by Remark 4.1. Let $r \in \{1, \dots, n\}$. We claim that $\Omega_r^*(F/v)$ is finite. If $r = 1$, then this follows by [2, Theorem 5.3]. Assume $r > 1$. Let w be a valuation on F such that $\mathcal{O}_w \in \Omega_r^*(F/v)$. Let $w_1 = \pi_1 \circ w$. It follows by Lemma 2.11 and by Lemma 2.8 that $\mathcal{O}_{w_1} \in \Omega_1^*(F/v)$ and that \mathcal{O}_{w_1} is an extension of \mathcal{O}_{v_1} . Since $\kappa_{w_1}/\kappa_{v_1}$ is a function field in one variable and \bar{v} is a \mathbb{Z}^{n-1} -valuation on κ_{v_1} , we have that $\Omega_{r-1}^*(\kappa_{w_1}/\bar{v})$ is finite. Furthermore, we have that \mathcal{O}_w is determined by the induced valuation ring $\mathcal{O}_{\bar{w}} \in \Omega_{r-1}^*(\kappa_{w_1}/\bar{v})$,

where \bar{w} is the residual valuation of w modulo w_1 , by Remark 2.3. Hence, we have that $|\Omega_r^*(F/v)| = \sum_{\mathcal{O} \in \Omega_1^*(F/v)} |\Omega_{r-1}^*(\kappa_{\mathcal{O}}/\bar{v})|$, where \bar{v} is the residual valuation of v modulo $v_{\mathcal{O}}$. Since for every \mathbb{Z} -valuation ν on F the set $\Omega_{r-1}(\kappa_{\nu}/\bar{v})$ is finite, we have that the set $\Omega_r^*(F/v)$ is finite, and since r was arbitrarily taken, we obtain the statement. \square

Let v be a \mathbb{Z} -valuation on K . We denote by (K^v, \hat{v}) the completion of (K, v) .

2.13. Lemma. *Let v be a henselian \mathbb{Z}^n -valuation on K , and let $v_1 = \pi_1 \circ v$. Let (K^{v_1}, \hat{v}_1) be the completion of (K, v_1) . Then v has a unique unramified extension v' to K^{v_1} , and this extension is a henselian \mathbb{Z}^n -valuation with $\kappa_{v'} = \kappa_v$.*

Proof. We observe that $\kappa_{v_1} = \kappa_{\hat{v}_1}$ by [8, Theorem 1.3.4]. Let \bar{v} be the residual valuation of v modulo v_1 , and let t be a uniformizer of \hat{v}_1 . We can consider the composition v' of \hat{v}_1 and \bar{v} with respect to t . Note that $\mathcal{O}_{v'} \cap K = \mathcal{O}_v$, and using the fact that \bar{v} and \hat{v}_1 are henselian \mathbb{Z}^{n-1} and \mathbb{Z} -valuation on κ_{v_1} and K^{v_1} , respectively, we have that v' is a henselian \mathbb{Z}^n -valuation on K^{v_1} by [8, Corollary 4.1.4], and with $\kappa_{v'} = \kappa_{\bar{v}} = \kappa_v$, by Remark 2.3. \square

Let $F/K, K'/K$ be two field extensions such that K is separably algebraically closed in K' . Then $F \otimes_K K'$ is a domain, see [11, Corollary 1, pag. 203]. We denote by FK' the fraction field of $F \otimes_K K'$ and we call it *the compositum of F and K' over K* . Note that FK' is an extension of F and of K' .

A field K that carries a henselian \mathbb{Z} -valuation v is separably algebraically closed in its completion K^v , by [19, Theorem 32.19], hence the compositum FK^v over K exists for any field extension F/K . In particular, if F/K is a function field in one variable, then FK^v/K^v is a function field in one variable.

2.14. Proposition (D. Harbater, J. Hartmann, D. Krashen). *Assume that K carries a henselian \mathbb{Z} -valuation v such that \mathcal{O}_v is excellent. Let F/K be a function field in one variable, and let E be the compositum of F and K^v over K . Let φ be a quadratic form over F of $\dim \varphi \geq 3$. If φ is isotropic over E , then φ is isotropic over F .*

Proof. See [10, Lemma 4.11]. \square

2.15. Corollary. *Assume that K carries a henselian \mathbb{Z} -valuation v such that \mathcal{O}_v is excellent. Let F/K be a function field in one variable, and let E be the compositum of F and K^v over K . Then*

$$F^\times \cap \mathbf{S}_k(E) = \mathbf{S}_k(F),$$

for any $k \geq 2$. In particular $p(F) \leq p(E)$.

Proof. Let $k \geq 2$. Let $\sigma \in F^\times \cap \mathbf{S}_k(E)$. Let $\varphi = k \times \langle 1 \rangle \perp \langle -\sigma \rangle$. Since φ is isotropic over E , then φ is isotropic over F by Proposition 2.14, hence $\sigma \in \mathbf{S}_k(F)$. The second statement follows trivially. \square

3. THE GROUP OF NONZERO SUMS OF SQUARES

Let K be a field. We define

$$s(K) = \inf\{d \in \mathbb{N} \mid -1 \in S_d(K)\} \in \mathbb{N} \cup \{\infty\},$$

the *level* of K . If $s(K) < \infty$, we say that K is *nonreal*, otherwise we say that K is real.

For a valued field (K, v) , a function field in one variable F/K , and $r \in \mathbb{N}$, we define

$$\mathcal{X}^r(F/v) = \{\mathcal{O} \in \Omega^*(F/v) \mid 2^r \leq s(\kappa_{\mathcal{O}}) < \infty\},$$

and

$$\begin{aligned} \mathcal{E}^r(F/v) &= \{x \in F^\times \mid x \in \mathcal{O}^\times F^{\times 2} \text{ for all } \mathcal{O} \in \mathcal{X}^r(F/v)\} \\ &= \{x \in F^\times \mid v_{\mathcal{O}}(x) \in 2\Gamma_{v_{\mathcal{O}}} \text{ for all } \mathcal{O} \in \mathcal{X}^r(F/v)\}. \end{aligned}$$

3.1. Lemma. *Let K be a field and $v \in V(K)$. Let F/K be a function field in one variable. Let $\mathcal{O} \in \Omega_1^*(F/v)$. Let $a \in F^\times$ and $r \in \mathbb{N}$. If $a \in \mathcal{E}^r(F/v) \cap \mathcal{O}^\times$, then $\bar{a} \in \mathcal{E}^r(\kappa_{\mathcal{O}}/\bar{v})$, where \bar{v} is the residual valuation of v modulo $v_{\mathcal{O}}|_K$.*

Proof. Let $v_1 = v_{\mathcal{O}}|_K$. It follows by Lemma 2.8 and Proposition 2.10 that v_1 is equivalent to a \mathbb{Z} -valuation. Let \bar{v} be the residual valuation of v modulo v_1 . Let $a \in \mathcal{E}^r(F/v) \cap \mathcal{O}^\times$. Let $\mathcal{O}' \in \mathcal{X}^r(\kappa_{\mathcal{O}}/\bar{v})$. We need to show that $\bar{a} \in \mathcal{O}'\kappa_{\mathcal{O}'}^{\times 2}$. Let $v' = v_{\mathcal{O}'}|_{\kappa_{v_1}}$. Let π, π' be uniformizers of $v_{\mathcal{O}}$ and v_1 respectively. Let w be a composition of $v_{\mathcal{O}}$ and $v_{\mathcal{O}'}$ with respect to π , and let ν be a composition of v_1 and v' with respect to π' . Since $\kappa_{\mathcal{O}'} = \kappa_w$ and $\kappa_{v'} = \kappa_\nu$ by Remark 2.3, we have that κ_w/κ_ν is a function field in one variable with $2^r \leq s(\kappa_w) < \infty$ and $\mathcal{O}_v \subseteq \mathcal{O}_{w|_K} = \mathcal{O}_\nu$, that is $\mathcal{O}_w \in \mathcal{X}^r(F/v)$. Then $a \in \mathcal{O}_w^\times F^{\times 2} \cap \mathcal{O}^\times$, because $a \in \mathcal{E}^r(F/v)$. This implies that $\bar{a} \in \mathcal{O}'^\times \kappa_{\mathcal{O}'}^{\times 2}$, because $(0, v_{\mathcal{O}'}(\bar{a})) = w(a) \in 2\Gamma_w$. \square

We say that a valuation v on K is real or nonreal, respectively, if κ_v has the corresponding property. We say that a valuation v on K is nondyadic if $\text{char}(\kappa_v) \neq 2$. We recall that an ordered abelian group that admits a minimal positive element is called *discrete*.

3.2. Lemma. *Let v be a nonreal and nondyadic valuation on K such that Γ_v is discrete. Then $s(\kappa_v) < p(K)$.*

Proof. Let $d \in \mathbb{N}$ be such that $d = s(\kappa_v)$. There exist $f \in \mathfrak{m}_v, x_1, \dots, x_d \in K$ such that $f = 1 + x_1^2 + \dots + x_d^2$. Let $b = (1 - \frac{f}{2})^2 + x_1^2 + \dots + x_d^2 = \frac{f^2}{4}$. Let $\gamma \in \Gamma_v$ be the minimal positive element and let $z \in K$ be such that $v(z) = \gamma$. Note that $\gamma \notin 2\Gamma_v$. Hence $0 < v(z) < 2v(f) = v(b)$. Let $\sigma = (z - (1 - \frac{f}{2}))^2 + x_1^2 + \dots + x_d^2$. Since $\sigma = z(z - 2(1 - \frac{f}{2})) + b$, $0 < v(z) < v(b)$, and since $v(2 - f) = 0 < v(z)$, we have that $v(\sigma) = \gamma$. It follows by Lemma [3, Lemma 4.1] that $\sigma \notin S_d(K)$, and hence $s(\kappa_v) < p(K)$. \square

3.3. Lemma. *Let $r \in \mathbb{N}^+$. Let K be a field and v a real henselian valuation in $V(K)$ such that $p(\kappa_v(X)) \leq 2^r$. Let $v_1 = \pi_1 \circ v$. Let \bar{v} be the residual valuation of v modulo v_1 . Let (K^{v_1}, \hat{v}_1) be the completion of (K, v_1) . Let v' be a composition of \hat{v}_1 and \bar{v} with respect to a uniformizer of \hat{v}_1 . Let F/K be a regular function field in one variable. Let E be the compositum of F and K^{v_1} over K . Then*

$$\mathbf{S}(F) \cap \mathcal{E}^r(F/v) \subseteq \mathbf{S}(E) \cap \mathcal{E}^r(E/v').$$

Proof. Let $\mathcal{O} \in \mathcal{X}^r(E/v')$. Let $\mathcal{O}_F = \mathcal{O} \cap F$ and $\mathcal{O}_K = \mathcal{O} \cap K$. We claim that $\mathcal{O}_F \in \mathcal{X}^r(F/v)$ or $\kappa_{\mathcal{O}_F}$ is real. By definition we have that $\mathcal{O}_{v'} \subseteq \mathcal{O} \cap K^{v_1}$, and since v is henselian, we have that $\mathcal{O} \cap K^{v_1} \subseteq \mathcal{O}_{\hat{v}_1} \subseteq K^{v_1}$, by Proposition 2.7. Hence $\mathcal{O}_v \subseteq \mathcal{O}_K \subseteq \mathcal{O}_{v_1} \subseteq K$. We denote by ν the residual valuation of v modulo $v_{\mathcal{O}_K}$. We recall that ν is a henselian valuation on $\kappa_{\mathcal{O}_K}$ such that $\kappa_\nu = \kappa_v$, by [8, Corollary 4.1.4]. Since $p(\kappa_v(X)) \leq 2^r$, it follows by [5, Theorem 3.5] that $s(L) \leq 2^{r-1}$ for every finite nonreal extension L/κ_v . Let $L'/\kappa_{\mathcal{O}_K}$ be a finite nonreal extension. Since ν is henselian, any extension of ν to L' is again henselian, and hence we obtain that $s(L') \leq 2^{r-1}$. Since $\kappa_{\mathcal{O}_F} \subseteq \kappa_{\mathcal{O}}$, we have that $s(\kappa_{\mathcal{O}_F}) \geq s(\kappa_{\mathcal{O}}) \geq 2^r$ and hence the extension $\kappa_{\mathcal{O}_F}/\kappa_{\mathcal{O}_K}$ cannot be algebraic. Hence $\kappa_{\mathcal{O}_F}/\kappa_{\mathcal{O}_K}$ is transcendental. If $\kappa_{\mathcal{O}_F}$ is nonreal, since $s(\kappa_{\mathcal{O}_F}) \geq 2^r$, then $2^r < p(\kappa_{\mathcal{O}_F})$, by Lemma 3.2, because $v_{\mathcal{O}|_F}$ is equivalent to a valuation in $V(F)$, by Proposition 2.10. This implies that $\kappa_{\mathcal{O}_F}/\kappa_{\mathcal{O}_K}$ cannot be ruled, by [5, Theorem 3.5], whenever $\kappa_{\mathcal{O}_F}$ is nonreal. Hence $\mathcal{O}_F \in \mathcal{X}^r(F/v)$ because $\mathcal{O}_v \subseteq \mathcal{O}_K$. We conclude that $s(\kappa_{\mathcal{O}_F}) \geq 2^r$.

Let $a \in \mathbf{S}(F) \cap \mathcal{E}^r(F/v)$. Hence $a \in \mathcal{O}_F^\times F^{\times 2}$ because $\mathcal{O}_F \in \mathcal{X}^r(F/v)$ or $\kappa_{\mathcal{O}_F}$ is real, and the latter follows by [3, Lemma 4.1]. Therefore $a \in \mathcal{O}_F^\times F^{\times 2} \subseteq \mathcal{O}^\times E^{\times 2}$. Since \mathcal{O} was arbitrarily chosen, we conclude that $a \in \mathbf{S}(E) \cap \mathcal{E}^r(E/v')$. \square

For a field K , we set

$$p'(K) = \begin{cases} p(K) & \text{if } K \text{ is real,} \\ s(K) + 1 & \text{if } K \text{ is nonreal.} \end{cases}$$

3.4. Lemma. *Let K be a perfect field. Let L/K be an algebraic extension. Then*

$$p'(L) \leq p(K(X)).$$

Proof. If the extension L/K is finite, then the result follows from [3, Lemma 6.3]. Assume that L/K is algebraic. If $p(K(X)) = \infty$, then the inequality is trivially satisfied. Assume $p(K(X)) < \infty$. Let $r = p(K(X))$, and let $\sigma \in \mathbf{S}_{r+1}(L)$. Let $x_0, \dots, x_r \in L$ be such that $\sigma = x_0^2 + \dots + x_r^2$, and let $K' = K(x_0, \dots, x_r)$. Hence K'/K is finite. Then $p(K') \leq r$, by [3, Lemma 6.3], and since $K' \subseteq L$, we have that $\sigma \in \mathbf{S}_r(L)$ because $\mathbf{S}_r(K') \subseteq \mathbf{S}_r(L)$. Therefore $p(L) \leq r$. \square

3.5. Proposition. *Let $r \in \mathbb{N}^+$. Let K be a field and v a real henselian valuation in $V(K)$ such that $p(\kappa_v(X)) \leq 2^r$. Let F/K be a regular function field in one variable. Then*

$$\mathbf{S}(F) \cap \mathcal{E}_r(F/v) = \mathbf{S}_{2^r}(F).$$

Proof. It follows from [3, Lemma 4.1] that $S_{2^r}(F) \subseteq S(F) \cap \mathcal{E}^r(F/v)$. Let $n = \text{rk}(v)$. Let us now show by induction over n that $S(F) \cap \mathcal{E}^r(F/v) \subseteq S_{2^r}(F)$. For $n = 0$, it follows trivially from the assumption.

Let now $n > 0$. Let $v_1 = \pi_1 \circ v$. Let (K^{v_1}, \hat{v}_1) be the completion of (K, v_1) . Let v' be a composition of \hat{v}_1 and \bar{v} with respect to a uniformizer of \hat{v}_1 . It follows by Lemma 2.13 that v' is a henselian valuation of rank n on K^{v_1} such that $\kappa_v = \kappa_{v'}$ and $\mathcal{O}_{v'} \cap K = \mathcal{O}_v$.

Let E be the compositum of F and K^{v_1} over K . Let $\sigma \in S(F) \cap \mathcal{E}^r(F/v)$. Then $\sigma \in S(E) \cap \mathcal{E}^r(E/v')$, by Lemma 3.3. Let φ be the quadratic form $2^r \times \langle 1 \rangle \perp \langle -\sigma \rangle$ over F . We first show that φ is isotropic over E .

Let $w \in V_1(E)$. We claim that $p(E^w) \leq 2^r$. If $w|_{K^{v_1}}$ is trivial, then κ_w/K^{v_1} is a finite field extension, and it follows that $p(E^w) \leq p'(\kappa_w)$, by [3, Proposition 4.3] and that $p'(\kappa_w) \leq 2^r$ by [5, Theorem 3.5]. Let us assume now that $\mathcal{O}_w \cap K^{v_1} = \mathcal{O}_{\hat{v}_1}$ and $\kappa_w/\kappa_{\hat{v}_1}$ is algebraic. Since the residual valuation \bar{v}' of v' modulo \hat{v}_1 is a henselian valuation on $\kappa_{\hat{v}_1}$ such that $p(\kappa_{\bar{v}'}(X)) \leq 2^r$, it follows by [5, Theorem 3.5] and Lemma 3.4 that $p'(\kappa_w) \leq p(\kappa_{\hat{v}_1}(X)) \leq 2^r$. Since $p(E^w) \leq p'(\kappa_w) \leq 2^r$ by Lemma 3.4 and by [3, Proposition 4.3], we have $p(E^w) \leq 2^r$. Therefore φ is isotropic over E^w in both cases.

Let us assume now that $\mathcal{O}_w \cap K^{v_1} = \mathcal{O}_{\hat{v}_1}$ and $\kappa_w/\kappa_{\hat{v}_1}$ is a function field in one variable. We claim that φ is isotropic over E^w . If $s(\kappa_w) \leq 2^{r-1}$, since $s(E^w) = s(\kappa_w)$, we have that φ is isotropic over E^w . Let $d \in \mathbb{N}$, $x_1, \dots, x_d \in F$ be such that $\sigma = x_1^2 + \dots + x_d^2$. If κ_w is real, then $w(\sigma) = \min\{2w(x_1), \dots, 2w(x_d)\}$, by [3, Lemma 4.1]. If $2^r \leq s(\kappa_w) < \infty$, then $2^r < p(\kappa_w)$ by Lemma 3.2. Hence $\kappa_w/\kappa_{\hat{v}_1}$ cannot be ruled by [5, Theorem 3.5], which implies that $\mathcal{O}_w \in \mathcal{X}^r(E/v')$. In any case $w(\sigma) \in 2\Gamma_w$, because $\sigma \in \mathcal{E}^r(E/v')$, and thus $w(\sigma) = 2w(y)$ for some $y \in E$. Let $\tau = \sigma y^{-2}$. Then $\bar{\tau} \in S(\kappa_w)$ and, if $\varphi' = 2^r \times \langle 1 \rangle \perp \langle -\tau \rangle$ is isotropic over E^w , then φ is isotropic over E^w . By the induction hypothesis, we have that $S_{2^r}(\kappa_w) = S(\kappa_w) \cap \mathcal{E}^r(\kappa_w/\bar{v}')$. Since $\tau \in \mathcal{E}^r(E/v')$, it follows by Lemma 3.1 that $\bar{\tau} \in \mathcal{E}^r(\kappa_w/\bar{v}')$, and hence $\bar{\tau} \in S_{2^r}(\kappa_w)$. Hence φ'_r is isotropic over κ_w , and then φ' is isotropic over E^w , because the extension \hat{w} of w to E^w is henselian, whereby φ is isotropic over E^w .

By [9, Theorem 6.1], φ is isotropic over E if and only if φ is isotropic over E^w for every rank one valuation w on E such that $w|_{K^{v_1}}$ is trivial or $\mathcal{O}_w \cap K^{v_1} = \mathcal{O}_{\hat{v}_1}$. If $\mathcal{O}_w \cap K^{v_1} = \mathcal{O}_{\hat{v}_1}$ then it follows by Lemma 2.8 that, either $\kappa_w/\kappa_{\hat{v}_1}$ is algebraic or $\kappa_w/\kappa_{\hat{v}_1}$ is a function field in one variable. By the above, in any case φ is isotropic over E , and it follows from Corollary 2.15 that φ is isotropic over F , whereby $\sigma \in S_{2^r}(F)$. \square

For a field K and $r \in \mathbb{N}$, we set $G_r(K) = S(K)/S_{2^r}(K)$.

3.6. Theorem. *Let $r \in \mathbb{N}^+$ and $n \in \mathbb{N}$. Let K be a field carrying a real henselian \mathbb{Z}^n -valuation v such that $p(\kappa_v(X)) \leq 2^r$. Let F/K be a function field in one variable. Then*

$$|G_r(F)| = 2^{|\mathcal{X}^r(F/v)|}.$$

In particular $|G_r(F)|$ is finite.

Proof. It follows by Theorem 2.12 that we may choose a finite and saturated set $W \subseteq \Omega(F)$ such that $\mathcal{X}^r(F/v) \subseteq W$, and it follows by Proposition 2.4 that we can choose a coherent subset $S' \subseteq V(F)$ such that $W = \{\mathcal{O}_w \mid w \in S'\}$. We consider $S = \{w \in S' \mid \mathcal{O}_w \in \mathcal{X}^r(F/v)\}$. Let $\Phi : S(F) \rightarrow \prod_{w \in S} \mathbb{Z}/2\mathbb{Z}$ be the map given by the composition of $\Phi_{S'}|_{S(F)}$, where $\Phi_{S'} : F^\times \rightarrow \prod_{w \in S'} \mathbb{Z}$ is defined in Equation (2.5), and the natural surjective map

$$\prod_{w \in S'} \mathbb{Z} \rightarrow \prod_{w \in S} \mathbb{Z} \rightarrow \prod_{w \in S} \mathbb{Z}/2\mathbb{Z}.$$

We claim that Φ is a surjective group homomorphism with $\ker(\Phi) = S_{2^r}(F)$.

First, we observe that Φ is a group homomorphism simply because valuations and projections are group homomorphisms. The inclusion $S_{2^r}(F) \subseteq \ker(\Phi)$ follows directly from Proposition 3.5. Let us show that $\ker(\Phi) \subseteq S_{2^r}(F)$. Let $\sigma \in \ker(\Phi)$ and we assume that $\sigma \notin S_{2^r}(F)$. By Proposition 3.5, $\sigma \notin \mathcal{O}^\times F^{\times 2}$, for some $\mathcal{O} \in \mathcal{X}^r(F/v)$. Let $w \in S$ be such that $\mathcal{O}_w = \mathcal{O}$, and let $a_1, \dots, a_{\text{rk}(w)} \in \mathbb{Z}$ be such that $w(\sigma) = (a_1, \dots, a_{\text{rk}(w)})$. Since $w(\sigma) \notin 2\mathbb{Z}$, there exists $d \leq \text{rk}(w)$ such that $a_d \notin 2\mathbb{Z}$. Let $w' = \pi_d \circ w$. Since $w'(\sigma) \notin 2\mathbb{Z}$, the residue field $\kappa_{w'}$ cannot be real by [3, Lemma 4.1]. Let \bar{w} be the residual valuation of w modulo w' . Then $2^r \leq s(\kappa_w) = s(\kappa_{\bar{w}}) \leq s(\kappa_{w'}) < \infty$, which implies that $\mathcal{O}_{w'} \in \mathcal{X}^r(F/v)$. Since S' is coherent, we obtain that $w' \in S'$, and since $\mathcal{O}_{w'} \in \mathcal{X}^r(F/v)$, we have $w' \in S$. But $\pi^1(w'(\sigma)) = a_d \notin 2\mathbb{Z}$, which contradicts the fact that $\sigma \in \ker(\Phi)$. This shows that $\ker(\Phi) = S_{2^r}(F)$.

We show now that Φ is surjective. Let $(e_w)_{w \in S}$ be the canonical basis of $\prod_{w \in S} \mathbb{Z}/2\mathbb{Z}$ as a $\mathbb{Z}/2\mathbb{Z}$ -module. Consider $w \in S$, and let $d = \text{rk}(w)$. We claim that there exists $\sigma \in S(F)$ with $\Phi(\sigma) = e_w$. Since $2^r \leq s(\kappa_w) < \infty$, there exists $f \in \mathfrak{m}_w$ and $x_1, \dots, x_m \in \mathcal{O}_w^\times$, for some $m \in \mathbb{N}$, such that $f = 1 + x_1^2 + \dots + x_m^2$. Let $b = (1 - \frac{f}{2})^2 + x_1^2 + \dots + x_m^2 = \frac{f^2}{4}$. Note that $w(b) = 2w(f) > e_1^d$, where e_1^d is the minimal positive element of \mathbb{Z}^d . By Proposition 2.6, there exists $z \in F$ such that for all $\nu \in S$ we have that $\nu(z) = e_{\text{rk}(\nu)+1-d}^{\text{rk}(\nu)}$ if ν is a refinement of w and $\nu(z) < \min\{0, \nu(f)\}$ otherwise. Let $\sigma = (z - (1 - \frac{f}{2}))^2 + x_1^2 + \dots + x_m^2$ in $S(F)$. Let $\nu \in S$ be a refinement of w . Since $\sigma = z(z - 2(1 - \frac{f}{2})) + b$ and $\nu(z) < \nu(b)$ we have that $\nu(\sigma) = \nu(z)$. Thus $\pi^1(\nu(\sigma)) = 0$ if \mathcal{O}_ν is a proper refinement of \mathcal{O}_w and $\pi^1(w(\sigma)) = \pi^1(e_1^d) = 1$. Consider $\nu \in S$ not a refinement of w . Then $\nu(\sigma) = \nu(z(z - 2(1 - \frac{f}{2}))) = 2\nu(z)$. Thus $\pi^1(\nu(\sigma)) \in 2\mathbb{Z}$. Hence $\Phi(\sigma) = e_w$. Therefore Φ is surjective and $|G_r(F)| = 2^{|\mathcal{X}^r(F/v)|}$. \square

A field K is called *hereditarily pythagorean* if K is real and $p(L) = 1$ for every finite real extension L of K .

3.7. Corollary. *Let $r \in \mathbb{N}^+$ and $n \in \mathbb{N}$. Assume that K carries a henselian \mathbb{Z}^n -valuation v such that κ_v is hereditarily pythagorean. Let F/K be a function field*

in one variable. If $|\kappa_v^\times / \kappa_v^{\times 2}| = 2^r$, then

$$|G_r(F)| = 2^{|\mathcal{X}_r(F/v)|}.$$

Proof. This follows directly from [1, Corollary 3.3] and Theorem 3.6. \square

4. HYPERELLIPTIC FUNCTION FIELDS

A hyperelliptic function field F/K is the function field generated by two variables X, Y subject to the relation $Y^2 = f(X)$, for some square-free polynomial f over K .

In this section, we will apply the valuation-theoretic description of the order of $G(F) := G_1(F)$ from the previous section in the case where F/K is a hyperelliptic function field where K carries a henselian \mathbb{Z}^n -valuation v such that κ_v is hereditarily pythagorean with $\kappa_v^\times = \kappa_v^{\times 2} \cup -\kappa_v^{\times 2}$, or equivalently, such that κ_v is hereditarily euclidean. Under this assumptions, it is a consequence of [1, Theorem 5.8.2] and [5, Corollary 4.6] that $p(F) \leq 3$ (even if F/K is not hyperelliptic). The particular case where $K = \mathbb{R}((t_1)) \dots ((t_n))$ was showed earlier in [3, Theorem 6.12]. Thus, in this case we have that

$$G(F) = \mathcal{S}_3(F)/\mathcal{S}_2(F).$$

Let F/K be a hyperelliptic function field. Let $f \in K[X]$ be a square-free polynomial such that $F = K(X)(\sqrt{f})$. Let $g = \lfloor \frac{\deg f - 1}{2} \rfloor$. The integer g is the *genus* of the function field F . See [14, Proposition 7.4.24]. In [5, Theorem 3.10], an upper bound for $|G(F)|$ in terms of the order of the square class group of the root fields of the distinct nonreal irreducible factors of f , is given in the general case where K is an arbitrary hereditarily pythagorean field. In the case where K is a field carrying a henselian \mathbb{Z}^n -valuation v with hereditarily euclidean residue field the bound becomes

$$|G(F)| \leq 2^{n(g+1)},$$

see Remark 4.1. We will then proceed to construct hyperelliptic function fields F/K such that the set $\mathcal{X}^1(F/v)$ satisfies $|\mathcal{X}^1(F/v)| \geq n(g+1)$, which then immediately yields the equalities $|G(F)| = 2^{n(g+1)}$ and $|\mathcal{X}^1(F/v)| = n(g+1)$. In particular the upper bound in [5, Theorem 3.10] is optimal.

4.1. Remark. Let $n \in \mathbb{N}$ and let K be a field with a henselian \mathbb{Z}^n -valuation v and hereditarily euclidean residue field. Then K is a hereditarily pythagorean field, by [7, Proposition 3.5]. Let $f \in K[X]$ be a square-free polynomial and let $F = K(X)(\sqrt{f})$. Let K_1, \dots, K_r be the root fields of the distinct nonreal irreducible factors of f . By [5, Theorem 3.10] we have that

$$|G(F)| \leq \prod_{i=1}^r |K_i^\times / K_i^{\times 2}|.$$

Thus $-1 \in K_i^{\times 2}$ for all $1 \leq i \leq r$, by [6, III, Theorem 1]. Since K_i/K is a finite extension, there exist a henselian \mathbb{Z}^n -valuation v' on K_i such that $\mathcal{O}_{v'} \cap K = \mathcal{O}_v$ and $-1 \in k_{v'}^{\times 2}$. It is well-known that $|K_i^\times/K_i^{\times 2}| = 2^n |\kappa_{v'}^\times/\kappa_{v'}^{\times 2}|$. By [13, Chap. VII, (7.15)] we have that $\kappa_{v'}$ is quadratically closed, whence $|K_i^\times/K_i^{\times 2}| = 2^n$, for all $1 \leq i \leq r$. Let $g \in \mathbb{N}$ be such that $\deg f = 2g+1$ or $2g+2$. Then $r \leq g+1$, hence

$$|G(F)| \leq 2^{n(g+1)}.$$

It follows that the unique situation where the bound can be optimal is when $f \in K[X]$ is a square-free polynomial of degree $2(g+1)$ and has $g+1$ nonreal irreducible factors.

4.2. Lemma. *Let K be a real field such that $p(K) = 1$. Let v be a valuation on K . Let $f \in \mathcal{O}_v[X]$ be a monic quadratic irreducible polynomial such that $K[X]/(f)$ is nonreal. Then there exist $a, b \in \mathcal{O}_v$ such that $f = (X - a)^2 + b^2$.*

Proof. Let $\alpha, \beta \in \mathcal{O}_v$ be such that $f = X^2 + \alpha X + \beta$. Thus, we can write $f = (X - \alpha/2)^2 + (\beta - \alpha^2/4)$. Since $K[X]/(f) = K(\sqrt{\alpha^2 - 4\beta})$ is nonreal, then $-(\alpha^2 - 4\beta) \in \mathbf{S}(K) = K^{\times 2}$, by [13, Chap. VIII, Lemma 1.4]. The statement follows considering $a = \alpha/2$ and $b = \sqrt{(4\beta - \alpha^2)/4}$. \square

4.3. Lemma. *Assume that K carries a \mathbb{Z} -valuation v . Let $f \in K[X]$ be a non-constant square-free polynomial of degree d , and let $F = K(X)(\sqrt{f})$. Then F is K -isomorphic to*

$$K(X)(\sqrt{\alpha \cdot q_1 \cdots q_r}),$$

for some $r \in \mathbb{N}$, $q_1, \dots, q_r \in \mathcal{O}_v[X]$ monic irreducible such that $d = \sum_{i=1}^r \deg q_i$, and where $\alpha = 1$ if d is odd and otherwise α is the leading coefficient of f .

Proof. Since $\text{Frac}(\mathcal{O}_v) = K$, we can assume that $f \in \mathcal{O}_v[X]$. Let $d = \deg f$. Let $a_0, \dots, a_d \in \mathcal{O}_v$ be such that $f = \sum_{i=0}^d a_i X^i$. Since F is the function field of $Y^2 = f(X)$, multiplying by $a_d^{2(d-1)}$, we have that $(a_d^{d-1} Y)^2 = a_d^{d-1} g(X)$, where $g = \sum_{i=0}^d b_i X^i$, where $b_d = a_d^d$ and $b_i = a_i a_d^{d-1}$ for all $0 \leq i \leq d-1$. Replacing $X' = a_d X$, and $Y' = a_d^{d-1} Y$, we have that F is K -isomorphic to $K(X') \left(\sqrt{a_d^{d-1} g(X')} \right)$. Write $g(X') = q_1(X') \cdots q_r(X')$, where $q_i(X') \in K[X']$ are monic irreducible polynomials, for some $r \in \mathbb{N}$. Since \mathcal{O}_v is a Unique Factorization Domain, by Gauss' Lemma, we may assume that $q_1(X'), \dots, q_r(X') \in \mathcal{O}_v[X']$, which concludes the proof. \square

Let v be a valuation on a field K . By [8, Corollary 2.2.2], there exists a unique extension w of v to the rational function field $K(X)$ such that $w(X) = 0$ and the residue $\overline{X} \in \kappa_w$ is transcendental over κ_v . A valuation w with these properties is called the *Gauss extension of v to F with respect to X* .

4.4. Lemma. *Let $n \in \mathbb{N}$. Assume that K is a field carrying a henselian \mathbb{Z}^n -valuation with hereditarily euclidean residue field. Let F/K be a regular function field of genus zero. Then*

$$|G(F)| = \begin{cases} 2^n & \text{if } F \text{ is nonreal,} \\ 1 & \text{if } F \text{ is real.} \end{cases}$$

Proof. We assume that F is nonreal. Then $F = K(X)(\sqrt{\alpha \cdot q})$, where q is a monic irreducible quadratic polynomial over K , by [12, Theorem 5.7.2, Theorem 5.7.3]. Then $q(X) = (X - a)^2 + b^2$, for some $a, b \in K$, by Lemma 4.2. Moreover $\alpha \in -K^{\times 2}$, because F is nonreal. Replacing $X' = X - a$, we have that F is isomorphic to $K(X)(\sqrt{-(X^2 + 1)})$. We prove that statement by induction on n . For $n = 0$, it follows from [5, Corollary 4.6] that $G(F)$ is trivial. Let $n \geq 1$. Let $v_1 = \pi_1 \circ v$. Let w' be the Gauss extension of v_1 to $K(X)$ with respect to X and let w be an extension of w' to F . Since κ_{v_1} is real, the polynomial $X^2 + 1$ is irreducible over κ_{v_1} . Then we have $\kappa_w = \kappa_{v_1}(\overline{X}) \left(\sqrt{-(\overline{X}^2 + 1)} \right)$, and $\mathcal{O}_w \in \mathcal{X}^1(F/v) \cap \Omega_1(F)$. It follows from [4, Corollary 3.6] that \mathcal{O}_w is the unique valuation ring in $\mathcal{X}^1(F/v) \cap \Omega_1(F)$, and hence every $\mathcal{O} \in \mathcal{X}^1(F/v)$ is a refinement of \mathcal{O}_w , by Remark 2.3. Let \overline{v} be the residual valuation of v modulo v_1 . Thus $|\mathcal{X}^1(F/v)| = 1 + |\{\mathcal{O} \in \Omega(\kappa_w) \mid \mathcal{O} \in \mathcal{X}^1(\kappa_w/\overline{v})\}|$, whereby $|G(F)| = 2|G(\kappa_w)|$, by Theorem 3.6. Note that \overline{v} is a henselian \mathbb{Z}^{n-1} -valuation on κ_{v_1} with hereditarily euclidean residue field. Thus, by the induction hypothesis $|G(\kappa_w)| = 2^{n-1}$. Therefore $|G(F)| = 2^n$.

If F is real, then $p(F) = 2$ by [18, Theorem 3], whereby $|G(F)| = 1$. \square

4.5. Example. Let $g, n \in \mathbb{N}$. Let F be the function field of the curve

$$Y^2 = - \prod_{i=0}^g (X^2 + t_n^{2i})$$

over $K_n := \mathbb{R}((t_1)) \dots ((t_n))$. We will show that $|G(F)| = 2^{n(g+1)}$. Let v_{t_n} be the t_n -adic valuation on K_n . Note that $\mathcal{O}_{v_{t_n}} = K_{n-1}[[t_n]]$. For $0 \leq i \leq g$, let v_i be the Gauss extension of v_{t_n} to $K_n(X)$ with respect to Xt_n^{-i} and let w_i be an extension of v_i to F .

We claim that $\kappa_{w_i} \simeq K_{n-1}(X)(\sqrt{-(X^2 + 1)})$, for all $0 \leq i \leq g$. Let $0 \leq i \leq g$, and let $Z := Xt_n^{-i}$. Hence $F = K_n(Z)(\sqrt{-g(Z)})$ since $K_n(X) = K_n(Z)$, where $g = (t_n^{2i}Z^2 + 1)(t_n^{2i-2}Z^2 + 1) \dots (Z^2 + 1) \dots (Z^2 + t_n^{2(g-i)})$ is a square-free polynomial over K_n . Then $g \in \mathcal{O}_{v_i}^\times$ and $\kappa_{w_i} = K_{n-1}(\overline{Z})(\sqrt{-\overline{g}}) = K_{n-1}(\overline{Z}) \left(\sqrt{-(\overline{Z}^2 + 1)} \right)$.

Let v be a henselian \mathbb{Z}^n -valuation on K_n . Let \overline{v} be the residual valuation of v modulo v_{t_n} . Since $|\mathcal{X}^1(\kappa_{w_i}/\overline{v})| = |\{\mathcal{O} \in \Omega(F) \mid \mathcal{O} \subsetneq \mathcal{O}_{w_i} \text{ and } \mathcal{O} \in \mathcal{X}^1(F/v)\}|$ by Remark 2.3 and $|\mathcal{X}^1(\kappa_{w_i}/\overline{v})| = n - 1$, by Lemma 4.4, for every $0 \leq i \leq g$, we have that $|\mathcal{X}^1(F/v)| \geq (g + 1) + (n - 1)(g + 1) = n(g + 1)$. Therefore $|G(F)| \geq 2^{n(g+1)}$.

by Theorem 3.6, whereby

$$|G(F)| = 2^{n(g+1)},$$

by Remark 4.1.

Let F/K be a function field in one variable of genus zero, where K is any hereditarily pythagorean field, that is $F = K(X)(\sqrt{aX^2 + b})$, for some $a, b \in K$. S. Tikhonov and V.I. Yanchevskii in [18, Theorem 3] showed that if F is real, then $p(F) = 2$. Hence, if we assume that K carries a henselian \mathbb{Z}^n -valuation with hereditarily euclidean residue field, for some $n > 0$, then $|G(F)| = 2^n$, implies that F is a nonreal field of level two, by Lemma 4.4. This motivates the following question.

4.6. Question. *Let $n, g \in \mathbb{N}$. Let K be a field carrying a henselian \mathbb{Z}^n -valuation with hereditarily euclidean residue field. Let F/K be a hyperelliptic function field of genus g . Does the equality $|G(F)| = 2^{n(g+1)}$ imply that F is a nonreal field ?*

We show in Corollary 4.9 that this question has an affirmative answer in the case where $n = 1$.

4.7. Proposition. *Let $g \in \mathbb{N}$. Assume that K carries a henselian \mathbb{Z} -valuation v with κ_v hereditarily pythagorean. Let $f \in K[X]$ be a square-free polynomial of degree $2g + 2$ with all roots in $K(\sqrt{-1}) \setminus K$. Set $F = K(X)(\sqrt{f})$. Let w be a residually transcendental extension of v to F . Assume that F is real. Then one of the following conditions holds:*

- (a) κ_w is nonreal with $s(\kappa_w) = 1$.
- (b) κ_w/κ_v is ruled.
- (c) $\kappa_w = \kappa_v(X)(\sqrt{h})$ is a real field with $h \in \kappa_v[X]$ a square-free polynomial with all roots in $\kappa_v(\sqrt{-1}) \setminus \kappa_v$.

Proof. By Lemma 4.2 and by Lemma 4.3 we may choose irreducible polynomials $q_i = (X - a_i)^2 + b_i^2$ with $a_i, b_i \in \mathcal{O}_v$, and $\alpha \in K^\times$ such that $f = \alpha \cdot q_0 \cdots q_g$. Since F is real, $\alpha \notin -K^{\times 2}$. We set $Y_i = (X - a_i)b_i^{-1}$.

Let w be a residually transcendental extension of v to F and let $w_0 = w|_{K(X)}$. As $[F : K(X)] = 2$ we have that $[\kappa_w : \kappa_{w_0}] \leq 2$. Let ℓ be the relative algebraic closure of κ_v in κ_w . If ℓ is nonreal, since κ_v is a hereditarily pythagorean, we have that $-1 \in \kappa_w^{\times 2}$ (case (a)).

Thus we assume now that ℓ is real. If $\kappa_w = \kappa_{w_0}$, since w is a residually transcendental extension of v , by Proposition 2.9 we have that $\kappa_w = \kappa_{w_0} = \ell(\overline{T})$, for some $T \in \mathcal{O}_{w_0}^\times$ with \overline{T} transcendental over κ_v (case (b)). Now assume that $[\kappa_w : \kappa_{w_0}] = 2$ and ℓ is real. Then $\Gamma_w = \Gamma_{w_0}$. Let $f' = q_0 \cdots q_g$. Since $f \in F^{\times 2}$, we have that $w_0(f) \in 2\Gamma_{w_0}$ and since $f' \in \mathcal{S}_2(K(X))$ we have $w_0(f') \in 2\Gamma_{w_0}$, by [3, Lemma 4.1]. Hence $w_0(\alpha) \in 2\Gamma_{w_0}$. Consider $i \in \{0, \dots, g\}$. Set

$$Z_i = \begin{cases} 1 + Y_i^2 & \text{if } w(Y_i) \geq 0, \\ 1 + Y_i^{-2} & \text{if } w(Y_i) < 0. \end{cases}$$

Note that, since ℓ is real, $Z_i \in \mathcal{O}_w^\times$ for all $0 \leq i \leq g$. Thus

$$\overline{Z}_i = \begin{cases} 1 + \overline{Y}_i^2 & \text{if } w(Y_i) = 0, \\ 1 & \text{otherwise.} \end{cases}$$

Let us first assume that $v(\alpha) \notin 2\mathbb{Z}$. This implies that w_0 is a ramified extension of v . Hence \overline{Y}_i is algebraic over κ_v for all $i \in \{0, \dots, g\}$, because otherwise w_0 must be a Gauss extension of v with respect to some Y_i , which would contradict the ramification. Since ℓ is pythagorean, we have $\overline{Z}_i \in \ell^{\times 2} \subseteq \kappa_w^{\times 2}$ because ℓ is the relative algebraic closure of κ_v in κ_w . By Proposition [4, Lemma 2.2] we have that $\kappa_w = \kappa_{w_0}(\sqrt{u})$, for any $u \in \alpha f' K(X)^{\times 2}$ with $w_0(u) = 0$. Since $\prod_{i=0}^g \overline{Z}_i \in \kappa_{w_0}^{\times 2}$, we have that $\kappa_w = \kappa_{w_0}(\sqrt{\alpha h^2})$ for any $h \in K(X)^\times$ such that $w_0(\alpha h^2) = 0$. Let $K' = K(\sqrt{\alpha})$. Let w'_0 be an extension of w_0 to $K'(X)$ and let $v' = w'_0|_{K'}$. We have that $\kappa_{w'_0} = \kappa_{w_0}(\sqrt{\alpha h^2})$, because $K'(X) = K(X)(\sqrt{\alpha})$ and $w_0(\alpha h^2) = 0$. Furthermore, w'_0 is a residually transcendental extension of v' to $K'(X)$. Therefore κ_w/κ_v is ruled, by Proposition 2.9 (case (b)).

Let us assume now that $v(\alpha) \in 2\mathbb{Z}$. Let J be the set of indices $i \in \{0, \dots, g\}$ such that $w(Y_i) = 0$ and \overline{Y}_i is transcendental over κ_v . Assume first that $J = \emptyset$. Then $\overline{Z}_i \in \kappa_w$ is algebraic over ℓ for all $0 \leq i \leq g$, and since ℓ is pythagorean, we have that $\kappa_w = \kappa_{w_0}(\sqrt{\alpha})$. Therefore κ_w/κ_v is ruled (case (b)). Now we assume that $J \neq \emptyset$. Without loss of generality we put $J = \{0, \dots, s\}$, for some $s \leq g$. For $i \in J$, let v_i be the Gauss extension of v to $K(X)$ with respect to Y_i . By [8, Corollary 2.2.2] we have that $w_0 = v_0 = \dots = v_s$, $\kappa_{w_0} = \kappa_v(\overline{Y}_0)$ and $\Gamma_{w_0} = \mathbb{Z}$. Thus $w_0(\alpha) \in 2\mathbb{Z}$, and we may consider some $\beta \in \mathcal{O}_v^\times$ such that $\alpha \in \beta K^{\times 2}$. Let $j \in J$. We have $Y_j^2 + 1 \in \mathcal{O}_{w_0}^\times$. We set $c_j = b_0 b_j^{-1}$, $d_j = (a_0 - a_j) b_j^{-1}$. Since $Y_j = c_j Y_0 + d_j$, we have $Y_j^2 + 1 = c_j^2 Y_0^2 + 2c_j d_j Y_0 + d_j^2 + 1$, and hence $f \in h(Y_0) \cdot K^{\times 2}$, where

$$h(Y_0) = \beta(Y_0^2 + 1) \cdots (c_s^2 Y_0^2 + 2d_s c_s Y_0 + d_s^2 + 1) \prod_{i=s+1}^g Z_i.$$

Therefore $F = K(Y_0)(\sqrt{h(Y_0)})$. Since v is henselian and $\beta \notin -K^{\times 2}$, we have $\overline{\beta} \notin -\kappa_v^{\times 2}$. Finally, since $h \in \mathcal{O}_{w_0}^\times$, we have that $h \in fK(Y_0)^{\times 2} \cap \mathcal{O}_{w_0}^\times$, and we can conclude that $\kappa_w = \kappa_v(\overline{Y}_0) \left(\sqrt{\overline{h(Y_0)}} \right)$ is a real field, where $\overline{h} \in \kappa_v[\overline{Y}_0]$ is a polynomial with all roots in $\kappa_v(\sqrt{-1}) \setminus \kappa_v$ by Proposition [4, Lemma 2.2] (case (c)). \square

4.8. Corollary. *Let $n \in \mathbb{N}^+$. Assume that K carries a henselian \mathbb{Z}^n -valuation v such that κ_v is hereditarily euclidean. Let $f \in K[X]$ be a nonconstant square-free polynomial with all roots in $K(\sqrt{-1}) \setminus K$. We set $F = K(X)(\sqrt{f})$. Assume that F is real. Then $p(F) = 2$.*

Proof. We prove the statement by induction on n . Note that by Corollary 3.7 we have that $p(F) = 2$ if and only if $\mathcal{X}^1(F/v)$ is empty. Assume $n = 1$. Let w be a residually transcendental extension of v . It follows by Proposition 4.7 that either $s(\kappa_w) = 1$ or κ_w is real. Hence $\mathcal{X}^1(F/v) = \emptyset$ and $p(F) = 2$. Assume now that $n > 1$. We show that $\mathcal{X}^1(F/v)$ is empty. Let $v_1 = \pi_1 \circ v$. Let \bar{v} be the residual valuation of v modulo v_1 . Then \bar{v} is a henselian \mathbb{Z}^{n-1} -valuation on κ_{v_1} such that $\kappa_v = \kappa_{\bar{v}}$. It follows by induction hypothesis and by Proposition 4.7 that all the residually transcendental extensions w of v_1 to F satisfy $p'(\kappa_w) = 2$. If we had some $\mathcal{O} \in \mathcal{X}^1(F/v)$, then we would obtain that the residue field of the rank-one coarsening \mathcal{O}' of \mathcal{O} would have $p'(\kappa_{\mathcal{O}'}) > 2$ by Proposition 3.2, which is a contradiction. Therefore $\mathcal{X}^1(F/v) = \emptyset$, whereby $p(F) = 2$ by Theorem 3.6. \square

Note that, if $f \in K[X]$ is assumed to be monic in Corollary 4.8, then $F/K(X)$ is a totally positive quadratic extension and hence the above result follows by [5, Corollary 4.10].

4.9. Corollary. *Let $g \in \mathbb{N}$. Assume that K carries a henselian \mathbb{Z} -valuation v with hereditarily euclidean residue field. Let F/K be a hyperelliptic function field of genus g . Let $f \in K[X]$ be a square-free polynomial such that $F = K(X)(\sqrt{f})$. If $|G(F)| = 2^{g+1}$, then F is nonreal.*

Proof. It follows by Proposition 4.1 and by Lemma 4.2 that we may choose irreducible polynomials $q_i = (X - a_i)^2 + b_i^2$, with $a_i, b_i \in \mathcal{O}_v$, and $\alpha \in K^\times$ such that $f = \alpha \cdot q_0 \cdots q_g$. For $i \in \{0, \dots, g\}$ let $Y_i := b_i^{-1}(X - a_i)$. If $\alpha \notin -K^{\times 2}$, then $p(F) = 2$, by Corollary 4.8. Therefore, under the assumption that $|G(F)| = 2^{g+1}$, we have that $\alpha \in -K^{\times 2}$. Since $\mathbf{S}_2(F)$ is a group, there exists $\eta_1, \eta_2 \in K(X)$ such that $F = K(X)(\sqrt{-(\eta_1^2 + \eta_2^2)})$, that is, F is nonreal. \square

4.10. Question. *Let $n, g \in \mathbb{N}$. Assume that K carries a henselian \mathbb{Z}^n -valuation with hereditarily euclidean residue field. Let F/K be a function field in one variable of genus $g \in \mathbb{N}$. Is $|G(F)| \leq 2^{n(g+1)}$ for every function field in one variable F/K of genus g ?*

Acknowledgments. I thank Marco Zaninelli, Parul Gupta and Nicolas Daans for useful conversation on these topics. I thank my thesis advisors David Grimm and Karim Johannes Becher.

This work was supported by CONICYT-PFCHA/Doctorado Nacional/2017-folio 21170477, and by Universidad de Santiago de Chile (Proyecto DICYT, código 041933G), by the FWO Odysseus Programme (project G0E6114N), and by the Bijzonder Onderzoeksfonds (BOF), Universiteit Antwerpen, (project BOF-DOCPRO4, 2533).

REFERENCES

- [1] K.J. Becher, N. Daans, D. Grimm, G. Manzano-Flores, M. Zaninelli. *The Pythagoras number of a rational function field in two variables*. <https://arxiv.org/abs/2302.11425> (2023).

- [2] K.J. Becher, D. Grimm, *Nonsplit conics in the reduction of an arithmetic curve*, <https://arxiv.org/abs/2005.11855>.
- [3] K.J. Becher, D. Grimm, and J. Van Geel, *Sums of squares in algebraic function fields over a complete discretely valued field*, Pacific J. of Math **267** (2014), 257–276.
- [4] K.J. Becher, P. Gupta, *Ruled residue theorem for function fields of conics*. Journal of Pure and Applied Algebra, **225(06)** (2021) 106638.
- [5] K.J. Becher and J. Van Geel, *Sums of squares in function fields of hyperelliptic curves*, Mathematische Zeitschrift, 261 (4): (2009) 829 - 844.
- [6] Becker, E. *Hereditarily pythagorean fields and orderings of higher level*. Monografias de Matemática Vol. 29. Instituto de matematica pura e aplicada, Rio de Janeiro (1978).
- [7] L. Bröcker, *Characterization of fans and hereditarily pythagorean fields*. Math. Z. **152**, (1976), 149-163.
- [8] A.J. Engler, A. Prestel, *Valued fields*, Springer-Verlag, 2005.
- [9] D. Grimm, *On an isotropy criterion for quadratic forms over function fields of curves over non-dyadic complete discrete valuation rings*. Algebra, logic and number theory, 95–103, Banach Center Publ., 108, Polish Acad. Sci. Inst. Math., Warsaw, (2016).
- [10] D. Harbater, J. Hartmann, and D. Krashen, *Applications of patching to quadratic forms and central simple algebras*, Inventiones Mathematicae **178** (2009), 231-269.
- [11] N. Jacobson, *Lectures in Abstract Algebra, III. Theory of Fields and Galois Theory*. Springer-Verlag, Berlin.
- [12] H. Koch. *Number theory: Algebraic numbers and functions*. Graduate Studies in Mathematics, 2000.
- [13] T.Y. Lam, *Introduction to quadratic forms over fields*, American Mathematical Society, 2005.
- [14] Q. Liu, *Algebraic Geometry and Arithmetic curves*, Oxford Graduate Texts in Mathematics, Oxford University Press, Oxford, 2002.
- [15] J. Ohm, *The ruled residue theorem for simple transcendental extensions of valued fields*, Proceedings of the American Mathematical Society **89** (1983), no. 1, 16–18.
- [16] P. Ribenboim, *Le théorème d'approximation pour les valuations de krull*. Math. Z. **68**, 1-18 (1957).
- [17] S.V. Tikhonov, J. Van Geel, V.I Yanchevskii, *Pythagoras number of function fields of hyperelliptic curves with good reduction*, Manuscripta Math. **119** (2006), 305-322.
- [18] S.V Tikhonov, V.I Yanchevskii, *Pythagoras number of function fields of conics over hereditarily pythagorean fields*. Dokl. Nats. Akad. Nauk Belarusi **47**, 5-8 (2003).
- [19] S. Warner, *Topological fields*. Mathematics studies **157**, North Holland, Amsterdam (1989).
- [20] E. Witt, *Zerlegung reeller algebraischer Funktionen in Quadrate. Schiefkörper über reellem Funktionenkörper*. J. Reine Angew. math **171** (1934), 4-11.

UNIVERSIDAD DE SANTIAGO DE CHILE, FACULTAD DE CIENCIAS, AVENIDA LIBERTADOR
BERNARDO O'HIGGINS N° 3363, ESTACIÓN CENTRAL, SANTIAGO, CHILE.

Email address: gonzalo.manzano@usach.cl