

Closed k -Schur Katalan functions as K -homology Schubert representatives of the affine Grassmannian

Dedicated to the memory of Bumsig Kim

Takeshi Ikeda, Shinsuke Iwao, Satoshi Naito

July 5, 2022

Abstract

Recently, Blasiak-Morse-Seelinger introduced symmetric functions called Katalan functions, and proved that the K -theoretic k -Schur functions due to Lam-Schilling-Shimozono form a subfamily of the Katalan functions. They conjectured that another subfamily of Katalan functions called the closed k -Schur Katalan functions are identified with the Schubert structure sheaves in the K -homology of the affine Grassmannian. The main result is a proof of the conjecture.

We also study a K -theoretic Peterson isomorphism that Ikeda, Iwao, and Maeno constructed, in a non-geometric manner, based on the unipotent solution of the relativistic Toda lattice of Ruijsenaars. We prove that the map sends a Schubert class of the quantum K -theory ring of the flag variety to a closed K - k -Schur Katalan function up to an explicit factor related to a translation element with respect to an anti-dominant coroot. In fact, we prove the above map coincides with a map whose existence was conjectured by Lam, Li, Mihalcea, Shimozono, and proved by Kato, and more recently by Chow and Leung.

1 Introduction

The study of K -theoretic Schubert calculus attracts much attention in the last few decades. In this paper, we focus on the K -theory version of the “quantum equals affine” phenomenon, which originally comes from an unpublished result by Peterson in 1997 for the case of (co)homology, and has been developed by many authors. See a text book by Lam, Lapointe, Morse, Schilling, Shimozono, and Zabrocki [17] on these topics.

Lam, Schilling, and Shimozono [19] identifies the K -homology $K_*(\text{Gr})$ of the affine Grassmannian $\text{Gr} = G(\mathbb{C}((t)))/G(\mathbb{C}[[t]])$ of $G = SL_{k+1}$, with a subring $\Lambda_{(k)} = \mathbb{C}[h_1, \dots, h_k]$ of the ring of symmetric functions, where h_i is the i -th complete symmetric function. In particular, the K -theoretic k -Schur functions $g_\lambda^{(k)}$, the K - k -Schur functions for short, were introduced in [19], which form a family of inhomogeneous symmetric functions in $\Lambda_{(k)}$. It was proven that the K -theoretic k -Schur functions, indexed by the partitions λ with $\lambda_1 \leq k$, are identified with a distinguished basis of $K_*(\text{Gr})$. These functions form a basis of $\Lambda_{(k)}$, and are indexed by the partitions λ with $\lambda_1 \leq k$.

Although the K - k -Schur functions can be characterized by a Pieri type formula (see Definition 2.6), there was no explicit combinatorial formula until recently. Blasiak, Morse, and Seelinger [4] proved a raising operator formula for the K - k -Schur functions. In fact, they introduced a family of inhomogeneous symmetric functions called the K -theoretic Catalan functions, the Katalan functions for short, and proved that the K -theoretic k -Schur functions form a subfamily of the Katalan functions.

The Catalan functions $K(\Psi; M; \gamma)$ are indexed by triples (Ψ, M, γ) , where Ψ is an upper order ideal in the set $\Delta_\ell^+ := \{\varepsilon_i - \varepsilon_j \mid 1 \leq i < j \leq \ell\}$ of positive roots of type $A_{\ell-1}$, M is a multiset supported on $\{1, \dots, \ell\}$, and $\gamma \in \mathbb{Z}^\ell$. For any root ideal $\mathcal{L} \subset \Delta_\ell^+$, set $L(\mathcal{L}) := \bigsqcup_{(i,j) \in \mathcal{L}} \{j\}$, where (i, j) is a short-hand notation for $\varepsilon_i - \varepsilon_j$. Let \mathcal{P}^k denote the set of all partitions λ such that $\lambda_1 \leq k$. For $\lambda \in \mathcal{P}^k$, let $g_\lambda^{(k)}$ denote the corresponding K - k -Schur functions (see § 2.5 for the definition). In [4], it was proved

$$g_\lambda^{(k)} = K(\Delta^k(\lambda); L(\Delta^{k+1}(\lambda)); \lambda), \quad (1.1)$$

where

$$\Delta^k(\lambda) := \{(i, j) \in \Delta_\ell^+ \mid \lambda_i + j - i > k\},$$

and $\ell \geq \ell(\lambda)$, the length of λ . As one of the consequences of (1.1), a long-standing conjecture by Morse [29] was verified in [4]: for $\lambda \in \mathcal{P}^k$,

$$\lambda_1 + \ell(\lambda) \leq k + 1 \implies g_\lambda^{(k)} = g_\lambda, \quad (1.2)$$

where g_λ is the *dual stable Grothendieck polynomial*.

In [4], they introduced another subfamily of Catalan functions, called the *closed k -Schur Catalan functions*, defined for $\lambda \in \mathcal{P}^k$ by

$$\tilde{g}_\lambda^{(k)} := K(\Delta^k(\lambda); L(\Delta^k(\lambda)); \lambda).$$

It is conjectured that the closed k -Schur Catalan function is related to the function

$$\tilde{g}_\lambda^{(k)} := \sum_{\substack{\mu \leq \lambda \\ \mu \in \mathcal{P}^k}} g_\mu^{(k)}, \quad (1.3)$$

where $\lambda, \mu \in \mathcal{P}^k$ and \leq denote the order on \mathcal{P}^k induced by the Bruhat order on the affine symmetric group \tilde{S}_{k+1} (see § 2.1 for details). We call $\tilde{g}_\lambda^{(k)}$ the *closed K - k -Schur function*. These functions are essential in the K -homology Schubert calculus because they are identified with the class of the Schubert structure sheaf $\mathcal{O}_\lambda^{\text{Gr}}$ for the affine Grassmannian, whereas $g_\lambda^{(k)}$ is identified with the class of ideal sheaf of the boundary of the Schubert variety; see [18, Theorem 1] and [19, Theorem 5.4 and 7.17(1)]. It should be noted that Takigiku [34] proved a Pieri type formula for $\tilde{g}_\lambda^{(k)}$. Another important result in [34] is called the *k -rectangle factorization formula*. For $1 \leq i \leq k$, define

$$R_i = \overbrace{(i, \dots, i)}^{k+1-i}.$$

Takigiku showed

$$\tilde{g}_{R_i} \cdot \tilde{g}_\lambda^{(k)} = \tilde{g}_{\lambda \cup R_i}^{(k)}, \quad (1.4)$$

where $\lambda \cup R_i$ is the partition made by combining the parts of λ and those of R_i and then sorting. This formula is natural from geometric point of view (see [18]), and play an important role in our constructions.

Let σ be a ring automorphism of Λ given by $\sigma(h_i) = \sum_{j=0}^i h_j$ ($i \geq 1$) with $h_0 = 1$. We can now state the main result of this paper confirming a conjecture [4, Conjecture 2.12 (a)] by Blasiak, Morse, and Seelinger, which enable us to have a better explicit knowledge of the structure sheaf $\mathcal{O}_\lambda^{\text{Gr}}$.

Theorem 1.1. For $\lambda \in \mathcal{P}^k$, we have

$$\tilde{g}_\lambda^{(k)} = \sigma(\tilde{g}_\lambda^{(k)}).$$

Knowing Takigiku's result (1.4), an immediate consequence is the following k -rectangle factorization formula for $\tilde{g}_\lambda^{(k)}$, which was first proved by Seelinger [32] by a more direct method involving generalized Catalan functions defined for any subset of Δ_ℓ^+ .

Corollary 1.2 (cf. [4] Conjecture 2.12 (f)). For $1 \leq i \leq k$, we have

$$\tilde{\mathfrak{g}}_{R_i}^{(k)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)} = \tilde{\mathfrak{g}}_{\lambda \cup R_i}^{(k)}.$$

Another motivation of our study is to clarify the connection between a map called a *K-theoretic Peterson isomorphism* constructed by Ikeda, Iwao, and Meano [11], and a map whose existence was conjectured by Lam, Li, Mihalcea, and Shimozono [18]; this conjecture was proved by Kato [13], and more recently by Chow and Leung [6] using different methods.

Let $QK(G/B)$ be the (small) quantum K -theory ring of the flag variety $G/B = \mathrm{SL}_{k+1}(\mathbb{C})/B$, a deformation of the Grothendieck ring of coherent sheaves on G/B studied by Givental and Lee [9] (see also the finiteness result [1] by Anderson, Chen, Tseng, and Iritani). This is a commutative associative algebra over the polynomial ring $\mathbb{C}[Q] := \mathbb{C}[Q_1, \dots, Q_k]$ in the variables Q_i 's called the Novikov variables. For $w \in S_{k+1}$, the symmetric group of degree $k+1$, the Schubert variety Ω^w in G/B is defined to be $\overline{B_- w B / B}$, where B_- is the opposite Borel subgroup. Let $\mathcal{O}_{G/B}^w$ denote the (class of) structure sheaf of Ω^w . As a $\mathbb{C}[Q]$ -module, $QK(G/B)$ has a basis consisting of $\mathcal{O}_{G/B}^w$ ($w \in S_{k+1}$). Let $K_*(\mathrm{Gr})$ be the K -homology group of the affine Grassmannian Gr , which has a ring structure by the Pontryagin product. The ring has a basis consisting of the (class of) Schubert structure sheaves $\mathcal{O}_\lambda^{\mathrm{Gr}}$ indexed by the k -bounded partitions (see [19] for more detailed description).

We represent $QK(G/B)$ as a quotient ring the the polynomial ring $\mathbb{C}[Q][z_1, \dots, z_{k+1}]$ (see § 4.1 for details). According to Lam, Schilling, and Shimozono [19], we can identify $K_*(\mathrm{Gr})$ with $\Lambda_{(k)}$. More precisely, $\mathcal{O}_\lambda^{\mathrm{Gr}}$ is identified with $\sigma^{-1}(\tilde{g}_\lambda^{(k)})$ in our convention (see § 4.3 for details). For any partition λ , let $g_\lambda \in \Lambda$ be the dual stable Grothendieck polynomial (see § 2.4). We also set $\tilde{g}_\lambda := \sum_{\mu \subset \lambda} g_\mu$ following Takigiku [35]. We define a localization $K_*(\mathrm{Gr})_{\mathrm{loc}}$ of $K_*(\mathrm{Gr}) \cong \Lambda_{(k)}$ as $\Lambda_{(k)}[g_{R_i}^{-1}, \tilde{g}_{R_i}^{-1} \ (1 \leq i \leq k)]$. We let $QK(G/B)_{\mathrm{loc}}$ be the localization of $QK(G/B)$ by the multiplicative set generated by Q_1, \dots, Q_k . There is a ring isomorphism [11]

$$\Phi_{k+1} : QK(G/B)_{\mathrm{loc}} \rightarrow K_*(\mathrm{Gr})_{\mathrm{loc}}$$

given by

$$\Phi_{k+1}(z_i) = \frac{\tau_i \tau_{i-1}^+}{\tau_i^+ \tau_{i-1}}, \quad \Phi_{k+1}(Q_i) = \frac{\tau_{i-1} \tau_{i+1}}{\tau_i^2},$$

where

$$\tau_i = g_{R_i}, \quad \tau_i^+ = \tilde{g}_{R_i}$$

for $1 \leq i \leq k$ and $\tau_0 = \tau_{k+1} = \tau_0^+ = \tau_{k+1}^+ = 1$. It should be emphasized that the nature of the construction of Φ_{k+1} is combinatorial rather than geometric, in the sense that it heavily depends on explicit presentations of the rings involved. In fact, they used a non-linear differential equation called the *relativistic Toda lattice* introduced by Ruijsenaars. The map Φ_{k+1} arises as a solution of the relativistic Toda lattice equation with its Lax matrix unipotent. For the affine side, the functions h_i ($1 \leq i \leq k$) can be thought of as the coordinates of a certain abelian centralizer subgroup in $PSL_{k+1}(\mathbb{C})$.

There is a map $\theta_k : S_{k+1} \rightarrow \mathcal{P}^k$ explicitly described by Lam and Shimozono [22]. For $w \in S_{k+1}$, define $\mathrm{Des}(w) = \{1 \leq i \leq k \mid w(i) > w(i+1)\}$. Next result is a generalization of Theorem 1 in [22] by Lam and Shimozono for the (co)homology case to K -theoretic setting.

Theorem 1.3. For $w \in S_{k+1}$, we have

$$\Phi_{k+1}(\mathcal{O}_{G/B}^w) = \frac{\sigma^{-1}(\tilde{g}_{\theta_k(w)})}{\prod_{i \in \mathrm{Des}(w)} \tau_i}.$$

As noted above $\sigma^{-1}(\tilde{g}_{\theta_k(w)}) = \tilde{\mathfrak{g}}_{\theta_k(w)}^{(k)}$ is identified with the structure sheaf $\mathcal{O}_{\theta_k(w)}^{\mathrm{Gr}}$, while τ_i is identified with the Schubert structure sheaf associated to an anti-dominant translation element in \hat{S}_{k+1} (see § 4.3 for a more precise statement). Note also that for $w \in S_{k+1}$ we

can take the *quantum Grothendieck polynomials* $\mathfrak{G}_w^Q \in \mathbb{C}[Q][z_1, \dots, z_k]$ of Lenart and Maeno [25] (with the change of variables $x_i = 1 - z_i$) as a representative for $\mathcal{O}_{G/B}^w$ (see § 4.1). The statement of Theorem 1.3 is a refinement of Conjecture 1.8 in [11]. See also Conjecture 2.12 (a), (b) in [4].

Future works and related results

Now we are able to say that the unipotent solution of the relativistic Toda lattice actually gives the canonical K -theoretic Peterson isomorphism, in type A . A natural question is to generalize this fact to any semisimple algebraic group G . Furthermore, we would like to describe the K -theoretic Peterson isomorphisms in various types at the level of concrete polynomial representatives of Schubert classes. This leads to a lot of interesting combinatorial problems related to geometry of K -theoretic Gromov-Witten theory. We know from Kim [14] that the quantum cohomology ring of G/B is identified with the quotient ring of a polynomial ring by the ideal generated by the conserved quantities of the Toda lattice for the Langlands-dual group G^\vee with the nilpotent initial condition. For our purpose, one of the central tasks is to obtain an analogue of Kim’s result in the context of quantum K -theory ring for G/B . Another possible clue in this direction is the work of Bezrukavnikov, Finkelberg, and Mirković [2] that studied a connection between the K -homology of affine Grassmannian and certain generalized Toda lattice equations, however, we still do not understand how this work fits into the framework of the K -Peterson isomorphism.

For the affine Grassmannian side, Lam, Schilling, and Shimozono [20] defined k -Schur functions for the symplectic groups. It is remarkable that Seelinger [32] made a conjecture that the symplectic k -Schur function of [20] can be expressed by a raising operator formula. Pon [31] studied the case of the orthogonal group. In particular, he gave a definition of “ k -Schur functions” as the Schubert representatives for homology of the affine Grassmannian in type B , and established the Pieri rule for type B and D . Further combinatorial researches on these functions are needed to explore the issue of giving explicit K -theoretic Peterson isomorphisms.

Even in type A , there are a lot of things to do. We have a Chevalley type formula for $QK(G/B)$ ([26], [27], [25], [30]) which describes the multiplication by $\mathcal{O}_{G/B}^{s_i}$ ($1 \leq i \leq k$), where s_i is the transposition $(i, i + 1)$. The image of $\mathcal{O}_{G/B}^{s_i}$ ($1 \leq i \leq k$) under Φ_{k+1} is $\sigma^{-1}(\tilde{g}_{R_i^*}^{(k)})/\tau_i$ where R_i^* is the partition obtained from R_i by removing the unique corner box. We expect a good combinatorial formula for $\tilde{g}_{R_i^*}^{(k)} \cdot \tilde{g}_\lambda^{(k)}$. Although we know a dictionary between quantum and affine Schubert classes, the problem of translating the Chevalley formula into affine counterpart seems not to be so simple. For the homology case, the analogous issue was pursued by Dalal and Morse ([8, Conjecture 39]). Another basic question is to find the formula for the quantum side corresponding to $\tilde{g}_{(i)}^{(k)} \cdot \tilde{g}_\lambda^{(k)}$ and $\tilde{g}_{(1^i)}^{(k)} \cdot \tilde{g}_\lambda^{(k)}$ ($1 < i < k - 1$). Furthermore, there is a conjecture by Lenart and Maeno of a Pieri type formula for \mathfrak{G}_w^Q ([25, Conjecture 6.7]). We hope the “quantum equals affine” phenomenon in K -theory would shed light on these questions. Also, it is natural to extend Theorem 1.3 torus equivariantly. In fact, Lam and Shimozono [21] proved the equivariant (co)homology version of the explicit Peterson isomorphism in [22]. It was shown that the double k -Schur functions of Lam-Shimozono ([23]), and the quantum double Schubert polynomials by Kirillov and Maeno [16] and by Ciocan-Fontanine and Fulton [7], can be obtained from each other by the map. There are arguments in [21, § 4] on the centralizer family for SL_{k+1} in connection with Peterson’s j -map, which would be useful in future studies.

Organization

In Sections 2, we present some basic definitions and preliminary facts. In Section 3, we prove Theorem 1.1. In Section 4.1, we discuss the K -theoretic Peterson isomorphism. In

Appendix A, we provide some results related to the parabolic quotient of a Coxeter group, which are used in the proof of Theorem 1.1 In Appendix B, we give a proof of a result (Lemma B.3) on a Grassmannian permutation. In Appendix C, we record the dual Pieri rule for the closed K - k -Schur functions.

Acknowledgements

We thank Hiroshi Naruse, Daisuke Sagaki, and Mark Shimozono for valuable discussions; in particular, S. N. thanks Daisuke Sagaki for helpful discussions about the material in Appendix A. We thank Cristian Lenart for valuable comments. T. I. was partly supported by JSPS Grant-in-Aid for Scientific Research 20H00119, 20K03571, 18K03261, 17H02838. S. N. was partly supported by JSPS Grant-in-Aid for Scientific Research (C) 21K03198. S. I. was partly supported by JSPS Grant-in-Aid for Scientific Research (C) 19K03605.

2 Basic definitions

Let k be a positive integer. In this section, we fix the notation and explain definitions and properties of basic notions needed to understand Theorem 1.1.

2.1 Affine symmetric groups

The *affine symmetric group* \tilde{S}_{k+1} is the group with generators $\{s_i \mid i \in I\}$ for $I = \{0, 1, \dots, k\}$ subject to the relations

$$s_i^2 = id, \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, \quad s_i s_j = s_j s_i \quad \text{for } i - j \neq 0, \pm 1,$$

with indices considered modulo $k+1$. The *length* $\ell(w)$ of $w \in \tilde{S}_{k+1}$ is the minimum number m such that $w = s_{i_1} \cdots s_{i_m}$ for some $i_j \in I$; any such expression for w with $\ell(w)$ generators is said to be *reduced*. The set of *affine Grassmannian elements* \tilde{S}_{k+1}^0 is the minimal length coset representatives of S_{k+1} in \tilde{S}_{k+1} , where $S_{k+1} = \langle s_1, \dots, s_k \rangle$.

The *Bruhat order* (or *strong order*) on \tilde{S}_{k+1} is denoted by \leq . It can be described by a subword property (see [3, § 2]).

2.2 k -bounded partitions and affine Grassmannian elements

Let $\mathcal{P}_\ell^k := \{(\lambda_1, \dots, \lambda_\ell) \in \mathbb{Z}^\ell \mid k \geq \lambda_1 \geq \dots \geq \lambda_\ell \geq 0\}$ denote the set of partitions contained in the $\ell \times k$ rectangle and let \mathcal{P}^k be the set of partitions λ with $\lambda_1 \leq k$. For a partition λ , the *length* $\ell(\lambda)$ is the number of nonzero parts of λ . There is a bijection $\mathcal{P}^k \rightarrow \tilde{S}_{k+1}^0$ ($\lambda \mapsto x_\lambda$) due to Lapointe and Morse [24, Definition 45, Corollary 48] given by

$$x_\lambda := (s_{\lambda_\ell - \ell} \cdots s_{-\ell+1}) \cdots (s_{\lambda_2 - 2} \cdots s_{-1}) (s_{\lambda_1 - 1} \cdots s_0), \quad (2.1)$$

where $\ell = \ell(\lambda)$. To make the notation simple, we are omitting the dependency of x_λ on k .

Example 2.1. Let $k = 4, \lambda = (4, 3, 2)$. The corresponding affine Grassmannian element in \tilde{S}_5^0 is $x_\lambda = s_4 s_3 \cdot s_1 s_0 s_4 \cdot s_3 s_2 s_1 s_0$. This is obtained by reading the $(k+1)$ -residues in each row of λ , from right-to-left, starting with the bottom row to top.

$$\begin{array}{|c|c|c|c|} \hline 0 & 1 & 2 & 3 \\ \hline 4 & 0 & 1 & \\ \hline 3 & 4 & & \\ \hline \end{array}.$$

For $\lambda, \mu \in \mathcal{P}^k$, we denote $\lambda \leq_k \mu$ if $x_\lambda \leq x_\mu$ holds, where \leq is the Bruhat order on \tilde{S}_{k+1} . The following fundamental fact is included in the proof of [4, Proposition 2.16].

Lemma 2.2. Suppose $\lambda \in \mathcal{P}^k$ satisfies $\lambda_1 + \ell(\lambda) \leq k+1$. For $\mu \in \mathcal{P}^k$, $\mu \leq_k \lambda \iff \mu \subset \lambda$.

Remark 2.3. For $\lambda, \mu \in \mathcal{P}^k$, $\lambda \subset \mu$ implies $\lambda \leq_k \mu$. The inverse implication is not true in general. For example, in \tilde{S}_3 we have $(2, 2) \leq_k (2, 1, 1)$ because the reduced expression $x_{(2,1,1,1)} = s_0 s_1 s_2 s_1 s_0$ has a subexpression $s_0 s_2 s_1 s_0 = x_{(2,2)}$.

2.3 Cyclically increasing elements in \tilde{S}_{k+1}^0

Let A be a proper subset of $I = \{0, 1, \dots, k\}$. Set $|A| = r$. There is a sequence (i_1, \dots, i_r) consisting of the elements of A such that an index i never occurs somewhere to the left of an index $i + 1 \pmod{k+1}$. For any such sequence, the element $s_{i_1} \cdots s_{i_r} \in \tilde{S}_{k+1}$ only depends on the set of indices $A = \{i_1, \dots, i_r\}$ (see also Remark B.2). Let us denote the element by u_A . Such elements in \tilde{S}_{k+1} is called a *cyclically increasing element*. Similarly, for any $A \subsetneq I$, we can define the corresponding *cyclically decreasing element* denoted by d_A . We choose a sequence (i_1, \dots, i_r) such that an index i never occurs somewhere to the right of an index $i + 1 \pmod{k+1}$, and set $d_A := s_{i_1} \cdots s_{i_r}$.

Example 2.4. Let $k = 4$. Then for $A = \{0, 1, 3, 4\}$, $u_A = s_3 s_4 s_0 s_1$, $d_A = s_1 s_0 s_4 s_3$, and for $A = \{0, 2, 4\}$, $u_A = s_4 s_0 s_2 = s_4 s_2 s_0 = s_2 s_4 s_0$, $d_A = s_2 s_0 s_4 = s_0 s_2 s_4 = s_0 s_4 s_2$.

2.4 Dual stable Grothendieck polynomials

We work in the ring $\Lambda = \mathbb{Z}[e_1, e_2, \dots] = \mathbb{Z}[h_1, h_2, \dots]$ of symmetric functions in infinitely many variables (x_1, x_2, \dots) , where $e_r = \sum_{i_1 < \dots < i_r} x_{i_1} \cdots x_{i_r}$ and $h_r = \sum_{i_1 \leq \dots \leq i_r} x_{i_1} \cdots x_{i_r}$. Set $h_0 = e_0 = 1$ and $h_r = 0$ for $r < 0$ by convention.

Let σ be the ring automorphism of Λ defined by $\sigma(h_i) = \tilde{h}_i$ ($i \geq 1$), where we set $\tilde{h}_i = \sum_{j=0}^i h_j$. Note that σ^{-1} sends h_i to $h_i - h_{i-1}$ for $i \geq 1$. For $i, m \in \mathbb{Z}$, define

$$h_i^{(m)} := \sigma^m(h_i) = \sum_{j=0}^i \binom{m+j-1}{j} h_{i-j},$$

where $\binom{n}{i} = n(n-1)\cdots(n-i+1)/i!$ for $n \in \mathbb{Z}$, $i \geq 1$ and $\binom{n}{0} = 1$ for $n \in \mathbb{Z}$. For $\gamma \in \mathbb{Z}^\ell$, define

$$g_\gamma := \det(h_{\gamma_i+j-i}^{(i-j)})_{1 \leq i, j \leq \ell}. \quad (2.2)$$

When λ is a partition, g_λ is the *dual stable Grothendieck polynomials*. The defining formula for g_γ in [4] is $\det(h_{\gamma_i+j-i}^{(i-1)})_{1 \leq i, j \leq \ell}$, however, by some column operations, we see their definition coincide with (2.2). Note that $h_i^{(m)}$ is denoted by $k_i^{(m)}$ in [4].

The following result is fundamental and used throughout the paper.

Proposition 2.5 (Takiguku [35]). For any partition λ , set $\tilde{g}_\lambda := \sum_{\mu \subset \lambda} g_\mu$. Then we have

$$\sigma(g_\lambda) = \tilde{g}_\lambda. \quad (2.3)$$

2.5 K - k -Schur functions

For $i \in I$, set for $w \in \tilde{S}_{k+1}$,

$$s_i * w = \begin{cases} s_i w & (s_i w > w) \\ w & (s_i w < w) \end{cases}. \quad (2.4)$$

If we denote $\phi_i : \tilde{S}_{k+1} \rightarrow \tilde{S}_{k+1}$ ($w \mapsto s_i * w$), then $\phi_i^2 = \phi_i$, $\phi_i \phi_{i+1} \phi_i = \phi_{i+1} \phi_i \phi_{i+1}$ (see the proof of [33, Proposition 2.1]). So for $v \in \tilde{S}_{k+1}$, we can define

$$v * w = s_{i_1} * (s_{i_2} * \cdots (s_{i_r} * w) \cdots) \quad (w \in \tilde{S}_{k+1})$$

where $v = s_{i_1} \cdots s_{i_r}$ is an arbitrary reduced expression.

For $x \in \tilde{S}_{k+1}^0$, we write $g_x^{(k)}$ for $g_\lambda^{(k)}$ with $x = x_\lambda$, $\lambda \in \mathcal{P}^k$.

Definition 2.6. The K - k -Schur functions $\{g_\lambda^{(k)}\}_{\lambda \in \mathcal{P}^k}$ are the family of elements of $\Lambda_{(k)}$ such that $g_\emptyset^{(k)} = 1$ and

$$h_r \cdot g_\lambda^{(k)} = \sum_{\substack{A \subset I, |A|=r \\ d_A * x_\lambda \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(d_A * x_\lambda) + \ell(\lambda)} g_{d_A * x_\lambda}^{(k)} \quad (2.5)$$

from $\lambda \in \mathcal{P}^k$ and $1 \leq r \leq k$.

Example 2.7. Let $k = 2$, $\lambda = (1, 1, 1)$, and $r = 2$. Then $x_\lambda = s_1 s_2 s_0$. There are three A 's: $\{1, 0\}$, $\{0, 2\}$, $\{1, 2\}$. We compute

$$\begin{aligned} d_{\{1,0\}} * x_\lambda &= (s_1 s_0) * (s_1 s_2 s_0) = s_1 s_0 s_1 s_2 s_0 = s_0 s_1 s_0 s_2 s_0 = s_0 s_1 s_2 s_0 s_2 \notin \tilde{S}_3^0, \\ d_{\{0,2\}} * x_\lambda &= (s_0 s_2) * (s_1 s_2 s_0) = s_0 s_2 s_1 s_2 s_0 = x_{(2,1,1,1)} \in \tilde{S}_3^0, \\ d_{\{1,2\}} * x_\lambda &= (s_2 s_1) * (s_1 s_2 s_0) = s_2 s_1 s_2 s_0 = x_{(2,1,1)} \in \tilde{S}_3^0, \end{aligned}$$

so

$$h_2 \cdot g_{(1,1,1)}^{(2)} = g_{(2,1,1,1)}^{(2)} - g_{(2,1,1)}^{(2)}.$$

It is known that $\{g_\lambda^{(k)}\}_{\lambda \in \mathcal{P}^k}$ is a basis of $\Lambda_{(k)}$ ([19]).

Proposition 2.8 ([19]). For $1 \leq r \leq k$, $g_{(r)}^{(k)} = h_r$.

Recall that we set $\tilde{g}_\lambda := \sum_{\mu \subset \lambda} g_\mu$ for any partition λ .

Proposition 2.9. For $\lambda \in \mathcal{P}^k$ such that $\lambda_1 + \ell(\lambda) \leq k + 1$,

$$\tilde{g}_\lambda^{(k)} = \tilde{g}_\lambda.$$

Proof. From Lemma 2.2, and (1.2), we have

$$\tilde{g}_\lambda^{(k)} = \sum_{\substack{\mu \leq \lambda \\ \bar{k}}} g_\mu^{(k)} = \sum_{\mu \subset \lambda} g_\mu^{(k)} = \sum_{\mu \subset \lambda} g_\mu = \tilde{g}_\lambda.$$

□

2.6 Catalan functions

Fix a positive integer ℓ . Let $\{\varepsilon_1, \dots, \varepsilon_\ell\}$ be the standard basis of \mathbb{Z}^ℓ . By a *positive root* β , we mean an element of the form $\varepsilon_i - \varepsilon_j \in \mathbb{Z}^\ell$ with $1 \leq i < j \leq \ell$, which is also denoted by (i, j) . The set of all positive roots is denoted by Δ_ℓ^+ . Although this is considered to be the set of positive roots of type $A_{\ell-1}$, we use this notation Δ_ℓ^+ according to [4] rather than $\Delta_{\ell-1}^+$.

A natural partial order \leq on Δ_ℓ^+ is defined by $\alpha \leq \beta$ if $\beta - \alpha$ is a linear combination of positive roots with coefficients in $\mathbb{Z}_{\geq 0}$. An upper order ideal Ψ of Δ_ℓ^+ is called a *root ideal*.

Given a root ideal $\Psi \subset \Delta_\ell^+$, a multiset M supported on $\{1, \dots, \ell\}$, and $\gamma \in \mathbb{Z}^\ell$, we say (Ψ, M, γ) is a *Katalan triple*. Let $m_M : \{1, \dots, \ell\} \rightarrow \mathbb{Z}_{\geq 0}$ denote the multiplicity function of M . Each Katalan triple (Ψ, M, γ) can be depicted by $\ell \times \ell$ grid of square boxes (labeled by matrix-style coordinates) with the boxes of Ψ shaded, $m_M(a)$ •'s in column a (assuming $m_M(a) < a$), and the entries of γ written along the diagonal boxes.

Example 2.10. Let $\Psi = \{(1, 2), (1, 3), (1, 4), (1, 5), (2, 4), (2, 5), (3, 5)\} \subset \Delta_5^+$, $M = \{3, 4, 5, 5\}$, and $\gamma = (3, 2, 0, 1, 0)$. The Katalan triple (Ψ, M, γ) is depicted by

3	•	•	•	•
2				•
	0			
		1		
				0

We define the *Katalan function* associated to the triple (Ψ, M, γ)

$$\begin{aligned} K(\Psi; M; \gamma) &:= \prod_{j \in M} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma, \\ k_\gamma &:= h_{\gamma_1}^{(0)} h_{\gamma_2}^{(1)} \cdots h_{\gamma_\ell}^{(\ell-1)}. \end{aligned}$$

Note that raising operators are not well-defined as linear transformations on Λ . They act on subscript γ of k_γ rather than the function k_γ . A rigorous formulation can be found in [4, § 3]. For any root ideal $\mathcal{L} \subset \Delta_\ell^+$, let

$$L(\mathcal{L}) = \bigsqcup_{(i,j) \in \mathcal{L}} \{j\}.$$

We also denote $K(\Psi; L(\mathcal{L}); \gamma)$ simply by $K(\Psi; \mathcal{L}; \gamma)$

In [4], Blasiak, Morse, and Seelinger introduced, for $\lambda \in \mathcal{P}_\ell^k$,

$$\begin{aligned} \mathfrak{g}_\lambda^{(k)} &:= K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda), \\ \tilde{\mathfrak{g}}_\lambda^{(k)} &:= K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda). \end{aligned}$$

$\mathfrak{g}_\lambda^{(k)}$ is called the *K-Schur Katalan function*, and $\tilde{\mathfrak{g}}_\lambda^{(k)}$ the *closed K-Schur Katalan function*. If we choose ℓ so that $\ell \geq \ell(\lambda)$, then $\mathfrak{g}_\lambda^{(k)}$ and $\tilde{\mathfrak{g}}_\lambda^{(k)}$ do not depend on ℓ ([4, Lemma 3.4, Remark 3.5]).

The following simplified formula is available.

Proposition 2.11. Let $\lambda \in \mathcal{P}_\ell^k$, then

$$\tilde{\mathfrak{g}}_\lambda^{(k)} = \prod_{(i,j) \in \Delta_\ell^+ \setminus \Delta^k(\lambda)} (1 - L_j)^{-1} (1 - R_{ij}) h_\lambda, \quad (2.6)$$

where $h_\lambda = h_{\lambda_1} \cdots h_{\lambda_\ell}$, and L_j, R_{ij} act on the suffix λ .

Proof. In the proof of [4, Proposition 2.3], it was shown that $\prod_{(i,j) \in \Delta_\ell^+} (1 - L_j) k_\gamma = h_\gamma$. (2.6) follows from this. \square

Example 2.12. For $k = 3$, $\lambda = (2, 1, 1)$, $\tilde{\mathfrak{g}}_\lambda^{(3)}$ is depicted by

2	1	1
1	1	1
1	1	1

. We have

$$\begin{aligned} \tilde{\mathfrak{g}}_{(2,1,1)}^{(3)} &= (1 - L_2)^{-1} (1 - L_3)^{-1} (1 - R_{12}) (1 - R_{23}) h_{211} \\ &= h_{211} + h_{201} - h_{301} - h_{220} + h_{310} \\ &= h_1^2 h_2 + h_1 h_2 + h_2^2 \\ &= h_2 (h_1^2 + h_1 - h_2). \end{aligned}$$

One of the main results in [4] is

$$\mathfrak{g}_\lambda^{(k)} = g_\lambda^{(k)}.$$

The main result of the present paper (Theorem 1.1) is

$$\tilde{\mathfrak{g}}_\lambda^{(k)} = \sigma^{-1}(\tilde{g}_\lambda^{(k)}). \quad (2.7)$$

A simple consequence of (2.7) is the following.

Corollary 2.13. If $\lambda_1 + \ell(\lambda) \leq k + 1$, then $\tilde{\mathfrak{g}}_\lambda^{(k)} = g_\lambda$.

Proof. By Lemma 2.9, and Proposition 2.5,

$$\tilde{\mathfrak{g}}_\lambda^{(k)} = \sigma^{-1}(\tilde{g}_\lambda^{(k)}) = \sigma^{-1}(g_\lambda) = g_\lambda.$$

\square

3 Proof of Theorem 1.1

Before we start the proof of Theorem 1.1, we gather some results on K - k -Schur functions and Katalan functions in the first three subsections. In § 3.1 we explain some results on the k -conjugation. In § 3.2, we correct some basic properties of Katalan functions used in § 3.5. In § 3.3, we introduce an action of the 0-Hecke algebras on $\Lambda_{(k)}$. The outline of the proof of Theorem 1.1 is given in § 3.4. The last section is devoted to the proof of a key lemma (Lemma 3.17). With the help of a general fact (Lemma 3.18) on the parabolic coset space of Coxeter group, we complete the proof of Theorem 1.1.

3.1 k -conjugation

There is an automorphism ω_k of \tilde{S}_{k+1} given by $\omega_k(s_i) = s_{-i} = s_{k+1-i}$ for $i \in I$. Note that ω_k fixes s_0 . In fact, ω_k is an automorphism of a Coxeter group. So it is easy to see

$$w \leq v \iff \omega_k(w) \leq \omega_k(v). \quad (3.1)$$

The left weak order \leq_L on \tilde{S}_{k+1} is defined by the covering relation

$$w \leq_L v \iff v = s_i w \text{ and } \ell(v) = \ell(w) + 1 \text{ for some } i \in I.$$

It is easy to see $w \leq_L v \iff \omega_k(w) \leq_L \omega_k(v)$, and therefore we have

$$w \leq_L v \iff \omega_k(w) \leq_L \omega_k(v). \quad (3.2)$$

For $A = \{i_1, \dots, i_m\} \subsetneq I$, we have

$$\omega_k(u_A) = d_{\bar{A}}, \quad \omega_k(d_A) = u_{\bar{A}},$$

where $\bar{A} := \{-i_1, \dots, -i_m\}$.

Definition 3.1. Let Ω be the ring homomorphism on Λ defined by

$$\Omega(h_i) = g_{(1^i)} \quad (i \geq 1). \quad (3.3)$$

Proposition 3.2. Ω is an involution on Λ and Ω commutes with σ .

Proof. A proof of the fact that Ω is an involution can be found in [29, § 8]. The commutativity follows from (2.3):

$$\Omega(\sigma(h_i)) = \Omega\left(\sum_{j=0}^i h_j\right) = \sum_{j=0}^i g_{(1^j)} = \sigma(g_{(1^i)}) = \sigma(\Omega(h_i)).$$

□

It is easy to see that ω_k preserves \tilde{S}_{k+1}^0 . Hence for $\lambda \in \mathcal{P}^k$, $\omega_k(x_\lambda) = x_\mu$ for some $\mu \in \mathcal{P}^k$. Then we define $\omega_k(\lambda) = \mu$. Explicit description of $\omega_k(\lambda)$, also denoted by λ^{ω_k} in [17], [29] is available (see [17, § 1.3]).

Theorem 3.3 ([29]). For $x \in \tilde{S}_{k+1}^0$, $\Omega(g_x^{(k)}) = g_{\omega_k(x)}^{(k)}$. Equivalently, for $\lambda \in \mathcal{P}^k$, $\Omega(g_\lambda^{(k)}) = g_{\omega_k(\lambda)}$.

Corollary 3.4. For $\lambda \in \mathcal{P}^k$, $\Omega(\tilde{g}_\lambda^{(k)}) = \tilde{g}_{\omega_k(\lambda)}^{(k)}$.

Proof. For $x \in \tilde{S}_{k+1}^0$, $\tilde{g}_x^{(k)} = \sum_{y \leq x} g_y^{(k)}$ where y are element in \tilde{S}_{k+1}^0 such that $y \leq x$ in the Bruhat order. From Theorem 3.3, and the fact (3.1), we have

$$\Omega(\tilde{g}_x^{(k)}) = \Omega\left(\sum_{y \leq x} g_y^{(k)}\right) = \sum_{y \leq x} \Omega(g_y^{(k)}) = \sum_{y \leq x} g_{\omega_k(y)}^{(k)} = \sum_{\omega_k(y) \leq \omega_k(x)} g_{\omega_k(y)}^{(k)} = \tilde{g}_{\omega_k(x)}^{(k)},$$

where the last equality holds because ω_k is an involution. Set $x = x_\lambda$. Then $\Omega(\tilde{g}_\lambda^{(k)}) = \tilde{g}_{\omega_k(\lambda)}^{(k)}$. □




Figure 1: Bounce graph of Ψ

3.2 Basic properties of Catalan functions

Let $\Psi \subset \Delta_\ell^+$ be a root ideal. A root $\alpha \in \Psi$ is a *removable root* of Ψ if $\Psi \setminus \alpha$ is a root ideal. A root $\beta \in \Delta_\ell^+$ is an *addable root* of Ψ if $\Psi \cup \alpha$ is a root ideal.

We define an oriented graph with $\{1, \dots, \ell\}$ as the vertex set and the oriented edges $j \rightarrow i$ if (i, j) is a removable root in Ψ (in [4] the bounce graph is not considered to be an oriented graph but here it is). An edge of the bounce graph of Ψ is called simply a *bounce edge* of Ψ . Let $p \in \{1, \dots, \ell\}$. If there is a bounce edge $p \rightarrow i$ of Ψ , then such i is unique by the construction, and we denote $i = \text{up}_\Psi(p)$.

Each connected component of the bounce graph of Ψ is called a *bounce path* of Ψ . Let p be a vertex of the bounce graph. The smallest element in the bounce path containing p is denoted by $\text{top}_\Psi(p)$. Note in [4], if there is no i such that $p \rightarrow i$, then $\text{top}_\Psi(p)$ is not defined, however in our definition $\text{top}_\Psi(p) = p$.

Example 3.5. For the root ideal Ψ in Figure 1, $\{1, 3\}$, $\{2, 5, 7\}$, $\{4, 6, 9\}$, $\{8\}$ are the bounce paths. We have, for example, $\text{up}_\Psi(3) = 1$, $\text{up}_\Psi(7) = 5$, $\text{top}_\Psi(7) = \text{top}_\Psi(5) = \text{top}_\Psi(2) = 2$.

Definition 3.6 (walls and ceilings). Let d be a positive integer. A root ideal Ψ is said to have a *wall* in rows $r, r+1, \dots, r+d$ if the rows $r, r+1, \dots, r+d$ of Ψ have the same length, a *ceiling* in columns $c, c+1, \dots, c+d$ if the columns $c, c+1, \dots, c+d$ of Ψ have same length.

Example 3.7. The root ideal Ψ in Figure 1 has a ceiling in columns 1, 2, in columns 3, 4, in columns 7, 8, and has a wall in rows 3, 4, in rows 7, 8, 9.

Let us begin with an obvious remark.

Lemma 3.8. Let Ψ be a root ideal of Δ_ℓ^+ , and $p \in \{2, \dots, \ell\}$. If $\text{top}_\Psi(p) = p$, i.e., there is no bounce edge starting from p , then Ψ has a ceiling in columns $p-1, p$.

The following lemmas are borrowed from [4, Lemma 4.4].

Lemma 3.9 (Adding or removing a root). Let (Ψ, M, γ) be a Catalan triple.

(i) for any addable root α of Ψ ,

$$K(\Psi; M; \gamma) = K(\Psi \cup \alpha; M; \gamma) - K(\Psi \cup \alpha; M; \gamma + \alpha).$$

(ii) for any removable root α of Ψ ,

$$K(\Psi; M; \gamma) = K(\Psi \setminus \alpha; M; \gamma) + K(\Psi; M; \gamma + \alpha).$$

Example 3.10. We apply Lemma 3.9 (i) to the following Catalan triple, with $\alpha = \varepsilon_2 - \varepsilon_3$:

$$\begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & 0 & & \\ \hline & & 1 & \\ \hline & & & 1 \\ \hline \end{array} = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & 0 & & \\ \hline & & 1 & \\ \hline & & & 1 \\ \hline \end{array} - \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & 1 & & \\ \hline & & 0 & \\ \hline & & & 1 \\ \hline \end{array}$$

We apply Lemma 3.9 (ii) to the following Katalan triple, with $\alpha = \varepsilon_3 - \varepsilon_4$:

$$\begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & 0 & & \\ \hline & & 1 & \\ \hline & & & 1 \\ \hline \end{array} = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & 0 & & \\ \hline & & 1 & \\ \hline & & & 1 \\ \hline \end{array} + \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & & 1 & \\ \hline & & & 2 \\ \hline & & & 0 \\ \hline \end{array}$$

Lemma 3.11 (Adding or removing a dot). Let (Ψ, M, γ) be a Katalan triple.

- (i) for any $j \in M$, $K(\Psi; M; \gamma) = K(\Psi; M \setminus \{j\}; \gamma) - K(\Psi; M \setminus \{j\}; \gamma - \varepsilon_j)$,
- (ii) for any $1 \leq j \leq \ell$, $K(\Psi; M; \gamma) = K(\Psi; M \sqcup \{j\}; \gamma) + K(\Psi; M; \gamma - \varepsilon_j)$.

The next lemma is [4, Lemma 3.3]

Lemma 3.12 (Alternating property). Let $\Psi \subset \Delta_\ell^+$ is a root ideal and, M a multiset on $\{1, \dots, \ell\}$. Suppose there is an index $1 \leq i \leq \ell - 1$ such that

- (a) Ψ has a ceiling in columns $i, i + 1$,
- (b) Ψ has a wall in rows $i, i + 1$,
- (c) $m_M(i + 1) = m_M(i) + 1$.

Then for any $\gamma \in \mathbb{Z}^\ell$,

$$K(\Psi; M; \gamma) = -K(\Psi; M; s_i \gamma - \varepsilon_i + \varepsilon_{i+1}). \quad (3.4)$$

In particular, if $\gamma_{i+1} = \gamma_i + 1$ holds, then

$$K(\Psi; M; \gamma) = 0. \quad (3.5)$$

3.3 The 0-Hecke algebra

The 0-Hecke algebra H_{k+1} is associative \mathbb{C} -algebra generated by $\{T_i \mid i \in I\}$ with the relations

$$T_i^2 = -T_i, \quad T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad T_i T_j = T_j T_i \quad \text{for } i - j \neq 0, \pm 1,$$

with indices considered modulo $k + 1$. For $w \in \tilde{S}_{k+1}$, define $T_w = T_{i_1} \cdots T_{i_m}$ for any reduced expression $w = s_{i_1} \cdots s_{i_m}$. The elements T_w ($w \in \tilde{S}_{k+1}$) form a basis of H_{k+1} .

We introduce a family of symmetric functions $\mathfrak{g}_\lambda^{(k)}$ ($\lambda \in \mathcal{P}^k$) so that $\tilde{\mathfrak{g}}_\lambda^{(k)} = \sum_{\mu \leq \lambda} \mathfrak{g}_\mu^{(k)}$ for all $\lambda \in \mathcal{P}^k$. Such functions uniquely exist since the transition matrix from $\{\tilde{\mathfrak{g}}_\lambda^{(k)}\}$ to $\{\mathfrak{g}_\lambda^{(k)}\}$ is upper uni-triangular.

Proposition 3.13. There is a left H_{k+1} -module structure on $\Lambda_{(k)}$ such that

$$T_i \cdot \mathfrak{g}_\lambda^{(k)} = \begin{cases} \mathfrak{g}_{s_i \lambda}^{(k)} & (s_i x_\lambda > x_\lambda \text{ and } s_i x_\lambda \in \tilde{S}_{k+1}^0) \\ -\mathfrak{g}_\lambda^{(k)} & (s_i x_\lambda < x_\lambda) \\ 0 & (\text{otherwise}) \end{cases},$$

for $\lambda \in \mathcal{P}^k$ and $i \in I$. Moreover, we have

$$D_i \cdot \tilde{\mathfrak{g}}_\lambda^{(k)} = \begin{cases} \tilde{\mathfrak{g}}_{s_i \lambda}^{(k)} & (s_i x_\lambda > x_\lambda \text{ and } s_i x_\lambda \in \tilde{S}_{k+1}^0) \\ \tilde{\mathfrak{g}}_\lambda^{(k)} & (\text{otherwise}) \end{cases},$$

for $i \in I$ and $\lambda \in \mathcal{P}^k$, and $D_i := T_i + 1$.

A proof of this proposition is given in § A.1.

Remark 3.14. Note that we will eventually show $\sigma(\mathfrak{g}_\lambda^{(k)}) = g_\lambda^{(k)}$. There is an action of H_{k+1} on $\Lambda_{(k)}$ used [4]. If we denote the action of T_i in [4, § 5.4] by T'_i , we have $T'_i = \sigma \circ T_i \circ \sigma^{-1}$.

3.4 Outline of Proof of Theorem 1.1

The following characterization property for the closed K - k -Schur functions is available. Note that for $1 \leq r \leq k$, we have $\tilde{g}_{(1^r)}^{(k)} = \tilde{g}_{(1^r)}$ and $\tilde{g}_{(r)}^{(k)} = \tilde{g}_{(r)} = \tilde{h}_r$ (Proposition 2.9).

Lemma 3.15. For $\lambda \in \mathcal{P}^k$, and $1 \leq r \leq k$, we have

$$\tilde{g}_{(1^r)} \cdot \tilde{g}_\lambda^{(k)} = \sum_{\substack{\mu \leq \lambda \\ k}} \sum_{\substack{A \subset I, |A| \leq r \\ u_A * x_\mu \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(u_A * \mu) + \ell(\mu)} g_{u_A * x_\mu}^{(k)}. \quad (3.6)$$

Moreover, the $\{\tilde{g}_\lambda^{(k)}\}_{\lambda \in \mathcal{P}^k}$ are the unique elements of $\Lambda_{(k)}$ satisfying (3.6) for $1 \leq r \leq k$.

Proof. By summing up (2.5) over $\mu \in \mathcal{P}^k$ such that $\mu \leq \lambda$ and integers i with $0 \leq i \leq r$, we have

$$\tilde{g}_{(r)} \cdot \tilde{g}_\lambda^{(k)} = \sum_{\substack{\mu \leq \lambda \\ k}} \sum_{\substack{A \subset I, |A| \leq r \\ d_A * x_\mu \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(d_A * \mu) + \ell(\mu)} g_{d_A * x_\mu}^{(k)}. \quad (3.7)$$

In fact, this identity appears in [34, p. 470]. We apply the k -conjugation to both hand sides of (3.7), and use Corollary 3.4, to have

$$\tilde{g}_{(1^r)} \cdot \tilde{g}_{\omega_k(\lambda)}^{(k)} = \sum_{\substack{\mu \leq \lambda \\ k}} \sum_{\substack{A \subset I, |A| \leq r \\ d_A * x_\mu \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(d_A * \mu) + \ell(\mu)} g_{u_{\overline{A}} * x_{\omega_k(\mu)}}^{(k)}.$$

This is equivalent to

$$\tilde{g}_{(1^r)} \cdot \tilde{g}_\lambda^{(k)} = \sum_{\substack{\omega_k(\mu) \leq \omega_k(\lambda) \\ k}} \sum_{\substack{A \subset I, |A| \leq r \\ d_A * x_{\omega_k(\mu)} \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(d_A * \omega_k(\mu)) + \ell(\omega_k(\mu))} g_{u_{\overline{A}} * x_\mu}^{(k)}. \quad (3.8)$$

Noting that $\omega_k(d_A * x_{\omega_k(\lambda)}) = u_{\overline{A}} * x_\mu$, and $|\overline{A}| = |A|$, the right hand side of (3.8) is equal to that of (3.6) because the involution ω_k preserves the Bruhat order and the length. \square

Theorem 1.1 will follow easily once we have the proposition below.

Proposition 3.16. For $\lambda \in \mathcal{P}^k$, and $1 \leq r \leq k$, we have

$$g_{(1^r)} \cdot \tilde{g}_\lambda^{(k)} = \sum_{\substack{\mu \leq \lambda \\ k}} \sum_{\substack{A \subset I, |A| \leq r \\ u_A * x_\mu \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(u_A * \mu) + \ell(\mu)} \mathfrak{g}_{u_A * x_\mu}^{\circ(k)}. \quad (3.9)$$

We apply the ring automorphism θ of $\Lambda_{(k)}$ on both hand sides of (3.9) to have

$$\tilde{g}_{(1^r)} \cdot \theta(\tilde{g}_\lambda^{(k)}) = \sum_{\substack{\mu \leq \lambda \\ k}} \sum_{\substack{A \subset I, |A| \leq r \\ u_A * x_\mu \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(u_A * \mu) + \ell(\mu)} \theta(\mathfrak{g}_{u_A * x_\mu}^{\circ(k)}). \quad (3.10)$$

Note that when we express $\theta(\mathfrak{g}_{u_A * x_\mu}^{\circ(k)})$ as a linear combination of $\{\theta(\tilde{g}_\lambda^{(k)})\}$, $g_{u_A * x_\mu}^{(k)}$ is also a linear combination of $\{\tilde{g}_\lambda^{(k)}\}$ with the same coefficients. Therefore we know that (3.6) and (3.10) are exactly the same equation via the correspondence $\theta(\mathfrak{g}_\lambda^{(k)}) \mapsto \tilde{g}_\lambda^{(k)}$, so we have $\theta(\mathfrak{g}_\lambda^{(k)}) = \tilde{g}_\lambda^{(k)}$ for all $\lambda \in \mathcal{P}^k$.

Here is the outline of the proof Proposition 3.16. We compute $g_{(1^r)} \cdot \tilde{g}_\lambda^{(k)}$ by using the combinatorial theory of Catalan functions developed in [4] to show the following result, which is the technical heart of this paper (see § 3.5 below for the proof).

Lemma 3.17 (Key lemma for Proposition 3.16).

$$g_{(1^r)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)} = \sum_{A \subset I, |A| \leq r} T_{u_A} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)} \quad (3.11)$$

By definition of $\mathfrak{g}_\lambda^{\circ(k)}$ the right hand side of (3.11) can be written as

$$\sum_{\substack{\mu \leq \lambda \\ k}} \sum_{A \subset I, |A| \leq r} T_{u_A} \cdot \mathfrak{g}_\mu^{\circ(k)} \quad (3.12)$$

Thus, in order to complete the proof of Proposition 3.16, it suffices to show

$$\sum_{A \subset I, |A| \leq r} T_{u_A} \cdot \mathfrak{g}_\mu^{\circ(k)} = \sum_{\substack{A \subset I, |A| \leq r \\ u_A * x_\mu \in \tilde{S}_{k+1}^0}} (-1)^{|A| - \ell(u_A * \mu) + \ell(\mu)} \mathfrak{g}_{u_A * \mu}^{\circ(k)}.$$

The final step of the proof of Proposition 3.16 is the following.

Lemma 3.18. Let $A \subsetneq I$. Then

$$T_{u_A} \cdot \mathfrak{g}_\mu^{\circ(k)} = \begin{cases} (-1)^{|A| - \ell(u_A * x_\mu) + \ell(\mu)} \mathfrak{g}_{u_A * \mu}^{\circ(k)} & (u_A * x_\mu \in \tilde{S}_{k+1}^0) \\ 0 & (\text{otherwise}) \end{cases}$$

The proof of Lemma 3.18 is given by a general statement (Proposition A.6) that holds in the context of an arbitrary Coxeter group (W, S) and its parabolic quotient. A detailed discussion is given in § A.2.

3.5 Proof of Lemma 3.17

This section is devoted to the proof of Lemma 3.17.

3.5.1 Basic straightening rule

Definition 3.19 (map τ). Define a map $\tau : \{1, \dots, \ell\} \rightarrow I = \mathbb{Z}/(k+1)\mathbb{Z}$ by $\tau(p) = -p + 1 \pmod{(k+1)}$.

The following is the crucial combinatorial result, whose proof will be given in § 3.5.4.

Lemma 3.20 (Basic straightening rule). For $\lambda \in \mathcal{P}_m^k$ and an integer $\ell - k < p \leq \ell$ with $p \geq m + 2$,

$$K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda + \varepsilon_p) = D_{\tau(p)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)}.$$

Furthermore, we have

$$\begin{aligned} D_{\tau(p)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)} = \tilde{\mathfrak{g}}_{s_{\tau(p)}x_\lambda}^{(k)} &\iff s_{\tau(p)}x_\lambda > x_\lambda \text{ and } s_{\tau(p)}x_\lambda \in \tilde{S}_{k+1}^0 \\ &\iff \text{top}_\Psi(p) < \text{top}_\Psi(p-1) \end{aligned} \quad (3.13)$$

with $\Psi = \Delta^k(\lambda)$.

Corollary 3.21. For $\lambda \in \mathcal{P}_m^k$ and an integer $\ell - k < p \leq \ell$ with $p \geq m + 2$,

$$K(\Delta^k(\lambda); L(\Delta^k(\lambda)) \sqcup \{p\}; \lambda + \varepsilon_p) = T_{\tau(p)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)}.$$

Proof. We apply Lemma 3.11 (i) to have

$$\begin{aligned}
& K(\Delta^k(\lambda); L(\Delta^k(\lambda)) \sqcup \{p\}; \lambda + \varepsilon_p) \\
&= K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda + \varepsilon_p) - K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda) \\
&= D_{\tau(i)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)} - \tilde{\mathfrak{g}}_\lambda^{(k)} \quad (\text{by Lemma 3.20}) \\
&= T_{\tau(i)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)}.
\end{aligned}$$

□

For the reader's convenience, we give some examples showing how the proof of Lemma 3.20 goes.

Example 3.22. Let $k = 4$, $\lambda = (3, 2, 1, 0, 0, 0)$, $\Psi := \Delta^k(\lambda)$, $M = L(\Psi)$. Note that $\beta = (2, 4)$ is an addable root of Ψ . Consider $\gamma = \lambda + \varepsilon_5 = (3, 2, 1, 0, 1, 0)$. By using Lemma 3.9 (i), we have

$$K(\Psi; M; \gamma) = K(\Psi \cup \beta; M; \gamma) - K(\Psi \cup \beta; M; \gamma + \varepsilon_2 - \varepsilon_4). \quad (3.14)$$

$$\begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 2 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 1 & \\ \hline & & & & & 0 \\ \hline \end{array} = \begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 2 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 1 & \\ \hline & & & & & 0 \\ \hline \end{array} - \begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 3 & & & & \\ \hline & & 1 & & & \\ \hline & & & -1 & & \\ \hline & & & & 1 & \\ \hline & & & & & 0 \\ \hline \end{array}$$

The first term of (3.14) vanishes by (3.5) with $i = 4$. The second term equals to

$$(-1)^2 K(\Psi \cup \beta; M; s_4(\gamma + \varepsilon_2 - \varepsilon_4) - \varepsilon_4 + \varepsilon_5) = K(\Psi \cup \beta; M; \mu)$$

with $\mu := (3, 3, 1, 0, 0, 0)$ by (3.4) with $i = 4$. So we have

$$K(\Psi; M; \lambda + \varepsilon_5) = K(\Psi \cup \beta; M; \mu) = \begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 3 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline \end{array}.$$

Next, we apply (iv) to it with $j = 4$ to have

$$K(\Psi \cup \beta; M \sqcup \{4\}; \mu) + K(\Psi \cup \beta; M; (3, 3, 1, -1, 0, 0)). \quad (3.15)$$

$$\begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 3 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline \end{array} + \begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 3 & & & & \\ \hline & & 1 & & & \\ \hline & & & -1 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline \end{array}$$

The second term of (3.15) vanishes by (3.5) with $i = 3$. Noting $\Psi \cup \beta = \Delta^4(\mu)$ and $M \sqcup \{4\} = L(\Delta^4(\mu))$, we finally obtain $K(\Psi \cup \beta; M \sqcup \{4\}; \mu) = K(\Delta^4(\mu); L(\Delta^4(\mu)); \mu) = \tilde{\mathfrak{g}}_{(3,3,1)}^{(4)}$.

$$\begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 3 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 0 \\ \hline \end{array}$$

Example 3.23. Let $k = 4$, and (Ψ, M, λ) the same as Example 3.22. Consider $\gamma = \lambda + \varepsilon_6 = (3, 2, 1, 0, 0, 1)$. We can apply Lemma 3.11 (i) with $j = 5$ to have

$$K(\Psi; M; \gamma) = K(\Psi; M \setminus \{5\}; \gamma) - K(\Psi; M \setminus \{5\}; \gamma - \varepsilon_5), \quad (3.16)$$

depicted by

$$\begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 2 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 1 \\ \hline \end{array} = \begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 2 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & 0 & \\ \hline & & & & & 1 \\ \hline \end{array} - \begin{array}{|c|c|c|c|c|c|} \hline 3 & & \bullet & \bullet & \bullet & \bullet \\ \hline & 2 & & & & \\ \hline & & 1 & & & \\ \hline & & & 0 & & \\ \hline & & & & -1 & \\ \hline & & & & & 1 \\ \hline \end{array}$$

- (c) $m_M(p) = m_M(p-1)$;
(d) Ψ has a wall in rows $p-1, p$.

Then,

$$K(\Psi; M; \gamma) = K(\Psi; M; \gamma - \varepsilon_p).$$

Proof. By (a), Ψ has a ceiling in columns $p-1, p$. From Lemma 3.11 (i) with $j = p-1$, we have

$$K(\Psi; M; \gamma) = K(\Psi; M \setminus \{p-1\}; \gamma) - K(\Psi; M \setminus \{p-1\}; \gamma - \varepsilon_{p-1}). \quad (3.20)$$

Let $M' = M \setminus \{p-1\}$. Then, we have $m_{M'}(p) = m_{M'}(p-1) + 1$, which implies that we can apply (3.5) with $i = p-1$ to the first term of (3.20). Thus we have

$$K(\Psi; M; \gamma) = -K(\Psi; M'; \gamma - \varepsilon_{p-1}) + K(\Psi; M'; \gamma - \varepsilon_p),$$

where we used (3.4) in the last equality with $i = p-1$. From Lemma 3.11 (ii) with $j = p-1$, we have

$$K(\Psi; M'; \gamma - \varepsilon_p) = K(\Psi; M; \gamma - \varepsilon_p) + K(\Psi; M'; \gamma - \varepsilon_p - \varepsilon_{p-1}),$$


in which the second term vanishes by (3.5) with $i = p-1$. \square

Example 3.27. From Lemma 3.26 with $p = 6$, we have

3.5.3 Mirror edges and mirror paths

Definition 3.28 (Mirror edge, Mirror path, Mirror top). Let Ψ be a root ideal, and $e := (p \rightarrow q)$ a bounce edge of Ψ . We say e is a *mirror edge* if $p-1 \rightarrow q-1$ also is a bounce edge of Ψ . A bounce path $p = p_0 \rightarrow p_1 \rightarrow \dots \rightarrow p_L$ is a *mirror path of length L* of Ψ if $p_i \rightarrow p_{i+1}$ is a mirror edge for each $0 \leq i \leq L-1$. If such L is maximal, we define $\text{mtop}_\Psi(p) = p_L$ called the *mirror top* of p . In particular, if p is not contained in any mirror path, we have $\text{mtop}_\Psi(p) = p$.

Example 3.29. For the root ideal Ψ illustrated in the following picture, $10 \rightarrow 6 \rightarrow 3$ is a mirror path having the maximal length 2. Then, we have $\text{mtop}_\Psi(10) = 3$.



It is easy to see the following.

Lemma 3.30. Let Ψ be a root ideal and $p \rightarrow q$ a mirror edge of Ψ . Then $\beta := (q, p-1)$ is an addable root of Ψ . Furthermore, $\Psi \cup \{\beta\}$ has a wall in rows $q-1, q$, and a ceiling in columns $p-1, p$.

Lemma 3.31. Let Ψ be a root ideal and z be the lowest nonempty row of Ψ . Assume that Ψ is wall-free in nonempty rows, *i.e.*, Ψ has no wall between the 1st and z -th rows. Then, for $p \geq 2$, we have

$$\text{mtop}_\Psi(p) = \max\{\text{top}_\Psi(p), \text{top}_\Psi(p-1) + 1\}.$$

Proof. Let $p \rightarrow q$ be a bounce edge of Ψ . Note that, whenever $\text{up}_\Psi(p-1)$ exists, it satisfies $\text{up}_\Psi(p-1) = q-1$; otherwise Ψ have a wall in rows $q-1, q$.

Let $p = p_0 \rightarrow p_1 \rightarrow \dots \rightarrow p_L$ and $p-1 = q_0 \rightarrow q_1 \rightarrow \dots \rightarrow q_{L'}$ be bounce paths of maximal lengths. By the above remark, we have $q_i = p_i - 1$ for $0 \leq i \leq \min(L, L')$. If $L > L'$ we have $\text{mtop}_\Psi(p) = p_{L'} = q_{L'} + 1 = \text{top}_\Psi(p-1) + 1$, while $\text{mtop}_\Psi(p) > \text{top}_\Psi(p)$. If $L = L'$ then $\text{mtop}_\Psi(p) = \text{top}_\Psi(p) = \text{top}_\Psi(p-1) + 1$. If $L < L'$ then $\text{mtop}_\Psi(p) = \text{top}_\Psi(p)$, while $\text{mtop}_\Psi(p) = \text{top}_\Psi(p) = q_L + 1 > \text{top}_\Psi(p-1) + 1$. \square

Note that $\Psi = \Delta^k(\lambda)$ is wall-free in nonempty rows, then we can apply the above result to it.

Lemma 3.32. Let $\lambda \in \mathcal{P}^k$. Set $\Psi := \Delta^k(\lambda)$, $M = L(\Psi)$. Let $e = (p \rightarrow q)$ be a bounce edge of Ψ such that $q \geq 2$.

- (1) e is a mirror edge of Ψ if and only if $\lambda_{q-1} = \lambda_q$.
- (2) If e is a mirror edge of Ψ then $m_M(p) = m_M(p-1) + 1$.
- (3) If e is not a mirror edge, then $\lambda_{q-1} > \lambda_q$.

Proof. (1), (2) are clear from the definition of Ψ, M . (3) If e is not a mirror edge then from (1) we have $\lambda_{q-1} \neq \lambda_q$. Since λ is a partition, we have $\lambda_{q-1} > \lambda_q$. \square

Lemma 3.33. Let $\lambda \in \mathcal{P}^k$, and $\Psi := \Delta^k(\lambda) \subset \Delta_\ell^+$. Then for $p \in \{1, \dots, \ell\}$,

$$\tau(\text{mtop}_\Psi(p)) = \tau(p).$$

Proof. This is [4, Lemma 5.7]. \square

In the final step of the proof of Lemma 3.20 we use the following.

Lemma 3.34 (Cleaning Lemma. [4, Lemma 4.7]). Let (Ψ, M, γ) be a Catalan triple such that

- (a) $\beta = (q, p-1)$ is a removable root of Ψ ;
- (b) Ψ has a wall in rows $p-1, p$;
- (c) Ψ has a ceiling in columns $p-1, p$;
- (d) $\gamma_q = \gamma_{q-1}$;
- (e) $m_M(p) = m_M(p-1) + 1$.

Then

$$K(\Psi; M; \gamma) = K(\Psi \setminus \beta; M; \gamma). \quad (3.21)$$

3.5.4 Proof of Lemma 3.20

Proof of Lemma 3.20. Let $\lambda \in \mathcal{P}^k$, and $\Psi := \Delta^k(\lambda) \subset \Delta_\ell^+$. Set $\mathcal{T} := (\Psi, M, \gamma) = (\Delta^k(\lambda), L(\Delta^k(\lambda)), \lambda + \varepsilon_p)$. Let $p = p_0 \rightarrow p_1 \rightarrow \dots \rightarrow p_L$ be the mirror path starting from p of maximal length $L \geq 0$. We will prove that we can successively apply Lemma 3.24 to \mathcal{T} and obtain the sequence of Catalan triples:

$$\mathcal{T} =: \mathcal{T}_0 \xrightarrow{p_0} \mathcal{T}_1 \xrightarrow{p_1} \dots \xrightarrow{p_{L-1}} \mathcal{T}_L =: \mathcal{T}'. \quad (3.22)$$

Here $\mathcal{T}_i \xrightarrow{p_i} \mathcal{T}_{i+1}$ means that \mathcal{T}_{i+1} is obtained from \mathcal{T}_i by applying Lemma 3.24 (3.17) with respect to the mirror edge $p_i \rightarrow p_{i+1}$ of \mathcal{T}_i . If $L = 0$, it suffices to put $\mathcal{T}' = \mathcal{T}$. Assume $L \geq 1$. We can check immediately that $\mathcal{T} = \mathcal{T}_0$ satisfies the assumptions (a–d) of Lemma 3.24; (a) holds by Lemma 3.30; (b) holds from $\lambda_{p-1} = \lambda_p = 0$ ($\because p \geq m+2$) and $\gamma = \lambda + \varepsilon_p$; (c) holds by Lemma 3.32 (2); (d) holds because there are no roots of $\Psi = \Delta^k(\lambda)$ in rows $p-1, p$ ($\because p > \ell - k$). Then, we obtain a new Catalan triple $\mathcal{T}_1 = (\Psi^{(1)}, M^{(1)}, \gamma^{(1)})$ by applying (3.17) to \mathcal{T}_0 . From Lemma 3.32 (1), we have $\lambda_{p_1-1} = \lambda_{p_1}$, which implies $\gamma_{p_1}^{(1)} = \gamma_{p_1-1}^{(1)} + 1$. By Lemma 3.30, $\Psi^{(1)}$ has a wall in rows p_1-1, p_1 . Hence, \mathcal{T}_1 satisfies (b), (d) of Lemma 3.24.

The remaining conditions (a) and (c) hold for \mathcal{T}_1 by the same reasons as the case of \mathcal{T}_0 ; the multiset $M^{(1)}$ is the same as M ; the root ideals $\Psi^{(1)}$ and Ψ coincide with each other in columns $\leq p_1$. Therefore, we can apply Lemma 3.24 (3.17) to \mathcal{T}_1 with respect to the mirror edge $p_1 \rightarrow p_2$ and obtain \mathcal{T}_2 . The above procedure can be repeated L times to obtain the sequence (3.22). Let $\mathcal{T}' = (\Psi', M', \gamma')$. Then, we have

$$\Psi' = \Psi \cup \{\beta_1, \dots, \beta_L\}, \quad M' = M, \quad \gamma' = \gamma + \varepsilon_{\text{mtop}_\Psi(p)} - \varepsilon_p (= \lambda + \varepsilon_{\text{mtop}_\Psi(p)}), \quad (3.23)$$

where $\beta_i := (p_i, p_{i-1} - 1)$ ($1 \leq i \leq L$). Note that Ψ' has a wall in rows $p_i - 1, p_i$ for each i . The following procedure differs depending on whether (i) $\text{top}_\Psi(p) < \text{top}_\Psi(p - 1)$ or (ii) $\text{top}_\Psi(p) > \text{top}_\Psi(p - 1)$.

If $\text{top}_\Psi(p) < \text{top}_\Psi(p - 1)$, we have $\text{mtop}_\Psi(p) = \text{top}_\Psi(p - 1) + 1 > \text{top}_\Psi(p)$ by Lemma 3.31, which implies that there is a bounce edge $\text{mtop}_\Psi(p) \rightarrow q$ of Ψ for some q . Note that $\text{mtop}_\Psi(p) \rightarrow q$ is also a bounce edge of Ψ' . Therefore, we can apply Lemma 3.24 to \mathcal{T}' with respect to this bounce edge. From (3.18), it follows that

$$K(\Psi'; M'; \gamma') = K(\Psi' \cup \beta; M' \sqcup \{\text{mtop}_\Psi(p) - 1\}; \gamma' + \varepsilon_q - \varepsilon_{\text{mtop}_\Psi(p)}),$$

where

$$\beta := (q, \text{mtop}_\Psi(p) - 1).$$

Note that

$$\lambda' := \gamma' + \varepsilon_q - \varepsilon_{\text{mtop}_\Psi(p)} = \lambda + \varepsilon_q - \varepsilon_p$$

and $M' \sqcup \{\text{mtop}_\Psi(p) - 1\} = L(\Delta^k(\lambda'))$. Moreover, λ' is a partition by Lemma 3.32 (3). To prove that λ' is k -bounded, it suffices to consider the case when $\lambda_1 = k$ and $q = 1$. However, one sees that this does not occur in view of Lemma 3.31. We also note that

$$\Psi \cup \beta = (\Psi' \cup \beta) \setminus \{\beta_1, \dots, \beta_L\} = \Delta^k(\lambda').$$

By using Lemma 3.34 recursively, we can remove $\beta_L, \dots, \beta_2, \beta_1$ from Ψ' and get

$$K(\Delta^k(\lambda'); L(\Delta^k(\lambda')); \lambda') = \tilde{\mathfrak{g}}_{\lambda'}^{(k)}.$$

From Lemma 3.33, we have

$$q = \text{up}_\Psi(\text{mtop}_{\mathcal{T}'}(p)) \equiv \text{mtop}_{\mathcal{T}'}(p) \equiv \tau(p) \pmod{k+1},$$

and $s_{\tau(p)}x_\lambda = x_{\lambda'}$.

Next assume $\text{top}_\Psi(p) > \text{top}_\Psi(p - 1)$. We check that \mathcal{T}' satisfies the conditions (a-d) of Lemma 3.26 with $p = \text{mtop}_\Psi(p)$. From Lemma 3.31, we have $\text{mtop}_\Psi(p) = \text{top}(p)$, which implies (a). (b) follows from Lemma 3.32 (1). (c) follows from the fact there is a ceiling in the columns in the rows $p - 1, p$. (d) follows from the definition of M' .

From Lemma 3.26, we have

$$K(\Psi'; M'; \gamma') = K(\Psi'; M'; \gamma' - \varepsilon_{\text{mtop}_\Psi(p)})$$

In view of (3.23), we can rewrite

$$K(\Psi'; M'; \gamma' - \varepsilon_{\text{mtop}_\Psi(p)}) = K(\Delta^k(\lambda) \cup \{\beta_1, \dots, \beta_L\}; \Delta^k(\lambda); \lambda).$$

By Lemma 3.34, we can remove $\beta_L, \dots, \beta_2, \beta_1$ successively to get

$$K(\Delta^k(\lambda); L(\Delta^k(\lambda)); \lambda) = \tilde{\mathfrak{g}}_\lambda^{(k)}.$$

Again by Lemma 3.33,


$$q = \text{up}_\Psi(\text{mtop}_{\mathcal{T}'}(p)) \equiv \text{mtop}_{\mathcal{T}'}(p) \equiv \tau(p) \pmod{k+1},$$

and $s_{\tau(p)}x_\lambda = x_\lambda$. □

Here are some examples.


Example 3.35. Case 1 ($p = 8$): $\text{top}(p) = \text{mtop}_\Psi(p) = 4 < \text{top}(p - 1) = 7$ with $p = 8$. This case we have $L = 0$.

1. Lemma 3.24 (3.18) with $p = 8$.




Example 3.36. Case 2 ($p = 11$): $\text{top}(p) = 11 > \text{top}(p - 1) = 6$. $L = 0$.

1. Lemma 3.26 with $p = 11$.




Example 3.37. Case 1 ($p = 9$): $\text{top}(p) = 2 < \text{top}(p - 1) = 4$

1. Lemma 3.24 (3.17) with $p = 9$.
2. Lemma 3.24 (3.18) with $p = 5$.
3. Cleaning Lemma $q = 5$.




Example 3.38. Case 1 ($p = 10$): $\text{top}(p) = 1 < \text{top}(p - 1) = 2$. $\tau(p) = 1$

1. Lemma 3.24 (3.17) with $p = 10$.
2. Lemma 3.24 (3.17) with $p = 6$.
3. Lemma 3.24 (3.18) with $p = 3$.
4. Cleaning Lemma twice $q = 3, 5$



Example 3.39. Case 2 ($p = 10$): $\text{top}(p) = 6 > \text{top}(p - 1) = 2$.

1. Lemma 3.24 (3.17) with $p = 10$.
2. Lemma 3.26 with $p = 6$.
3. Cleaning $q = 6$



Remark 3.40. From the proof of Lemma 3.20 given above, we can slightly improve the statement of the lemma as follows: For $\lambda \in \mathcal{P}_m^k$, an integer $\ell - k < p \leq \ell$ with $p \geq m + 2$, and $\kappa \in \mathbb{Z}^\ell$ with $p < \text{supp}(\kappa) \leq \ell$, we have

$$K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda + \varepsilon_p + \kappa) = K(\Delta^k(\mu); \Delta^k(\mu); \mu + \kappa),$$

where $\tilde{\mathfrak{g}}_\mu^{(k)} = D_{\tau(p)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)}$. In fact, throughout the straightening process introduced in § 3.5.4, κ is left unchanged because the process affects only topmost p rows of the diagram.

From Remark 3.40, we obtain the following key formula: For $\lambda \in \mathcal{P}_m^k$, and a set of integers $\ell - k < p_1 < \dots < p_r \leq \ell$ with $p_1 \geq m + 2$, we have

$$K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda + \varepsilon_{p_1} + \dots + \varepsilon_{p_r}) = D_{\tau(p_r)} \cdots D_{\tau(p_1)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)}. \quad (3.24)$$

3.5.5 Proof of Lemma 3.17

Let Ψ_1, Ψ_2 be root ideals of $\Delta_{\ell_1}^+$ and $\Delta_{\ell_2}^+$ respectively. Let $\Psi_1 \uplus \Psi_2$ be a subset of $\Delta_{\ell_1 + \ell_2}^+$ defined by

$$\{(i, j) \in \Delta_{\ell_1 + \ell_2}^+ \mid (i, j) \in \Psi_1 \text{ or } (i - \ell_1, j - \ell_1) \in \Psi_2 \text{ or } (i \leq \ell_1 \text{ and } j > \ell_1)\}$$

Let $\lambda \in \mathcal{P}_\ell^k$. Set $\Psi = \Delta(\lambda)^k$. By the product rule [4, Lemma 3.8], we have

$$g_{(1^r)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)} = K(\Psi \uplus \emptyset_r; \Psi \uplus \emptyset_r; \lambda + \varepsilon_{\{\ell+1, \dots, \ell+r\}}). \quad (3.25)$$

By applying Diagonal Removable Lemma [4, Lemma 4.13], one shows that the right hand side of (3.25) is equal to

$$K(\Psi'; \Psi'; \lambda + \varepsilon_{\{\ell+1, \dots, \ell+r\}}) \quad (3.26)$$

where $\Psi' = (\Delta^k(\lambda) \uplus \emptyset_r) \setminus \{(i, j) \in \Delta_{\ell+r}^+ \mid i \leq \ell, j \geq \ell, j - i \leq r - 1\}$. We can apply [4, Lemma 5.3] with $x = \ell - r + 1, h = \ell - k + 1$ (in the notation there) to show that (3.26) is equal to

$$\sum_{a=0}^r \sum_{\substack{S \subset \{\ell-k+1+r-a, \dots, \ell\} \\ |S|=a}} K(\Psi''; L(\Psi'') \sqcup S; \lambda + \varepsilon_S + \varepsilon_{\{\ell+1, \dots, \ell+r-a\}}), \quad (3.27)$$

where $\Psi'' = \Psi' \setminus \{(i, j) \mid i \leq \ell, j \geq \ell, j - i \leq k\}$ (see also the last part of the proof of [4, Proposition 5.4]). By using Lemma 3.20, Corollary 3.21, and (3.24), we write the right hand side of (3.27) as

$$\sum_{a=0}^r \sum_{\ell-k+1+r-a \leq p_1 < \dots < p_a \leq \ell} D_{\tau(\ell+r-a)} \cdots D_{\tau(\ell+1)} T_{\tau(p_a)} \cdots T_{\tau(p_1)} \cdot \tilde{\mathfrak{g}}_\lambda^{(k)}. \quad (3.28)$$

Proposition 3.41. For $1 \leq r \leq k$, and $\ell \in \mathbb{Z}$, we have

$$\sum_{a=0}^r \sum_{\ell-k+1+r-a \leq p_1 < \dots < p_a \leq \ell} D_{\tau(\ell+r-a)} \cdots D_{\tau(\ell+1)} T_{\tau(p_a)} \cdots T_{\tau(p_1)} = \sum_{A \subset I, |A| \leq r} T_{u_A}.$$

Proof. We first note that for any subsequence A_1 of $(\tau(\ell+r-a), \dots, \tau(\ell+1))$ such that $0 \leq |A_1| \leq r-a$, and A_2 of $(\tau(\ell-k+r-a), \dots, \tau(\ell))$ such that $|A_2| = a$, the concatenation $A_1 \cdot A_2$ is a cyclically increasing sequence of length less than or equal to r .

For $0 \leq a \leq n \leq r$, let

$$X_a^n = \left\{ (p_1, \dots, p_n) \in \mathbb{Z}^n; \begin{array}{l} \ell - k + r - a < p_1 < \dots < p_a \leq \ell \\ < p_{a+1} < \dots < p_n \leq \ell + r - a \end{array} \right\}.$$

Substituting $D_p = T_p + 1$ to the left hand side of the proposition, we have

$$\begin{aligned} & \sum_{a=0}^r \sum_{\ell-k+1+r-a \leq p_1 < \dots < p_a \leq \ell} (T_{\tau(\ell+r-a)} + 1) \cdots (T_{\tau(\ell+1)} + 1) T_{\tau(p_a)} \cdots T_{\tau(p_1)} \\ &= \sum_{a=0}^r \sum_{n=a}^r \sum_{(p_1, \dots, p_n) \in X_a^n} T_{u_{\{\tau(p_1), \dots, \tau(p_n)\}}} \\ &= \sum_{n=0}^r \sum_{a=0}^n \sum_{(p_1, \dots, p_n) \in X_a^n} T_{u_{\{\tau(p_1), \dots, \tau(p_n)\}}}. \end{aligned}$$

Then it suffices to show that the map

$$\text{rm} : \bigsqcup_{a=0}^n X_a^n \rightarrow \{A \subset I; |A| = n\}; \quad (p_1, \dots, p_n) \mapsto \{\overline{p_1}, \dots, \overline{p_n}\}$$

is bijective. For this, we identify a subset $A \subset I$ with a 01-sequence $\eta_1 \eta_2 \dots \eta_{k+1} \in \{0, 1\}^{k+1}$ by letting $\eta_p = 0$ if $\overline{p} \in A$ and $\eta_p = 1$ otherwise. Then the image of X_a^n by rm is contained in the set

$$S_a^n := \left\{ \eta_1 \eta_2 \dots \eta_{k+1} \in \{0, 1\}^{k+1}; \begin{array}{l} \sum_{p=1}^{k+1} \eta_p = k+1-n, \\ \sum_{i=1}^{r-a} \eta_{\ell+i} = r-n \\ \sum_{i=1}^{r-a+1} \eta_{\ell+i} = r-n+1 \end{array} \right\}.$$

Obviously, we have $S_a^n \cap S_{a'}^n \neq \emptyset \iff a = a'$, which implies the fact that rm is injective because its restriction on X_a^n is injective. Since $\sum_{a=0}^n |X_a^n| = \sum_{a=0}^n \binom{k-r+a}{a} \binom{r-a}{n-a} = \binom{k+1}{n} = \#\{A \subset I; |A| = n\}$, rm is also surjective. \square

This completes the proof of Lemma 3.17.

4 Relation to quantum K -theory

4.1 Quantum K -theory ring of the flag variety

Let $QK(G/B)$ denote the finite version of the quantum K -theory ring of the flag variety G/B . This is a free $\mathbb{C}[Q]$ -module with basis $\mathcal{O}_{G/B}^w$ ($w \in S_{k+1}$) inside the (small) quantum

K -theory ring $\widehat{QK}(G/B)$ of G/B defined as a free $\mathbb{C}[[Q]]$ -module ([9]). Anderson, Chen, Tseng, and Iritani [1] proved that $QK(G/B)$ forms a subring.

Theorem 4.1 ([15],[27],[25],[26]). $QK(G/B)$ can be identified with the quotient ring

$$A_{k+1} := \mathbb{C}[Q][z_1, \dots, z_{k+1}]/I_{k+1}, \quad (4.1)$$

$$I_{k+1} := \left\langle \sum_{\substack{I \subset \{1, \dots, k+1\} \\ |I|=i}} \prod_{j \in I} z_j \prod_{\substack{j \in I \\ j+1 \notin I}} (1 - Q_j) - \binom{k+1}{i} \mid 1 \leq i \leq k+1 \right\rangle, \quad (4.2)$$

such that for $w \in S_{k+1}$ the *quantum Grothendieck polynomial* \mathfrak{G}_w^Q of Lenert and Maeno, with the change of variables $x_i = 1 - z_i$, represents $\mathcal{O}_{G/B}^w$ in $QK(G/B_{k+1})$.

The presentation for $QK(G/B)$ given by (4.1), (4.2) was announced by Kirillov and Maeno [15] (see [25, Theorem 3.1]). Lenert and Maeno proved that the quantum Grothendieck polynomials \mathfrak{G}_w^Q satisfy the conjectural Chevalley type formula [25, Theorem 6.1] due to Lenart and Postnikov [27], which was proved to be true by Lenart, Naito, and Sagaki [26].

4.2 Peterson isomorphism in K -theory

Recall that $\tau_i := g_{R_i}$ with $R_i = \overbrace{(i, \dots, i)}^{k+1-i}$ for $1 \leq i \leq k$. Note that the notation R_i is different from the one used in [11]. So the indices of τ_i is switched from the one in [11] by $i \mapsto k+1-i$ too (see Remark 4.2 below).

There is a ring isomorphism

$$\Phi_{k+1} : A_{k+1}[Q_1^{-1}, \dots, Q_k^{-1}] \longrightarrow \Lambda_{(k)}[\tau_i^{-1}, (\tau_i^+)^{-1} \ (1 \leq i \leq k)], \quad (4.3)$$

$$z_i \mapsto \frac{\tau_i \tau_{i-1}^+}{\tau_i^+ \tau_{i-1}} \ (1 \leq i \leq k+1), \quad Q_i \mapsto \frac{\tau_{i-1} \tau_{i+1}}{\tau_i^2} \ (1 \leq i \leq k), \quad (4.4)$$

given by Ikeda, Iwao, and Maeno in [11]. The map Φ_{k+1} was constructed by solving the relativistic Toda lattice equation with the initial condition that the Lax matrix is unipotent. Although the construction of Φ_{k+1} has no apparent geometric meaning, it is expected that the map sends a Schubert structure sheaf $\mathcal{O}_{G/B}^w$ to an element in $K_*(\text{Gr})_{\text{loc}}$ related to a Schubert class.

Remark 4.2. Our convention for Φ_{k+1} is slightly different from the one in [11]. Let $\overline{\Omega} \in \text{Aut}(\Lambda_{(k)}[\tau_i^{-1}, (\tau_i^+)^{-1}])$ be the natural extension of Ω . One can check that $\Phi'_{k+1} = \overline{\Omega} \circ \Phi_{k+1}$ coincides with the map introduced in [11] by replacing σ_i in their notation with τ_i^+ .

4.3 K -homology of the affine Grassmannian

For $x \in \tilde{S}_{k+1}^0$, let $\xi_x^0 \in K_*(\text{Gr})$ be the element defined [19, § 6.3]. The non-equivariant K -theoretic k -Schur function $g_x^{(k)}$ [19, Theorem 7.17 (2)] corresponds ξ_x^0 . Let us denote this isomorphism by $\alpha_\circ : K_*(\text{Gr}) \rightarrow \Lambda_{(k)}$. It holds that $\alpha_\circ(\mathcal{O}_x) = \tilde{g}_x^{(k)}$ ([34], [18, Lemma 2 (ii)]). We define a twisted isomorphism $\alpha : K_*(\text{Gr}) \rightarrow \Lambda_{(k)}$ given by $\alpha := \sigma^{-1} \circ \alpha_\circ$. So we have

$$\alpha(\mathcal{O}_x^{\text{Gr}}) = \sigma^{-1}(\tilde{g}_x^{(k)}). \quad (4.5)$$

Proposition 4.3. If $\lambda \subset R_i$ for some $1 \leq i \leq k$, equivalently $\lambda_1 + \ell(\lambda) \leq k+1$, we have

$$\alpha(\mathcal{O}_\lambda^{\text{Gr}}) = g_\lambda. \quad (4.6)$$

In particular we have

$$\alpha(\mathcal{O}_{R_i}^{\text{Gr}}) = \tau_i. \quad (4.7)$$

Proof. We know from Proposition 2.9 that $\tilde{g}_\lambda^{(k)} = \tilde{g}_\lambda$. Therefore using Proposition 2.5, we have

$$\alpha(\mathcal{O}_\lambda^{\text{Gr}}) = \sigma^{-1}(\tilde{g}_\lambda^{(k)}) = \sigma^{-1}(\tilde{g}_\lambda) = g_\lambda. \quad \square$$

4.4 Correspondence of Schubert bases

We consider the extended affine symmetric group \hat{S}_{k+1} , which is generated by $\{s_i \mid i \in I\} \cup \{\pi\}$ satisfying the same relations among s_i ' and

$$\pi^{k+1} = id, \quad \pi s_i = s_{i+1} \pi.$$

We have $\hat{S}_{k+1} \cong S_{k+1} \ltimes P^\vee$, where P^\vee is the coweight lattice of $SL_{k+1}(\mathbb{C})$. The translation element associated to $-\varpi_i^\vee \in P^\vee$ is explicitly given by

$$t_{-\varpi_i^\vee} = \pi^{-i} x_{R_i}.$$

Example 4.4. For $k = 3$,

$$t_{-\varpi_1^\vee} = \pi^{-1} s_2 s_3 s_1, \quad t_{-\varpi_2^\vee} = \pi^{-2} s_0 s_3 s_1 s_0, \quad t_{-\varpi_3^\vee} = \pi^{-3} s_2 s_1 s_0. \quad (4.8)$$

Let $w \in S_{k+1}$ be an i -Grassmannian permutation, i.e., $\text{Des}(w) = \{i\}$. The set of all i -Grassmannian permutations in S_{k+1} is in bijection with \mathcal{P}_i^{k+1-i} . Explicitly, for an i -Grassmannian permutation in S_{k+1} the corresponding partition $\lambda \in \mathcal{P}_i^{k+1-i}$ is given by

$$\lambda_{i+1-j} = w(j) - j \quad (1 \leq j \leq i). \quad (4.9)$$

For each partition λ in \mathcal{P}_i^{k+1-i} , we denote the corresponding i -Grassmannian permutation by $w_{\lambda,i}$. For $\lambda \in \mathcal{P}_i^{k+1-i}$, the dual partition of λ is the element λ^\vee in \mathcal{P}_i^{k+1-i} defined by $\lambda_j^\vee = k+1-i-\lambda_{i+1-j}$ ($1 \leq j \leq i$).

Proposition 4.5 ([11]). Let $w_{\lambda,i} \in S_{k+1}$ be an i -Grassmannian permutation. Then

$$\Phi_{k+1}(\mathcal{O}_{G/B}^{w_{\lambda,i}}) = \frac{g^{(\lambda^\vee)'}}{\tau_i}, \quad (4.10)$$

where $(\lambda^\vee)'$ is the conjugate of λ^\vee .

Proof. Let Φ'_{k+1} be the map in [11]. Recall that for $w \in S_{k+1}$, $\mathcal{O}_{G/B}^w$ is identified with $\mathfrak{G}_w^Q \bmod I_{k+1}$. Theorem 7.1 [11] reads $\Phi'_{k+1}(\mathfrak{G}_{w_{\lambda,i}}^Q \bmod I_{k+1}) = g^{\lambda^\vee} / g^{(k+1-i)i}$. From Remark 4.2 we have (4.10). \square

In order to describe the images $\Phi_{k+1}(\mathcal{O}_{G/B}^w)$ we need a map $S_{k+1} \rightarrow \mathcal{P}^k$, $w \mapsto \theta_k(w)$ due to Lam and Shimozono [22, Lemma 11].

Proposition 4.6 ([22]). Let $w \in S_{k+1}$. There is a k -bounded partition $\theta_k(w)$ such that

$$w t_{-\sum_{i \in \text{Des}(w)} \varpi_i^\vee} = \pi^{-\sum_{i \in \text{Des}(w)} i} \cdot x_{\theta_k(w)}. \quad (4.11)$$

Proof. One can show that the left hand side of (4.11) is an affine Grassmannian element in $\hat{S}_{k+1} = \langle \pi \rangle \ltimes \tilde{S}_{k+1}$ (see the first part of the proof of Lemma 11 in [22]). Then such an element can be uniquely written in the form of the right hand side of (4.11) for a k -bounded partition, which we denote by $\theta_k(w)$. \square

For an i -Grassmannian permutation $w = w_{\lambda,i}$ we have $\theta_k(w) = (\lambda^\vee)'$ ([11, Lemma 7.1]). See § B, for another direct proof of this fact.

Next result is a refined version of [11, Conjecture 1.8].

Theorem 4.7. For $w \in S_{k+1}$, we have

$$\Phi_{k+1}(\mathcal{O}_{G/B}^w) = \frac{\sigma^{-1}(\tilde{g}_{\theta_k(w)}^{(k)})}{\prod_{i \in \text{Des}(w)} \tau_i}. \quad (4.12)$$

Proof. Let Q^\vee denote the coroot lattice of $\mathrm{SL}_{k+1}(\mathbb{C})$. Then we have $\tilde{S}_{k+1} \cong S_{k+1} \times Q^\vee$. We denote $t_\beta \in \tilde{S}_{k+1}$ be the translation element corresponding to $\beta \in Q^\vee$. Kato [13] constructed an injective ring homomorphism

$$\kappa : K_*(\mathrm{Gr}_{\mathrm{SL}_{k+1}})_{\mathrm{loc}} \hookrightarrow \widehat{QK}(G/B_{k+1})_{\mathrm{loc}}$$

such that

$$\mathcal{O}_{wt_\beta}^{\mathrm{Gr}} \cdot (\mathcal{O}_{t_\beta}^{\mathrm{Gr}})^{-1} \mapsto \mathcal{O}_{G/B}^w \quad (w \in S_{k+1}), \quad (4.13)$$

where $\beta \in -Q^\vee$ so that $wt_\beta \in \tilde{S}_{k+1}^0$.

The localization $K_*(\mathrm{Gr})_{\mathrm{loc}}$ in [13] is defined by the multiplicative set generated by $\mathcal{O}_{t_\beta}^{\mathrm{Gr}}$ ($\beta \in \tilde{Q}$). Since $\mathcal{O}_{R_i}^{\mathrm{Gr}}$ corresponds to $\tau_i = g_{R_i}$ by α (Proposition 4.3), the isomorphism α yields $K_*(\mathrm{Gr})_{\mathrm{loc}} \cong \Lambda_{(k)}[\tau_i^{-1} \ (1 \leq i \leq k)]$. Recall A_{k+1} is the quotient ring defined by (4.1), (4.2). Let κ' be the composition

$$\begin{aligned} K_*(\mathrm{Gr})_{\mathrm{loc}} &\cong \Lambda_{(k)}[\tau_i^{-1} \ (1 \leq i \leq k)] \\ &\hookrightarrow \Lambda_{(k)}[\tau_i^{-1}, (\tau_i^+)^{-1} \ (1 \leq i \leq k)] \\ &\cong A_{k+1}[\mathcal{Q}_i^{-1} \ (1 \leq i \leq k)] \\ &\cong QK(G/B_{k+1})_{\mathrm{loc}} \\ &\hookrightarrow \widehat{QK}(G/B_{k+1})_{\mathrm{loc}}. \end{aligned}$$

We claim that $\kappa' = \kappa$.

Note that $K_*(\mathrm{Gr})$ is generated by $\mathcal{O}_{s_{i-1} \dots s_1 s_0}^{\mathrm{Gr}} = \mathcal{O}_{(i)}^{\mathrm{Gr}}$ ($1 \leq i \leq k$), and that $\alpha(\mathcal{O}_{(i)}^{\mathrm{Gr}}) = g_{(i)}$. For $1 \leq i \leq k$, let $u_i \in S_{k+1}$ be the k -Grassmannian permutation of shape (1^{k-i}) . So by Proposition 4.5 we have

$$\Phi_{k+1}(\mathcal{O}_{G/B}^{u_i}) = \frac{g_{(i)}}{\tau_k}.$$

Thus we have

$$\kappa'(\mathcal{O}_{(i)}^{\mathrm{Gr}} \cdot (\mathcal{O}_{R_k}^{\mathrm{Gr}})^{-1}) = \mathcal{O}_{G/B}^{u_i}.$$

From Lemma B.3, we have $\theta_k(u_i) = (i)$. Therefore $\mathcal{O}_{u_i t_{-\varpi_k^\vee}}^{\mathrm{Gr}} = \mathcal{O}_{(i)}^{\mathrm{Gr}}$ so by the definition of κ we have

$$\kappa(\mathcal{O}_{(i)}^{\mathrm{Gr}} \cdot (\mathcal{O}_{R_k}^{\mathrm{Gr}})^{-1}) = \mathcal{O}_{G/B}^{u_i}.$$

Because $(\mathcal{O}_{R_j}^{\mathrm{Gr}})^{-1}$ is an invertible element, we have $\kappa'(\mathcal{O}_{(i)}^{\mathrm{Gr}}) = \kappa(\mathcal{O}_{(i)}^{\mathrm{Gr}})$, and therefore $\kappa' = \kappa$.

For $w \in S_{k+1}$, we have $\mathcal{O}_{wt_{-\sum_{i \in \mathrm{Des}(w)} \varpi_i^\vee}}^{\mathrm{Gr}} = \mathcal{O}_{\theta_k(w)}^{\mathrm{Gr}}$ from Proposition 4.6, so

$$\begin{aligned} \mathcal{O}_{G/B}^w &= \kappa \left(\mathcal{O}_{\theta_k(x)}^{\mathrm{Gr}} \cdot \prod_{i \in \mathrm{Des}(w)} (\mathcal{O}_{R_i}^{\mathrm{Gr}})^{-1} \right) = \kappa' \left(\mathcal{O}_{\theta_k(x)}^{\mathrm{Gr}} \cdot \prod_{i \in \mathrm{Des}(w)} (\mathcal{O}_{R_i}^{\mathrm{Gr}})^{-1} \right) \\ &= \Phi_{k+1}^{-1} \left(\sigma^{-1}(g_{\theta_k(x)}) \cdot \prod_{i \in \mathrm{Des}(w)} \tau_i^{-1} \right) \end{aligned}$$

Thus we have (4.12). \square

4.5 Localizations of $K_*(\mathrm{Gr})$

The isomorphism conjectured by Lam, Li, Mihalcea, and Shimozono in [18] is different from Φ_{k+1} in the way of localization of $K_*(\mathrm{Gr})$. The localization of $K_*(\mathrm{Gr})$ in [18] can be identified with $\Lambda_{(k)}[\tau_i^{-1} \ (1 \leq i \leq k)]$, while our version is $\Lambda_{(k)}[\tau_i^{-1}, (\tau_i^+)^{-1} \ (1 \leq i \leq k)]$. The aim of this section is clarify a geometric meaning of $\tau_i^+ = \tilde{g}_{R_i} \in \Lambda_{(k)}$.

Lemma 4.8. $\tau_i^2 - \tau_{i-1} \tau_{i+1} = \tau_i^+ \cdot \tau_i^-$.

Proof. This is the discrete Toda equation given by Hirota [10]. We can show this by comparing the construction of Φ_{k+1} [11] and the Lax formalism for the discrete Toda equation given in [12, Section 1]. \square

Lemma 4.9. For $1 \leq i \leq k$, $\Phi_{k+1}(1 - Q_i) = \frac{\tau_i^+ \cdot \tau_i^-}{\tau_i^2}$.

Proof. This follows from Lemma 4.8. \square

Lemma 4.10. For $1 \leq i \leq k+1$

$$\tau_i = g_{R_i^*} + \tau_i^-.$$

Proof. It suffices to show $\sigma(\tau_i - g_{R_i^*}) = \tau_i$. This follows from (2.3), because R_i^* is the unique maximal proper element among the partitions $\mu \subset R_i$. \square

Proposition 4.11. For $1 \leq i \leq k$,

$$z_1 \cdots z_i = \frac{1}{1 - Q_i} (1 - \mathcal{O}_{G/B}^{s_i}).$$

Proof. By $\Phi_{k+1}(\mathcal{O}^{s_i}) = g_{R_i^*}/\tau_i$ ([11, Theorem 1.7]), and Lemma 4.10, the image of the right hand side is

$$\frac{\tau_i^2}{\tau_i^+ \cdot \tau_i^-} \left(1 - \frac{g_{R_i^*}}{\tau_i} \right) = \frac{\tau_i^2}{\tau_i^+ \cdot \tau_i^-} \left(\frac{\tau_i - g_{R_i^*}}{\tau_i} \right) = \frac{\tau_i}{\tau_i^+}.$$

On the other hand

$$\Phi_{k+1}(z_1 \cdots z_i) = \prod_{j=1}^i \frac{\tau_j \tau_{j-1}^+}{\tau_j^+ \tau_{j-1}} = \frac{\tau_i}{\tau_i^+}.$$

\square

Remark 4.12. The result corresponds to Corollary 3.33 in [25]. Note also that

$$1 - \mathcal{O}_{G/B}^{s_i} = \mathcal{O}_{G/B}(-\varpi_i).$$

The result is due to [30].

Corollary 4.13. The following element of $QK(G/B_{k+1})_{\text{loc}}$ is sent to $1/\tau_i^+$

$$Q_{t_{\varpi_i^\vee}} (1 - Q_i)^{-1} (1 - \mathcal{O}_{G/B}^{s_i})$$

by Φ_{k+1} , where $Q_{t_{\varpi_i^\vee}}$ is the element such that $\Phi_{k+1}(Q_{t_{\varpi_i^\vee}}) = \tau_i$.

Note the factor $Q_{t_{\varpi_i^\vee}}$ is invertible in $QK(G/B)_{\text{loc}}$. So up to an invertible factor, $1/\tau_i^+$ corresponds to the element $(1 - Q_i)^{-1} (1 - \mathcal{O}_{G/B}^{s_i})$ of $QK(G/B) \subset QK(G/B)_{\text{loc}}$.

A Parabolic quotient of Coxeter groups

We discuss some properties of a coset space of a Coxeter group. Our basic reference is Bjorner-Brenti [3]. Let (W, S) be a *Coxeter system* (see [3] for the definition), where W is the group generated by $S = \{s_i \mid i \in I\}$ with index set I . The *Bruhat order* on W (see [3, Chapter 2]) is denoted by \leq . Let J be any subset of I . Let W_J be the subgroup of W generated by s_i ($i \in J$). The *minimal coset representatives* W^J of the quotient W/W_J is defined to be $W^J := \{w \in W \mid ws_i > w \text{ for all } i \in J\}$. Any element w of W is expressed uniquely as $w = w^J w_J$, with $w_J \in W_J$ and $w^J \in W^J$ ([3, Proposition 2.4.4]).

The following result is well-known and used throughout this section.

Lemma A.1. Let $x \in W^J$, and $i \in I$. Then $s_i x < x \implies s_i x \in W^J$.

A.1 Proof of Proposition 3.13

The 0-Hecke algebra H_W is the associative \mathbb{C} -algebra generated by $\{T_i \mid i \in I\}$ with the same relations as W except $T_i^2 = -T_i$ in place of $s_i^2 = id$. For $w \in W$, define $T_w = T_{i_1} \cdots T_{i_m}$ for any reduced expression $w = s_{i_1} \cdots s_{i_m}$. The elements T_w ($w \in W$) form a basis of H_W .

Proposition A.2. Let V^J be a left H_W module given by

$$V^J = H_W e_J, \quad e_J := \sum_{w \in W_J} T_w.$$

Then $V^J = \bigoplus_{x \in W^J} \mathbb{C} a_x$ with $a_x := T_x e_J$, and for $i \in I$, we have

$$T_i \cdot a_x = \begin{cases} a_{s_i x} & (s_i x > x \text{ and } s_i x \in W^J) \\ -a_x & (s_i x < x) \\ 0 & (s_i x > x \text{ and } s_i x \notin W^J) \end{cases}. \quad (\text{A.1})$$

Proof. For $i \in J$, we will prove $T_i e_J = 0$. Let $X^+ = \{v \in W \mid s_i v > v\}$ and $X^- = \{v \in W \mid s_i v < v\}$. We have $s_i(X^\pm) = X^\mp$. If $v \in X^+$, then $T_i T_v = T_{s_i v}$. If $v \in X^-$, then $T_v = T_i T_{s_i v}$, and so $T_i T_v = T_i^2 T_{s_i v} = -T_i T_{s_i v} = -T_v$. Thus

$$T_i e_J = \sum_{v \in X^+} T_i T_v + \sum_{v \in X^-} T_i T_v = \sum_{v \in X^+} T_{s_i v} - \sum_{v \in X^-} T_v = 0.$$

If $v \notin W^J$, then it is easy to see that there is a reduced expression $v = s_{i_1} \cdots s_{i_m}$ with $i_m \in J$. Hence $T_v e_J = T_{s_{i_1} \cdots s_{i_{m-1}}} T_{i_m} e_J = 0$. It follows that $\{a_x\}_{x \in W^J}$ spans $V^J = H_W e_J$. Note that for $x \in W^J$ and $v \in W_J$, we have $T_x T_v = T_{xv}$ since $\ell(xv) = \ell(x) + \ell(v)$ ([3, Proposition 2.4.4 (2)]); the linear independence of $\{a_x\}_{x \in W^J}$ follows from this fact.

Let $x \in W^J$. If $s_i x > x$, then

$$T_i \cdot a_x = T_i \cdot T_x e_J = T_{s_i x} e_J = \begin{cases} a_{s_i x} & (s_i x \in W^J) \\ 0 & (s_i x \notin W^J) \end{cases}.$$

If $s_i x < x$, then $T_i \cdot a_x = T_i T_x e_J = -T_x e_J = -a_x$. \square

Proposition A.3. For $x \in W^J$, let $b_x := \sum_{y \leq x} a_y$. Set $D_i = T_i + 1$ ($i \in I$). Then

$$D_i \cdot b_x = \begin{cases} b_{s_i x} & (s_i x \in W^J, s_i x > x) \\ b_x & (\text{otherwise}) \end{cases}. \quad (\text{A.2})$$

Lemma A.4 (Z-lemma). Let $w, v \in W$ and $i \in I$. Suppose $s_i w > w$ and $s_i v > v$. Then the following conditions are equivalent:

$$(1) w \leq v, \quad (2) s_i w \leq s_i v, \quad (3) w \leq s_i v.$$

Proof. [28, Proposition 5.4.3]. \square

Lemma A.5. Let $x \in W^J$ and $i \in I$. We set

$$\begin{aligned} X_{\leq x}^+ &:= \{y \in W^J \mid y \leq x, s_i y \in W^J, s_i y > y\}, \\ X_{\leq x}^- &:= \{y \in W^J \mid y \leq x, s_i y \in W^J, s_i y < y\}, \\ X_{\leq x}^0 &:= \{y \in W^J \mid y \leq x, s_i y \notin W^J\}. \end{aligned}$$

- (1) If $s_i x \in W^J$, $s_i x > x$, then $s_i(X_{\leq x}^+) = X_{\leq s_i x}^-$, $X_{\leq x}^+ = X_{\leq s_i x}^+$, $X_{\leq x}^0 = X_{\leq s_i x}^0$.
- (2) If $s_i x \in W^J$, $s_i x < x$, then $s_i(X_{\leq x}^+) = X_{\leq x}^-$.

(3) If $s_i x \notin W^J$, then $s_i(X_{\leq x}^+) = X_{\leq x}^-$.

Proof. (1) and (2) follow immediately from Lemma A.4. (3) The inclusion $s_i(X_{\leq x}^+) \supset X_{\leq x}^-$ follows from Lemma A.4. We will show $s_i(X_{\leq x}^+) \subset X_{\leq x}^-$. Take arbitrary $z \in s_i(X_{\leq x}^+)$. Write $z = s_i y$ with $y \in X_{\leq x}^+$. Since $s_i z = s_i^2 y = y < s_i y = z$, it suffices to prove $s_i y \leq x$. From Lemma A.4, we have $s_i x \geq s_i y$. Since $s_i x \notin W^J$, we have $s_i x > x$ by Lemma A.1. Then there is $v \in W_J$ such that $s_i x = xv$ ([3, Corollary 2.5.2]). Hence we have $xv \geq s_i y$. Here note that $s_i y \in W^J$ as $y \in X_{\leq x}^+$. We have

$$s_i y = (s_i y)^J \leq (xv)^J = x^J = x;$$

here we used the fact that $w \leq v$ for $w, v \in W$ implies $w^J \leq v^J$ ([3, Proposition 2.5.1]). \square

Proof of Proposition A.3. We have $b_x = \sum_{y \in X_{\leq x}^+} a_y + \sum_{y \in X_{\leq x}^-} a_y + \sum_{y \in X_{\leq x}^0} a_y$. Using (A.1), it is straightforward to verify

$$D_i \cdot b_x = \sum_{y \in X_{\leq x}^+} a_{s_i y} + \sum_{y \in X_{\leq x}^+} a_y + \sum_{y \in X_{\leq x}^0} a_y.$$

Consider first the case when $s_i x \in W^J$, $s_i x > x$. From Lemma A.5 (1), we have

$$D_i \cdot b_x = \sum_{y \in X_{\leq s_i x}^-} a_y + \sum_{y \in X_{\leq s_i x}^+} a_y + \sum_{y \in X_{\leq s_i x}^0} a_y = b_{s_i x}.$$

Next consider the case when $s_i x \in W^J$, $s_i x < x$. We have $s_i(X_{\leq x}^+) = X_{\leq x}^-$ by Lemma A.5 (2), and so we have

$$D_i \cdot b_x = \sum_{y \in X_{\leq x}^-} a_y + \sum_{y \in X_{\leq x}^+} a_y + \sum_{y \in X_{\leq x}^0} a_y = b_x.$$

Finally we consider the case when $s_i x \notin W^J$. We have $s_i(X_{\leq x}^+) = X_{\leq x}^-$ by Lemma A.5 (3), so we have $D_i \cdot b_x = b_x$ by exactly the same reasoning as in the previous case. \square

Proof of Proposition 3.13. We apply Proposition A.2 to $W = \tilde{S}_{k+1} = \langle s_i \mid i = 0, 1, \dots, k \rangle$, $W_J = S_{k+1}$, $W^J = \tilde{S}_{k+1}^0$ with $J = \{1, 2, \dots, k\}$, and define an isomorphism $V^J \rightarrow \Lambda_{(k)}$ of vector spaces by $a_x \mapsto \mathfrak{g}_\lambda^{(k)}$, $b_x \mapsto \tilde{\mathfrak{g}}_\lambda^{(k)}$ under the bijection $W^J = \tilde{S}_{k+1}^0 \ni x \mapsto \lambda \in \mathcal{P}^k$. Then we obtain Proposition 3.13. \square

A.2 Proof of Lemma 3.18

For $w \in W$, and $i \in I$, we define $s_i * w$ in the same way as (2.4). Let $\mathcal{S} = (i_1, \dots, i_r)$ be a sequence of elements of I , and $w \in S$. We define

$$\mathcal{S} * w = s_{i_1} * (s_{i_2} * \dots * (s_{i_r} * w) \dots). \quad (\text{A.3})$$

Proposition A.6. Let $\mathcal{S} = (i_1, \dots, i_r)$ be a sequence of elements of I . Set $|\mathcal{S}| = r$, and $T_{\mathcal{S}} := T_{i_1} \dots T_{i_r} \in H_W$. Then for $x \in W^J$, we have

$$T_{\mathcal{S}} \cdot a_x = \begin{cases} (-1)^{|\mathcal{S}| - \ell(\mathcal{S} * x) + \ell(x)} a_{\mathcal{S} * x} & (\mathcal{S} * x \in W^J) \\ 0 & (\mathcal{S} * x \notin W^J) \end{cases}.$$

Proof. For $1 \leq p \leq r$, set $\mathcal{S}_p = (i_p, \dots, i_r)$, and $\mathcal{S}_{r+1} = \emptyset$. We use decreasing induction on p . If $p = r + 1$, then the assertion is obvious. Suppose $1 \leq p < r + 1$. We first consider the case when $\mathcal{S}_p * x \notin W^J$. Let $q (\geq p)$ be the maximal integer such that $\mathcal{S}_q * x \notin W^J$. Then, we have $\mathcal{S}_{q+1} * x \in W^J$, $\mathcal{S}_q * x = s_{i_q} * (\mathcal{S}_{q+1} * x) = s_{i_q}(\mathcal{S}_{q+1} * x) \notin W^J$, and so $s_{i_q}(\mathcal{S}_{q+1} * x) > \mathcal{S}_{q+1} * x$ by Lemma A.1. Therefore, $T_{i_q} \cdot a_{\mathcal{S}_{q+1} * x} = 0$. By the inductive hypothesis, we have $T_{i_{q+1}} \cdots T_{i_r} \cdot a_x = \pm a_{\mathcal{S}_{q+1} * x}$, and hence

$$T_{\mathcal{S}_p} \cdot a_x = T_{i_p} \cdots T_{i_q} (T_{i_{q+1}} \cdots T_{i_r} \cdot a_x) = \pm T_{i_p} \cdots T_{i_q} \cdot a_{\mathcal{S}_{q+1} * x} = 0.$$

Next we consider the case when $\mathcal{S}_p * x \in W^J$. Note that, in view of Lemma A.1, we have $\mathcal{S}_q * x \in W^J$ for $p \leq q \leq r + 1$. By the inductive hypothesis, we have $T_{i_{p+1}} \cdots T_{i_r} a_x = (-1)^{r-p-\ell(\mathcal{S}_{p+1} * x) + \ell(x)} \cdot a_{\mathcal{S}_{p+1} * x}$, so

$$T_{\mathcal{S}_p} \cdot a_x = T_{i_p} \cdot (T_{i_{p+1}} \cdots T_{i_r} a_x) = (-1)^{r-p-\ell(\mathcal{S}_{p+1} * x) + \ell(x)} T_{i_p} \cdot a_{\mathcal{S}_{p+1} * x}. \quad (\text{A.4})$$

Now we consider two cases : (a) $s_{i_p}(\mathcal{S}_{p+1} * x) > \mathcal{S}_{p+1} * x$, (b) $s_{i_p}(\mathcal{S}_{p+1} * x) < \mathcal{S}_{p+1} * x$. If (a) holds, then $s_{i_p}(\mathcal{S}_{p+1} * x) = s_{i_p} * (\mathcal{S}_{p+1} * x) = \mathcal{S}_p * x \in W^J$. Therefore

$$T_{i_p} \cdot a_{\mathcal{S}_{p+1} * x} = a_{s_{i_p}(\mathcal{S}_{p+1} * x)} = a_{\mathcal{S}_p * x},$$

and hence by (A.4),

$$T_{\mathcal{S}_p} \cdot a_x = (-1)^{r-p-\ell(\mathcal{S}_{p+1} * x) + \ell(x)} T_{i_p} a_{\mathcal{S}_{p+1} * x} = (-1)^{r-p-\ell(\mathcal{S}_{p+1} * x) + \ell(x)} a_{\mathcal{S}_p * x};$$

the sign is correct because we have $\ell(\mathcal{S}_{p+1} * x) = \ell(\mathcal{S}_p * x) - 1$. Suppose (b) holds. Note that $\mathcal{S}_p * x = \mathcal{S}_{p+1} * x \in W^J$ by (A.3). By the definition of T_{i_p} ,

$$T_{i_p} \cdot a_{\mathcal{S}_{p+1} * x} = -a_{\mathcal{S}_{p+1} * x} = -a_{\mathcal{S}_p * x}.$$

So we have

$$T_{\mathcal{S}_p} \cdot a_x = (-1)^{r-p-\ell(\mathcal{S}_{p+1} * x) + \ell(x)} (-a_{\mathcal{S}_p * x}) = (-1)^{r-p+1-\ell(\mathcal{S}_p * x) + \ell(x)} a_{\mathcal{S}_p * x},$$

where we again used $\mathcal{S}_p * x = \mathcal{S}_{p+1} * x$. This completes the proof. \square

Proof of Lemma 3.18. Let $A \subsetneq I$, with $|A| = r$, and take a reduced expression $s_{i_1} \cdots s_{i_r}$ of u_A . We apply Proposition A.6 to the sequence $\mathcal{S} = (i_1, \dots, i_r)$. Then $T_{\mathcal{S}} = T_{u_A}$ and we obtain Lemma 3.18. \square

B Grassmannian permutations

An explicit description of $\theta_k(w)$ is available (see [22, §6]). For the readers' convenience, we include a simple direct proof when w is a Grassmann element $w_{\lambda, i}$ (Lemma B.3).

Proposition B.1. Let $\lambda \in \mathcal{P}^k$ be a k -bounded partition of size r such that $\lambda_1 + \ell(\lambda) \leq k + 1$. Take any standard tableau T of shape λ . We denote the box of λ with entry i in T by $b_T(i)$. Then $s_{\text{res}(b_T(r))} \cdots s_{\text{res}(b_T(2))} s_{\text{res}(b_T(1))}$ is a reduced expression for x_λ .

Proof. We note first that the corresponding fact is well-known for i -Grassmannian permutation $w_{\lambda, i}$. The reader can consult [5, § 3.1] for an exposition of this fact in a more general setting.

Since $\lambda_1 + \ell(\lambda) \leq k + 1$, there is $1 \leq i \leq k$ such that $\lambda \subset R_{k+1-i}$. Then in the reduced expression of x_λ given by (2.1), s_{k+1-i} does not appear. So $x_\lambda \in \langle s_{-i+1}, \dots, s_{-i+k} \rangle \cong S_{k+1}$. An isomorphism of group $\phi : S_{k+1} \rightarrow \langle s_{-i+1}, \dots, s_{-i+k} \rangle$ is given by $\phi(s_j) = s_{-i+j}$. Then the result follows from the case of i -Grassmannian permutation. \square

Remark B.2. An element in a Coxeter group W is *fully commutative* if any two of its reduced expressions are related by a series of transpositions of adjacent commuting generators. It is well-known that any i -Grassmannian element in S_{k+1} is fully commutative. From the above proof, we see that if $\lambda_1 + \ell(\lambda) \leq k + 1$, then x_λ is fully commutative.

Lemma B.3. Let λ be a partition contained in $R_{k+1-i} = (k+1-i)^i$, and $w_{\lambda,i} \in S_{k+1}$ the corresponding i -Grassmannian permutation. Then $\theta_k(w_{\lambda,i}) = (\lambda^\vee)'$.

Proof. In the extended affine symmetric group \hat{S}_{k+1} , we compute $w_{\lambda,i} t_{-\varpi_i} = w_{\lambda,i} \pi^{-i} x_{R_i} = \pi^{-i} (\pi^i w_{\lambda,i} \pi^{-i}) x_{R_i}$. A reduced expression of $y_\lambda := \pi^i w_{\lambda,i} \pi^{-i}$ is obtained by replacing s_j by s_{j+i} in $w_{\lambda,i}$. It is straightforward to see when we reflect the tableau of y_λ along the line with a slope of 1, it fits inside the tableau with shape R_i filled with $(k+1)$ -residues adjusted to the south-east corner (see Example B.4 below). We can apply Proposition B.1 to the element x_{R_i} . If we take T to be the standard tableau with shape R_i that is obtained by filling positive integers into each row of $(\lambda^\vee)'$ left to right, taking rows from top to bottom, and then each column of the remaining boxes from top to bottom, taking columns from left to right. The obtained reduced expression of x_{R_i} shows $x_{R_i} = y_\lambda^{-1} \cdot x_{(\lambda^\vee)'}$, so $w_{\lambda,i} t_{-\varpi_i} = \pi^{-i} x_{(\lambda^\vee)'}$ showing $\theta_k(w_{\lambda,i}) = (\lambda^\vee)'$. \square

Example B.4. For $k = 6$, $i = 3$, and $\lambda = (3, 2)$, the corresponding 3-Grassmannian element is $w_{\lambda,3} = s_3 s_2 \cdot s_5 s_4 s_3$. We have

$$y_\lambda := \pi^3(w_{\lambda,3})\pi^{-3} = s_6 s_5 \cdot s_1 s_0 s_6 : \begin{array}{|c|c|c|} \hline 6 & 0 & 1 \\ \hline 5 & 6 & \\ \hline \end{array}.$$

We read the entries of the tableau with shape R_3 filled with the 7-residues $\begin{array}{|c|c|c|} \hline 0 & 1 & 2 \\ \hline 6 & 0 & 1 \\ \hline 5 & 6 & 0 \\ \hline 4 & 5 & 6 \\ \hline \end{array}$ according

to the order given by $T = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & 10 \\ \hline 6 & 8 & 11 \\ \hline 7 & 9 & 12 \\ \hline \end{array}$ to have

$$x_{R_3} = (s_6 s_0 s_1 \cdot s_5 s_6) \cdot s_4 s_5 s_0 s_6 s_2 s_1 s_0.$$

This is $y_\lambda^{-1} x_{(\lambda^\vee)'}$. The shaded boxes corresponds to y_λ .

C Dual Pieri rule for the closed K - k -Schur functions

A Pieri rule for $\tilde{g}_\lambda^{(k)}$ was proved by Takigiku [35]. We record here the dual version of Takigiku's formula, which should be known for experts but is missing from the literature. We do not use these result in the main part of this paper.

For $\lambda \in \mathcal{P}^k$, $1 \leq r \leq k$, define

$$\begin{aligned} \mathcal{H}_{\lambda,r}^{(k)} &:= \{A \subsetneq I : |A| = r, d_A x_\lambda \in \tilde{S}_{k+1}^0, d_A x_\lambda \geq_L x_\lambda\}, \\ \mathcal{V}_{\lambda,r}^{(k)} &:= \{A \subsetneq I : |A| = r, u_A x_\lambda \in \tilde{S}_{k+1}^0, u_A x_\lambda \geq_L x_\lambda\}. \end{aligned}$$

For an element A in $\mathcal{H}_{\lambda,r}^{(k)}$, we write $d_A x_\lambda = x_\kappa$ for an element $\kappa \in \mathcal{P}^k$, and denote this κ by $d_A \lambda$. Similarly, for an element A in $\mathcal{V}_{\lambda,r}^{(k)}$, we write $u_A x_\lambda = x_\kappa$ for an element $\kappa \in \mathcal{P}^k$, and denote this κ by $u_A \lambda$.

Example C.1. For $k = 3$, $\lambda = (2, 1) \in \mathcal{P}^3$, and $r = 2$, we have (see Figure 2)

$$\begin{aligned} \mathcal{H}_{\lambda,r}^{(k)} &= \{\{2, 3\}, \{0, 2\}\}, & d_{\{2,3\}} &= s_3 s_2, & d_{\{0,2\}} &= s_0 s_2 = s_2 s_0, \\ \mathcal{V}_{\lambda,r}^{(k)} &= \{\{1, 2\}, \{0, 2\}\}, & u_{\{1,2\}} &= s_1 s_2, & u_{\{0,2\}} &= s_0 s_2 = s_2 s_0. \end{aligned}$$

The corresponding weak (horizontal and vertical) strips are given by

$$d_{\{2,3\}}\lambda = (3, 1, 1), \quad d_{\{0,2\}}\lambda = u_{\{0,2\}}\lambda = (2, 1, 1), \quad u_{\{1,2\}}\lambda = (2, 1, 1, 1).$$

For $A_1, \dots, A_m \in \mathcal{H}_{\lambda, r}^{(k)}$, it is known by Takigiku [35, Corollary 4.8] that $A_1 \cap \dots \cap A_m \in \mathcal{H}_{\lambda, r'}^{(k)}$, with $r' = |A_1 \cap \dots \cap A_m|$ so $d_{A_1 \cap \dots \cap A_m} \lambda \in \mathcal{P}^k$ is defined.

Proposition C.2. Let $A \mapsto \overline{A}$ be the map given by sending $i \in A$ to $-i \in \overline{A}$. Then for $\lambda \in \mathcal{P}^k$, and $1 \leq r \leq k$,

$$\overline{\mathcal{H}_{\lambda, r}^{(k)}} = \mathcal{V}_{\lambda^{\omega_k}, r}^{(k)}.$$

Proof. Let $A \in \mathcal{H}_{\lambda, r}^{(k)}$. Recall that ω_k is an automorphism the group \tilde{S}_{k+1} preserving the left weak order,

$$u_{\overline{A}} x_{\lambda^{\omega_k}} = \omega_k(d_A) \omega_k(x_\lambda) = \omega_k(d_A x_\lambda) \geq_L \omega_k(x_\lambda) = x_{\lambda^{\omega_k}}$$

so $\overline{A} \in \mathcal{V}_{\lambda^{\omega_k}, r}^{(k)}$. Thus $\overline{\mathcal{H}_{\lambda, r}^{(k)}} \subset \mathcal{V}_{\lambda^{\omega_k}, r}^{(k)}$. Similarly we have $\overline{\mathcal{H}_{\lambda, r}^{(k)}} \supset \mathcal{V}_{\lambda^{\omega_k}, r}^{(k)}$. \square

Theorem C.3 ([34]). Let $\lambda \in \mathcal{P}^k$, and $1 \leq r \leq k$. Let

$$\mathcal{H}_{\lambda, r}^{(k)} = \{A_1, \dots, A_m\}, \quad \mathcal{V}_{\lambda, r}^{(k)} = \{B_1, \dots, B_n\}.$$

Then

$$\begin{aligned} \tilde{g}_{(r)} \cdot \tilde{g}_\lambda^{(k)} &= \sum_{i=1}^m (-1)^{i-1} \sum_{1 \leq a_1 < \dots < a_i \leq m} \tilde{g}_{d_{A_{a_1} \cap \dots \cap A_{a_i}} \lambda}^{(k)}, \\ \tilde{g}_{(1^r)} \cdot \tilde{g}_\lambda^{(k)} &= \sum_{i=1}^n (-1)^{i-1} \sum_{1 \leq a_1 < \dots < a_i \leq n} \tilde{g}_{u_{B_{a_1} \cap \dots \cap B_{a_i}} \lambda}^{(k)}. \end{aligned}$$

Proof. (C.1) is due to Takigiku [34]. From Proposition C.2, we have $\{\overline{A_1}, \dots, \overline{A_m}\} = \mathcal{V}_{\lambda^{\omega_k}, r}^{(k)}$. By applying Ω to both hand sides, we have

$$\tilde{g}_{(1^r)} \cdot \tilde{g}_{\lambda^{\omega_k}}^{(k)} = \sum_{i=1}^m (-1)^{i-1} \sum_{1 \leq a_1 < \dots < a_i \leq m} \tilde{g}_{u_{\overline{A_{a_1} \cap \dots \cap A_{a_i}}} \lambda^{\omega_k}}^{(k)},$$

where we used $\Omega(\tilde{g}_{(r)}) = \tilde{g}_{(1^r)}$ and

$$\Omega(\tilde{g}_{d_{A_{a_1} \cap \dots \cap A_{a_i}} \lambda}^{(k)}) = \tilde{g}_{u_{\overline{A_{a_1} \cap \dots \cap A_{a_i}}} \lambda^{\omega_k}}^{(k)} = \tilde{g}_{u_{\overline{A_{a_1} \cap \dots \cap A_{a_i}}} \lambda^{\omega_k}}^{(k)}.$$

Thus we have (C.1). \square

Example C.4. According to Example C.1, we have

$$\begin{aligned} \tilde{g}_{(2)}^{(3)} \tilde{g}_{(2,1)}^{(3)} &= \tilde{g}_{(3,1,1)}^{(3)} + \tilde{g}_{(2,2,1)}^{(3)} - \tilde{g}_{(2,1,1)}^{(3)}, \\ \tilde{g}_{(1,1)}^{(3)} \tilde{g}_{(2,1)}^{(3)} &= \tilde{g}_{(2,1,1,1)}^{(3)} + \tilde{g}_{(2,2,1)}^{(3)} - \tilde{g}_{(2,1,1)}^{(3)}. \end{aligned}$$




Figure 2: $k = 3$, $\lambda = (2, 1)$, $r = 2$.

References

- [1] D. Anderson, L. Chen, H.-H. Tseng (with Appendix by H. Iritani), On the Finiteness of Quantum K -Theory of a Homogeneous Space, *Int. Math. Res. Not.* Vol. **2022**, 1313–1349.
- [2] R. Bezrukavnikov, M. Finkelberg, I. Mirković, Equivariant homology and K -theory of affine Grassmannians and Toda lattices, *Compos. Math.* **141** (2005), 746–768.
- [3] A. Bjorner and F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics vol. 231, Springer (2005).
- [4] J. Blasiak, J. Morse, and G. H. Seelinger, K -theoretic Catalan functions (2020), preprint. arXiv:2010.01759
- [5] A. S. Buch, P.-E. Chaput, L. C. Mihalea, N. Perrin, A Chevalley formula for the equivariant quantum K -theory of cominuscule varieties, *Alg. Geom.* **5** (2018), 568–595.
- [6] C. H. Chow and N. C. Leung, Quantum K -theory of G/P and K -homology of affine Grassmannian, preprint arXiv:2201.12951.
- [7] I. Ciocan-Fontanine, W. Fulton, Quantum double Schubert polynomials, Appendix J in *Schubert Varieties and Degeneracy Loci* by W. Fulton and P. Pragacz, Lecture Notes in Math 1689 (1998), 134–138.
- [8] A. J. Dalal, J. Morse, Quantum and affine Schubert calculus and Macdonald polynomials, *Adv. Math.* **312** (2017), 425–458.
- [9] A. Givental, Y.-P. Lee, Quantum K -theory on flag manifolds, finite-difference Toda lattices and quantum groups, *Invent. Math.* **151** (2003), 193–219.
- [10] R. Hirota, Nonlinear Partial Difference Equations. II. Discrete-Time Toda Equation, *J. Phys. Soc. Jap.* **43** (1977), 2074–2078.
- [11] T. Ikeda, S. Iwao, and T. Maeno, Peterson isomorphism in K -theory and Relativistic Toda lattice, *Int. Math. Res. Not.* **2020** (2020), 6421–6462.
- [12] S. Iwao, H. Nagai, The discrete Toda equation revisited:dual β -Grothendieck polynomial, ultradiscretization and static soliton, *J. Phys. A: Math. Theor.* **51**, 134002.
- [13] S. Kato, Loop structure on equivariant K -theory of semi-infinite flag manifolds, preprint arXiv:1805.01718
- [14] B. Kim, Quantum Cohomology of Flag Manifolds G/B and Quantum Toda Lattices, *Ann. Math.* **149** (1999), 129–148.
- [15] A. Kirillov, T. Maeno, A note on quantum K -theory of flag varieties, in preparation.
- [16] A. N. Kirillov, T. Maeno, Quantum double Schubert polynomials, quantum Schubert polynomials and the Vafa-Intriligator formula, *Discrete Math.* **217** (200), 191–223.

- [17] T. Lam, L. Lapointe, J. Morse, A. Schilling, M. Shimozono, M. Zabrocki, *k-Schur Functions and Affine Schubert Calculus*, Fields Institute Monographs 33, Springer (2016).
- [18] T. Lam, C. Li, L. C. Mihalea, M. Shimozono, A conjectural Peterson isomorphism in K -theory, *J. Alg.* **513** (2018) 326–343.
- [19] T. Lam, A. Schilling, and M. Shimozono, K -theory Schubert calculus of the affine Grassmannian, *Compos. Math.* **146** (2010) 811–852.
- [20] T. Lam, A. Schilling, M. Shimozono, Schubert polynomials for the affine Grassmannian of the symplectic group, *Math. Z.* **264** (2010), 765–811.
- [21] T. Lam, M. Shimozono, From double quantum Schubert polynomials to k -double Schur functions via the Toda lattice, Preprint arXiv:1109.2193v1
- [22] T. Lam, M. Shimozono, From quantum Schubert polynomials to k -Schur functions via the Toda lattice, *Math. Res. Lett.* 19 (2012), no. 01, 81–93.
- [23] T. Lam, M. Shimozono, k -double Scour functions and equivariant (co)homology of the affine Grassmannian, *Math. Ann.* **356** (2013), 1379–1404.
- [24] L. Lapointe, J. Morse, Tableaux on $k+1$ -cores, reduced words for affine permutations, and k -Schur expansions, *J. Combinat. Theory, Ser. A* **112** (2005) 44–81.
- [25] C. Lenart, T. Maeno, Quantum Grothendieck polynomials, preprint arXiv:068232
- [26] C. Lenart, S. Naito, D. Sagaki, A general Chevalley formula for semi-infinite flag manifolds and quantum K -theory, preprint arXiv:2010.06143
- [27] C. Lenart, A. Postkinov, Affine Weyl Groups in K -Theory and Representation Theory, *Int. Math. Res. Not. Volume* **2007** (2007), rnm038.
- [28] R. V. Moody and A. Pianzola, *Lie Algebras with Triangular Decompositions* (Wiley-Interscience and Canadian Mathematics Series of Monographs and Texts)
- [29] J. Morse, Combinatorics of the K -theory of affine Grassmannians, *Adv. Math.* 229 (2012) 2950–2984.
- [30] S. Naito, D. Orr, and D. Sagaki, Chevalley formula for anti-dominant weights in the equivariant K -theory of semi-infinite flag manifolds, *Adv. Math.* **387** (2021) 107828.
- [31] S. Pon, Affine Stanley symmetric functions for classical types, *J. Algebr. Comb.* **36** (2012), 595–622.
- [32] G. H. Seelinger, K -theoretic Catalan functions, Dissertation, The university of Virginia, 2021.
- [33] J. Stembridge, A short derivation of the Möbius function for the Bruhat order, *J. Algebr. Comb.* **25** (2007), 144–148.
- [34] M. Takigiku, A Pieri formula and a factorization formula for sums of K -theoretic k -Schur functions, *Algebr. Comb.* **2** (2019), no. 4, 447–480.
- [35] M. Takigiku, On the Pieri rules of stable and dual stable Grothendieck, preprint arXiv:1806.06369