

ON VIRTUAL EMBEDDINGS OF BRAID GROUPS INTO MAPPING CLASS GROUPS OF SURFACES

TAKUYA KATAYAMA AND ERIKA KUNO

ABSTRACT. In this article, we give a necessary and sufficient condition for embedding a finite index subgroup of Artin's braid group into the mapping class group of a connected orientable surface.

1. INTRODUCTION

Let $S_{g,p}^b$ be a connected orientable surface of genus g with p punctures and b boundary components. In the case where $b = 0$ or $p = 0$, we drop the suffix that denotes 0, excepting g , from $S_{g,p}^b$. For example, $S_{g,p}^0$ will simply be denoted as $S_{g,p}$. The *mapping class group* $\text{Mod}(S_{g,p}^b)$ of $S_{g,p}^b$ is the group of orientation-preserving homeomorphisms of $S_{g,p}^b$, fixing the boundary pointwise, up to isotopy relative to the boundary. We write B_n for the *braid group* on n strands, which is identified with $\text{Mod}(S_{0,n}^1)$. We define the Euler characteristic of $S_{g,p}^b$ as

$$\chi_{g,p}^b := 2 - 2g - p - b.$$

We say that a group H is *virtually embedded* in a group G if H has a finite index subgroup that injects into G . An injective homomorphism from a finite index subgroup of H to G is called a *virtual embedding* of H (into G). In this paper we give a necessary and sufficient condition for the existence of virtual embeddings of braid groups into the mapping class groups of orientable surfaces.

Theorem 1.1. *Suppose $b = 0$. Then the braid group B_n is virtually embedded in $\text{Mod}(S_{g,p})$ if and only if*

$$n \leq \begin{cases} 1 & ((g,p) \in \{(0,0), (0,1), (0,2), (0,3)\}) \\ 2 & ((g,p) \in \{(0,4), (1,0), (1,1)\}) \\ -\chi_{g,p} & (g=0, p \geq 5) \\ 2 - \chi_{g,p} & (g \geq 2, p=0) \\ 1 - \chi_{g,p} & (\text{otherwise}). \end{cases}$$

Theorem 1.2. *Suppose $b \geq 1$. Then B_n is virtually embedded in $\text{Mod}(S_{g,p}^b)$ if and only if*

$$n \leq \begin{cases} 2 - \chi_{g,p}^b & (g \geq 1, p+b \leq 2) \\ 1 - \chi_{g,p}^b & (\text{otherwise}). \end{cases}$$

Date: March 29, 2022.

2020 Mathematics Subject Classification. 20F36, 20F65, 57K20.

Key words and phrases. Mapping class group; braid group; virtual embeddings; right-angled Artin groups.

If a group L is virtually embedded in a group H and H is virtually embedded in a group G , then L is virtually embedded in G . Note as well that virtual embeddings are closely related to homomorphisms with finite kernel. In fact, the mapping class group $\text{Mod}(S)$ is virtually embedded in a group G if and only if there is a homomorphism with finite kernel from a finite index subgroup of $\text{Mod}(S)$ to G , because $\text{Mod}(S)$ is virtually torsion-free by Serre's theorem. We are often interested in the *pure mapping class group*, denoted by $\text{PMod}(S_{g,p}^b)$, which is a subgroup of $\text{Mod}(S_{g,p}^b)$ and consists of the mapping classes fixing the punctures. We denote the *pure braid group* $\text{PMod}(S_{0,n}^1)$ by PB_n ; this is an index $n!$ subgroup of B_n .

The study of injective homomorphisms between the mapping class groups of surfaces has been extensively developed by various researchers. Aramayona–Leininger–Souto [1], Birman–Hilden [5] and Paris–Rolfsen [17] gave injective homomorphisms induced by inclusion maps and (possibly branched) coverings of surfaces in an ingenious manner. Aramayona–Souto [2] proved that $\text{PMod}(S_{g,p}^b)$ cannot be embedded in $\text{PMod}(S_{g',p'}^{b'})$ if $g \geq 6$ and either $g' \leq 2g-2$ or $g' = 2g-1$ and $p'+b' \geq 1$. Ivanov–McCarthy [11] proved that every injective homomorphism between the mapping class groups of surfaces with almost the same topological complexity is necessarily an isomorphism. Shackleton [19] proved that Ivanov–McCarthy's result is also true for virtual embeddings. The topological complexity of surfaces [6] and virtual cohomological dimension [10] are known to be obstructions to the existence of virtual embeddings. It follows from these results that, apart from a few exceptions, two mapping class groups are abstractly commensurable if and only if the defining surfaces are homeomorphic. Moreover, the rigidity result of the mapping class groups due to Behrstock–Kleiner–Minsky–Mosher [4] and Hamenstädt [9] implies that two mapping class groups are quasi-isometric if and only if they are abstractly commensurable. Therefore, finite index subgroups of the mapping class groups of surfaces reflect the homeomorphism types of the surfaces. As for braid groups, Castel [7] characterized homomorphisms from braid groups to the mapping class groups of surfaces without punctures and proved that every injective homomorphism is a transvection of a geometric monodromy.

Remark 1.3. From Castel's work [7], it follows that B_{2g+3} is not embedded in $\text{Mod}(S_g^b)$ for any $g \geq 0$ and $b \geq 0$. On the other hand, Theorem 1.2 says that B_{2g+3} has a finite index subgroup embedded in $\text{Mod}(S_g^b)$ if $b \geq 4$. Therefore, if $b \geq 4$, then there is a virtual embedding of B_{2g+3} into $\text{Mod}(S_g^b)$ that does not extend to an embedding of the entire group B_{2g+3} .

Recently, Chen–Mukherjea in [15] proved that any non-trivial homomorphism from a braid group to the pure mapping class groups of a surface with punctures is a transvection of a geometric monodromy if the genus of the surface lies in a certain range.

This paper is organized as follows. In Section 2, we review the embeddings between the mapping class groups of surfaces that have been obtained from the Birman–Hilden theory and Paris–Rolfsen's result. We then introduce the concept of “pseudo-annular extensions” of surfaces and prove that “pseudo-annular extensions” of spheres induce virtual embeddings of the mapping class groups of the spheres. These virtual embeddings of the mapping class groups of surfaces enable us to prove the “if part” of Theorems 1.1 and 1.2 in the last section of this paper. On the other hand, the “only if part” of Theorems 1.1 and 1.2 will be proved by

comparing right-angled Artin subgroups in the mapping class groups of surfaces. In Sections 3 and 4, we determine whether a right-angled Artin group A of specific type is embedded in the mapping class group of a surface S . More precisely, in Section 3, we decide whether the defining graph of A and its companions are full subgraphs of the 1-skeleton of the Harvey's curve complex of S . In Section 4, we prove that A is embedded in the mapping class group of S if and only if the defining graph of A is a full subgraph of the 1-skeleton of Harvey's the curve complex of S (Remark 4.9). The last section of this paper is devoted to proving Theorems 1.1 and 1.2.

Acknowledgements. The authors are grateful to Donggyun Seo and Dan Margalit for helpful comments. The first author was supported by JSPS KAKENHI through a grant, number 20J01431. The second author was supported by JSPS KAKENHI Grant-in-Aid for Young Scientists, number 21K13791.

2. CONSTRUCTIONS OF VIRTUAL EMBEDDINGS

In this section, we explain how to obtain virtual embeddings of braid groups into the mapping class groups of surfaces. Our constructions of the virtual embeddings are based on extensions and coverings of surfaces.

First, let us review homomorphisms induced by embeddings of surfaces. Let $i: S \rightarrow F$ be an inclusion map between two connected orientable surfaces of finite type. We call i an *extension* of S for convenience sake. The extension i is said to be *admissible* if $i(S)$ is a closed subset of F and no component of $i(\partial S)$ is parallel to a component of ∂F . Additionally, we say that an admissible extension i is *annular* if $F \setminus \text{Int}(i(S))$ is a disjoint union of annuli.

Proposition 2.1. *Suppose that $i: S \rightarrow F$ is an admissible extension.*

- (1) *If the extension i is annular, then the kernel of the induced homomorphism $\text{Mod}(S) \rightarrow \text{Mod}(F)$ is a free abelian group generated by*

$$\mathcal{A} = \{[T_{c_1}][T_{c_2}]^{-1} \mid c_1, c_2 \text{ are the boundary components of an outer annulus}\}.$$

- (2) *If $F \setminus \text{Int}(i(S))$ is a disjoint union of annuli and once-punctured disks, then the kernel of the induced homomorphism $\text{Mod}(S) \rightarrow \text{Mod}(F)$ is a free abelian group generated by \mathcal{A} and*

$$\{[T_c] \mid c \text{ is the boundary component of an outer once-punctured disk}\}.$$

Here, T_c is a Dehn twist about a curve c .

When each component of the exterior of an admissible extension is sufficiently complicated, the extension induces an embedding of the mapping class group.

Proposition 2.2. *Suppose that $F \setminus i(S)$ is a disjoint union of surfaces of negative Euler characteristics. Then $\text{Mod}(S) \hookrightarrow \text{Mod}(F)$.*

Propositions 2.1 and 2.2 were proved by Paris–Rolfsen in [17]. We say that an admissible extension i is *hyperbolic* if i satisfies the assumption in Proposition 2.2. The embedding given in the following corollary is induced by a hyperbolic extension $S_{0,n}^1 \rightarrow S_{0,m}^1$.

Corollary 2.3. *If $n \leq m$, then $B_n \hookrightarrow B_m$.*

Now, let us recall some injective homomorphisms induced by coverings of surfaces. Consider the double-branched covering $S_{g-1,0}^2 \rightarrow S_{0,2g}^1$ induced by the hyperelliptic involution of $S_{g-1,0}^2$. According to Birman–Hilden theory, it follows that B_{2g} is embedded in $\text{Mod}(S_{g-1}^2)$ as the symmetric mapping class subgroup with respect to the hyperelliptic involution. By gluing S_{g-1}^2 and $S_{0,p}^{b+2}$ along their boundaries, we obtain an admissible extension $S_{g-1}^2 \rightarrow S_{g,p}^b$. This extension is either annular or hyperbolic, thereby inducing a homomorphism $\text{Mod}(S_{g-1}^2) \rightarrow \text{Mod}(S_{g,p}^b)$, whose restriction to the subgroup B_{2g} is injective. Therefore, B_{2g} is embedded in $\text{Mod}(S_{g,p}^b)$. For more information about Birman–Hilden theory, see [5] and [16].

Proposition 2.4. *Suppose $g \geq 1$, $p \geq 0$ and $b \geq 0$. Then $B_{2g} \hookrightarrow \text{Mod}(S_{g,p}^b)$.*

Similarly, from the double-branched covering $S_g^1 \rightarrow S_{0,2g+1}^1$, we obtain $B_{2g+1} \hookrightarrow \text{Mod}(S_g^1)$. The target of this embedding can be extended as follows.

Proposition 2.5. *Suppose $g \geq 1$ and $b + p \geq 2$. Then $B_{2g+1} \hookrightarrow \text{Mod}(S_{g,p}^b)$.*

Proof. Since $b + p \geq 2$, an extension $S_g^1 \rightarrow S_{g,p}^b$ is hyperbolic. Hence, we have $B_{2g+1} \hookrightarrow \text{Mod}(S_g^1) \hookrightarrow \text{Mod}(S_{g,p}^b)$. \square

In the case where $g \geq 1$, $b = 1$ and $p = 1$, we can find a larger braid group in $\text{Mod}(S_{g,1}^1)$.

Proposition 2.6. *Suppose $g \geq 1$. Then $B_{2g+2} \hookrightarrow \text{Mod}(S_{g,1}^1)$.*

Proof. We use the embedding $B_{2g+2} \hookrightarrow \text{Mod}(S_g^2)$ induced by the covering described above and the restriction of the capping homomorphism $\varphi: \text{Mod}(S_g^2) \rightarrow \text{Mod}(S_{g,1}^1)$ to B_{2g+2} . Since no non-trivial element of $\text{Ker}(\varphi)$ preserves the fiber of a point in $S_{0,2g+1}^1$, the restriction is injective. \square

The following concept is fairly important for finding the largest virtual embeddings of braid groups into mapping class groups.

Definition 2.7. Let S be a surface with punctures and F a surface. By \bar{S} , we denote the compactification of S that has circles at infinity (if $S \cong S_{g,p}^b$, then $\bar{S} \cong S_{g,p}^{b+p}$). An extension $i: S \rightarrow F$ is said to be *pseudo-annular* if i is obtained from a disjoint union of \bar{S} and copies of an annulus and a once-punctured disk by gluing each annulus A (resp. once-punctured disk D) to \bar{S} along their boundaries so that one boundary component of A (resp. the boundary of D) is identified with a component of $\bar{S} - S$ and that another boundary component of A is either identified with a component of $\partial\bar{S}$ or is unattached. Note that no pseudo-annular extension is admissible, because its image is not a closed subset. However, $\bar{S} \subset F$ is admissible.

The pure mapping class group of a sphere is naturally embedded in the mapping class group of its compactification as follows.

Lemma 2.8. *Suppose $p + b \geq 3$. Then $\text{PMod}(S_{0,p}^b) \times \mathbb{Z}^p \cong \text{PMod}(S_0^{p+b})$. Moreover, each standard generator of the direct factor \mathbb{Z}^p corresponds to a Dehn twist about a closed curve parallel to a boundary component of S_0^{p+b} .*

Proof. See [8, Theorem 10]. \square

A pseudo-annular extension of a sphere induces an embedding of the pure mapping class group of the sphere.

Proposition 2.9. *Let $S \rightarrow F$ be a pseudo-annular extension. If S is a sphere, then $\text{PMod}(S)$ is embedded in $\text{PMod}(F)$.*

Proof. The admissible extension $\bar{S} \subset F$ induces an embedding $\phi: \text{Homeo}(\bar{S}) \hookrightarrow \text{Homeo}(F)$. Then ϕ gives a well-defined homomorphism $\varphi: \text{Mod}(\bar{S}) \rightarrow \text{Mod}(F)$ whose kernel is generated by Dehn twists corresponding to the attached once-punctured disks and products of Dehn twists corresponding to the attached annuli. Since S is a sphere, Lemma 2.8 implies that $\text{PMod}(S) \hookrightarrow \text{Mod}(\bar{S})$. Since $S \rightarrow F$ is pseudo-annular, Proposition 2.1 together with Lemma 2.8 implies that the intersection of $\text{Ker}(\varphi)$ and $\text{PMod}(S)$ is trivial. Therefore, the restriction of φ to $\text{PMod}(S)$ is injective. \square

3. CYCLIC CHAINS ON SURFACES

In this section we investigate combinatorial obstructions, contained in the 1-skeleta of Harvey's curve complexes, to embedding certain right-angled Artin groups into the mapping class groups of surfaces. More precisely, we give a necessary and sufficient condition for embedding C_m^c , $C_m^c * K_1$ and some graphs similar to these into $\mathcal{C}(S)$ as "full subgraphs"; see Lemmas 3.27–3.32. Here, C_m is a cyclic graph of length m (the geometric realization is homeomorphic to the unit circle), Γ^c is the complement graph of a given graph Γ , K_m is a complete graph on m vertices, and $\mathcal{C}(S)$ is the curve graph of a surface S that is the 1-skeleton of Harvey's curve complex.

An arc or closed curve on a surface is called *simple* if it has no self-intersection. An arc δ on a surface S is *properly embedded* if $\partial\delta \subseteq \partial S$ and δ is a transversal to ∂S . A properly embedded arc δ on S is called *essential* if it is not isotopic to ∂S . A simple closed curve α on S is called *essential* if it does not bound a disk and it is not isotopic to a boundary component of S . For short, we use will the term arc (resp. closed curve) to mean a simple, properly embedded and essential arc (resp. closed curve) unless otherwise noted.

Definition 3.1. A sequence $(\alpha_1, \dots, \alpha_m)$ of closed curves on a surface S is called a *semi-cyclic chain* if any two consecutive curves α_i and α_{i+1} are in minimal position and not disjoint ($i = 1, \dots, m \bmod m$). If $(\alpha_1, \dots, \alpha_m)$ is a semi-cyclic chain, we call m its *length*. We say that a semi-cyclic chain $(\alpha_1, \dots, \alpha_m)$ on a surface has a *chord* if there are non-consecutive curves α_i and α_j that are in minimal position and not disjoint ($j \neq i, j \neq i \pm 1 \bmod m$). A semi-cyclic chain without a chord is called a *cyclic chain*. Note that if $(\alpha_1, \dots, \alpha_m)$ is a cyclic chain, then every length $(m-1)$ subsequence of the form $(\alpha_{i+1}, \dots, \alpha_m, \alpha_1, \dots, \alpha_{i-1})$ is a linear chain ($i = 1, \dots, m \bmod m$) as defined in [12, Section 2].

Let Λ and Γ be graphs. We denote the vertex set of the graph Λ (resp. the edge set of Λ) by $V(\Lambda)$ (resp. $E(\Lambda)$). We use the term map, from a graph Λ to a graph Γ , to mean a map from $V(\Lambda)$ to $V(\Gamma)$. A map $\phi: \Lambda \rightarrow \Gamma$ is called a (*graph*) *homomorphism* if it is a product map from $\Lambda = (V(\Lambda), E(\Lambda))$ to $\Gamma = (V(\Gamma), E(\Gamma))$ such that $\phi([v_1, v_2]) = [\phi(v_1), \phi(v_2)]$ for all pair of adjacent vertices v_1, v_2 of Λ . Here, we use the notation $[\bullet, \bullet]$ to mean the edge joining two vertices in the bracket. A (*graph*) *embedding* is an injective graph homomorphism. The image of a graph embedding is called a *subgraph*. A map $\phi: \Lambda \rightarrow \Gamma$ is said to be *full* if it is a product map from Λ to Γ such that $\phi([v_1, v_2]) = [u_1, u_2]$ for all pairs of adjacent vertices u_1, u_2 of Γ and for all $v_1 \in \phi^{-1}(u_1)$ and $v_2 \in \phi^{-1}(u_2)$. We write $\Lambda \leq \Gamma$ if there

is a full embedding of Λ into Γ . In this case, both Λ and the image of the full embedding are called a *full subgraph* of Γ . Note that any full subgraph of a graph is uniquely determined by its vertex set. For a subset $G \subset \Gamma$, the *full subgraph induced by G* is the full subgraph of Γ , whose vertex set is identical to that of G .

By definition, we have the following.

Proposition 3.2. *There is a cyclic chain of length m on a surface S of negative Euler characteristic if and only if $C_m^c \leq \mathcal{C}(S)$.*

For the purpose of proving the non-existence of cyclic chains on surfaces, we introduce the following concept.

Definition 3.3. Let δ_a and δ_b be arcs on a surface S , and $(\alpha_1, \dots, \alpha_m)$ a linear chain on S . A triple $(\delta_a, (\alpha_1, \dots, \alpha_m), \delta_b)$ is called *chained* if the triple satisfies the following conditions:

- (1) δ_a intersects α_1 non-trivially in minimal position but is disjoint from $\alpha_2, \dots, \alpha_m$ and δ_b .
- (2) δ_b intersects α_m non-trivially in minimal position but is disjoint from $\alpha_1, \dots, \alpha_{m-1}$ and δ_a .

We call m the *length* of the chained triple. We say that an arc on S is *recursive* if the arc joins a single boundary component of S to itself. Note that a separating arc on a surface (i.e., the surface minus the arc is not connected) is recursive. A chained triple $(\delta_a, (\alpha_1, \dots, \alpha_m), \delta_b)$ on a surface S satisfies condition $(*)$ if the following holds: for any boundary component C of S , we have either $\delta_a \cap C = \emptyset$ or $\delta_b \cap C = \emptyset$.

We will frequently use the following two lemmas.

Lemma 3.4. *If δ is an arc on S_g^p , then*

$$S_g^p \setminus \text{Int}N(\delta) \cong \begin{cases} S_{g-1}^{p+1} & (\delta : \text{non-separating}), \\ S_{g_1}^{p_1} \sqcup S_{g_2}^{p_2} & (\delta : \text{separating}), \end{cases}$$

where g_1, p_1, g_2 and p_2 are natural numbers satisfying

- $g_1 + g_2 = g, g_1 \geq 0, g_2 \geq 0,$
- $p_1 + p_2 = p + 1, p_1 \geq 1, p_2 \geq 1,$
- if $g_i = 0$, then $p_i \geq 2$ ($i = 1, 2$).

Lemma 3.5. *Let $\mathcal{T} = (\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ be a chained triple on S_g^p . Suppose that S' is a connected component of $S_g^p \setminus \text{Int}N(\delta_2)$, which contains a (possibly not chained) triple $\mathcal{T}' = (\delta_1, (\alpha_1, \dots, \alpha_{m-1}), \delta'_2)$, where δ'_2 is a connected component of $\alpha_m \cap S'$ intersecting α_{m-1} . Then, we have the following.*

- (1) *If there is a closed curve α_i which is parallel to a boundary component of S' , then $i = 1$ and $m = 2$.*
- (2) *If $m \geq 3$, then $\mathcal{T}' = (\delta_1, (\alpha_1, \dots, \alpha_{m-1}), \delta'_2)$ is a chained triple on S' .*
- (3) *If δ_1 is recursive and δ_2 is either separating (i.e., $S_g^p \setminus \text{Int}N(\delta_2)$ is not connected) or non-separating and not recursive on S_g^p , then δ_1 is recursive on S' .*
- (4) *If δ_1 is either separating or non-separating and not recursive, then δ'_2 is recursive on S' .*
- (5) *If \mathcal{T} is chained triple satisfying condition $(*)$, then for every boundary component C of S' , we have either $\delta_1 \cap C = \emptyset$ or $\delta'_2 \cap C = \emptyset$.*




FIGURE 1. (a) the arc δ_2 joins C to itself, (b) δ_2 joins two distinct boundary components.

Proof. (1) Suppose that a curve α_i is isotopic to a boundary component of S' . Then, since α_i is essential in S_g^p , α_i is isotopic to $\partial S_g^p \cup \delta_2$. Then, depending on whether $\partial\delta_2$ lies in a single boundary component of S_g^p or $\partial\delta_2$ is not contained in a single boundary component, we have the following.

- (a) If δ_2 joins a boundary component C to itself, then there is an annulus A in S_g^p such that $\partial A \subset \alpha_i \sqcup (\delta_2 \cup \partial S_g^p)$ (see Figure 1 (a)).
- (b) If δ_2 joins a boundary component C_1 to another boundary component C_2 , then there is a two-holed disk D in S_g^p such that $\partial D = \alpha_i \sqcup C_1 \sqcup C_2$ (see Figure 1 (b)).

In each case, any closed curve in S_g^p that intersects α_i non-trivially and minimally must intersect with δ_2 . Hence, we have $i = 1$ and $m = 2$.

(2) Suppose that $m \geq 3$. Then, any closed curve in $(\alpha_1, \dots, \alpha_{m-1})$ is not boundary parallel (by (1)) and bounds no disk, and hence, it is essential in S' . In addition, consecutive closed curves α_i and α_{i+1} intersect minimally and non-trivially, and non-consecutive closed curves are disjoint. Therefore, $(\alpha_1, \dots, \alpha_{m-1})$ is a linear chain on S' . From our assumption that $(\delta_a, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple and $\delta'_b \subset \alpha_m$, the arc δ_1 intersects with α_1 minimally and non-trivially but does not intersect $\alpha_2, \dots, \alpha_{m-1}$ or δ'_2 . Similarly, the arc δ'_2 intersects α_{m-1} minimally and non-trivially but does not intersect $\alpha_1, \dots, \alpha_{m-2}$ or δ_1 . Thus, $(\delta_1, (\alpha_1, \dots, \alpha_{m-1}), \delta'_2)$ is a chained triple on S' .

(3) Suppose that δ_2 is either separating or non-separating and not recursive on S_g^p . Then, in either case, for each component S'' of $S_g^p \setminus \text{Int}N(\delta_2)$, the number of boundary components of S'' that are derived from δ_2 (and a boundary component of S_g^p) is exactly one. Hence, this fact together with the assumption that δ_1 is recursive on S_g^p implies that δ_1 is recursive on S' .

(4) Suppose that δ_2 is either separating or non-separating and not recursive on S_g^p . Then, for each component S'' of $S_g^p \setminus \text{Int}N(\delta_2)$, the number of boundary components of S'' that are derived from δ_2 (and a boundary component of S_g^p) is one. Hence, since $\partial\delta'_2 \subset \delta_2$, it follows that $\partial\delta'_2$ is contained in a single boundary component. Therefore δ'_2 is recursive on S' .

(5) Suppose that \mathcal{T} satisfies condition (*). Pick a boundary component C of S' . Then, the following holds.

C has an endpoint of δ'_2 .

$\Leftrightarrow C$ is derived from δ_2 and the boundary component(s) of S_g^p .

Condition (*) implies that C is not derived from δ_2 or the boundary component(s) of S_g^p if C contains an endpoint of δ_1 . Hence, either $\delta_1 \cap C = \emptyset$ or $\delta_2' \cap C = \emptyset$ holds. \square

Now, let us compute the maximal lengths of chained triples on holed spheres.

Lemma 3.6. *Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple on S_0^3 . Then $m = 0$.*

Proof. There is no essential simple closed curve in S_0^3 . Hence, m must be 0. \square

Lemma 3.7. *Let p be an integer ≥ 4 . Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple on S_0^p . Then $m \leq p - 2$.*

Proof. Note that $((\alpha_1, \dots, \alpha_m), \delta_2)$ is a “chained pair” on S_0^p (a chained pair is defined in [12, Section 2]). By [12, Lemmas 2.5 and 2.6], we have $m \leq p - 2$. \square

Next, we compute the maximal lengths of chained triples satisfying condition (*) on spheres with holes in Lemmas 3–3.10 below.

Lemma 3.8. *Suppose that $2 \leq p \leq 3$. Then there is no chained triple satisfying condition (*) on S_0^p .*

Proof. We will prove this lemma in the case where $p = 3$. Let δ_1 and δ_2 be mutually non-isotopic disjoint arcs on S_0^3 . If $\partial\delta_1$ is contained in a single boundary component, then δ_2 cannot be essential. Hence, δ_1 connects two boundary components. Similarly, δ_2 connects two boundary components. Since $S_0^3 \setminus \text{Int}N(\delta_1)$ is an annulus, a boundary component of S_0^3 intersects both $\partial\delta_1$ and $\partial\delta_2$. Hence, no chained triple satisfies condition (*) on S_0^3 . \square

Lemma 3.9. *Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple satisfying condition (*) on S_0^4 . Then $m \leq 1$.*

Proof. Suppose, on the contrary, that $m \geq 2$. Then, a component S' of $S_0^4 \setminus \text{Int}N(\delta_b)$ contains δ_1 . Pick a connected component δ_2' of $\alpha_m \cap S'$ intersecting α_{m-1} . Since $m \geq 2$, the arc δ_1 is disjoint from δ_2' . By Lemma 3.5 (5), it follows that δ_1 is disjoint from δ_2' and is not isotopic to δ_2' . This implies that S' has $(\delta_1, \emptyset, \delta_2')$, a chained triple of length 0 satisfying condition (*). This is absurd because S' is homeomorphic to S_0^2 or S_0^3 and does not contain such a chained triple by Lemma 3. Thus, we have $m \leq 1$. \square

Lemma 3.10. *Let p be an integer ≥ 5 . Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple satisfying condition (*) on S_0^p . Then $m \leq p - 3$.*

Proof. **Case of $p = 5$.** Suppose, on the contrary, that $m \geq 3$. Then, by Lemma 3.5 (2) and (5), a connected component S' of $S_0^5 \setminus \text{Int}N(\delta_2)$ contains a chained triple $\mathcal{T}' = (\delta_1, (\alpha_1, \dots, \alpha_{m-1}), \delta_2')$ satisfying condition (*). Hence, S' contains a length $m - 1 \geq 2$ chained triple satisfying condition (*). This is a contradiction by Lemmas 3.4, and 3.9. Thus, we have $m \leq 2$.

Case of $p \geq 6$. We may assume that $m \geq 3$. A similar argument to the $p = 5$ case, together with the induction hypothesis, completes the proof. \square

Next, we compute the maximal length of chained triples on a one-holed torus.

Lemma 3.11. *Suppose that $(\delta_1, \{\alpha_1, \dots, \alpha_m\}, \delta_2)$ is a chained triple on S_1^1 . Then $m \leq 2$.*

Proof. Annuli do not have a chained triple of length ≥ 0 . So Lemma 3.5 (2) immediately implies the assertion. \square

Next, we compute the maximal lengths of a few different chained triples on a two-holed torus.

Lemma 3.12. *Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple on S_1^2 satisfying condition (*). Then $m \leq 2$.*

Proof. Now, let us assume that $m \geq 3$ and derive a contradiction. It follows that either δ_1 or δ_2 is non-separating. Since δ_2 is non-separating, $S' = S_1^2 \setminus \text{Int}N(\delta_2)$ is homeomorphic to S_0^3 . Since $m \geq 3$, S' contains a chained triple of length $m - 1$ by Lemma 3.5 (2). By Lemma 3.6, we have $m - 1 \leq 0$, which is a contradiction. Thus, $m \leq 2$. \square

Lemma 3.13. *Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple on S_1^2 such that δ_2 is recursive. Then $m \leq 2$.*

Proof. To the contrary, suppose that $m \geq 3$. Since δ_2 is recursive, δ_1 must be non-separating on S_1^2 . Then $S_0^3 \cong S_1^2 \setminus \text{Int}N(\delta_1)$ contains a chained triple of length $m - 1 \geq 2$ by Lemma 3.5 (2), this is a contradiction. Thus we have $m \leq 2$. \square

Now, let us compute the maximal lengths of chained triples on holed tori.

Lemma 3.14. *Let p be an integer ≥ 3 . Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple on S_1^p . Then $m \leq p$.*

Proof. Proof by induction on p .

Case of $p = 3$. Assuming $m \geq p = 3$, we will prove that $m = 3$. First, we treat the case where δ_2 is non-separating. Since $m \geq 3$, the surface $S' = S_1^3 \setminus \text{Int}N(\delta_2)$ contains a chained triple of length $m - 1$ by Lemma 3.5 (2), and is homeomorphic to $S_0^{p'}$, where $p' \leq 4$. Hence, by Lemmas 3.6 and 3.7 we have $m - 1 \leq 2$. Therefore $m = 3$. Next, we treat the case where δ_2 is separating. By Lemma 3.5 (2) and (4), there is a connected component S' of $S_1^3 \setminus \text{Int}N(\delta_2)$ that contains a chained triple $\mathcal{T}' = (\delta_1, (\alpha_1, \dots, \alpha_{m-1}), \delta_2')$ such that δ_2' is recursive on S' . Since a sphere with at most three boundary components does not have a chained triple of positive length, S' must be a torus with at most two boundary components. Hence, Lemma 3.13 implies $m - 1 \leq 2$ i.e., $m \leq 3$.

Case of $p \geq 4$. We may assume that $m \geq p > 3$. First, we treat the case where δ_2 is non-separating. Since $m \geq 3$, the surface $S' = S_1^p \setminus \text{Int}N(\delta_2)$ contains a chained triple of length $m - 1$ by Lemma 3.5 (2), and is homeomorphic to $S_0^{p'}$, where $p' \leq p + 1$. Hence, by Lemmas 3.6 and 3.7 we have $m - 1 \leq (p + 1) - 2$, namely, $m \leq p$. Next, we treat the case where δ_2 is separating. Then, a connected component S' of $S_1^p \setminus \text{Int}N(\delta_2)$ contains a chained triple of length $m - 1$ by Lemma 3.5 (2). When S' is a sphere with a boundary, an argument similar to the case where δ_2 is non-separating implies $m \leq p - 1$. When S' is a torus with a boundary, the induction hypothesis implies $m \leq p$. \square

Remark 3.15. Hyperbolic extensions of surfaces preserves chained triples. Hence, Lemma 3.14 also implies that the maximal length of chained triples on S_1^2 is at most 3.

Now, let us compute the maximal lengths of chained triples on higher genera surfaces.

Lemma 3.16. *Let g be an integer ≥ 2 and p be an integer ≥ 1 . Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple on S_g^p . Then*

$$m \leq \begin{cases} 2g + p - 1 & (p \leq 2) \\ 2g + p - 2 & (p \geq 3) \end{cases}$$

Furthermore, if $p = 2$ and δ_2 is recursive, then $m \leq 2g$.

Proof. We give a proof by induction. Assume that $m \leq 3$.

Case of $(g, p) = (2, 1)$. In the case where δ_2 is non-separating, $S_2^1 \setminus \text{Int}N(\delta_2)$ is homeomorphic to S_1^2 and contains a chained triple of length $m - 1$ by Lemma 3.5 (2). By Remark 3.15, we have $m - 1 \leq 3$ i.e., $m \leq 4 = 2g + p - 1$. In the case where δ_2 is separating, a connected component of $S_2^1 \setminus \text{Int}N(\delta_2)$ is homeomorphic to S_1^1 and contains a chained triple of length $m - 1$. By Lemma 3.11, we have $m - 1 \leq 2$ i.e., $m \leq 3 < 2g + p - 1$.

Case of $g \geq 2$, $p = 2$ and δ_2 is recursive. In order to prove the assertion we further divide the proof into two cases: (i) δ_2 is non-separating and (ii) δ_2 is separating.

(i) Suppose that δ_2 is non-separating. Then $S_g^2 \setminus \text{Int}N(\delta_2)$ is homeomorphic to S_{g-1}^3 and contains a chained triple of length $m - 1$ by Lemma 3.5 (2). By Lemma 3.14 and the induction hypothesis, we have $m - 1 \leq 2(g - 1) + 3 - 2$ i.e., $m \leq 2g$.

(ii) Suppose that δ_2 is separating. We claim that either δ_1 or δ_2 does not bound an annulus. To see this, assume that δ_2 bounds an annulus, i.e., $S_g^2 \setminus \text{Int}N(\delta_2) \cong S_g^1 \cup S_0^2$. Since any arc that is not recursive and does not intersect δ_2 should be contained in the annulus and since α_m must intersect such an arc, the inequality $m \geq 3$ implies that δ_1 is not such an arc; δ_1 should be recursive. Moreover, by (i) we may assume that δ_1 is separating. Then δ_1 does not bound an annulus, because δ_1 and δ_2 are not isotopic and that δ_1 is boundary parallel on $S_g^2 \setminus \text{Int}N(\delta_2) (\cong S_g^1)$. Hence, either δ_1 or δ_2 does not bound an annulus. So we may assume that δ_2 does not bound an annulus. Pick a connected component S' of $S_g^2 \setminus \text{Int}N(\delta_2)$ which contains a chained triple $(\delta_1, (\alpha_1, \dots, \alpha_{m-1}), \delta_2')$ obtained from Lemma 3.5 (2). Since δ_2 does not bound an annulus, the component S' is homeomorphic to a surface of genus $\leq g - 1$ with at most two boundary components. Moreover, Lemma 3.5 (2) implies that the arc δ_2' derived from α_m is recursive, because δ_2 is separating. In the case where the number of boundary components of S' is 2, by Lemma 3.13 and the induction hypothesis we have $m - 1 \leq 2(g - 1)$, i.e., $m \leq 2g - 1$. Since a hyperbolic extension of a surface preserves chained triple, the case where the number of boundary components of S' is 1 can be reduced to the case where the number of boundary components of S' is 2. Thus we have $m \leq 2g - 1$.

Case of $(g, p) = (2, 2)$. By Lemma 3.5 (2), a connected component S' of $S_2^2 \setminus \text{Int}N(\delta_2)$ contains a chained triple of length $m - 1$. Then S' is homeomorphic to S_2^1 ; otherwise the genus of S' is at most 1 and the number of boundary components of S' is at most 3. In both cases we have $m - 1 \leq 4$ by Lemmas 3.6, 3.14 and the assertion in the case where $(g, p) = (2, 1)$. Hence $m \leq 5 = 2g + p - 1$.

Case of $g \geq 3$ and $p = 1$. In the case where δ_2 is non-separating, $S_g^1 \setminus \text{Int}N(\delta_2)$ is homeomorphic to S_{g-1}^2 and contains a chained triple of length $m - 1$. By the induction hypothesis, we have $m - 1 \leq 2(g - 1) + 2 - 1$ i.e., $m \leq 2g = 2g + p - 1$. In the case where δ_2 is separating, a connected component of $S_g^1 \setminus \text{Int}N(\delta_2)$ is homeomorphic to $S_{g'}^1$ ($g' \leq g - 1$) and contains a chained triple of length $m - 1$. The induction hypothesis gives $m - 1 \leq 2(g - 1)$, i.e., $m \leq 2g - 1 < 2g + p - 1$.

Case of $g \geq 3$ and $p = 2$. A connected component S' of $S_g^2 \setminus \text{Int}N(\delta_2)$ contains a chained triple of length $m-1$. Then either S' is homeomorphic to S_g^1 or the genus of S' is at most $g-1$ and the number of boundary components of S' is at most 3. By the induction hypothesis, we have $m-1 \leq 2g$. Hence, $m \leq 2g+1 = 2g+p-1$.

Case of $g \geq 2$ and $p \geq 3$. A connected component S' of $S_g^p \setminus \text{Int}N(\delta_2)$ contains a chained triple of $(\delta_1, (\alpha_1, \dots, \alpha_{m-1}), \delta_2')$. Suppose that δ_2 is separating. Then δ_2' is recursive. When the genus of S' is g , the induction hypothesis gives $m-1 \leq 2g+(p-1)-2$, i.e., $m \leq 2g+p-2$. When the genus of S' is at most $g-1$, the induction hypothesis gives $m-1 \leq 2(g-1)+(p+1)-2$, i.e., $m \leq 2g+p-2$. Next, let us consider the case where δ_2 is non-separating. Then, S' is homeomorphic to S_{g-1}^{p+1} . The induction hypothesis gives $m-1 \leq 2(g-1)+(p+1)-2$. Thus, the assertion holds. \square

We will also use the following sequence on surfaces.

Definition 3.17. A quadruple $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2, \delta_3)$ of arcs δ_1, δ_2 and δ_3 , and a sequence $(\alpha_1, \dots, \alpha_m)$ of closed curves on a surface S is said to be *Y-chained* if the quadruple satisfies the following conditions:

- $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2)$ is a chained triple on S ,
- $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_3)$ is a chained triple on S ,
- δ_2, δ_3 are disjoint and non-isotopic, and
- there is a boundary component C of S such that $\partial\delta_2 \cup \partial\delta_3 \subset C$.

We prepare a lemma for Y-chained quadruples, which is similar to Lemma 3.5.

Lemma 3.18. *Suppose $m \geq 3$. Suppose also that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2, \delta_3)$ is a Y-chained quadruple on a surface S . Then, there is an arc δ_1' on $S \setminus \text{Int}N(\delta_1)$ derived from α_1 such that $(\delta_1', (\alpha_2, \dots, \alpha_m), \delta_2, \delta_3)$ is Y-chained quadruple on $S \setminus \text{Int}N(\delta_1)$.*

Proof. Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2, \delta_3)$ is a Y-chained quadruple on a surface S . Pick a proper arc δ_1' on $S \setminus \text{Int}N(\delta_1)$ which is derived from α_1 such that $\delta_1' \cap \alpha_2 = \emptyset$. By [12], both $(\delta_1, (\alpha_2, \dots, \alpha_m), \delta_2)$ and $(\delta_1, (\alpha_2, \dots, \alpha_m), \delta_3)$ are chained triples on $S \setminus \text{Int}N(\delta_1)$. Obviously, δ_2 and δ_3 are disjoint on $S \setminus \text{Int}N(\delta_1)$. Since any isotopy between δ_2 and δ_3 on $S \setminus \text{Int}N(\delta_1)$ gives rise to an isotopy between δ_2 and δ_3 on S relative to $\text{Int}N(\delta_1)$, it holds that δ_2 and δ_3 are non-isotopic on $S \setminus \text{Int}N(\delta_1)$. Since the boundary component of S containing $\partial\delta_2 \cup \partial\delta_3$ naturally becomes a boundary component of $S \setminus \text{Int}N(\delta_1)$, $(\delta_1', (\alpha_2, \dots, \alpha_m), \delta_2, \delta_3)$ is a Y-chained quadruple on $S \setminus \text{Int}N(\delta_1)$. \square

Let us compute the maximal lengths of Y-chained quadruples on surfaces. First, we treat holed spheres.

Lemma 3.19. *Any two disjoint arcs δ_2, δ_3 such that $\partial\delta_2 \cup \partial\delta_3$ is contained in a single boundary component of S_0^3 are isotopic.*

Proof. Since $S_0^3 \setminus \text{Int}N(\delta_3)$ consists of two copies of annuli, δ_2 is boundary parallel on $S_0^3 \setminus \text{Int}N(\delta_3)$. This implies that δ_2 and δ_3 are isotopic on S_0^3 . \square

Lemma 3.20. *Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2, \delta_3)$ is a Y-chained quadruple on S_0^4 . Then $m \leq 1$.*

Proof. Suppose, on the contrary, that $m \geq 2$. Then a connected component S' of $S_0^4 \setminus \text{Int}N(\alpha_1)$ contains δ_2 and δ_3 and is homeomorphic to S_0^3 . By Lemma 3.19, there is an isotopy between δ_2 and δ_3 on S' . However, this isotopy in turn gives an isotopy between δ_2 and δ_3 on the original surface S_0^4 , which is a contradiction. \square

Lemma 3.21. *Let p be an integer ≥ 5 . Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2, \delta_3)$ is a Y -chained quadruple on S_0^p . Then $m \leq p - 3$.*

Proof. Proof by induction on p .

Case of $p = 5$. Suppose, on the contrary, that $m \geq 3 > p - 3$. Then, a connected component S' of $S \setminus \text{Int}N(\alpha_1)$ contains a length $m - 1$ chained quadruple by Lemma 3.18. Since S' is homeomorphic to a sphere with at most four boundary components, by Lemmas 3.19 and 3.20 $m - 1 \geq 2$ is impossible. This is a contradiction, so $m \leq 2$.

Case of $p \geq 6$. An argument similar to the $p = 5$ case, together with the induction hypothesis, completes the proof. \square

Next, let us consider holed tori.

Lemma 3.22. *No Y -chained quadruple is contained in S_1^1 .*

Proof. Any triple of mutually non-isotopic arcs on S_1^1 intersects minimally non-trivially. \square

Lemma 3.23. *Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2, \delta_3)$ is a Y -chained quadruple on S_1^2 . Then $m \leq 2$. Furthermore, if $(\delta_1, (\alpha_1, \alpha_2), \delta_2, \delta_3)$ is a Y -chained quadruple on S_1^2 , then δ_2 is separating on S_1^2 .*

Proof. First, we treat the case where δ_2 is separating. In this case $S_1^2 \setminus \text{Int}N(\delta_2)$ is homeomorphic to a disjoint union of a one-holed torus T and an annulus A . If δ_1 is contained in A , then any essential simple closed curve on S_1^2 intersecting δ_1 must also intersect δ_2 , so $m \leq 1$. Thus, we may assume that δ_1 is contained in T . The arc δ_3 is contained in either T or A , because δ_3 is disjoint from δ_2 . Since $\partial\delta_2$ and $\partial\delta_3$ are contained in the same boundary component and since δ_3 is not isotopic to δ_2 , the arc δ_3 must be contained in T . Since δ_1 is not isotopic to δ_3 , it follows that any essential simple closed curve in T must intersect either δ_1 or δ_3 . This implies that $m \leq 2$ (if $m \geq 2$, α_2 must intersect δ_3).

Next, we treat the case where δ_2 is not separating. We may assume that δ_3 is not separating either. Then $S' := S_1^2 \setminus \text{Int}N(\delta_2)$ is homeomorphic to S_0^3 and δ_3 connects the two boundary components C_1 and C_2 which are derived from δ_2 on S' . Any essential simple closed curve on S_1^2 that is disjoint from δ_2 is isotopic to either C_1 or C_2 on S' and thereby intersecting δ_3 on the original surface S_1^2 . Hence, we have $m \leq 1$.

The last assertion of this lemma immediately follows from the above. \square

The maximal lengths of Y -chained quadruples on higher genera surfaces can be computed as follows.

Lemma 3.24. *Suppose that $(\delta_1, (\alpha_1, \dots, \alpha_m), \delta_2, \delta_3)$ is a Y -chained quadruple on S_g^p . If the pair (g, p) satisfies either $g = 1$ and $p \geq 3$ or $g \geq 2$ and $p \geq 2$, then $m \leq 2g + p - 3$.*

Proof. Proof by induction on (g, p) .

Case of $(g, p) = (1, 3)$. In the case where δ_1 is non-separating, $S_1^3 \setminus \text{Int}N(\delta_1)$ is homeomorphic to S_0^4 , so we have $m \leq 2$ by Lemma 3.20. Next, let us consider

the case where δ_1 is separating. Suppose, on the contrary, that $m \geq 3$. By Lemma 3.18, $S_g^p \setminus \text{Int}N(\delta_1)$ contains a Y -chained quadruple L of length $m - 1 \geq 2$. By Lemmas 3.20 and 3.22, $S \setminus \text{Int}N(\delta_1)$ must be homeomorphic to a disjoint union of a two-holed torus T and an annulus A , and L is contained in T . Lemma 3.23 implies that m must be equal to 3 and δ_2 is separating on T . It follows that δ_2 is separating on the original surface S_1^3 . Moreover, δ_3 is non-separating on S_1^3 . Therefore $S_1^3 \setminus \text{Int}N(\delta_3)$ is homeomorphic to S_0^4 . Obviously, α_2 and δ_2 are essential on $S_1^3 \setminus \text{Int}N(\delta_3)$. Every essential simple closed curve on $S_1^3 \setminus \text{Int}N(\delta_3)$ intersecting α_2 but disjoint from δ_2 forms a bigon with α_2 . Since α_1 does not form a bigon with α_2 on S_1^3 , α_3 is isotopic to a boundary component C of $S_1^3 \setminus \text{Int}N(\delta_3)$. However, this implies that α_1 intersects α_2 trivially on S_1^3 , which is a contradiction. Hence, we have $m \leq 2$.

In the following cases we may assume that $m \geq 3$. By Lemma 3.18, $S_g^p \setminus \text{Int}N(\delta_1)$ contains a Y -chained quadruple of length $m - 1$.

Case of $g = 1$ and $p \geq 4$. Lemma 3.21 and the induction hypothesis give $m - 1 \leq p - 3$.

Case of $g \geq 2$ and $p \geq 1$. Lemma 3.21 and the induction hypothesis give $m - 1 \leq 2(g - 1) + (p + 1) - 3$, i.e., $m \leq 2g + p - 3$. \square

We are now ready to decide whether there is a cyclic chain of a given length on surfaces. To begin with, we compute the maximal lengths of cyclic chains whose two consecutive curves intersect exactly once on surfaces.

Lemma 3.25. *Let $(\alpha_1, \dots, \alpha_m)$ be a cyclic chain on S_g^p . If $\text{int}(\alpha_i, \alpha_{i+1}) = 1$ ($1 \leq i \leq m$), then $m \leq 2g + 2$.*

Proof. We will prove that the genus g' of a subsurface $N(\alpha_1 \cup \dots \cup \alpha_m)$ is at least $\frac{m}{2} - 1$. This fact directly implies that $m \leq 2g + 2$, because the genus of a subsurface does not exceed that of the ambient surface. First, we show that the number n of boundary components of $N(\alpha_1 \cup \dots \cup \alpha_m)$ is either at most 3 or at most 4 according to whether m is odd or even. Consider the case where m is odd. The number of the boundary components of a sub-subsurface $N(\alpha_1 \cup \dots \cup \alpha_{m-1})$ is 1. Note that the subsurface $N(\alpha_1 \cup \dots \cup \alpha_m)$ can be obtained from $(\alpha_1, \dots, \alpha_{m-1})$ by attaching two bands to its two boundary components so that the bands connect the boundary components. Since these increase at most 1 boundary component, n is at most 3. Next, we consider the case where m is even. The number of boundary components of a sub-subsurface $N(\alpha_1 \cup \dots \cup \alpha_{m-1})$ is 2. Hence, n is at most 4.

Using the Euler characteristic we obtain the equality $2 - 2g' - n = -m$, i.e., $g' = \frac{2+m-n}{2}$. Thus $g' \geq \frac{m}{2} - 1$. \square

Note that a cyclic chain of length m on a surface gives a shorter cyclic chain as follows.

Lemma 3.26. *Let m be an integer greater than 3, S a surface with negative Euler characteristic. If S has a cyclic chain of length m , then S has a cyclic chain of length $m - 1$.*

Proof. Let $(\alpha_1, \dots, \alpha_m)$ be a cyclic chain on S , T_{α_m} be a Dehn twist along a closed curve α_m . Then $(\alpha_1, \dots, \alpha_{m-2}, T_{\alpha_m}(\alpha_{m-1}))$ is a cyclic chain of length $m - 1$. \square

Let us compute the maximal lengths of cyclic chains on surfaces.




FIGURE 2. Longest cyclic chains on tori with at most two boundary components.




FIGURE 3. Longest cyclic chains on tori with at least three boundary components.

Lemma 3.27. *The complement graph C_m^c of the cyclic graph on m vertices is a full subgraph of $\mathcal{C}(S_g^p)$ if and only if*

$$m \leq \begin{cases} p & (g = 0, p \geq 5) \\ 3 & (g, p) = (1, 1) \\ 5 & (g, p) = (1, 2) \\ p + 2 & (g = 1, p \geq 3) \\ 2g + 2 & (g \geq 2, p = 0) \\ 2g + p + 1 & (g \geq 2, 1 \leq p \leq 2). \\ 2g + p & (g \geq 2, p \geq 3). \end{cases}$$

Proof. **Case of $g = 0$ and $p \geq 5$.** From [12, Figure 7], we can see that S_0^p contains a cyclic chain of length p . By Lemma 3.26, we can obtain full embeddings of shorter cyclic graphs into $\mathcal{C}(S_0^p)$. On the other hand, S_0^p does not contain a linear chain of length $\geq p$ by [12, Theorem 2.2]. Since any cyclic chain of length m induces a linear chain of length $m - 1$, we have $C_m^c \not\leq \mathcal{C}(S_0^p)$ for each $m \geq p + 1$.

Case of $(g, p) = (1, 1)$. The left picture in Figure 2 shows $C_3^c \leq \mathcal{C}(S_1^1)$. Note that a cyclic chain of length ≥ 4 contains a pair of non-isotopic disjoint closed curves. A one-holed torus does not contain such a pair.

Case of $(g, p) = (1, 2)$. The right picture in Figure 2 shows $C_5^c \leq \mathcal{C}(S_1^2)$. By Lemma ??, we can obtain full embeddings of C_3^c and C_4^c into $\mathcal{C}(S_1^2)$. A cyclic chain of length ≥ 6 contains a triple of pairwise non-isotopic disjoint closed curves. A two-holed torus does not contain such a triple.

Case of $g = 1, p \geq 3$. Figure 3 shows $C_{p+2}^c \leq \mathcal{C}(S_1^p)$. Suppose that $(\alpha_1, \dots, \alpha_m)$ is a cyclic chain on S_1^p . We may assume that $m \geq p + 2 \geq 5$. By Lemma 3.25, $(\alpha_1, \dots, \alpha_m)$ contains a pair of closed curves whose intersection number is greater than 1. Hence, without loss of generality we may assume that the intersection number of α_{m-1} and α_m is greater than 1. Thus, a connected component S' of




FIGURE 4. Longest cyclic chains on surfaces of genus ≥ 2 with at most two boundary components.

$S_1^p \setminus \text{Int}N(\alpha_m)$ contains a quadruple $L := (\delta_1, (\alpha_2, \dots, \alpha_{m-2}), \delta_2, \delta_3)$. The quadruple L is Y -chained or not according to whether δ_2 intersects α_{m-2} or not.

(i) Suppose that δ_2 intersects α_{m-2} . Then, L is Y -chained on S' . By Lemmas 3.21, 3.23 and 3.24 we have $m - 3 \leq (p + 2) - 3$ (resp. $m - 3 \leq 2$) if $p \geq 4$ (resp. $p = 3$).

(ii) Suppose that δ_2 is disjoint from α_2 . Then δ_2 is disjoint from a chained triple $(\delta_1, (\alpha_2, \dots, \alpha_{m-2}), \delta_3)$. Since δ_2 is isotopic to neither δ_1 nor δ_3 , a connected component of $S_1^p \setminus \text{Int}(N(\alpha_m) \cup N(\delta_2))$ contains $(\delta_1, (\alpha_2, \dots, \alpha_{m-2}), \delta_3)$. By Lemmas 3.7, 3.11 and 3.14, we have $m - 3 \leq p - 1$. Thus, we have $m \leq p + 2$.

Case of $g \geq 2$ and $p = 0$. From [12, Figure 7], we can see that S_g contains a cyclic chain of length $2g + 2$. On the other hand, S_0^p does not contain a linear chain of length $\geq 2g + 2$ by [12, Theorem 2.2]. Hence, $C_m^c \not\leq \mathcal{C}(S_0^p)$ for each $m \geq 2g + 3$.

Case of $g \geq 2$, $1 \leq p \leq 2$. The left picture in Figure 4 shows $C_{2g+3}^c \leq \mathcal{C}(S_g^1)$ and the right picture shows $C_{2g+3}^c \leq \mathcal{C}(S_g^2)$. Suppose that $(\alpha_1, \dots, \alpha_m)$ is a cyclic chain on S_g^p . We may assume that $m \geq 2g + p + 1 > 3$. Then a connected component S' of $S_g^p \setminus \text{Int}N(\alpha_m)$ contains a chained triple $(\delta_a, \{\alpha_2, \dots, \alpha_{m-2}\}, \delta_b)$. Hence, by Lemma 3.16 we have $m - 3 \leq 2(g - 1) + (p + 2) - 2$ i.e., $m \leq 2g + p + 1$.

Case of $g \geq 2$, $p \geq 3$. Figure 5 shows $C_{2g+p}^c \leq \mathcal{C}(S_g^p)$. Suppose that $(\alpha_1, \dots, \alpha_m)$ is a cyclic chain on S_g^p . We may assume that $m \geq 2g + p > 2g + 2 = 6$. By Lemma 3.25, $(\alpha_1, \dots, \alpha_m)$ contains a pair of closed curves whose intersection number is greater than 1. Hence, without loss of generality we may assume that the intersection number of α_{m-1} and α_m is greater than 1. Then a connected component S' of $S_g^p \setminus \text{Int}N(\alpha_m)$ contains a quadruple $L := (\delta_1, (\alpha_2, \dots, \alpha_{m-2}), \delta_2, \delta_3)$. The quadruple L is Y -chained or not according to whether δ_2 intersects α_{m-2} or not.

(i) Suppose that δ_2 intersects α_{m-2} . Then L is Y -chained on S' . By Lemmas 3.21 and 3.24, we have $m - 3 \leq 2(g - 1) + (p + 2) - 3$, i.e., $m \leq 2g + p$.

(ii) Suppose that δ_2 is disjoint from α_2 . Then δ_2 is disjoint from a chained triple $(\delta_1, (\alpha_2, \dots, \alpha_{m-2}), \delta_3)$. Since δ_2 is neither isotopic to δ_1 nor δ_3 , a connected component of $S_g^p \setminus \text{Int}N(\alpha_m) \cup N(\delta_2)$ contains $(\delta_1, (\alpha_2, \dots, \alpha_{m-2}), \delta_3)$. From Lemmas 3.7, 3.11, 3.14 and 3.16, it holds that $m - 3 \leq 2(g - 2) + (p + 3) - 2$, i.e., $m \leq 2g + p$. \square

For a pair of graphs Λ_1 and Λ_2 , the *graph-join* $*_{i=1}^2 \Lambda_i = \Lambda_1 * \Lambda_2$ is a graph such that the vertex set is a disjoint union of $V(\Lambda_1)$ and $V(\Lambda_2)$, and that the edge set is a disjoint union of $E(\Lambda_1)$, $E(\Lambda_2)$ and $\{[u, v] \mid u \in V(\Lambda_1), v \in V(\Lambda_2)\}$.




FIGURE 5. Longest cyclic chains on surfaces of genus ≥ 2 with at least three boundary components.

Lemma 3.28. $C_m^c * K_1$ is a full subgraph of $\mathcal{C}(S_g^p)$ if and only if

$$m \leq \begin{cases} 3 & (g, p) \in \{(0, 5), (1, 2)\} \\ p - 1 & (g = 0, p \geq 6) \\ p + 2 & (g = 1, p \geq 3) \\ 2g + 1 & (g \geq 2, p = 0) \\ 2g + p & (g \geq 2, p \geq 1) \end{cases}$$

Proof. We will reduce the “if part” of Lemma 3.28 to Lemma 3.27. Suppose $g \geq 2$ and $p \leq 1$. Then, any cyclic chain of length m on S_{g-1}^{p+2} induces a cyclic chain of length m on S_g^p via an annular extension $S_{g-1}^{p+2} \rightarrow S_g^p$. Hence, the disjoint union of the core of the attached annulus and the induced cyclic chain of length m induces a full embedding $C_m^c * K_1 \leq \mathcal{C}(S_g^p)$. Consequently, $C_m^c \leq \mathcal{C}(S_{g-1}^{p+2})$ implies $C_m^c * K_1 \leq \mathcal{C}(S_g^p)$. Next, we consider the case where $p \geq 2$ (and $g \geq 0$). Use a hyperbolic extension $S_{g-1}^{p-1} \rightarrow S_g^p$. Then $C_m^c \leq \mathcal{C}(S_{g-1}^{p-1})$ implies $C_m^c * K_1 \leq \mathcal{C}(S_g^p)$. Hence, Lemma 3.27 together with [12, Figure 7] completes the “if part” of Lemma 3.28.

Now, we prove the “only if part” of Lemma 3.28.

Case of $g = 0, p \geq 5$. Suppose that $C_m^c * K_1 \leq \mathcal{C}(S_0^p)$. Then, a cyclic chain of length m is contained in a connected component S' of the surface $S_0^p \setminus \text{Int}N(\alpha)$, where α is a closed curve corresponding to the graph K_1 . Since S' is a sphere with $\leq p - 1$ holes, m must be not more than $p - 1$.

Case of $(g, p) = (1, 2)$. Suppose that $C_m^c * K_1 \leq \mathcal{C}(S_1^2)$. Then, a cyclic chain of length m is contained in a connected component S' of the surface $S_1^2 \setminus \text{Int}N(\alpha)$, where α is a closed curve corresponding to the graph K_1 . Since S' is homeomorphic to either S_0^4 or S_1^1 , we have $m \leq 3$.

Case of $g = 1, p \geq 3$. Suppose that $C_m^c * K_1 \leq \mathcal{C}(S_1^p)$. Then, a cyclic chain of length m is contained in a connected component S' of the surface $S_1^p \setminus \text{Int}N(\alpha)$, where α is a closed curve corresponding to the graph K_1 . The component S' is homeomorphic to either $S_0^{p'}$ or $S_1^{p''}$, where $p' \leq p + 2$ and $p'' \leq p - 1$. Hence, by Lemma 3.27, we have $m \leq p + 2$.

Case of $g \geq 2, p = 0$. Suppose that $C_m^c * K_1 \leq \mathcal{C}(S_g^0)$. Then, a cyclic chain of length m is contained in a connected component S' of the surface $S_g^0 \setminus \text{Int}N(\alpha)$, where α is a closed curve corresponding to the graph K_1 . The component S' is homeomorphic to $S_{g'}^{p'}$, where $g' = g - 1$ and $p' \leq 2$. Hence, we have $m \leq 2(g - 1) + 2 + 1 = 2g + 1$ by Lemma 3.27.

Case of $g \geq 2$, $p \geq 1$. Suppose that $C_m^c * K_1 \leq \mathcal{C}(S_g^p)$. Then, a cyclic chain of length m is contained in a connected component S' of the surface $S_g^p \setminus \text{Int}N(\alpha)$, where α is a closed curve corresponding to the graph K_1 . The component S' is homeomorphic to $S_{g'}^{p'}$ with one of the following:

- (a) $g' = g$ and $1 \leq p' \leq p - 1$,
- (b) $g' \leq g - 1$ and $p' \leq p + 2$.

Hence, we have $m \leq 2(g - 1) + (p + 2) = 2g + p$ by Lemma 3.27. \square

We denote the path graph on m vertices by P_m (the geometric realization is homeomorphic to the unit interval).

Lemma 3.29. *Suppose that $p \geq 5$. If $P_m^c * K_2 \leq \mathcal{C}(S_0^p)$, then $m \leq p - 3$.*

Proof. Suppose that $P_m^c * K_2 \leq \mathcal{C}(S_0^p)$. Then, there is a linear chain $(\alpha_1, \dots, \alpha_m)$ and closed curves β_1, β_2 on $S_{0,p}$ such that each β_i is isotopic to none of $\{\alpha_1, \dots, \alpha_m, \beta_j\}$ and is disjoint from $\{\alpha_1, \dots, \alpha_m, \beta_j\}$ for $j \neq i$. A connected component S' of $S_{0,p} \setminus \text{Int}N(\beta_1) \cup \text{Int}N(\beta_2)$ contains the linear chain $(\alpha_1, \dots, \alpha_m)$. Note that the connected component S' is a sphere with at most $p - 2$ holes. Hence, by [12, Theorem 2.2], the length of any linear chain on $S_{0,p} \setminus \text{Int}N(\beta_1) \cup \text{Int}N(\beta_2)$ is at most $p - 3$. Therefore, $m \leq p - 3$. \square

Lemma 3.30. *Suppose that $p \geq 2$. If $P_m^c * K_2 \leq \mathcal{C}(S_1^p)$, then $m \leq p$.*

Proof. We may assume that $m \geq p + 1 \geq 3$. Suppose that $P_m^c * K_2 \leq \mathcal{C}(S_1^p)$. Then, there is a linear chain $(\alpha_1, \dots, \alpha_m)$ and closed curves β_1, β_2 on $S_{1,p}$ such that β_i is isotopic to none of $\{\alpha_1, \dots, \alpha_m, \beta_j\}$ and is disjoint from $\{\alpha_1, \dots, \alpha_m, \beta_j\}$ for $i = 1, 2$ and $j \neq i$. A connected component of $S_{1,p} \setminus \text{Int}N(\beta_1) \cup \text{Int}N(\beta_2)$ contains the linear chain $(\alpha_1, \dots, \alpha_m)$. From results in [12, Theorem 2.2], it follows that the length of any linear chain on $S_{0,p} \setminus \text{Int}N(\beta_1) \cup \text{Int}N(\beta_2)$ is at most p . Therefore, we have $m \leq p$. \square

Lemma 3.31. *Suppose that $m \geq 1$, $g \geq 2$ and $p \geq 0$. If $P_{2g+p+1}^c * K_m \leq \mathcal{C}(S_g^p)$, then $m \leq g - 2$.*

Proof. We give a proof by induction on the genus g . Suppose that $P_{2g+p+1}^c * K_m \leq \mathcal{C}(S_g^p)$. Then, there are mutually non-isotopic closed curves $\alpha_1, \dots, \alpha_{2g+p+1}$, β_1, \dots, β_m satisfying the following conditions: (i) $\{\alpha_1, \dots, \alpha_{2g+p+1}\}$ is a linear chain of length $2g + p + 1$ and (ii) β_j is disjoint from the other curves $\alpha_1, \dots, \alpha_{2g+p+1}$ and $\beta_1, \dots, \beta_{j-1}, \beta_{j+1}, \dots, \beta_m$.

Case of $g = 2$, $p \geq 0$. Suppose, on the contrary, that $m \geq 1$. By conditions (i) and (ii), there is a connected component S' of $S_2^p \setminus \text{Int}N(\beta_1)$ containing a linear chain $(\alpha_1, \dots, \alpha_{p+5})$. However, [12, Theorem 2.2] implies that the length of any linear chain on S' is at most $p + 4$. Therefore, the initial assumption $m \geq 1$ must be false.

Case of $g \geq 3$, $p \geq 0$. Assuming $m \geq g - 2 \geq 1$, we will prove that $m = g - 2$. A connected component S' of $S_g^p \setminus \text{Int}N(\beta_1)$ contains a linear chain $(\alpha_1, \dots, \alpha_{2g+p+1})$. By [12, Theorem 2.2], we may assume that S' is homeomorphic to S_{g-1}^{p+2} . Hence, $P_{2g+p+1}^c * K_{m-1} \leq \mathcal{C}(S_{g-1}^{p+2})$. The induction hypothesis implies $m - 1 = (g - 1) - 2$, as desired. \square

Lemma 3.32. *Suppose that $m \geq 1$, $g \geq 2$ and $p \geq 0$. If $P_{2g+p}^c * K_m \leq \mathcal{C}(S_g^p)$, then $m \leq g - 1$.*

Proof. Suppose that $P_{2g+p} * K_m \leq \mathcal{C}(S_g^p)$. Then, we have mutually non-isotopic closed curves $\alpha_1, \dots, \alpha_{2g+p}, \beta_1, \dots, \beta_m$ satisfying the following conditions:

- (i) $(\alpha_1, \dots, \alpha_{2g+p})$ is a linear chain and
- (ii) β_j is disjoint from the other curves $\alpha_1, \dots, \alpha_{2g+p}, \beta_1, \dots, \beta_{j-1}, \beta_{j+1}, \dots, \beta_m$.

We give a proof by induction on the genus g .

Case of $g = 2, p \geq 0$. Suppose, on the contrary, that $m \geq 2 (> g - 1)$. From properties (i) and (ii), it follows that there is a connected component S' of $S_{2,p} \setminus \text{Int}N(\beta_1 \cup \beta_2)$ containing a linear chain $\{\alpha_1, \dots, \alpha_{p+4}\}$. We may assume that S' is homeomorphic to either S_2^{p-2} or S_1^{p+1} . However, neither of the surfaces contain a linear chain of length $p + 4$. Therefore, the initial assumption $m \geq 2$ must be false, and so $m \leq 1$.

Case of $g \geq 3, p \geq 0$. Assuming $m \geq g - 1 \geq 2$, we will prove that $m = g - 1$. From properties (i) and (ii), it follows that there is a connected component S' of $S_g^p \setminus \text{Int}N(\beta_1)$ containing a linear chain $(\alpha_1, \dots, \alpha_{2g+p})$. In the case where S' is a surface of genus g with $\leq p - 1$ holes, Lemma 3.31 implies $m - 1 \leq g - 2$, i.e., $m \leq g - 1$. Assume that S' is a surface of genus $\leq g - 1$ with $\leq p + 2$ holes. Since $P_{2g+p}^c * K_{m-1} \leq \mathcal{C}(S')$, we have that $m - 1 \leq (g - 1) - 1$ by the induction hypothesis. In every case, we have the desired inequality $m \leq g - 1$. \square

4. RIGHT-ANGLED ARTIN SUBGROUPS IN MAPPING CLASS GROUPS

For a simple finite graph Γ , the *right-angled Artin group* $A(\Gamma)$ is the group with the following finite presentation:

$$A(\Gamma) := \langle V(\Gamma) \mid v_i v_j = v_j v_i \text{ if } \{v_i, v_j\} \in E(\Gamma) \rangle.$$

Here, $V(\Gamma)$ is the vertex set of Γ and $E(\Gamma)$ is the edge set of Γ . The main purpose of this section is to prove Theorems 4.7 and 4.8. Deciding whether a right-angled Artin group can be embedded in the mapping class group of a surface is supposed to be difficult; at the moment the authors do not have an algorithm that solves the problem. Koberda in [14] proved that, for any family of mutually non-isotopic essential simple closed curves on a surface, sufficiently high powers of Dehn twists about the curves generate a right-angled Artin group in the mapping class group of the surface. This theorem is phrased as follows.

Theorem 4.1 (Koberda's embedding theorem). *Let S be a surface of negative Euler characteristic and let Γ be a finite graph. If $\Gamma \leq \mathcal{C}(S)$, then $A(\Gamma) \hookrightarrow \text{Mod}(S)$.*

Kim–Koberda in [13] introduced combinatorial embeddings of right-angled Artin groups into the mapping class groups of surfaces and proved that such embeddings are normal (Lemma 4.2 below). We say that a group homomorphism $\psi: A(\Lambda) \rightarrow \text{Mod}(S)$ satisfies *condition (KK)* if $\psi(v)$ is a product of mutually commutative Dehn twists for every $v \in V(\Lambda)$. Note that, if ψ satisfies condition (KK), the support curves of Dehn twists appearing in $\psi(v)$ induce a clique in $\mathcal{C}(S)$ if ψ satisfies condition (KK).

Lemma 4.2. *Let S be a surface of negative Euler characteristic and without a boundary and let Γ be a finite graph. If $A(\Gamma) \hookrightarrow \text{Mod}(S)$, then there is an embedding $A(\Gamma) \hookrightarrow \text{Mod}(S)$ that satisfies condition (KK).*

For an embedding $\psi: A(\Lambda) \hookrightarrow \text{Mod}(S)$ satisfying condition (KK) and a vertex u of Λ , $\text{supp}(\psi(u))$ denotes the subset of $V(\mathcal{C}(S))$ corresponding to the supports of

Dehn twists that appears in $\psi(u)$. To simplify the notation, we write $\text{supp}(\psi)$ for $\cup_{u \in V(\Lambda)} \text{supp}(\psi(u))$.

Theorem 4.3. *Let m be an integer ≥ 5 , S a surface of negative Euler characteristic and without a boundary, and Λ a graph isomorphic to C_m^c . Suppose that there is an embedding $\psi: A(\Lambda) \hookrightarrow \text{Mod}(S)$ with condition (KK). Then there is a full subgraph $\Delta \leq \mathcal{C}(S)$ that is induced by some vertices in $\text{supp}(\psi)$ and has exactly one of the following properties:*

- (1) $\Delta \cong C_n^c$ (for some $n \geq m$),
- (2) $\Delta \cong P_n^c$ (for some $n \geq m+1$),
- (3) $\Delta \cong P_{m-1}^c * K_m$. Furthermore, for each vertex v of Δ , there is a vertex in $\text{supp}(\psi)$ that is not adjacent to v .

Proof. By Lemma 4.2, we may assume that the embedding $\psi: A(\Lambda) \hookrightarrow \text{Mod}(S)$ satisfies condition (KK). We name the vertices u_1, u_2, \dots, u_m of $\Lambda \cong C_m^c$ to indicate that u_i is not adjacent to u_{i+1} ($1 \leq i \leq m-1$) and that u_m is not adjacent to u_1 . Let $\Gamma[\psi]$ denote the full subgraph induced by $\text{supp}(\psi)$ in $\mathcal{C}(S)$. There are m mutually distinct full embeddings $\iota_i: P_{m-1}^c \rightarrow C_m^c$, up to inversion. By [12, Proposition 3.9], each ι_i induces a full embedding $\tilde{\iota}_i: P_{m-1}^c \rightarrow \Gamma[\psi]$. We denote the image of $\tilde{\iota}_i$ by J_i . Since $J_i \cong P_{m-1}^c$,

- (*) for each vertex v of J_i , there is a vertex of J_i that is not adjacent to v in $\Gamma[\psi]$.

Assuming that neither assertion (1) nor (2) holds, we shall prove that assertion (3) holds. First, consider the case where J_1, \dots, J_m induces a full subgraph isomorphic to $*^m P_{m-1}^c$ in $\Gamma[\psi]$. Since $m \geq 5$, P_{m-1}^c contains a full subgraph isomorphic to K_2 . This implies that $P_{m-1}^c * K_{2m-2} \leq P_{m-1} * (*^{m-1} P_{m-1}^c) = *^m P_{m-1}$. Hence, we have an isomorphic copy Δ of $P_{m-1}^c * K_{2m-2}$ in $\Gamma[\psi]$ ($m \geq 5$ implies $2m-2 \geq m$). Moreover, since each J_i has the above property (*), for each vertex v of Δ , there is a vertex of $*_{i=1}^m J_i \leq \Gamma[\psi]$ that is not adjacent to v in $\Gamma[\psi]$. Therefore, assertion (3) holds.

Now, let us consider the case where some pair of J_1, \dots, J_m does not induce a graph-join in $\Gamma[\psi]$. Let ϕ be the correspondence between Λ and $\Gamma[\psi]$ such that $\phi \circ \iota_i = \iota_i$ given in [12, Proposition 3.9]. Then, the restriction of ϕ to $\cup_{i=1}^m J_i$ is a (surjective) map onto Λ . To simplify the notation, let ϕ denote this restriction map $\phi|_{\cup_{i=1}^m J_i}$ (we will use the restriction rather than the original correspondence). Moreover, ϕ is a graph embedding of $J_i^c \cong P_{m-1}$ into Λ . Also, by changing indices, if necessary, we may assume that J_1 and J_i ($i \neq 1$) does not induce a graph-join. In other words, there is a pair of vertices of J_1 and J_i that are not adjacent in $\Gamma[\psi]$. To prove assertion (3), we have to prove the following claim.

Claim 4.4. *The full subgraph induced by J_1 and J_i in $\Gamma[\psi]$ contains a full subgraph isomorphic to P_m^c .*

Proof of Claim 4.4. Let u_0 be the element of a singleton set $V(\iota_1(P_{m-1})) \setminus V(\iota_i(P_{m-1}))$. Pick a vertex v_0 of J_1 so that $\phi(v_0) = u_0$. Since $\phi \circ \tilde{\iota}_i = \iota_i$, it follows that $u_0 \notin \phi(V(J_i))$. Therefore, we have that $v_0 \notin J_i$. In the following, we use both $\Gamma[\psi]$ and its complement graph, $\Gamma[\psi]^c$, for convenience. Pick a sequence $J := (v_0, v_1, v_2, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ of vertices in $\Gamma[\psi]$ satisfying the following conditions.

- (a) $v_0, v_1, v_2, \dots, v_s \in V(J_1)$ and $v_{s+1}, v_{s+2}, \dots, v_{s+t} \in V(J_i)$,

- (b) v_j and v_{j+1} are adjacent in $\Gamma[\psi]^c$ (for all $0 \leq j \leq s+t-1$),
- (c) the vertices $v_0, v_1, v_2, \dots, v_s, v_{s+1}, \dots, v_{s+t}$ are mutually distinct,
- (d) $\phi(\{v_0, v_1, v_2, \dots, v_s, v_{s+1}, \dots, v_{s+t}\}) = V(\Lambda)$ and $\phi(v_0) \notin \phi(J_i)$,
- (e) $s = 0$, or J_i and the full subgraph induced by $\{v_0, v_1, \dots, v_{s-1}\}$ make a graph-join in $\Gamma[\psi]$,
- (f) J_1 and the full subgraph induced by $\{v_{s+2}, \dots, v_{s+t}\}$ make a graph-join in $\Gamma[\psi]$.

Before proving the existence of a sequence of vertices satisfying conditions (a), (b), (c), (d), (e) and (f), we prove that such a sequence induces a full subgraph isomorphic to P_m^c in $\Gamma[\psi]$. Assume that we have a sequence J as above. Conditions (a) and (d) imply $r := s+t+1 \geq m$. Hence, from conditions (b) and (c), we obtain a graph embedding τ of P_r into $\Gamma[\psi]^c$ that maps the vertices of P_r to J . Now, we prove that this embedding is full. Pick an edge e of $\Gamma[\psi]^c$ (not $\Gamma[\psi]$) joining two vertices of J . Consider the case where $\partial e \subset \{v_0, v_1, \dots, v_s\}$. Since J_1^c is a full subgraph of $\Gamma[\psi]^c$, we have that $e \in E(J_1^c)$. Then, conditions (a) and (b) imply $\partial e = \{v_j, v_{j+1}\}$ for some $0 \leq j \leq s$. Hence, e is contained in $\tau(P_r)$. Similarly, in the case where $\partial e \subset V(J_i^c)$, e is contained in $\tau(P_r)$. The remaining case is that $\partial e = \{v_x, v_y\}$ for some $0 \leq x \leq s$ and $s+1 \leq y \leq s+t$. Notice that $0 \leq x < s$ together with condition (e) implies $s = 0$ and $x = 0$, because $e \in \Gamma[\psi]^c$. Therefore we have $x = s$. Similarly, condition (f) implies $y = s+1$. Thus, e is contained in $\tau(P_r)$, and therefore τ is a full embedding. Consequently, τ induces a full embedding of P_r^c into $\Gamma[\psi]$ whose image is J .

Now, let us construct a sequence satisfying the desired conditions. Pick a vertex v_\star of $J_1^c \cong P_{m-1}$ such that v_\star is adjacent to some vertex of J_i^c in $\Gamma[\psi]^c$, and that the shortest path from v_0 to v_\star in J_1^c has no vertex adjacent to a vertex of J_i^c in $\Gamma[\psi]^c$ other than v_0 and v_\star (v_\star might be v_0). Let v_1, \dots, v_{s-1} be the vertices of J_1^c such that $(v_0, v_1, \dots, v_{s-1}, v_\star)$ is the shortest path from v_0 to v_\star in J_1^c . It follows that $s+1 \leq m-1$. We put $v_s := v_\star$. Since $\phi(\{v_0, v_1, \dots, v_{s-1}, v_s\})$ induces a path-graph on $\leq m-1$ vertices in $\Lambda^c = C_m$ and since $\phi(v_0) \notin \phi(V(J_i))$, at least one end point v^{end} of the path-graph J_i^c satisfies $\phi(v^{\text{end}}) \notin \{\phi(v_0), \phi(v_1), \dots, \phi(v_{s-1}), \phi(v_s)\}$. We should note that $\phi(v^{\text{end}})$ is adjacent to $\phi(v_0)$ in Λ^c . Let v^\star be the vertex of J_i^c which is adjacent to v_s in $\Gamma[\psi]^c$ and is nearest to v^{end} in J_i^c . Put $v_{s+1} := v^\star$ and pick the vertices $v_{s+2}, \dots, v_{s+t-1}$ of J_i^c in such a that $(v_{s+1}, v_{s+2}, \dots, v_{s+t-1}, v^{\text{end}})$ is the shortest path from v^{end} to v_{s+1} . Let us show that the sequence

$$(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t-1}, v_{s+t} := v^{\text{end}})$$

satisfies conditions (a), (b), (c), (d), (e) and (f).

Conditions (a) and (b): Obvious.

Condition (c): Since (v_0, v_1, \dots, v_s) is the shortest path from v_0 to v_s in J_1^c , any two vertices in $\{v_0, v_1, \dots, v_s\}$ are distinct. Similarly, any two vertices in $\{v_{s+1}, \dots, v_{s+t}\}$ are distinct. Moreover, since the only edge joining $\{v_1, \dots, v_s\}$ to $\{v_{s+1}, \dots, v_{s+t}\}$ in $\Gamma[\psi]^c$ is $[v_s, v_{s+1}]$, we have that $\{v_1, \dots, v_s\} \cap \{v_{s+1}, \dots, v_{s+t}\} = \emptyset$. Furthermore, v_0 is not contained in $\{v_{s+1}, \dots, v_{s+t}\}$, because $\phi(v_0)$ is not contained in $\phi(\{v_{s+1}, \dots, v_{s+t}\}) \subset \phi(V(J_i))$. Therefore, the vertices in

$$\{v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t-1}, v_{s+t}\}$$

are mutually distinct.

Condition (d): By definition, we have $\phi(v_0) = u_0 \notin \phi(J_i)$. So we have to prove that $\phi(\{v_0, v_1, v_2, \dots, v_s, v_{s+1}, \dots, v_{s+t}\}) = V(\Lambda)$. Since $\phi(v_0) = u_0$, it is enough

to show that $\phi(\{v_1, v_2, \dots, v_{s+t}\}) = V(\iota_i(P_{m-1}^c)) (= V(\Lambda) \setminus \{u_0\})$. Since v_{s+t} is an end point of J_i and since $v_0 \in V(\Lambda^c \setminus V(\iota_i(P_{m-1}^c)))$, $\phi(v_{s+t})$ is not adjacent to $\phi(v_0)$ in Λ . In addition, $\phi(v_1)$ is not adjacent to $\phi(v_0)$, because the restriction of ϕ to J_1^c is a graph homomorphism and that v_1 is not adjacent to v_0 . Therefore, $\phi(v_1)$ and $\phi(v_{s+t})$ are the end points of the path graph $\iota_i(P_{m-1})$. Hence, the path $(\phi(v_1), \phi(v_2), \dots, \phi(v_{s+t}))$ joins the end points of $\iota_i(P_{m-1})$ in $\Lambda^c \cong C_m$. Since the restriction of ϕ to J_1^c is a graph embedding and since $\phi(J_i) (= \iota(P_{m-1}))$ does not contain $u_0 = \phi(v_0)$, it holds that the path $(\phi(v_1), \phi(v_2), \dots, \phi(v_{s+t}))$ in Λ^c does not pass $\phi(v_0)$. Thus, $(\phi(v_1), \phi(v_2), \dots, \phi(v_{s+t}))$ covers $V(\iota(P_{m-1}^c))$, as desired.

Conditions (e) and (f): It is enough to show that v_{s+t} is adjacent to v_0 in $\Gamma[\psi]$. To this end, we shall show that $(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ induces a full subgraph isomorphic to C_{s+t+1}^c in $\Gamma[\psi]$ (assertion (1)) by assuming that v_{s+t} is not adjacent to v_0 in $\Gamma[\psi]$ (hence, they are adjacent in $\Gamma[\psi]^c$). This assumption together with the fact that $(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ satisfies conditions (b) and (c) implies that there is a natural graph embedding $\alpha: C_{s+t+1}^c \rightarrow \Gamma[\psi]^c$. Moreover, it follows that α is full from the choice of paths (v_0, v_1, \dots, v_s) and $(v_{s+t}, v_{s+t-1}, \dots, v_{s+1})$. Hence, $(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ induces a full subgraph isomorphic to C_{s+t+1}^c in $\Gamma[\psi]^c$. However, by assumption, $\Gamma[\psi]$ has no full subgraph isomorphic to C_{s+t+1}^c , and hence v_{s+t} is adjacent to v_0 in $\Gamma[\psi]$. Thus, the sequence $(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ satisfies conditions (e) and (f).

As shown above, the sequence $(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ with the above conditions induces a full subgraph isomorphic to P_{s+t+1}^c ($s+t+1 \geq m$) in $\Gamma[\psi]$. \square

We also need to prove the following claim.

Claim 4.5. *If $J_1^c \cup J_i^c \cup J_j^c$ ($j \neq 1$ and $j \neq i$) induces a connected full subgraph in $\Gamma[\psi]^c$, then the full subgraph of $\Gamma[\psi]$ induced by $J_1 \cup J_i \cup J_j$ contains a full subgraph isomorphic to P_{m+1}^c .*

Proof of Claim 4.5. Suppose that $J_1^c \cup J_i^c \cup J_j^c$ induces a connected full subgraph in $\Gamma[\psi]^c$. By changing indices, if necessary, we may assume that J_1^c and J_i^c are connected by an edge in $\Gamma[\psi]^c$, and that J_1^c and J_j^c are connected by an edge in $\Gamma[\psi]^c$. Let $(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ be the sequence of vertices in $V(J_1) \cup V(J_i)$ constructed in the proof of Claim 4.4. If $s+t+1 \geq m+1$; then, the claim holds. Thus, we may assume that $s+t+1 = m$. Then, condition (c) implies that the restriction of ϕ to the sequence $(v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ is a graph embedding of P_m^c into C_m^c . Moreover, as in the proof of Claim 4.4, $\phi(v_{s+t})$ is adjacent to $\phi(v_0)$ in $\Gamma[\psi]^c$.

(i) Case where J_1 has a vertex v_a such that $\phi(v_a) = \phi(v_{s+t})$. A key observation is that $\phi(E(J_1^c) \cup E(J_i^c)) = E(\Lambda^c) = E(C_m)$. We claim that $(v_a, v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t})$ induces a full subgraph isomorphic to P_{s+t+2}^c ($s+t+2 \geq m+1$) in $\Gamma[\psi]$. To see this, we will show the following:

- v_a is adjacent to v_0 in $\Gamma[\psi]^c$,
- $v_a \notin \{v_0, \dots, v_{s+t}\}$, and
- each vertex in $\{v_1, \dots, v_{s+t}\}$ is adjacent to v_a in $\Gamma[\psi]$.

These facts together with the fact that the sequence (v_0, \dots, v_{s+t}) induces a full subgraph isomorphic to P_m^c imply that the longer sequence $(v_a, v_0, \dots, v_{s+t})$ induces a full subgraph isomorphic to P_{m+1}^c in $\Gamma[\psi]$. First, we show that v_a is not adjacent to v_0 in $\Gamma[\psi]$. Since $\phi(v_a) = \phi(v_{s+t})$ and since $\phi(v_{s+t})$ is adjacent to $\phi(v_0)$ in Λ^c , $\phi(v_a)$ is adjacent to $\phi(v_0)$ in Λ^c . The restriction of ϕ to J_1^c is a

full embedding of J_1^c into Λ^c , because $J_1^c \cong P_{m-1}$ and $\phi \circ \tilde{\iota}_1 = \iota_1$. Therefore, v_a must be adjacent to v_0 . Next, we show that $v_a \notin \{v_0, \dots, v_{s+t}\}$. The path $(\phi(v_0), \dots, \phi(v_s))$ ($s+1 \leq m-1$) in Λ^c cannot reach $\phi(v_{s+t})$, because $\phi(v_0)$ is adjacent to $\phi(v_{s+t})$ in Λ^c and $\phi(v_1) \neq \phi(v_{s+t})$. Hence, we have $v_a \notin \{v_0, v_1, \dots, v_s\}$. In addition, no vertex in $\{v_{s+1}, v_{s+2}, \dots, v_{s+t-1}\}$ is mapped to $\phi(v_{s+t})$. Therefore, $v_a \notin \{v_{s+1}, v_{s+2}, \dots, v_{s+t-1}\}$. By assumption, $\Gamma[\psi]$ has no full subgraph isomorphic to C_m^c , and hence $v_a \neq v_{s+t}$ (if not, the path $(v_0, v_1, \dots, v_{s+t} = v_a)$ induces $C_{s+t+1} = C_m$ in $\Gamma[\psi]^c$). As a result, we have $v_a \notin \{v_0, v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t}\}$. Next, we show that each vertex in $\{v_1, \dots, v_s, v_{s+1}, \dots, v_{s+t}\}$ is adjacent to v_a in $\Gamma[\psi]$. Since $(\phi(v_0), \dots, \phi(v_{s+t}))$ induces an embedding of P_m into $\Lambda^c \cong C_m$, no vertex in $(\phi(v_1), \dots, \phi(v_{s+t-2}))$ is adjacent to $\phi(v_a)$ in Λ^c . In other words, each vertex in $(\phi(v_1), \dots, \phi(v_{s+t-2}))$ is adjacent to $\phi(v_a)$ in Λ . Since ϕ is a graph homomorphism from the full subgraph induced by $J_1^c \cup J_i^c$ in $\Gamma[\psi]^c$ to Λ^c , no vertex in (v_1, \dots, v_{s+t-2}) is adjacent to v_a in $\Gamma[\psi]^c$. Therefore, each vertex in (v_1, \dots, v_{s+t-2}) is adjacent to v_a in $\Gamma[\psi]$. Note that $(v_a, v_0, v_1, \dots, v_{s+t-2})$ induces a full subgraph isomorphic to P_{m-1}^c in $\Gamma[\psi]$. This implies that v_a is adjacent to v_{s+t-1} in $\Gamma[\psi]$, because there is no full subgraph isomorphic to C_m^c in $\Gamma[\psi]$. Also, v_a is adjacent to v_{s+t} , because ψ satisfies condition (KK) and $\phi(v_a) = \phi(v_{s+t})$. Thus each vertex in $\{v_1, \dots, v_{s+t}\}$ is adjacent to v_a in $\Gamma[\psi]$. The above argument shows that $(v_a, v_0, \dots, v_{s+t})$ induces a full subgraph isomorphic to P_{m+1}^c in $\Gamma[\psi]$.

(ii) Case where J_1 has no vertex mapped to $\phi(v_{s+t})$. In this case, it holds that $\phi(v_{s+t}) \notin \phi(J_1^c)$ and $\phi(v_0) \notin \phi(J_i^c)$. Hence, $\phi(E(J_1^c) \cup E(J_i^c)) \subsetneq E(\Lambda^c) = E(C_m)$. Note that J_j^c has an edge $[v_b, v_c]$ such that $\phi(v_b) = \phi(v_{s+t})$ and $\phi(v_c) = \phi(v_0)$. Suppose that J_j^c contains a vertex mapped to $\phi(v_{s+t-1})$. Accordingly, we have $\phi(E(J_1^c)) \cup \phi(E(J_j^c)) = E(\Lambda^c)$. By applying Claim 4.4 and the argument in (i) to $J_1 \cup J_j$ instead of $J_1 \cup J_i$, we have that the full subgraph induced by $J_1 \cup J_j$ contains P_{m+1}^c . Hence, we may assume that $\phi(J_j^c)$ does not contain $\phi(v_{s+t-1})$. Then we have that v_0 is an endpoint of J_1^c , v_b is an endpoint of J_j^c , and that $\phi(E(J_i^c)) \cup \phi(E(J_j^c)) = E(\Lambda^c)$. So again by Claim 4.4 and (i), we may assume that there is no edge between J_i^c and J_j^c in $\Gamma[\psi]^c$. We can now construct a path γ from $v_{s+t} \in J_i^c$ to an endpoint of J_j^c which passes J_1^c and $[v_b, v_c]$ in this order as follows.

Case (ii)-(A). Let v_e be an endpoint v_e of $J_1^c \cong P_{m-1}$ other than v_0 . If v_b is not adjacent to v_e in $\Gamma[\psi]^c$, then γ can be obtained by connecting $(v_{s+t}, \dots, v_{s+1}, v_s, \dots, v_d)$ with a subpath S of J_j^c starting from an internal vertex of J_j^c and ending at v_b , where $v_d \in \{v_0, v_1, \dots, v_s\}$ and no vertex between v_s and v_d is adjacent to a vertex of J_j^c in $\Gamma[\psi]^c$.

Case (ii)-(B). If v_b is adjacent to v_e in $\Gamma[\psi]^c$, then γ can be obtained by connecting the shortest path from v_{s+t} to v_e in the full subgraph induced by $J_1^c \cup J_i^c$ with the entire J_j^c .

Let us show that γ induces a full subgraph P isomorphic to a path graph of length at least $m+1$ in $\Gamma[\psi]^c$. By our construction, $\phi(\gamma)$ passes $\phi(v_{s+t})$ twice and all of the vertices in Λ^c , so the number of vertices contained in γ is at least $m+1$. It is easy to see that γ does not have an overlap. Moreover, we can show that γ has no shortcut as follows.

Case (ii)-(A). Both $(v_{s+t}, \dots, v_{s+1}, v_s, \dots, v_c)$ and S have no shortcut. Since there is no edge between J_i^c and J_j^c , no vertex of S is adjacent to $(v_{s+t}, \dots, v_{s+1})$ in $\Gamma[\psi]^c$. By the properties of v_d , there is no edge between S and (v_s, \dots, v_d) .

Case (ii)-(B). Suppose, on the contrary, that there is an edge e connecting a vertex of J_1^c and a vertex of J_j^c other than v_b . Then $J_1^c \cup J_j^c$ contains a full subgraph isomorphic to C_n for some $n \geq m$, because $[v_e, v_b]$ is an edge in $\Gamma[\psi]^c$. This does not happen, so there is exactly one edge, $[v_e, v_b]$, between J_1^c and J_j^c . This implies that there is no shortcut in γ , because there is no edge between J_i^c and J_j^c .

By the above argument, γ induces a full subgraph isomorphic to P_n for some $n \geq m + 1$, as desired. \square

We denote by F the full subgraph of $\Gamma[\psi]^c$ induced by J_1^c, \dots, J_m^c . Since two of J_1^c, \dots, J_m^c , say J_1^c and J_i^c , are connected by an edge in $\Gamma[\psi]^c$, Claim 4.4 implies that F contains a full subgraph $P_m \sqcup \check{F}$, where \check{F} is a full subgraph of $\Gamma[\psi]^c$ induced by $J_2^c, \dots, J_{i-1}^c, J_{i+1}^c, \dots, J_m^c$. Claim 4.5 together with the assumption $P_{m+1}^c \not\leq \Gamma[\psi]$ ensures that no triple of J_1^c, \dots, J_m^c is connected in $\Gamma[\psi]^c$. Hence, there is no edge between the isomorphic copy of P_m and \check{F} in the above union. Let x denote the number of connected components of F induced by pairs of J_1^c, \dots, J_m^c , and let y denote the number of other connected components of F . Then, we have $2x + y = m$. In order to prove that there is a full subgraph isomorphic to $P_m \sqcup (\sqcup^m K_1)$ in F , we will find $\sqcup^m K_1$ in \check{F} . Note that \check{F} has $x - 1$ connected components that contain full subgraphs isomorphic to P_m and y connected components that do not contain P_m (but contain P_{m-1}). Since $m \geq 5$, P_m contains a full subgraph isomorphic to $\sqcup^3 K_1$. On the other hand, P_{m-1} contains a full subgraph isomorphic to $\sqcup^2 K_1$. Therefore, \check{F} contains a full subgraph isomorphic to $\sqcup^{3(x-1)+2y} K_1$. The equality $2x + y = m$ implies $3(x-1) + 2y = \frac{3m+y-6}{2}$. We now claim that $\frac{3m+y-6}{2} \geq m$. To see this, assume that $m = 5$. Then, at least one component should be induced by a single J_k , and hence $y \geq 1$. Therefore, we have $\frac{3m+y-6}{2} \geq m$. Next, we assume that $m \geq 6$. Then we have $3m - 6 \geq 2m$, and hence, $\frac{3m+y-6}{2} \geq m$. Thus, $\frac{3m+y-6}{2} \geq m$ for all $m \geq 5$. Now it follows that $P_m \sqcup (\sqcup^m K_1) \leq P_m \sqcup (\sqcup^{\frac{3m+y-6}{2}} K_1) \leq P_m \sqcup \check{F} \leq F \leq \Gamma[\psi]^c$. Namely, we have $P_m^c * K_m \leq \Gamma[\psi]$. Property (*) implies that for each vertex v of the full subgraph $P_m^c * K_m \leq \Gamma[\psi]$, there is a vertex of $\Gamma[\psi]$, which is not adjacent to v in $\Gamma[\psi]$, and therefore assertion (3) holds. \square

The above theorem immediately implies the following.

Corollary 4.6. *Let m be an integer ≥ 5 , S a surface with negative Euler characteristic and without boundary, and Γ a finite graph with $P_{m+1}^c \not\leq \mathcal{C}(S)$. Suppose that there is an embedding $\psi: A(C_m^c) \times A(K_1) \hookrightarrow \text{Mod}(S)$. Then at least one of the following holds:*

- (1) $C_n * K_1 \leq \mathcal{C}(S)$ (for some $n \geq m$),
- (2) $P_{m-1}^c * K_{m+1} \leq \mathcal{C}(S)$.

Proof. We may assume that ψ satisfies condition (KK). Restricting ψ to $A(C_m^c)$ and $A(K_1)$, we have two embeddings $\psi_1 := \psi|_{A(C_m^c)}: A(C_m^c) \hookrightarrow \text{Mod}(S)$ and $\psi_2 := \psi|_{A(K_1)}: A(K_1) \hookrightarrow \text{Mod}(S)$ with condition (KK). Applying Theorem 4.3 to ψ_1 , we have that there is a full subgraph Δ of $\mathcal{C}(S)$ that is induced by some vertices in $\text{supp}(\psi_1(V(C_m^c)))$ and is isomorphic to one of the following:

- (1') C_n (for some $n \geq m$) or
- (2') $P_{m-1} * K_m$. Furthermore, for each vertex v of $P_{m-1} * K_m$, there is a vertex in $\text{supp}(\psi_1(V(C_m^c)))$ that is not adjacent to v .

We only treat the case where Δ is isomorphic to $P_{m-1} * K_m$, because another case can be treated similarly. Suppose that Δ is isomorphic to $P_{m-1} * K_m$. Pick a vertex u of $\text{supp}(\psi_2(V(K_1)))$. Then every element of $\text{supp}(\psi_1(V(C_m^c)))$ is adjacent to u in $\mathcal{C}(S)$, because the images of ψ_1 and ψ_2 are commutative. So we have $u \notin V(\Delta)$, and hence, $P_{m-1}^c * K_{m+1} \cong P_{m-1}^c * K_m * \{u\} \leq \mathcal{C}(S)$. Thus, assertion (2) holds. \square

Using the above results, we can decide whether $A(C_m^c)$ is embedded in the mapping class group of a surface.

Theorem 4.7. *$A(C_m^c)$ is embedded in $\text{Mod}(S_{g,p})$ if and only if m satisfies*

$$m \leq \begin{cases} 0 & ((g, p) \in \{(0, 0), (0, 1), (0, 2), (0, 3)\}) \\ 3 & ((g, p) \in \{(0, 4), (1, 0), (1, 1)\}) \\ 5 & ((g, p) = (1, 2), (0, 5)) \\ 2g + 2 & (g \geq 2, p = 0) \\ 2g + p + 1 & (g \geq 2, 1 \leq p \leq 2) \\ 2g + p & (\text{otherwise}). \end{cases}$$

Proof. **Case of $(g, p) \in \{(0, 0), (0, 1), (0, 2), (0, 3)\}$.** In this case, the mapping class group $\text{Mod}(S_{g,p})$ is a finite group, and so the right-angled Artin group embedded in $\text{Mod}(S_{g,p})$ should be trivial.

Case of $(g, p) = (1, 1)$. Note that $A(C_3^c) \cong F_3$ is embedded in $\text{Mod}(S_{1,1}) \cong \text{SL}(2, \mathbb{Z})$. However, if $m \geq 4$, then $A(C_m^c)$ contains \mathbb{Z}^2 . On the other hand, $\text{SL}(2, \mathbb{Z})$ does not contain \mathbb{Z}^2 . Hence, the assertion holds.

Case of $(g, p) \in \{(0, 5), (1, 2)\}$. The assertion in the case where $(g, p) = (0, 5)$, was proved in [12, Proposition 4.1]. Suppose that $(g, p) = (1, 2)$. Koberda's embedding theorem together with Lemma 3.27 shows that $A(C_5^c)$ is embedded in $\text{Mod}(S_{1,2})$. Next, we prove that $A(C_m^c) \hookrightarrow \text{Mod}(S_{1,2})$ implies $m \leq 5$. Note that $A(C_6^c)$ contains a subgroup isomorphic to \mathbb{Z}^3 . On the other hand, the topological complexity of $S_{1,2}$ is two. Hence, $A(C_6^c)$ is cannot be embedded in $\text{Mod}(S_{1,2})$.

Case of $g \geq 2, p = 0$. This is shown in [12, Proposition 4.1].

Case of $g \geq 2, 1 \leq p \leq 2$. Koberda's embedding theorem together with Lemma 3.27 shows that $A(C_{2g+p+1}^c)$ is embedded in $\text{Mod}(S_{g,p})$. Suppose that $A(C_m^c) \hookrightarrow \text{Mod}(S_{g,p})$. Assuming $m \geq 2g + p + 1$, we shall prove $m = 2g + p + 1$. By [12], P_{m+1}^c is not a full subgraph of $\mathcal{C}(S_g^p)$. Hence, from Theorem 4.3, we have either $C_m^c \leq \mathcal{C}(S_{g,p})$ or $P_{m-1}^c * K_m \leq \mathcal{C}(S_g^p)$. Lemma 3.27 together with Lemma 3.32 implies that $m = 2g + p + 1$.

Case of $g = 0, p \geq 6$. See [12, Proposition 4.1].

In the rest of the cases, we can give proofs similar to the case where $g \geq 2, 1 \leq p \leq 2$.

Case of $g = 1, p \geq 3$. Koberda's embedding theorem together with Lemma 3.27 shows that $A(C_{p+2}^c)$ is embedded in $\text{Mod}(S_{1,p})$. Next, we suppose that $A(C_m^c) \hookrightarrow \text{Mod}(S_{1,p})$. Furthermore, assume that $m \geq p + 2 \geq 5$. Since $P_{p+3}^c \not\leq \mathcal{C}(S_1^p)$ by [12, Theorem 2.2], either $C_m^c \leq \mathcal{C}(S_{1,p})$ or $P_{m-1}^c * K_m \leq \mathcal{C}(S_{1,p})$ holds. Hence, $m = p + 2$.

Case of $g \geq 2, p \geq 3$. Koberda's embedding theorem together with Lemma 3.27 shows that $A(C_{2g+p}^c)$ is embedded in $\text{Mod}(S_{g,p})$. Suppose that $A(C_m^c) \hookrightarrow \text{Mod}(S_{g,p})$. To the contrary, assume that $m \geq 2g + p + 1$. Then, one of the following holds: $C_m^c \leq \mathcal{C}(S_{g,p})$ or $P_{2g+p+1}^c \leq \mathcal{C}(S)$ or $P_{2g+p}^c * K_m \leq \mathcal{C}(S_g^p)$. However, this is impossible by [12] and Lemmas 3.27 and 3.32. Thus, $m \leq 2g + p$. \square

Similarly, we can decide whether $A(C_m^c) \times \mathbb{Z}$ is embedded in the mapping class group of a surface.

Theorem 4.8. $A(C_m^c) \times \mathbb{Z}$ is embedded in $\text{Mod}(S_{g,p})$ if and only if m satisfies

$$m \leq \begin{cases} 0 & (g, p) \in \{(0, 4), (1, 1)\} \\ 3 & (g, p) \in \{(0, 5), (1, 2)\} \\ p-1 & (g=0, p \geq 6) \\ p+2 & (g=1, p \geq 3) \\ 2g+1 & (g \geq 2, p=0) \\ 2g+p & (g \geq 2, p \geq 1). \end{cases}$$

Proof. Koberda's embedding theorem together with Lemma 3.28 and [12, Proposition 4.1] implies the “if part” of the theorem. Thus, we will prove the “only if part”.

Case of $(g, p) \in \{(0, 4), (1, 1)\}$. If $m \geq 1$, then $A(C_m^c) \times \mathbb{Z}$ contains \mathbb{Z}^2 . On the other hand, $\text{Mod}(S_{0,4})$ and $\text{Mod}(S_{1,1})$ do not contain \mathbb{Z}^2 . Hence, $A(C_m^c) \times \mathbb{Z}$ is embedded in neither $\text{Mod}(S_{0,4})$ nor $\text{Mod}(S_{1,1})$ if $m \geq 1$.

Case of $(g, p) \in \{(0, 5), (1, 2)\}$. Since the topological complexity of $S_{g,p}$ is two, the right-angled Artin group $A(C_4^c) \times \mathbb{Z}$ is not embedded in $\text{Mod}(S_{g,p})$.

Case of $g=0, p \geq 6$. Suppose that $A(C_m^c) \times \mathbb{Z}$ is embedded in $\text{Mod}(S_{0,p})$. We may assume that $m \geq p-1 \geq 5$. Then, there is an embedding $A(C_m^c) \times \mathbb{Z} \hookrightarrow \text{Mod}(S_{0,p})$ satisfying condition (KK). As shown in [12], we have $P_{m+1}^c \not\leq \mathcal{C}(S_0^p)$. Then, from Corollary 4.6, either $C_m^c * K_1 \leq \mathcal{C}(S_{0,p})$ or $P_{p-2}^c * K_{m+1} \leq \mathcal{C}(S_0^p)$ holds. Hence, by Lemmas 3.28 and 3.29, we have $m \leq p-1$.

The remaining cases can be treated similarly.

Case of $g=1, p \geq 3$. Suppose that $A(C_m^c) \times \mathbb{Z}$ is embedded in $\text{Mod}(S_{1,p})$. We may assume that $m \geq p+2 \geq 5$. Then there is an embedding $A(C_m^c) \times \mathbb{Z} \hookrightarrow \text{Mod}(S_{0,p})$ satisfying condition (KK). By [12], we have $P_{m+1}^c \not\leq \mathcal{C}(S_g^p)$. From Corollary 4.6, it follows that either $C_m^c * K_1 \leq \mathcal{C}(S_1^p)$ or $P_{p+1}^c * K_{m+1} \leq \mathcal{C}(S_1^p)$. By Lemmas 3.28 and 3.30, we have $m \leq p+2$.

Case of $g \geq 2, p=0$. From Lemma 3.31, it routinely follows that $P_{2g} * K_{2g+1} \not\leq \mathcal{C}(S_g)$. In addition, $C_m^c * K_1 \leq \mathcal{C}(S_g)$ only if $m \leq 2g+1$.

Case of $g \geq 2, p \geq 1$. Suppose that $A(C_m^c) \times \mathbb{Z}$ is embedded in $\text{Mod}(S_{g,p})$. Assume to the contrary that $m \geq 2g+p+1$. There is an embedding $A(C_m^c) \times \mathbb{Z} \hookrightarrow \text{Mod}(S_{g,p})$ satisfying condition (KK). Since $P_{2g+p+2}^c \not\leq \mathcal{C}(S_g^p)$ by [12, Theorem 2.2], either $C_m^c * K_1 \leq \mathcal{C}(S_g^p)$ or $P_{2g+p}^c * K_{2g+p+2} \leq \mathcal{C}(S_g^p)$ holds. From Lemma 3.31, it follows that $P_{2g+p}^c * K_{2g+p+1} \not\leq \mathcal{C}(S_g^p)$. In addition, $C_m^c * K_1 \not\leq \mathcal{C}(S_g^p)$. This is a contradiction. Thus, we have $m \leq 2g+p$. \square

Remark 4.9. Apart from a few exceptions, it holds that $A(C_m^c) \hookrightarrow \text{Mod}(S_{g,p})$ if and only if $C_m^c \leq \mathcal{C}(S_g^p)$. Similarly, $A(C_m^c) \times \mathbb{Z} \hookrightarrow \text{Mod}(S_{g,p})$ if and only if $C_m^c * K_1 \leq \mathcal{C}(S_g^p)$.

We end this section by noting a result for braid groups.

Theorem 4.10 ([12]). Suppose $n \geq 3$. $A(C_{n+1}^c) \times \mathbb{Z} \hookrightarrow B_n$.

5. PROOFS OF THEOREMS 1.1 AND 1.2

In this section, we prove Theorems 1.1 and 1.2. Let us start with the following lemma.

Lemma 5.1 ([3]). *Let A be a right-angled Artin group and G a group. If A is embedded in G , then A is also embedded in each finite index subgroup of G .*

The following is the “only if part” of Theorem 1.1.

Theorem 5.2. *If the braid group B_n is virtually embedded in $\text{Mod}(S_{g,p})$, then*

$$n \leq \begin{cases} 1 & ((g,p) \in \{(0,0), (0,1), (0,2), (0,3)\}) \\ 2 & ((g,p) \in \{(0,4), (1,0), (1,1)\}) \\ 3 & ((g,p) \in \{(0,5), (1,2)\}) \\ p-2 & (g=0, p \geq 6) \\ p+1 & (g=1, p \geq 3) \\ 2g & (g \geq 2, p=0) \\ 2g+p-1 & (g \geq 2, p \geq 1). \end{cases}$$

Proof. Suppose that B_n is virtually embedded in $\text{Mod}(S_{g,p})$. First, we treat the cases where surfaces have complexities less than 3.

Case of $(g,p) \in \{(0,0), (0,1), (0,2), (0,3)\}$. In this case $\text{Mod}(S_{g,p})$ is a finite group, so there is no virtual embedding of B_2 into $\text{Mod}(S_{g,p})$.

Case of $(g,p) \in \{(0,4), (1,0), (1,1)\}$. Since the mapping class group of $S_{g,p}^b$ is Gromov hyperbolic and since every finite index subgroup of B_3 contains a free abelian group of rank 2, B_3 has no finite index subgroup embedded in the mapping class group.

Case of $(g,p) \in \{(0,5), (1,2)\}$. Since \mathbb{Z}^3 is not embedded in $\text{Mod}(S_{g,p})$, B_4 has no finite index subgroup embedded in $\text{Mod}(S_{g,p})$. Hence, the desired inequality holds.

In the remaining cases, we may assume that $n \geq 3$. Then, since $A(C_{n+1}^c) \times \mathbb{Z}$ is embedded in all finite index subgroups of B_n , $A(C_{n+1}^c) \times \mathbb{Z}$ is embedded in $\text{Mod}(S_{g,p})$. We use Theorem 4.8 to obtain the desired inequalities.

Case of $g=0$ and $p \geq 6$. We have $n+1 \leq p-1$.

Case of $g=1$ and $p \geq 3$. We have $n+1 \leq p+2$.

Case of $g \geq 2$ and $p=0$. We have $n+1 \leq 2g+1$.

Case of $g \geq 2$ and $p \geq 1$. We have $n+1 \leq 2g+p$. □

The following lemma is convenient.

Lemma 5.3. *Suppose $n \geq 4$. Suppose also that B_n is virtually embedded in $\text{Mod}(S_{g,p}^b)$. Then, $A(C_{n+1}^c)$ is embedded in $\text{Mod}(S_{g,p+b})$.*

Proof. The kernel of the capping homomorphism $\text{Mod}(S_{g,p}^b) \rightarrow \text{Mod}(S_{g,p+b})$ is generated by the center. Since C_{n+1}^c has no vertex adjacent to the other vertices, the center of $A(C_{n+1}^c)$ is trivial. This is an immediate corollary of Servatius’ centralizer theorem [18] which implies the centralizer of a vertex v is generated by v and the vertices adjacent to v . Hence, we obtain an embedding of $A(C_{n+1}^c)$ into $\text{Mod}(S_{g,p+b})$. □

The following is the “only if part” of Theorem 1.2.

Theorem 5.4. *Suppose that $b \geq 1$. Suppose also that the braid group B_n is virtually embedded in $\text{Mod}(S_{g,p}^b)$. Then,*

$$n \leq \begin{cases} 2g+p+b & (g \geq 1, p+b \leq 2) \\ 2g+p+b-1 & (\text{otherwise}). \end{cases}$$

Proof. Suppose that B_n is virtually embedded in $\text{Mod}(S_{g,p}^b)$.

Case of $g = 0, p \geq 0, b = 1$. In this case $\text{Mod}(S_{g,p}^1)$ is isomorphic to B_p . The inequality $n \leq p (= 2g + p + b - 1)$ immediately follows from the fact that $\mathbb{Z}^l \hookrightarrow B_m$ if and only if $l \leq m - 1$.

Case of $g = 0$ and $p \geq 0, b \geq 2$. Suppose that $p + b \leq 3$. Then, $(p, b) \in \{(0, 2), (1, 2), (0, 3)\}$, and so $\text{Mod}(S_{0,p}^b)$ is a free abelian group. Since B_3 has no finite index abelian subgroup, B_3 is not embedded in $\text{Mod}(S_{0,p}^b)$ even virtually. Therefore $n \leq 2 (= 2g + p + b - 1)$. Next, consider the case where $p + b = 4$. Since $\text{Mod}(S_{0,4})$ is virtually free, $\text{Mod}(S_{0,p}^b)$ is a central extension of a virtually free group. This implies that any centerless right-angled Artin subgroup of B_n (embedded in $\text{Mod}(S_{0,p}^b)$) must be free. Since B_4 contains $A(P_4^c)$ which is centerless but not free, we have $n \leq 3$. Suppose that $p + b \geq 5$. Assuming $n \geq 4$, we shall prove that $n \leq p + b - 1$. By Lemma 5.3, $A(C_{n+1}^c)$ is embedded in $\text{Mod}(S_{0,p+b})$. Theorem 4.7 implies that $n + 1 \leq p + b$.

Case of $g = 1, p = 0, b \leq 2$. In this case \mathbb{Z}^{b+2} cannot be embedded in $\text{Mod}(S_1^b)$. Hence, no finite index subgroup of B_{b+3} is embedded in $\text{Mod}(S_1^b)$.

Case of $g \geq 2, p = 0, b \leq 2$. We may assume that $n \geq 4$. By Lemma 5.3, $A(C_{n+1}^c)$ is embedded in $\text{Mod}(S_{g,b})$. Since $1 \leq b \leq 2$, we have $n + 1 \leq 2g + b + 1$ by Theorem 4.7. When $b \geq 3$, we have $n + 1 \leq 2g + b$.

Case of $g \geq 1, p = 1, b = 1$. We may assume that $n \geq 4$. By Lemma 5.3 and Theorem 4.7, we have $n + 1 \leq 2g + p + b + 1$.

Case of $g \geq 1, p + b \geq 3$. We may assume that $n \geq 4$. By Lemma 5.3 and Theorem 4.7, we have $n + 1 \leq 2g + p + b$. \square

Below we summarize topological methods for constructing virtual embeddings of braid groups into the mapping class groups of surfaces in Theorems 5.5 and 5.6.

Theorem 5.5. *Suppose that either $b \neq 1$ or $p \neq 1$ holds. Let S denote the surface $S_{g,p}^b$. Then the inequality in Theorems 1.1 and 1.2 holds if and only if there is at least one of the following:*

- (1) *an extension $D \rightarrow S$ which is either trivial or hyperbolic,*
- (2) *an extension $\tilde{D} \rightarrow S$ which is trivial, annular, or hyperbolic,*
- (3) *a composition of extensions $D \rightarrow P \rightarrow S$, where the middle surface P is a sphere, the first extension is hyperbolic and the second is pseudo-annular.*

Here, D is the defining surface $S_{0,n}^1$ of B_n and \tilde{D} is a double branched cover of D , which is homeomorphic to either S_k^2 ($k = \frac{n}{2} - 1$) or S_l^1 ($l = \frac{n-1}{2}$) according to whether n is even or odd.

Proof. Suppose that there is at least one of (1), (2) or (3). Then we can find a virtual embedding of the braid group B_n described in Section 2. Hence, Theorems 5.2 and 5.4 imply that the inequality on n in Theorems 1.1 and 1.2 holds.

We shall prove the “only if part” of this theorem.

Case of $S \cong S_{0,p}$. Suppose $n \leq p - 2$. Then, there is a hyperbolic extension of D to S , so we have (1).

Case of $S \cong S_{1,p}$. Suppose $n \leq p + 1$. Consider a pseudo-annular extension $S_{0,p+1}^1 \rightarrow S$. This is obtained from the compactification S_0^{p+2} of $S_{0,p+1}^1$, a single annulus and p copies of a once-punctured disks. Then, this pseudo-annular extension together with a hyperbolic extension $D \rightarrow S_{0,p+1}^1$ satisfies (3).

Case of $S \cong S_g$ with $g \geq 2$. Suppose $n \leq 2g$. In this case we have an annular extension $S_{g-1}^2 \rightarrow S$. Since $n \leq 2g$, the natural numbers k and l defined in the assertion are both less than g . Hence, we have an extension $\tilde{D} \rightarrow S_{g-1}^2$ which is either trivial (this case occurs if $n = 2g$) or hyperbolic. So we have (2).

Case of $S \cong S_{g,p}$ with $g \geq 2$ and with $p \geq 1$. Suppose $n \leq 2g+p-1$. Consider a pseudo-annular extension $S_{0,2g+p-1}^1 \rightarrow S$ obtained from the compactification S_0^{2g+p} , g copies of an annulus and p copies of a once-punctured disk. Since $n \leq 2g+p-1$, there is a hyperbolic extension $D \rightarrow S_{0,2g+p-1}^1$. So we have (3).

Case of $S \cong S_{0,p}^1$. Suppose $n \leq p$. Then we can find an extension $D \rightarrow S$ that is either trivial or hyperbolic. So we have (1).

Case of $S \cong S_{0,p}^b$ with $b \geq 2$. Suppose $n \leq p+b-1$. Then, there is a pseudo-annular extension $S_{0,p+b-1}^1 \rightarrow S$ obtained from S_0^{p+b} , $b-1$ copies of an annulus, and p copies of a once-punctured disks. Since $n \leq p+b-1$, there is a hyperbolic (or trivial) extension $D \rightarrow S_{0,p+b-1}^1$. So we have (3).

Case of $S \cong S_g^b$ with $g \geq 1$ and with $1 \leq b \leq 2$. Suppose $n \leq 2g+b$. Then, S is a double branched covering of $S_{0,2g+b}^1$. From a hyperbolic (resp. trivial) extension $D \rightarrow S_{0,2g+b}^1$, we have a hyperbolic (resp. trivial) extension $\tilde{D} \rightarrow S$. So we have (2).

Case of $S \cong S_{g,p}^b$ with $g \geq 1$ and with $p+b \geq 3$. Suppose $n \leq 2g+p+b-1$. Consider a pseudo-annular extension $S_{0,2g+p+b-1}^1 \rightarrow S$ obtained from S_0^{2g+p+b} , g copies of an annulus and p copies of a once-punctured disk. A combination of this pseudo-annular extension and a hyperbolic extension $D \rightarrow S_{0,2g+p+b-1}^1$ satisfies (3) and completes the proof. \square

Note that Theorem 5.5 proves the “if part” of Theorems 1.1 and 1.2 when $S \not\cong S_{g,1}^1$.

Theorem 5.6. *Suppose that $g \geq 1$. Then $n \leq 2g+2$ if and only if one of the following holds:*

- (1) $\tilde{D} \cong S_g^2 \xrightarrow{\text{cap}} S_{g,1}^1$ or
- (2) *there is a hyperbolic extension $\tilde{D} \rightarrow S_g^2$ together with $S_g^2 \xrightarrow{\text{cap}} S_{g,1}^1$.*

Here, \tilde{D} is a Birman–Hilden cover of $S_{0,n}^1$, which is homeomorphic to either S_k^2 ($k = \frac{n}{2} - 1$) or S_l^1 ($l = \frac{n-1}{2}$) according to whether n is even or odd, and $S_g^2 \xrightarrow{\text{cap}} S_{g,1}^1$ is an admissible extension obtained from S_g^2 and a once-punctured disk by gluing along a boundary component of S_g^2 and the boundary of the once-punctured disk.

Proof. Suppose that (1) or (2) holds. Then the entire braid group B_n is embedded in $\text{Mod}(S_g^2)$. Hence, Theorem 5.4 implies $n \leq 2g+2$.

Next, let us assume that $n \leq 2g+2$ and $\tilde{D} \not\cong S_g^2$ (so (1) does not hold). Since $\tilde{D} \not\cong S_g^2$, we have $n \leq 2g+1$. So we can find a hyperbolic extension $\tilde{D} \rightarrow S_g^2$. \square

Recall that B_n is virtually embedded in $\text{Mod}(S_{g,1}^1)$ only if $n \leq 2g+2$ by Theorem 5.4. Let us prove the converse in accordance with Theorem 5.6.

Theorem 5.7. *B_n is virtually embedded in $\text{Mod}(S_{g,1}^1)$ if $n \leq 2g+2$.*

Proof. Suppose $n \leq 2g+2$. Then we have an embedding $B_{2g+2} \hookrightarrow \text{Mod}(S_g^2)$ induced by a double branched covering. Moreover, from Proposition 2.6 we have

an embedding $B_{2g+2} \rightarrow \text{Mod}(S_{g,1}^1)$. Now the embedding $B_n \hookrightarrow B_{2g+2}$ induced by a hyperbolic extension completes the proof. \square

Consequently, the “if part” of Theorems 1.1 and 1.2 is also true.

Furthermore, our argument allows us to obtain the following results on embeddings of pure braid groups.

Theorem 5.8. $PB_n \hookrightarrow \text{Mod}(S_{g,p})$ if and only if

$$n \leq \begin{cases} 1 & ((g,p) \in \{(0,0), (0,1), (0,2), (0,3)\}) \\ 2 & ((g,p) \in \{(0,4), (1,0), (1,1)\}) \\ -\chi_{g,p} & (g=0, p \geq 5) \\ 2 - \chi_{g,p} & (g \geq 2, p=0) \\ 1 - \chi_{g,p} & (\text{otherwise}). \end{cases}$$

Theorem 5.9. Suppose $b \geq 1$. Then $PB_n \hookrightarrow \text{Mod}(S_{g,p}^b)$ if and only if

$$n \leq \begin{cases} 2 - \chi_{g,p}^b & (g \geq 1, p+b \leq 2) \\ 1 - \chi_{g,p}^b & (\text{otherwise}). \end{cases}$$

REFERENCES

- [1] J. Aramayona, C. Leininger and J. Souto, *Injections of mapping class groups*, Geom. Topol. **13** (2009), no. 5, 2523–2541.
- [2] J. Aramayona and J. Souto, *Homomorphisms between mapping class groups*, Geom. Topol. **16** (2012), no. 4, 2285–2341.
- [3] H. Baik, S. Kim and T. Koberda, *Unsmoothable group actions on compact one-manifolds*, J. Eur. Math. Soc. **21** (2019), 2333–2353.
- [4] J. Behrstock, B. Kleiner, Y. Minsky and L. Mosher, *Geometry and rigidity of mapping class groups*, Geom. Topol. **16** (2012), no. 2, 781–888.
- [5] J. Birman and H. Hilden, *On isotopies of homeomorphisms of Riemann surfaces*, Ann. of Math. (2) **97** (1973), no. 3, 424–439.
- [6] J. Birman, A. Lubotzky and J. McCarthy, *Abelian and solvable subgroups of the mapping class groups*, Duke Math. J. **50** (1983), no. 4, 1107–1120.
- [7] F. Castel, *Geometric representations of the braid groups*, Asterisque, No. 378 (2016), vi+175 pp. ISBN: 978-2-85629-835-0.
- [8] M. Clay, C. Leininger and D. Margalit, *Abstract commensurators of right-angled Artin groups and mapping class groups*, Math. Res. Lett. **21** (2014), no. 3, 461–467.
- [9] U. Hamenstädt, *Geometry of the mapping class groups III: Quasi-isometric rigidity*, preprint, arXiv: math/051242.
- [10] J. Harer, *The virtual cohomological dimension of the mapping class group of an orientable surface*, Invent. Math. **84** (1986), 157–176.
- [11] N. Ivanov and J. McCarthy, *On injective homomorphisms between Teichmüller modular groups. I*, Invent. Math. **135** (1999), 425–486.
- [12] T. Katayama and E. Kuno, *The RAAGs on the complement graphs of path graphs in mapping class groups*, preprint, available at arXiv: 1804.03470v2.
- [13] S. Kim and T. Koberda, *An obstruction to embedding right-angled Artin groups in mapping class groups*, Int. Math. Res. Not. **2014** (2014), no. 14, 3912–3918.
- [14] T. Koberda, *Right-angled Artin groups and a generalized isomorphism problem for finitely generated subgroups of mapping class groups*, Geom. Funct. Anal. **22** (2012), 1541–1590.
- [15] L. Chen and A. Mukherjee, *From braid groups to mapping class groups*, preprint, available at arXiv: 2011.13020.
- [16] D. Margalit and R. Winarski, *The Birman–Hilden theory*, Celebratio Mathematica (2017) and Bulletin of the London Mathematical Society, **53** (2021), no. 3, 643–659.
- [17] L. Paris and D. Rolfsen, *Geometric subgroups of mapping class groups*, J. Reine Angew. Math. **521** (2000), 47–83.

- [18] H. Servatius, *Automorphisms of graph groups*, J. Algebra, **126** (1989), 34–60
- [19] K. Shackleton, *Combinatorial rigidity in curve complexes and mapping class groups*, Pac. J. Math. **230** (2007), no. 2, 217–232.

(TAKUYA KATAYAMA) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, GAKUSHUIN UNIVERSITY, 1-5-1 MEJIRO, TOSHIMA-KU, TOKYO 171-8588, JAPAN
Email address: `katayama@math.gakushuin.ac.jp`

(ERIKA KUNO) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, 1-1 MACHIKANEYAMA-CHO TOYONAKA, OSAKA 560-0043, JAPAN
Email address: `e-kuno@math.sci.osaka-u.ac.jp`