

THE MANIN–PEYRE CONJECTURE FOR SMOOTH SPHERICAL FANO THREEFOLDS

VALENTIN BLOMER, JÖRG BRÜDERN, ULRICH DERENTHAL, AND GIULIANO GAGLIARDI

ABSTRACT. The Manin–Peyre conjecture is established for smooth spherical Fano threefolds of semisimple rank one and type N . Together with the previously solved case T and the toric cases, this covers all types of smooth spherical Fano threefolds. The case N features a number of structural novelties; most notably, one may lose regularity of the ambient toric variety, the height conditions may contain fractional exponents, and it may be necessary to exclude a thin subset with exceptionally many rational points from the count, as otherwise Manin’s conjecture in its original form would turn out to be incorrect.

CONTENTS

1. Introduction	1
Part 1. Desingularization of the ambient toric variety	4
2. Heights and parametrization	4
3. Peyre’s constant	9
Part 2. Analysis of a diophantine equation	10
4. Elementary bounds	11
5. Character sums	12
6. Upper bound estimates	15
7. An asymptotic formula	22
8. Introducing the height conditions	27
Part 3. Proof of Theorem 1.1	28
9. Geometry	28
10. Cox rings and torsors	30
11. Counting problems	32
12. Application: Proof of Theorem 1.1	34
References	39

1. INTRODUCTION

Let G be a connected reductive group. A normal G -variety X is called spherical if a Borel subgroup of G has a dense orbit in X . Spherical varieties are a very large and interesting class of varieties that admit a combinatorial description by spherical systems and colored fans [Lu, BP, LV] generalizing the combinatorial description of toric varieties. Indeed, if the acting group G has semi-simple rank 0, then G is a torus.

2010 *Mathematics Subject Classification.* Primary 11D45; Secondary 14M27, 14G05, 11G35.

Key words and phrases. rational points, spherical varieties, Fano threefolds, Manin–Peyre conjecture, Cox rings, harmonic analysis.

The first author was supported by the DFG through Germany’s Excellence Strategy - EXC-2047/1 - 390685813. The third author was supported by DFG grant DE 1646/4-2.

If G has semi-simple rank 1, we may assume that $G = \mathrm{SL}_2 \times \mathbb{G}_m^r$ for some $r \geq 0$. Let $G/H = (\mathrm{SL}_2 \times \mathbb{G}_m^r)/H$ be the open orbit in X and define H' by $H' \times \mathbb{G}_m^r = H \cdot \mathbb{G}_m^r \subseteq \mathrm{SL}_2 \times \mathbb{G}_m^r$. Then the homogeneous space SL_2/H' is spherical, and there are three possible cases:

- either H' is a maximal torus (*the case T*),
- or H' is the normalizer of a maximal torus in SL_2 (*the case N*),
- or the homogeneous space SL_2/H' is horospherical, in which case X is isomorphic (as an abstract variety) to a toric variety.

In a recent paper [BBDG], the authors initiated a program to establish Manin's conjecture for smooth spherical Fano varieties, based on (a) the combinatorial description of spherical varieties, (b) the universal torsor method, and (c) techniques from analytic number theory including the Hardy-Littlewood circle method and multiple Dirichlet series. In particular, in case T , we confirmed Manin's conjecture for threefolds as well as for some higher-dimensional varieties. Here we relied on Hofscheier's classification [Ho] of smooth spherical Fano threefolds over $\overline{\mathbb{Q}}$, which identifies four such cases having natural split models over \mathbb{Q} .

In this paper we fully resolve the harder case N . This is achieved by a further development of the methods employed in [BBDG], and we proceed to describe the major new ingredients.

The first point concerns a general reformulation of Manin's conjecture as a counting problem. Let X be a smooth split projective variety over \mathbb{Q} with big and semiample anticanonical class ω_X^\vee whose Picard group is free of finite rank. Assume that its Cox ring $\mathcal{R}(X)$ is finitely generated with precisely one relation with integral coefficients. This defines a canonical embedding of X into a (not necessarily complete) ambient toric variety Y° ; it can be completed to a projective toric variety Y such that the natural map $\mathrm{Cl} Y \rightarrow \mathrm{Cl} X = \mathrm{Pic} X$ is an isomorphism and $-K_X$ is big and semiample on Y . Under the assumption that Y is regular, Sections 2 – 4 in [BBDG] (culminating in [BBDG, Propositions 3.7 & 4.11]) provide a general scheme to parametrize the rational points on X in terms of the universal torsor and to express the Manin-Peyre constant in terms of the Cox coordinates. The corresponding counting problem is in many cases amenable to techniques from analytic number theory. The assumption that Y is regular holds for smooth Fano threefolds of type T , but may fail in the case of type N . Part 1 of the present paper generalizes the passage to the universal torsor to varieties X for which Y is not necessarily regular. This result is independent of the theory of spherical varieties and should therefore have applications elsewhere.

The second point is of analytic nature. The universal torsor of spherical varieties of semi-simple rank 1 and type T has a defining equation of the form

$$x_{11}x_{12} - x_{21}x_{22} - \text{some monomial} = 0$$

which needs to be analyzed subject to rather complicated height conditions. The fact that we have a decoupled bilinear form in four variables is crucial for the method and allows in particular an auxiliary soft argument based on lattice considerations. For type N , the equation takes the form

$$x_{11}x_{12} - x_{21}^2 - \text{some monomial} = 0$$

which can analytically become very delicate. A considerable portion of this paper is devoted to the investigation of the particular equation

$$(1.1) \quad x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}x_{33}^2 = 0$$

in dyadic boxes

$$|x_{11}| \asymp |x_{12}| \asymp |x_{21}| \asymp X, \quad |x_{31}| \asymp |x_{32}| \asymp Y, \quad x_{33} \asymp X/Y$$

for some large quantity X and $1 \leq Y \leq X$, and we need an asymptotic formula with a power saving error term in $\min(Y, X/Y)$. It is conceivable that a modern variant of the circle method (like [HB]) can handle this, but this is not straightforward. There are considerable uniformity issues, since we need to deal simultaneously with the cases when Y is small, say $Y = \exp((\log \log X)^2)$ (in which case the equation looks roughly like a sum of two squares and a product), and Y is large, say $Y = X/\exp((\log \log X)^2)$ (in which case the equation looks roughly like a sum of a square and two

products). To keep track of uniformity, we will not use the circle method directly, but instead apply Poisson summation to selected variables depending on the ranges of parameters. This is ultimately more or less equivalent to the delta-symbol method of Duke–Friedlander–Iwaniec [DFI], but it is in this case a more convenient packaging.

The shape of equation (1.1) offers a new feature that was not present in [BBDG], and for which in fact very few examples are known. In the special case where $-x_{31}x_{32}$ is a square, the equation (1.1) describes a split quaternary quadratic form over \mathbb{Q} of signature $(2, 2)$, i.e., the sum of two hyperbolic planes. In this case it is well-known (see e.g. [HB]) that the asymptotic formula contains an additional logarithm. In particular, Manin’s conjecture in its original form turns out to be wrong, and instead we need to prove a “thin subset version” of Manin’s conjecture: we first remove a certain portion from the variety with exceptionally many points and then confirm the conjecture for the remaining set. More precisely, there should be a thin subset T of the set of rational points $X(\mathbb{Q})$ such that, for an anticanonical height H ,

$$N_{X(\mathbb{Q}) \setminus T, H}(B) := \#\{x \in X(\mathbb{Q}) \setminus T \mid H(x) \leq B\} = (1 + o(1))cB(\log B)^{\text{rk Pic } X - 1}$$

as in Manin’s conjecture with Peyre’s constant c . We refer the reader to the general discussion of this phenomenon in [LST] and to the (to our knowledge) only example of a smooth Fano variety [BHB] for which a thin version of Manin’s conjecture has been proved.

We now describe in more detail our results. As explained in [BBDG, §11], there are three smooth spherical Fano threefolds over N over $\overline{\mathbb{Q}}$ that are neither horospherical nor equivariant compactifications of \mathbb{G}_a^3 ; we construct split models X_1, X_2, X_3 over \mathbb{Q} as follows:

rk Pic	Hofscheier	Mori–Mukai	torsor equation	label
2	$N_1 8$	II.29	$x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}x_{33}^2$	variety X_1
3	$N_0 3$	III.22	$x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}$	variety X_2
3	$N_1 9$	III.19	$x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}$	variety X_3

TABLE 1. Smooth Fano threefolds of type N that are spherical, but not horospherical

More precisely, let X_1 the blow-up of the quadric $Q = \mathbb{V}(z_{11}z_{12} - z_{21}^2 - z_{31}z_{32}) \subset \mathbb{P}_{\mathbb{Q}}^4$ in the conic $C_{33} = \mathbb{V}(z_{31}, z_{32})$. This is a smooth Fano threefold of type II.29 in the Mori–Mukai classification. We have a fibration $X_1 \rightarrow \mathbb{P}^1$ that is defined in Cox coordinates by $(x_{11} : \cdots : x_{33}) \mapsto (x_{31} : x_{32})$. As explained above, we must remove the thin subset

$$T_1 = \{(x_{11} : \cdots : x_{33}) \in X_1(\mathbb{Q}) \mid x_{31}x_{32} = -\square \text{ or } x_{11}x_{12}x_{21}x_{31}x_{32}x_{33} = 0\}.$$

Let $W_2 = \mathbb{P}_{\mathbb{Q}}^1 \times \mathbb{P}_{\mathbb{Q}}^2$ with coordinates $(z_{01} : z_{02})$ and $(z_{11} : z_{12} : z_{21})$. Let C_{32} be the curve $\mathbb{V}(z_{02}, z_{11}z_{12} - z_{21}^2)$ on W_2 and let X_2 be the blow-up of W_2 in C_{32} . This is a smooth Fano threefold of type III.22. We will see later that the height conditions contain fractional exponents $\alpha_{ij, \nu}$. To our knowledge, no such case has been treated hitherto. Let $T_2 \subset X_2(\mathbb{Q})$ be the set of rational points where at least one Cox coordinate is zero.

Let X_3 be the blow-up of the quadric $Q = \mathbb{V}(z_{11}z_{12} - z_{21}^2 - z_{31}z_{32}) \subset \mathbb{P}_{\mathbb{Q}}^4$ in the points $P_{01} = \mathbb{V}(z_{11}, z_{12}, z_{21}, z_{31})$ and $P_{02} = \mathbb{V}(z_{11}, z_{12}, z_{21}, z_{32})$. Its type is III.19. As before, let $T_3 \subset X_3(\mathbb{Q})$ be the set of rational points where at least one Cox coordinate is zero.

In Sections 2.3 and 11, we will define natural anticanonical height functions $H_j : X_j(\mathbb{Q}) \rightarrow \mathbb{R}$, $j = 1, 2, 3$, using the anticanonical monomials in their Cox rings. We write $N_j(B) = N_{X_j(\mathbb{Q}) \setminus T_j, H_j}(B)$.

Theorem 1.1. *The Manin–Peyre conjecture holds for the smooth spherical Fano threefolds X_2, X_3 of semisimple rank one and type N , and a thin version of the Manin–Peyre conjecture holds for X_1 . More precisely, there exist explicit constants C_1, C_2, C_3 such that*

$$N_j(B) = (1 + o(1))C_j B(\log B)^{\text{rk Pic } X_j - 1}$$

for $1 \leq j \leq 3$. The values of C_j are the ones predicted by Peyre.

Together with previous results, this covers all types of smooth spherical Fano threefolds.

Part 1. Desingularization of the ambient toric variety

If a variety X has a finitely generated Cox ring with one relation, this gives a description of X as a hypersurface in a toric variety Y . The description of height and Tamagawa measures in [BBDG, Part 1] relies on the assumption [BBDG, (2.3)] that Y can be chosen to be regular, which does not hold in our examples X_2, X_3 . Here, we describe how to circumvent this problem.

2. HEIGHTS AND PARAMETRIZATION

We start with a situation similar to (but in several respects more general than) [BBDG, Section 2]. Let Y° be a smooth split toric variety over \mathbb{Q} . We do not assume that Y° is complete, but we still assume the weaker property that Y° has only constant regular functions. Let

$$(2.1) \quad \mathcal{R}(Y^\circ) \cong \mathbb{Q}[x_1, \dots, x_J]$$

be its Cox ring, where x_1, \dots, x_J correspond to the torus invariant prime divisors in Y° . Let $0 \neq \Phi \in \mathcal{R}(Y^\circ)$ be a homogeneous equation, and let $X \subseteq Y^\circ$ be the corresponding subvariety.

We assume that X is smooth and projective, with big and semiample anticanonical class $-K_X$. Moreover, we assume that every torus orbit in Y° meets X . We also assume that the pullback map $\text{Pic } Y^\circ \rightarrow \text{Pic } X$ sends a big and semiample divisor class L° to $-lK_X$ for some $l \in \mathbb{Z}_{\geq 0}$. Finally, we assume $\Phi \in \mathbb{Z}[x_1, \dots, x_J]$.

Remark 2.1. A situation as above can be naturally obtained starting from a smooth split projective variety over \mathbb{Q} with big and semiample anticanonical class $-K_X$ and finitely generated Cox ring

$$\mathcal{R}(X) \cong \mathbb{Q}[x_1, \dots, x_J]/(\Phi),$$

where x_1, \dots, x_J is a system of pairwise nonassociated Pic X -prime generators and $\Phi \neq 0$. By [ADHL, 3.2.5], there exists a canonical embedding into an ambient toric variety Y° , and it satisfies all the above assumptions. Moreover, the pullback map $\text{Pic } Y^\circ \rightarrow \text{Pic } X$ is an isomorphism, and we may take $L^\circ = -K_X$ under this identification.

If Σ is the fan of any toric variety, we write Σ_{\max} for the set of maximal cones and $\Sigma(1)$ for the set of rays.

Let Σ° be the fan of Y° . The generators $x_1, \dots, x_J \in \mathcal{R}(Y^\circ)$ are in bijection to the rays $\rho \in \Sigma^\circ(1)$; we also write x_ρ for x_i corresponding to ρ .

Let Y be a completion of Y° such that the pullback map $\text{Cl } Y \rightarrow \text{Cl } Y^\circ = \text{Pic } Y^\circ$ is an isomorphism and L° is big and semiample on Y ; let Σ be the fan of this toric variety Y . We have $\Sigma(1) = \Sigma^\circ(1)$ and $\mathcal{R}(Y) = \mathcal{R}(Y^\circ)$. For example, we may choose the unique Y such that L° is ample on Y , which exists by [ADHL, Proposition 2.4.2.6]; we call this Y the *standard small completion* of Y° .

We do not assume that Y is regular (in contrast to [BBDG, Section 2]). Let

$$\rho : Y'' \rightarrow Y$$

be a toric desingularization of Y that does not change the smooth locus of Y . Such a desingularization can be obtained by suitably subdividing the singular cones in the fan Σ of Y into a smooth fan Σ'' .

Let $Y' \subset Y''$ be a toric subvariety with $Y^\circ \subset Y'$. This means that the fan Σ' of Y' is a subfan of Σ'' and contains Σ° . If we write $\rho_{J+1}, \dots, \rho_{J'}$ for the rays in $\Sigma'(1) \setminus \Sigma(1)$, we can write

$$\mathcal{R}(Y') = \mathbb{Q}[y_1, \dots, y_J, y_{J+1}, \dots, y_{J'}]$$

for the Cox ring of Y' , again with the correspondence between rays ρ_i and generators y_i .

Since X is smooth, we can identify it with its strict transform under $\rho : Y'' \rightarrow Y$; it is a hypersurface in Y' defined by a homogeneous equation Φ' that is obtained from Φ by homogenizing (therefore, $y_i \mapsto x_i$ for $i \leq J$ and $y_i \mapsto 1$ for $i > J$ turns Φ' into Φ). We obtain the following commutative diagram:

$$\begin{array}{ccccc} X & \hookrightarrow & Y' & \hookrightarrow & Y'' \\ \parallel & & \uparrow & & \downarrow \\ X & \hookrightarrow & Y^\circ & \hookrightarrow & Y. \end{array}$$

Below, we will regard X mostly as embedded into Y' .

For simplicity, we assume that every cone in Σ' is the face of a maximal cone.

Let U be the open torus in Y' . For each $\rho \in \Sigma'(1)$, we have a U -invariant Weil divisor D_ρ defined by y_ρ of class $[D_\rho] = \deg(y_\rho) \in \text{Pic } Y'$. Let $D_0 := \sum_{\rho \in \Sigma'(1)} D_\rho$, which is an effective divisor of class $[D_0] = -K_{Y'}$.

Let $S \subset \Sigma'(1)$ be such that $\{\deg(y_\rho) \mid \rho \notin S\}$ is a basis of $\text{Pic } Y'$. This is true if and only if S is a basis of N . Therefore, we can write $\deg(y_{\rho'}) = \sum_{\rho \notin S} a_{\rho', \rho}^S \deg(y_\rho)$ for each $\rho' \in \Sigma'(1)$ with certain $a_{\rho', \rho}^S \in \mathbb{Z}$. We define the rational section

$$z_{\rho'}^S := y_{\rho'} / \prod_{\rho \notin S} y_\rho^{a_{\rho', \rho}^S}$$

of degree 0 $\in \text{Pic } Y'$, with $z_\rho^S = 1$ for $\rho \notin S$. The z_ρ^S for $\rho \in S$ define a chart

$$f^S : U^S \rightarrow \mathbb{A}_{\mathbb{Q}}^S, \quad P \mapsto (z_\rho^S(P))_{\rho \in S}$$

where U^S is the open subset of Y' where $y_\rho \neq 0$ for all $\rho \notin S$ (i.e., the complement of $\bigcup_{\rho \notin S} D_\rho$ in Y'). Note that f^S is an isomorphism if and only if $S = \sigma(1)$ for some $\sigma \in \Sigma'_{\max}$. For the open subset $X^S := X \cap U^S$ of X , the image $f^S(X^S) \subset \mathbb{A}_{\mathbb{Q}}^S$ is defined by

$$\Phi^S := \Phi'(z_\rho^S) = \Phi'(y_\rho) / \prod_{\rho \notin S} y_\rho^{b_\rho^S}$$

(with $b_\rho^S \in \mathbb{Z}$ satisfy $\deg \Phi' = \sum_{\rho \notin S} b_\rho^S \deg(y_\rho)$) since $y_\rho \neq 0$ on U^S for $\rho \notin S$.

2.1. Universal torsors and models. Let $\pi' : Y'_0 \rightarrow Y'$ be the universal torsor as in [Sal]. Then the restriction $\pi' : X_0 \rightarrow X$ to the preimage of $X \subset Y'$ is a torsor, but not necessarily universal since the acting torus (dual to $\text{Pic } Y'$) may be too large. The toric varieties $\mathbb{A}_{\mathbb{Q}}^{J'} = \mathbb{A}_{\mathbb{Q}}^{\Sigma'(1)} =: Y'_1 \supset Y'_0 \rightarrow Y'$ have the fans $\Sigma'_1 \supset \Sigma'_0 \rightarrow \Sigma'$. Here, the sets of rays $\Sigma'_1(1) = \Sigma'_0(1)$ are in natural bijection to $\Sigma'(1)$. The r' irreducible components of $Z'_Y = Y'_1 \setminus Y'_0$ are defined by the vanishing of x_ρ for all $\rho \in S'_j$, where the *primitive collections*

$$(2.2) \quad S'_1, \dots, S'_{r'} \subseteq \Sigma'(1)$$

are all sets with the following property: $S'_j \not\subseteq \sigma(1)$ for all $\sigma \in \Sigma'$, but for every proper subset S''_j of S'_j , there is a $\sigma \in \Sigma'$ with $S''_j \subseteq \sigma(1)$.

From the fans and their maps, we may construct \mathbb{Z} -models $\tilde{\pi}' : \tilde{Y}'_1 \setminus \tilde{Z}'_Y = \tilde{Y}'_0 \rightarrow \tilde{Y}'$, again as in [Sal]. By our assumption that Φ has integral coefficients, we obtain \mathbb{Z} -models $\tilde{\pi}' : \tilde{X}_1 \setminus \tilde{Z}_X = \tilde{X}_0 \rightarrow \tilde{X}$ of $\pi' : X_1 \setminus Z_X = X_0 \rightarrow X$ by restricting everything to $\Phi' = 0$.

We assume:

$$(2.3) \quad \text{The toric variety } Y' \text{ is chosen such that } \tilde{X} \text{ is proper over } \text{Spec } \mathbb{Z}.$$

This is always possible since for $Y' = Y''$ the scheme \tilde{Y}' is projective over $\text{Spec } \mathbb{Z}$.

Proposition 2.2. *We have*

$$\begin{aligned} \tilde{X}_0(\mathbb{Z}) &= \{\mathbf{x} = (x_\rho)_{\rho \in \Sigma'(1)} \in \mathbb{Z}^{\Sigma'(1)} : \Phi(\mathbf{x}) = 0, \gcd\{x_\rho : \rho \in S'_j\} = 1 \text{ for all } j = 1, \dots, r'\}, \\ \tilde{X}_0(\mathbb{Z}_p) &= \{\mathbf{x} = (x_\rho)_{\rho \in \Sigma'(1)} \in \mathbb{Z}_p^{\Sigma'(1)} : \Phi(\mathbf{x}) = 0, p \nmid \gcd\{x_\rho : \rho \in S'_j\} \text{ for all } j = 1, \dots, r'\}. \end{aligned}$$

The map $\tilde{\pi}'$ induces a $2^{\text{rk Pic } Y'}$: 1-map $\tilde{X}_0(\mathbb{Z}) \rightarrow \tilde{X}(\mathbb{Z}) = X(\mathbb{Q})$.

Proof. The proof is as in [BBDG, Proposition 2.2] by (2.3). □

2.2. Metrization of ω_X^{-1} via Poincaré residues. Let L be any big and semiample divisor class on Y' such that $L|_X = -lK_X$. Then there exists a uniquely determined divisor E with support $\text{supp } E$ in the exceptional locus of ρ such that

$$L = -lK_{Y'} - l[X] + [E].$$

The following lemma shows that (after possibly enlarging l) such an L can always be found.

Lemma 2.3. *After suitably enlarging l , there exists a big and semiample divisor class on Y' such that $L|_X = -lK_X$.*

Proof. Since L° is ample, it is \mathbb{Q} -Cartier on Y . Hence, after replacing l by a positive multiple, we may assume that L° is Cartier on Y . Let L'' be the pullback $\rho^*(L^\circ)$. Then L'' is big and semiample on Y'' , and moreover $L''|_X = -lK_X$. The same is then true for $L = L''|_{Y'}$. \square

As before, let $S \subset \Sigma'(1)$ be such that $\{\deg(y_\rho) \mid \rho \notin S\}$ is a basis of $\text{Pic } Y'$. Hence there is a unique (not necessarily effective) Weil divisor $D(S) = \sum_{\rho \notin S} a_\rho^S D_\rho$ of class $-K_{Y'} - [X]$ whose support is contained in $\bigcup_{\rho \notin S} D_\rho$. Let $y^{D(S)}$ be the corresponding rational monomial in $\mathcal{R}(Y')$ of degree $-K_{Y'} - [X]$. The characters defined by z_ρ^S for $\rho \in S$ form a basis of $M = \text{Hom}(U, \mathbb{G}_m)$.

By [CLS, Proposition 8.2.3], we have a global nowhere vanishing section $s_{Y'}$ of $\omega_{Y'}(D_0)$ (defined up to sign, independent of S ; we have $s_{Y'} = \Omega_0 / \prod_{\rho \in \Sigma'(1)} y_\rho$ for Ω_0 from [CLS, (8.2.3)]) whose restriction to the open subset U^S is $\pm \bigwedge_{\rho \in S} \frac{dz_\rho^S}{z_\rho^S}$.

With $y^{D_0} = \prod_{\rho \in \Sigma'(1)} y_\rho$, for each S as above,

$$\varpi^S := \frac{y^{D_0}}{y^{D(S)} \Phi} s_{Y'} \in \Gamma(Y', \omega_{Y'}(D(S) + X))$$

defines a nowhere vanishing global section. On U^S , we have

$$\varpi^S = \frac{\pm 1}{\Phi^S} \bigwedge_{\rho \in S} dz_\rho^S \in \Gamma(U^S, \omega_{Y'}(X)).$$

Let \mathcal{P}^l be a set of polynomials $F \in \mathbb{Q}[y_1, \dots, y_{J'}]$ of degree L . For each polynomial $F \in \mathcal{P}^l$ of degree L , let $D(F)$ be the effective divisor on Y' of class $\deg F = L$ defined by F (in Cox coordinates). If $X \subset \text{supp } D(F)$, then we remove F from our set \mathcal{P}^l ; clearly, this does not change the results that we want to prove. For $X \not\subset \text{supp } D(F)$, we define

$$\varpi_F := \frac{y^{lD_0+E}}{F \cdot \Phi^l} s_{Y'}^l \in \Gamma(Y', \omega_{Y'}^l(D(F) + lX - E)).$$

We have the Poincaré residue map $\text{Res} : \omega_{Y'}(X) \rightarrow \iota'_* \omega_X$ of $\mathcal{O}_{Y'}$ -modules (where $\iota' : X \rightarrow Y'$ is the inclusion). On the smooth open subset U^S of Y' , it maps $\varpi^S \in \Gamma(U^S, \omega_{Y'}(X))$ to $\text{Res}(\varpi^S) \in \Gamma(U^S, \iota'_* \omega_X) = \Gamma(X^S, \omega_X)$, which is

$$(2.4) \quad \text{Res}(\varpi^S) = \frac{\pm 1}{\partial \Phi^S / \partial z_{\rho_0}^S} \bigwedge_{\rho \in S \setminus \{\rho_0\}} dz_\rho^S$$

on the open subset of X^S where $\partial \Phi_\rho / \partial z_{\rho_0}^S \neq 0$, for any $\rho_0 \in S$. Furthermore, $\text{Res}^l : \omega_{Y'}^l(lX) \rightarrow \iota'_* \omega_X^l$ sends $\varpi_F \in \Gamma(U_F, \omega_{Y'}^l(lX - E))$ to $\text{Res}^l(\varpi_F) \in \Gamma(U_F, \iota'_* \omega_X^l) = \Gamma(X_F, \omega_X^l)$, where U_F is the complement of $D(F)$ in Y' , and $X_F = X \cap U_F$.

Lemma 2.4. *The section $\text{Res}(\varpi^S)$, $\text{Res}^l(\varpi_F)$ extends uniquely to nowhere vanishing global section of $\omega_X(D(S) \cap X)$, $\omega_X^l(D(F) \cap X)$, respectively.*

Proof. For $\text{Res}(\varpi^S)$, this is as in [BBDG, Lemma 2.3], i.e., similar to [BBS, Lemma 13]. The computation for $\text{Res}^l(\varpi_F)$ is analogous, using $X \not\subset \text{supp } D(F)$ and $E \cap X = \emptyset$. \square

Therefore, $\tau^S := \text{Res}(\varpi^S)^{-1}$, $\tau_F := \text{Res}^l(\varpi_F)^{-1}$ are global nowhere vanishing sections of $\omega_X^{-1}(-D(S) \cap X)$, $\omega_X^{-l}(-D(F) \cap X)$, which we can also view as a global section of ω_X^{-1} , ω_X^{-l} , respectively.

Lemma 2.5. *The sections $\tau^S \in \Gamma(X, \omega_X^{-1})$, $\tau_F \in \Gamma(X, \omega_X^{-l})$ do not vanish anywhere on X^S , X_F , respectively.*

Proof. The support of $D(S) \cap X$ is contained in $X \cap \bigcup_{\rho \notin S} D_\rho$, which is the complement of X^S . Moreover, $D_F \cap X$ is the complement of X_F . \square

From now on, we assume:

(2.5) \mathcal{P}^l only contains monic monomials (of degree L), and for each $\sigma \in \Sigma'_{\max}$ there exists $m \in \mathcal{P}^l$ such that $\text{supp div } m$ does not meet U_σ .

Then the set \mathcal{P}^l is in particular basepoint-free.

We define a v -adic norm/metric on ω_X^{-1} by

$$\|\tau(P)\|_v := \min_{F \in \mathcal{P}^l: P \notin D(F)} \left| \frac{\tau^l}{\tau_F}(P) \right|_v^{1/l}$$

for a local section τ of ω_X^{-1} not vanishing in $P \in X(\mathbb{Q}_v)$.

Lemma 2.6. *Let p be a prime such that \tilde{X} is smooth over \mathbb{Z}_p . On ω_X^{-l} , the p -adic norm $\|\cdot\|_p$ defined by*

$$\|\tau(P)\|_v := \min_{F \in \mathcal{P}^l: P \notin D(F)} \left| \frac{\tau}{\tau_F}(P) \right|_v$$

for a local section τ of ω_X^{-l} not vanishing in $P \in X(\mathbb{Q}_v)$ coincides with the model norm $\|\cdot\|_p^*$ determined by \tilde{X} over \mathbb{Z}_p as in [Sal, Definition 2.9].

Proof. See [BBDG, Lemma 3.3]. For such a τ not vanishing in P , let $Q \in \mathcal{P}^l$ be such that $|(\tau^Q/\tau)(P)|_p = \max_{F \in \mathcal{P}^l} |(\tau^F/\tau)(P)|_p$, which is positive by Lemma 2.5 and the fact that the set \mathcal{P}^l is basepoint-free. Hence τ^Q does not vanish in P , and

$$\|\tau^Q(P)\|_p^{-1} = \max_{F \in \mathcal{P}^l} \left| \frac{\tau^F}{\tau^Q}(P) \right|_p = \max_{F \in \mathcal{P}^l} \frac{|(\tau^F/\tau)(P)|_p}{|(\tau^Q/\tau)(P)|_p} = 1.$$

For each $F \in \mathcal{P}^l$, the section τ^F extends to a global section $\tilde{\tau}^F$ of $\omega_{\tilde{X}/\mathbb{Z}_p}^{-l}$, and $\omega_{\tilde{X}/\mathbb{Z}_p}^{-l}$ is generated by the set of all these $\tilde{\tau}^F$ as an $\mathcal{O}_{\tilde{X}}$ -module. The reason is that everything required for the definition of τ^F above can also be defined over \mathbb{Z}_p . For the existence of the Poincaré residue map in this case, we refer to [KK, Definition 4.1].

For every $F \in \mathcal{P}^l$, we have $\left| \frac{\tau^F}{\tau^Q}(P) \right|_p \leq 1$ as in the computation above. This implies $\tau^F(P) = a_F \tau^Q(P)$ for some $a_F \in \mathbb{Z}_p$ in the \mathbb{Q}_p -module $\omega_X^{-1}(P)$, and hence also $\tilde{\tau}^F(P) = a_F \tilde{\tau}^Q(P)$ in the \mathbb{Z}_p -module $P^*(\omega_{\tilde{X}/\mathbb{Z}_p}^{-1})$, which shows that $P^*(\omega_{\tilde{X}/\mathbb{Z}_p}^{-1})$ is generated by $\tilde{\tau}^Q(P)$. Hence $\|\tau^Q(P)\|_p^* = 1$ by definition of the model norm. We conclude

$$\|\tau(P)\|_p = |(\tau/\tau^Q)(P)|_p \cdot \|\tau^Q(P)\|_p = |(\tau/\tau^Q)(P)|_p \cdot \|\tau^Q(P)\|_p^* = \|\tau(P)\|_p^*. \quad \square$$

2.3. Height functions. For $P \in X(\mathbb{Q})$, we define

$$(2.6) \quad H(P) := \prod_v \|\tau(P)\|_v^{-1}$$

for a local section τ of ω_X^{-1} not vanishing in P .

Remark 2.7. Let F, F_0 be homogeneous elements of the same degree in the Cox ring of Y' . If F_0 does not vanish in P , then F/F_0 can be regarded as a rational function on X that can be evaluated in $P \in X(\mathbb{Q})$.

Lemma 2.8. *We have*

$$H(P) = \left(\prod_v \max_{F \in \mathcal{P}^l} \left| \frac{F}{F_0}(P) \right|_v \right)^{1/l}$$

for any polynomial F_0 of degree L not vanishing in P .

Proof. We have $P \in X^S(\mathbb{Q})$ for some S as above. We can compute $H(P)$ with $\tau := \tau^S$ by Lemma 2.5. Applying the $\mathcal{O}_{Y'}$ -module homomorphism Res to $\varpi_F = F^{-1}y^{lD(S)+E}(\varpi^S)^l$ shows $\tau_F = Fy^{-lD(S)-E}(\tau^S)^l$, hence

$$(2.7) \quad \|\tau^S(P)\|_v^{-1} = \max_{F \in \mathcal{P}^l} \left| \frac{\tau_F}{(\tau^S)^l}(P) \right|_v^{1/l} = \max_{F \in \mathcal{P}^l} \left| \frac{F}{y^{lD(S)+E}}(P) \right|_v^{1/l},$$

which is our claim in the case $F_0 := y^{lD(S)+E}$. The general case follows using the product formula. \square

We lift the height function H to X_0 by composing it with $\pi: X_0(\mathbb{Q}) \rightarrow X(\mathbb{Q})$, giving

$$H_0: X_0(\mathbb{Q}) \rightarrow \mathbb{R}_{>0}.$$

Lemma 2.9. *For $P_0 \in X_0(\mathbb{Q})$, we have*

$$H_0(P_0) = \left(\prod_v \max_{F \in \mathcal{P}^l} |F(P_0)|_v \right)^{1/l}.$$

Proof. Let $P = \pi(P_0) \in X(\mathbb{Q})$. As in the proof of [BBDG, Lemma 3.5], for F_0 of degree L not vanishing in P and $F \in \mathcal{P}^l$, we have $(F/F_0)(P) = F(P_0)/F_0(P_0)$ if we compute $(F/F_0)(P)$ as in Remark 2.7 and also regard F, F_0 as regular functions on $X_0(\mathbb{Q})$ that can be evaluated in P_0 . We apply this to Lemma 2.8 to obtain

$$H_0(P_0) = H(P) = \left(\prod_v \max_{F \in \mathcal{P}^l} \left| \frac{F}{F_0}(P) \right|_v \right)^{1/l} = \left(\prod_v \max_{F \in \mathcal{P}^l} \left| \frac{F(P_0)}{F_0(P_0)} \right| \right)^{1/l}.$$

Then we use the product formula. \square

In its integral model, this simplifies as follows.

Corollary 2.10. *For $P_0 \in \tilde{X}_0(\mathbb{Z})$, we have*

$$H_0(P_0) = \max_{F \in \mathcal{P}^l} |F(P_0)|_\infty^{1/l}.$$

Proof. This is analogous to [Sal, Proposition 11.3] and [BBDG, Corollary 3.6]. For a prime p , we have $P_0 \pmod{p}$ in $\tilde{X}_0(\mathbb{F}_p)$. There is a $\sigma \in \Sigma'_{\max}$ such that $y_\rho(P_0 \pmod{p}) \neq 0 \in \mathbb{F}_p$ for all $\rho \notin \sigma(1)$ since \tilde{X}_0 is defined by the irrelevant ideal in \tilde{X}_1 . Choose $Q \in \mathcal{P}^l$ such that $\text{supp div } Q$ does not meet U_σ . Then we have $Q(P_0 \pmod{p}) \neq 0 \in \mathbb{F}_p$, and hence $|Q(P_0)|_p = 1$. Therefore, we have $\max_{F \in \mathcal{P}^l} |F(P_0)|_p^{1/l} = 1$ and only the archimedean factor in Lemma 2.9 remains. \square

2.4. Counting problem. The following result parametrizes the set $N_{X(\mathbb{Q}) \setminus T, H}(B)$ in terms of integral points on the universal torsor of the ambient toric variety Y' (which is given by its Cox ring (2.1) and the primitive collections (2.2)), the equation Φ' , and the monomials in \mathcal{P}^l . The resulting counting problem is amenable to methods of analytic number theory.

Proposition 2.11. *Let X be a variety as in Section 2, let Y' be a toric variety satisfying (2.3), let L be a divisor class as in Section 2.2, and let \mathcal{P}^l be a set of monomials satisfying (2.5).*

Let T be an arbitrary subset of $X(\mathbb{Q})$. Then

$$N_{X(\mathbb{Q}) \setminus T, H}(B) = \frac{1}{2^{\text{rk Pic } Y'}} \# \left\{ \mathbf{y} \in \mathbb{Z}^{\Sigma'(1)} : \begin{array}{l} \Phi'(\mathbf{y}) = 0, \max_{F \in \mathcal{P}^l} |F(\mathbf{y})|_\infty^{1/l} \leq B, \pi'(\mathbf{y}) \notin T, \\ \text{gcd}\{y_\rho : \rho \in S'_j\} = 1 \text{ for every } j = 1, \dots, r' \end{array} \right\},$$

using the notation (2.1) and (2.2).

Proof. This follows from Proposition 2.2 and Corollary 2.10. \square

3. PEYRE’S CONSTANT

We keep the notation and assumptions of Section 2. In addition, we assume from now on that we are in the situation of Remark 2.1. In particular, the pullback map $\text{Pic } Y^\circ \rightarrow \text{Pic } X$ is an isomorphism and X is split.

3.1. Tamagawa measures. By [Pey, (2.2.1)] and [Sal, Theorem 1.10], the v -adic norm $\|\cdot\|_v$ on ω_X^{-1} defined above induces a Tamagawa measure μ_v on $X(\mathbb{Q}_v)$.

Lemma 3.1. *Let $S = \sigma(1)$ for some $\sigma \in \Sigma'_{\max}$. For a Borel subset N_v of $X^S(\mathbb{Q}_v)$, we have*

$$(3.1) \quad \mu_v(N_v) = \int_{N_v} \frac{|\text{Res}(\varpi^S)|_v}{\max_{F \in \mathcal{P}^l} |\tau_F \text{Res}(\varpi^S)|_v^{1/l}} = \int_{N_v} \frac{|\text{Res}(\varpi^S)|_v}{\max_{F \in \mathcal{P}^l} |F/y^{lD(S)+E}|_v^{1/l}},$$

where $|\text{Res}(\varpi^S)|_v$ is the v -adic density on $X^S(\mathbb{Q}_v)$ of the volume form $\text{Res}(\varpi^S)$ on X^S .

If N_v is contained in a sufficiently small neighborhood of P in $X^S(\mathbb{Q}_v)$ with $\partial\Phi^S/\partial z_{\rho_0}^S(P) \neq 0$, then

$$(3.2) \quad \mu_v(N_v) = \int_{\pi_{\rho_0}^S(N_v)} \frac{\bigwedge_{\rho \in S \setminus \{\rho_0\}} dz_\rho^S}{|\partial\Phi^S/\partial z_{\rho_0}^S(\mathbf{z}^S)|_v \max_{F \in \mathcal{P}^l} |F(\mathbf{z}^S)|_v^{1/l}}$$

in the affine coordinates $\mathbf{z}^S = (z_\rho^S)_{\rho \in S}$, where $\pi_{\rho_0}^S : U^S(\mathbb{Q}_v) = \mathbb{Q}_v^S \rightarrow \mathbb{Q}_v^{S \setminus \{\rho_0\}}$ is the natural projection and $z_{\rho_0}^S$ is expressed in terms of the other coordinates using the implicit function for Φ^S .

Proof. This is analogous to the proof of [BBDG, Proposition 4.1]. However, we work with $\|\tau_{F_0}(P)\|_v$ for $F_0 = x^{lD(S)+E}$ and use $F_0(\mathbf{z}^S) = 1$ in our affine coordinates on $X^S(\mathbb{Q}_v)$. At the end, comparing the definitions of ϖ^S and ϖ_F shows $\varpi_F/(\varpi^S)^l = y^{lD(S)+E}/F$, hence $\tau_F \text{Res}(\varpi^S)^l = \tau_F/(\tau^S)^l = F/y^{lD(S)+E}$, and hence the integrals in (3.1) are equal. \square

Remark 3.2. Since we have assumed that every cone in Σ' is the face of a maximal cone, the open subvarieties X^S for $S = \sigma(1)$ with $\sigma \in \Sigma'_{\max}$ cover X .

Remark 3.3. If we are in the special case where X is covered by open subvarieties X^S with $S \subset \Sigma(1)$ for $S = \sigma(1)$ with $\sigma \in \Sigma'_{\max}$, then (3.2) gives the same formula for the v -adic density as we would have obtained by working with $\mathcal{R}(X)$ directly (since the additional coordinates z_ρ^S are 1 for all $\rho \in \Sigma'(1) \setminus \Sigma(1)$; up to the description of the height function defined via the monomials $F \in \mathcal{P}^l \subset \mathbb{Q}[y_1, \dots, y_{J'}]$, in which the additional variables $y_{J+1}, \dots, y_{J'}$ corresponding to $\rho \in \Sigma'(1) \setminus \Sigma(1)$ are set to $z_\rho^S = 1$). For example, this is the case if X is covered by the open subvarieties X^S for $S \in \Sigma'_{\max}$.

3.2. Measures on the torsor. Analogously to [BBDG, (4.4)], we obtain a v -adic measure m_v on $X_0(\mathbb{Q}_v)$, which is explicitly (for sufficiently small subsets M_v)

$$m_v(M_v) = \int_{\pi_{\rho_0}(M_v)} \frac{\bigwedge_{\rho \in \Sigma'_0(1) \setminus \{\rho_0\}} dy_\rho}{|\partial\Phi'/\partial x_{\rho_0}(\mathbf{y})|_v \max_{F \in \mathcal{P}^l} |F(\mathbf{y})|_v^{1/l}}$$

in the coordinates $\mathbf{y} = (y_\rho)_{\rho \in \Sigma'_0(1)}$, where π_{ρ_0} is the projection to all coordinates y_ρ with $\rho \neq \rho_0$ and where y_{ρ_0} is expressed in terms of these coordinates using the implicit function theorem.

Lemma 3.4. *For any prime p , we have $m_p(\tilde{X}_0(\mathbb{Z}_p)) = (1 - p^{-1})^{\text{rk Pic } Y'} \mu_p(X(\mathbb{Q}_p))$.*

Proof. This is analogous to [BBDG, Lemma 4.2, Lemma 4.3]. \square

3.3. Comparison to the number of points modulo p^ℓ . As in [BBDG, §4.4], for any prime p and $\ell \in \mathbb{Z}_{>0}$, we have

$$\tilde{X}_0(\mathbb{Z}/p^\ell\mathbb{Z}) = \{\mathbf{x} \in (\mathbb{Z}/p^\ell\mathbb{Z})^{\Sigma'(1)} : \Phi'(\mathbf{x}) = 0 \in \mathbb{Z}/p^\ell\mathbb{Z}, p \nmid \gcd\{x_\rho : \rho \in S'_j\} \text{ for all } j = 1, \dots, r'\}.$$

In particular, the additional variables x_ρ indexed by $\rho \in \Sigma'(1) \setminus \Sigma(1)$ (obtained via the desingularization of the ambient toric variety) appear here.

Proposition 3.5. *For every prime p , there is an $\ell_0 \in \mathbb{Z}_{>0}$ such that for all $\ell \geq \ell_0$ we have*

$$m_p(\tilde{X}_0(\mathbb{Z}_p)) = \frac{\#\tilde{X}_0(\mathbb{Z}/p^\ell\mathbb{Z})}{(p^\ell)^{\Sigma'(1)-1}}.$$

Furthermore,

$$(1 - p^{-1})^{\text{rk Pic } X} \mu_p(X(\mathbb{Q}_p)) = (1 - p^{-1})^{\#\Sigma(1) - \#\Sigma'(1)} \lim_{\ell \rightarrow \infty} \frac{\#\tilde{X}_0(\mathbb{Z}/p^\ell\mathbb{Z})}{(p^\ell)^{\dim X_0}}.$$

Proof. This relies on the regularity of Y' and is otherwise analogous to [BBDG, Lemma 4.4, Proposition 4.5, Proposition 4.6]. \square

3.4. The real density. In this section, we compute the real density and Peyre's α -constant as in [BBDG, §4.5].

We choose $\sigma \in \Sigma'_{\max}$, $\rho_0 \in \sigma(1)$, and $\rho_1 \in \Sigma(1) \setminus \sigma(1)$ subject to the following conditions analogous to [BBDG, (4.7)]:

- $\sigma \in \Sigma'_{\max}$ also appears in Σ_{\max} .
- Every variable x_ρ for $\rho \in \Sigma(1)$ appears in at most one monomial of Φ .
- (3.3) Writing $-K_X = \sum_{\rho \in \Sigma(1) \setminus \sigma(1)} \alpha_\rho^\sigma \deg(x_\rho)$ in $\text{Pic } X$, we have $\alpha_{\rho_1}^\sigma \neq 0$.
- The variable x_{ρ_0} appears with exponent 1 in Φ .
- No x_ρ with $\rho \in \sigma(1) \cup \{\rho_1\} \setminus \{\rho_0\}$ appears in the same monomial of Φ as x_{ρ_0} .

We define the numbers $b_{\rho, \rho'}$ and $b_{\rho'}$ as in [BBDG, §4.5], computed in $\text{Pic } X$, for $\rho' \in \sigma(1)' = \sigma(1) \cup \{\rho_1\}$ and $\rho \in \Sigma(1) \setminus \sigma(1)'$. Then we define c^*, c_∞ analogously to [BBDG, (4.9), (4.11)]; here, H_∞ is defined using only the monomials in \mathcal{P}^l , and the additional coordinates indexed by $\Sigma'(1) \setminus \Sigma(1)$ do not appear in c^* and are set to 1 in c_∞ .

Proposition 3.6. *In the notation above and assuming (3.3), we have*

$$\alpha(X) \mu_\infty(X(\mathbb{R})) = \frac{1}{2^{\text{rk Pic } X}} c^* c_\infty.$$

Proof. We note that [BBDG, Lemma 4.7, Lemma 4.8] are valid without assumption [BBDG, (2.3)], with c^* computed with respect to $\text{Pic } X$, not $\text{Pic } Y'$.

We observe that the additional variables z_ρ^S for $\rho \in \Sigma'(1) \setminus \Sigma(1)$ are 1 in the expression (3.2) for $\mu_\infty(X(\mathbb{R}))$ for $S = \sigma(1)$. Therefore, the expected real density $\omega_\infty = \mu_\infty(X(\mathbb{R}))$ has the same description as in [BBDG, (4.10)] (except for the choice of monomials in the height function, which does not cause problems), and also c_∞ in [BBDG, (4.11)] is essentially unchanged (both expressed without variables corresponding to $\rho \in \Sigma'(1) \setminus \Sigma(1)$, and computing in $\text{Pic } X$, not $\text{Pic } Y'$). Therefore, the statement and proof of [BBDG, Proposition 4.10] stay the same. \square

Part 2. Analysis of a diophantine equation

The counting problem in Corollary 11.1(a) corresponding to the variety X_1 is rather delicate and not covered by the general method of [BBDG]. This part of the paper is devoted to a detailed investigation.

4. ELEMENTARY BOUNDS

For $\xi \in \mathbb{Z} \setminus \{0\}$ we consider the equation

$$(4.1) \quad x_{11}x_{12} + x_{21}^2 + \xi^2 x_{31}x_{32}x_{33}^2 = 0, \quad x_{31}x_{32} \neq -\square$$

where all variables are non-zero integers. For $\mathbf{X} = (X_{ij})$ with $X_{ij} \geq 1$ let $N_\xi(\mathbf{X})$ be the number of solutions to (4.1) in boxes $\frac{1}{2}X_{ij} \leq |x_{ij}| \leq X_{ij}$. In many cases it will improve the readability considerably to relabel the variables, and it will be convenient to refer to (4.1) in the form

$$ab + c^2 + \xi^2 ywz^2 = 0, \quad yw \neq \square.$$

Consequently, we will write $\mathbf{X} = (A, B, C, Y, W, Z)$. We generally write

$$\|\mathbf{X}\| = \max(A, B, C, Y, W, Z)$$

Using this notation, we start with some elementary bounds.

Proposition 4.1. *We have*

$$N_\xi(\mathbf{X}) \ll \|\mathbf{X}\|^\varepsilon (ABC)^{1/2} (YW)^{3/4} Z^{1/2}.$$

Proof. The bound

$$(4.2) \quad N_\xi(\mathbf{X}) \ll \|\mathbf{X}\|^\varepsilon \min(ABC, CYWZ)$$

follows from a simple divisor argument. Alternatively, we can fix y, w, z so that

$$c^2 = -ywz^2\xi^2 + O(AB),$$

which then determines a, b up to a divisor function. The equation for c defines an interval of length

$$\ll \frac{AB}{\sqrt{YWZ}|\xi|}$$

so that

$$(4.3) \quad N_\xi(\mathbf{X}) \ll \|\mathbf{X}\|^\varepsilon YWZ \left(\frac{AB}{Z\sqrt{YW}|\xi|} + 1 \right) = \|\mathbf{X}\|^\varepsilon \left(\frac{AB\sqrt{YW}}{|\xi|} + YWZ \right).$$

The claim follows now from (4.2) and (4.3) after taking suitable geometric means, namely

$$\min(AB\sqrt{YW}, CYWZ) \leq (ABC)^{1/2} (YW)^{3/4} Z^{1/2}, \quad \min(ABC, YWZ) \leq \sqrt{ABCYWZ}.$$

□

We refine this argument a bit as follows. For $0 < \Delta \leq 1$ and $\mathbf{X} = (X_{ij})$ let $N_\xi^*(\mathbf{X}, \Delta)$ be the set of solutions to (4.1) with the same size conditions as $N_\xi(\mathbf{X})$ except that for one index

$$(ij) \in \{(21), (31), (32), (33)\}$$

the condition $\frac{1}{2}X_{ij} \leq |x_{ij}| \leq X_{ij}$ is replaced with $X_{ij} \leq |x_{ij}| \leq X_{ij}(1 + \Delta)$. We also define $N_\xi(\mathbf{X}, \Delta)$ to be the same size conditions as $N_\xi(\mathbf{X})$ except that $\frac{1}{2}X_{11} \leq |x_{11}| \leq X_{11}$ is replaced with $X_{11} \leq |x_{11}| \leq X_{11}(1 + \Delta)$. The following proposition investigates $N_\xi^*(\mathbf{X}, \Delta)$, while $N_\xi(\mathbf{X}, \Delta)$ comes up in Proposition 6.2.

Proposition 4.2. *We have*

$$N_\xi^*(\mathbf{X}, \Delta) \ll \|\mathbf{X}\|^\varepsilon \Delta^{1/2} (ABC)^{1/2} (YW)^{3/4} Z^{1/2}.$$

Proof. We distinguish two cases. If it is not the c -variable that is restricted, then by the same argument as above we have

$$\begin{aligned} N_\xi^*(\mathbf{X}, \Delta) &\ll \|\mathbf{X}\|^\varepsilon \min \left(ABC, \Delta CYWZ, \Delta(AB\sqrt{YW} + YWZ) \right) \\ &\ll \|\mathbf{X}\|^\varepsilon \left(\Delta(ABC)^{1/2} (YW)^{3/4} Z^{1/2} + \Delta^{1/2} (ABCYWZ)^{1/2} \right) \end{aligned}$$

and the claim follows.

If the restricted variable is c , then we have similarly

$$\begin{aligned} N_\xi^*(\mathbf{X}, \Delta) &\ll \|\mathbf{X}\|^\varepsilon \min \left(\Delta ABC, \Delta CYWZ, AB\sqrt{YW} + YWZ \right) \\ &\ll \|\mathbf{X}\|^\varepsilon (\Delta^{1/2}(ABC)^{1/2}(YW)^{3/4}Z^{1/2} + \Delta^{1/2}(ABCYWZ)^{1/2}). \end{aligned}$$

This completes the proof. \square

Remarks: 1) It will be convenient to introduce the following short-hand notation for expressions like those in Proposition 4.2. For $\zeta = (\zeta_1, \zeta_2, \zeta_3) \in \mathbb{R}^3$ and $\mathbf{X} = (A, B, C, Y, W, Z)$ we define

$$(4.4) \quad \mathbf{X}^{(\zeta)} = (AB)^{1-\zeta_1} C^{1-2\zeta_2} (YW)^{1-\zeta_3} Z^{1-2\zeta_3}.$$

With this notation, the bounds in Propositions 4.1 and 4.2 involve $\mathbf{X}^{(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})}$.

2) In order for $N_\xi(\mathbf{X})$ to be non-zero, we must have

$$(4.5) \quad C \ll (AB)^{1/2} + |\xi|Z\sqrt{YW}.$$

5. CHARACTER SUMS

We consider the following two character sums. For $a, c, z, \xi \in \mathbb{N}$, $h_1, h_2 \in \mathbb{Z}$ let

$$(5.1) \quad S_\xi(h_1, h_2, a, c, z) = \sum_{\substack{y, w \pmod{a} \\ a|c^2 + \xi^2 ywz^2}} e\left(\frac{h_1 y + h_2 w}{a}\right).$$

For $x, a \in \mathbb{N}$, $k_1, k_2 \in \mathbb{Z}$ let

$$(5.2) \quad T(k_1, k_2, x, a) = \sum_{\gamma^2 + x\xi^2 \equiv 0 \pmod{a}} e\left(\frac{h_1 \gamma + h_2 \xi}{a}\right).$$

Let τ denote the divisor function.

Lemma 5.1. *We have*

$$S_\xi(0, 0, a, c, z) = \sum_{\substack{a_1 a_2 a_3 = a \\ a_3 | c^2}} a_1(\xi^2 z^2, a_2 a_3) a_3 \mu(a_2)$$

and

$$|S_\xi(h_1, h_2, a, c, z)| \leq \tau(a)(a, h_1, h_2) a^{1/2} (a, c^2)^{1/2}.$$

Proof. We have

$$\begin{aligned} S_\xi(h_1, h_2, a, c, z) &= \frac{1}{a} \sum_{\alpha, y, w \pmod{a}} e\left(\frac{\alpha(c^2 + ywz^2\xi^2) + h_1 y + h_2 w}{a}\right) \\ &= \sum_{\substack{\alpha, w \pmod{a} \\ \alpha w z^2 \xi^2 + h_1 \equiv 0 \pmod{a}}} e\left(\frac{\alpha c^2 + h_2 w}{a}\right) \\ &= \sum_{\substack{a_1 a_2 = a \\ a_1 | h_1}} \sum_{\alpha \pmod{a_2}}^* \sum_{\substack{w \pmod{a} \\ w z^2 \xi^2 \equiv -\bar{\alpha} \frac{h_1}{a} \pmod{a_2}}} e\left(\frac{\alpha a_1 c^2 + h_2 w}{a}\right) \\ &= \sum_{\substack{a_1 a_2 = |a| \\ a_1(z^2 \xi^2, a_2) | h_1}} \sum_{\alpha \pmod{a_2}}^* \sum_{\substack{w \pmod{a} \\ w \equiv -\bar{\alpha} \frac{z^2 \xi^2}{(z^2, a_2)} \frac{h_1}{a_1(z^2 \xi^2, a_2)} \pmod{\frac{a_2}{(z^2 \xi^2, a_2)}}}} e\left(\frac{\alpha a_1 c^2 + h_2 w}{a}\right). \end{aligned}$$

The w -sum vanishes unless $\frac{a(z^2\xi^2, a_2)}{a_2} \mid h_2$, so we obtain

$$\begin{aligned} & \sum_{\substack{a_1 a_2 = a \\ a_1(z^2\xi^2, a_2) \mid (h_1, h_2)}} \sum_{\alpha \pmod{a_2}}^* e\left(\frac{bc^2}{a_2}\right) a_1(z^2\xi^2, a_2) e\left(-\frac{h_2 \overline{\frac{z^2\xi^2}{(z^2\xi^2, a_2)} \frac{h_1}{a_1(z^2\xi^2, a_2)}}}{a}\right) \\ &= \sum_{\substack{a_1 a_2 = a \\ a_1(z^2\xi^2, a_2) \mid (h_1, h_2)}} a_1(z^2\xi^2, a_2) S\left(c^2, -\frac{h_2}{a_1} \frac{h_1}{a_1(z^2\xi^2, a_2)} \frac{\overline{z^2\xi^2}}{(z^2\xi^2, a_2)}, a_2\right) \end{aligned}$$

where $S(., ., .)$ is the Kloosterman sum. If $h_1 = h_2 = 0$, then the claim follows by the formula

$$S(c^2, 0, a_2) = \sum_{d \mid (a_2, c^2)} d\mu\left(\frac{a_2}{d}\right).$$

In general we use Weil's bound $|S(c^2, *, a_2)| \leq \tau(a_2) a_2^{1/2} (c^2, a_2)^{1/2}$ to complete the proof of the lemma. \square

A number $D \in \mathbb{Z} \setminus \{0\}$ is a discriminant if $D \equiv 0, 1 \pmod{4}$. For each discriminant we denote by $\chi_D = (D/.)$ the corresponding quadratic character. It is primitive if and only if D is a fundamental discriminant. If $d \in \mathbb{N}$ is odd we write d^* for the unique discriminant with $|d^*| = d$. For an odd number d we write $\epsilon_d = \sqrt{\chi_{-4}(d)} \in \{1, i\}$.

Lemma 5.2. *We have*

$$T(0, 0, x, a) = \sum_{\substack{d_1 d_2 = a \\ (x, d_2) = \square}} d_1 \phi(d_2) \chi_{\frac{d_2}{(x, d_2)}}\left(\frac{-x}{(x, d_2)}\right) (x, d_2)^{1/2}$$

where d_2 runs over all discriminants (positive or negative). If a is odd, then

$$|T(k_1, k_2, x, a)| \leq \tau(a) (a, k_1^2 x + k_2^2) (a, x)^{1/2}.$$

Proof. We have

$$\begin{aligned} T(k_1, k_2, x, a) &= \frac{1}{a} \sum_{\alpha, c, z \pmod{a}} e\left(\frac{\alpha(c^2 + xz^2) + k_1 c + k_2 z}{a}\right) \\ &= \frac{1}{a} \sum_{\substack{d_1 d_2 = a \\ d_1 \mid (k_1, k_2)}} d_1^2 \sum_{\alpha \pmod{d_2}}^* \sum_{c, z \pmod{d_2}} e\left(\frac{\alpha(c^2 + xz^2) + \frac{k_1}{d_1} c + \frac{k_2}{d_1} z}{d_2}\right). \end{aligned}$$

Let us first consider the case $k_1 = k_2 = 0$. We split the modulus $d_2 = u2^\rho$ into an odd part and a power of two. The α, c, z -sum becomes

$$\sum_{\alpha \pmod{u}}^* \sum_{c, z \pmod{u}} e\left(\frac{\overline{2^\rho} \alpha(c^2 + xz^2)}{u}\right) \sum_{\alpha \pmod{2^\rho}}^* \sum_{c, z \pmod{2^\rho}} e\left(\frac{\overline{u} \alpha(c^2 + xz^2)}{2^\rho}\right).$$

By the well-known evaluation of the Gauß sum, the first c, z -sum equals

$$d_2 \epsilon_u \epsilon_{\frac{u}{(x, u)}}(x, u)^{1/2} \chi_{u^*}(\alpha) \chi_{\left(\frac{u}{(x, u)}\right)^*} \left(\frac{x}{(x, u)} \alpha\right).$$

Summing this over α , we see that only the contribution of $(x, u) = \square$ survives, and the first α, c, z -sum equals

$$\delta_{(x, u) = \square} u \phi(u) (x, u)^{1/2} \chi_{\frac{u^*}{(x, u)}} \left(-\frac{x}{(x, u)}\right).$$

For the second α, c, z -sum modulo powers of 2 we argue similarly, but we have to distinguish a few cases. Recall first that for odd α we have

$$\sum_{d \pmod{2^\rho}} e\left(\frac{\alpha d^2}{2^\rho}\right) = \begin{cases} 1, & \rho = 0, \\ 0, & \rho = 1, \\ 2^{\rho/2}(\chi_{2^\rho}(\alpha) + i\chi_{-2^\rho}(\alpha)), & \rho \geq 2. \end{cases}$$

If $4 \mid x/(2^\rho, x)$ and $4 \mid 2^\rho$, we obtain

$$\sum_{\alpha \pmod{2^\rho}}^* 2^\rho (2^\rho, x)^{1/2} (\chi_{2^\rho}(\alpha) + i\chi_{-2^\rho}(\alpha)) \left(\chi_{\frac{2^\rho}{(x, 2^\rho)}}\left(\frac{x}{(x, 2^\rho)}\alpha\right) + i\chi_{-\frac{2^\rho}{(x, 2^\rho)}}\left(\frac{x}{(x, 2^\rho)}\alpha\right) \right).$$

This vanishes, unless $(x, 2^\rho) = \square$ in which case it equals

$$2^\rho \phi(2^\rho)(x, 2^\rho)^{1/2} \sum_{\pm} \chi_{\pm \frac{2^\rho}{(x, 2^\rho)}}\left(-\frac{x}{(x, 2^\rho)}\right).$$

The remaining cases are simple: if $\rho = 1$, the sum vanishes and the case $\rho = 0$ is trivial. If $2^\rho \mid x$, the z -sum is equals $2^\rho = 2^{\rho/2}(x, 2^\rho)^{1/2}$, and evaluating the c -sum, we see that the α -sum vanishes unless ρ is even, i.e. $2^\rho = (x, 2^\rho) = \square$. If $x/(2^\rho, x) = 2$, the z -sum vanishes. In this way we confirm in all cases that the second α, c, z -sum equals

$$\delta_{(x, 2^\rho) = \square} 2^\rho \phi(2^\rho)(x, 2^\rho)^{1/2} \left(\delta_{4 \mid (x, 2^\rho)} \sum_{\pm} \chi_{\pm \frac{2^\rho}{(x, 2^\rho)}}\left(-\frac{x}{(x, 2^\rho)}\right) + \delta_{2^\rho \mid x} \right).$$

Combining this with the evaluation for odd moduli, we confirm that $T(0, 0, x, a)$ equals

$$\frac{1}{a} \sum_{\substack{d_1 d_2 = a \\ (d_2, x) = \square}} d_1^2 d_2 \phi(d_2)(x, d_2)^{1/2} \left(\delta_{4 \mid (x, d_2)} \sum_{\pm} \chi_{\pm \frac{d_2}{(x, d_2)}}\left(-\frac{x}{(x, d_2)}\right) + \delta_{2 \mid (x, d_2)} \chi_{\frac{d_2}{(x, d_2)}}\left(-\frac{x}{(x, d_2)}\right) \right).$$

This is equivalent to the formula given in the lemma.

We now turn to the estimation for general k_1, k_2 and for simplicity restrict ourselves to odd a , as in the statement of the lemma. In this case we can evaluate the two Gauß sums in c, z simply by completing the square, and we obtain

$$\frac{1}{a} \sum_{\substack{d_1 d_2 = a \\ d_1 \mid k_1 \\ d_1(x, d_2) \mid k_2}} \sum_{\alpha \pmod{d_2}}^* d_1^2 d_2 \epsilon_{d_2} \chi_{(x, d_2)^*}(\alpha) \chi_{\left(\frac{d_2}{(x, d_2)}\right)^*}\left(\frac{x}{(x, d_2)}\right) e\left(\frac{-4\alpha \left(\frac{k_1^2 x + k_2^2}{d_1^2(x, d_2)}\right)}{d_2}\right) (x, d_2)^{1/2} \epsilon_{\frac{d_2}{(x, d_2)}}.$$

Let δ denote the conductor of $\chi_{(x, d_2)^*}$ and write $d_2 = \delta \delta_1 \delta_2$ with $(\delta_2, \delta) = 1$, $\delta_1 \mid \delta^\infty$. Then by the well-known evaluation of quadratic character sums, the α -sum is bounded by

$$\delta_1 \delta^{1/2} \left(\delta_2, \frac{k_1^2 x + k_2^2}{d_1^2(x, d_2)} \right).$$

We decompose uniquely $(x, d_2) = t_1 t_2$ such that t_1 is the largest square coprime to t_2 . Then $\delta = \text{rad}(t_2)$, $\delta_1 = t_2/\text{rad}(t_2)$, $\delta_2 = t_1$, so that we obtain the upper bound

$$\begin{aligned} & \sum_{\substack{d_1 d_2 = a \\ d_1 \mid k_1 \\ d_1(x, d_2) \mid k_2}} d_1(x, d_2)^{1/2} \frac{t_2}{\text{rad}(t_2)^{1/2}} \left(t_1, \frac{k_1^2 x + k_2^2}{d_1^2(x, d_2)} \right) \\ &= \sum_{\substack{d_1 d_2 = a \\ d_1 \mid k_1 \\ d_1(x, d_2) \mid k_2}} \frac{(x, d_2)^{1/2}}{\text{rad}(t_2)^{1/2}} \left(t_1 d_1 t_2, \frac{k_1^2 x + k_2^2}{d_1 t_1} \right) \leq \sum_{d_1 d_2 = a} (x, d_2)^{1/2} (a, k_1^2 x + k_2^2), \end{aligned}$$

and the claim follows. \square

We also recall the following standard estimate.

Lemma 5.3. *Let V be a smooth function with compact support in $[-2, -1/3] \cup [1/3, 2]$ such that $V^{(j)} \ll_j \Omega^j$ for some $\Omega \geq 1$. If $\Delta \neq \square$ is a discriminant, $g \in \mathbb{N}$ and $N \geq 1$, then*

$$\sum_{(n,g)=1} \chi_\Delta(n) V\left(\frac{n}{N}\right) \ll \tau(g) N^{1/2} (\Omega|\Delta|)^{1/4+\varepsilon}.$$

In addition, if $n \in \mathbb{Z} \setminus \{0\}$ is not a square and $D \geq 1$, then

$$\sum_{(\Delta,g)=1} \chi_\Delta(n) V\left(\frac{\Delta}{D}\right) \ll \tau(g) D^{1/2} (\Omega|n|)^{1/4+\varepsilon}$$

where the sum runs over all discriminants. The implied constants depend only on ε .

Proof. This is standard by Mellin inversion and the convexity bound for Dirichlet L -functions (with Euler factors at primes dividing g removed)

$$L^{(g)}(s, \chi_\Delta) \ll \tau(g) (|\Delta|(1+|s|))^{1/4+\varepsilon}, \quad \Re s = 1/2.$$

Suffice it to say that the (two-sided) Mellin transform of V is entire and satisfies

$$\widehat{V}(s) \ll_{\Re s, A} \left(1 + \frac{|\Im s|}{\Omega}\right)^{-A}$$

for all $A > 0$. The second bound follows from quadratic reciprocity as follows: By quadratic reciprocity we have $\chi_\Delta(n) = \chi_{\tilde{n}}(\Delta)$ for $n > 0$ where \tilde{n} is the discriminant computed as follows: write $n = 2^a y$ with y odd, and let y^* denote the discriminant satisfying $|y^*| = y$. Let $a' = a + 2$ if $a = 2$ and $a' = a$ otherwise. Then $\tilde{n} = 2^{a'} y^*$. If $n < 0$ then $\chi_\Delta(n) = \chi_{\tilde{n}}(\Delta) \chi_\Delta(-1)$. In this way, the second bound follows from the first by detecting the condition $\Delta \equiv 0, 1 \pmod{4}$ by characters. \square

Remark: Using the strongest available uniform subconvexity bounds, the exponents $1/4$ can be replaced with $1/6$.

6. UPPER BOUND ESTIMATES

For $\mathbf{X} = (A, B, C, Y, W, Z)$ and $\xi \in \mathbb{Z} \setminus \{0\}$ we recall the definition of $N_\xi(\mathbf{X})$ and $N_\xi(\mathbf{X}, \Delta)$ from Section 4, and we now define a smooth version of the former. Let $\Omega \geq 3$ be a parameter. We choose an even, smooth, non-negative test function V with support in $[-1 - \Omega^{-1}, -1/2 + \Omega^{-1}] \cup [1/2 - \Omega^{-1}, 1 + \Omega]$ and $V(x) = 1$ on $[-1, -1/2] \cup [1/2, 1]$ satisfying $V^{(j)}(x) \ll_j \Omega^j$ for all $j \in \mathbb{N}_0$. We define

$$\tilde{N}_\xi(\mathbf{X}, \Omega) = \sum_{\substack{ab+c^2+\xi^2 ywz^2=0 \\ yw \neq -\square}} V\left(\frac{a}{A}\right) V\left(\frac{b}{B}\right) V\left(\frac{c}{C}\right) V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) V\left(\frac{z}{Z}\right).$$

(Strictly speaking, $\tilde{N}_\xi(\mathbf{X}, \Omega)$ depends on V and not only on Ω , but this small abuse of notation is convenient.) Fix a small constant $\lambda \leq 10^{-6}$ and consider \mathbf{X}, ξ satisfying

$$\begin{aligned} (6.1) \quad & \min(A, B, C) \geq \max(A, B, C)^{1-\lambda}, \\ & \min(Y, W) \geq \max(Y, W) \|\mathbf{X}\|^{-\lambda}, \\ & \min(AB, C^2, YWZ^2) \geq \max(AB, C^2, YWZ^2)^{1-\lambda}. \\ & |\xi| \leq \|\mathbf{X}\|^\lambda. \end{aligned}$$

We will see later that these are the critical size conditions.

Our aim in this section is to establish the following three estimates that we will prove simultaneously. Recall the notation (4.4).

Proposition 6.1. *For \mathbf{X}, ξ satisfying (6.1) we have*

$$N_\xi(\mathbf{X}) \ll \mathbf{X}^{(\frac{3}{4}, 0, \frac{1}{4})} + \mathbf{X}^{(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})}.$$

Proposition 6.2. *For \mathbf{X}, ξ satisfying (6.1) and $0 < \Delta < 1$ we have*

$$N_\xi(\mathbf{X}, \Delta) \ll (\mathbf{X}^{(\frac{3}{4}, \frac{1}{8}, \frac{1}{8})} + \mathbf{X}^{(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})})(\Delta \|\mathbf{X}\|^\varepsilon + \|\mathbf{X}\|^{-1/5}).$$

Proposition 6.3. *For \mathbf{X}, ξ satisfying (6.1) and $\Omega \geq 3$ we have*

$$\tilde{N}_\xi(\mathbf{X}, \Omega) = M_2 + O\left(\left(\mathbf{X}^{(\frac{5}{8}, \frac{1}{8}, \frac{1}{4})} + \mathbf{X}^{(\frac{3}{4}, \frac{1}{8}, \frac{1}{8})}\right)\Omega \|\mathbf{X}\|^{11\lambda} \frac{A^{1/2}}{Y}\right)$$

where M_2 is given by (6.4) below. We also have

$$\tilde{N}_\xi(\mathbf{X}, \Omega) = \tilde{M}_3 + O\left(\mathbf{X}^{(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})}\Omega^4 \left(\frac{\|\mathbf{X}\|^{10\lambda}}{Z} + \frac{1}{(YW)^{1/21}}\right)\right)$$

where \tilde{M}_3 is defined in (6.12) below.

The rest of this section is devoted to the proof. We will occasionally use the following notation: for a positive integer n let \sqrt{n}^+ denote the smallest integer whose square is a multiple of n .

We have trivially $N_\xi(\mathbf{X}) \leq \tilde{N}_\xi(\mathbf{X}, \Omega)$ for every $\Omega \geq 3$ and

$$(6.2) \quad \tilde{N}_\xi(\mathbf{X}, \Omega) = \sum_{\substack{a, c, y, w, z \\ a|c^2 + \xi^2 y w z^2 \\ y w \neq -\square}} V\left(\frac{a}{A}\right) V\left(\frac{-\xi^2 y w z^2 - c^2}{aB}\right) V\left(\frac{c}{C}\right) V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) V\left(\frac{z}{Z}\right).$$

We apply two different strategies. We first apply Poisson summation in y, w and then Poisson summation in c, z . In order to obtain Proposition 6.2, we will occasionally replace $V(a/A)$ with the characteristic function on $A \leq |a| \leq A(1 + \Delta)$. In the interest of a reasonably compact presentation, we will not introduce extra notation for this.

6.1. Poisson summation in y, w . We first add the contribution of $y w = -\square$ to (6.2). As in the proof of Proposition 4.1, we see by a divisor argument that this infers an error of at most $O(C(YW)^{1/2} Z \|\mathbf{X}\|^\varepsilon)$. Then we apply Poisson summation y, w to obtain

$$(6.3) \quad \begin{aligned} \tilde{N}_\xi(\mathbf{X}, \Omega) &= \sum_{a, c, z} V\left(\frac{a}{A}\right) V\left(\frac{c}{C}\right) V\left(\frac{z}{Z}\right) \frac{1}{a^2} \sum_{h_1, h_2} S_\xi(h_1, h_2, a, c, z) \Phi_\xi(h_1, h_2, a, c, z) \\ &\quad + O(C(YW)^{1/2} Z \|\mathbf{X}\|^\varepsilon) \end{aligned}$$

with S_ξ as in (5.1) and

$$\Phi_\xi(h_1, h_2, a, c, z) = \int_{\mathbb{R}} \int_{\mathbb{R}} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) V\left(\frac{-\xi^2 y w z^2 - c^2}{aB}\right) e\left(-\frac{h_1 y + h_2 w}{|a|}\right) dy dw.$$

Let

$$H_1 = \Omega\left(\frac{A}{Y} + \frac{\xi^2 W Z^2}{B}\right), \quad H_2 = \Omega\left(\frac{A}{W} + \frac{\xi^2 Y Z^2}{B}\right).$$

Lemma 6.4. *For $|a| \asymp A$, $|z| \asymp Z$ we have*

$$\Phi_\xi(h_1, h_2, a, c, z) \ll_N \frac{(AB)^{1/4} (YW)^{3/4}}{(Z|\xi|)^{1/2}} \left(1 + \frac{|h_1|}{H_1} + \frac{|h_2|}{H_2}\right)^{-N}$$

for arbitrary $N > 0$ and

$$(a\partial_a)^{j_1} (c\partial_c)^{j_2} (z\partial_z)^{j_3} \Phi_\xi(0, 0, a, c, z) \ll_j \frac{(AB)^{1/4} (YW)^{3/4}}{(Z|\xi|)^{1/2}} \left(1 + \frac{C^2}{AB}\right)^{j_2} \left(1 + \frac{\xi^2 Y W Z^2}{AB}\right)^{j_1} \Omega^{j_1 + j_2 + j_3}$$

for arbitrary $\mathbf{j} \in \mathbb{N}_0^3$.

Proof. We observe that the volume of the (y, w) -region defined by

$$|y| \asymp Y, \quad |w| \asymp W, \quad |c^2 + \xi^2 y w z^2| \asymp AB$$

is trivially bounded by $O(YW)$, but also by $O(AB/z^2 \xi^2)$ and so by $O((AB)^{1/4}(YW)^{3/4}/|z\xi|^{1/2})$. The first claim follow now by repeated integration by parts, the second by differentiating under the integral sign. There is an important subtlety: we combine each application of the operator $z\partial_z$ with a partial integration in y , i.e.

$$\int_{\mathbb{R}} z\partial_z \left[V\left(\frac{y}{Y}\right) V\left(\frac{-\xi^2 y w z^2 - c^2}{aB}\right) \right] dy = -2 \int_{\mathbb{R}} \partial_y \left(y V\left(\frac{y}{Y}\right) \right) V\left(\frac{-\xi^2 y w z^2 - c^2}{aB}\right) dy.$$

This completes the proof. \square

We now write the right hand side of (6.3) as $M_1 + M_2$ where M_1 is the off-diagonal contribution $(h_1, h_2) \neq (0, 0)$ and M_2 is the diagonal contribution $h_1 = h_2 = 0$. By Lemma 5.1 we have

$$(6.4) \quad M_2 = \sum_{a, c, z} V\left(\frac{a}{A}\right) V\left(\frac{c}{C}\right) V\left(\frac{z}{Z}\right) \frac{1}{a^2} \sum_{\substack{a_1 a_2 a_3 = |a| \\ a_3 |c|^2}} a_1 (\xi^2 z^2, a_2 a_3) a_3 \mu(a_2) \Phi_\xi(0, 0, a, c, z).$$

We postpone the analysis of M_2 and investigate first M_1 . By Lemma 5.1 and Lemma 6.4 we have

$$\begin{aligned} M_1 &\ll \sum_{a, c, z} V\left(\frac{a}{A}\right) V\left(\frac{c}{C}\right) V\left(\frac{z}{Z}\right) \frac{(AB)^{1/4}(YW)^{3/4}}{A^2(Z|\xi|)^{1/2}} \\ &\quad \times \sum_{(h_1, h_2) \neq (0, 0)} (a, h_1, h_2) \tau(a) |a|^{1/2} (a, c) \left(1 + \frac{|h_1|}{H_1} + \frac{|h_2|}{H_2}\right)^{-10} \\ &\ll \sum_a V\left(\frac{a}{A}\right) \frac{CZ(AB)^{1/4}(YW)^{3/4}}{A^{3/2}(Z|\xi|)^{1/2}} \sum_{(h_1, h_2) \neq (0, 0)} (a, h_1, h_2) \tau(a)^2 \left(1 + \frac{|h_1|}{H_1} + \frac{|h_2|}{H_2}\right)^{-10} \\ &\ll \sum_a V\left(\frac{a}{A}\right) \frac{CZ(AB)^{1/4}(YW)^{3/4}}{A^{3/2}(Z|\xi|)^{1/2}} \tau(a)^3 (1 + H_1 + H_2). \end{aligned}$$

By several applications of (6.1), we see that

$$\frac{1 + H_1 + H_2}{A^{1/2-\varepsilon}} \ll \Omega \|\mathbf{X}\|^{10\lambda} \frac{A^{1/2}}{Y},$$

so that

$$(6.5) \quad M_1 \ll \frac{C(AB)^{1/4}(YW)^{3/4} Z^{1/2}}{|\xi|^{1/2}} \Omega \|\mathbf{X}\|^{10\lambda} \frac{A^{1/2}}{Y}.$$

Note that up until now we have not used any property of the weight $V(a/A)$ except that it restricts $a \ll A$. In particular, (6.5) continues to hold with any Ω if $V(a/A)$ is replaced by the characteristic function on $A \leq |a| \leq A(1 + \Delta)$.

We can now complete the proof of the first half of Proposition 6.3 by recalling (6.3) and noting that (6.1) implies

$$CZ\sqrt{YW}(ABYW)^\varepsilon \ll (AB)^{1/4} C(YW)^{3/4} Z^{1/2} \cdot \frac{\|\mathbf{X}\|^{10\lambda}}{Y},$$

so that

$$N_\xi(\mathbf{X}, \Omega) = M_2 + O\left((AB)^{1/4} C(YW)^{3/4} Z^{1/2} \cdot \Omega \frac{\|\mathbf{X}\|^{10\lambda}}{Y}\right).$$

As in the proof of Proposition 4.2 we reduce the power of C using (4.5), which completes the proof of the first half of Proposition 6.3.

We now turn to M_2 as given in (6.4). We will evaluate this asymptotically later, but for now content ourselves with an upper bound given by

$$(6.6) \quad M_2 \ll \frac{C}{A^2} \cdot \frac{(AB)^{1/4}(YW)^{3/4}}{(Z|\xi|)^{1/2}} \sum_{a_1, a_2, a_3, z} V\left(\frac{a_1 a_2 a_3}{A}\right) V\left(\frac{z}{Z}\right) a_1 a_3^{1/2} (z^2 \xi^2, a_2 a_3).$$

If we replace $V(a/A)$ with the characteristic function on $A \leq |a| \leq A(1 + \Delta)$, a very soft bound is given by

$$\begin{aligned} &\ll \frac{C}{A^2} \cdot \frac{(AB)^{1/4}(YW)^{3/4}}{(Z|\xi|)^{1/2}} \sum_z \sum_{A \leq a_1 a_2 a_3 \leq A(1+\Delta)} V\left(\frac{z}{Z}\right) a_1 a_2 a_3 (z \xi, a_3) \\ &\ll \frac{C}{A^2} \cdot \frac{(AB)^{1/4}(YW)^{3/4}}{(Z|\xi|)^{1/2}} \Delta A^{2+\varepsilon} Z \tau(\xi) \ll \Delta (AB)^{1/4} C (YW)^{3/4} Z^{1/2} A^\varepsilon. \end{aligned}$$

Together with (6.5) for $\Omega = 3$ we obtain

$$(6.7) \quad N_\xi(\mathbf{X}, \Delta) \ll (AB)^{1/4} C (YW)^{3/4} Z^{1/2} \left(\Delta \|\mathbf{X}\|^\varepsilon + \|\mathbf{X}\|^{10\lambda} \frac{A^{1/2}}{Y} \right).$$

After this interlude we now return to (6.6) and estimate the right hand side by

$$\leq \frac{C}{A^2} \frac{(AB)^{1/4}(YW)^{3/4}}{(Z|\xi|)^{1/2}} \sum_d \sum_{\substack{a_1, a_2, a_3 \\ d|a_2 a_3}} \sum_{\substack{a_1, a_2, a_3 \\ d|(a_1 a_2 a_3)}} V\left(\frac{a_1 a_2 a_3}{A}\right) V\left(\frac{z}{Z}\right) a_1 a_3^{1/2} d.$$

For notational simplicity let us write $d_\xi = d/(d, \xi^2)$. Then we can continue to estimate

$$\begin{aligned} M_2 &\ll \frac{C}{A^2} \frac{(AB)^{1/4}(YW)^{3/4}}{(Z|\xi|)^{1/2}} \sum_d \sum_{d_2 d_3 = d} \sum_{\substack{a_1, a_2, a_3 \\ d_2 | a_2, d_3 | a_3}} V\left(\frac{a_1 a_2 a_3}{A}\right) \frac{Z}{\sqrt{d_\xi}^+} a_1 a_3^{1/2} d \\ &\ll \frac{C}{A^2} \frac{(AB)^{1/4}(YW)^{3/4}}{(Z|\xi|)^{1/2}} \sum_d d \sum_{d_2 d_3 = d} \frac{A^2}{d_2^2 d_3^{3/2}} \frac{Z}{\sqrt{d_\xi}^+}. \end{aligned}$$

The d -sum is

$$\leq \sum_d \frac{\tau(d)}{d^{1/2} \sqrt{d_\xi}^+} \leq \sum_{\eta|\xi^2} \frac{\tau(\eta)}{\eta^{1/2}} \sum_d \frac{\tau(d)}{d^{1/2} \sqrt{d}^+} \ll \tau(\xi^2).$$

Combining this off-diagonal contribution (6.5) and choosing $\Omega = 3$, we have shown our first important bound

$$(6.8) \quad N_\xi(\mathbf{X}) \leq \tilde{N}_\xi(\mathbf{X}, 3) \ll \frac{C(AB)^{1/4}(YW)^{3/4} Z^{1/2}}{|\xi|^{1/2}} \left(\tau(\xi^2) + \|\mathbf{X}\|^{10\lambda} \frac{A^{1/2}}{Y} \right).$$

6.2. Poisson summation in c, z . We now return to (6.2) and apply Poisson summation in c, z getting

$$N_\xi(\mathbf{X}, \Omega) = \sum_{wy \neq -\square} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) \sum_a V\left(\frac{a}{A}\right) \frac{CZ}{a^2} \sum_{k_1, k_2} T(k_1, k_2, yw\xi^2, a) \Psi(k_1, k_2, yw\xi^2, a)$$

with T as in (5.2) and

$$\Psi(k_1, k_2, x, a) = \int_{\mathbb{R}} \int_{\mathbb{R}} V\left(\frac{c}{C}\right) V\left(\frac{z}{Z}\right) V\left(-\frac{c^2 + xz^2}{aB}\right) e\left(\frac{-k_1 c - k_2 z}{|a|}\right) dc dz.$$

Let

$$K_1 = \Omega\left(\frac{A}{C} + \frac{C}{B}\right), \quad K_2 = \Omega\left(\frac{A}{Z} + \frac{\xi^2 YWZ}{B}\right).$$

Lemma 6.5. *For $|a| \asymp A$ we have*

$$\Psi(k_1, k_2, x, a) \ll_N \frac{(CZAB)^{1/2}}{|x|^{1/4}} \left(1 + \frac{|k_1|}{K_1} + \frac{|k_2|}{K_2}\right)^{-N}$$

and

$$(x\partial_x)^{j_1} (a\partial_a)^{j_2} \Psi(0, 0, x, a) \ll_{j_1, j_2} \frac{(CZAB)^{1/2}}{|x|^{1/4}} \Omega^{j_1+j_2}$$

for arbitrary $N, j_1, j_2 \in \mathbb{N}_0$.

Proof. As in Lemma 6.4 we observe that the volume of the (c, z) -region defined by $|c| \asymp C$, $|z| \asymp Z$, $|c^2 + xz^2| \asymp AB$ is trivially bounded by $O(CZ)$, but also by $O(AB/|x|^{1/2})$. Indeed, if $x > 0$, this is the volume of the ellipse $c^2 + xz^2 \ll AB$, while for $x < 0$ we have $c = \sqrt{|x|z^2 + O(AB)}$ for fixed $|z| \asymp Z$, which has volume $\ll AB/(|x|^{1/2}Z)$. Taking the geometric mean, we bound (c, z) -volume by $O((CZAB)^{1/2}/|x|^{1/4})$. The claims follow now by repeated partial integration and differentiation under the integral sign. As in the proof of Lemma 6.4, each application of $x\partial_x$ is coupled with an integration by parts in z . \square

As before we decompose

$$N_\xi(\mathbf{X}, \Omega) = \tilde{M}_1 + \tilde{M}_2$$

where \tilde{M}_1 is the off-diagonal contribution $(k_1, k_2) \neq 0$ and \tilde{M}_2 is the diagonal contribution $k_1 = k_2 = 0$. For notational simplicity let

$$\Xi(k_1, k_2) = \frac{\sqrt{CZAB}}{|\xi|^{1/2}(YW)^{1/4}} \left(1 + \frac{|k_1|}{K_1} + \frac{|k_2|}{K_2}\right)^{-10}.$$

Using Lemma 5.2 and Lemma 6.5 we obtain

$$\begin{aligned} \tilde{M}_1 &\ll \sum_{wy \neq -\square} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) \sum_a V\left(\frac{a}{A}\right) \frac{1}{A^2} \sum_{(k_1, k_2) \neq (0,0)} (a, \xi^2 yw)^{1/2} (a, 2^\infty (k_1^2 \xi^2 yw + k_2^2)) \Xi(k_1, k_2) \\ &\ll \sum_{d_1, d_2 \ll A} \sum_{\substack{wy \neq -\square \\ d_1 | \xi^2 yw}} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) \sum_{[d_1, d_2] | a} V\left(\frac{a}{A}\right) \frac{1}{A^2} \sum_{\substack{(k_1, k_2) \neq (0,0) \\ d_2 | 2^\infty (k_1^2 \xi^2 yw + k_2^2)}} d_1^{1/2} d_2 \Xi(k_1, k_2) \\ &\ll \sum_{d_1, d_2 \ll A} \sum_{\substack{wy \neq -\square \\ d_1 | \xi^2 yw}} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) \frac{\log A}{A} \sum_{\substack{(k_1, k_2) \neq (0,0) \\ d_2 | k_1^2 \xi^2 yw + k_2^2}} \frac{d_1^{1/2} d_2}{[d_1, d_2]} \Xi(k_1, k_2). \end{aligned}$$

We write $k_1^2 \xi^2 yw = -k_2^2 + \alpha d_2$. Since $yw \neq -\square$ and $(k_1, k_2) \neq (0, 0)$, we have $\alpha \neq 0$, and moreover $\alpha \equiv -k_2^2 \pmod{d_1}$. Once α and k_2 are chosen, the variables y, w, k_1 are determined up to a divisor function. We conclude the upper bound

$$\tilde{M}_1 \ll \sum_{d_1, d_2 \ll A} \frac{\log A}{A} \left(1 + \frac{K_2}{\sqrt{d_1^+}}\right) \frac{K_2^2 + K_1^2 \xi^2 YW}{d_2} \frac{d_1^{1/2} d_2}{[d_1, d_2]} \frac{\sqrt{CZAB}}{|\xi|^{1/2}(YW)^{1/4}} \|\mathbf{X}\|^\varepsilon$$

and so

$$\tilde{M}_1 \ll \frac{1}{A} (A^{1/2} + K_2) (K_2^2 + K_1^2 \xi^2 YW) \frac{\sqrt{CZAB}}{|\xi|^{1/2}(YW)^{1/4}} \|\mathbf{X}\|^\varepsilon.$$

We now invoke (6.1) several times to conclude that

$$\frac{(A^{1/2} + K_2)}{A} \left(\frac{K_2^2}{YW} + K_1^2 \xi^2\right) \|\mathbf{X}\|^\varepsilon \ll \Omega^4 \frac{\|\mathbf{X}\|^{10\lambda}}{Z}.$$

Thus we obtain the simplified bound

$$(6.9) \quad \tilde{M}_1 \ll \frac{\sqrt{CZAB}(YW)^{3/4}}{|\xi|^{1/2}} \Omega^4 \frac{\|\mathbf{X}\|^{10\lambda}}{Z}.$$

Note that in order to derive this bound we did not use any property of the weight $V(a/A)$, except that it bounds $a \ll A$. In particular, (6.9) continues to hold for all Ω , if $V(a/A)$ is replaced with the characteristic function on $A \leq |a| \leq A(1 + \Delta)$.

On the other hand, again by Lemma 5.2 we have

$$(6.10) \quad \begin{aligned} \tilde{M}_2 &= 2 \sum_{yw \neq -\square} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) \sum_{\substack{d_1, d_2 \\ (\xi^2 yw, d_2) = \square}} V\left(\frac{d_1 d_2}{A}\right) \frac{\phi(d_2)}{d_1 d_2^2} \\ &\quad \times \chi_{\frac{d_2}{(\xi^2 yw, d_2)}} \left(-\frac{\xi^2 yw}{(\xi^2 yw, d_2)} \right) (\xi^2 yw, d_2)^{1/2} \Psi(0, 0, \xi^2 yw, d_1 d_2) \end{aligned}$$

where d_2 runs over all (positive or negative) discriminants and the factor 2 comes from the fact that V is even and we have used the decomposition $|a| = d_1 |d_2|$ from Lemma 5.2.

Before we manipulate this further, we complete the proof of Proposition 6.2. Replacing $V(d_1 d_2/A)$ in the previous display by the characteristic function on $A \leq |d_1 d_2| \leq A(1 + \Delta)$, we obtain by a simple divisor estimate the upper bound

$$\ll \tau(\xi^2) YW \Delta \|\mathbf{X}\|^\varepsilon \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}}$$

for the right hand side of (6.10). By (6.1), the factor $\tau(\xi^2)$ can be absorbed into $\|\mathbf{X}\|^\varepsilon$. Combining this with (6.9), we obtain

$$N_\xi(\mathbf{X}, \Delta) \ll (ABC)^{1/2} (YW)^{3/4} Z^{1/2} \left(\frac{\Delta \|\mathbf{X}\|^\varepsilon}{|\xi|^{1/2}} + \frac{\|\mathbf{X}\|^{10\lambda}}{Z} \right).$$

Together with (6.7) we obtain

$$N_\xi(\mathbf{X}, \Delta) \ll (AB)^{1/4} C^{1/2} ((AB)^{1/4} + C^{1/2}) (YW)^{3/4} Z^{1/2} \left(\frac{\Delta \|\mathbf{X}\|^\varepsilon}{|\xi|^{1/2}} + \|\mathbf{X}\|^{10\lambda} \min\left(\frac{A^{1/2}}{Y}, \frac{1}{Z}\right) \right).$$

Using (4.5), we replace the second appearance of $C^{1/2}$ with $(AB)^{1/4} + C^{1/4}(|\xi|Z)^{1/4} (YW)^{1/8}$. By several applications of (6.1) we have

$$(6.11) \quad \min\left(\frac{A^{1/2}}{Y}, \frac{1}{Z}\right) \leq \frac{A^{1/4}}{(YZ)^{1/2}} \leq \frac{Y^\lambda}{B^{1/4}} \frac{(AB)^{1/4}}{(YWZ^2)^{1/4}} \leq \frac{(YAB)^\lambda}{B^{1/4}} \ll \|\mathbf{X}\|^{-1/5-15\lambda}$$

which completes the proof of Proposition 6.2.

After this interlude we return to (6.10). With $\delta^2 = (\xi^2 yw, d_2)$ we re-write this as

$$\tilde{M}_2 = 2 \sum_{\delta} \sum_{\substack{\xi^2 yw \equiv 0 \pmod{\delta^2} \\ yw \neq -\square}} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) \sum_{d_1, d_2} V\left(\frac{d_1 d_2 \delta^2}{A}\right) \frac{\phi(d_2 \delta^2)}{d_1 d_2^2 \delta^3} \chi_{d_2} \left(-\frac{\xi^2 yw}{\delta^2} \right) \Psi(0, 0, \xi^2 yw, d_1 d_2 \delta^2)$$

where the condition $(\xi^2 yw/\delta^2, d_2) = 1$ is automatic from the Jacobi symbol. Write $\delta' = \delta/(\delta, \xi)$, so $\delta'^2 \mid yw$. We write $(\delta'^2, y) = f$, so that $g = \frac{\delta'^2}{f} \mid w$, and we obtain

$$2 \sum_{\delta} \sum_{fg = \delta'^2} \sum_{\substack{yw \neq -\square \\ (g, y) = 1}} V\left(\frac{fy}{Y}\right) V\left(\frac{gw}{W}\right) \sum_{d_1, d_2} V\left(\frac{d_1 d_2 \delta'^2}{A}\right) \frac{\phi(d_2 \delta'^2)}{d_1 d_2^2 \delta'^3} \chi_{d_2} \left(-\frac{\xi^2 yw}{(\delta, \xi)^2} \right) \Psi(0, 0, \xi^2 \delta'^2 yw, d_1 d_2 \delta'^2).$$

We write $\delta = \delta' \xi_1$ with $\xi_1 \xi_2 = \xi$ and $(\xi_2, \delta') = 1$ getting

$$\begin{aligned} \tilde{M}_2 &= 2 \sum_{\substack{\xi_1 \xi_2 = \xi \\ (\xi_2, fg) = 1}} \sum_{fg = \square} \sum_{\substack{yw \neq -\square \\ (g, y) = 1}} V\left(\frac{fy}{Y}\right) V\left(\frac{gw}{W}\right) \sum_{d_1, d_2} V\left(\frac{d_1 d_2 fg \xi_1^2}{A}\right) \frac{\phi(d_2 fg \xi_1^2)}{d_1 d_2^2 (fg)^{3/2} \xi_1^3} \\ &\quad \times \chi_{d_2} (-\xi_2^2 yw) \Psi(0, 0, \xi^2 fg yw, d_1 d_2 fg \xi_1^2). \end{aligned}$$

We further decompose

$$\tilde{M}_2 = \tilde{M}_3 + \tilde{M}_4$$

where \tilde{M}_3 is the contribution $d_2 = \square$ and \tilde{M}_4 is the contribution $d_2 \neq \square$. We have

$$(6.12) \quad \begin{aligned} \tilde{M}_3 = 2 \sum_{\substack{\xi_1 \xi_2 = \xi \\ (\xi_2, fg) = 1}} \sum_{fg = \square} \sum_{\substack{wy \neq -\square \\ (g, y) = 1}} V\left(\frac{fy}{Y}\right) V\left(\frac{gw}{W}\right) \sum_{\substack{d_1, d_2 \\ (d_2, \xi_2 yw) = 1}} V\left(\frac{d_1 d_2^2 fg F_1^2}{A}\right) \frac{\phi(d_2^2 fg \xi_1^2)}{d_1 d_2^4 (fg)^{3/2} \xi_1^3} \\ \times \Psi(0, 0, \xi^2 fg yw, d_1 d_2^2 fg \xi_1^2). \end{aligned}$$

We will later evaluate this asymptotically, but for now we content ourselves with the upper bound

$$(6.13) \quad \begin{aligned} \tilde{M}_3 &\ll \sum_{\xi_1 \xi_2 = \xi} \sum_{fg = \square} \sum_{w, y} V\left(\frac{fy}{Y}\right) V\left(\frac{gw}{W}\right) \sum_{d_1, d_2} V\left(\frac{d_1 d_2^2 fg \xi_1^2}{A}\right) \frac{1}{d_1 d_2^2 (fg)^{1/2} \xi_1} \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}} \\ &\ll \tau(\xi) \frac{\sqrt{CZAB} (YW)^{3/4}}{|\xi|^{1/2}}. \end{aligned}$$

To bound \tilde{M}_4 , we insert a smooth partition of unity localizing $|d_2| \asymp D_2$, say, so that

$$\tilde{M}_4 = \sum_{D_2} \tilde{M}_4(D_2)$$

where $D_2 \ll A$ runs over powers of 2 and

$$\begin{aligned} \tilde{M}_4(D_2) &= \sum_{\substack{\xi_1 \xi_2 = \xi \\ (\xi_2, fg) = 1}} \sum_{fg = \square} \sum_{\substack{wy \neq -\square \\ (g, y) = 1}} V\left(\frac{fy}{Y}\right) V\left(\frac{gw}{W}\right) \sum_{d_1} \sum_{\substack{d_2 \neq \square \\ (d_2, \xi) = 1}} V\left(\frac{d_1 d_2 fg \xi_1^2}{A}\right) W\left(\frac{d_2}{D_2}\right) \\ &\quad \times \frac{\phi(d_2 fg \xi_1^2)}{d_1 d_2^2 (fg)^{3/2} \xi_1^3} \chi_{d_2}(-yw) \Psi(0, 0, \xi^2 fg yw, d_1 d_2 fg \xi_1^2) \end{aligned}$$

for a suitable smooth compactly supported function W . We estimate $\tilde{M}_4(D_2)$ in two ways. First we re-insert the contribution of $yw = -\square$ at the cost of an error

$$\ll \tau(\xi) (YW)^{1/2} \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}}$$

and then sum over y, w with Lemma 5.3 and 6.5 getting a bound

$$(6.14) \quad \ll \tau(\xi) (YW)^{1/2} \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}} (\Omega D_2)^{1/2+\varepsilon}.$$

Obviously this majorizes the previous error term.

Alternatively, we can also re-insert the contribution of $d_2 = \square$ at the cost of an error

$$\ll \tau(\xi) \frac{YW}{D_2^{1/2}} \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}}$$

and then sum over d_2 with Lemma 5.3 and 6.5 getting a bound

$$(6.15) \quad \ll \tau(\xi)^2 \frac{YW}{D_2^{1/2}} \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}} (\Omega YW)^{1/4+\varepsilon},$$

which again majorizes the previous error.

Combining (6.14) and (6.15), we bound $\tilde{M}_4(D_2)$ by

$$\begin{aligned} &\ll \tau(\xi)^2 \frac{\sqrt{CZAB} (YW)^{3/4}}{|\xi|^{1/2}} \min\left(\frac{(\Omega D_2)^{1/2+\varepsilon}}{(YW)^{1/2}}, \frac{(\Omega YW)^{1/4+\varepsilon}}{D_2^{1/2}}\right) \\ &\ll \tau(\xi)^2 \frac{\sqrt{CZAB} (YW)^{3/4}}{|\xi|^{1/2}} \frac{\Omega^{1/2+\varepsilon}}{(YW)^{1/20-\varepsilon} D_2^{1/10-\varepsilon}}. \end{aligned}$$

Summing over D_2 , we finally obtain

$$\tilde{M}_4 \ll \sqrt{CZAB}(YW)^{3/4} \frac{\Omega^{1/2+\varepsilon}}{(YW)^{1/20-\varepsilon}}.$$

Combining this with (6.9) we have

$$\begin{aligned} \tilde{M}_1 + \tilde{M}_4 &\ll \sqrt{CZAB}(YW)^{3/4} \left(\frac{\Omega^{2/3}}{(YW)^{1/21}} + \Omega^4 \frac{\|\mathbf{X}\|^{10\lambda}}{Z} \right) \\ &\ll \sqrt{CZAB}(YW)^{3/4} \Omega^4 \left(\frac{\|\mathbf{X}\|^{10\lambda}}{Z} + \frac{1}{(YW)^{1/21}} \right). \end{aligned}$$

Since $N_\xi(\mathbf{X}, \Omega) = \tilde{M}_1 + \tilde{M}_3 + \tilde{M}_4$, this completes the proof of the second half of Proposition 6.3.

On the other hand, choosing $\Omega = 3$, we also invoke (6.13) to obtain our second important bound

$$N_\xi(\mathbf{X}) \leq N_\xi(\mathbf{X}, 3) = \tilde{M}_1 + \tilde{M}_3 + \tilde{M}_4 \ll \frac{(ABC)^{1/2}(YW)^{3/4}Z^{1/2}}{|\xi|^{1/2}} \left(\tau(\xi)^2 + \frac{\|\mathbf{X}\|^{10\lambda}}{Z} \right).$$

We now combine this with (6.8) to conclude

$$N_\xi(\mathbf{X}) \ll \left((AB)^{1/4} C^{1/2} (C^{1/2} + (AB)^{1/4}) \right) (YW)^{3/4} Z^{1/2} \left(1 + \|\mathbf{X}\|^{10\lambda} \min \left(\frac{A^{1/2}}{Y}, \frac{1}{Z} \right) \right).$$

The proof of Proposition 6.1 is now completed by (6.11).

7. AN ASYMPTOTIC FORMULA

We now upgrade the previous upper bound to an asymptotic formula for $N_\xi(\mathbf{X})$. The main term features the singular series and the singular integral that would follow from a formal application of the circle method. With this in mind we define

$$(7.1) \quad \mathcal{E}_\xi = \sum_q \frac{1}{q^6} \sum_{d \pmod q}^* \sum_{a,b,c,y,w,z \pmod q} e \left(\frac{d(ab + c^2 + \xi^2 wyz^2)}{q} \right).$$

For later purposes we compute this as an Euler product. We have

$$\mathcal{E}_\xi = \sum_q \sum_{d \pmod q}^* \frac{1}{q^5} \sum_{c,y,w,z \pmod q} e \left(\frac{d(c^2 + \xi^2 wyz^2)}{q} \right) = \sum_q \frac{1}{q^4} \sum_{d \pmod q}^* \sum_{c \pmod q} \sum_{\xi^2 yz^2 \equiv 0 \pmod q} e \left(\frac{dc^2}{q} \right).$$

We have

$$\sum_{d \pmod q}^* \sum_{c \pmod q} e \left(\frac{dc^2}{q} \right) = \begin{cases} q^{1/2} \phi(q), & q = \square, \\ 0, & q \neq \square, \end{cases}$$

so we conclude

$$\mathcal{E}_\xi = \sum_q \frac{\phi(q^2)}{q^7} \sum_{\xi^2 yz^2 \equiv 0 \pmod{q^2}} 1 = \sum_q \frac{\phi(q)(\xi, q)^4}{q^6} \sum_{yz^2 \equiv 0 \pmod{(\frac{q}{(\xi, q)})^2}} 1.$$

If p is a prime and $n \geq 1$, then a simple combinatorial argument shows that the number of pairs $y, z \pmod{p^{2n}}$ with $p^{2n} \mid yz^2$ equals

$$p^{3n} + p^{3n-1} - p^{2n-1}.$$

For a prime p let $r_p = v_p(\xi)$ denote the p -adic valuation of ξ . Evaluating geometric sums, we finally conclude

$$\begin{aligned} \mathcal{E}_\xi &= \prod_p \left(1 + \sum_{n=1}^{r_p} \frac{(p-1)p^{n-1+4n}}{p^{6n}} + \sum_{n=r_p+1}^{\infty} \frac{(p-1)p^{n-1+4r_p}}{p^{6n}} (p^{3(n-r_p)} + p^{3(n-r_p)-1} - p^{2(n-r_p)-1}) \right) \\ &= \prod_p \left(1 + \frac{1+p+p^2-p^{2-r_p}}{p(1+p+p^2)} \right). \end{aligned}$$

In particular, the Euler product is absolutely convergent and satisfies

$$(7.2) \quad \mathcal{E}_\xi \ll \xi^\varepsilon.$$

We also define the singular integral for a tuple $\mathbf{X} = (X_{11}, X_{12}, X_{22}, X_{31}, X_{32}, X_{33})$ with $X_{ij} \geq 1$ as

$$\mathcal{I}_\xi(\mathbf{X}) = \int_{\mathbb{R}} \int_{\frac{1}{2}X_{ij} \leq |x_{ij}| \leq X_{ij}} e((x_{11}x_{12} + x_{21}^2 + \xi^2 x_{31}x_{32}x_{33}^2)\alpha) d\mathbf{x} d\alpha.$$

That the α -integral is absolutely convergent follows from the estimates

$$(7.3) \quad \begin{aligned} \int \int_{\substack{\frac{1}{2}A \leq |a| \leq A \\ \frac{1}{2}B \leq |b| \leq B}} e(ab\alpha) da db &\ll \min(AB, |\alpha|^{-1}) \ll \frac{\|\mathbf{X}\|^\varepsilon (AB)^{1/2}}{|\alpha|^{1/2}(|\alpha|^\varepsilon + |\alpha|^{-\varepsilon})}, \\ \int_{\frac{1}{2}C \leq |c| \leq C} e(c^2\alpha) dc &\ll \min(C, |C\alpha|^{-1}) \ll \frac{\|\mathbf{X}\|^\varepsilon C^{1/2}}{|\alpha|^{1/4}(|\alpha|^\varepsilon + |\alpha|^{-\varepsilon})}, \\ \int \int \int_{\substack{\frac{1}{2}Y \leq |y| \leq Y \\ \frac{1}{2}W \leq |w| \leq W \\ \frac{1}{2}Z \leq |z| \leq Z}} e(\xi^2 ywz^2\alpha) dy dw dz &\ll \min(YWZ, |\xi^2 Z\alpha|^{-1}) \ll \frac{\|\mathbf{X}\|^\varepsilon (YW)^{3/4} X^{1/2}}{|\alpha|^{1/4}(|\alpha|^\varepsilon + |\alpha|^{-\varepsilon})}. \end{aligned}$$

For future reference we state the similar bounds

$$(7.4) \quad \begin{aligned} \int \int_{\substack{\frac{1}{2}A \leq |a| \leq A \\ \frac{1}{2}B \leq |b| \leq B}} e(ab\alpha) da db - \int \int_{\mathbb{R}^2} V\left(\frac{a}{A}\right) V\left(\frac{b}{B}\right) e(ab\alpha) da db \\ &\ll \min(\Omega^{-1}AB, |\alpha|^{-1}) \ll \frac{\Omega^{-1/2} \|\mathbf{X}\|^\varepsilon (AB)^{1/2}}{|\alpha|^{1/2}(|\alpha|^\varepsilon + |\alpha|^{-\varepsilon})}, \\ \int_{\frac{1}{2}C \leq |c| \leq C} e(c^2\alpha) dc - \int_{\mathbb{R}} V\left(\frac{c}{C}\right) &\ll \min(\Omega^{-1}C, |C\alpha|^{-1}) \ll \frac{\Omega^{-3/4} \|\mathbf{X}\|^\varepsilon C^{1/2}}{|\alpha|^{1/4}(|\alpha|^\varepsilon + |\alpha|^{-\varepsilon})}, \\ \int \int \int_{\substack{\frac{1}{2}Y \leq |y| \leq Y \\ \frac{1}{2}W \leq |w| \leq W \\ \frac{1}{2}Z \leq |z| \leq Z}} e(\xi^2 ywz^2\alpha) dy dw dz - \int \int \int_{\mathbb{R}^3} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) V\left(\frac{z}{Z}\right) \\ &\ll \min(\Omega^{-1}YWZ, |\xi^2 Z\alpha|^{-1}) \ll \frac{\Omega^{-3/4} \|\mathbf{X}\|^\varepsilon (YW)^{3/4} Z^{1/2}}{|\alpha|^{1/4}(|\alpha|^\varepsilon + |\alpha|^{-\varepsilon})}. \end{aligned}$$

Our aim in this section is to prove the following asymptotic formula. Let

$$(7.5) \quad \zeta^{(1)} = (1/2, 1/4, 1/4), \quad \zeta^{(2)} = (3/4, 1/8, 1/8), \quad \zeta^{(3)} = (5/8, 1/8, 1/4).$$

Proposition 7.1. *The exists $\delta > 0$ with the following property. For $\mathbf{X} = (X_{ij}), \xi$ satisfying (6.1) and any $\varepsilon > 0$ we have*

$$N_\xi(\mathbf{X}) - \mathcal{E}_\xi \mathcal{I}_\xi(\mathbf{X}) \ll_\varepsilon (\min_{ij} X_{ij})^{-\delta} \|\mathbf{X}\|^\varepsilon \sum_{i=1}^3 \mathbf{X}^{(\zeta^{(i)})}.$$

The rest of this section is devoted to the proof. As a first step we estimate the effect of smoothing. To this end we recall the definition of $N_\xi(\mathbf{X}, \Delta)$ and $N_\xi^*(\mathbf{X}, \Delta)$: the former restricts one of the variables c, y, w, z to a small interval, the latter the variable a . By symmetry and (6.1), the bound in Proposition 6.2 holds also when b is restricted to a small interval. Combining Propositions 4.2 and 6.2, we obtain

$$(7.6) \quad N_\xi(\mathbf{X}) = N_\xi(\mathbf{X}, \Omega) + O\left((\|\mathbf{X}\|^\varepsilon \Omega^{-1/2} + \|\mathbf{X}\|^{-1/5}) \sum_{i=1}^3 \mathbf{X}^{(\zeta^{(i)})}\right).$$

An evaluation of $N_\xi(\mathbf{X}, \Omega)$ is given in Proposition 6.3, and we proceed to evaluate the main terms M_2 and \tilde{M}_3 defined in (6.4) and (6.12).

7.1. **Computation of M_2 .** Recall that

$$M_2 = \sum_{a,c,z} V\left(\frac{a}{A}\right) V\left(\frac{c}{C}\right) V\left(\frac{z}{Z}\right) \frac{1}{a^2} \sum_{\substack{a_1 a_2 a_3 = |a| \\ a_3 |c|^2}} a_1(\xi^2 z^2, a_2 a_3) a_3 \mu(a_2) \Phi_\xi(0, 0, a, c, z)$$

where

$$\Phi_\xi(0, 0, a, c, z) = \int_{\mathbb{R}^2} V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) V\left(\frac{-(c^2 + \xi^2 y w z^2)}{aB}\right) dy dw.$$

Let

$$F_\xi(a, c, z) = V\left(\frac{a}{A}\right) V\left(\frac{z}{Z}\right) V\left(\frac{c}{C}\right) \Phi_\xi(0, 0, a, c, z).$$

By symmetry we can restrict a, c, z to be positive at the cost of a factor 8. We denote by

$$\widehat{F}_\xi(s, u, v) = 8 \int_{[0, \infty)^3} F_\xi(a, c, z) a^{s-1} c^{u-1} z^{v-1} da dc dz$$

the Mellin transform of F_ξ . It is holomorphic in all three variables (since V has compact support), and by Lemma 6.4 and partial integration we have

$$\widehat{F}_\xi(s, u, v) \ll_N \frac{(AB)^{1/4} (YW)^{3/4}}{(Z|\xi|)^{1/2}} A^{\Re s} C^{\Re u} Z^{\Re v} \left(1 + \frac{|\Im s|}{S} + \frac{|\Im u|}{U} + \frac{|\Im v|}{V}\right)^{-N}$$

for all $N > 0$ where

$$S = \Omega\left(1 + \frac{|\xi|^2 YWZ^2}{AB}\right), \quad U = \Omega\left(1 + \frac{C^2}{AB}\right), \quad V = \Omega.$$

By (6.1) we have $S, U \ll \Omega\|\mathbf{X}\|^{10\lambda}$. Let us also define

$$L_\xi(s, u, v) = \sum_{a_1, a_2, a_3} \sum_z \sum_{a_3 | c^2} \frac{(\xi^2 z^2, a_2 a_3) \mu(a_2)}{a_1^{1+s} a_2^{2+s} a_3^{1+s} c^u z^v}, \quad \Re s > 0, \quad \Re u > 1, \quad \Re v > 1.$$

By Mellin inversion we have

$$(7.7) \quad M_2 = \int_{(2)} \int_{(2)} \int_{(2)} L_\xi(s, u, v) \widehat{F}_\xi(s, u, v) \frac{ds du dv}{(2\pi i)^3}.$$

By a long, cumbersome and uninspiring, but completely straightforward computation based on geometric series we can compute $L_\xi(s, u, v)$ as an Euler product. If $v_p(\xi) = r_p$, then

$$L_\xi(s, u, v) = \zeta(s+1) \zeta(u) \zeta(v) \zeta(2+2s+u) \zeta(2s+u+v) H_\xi(s, u, v)$$

where the p -Euler factor of $H_\xi(s, u, v)$ is given by

$$\begin{aligned} & \left(1 - p^{-3-6s-3u-v} - p^{-2-s-(2s+u)r_p} + p^{-1-s-(2s+u)r_p} + p^{-2-3s-2u-(2s+u)r_p} - p^{-1-3s-2u-(2s+u)r_p} \right. \\ & - p^{-3-3s-u-(2s+u)r_p} + p^{-2-3s-u-(2s+u)r_p} - p^{-3-2s-u-(2s+u)r_p} + p^{-2-2s-u-(2s+u)r_p} + p^{-1-2s-u-(2s+u)r_p} \\ & - p^{-2-s-u-(2s+u)r_p} + p^{-1-s-u-(2s+u)r_p} - p^{-s-u-(2s+u)r_p} + p^{-2-s-v-(2s+u)r_p} - p^{-1-s-v-(2s+u)r_p} \\ & - p^{-2-3s-2u-v-(2s+u)r_p} + p^{-1-3s-2u-v-(2s+u)r_p} + p^{-3-3s-u-v-(2s+u)r_p} - p^{-2-3s-u-v-(2s+u)r_p} \\ & - p^{-2-2s-u-v-(2s+u)r_p} - p^{-1-2s-u-v-(2s+u)r_p} + p^{-2-s-u-v-(2s+u)r_p} - p^{-1-s-u-v-(2s+u)r_p} + p^{-s-u-v-(2s+u)r_p} \\ & + p^{-3-2s-u-v-(2s+u)r_p} - p^{-2-3s-2u} + p^{-2-5s-3u-v} - p^{-3-5s-2u-v} - p^{-3s-2u-v} + p^{-3-4s-2u} + p^{-2-4s-2u-v} \\ & \left. + p^{-1-4s-2u-v} - p^{-2-2s-u} - p^{-1-2s-u} - p^{-2s-u-v} + p^{-3-3s-u} + p^{-1-3s-u-v} + p^{-s-u} - p^{-1-s} \right) \left(1 - \frac{1}{p^{2s+u}} \right)^{-1}. \end{aligned}$$

For $p \nmid \xi$ (i.e. $r_p = 0$) this simplifies considerably as

$$\begin{aligned} & 1 - p^{-2-s} - p^{-3-2s-u} + p^{-1-s-u} + p^{-2-s-v} - p^{-1-s-v} + p^{-3-4s-2u-v} - p^{-2-3s-2u-v} \\ & + p^{-3-3s-u-v} + p^{-3-2s-u-v} - p^{-2-2s-u-v} - p^{-1-2s-u-v} - p^{-1-s-u-v} + p^{-s-u-v}. \end{aligned}$$

In particular, we see that $L_\xi(s, u, v)$ is holomorphic in

$$(7.8) \quad \Re s \geq -1/5, \quad \Re v \geq 4/5, \quad \Re u \geq 4/5$$

except for polar divisors at $s = 0$, $u = 1$, $v = 1$; away from the polar divisors it is (crudely) bounded by $\tau(\xi)((1 + |s|)(1 + |u|)(1 + |v|))^{1/8}$ in this region. Another computation shows that

$$\operatorname{res}_{s=0} \operatorname{res}_{u=1} \operatorname{res}_{v=1} L_\xi(s, u, v) = \prod_p \left(1 + \frac{1 + p + p^2 - p^{2-r_p}}{p(1 + p + p^2)} \right) = \mathcal{E}_\xi.$$

Shifting contours to the left in (7.7) we conclude that

$$M_2 = \mathcal{E}_\xi \widehat{F}_\xi(0, 1, 1) + O\left(\Omega^3 |\xi|^\varepsilon (AB)^{1/4} C(YW)^{3/4} Z^{1/2} \left(\|\mathbf{X}\|^{30\lambda} (A^{-1/8} + C^{-1/8}) + Z^{-1/8} \right)\right).$$

Using again (6.1) and the now familiar device based on (4.5), we can write the error term as

$$(7.9) \quad \|\mathbf{X}\|^\varepsilon \Omega^3 (\mathbf{X}^{(\frac{5}{8}, \frac{1}{8}, \frac{1}{4})} + \mathbf{X}^{(\frac{3}{4}, \frac{1}{8}, \frac{1}{8})}) \min(A, C, Z)^{-1/10}.$$

By definition and using symmetry again to remove the factor 8, we have

$$\widehat{F}_\xi(0, 1, 1) = \int_{\mathbb{R}^5} V\left(\frac{a}{A}\right) V\left(\frac{z}{Z}\right) V\left(\frac{c}{C}\right) V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) V\left(\frac{-(c^2 + \xi^2 y w z^2)}{aB}\right) dy dw \frac{da}{|a|} dz dc$$

By Fourier inversion we have

$$\begin{aligned} V\left(\frac{-(c^2 + \xi^2 y w z^2)}{aB}\right) &= \int_{\mathbb{R}} \int_{\mathbb{R}} V(b) e(b\alpha) db e\left(\alpha \frac{c^2 + \xi^2 y w z^2}{aB}\right) d\alpha \\ &= |a| \int_{\mathbb{R}} \int_{\mathbb{R}} V\left(\frac{b}{B}\right) e(ba\alpha) db e(\alpha(c^2 + \xi^2 y w z^2)) d\alpha, \end{aligned}$$

so that $\widehat{F}_f(0, 1, 1)$ equals

$$\int_{\mathbb{R}} \int_{\mathbb{R}^6} V\left(\frac{a}{A}\right) V\left(\frac{z}{Z}\right) V\left(\frac{c}{C}\right) V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) V\left(\frac{b}{B}\right) e(\alpha(ab + c^2 + \xi^2 y w z^2)) db dy dw da dz dc d\alpha.$$

It remains to remove the smoothing and quantify the error from replacing V with the characteristic function on $[1/2, 1]$. By (7.3) and (7.4), we see that

$$(7.10) \quad \widehat{F}_f(0, 1, 1) - \mathcal{I}_\xi(\mathbf{X}) \ll \Omega^{-1/2} \mathbf{X}^{(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})} \|\mathbf{X}\|^\varepsilon.$$

Combining this with (7.9), (7.6), the first part of Proposition 6.3 and choosing $\Omega = \min(A, C, Z)^{1/50}$, we have shown

$$(7.11) \quad N_\xi(\mathbf{X}) - \mathcal{E}_\xi \mathcal{I}_\xi \ll \|\mathbf{X}\|^\varepsilon \sum_{i=1}^3 \mathbf{X}^{(\zeta^{(i)})} \left(\min(\mathbf{X}_{ij})^{-1/100} + \|\mathbf{X}\|^{11\lambda} \frac{A^{1/2+1/50}}{Y} \right).$$

7.2. Computation of \tilde{M}_3 . The argument for \tilde{M}_3 is similar. Recall from (6.12) that

$$\begin{aligned} \tilde{M}_3 &= 2 \sum_{\substack{\xi_1 \xi_2 = \xi \\ (\xi_2, fg) = 1}} \sum_{\substack{fg = \square \\ (g, y) = 1}} \sum_{\substack{wy \neq -\square \\ (g, y) = 1}} V\left(\frac{fy}{Y}\right) V\left(\frac{gw}{W}\right) \sum_{\substack{d_1, d_2 \\ (d_2, \xi_2 y w) = 1}} V\left(\frac{d_1 d_2^2 fg \xi_1^2}{A}\right) \frac{\phi(d_2^2 fg \xi_1^2)}{d_1 d_2^4 (fg)^{3/2} \xi_1^3} \\ &\quad \times \Psi(0, 0, \xi^2 f g y w, d_1 d_2^2 fg \xi_1^2) \end{aligned}$$

where Ψ satisfies the bounds of Lemma 6.5. Recall that all variables run over positive integers, except for y, w that run over positive and negative integers. We first add back the contribution $wy = -\square$ at the cost of an error

$$(7.12) \quad \ll \tau(\xi) \sqrt{YW} \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}}$$

by estimating trivially the contribution of all variables.

Since V is even, we can re-write \tilde{M}_3 resulting expression, up to the error (7.12), as

$$4 \sum_{\pm} \sum_{\xi_1 \xi_2 = \xi} \sum_{\substack{fg = \square \\ (\xi_2, fg) = 1}} \sum_{\substack{wy \neq -\square \\ (g, y) = 1}} V\left(\frac{fy}{Y}\right) V\left(\frac{gw}{W}\right) \sum_{\substack{d_1, d_2 \\ (d_2, \xi_2 yw) = 1}} V\left(\frac{d_1 d_2^2 fg \xi_1^2}{A}\right) \frac{\phi(d_2^2 fg \xi_1^2)}{d_1 d_2^4 (fg)^{3/2} \xi_1^3} \\ \times \Psi(0, 0, \pm \xi^2 fg yw, d_1 d_2^2 fg \xi_1^2)$$

where now all variables run over positive integers.

Let

$$G_\xi(a, y, w) = 4 \sum_{\pm} V\left(\frac{a}{A}\right) V\left(\frac{y}{Y}\right) V\left(\frac{w}{W}\right) \Psi(0, 0, \pm \xi^2 yw, a).$$

Then

$$(7.13) \quad \tilde{M}_3 = \sum_{\xi_1 \xi_2 = \xi} \sum_{\substack{fg = \square \\ (\xi_2, fg) = 1}} \sum_{(g, y) = 1} \sum_{\substack{d_1, d_2 \\ (d_2, \xi_2 yw) = 1}} \frac{\phi(d_2^2 fg \xi_1^2)}{d_1 d_2^4 (fg)^{3/2} \xi_1^3} G(fy, gw, d_1 d_2^2 fg \xi_1^2) \\ + O((CZAB)^{1/2} (YW)^{1/4}).$$

As before, we denote by $\hat{G}_\xi(s, u, v)$ the Mellin transform of $G_\xi(a, y, w)$; it is entire in all three variables and by Lemma 6.5 satisfies

$$\hat{G}_\xi(s, u, v) \ll_N \frac{\sqrt{CZAB}}{|\xi|^{1/2} (YW)^{1/4}} A^{\Re s - 1} Y^{\Re u - 1} W^{\Re v - 1} (1 + |s| + |u| + |v|)^{-N}.$$

Next we define

$$\tilde{L}_\xi(s, u, v) = \sum_{\xi_1 \xi_2 = \xi} \sum_{\substack{fg = \square \\ (\xi_2, fg) = 1}} \sum_{(g, y) = 1} \sum_{(d_2, \xi_2 yw) = 1} \frac{\phi(d_2^2 fg \xi_1^2)}{d_2^4 (fg)^{3/2} \xi_1^3 (d_2^2 fg \xi_1^2)^s (fy)^u (gw)^v}$$

which is absolutely convergent in $\Re u, \Re v > 1$, $\Re s > -1/2$. Then by Mellin inversion we have

$$\tilde{M}_3 = \int_{(2)} \int_{(2)} \int_{(2)} \tilde{L}_\xi(s, u, v) \zeta(s+1) \hat{G}_\xi(s, u, v) \frac{ds du dv}{(2\pi i)^3}.$$

As before, we analyze $\tilde{L}_\xi(s, u, v)$ by computing its Euler product expansion. A similarly long and cumbersome computation yields

$$\tilde{L}_\xi(s, u, v) = \zeta(u) \zeta(v) H_\xi(s, u, v)$$

where $H_\xi(s, u, v) \ll \tau(\xi)$ is holomorphic in the same region (7.8) and $H_\xi(0, 1, 1) = \mathcal{E}_\xi$. Shifting contours, we conclude as before

$$\tilde{M}_3 - \mathcal{E}_\xi \hat{G}_\xi(0, 1, 1) \ll \|\mathbf{X}\|^\varepsilon \Omega^3 (CZAB)^{1/2} (YW)^{3/4} \min(Y, W, A)^{-1/10}$$

which also contains the error term from (7.13). Unraveling the definitions and using symmetry to remove the factor 4 and the \pm -sign, we see that $\hat{G}_\xi(0, 1, 1) = \hat{F}_\xi(0, 1, 1)$, so that by (7.10) we get

$$\tilde{M}_3 - \mathcal{E}_\xi \mathcal{I}_\xi \ll \|\mathbf{X}\|^\varepsilon \mathbf{X}^{(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})} (\Omega^3 \min(Y, W, A)^{-1/10} + \Omega^{-1/2}).$$

Combining this with (7.6) and the second part of Proposition 6.3 and choosing $\Omega = \min(Y, W, A)^{1/50}$, we have shown

$$(7.14) \quad N_\xi(\mathbf{X}) - \mathcal{E}_\xi \mathcal{I}_\xi \ll \|\mathbf{X}\|^\varepsilon \sum_{i=1}^3 \mathbf{X}^{(\mathfrak{c}^{(i)})} \left(\min(\mathbf{X}_{ij})^{-1/100} + \|\mathbf{X}\|^{10\lambda} \frac{Y^{4/50}}{Z} \right).$$

It remains to combine (7.11) and (7.14). As in (6.11) we conclude

$$\min\left(\frac{A^{1/2+1/50}}{Y}, \frac{Y^{4/50}}{Z}\right) \leq \frac{(YAB)^\lambda A^{1/100} Y^{2/50}}{B^{1/4}} \ll \|\mathbf{X}\|^{-1/100-15\lambda}.$$

This completes the proof of Proposition 7.1.

8. INTRODUCING THE HEIGHT CONDITIONS

Let $P > 1$ be a large parameter. Let as before $\mathbf{X} = (A, B, C, Y, W, Z)$, where all entries are restricted to powers of 2. The condition that the entries of \mathbf{X} are powers of 2 will remain in force throughout this section. Let $\mathbf{x} = (a, b, c, y, w, z) \in \mathbb{N}^6$, $\mathbf{g} = (\eta, \xi) \in \mathbb{N}^2$. For given \mathbf{x} , \mathbf{g} let $\mathcal{X} = \mathcal{X}(P, \mathbf{g}, \mathbf{x})$ be the set of tuples \mathbf{X} satisfying

$$(8.1) \quad \max(aA, bB, cC)^2 \max(yY, wW) \leq \frac{P}{\eta^2 \xi}, \quad \max(yY, wW)^3 (zZ)^2 \leq \frac{P}{\eta^2 \xi^3}.$$

Fix some sufficiently small $\lambda > 0$. We call a pair $(\mathbf{X}, \mathbf{g} = (\eta, \xi))$ *bad* if (6.1) is violated (i.e., one of these inequalities does not hold), otherwise we call it *good*. We call it *very good* if the following stronger version of (6.1) holds:

$$(8.2) \quad \begin{aligned} \min(A, B, C) &\geq \max(A, B, C)(\log P)^{-100}, \\ \min(Y, W) &\geq \max(Y, W)(\log P)^{-100}, \\ \min(AB, C^2, YWZ^2) &\geq \max(AB, C^2, YWZ^2)(\log P)^{-100}, \\ |\xi| &\leq (\log P)^{100}, \\ P(\log P)^{-100} &\leq CYWZ \leq P. \end{aligned}$$

Clearly there are at most $O((\log P)^6)$ tuples \mathbf{X} (the entries being powers of 2) satisfying (8.1). Moreover, it is easy to see that there are at most

$$(8.3) \quad \ll \log H (\log \log P)^6$$

such tuples satisfying (8.1), (8.2) and

$$(8.4) \quad \min(A, B, C, Y, W, Z) \leq H.$$

Let $\mathcal{X}_{\text{bad}}(P, \mathbf{g}, \mathbf{x})$ be the set of $\mathbf{X} \in \mathcal{X}(P, \mathbf{g}, \mathbf{x})$ such that (\mathbf{X}, \mathbf{g}) is bad. Let $\mathcal{X}_H(P, \mathbf{g}, \mathbf{x})$ be the set of $\mathbf{X} \in \mathcal{X}(P, \mathbf{g}, \mathbf{x})$ such that (8.4) holds. Finally let

$$\mathcal{X}^*(P, \mathbf{g}, \mathbf{x}) = \mathcal{X}_{\text{bad}}(P, \mathbf{g}, \mathbf{x}) \cup \mathcal{X}_H(P, \mathbf{g}, \mathbf{x}).$$

We write

$$N(P, \mathbf{g}, \mathbf{x}) = \sum_{\mathbf{X} \in \mathcal{X}^*(P, \mathbf{g}, \mathbf{x})} N_{\xi}(\mathbf{X}).$$

Both $\mathcal{X}^*(P, \mathbf{g}, \mathbf{x})$ and $N(P, \mathbf{g}, \mathbf{x})$ depend on H , but this is not displayed in the notation. Our main result in this section is

Proposition 8.1. *For $1 \ll H \leq P$ and $0 < \lambda < 1$ we have*

$$(8.5) \quad N(P, \mathbf{g}, \mathbf{x}) \ll \frac{P(1 + \log H)(\log P)^{\varepsilon}}{(\xi^{3/2} \eta^2)^{99/100} (abcywz)^{1/4}}.$$

Proof. We first claim that

$$(8.6) \quad \sum_{\mathbf{X} \in \mathcal{X}_{\text{bad}}(P, \mathbf{g}, \mathbf{x})} N_{\xi}(\mathbf{X}) \ll P^{1-\lambda/100+2\varepsilon}.$$

Let $\delta = \lambda/100$ and suppose that

$$(8.7) \quad N_{\xi}(\mathbf{X}) \geq P^{1-\delta+\varepsilon}.$$

We show that this implies that (6.1) holds, so \mathbf{X} is good, and since there are only $O(P^{\varepsilon})$ tuples \mathbf{X} , this implies the claim.

From Proposition 4.1 and (8.1) we have

$$\begin{aligned} N_{\xi}(\mathbf{X}) &\ll P^{\varepsilon} (ABC)^{1/2} (YW)^{3/4} Z^{1/2} \\ &\leq P^{\varepsilon} \max(A, B, C)^{3/2} \max(Y, W)^{3/4} \cdot \max(Y, W)^{3/4} Z^{1/2} \ll \frac{P^{1+\varepsilon}}{\eta^2 \xi^{3/2}}. \end{aligned}$$

Contemplating this sequence of inequalities, we conclude from (8.7) that

$$(8.8) \quad \begin{aligned} \xi &\ll P^{5\delta}, \quad \min(Y, W) \gg \max(Y, W)P^{-5\delta}, \\ \min(A, B, C) &\gg \max(A, B, C)P^{-5\delta}, \\ C^2Y &\gg P^{1-5\delta}, \quad Y^3Z^2 \gg P^{1-5\delta}, \end{aligned}$$

which also implies that all of the three blocks AB , C^2 , YWZ^2 must be within $B^{20\delta}$. This shows our claim \mathbf{X} must satisfy (6.1) with $\lambda = \delta/100$. This establishes (8.6). We complement this with a second bound. From Proposition 4.1 and (8.1) we have

$$N_\xi(\mathbf{X}) \ll P^\varepsilon (ABC)^{1/2} Y^{3/4} \cdot W^{3/4} Z^{1/2} \ll P^\varepsilon \frac{P}{\xi^{3/2} \eta^2 (abcz)^{1/2} (yw)^{3/4}}.$$

Combining this with (8.6), we conclude

$$\sum_{\mathbf{X} \in \mathcal{X}_{\text{bad}}(P, \mathbf{g}, \mathbf{x})} N_\xi(\mathbf{X}) \ll P \min \left(P^{-\lambda/200}, \frac{P^\varepsilon}{\xi^{3/2} \eta^2 (abcyz)^{1/2}} \right) \ll \frac{P}{(\xi^{3/2} \eta^2 (abcyz)^{1/2})^{99/100}}.$$

This is acceptable for (8.5).

For the contribution of $\mathbf{X} \in \mathcal{X}_H(P, \mathbf{g}, \mathbf{x}) \setminus \mathcal{X}_{\text{bad}}(P, \mathbf{g}, \mathbf{x})$ we observe that Proposition 6.1 is available. We note that

$$\mathbf{X}^{(\frac{3}{4}, 0, \frac{1}{4})} + \mathbf{X}^{(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})} = (AB)^{1/4} C(YW)^{3/4} Z^{1/2} + (ABC)^{1/2} (YW)^{3/4} Z^{1/2} \ll \frac{P}{\eta^2 \xi^{3/2} (abcyz)^{1/4}}$$

upon using (8.1). By (8.3), this is acceptable for the very good tuples. For tuples that are good, but not very good, the previous inequality along with the same argument as leading to (8.8) shows

$$N_\xi(\mathbf{X}) \ll \frac{P(\log P)^{-10}}{\eta^2 \xi^{3/2} (abcyz)^{1/4}}$$

for such \mathbf{X} . Since there are at most $O((\log P)^6)$ such tuples, the proof is complete. \square

Part 3. Proof of Theorem 1.1

9. GEOMETRY

The following Table 2 contains the nine smooth spherical Fano threefolds over $\overline{\mathbb{Q}}$ that are not horospherical (since the horospherical smooth Fano threefolds are all either toric or flag varieties; see [Ho, §6.3]). The notation T and N in [Ho, Table 6.5] and in our Table 2 refers to the cases described at the beginning of the introduction (Section 1) and in [BBDG, §10.2].

rk Pic	Hofscheier	Mori–Mukai	torsor equation	remark
2	$T_1 12$	II.31	$x_{11}x_{12} - x_{21}x_{22} - x_{31}x_{32}^2$	eq. \mathbb{G}_a^3 -cpct.
2	$N_1 6, N_1 7$	II.30	$x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}$	eq. \mathbb{G}_a^3 -cpct.
2	$N_1 8$	II.29	$x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}x_{33}^2$	variety X_1
3	$T_1 18$	III.24	$x_{11}x_{12} - x_{21}x_{22} - x_{31}x_{32}$	[BBDG]
3	$T_1 21$	III.20	$x_{11}x_{12} - x_{21}x_{22} - x_{31}x_{32}x_{33}^2$	[BBDG]
3	$N_0 3$	III.22	$x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}$	variety X_2
3	$N_1 9$	III.19	$x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}$	variety X_3
4	$T_0 3$	IV.8	$x_{11}x_{12} - x_{21}x_{22} - x_{31}x_{32}$	[BBDG]
4	$T_1 22$	IV.7	$x_{11}x_{12} - x_{21}x_{22} - x_{31}x_{32}$	[BBDG]

TABLE 2. Smooth Fano threefolds that are spherical, but not horospherical

We proceed to describe the three N cases X_1, X_2, X_3 in Table 2 that are not equivariant \mathbb{G}_a^3 -compactifications [HM] in more detail. From the description in the Mori–Mukai classification, we

can construct a split form over \mathbb{Q} in each case. We then recall from Hofscheier’s list the description using the Luna–Vust theory of spherical embeddings.

The three varieties will be equipped with an action of $G = \mathrm{SL}_2 \times \mathbb{G}_m$. Let $\varepsilon_1 \in \mathfrak{X}(B)$ always be a primitive character of \mathbb{G}_m composed with the natural inclusion $\mathfrak{X}(\mathbb{G}_m) \rightarrow \mathfrak{X}(B)$.


9.1. X_1 of type II.29. Consider $\mathbb{P}_{\mathbb{Q}}^4$ with coordinates $(z_{11} : z_{12} : z_{21} : z_{31} : z_{32})$ and the hypersurface $Q = \mathbb{V}(z_{11}z_{12} - z_{21}^2 - z_{31}z_{32}) \subset \mathbb{P}_{\mathbb{Q}}^4$. It contains the conic

$$C_{33} = \mathbb{V}(z_{31}, z_{32}).$$

Let X_1 be the blow-up of Q in C_{33} . This is a smooth Fano threefold of type II.29. We may define an action of $G = \mathrm{SL}_2 \times \mathbb{G}_m$ on Q by

$$(A, t) \cdot \left(\begin{pmatrix} z_{11} & z_{21} \\ z_{21} & z_{12} \end{pmatrix}, z_{31}, z_{32} \right) = \left(A \cdot \begin{pmatrix} z_{11} & z_{21} \\ z_{21} & z_{12} \end{pmatrix} \cdot A^{\top}, t \cdot z_{31}, t^{-1} \cdot z_{32} \right),$$

which turns Q into a spherical variety. The following description using the Luna–Vust theory of spherical embeddings can be easily verified. The lattice \mathcal{M} has basis $(\alpha + \varepsilon_1, \alpha - \varepsilon_1)$. We denote the corresponding dual basis of the lattice \mathcal{N} by (d_1, d_2) . Then there is one color with valuation $d = d_1 + d_2$, and the valuation cone is given by $\mathcal{V} = \{v \in \mathcal{N}_{\mathbb{Q}} : \langle v, \alpha \rangle \leq 0\}$. Since C_{32} is G -invariant, the variety X_1 is a spherical G -variety and the blow-up morphism $X_1 \rightarrow Q$ can be described by a map of colored fans. The following figure illustrates this.




Here, $u_{31} = -d_1$ and $u_{32} = -d_2$ are the valuations of the G -invariant prime divisors $\mathbb{V}(z_{31})$ and $\mathbb{V}(z_{32})$, respectively, and $u_{33} = -d$ is the valuation of the exceptional divisor E_{33} over C_{33} .

We obtain a smooth projective ambient toric variety $Y_1^\circ = Y_1 = Y_1''$. From the description of Σ_{\max} in [BBDG, §10.3], we deduce that Y_1 is smooth and that $-K_{X_1}$ is ample on Y_1 . Hence assumption [BBDG, (2.3)] holds.

9.2. X_2 of type III.22. Let $W = \mathbb{P}_{\mathbb{Q}}^1 \times \mathbb{P}_{\mathbb{Q}}^2$ with coordinates $(z_{01} : z_{02})$ and $(z_{11} : z_{12} : z_{21})$. Let C_{32} be the curve $\mathbb{V}(z_{02}, z_{11}z_{12} - z_{21}^2)$ on W . Let X_2 be the blow-up of W in C_{32} . This is a smooth Fano threefold of type III.22. We may define an action of $G = \mathrm{SL}_2 \times \mathbb{G}_m$ on W by

$$(A, t) \cdot \left(z_{01}, z_{02}, \begin{pmatrix} z_{11} & z_{21} \\ z_{21} & z_{12} \end{pmatrix} \right) = \left(t \cdot z_{01}, z_{02}, A \cdot \begin{pmatrix} z_{11} & z_{21} \\ z_{21} & z_{12} \end{pmatrix} \cdot A^{\top} \right),$$

which turns W into a spherical variety with the following Luna–Vust description. The lattice \mathcal{M} has basis $(2\alpha, \varepsilon)$. We denote the corresponding dual basis of the lattice \mathcal{N} by (d, ε^*) . Then there is one color with valuation $2d = \frac{1}{2}\alpha^\vee$, and the valuation cone is given by $\mathcal{V} = \{v \in \mathcal{N}_{\mathbb{Q}} : \langle v, \alpha \rangle \leq 0\}$. Since the curve C_{32} is G -invariant, the variety X_2 is a spherical G -variety, and the blow-up morphism $X_2 \rightarrow W$ can be described a map of colored fans. The right-hand arrow in the following figure illustrates this.



Here, $u_{01} = -\varepsilon^*$, $u_{02} = \varepsilon^*$, $u_{31} = -d$ are the valuations of $\mathbb{V}(z_{01})$, $\mathbb{V}(z_{02})$, $\mathbb{V}(z_{11}z_{12} - z_{21}^2)$, respectively, and $u_{32} = -d + \varepsilon^*$ is the valuation of the exceptional divisor E_{32} over C_{32} .


The dotted circles in the colored fan of X_2 specify the standard small completion Y_2 of the ambient toric variety Y_2° . We note that Y_2 is singular and that it is not possible to obtain a smooth small completion with the construction from [BBDG, §10.3] in this case. We may, however, construct a resolution of singularities $Y_2'' \rightarrow Y_2$ which does not affect X_2 . This is illustrated by the left-hand arrow in the figure above.

9.3. X_3 of type III.19. Consider $\mathbb{P}_{\mathbb{Q}}^4$ with coordinates $(z_{11} : z_{12} : z_{21} : z_{31} : z_{32})$ and the hypersurface $Q = \mathbb{V}(z_{11}z_{12} - z_{21}^2 - z_{31}z_{32}) \subset \mathbb{P}_{\mathbb{Q}}^4$. It contains the points

$$P_{01} = \mathbb{V}(z_{11}, z_{12}, z_{21}, z_{31}),$$

$$P_{02} = \mathbb{V}(z_{11}, z_{12}, z_{21}, z_{32}).$$

Let X_3 be the blow-up of Q in P_{01} and P_{02} . This is a smooth Fano threefold of type III.19. Since P_{01} and P_{02} are G -invariant, the variety X_3 is a spherical G -variety and the blow-up morphism $X_3 \rightarrow Q$ can be described by a map of colored fans. The right-hand arrow in the following figure illustrates this.



Here, $u_{31} = -d_1$ and $u_{32} = -d_2$ are the valuations of $\mathbb{V}(z_{31})$ and $\mathbb{V}(z_{32})$, respectively, and u_{01} and u_{02} are the valuations of the exceptional divisors E_{01} over P_{01} and E_{02} over P_{02} , respectively.

The dotted circles in the colored fan of X_3 specify the singular standard small completion Y_3 of the ambient toric variety Y_3° . Again, it is not possible to obtain a smooth small completion with the construction from [BBDG, §10.3]. The left-hand arrow in the figure above describes a resolution of singularities $Y_3'' \rightarrow Y_3$ which does not affect X_3 .

10. COX RINGS AND TORSORS

10.1. Type II.29. A Cox ring for X_1 is given by

$$\mathcal{R}(X_1) = \mathbb{Q}[x_{11}, x_{12}, x_{21}, x_{31}, x_{32}, x_{33}] / (x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}x_{33}^2)$$

with $\text{Pic } X \cong \mathbb{Z}^2$ where

$$\begin{aligned} \deg(x_{11}) &= \deg(x_{12}) = \deg(x_{22}) = (1, 0), \\ \deg(x_{31}) &= \deg(x_{32}) = (0, 1), \quad \deg(x_{33}) = (1, -1). \end{aligned}$$

The anticanonical class is $-K_X = (2, 1)$. A universal torsor over X_1 is

$$\mathcal{T}_1 = \text{Spec } \mathcal{R}(X_1) \setminus Z_{X_1},$$

where

$$Z_{X_1} = \mathbb{V}(x_{11}, x_{12}, x_{21}, x_{33}) \cup \mathbb{V}(x_{31}, x_{32}).$$

10.2. **Type III.22.** A Cox ring for X_2 is given by

$$\mathcal{R}(X_2) = \mathbb{Q}[x_{01}, x_{02}, x_{11}, x_{12}, x_{21}, x_{31}, x_{32}] / (x_{11}x_{12} - x_{21}^2 - x_{31}x_{32})$$

with $\text{Pic } X_2 \cong \mathbb{Z}^3$ where

$$\begin{aligned} \deg(x_{01}) &= (1, 0, 0), & \deg(x_{02}) &= (1, 0, -1), \\ \deg(x_{11}) &= \deg(x_{12}) = \deg(x_{21}) = (0, 1, 0), \\ \deg(x_{31}) &= (0, 2, -1), & \deg(x_{32}) &= (0, 0, 1). \end{aligned}$$

The anticanonical class is $-K_{X_2} = (2, 3, -1)$. A universal torsor over X_2 is

$$\mathcal{T}_2 = \text{Spec } \mathcal{R}(X_2) \setminus Z_{X_2},$$

where

$$Z_{X_2} = \mathbb{V}(x_{11}, x_{12}, x_{21}) \cup \mathbb{V}(x_{01}, x_{32}) \cup \mathbb{V}(x_{02}, x_{31}) \cup \mathbb{V}(x_{01}, x_{02}).$$

Let Y_2° be the ambient toric variety of X_2 ; its rays are

$$(0, 0, 1, -1), (0, 0, -1, 1), (-1, -1, -2, 0), (1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1),$$

corresponding to x_{01}, \dots, x_{32} , respectively, and each of its nine maximal cones is generated by the four rays corresponding to two of x_{11}, x_{12}, x_{21} and one of the pairs x_{02}, x_{32} or x_{31}, x_{32} or x_{01}, x_{31} from the maximal cones of the spherical fan. We have

$$Z_{Y_2^\circ} = \mathbb{V}(x_{11}, x_{12}, x_{21}) \cup \mathbb{V}(x_{01}, x_{32}) \cup \mathbb{V}(x_{02}, x_{31}) \cup \mathbb{V}(x_{01}, x_{02}).$$

Let Y_2 be the standard small completion of Y_2° . It has the same rays as Y_2° , and the two additional maximal cones generated by the four rays corresponding to x_{11}, x_{12}, x_{21} and x_{01} or x_{02} (see the spherical fan), which are both singular; the singular locus of Y_2° corresponds to their intersection, the cone generated by $(-1, -1, -2, 0)$, $(1, 0, 0, 0)$, $(0, 1, 0, 0)$. We have

$$Z_{Y_2} = \mathbb{V}(x_{11}, x_{12}, x_{21}, x_{31}) \cup \mathbb{V}(x_{11}, x_{12}, x_{21}, x_{32}) \cup \mathbb{V}(x_{01}, x_{02}) \cup \mathbb{V}(x_{01}, x_{32}) \cup \mathbb{V}(x_{02}, x_{31}).$$

A toric desingularization $\rho : Y_2'' \rightarrow Y_2$ is obtained by adding the ray $(0, 0, -1, 0)$ (the primitive multiple of the sum of the rays of the singular cone). Its Cox ring is

$$\mathcal{R}(Y_2'') = \mathbb{Q}[x_{01}, \dots, x_{32}, z_1]$$

with

$$\begin{aligned} \deg(x_{01}) &= (0, 0, 1, 1), & \deg(x_{02}) &= (0, 0, 0, 1), \\ \deg(x_{11}) &= \deg(x_{12}) = \deg(x_{21}) = (0, 1, 0, 0), \\ \deg(x_{31}) &= (1, 2, 0, 0), & \deg(x_{32}) &= (0, 0, 1, 0), & \deg(z_1) &= (1, 0, 1, 0) \end{aligned}$$

and the irrelevant ideal

$$Z_{Y_2''} = \mathbb{V}(x_{01}, x_{02}) \cup \mathbb{V}(x_{01}, x_{32}) \cup \mathbb{V}(x_{02}, x_{31}) \cup \mathbb{V}(x_{31}, z_1) \cup \mathbb{V}(x_{32}, z_1) \cup \mathbb{V}(x_{11}, x_{12}, x_{21}).$$

Now let $Y_2' = Y_2''$. Then $X_2 \subset Y_2'$ is given in Cox coordinates by the homogeneous equation

$$x_{11}x_{12}z_1 - x_{21}^2z_1 - x_{31}x_{32} = 0,$$

and we have $\rho^*(-K_{X_2}) = (\frac{3}{2}, 3, \frac{5}{2}, 2)$.

10.3. **Type III.19.** A Cox ring for X_3 is given by

$$\mathcal{R}(X_3) = \mathbb{Q}[x_{01}, x_{02}, x_{11}, x_{12}, x_{21}, x_{31}, x_{32}] / (x_{11}x_{12} - x_{21}^2 - x_{31}x_{32})$$

with $\text{Pic } X_3 \cong \mathbb{Z}^3$ where

$$\begin{aligned} \deg(x_{01}) &= (0, 1, 0), & \deg(x_{02}) &= (0, 0, 1), \\ \deg(x_{11}) &= \deg(x_{12}) = \deg(x_{21}) = (1, 0, 0), \\ \deg(x_{31}) &= (1, -1, 1), & \deg(x_{32}) &= (1, 1, -1). \end{aligned}$$

The anticanonical class is $-K_{X_3} = (3, 1, 1)$. A universal torsor over X_3 is

$$\mathcal{T}_3 = \text{Spec } \mathcal{R}(X_3) \setminus Z_{X_3},$$

where

$$Z_{X_3} = \mathbb{V}(x_{11}, x_{12}, x_{21}) \cup \mathbb{V}(x_{01}, x_{02}) \cup \mathbb{V}(x_{01}, x_{32}) \cup \mathbb{V}(x_{02}, x_{31}).$$

Its ambient toric variety Y_3° has the rays

$$(1, -1, 0, 0), (-1, 1, 0, 0), (-1, -1, -1, -1), (0, 0, 0, 1), (0, 0, 1, 0), (1, 0, 0, 0), (0, 1, 0, 0);$$

we have

$$Z_{Y_3^\circ} = \mathbb{V}(x_{11}, x_{12}, x_{21}) \cup \mathbb{V}(x_{01}, x_{02}) \cup \mathbb{V}(x_{01}, x_{32}) \cup \mathbb{V}(x_{02}, x_{31}).$$

Its standard small completion Y_3 has two singular maximal cones corresponding to $x_{0i}, x_{11}, x_{12}, x_{21}$, but their intersection is smooth, hence Y_3 has two isolated singularities; we have

$$Z_{Y_3} = \mathbb{V}(x_{11}, x_{12}, x_{21}, x_{31}) \cup \mathbb{V}(x_{11}, x_{12}, x_{21}, x_{32}) \cup \mathbb{V}(x_{01}, x_{02}) \cup \mathbb{V}(x_{01}, x_{32}) \cup \mathbb{V}(x_{02}, x_{31}).$$

Blowing up the singularities adds the rays $(-1, 0, 0, 0)$, $(0, -1, 0, 0)$ (half of the sum of the corresponding four rays), giving Y_3'' . The Cox ring of Y_3'' has two extra generators z_1, z_2 , with degrees

$$\begin{aligned} \deg(x_{01}) &= (0, 0, 0, 0, 1), & \deg(x_{02}) &= (0, 0, 0, 1, 0), \\ \deg(x_{11}) &= \deg(x_{12}) = \deg(x_{21}) = (0, 0, 1, 0, 0), \\ \deg(x_{31}) &= (0, 1, 1, 1, 0), & \deg(x_{32}) &= (1, 0, 1, 0, 1), \\ \deg(z_1) &= (0, 1, 0, 0, 1), & \deg(z_2) &= (1, 0, 0, 1, 0). \end{aligned}$$

We have

$$\begin{aligned} Z_{Y_3''} &= \mathbb{V}(z_2, z_1) \cup \mathbb{V}(z_2, x_{32}) \cup \mathbb{V}(z_1, x_{32}) \cup \mathbb{V}(z_2, x_{31}) \cup \mathbb{V}(z_1, x_{31}) \cup \mathbb{V}(x_{32}, x_{21}, x_{12}, x_{11}) \\ &\cup \mathbb{V}(x_{31}, x_{21}, x_{12}, x_{11}) \cup \mathbb{V}(z_2, x_{02}) \cup \mathbb{V}(x_{31}, x_{02}) \cup \mathbb{V}(x_{21}, x_{12}, x_{11}, x_{02}) \\ &\cup \mathbb{V}(z_1, x_{01}) \cup \mathbb{V}(x_{32}, x_{01}) \cup \mathbb{V}(x_{21}, x_{12}, x_{11}, x_{01}) \cup \mathbb{V}(x_{02}, x_{01}). \end{aligned}$$

Let $Y_3' = Y_3''$. Then the equation for $X_3 \subset Y_3'$ is

$$x_{11}x_{12}z_1z_2 - x_{21}^2z_1z_2 - x_{31}x_{32} = 0,$$

and we have $\rho^*(-K_{X_3}) = (2, 2, 3, 3, 3)$.

11. COUNTING PROBLEMS

Applying part one of this paper, we obtain the following counting problems, in which T_j is always the subset of $X_j(\mathbb{Q})$ where all Cox coordinates are nonzero and, in case of X_1 , where $x_{31}x_{33} \neq -\square$. For simplicity, we write $N_j(B)$ for $N_{X_j(\mathbb{Q}) \setminus T_j, H_j}(B)$ as in the introduction, and we write $\{x, y\}$ to mean x or y .

Corollary 11.1. (a) *We have*

$$N_1(B) = \frac{1}{4} \# \left\{ \mathbf{x} \in \mathbb{Z}_{\neq 0}^6 : \begin{aligned} &x_{11}x_{12} - x_{21}^2 - x_{31}x_{32}x_{33}^3 = 0, \quad \max |\mathcal{P}_1(\mathbf{x})| \leq B \\ &\gcd(x_{11}, x_{12}, x_{21}, x_{33}) = \gcd(x_{31}, x_{32}) = 1, \quad x_{31}x_{32} \neq -\square \end{aligned} \right\},$$

with

$$\mathcal{P}_1(\mathbf{x}) = \{ \{x_{11}, x_{12}, x_{21}\}^2 \{x_{31}, x_{32}\}, \{x_{31}, x_{32}\}^3 x_{33}^2 \}.$$

(b) We have

$$N_2(B) = \frac{1}{8} \# \left\{ \mathbf{x} \in \mathbb{Z}_{\neq 0}^7 : \begin{array}{l} x_{11}x_{12} - x_{21}^2 - x_{31}x_{32} = 0, \quad \max |\mathcal{P}_2(\mathbf{x})| \leq B \\ \gcd(x_{11}, x_{12}, x_{21}) = \gcd(x_{01}, x_{32}) = \gcd(x_{02}, x_{31}) = \gcd(x_{01}, x_{02}) = 1 \end{array} \right\},$$

with

$$\mathcal{P}_2(\mathbf{x}) = \left\{ x_{01}^2 \{x_{11}, x_{12}, x_{21}\} x_{31}, x_{02}^2 \{x_{11}, x_{12}, x_{21}\}^3 x_{32}, \right. \\ \left. x_{01}x_{02} \{x_{11}, x_{12}, x_{21}\}^3, x_{02}^2 x_{31}^{3/2} x_{32}^{5/2}, x_{01}^2 x_{31}^{3/2} x_{32}^{1/2} \right\}.$$

(c) We have

$$N_3(B) = \frac{1}{8} \# \left\{ \mathbf{x} \in \mathbb{Z}_{\neq 0}^7 : \begin{array}{l} x_{11}x_{12} - x_{21}^2 - x_{31}x_{32} = 0, \quad \max |\mathcal{P}_3(\mathbf{x})| \leq B \\ \gcd(x_{11}, x_{12}, x_{21}) = \gcd(x_{01}, x_{32}) = \gcd(x_{02}, x_{31}) = \gcd(x_{01}, x_{02}) = 1 \end{array} \right\},$$

with

$$\mathcal{P}_3(\mathbf{x}) = \left\{ x_{01}^2 \{x_{11}, x_{12}, x_{21}\}^2 x_{31}, x_{02}^2 \{x_{11}, x_{12}, x_{21}\}^2 x_{32}, \right. \\ \left. x_{01}x_{02} \{x_{11}, x_{12}, x_{21}\}^3, x_{01}^2 x_{31}^2 x_{32}, x_{02}^2 x_{31} x_{32}^2 \right\}.$$

Proof. For X_1 , we argue as in [BBDG] since the ambient toric variety Y_1 is regular.

For X_2 , we apply Proposition 2.11 and obtain the counting problem

$$N_2(B) = \frac{1}{16} \# \left\{ \begin{array}{l} x_{11}x_{12}z_1 - x_{21}^2z_1 - x_{31}x_{32} = 0, \quad \max |\mathcal{P}'_2(\mathbf{x}, z_1)| \leq B \\ (\mathbf{x}, z_1) \in \mathbb{Z}_{\neq 0}^8 : \gcd(x_{11}, x_{12}, x_{21}) = \gcd(x_{01}, x_{32}) = \gcd(x_{02}, x_{31}) = 1 \\ \gcd(x_{01}, x_{02}) = \gcd(x_{31}, z_1) = \gcd(x_{32}, z_1) = 1 \end{array} \right\}$$

with

$$\mathcal{P}'_2(\mathbf{x}, z_1) = \left\{ x_{01}^2 \{x_{11}, x_{12}, x_{21}\} x_{31} z_1^{1/2}, x_{02}^2 \{x_{11}, x_{12}, x_{21}\}^3 x_{32} z_1^{3/2}, x_{01}x_{02} \{x_{11}, x_{12}, x_{21}\}^3 z_1^{3/2}, \right. \\ \left. x_{02}^2 x_{31}^{3/2} x_{32}^{5/2}, x_{01}^2 x_{31}^{3/2} x_{32}^{1/2} \right\}.$$

But the equation together with $\gcd(x_{31}x_{32}, z_1) = 1$ implies $z_1 = \pm 1$. After multiplying the equation with z_1 , the substitution of $(x_{11}z_1, x_{12}z_1, x_{21}z_1, x_{31}z_1)$ by $(x_{11}, x_{12}, x_{21}, x_{31})$ lead to our counting problem.

For X_3 , we similarly obtain

$$N_3(B) = \frac{1}{16} \# \left\{ (\mathbf{x}, z_1, z_2) \in \mathbb{Z}_{\neq 0}^9 : \begin{array}{l} x_{11}x_{12}z_1z_2 - x_{21}^2z_1z_2 - x_{31}x_{32} = 0, \quad \max |\mathcal{P}'_3(\mathbf{x}, z_1)| \leq B \\ \gcd(z_2, z_1) = \gcd(z_2, x_{32}) = \gcd(z_1, x_{32}) = \gcd(z_2, x_{31}) = 1 \\ \gcd(z_1, x_{31}) = \gcd(x_{32}, x_{21}, x_{12}, x_{11}) = 1 \\ \gcd(x_{31}, x_{21}, x_{12}, x_{11}) = \gcd(z_2, x_{02}) = \gcd(x_{31}, x_{02}) = 1 \\ \gcd(x_{21}, x_{12}, x_{11}, x_{02}) = \gcd(z_1, x_{01}) = \gcd(x_{32}, x_{01}) = 1 \\ \gcd(x_{21}, x_{12}, x_{11}, x_{01}) = \gcd(x_{02}, x_{01}) = 1 \end{array} \right\},$$

The height condition is given by the monomials

$$\mathcal{P}'_3(\mathbf{x}, z_1) = \left\{ x_{01}^2 \{x_{11}, x_{12}, x_{21}\}^2 x_{31} z_1 z_2^2, x_{02}^2 \{x_{11}, x_{12}, x_{21}\}^2 x_{32} z_1^2 z_2, \right. \\ \left. x_{01}x_{02} \{x_{11}, x_{12}, x_{21}\}^3 z_1^2 z_2^2, x_{02}^2 x_{31} x_{32}^2 z_1, x_{01}^2 x_{31}^2 x_{32} z_2 \right. \\ \left. x_{01}^2 x_{31}^2 x_{32} z_2, x_{02}^2 x_{31} x_{32}^2 z_1 \right\}.$$

In this counting problem, we observe that this equation together with $\gcd(z_1z_2, x_{31}x_{32}) = 1$ implies $z_1 = \pm 1$ and $z_2 = \pm 1$. The torsor equation also allows us to simplify the coprimality conditions. \square

Remark 11.2. The varieties X_1 , X_2 and X_3 are as in Remark 2.1 and in each case we have chosen L as in the proof of Lemma 2.3. After having eliminated the additional variables in the proof of Corollary 11.1, we have obtained the same monomials $\mathcal{P}_1(\mathbf{x})$, $\mathcal{P}_2(\mathbf{x})$, $\mathcal{P}_3(\mathbf{x})$ as if we had directly

applied [BBDG] (disregarding that Y is singular). In this case, [BBDG, Lemmas 3.8, 3.9, and 3.10] still apply, with the difference that the vertices of the polytopes are not necessarily integral.

12. APPLICATION: PROOF OF THEOREM 1.1

12.1. The analytic machinery. Theorems 8.4, 9.2 and 10.1 in [BBDG] provide an asymptotic formula for counting problems as in Corollary 11.1 under various assumptions and show in addition that the shape of the asymptotic formula agrees with the Manin-Peyre prediction. We need a small variation of these results that we state in full detail for the reader's convenience.

Suppose that we are given a diophantine equation

$$(12.1) \quad \sum_{i=1}^k \prod_{j=1}^{J_i} x_{ij}^{h_{ij}} = 0$$

with certain $h_{ij} \in \mathbb{N}$ and height conditions

$$(12.2) \quad \prod_{i=0}^k \prod_{j=1}^{J_i} |x_{ij}|^{\alpha_{ij}^\nu} \leq B \quad (1 \leq \nu \leq N)$$

for certain nonnegative exponents α_{ij}^ν , whose variables are restricted by coprimality conditions

$$(12.3) \quad \gcd\{x_{ij} : (i, j) \in S_\rho\} = 1 \quad (1 \leq \rho \leq r)$$

for certain $S_\rho \subseteq \{(i, j) : i = 0, \dots, k, j = 1, \dots, J_i\}$, cf. [BBDG, (1.2) – (1.4)]. We write $J = J_0 + \dots + J_k$. Let $N(B)$ the number of integral solutions to (12.1) with nonzero variables \mathbf{x}_j subject to (12.2) and (12.3).

With these data, we define the following quantities. For $\mathbf{g} \in \mathbb{N}^r$ write $\boldsymbol{\gamma} = (\gamma_{ij}) \in \mathbb{N}^J$, $\gamma_{ij} = \text{lcm}\{g_\rho : (i, j) \in S_\rho\}$ and

$$\boldsymbol{\gamma}^* = \left(\prod_{j=1}^{J_i} \gamma_{ij}^{h_{ij}} \right)_{1 \leq i \leq k} \in \mathbb{N}^k$$

as in [BBDG, (8.11), (8.14)].

As in [BBDG, (5.1) – (5.6)], for $\mathbf{b} \in \mathbb{N}^k$ define the (formal) singular series

$$\mathcal{E}_{\mathbf{b}} = \sum_{q=1}^{\infty} \sum_{a \bmod q}^* \prod_{i=1}^k E_i(q, ab_i, \mathbf{h}_i), \quad E(q, a; \mathbf{h}) = q^{-n} \sum_{\substack{1 \leq x_j \leq q \\ 1 \leq j \leq n}} e\left(\frac{ax_1^{h_1} x_2^{h_2} \dots x_n^{h_n}}{q}\right),$$

the (absolutely convergent) singular integral

$$\mathcal{J}_{\mathbf{b}}(\mathbf{X}) = \langle \mathbf{X}_0 \rangle \int_{-\infty}^{\infty} \prod_{i=1}^k I_1(b_i \beta, \mathbf{X}_i; \mathbf{h}_i) d\beta, \quad I(\beta, \mathbf{Y}; \mathbf{h}) = \int_{\mathcal{Y}} e(\beta y_1^{h_1} y_2^{h_2} \dots y_n^{h_n}) d\mathbf{y}$$

and the number $\mathcal{N}_{\mathbf{b}}(\mathbf{X})$ of solutions $\mathbf{x} \in \mathbb{Z}^J$ to (12.1) satisfying $\frac{1}{2}X_{ij} \leq |x_{ij}| \leq X_{ij}$.

As in [BBDG, (3.6), (7.1) – (7.3)] define the block matrix

$$\mathcal{A} = \begin{pmatrix} \mathcal{A}_1 & \mathcal{A}_2 \\ \mathcal{A}_3 & \mathcal{A}_4 \end{pmatrix} \in \mathbb{R}^{(J+1) \times (N+k)}$$

where $\mathcal{A}_1 = (\alpha_{ij}^\nu) \in \mathbb{R}_{\geq 0}^{J \times N}$ with $0 \leq i \leq k$, $1 \leq j \leq J_i$, $1 \leq \nu \leq N$,

$$\mathcal{A}_2 = (e_{ij}^\mu) \in \mathbb{R}^{J \times k} \text{ with } e_{ij}^\mu = \begin{cases} \delta_{\mu=i} h_{ij} & i < k, \mu < k, \\ -h_{kj} & i = k, \mu < k, \\ -1 & i < k, \mu = k, \\ h_{kj} - 1 & i = k, \mu = k, \end{cases}$$

$\mathcal{A}_3 = (1, \dots, 1) \in \mathbb{R}^{1 \times N}$ and $\mathcal{A}_4 = (0, \dots, 0, -1) \in \mathbb{R}^{1 \times k}$. Let $R = \text{rk}(\mathcal{A}_1)$ and $c_2 = J - R$ as in [BBDG, (7.5)]. Let \mathcal{J} be a set of subsets of pairs (i, j) with $0 \leq i \leq k$, $1 \leq j \leq J_i$. For $H \geq 1$,

some small fixed constant $0 < \lambda < 1$ and $\mathbf{b}, \mathbf{y} \in \mathbb{N}^J$ let $N_{\mathbf{b}, \mathbf{y}}(B, H, \mathcal{J})$ be the number of solutions $\mathbf{x} \in (\mathbb{Z} \setminus \{0\})^J$ satisfying the conditions

$$(12.4) \quad \sum_{i=1}^k \prod_{j=1}^{J_i} (b_{ij} x_{ij})^{h_{ij}} = 0, \quad \prod_{i=0}^k \prod_{j=1}^{J_i} |y_{ij} x_{ij}|^{\alpha_{ij}^\nu} \leq B \quad (1 \leq \nu \leq N),$$

and at least one of the conditions

$$(12.5) \quad \begin{aligned} \min_{ij} |x_{ij}| &\leq H, \\ \min_{1 \leq i \leq k} \prod_{j=1}^{J_i} |x_{ij}|^{h_{ij}} &< \left(\max_{1 \leq i \leq k} \prod_{j=1}^{J_i} |2x_{ij}|^{h_{ij}} \right)^{1-\lambda}, \\ \min_{(ij) \in J} (|x_{ij}|) &\leq \max_{(ij) \in J} (|2x_{ij}|) \max_{(ij) \in J} (|x_{ij}|)^{-\lambda}, \quad J \in \mathcal{J}. \end{aligned}$$

Let $\mathcal{S}_{\mathbf{y}}(B, H, \mathcal{J})$ denote the set of all $\mathbf{x} \in [1, \infty)^J$ that satisfy (12.5) and the N inequalities in the second part of (12.4).

We choose a maximal linearly independent set of R rows Z_1, \dots, Z_R of the matrix $(\mathcal{A}_1 \mathcal{A}_2)$. Let Z_{R+1}, \dots, Z_J be the remaining rows of $(\mathcal{A}_1 \mathcal{A}_2)$. As in [BBDG, (8.23), (8.24)] let $\mathcal{B} = (b_{kl}) \in \mathbb{R}^{(J-R) \times R}$ be the unique matrix with

$$\mathcal{B} \begin{pmatrix} Z_1 \\ \vdots \\ Z_R \end{pmatrix} = \begin{pmatrix} Z_{R+1} \\ \vdots \\ Z_J \end{pmatrix}.$$

Under Hypothesis 2 below (cf. (12.10)), the last row $(\mathcal{A}_3 \mathcal{A}_4)$ of \mathcal{A} can be written as a linear combination of Z_1, \dots, Z_R , say

$$\sum_{\ell=1}^R b_\ell Z_\ell = (\mathcal{A}_3 \mathcal{A}_4).$$

Suppose these R rows are indexed by a set I of pairs (i, j) with $0 \leq i \leq k$, $1 \leq j \leq J_i$ with $|I| = R$. As in [BBDG, (9.1)] let

$$\Phi^*(\mathbf{t}) = \sum_{i=1}^k \prod_{(i,j) \in I} t_{ij}^{h_{ij}},$$

and let \mathcal{F} be the affine $(R-1)$ -dimensional hypersurface $\Phi^*(\mathbf{t}) = 0$ over \mathbb{R} . Let χ_I be the characteristic function on the set

$$\prod_{(i,j) \in I} |t_{ij}|^{\alpha_{ij}^\mu} \leq 1, \quad 1 \leq \mu \leq N.$$

and define the surface integral

$$c_\infty = 2^{J-R} \int_{\mathcal{F}} \frac{\chi_I(\mathbf{t})}{\|\nabla \Phi^*(\mathbf{t})\|} d\mathcal{F} \mathbf{t}.$$

Let

$$c_{\text{fin}} = \prod_p \lim_{L \rightarrow \infty} \frac{1}{p^{L(J-1)}} \# \left\{ \mathbf{x} \bmod p^L : \begin{aligned} &\sum_{i=1}^k \prod_{j=1}^{J_i} x_{ij}^{h_{ij}} \equiv 0 \bmod p^L, \\ &(\{x_{ij} : (i, j) \in S_\rho\}, p) = 1 \text{ for } 1 \leq \rho \leq r \end{aligned} \right\}$$

and

$$c^* = \text{vol} \left\{ \mathbf{r} \in [0, \infty]^{J-R} : b_\ell - \sum_{\iota=1}^{J-R} r_\iota b_{\iota\ell} \geq 0 \text{ for all } 1 \leq \ell \leq R \right\},$$

cf. [BBDG, (9.3), (8.36), (8.34)].

Suppose that the following three hypotheses hold:

1) [BBDG, Hypothesis 1] The singular series \mathcal{E}_{γ^*} is absolutely convergent and moreover

$$(12.6) \quad \mathcal{E}_{\gamma^*} \ll \gamma_1^{\beta_1} \gamma_2^{\beta_2} \cdots \gamma_k^{\beta_k}$$

for some $\beta_1, \dots, \beta_k \leq 1$. Further, there exist finitely many $\zeta_1 = (\zeta_{11}, \dots, \zeta_{1k}), \dots, \zeta_t = (\zeta_{t1}, \dots, \zeta_{tk}) \in \mathbb{R}^k$ satisfying

$$(12.7) \quad \zeta_{\nu i} > 0 \text{ for all } 1 \leq i \leq k, \quad h_{ij} \zeta_{\nu i} < 1 \text{ for all } i, j, \quad \sum_{i=1}^k \zeta_{\nu i} = 1$$

for all $1 \leq \nu \leq t$, real numbers $0 < \lambda \leq 1$, $\delta_1 > 0$ and $C \geq 0$ and a set \mathcal{J} of subsets of pairs (i, j) with $0 \leq i \leq k$, $1 \leq j \leq J_i$, such that whenever $\mathbf{X} \in [1, \infty)^J$ obeys the conditions

$$(12.8) \quad \begin{aligned} \min_{0 \leq i \leq k} \mathbf{X}_i^{\mathbf{h}_i} &\geq \left(\max_{1 \leq i \leq k} \mathbf{X}_i^{\mathbf{h}_i} \right)^{1-\lambda}, \\ \min_{(ij) \in J} X_{ij} &\geq \max_{(ij) \in J} X_{ij} \cdot \max(X_{ij})^{-\lambda}, \quad J \in \mathcal{J}. \end{aligned}$$

then uniformly in $\mathbf{b} \in (\mathbb{Z} \setminus \{0\})^k$, one has

$$(12.9) \quad \mathcal{N}_{\gamma^*}(\mathbf{X}) - \mathcal{E}_{\gamma^*} \mathcal{I}_{\gamma^*}(\mathbf{X}) \ll (\gamma_1 \cdots \gamma_k)^C (\min_{ij} X_{ij})^{-\delta_1} \sum_{\nu=1}^t \prod_{i=0}^k \prod_{j=1}^{J_i} X_{ij}^{1-h_{ij}\zeta_{\nu i}+\varepsilon}.$$

2) [BBDG, (7.4), (7.6), (8.5), (8.6), Hypothesis 2] Suppose that

$$(12.10) \quad \text{rk}(\mathcal{A}_1) = \text{rk}(\mathcal{A}).$$

For λ and \mathcal{J} as above suppose that there exist $\boldsymbol{\eta} = (\eta_{ij}) \in \mathbb{R}_{>0}^J$, ζ as¹ in (12.7) and $\delta_2, \delta_2^* > 0$ with the following properties:

$$(12.11) \quad C_1(\boldsymbol{\eta}): \quad \sum_{(i,j) \in S_\rho} \eta_{ij} \geq 1 + \delta_2 \quad \text{for all } 1 \leq \rho \leq r,$$

$$(12.12) \quad N_{\gamma, \gamma \cdot \mathbf{y}}(B, H, \mathcal{J}) \ll B(\log B)^{c_2-1+\varepsilon} (1 + \log H) \gamma^{-\boldsymbol{\eta}} \langle \mathbf{y} \rangle^{-\delta_2^*}$$

and

$$(12.13) \quad \sum_{\nu=1}^t \int_{\mathcal{J}_{\mathbf{y}}(B, H, \mathcal{J})} \prod_{ij} x_{ij}^{-h_{ij}\zeta_{\nu i}} d\mathbf{x} \ll B(\log B)^{c_2-1+\varepsilon} (1 + \log H) \langle \mathbf{y} \rangle^{-\delta_2^*}$$

for any $\varepsilon > 0$. In addition, suppose that there is some $\delta_4 > 0$ with

$$(12.14) \quad \sum_{(i,j) \in S_\rho} (1 - \beta_i h_{ij}) \geq 1 + \delta_4 \quad (1 \leq \rho \leq r) \quad \text{and} \quad \beta_i h_{ij} \leq 1 \quad (1 \leq i \leq k, 1 \leq j \leq J_i).$$

and

$$(12.15) \quad J_i \geq 2 \quad \text{whenever } \zeta_i \geq 1/2.$$

For any vector ζ satisfying (12.7), where we allow more generally $\zeta_i \geq 0$, and for arbitrary $\zeta_0 > 0$, we also assume that the system of $J+1$ linear equations

$$(12.16) \quad \begin{pmatrix} \mathcal{A}_1 \\ \mathcal{A}_3 \end{pmatrix} \boldsymbol{\sigma} = \left(1 - h_{01}\zeta_0, \dots, 1 - h_{kJ_k}\zeta_k, 1 \right)^\top$$

in N variables has a solution $\boldsymbol{\sigma} \in \mathbb{R}_{>0}^N$.

3) [BBDG, (9.2)] Assume that one of the k monomials in Φ^* consists of only one variable, which has exponent 1.

¹Note that in [BBDG, (7.11)], any ζ does the job, contrary to the statement this has nothing to do with the ζ in Hypothesis 5.1.

Then

$$(12.17) \quad N(B) \sim c^* c_{\text{fin}} c_{\infty} B(\log B)^{c_2}, \quad B \rightarrow \infty.$$

Proof. This is [BBDG, Theorem 8.4 & Lemma 9.1]. The only changes are

- the bound (12.6) is only needed for $\mathbf{b} = \gamma^*$, see the displays before (8.19) and (8.35), and the display (8.30) in [BBDG];
- the bound (12.12) is only needed for $\mathbf{b} = \gamma$, see (8.12), and the last display in Section 8.2 in [BBDG];
- in (12.9) we can afford a sum over finitely many values of ζ . Obviously this has no influence on the argument in [BBDG, Section 8.3];
- we inserted some additional inequalities in (12.5) and the corresponding inequalities in (12.8) (in [BBDG, (5.11), (7.8)] we had $\mathcal{J} = \emptyset$). This has no impact on the argument in [BBDG, Section 8]. Only the set $\mathcal{R}_{\delta, \lambda}$ in [BBDG, Section 8.2] needs to be redefined as

$$\mathcal{R}_{\delta, \lambda} = \left\{ \mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_k) \in [1, \infty)^J : \begin{array}{l} \min_{i,j} X_{ij} \geq \max X_{ij}^{\delta}, \\ \min_{1 \leq i \leq k} \mathbf{X}_i^{\mathbf{h}_i} \geq (\max_{1 \leq i \leq k} \mathbf{X}_i^{\mathbf{h}_i})^{1-\lambda}, \\ \min_{(ij) \in J} X_{ij} \geq \max_{(ij) \in J} X_{ij} \cdot \max(X_{ij})^{-\lambda}, J \in \mathcal{J} \end{array} \right\}.$$

The conditions under which (12.17) holds, look very complicated, but much of this will be automatic in our applications. All counting problems given in Corollary 11.1 are of the form (12.1) – (12.3). In particular, for X_1 we have $r = 2$ and

$$S_1 = \{(1, 1), (1, 2), (2, 1), (3, 3)\}, \quad S_2 = \{(3, 1), (3, 2)\},$$

so for $\mathbf{g} = (\eta, \xi) \in \mathbb{N}^2$ we have

$$\gamma = (\eta, \eta, \eta, \xi, \xi, \eta) \in \mathbb{N}^6, \quad \gamma^* = (\eta^2, \eta^2, \eta^2 \xi^2) \in \mathbb{N}^3,$$

and hence $\mathcal{E}_{\gamma^*} = \mathcal{E}_{\xi}$ in the notation of (7.1).

Hypothesis 1 with $\mathcal{J} = \emptyset$ along with (12.14) and (12.15) follows from [BBDG, Proposition 5.1] exactly as in the proof of [BBDG, Theorem 10.1] for X_2, X_3 . For X_1 with its special torsor equation, we choose

$$(12.18) \quad \mathcal{J} = \{\{(1, 1), (1, 2), (2, 1)\}, \{(3, 1), (3, 2)\}\}$$

reflecting the first two inequalities in (6.1). Then Hypothesis 1 follows from Proposition 7.1 and (7.2), and (12.14) and (12.15) are obvious.

In all cases, (12.10) and (12.16) follow from Remark 11.2 and Hypothesis 3 is satisfied by (3.3), which we verify by inspection of $\Sigma'(1) \supset \Sigma(1)$ in each case.

Thus in order to prove Theorem 1.1 it only remains to check (12.11) – (12.13) and that the constant $c^* c_{\text{fin}} c_{\infty}$ in (12.17) agrees with Peyre’s prediction. For the latter, we apply Proposition 3.5 (where transformations similar to those in Corollary 11.1 will be necessary for X_2, X_3 since Y_2, Y_3 are singular) and Proposition 3.6 with (3.3) since we are in the situation of Remark 2.1.

12.2. The variety X_1 . We are given the equation (4.1) with $J = 6$ variables, with $r = 2$ coprimality conditions

$$(x_{11}, x_{12}, x_{21}, x_{33}) = (x_{31}, x_{32}) = 1$$

and with $N = 8$ height conditions given by the exponent matrix

$$\mathcal{A}_1 = (\alpha_{ij, \nu}) = \begin{pmatrix} 2 & 2 & 2 & 2 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \end{pmatrix} \in \mathbb{R}^{6 \times 8}, \quad \mathcal{A}_2 = \begin{pmatrix} 1 & -1 \\ 1 & -1 \\ -1 & 2 \\ -1 & -1 \\ -1 & -1 \\ -2 & -2 \end{pmatrix} \in \mathbb{R}^{6 \times 3}.$$

We are going to check (12.11) – (12.13). With \mathcal{J} as in (12.18), (12.12) follows from Proposition 8.1 with $\delta_2^* = 1/4 > 0$ and $\boldsymbol{\eta} = \frac{99}{100}(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}, \frac{3}{4}, \frac{1}{2})$ satisfying (12.11). The continuous version (12.13) of (12.12) for the three ζ values given in (7.5) proceeds now in exactly in the same way as the

proof of Proposition 8.1. The resulting leading constant is Peyre's constant by Proposition 3.5 and Proposition 3.6.

12.3. The variety X_2 . For X_2 and X_3 we choose $\mathcal{J} = \emptyset$ as in [BBDG] and apply [BBDG, Proposition 7.6] (as in [BBDG, Sections 11.4 and 12.4]) to check (12.11) – (12.13).

By Corollary 11.1(b) we have $J = 7$ torsor variables x_{ij} with $0 \leq i \leq 3$ satisfying the equation

$$(12.19) \quad x_{11}x_{12} + x_{21}^2 + x_{31}x_{32} = 0,$$

after changing signs. We have $r = 4$ coprimality conditions with

$$(12.20) \quad S_1 = \{(1, 1), (1, 2), (2, 1)\}, \quad S_2 = \{(0, 1), (3, 2)\}, \quad S_3 = \{(0, 2), (3, 1)\}, \quad S_4 = \{(0, 1), (0, 2)\},$$

We have $N = 11$ height conditions with corresponding exponent matrix

$$\mathcal{A}_1 = \begin{pmatrix} 2 & 2 & 2 & 1 & 1 & 1 & 2 \\ 1 & 1 & 3 & 3 & 3 & 3 & 2 \\ 1 & 1 & 1 & 3 & 3 & 3 & 3/2 \\ 1 & 1 & 1 & 1 & 1 & 5/2 & 3/2 \end{pmatrix} \in \mathbb{R}_{\geq 0}^{7 \times 11}, \quad \mathcal{A}_2 = \begin{pmatrix} -1 \\ -1 \\ -1 \\ -1 \\ 1 \\ -2 \\ 1 \end{pmatrix} \in \mathbb{R}^{7 \times 3}.$$

We choose $\zeta = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ satisfying [BBDG, (5.10), (8.6)], $\lambda = 1/25200$ as in [BBDG, (5.13)], and $\tau^{(2)} = (1, 1, \frac{1}{2}, \frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2})$ satisfying [BBDG, (7.18) using the notation (7.13)]. Using a computer algebra system and the notation [BBDG, (7.30) – (7.32)], we confirm the conditions $C_2(\tau^{(2)})$, $C_2((1 - h_{ij}/3)_{ij})$ from [BBDG, Proposition 7.6]. With $c_2 = 2$ we obtain

$$\dim(\mathcal{H} \cap \mathcal{P}) = 2,$$

$$\dim(\mathcal{H} \cap \mathcal{P}_{ij}) = \begin{cases} 1, & (i, j) = (0, 2), (3, 1), (3, 2), \\ 0, & \text{otherwise,} \end{cases}$$

$$\dim(\mathcal{H} \cap \mathcal{P}(1/25200, \pi)) = 0$$

for both vectors $(1 - h_{ij}/3)_{ij}$ and $\tau^{(2)}$, confirming the two remaining conditions $C_3(\tau^{(2)})$, $C_3((1 - h_{ij}/3)_{ij})$. Thus [BBDG, Proposition 7.6] is available and yields the conditions (12.11) – (12.13).

It remains to verify that the leading constant is of the correct shape, which follows from Proposition 3.5 and Proposition 3.6.

More precisely, for the p -adic densities, we have $c_{\text{fin}} = \prod_p c_p$ as in Section 12.1 with

$$c_p = \lim_{L \rightarrow \infty} \frac{1}{p^{6L}} \# \left\{ \mathbf{x} \bmod p^L : \begin{array}{l} x_{11}x_{12} + x_{21}^2 + x_{31}x_{32} \equiv 0 \bmod p^L, \\ (\{x_{ij} : (i, j) \in S_\rho\}, p) = 1 \text{ for } 1 \leq \rho \leq 4 \end{array} \right\}$$

We note that

$$c_p = \left(1 - \frac{1}{p}\right)^{-1} \lim_{L \rightarrow \infty} \frac{1}{p^{7L}} \# \left\{ (\mathbf{x}, z_1) \bmod p^L : \begin{array}{l} x_{11}x_{12}z_1 - x_{21}^2z_1 - x_{31}x_{32} \equiv 0 \bmod p^L, \\ (\{x_{ij} : (i, j) \in S_\rho\}, p) = 1 \text{ for } 1 \leq \rho \leq 4, \\ (x_{31}x_{32}, z_1, p) = 1 \end{array} \right\}$$

because we have the $(\#(\mathbb{Z}/p^L\mathbb{Z})^\times : 1)$ -surjection

$$(x_{01}, \dots, x_{31}, x_{32}, z_1) \mapsto (x_{01}, x_{02}, x_{11}, -x_{12}, x_{21}, x_{31}, x_{32}z_1^{-1})$$

between the two sets since the final coprimality condition and the congruence imply $z_1 \in (\mathbb{Z}/p^L\mathbb{Z})^\times$. By Proposition 3.5, this shows that $c_p = (1 - p^{-1})^{\text{rk Pic } X} \mu_p(X(\mathbb{Q}_p))$, as expected.

12.4. The variety X_3 . This is very similar to the treatment of X_2 . We have the same torsor variables and the same torsor equation (12.19) and the same coprimality conditions (12.20). We have $N = 11$ height conditions with corresponding exponent matrix

$$\mathcal{A}_1 = \begin{pmatrix} 2 & 2 & 2 & 2 & 1 & 1 & 1 & 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 & 1 & 1 & 1 & 3 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 & 1 & 1 & 1 & 3 & 2 & 2 & 2 \\ 1 & 2 & 2 & 2 & 1 & 1 & 1 & 3 & 2 & 2 & 2 \\ 2 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix} \in \mathbb{R}_{\geq 0}^{7 \times 11}$$

and the same \mathcal{A}_2 as before.

We work with the same $\zeta, \lambda, \tau^{(2)}, \beta, \delta_4$. Everything is identical except [BBDG, (7.35)] because of the different \mathcal{A}_1 . We confirm $C_2(\tau^{(2)}), C_2((1 - h_{ij}/3)_{ij})$ and compute $c_2 = 2$ and

$$\begin{aligned} \dim(\mathcal{H} \cap \mathcal{P}) &= 2, \\ \dim(\mathcal{H} \cap \mathcal{P}_{ij}) &= \begin{cases} 1, & (i, j) = (0, 1), (0, 2), (3, 1), (3, 2), \\ 0, & \text{otherwise,} \end{cases} \\ \dim(\mathcal{H} \cap \mathcal{P}(1/25200, \pi)) &= 0 \end{aligned}$$

for the vector $(1 - h_{ij}/3)_{ij}$, and the same for the vector $\tau^{(2)}$.

REFERENCES

- [ADHL] I. Arzhantsev, U. Derenthal, J. Hausen, A. Laface, *Cox rings*, Cambridge Studies in Advanced Mathematics **144**, Cambridge University Press, Cambridge, 2015.
- [BBDG] V. Blomer, J. Brüdern, U. Derenthal, G. Gagliardi, *The Manin–Peyre conjecture for smooth spherical Fano varieties of semisimple rank one*, [arXiv:2004.09357](https://arxiv.org/abs/2004.09357).
- [BBS] V. Blomer, J. Brüdern, P. Salberger, *On a certain senary cubic form*, Proc. Lond. Math. Soc. **108** (2014), 911–964.
- [BP] P. Bravi, G. Pezzini, *Primitive wonderful varieties*, Math. Z. **282** (2016), 1067–1096.
- [BHB] T. D. Browning, D. R. Heath-Brown, *Density of rational points on a quadric bundle in $\mathbb{P}^3 \times \mathbb{P}^3$* , Duke Math. J. **169** (2020), no. 16, 3099–3165.
- [CLS] D. Cox, J. B. Little, H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics **124**, American Mathematical Society, Providence, RI, 2011.
- [DFI] W. Duke, J. Friedlander, H. Iwaniec, *Bounds for automorphic L-functions*, Invent. Math. **112** (1993), 1–8.
- [HB] D. R. Heath-Brown, *A new form of the circle method, and its application to quadratic forms*, J. Reine Angew. Math. **481** (1996), 149–206.
- [Ho] J. Hofschneider, *Spherical Fano varieties*, Ph.D. thesis, Universität Tübingen, 2015.
- [HM] Z. Huang, P. Montero, *Fano threefolds as equivariant compactifications of the vector group*, Michigan Math. J. **69** (2020), no. 2, 341–368.
- [KK] J. Kollár, S. Kovács, *Singularities of the minimal model program*, Cambridge University Press, Cambridge, 2013.
- [LST] B. Lehmann, A. Sengupta, S. Tanimoto, *Geometric consistency of Manin’s conjecture*, [arXiv:1805.10580](https://arxiv.org/abs/1805.10580).
- [Lu] D. Luna, *Variétés sphériques de type A*, Publ. Math. Inst. Hautes Études Sci. (2001), 161–226.
- [LV] D. Luna, Th. Vust, *Plongements d’espaces homogènes*, Comment. Math. Helv. **58** (1983), 186–245.
- [Pey] E. Peyre, *Hauteurs et mesures de Tamagawa sur les variétés de Fano*, Duke Math. J. **79** (1995), 101–218.
- [Sal] P. Salberger, *Tamagawa measures on universal torsors and points of bounded height on Fano varieties*, Astérisque **251** (1998), 91–258. Nombre et répartition de points de hauteur bornée (Paris, 1996).

UNIVERSITÄT BONN, MATHEMATISCHES INSTITUT, ENDENICHER ALLEE 60, 53115 BONN, GERMANY
Email address: blomer@math.uni-bonn.de

UNIVERSITÄT GÖTTINGEN, MATHEMATISCHES INSTITUT, BUNSENSTRASSE 3–5, 37073 GÖTTINGEN, GERMANY
Email address: jbruede@gwdg.de

LEIBNIZ UNIVERSITÄT HANNOVER, INSTITUT FÜR ALGEBRA, ZAHLENTHEORIE UND DISKRETE MATHEMATIK, WELFENGARTEN 1, 30167 HANNOVER, GERMANY
Email address: derenthal@math.uni-hannover.de

LEIBNIZ UNIVERSITÄT HANNOVER, INSTITUT FÜR ALGEBRA, ZAHLENTHEORIE UND DISKRETE MATHEMATIK, WELFENGARTEN 1, 30167 HANNOVER, GERMANY
Email address: gagliardi@math.uni-hannover.de