

FANS AND POLYTOPES IN TILTING THEORY I: FOUNDATIONS

TOSHITAKA AOKI, AKIHIRO HIGASHITANI, OSAMU IYAMA, RYOICHI KASE, AND YUYA MIZUNO

ABSTRACT. For a finite dimensional algebra A over a field k , the 2-term silting complexes of A gives a simplicial complex $\Delta(A)$ called the *g-simplicial complex*. We give tilting theoretic interpretations of the h -vectors and Dehn-Sommerville equations of $\Delta(A)$. Using g -vectors of 2-term silting complexes, $\Delta(A)$ gives a nonsingular fan $\Sigma(A)$ in the real Grothendieck group $K_0(\text{proj } A)_{\mathbb{R}}$ called the *g-fan*. We give several basic properties of $\Sigma(A)$ including sign-coherence, sign decomposition, idempotent reductions, Jasso reductions, pairwise positivity and a connection with Newton polytopes of A -modules. Moreover, $\Sigma(A)$ gives a (possibly infinite and non-convex) polytope $P(A)$ in $K_0(\text{proj } A)_{\mathbb{R}}$ called the *g-polytope* of A . We call A *g-convex* if $P(A)$ is convex. In this case, we show that it is a reflexive polytope, and that the dual polytope is given by the 2-term simple minded collections of A . There are precisely 7 convex g -polytopes up to isomorphism. We give a classification of algebras whose g -polytopes are smooth Fano.

We study g -fans and g -polytopes of 3 classes of algebras. First, the g -fan of a classical or generalized preprojective algebra is given by the Coxeter fan. It is g -convex if and only if it is of type A or B , and in this case, its g -polytope is the dual polytope of the short root polytope. Secondly, the g -fan of a Jacobian algebra of a non-degenerate quiver with potential is given by the g -fan of the corresponding cluster algebra. It is g -convex if and only if it is mutation equivalent to a quiver of type A . Thirdly, we classify Brauer graph algebras which are g -convex, and describe their g -polytopes as the root polytopes of type A or C .

CONTENTS

1. Introduction	1
2. Preliminaries	6
3. g -simplicial complexes	9
4. g -fans	14
5. g -polytopes, c -polytopes and Newton polytopes	24
6. Convex g -polygons and Smooth Fano g -polytopes	32
7. Preprojective algebras and Coxeter fans	36
8. Jacobian algebras and Cluster algebras	45
9. Brauer graph algebras and Root polytopes	51
Acknowledgments	63
References	63


1. INTRODUCTION

The notion of tilting objects is basic to study the structure of a given derived category. The set of partial tilting modules over a finite dimensional algebra has a structure of a simplicial complex, and gives rise to a fan in the Grothendieck group. Their structure has been studied by a number of authors including [RiS, U, H1]. The class of silting objects gives a completion of the class of tilting objects from a point of view of mutation [KV, AiI]. Silting objects correspond bijectively with other important objects in the derived category, including (co-)t-structures and simple-minded collections [AIR, IT, KY]. The subset of 2-term silting complexes enjoys especially nice properties, e.g. [An, As1, AMV, BY, DIRRT]. It plays an important role in categorification of cluster algebras

of Fomin and Zelevinsky. The 2-term silting version of the simplicial complex and the fan as well as their applications to cluster algebras have been studied e.g. in [Pl2, DF, B, DIJ, BST, As2].

One of main problems in tilting theory is to classify algebras which are *g-finite* (that is, there are only finitely many isomorphism classes of basic 2-term silting complexes). Giving an explicit classification up to algebra isomorphism is not a reasonable problem to study (e.g. any local algebra is *g-finite*). Thus the Hasse quivers of basic 2-term silting complexes have been studied by a number of researchers. As was pointed out in [DIJ], the associated fan is a stronger combinatorial invariant than the Hasse quiver. The aim of this paper is to make a systematic use of the simplicial complexes and the fans as essential combinatorial invariants of tilting theory of given algebras.

To explain more details, let A be a finite dimensional algebra over a field k , and let $n := |A|$, where $|X|$ is the number of non-isomorphic indecomposable direct summands of $X \in \text{mod } A$. The 2-term presilting complexes of A give a simplicial complex $\Delta(A)$ called the *g-simplicial complex*. Moreover, each 2-term presilting complex T of A gives a simplicial cone in the real Grothendieck group $K_0(\text{proj } A)_{\mathbb{R}} \simeq \mathbb{R}^n$ generated by the *g-vectors* of the indecomposable direct summands of T , and we obtain a nonsingular fan $\Sigma(A)$ in $K_0(\text{proj } A)_{\mathbb{R}}$ called the *g-fan* of A . Each 2-term silting complex T also gives an n -simplex $C_{\leq 1}(T)$ as the convex hull of the origin and the *g-vectors* of the indecomposable direct summands of T . Gluing them together, we obtain the *g-polytope* $P(A)$. We study the *g-simplicial complex* $\Delta(A)$, the *g-fan* $\Sigma(A)$ and the *g-polytope* $P(A)$ of a finite dimensional algebra A mainly in the case A is *g-finite*. We give some examples of $P(A)$.



We denote by $2\text{-silt } A$ the set of isomorphism classes of basic 2-term silting complexes of A , which has a natural partial order such that the Hasse quiver $\text{Hasse}(2\text{-silt } A)$ is n -regular (see Section 2). The *f-vector* of $\Delta(A)$ gives the number of isomorphism classes of basic 2-term presilting complexes with a fixed number of indecomposable direct summands. Our first main result gives the following representation theoretic interpretation of the *h-vector* of $\Delta(A)$.

Theorem 1.1 (Theorem 3.4). *Let A be a finite dimensional algebra over a field k which is *g-finite*, $n := |A|$ and (h_0, \dots, h_n) the *h-vector* of $\Delta(A)$. Then, for each $0 \leq j \leq n$, we have*

$$h_j = \#2\text{-silt}_j A = \#\text{sbrick}_j A,$$

where $2\text{-silt}_j A$ is the set of isomorphism classes of basic 2-term silting complexes T such that precisely j arrows start at T in $\text{Hasse}(2\text{-silt } A)$, and $\text{sbrick}_j A$ is the set of isomorphism classes of basic semibricks S of A satisfying $|S| = j$.

It was shown in [DIRRT] that there is a canonical bijection $2\text{-silt}_1 A \simeq 2\text{-silt}_{n-1} A$ between join-irreducible elements in $2\text{-silt } A$ and meet-irreducible elements in $2\text{-silt } A$. We give the following generalization, which categorifies Dehn-Sommerville equations $h_j = h_{n-j}$ for *h-vectors*.

Theorem 1.2 (Theorem 3.14). *Let A be a finite dimensional algebra over a field k which is *g-finite*, $n := |A|$ and (h_0, \dots, h_n) the *h-vector* of $\Delta(A)$. For $0 \leq j \leq n$, there are canonical bijections*

$$\text{sbrick}_j A \simeq \text{sbrick}_{n-j} A \text{ and } 2\text{-silt}_j A \simeq 2\text{-silt}_{n-j} A.$$

In particular, we have $h_j = h_{n-j}$.

The *h-vectors* of boundary complexes of simplicial polytopes are known to be unimodal (see [Zi, Section 8]). Recently, this was generalized to simplicial spheres [Adi]. As an application, we obtain the unimodality result below. It will be interesting to give a representation theoretic proof.

Corollary 1.3 (Corollary 3.16). *Let A be a finite dimensional algebra over a field k which is g -finite, and $n := |A|$. Then we have*

$$\begin{aligned} \#sbrick_1 A &\leq \#sbrick_2 A \leq \cdots \leq \#sbrick_{\lfloor \frac{n}{2} \rfloor - 1} A \leq \#sbrick_{\lfloor \frac{n}{2} \rfloor} A, \\ \#sbrick_{\lceil \frac{n}{2} \rceil} A &\geq \#sbrick_{\lceil \frac{n}{2} \rceil + 1} A \geq \cdots \geq \#sbrick_{n-1} A \geq \#sbrick_n A. \end{aligned}$$

In Section 4, we study g -fans of finite dimensional algebras. The following are straightforward consequences of known results in tilting theory.

Proposition 1.4 (Propositions 4.2, 4.12, 4.14, 5.8). *Let A be a finite dimensional algebra over a field k and $n := |A|$.*

- (a) $\Sigma(A)$ is a nonsingular fan in $K_0(\text{proj } A)_{\mathbb{R}}$.
- (b) Any cone in $\Sigma(A)$ is a face of a cone of dimension n .
- (c) Any cone in $\Sigma(A)$ of dimension $n - 1$ is a face of precisely two cones of dimension n .
- (d) A is g -finite (or equivalently, $\Sigma(A)$ is finite) if and only if $\Sigma(A)$ is complete.
- (e) $\Sigma(A)$ is sign-coherent (see Definition 4.11), ordered (see Definition 4.13), and pairwise positive (see Definition 5.6).

For each idempotent $e \in A$ and the corresponding subalgebra eAe of A , we regard $K_0(\text{proj } eAe)_{\mathbb{R}}$ as a subspace of $K_0(\text{proj } A)_{\mathbb{R}}$. Thanks to sign-coherence, one can restrict $\Sigma(A)$ to $K_0(\text{proj } eAe)_{\mathbb{R}}$ to get a subfan. It has the following representation theoretic meaning.

Theorem 1.5 (Theorem 4.16). *There exists an isomorphism of fans*

$$\Sigma(eAe) \simeq \{\sigma \in \Sigma(A) \mid \sigma \subset K_0(\text{proj } eAe)_{\mathbb{R}}\}.$$

We also show that the restriction of $\Sigma(A)$ to each orthant can be described by a simpler algebra (Theorem 4.23) as an analog of the sign decomposition [Ao].

For each 2-term presilting complex U of A , we obtain a new fan $\Sigma(A)/C(U)$ (Definition 4.8). Also there exists a finite dimensional algebra B (called *Jasso reduction* [J]) such that there exist a canonical bijection

$$\{T \in 2\text{-silt } A \mid U \in \text{add } T\} \simeq 2\text{-silt } B$$

and an isomorphism $K_0(\text{proj } A)_{\mathbb{R}}/\mathbb{R}C(U) \simeq K_0(\text{proj } B)_{\mathbb{R}}$. These two constructions are closely related to each other.

Theorem 1.6 (Theorem 4.9). *There exists an isomorphism of fans*

$$\Sigma(A)/C(U) \simeq \Sigma(B)$$

induced by a natural isomorphism $K_0(\text{proj } A)_{\mathbb{R}} \rightarrow K_0(\text{proj } A)_{\mathbb{R}}/\mathbb{R}C(U) \simeq K_0(\text{proj } B)_{\mathbb{R}}$.

Even in rank 2, there are infinitely many g -fans (Example 4.3). It is interesting to classify all possible g -fans in \mathbb{R}^d . In a forthcoming paper [AHKM1], we will give a complete answer to the following problem for $d = 2$.

Problem 1.7. *Characterize sign-coherent fans in \mathbb{R}^d which can be realized as a g -fan of some finite dimensional algebra.*

In Section 5, using the g -fans, we introduce the g -polytopes $P(A)$ of finite dimensional algebras A . We study the Ehrhart series $\text{Ehr}_A(x)$ of $P(A)$, which is the generating function of the number of isomorphism classes of 2-term presilting complexes of A with at most ℓ (possibly isomorphic) indecomposable direct summands. Using the h -vector, we will give the following formula.

Theorem 1.8 (Theorem 5.5). *Let A be a finite dimensional algebra over a field k which is g -finite, $n := |A|$ and (h_0, \dots, h_n) the h -vector of $\Delta(A)$. Then the Ehrhart series of A is given by*

$$\text{Ehr}_A(x) = \frac{\sum_{i=0}^n h_i x^i}{(1-x)^{n+1}}.$$

Using tilting theory, we give explicit connections between the g -fan $\Sigma(A)$ and the normal fans $\Sigma(N(X))$ of the Newton polytopes $N(X)$ of A -modules X (Definitions 5.17, 2.4), and between the Hasse quiver $\text{Hasse}(2\text{-silt}A)$ and the 1-skeleton of $N(X)$. In particular, the following result recovers results in [Fe1] for g -finite case (see also [BCDMTY, PPPP]).

Theorem 1.9 (Theorem 5.22, Corollary 5.24). *Let A be a finite dimensional algebra over a field k which is g -finite. For each $X \in \text{mod } A$, we have*

$$\Sigma(N(X)) = \Sigma(A) / \sim_X \quad \text{and} \quad \vec{N}_1(X) \simeq \text{Hasse}(2\text{-silt}A) / \sim_X$$

(see Definition 5.21). Moreover, there exists $X \in \text{mod } A$ such that $\Sigma(N(X)) = \Sigma(A)$ and $\vec{N}_1(X) \simeq \text{Hasse}(2\text{-silt}A)$.

The rest of this paper is devoted to study finite dimensional algebras whose g -polytopes are convex. We call A g -convex if $P(A)$ is convex. We give characterizations of g -convexity (Theorem 5.10). In particular, g -convexity implies g -finiteness. We introduce the c -polytope $P^c(A)$ by using 2-simple-minded collections (Definition 5.13). Using tilting- t -structure correspondence, we prove the following result.

Theorem 1.10 (Theorem 5.14). *Let A be a finite dimensional algebra over a field k . Then A is g -convex if and only if*


$$P(A) = (P^c(A))^*.$$

In this case, both $P(A)$ and $P^c(A)$ are reflexive polytopes.


For each $n \geq 1$, there exists only finitely many convex g -polytopes of dimension n up to isomorphisms of g -polytopes (see Definition 6.1, Proposition 6.2). It is interesting to know the maximal number of $\#2\text{-psilt}A$ for g -convex algebras A with $|A| = n$, see Problem 1.15 below.

In Section 6, we give the following classification of convex g -polygons by using the well-known list of 16 reflexive polygons [PR].

Corollary 1.11 (Theorem 6.3). *There are precisely 7 convex g -polygons up to isomorphism of g -polytopes.*



The corresponding c -polygons are the following.



It is well-known that there are 4319 reflexive polytopes in dimension 3 [KS]. It is interesting to know which one can be realized as a g -polytope. In a forthcoming paper [AHKM2], we will give a complete answer to the following problem for $d = 3$.

Problem 1.12. *Classify convex g -polytopes in \mathbb{R}^d .*

We also give a classification of algebras whose g -polytopes are smooth Fano (Definition 6.4), a much stronger notion than convexity.

Theorem 1.13 (Theorem 6.6). *Let A be a finite dimensional algebra over a field k . Then $P(A)$ is a smooth Fano polytope if and only if A is a product of local algebras, algebras of pentagon type and algebras of hexagon type.*

In Sections 7–9, we give explicit descriptions of the g -fans and/or g -polytopes of certain important classes of algebras. In Section 7, we describe g -polytopes for classical and generalized preprojective algebras due to Geiss-Leclerc-Schröer [GLS] by using the root polytope.

Theorem 1.14 (Theorem 7.4). *Let Π be a classical or generalized preprojective algebra of Dynkin type.*

- (a) $\Sigma(\Pi)$ is the Coxeter fan.
- (b) Π is g -convex if and only if it is either of type A_n or B_n . In this case, $P(\Pi)$ is the dual polytope of the short root polytope of type A_n or B_n respectively.

In particular, $\#2\text{-silt}\Pi$ is the order of the Weyl group. In type B_n , it is $2^n n!$ and hence the volume of $P(\Pi)$ is 2^n . The following is a list of natural questions to study.

- Problem 1.15.** (a) *Is the volume of convex g -polytopes of rank n at most 2^n ?*
 (b) *Classify symmetric convex g -polytopes.*

Note that the part (a) is true for centrally symmetric convex polytopes by Minkowski's convex body Theorem [C, Chapter III].

In Section 8, we apply silting theory to give a simple proof of some properties of g -vectors and c -vectors of skew-symmetric cluster algebras (Theorem 8.4), which were conjectured in [FZ3] and proved in [DWZ, Nag, NZ, Pl1, GHKK]. In particular, the g -vectors of the cluster algebra associated with a skew-symmetric matrix B gives a sign-coherent nonsingular fan $\Sigma(B)$. We show that $\Sigma(B)$ gives the g -fans of Jacobian algebras A of non-degenerate quivers with potential. More precisely, the following result holds for the fan $\Sigma^\circ(A)$ corresponding to the subset $2\text{-silt}^\circ A$ of $2\text{-silt} A$ consisting of all iterated mutations of A .

Theorem 1.16 (Theorem 8.5, Corollary 8.7). *Let (Q, W) be a non-degenerate quiver with potential with $Q_0 = \{1, \dots, n\}$, and $A := P(Q, W)$ the Jacobian algebra, and $B := B(Q)$ the skew-symmetric matrix. Then we have an isomorphism of fans*

$$\Sigma(B) \simeq \Sigma^\circ(A).$$

Moreover, A is g -convex if and only if Q is mutation equivalent to a quiver of type A .

In Section 9, we study g -polytopes of Brauer graph algebras. In particular, we give the following characterization of g -convex Brauer graph algebras.

Theorem 1.17 (Theorem 9.4). *Let Γ be a connected Brauer graph with n edges and B_Γ the Brauer graph algebra associated to Γ .*

- (a) B_Γ is g -finite if and only if B_Γ is g -convex if and only if Γ is either a Brauer tree or a Brauer odd-cycle.
- (b) If Γ is a Brauer tree (respectively, Brauer odd-cycle), then $P(B_\Gamma)$ is the root polytope of type A_n (respectively, C_n).

We remark that all of our results are valid for a dg algebra A over a field k which is non-positive and proper (see Remark 1.18 for more general setting). In fact, $B := H^0(A)$ is a finite dimensional k -algebra such that $\Delta(A) = \Delta(B)$, $\Sigma(A) = \Sigma(B)$ and $P(A) = P(B)$, see Proposition 2.14. For simplicity, we will work on finite dimensional algebras over fields throughout this paper.

Remark 1.18. The most of our results are valid for a dg algebra A over a field k satisfying the following conditions.

- (A) A is *non-positive*, that is, $H^i(A) = 0$ holds for each $i \geq 1$.
- (B) For each $T \in 2\text{-silt} A$, $\text{add } T$ is a Krull-Schmidt category.
- (C) For each $T \in 2\text{-silt} A$ and an indecomposable direct summand X of T , A has a left $\text{add}(T/X)$ -approximation and $A[1]$ has a right $\text{add}(T/X)$ -approximation.

The assumptions (B) and (C) are satisfied if A is *proper*, that is, $\sum_{i \in \mathbb{Z}} \dim_k H^i(A)$ is finite.

Conventions All modules are right modules. The composition of morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is denote by $gf : X \rightarrow Z$. The composition of arrows $a : i \rightarrow j$ and $b : j \rightarrow k$ in a quiver is denote by $ab : i \rightarrow k$. Let A be a finite dimensional algebra over a field k . We denote by $\text{mod } A$ the category of finitely generated right A -modules and by $\text{proj } A$ the category of finitely

generated projective A -modules. We denote by $D^b(\text{mod } A)$ the bounded derived category of $\text{mod } A$ and by $K^b(\text{proj } A)$ the bounded homotopy category of $\text{proj } A$. For an object in a Krull-Schmidt category (e.g. $\text{mod } A$, $K^b(\text{proj } A)$, $D^b(\text{mod } A)$), we denote by $|X|$ the number of non-isomorphic indecomposable direct summands of X .

2. PRELIMINARIES

2.1. Preliminaries on fans and polytopes. We recall some fundamental materials on fans and polytopes. We refer the reader to e.g. [Fu, BR, BP] for these materials.

A *convex polyhedral cone* σ is a set of the form $\sigma = \{\sum_{i=1}^s r_i v_i \mid r_i \geq 0\}$, where $v_1, \dots, v_s \in \mathbb{R}^d$. We denote it by $\sigma = \text{cone}\{v_1, \dots, v_s\}$. Note that $\{0\}$ is regarded as a convex polyhedral cone. We collect some notions concerning convex polyhedral cones. Let σ be a convex polyhedral cone.

- The dimension of σ is the dimension of the linear space generated by σ .
- We say that σ is *strongly convex* if $\sigma \cap (-\sigma) = \{0\}$ holds, i.e., σ does not contain a linear subspace of positive dimension.
- We call σ *rational* if each v_i can be taken from \mathbb{Q}^d .
- We denote by $\langle \cdot, \cdot \rangle$ the usual inner product. A *supporting hyperplane* of σ is a hyperplane $\{v \in \sigma \mid \langle u, v \rangle = 0\}$ in \mathbb{R}^d given by some $u \in \mathbb{R}^d$ satisfying $\sigma \subset \{v \in \mathbb{R}^d \mid \langle u, v \rangle \geq 0\}$.
- A *face* τ of σ is the intersection of σ with a supporting hyperplane of σ .

In what follows, a cone means a strongly convex rational polyhedral cone for short.

Definition 2.1. A *fan* Σ in \mathbb{R}^d is a collection of cones in \mathbb{R}^d such that

- (a) each face of a cone in Σ is also contained in Σ , and
- (b) the intersection of two cones in Σ is a face of each of those two cones.

In this case, for each $0 \leq i \leq d$, we denote by Σ_i the subset of cones of dimension i . For example, Σ_0 consists of the trivial cone $\{0\}$. We call each element in Σ_1 a *ray* of Σ .

We collect some notions concerning fans used in this paper. Let Σ be a fan in \mathbb{R}^d .

- We call Σ *finite* if it consists of a finite number of cones.
- We call Σ *complete* if $\bigcup_{\sigma \in \Sigma} \sigma = \mathbb{R}^d$.
- We call Σ *nonsingular* (or *smooth*) if each maximal cone in Σ is generated by a \mathbb{Z} -basis for \mathbb{Z}^d .

We prepare some notions which will be used in this paper.

Definition 2.2. Let Σ and Σ' be fans in \mathbb{R}^d and $\mathbb{R}^{d'}$ respectively. An *isomorphism* $\Sigma \simeq \Sigma'$ of fans is an isomorphism $\mathbb{Z}^d \simeq \mathbb{Z}^{d'}$ of abelian groups such that the induced linear isomorphism $\mathbb{R}^d \rightarrow \mathbb{R}^{d'}$ gives a bijection $\Sigma \simeq \Sigma'$ between cones.

Definition 2.3. Let Σ be a finite fan in \mathbb{R}^d , and let \sim be an equivalence relation on Σ_d . We say that \sim *coarsens* Σ if, for each $\sigma \in \Sigma_d$, the set $[\sigma] := \bigcup_{\tau \sim \sigma} \tau$ is convex. In this case, we define a fan Σ / \sim called the *coarsening* of Σ by

$$\Sigma / \sim := \{[\sigma_1] \cap \dots \cap [\sigma_s] \mid s \geq 1, \sigma_1, \dots, \sigma_s \in \Sigma_d\}.$$

A *polytope* P is a convex hull of a finite subset K of \mathbb{R}^d . It is called *lattice polytope* if K is contained in \mathbb{Z}^d . For lattice polytopes P and P' in \mathbb{Z}^d and $\mathbb{Z}^{d'}$ respectively, an *isomorphism* $P \simeq P'$ of lattice polytopes is an isomorphism $\mathbb{Z}^d \simeq \mathbb{Z}^{d'}$ of abelian groups such that the induced linear isomorphism $\mathbb{R}^d \rightarrow \mathbb{R}^{d'}$ gives a bijection $P \simeq P'$.

Definition 2.4. Let P be a polytope in $V = \mathbb{R}^d$, and V^* the dual space of V . For each face F of P , let F_1, \dots, F_s be all facets of P containing F . For each $1 \leq i \leq s$, let $v_i \in V^*$ be an outer normal vector of F_i , and let

$$\sigma_F := \text{cone}\{v_1, \dots, v_s\}.$$

The *normal fan* of P is

$$\Sigma(P) := \{\sigma_F \mid F \text{ is a face of } P\}.$$

If P has dimension d , then $\Sigma(P)$ is a finite complete fan in \mathbb{R}^d . Otherwise, the cones of $\Sigma(P)$ is not strongly convex.

Each element $f \in V^*$ gives a face of P :

$$P_f := \{v \in P \mid f(v) = \max f(P)\}.$$

For each face F of P , the corresponding cone $\sigma_F \in \Sigma(P)$ can be written as

$$\sigma_F = \{f \in V^* \mid P_f \supseteq F\}.$$

Then each $f \in \sigma_F^\circ$ satisfies $P_f = F$.

2.2. Preliminaries on tilting theory. Let A be a finite dimensional algebra over a field k . We recall basic results on silting theory from [AiI, AIR]. First we recall the definition of 2-term silting complexes.

Definition 2.5. Let $T = (T^i, d^i) \in \mathbf{K}^b(\text{proj } A)$.

- (a) T is called *presilting* if $\text{Hom}_{\mathbf{K}^b(\text{proj } A)}(T, T[\ell]) = 0$ for all positive integers ℓ .
- (b) T is called *silting* if it is presilting and $\mathbf{K}^b(\text{proj } A) = \text{thick } T$.
- (c) T is called *2-term* if $T^i = 0$ for all $i \neq 0, -1$.

We denote by $\text{silt } A$ (respectively, $\text{psilt } A$, $2\text{-silt } A$, $2\text{-psilt } A$) the set of isomorphism classes of basic silting (respectively, presilting, 2-term silting, 2-term presilting) complexes of $\mathbf{K}^b(\text{proj } A)$. Note that a 2-term presilting complex T is silting if and only if $|T| = |A|$ holds.

For $T, U \in \text{silt } A$, we write $T \geq U$ if $\text{Hom}_{\mathbf{K}^b(\text{proj } A)}(T, U[\ell]) = 0$ holds for all positive integers ℓ . Then $(\text{silt } A, \geq)$ is a partially ordered set [AiI].

Recall that the *Hasse quiver* $\text{Hasse } P$ of a poset P has the set P of vertices, and an arrow $x \rightarrow y$ if $x > y$ and there does not exist $z \in P$ satisfying $x > z > y$.

In this paper, the subposet $(2\text{-silt } A, \geq)$ of $(\text{silt } A, \geq)$ plays a central role. It is known that $\text{Hasse}(2\text{-silt } A)$ is n -regular for $n := |A|$. More precisely, let $T = T_1 \oplus \cdots \oplus T_n \in 2\text{-silt } A$ with indecomposable T_i . For each $1 \leq i \leq n$, there exists precisely one $T' \in 2\text{-silt } A$ such that $T' = T'_i \oplus (\bigoplus_{j \neq i} T_j)$ for some $T'_i \neq T_i$. In this case, we call T' *mutation* of T at T_i and write

$$T' = \mu_i(T).$$

In this case, either $T > T'$ or $T' < T$ holds. The following result is fundamental in silting theory.

Proposition 2.6. Let $T, T' \in 2\text{-silt } A$. Take a decomposition $T = T_1 \oplus \cdots \oplus T_n$ with indecomposable T_i . Then the following conditions are equivalent.

- (a) $T > T'$, and T and T' are mutation of each other.
- (b) There is an arrow $T \rightarrow T'$ in $\text{Hasse}(2\text{-silt } A)$.
- (c) $T' = T'_i \oplus (\bigoplus_{j \neq i} T_j)$ and there is a triangle

$$T_i \xrightarrow{f} U_i \rightarrow T'_i \rightarrow T_i[1]$$

such that f is a minimal left $(\text{add } \bigoplus_{j \neq i} T_j)$ -approximation.

- (d) $T' = T'_i \oplus (\bigoplus_{j \neq i} T_j)$ and there is a triangle

$$T_i \rightarrow U_i \xrightarrow{g} T'_i \rightarrow T_i[1]$$

such that g is a minimal right $(\text{add } \bigoplus_{j \neq i} T_j)$ -approximation.

The triangles in (c) and (d) are isomorphic, and called an exchange triangle.

For a class \mathcal{C} of objects in $\text{mod } A$, let

$$\begin{aligned} \mathcal{C}^\perp &:= \{X \in \text{mod } A \mid \text{Hom}_A(\mathcal{C}, X) = 0\}, \\ {}^\perp \mathcal{C} &:= \{X \in \text{mod } A \mid \text{Hom}_A(X, \mathcal{C}) = 0\}. \end{aligned}$$

There is a strong connection between 2-term silting complexes and some important subcategories defined as follows. See section 3 for more details.

Definition 2.7. A full subcategory \mathcal{C} of $\text{mod } A$ is called a *torsion class* (respectively, *torsionfree class*) if it is closed under extensions and factor modules (respectively, submodules). It is called *functorially finite* if there exists $M \in \mathcal{C}$ satisfying $\mathcal{C} = \text{Fac } M$ (respectively, $\mathcal{C} = \text{Sub } M$). We denote by $\text{tors } A$ (respectively, $\text{f-tors } A$, $\text{torf } A$, $\text{f-torf } A$) the set of torsion classes (respectively, functorially finite torsion classes, torsionfree classes, functorially finite torsionfree classes) in $\text{mod } A$.

We have mutually inverse bijections

$$\text{tors } A \simeq \text{torf } A, \mathcal{C} \mapsto \mathcal{C}^\perp \text{ and } \text{torf } A \simeq \text{tors } A, \mathcal{C} \mapsto {}^\perp \mathcal{C}. \quad (2.1)$$

A pair $(\mathcal{T}, \mathcal{F}) \in \text{tors } A \times \text{torf } A$ is called a *torsion pair* if $\mathcal{F} = \mathcal{T}^\perp$, or equivalently, $\mathcal{T} = {}^\perp \mathcal{F}$. In this case, we have functors

$$t_{\mathcal{T}} : \text{mod } A \rightarrow \mathcal{T} \text{ and } f_{\mathcal{F}} : \text{mod } A \rightarrow \mathcal{F}$$

such that each $X \in \text{mod } A$ admits an exact sequence

$$0 \rightarrow t_{\mathcal{T}} X \rightarrow X \rightarrow f_{\mathcal{F}} X \rightarrow 0. \quad (2.2)$$

The bijections (2.1) restrict to bijections $[\text{Sm}]$

$$\text{f-tors } A \simeq \text{f-torf } A \text{ and } \text{f-torf } A \simeq \text{f-tors } A.$$

The following bijection is also important.

Definition-Proposition 2.8. [AIR, Proposition 1.2(b), Lemma 3.4]

(a) *We have surjections*

$$\begin{aligned} 2\text{-psilt } A &\rightarrow \text{f-tors } A, U \mapsto \mathcal{T}_U := \text{Fac } H^0(U), \\ 2\text{-psilt } A &\rightarrow \text{f-tors } A, U \mapsto \overline{\mathcal{T}}_U := {}^\perp H^{-1}(\nu U), \\ 2\text{-psilt } A &\rightarrow \text{f-torf } A, U \mapsto \mathcal{F}_U := \text{Sub } H^{-1}(\nu U), \\ 2\text{-psilt } A &\rightarrow \text{f-torf } A, U \mapsto \overline{\mathcal{F}}_U := H^0(U)^\perp \end{aligned}$$

such that $(\mathcal{T}_U, \overline{\mathcal{F}}_U)$ and $(\overline{\mathcal{T}}_U, \mathcal{F}_U)$ form torsion pairs. Thus each $X \in \text{mod } A$ admits exact sequences

$$\begin{aligned} 0 \rightarrow t_U X \rightarrow X \rightarrow \bar{f}_U X \rightarrow 0 \text{ for } t_U X &:= t_{\mathcal{T}_U} X \text{ and } \bar{f}_U X := f_{\overline{\mathcal{F}}_U} X, \\ 0 \rightarrow \bar{t}_U X \rightarrow X \rightarrow f_U X \rightarrow 0 \text{ for } \bar{t}_U X &:= t_{\overline{\mathcal{T}}_U} X \text{ and } f_U X := f_{\mathcal{F}_U} X. \end{aligned}$$

(b) *The first two surjections in (a) restrict to the same bijection*

$$2\text{-silt } A \simeq \text{f-tors } A, T \mapsto \mathcal{T}_T = \overline{\mathcal{T}}_T.$$

The last two surjections in (a) restrict to the same bijection

$$2\text{-silt } A \simeq \text{f-torf } A, T \mapsto \mathcal{F}_T = \overline{\mathcal{F}}_T.$$

The following finiteness condition plays a central role in this paper.

Definition 2.9. Let A be a finite dimensional algebra over a field k . We say that A is *g-finite* if $\#2\text{-silt } A < \infty$. (This is called *τ -tilting finite* in [DIJ].)

We give a characterization of *g-finiteness*.

Proposition 2.10. [DIJ, Theorem 1.2] *Let A be a finite dimensional algebra over a field k . Then A is g-finite if and only if $\text{tors } A = \text{f-tors } A$ if and only if $\text{torf } A = \text{f-torf } A$.*

There is a strong connection between 2-term silting complexes and the following class of modules.

Definition 2.11. Let A be a finite dimensional algebra over a field k .

(a) An object $X = X_1 \oplus \cdots \oplus X_r \in \text{mod } A$ is called a *semibrick* if

$$\text{Hom}_A(X_i, X_j) = \begin{cases} \text{division ring} & (i = j) \\ 0 & (j \neq i). \end{cases} \quad (2.3)$$

We denote by $\text{sbrick } A$ the set of isomorphism classes of semibricks in $\text{mod } A$.

- (b) A full subcategory \mathcal{C} of $\text{mod } A$ is called *wide* if it is closed under extensions, kernels and cokernels. We denote by $\text{wide } A$ the set of wide subcategories of $\text{mod } A$.

Note that the usual definition of a semibrick is more general: a (possibly infinite) set of modules satisfying (2.3) [As1]. In this paper, we only need to consider semibricks in the sense above. It is basic that there is a bijection

$$\text{sbrick } A \simeq \text{wide } A$$

sending X to the smallest extension closed subcategory containing $\text{add } X$ [Ri].

The following notion is a derived category version of semibricks.

Definition 2.12. An object $X = X_1 \oplus \cdots \oplus X_r \in \text{D}^b(\text{mod } A)$ is called a *simple-minded collection* if the following conditions hold.

- $\text{Hom}_{\text{D}^b(\text{mod } A)}(X, X[\ell]) = 0$ for all negative integers ℓ .
- For $1 \leq i, j \leq r$, $\text{Hom}_{\text{D}^b(\text{mod } A)}(X_i, X_j) = \begin{cases} \text{division ring} & (i = j) \\ 0 & (j \neq i). \end{cases}$
- $\text{D}^b(\text{mod } A) = \text{thick } X$.

Note that $r = |A|$ holds in this case. A simple-minded collection X is called *2-term* if $H^i(X) = 0$ holds for all integers $i \neq -1, 0$. We denote by $\text{smc } A$ (respectively, $2\text{-smc } A$) the set of isomorphism classes of simple-minded collections (respectively, 2-term simple-minded collections) on $\text{D}^b(\text{mod } A)$.

We have the following silting-t-structure correspondence.

Proposition 2.13. [KY] *Let A be a finite dimensional k -algebra over a field k and $n = |A|$. Then there exists a bijection between $\text{silt } A$ and $\text{smc } A$ such that $T = T_1 \oplus \cdots \oplus T_n \in \text{silt } A$ and the corresponding $S = S_1 \oplus \cdots \oplus S_n \in \text{smc } A$ satisfy*

$$\text{Hom}_{\text{D}^b(\text{mod } A)}(T_i, S_j[p]) = \begin{cases} \text{End}_{\text{D}^b(\text{mod } A)}(S_i) & (i = j \text{ and } p = 0) \\ 0 & (\text{otherwise}). \end{cases} \quad (2.4)$$

In particular, we have

$$([T_i], [S_j]) = \delta_{ij} \cdot \dim_k \text{End}_{\text{D}^b(\text{mod } A)}(S_j). \quad (2.5)$$

Let \mathcal{T} be a triangulated category, and $\text{silt } \mathcal{T}$ (respectively, $\text{psilt } \mathcal{T}$) be the set of isomorphism classes of basic silting (respectively, presilting) objects in \mathcal{T} . Then $\text{silt } \mathcal{T}$ has a partial order \geq as in Definition 2.5. For $T \in \text{silt } \mathcal{T}$, let

$$\begin{aligned} 2_T\text{-silt } \mathcal{T} &:= \{U \in \text{silt } \mathcal{T} \mid T \geq U \geq T[1]\}, \\ 2_T\text{-psilt } \mathcal{T} &:= \{V \in \mathcal{T} \mid \exists U \in 2_T\text{-silt } \mathcal{T} \text{ such that } V \in \text{add } U\}, \end{aligned}$$

Later we use the following basic fact.

Proposition 2.14. [BY, Theorem A.7] *Let \mathcal{T} be an idempotent-complete algebraic triangulated category. For $T \in \text{silt } \mathcal{T}$, let $A := \text{End}_{\mathcal{T}}(T)$. Then there exists a triangle functor $\mathcal{T} \rightarrow \text{K}^b(\text{proj } A)$ which sends T to A and gives an isomorphism $K_0(\mathcal{T}) \simeq K_0(\text{proj } A)$ and bijections*

$$2_T\text{-silt } \mathcal{T} \simeq 2\text{-silt } A \quad \text{and} \quad 2_T\text{-psilt } \mathcal{T} \simeq 2\text{-psilt } A.$$

3. g -SIMPLICIAL COMPLEXES

In this section, we introduce g -simplicial complexes and study their basic properties. In particular, we give a representation theoretic interpretations of their h -vectors. Moreover we give a proof of Dehn-Sommerville equations in terms of the representation theory.

Throughout this section, let A be a finite dimensional algebra over a field k .

Definition 3.1. For $j \geq 0$, let $2\text{-psilt}^j A$ be the set of isomorphism classes of basic 2-term presilting complexes T such that $|T| = j$. We define a simplicial complex $\Delta(A)$ called the *g -simplicial complex* of A as follows: The set of j -simplices is $2\text{-silt}^{j+1} A$.

We give the following basic properties.

Proposition 3.2. *Let A be a finite dimensional algebra over a field k and $n := |A|$.*

- (a) *$\Delta(A)$ is pure of dimension $n - 1$, that is, all facets of $\Delta(A)$ have dimension $n - 1$.*
- (b) *Each face of dimension $n - 2$ in $\Delta(A)$ is contained in precisely two facets.*
- (c) *$\Delta(A)$ is flag, that is, each minimal subset which is not a face of $\Delta(A)$ consists of two points.*

Proof. For (a) and (b), see [DIJ]. (c) is clear. In fact, for distinct elements $T_1, \dots, T_j \in 2\text{-psilt}^1 A$, their direct sum belongs to $2\text{-psilt}^j A$ if and only if $\text{Hom}_{K^b(\text{proj } A)}(T_i, T_{i'}[1]) = 0$ for each $1 \leq i \neq i' \leq j$ if and only if $T_i \oplus T_{i'} \in 2\text{-psilt}^2 A$ for each $1 \leq i \neq i' \leq j$. \square

Now we assume that A is g -finite. For $n := |A|$, we denote by

$$(f_{-1}, f_0, \dots, f_{n-1}) \quad \text{and} \quad (h_0, h_1, \dots, h_n)$$

the f -vector and the h -vector of the g -simplicial complex $\Delta(A)$. Thus

$$f_{-1} := 1 \quad \text{and} \quad f_j := \#2\text{-psilt}^{j+1} A$$

is the number of the j -simplices in $\Delta(A)$ for $j \geq 0$, and h_j is defined by

$$h_j = \sum_{i=0}^j (-1)^{j-i} \binom{n-i}{j-i} f_{i-1} \quad \text{for } 0 \leq j \leq n. \quad (3.1)$$

In other words, the f -polynomial and h -polynomial

$$f(x) := \sum_{i=0}^n f_{i-1} x^{n-i} \quad \text{and} \quad h(x) := \sum_{i=0}^n h_i x^{n-i}$$

are related by $h(x) = f(x-1)$. Thus the equations (3.1) are equivalent to

$$f_{j-1} = \sum_{i=0}^j \binom{n-i}{j-i} h_i \quad \text{for } 0 \leq j \leq n, \quad (3.2)$$

which recover the f -vector from the h -vector. We give a representation theoretic meaning of the f - and h -vectors.

Definition 3.3. Let A be a finite dimensional algebra over a field k and $j \geq 0$.

- (a) Let $2\text{-silt}_j A$ be the set of all $T \in 2\text{-silt} A$ such that there exist precisely j arrows starting at T in $\text{Hasse}(2\text{-silt} A)$.
- (b) Let $\text{sbrick}_j A$ be the set of all $S \in \text{sbrick} A$ such that $|S| = j$.

The following is the first main result in this section.

Theorem 3.4. *Let A be a finite dimensional algebra over a field k which is g -finite. Then for each $0 \leq j \leq |A|$, we have*

$$h_j = \#2\text{-silt}_j A = \#\text{sbrick}_j A.$$

To prove Theorem 3.4, we need preparations. In the rest, let A be a finite dimensional algebra over a field k (which is not necessarily g -finite).

First, we recall the following classical fact (a), see e.g. [AS, Corollary 2.4(a)].

- Definition-Proposition 3.5.** (a) *Each $X \in \text{mod } A$ has a direct summand X_{gen} such that a direct summand Y of X satisfies $\text{Fac } X = \text{Fac } Y$ if and only if Y has X_{gen} as a direct summand.*
- (b) *For a 2-term complex $T \in K^b(\text{proj } A)$, we denote by T_{gen} a minimal direct summand of T such that $H^0(T)_{\text{gen}} = H^0(T_{\text{gen}})$.*

Clearly X_{gen} and T_{gen} are unique up to isomorphism.

For $T \in 2\text{-silt} A$, we have the following description of T_{gen} in terms of $\text{Hasse}(2\text{-silt} A)$.

Lemma 3.6. (a) For $T = T_1 \oplus \cdots \oplus T_n \in 2\text{-silt}A$ with indecomposable T_i , we have

$$T_{\text{gen}} = \bigoplus_{T > \mu_i(T)} T_i.$$

(b) $2\text{-silt}_j A = \{T \in 2\text{-silt}A \mid |T_{\text{gen}}| = j\}$.

Proof. (a) It is immediate from definition of T_{gen} that T_i is a direct summand of T_{gen} if and only if $H^0(T_i) \notin \text{Fac } H^0(T/T_i)$. This is clearly equivalent to $T > \mu_i(T)$ (see the proof of [IRRT, Theorem 2.7]). Thus the assertion holds.

(b) Immediate from (a). \square

Recall from Definition-Proposition 2.8 that each $U \in 2\text{-psilt}A$ gives torsion classes

$$\mathcal{T}_U = \text{Fac } H^0(U) \subseteq \overline{\mathcal{T}}_U = {}^\perp H^{-1}(\nu U)$$

such that the equality holds if $U \in 2\text{-silt}A$. We need the following notions.

Definition-Proposition 3.7. Let $U \in 2\text{-psilt}A$.

- (a) We call $T \in 2\text{-silt}A$ a completion of U if it satisfies $U \in \text{add } T$.
(b) A completion T of U is called minimal (or co-Bongartz) if

$$\mathcal{T}_T = \mathcal{T}_U.$$

Then U has a unique minimal completion up to isomorphism, which we denote by U_{\min} .

- (c) A completion T of U is called maximal (or Bongartz) if

$$\mathcal{T}_T = \overline{\mathcal{T}}_U.$$

Then U has a unique maximal completion up to isomorphism, which we denote by U_{\max} .

- (d) (Jasso reduction) We have

$$\mathcal{W}_U := \overline{\mathcal{T}}_U \cap \overline{\mathcal{F}}_U \in \text{wide}A.$$

Define a functor

$$\text{w}_U : \text{mod } A \rightarrow \mathcal{W}_U \quad \text{by} \quad \text{w}_U X := \overline{\text{t}}_U X / \text{t}_U X = \text{t}_{U_{\max}} X / \text{t}_{U_{\min}} X.$$

Let $[U]$ be an ideal of $\text{End}_{\text{K}^b(\text{proj } A)}(U_{\max})$ consisting of all morphisms factoring through objects in $\text{add } U$, and

$$B := \text{End}_{\text{K}^b(\text{proj } A)}(U_{\max})/[U].$$

Then $|B| = |A| - |U|$ holds, and there exists an equivalence

$$\text{mod } B \simeq \mathcal{W}_U \subseteq \text{mod } A$$

which induces an injective homomorphism $K_0(\text{mod } B) \subseteq K_0(\text{mod } A)$.

Proof. (a) By Definition-Proposition 2.8(a), $\mathcal{T}_U \in \text{f-tors}A$ holds. By Definition-Proposition 2.8(c), there exists unique $T \in 2\text{-silt}A$ satisfying $\mathcal{T}_T = \mathcal{T}_U$.

(b) This is shown similarly.

(c) This is [J, Theorem 3.8], see also [DIRRT, Theorem 4.12]. \square

Lemma 3.8. Let $U \in 2\text{-psilt}A$. Then we have

$$(U_{\min})_{\text{gen}} \simeq U_{\text{gen}} \quad \text{and} \quad (U_{\text{gen}})_{\min} \simeq U_{\min}.$$

Proof. Since U_{\min} has U as a direct summand and $\mathcal{T}_{U_{\min}} = \mathcal{T}_U$, we have $(U_{\min})_{\text{gen}} \simeq U_{\text{gen}}$. Since $\mathcal{T}_{(U_{\text{gen}})_{\min}} = \mathcal{T}_{U_{\text{gen}}} = \mathcal{T}_U = \mathcal{T}_{U_{\min}}$ holds, we have $(U_{\text{gen}})_{\min} \simeq U_{\min}$ by Definition-Proposition 2.8(c). \square

Definition-Proposition 3.9. For $0 \leq j \leq |A|$, let

$$2\text{-psilt}_j A := \{U \in 2\text{-psilt}^j A \mid U_{\min} \in 2\text{-silt}_j A\} \quad (3.3)$$

$$= \{U \in 2\text{-psilt}^j A \mid U_{\text{gen}} = U\}. \quad (3.4)$$

Proof. By Lemma 3.6, $U_{\min} \in 2\text{-silt}_j A$ holds if and only if $|(U_{\min})_{\text{gen}}| = j$ holds. By Lemma 3.8, this is equivalent to $|U_{\text{gen}}| = j$. Since $|U| = j$ holds and U has U_{gen} as a direct summand, this is equivalent to $U_{\text{gen}} \simeq U$. \square

Now we are ready to prove two key results. The first one is the following.

Theorem 3.10. *Let A be a finite dimensional algebra over a field k and $0 \leq j \leq |A|$. Then we have bijections*

$$(-)_{\min} : 2\text{-psilt}_j A \simeq 2\text{-silt}_j A,$$

whose inverse is given by $(-)_{\text{gen}}$.

Proof. For each $U \in 2\text{-psilt}_j A$, we have $U_{\min} \in 2\text{-silt}_j A$ by (3.3). Thus the map $(-)_{\min} : 2\text{-psilt}_j A \rightarrow 2\text{-silt}_j A$ is well-defined. Moreover, $(U_{\min})_{\text{gen}} \simeq U_{\text{gen}} \simeq U$ holds by Lemma 3.8 and (3.4).

For each $T \in 2\text{-silt}_j A$, let $U := T_{\text{gen}}$. Then $U_{\text{gen}} = U$ holds, and moreover $|U| = j$ holds by Lemma 3.6(b). Thus $U \in 2\text{-psilt}_j A$ holds by (3.4), and the map $(-)_{\text{gen}} : 2\text{-silt}_j A \rightarrow 2\text{-psilt}_j A$ is well-defined. By Lemma 3.8, we have $(T_{\text{gen}})_{\min} \simeq T_{\min} = T$. Thus the assertion follows. \square

The second one is the following.

Theorem 3.11. *Let A be a finite dimensional algebra over a field k and $0 \leq j \leq |A|$. Then we have a bijection*

$$2\text{-psilt}^j A \simeq \bigsqcup_{i=0}^j \{(V, W) \in 2\text{-psilt}_i A \times 2\text{-psilt}^{j-i} A \mid W \text{ is a direct summand of } V_{\min}/V\} \quad (3.5)$$

given by $U \mapsto (U_{\text{gen}}, U/U_{\text{gen}})$, and the converse is given by $(V, W) \mapsto V \oplus W$.

Proof. For $U \in 2\text{-psilt}^j A$, let $(V, W) := (U_{\text{gen}}, U/U_{\text{gen}})$ and $i := |U_{\text{gen}}|$. Then $0 \leq i \leq j$ and $V_{\text{gen}} = V$ hold, and hence $V \in 2\text{-psilt}_i A$. Clearly $W \in 2\text{-psilt}^{j-i} A$ holds. Since $V_{\min} = (U_{\text{gen}})_{\min} \simeq U_{\min}$ holds by Lemma 3.8, $V_{\min}/V \simeq U_{\min}/U_{\text{gen}}$ has $W = U/U_{\text{gen}}$ as a direct summand. Thus the map $U \mapsto (U_{\text{gen}}, U/U_{\text{gen}})$ is well-defined. It is injective since $U \simeq U_{\text{gen}} \oplus (U/U_{\text{gen}})$ holds.

To prove surjectivity, take $(V, W) \in 2\text{-psilt}_i A \times 2\text{-psilt}^{j-i} A$ such that W is a direct summand of V_{\min}/V , and let $U := V \oplus W \in 2\text{-psilt}^j A$. Since $\mathcal{T}_V \subseteq \mathcal{T}_U \subseteq \mathcal{T}_{V_{\min}} = \mathcal{T}_V$ holds, we have $\mathcal{T}_V = \mathcal{T}_U$. Since $V_{\text{gen}} = V$ holds by (3.4), we have $U_{\text{gen}} \simeq V$ and hence $(U_{\text{gen}}, U/U_{\text{gen}}) \simeq (V, W)$. \square

We need the following preparation on semibricks.

Definition 3.12. Let S be a semibrick of A .

- (a) We call S *left-finite* if the smallest torsion class containing S is functorially finite. We denote by $\text{f}_L\text{-sbrick} A$ the set of isomorphism classes of left-finite semibricks of A .
- (b) We call S *right-finite* if the smallest torsionfree class containing S is functorially finite. We denote by $\text{f}_R\text{-sbrick} A$ the set of isomorphism classes of right-finite semibricks of A .

If A is g -finite, then all semibricks of A are left-finite and right-finite by Proposition 2.10.

Proposition 3.13. [As1, Theorem 2.3] *Let A be a finite dimensional algebra over a field k with $n := |A|$. Then we have the following bijections, where $\nu = - \otimes_A DA : \text{K}^b(\text{proj } A) \simeq \text{K}^b(\text{inj } A)$.*

$$\begin{aligned} 2\text{-silt}_j A &\simeq \text{f}_L\text{-sbrick}_j A, & T &\mapsto H^0(T) / \text{rad}_{\text{End}_A(H^0(T))} H^0(T). \\ 2\text{-silt}_j A &\simeq \text{f}_R\text{-sbrick}_{n-j} A, & T &\mapsto \text{soc}_{\text{End}_A(H^{-1}(\nu T))} H^{-1}(\nu T). \end{aligned}$$

We are ready to prove Theorem 3.4.

Proof of Theorem 3.4. By Theorem 3.11, $f_{j-1} = \#2\text{-psilt}^j A$ is equal to the cardinality of the right-hand side of (3.5). For each $V \in 2\text{-psilt}_i A$, there are $\binom{n-i}{j-i}$ choices of W . Thus the equality

$$f_{j-1} = \#2\text{-psilt}^j A = \sum_{i=0}^j \binom{n-i}{j-i} \#2\text{-psilt}_i A$$

holds. Comparing with (3.2), we obtain

$$h_j = \#2\text{-psilt}_j A,$$

which is equal to $\#2\text{-silt}_j A$ by Theorem 3.10. Finally, $\#2\text{-silt}_j A = \#f_L\text{-sbrick}_j A = \#\text{sbrick}_j A$ holds by Proposition 3.13. \square

Since the g -simplicial complex $\Delta(A)$ is a simplicial sphere [DIJ], Dehn-Sommerville equations

$$h_j = h_{n-j}$$

are satisfied [V, Theorem 6.8.8] (see also [Zi, Theorem 8.21]). Our next result categorifies these equations by giving a symmetry of the set $2\text{-silt}A$.

Theorem 3.14. *Let A be a finite dimensional algebra over a field k which is g -finite. For $0 \leq j \leq n := |A|$, there is a canonical bijection*

$$\text{sbrick}_j A \simeq \text{sbrick}_{n-j} A \text{ and } 2\text{-silt}_j A \simeq 2\text{-silt}_{n-j} A.$$

In particular, we have $h_j = h_{n-j}$.

For the case $j = 1$, the bijection $2\text{-silt}_1 A \simeq 2\text{-silt}_{n-1} A$ between join-irreducible elements in $2\text{-silt}A$ and meet-irreducible elements in $2\text{-silt}A$ was shown in [DIRRT] (see also [IRRT]). To prove Theorem 3.14, we need the following result.

Proposition 3.15. [As1] *Let A be a finite dimensional algebra over a field k with $n := |A|$.*

(a) *There exist bijections*

$$H^0 : 2\text{-smc}A \simeq f_L\text{-sbrick}A \text{ and } H^{-1} : 2\text{-smc}A \simeq f_R\text{-sbrick}A$$

such that $S = H^0(S) \oplus H^{-1}(S)[1]$ holds for each $S \in 2\text{-smc}A$.

(b) *The following diagram commutes.*

$$\begin{array}{ccccc} & \text{Prop. 3.13} & 2\text{-silt}A & \xleftarrow{\text{Prop. 3.13}} & \\ & \sim & \downarrow \wr_{\text{Prop. 2.13}} & \sim & \\ f_L\text{-sbrick}A & \xleftarrow{H^0} & 2\text{-smc}A & \xrightarrow{H^{-1}} & f_R\text{-sbrick}A. \end{array}$$

(c) *Assume that $T = T_1 \oplus \cdots \oplus T_n \in 2\text{-silt}A$ and $S = S_1 \oplus \cdots \oplus S_n \in 2\text{-smc}A$ correspond to each other by the bijection in Proposition 3.13. For each $1 \leq i \leq n$, $\mu_i(T) < T$ if and only if $S_i \in \text{mod } A$, and $\mu_i(T) > T$ if and only if $S_i \in (\text{mod } A)[1]$.*

Proof. (a) and (b) are [As1, Theorem 3.3]. To prove (c), it suffices to show the first equivalence. By the left part of the commutative diagram in (b), we have $H^0(S) = H^0(T)/\text{rad}_{\text{End}_A(H^0(T))} H^0(T)$ and hence

$$H^0(S_i) = H^0(T_i) / \sum_{f \in \text{rad}_{\text{mod } A}(H^0(T), H^0(T_i))} \text{Im } f.$$

Thus $S_i \in \text{mod } A$ if and only if $H^0(S_i) \neq 0$ if and only if $H^0(T_i) \notin \mathcal{T}_{T/T_i}$. By Lemma 3.6(a), this is equivalent to $\mu_i(T) < T$. \square

We are ready to prove Theorem 3.14.

Proof of Theorem 3.14. It suffices to give a bijection $\text{sbrick}_j A \simeq \text{sbrick}_{n-j} A$. Since A is g -finite, Proposition 3.15(a) gives bijections $H^0 : 2\text{-smc}A \simeq \text{sbrick}A$ and $H^{-1} : 2\text{-smc}A \simeq \text{sbrick}A$ such that $S = H^0(S) \oplus H^{-1}(S)[1]$ for each $S \in 2\text{-smc}A$. Since $|H^0(S)| + |H^{-1}(S)| = |S| = n$ holds, they give a bijection $\text{sbrick}_j A \simeq \text{sbrick}_{n-j} A$ for each j . \square

Using unimodality results of h -vectors in combinatorics, we obtain the following result as an application. It will be an interesting question if there is a direct proof using tilting theory.

Corollary 3.16. *Let A be a finite dimensional algebra over a field k which is g -finite, and $n := |A|$. Then we have*

$$\begin{aligned} \#sbrick_1 A &\leq \#sbrick_2 A \leq \cdots \leq \#sbrick_{\lfloor \frac{n}{2} \rfloor - 1} A \leq \#sbrick_{\lfloor \frac{n}{2} \rfloor} A, \\ \#sbrick_{\lceil \frac{n}{2} \rceil} A &\geq \#sbrick_{\lceil \frac{n}{2} \rceil + 1} A \geq \cdots \geq \#sbrick_{n-1} A \geq \#sbrick_n A. \end{aligned}$$

Proof. The unimodality of h -vectors was originally proved for boundary complexes of simplicial polytopes, and this is generalized for simplicial spheres (see [Adi]). Since $\Delta(A)$ gives a simplicial sphere (see Definition ?? or [DIJ, Theorem 5.4]), the result follows from Theorem 3.4. \square

There are several natural problems [Pe] in view of Corollary 3.16 and the fact that $\Delta(A)$ is flag (Proposition 3.2(c)). The γ -vector $(\gamma_0, \dots, \gamma_{\lfloor \frac{n}{2} \rfloor})$ is defined by the equality

$$h(x) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_i x^i (1+x)^{n-2i}.$$

Problem 3.17. *Let A be a finite dimensional algebra over a field k which is g -finite, and let $h(x)$ be the h -polynomial of $\Delta(A)$.*

- (a) *Is $h(x)$ real-rooted (that is, all roots are real numbers)?*
- (b) *Is $h(x)$ log-concave (that is, $h_i^2 \geq h_{i-1}h_{i+1}$ holds for each i)?*
- (c) *Is $h(x)$ γ -nonnegative (that is, $\gamma_i \geq 0$ for each i)?*

Gal's conjecture asks if each flag simplicial complex is γ -nonnegative. The following implications are known.

$$\begin{array}{ccc} \text{real-rooted} & \implies & \text{log-concave} \\ \Downarrow & & \Downarrow \\ \gamma\text{-nonnegative} & \implies & \text{unimodal} \end{array}$$

4. g -FANS

4.1. Definition and basic properties. We introduce the g -fan of a finite dimensional algebra. Let A be a finite dimensional algebra over a field k . Let $K_0(\text{proj } A)$ be the Grothendieck group of $K^b(\text{proj } A)$ and

$$K_0(\text{proj } A)_{\mathbb{R}} := K_0(\text{proj } A) \otimes_{\mathbb{Z}} \mathbb{R} \simeq \mathbb{R}^{|A|}.$$

The g -simplicial complex has a canonical geometric realization in the real Grothendieck group $K_0(\text{proj } A)_{\mathbb{R}}$ [DIJ]. Now we introduce its fan version.

Definition 4.1. For $T = T_1 \oplus \cdots \oplus T_j \in 2\text{-psilt } A$ with indecomposable T_i , let

$$C(T) := \left\{ \sum_{i=1}^j a_i [T_i] \mid a_1, \dots, a_j \geq 0 \right\} \subset K_0(\text{proj } A)_{\mathbb{R}}.$$

We call the set

$$\Sigma(A) := \{C(T) \mid T \in 2\text{-psilt } A\}$$

of cones the g -fan of A .

Notice that $\Sigma(A)$ can be an infinite set. For each $0 \leq i \leq |A|$, the subset $\Sigma_i(A)$ of cones of dimension i is given by

$$\Sigma_i(A) = \{C(U) \mid U \in 2\text{-psilt}^i A\}.$$


We give the following basic properties of g -fans.

Proposition 4.2. *Let A be a finite dimensional algebra over a field k and $n := |A|$.*

- (a) *$\Sigma(A)$ is a nonsingular fan in $K_0(\text{proj } A)_{\mathbb{R}}$.*
- (b) *Any cone in $\Sigma(A)$ is a face of a cone of dimension n .*
- (c) *Any cone in $\Sigma(A)$ of dimension $n-1$ is a face of precisely two cones of dimension n .*
- (d) *A is g -finite (or equivalently, $\Sigma(A)$ is finite) if and only if $\Sigma(A)$ is complete.*

Proof. For (a), (b) and (c), see [DIJ] and Proposition 3.2. (d) is [As2, Theorem 4.7]. \square

Example 4.3. [Ka, Proposition 6.1] For positive integers $\ell \geq 1, m \geq 1$, we define an algebra $A := kQ/I$ as follows. The quiver Q is the following



The ideal I of kQ is generated by the following elements for all possible i, j :

$$a_i a_j \ (i - j \neq 1), \ b_i b_j \ (i - j \neq 1), \ a_i b_j \text{ and } b_i a_j.$$

Then $\text{Hase}(2\text{-silt}A_{\ell,m})$ is

$$A = P^{(0)} \oplus Q^{(0)} \begin{array}{c} \nearrow P^{(0)} \oplus P^{(1)} \longrightarrow P^{(1)} \oplus P^{(2)} \longrightarrow \dots \longrightarrow P^{(\ell-1)} \oplus P^{(\ell)} \\ \searrow Q^{(0)} \oplus Q^{(1)} \longrightarrow Q^{(1)} \oplus Q^{(2)} \longrightarrow \dots \longrightarrow Q^{(m-1)} \oplus Q^{(m)} \end{array} \begin{array}{c} \nearrow A[1] = P^{(\ell)} \oplus Q^{(m)} \\ \searrow \end{array}$$


with $P^{(0)} = P_1 := e_1 A$, $P^{(\ell)} = P_1[1]$, $Q^{(0)} = P_2 := e_2 A$ and $Q^{(m)} = P_2[1]$. Since

$$P^{(i)} = [P_1] - i[P_2] \ (0 \leq i \leq \ell - 1) \text{ and } Q^{(j)} = [P_2] - j[P_1] \ (0 \leq j \leq m - 1),$$

the g -fan $\Sigma(A)$ consists of

$$\begin{aligned} & \text{cone}\{[P_1], [P_2]\}, \text{cone}\{-[P_1], -[P_2]\}, \\ & \text{cone}\{[P_1] - (i-1)[P_2], [P_1] - i[P_2]\} \ (1 \leq i \leq \ell - 1), \text{cone}\{[P_1] - (\ell-1)[P_2], -[P_2]\}, \\ & \text{cone}\{[P_2] - (j-1)[P_1], [P_2] - j[P_1]\} \ (1 \leq j \leq m - 1), \text{cone}\{[P_2] - (m-1)[P_1], -[P_1]\}. \end{aligned}$$


For example, if $(\ell, m) = (4, 5)$, then $\Sigma(A)$ is



Example 4.4. Consider a finite dimensional algebra A given by

$$Q = \left[\begin{array}{c} 1 \\ a \downarrow \\ 2 \end{array} \begin{array}{c} \nearrow d \\ \searrow b \end{array} 3 \begin{array}{c} \curvearrowright c \end{array} \right] \text{ and } A := kQ / \langle c^2, bd, abcd, cdab - dabc \rangle$$

which is a Brauer graph algebra (see Example 9.14). Then $A = \begin{smallmatrix} 1 \\ 2 \\ 3 \\ 1 \end{smallmatrix} \oplus \begin{smallmatrix} 2 \\ 3 \\ 3 \\ 1 \end{smallmatrix} \oplus \begin{smallmatrix} 3 \\ 2 \\ 3 \\ 1 \end{smallmatrix}$, and $\Sigma(A)$ is given by the following.



The following general results give canonical isomorphisms between the g -fans of two algebras.

Proposition 4.5. *Let A be a finite dimensional algebra over a field k .*

(a) *The isomorphism $-\operatorname{Hom}_A(-, A) : K_0(\operatorname{proj} A) \simeq K_0(\operatorname{proj} A^{\operatorname{op}})$ gives an isomorphism of g -fans:*

$$\Sigma(A) \simeq \Sigma(A^{\operatorname{op}}).$$

(b) *Let I be an ideal of A which annihilates all bricks of A . Then the isomorphism $-\otimes_A (A/I) : K_0(\operatorname{proj} A) \simeq K_0(\operatorname{proj} A/I)$ gives an isomorphism of g -fans:*

$$\Sigma(A) \simeq \Sigma(A/I).$$

For example, if I is generated by a set of central nilpotent elements of A , then we have the isomorphism of g -fans above.

(c) *Let K/k be a field extension such that the functor $K \otimes_k - : \operatorname{proj} A \rightarrow \operatorname{proj} K \otimes_k A$ preserves the indecomposability. Then the isomorphism $K_0(\operatorname{proj} A) \simeq K_0(\operatorname{proj} K \otimes_k A)$ gives an isomorphism of g -fans:*

$$\Sigma(A) \simeq \Sigma(K \otimes_k A).$$

Proof. (a) We have a duality $\mathbf{R}\operatorname{Hom}_A(-, A)[1] : K^b(\operatorname{proj} A) \simeq K^b(\operatorname{proj} A^{\operatorname{op}})$, which gives the isomorphism $-\operatorname{Hom}_A(-, A) : K_0(\operatorname{proj} A) \simeq K_0(\operatorname{proj} A^{\operatorname{op}})$ and bijections $2\text{-silt} A \simeq 2\text{-silt} A^{\operatorname{op}}$ and $2\text{-psilt} A \simeq 2\text{-psilt} A^{\operatorname{op}}$. Thus the assertion follows.

(b) The assertion follows from [DIRRT] and [EJR].

(c) The assertion follows from [IK]. □

One of the importance of the g -fan is that it gives an explicit description of torsion classes given by stability conditions. For each $\theta \in K_0(\operatorname{proj} A)_{\mathbb{R}}$, let

$$\begin{aligned} \mathcal{T}_{\theta} &:= \{X \in \operatorname{mod} A \mid \theta(X') > 0 \text{ for all factor modules } X' \neq 0 \text{ of } X\}, \\ \overline{\mathcal{T}}_{\theta} &:= \{X \in \operatorname{mod} A \mid \theta(X') \geq 0 \text{ for all factor modules } X' \text{ of } X\}, \\ \mathcal{F}_{\theta} &:= \{X \in \operatorname{mod} A \mid \theta(X') < 0 \text{ for all submodules } X' \neq 0 \text{ of } X\}, \\ \overline{\mathcal{F}}_{\theta} &:= \{X \in \operatorname{mod} A \mid \theta(X') \leq 0 \text{ for all submodules } X' \text{ of } X\}, \\ \mathcal{W}_{\theta} &:= \overline{\mathcal{T}}_{\theta} \cap \overline{\mathcal{F}}_{\theta}. \end{aligned}$$

The following properties are elementary.

Definition-Proposition 4.6. [BKT, Ki] *Let $\theta \in K_0(\operatorname{proj} A)_{\mathbb{R}}$.*

(a) *$(\mathcal{T}_{\theta}, \overline{\mathcal{F}}_{\theta})$ and $(\overline{\mathcal{T}}_{\theta}, \mathcal{F}_{\theta})$ are torsion pairs in $\operatorname{mod} A$ satisfying $\mathcal{T}_{\theta} \subseteq \overline{\mathcal{T}}_{\theta}$ and $\mathcal{F}_{\theta} \subseteq \overline{\mathcal{F}}_{\theta}$. In particular, for $\mathbf{t}_{\theta} := \mathbf{t}_{\mathcal{T}_{\theta}}$, $\overline{\mathbf{t}}_{\theta} := \mathbf{t}_{\overline{\mathcal{T}}_{\theta}}$, $\mathbf{f}_{\theta} := \mathbf{f}_{\mathcal{F}_{\theta}}$, $\overline{\mathbf{f}}_{\theta} := \mathbf{f}_{\overline{\mathcal{F}}_{\theta}}$ in (2.2), each $X \in \operatorname{mod} A$ admits exact sequences*

$$\begin{aligned} 0 \rightarrow \mathbf{t}_{\theta} X \rightarrow X \rightarrow \overline{\mathbf{f}}_{\theta} X \rightarrow 0 \text{ with } \mathbf{t}_{\theta} X \in \mathcal{T}_{\theta} \text{ and } \overline{\mathbf{f}}_{\theta} X \in \overline{\mathcal{F}}_{\theta}, \\ 0 \rightarrow \overline{\mathbf{t}}_{\theta} X \rightarrow X \rightarrow \mathbf{f}_{\theta} X \rightarrow 0 \text{ with } \overline{\mathbf{t}}_{\theta} X \in \overline{\mathcal{T}}_{\theta} \text{ and } \mathbf{f}_{\theta} X \in \mathcal{F}_{\theta}. \end{aligned}$$

(b) \mathcal{W}_θ is a wide subcategory of $\text{mod } A$. Moreover, each $X \in \text{mod } A$ admits a filtration

$$0 \subseteq t_\theta X \subseteq \bar{t}_\theta X \subseteq X \quad \text{such that} \quad w_\theta X := \bar{t}_\theta X / t_\theta X \in \mathcal{W}_\theta.$$

For $T = T_1 \oplus \cdots \oplus T_j \in 2\text{-psilt } A$ with indecomposable T_i , let

$$C^+(T) := \left\{ \sum_{i=1}^j a_i [T_i] \mid a_1, \dots, a_j > 0 \right\} \subset C(T).$$

We have the following descriptions of torsion pairs given by stability conditions in terms of 2-term presilting complexes (see Definition-Propositions 2.8 and 3.7 for functors f_U , w_U and so on). We refer to [As2, AsI] for more explicit results.

Proposition 4.7. [Y1, Proposition 3.3][BST, Proposition 3.27] *Let $U \in 2\text{-psilt } A$ and $\theta \in C^+(U)$. Then we have*

$$\mathcal{T}_\theta = \mathcal{T}_U, \quad \overline{\mathcal{T}}_\theta = \overline{\mathcal{T}}_U, \quad \mathcal{F}_\theta = \mathcal{F}_U, \quad \overline{\mathcal{F}}_\theta = \overline{\mathcal{F}}_U \quad \text{and} \quad \mathcal{W}_\theta = \mathcal{W}_U.$$

Therefore we have

$$t_\theta = t_U, \quad \bar{t}_\theta = \bar{t}_U, \quad f_\theta = f_U, \quad \bar{f}_\theta = \bar{f}_U, \quad \text{and} \quad w_\theta = w_U.$$

To explain a remarkable property of g -fans, we introduce the following notion.

Definition 4.8. For a fan Σ in \mathbb{R}^d and $\sigma \in \Sigma$, we define the *reduction* of Σ at σ by

$$\Sigma/\sigma := \{\pi(\tau) \mid \tau \in \Sigma, \sigma \subseteq \tau\},$$

where $\pi : \mathbb{R}^d \rightarrow \mathbb{R}^d/\mathbb{R}\sigma$ is a natural projection. Then Σ/σ is a fan in $\mathbb{R}^d/\mathbb{R}\sigma$.

This reduction process of fans corresponds to Jasso reduction given in Definition 3.7(d), see also [As2, Theorem 4.5].

Theorem 4.9. *Let A be a finite dimensional algebra over a field k , and $U \in 2\text{-psilt } A$. For the Bongartz completion U_{\max} of U , let $B := \text{End}_{K^b(\text{proj } A)}(U_{\max})/[U]$.*

(a) *There exists a triangle functor $K^b(\text{proj } A) \rightarrow K^b(\text{proj } B)$ which sends U_{\max} to B and gives an isomorphism $K_0(\text{proj } A)/K_0(\text{add } U) \simeq K_0(\text{proj } B)$ and bijections*

$$\{T \in 2\text{-silt } A \mid U \in \text{add } T\} \simeq 2\text{-silt } B \quad \text{and} \quad \{T \in 2\text{-psilt } A \mid U \in \text{add } T\} \simeq 2\text{-psilt } B.$$

(b) *The isomorphism $K_0(\text{proj } A)_{\mathbb{R}}/K_0(\text{add } U)_{\mathbb{R}} \simeq K_0(\text{proj } B)_{\mathbb{R}}$ gives an isomorphism of fans*

$$\Sigma(A)/C(U) \simeq \Sigma(B).$$

Proof. (a) Let $\mathcal{T} := K^b(\text{proj } A)/\text{thick } U$ be the Verdier quotient, $\pi : K^b(\text{proj } A) \rightarrow \mathcal{T}$ the canonical functor, and $V := \pi(U_{\max}) \in \mathcal{T}$. Then we have an isomorphism $\pi : K_0(\text{proj } A)/K_0(\text{add } U) \simeq K_0(\mathcal{T})$. By [IY, Theorems 3.1, 3.7, Corollary 3.8], \mathcal{T} is a k -linear Hom-finite Krull-Schmidt triangulated category such that π gives an isomorphism

$$\{T \in \text{silt } A \mid U \in \text{add } T\} \simeq \text{silt } \mathcal{T}$$

of posets, a bijection

$$\{T \in \text{psilt } A \mid U \in \text{add } T\} \simeq \text{psilt } \mathcal{T}$$

and an isomorphism

$$B \simeq \text{End}_{\mathcal{T}}(V)$$

of k -algebras. Since $\pi(A) \simeq V$ holds by [J, Propositions 4.10], π gives bijections

$$\{T \in 2\text{-silt } A \mid U \in \text{add } T\} \simeq 2_V\text{-silt } \mathcal{T} \quad \text{and} \quad \{T \in 2\text{-psilt } A \mid U \in \text{add } T\} \simeq 2_V\text{-psilt } \mathcal{T}.$$

On the other hand, \mathcal{T} is algebraic [Ke2, D]. Applying Proposition 2.14 to \mathcal{T} and V , there is a triangle functor $F : \mathcal{T} \rightarrow K^b(\text{proj } B)$ which sends V to B and gives an isomorphism $K_0(\mathcal{T}) \simeq K_0(\text{proj } B)$ and bijections

$$2_V\text{-silt } \mathcal{T} \simeq 2\text{-silt } B \quad \text{and} \quad 2_V\text{-psilt } \mathcal{T} \simeq 2\text{-psilt } B.$$


Thus the composition $F \circ \pi : K^b(\text{proj } A) \rightarrow K^b(\text{proj } B)$ gives the desired triangle functor.

(b) For $\ell := |U| \leq j \leq |A|$, the triangle functors π and F give bijections

$$\{T \in 2\text{-psilt}^j A \mid U \in \text{add } T\} \simeq 2_V\text{-psilt}^{j-\ell} \mathcal{T} \simeq 2\text{-psilt}^{j-\ell} B$$

such that $F \circ \pi(C(T)) = C(F \circ \pi(T))$ as cones in $K_0(\text{proj } B)_{\mathbb{R}}$. Thus the assertion follows. \square

Example 4.10. Let A be the algebra in Example 4.4. For $\sigma := C(e_1 A)$, the cones $\tau \in \Sigma(A)$ satisfying $\sigma \subset \tau$ is shown by the left picture below. Moreover, $\Sigma(A)/\sigma$ is the fan shown by the right picture below.



4.2. Idempotents and g -fans. In this subsection, we observe that g -fans of finite dimensional algebras are rather special.

Definition 4.11. A *sign-coherent fan* is a pair (Σ, σ_+) satisfying the following conditions.

- (a) Σ is a nonsingular fan in \mathbb{R}^d .
- (b) $\sigma_+, -\sigma_+ \in \Sigma_d$.
- (c) Take a \mathbb{Z} -basis e_1, \dots, e_d of \mathbb{Z}^d such that $\sigma_+ = \text{cone}\{e_i \mid 1 \leq i \leq d\}$, and denote the orthant corresponding to $\epsilon \in \{\pm 1\}^d$ by

$$\mathbb{R}_\epsilon^d := \text{cone}\{\epsilon(1)e_1, \dots, \epsilon(d)e_d\} = \{x_1 e_1 + \dots + x_d e_d \mid \epsilon(i)x_i \geq 0 \text{ for each } 1 \leq i \leq d\}.$$

Then for each $\sigma \in \Sigma$, there exists $\epsilon \in \{\pm 1\}^d$ such that $\sigma \subseteq \mathbb{R}_\epsilon^d$.

The following property of g -fans is basic.

Proposition 4.12. For a finite dimensional algebra A over a field k , $(\Sigma(A), C(A))$ is a sign-coherent fan. In this case, the isomorphism classes of indecomposable projective A -modules gives the \mathbb{Z} -basis which generate $C(A)$ as a cone.

Proof. The conditions (a) and (b) are clear. The condition (c) is the sign-coherence of g -vectors [AIR]. \square

Now we show that the g -fans form a very special class of sign-coherent fans.

Definition 4.13. An *ordered fan* is a sign-coherent fan (Σ, σ_+) satisfying the following conditions.

- (a) For each adjacent cones $\sigma, \tau \in \Sigma_d$, a normal vector of $\sigma \cap \tau$ belongs to the interior of σ_+ .
- (b) There exists a partial order \leq on Σ_d such that (Σ_d, \leq) has the maximum element σ_+ and the minimum element $-\sigma_+$.
- (c) For any $\sigma, \tau \in \Sigma_d$, the following conditions are equivalent.
 - There is an arrow $\sigma \rightarrow \tau$ in the Hasse quiver.
 - σ and τ are adjacent. Moreover, consider the hyperplane $H := \mathbb{R}(\sigma \cap \tau) \subset \mathbb{R}^d$. Then σ and σ_+ belong to the same connected component of $\mathbb{R}^d \setminus H$.

For $d = 2$, it is easy to check that any sign-coherent fan is ordered. For $d \geq 3$, there are many sign-coherent fans which are not ordered.

Notice that, if A is g -finite, then the partial order \leq is uniquely determined by the condition (c). Therefore the following property of $\Sigma(A)$ is important since it claims that the partial order on $2\text{-silt } A$ can be recovered from $\Sigma(A)$ if A is g -finite.

Proposition 4.14. For a finite dimensional algebra A over a field k , $(\Sigma(A), C(A))$ is an ordered fan.

Proof. This is shown in [DIJ, Lemma 6.9]. \square

Now we explain idempotent reductions of g -fans. We prepare the following general notion.

Definition 4.15. Let (Σ, σ_+) be a sign-coherent fan, and e_1, \dots, e_d the basis in Definition 4.11(c). For each subset $I \subset \{1, \dots, d\}$, consider the subspace

$$\mathbb{R}^I = \bigoplus_{i \in I} \mathbb{R}e_i \subset \mathbb{R}^d.$$

Then the subset

$$\Sigma^I := \{\sigma \in \Sigma \mid \sigma \subset \mathbb{R}^I\}$$

is a subfan of Σ thanks to the condition Definition 4.11(c).

Now let A be a basic finite dimensional algebra over a field k , and $1 = e_1 + \dots + e_n$ the orthogonal primitive idempotents. As in Definition 4.15, for each subset $I \subset \{1, \dots, n\}$, we obtain a subspace

$$K_0(\text{proj } A)_{\mathbb{R}}^I := \bigoplus_{i \in I} \mathbb{R}[e_i A] \subset K_0(\text{proj } A)_{\mathbb{R}}$$

and a subfan of $\Sigma(A)$ given by

$$\Sigma^I(A) := \{\sigma \in \Sigma(A) \mid \sigma \subset K_0(\text{proj } A)_{\mathbb{R}}^I\}.$$

On the other hand, we consider the idempotent

$$e = e^I := \sum_{i \in I} e_i \in A$$

and the corresponding subalgebra eAe of A . Then we have a fully faithful functor

$$- \otimes_{eAe} eA : \text{proj } eAe \rightarrow \text{proj } A$$

which induces an isomorphism

$$- \otimes_{eAe} eA : K_0(\text{proj } eAe)_{\mathbb{R}} \simeq K_0(\text{proj } A)_{\mathbb{R}}^I. \quad (4.1)$$

We are ready to state the following result.

Theorem 4.16. *Let A be a finite dimensional algebra over a field k , and $1 = e_1 + \dots + e_n$ the orthogonal primitive idempotents. For each subset $I \subset \{1, \dots, n\}$ and $e := e^I$, the isomorphism (4.1) gives an isomorphism of fans*

$$\Sigma(eAe) \simeq \Sigma^I(A).$$

Proof. For the thick subcategory $\mathbf{K}^b(\text{add } eA)$ of $\mathbf{K}^b(\text{proj } A)$, we have a triangle equivalence


$$- \otimes_{eAe} eA : \mathbf{K}^b(\text{proj } eAe) \simeq \mathbf{K}^b(\text{add } eA) \subset \mathbf{K}^b(\text{proj } A).$$

Therefore we have a bijection

$$- \otimes_{eAe} eA : 2\text{-psilt}(eAe) \simeq \{T \in 2\text{-psilt } A \mid T^0, T^{-1} \in \text{add } eA\}.$$

Since $K_0(\text{add } eA)_{\mathbb{R}} = K_0(\text{proj } A)_{\mathbb{R}}^I$ holds, the assertion follows immediately. \square

Example 4.17. Let A be the algebra in Example 4.4. Then the subfans $\Sigma^{\{1,2\}}(A)$, $\Sigma^{\{1,3\}}(A)$, $\Sigma^{\{2,3\}}(A)$ are the following.



We show that the product of fans corresponds to the product of algebras.

Definition 4.18. [Fu] Let Σ and Σ' be fans in \mathbb{R}^d and $\mathbb{R}^{d'}$ respectively. We define a *product fan* $\Sigma \times \Sigma'$ in $\mathbb{R}^{d+d'}$ by

$$\Sigma \times \Sigma' := \{\sigma \times \sigma' \mid \sigma \in \Sigma, \sigma' \in \Sigma'\}.$$

We say that Σ is *indecomposable* if it can not be written as a product of two fans.

The following result shows that the decomposition of g -fans precisely corresponds to the decomposition of algebras.

Theorem 4.19. Let A be a finite dimensional algebra over a field k .

(a) If $A = A_1 \times \cdots \times A_\ell$ for a finite dimensional algebra A_i , then we have

$$\Sigma(A) = \Sigma(A_1) \times \cdots \times \Sigma(A_\ell).$$

(b) In (a), assume that each A_i is ring-indecomposable. Then, for each decomposition $\Sigma(A) = \Sigma^1 \times \cdots \times \Sigma^m$, there exists a decomposition $\{1, \dots, \ell\} = \bigsqcup_{j=1}^m I_j$ such that $\Sigma^j = \Sigma(\prod_{i \in I_j} A_i)$ for each $1 \leq j \leq m$.

(c) If A is ring-indecomposable, then the fan $\Sigma(A)$ is indecomposable.

Proof of (a). There is a bijection

$$2\text{-psilt} A_1 \times \cdots \times 2\text{-psilt} A_\ell \simeq 2\text{-psilt}(A_1 \times \cdots \times A_\ell) \text{ given by } (T_1, \dots, T_\ell) \mapsto T_1 \oplus \cdots \oplus T_\ell.$$

Since $C(T_1 \oplus \cdots \oplus T_\ell) = C(T_1) \times \cdots \times C(T_\ell)$, we obtain the desired equation. \square

To prove Theorem 4.19(b)(c), we need the following observation.

Lemma 4.20. Let $e \in A$ be an idempotent and $f := 1 - e$.

(a) We have an injective map

$$2\text{-psilt}^1(eAe) \sqcup 2\text{-psilt}^1(fAf) \subset 2\text{-psilt}^1 A$$

sending $P \in 2\text{-psilt}^1(eAe)$ to $P \otimes_{eAe} eA$ and $Q \in 2\text{-psilt}^1(fAf)$ to $Q \otimes_{fAf} fA$.

(b) The equality holds in (a) if and only if e is a central idempotent.

Proof. (a) Immediate from Theorem 4.16.

(b) It suffices to prove “only if” part. Assume that the equality holds. If e is not central, then at least one of $\text{Hom}_A(eA, fA)$ and $\text{Hom}_A(fA, eA)$ is non-zero. Without loss of generality, assume $\text{Hom}_A(eA, fA) \neq 0$. Take an indecomposable direct summand $e_i A$ of eA such that $\text{Hom}_A(e_i A, fA) \neq 0$, and take a minimal left (add fA)-approximation of $e_i A$

$$e_i A \rightarrow Q \rightarrow X \rightarrow e_i A[1].$$

Then X is an indecomposable direct summand of $\mu_{eA}^-(A) \in 2\text{-silt} A$, and thus $X \in 2\text{-psilt}^1 A$. Moreover Q is non-zero by our choice, and hence X is not contained in the image of the map in (a). Thus e has to be central. \square

We are ready to prove Theorem 4.19(b)(c).

Proof of Theorem 4.19(b)(c). It suffices to prove (b). Without loss of generality, we can assume that A is basic. Let $1 = e_1 + \cdots + e_n$ be a primitive orthogonal idempotents and $P_i := e_i A$. By definition, there exist $\sigma^j \in \Sigma^j$ for each $1 \leq j \leq m$ such that

$$C(A) = \sigma^1 \times \cdots \times \sigma^m.$$

Since $C(A) = \text{cone}\{[P_1], \dots, [P_n]\}$, there exists a decomposition $\{1, \dots, n\} = \bigsqcup_{j=1}^m J_j$ such that

$$\sigma^j = \text{cone}\{[P_i] \mid i \in J_j\}$$

for each $1 \leq j \leq m$. Let

$$e^j := \sum_{i \in J_j} e_i \in A.$$

Then Σ^j is a fan in $K_0(\text{proj } A)_{\mathbb{R}}^{J_j} = K_0(\text{proj } e^j A e^j)_{\mathbb{R}}$, and hence each ray of $\Sigma(A)$ belongs to $K_0(\text{proj } e^j A e^j)$ for some $1 \leq j \leq m$. Thus the map

$$\bigsqcup_{j=1}^m 2\text{-psilt}^1(e^j A e^j) \rightarrow 2\text{-psilt}^1 A$$

sending $P \in 2\text{-psilt}^1(e^j A e^j)$ to $P \otimes_{e^j A e^j} e^j A$ is bijective. By Lemma 4.20, e^j is a central idempotent of A . Thus there exists a decomposition $\{1, \dots, \ell\} = \bigsqcup_{j=1}^m I_j$ such that $A e^j = \prod_{i \in I_j} A_i$, and the assertion follows. \square

Next we explain the sign decomposition of g -fans [Ao]. We prepare the following general notion.

Definition 4.21. Let (Σ, σ_+) be a sign-coherent fan in \mathbb{R}^d and $\epsilon \in \{\pm 1\}^d$. Consider the basis e_1, \dots, e_d of \mathbb{R}^d and the orthant \mathbb{R}_ϵ^d given in Definition 4.11(c). Define a subfan of Σ by

$$\Sigma_\epsilon := \{\sigma \in \Sigma \mid \sigma \subset \mathbb{R}_\epsilon^d\}.$$

Thanks to the condition Definition 4.11(c), we have $\Sigma = \bigcup_{\epsilon \in \{\pm 1\}^d} \Sigma_\epsilon$.

Now let A be a basic finite dimensional algebra over a field k with $|A| = n$, and $1 = e_1 + \cdots + e_n$ the orthogonal primitive idempotents. For $\epsilon \in \{\pm 1\}^n$, as in Definition 4.21, we obtain an orthant

$$K_0(\text{proj } A)_{\epsilon, \mathbb{R}} := \text{cone}(\epsilon_i [e_i A] \mid i \in \{1, \dots, n\})$$

and a subfan of $\Sigma(A)$ given by

$$\Sigma_\epsilon(A) := \{\sigma \in \Sigma(A) \mid \sigma \subset K_0(\text{proj } A)_{\epsilon, \mathbb{R}}\}.$$

We can describe the fan $\Sigma_\epsilon(A)$ by a simpler algebra defined as follows.

Definition 4.22. For $\epsilon \in \{\pm 1\}^n$, let

$$e_\epsilon^+ := \sum_{\epsilon_i=1} e_i \quad \text{and} \quad e_\epsilon^- := \sum_{\epsilon_i=-1} e_i.$$

We denote by A_ϵ the subalgebra of A given by

$$A_\epsilon := \begin{bmatrix} e_\epsilon^+ A e_\epsilon^+ & e_\epsilon^+ A e_\epsilon^- \\ 0 & e_\epsilon^- A e_\epsilon^- \end{bmatrix}.$$

The functor $- \otimes_{A_\epsilon} A : \text{proj } A_\epsilon \rightarrow \text{proj } A$ gives an isomorphism

$$- \otimes_{A_\epsilon} A : K_0(\text{proj } A_\epsilon)_{\mathbb{R}} \simeq K_0(\text{proj } A)_{\mathbb{R}}. \quad (4.2)$$

and a functor

$$- \otimes_{A_\epsilon} A : K^\epsilon(\text{proj } A_\epsilon) \rightarrow K^\epsilon(\text{proj } A). \quad (4.3)$$

We denote by $K^\epsilon(\text{proj } A)$ the full subcategory of $K^b(\text{proj } A)$ consisting of 2-term complexes of the form $P^{-1} \rightarrow P^0$ with $P^{-1} \in \text{add } e_\epsilon^- A$ and $P^0 \in \text{add } e_\epsilon^+ A$. Let

$$2\text{-psilt}_\epsilon A := 2\text{-psilt } A \cap K^\epsilon(\text{proj } A) \quad \text{and} \quad 2\text{-silt}_\epsilon A := 2\text{-silt } A \cap K^\epsilon(\text{proj } A).$$

Thus the cones in $\Sigma_\epsilon(A)$ are given by the elements in $2\text{-psilt}_\epsilon A$. Now we state main properties of sign decomposition, where the part (b) is a generalization of [Ao, Theorem 4.5].

Theorem 4.23. *Let A be a basic finite dimensional algebra algebra over a field k , and $1 = e_1 + \cdots + e_n$ the orthogonal primitive idempotents. Let $\epsilon \in \{\pm 1\}^n$.*

- (a) [Ao, Proposition 3.2] *The subset $2\text{-silt}_\epsilon A$ of the poset $2\text{-silt} A$ is an interval with a maximal element $(e_+ A)_{\min}$ and a minimal element $(e_- A)_{\max}$.*
- (b) *The functor (4.3) gives bijections $2\text{-psilt}_\epsilon A_\epsilon \simeq 2\text{-psilt}_\epsilon A$ and $2\text{-silt}_\epsilon A_\epsilon \simeq 2\text{-silt}_\epsilon A$, and the isomorphism (4.2) gives an isomorphism of fans*

$$\Sigma_\epsilon(A_\epsilon) \simeq \Sigma_\epsilon(A).$$

Proof. (b) The following properties of the functor $F : K^\epsilon(\text{proj } A_\epsilon) \rightarrow K^\epsilon(\text{proj } A)$ in (4.3) can be checked easily.


- (i) F is full and dense.
- (ii) If $P \in K^\epsilon(\text{proj } A_\epsilon)$ satisfies $F(P) \simeq 0$, then $P \simeq 0$.
- (iii) $\text{Hom}_{K^b(\text{proj } A_\epsilon)}(P, Q[1]) \simeq \text{Hom}_{K^b(\text{proj } A)}(F(P), F(Q)[1])$.

By (i) and (ii), F gives a bijection between the isomorphism classes of indecomposable objects in $K^\epsilon(\text{proj } A_\epsilon)$ and those in $K^\epsilon(\text{proj } A)$. Therefore by (iii), F gives a bijection

$$F : 2\text{-psilt}_\epsilon A_\epsilon \simeq 2\text{-psilt}_\epsilon A.$$

Thus the assertion follows. □

Example 4.24. Let A be the algebra in Example 4.4. Then $\Sigma_\epsilon(A)$ is the following.



We give another condition for two algebras A and B such that $\Sigma_\epsilon(A)$ and $\Sigma_\epsilon(B)$ are isomorphic. For each ideal I of A contained in $\text{rad } A$, the functor $- \otimes_A (A/I) : \text{proj } A \rightarrow \text{proj } A/I$ induces an isomorphism

$$- \otimes_A (A/I) : K_0(\text{proj } A)_\mathbb{R} \simeq K_0(\text{proj } A/I)_\mathbb{R} \quad (4.4)$$

and a functor

$$- \otimes_A (A/I) : K^\epsilon(\text{proj } A) \rightarrow K^\epsilon(\text{proj } A/I). \quad (4.5)$$

We give a sufficient condition for I such that (4.4) induces an isomorphism $\Sigma_\epsilon(A) \simeq \Sigma_\epsilon(A/I)$.

Proposition 4.25. *Let A be a basic finite dimensional algebra over a field k , $1 = e_1 + \cdots + e_n$ the orthogonal primitive idempotents, and $\epsilon \in \{\pm 1\}^n$. Assume that an ideal I of A satisfies*

$$I \subseteq \text{rad } A, \quad e_\epsilon^+ I e_\epsilon^- = 0, \quad (e_\epsilon^+ I e_\epsilon^+)(e_\epsilon^+ A e_\epsilon^-) = 0 \quad \text{and} \quad (e_\epsilon^+ A e_\epsilon^-)(e_\epsilon^- I e_\epsilon^-) = 0.$$

Then the functor (4.5) gives bijections $2\text{-psilt}_\epsilon A \simeq 2\text{-psilt}_\epsilon A/I$ and $2\text{-silt}_\epsilon A \simeq 2\text{-silt}_\epsilon A/I$, and Then the isomorphism (4.4) gives an isomorphism of fans

$$\Sigma_\epsilon(A) \simeq \Sigma_\epsilon(A/I).$$

Proof. By our assumptions on I , the functor $F := - \otimes_A (A/I) : \mathcal{K}^\epsilon(\text{proj } A) \rightarrow \mathcal{K}^\epsilon(\text{proj } A/I)$ satisfies the following conditions

- (i) F is full and dense.
- (ii) If $P \in \mathcal{K}^\epsilon(\text{proj } A)$ satisfies $F(P) \simeq 0$, then $P \simeq 0$.
- (iii) $\text{Hom}_{\mathcal{K}^b(\text{proj } A)}(P, Q[1]) \simeq \text{Hom}_{\mathcal{K}^b(\text{proj } A/I)}(F(P), F(Q)[1])$.

Thus F gives a bijection $F : 2\text{-psilt } A \cap \mathcal{K}^\epsilon(\text{proj } A) \simeq 2\text{-psilt } A/I \cap \mathcal{K}^\epsilon(\text{proj } A/I)$, and the assertion follows. \square

We record the following useful observation.

Example 4.26. In the setting of Definition 4.22, define an ideal I_ϵ of A_ϵ by

$$I_\epsilon := \begin{bmatrix} \text{rad}(e_\epsilon^+ A e_\epsilon^+) \cap \text{Ann}_{e_\epsilon^+ A e_\epsilon^+}(e_\epsilon^+ A e_\epsilon^-) & 0 \\ 0 & \text{rad}(e_\epsilon^- A e_\epsilon^-) \cap \text{Ann}_{e_\epsilon^- A e_\epsilon^-}(e_\epsilon^+ A e_\epsilon^-) \end{bmatrix}.$$

For each ideal I of A_ϵ contained in I_ϵ , we have an isomorphism of fans

$$\Sigma_\epsilon(A) \simeq \Sigma_\epsilon(A_\epsilon) \simeq \Sigma_\epsilon(A_\epsilon/I)$$

by Theorem 4.23 and Proposition 4.25.

Definition 4.27. Let (Σ, σ_+) be a sign-coherent fan in \mathbb{R}^d , and e_1, \dots, e_d the basis of \mathbb{R}^d given in Definition 4.11(c). For $1 \leq i \leq d$ and $\delta \in \{\pm 1\}$, consider a half space

$$\mathbb{R}_{i,\delta}^d := \{x_1 e_1 + \cdots + x_d e_d \in \mathbb{R}^d \mid \delta x_i \geq 0\}$$

and define a subfan of Σ by

$$\Sigma_{i,\delta} := \{\sigma \in \Sigma \mid \sigma \subset \mathbb{R}_{i,\delta}^d\}.$$

Thanks to the condition Definition 4.11(c), we have $\Sigma = \Sigma_{i,+} \cup \Sigma_{i,-}$.

Let A be a basic finite dimensional algebra over a field k with $|A| = n$, and $1 = e_1 + \cdots + e_n$ the orthogonal primitive idempotents. For $1 \leq i \leq d$ and $\delta \in \{\pm 1\}$, as in Definition 4.27, we obtain a half space

$$K_0(\text{proj } A)_{i,\delta,\mathbb{R}} := \{x_1[e_1 A] + \cdots + x_d[e_d A] \in K_0(\text{proj } A)_\mathbb{R} \mid \delta x_i \geq 0\}$$

and a subfan of $\Sigma(A)$ given by

$$\Sigma_{i,\delta}(A) := \{\sigma \in \Sigma(A) \mid \sigma \subset K_0(\text{proj } A)_{i,\delta,\mathbb{R}}\}.$$

For elements $T \geq T'$ in $\text{silt } A$, we consider the interval

$$[T', T] := \{U \in \text{silt } A \mid T \geq U \geq T'\}.$$

The following result gives information how g -fans change under mutation.

Theorem 4.28. *Let A be a basic finite dimensional algebra over a field k with $|A| = n$, and $1 = e_1 + \cdots + e_n$ the orthogonal primitive idempotents. For $1 \leq i \leq n$, let $B := \text{End}_A(\mu_i(A))$. Then there exists a triangle functor $\mathcal{K}^b(\text{proj } A) \rightarrow \mathcal{K}^b(\text{proj } B)$ which sends $\mu_i(A)$ to B and gives an isomorphism $K_0(\text{proj } A) \simeq K_0(\text{proj } B)$, a bijection*

$$[A[1], \mu_i(A)] \simeq [\mu_i(B[1]), B]$$

and an isomorphism of fans

$$\Sigma_{i,-}(A) \simeq \Sigma_{i,+}(B).$$

Proof. Applying Proposition 2.14 to $\mathcal{T} := K^b(\text{proj } A)$ and $T := \mu_i(A)$, we obtain a triangle functor $F : K^b(\text{proj } A) \rightarrow K^b(\text{proj } B)$ satisfying $F(T) = B$ and giving an isomorphism $K_0(\text{proj } A) \simeq K_0(\text{proj } B)$ and bijections

$$2_T\text{-silt } A \simeq 2\text{-silt } B \text{ and } 2_{T\text{-psilt}} A \simeq 2\text{-psilt } B.$$

Since $F(A[1])$ and $F(T[1]) = B[1]$ have the same direct summands except the i -th one, we have $F(A[1]) = \mu_i(B[1])$. Thus the bijection $F : 2_T\text{-silt } A \simeq 2\text{-silt } B$ restricts to the desired bijection

$$[A[1], T] \simeq [\mu_i(B[1]), B].$$

Since the subfan $\Sigma_{i,-}(A)$ (respectively, $\Sigma_{i,+}(B)$) consists of the cones corresponding to the interval $[A[1], T]$ (respectively, $[\mu_i(B[1]), B]$), we obtain the desired isomorphism $\Sigma_{i,-}(A) \simeq \Sigma_{i,+}(B)$ of fans. \square

5. g -POLYTOPES, c -POLYTOPES AND NEWTON POLYTOPES

In this section, we introduce g -polytopes of finite dimensional algebras and characterize when they are convex. We show that convex g -polytopes are reflexive polytopes, and describe the dual polytopes as c -polytopes associated with the set of the 2-term simple minded collections. We also describe Newton polytopes of A -modules by using the g -fan of A .

5.1. Definition and basic properties. With each nonsingular fan, we associate a (not necessarily convex) polytope as follows.

Definition 5.1. Let Σ be a nonsingular fan in \mathbb{R}^d . For each $\sigma \in \Sigma_d$, take a basis v_1, \dots, v_d of \mathbb{Z}^d such that $\sigma = \text{cone}\{v_1, \dots, v_d\}$, and let

$$\sigma_{\leq 1} := \text{conv}\{0, v_1, \dots, v_d\} \subset \mathbb{R}^d.$$

Define a (not necessarily convex) polytope in \mathbb{R}^d by

$$P(\Sigma) := \bigcup_{\sigma \in \Sigma_d} \sigma_{\leq 1}.$$

We say that Σ is *convex* if $P(\Sigma)$ is convex.

Applying this construction to g -fans, we obtain the following notion.

Definition 5.2. Let A be a finite dimensional algebra over a field k .

(a) We call $P(A) := P(\Sigma(A))$ the g -polytope of A . More precisely, for $T = T_1 \oplus \dots \oplus T_\ell \in 2\text{-psilt } A$ with indecomposable T_i , let


$$C_{\leq 1}(T) := \left\{ \sum_{i=1}^{\ell} a_i [T_i] \mid a_i \geq 0, \sum_{i=1}^{\ell} a_i \leq 1 \right\} \subset K_0(\text{proj } A)_{\mathbb{R}}.$$

Then $P(A)$ is defined by

$$P(A) := \bigcup_{T \in 2\text{-silt } A} C_{\leq 1}(T).$$

(b) We say that A is g -convex if the g -polytope $P(A)$ is convex (i.e. $\Sigma(A)$ is convex).

Example 5.3. For integers $\ell, m \geq 1$, let $A = A_{\ell, m}$ be the algebra in Example 4.3. Then A is g -convex if and only if $\ell \leq 3$ and $m \leq 3$. For example, if $(\ell, m) = (4, 5)$, then $P(A)$ is



Now we study the Ehrhart series of A , which is the generating function of the number of 2-term presilting complexes.

Definition 5.4. For each integer $\ell \geq 0$, we denote by

$$2\text{-psilt}_{\oplus}^{\leq \ell} A$$

the set of isomorphism classes of (not necessarily basic) 2-term presilting complexes of A which have at most ℓ indecomposable direct summands. We define the *Ehrhart series* of A by

$$\text{Ehr}_A(x) := 1 + \sum_{\ell \geq 1} \#(2\text{-psilt}_{\oplus}^{\leq \ell} A) x^\ell.$$

Since there is a canonical bijection

$$2\text{-psilt}_{\oplus}^{\leq \ell} A \simeq P(A) \cap \frac{1}{\ell} K_0(\text{proj } A) \quad \text{given by } U \mapsto \frac{[U]}{\ell},$$

the Ehrhart series $\text{Ehr}_A(x)$ of A coincides with the Ehrhart series of the g -polytope $P(A)$ though $P(A)$ is not necessarily convex. We give the following explicit description of the Ehrhart series by using the h -vector.

Theorem 5.5. *Let A be a finite dimensional algebra over a field k which is g -finite, $n := |A|$ and (h_0, \dots, h_n) the h -vector of $\Delta(A)$. Then the Ehrhart series of A is given by*

$$\text{Ehr}_A(x) = \frac{\sum_{i=0}^n h_i x^i}{(1-x)^{n+1}}.$$

In other words, for each $\ell \geq 0$, we have

$$\#2\text{-psilt}_{\oplus}^{\leq \ell} A = \sum_{j=0}^n \binom{n+\ell-j}{n} h_j.$$

Proof. Since $P(A)$ has a unimodular triangulation, the proof of [BR, Theorem 10.3] works (though $P(A)$ is not necessarily convex). \square

To characterize g -convexity, we introduce two conditions below. The first one is combinatorial.

Definition 5.6. Let Σ be a nonsingular fan in \mathbb{R}^d . We call Σ *pairwise positive* if the following condition is satisfied.

- For each two adjacent maximal cones $\sigma, \tau \in \Sigma_d$, take \mathbb{Z} -basis $\{v_1, \dots, v_{d-1}, v_d\}$ and $\{v_1, \dots, v_{d-1}, v'_d\}$ of \mathbb{Z}^d such that $\sigma = \text{cone}\{v_1, \dots, v_{d-1}, v_d\}$ and $\tau = \text{cone}\{v_1, \dots, v_{d-1}, v'_d\}$. Then $v_d + v'_d$ belongs to $\text{cone}\{v_1, \dots, v_{d-1}\}$.

We call Σ *pairwise convex* if the following condition is satisfied.

- Define v_d and v'_d as above. Then $v_d + v'_d$ is either 0, v_i for some $1 \leq i \leq d-1$ or $v_i + v_j$ for some $1 \leq i, j \leq d-1$.

This notion in fact characterizes convexity of $P(\Sigma)$ in the following sense.

Proposition 5.7. *Let Σ be a nonsingular fan in \mathbb{R}^d . Then the following conditions are equivalent.*

- (a) Σ is pairwise convex.
 - (b) For each two adjacent maximal cones σ and τ , the union $\sigma_{\leq 1} \cup \tau_{\leq 1}$ is convex.
- Moreover, if Σ is finite and complete, then the following condition is also equivalent.
- (c) Σ is pairwise positive and $P(\Sigma)$ is convex.

Proof. (a) \Rightarrow (b) This is [AMN, Lemma 2.17].

(b) \Rightarrow (a) Since both $\{v_1, \dots, v_{d-1}, v_d\}$ and $\{v_1, \dots, v_{d-1}, v'_d\}$ are \mathbb{Z} -basis of \mathbb{Z}^d , it follows that $v_d + v'_d$ belongs to $\sum_{i=1}^{d-1} \mathbb{Z}v_i$. Thus $(v_d + v'_d)/2$ belongs to $\sigma_{\leq 1} \cup \tau_{\leq 1}$ since it is convex. Thus

$$v_d + v'_d \in (\sigma \cup \tau) \cap \left(\sum_{i=1}^{d-1} \mathbb{Z}v_i \right) = \sum_{i=1}^{d-1} \mathbb{Z}_{\geq 0} v_i \quad (5.1)$$

holds. Now take an \mathbb{R} -linear form $f : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $f(v_i) = 1$ holds for each $1 \leq i \leq d$. Then the hyperplane $f^{-1}(1)$ contains a facet $\text{conv}\{v_i \mid 1 \leq i \leq d\}$ of $\sigma_{\leq 1}$. Since $\sigma_{\leq 1} \cup \tau_{\leq 1}$ is convex, it is

contained in the half space $\{x \in \mathbb{R}^d \mid f(x) \leq 1\}$. In particular, $f(v_d + v'_d) \leq 2$ holds. Using (5.1), we obtain the assertion.

(a) \Rightarrow (c) Clearly Σ is pairwise positive. Moreover $P(\Sigma)$ is convex by the same argument as in [AMN, Proposition 2.23].

(c) \Rightarrow (a) Since Σ is pairwise positive, we have $v_d + v'_d \in \sum_{i=1}^{d-1} \mathbb{Z}_{\geq 0} v_i$. Using the latter half of the proof of (b) \Rightarrow (a), we obtain that Σ is pairwise convex. \square

Notice that g -fans are always pairwise positive.

Proposition 5.8. *Let A be a finite dimensional algebra over a field k . Then $\Sigma(A)$ is pairwise positive.*

Proof. Let $T = T_1 \oplus \cdots \oplus T_n \in 2\text{-silt} A$ and $T_n \rightarrow U_n \rightarrow T'_n \rightarrow T_n[1]$ be an exchange triangle. Then $[T_n] + [T'_n] = [U_n]$ belongs to $\sum_{i=1}^{n-1} \mathbb{Z}_{\geq 0} [T_i]$. \square

The pairwise convexity of g -fans can be stated as follows.

Definition 5.9. Let A be a finite dimensional algebra over a field k . We say that A is *pairwise g -convex* if the following condition is satisfied.

- For any $T \in 2\text{-silt} A$ and an indecomposable direct summand T_i of T , let

$$T_i \rightarrow U_i \rightarrow T'_i \rightarrow T_i[1],$$

be an exchange triangle. Then U_i has at most two indecomposable direct summands.

Inspired by [H2], we give characterizations of the convexity of g -polytope.

Theorem 5.10. *Let A be a finite dimensional algebra over a field k .*

- (a) *A is g -finite if and only if $P(A)$ contains the origin in its interior.*
- (b) *The following conditions are equivalent.*
 - (i) *A is g -convex.*
 - (ii) $P(A) = \text{conv}\{[U] \mid U \in 2\text{-psilt}^1 A\}$.
 - (iii) $\Sigma(A)$ is finite and pairwise convex.
 - (iv) A is g -finite and pairwise g -convex.

Proof. (a) Immediate from Proposition 4.2(d).

(b) (i) \Leftrightarrow (ii) and (iii) \Leftrightarrow (iv) are clear.

(i) \Rightarrow (iii) Since the convex hull of $C_{\leq 1}(A) \cup C_{\leq 1}(A[1])$ contains the origin in its interior, so does $P(A)$. Thus $\Sigma(A)$ is complete and finite by (a), and pairwise positive by Proposition 5.8. By Proposition 5.7(c) \Rightarrow (a), $\Sigma(A)$ is pairwise convex.

(iii) \Rightarrow (i) By Proposition 4.2(d), $\Sigma(A)$ is complete. Thus the assertion follows from Proposition 5.7(a) \Rightarrow (c). \square

The following is a polytope analog of Definition 4.18.

Definition 5.11. [E, VK] Let P and Q be polytopes in \mathbb{R}^d and $\mathbb{R}^{d'}$ containing 0 in its interior respectively. We define a polytope $P \oplus Q$ in $\mathbb{R}^{d+d'}$ by

$$P \oplus Q := \text{conv}(P \cup Q).$$

We give the following easy observations.

Proposition 5.12. *Let A be a finite dimensional algebra over a field k .*

- (a) *For each idempotent $e \in A$, we have*

$$P(eAe) = P(A) \cap K_0(\text{proj } eAe)_{\mathbb{R}}.$$

In particular, if A is g -convex, then so is eAe .

- (b) Assume $A = A_1 \times \cdots \times A_\ell$ for a finite dimensional k -algebra A_i . Then A is g -convex if and only if each A_i is g -convex. In this case, we have

$$P(A) = P(A_1) \oplus \cdots \oplus P(A_\ell),$$

where we identify $K_0(\text{proj } A)_\mathbb{R}$ and $\bigoplus_{i=1}^\ell K_0(\text{proj } A_i)_\mathbb{R}$ by (4.1).

Proof. (a) This is immediate from Theorem 4.16.

(b) We have a bijection

$$2\text{-silt } A_1 \times \cdots \times 2\text{-silt } A_\ell \simeq 2\text{-silt } A \text{ given by } (T_1, \dots, T_\ell) \mapsto T_1 \oplus \cdots \oplus T_\ell.$$

Moreover the exchange sequences for A are given by the exchange sequences for some A_i . Thus A is pairwise g -convex if and only if each A_i is pairwise g -convex. By Theorem 5.10(b)(iii) \Rightarrow (i), we obtain the first assertion. Since $2\text{-psilt}^1 A = 2\text{-psilt}^1 A_1 \sqcup \cdots \sqcup 2\text{-psilt}^1 A_\ell$ holds, Theorem 5.10(b)(ii) implies

$$\begin{aligned} P(A) &= \text{conv}\{[U] \mid U \in 2\text{-psilt}^1 A\} = \text{conv}(\text{conv}\{[U] \mid U_i \in 2\text{-psilt}^1 A_i\} \mid 1 \leq i \leq \ell) \\ &= \text{conv}(P(A_i) \mid 1 \leq i \leq \ell) = \bigoplus_{i=1}^\ell P(A_i). \end{aligned}$$

Thus the second assertion follows. \square

5.2. Dual polytopes and c -polytopes. Let P be a convex polytope P in $V = \mathbb{R}^n$ containing the origin in its interior. For the dual \mathbb{R} -vector space $V^* \simeq \mathbb{R}^n$, we denote by $(-, -) : V^* \times V \rightarrow \mathbb{R}$ the natural pairing. Recall that the *dual polytope* is defined by

$$P^* := \{x \in V^* \mid \forall y \in P, (x, y) \leq 1\}.$$

Then P^* is again a convex polytope in V^* containing the origin in its interior, and $P^{**} = P$ holds. We call P *reflexive* if P and P^* are lattice polytopes.

Throughout this subsection, let A be a finite dimensional algebra over a field k . We introduce the notion of c -polytope as follows.

Definition 5.13. For $X \in D^b(\text{mod } A)$, let

$$[X]' := (\dim_k \text{End}_{D^b(\text{mod } A)}(X))^{-1} [X] \in K_0(\text{mod } A)_\mathbb{R}.$$

For the simple A -modules S_1, \dots, S_n , we define the k -Grothendieck group $K_0(\text{mod } A, k)$ of $\text{mod } A$ as the subgroup of $K_0(\text{mod } A)_\mathbb{R}$ generated by $[S_1]', \dots, [S_n]'$.

For $S = S_1 \oplus \cdots \oplus S_n \in \text{smc } A$, let

$$v_S := \sum_{i=1}^n [S_i]' \in K_0(\text{mod } A)_\mathbb{R}.$$

We define the c -polytope $P^c(A)$ of A as the convex hull

$$P^c(A) := \text{conv}\{v_S \mid S \in 2\text{-smc } A\} \subset K_0(\text{mod } A)_\mathbb{R}.$$

Using the Euler form

$$K_0(\text{proj } A)_\mathbb{R} \times K_0(\text{mod } A)_\mathbb{R} \rightarrow \mathbb{R} = K_0(\text{mod } k)_\mathbb{R} \text{ given by } (X, Y) \mapsto [\mathbf{R}\text{Hom}_A(X, Y)],$$

we identify $(K_0(\text{proj } A)_\mathbb{R})^*$ with $K_0(\text{mod } A)_\mathbb{R}$. Using this identification, we can state the following main result in this subsection.

Theorem 5.14. Let A be a finite dimensional algebra over a field k . Then A is g -convex if and only if

$$P(A) = (P^c(A))^*.$$

In this case, both $P(A)$ and $P^c(A)$ are reflexive polytopes.

We need the following information on silting- t -structure correspondence (Proposition 2.13).

Proposition 5.15. Let A be a finite dimensional algebra over a field k .

- (a) The abelian groups $K_0(\text{mod } A, k)$ and $K_0(\text{proj } A)$ are dual to each other with respect to the Euler form.
- (b) For any $S = S_1 \oplus \cdots \oplus S_n \in \text{smc } A$, the elements $[S_1]', \dots, [S_n]'$ are the basis of $K_0(\text{mod } A, k)$.

Proof. (a) This is immediate from Proposition 2.13.

(b) Take $T = T_1 \oplus \cdots \oplus T_n \in \text{silt } A$ corresponding to S . Then $[T_1], \dots, [T_n]$ give a basis of $K_0(\text{proj } A)$ by [AiI]. Thus the claim follows from (a). \square

The following simple observation is crucial.

Proposition 5.16. *Keep the setting in Proposition 2.13. Then the element $v_S \in K_0(\text{mod } A)_{\mathbb{R}}$ gives a normal vector of the facet of $P(A)$ corresponding to T .*

Proof. For each $1 \leq i, j \leq n$, (2.5) implies

$$([T_i] - [T_j], v_S) = ([T_i], v_S) - ([T_j], v_S) = 1 - 1 = 0.$$

This shows the assertion. \square

Now we are ready to prove Theorem 5.14.

Proof of Theorem 5.14. It suffices to show the “only if” part. Assume that A is g -convex. For $x \in K_0(\text{mod } A)_{\mathbb{R}}$, we consider the half space

$$H_x^{\leq 1} := \{y \in K_0(\text{proj } A)_{\mathbb{R}} \mid (y, x) \leq 1\} \subset K_0(\text{proj } A)_{\mathbb{R}}.$$

Since $P(A)$ is a convex polytope by assumption, Proposition 5.16 shows

$$P(A) = \bigcap_{S \in 2\text{-smc } A} H_{v_S}^{\leq 1} = (P^c(A))^*,$$

where the first equality follows from [Zi, Theorem 2.15(7)]. \square

We end this subsection by a remark. Using $2\text{-smc } A$, we can define the c -simplicial complex of A by the completely same way as in the definition of g -simplicial complex. However, it is not clear if the c -simplicial complex enjoys some nice properties. For example, the c -polytope $P^c(A)$ is not a geometric realization of the c -simplicial complex.

5.3. Newton polytopes. Throughout this subsection, let A be a finite dimensional algebra over a field k . We study a connection between g -fans $\Sigma(A)$ and Newton polytopes (also known as *Harder-Narasimhan polytopes*) of A -modules. In particular, our results recover some of results in [Fe1] (see also [BCDMTY, PPPP]).

Definition 5.17. [BKT, Fe1] The *Newton polytope* of $X \in \mathcal{A}$ is the convex hull

$$\begin{aligned} n(X) &:= \{[Y] \in K_0(\text{mod } A) \mid Y \text{ is a submodule of } X\} \subset K_0(\text{mod } A), \\ N(X) &:= \text{conv } n(X) \subset K_0(\text{mod } A)_{\mathbb{R}}. \end{aligned}$$

The dimension of $N(X)$ clearly equals the number of isomorphism classes of simple A -modules appearing in X as composition factors. Let $0 \neq \theta \in K_0(\text{proj } A)_{\mathbb{R}}$. Each $r \in \mathbb{R}$ give a hyperplane

$$H_{\theta}^r := \{x \in K_0(\text{mod } A)_{\mathbb{R}} \mid \theta(x) = r\} \subset K_0(\text{proj } A)_{\mathbb{R}}$$

and a half-space

$$H_{\theta}^{\leq r} := \{x \in K_0(\text{mod } A)_{\mathbb{R}} \mid \theta(x) \leq r\} \subset K_0(\text{proj } A)_{\mathbb{R}}.$$

Let $X \in \text{mod } A$. Clearly we have

$$N(X) = \bigcap_{0 \neq \theta \in K_0(\text{proj } A)} H_{\theta}^{\leq \max \theta(N(X))}.$$

Each $0 \neq \theta \in K_0(\text{proj } A)_{\mathbb{R}}$ gives a face of $N(X)$:

$$N(X)_{\theta} := N(X) \cap H_{\theta}^{\max \theta(N(X))}.$$

Conversely each face of $N(X)$ has a form $N(X)_{\theta}$ for some $0 \neq \theta \in K_0(\text{proj } A)_{\mathbb{R}}$.

The following properties are basic to study $N(X)$, see Definition-Proposition 4.6 for t_θ and \bar{t}_θ .

Lemma 5.18. *For each $\theta \in K_0(\text{proj } A)_\mathbb{R}$ and $X \in \text{mod } A$, the following assertions hold.*

- (a) $\theta(X) \leq \theta(t_\theta X) = \theta(\bar{t}_\theta X) = \max \theta(N(X))$.
- (b) *For each submodule Y of X , we have $\theta(Y) \leq \theta(t_\theta X)$.*
- (c) *Let Y be a submodule of X . Then $\theta(Y) = \theta(t_\theta X)$ holds if and only if $t_\theta X \subseteq Y \subseteq \bar{t}_\theta X$ and $Y/t_\theta X \in \mathcal{W}_\theta$ hold.*
- (d) *If a submodule Y of X satisfies $[Y] = [t_\theta X]$ (respectively, $[Y] = [\bar{t}_\theta X]$), then $Y = t_\theta X$ (respectively, $Y = \bar{t}_\theta X$).*

Proof. (a) The exact sequence $0 \rightarrow t_\theta X \rightarrow X \rightarrow \bar{f}_\theta X \rightarrow 0$ shows $\theta(X) = \theta(t_\theta X) + \theta(\bar{f}_\theta X)$. Since $\theta(\bar{f}_\theta X) \leq 0$, we obtain the left inequality. Similarly, the exact sequence $0 \rightarrow t_\theta X \rightarrow \bar{t}_\theta X \rightarrow w_\theta X \rightarrow 0$ and the equality $\theta(w_\theta X) = 0$ implies the middle equality. To prove the right equality, it suffices to show (b).

(b) By (a), $\theta(Y) \leq \theta(t_\theta Y)$ holds. The exact sequence $0 \rightarrow t_\theta Y \rightarrow t_\theta X \rightarrow t_\theta X/t_\theta Y \rightarrow 0$ and the inequality $\theta(t_\theta X/t_\theta Y) \geq 0$ imply $\theta(t_\theta Y) \leq \theta(t_\theta X)$. Thus the assertion follows.

(c) The “if” part is clear from $\theta(Y/t_\theta X) = 0$. We prove the “only if” part. Since $\theta(t_\theta X/t_\theta Y) = \theta(Y/t_\theta Y) = \theta(\bar{f}_\theta X) \leq 0$ and $t_\theta X \in \mathcal{T}_\theta$ hold, we have $t_\theta X = t_\theta Y \subseteq Y$. Since $\theta(Y/\bar{t}_\theta Y) = \theta(f_\theta Y) \leq 0$ and $\theta(\bar{t}_\theta X/\bar{t}_\theta Y) \geq 0$ hold, we have $\theta(Y) \leq \theta(\bar{t}_\theta Y) \leq \theta(\bar{t}_\theta X) = \theta(t_\theta X)$. Thus the equalities hold. Since $f_\theta Y \in \mathcal{F}_\theta$, we have $f_\theta Y = 0$ and hence $Y = \bar{t}_\theta Y \subseteq \bar{t}_\theta X$. The claim $Y/t_\theta X \in \mathcal{W}_\theta$ follows from a general fact: A submodule Z' of $Z \in \mathcal{W}_\theta$ belongs to \mathcal{W}_θ if and only if $\theta(Z') = 0$.

(d) Since the assumption implies $\theta(Y) = \theta(t_\theta X)$, we have $t_\theta X \subseteq Y \subseteq \bar{t}_\theta X$ by (c). Thus the assertion clearly holds. \square

Let $\theta \in K_0(\text{proj } A)_\mathbb{R}$. Then the inclusion functor $\mathcal{W}_\theta \rightarrow \text{mod } A$ induces morphisms

$$\iota : K_0(\mathcal{W}_\theta) \rightarrow K_0(\text{mod } A) \quad \text{and} \quad \iota : K_0(\mathcal{W}_\theta)_\mathbb{R} \rightarrow K_0(\text{mod } A)_\mathbb{R}.$$

As in Definition 5.17, for each $X \in \mathcal{W}_\theta$, let

$$\begin{aligned} n_{\mathcal{W}_\theta}(X) &:= \{[Y] \in K_0(\mathcal{W}_\theta) \mid Y \text{ is a subobject of } X\} \subset K_0(\mathcal{W}_\theta), \\ N_{\mathcal{W}_\theta}(X) &:= \text{conv } n_{\mathcal{W}_\theta}(X) \subset K_0(\mathcal{W}_\theta)_\mathbb{R}. \end{aligned}$$

We have the following descriptions of the faces of $N(X)$.

Lemma 5.19. *Let $X \in \text{mod } A$ and $0 \neq \theta \in K_0(\text{proj } A)_\mathbb{R}$. Then we have*

$$\begin{aligned} N(X)_\theta &= [t_\theta X] + \iota(N_{\mathcal{W}_\theta}(w_\theta X)), \\ \{\text{vertices of } N(X)\} &= \{[t_\theta X] \mid 0 \neq \theta \in K_0(\text{proj } A)\}. \end{aligned}$$

Moreover, each edge of $N(X)$ has a form $\text{conv}\{[t_\theta X], [\bar{t}_\theta X]\}$ for some $0 \neq \theta \in K_0(\text{proj } A)$.

Proof. By Lemma 5.18(a)(c), we have

$$n(X) \cap H_\theta^{\max \theta(N(X))} = [t_\theta X] + \iota(n_{\mathcal{W}_\theta}(w_\theta X)).$$

Taking the convex hull, we obtain the first equality. In particular, $[t_\theta X]$ is a vertex of $N(X)_\theta$ and hence of $N(X)$. Conversely, for each vertex v of $N(X)$, there exists $0 \neq \theta \in K_0(\text{proj } A)$ such that $N(X)_\theta = \{v\}$. Then $v = [t_\theta X]$ holds by the first equality.

Since each face of $N(X)$ has a form $N(X)_\theta$ for some $0 \neq \theta \in K_0(\text{proj } A)$ and contains $[t_\theta X]$ and $[\bar{t}_\theta X]$, the last assertion follows. \square

To give a more explicit description of faces of $N(X)$, we use 2-term presilting complexes. For $0 \neq U \in 2\text{-psilt } A$, define a face of $N(X)$ by

$$N(X)_U := N(X)_{[U]}.$$

We have torsion classes $\mathcal{T}_U, \overline{\mathcal{T}}_U$, a wide subcategory \mathcal{W}_U and functors t_U, \bar{t}_U and w_U (see Definition-Propositions 2.8 and 3.7). For each $Y \in \mathcal{W}_U$, we denote by $s_{\mathcal{W}_U}(Y)$ the number of isomorphism classes of simple objects in \mathcal{W}_U appearing in Y as a composition factor.

Lemma 5.20. *Let $X \in \text{mod } A$ and $0 \neq U \in 2\text{-psilt } A$.*

(a) *We have*

$$\dim N(X)_U = s_{\mathcal{W}_U}(w_U X) \leq |A| - |U|.$$

(b) *For each $\theta \in C^+(U)$, we have*

$$N(X)_\theta = N(X)_U.$$

Proof. (a) By Definition-Proposition 3.7(d), there exists a finite dimensional algebra B such that $\mathcal{W}_U \simeq \text{mod } B$ and $|B| = |A| - |U|$ and $\iota : K_0(\mathcal{W}_U) \simeq K_0(\text{mod } B) \rightarrow K_0(\text{mod } A)$ is injective. Thus we have

$$\dim N(X)_\theta \stackrel{5.19}{=} \dim N_{\mathcal{W}_U}(w_U X) = s_{\mathcal{W}_U}(w_U X) \leq |B| = |A| - |U|.$$

(b) By Proposition 4.7, we have $t_\theta = t_U$, $\mathcal{W}_\theta = \mathcal{W}_U$ and $w_\theta = w_U$. By Lemma 5.19, we obtain

$$N(X)_\theta = [t_\theta X] + \iota(N_{\mathcal{W}_\theta}(w_\theta X)) = [t_U X] + \iota(N_{\mathcal{W}_U}(w_U X)) = N(X)_U. \quad \square$$

To state our main result, we introduce some notions. Recall that $2\text{-psilt}^i A$ is the set of isomorphism classes of basic 2-term presilting complex $U \in K^b(\text{proj } A)$ with $|U| = i$ (Definition 3.1).

Definition 5.21. Let A be a finite dimensional algebra over a field k , $n := |A|$, and $X \in \text{mod } A$.

(a) An object X in an abelian length category \mathcal{W} is called *sincere* if each simple object in \mathcal{W} appears in X as a composition factor. For $0 \leq i \leq n$, let

$$\begin{aligned} 2\text{-psilt}_X^i A &:= \{U \in 2\text{-psilt}^i A \mid w_U X \text{ is sincere in } \mathcal{W}_U\}, \\ \Sigma_{i,X}(A) &:= \{C(U) \mid U \in 2\text{-psilt}_X^i A\} \subseteq \Sigma_i(A). \end{aligned}$$

By Lemma 5.20(a), $U \in 2\text{-psilt}^i A$ belongs to $2\text{-psilt}_X^i A$ if and only if $\dim N(X)_U = n - i$.

(b) We have an order preserving map

$$2\text{-silt } A \rightarrow \{\text{submodules of } X\} \text{ given by } U \mapsto t_U X. \quad (5.2)$$

We define an equivalence relation \sim_X on $2\text{-silt } A$ by

$$T \sim_X U \iff t_T X = t_U X \stackrel{5.18(d)}{\iff} [t_T X] = [t_U X].$$

This gives an equivalence relation \sim_X on $\Sigma_n(A)$ since we have a bijection $2\text{-silt } A \simeq \Sigma_n(A)$ given by $T \mapsto C(T)$.

- (c) We regard the vertices and the edges of $N(X)$ as a graph. Define a quiver $\vec{N}_1(X)$ by regarding each edge $\text{conv}\{[t_\theta X], [\bar{t}_\theta X]\}$ (see Lemma 5.19) as an arrow $[\bar{t}_\theta X] \rightarrow [t_\theta X]$.
 (d) Define a *contraction* $\text{Hasse}(2\text{-silt } A) / \sim_X$ of $\text{Hasse}(2\text{-silt } A)$ by identifying all vertices in each equivalence class of \sim_X and removing all loops.

Now we give explicit descriptions of the normal fan $\Sigma(N(X))$ (see Definition 2.4) as a coarsening fan of the g -fan $\Sigma(A)$ (see Definition 2.3), and of the 1-skeleton $\vec{N}_1(X)$ as a contraction of the Hasse quiver $\text{Hasse}(2\text{-silt } A)$. The following result is an explicit version of [Fe2, Propositions 7.4, 8.5] for g -finite case.

Theorem 5.22. *Let A be a finite dimensional algebra over a field k which is g -finite, and $n := |A|$.*

(a) *The equivalence relation \sim_X coarsens $\Sigma(A)$, and we have*

$$\Sigma(N(X)) = \Sigma(A) / \sim_X.$$

(b) *For each $0 \leq i \leq n$, there is a surjection*

$$2\text{-psilt}_X^i A \simeq \Sigma_{i,X}(A) \rightarrow \{\text{faces of } N(X) \text{ of dimension } n - i\} \quad (5.3)$$

given by $U \mapsto C(U) \mapsto N(X)_U$, which induces bijections

$$\begin{aligned} 2\text{-silt } A / \sim_X &\simeq \{\text{vertices of } N(X)\}, \\ 2\text{-psilt}_X^1 A &\simeq \{\text{facets of } N(X)\}. \end{aligned}$$

(c) We have an isomorphism of quivers

$$\vec{N}_1(X) \simeq \text{Hasse}(2\text{-silt}A)/\sim_X.$$

To prove this, we need the following preparation.

Lemma 5.23. *Let $n := |A|$ and $X \in \text{mod } A$.*

- (a) *Each cone of $\Sigma(A)$ is contained in a cone of $\Sigma(N(X))$.*
- (b) *Let $\sigma, \tau \in \Sigma_n(A)$. Then $\sigma \sim_X \tau$ if and only if σ and τ are contained in the same cone of $\Sigma_n(N(X))$.*

Proof. (a) Immediate from Lemma 5.20(b).

(b) Take $T, U \in 2\text{-silt}A$ such that $\sigma = C(T)$ and $\tau = C(U)$. Then $N(X)_T = \{[t_TX]\}$ and $N(X)_U = \{[t_UX]\}$ hold by Lemmas 5.19 and 5.20(a). Thus σ and τ are contained in the same cone of $\Sigma_n(N(X))$ if and only if $[t_TX] = [t_UX]$, that is, $\sigma \sim_X \tau$. \square

We are ready to prove Theorem 5.22.

Proof of Theorem 5.22. (a) The assertion follows from Lemma 5.23(a)(b).

(b) The map (5.3) is well-defined by the definition of $2\text{-psilt}_X^i A$. We prove that it is surjective. Let F be a face of $N(X)$ of dimension $n - i$. Since $\Sigma(A)$ is complete, Lemma 5.23(a) implies that the cone σ_F of $\Sigma(N(X))$ of dimension i is a union of cones in $\Sigma(A)$. Thus there exists $U \in 2\text{-psilt}_X^i A$ such that $C(U) \subseteq \sigma_F$. Since $[U] \in \sigma_F^\circ$, we have $N(X)_U = F$. Thus $U \in 2\text{-psilt}_X^i A$ holds, and the map (5.3) is surjective.

Since $2\text{-psilt}_X^n A = 2\text{-silt}A$, the map (5.3) gives the first bijection by Lemma 5.23(b). It also gives the second bijection since different elements in $\Sigma_{1,X}(A)$ give different facets of $N(X)$.

(c) By (b), both quivers have the set $2\text{-silt}A/\sim_X$ of vertices. We compare their arrows. By Lemma 5.19, we have a surjection

$$2\text{-psilt}_X^{n-1}A \rightarrow \{\text{arrows in } \vec{N}_1(X)\} \text{ given by } U \mapsto [\bar{t}_U X \rightarrow t_U X].$$

On the other hand, by Proposition 2.6, each arrow in $\text{Hasse}(2\text{-silt}A)$ can be written as $U_{\max} \rightarrow U_{\min}$ for some $U \in 2\text{-psilt}_X^{n-1}A$. It is still an arrow in $\text{Hasse}(2\text{-silt}A)/\sim_X$ if and only if $U_{\max} \not\sim_X U_{\min}$ if and only if $\bar{t}_U X \neq t_U X$ (since $t_U = t_{U_{\min}}$ and $\bar{t}_U = t_{U_{\max}}$ hold by Definition-Proposition 3.7) if and only if $U \in 2\text{-psilt}_X^{n-1}A$. Thus we have a surjection

$$2\text{-psilt}_X^{n-1}A \rightarrow \{\text{arrows in } \text{Hasse}(2\text{-silt}A)/\sim_X\} \text{ given by } U \mapsto [U_{\max} \rightarrow U_{\min}].$$

Moreover, $U, V \in 2\text{-psilt}_X^{n-1}A$ give the same arrow in $\vec{N}_1(X)$ if and only if $\bar{t}_U X = \bar{t}_V X$ and $t_U X = t_V X$ hold if and only if $U_{\max} \sim_X V_{\max}$ and $U_{\min} \sim_X V_{\min}$ hold if and only if U and V give the same arrow in $\text{Hasse}(2\text{-silt}A)/\sim_X$. Thus the assertion follows. \square

As an application of Theorem 5.22, we obtain the following result. The part (b) was obtained in [Fe2, Theorem 4.17, Corollary 6.9].

Corollary 5.24. *Let A be a finite dimensional algebra over a field k which is g -finite.*

- (a) *For $X \in \text{mod } A$, the following conditions are equivalent.*
 - (i) $\Sigma(N(X)) = \Sigma(A)$ holds.
 - (ii) $\vec{N}_1(X) \simeq \text{Hasse}(2\text{-silt}A)$ holds.
 - (iii) *For each $U \in 2\text{-psilt}_X^{n-1}A$, $w_U X \neq 0$ holds.*
- (b) *If one of the following conditions is satisfied, then the conditions in (a) are satisfied.*
 - (i) *Each brick of A is a direct summand of X .*
 - (ii) *For each $V \in 2\text{-psilt}^1 A$, $H^0(V)$ is a direct summand of X .*

Proof. (a) By Theorem 5.22(a)(c), the conditions (i) and (ii) are equivalent to that the map (5.2) is injective. This is equivalent to that $t_T X \neq t_{T'} X$ holds for each arrow $T \rightarrow T'$ in $\text{Hasse}(2\text{-silt}A)$.

Recall that there is a bijection between $2\text{-psilt}^{n-1}A$ and the set of arrows of $\text{Hasse}(2\text{-silt}A)$ given by $U \mapsto [U_{\max} \rightarrow U_{\min}]$, and we have

$$w_U X = t_{U_{\max}} X / t_{U_{\min}} X,$$

see Definition-Proposition 3.7(d). Thus the map (5.2) is injective if and only if (iii) holds.

(b) In each case, it suffices to check the condition (a)(iii).


Consider the case (i). For each $U \in 2\text{-psilt}^{n-1}A$, the wide subcategory $\mathcal{W}_U = \overline{\mathcal{T}}_U \cap \overline{\mathcal{F}}_U$ has a unique simple object S . Then $w_U S = \bar{t}_U S / t_U S = S \neq 0$ holds. Since S is a brick and hence a direct summand of X , we have $w_U X \neq 0$.

Consider the case (ii). For each $U \in 2\text{-psilt}^{n-1}A$, let $U_{\max} = U \oplus V$ for some $V \in 2\text{-psilt}^1 A$. Then $H^0(V)$ belongs to $\mathcal{T}_{U_{\max}} = \overline{\mathcal{T}}_U$ and does not belong to $\mathcal{T}_{U_{\min}} = \mathcal{T}_U$. Thus $\bar{t}_U H^0(V) = H^0(V) \neq t_U H^0(V)$ and hence $w_U H^0(V) \neq 0$. Since $H^0(V)$ is a direct summand of X , we have $w_U X \neq 0$. \square

Example 5.25. Consider a finite dimensional algebra A given by

$$Q = \left[1 \begin{smallmatrix} \xrightarrow{a} \\ \xleftarrow{c} \end{smallmatrix} 2 \begin{smallmatrix} \xrightarrow{b} \\ \xleftarrow{d} \end{smallmatrix} 3 \right] \quad \text{and} \quad A := kQ / \langle ab, dc, ca - bd \rangle.$$

Then $A = \frac{1}{2} \oplus 1 \oplus \frac{2}{2} \oplus \frac{3}{3}$. For $X = P_1^{\oplus p} \oplus P_2^{\oplus q} \oplus P_3^{\oplus r} \oplus S_2^{\oplus s}$ with $p, q, r, s \geq 0$, the Newton polytope $N(X)$ is given by the following.



In particular, $\vec{N}_1(X) \simeq \text{Hasse}(2\text{-silt}A)$ holds if all of p, q, r, s are positive.

6. CONVEX g -POLYGONS AND SMOOTH FANO g -POLYTOPES

In this section, we give complete classifications of convex g -polygons and smooth Fano g -polytopes. To state our result explicitly, we need the following general notion.


- Definition 6.1.** (a) Let (Σ, σ_+) and (Σ', σ'_+) be sign-coherent fans in \mathbb{R}^d and $\mathbb{R}^{d'}$ respectively. An *isomorphism of sign-coherent fans* is an isomorphism $f : \Sigma \simeq \Sigma'$ of fans (Definition 2.2) such that $\{f(\sigma_+), f(-\sigma_+)\} = \{\sigma'_+, -\sigma'_+\}$.
- (b) Let A and A' be finite dimensional algebras. An *isomorphism of g -fans* is an isomorphism $\Sigma(A) \simeq \Sigma(A')$ of sign-coherent fans. The induced isomorphism $P(A) \simeq P(A')$ is called an *isomorphism of g -polytopes*.

Proposition 6.2. *For each $n \geq 1$, there exist only finitely many convex g -polytopes (respectively, convex g -fans) of dimension n up to isomorphisms of g -polytopes (respectively, g -fans).*


Proof. The origin is a unique lattice point in the interior of $P(A)$. By [LZ], there exists only finitely many convex g -polytopes of dimension n up to isomorphisms. Each polytope has only finitely many unimodular triangulations, and each nonsingular fan has only finitely many choices of σ_+ . Thus the assertion follows. \square

6.1. Convex g -polygons. We give the following classification of convex g -polygons.


Theorem 6.3. *There are precisely 7 convex g -polygons up to isomorphism of g -polytopes (Definition 6.1).*



The corresponding c -polygons are the following.



Proof. By Theorem 5.14, g -polytopes are reflexive. It is well-known that there are precisely 16 reflexive polytopes [PR].




On the other hand, g -fans are sign-coherent by Proposition 4.12. One can easily check by case-by-case analysis that, among 16 fans given by these 16 reflexive polygons, there are exactly 7 sign-coherent fans listed above up to isomorphism of sign-coherent fans. \square

6.2. Smooth Fano g -polytopes. The aim of this subsection is to characterize finite dimensional algebras whose g -polytopes satisfy the following property.


Definition 6.4. [O] Let d be a positive integer. A lattice polytope P in \mathbb{R}^d containing the origin in its interior is called a *smooth Fano d -polytope* if the vertices of every facet F of P is \mathbb{Z}^d -basis of the lattice \mathbb{Z}^d .

Among 16 reflexive polygons, there are 5 smooth Fano polygons.



The following special class of algebras plays an important role.

Definition 6.5. A finite dimensional algebra A over a field k is *pentagon type* (respectively, *hexagon type*) if $P(A)$ is the left (respectively, right) polygon below.



We refer to [AHIKM1] for characterizations of these algebras.

The following is a main result of this section.

Theorem 6.6. *Let A be a finite dimensional algebra over a field k . Then $P(A)$ is a smooth Fano polytope if and only if A is a product of local algebras, algebras of pentagon type and algebras of hexagon type.*

The rest of this section is devoted to a proof of Theorem 6.6. We start with the following, where we refer to Definition 5.11 for the definition of sums of polytopes.

Lemma 6.7. *The following assertions hold.*

- (a) *Let P_i be a lattice polytope in \mathbb{R}^{d_i} for $1 \leq i \leq \ell$. Then $P_1 \oplus \cdots \oplus P_\ell$ is smooth Fano if and only if P_i is smooth Fano for each $1 \leq i \leq \ell$.*
- (b) *Let A_i be a finite dimensional algebra over a field k for $1 \leq i \leq \ell$ and $A = A_1 \times \cdots \times A_\ell$. Then $P(A)$ is smooth Fano if and only if $P(A_i)$ is smooth Fano for each $1 \leq i \leq \ell$.*

Proof. (a) This is clear.

(b) This is immediate from (a) and Proposition 5.12(b). \square

The following technical notion plays a central role.

Definition 6.8. Let $k \geq 1$. A *del Pezzo $2k$ -polytope* (respectively, *pseudo del Pezzo $2k$ -polytope*) is

$$V_{2k} := \text{conv}\{\pm e_1, \dots, \pm e_{2k}, \pm \sum_{i=1}^{2k} e_i\} \quad (\text{respectively, } \tilde{V}_{2k} := \text{conv}\{\pm e_1, \dots, \pm e_{2k}, \sum_{i=1}^{2k} e_i\}).$$

The following is clear.

Lemma 6.9. *Let A be a finite dimensional algebra.*

- (a) *A is local if and only if $P(A)$ is isomorphic to a line segment $[e_1, -e_1]$.*
- (b) *A is pentagon type if and only if $P(A) \simeq V_2$.*
- (c) *A is hexagon type if and only if $P(A) \simeq \tilde{V}_2$.*

Recall that a polytope P in \mathbb{R}^d is called a *symmetric* if $v \in P$ implies $-v \in P$. More generally, a polytope P in \mathbb{R}^d is called a *pseudo-symmetric* [O] if there exists a facet F of P such that $-F$ is also a facet.

The following result provides a classification of pseudo-symmetric smooth Fano d -polytopes.

Proposition 6.10. [E, VK] *Any pseudo-symmetric smooth Fano d -polytope P splits into copies of line segments, del Pezzo polytopes and pseudo del Pezzo polytopes. That is, P is isomorphic to a convex polytope*

$$\overbrace{L \oplus \cdots \oplus L}^{i\text{-times}} \oplus V_{2a_1} \oplus \cdots \oplus V_{2a_j} \oplus \tilde{V}_{2b_1} \oplus \cdots \oplus \tilde{V}_{2b_k},$$

where each L is a lattice convex polytope isomorphic to the line segment $[e_1, -e_1]$ and $i + 2(a_1 + \cdots + a_j + b_1 + \cdots + b_k) = d$.

We also need the following description of the facets of (pseudo) del Pezzo polytopes.

Lemma 6.11. *Let $s := \sum_{i=1}^{2k} e_i$.*

- (a) *The facets of a del Pezzo $2k$ -polytope V_{2k} consist of*

$$\text{conv}\{e_{i_1}, \dots, e_{i_k}, -e_{j_1}, \dots, -e_{j_k}\}, \quad (6.1)$$

where $1 \leq i_1 < \dots < i_k \leq 2k$, $1 \leq j_1 < \dots < j_k \leq 2k$ and $\{i_1, \dots, i_k\} \cap \{j_1, \dots, j_k\} = \emptyset$, and

$$\text{conv}\{e_{i_1}, \dots, e_{i_k}, -e_{j_1}, \dots, -e_{j_{k-1}}, s\} \text{ and} \quad (6.2)$$

$$\text{conv}\{-e_{i_1}, \dots, -e_{i_k}, e_{j_1}, \dots, e_{j_{k-1}}, -s\}, \quad (6.3)$$

where $1 \leq i_1 < \dots < i_k \leq 2k$, $1 \leq j_1 < \dots < j_{k-1} \leq 2k$ and $\{i_1, \dots, i_k\} \cap \{j_1, \dots, j_{k-1}\} = \emptyset$.

(b) The facets of a pseudo del Pezzo $2k$ -polytope \tilde{V}_{2k} consist of (6.2) and

$$\text{conv}\{e_{i_1}, \dots, e_{i_\ell}, -e_{j_1}, \dots, -e_{j_{2k-\ell}}\}, \quad (6.4)$$

where $1 \leq i_1 < \dots < i_\ell \leq 2k$, $1 \leq j_1 < \dots < j_{2k-\ell} \leq 2k$, $\{i_1, \dots, i_\ell\} \cap \{j_1, \dots, j_{2k-\ell}\} = \emptyset$ and $\ell \leq k$.

Proof. Although the proof is routine, we include it for convenience of the reader.

(a) Let F be a facet of V_{2k} and let H be the supporting hyperplane of F . Then $\pm e_i$ is not contained in F at the same time, and so is $\pm s$.

- When F does not contain $\pm s$, the vertices of F look like, say, $e_1, \dots, e_\ell, -e_{\ell+1}, \dots, -e_{2k}$. Then H is defined by the equality $f(x) = 1$ for

$$f(x) := x_1 + \dots + x_\ell - x_{\ell+1} - \dots - x_{2k}$$

and V_{2k} is contained in the half space $\{x \in \mathbb{R}^d \mid f(x) \leq 1\}$. If $\ell \neq k$, then one of $f(s)$ and $f(-s)$ is greater than 1, a contradiction. Hence, $\ell = k$, i.e., F is of the form (6.1).

- When F contains $\pm s$, the vertices of F look like $\pm(e_1, \dots, e_\ell, -e_{\ell+1}, \dots, -e_{2k-1}, s)$. Then H is defined by the equality $f(x) = 1$ for

$$f(x) := x_1 + \dots + x_\ell - x_{\ell+1} - \dots - x_{2k-1} + (2k - 2\ell)x_{2k}$$

and V_{2k} is contained in the half space $\{x \in \mathbb{R}^d \mid f(x) \leq 1\}$. If $\ell \neq k$, then one of $f(e_{2k})$ and $f(-e_{2k})$ is greater than 1, a contradiction. Hence, $\ell = k$, i.e., F is of the forms (6.2) and (6.3).

(b) Let F be a facet of \tilde{V}_{2k} and let H be the supporting hyperplane of F . When F does not contain s , the vertices of F look like $e_1, \dots, e_\ell, -e_{\ell+1}, \dots, -e_{2k}$. Then H is defined by the equality $f(x) = 1$ for

$$f(x) := x_1 + \dots + x_\ell - x_{\ell+1} - \dots - x_{2k}.$$

and \tilde{V}_{2k} is contained in the half space $\{x \in \mathbb{R}^d \mid f(x) \leq 1\}$. If $\ell > k$, then $f(s) > 1$, a contradiction. Hence, $\ell \leq k$, i.e., F is of the form (6.4). When F contains s , the same proof as above can be applied, and we conclude that F is of the form (6.2). \square

Let P be a smooth Fano polytope in \mathbb{R}^d . Then there exists a unique fan Σ in \mathbb{R}^d such that $P = P(\Sigma)$, see Definition 5.1. We call P *ordered* if there exists $\sigma_+ \in \Sigma_d$ such that (Σ, σ_+) is an ordered fan in the sense of Definition 4.13.

Now we prove the following key observation.

Proposition 6.12. (a) A del Pezzo $2k$ -polytope V_{2k} is ordered if and only if $k = 1$.

(b) A pseudo del Pezzo $2k$ -polytope \tilde{V}_{2k} is ordered if and only if $k = 1$.

Proof. Recall that an ordered polytope P has a facet F_+ such that $(\text{cone } F_+)^\circ \cap \text{span } F' = \emptyset$ for each non-maximal face F' of P .

(a) ‘if’ part is clear. To prove ‘only if’ part, we assume that V_{2k} is ordered for $k \geq 2$. By Lemma 6.11 (a), we may assume that

$$F_+ = \text{conv}\{e_1, \dots, e_k, -e_{k+1}, \dots, -e_{2k}\} \text{ or } F_+ = \text{conv}\{e_1, \dots, e_k, -e_{k+1}, \dots, -e_{2k-1}, s\}$$

without loss of generality. In the first case, since $k \geq 2$, we have

$$(\text{cone } F_+)^\circ \ni e_1 + \dots + e_k - e_{k+1} - \dots - e_{2k} = s + 2(-e_{k+1} - \dots - e_{2k}) \in \text{span } F'$$

for the non-maximal face $F' = \text{conv}\{e_{k+1}, \dots, e_{2k}, s\}$, a contradiction. In the second case, we have

$$(\text{cone } F_+)^\circ \ni e_1 + \dots + e_k - e_{k+1} - \dots - e_{2k-1} + s = 2(e_1 + \dots + e_k) + e_{2k} \in \text{span } F'$$

for the non-maximal face $F' = \text{conv}\{e_1, \dots, e_k, -e_{2k}\}$, a contradiction.

(b) ‘if’ part is clear. To prove ‘only if’ part, we assume that \tilde{V}_{2k} is ordered for $k \geq 2$. By Lemma 6.11 (b), we may assume that

$$F_+ = \text{conv}\{e_1, \dots, e_\ell, -e_{\ell+1}, \dots, -e_{2k}\} \text{ or } F_+ = \text{conv}\{e_1, \dots, e_k, -e_{k+1}, \dots, -e_{2k-1}, s\}$$

where $\ell \leq k$. In the first case, we have

$$(\text{cone } F_+)^{\circ} \ni e_1 + \dots + e_\ell - e_{\ell+1} - \dots - e_{2k} = 2(e_1 + \dots + e_\ell) - s \in \text{span } F'$$

for the non-maximal face $F' = \text{conv}\{e_1, \dots, e_\ell, s\}$, a contradiction. In the second case, we get a contradiction since $-F_+$ is not a facet of \tilde{V}_{2k} . \square

Now we are ready to give a proof of Theorem 6.6.

Proof of Theorem 6.6. “If” part is clear. In fact, each of local algebras, algebras of pentagon type and algebras of hexagon type is smooth Fano, and therefore their product is also smooth Fano by Lemma 6.7(b).

To prove “only if” part, assume that $P(A)$ is smooth Fano. By Lemma 6.7(b), we only have to consider the case A is ring-indecomposable. Then $\Sigma(A)$ is indecomposable by Theorem 4.19(c). By Propositions 6.10 and 6.12, $P(A)$ is one of the line segments, del Pezzo 2-polytopes and pseudo del Pezzo 2-polytopes. Hence A is the one of the local algebras, algebras of pentagon type and algebras of hexagon type by Lemma 6.9. \square

7. PREPROJECTIVE ALGEBRAS AND COXETER FANS

The aim of this section is to show that g -fan of a preprojective algebra Π of Dynkin type is the Coxeter fans, and the c -polytope $\Pi^c(\Pi)$ is the short root polytope. In particular, Π is g -convex if and only if it is of type A_n or B_n . In this case, $P(\Pi)$ is the dual polytope of the short root polytope.

7.1. Classical preprojective algebras. To recall the definition of preprojective algebras, we start with the definition of hereditary algebras.

Definition 7.1. (a) We call a pair $(D_i, {}_iM_j)_{1 \leq i, j \leq n}$ k -species if

- (i) D_i is a finite dimensional division k -algebra.
- (ii) ${}_iM_j$ is a finitely generated $D_j \otimes_k D_i^{\text{op}}$ -module. In other words, ${}_iM_j$ is a (D_i, D_j) -module and k acts centrally on ${}_iM_j$.

(b) We call a k -species $(D_i, {}_iM_j)$ *acyclic* if there does not exist a sequence $i_1, i_2, \dots, i_\ell, i_{\ell+1} = i_1$ such that ${}_iM_{i_{j+1}} \neq 0$ for $1 \leq j \leq n$.

Let $(D_i, {}_iM_j)$ be a k -species, $D := \prod_{i=1}^n D_i$ and $M := \bigoplus_{1 \leq i, j \leq n} {}_iM_j$. We define the tensor algebra by

$$T_D(M) := \bigoplus_{n=0}^{\infty} T^n(M),$$

where $T^0(M) = D$ and $T^n(M) := M^{\otimes n} := M \otimes_D \dots \otimes_D M$ (n times). This is a k -algebra, which is finite dimensional if and only if $(D_i, {}_iM_j)$ is acyclic.

If $(D_i, {}_iM_j)$ is an acyclic k -species, then $T_D(M)$ is a finite dimensional hereditary k -algebra [L, Theorem 2.35].

Definition 7.2. (a) A matrix $C = (c_{ij}) \in M_n(\mathbb{Z})$ is a *symmetrizable generalized Cartan matrix* if the following conditions hold.

- (C1) $c_{ii} = 2$ for all i ;
- (C2) $c_{ij} \leq 0$ for all $i \neq j$;
- (C3) $c_{ij} \neq 0$ if and only if $c_{ji} \neq 0$.

(C4) There is a diagonal integer matrix $D = \text{diag}(c_1, \dots, c_n)$ with $c_i \geq 1$ for all i such that CD is symmetric. It is called a *symmetrizer* of C .

(b) Let $(D_i, {}_iM_j)$ be an acyclic k -species. We define the matrix $C = (c_{i,j})_{1 \leq i,j \leq n}$ associated to it as follows: We define $c_{ii} = 2$ for any i . If $i \neq j$, then we define c_{ij} and c_{ji} as follows.

- (i) If ${}_iM_j = 0 = {}_jM_i$, then $c_{ij} = 0 = c_{ji}$.
- (ii) If ${}_iM_j \neq 0$, then $c_{ij} := -\dim({}_iM_j)_{D_j}$ and $c_{ji} := -\dim_{D_i}({}_iM_j)$.
- (iii) If ${}_jM_i \neq 0$, then $c_{ij} := -\dim_{D_j}({}_jM_i)$ and $c_{ji} := -\dim({}_jM_i)_{D_i}$.

This is well-defined since the acyclicity implies that at least one of the ${}_iM_j$ or ${}_jM_i$ is zero.

Then the matrix C is symmetrizable generalized Cartan matrix. Indeed, let $c_i := \dim_k D_i$ and $D := \text{diag}(c_1, \dots, c_n)$. Then if ${}_iM_j \neq 0$, then we have $c_{ij}c_j = -\dim_k({}_iM_j) = c_{ji}c_i$ and hence CD is symmetric. The other cases are similar.

(c) Let $C = (c_{i,j})$ be the symmetrizable generalized Cartan matrix corresponding an acyclic k -species $(D_i, {}_iM_j)$. We call $H := T_D(M)$ is Dynkin type if the matrix C is Dynkin, that is, one of the type A_n, B_n, \dots, G_2 .

For example, the Cartan matrix of type A_n or B_n is

$$\begin{pmatrix} 2 & -1 & 0 & \dots & \dots & 0 \\ -1 & 2 & -1 & \ddots & & \vdots \\ 0 & -1 & \ddots & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & 2 & -1 \\ \vdots & & & & -1 & 2 \\ 0 & \dots & 0 & 0 & -1 & 2 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 2 & -1 & 0 & \dots & \dots & 0 \\ -1 & 2 & -1 & \ddots & & \vdots \\ 0 & -1 & \ddots & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & 2 & -1 \\ \vdots & & & & -1 & 2 \\ 0 & \dots & 0 & 0 & -2 & 2 \end{pmatrix}$$

Definition 7.3. Let $H^e := H \otimes H^{\text{op}}$ and $E := \text{Ext}_{H^e}^1(H, H^e) \in \text{mod } H^e$. We define the tensor algebra $\Pi := T_{H^e}(E)$ and call it the *preprojective algebra* of H . We remark that it is a \mathbb{Z} -graded algebra with $\Pi_i = E^{\otimes i}$, and $(\Pi_i)_H \simeq \tau^{-i}H$ in $\text{mod } H$ and ${}_H(\Pi_i) \simeq \tau^{-i}H$ in $\text{mod } H^{\text{op}}$.

Let C be a symmetrizable Cartan matrix of Dynkin type of rank n , that is, one of the type A_n, B_n, \dots, G_2 . Let $\Phi = \Phi(C)$ be the root system of C . Let $\{\alpha_1, \dots, \alpha_n\} \subset \Phi$ be a set of simple roots and L the root lattice. We let $V := L \otimes_{\mathbb{Z}} \mathbb{R}$ and denote by V^* the dual of V with the basis $\alpha_1^*, \alpha_2^*, \dots, \alpha_n^*$. We denote by the natural pairing (v^*, v) for $v \in V$ and $v^* \in V^*$.

Define a reflection $s_i : V \rightarrow V$ by

$$s_i(\alpha_j) := \alpha_j - c_{ij}\alpha_i.$$

The *Weyl group* is defined as a subgroup

$$W = W(C) = \langle s_1, \dots, s_n \rangle$$

of $\text{GL}(V)$. Then W acts also on V^* by

$$(wf)(v) = (f, w^{-1}v) \quad \text{for } f \in V^*, v \in V.$$

Define the *dominant chamber* as follows:

$$D := \bigcap_{i \in Q_0} \{v^* \in V^* \mid (v^*, \alpha_i) \geq 0\} = \left\{ \sum_{i=1}^n a_i \alpha_i^* \mid a_i \geq 0 \right\}.$$

Then the set $\{wD \mid w \in W\}$ of all cones wD and their faces is a fan in V^* and we call the *Coxeter fan*, see e.g. [ReS2].

Recall from Definition 5.13 that, for $X \in \text{D}^b(\text{mod } \Pi)$, let

$$[X]' := (\dim_k \text{End}_{\text{D}^b(\text{mod } \Pi)}(X))^{-1}[X] \in K_0(\text{mod } \Pi)_{\mathbb{R}}.$$

For the simple Π -modules D_1, \dots, D_n , we define the isomorphism

$$\iota : K_0(\text{mod } \Pi)_{\mathbb{R}} \simeq V \quad \text{by } \iota([D_i]') = \alpha_i,$$

and we identify $K_0(\text{mod } \Pi)_{\mathbb{R}}$ with V via ι . Moreover, for $X = X_1 \oplus \cdots \oplus X_n \in \text{smc}\Pi$, let

$$v_X := \iota\left(\sum_{i=1}^n [X_i]'\right) \in V.$$

The following is the main result of this subsection.

Theorem 7.4. *Let H be a finite dimensional hereditary k -algebra of Dynkin type and Π the preprojective algebra of H .*

(a) $\Sigma(\Pi)$ is the Coxeter fan and we have

$$P(\Pi) = \bigcup_{w \in W} wC_{\leq 1}(\Pi).$$

(b) $P(\Pi)$ is convex if and only if the Cartan matrix C is type A_n or B_n .

(c) (i) Let $\text{ind-2-smc}\Pi$ be the set of indecomposable direct summands of 2-simple minded collections of Π . We have

$$\{[S]' \mid S \in \text{ind-2-smc}\Pi\} = \Phi.$$

(ii) Let Φ_{short} be the set of short roots of Φ . We have

$$\{v_X \mid X \in \text{2-smc}\Pi\} = \Phi_{\text{short}} \quad \text{and} \quad P^c(\Pi) = \text{conv}(\Phi_{\text{short}}).$$

(iii) If the Cartan matrix C is type A_n or B_n , then $P(\Pi)$ is the dual polytope of $P^c(\Pi) = \text{conv}(\Phi_{\text{short}})$.

Remark 7.5. We remark that the g -simplicial complex $\Delta(\Pi)$ is nothing but the *Coxeter complex*. It is known that the h -vectors (they are the number of semibricks of $\text{mod } \Pi$ by Theorem 3.4) can be calculated as *W-Eulerian numbers* [Pe, Theorem 11.3]. More precisely, for $w \in W$, we define

$$\text{Des}(w) = \{s_i \mid \ell(w) > \ell(s_i w)\} \quad \text{and} \quad \text{des}(w) = |\text{Des}(w)|.$$

Then we call

$$\binom{W}{k} = |\{w \in W \mid \text{des}(w) = k\}|$$

the *W-Eulerian numbers* and define the *W-Eulerian polynomial* by

$$W(x) = \sum_{w \in W} x^{\text{des}(w)} = \sum_{k=0}^n \binom{W}{k} x^k.$$

Thus, the h -polynomial of $\Delta(\Pi)$ is given as *W-Eulerian polynomial*, and their real-rootedness are already known [Pe, Theorem 11.2].

TABLE 1. The Eulerian numbers of type A_n .

$n \setminus k$	0	1	2	3	4	5	6
1	1	1					
2	1	4	1				
3	1	11	11	1			
4	1	26	66	26	1		
5	1	57	302	302	57	1	
6	1	120	1191	2416	1191	120	1

From now on, we will give a proof of Theorem 7.4. For this purpose, following [ARS], we introduce some notations. Let $X, Y \in \text{mod } H$ be indecomposable modules. In the Auslander-Reiten quiver (AR quiver, for short), we write

$$X \xrightarrow{(a,b)} Y$$

TABLE 2. The Eulerian numbers of type B_n .

$n \setminus k$	0	1	2	3	4	5	6
2	1	6	1				
3	1	23	23	1			
4	1	76	230	76	1		
5	1	237	1682	1682	237	1	
6	1	722	10543	23548	10543	722	1

if there is a minimal right almost split $X^a \oplus M \rightarrow Y$ such that M contains no summands isomorphic to X , and there is a left right almost split $X \rightarrow Y^b \oplus N$ such that N contains no summands isomorphic to Y . Let $D_X := \text{End}_H(X)/\text{rad}_H(X, X)$ and $D_Y := \text{End}_H(Y)/\text{rad}_H(Y, Y)$. Then we have the following basic result.

Proposition 7.6. [ARS, section VII]

(a) If there exists $X \xrightarrow{(a,b)} Y$ in the AR quiver of $\text{mod } H$, then we have

$$\dim(\text{rad}_H(X, Y)/\text{rad}_H^2(X, Y))_{D_X} = a \quad \text{and} \quad \dim_{D_Y}(\text{rad}_H(X, Y)/\text{rad}_H^2(X, Y)) = b.$$

Moreover we have

$$a \dim_k(D_X) = b \dim_k(D_Y) = \dim_k(\text{rad}_H(X, Y)/\text{rad}_H^2(X, Y)).$$

(b) If there exists an arrow $Y \xrightarrow{(b,a)} Z$ in the AR quiver with non-projective Z , then there exists the arrow $\tau Z \xrightarrow{(a,b)} Y$.

Then we show the following proposition.

Proposition 7.7. Let H be a finite dimensional hereditary k -algebra and e_i a primitive idempotent of H and $Q_i := e_i H$ the indecomposable projective H -module. Then there exists the following almost split sequence

$$0 \longrightarrow Q_i \longrightarrow \bigoplus_{j: M_i \neq 0} Q_j^{-c_{ij}} \oplus \bigoplus_{i: M_k \neq 0} \tau^- Q_k^{-c_{ik}} \longrightarrow \tau^- Q_i \longrightarrow 0.$$

Proof. We denote the multiplicity of X in M by $m_X(M)$. Let $Q_i \rightarrow E$ be a minimal left almost split. By Proposition 7.6, if we have $Q_i \xrightarrow{(b,a)} \tau^- Q_k$, then we have $Q_k \xrightarrow{(a,b)} Q_i$ in the AR quiver. Since $\text{rad } Q_i \rightarrow Q_i$ is a minimal right almost split, we have $m_{\tau^- Q_k}(E) = m_{Q_k}(\text{rad } Q_i)$. Thus we have

$$m_{Q_k}(\text{rad } Q_i) = \dim((\text{rad } Q_i / \text{rad}^2 Q_i) e_k)_{D_k} = \dim({}_i M_k)_{D_k} = -c_{ik}.$$

On the other hand, Proposition 7.6 implies that, for $Q_i \xrightarrow{(a,b)} Q_j$, we have $a = m_{Q_i}(\text{rad } Q_j)$, $b = m_{Q_j}(E)$ and

$$\dim_k(\text{rad}_H(Q_i, Q_j)/\text{rad}_H^2(Q_i, Q_j)) = a \dim_k(D_{Q_i}) = b \dim_k(D_{Q_j}).$$

Since H is acyclic, we have $\dim_k(D_{Q_i}) = \dim_k(D_i) = c_i$ and $\dim_k(D_{Q_j}) = \dim_k(D_j) = c_j$. Thus, Proposition 7.6 also shows that we have

$$m_{Q_j}(E) = m_{Q_i}(\text{rad } Q_j) c_i / c_j = \dim({}_j M_i)_{D_i} c_i / c_j = -c_{ji} c_i / c_j = -c_{ij}.$$

and the conclusion follows. \square

Let Π be the preprojective algebra of H and we denote by $\{e_1, \dots, e_n\}$ be a complete set of primitive orthogonal idempotents of Π . For $1 \leq i \leq n$, we denote by $i^\pm := \{1 \leq j \leq n \mid {}_j M_i \neq 0 \text{ or } {}_i M_j \neq 0\}$.

Proposition 7.8. *Let H be a finite dimensional hereditary k -algebra of non-Dynkin type and Π the preprojective algebra of H . For any $1 \leq i \leq n$, we have the following minimal projective resolution*

$$0 \longrightarrow e_i \Pi \longrightarrow \bigoplus_{j \in i^\pm} e_j \Pi^{-c_{ij}} \longrightarrow e_i \Pi \longrightarrow D_i \longrightarrow 0.$$

Although this is quite standard fact, we give a sketch for the convenience of the reader.

Proof. Since the preprojective algebra Π does not depend on the orientation of $(D_i, {}_i M_j)$, we choose an orientation such that ${}_i M_k = 0$ for any k and hence Q_i is simple. Then, a minimal projective resolution of $Q_i = D_i$ is obtained as \mathbb{Z} -graded modules and its n -degree is given by applying τ^{-n} to the exact sequence of Proposition 7.7, see e.g. [MY, Proposition 3.7], [GI, Theorem 4.12], [Sö, Proposition 6.8]. Since $(e_i \Pi)_n \cong \tau^{-n}(Q_i)$, we have an exact sequence

$$0 \longrightarrow e_i \Pi \longrightarrow \bigoplus_{j \in i^\pm} e_j \Pi^{-c_{ij}} \longrightarrow e_i \Pi(1) \longrightarrow D_i(1) \longrightarrow 0$$

as \mathbb{Z} -graded modules and this is a minimal projective resolution of D_i . \square

Lemma 7.9. *Let H be a finite dimensional hereditary k -algebra of non-Dynkin type, Π the preprojective algebra of H and $I_i := \Pi(1 - e_i)\Pi$.*

(a) *I_i is a classical tilting Π -module and we have an equivalence*

$$-\otimes_{\Pi}^{\mathbf{L}} I_i : \mathbf{D}^b(\text{mod } \Pi) \rightarrow \mathbf{D}^b(\text{mod } \Pi).$$

(b) *The action $R_i := -\otimes_{\Pi} I_i : K_0(\text{proj } \Pi) \rightarrow K_0(\text{proj } \Pi)$, $[P] \mapsto [P \otimes_{\Pi} I_i]$ satisfies the following*

$$R_i(e_\ell \Pi) := \begin{cases} [e_\ell \Pi] & \ell \neq i, \\ -[e_i \Pi] + \sum_{j=1}^n c_{ij} [e_j \Pi] & \ell = i. \end{cases}$$

Proof. By Proposition 7.8, we have an exact sequence

$$0 \longrightarrow e_i \Pi \longrightarrow \bigoplus_{j \in i^\pm} e_j \Pi^{-c_{ij}} \longrightarrow e_i I_i \longrightarrow 0.$$

From this sequence, the results follow from by the same argument of [BIRS, section II.1] (some general situation is also discussed in [IR, section 6]). \square

From now on, let $H := T_D(M)$ be a finite dimensional hereditary k -algebra of Dynkin type and we denote by the corresponding matrix by C . Let Π be the preprojective algebra of H and $I_i := \Pi(1 - e_i)\Pi$, where e_i the primitive idempotent of Π . We denote by $\langle I_1, \dots, I_n \rangle$ the set of ideals of Π which can be written as

$$I_{i_1} I_{i_2} \cdots I_{i_\ell}$$

for some $k \geq 0$ and $i_1, \dots, i_\ell \in Q_0$. Then we have the following results.

Theorem 7.10. *There exists a bijection $W \rightarrow \langle I_1, \dots, I_n \rangle$, which is given by $w \mapsto I_w = I_{i_\ell} I_{i_{\ell-1}} \cdots I_{i_1}$ for any reduced expression $w = s_{i_1} \cdots s_{i_\ell}$. Moreover the map gives a bijection between*

$$W \longrightarrow \text{s}\tau\text{-tilt } \Pi.$$

Proof. In simply-laced case, the result is [M, Theorem 2.21]. In non-simply-laced case, results of [IR, BIRS] implies that the map is bijection. Using Proposition 7.8, the same argument of [M] shows that they are exactly support τ -tilting modules. \square

Moreover, we prepare the following set up. Let \tilde{H} be a finite dimensional hereditary k -algebra of affine type \tilde{C} whose restriction to 1 to n columns and 1 to n rows is C . Let $\tilde{\Pi}$ be the preprojective algebra of \tilde{H} such that $\tilde{\Pi}/\langle e_{n+1} \rangle \simeq \Pi$ and \tilde{W} the Coxeter group of \tilde{C} . As same as Dynkin type, we define $\tilde{I}_i := \tilde{\Pi}(1 - e_i)\tilde{\Pi}$ and \tilde{I}_w for $w \in \tilde{W}$. We naturally regard $W = \langle s_1, \dots, s_n \rangle$ as a subgroup of $\tilde{W} = \langle s_1, \dots, s_n, s_{n+1} \rangle$. Then we recall the following lemma.

Proposition 7.11. (a) *We have*

$$2\text{-silt}\Pi = \{\tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi \mid w \in W\}.$$

(b) *For any $w \in W$, we have*

$$[e_i \tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi] = w\alpha_i^*$$

Proof. (a) Theorem 7.10 shows $s\tau\text{-tilt}\Pi = \{I_w \mid w \in W\}$. Then [M, Proposition 5.2] (which works for arbitrary Dynkin type) implies that the correspondence $I_w \mapsto \tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi$ gives a bijective map $s\tau\text{-tilt}\Pi \rightarrow 2\text{-silt}\Pi$ of [AIR, Theorem 3.2].

(b) Since \tilde{I}_w is a classical tilting module, we have the following projective resolution

$$0 \longrightarrow \tilde{P}^1 \longrightarrow \tilde{P}^0 \longrightarrow e_i \tilde{I}_w \longrightarrow 0,$$

where $\tilde{P}^0, \tilde{P}^1 \in \text{proj } \tilde{\Pi}$. Therefore, we have $[e_i \tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi] = [\tilde{P}^0 \otimes_{\tilde{\Pi}} \Pi] - [\tilde{P}^1 \otimes_{\tilde{\Pi}} \Pi]$. Since $\Pi \simeq \tilde{\Pi}/\langle e_{n+1} \rangle$, $[\tilde{P}^j \otimes_{\tilde{\Pi}} \Pi]$ is given by the restriction of $[\tilde{P}^j]$ to $K_0(\text{proj } \Pi)_{\mathbb{R}}$ for $j \in \{0, 1\}$. Hence, Lemma 7.9 implies that we have $[e_i \tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi] = w\alpha_i^*$ in $K_0(\text{proj } \Pi)_{\mathbb{R}}$. \square

Then we are ready to give a proof Theorem 7.4 (a) and (b).

Proof of Theorem 7.4 (a) and (b). (a) Recall that $C(T) := \{\sum_{i=1}^j a_i [T_i] \mid a_i \geq 0\}$ for $T = T_1 \oplus \cdots \oplus T_n \in 2\text{-silt}\Pi$ with indecomposable T_i . Hence we have $C(\Pi) = \{\sum_{i=1}^j a_i [P_i] \mid a_i \geq 0\} = \{\sum_{i=1}^n a_i \alpha_i^* \mid a_i \geq 0\} = D$. Then, by Proposition 7.11 (b), we have

$$\begin{aligned} C(\tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi) &= \{\sum_{i=1}^n a_i [e_i \tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi] \mid a_i \geq 0\} \\ &= \{\sum_{i=1}^n a_i w\alpha_i^* \mid a_i \geq 0\} \\ &= w\{\sum_{i=1}^n a_i \alpha_i^* \mid a_i \geq 0\} \\ &= wD. \end{aligned}$$

Therefore, Proposition 7.11 (a) implies

$$\bigcup_{T \in 2\text{-silt}\Pi} C(T) = \bigcup_{w \in W} wD.$$

As a consequence, $\Sigma(\Pi) := \{C(T) \mid T \in 2\text{-psilt}\Pi\}$ is the set of all cones of wD and their faces, and we are done. By the same argument, we have $C_{\leq 1}(\tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi) = wC_{\leq 1}(\Pi)$ and hence

$$P(\Pi) = \bigcup_{T \in 2\text{-silt}\Pi} C_{\leq 1}(T) = \bigcup_{w \in W} wC_{\leq 1}(\Pi).$$

(b) Since

$$\pi := - \otimes_{\tilde{\Pi}} \Pi : K_0(\text{proj } \tilde{\Pi}) \rightarrow K_0(\text{proj } \Pi)$$

is compatible with the action of W , Theorem 5.10 and Proposition 7.11 implies that $P(\Pi)$ is convex if and only if $\sum_{j \in i^{\pm}} |c_{ij}| \leq 2$. This is equivalent to saying that the Cartan matrix C is type A_n or B_n . \square

For a proof of Theorem 7.4 (c), we use the following well-known facts.

Lemma 7.12. (a) *All roots of a given length are conjugate under W .*

(b) *The sum of all simple roots is a short root.*

Proof. (a) is [H2, III.10.4 Lemma C]. (b) can be checked by case-by-case analysis. \square

Then we give a proof of Theorem 7.4 (c).

Proof of Theorem 7.4 (c). Recall that $2\text{-silt}\Pi = \{\tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi \mid w \in W\}$ from Proposition 7.11. For simplicity, we let $P(w) := \tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi$ and $P(w)_i := e_i \tilde{I}_w \otimes_{\tilde{\Pi}}^{\mathbf{L}} \Pi$. By Proposition 2.13, there exists $S(w) \in 2\text{-smc}\Pi$ and $2\text{-smc}\Pi = \{S(w) \mid w \in W\}$ such that $S(w) = S(w)_1 \oplus \cdots \oplus S(w)_n$ satisfying

$$([P(w)_i], [S(w)_j]') = \delta_{ij}.$$

On the other hand, Proposition 7.11 implies that we have $[P(w)_i] = w\alpha_i^*$. Therefore, we have

$$([w\alpha_i^*, [S(w)_j]']) = \delta_{ij}.$$

Then, since the bilinear form $(-, -)$ is non-degenerate, we have $[S(w)_j]' = w\alpha_j$ and hence


$$\{[S(w)_j]' \mid 1 \leq j \leq n, w \in W\} = \{w\alpha_j \mid 1 \leq j \leq n, w \in W\}.$$

Thus (i) follows. Moreover, by this argument, we have


$$v_{S(w)} = \sum_{i=1}^n [S(w)_i]' = w \sum_{i=1}^n \alpha_i.$$

Then, Lemma 7.12 shows that $\{w(\sum_{i=1}^n \alpha_i) \mid w \in W\}$ coincide with all short roots of Φ and we get (ii). Finally, (iii) immediately follows from (b) and Theorem 5.14. \square


Example 7.13. (a) Let $H = \begin{pmatrix} \mathbb{R} & 0 \\ \mathbb{C} & \mathbb{C} \end{pmatrix}$. Then the corresponding Cartan matrix is $\begin{pmatrix} 2 & -1 \\ -2 & 2 \end{pmatrix}$. Let Π be the preprojective algebra of H . We obtain the Hasse quiver of $\text{s}\tau\text{-tilt}\Pi$ as follows.



On the other hand, the corresponding 2-simple minded collections are shown as follows.




Then, by taking $[-]'$, we have the following set of roots as follows



and hence we have


$$\{v_X \mid X \in 2\text{-smc}\Pi\} = \{\pm(\alpha_2, \alpha_1 + \alpha_2)\}.$$

Thus c -polytope $P^c(\Pi)$ is illustrated as a dotted line in the left picture below and the g -polytope $P(\Pi)$ is illustrated in the right picture below.



- (b) Let $H = \begin{pmatrix} \mathbb{R} & 0 & 0 \\ \mathbb{R} & \mathbb{R} & 0 \\ \mathbb{C} & \mathbb{C} & \mathbb{C} \end{pmatrix}$. Then the corresponding Cartan matrix is $\begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{pmatrix}$. Let Π be the preprojective algebra of H .

The root system of type B_3 is illustrated in the left picture below and the g -polytope $P(\Pi(B_3)) = (P^c(\Pi(B_3)))^*$ is illustrated in the right picture below.



7.2. Generalized preprojective algebras. In this subsection, we study g -polytopes of generalized preprojective algebras introduced by [GLS]. We show that the g -polytope is convex if and only if the Cartan matrices is type A_n or B_n .

First we introduce the notion of generalized preprojective algebras associated with symmetrizable generalized Cartan matrices [GLS].

Let $C = (c_{ij}) \in M_n(\mathbb{Z})$ be a symmetrizable generalized Cartan matrix. We denote by $g_{ij} := |\gcd(c_{ij}, c_{ji})|$ and $f_{ij} := |c_{ij}|/g_{ij}$.

An *orientation* of C is a subset Ω of $\{1, 2, \dots, n\} \times \{1, 2, \dots, n\}$ such that the followings hold:

- (i) $\{(i, j), (j, i)\} \cap \Omega \neq \emptyset$ if and only if $c_{ij} < 0$;
- (ii) For each sequence $((i_1, i_2), (i_2, i_3), \dots, (i_t, i_{t+1}))$ with $t \geq 1$ and $(i_s, i_{s+1}) \in \Omega$ for all $1 \leq s \leq t$, we have $i_1 \neq i_{t+1}$.

The *opposite orientation* of an orientation Ω is defined as $\Omega^* := \{(j, i) \mid (i, j) \in \Omega\}$. Let $\overline{\Omega} := \Omega \cup \Omega^*$ and we define

$$\overline{\Omega}(-, i) := \{j \in Q_0 \mid (j, i) \in \overline{\Omega}\}$$

For an orientation Ω of C , define the quiver $Q := Q(C, \Omega) := (Q_0, Q_1)$ with the set of vertices $Q_0 := \{1, \dots, n\}$ and with the set of arrows

$$Q_1 := \{\alpha_{ij}^{(g)} : j \rightarrow i \mid (i, j) \in \Omega, 1 \leq g \leq g_{ij}\} \cup \{\epsilon_i : i \rightarrow i \mid 1 \leq i \leq n\}.$$

We call Q a *quiver of type C* . Let $Q^\circ := Q^\circ(C, \Omega)$ be the quiver obtained from Q by deleting all loops ϵ_i .

Then we define the generalized preprojective algebra associated to C as follows.

Definition 7.14. Let C be a symmetrizable Cartan matrix with a symmetrizer D .

For $(i, j) \in \overline{\Omega}$, define

$$\text{sgn}(i, j) := \begin{cases} 1 & \text{if } (i, j) \in \Omega, \\ -1 & \text{if } (i, j) \in \Omega^*. \end{cases}$$

For $Q = Q(C, \Omega)$ and a symmetrizer $D = \text{diag}(c_1, \dots, c_n)$ of C , we define an algebra

$$\Pi = \Pi(C, D, \Omega) := K\overline{Q}/\overline{I}$$

as follows. The *double quiver* $\overline{Q} = \overline{Q}(C)$ is obtained from Q by adding a new arrow $\alpha_{ji}^{(g)} : j \rightarrow i$ for each arrow $\alpha_{ij}^{(g)} : i \rightarrow j$ of Q° .

The ideal \overline{I} of the path algebra $K\overline{Q}$ is defined by the following relations:

- (P1) For each $i \in Q_0$, we have the *nilpotency relation*

$$\epsilon_i^{c_i} = 0.$$

- (P2) For each $(i, j) \in \overline{\Omega}$ and each $1 \leq g \leq g_{ij}$, we have the *commutativity relation*

$$\alpha_{ij}^{(g)} \epsilon_i^{f_{ij}} = \epsilon_j^{f_{ji}} \alpha_{ij}^{(g)}.$$

(P3) For each $i \in Q_0$, we have the *mesh relation*

$$\sum_{j \in \overline{\Omega}(-, i)} \sum_{g=1}^{g_{ji}} \sum_{f=0}^{f_{ij}-1} \operatorname{sgn}(i, j) \epsilon_i^{f_{ij}-1-f} \alpha_{ji}^{(g)} \alpha_{ij}^{(g)} \epsilon_i^f = 0.$$

Remark 7.15. Our definition of the preprojective algebra is slightly different from the original one given by [GLS] (which we denote by Π^{GLS}), but two definitions essentially give same objects. More precisely, we have $\Pi(C, D) = \Pi^{GLS}({}^t C, D)$, where ${}^t C$ is the transposed Cartan matrix.

We remark that Π does not depend on the orientation Ω of C , so that we can write $\Pi = \Pi(C, D)$.

For the one dimensional simple Π -modules S_1, \dots, S_n , we define the $\iota : K_0(\operatorname{mod} \Pi)_{\mathbb{R}} \simeq V$ by $\iota([S_i]) = \alpha_i$ and, for $X = X_1 \oplus \dots \oplus X_n \in \operatorname{smc} \Pi$, let

$$v_X := \iota\left(\sum_{i=1}^n [X_i]\right) \in V.$$

Then we have the following analogous result of Theorem 7.4, where the only difference is that $[S]'$ in Theorem 7.4(c)(i) is replaced by $[S]$.

Theorem 7.16. *Let C be a symmetrizable Cartan matrix of Dynkin type with a symmetrizer D and $\Pi = \Pi(C, D)$.*

(a) $\Sigma(\Pi)$ is the Coxeter fan and we have

$$P(\Pi) = \bigcup_{w \in W} wC_{\leq 1}(\Pi).$$

(b) $P(\Pi)$ is convex if and only if the Cartan matrix C is type A_n or B_n .

(c) (i) We have

$$\{[S] \mid S \in \operatorname{ind-2-smc} \Pi\} = \Phi.$$

(ii) Let Φ_{short} be the set of short roots of Φ . We have

$$\{v_X \mid X \in 2\text{-smc} \Pi\} = \Phi_{\text{short}} \quad \text{and} \quad P^c(\Pi) = \operatorname{conv}(\Phi_{\text{short}}).$$

(iii) If the Cartan matrix C is type A_n or B_n , then $P(\Pi)$ is the dual polytope of $P^c(\Pi) = \operatorname{conv}(\Phi_{\text{short}})$.

Since a proof of Theorem 7.16 follows from an analogous argument of Theorem 7.4, we only emphasis on the difference.

Instead of Proposition 7.8, we use the following result.

Proposition 7.17. [FG] *Let C be a symmetrizable Cartan matrix of non-Dynkin type with a symmetrizer D and $\Pi = \Pi(C, D)$. For any $1 \leq i \leq n$, we have the following minimal projective resolution¹*

$$0 \longrightarrow e_i \Pi \longrightarrow \bigoplus_{j \in i^{\pm}} e_j \Pi^{-c_{ij}} \longrightarrow e_i \Pi \longrightarrow \hat{S}_i \longrightarrow 0,$$

where \hat{S}_i denotes by the generalized simple module. In particular, we have an exact sequence

$$0 \longrightarrow e_i \Pi \longrightarrow \bigoplus_{j \in i^{\pm}} e_j \Pi^{-c_{ij}} \longrightarrow I_i \longrightarrow 0.$$

Then, by combining with results of [M, FG], Theorem 7.16 (a) and (b) follows from the same argument of subsection 7.1. To show Theorem 7.16 (c), it is enough to show the following result.

¹Because of our definition, we use the index c_{ij} instead of c_{ji} used in [FG]

Lemma 7.18. *For any $S \in \text{ind-2-smc}\tilde{\Pi}$, we have*

$$\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(S) \cong k.$$

In particular,

$$[S] = [S]'$$

For a proof, we recall results of [KM] and prepare the following setting. Let \tilde{C} be a affine Cartan matrix whose restriction to 1 to n columns and 1 to n rows is C . Let $\tilde{\Pi}$ be the complete preprojective algebra of \tilde{C} with a symmetrizer. Then we have the following result.

Proposition 7.19. *For any $T \in 2\text{-silt}\tilde{\Pi}$, we have*

$$\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(T) \cong \tilde{\Pi}.$$

Moreover, for any brick S , we have $\text{End}_{\tilde{\Pi}}(S) \cong k$.

Proof. Recall from [KM] that any 2-silting complex is tilting complex and it belongs to one of the two connected components of mutation (which also works for non-simply laced cases by [FG]). One component contains $\tilde{\Pi}$ and the other contains $\tilde{\Pi}[1]$. Then by [FG, BIRS], the endomorphism algebra of them is isomorphic to $\tilde{\Pi}$. Thus, for $T_i \in 2\text{-psilt}^1\tilde{\Pi}$, we have $\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(T_i) \cong e_i\tilde{\Pi}e_i$ for an idempotent e_i of $\tilde{\Pi}$. Hence $\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(T_i)/\text{rad}(\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(T_i)) \simeq k$. On the other hand, for any $S \in \text{ind-2-smc}\tilde{\Pi}$, there exists $T_i \in 2\text{-psilt}^1\tilde{\Pi}$ such that

$$\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(T_i)/\text{rad}(\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(T_i)) \cong \text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(S).$$

Therefore, we get $\text{End}_{\mathcal{D}^b(\text{mod } \tilde{\Pi})}(S) \cong k$. \square

We give a proof of Lemma 7.18.

Lemma 7.18. Because $\tilde{\Pi} = \tilde{\Pi}/\langle e_{n+1} \rangle$, $\text{mod } \tilde{\Pi}$ is a full subcategory of $\text{mod } \tilde{\Pi}$. Therefore, Proposition 7.19 implies that we conclude $\text{End}_{\tilde{\Pi}}(S) \cong k$ for any brick S of $\text{mod } \tilde{\Pi}$. \square

8. JACOBIAN ALGEBRAS AND CLUSTER ALGEBRAS

Path algebras of quivers and more generally, Jacobian algebras of quivers with potential are important classes of algebras. They are used to categorify cluster algebras of Fomin and Zelevinsky. Cluster variables of a fixed cluster algebra $\mathcal{A}(B)$ are parametrized by integer vectors called g -vectors, defined inductively by a purely combinatorial formula. They give a fan $\Sigma(B)$ which we call the *cluster fan*. The aim of this section is to show that the g -fans of Jacobian algebras coincide with the cluster fans.

8.1. g -vectors and c -vectors of cluster algebras. To recall the notion of g -fans of cluster algebras, we prepare some notations. For $x \in \mathbb{R}$, we write

$$[x]_+ := \min\{x, 0\}.$$

Let $B = (b_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$ be an $m \times n$ matrix with $m \geq n$. For each $1 \leq k \leq n$, we define a new $m \times n$ matrix $\mu_k(B) = (b'_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$ by

$$b'_{ij} := \begin{cases} -b_{ij} & i = k \text{ or } j = k \\ b_{ij} + [b_{ik}]_+ [b_{kj}]_+ - [-b_{ik}]_+ [-b_{kj}]_+ & \text{else.} \end{cases}$$

We denote by \mathbb{T}_n the n -regular tree whose edges are labeled by the numbers $1, \dots, n$, so that the n edges adjacent to each vertex have different labels, and fix a vertex $t_0 \in \mathbb{T}_n$.

The g -fan is defined for each fixed $n \times n$ skew-symmetric integer matrix

$$B = [b_1^0, \dots, b_n^0].$$

Recall that $n \times n$ skew-symmetric integer matrices B correspond bijectively with quivers Q with vertices $Q_0 = \{1, \dots, n\}$ which do not have loops and 2-cycles. The matrix corresponding to such

a quiver Q is given by $B(Q) := (b_{ij})_{1 \leq i \leq n, 1 \leq j \leq n}$, where b_{ij} is the number of arrows from i to j in Q minus the number of arrows from j to i in Q .

We define a $2n \times n$ matrix $\tilde{B}_t = (b_{ij}^t)_{1 \leq i \leq 2n, 1 \leq j \leq n}$ for each $t \in \mathbb{T}_n$ inductively by equalities

$$\tilde{B}_{t_0} := \begin{bmatrix} B \\ I_n \end{bmatrix} \quad \text{and} \quad \mu_k(\tilde{B}_t) = \tilde{B}_{t'} \quad \text{for each } t, t' \in \mathbb{T}_n \text{ satisfying } t \xrightarrow{k} t'.$$

Definition 8.1. Let e_1, \dots, e_n be the standard basis of \mathbb{Z}^n .

(a) For $t \in \mathbb{T}_n$, we define the c -vectors $c_{1t}, \dots, c_{nt} \in \mathbb{Z}^n$ as the lower n entries of the columns of \tilde{B}_t :

$$c_{jt} = (c_{ijt})_{1 \leq i \leq n} := (b_{n+i,j}^t)_{1 \leq i \leq n} \quad \text{for } 1 \leq j \leq n.$$

In other words, let $c_{it_0} := e_i$ for $1 \leq i \leq n$, and for each $t, t' \in \mathbb{T}_n$ satisfying $t \xrightarrow{k} t'$, we define

$$c_{ijt'} := \begin{cases} -c_{ijt} & \text{if } j = k, \\ c_{ijt} + [c_{ikt}]_+ [b_{kj}^t]_+ - [-c_{ikt}]_+ [-b_{kj}^t]_+ & \text{otherwise.} \end{cases} \quad (8.1)$$

The c -matrix is $C_t := [c_{1t}, \dots, c_{nt}]$.

(b) For $t \in \mathbb{T}_n$, we define the g -vectors $g_{1t}, \dots, g_{nt} \in \mathbb{Z}^n$ inductively as follows: Let

$$g_{it_0} := e_i \quad \text{for } 1 \leq i \leq n.$$


For each $t, t' \in \mathbb{T}_n$ satisfying $t \xrightarrow{k} t'$, we define

$$\begin{aligned} g_{jt'} &:= g_{jt} \quad \text{for each } j \neq k, \\ g_{kt'} &:= -g_{kt} + \sum_{i=1}^n [b_{ik}^t]_+ g_{it} - \sum_{i=1}^n [b_{n+i,k}^t]_+ b_i^0 \\ &= -g_{kt} + \sum_{i=1}^n [-b_{ik}^t]_+ g_{it} - \sum_{i=1}^n [-b_{n+i,k}^t]_+ b_i^0, \end{aligned} \quad (8.2)$$

where the last equality follows from [FZ3, (6.13)]. The g -matrix is $G_t := [g_{1t}, \dots, g_{nt}]$.

We omit the definition of the cluster algebra $\mathcal{A}(B)$ with principal coefficients and the x -variable x_{it} since we will not use them in this article. We note that $\mathcal{A}(B)$ has a \mathbb{Z}^n -grading, and the g -vector g_{it} is a degree of x_{it} .

Example 8.2. Let $B = \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$. Then the matrices \tilde{B}_t (written as a quiver), the c -matrices (written above) and the g -matrices (written below) are the following.



8.2. Jacobian algebras. Let Q be a quiver, and \widehat{kQ} the complete path algebra over a field k equipped with the $\text{rad } \widehat{kQ}$ -adic topology. We call (Q, W) a quiver with *potential* (or *QP* in short) if $W \in \widehat{kQ}$ is a formal linear combination of cycles of length at least one. For an arrow $a \in Q_1$, the *cyclic derivative* $\partial_a W \in \widehat{kQ}$ is defined by

$$\partial_a(a_1 a_2 \cdots a_\ell) := \sum_{a_i=a} a_{i+1} a_{i+2} \cdots a_{i-1}$$

for a cycle $a_1 a_2 \cdots a_\ell$ and extended linearly and continuously.

Definition 8.3. Let (Q, W) be a QP. The *Jacobian ideal* of (Q, W) is the ideal of \widehat{kQ} defined by

$$J(Q, W) := \overline{\langle \partial_a W \mid a \in Q_1 \rangle},$$

where $\overline{(-)}$ is the closure with respect to the $\text{rad } \widehat{kQ}$ -adic topology. The *Jacobian algebra* of (Q, W) is defined by

$$P(Q, W) := \widehat{kQ} / J(Q, W).$$

In the rest of this subsection, let (Q, W) be a QP with $Q_0 = \{1, \dots, n\}$, and $A := P(Q, W)$. Although $\dim_k A$ is infinite in general, we can apply mutation to A repeatedly, see Proposition 8.10 below. For each $t \in \mathbb{T}_n$, we define $T_t = \bigoplus_{i=1}^n T_{it} \in 2\text{-silt } A$ and $S_t = \bigoplus_{i=1}^n S_{it} \in 2\text{-smc } A$ by

$$T_{it_0} = e_i A \quad \text{and} \quad S_{it_0} = e_i A / \text{rad } e_i A \quad \text{for } 1 \leq i \leq n$$

and for each $t, t' \in \mathbb{T}_n$ satisfying $t \xrightarrow{k} t'$, let

$$T_{t'} := \mu_k(T_t) \quad \text{and} \quad S_{t'} = \mu_k(S_t).$$

The indices of the direct summands of $T_{t'}$ and $S_{t'}$ are defined naturally: $T_{it'} = T_{it}$ for each $i \neq k$ and $T_{kt'}$ is the unique indecomposable direct summand of $T_{t'}$ which does not appear in T_t . Moreover, $\dim_k \text{Hom}_{\mathbb{D}^b(\text{mod } A)}(S_{it'}, S_{jt'}[\ell]) = \delta_{ij} \delta_{\ell 0}$ for each $1 \leq i, j \leq n$ and $\ell \in \mathbb{Z}$. Then

$$2\text{-silt}^\circ A := \{T_t \mid t \in \mathbb{T}_0\}$$

consists of all iterated mutations of A . We consider isomorphisms

$$\begin{aligned} \iota_p : \mathbb{Z}^n &\simeq K_0(\text{proj } A) & \text{given by} & \quad \iota_p(e_i) := [e_i A], \\ \iota_s : \mathbb{Z}^n &\simeq K_0(\text{mod } A) & \text{given by} & \quad \iota_s(e_i) := [e_i A / \text{rad } e_i A]. \end{aligned}$$

Theorem 8.4 below is fundamental in categorification of cluster algebras, where the definition of non-degenerate QPs will be given in the next subsection (Definition 8.8). Notice that (b),(c) and (d) were conjectured in [FZ3, Conjecture 7.10], [FZ3, Conjecture 5.4, Proposition 5.6] and [FZ3, Conjecture 6.13] respectively. They were proved in [DWZ, Nag, NZ, P11] (see also [GHKK] for skew-symmetrizable case). In the next subsection, we give a proof of Theorem 8.4 as easy consequences of basic results in silting theory.

Theorem 8.4. *Let (Q, W) be a non-degenerate QP with $Q_0 = \{1, \dots, n\}$, $A := P(Q, W)$ the Jacobian algebra, and $B := B(Q)$ the skew-symmetric matrix. For each $t \in \mathbb{T}_n$, the following assertions hold.*

(a) *For each $1 \leq i \leq n$, we have*

$$\iota_p(g_{it}) = [T_{it}] \quad \text{and} \quad \iota_s(c_{it}) = [S_{it}].$$

- (b) *The g -vectors g_{1t}, \dots, g_{nt} and the c -vectors c_{1t}, \dots, c_{nt} are the \mathbb{Z} -basis of \mathbb{Z}^n . They are dual to each other with respect to the canonical pairing $\mathbb{Z}^n \times \mathbb{Z}^n \rightarrow \mathbb{Z}$.*
- (c) *For each $1 \leq i \leq n$, the c -vector c_{it} belongs to precisely one of $\mathbb{Z}_{\geq 0}^n$ and $\mathbb{Z}_{\leq 0}^n$.*
- (d) *The g -vectors g_{1t}, \dots, g_{nt} are sign-coherent, that is, for each $1 \leq i \leq n$, the vector $(g_{i1t}, \dots, g_{int})$ belongs to precisely one of $\mathbb{Z}_{\geq 0}^n$ and $\mathbb{Z}_{\leq 0}^n$.*

(e) (8.1) and (8.2) are simplified as follows:

$$\begin{aligned} c_{jt'} &:= \begin{cases} -c_{jt} & \text{if } j = k, \\ c_{jt} + [b_{kj}^t] + c_{kt} & \text{if } j \neq k \text{ and } c_{kt} \in \mathbb{Z}_{\geq 0}^n, \\ c_{jt} + [-b_{kj}^t] + c_{kt} & \text{if } j \neq k \text{ and } c_{kt} \in \mathbb{Z}_{\leq 0}^n. \end{cases} \\ g_{jt'} &:= \begin{cases} g_{jt} & \text{if } j \neq k, \\ -g_{kt} + \sum_{i=1}^n [-b_{ik}^t] + g_{it} & \text{if } j = k \text{ and } c_{kt} \in \mathbb{Z}_{\geq 0}^n, \\ -g_{kt} + \sum_{i=1}^n [b_{ik}^t] + g_{it} & \text{if } j = k \text{ and } c_{kt} \in \mathbb{Z}_{\leq 0}^n. \end{cases} \end{aligned}$$

We denote by $\Sigma^\circ(A)$ the fan corresponding to the subset $2\text{-silt}^\circ A$ of $2\text{-silt} A$. The following main result of this section shows that $\Sigma^\circ(A)$ is described by the g -vectors of the corresponding cluster algebra, where Σ_n denotes the set of cones of dimension n . It also gives a conceptual reason why the g -vectors of the cluster algebra gives a non-singular cluster fan $\Sigma(B)$.

Theorem 8.5. *Let (Q, W) be a non-degenerate quiver with potential with $Q_0 = \{1, \dots, n\}$, $A := P(Q, W)$ the Jacobian algebra, and $B := B(Q)$ the skew-symmetric matrix.*

(a) *A sign-coherent nonsingular fan in \mathbb{R}^n is given by*

$$\Sigma(B) := \{\text{cone}\{g_{jt} \mid j \in I\} \mid I \subset \{1, \dots, n\}, t \in \mathbb{T}_n\}.$$

(b) *The isomorphism $\iota_p : \mathbb{Z}^n \simeq K_0(\text{proj } A)$ given by $e_i \mapsto e_i A$ gives an isomorphism of fans*


$$\Sigma(B) \simeq \Sigma^\circ(A).$$

(c) *There are bijections*

$$\{\text{clusters of } \mathcal{A}(B)\} \simeq \Sigma_n(B) \simeq \Sigma_n^\circ(A).$$

For more information on the cluster fan $\Sigma(B)$, we refer to [Nak, NZ, Re, Y2] and also [BCDMTY, BFMN, CFZ, Fe2, FZ1, HLT, HPS, INT, PPPP].

Example 8.6. Let $B = \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$ be as in Example 8.2. Then $A = kQ$ for $Q = [1 \Rightarrow 2]$, and $\Sigma(B) = \Sigma(A)$ is the following:



Applying Theorem 8.5, we obtain the following results.

Corollary 8.7. *Let (Q, W) be a non-degenerate quiver with potential, $A := P(Q, W)$ the Jacobian algebra, and $B := B(Q)$ the skew-symmetric matrix.*

(a) *A is g -finite if and only if Q is mutation equivalent to a Dynkin quiver.*

(b) *A is g -convex if and only if Q is mutation equivalent to a quiver of type A.*

8.3. Proof of Theorems 8.4, 8.5 and Corollary 8.7. We start with reminding relevant notions. Let Q be a quiver. We call two potentials W and W' on Q *cyclically equivalent* if $W - W'$ belongs to the closure of the space $\langle [x, y] \mid x, y \in \widehat{kQ} \rangle_k$ spanned by the commutators. In this case, $J(Q, W) = J(Q, W')$ and hence $P(Q, W) = P(Q, W')$ hold.

Let (Q, W) and (Q', W') be QPs satisfying $Q_0 = Q'_0$. They are called *right-equivalent* if there exists an isomorphism $\phi : \widehat{kQ} \simeq \widehat{kQ'}$ of k -algebras such that ϕ is the identity on Q_0 and $\phi(W)$ is cyclically equivalent to W' . In this case, ϕ induces an isomorphism $P(Q, W) \simeq P(Q', W')$. On the other hand, we define a *direct sum* of (Q, W) and (Q', W') by

$$(Q, W) \oplus (Q', W') := (Q'', W + W'),$$

where Q'' is the quiver $(Q_0, Q_1 \sqcup Q'_1)$.

A QP (Q, W) is called *reduced* if all cycles appearing in W have length at least 3, and *trivial* if $J(Q, W) = \text{rad } \widehat{kQ}$. It is shown in [DWZ, Theorem 4.6] that, for each QP (Q, W) , there exists

a reduced QP $(Q_{\text{red}}, W_{\text{red}})$ (called the *reduced part* of (Q, W)) and a trivial QP $(Q_{\text{triv}}, W_{\text{triv}})$ such that (Q, W) is right equivalent to $(Q_{\text{red}}, W_{\text{red}}) \oplus (Q_{\text{triv}}, W_{\text{triv}})$, and that $(Q_{\text{red}}, W_{\text{red}})$ and $(Q_{\text{triv}}, W_{\text{triv}})$ are unique up to right equivalence.

Using these notions, we are now able to introduce mutation of QPs. For a QP (Q, W) and a vertex $k \in Q_0$, we define new QPs $\tilde{\mu}_k(Q, W) = (Q', W')$ and $\mu_k(Q, W)$ as follows.

- Q' is a quiver obtained from Q by the following changes.
 - (i) Add a new arrow $[ab] : i \rightarrow j$ for each pair of arrows $a : i \rightarrow k$ and $b : k \rightarrow j$.
 - (ii) Replace each arrow $a : i \rightarrow k$ in Q by a new arrow $a^* : k \rightarrow i$.
 - (iii) Replace each arrow $b : k \rightarrow j$ in Q by a new arrow $b^* : j \rightarrow k$.
- $W' = [W] + \Delta$, where $[W]$ and Δ are the following.
 - (i) $[W]$ is obtained by substituting $[ab]$ for each factor ab in W with $a : i \rightarrow k$ and $j : k \rightarrow j$.
 - (ii) $\Delta = \sum_{a, b \in Q_1, t(a)=k=s(b)} a^*[ab]b^*$.
- $\mu_k(Q, W)$ is the reduced part of $\tilde{\mu}_k(Q, W)$.

We are ready to define non-degenerate QPs.

Definition 8.8. Let (Q, W) be a QP with $Q_0 = \{1, \dots, n\}$. For each $t \in \mathbb{T}_n$, we define a QP (Q_t, W_t) by $(Q_{t_0}, W_{t_0}) := (Q, W)$ and for each $t, t' \in \mathbb{T}_n$ satisfying $t \xrightarrow{k} t'$, let

$$(Q_{t'}, W_{t'}) := \mu_k(Q_t, W_t).$$

We call a QP (Q, W) *non-degenerate* if Q_t does not have 2-cycles for each $t \in \mathbb{T}_n$. In this case, $B(Q_t)$ is the $n \times n$ upper part of \tilde{B}_t for each $t \in \mathbb{T}_n$.

Now we introduce the Ginzburg dg algebra of a QP.

Definition 8.9. Let (Q, W) be a QP. Let \tilde{Q} be a graded quiver with the same vertices as Q and the following three kinds of arrows.

- All arrows of Q . Their degrees are defined to be zero.
- An arrow $a^* : j \rightarrow i$ of degree -1 for each arrow $a : i \rightarrow j$ in Q .
- A loop $t_i : i \rightarrow i$ of degree -2 for each $i \in Q_0$.

We denote by $\widehat{k\tilde{Q}}$ the complete graded path algebra of \tilde{Q} . The (complete) *Ginzburg dg algebra* of (Q, W) is $(\widehat{k\tilde{Q}}, d)$, where the differential $d : \widehat{k\tilde{Q}} \rightarrow \widehat{k\tilde{Q}}$ is the continuous linear map satisfying

- $d(a) = 0$ for each $a \in Q_1$,
- $d(a^*) = \partial_a W$ for each $a \in Q_1$,
- $d(t_i) = e_i(\sum_{a \in Q_1} (aa^* - a^*a))e_i$ for each $i \in Q_0$.

We collect basic properties of Ginzburg dg algebras. Let (Q, W) be a QP with $Q_0 = \{1, \dots, n\}$, and $\Gamma := \Gamma(Q, W)$. For each $t \in \mathbb{T}_n$, we define $T_t \in 2\text{-silt}\Gamma$ by $T_{t_0} := \Gamma$ and for each $t, t' \in \mathbb{T}_n$ satisfying $t \xrightarrow{k} t'$, let $T_{t'} := \mu_k(T_t)$.

Proposition 8.10. *Let (Q, W) be a QP with $Q_0 = \{1, \dots, n\}$. Then the following assertions hold.*

- (a) $\Gamma(Q, W)$ is a non-positive dg algebra such that $H^0(\Gamma(Q, W)) \simeq P(Q, W)$ as k -algebras.
- (b) $P(Q, W)$ and $\Gamma(Q, W)$ satisfy (A), (B) in Assumption 1.18 and the following condition.
 - (C') For each iterated mutation $T \in 2\text{-silt}A$ of A and an indecomposable direct summand X of T , A has a left $\text{add}(T/X)$ -approximation and $A[1]$ has a right $\text{add}(T/X)$ -approximation.
- (c) For each $t \in \mathbb{T}_n$, the DG endomorphism algebra of T_t is quasi-isomorphic to $\Gamma(Q_t, W_t)$, and $\text{End}_{\mathcal{D}(P(Q, W))}(T_t) \simeq P(Q_t, W_t)$.
- (d) If (Q, W) is non-degenerate, then $B_t = B(Q_t)$ for each $t \in \mathbb{T}_n$.

Proof. (a) is clear, and (c) follows from [KeY, Theorem 3.2] inductively. Thus (d) is immediate from (c). Now (b) follows from [KeY, Lemma 2.17] and (c). \square

Now we are ready to prove Theorem 8.4.

Proof of Theorem 8.4. For a fixed $t \in \mathbb{T}_n$, we denote by $(a)_t, \dots, (e)_t$ the assertions (a), \dots , (e) for t . Using induction, we prove $(a)_t, \dots, (e)_t$ for each $t \in \mathbb{T}_n$.

Since $g_{it_0} = e_i = c_{it_0}$ for $1 \leq i \leq n$, each of $(a)_{t_0}, \dots, (e)_{t_0}$ hold true.

Take $t, t' \in \mathbb{T}_n$ satisfying $t \xrightarrow{k} t'$, and assume that $(a)_t, \dots, (e)_t$ hold true. We will prove that $(a)_{t'}, \dots, (e)_{t'}$ hold true.

It suffices to prove $(a)_{t'}$. In fact, $(b)_{t'}, (c)_{t'}$ and $(d)_{t'}$ are basic properties of 2-term sifting complexes and 2-term simple-minded collections. $(b)_{t'}$ follows from Proposition 2.13. Also $(e)_{t'}$ follows immediately from $(c)_{t'}$, (8.1) and (8.2).

$(a)_{t'}$ For each $j \neq k$, we have $T_{jt} = T_{jt'}$ and hence

$$\iota_p(g_{jt'}) \stackrel{(8.2)}{=} \iota_p(g_{jt}) \stackrel{(a)_t}{=} [T_{jt}] = [T_{jt'}].$$

By $(c)_t$, precisely one of the following statements holds.

$$(i) \ c_{kt} \in \mathbb{Z}_{\geq 0}^n, \quad (ii) \ c_{kt} \in \mathbb{Z}_{\leq 0}^n.$$

Since $[S_{kt}] = \iota_s(c_{kt})$ holds by $(a)_t$, we have $S_{kt} \in \mathbf{mod} A$ in case (i) and $S_{kt} \in (\mathbf{mod} A)[1]$ in case (ii). By Proposition 3.15(c), $T_t > T_{t'}$ holds in case (i) and $T_t < T_{t'}$ holds in case (ii). By Proposition 8.10(d), the quiver of $\text{End}_{\mathbf{D}(A)}(T_t)$ is Q_t , and the exchange sequence has the form

$$\begin{aligned} T_{kt} &\rightarrow \bigoplus_{1 \leq i \leq n} T_{it}^{\oplus [b_{ki}^t]_+} \rightarrow T_{kt'} \rightarrow T_{kt}[1] \quad \text{in case (i),} \\ T_{kt'} &\rightarrow \bigoplus_{1 \leq i \leq n} T_{it}^{\oplus [b_{ik}^t]_+} \rightarrow T_{kt} \rightarrow T_{kt'}[1] \quad \text{in case (ii).} \end{aligned}$$

In particular, we have

$$\begin{aligned} [T_{kt'}] &= -[T_{kt}] + \sum_{i=1}^n [b_{ki}^t]_+ [T_{it}] \stackrel{(a)_t}{=} \iota_p(-g_{kt} + \sum_{i=1}^n [-b_{ik}^t]_+ g_{it}) \stackrel{(e)_t}{=} \iota_p(g_{kt'}) \quad \text{in case (i),} \\ [T_{kt'}] &= -[T_{kt}] + \sum_{i=1}^n [b_{ik}^t]_+ [T_{it}] \stackrel{(a)_t}{=} \iota_p(-g_{kt} + \sum_{i=1}^n [b_{ik}^t]_+ g_{it}) \stackrel{(e)_t}{=} \iota_p(g_{kt'}) \quad \text{in case (ii),} \end{aligned}$$

as desired.

By $(b)_t$ and the equalities in $(e)_t$, we obtain $(b)_{t'}$, that is, $g_{1t'}, \dots, g_{nt'}$ and $c_{1t'}, \dots, c_{nt'}$ are dual basis of each other. Since $g_{jt'} = [T_{jt'}]$ holds for $1 \leq j \leq n$, Proposition 2.13 implies that $c_{jt'} = [S_{jt'}]$ holds for $1 \leq j \leq n$. Thus $(a)_{t'}$ holds, and we completed the proof. \square

We are ready to prove Theorem 8.5.

Proof of Theorem 8.5. For each $t \in \mathbb{T}_n$ and $I \subset \{1, \dots, n\}$, Theorem 8.4(a) implies

$$C(\bigoplus_{i \in I} T_{it}) = \text{cone}\{g_{jt} \mid j \in I\}.$$

Thus the assertion (b) holds.

Now (a) follows immediately from (b), and (c) follows from (b) and [CKLP, Corollary 3.5]. \square

Corollary 8.7 is an easy consequence.

Proof of Corollary 8.7. (a) By Theorem 8.5(c), A is g -finite if and only if there are only finitely many clusters. By the classification of finite type cluster algebras [FZ2], this is equivalent to that Q is mutation equivalent to a Dynkin quiver.

(b) By Theorem 5.10, A is g -convex if and only if A is g -finite and pairwise g -convex. Thus Q is mutation equivalent to a Dynkin quiver, and each quiver Q' which is mutation equivalent to Q satisfies that for each $i \in Q'_0$, at most two arrows start at i and at most two arrow end at i . This implies that Q is mutation equivalent to a quiver of type A . \square

9. BRAUER GRAPH ALGEBRAS AND ROOT POLYTOPES

In this section, we study g -polytopes of Brauer graph algebras. We show that every g -finite Brauer graph algebra is g -convex and the g -polytope is isomorphic to the root polytope of type A_n or C_n (Theorem 9.4), where n is the number of edges of the associated Brauer graph.


9.1. Definition and main result. A Brauer graph algebra is defined from a combinatorial object called Brauer graph. We refer to a survey paper [Sc] for the background and basic properties of Brauer graph algebras.

Definition 9.1. A *ribbon graph* is a triple $\Gamma = (H, \sigma, \overline{})$, where H is a non-empty finite set, σ is a permutation on H , and $\overline{}: H \rightarrow H$ is a fixed-point free involution.

- Each element of H is called *half-edge* of Γ .
- Let $H \rightarrow H/\overline{}$ be a canonical surjection. Each element of $H/\overline{}$ is called *edge* of Γ . We denote by $[h]$ the edge containing $h \in H$. Then, we have $[h] = [\overline{h}]$.
- Let $s: H \rightarrow H/\langle \sigma \rangle$ be a canonical surjection. Each element of $H/\langle \sigma \rangle$ is called *vertex* of Γ .
- For each vertex v of Γ , the σ -orbit $(h, \sigma(h), \dots, \sigma^{\ell-1}(h))$ incident to v is called the *cyclic ordering* around v , where ℓ is the cardinality of this orbit.

A *Brauer graph* is a ribbon graph Γ equipped with a *multiplicity function*, which assigns a positive integer $\mathbf{m}(v) > 0$, called *multiplicity*, for every vertex v of Γ .

In order to describe a given Brauer/ribbon graph $\Gamma = (H, \sigma, \overline{})$, we usually use its geometric realization, that is, a graph obtained from Γ by gluing half-edges h and \overline{h} together to form a line whose endpoints are $s(h)$ and $s(\overline{h})$. When we describe the cyclic ordering $(h, \sigma(h), \dots, \sigma^{\ell-1}(h))$ of half-edges around a vertex v , we draw them in the plane locally so that the half-edges appear around v in this order counterclockwisely. See the following figure.



We also notice that a geometric realization naturally provides an undirected graph, which we call the *underlying graph* of Γ . From now on, we assume that every ribbon graph is connected, that is, the underlying graph is connected.

Definition 9.2. Let $\Gamma = (H, \sigma, \overline{})$ be a Brauer graph with multiplicity function \mathbf{m} . Let Q_Γ be a finite quiver given as follows:

- The set of vertices is the set E of edges of Γ .
- We draw an arrow $a_h: [h] \rightarrow [\sigma(h)]$ for every $h \in H$.

We define the algebra $B_\Gamma := kQ_\Gamma/I_\Gamma$, where I_Γ is a two-sided ideal in the path algebra kQ_Γ generated by all relations of the following forms:

$$a_{\sigma^{-1}(h)} a_{\overline{h}}: [\sigma^{-1}(h)] \xrightarrow{a_{\sigma^{-1}(h)}} [h] = [\overline{h}] \xrightarrow{a_{\overline{h}}} [\sigma(\overline{h})] \quad \text{and}$$

$$C_h^{\mathbf{m}(s(h))} - C_{\overline{h}}^{\mathbf{m}(s(\overline{h}))}$$

for all $h \in H$. Here, C_h denotes a cycle

$$C_h: [h] \xrightarrow{a_h} [\sigma(h)] \xrightarrow{a_{\sigma(h)}} [\sigma^2(h)] \longrightarrow \dots \longrightarrow [\sigma^{\ell-1}(h)] \xrightarrow{a_{\sigma^{\ell-1}(h)}} [\sigma^\ell(h)] = [h]$$

in Q_Γ and ℓ denotes the cardinality of the σ -orbit incident to $s(h)$. We call B_Γ the *Brauer graph algebra* associated to Γ .




FIGURE 1. Examples of a Brauer tree (in the left) and a Brauer odd-cycle (in the right).

It is well-known (see [Sc] for example) that Brauer graph algebras are finite dimensional symmetric algebras, which are special biserial. We give an example of some Brauer graphs and their associated Brauer graph algebras in Example 9.14.

The following special classes of Brauer graph algebras play a central role in this section. Now, we say that a *cycle graph* of length ℓ is an undirected graph which consists of a single cycle having ℓ edges. An *odd cycle* (respectively, *even cycle*) is a cycle graph of odd (respectively, even) length.

Definition 9.3. A Brauer graph Γ is called *Brauer tree* if its underlying graph is a tree, and called *Brauer odd-cycle* if its underlying graph contains precisely one odd cycle and no even cycles. In this case, B_Γ is called *Brauer tree algebra* and *Brauer odd-cycle algebra* respectively.

Brauer trees and Brauer odd-cycles have canonical embeddings into the plane so that the cyclic ordering of the half-edges incident to each vertex is described in counterclockwise direction. See Figure 1.

The following is a main result in this section, where the statement (b) for Brauer tree algebras was shown in [AMN].

Theorem 9.4. Let Γ be a Brauer graph with n edges and B_Γ the Brauer graph algebra of Γ .

- (a) B_Γ is pairwise g -convex. Moreover, the following conditions are equivalent.
 - (i) B_Γ is g -finite,
 - (ii) B_Γ is g -convex,
 - (iii) Γ is either a Brauer tree or a Brauer odd-cycle.
- (b) Let Φ be a root system of type A_n or C_n and L the root lattice of Φ . If one of the equivalent conditions of (a) hold, then we have an isomorphism $K_0(\text{proj } B_\Gamma) \simeq L$ which restricts to a bijection $[2\text{-psilt}^1 B_\Gamma] \simeq \Phi$:

$$\begin{array}{ccc} K_0(\text{proj } B_\Gamma) & \xrightarrow{\sim} & L \\ \cup & & \cup \\ [2\text{-psilt}^1 B_\Gamma] & \xrightarrow{\sim} & \Phi, \end{array} \quad (9.1)$$

where Φ is of type A_n (respectively, C_n) if Γ is a Brauer tree (respectively, Brauer odd-cycle). Therefore, it induces an isomorphism

$$P(B_\Gamma) \cong P_\Phi$$

of lattice polytopes, and the f -vector and the h -vector are given by

$$\begin{aligned} f_{A_n}(x) &:= \sum_{m=0}^n \frac{(n+m)!}{m!m!(n-m)!} x^{n-m}, & h_{A_n}(x) &:= \sum_{k=0}^n \binom{n}{k}^2 x^k \\ \left(\text{respectively, } f_{C_n}(x) &:= \sum_{m=0}^n \frac{n2^{2m}}{n+m} \binom{n+m}{2m} x^{n-m}, & h_{C_n}(x) &:= \sum_{k=0}^n \binom{2n}{2k} x^k \right). \end{aligned}$$

We note that two g -convex Brauer graph algebras may have the same g -polytope even though they are not derived equivalent, see Example 9.14. We also note that one can describe the g -fan $\Sigma(B_\Gamma)$ in terms of shear coordinate of laminations corresponding to Γ [AY] (see also [FT]).

To prove Theorem 9.4, we need some preparations. It is mentioned in [EJR] (see also [AAC]) that the set $2\text{-silt}B_\Gamma$ only depends on the ribbon graph of Γ and is independent of the multiplicity function \mathbf{m} . For simplicity, we may replace \mathbf{m} to a multiplicity function identically equal to 1, and identify Γ with its ribbon graph naturally.

9.2. Signed half-walks. In this subsection, we recall a classification of 2-term silting complexes over Brauer graph algebras due to [AAC]. Let $\Gamma = (H, \sigma, \overline{})$ be an arbitrary Brauer graph with n edges.


Definition 9.5. (1) A *half-walk* of Γ is a sequence of half-edges $w := (h_1, \dots, h_m)$ such that $s(\overline{h_i}) = s(h_{i+1})$ and $\overline{h_i} \neq h_{i+1}$ for all $i \in \{1, \dots, m-1\}$. Defining $\overline{w} = (\overline{h_m}, \dots, \overline{h_1})$ makes $\overline{}$ an involution on the set of signed half-walks.
(2) A *walk* of Γ is the unordered pair $W = \{w, \overline{w}\}$ of half-walks. For a walk $W = \{w = (h_1, \dots, h_m), \overline{w}\}$, we define $s(w) := s(h_1)$ and $s(\overline{w}) = s(\overline{h_m})$ as the *endpoints* of W .
(3) A *signature* on a walk $W = \{w := (h_1, \dots, h_m), \overline{w}\}$ is an assignment of signs $\epsilon(h) = \epsilon(\overline{h}) \in \{\pm 1\}$ for the half-edges appearing in W such that $\epsilon(h_i) \neq \epsilon(h_{i+1})$ for all $i \in \{1, \dots, m-1\}$. A half-walk w equipped with a signature ϵ on a walk $w \in W$ is called *signed half-walk* and written by w^ϵ or $(h_1, \dots, h_m; \epsilon)$. A *signed walk* is the unordered pair $W^\epsilon = \{w^\epsilon, \overline{w}^\epsilon\}$ of signed half-walks.

Remark that one can not always find a signature for a given walk $W = \{w := (h_1, \dots, h_m), \overline{w}\}$. However, if one finds a signature on W , then there are precisely two signatures $\pm\epsilon$, giving signed half-walks $w^\epsilon = (h_1, \dots, h_m; \epsilon)$ and $w^{-\epsilon} = (h_1, \dots, h_m; -\epsilon)$.

Similar to a walk on undirected graphs, we can describe a given (signed) half-walk graphically as follows:

$$w := (h_1, h_2, \dots, h_m) \\ s(w) \bullet \xrightarrow{h_1} \bullet \xrightarrow{h_2} \bullet \cdots \bullet \xrightarrow{h_m} \bullet s(\overline{w})$$

Here, we need to take in account of loops carefully. For example, $w_1 \neq w_2$ as half-walks in the next example:



We need to attach some extra data to each endpoint of a signed walk, which is uniquely determined by its signature.

Definition 9.6. A *virtual (half-)edge* is an element of the set of the symbols $\{\text{vr}_+(h), \text{vr}_-(h) \mid h \in H\}$. We extend the map s to virtual edges by $s(\text{vr}_\pm(h)) = s(h)$ for all $h \in H$. For a vertex v , we extend the cyclic ordering $(h, \sigma(h), \dots, \sigma^{\ell-1}(h))$ around v to the *cyclic ordering around v accounting the virtual edges* by

$$(\text{vr}_-(h), h, \text{vr}_+(h), \text{vr}_-(\sigma(h)), \sigma(h), \text{vr}_+(\sigma(h)), \dots, \text{vr}_-(\sigma^{\ell-1}(h)), \sigma^{\ell-1}(h), \text{vr}_+(\sigma^{\ell-1}(h))). \quad (9.2)$$

A subsequence of (9.2) is called a *cyclic subordering around v accounting the virtual edges*. To each signed walk $W^\epsilon = \{w^\epsilon = (h_1, \dots, h_m; \epsilon), \overline{w}^\epsilon\}$, we attach virtual edges given by

$$h_0 = \overline{h_0} := \text{vr}_{-\epsilon(h_1)}(h_1) \quad \text{and} \quad h_{r+1} = \overline{h_{r+1}} := \text{vr}_{-\epsilon(h_m)}(\overline{h_m})$$

with signs $\epsilon(h_0) = \epsilon(\overline{h_0}) = -\epsilon(h_1)$ and $\epsilon(h_{m+1}) = \epsilon(\overline{h_{m+1}}) = -\epsilon(h_m)$.

Now we define the admissibility of signed walks by the non-crossing conditions (NC0)-(NC3) below. Fix a pair of (not necessarily distinct) signed walks $W^\epsilon = \{w^\epsilon = (h_1, \dots, h_m; \epsilon), \overline{w}^\epsilon\}$ and $W'^{\epsilon'} = \{w'^{\epsilon'} = (h'_1, \dots, h'_\ell; \epsilon'), \overline{w}'^{\epsilon'}\}$. Notice that there are virtual edges $h_0, h_{\ell+1}, h'_0, h'_{m+1}$, etc. with signs attached to their endpoints.

Definition 9.7. [AAC, Definition 2.7] We say that W^ϵ and $W'^{\epsilon'}$ satisfies (NC0) (it was called the sign condition in [AAC]) if the following condition is satisfied:


(NC0) Whenever W^ϵ and $W'^{\epsilon'}$ have a common endpoint v , the signatures on the half-edges of W^ϵ and of $W'^{\epsilon'}$ incident to v are the same.

Next, we say that a *maximal common subwalk* of W and W' is a walk $Z = \{z, \bar{z}\}$ such that z is a common continuous subsequence of some $x \in W$ and $y \in W'$, and there are no common continuous subsequences of x and y properly containing z as a continuous subsequence. It is said to be *proper* if no endpoints of Z are the common endpoint(s) of W and W' . Notice that there are several maximal common subwalks of W and W' in general.

Definition 9.8. [AAC, Definition 2.8] Let Z be a maximal common subwalk of W and W' given by a half-walk $z = (t_1, \dots, t_r)$ so that $t_k = h_{i+k-1} = h'_{j+k-1}$ for all $k \in \{1, \dots, r\}$. Let $u := s(t_1)$ and $v := s(\bar{t}_r)$ be endpoints of Z . We call the pair of cyclic subordering on $\{\overline{h_{i-1}}, \overline{h'_{j-1}}, t_1\}$ around u and $\{h_{i+r}, h'_{j+r}, \bar{t}_r\}$ around v accounting virtual edges the *neighbourhood cyclic ordering* of Z . Then, we say that W^ϵ and $W'^{\epsilon'}$ satisfies (NC1) and (NC2) at Z if the following condition is satisfied respectively.

(NC1) $\epsilon(t_k) = \epsilon'(t_k)$ for all $k \in \{1, \dots, r\}$.

(NC2) If Z is proper, then the neighbourhood cyclic ordering of Z is either




Notice that the condition (NC1) is automatically satisfied for any signed walk with itself by the definition of the signature (Definition 9.5(3)).

Lastly, for a vertex v and an integer $i \in \{1, \dots, r\}$ such that $v = s(h_i)$, we refer to the set $\{\overline{h_{i-1}}, h_i\}$ as a *neighbourhood* of v in W . Notice that half-edges appearing in the neighbourhood at a vertex can be virtual.

Definition 9.9. [AAC, Definition 2.9] For a vertex v , suppose that $\{a, b\}$ and $\{c, d\}$ are neighbourhoods of v in W and in W' respectively. Then, we say that v is the *intersecting vertex* of W and W' if a, b, c, d are pairwise distinct. We say that W and W' satisfies (NC3) at the intersecting vertex v if the following condition is satisfied.

(NC3) If v is an intersecting vertex with respect to the neighbourhoods $\{a, b\}$ in W and $\{c, d\}$ in W' , and at most one of a, b, c, d is virtual, then the cyclic subordering around v accounting virtual edges and signatures are either



Definition 9.10. We say that two signed walks W^ϵ and $W'^{\epsilon'}$ are *admissible* if they satisfy (NC0), (NC1) and (NC2) at all maximal common subwalks, and (NC3) at all intersecting vertices. An *admissible signed walk* is a signed walk that is admissible with itself.

Now, we denote by $\text{AW}(\Gamma)$ the set of admissible signed walks of Γ , by $\text{CW}(\Gamma)$ the set of maximal collections consisting of admissible signed walks of Γ which are pairwise admissible.

Theorem 9.11. [AAC, Theorem 4.6] *Let Γ be a Brauer graph and B_Γ the Brauer graph algebra of Γ . Then, there are bijections*

$$2\text{-psilt}^1 B_\Gamma \xrightarrow{\sim} \text{AW}(\Gamma) \quad \text{and} \quad 2\text{-silt} B_\Gamma \xrightarrow{\sim} \text{CW}(\Gamma). \quad (9.3)$$

From the definition of Brauer graph algebras, the Grothendieck group $K_0(\text{proj } B_\Gamma)$ is canonically isomorphic to the free \mathbb{Z} -module $\mathbb{Z}E$ over the set E of edges of Γ . For a signed walk $W^\epsilon = \{w^\epsilon = (h_1, \dots, h_m; \epsilon), \bar{w}^\epsilon\}$, one can define

$$[W^\epsilon] := \sum_{i=1}^r \epsilon(h_i)[h_i] \in \mathbb{Z}E. \quad (9.4)$$

In fact, it is independent of a choice of $w \in W$. In addition, let

$$[\text{AW}(\Gamma)] := \{[W^\epsilon] \mid W^\epsilon \in \text{AW}(\Gamma)\} \subset \mathbb{Z}E.$$

Then we have the following.

Proposition 9.12. *We have the following commutative diagram*

$$\begin{array}{ccc} K_0(\text{proj } B_\Gamma) & \xrightarrow{\sim} & \mathbb{Z}E \\ \cup & & \cup \\ [2\text{-psilt}^1 B_\Gamma] & \xrightarrow{\sim} & [\text{AW}(\Gamma)]. \end{array} \quad (9.5)$$

Proof. Let $T \in 2\text{-psilt}^1 B_\Gamma$ and $W^\epsilon \in \text{AW}(\Gamma)$ the signed walk corresponding to T under the bijection $2\text{-psilt}^1 B_\Gamma \simeq \text{AW}(\Gamma)$ in (9.3). According to [AAC, Section 4], one can see that the class $[T]$ is sent to $[W^\epsilon]$ under the canonical isomorphism $K_0(\text{proj } B_\Gamma) \simeq \mathbb{Z}E$. \square

Example 9.13. For an edge $X = \{h, \bar{h}\} \in E$, the element $X \in \mathbb{Z}E$ belongs to $[\text{AW}(\Gamma)]$. In fact, a signed half-walk $W_X^+ = \{(h; +), (\bar{h}; +)\}$ is clearly admissible and satisfies $[W_X^+] = X$ in $\mathbb{Z}E$. On the other hand, let P_X be the indecomposable projective B_Γ -module corresponding to X . Then $P_X \in 2\text{-psilt}^1 B_\Gamma$ is sent to W_X^+ by the bijection in Theorem 9.11.

Example 9.14. We depict the g -polytope and the corresponding root polytope for a class of Brauer graph algebras with small number of edges.


(a) For $n = 2$, there are only one Brauer tree and one Brauer odd-cycle described as follows.



Then the associated Brauer graph algebras are presented by the following quivers with relations:


$$k\left(1 \xrightleftharpoons[a]{b} 2\right) / \langle aba, bab \rangle \quad \text{and} \quad k\left(1 \xrightleftharpoons[a]{c} 2 \begin{array}{c} \curvearrowright \\ b \end{array}\right) / \langle ac, b^2, abca, bca - cab \rangle$$

The g -polytopes of their associated Brauer graph algebras are



which clearly correspond to root polytopes of type A_2 and C_2 respectively.


(b) Let Γ_1 be the following Brauer tree having 3 edges.



Then B_{Γ_1} is the algebra given in Example 5.25. There are 12 elements in $2\text{-psilt}^1 B_{\Gamma_1}$. We describe the corresponding g -vectors and admissible signed walks in the following table.


$[W^\epsilon]$	$(1, 0, 0)$	$(0, 1, 0)$	$(0, 0, 1)$	$(1, -1, 0)$	$(0, 1, -1)$	$(1, -1, 1)$
W^ϵ	$\bullet \xrightarrow{+} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{+} \bullet$	$\bullet \cdots \bullet \xrightarrow{+} \bullet$	$\bullet \xrightarrow{+} \bullet \xrightarrow{-} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{+} \bullet \xrightarrow{-} \bullet$	$\bullet \xrightarrow{+} \bullet \xrightarrow{-} \bullet \xrightarrow{+} \bullet$
	$(-1, 0, 0)$	$(0, -1, 0)$	$(0, 0, -1)$	$(-1, 1, 0)$	$(0, -1, 1)$	$(-1, 1, -1)$
	$\bullet \xrightarrow{-} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{-} \bullet$	$\bullet \cdots \bullet \xrightarrow{-} \bullet$	$\bullet \xrightarrow{-} \bullet \xrightarrow{+} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{-} \bullet \xrightarrow{+} \bullet$	$\bullet \xrightarrow{-} \bullet \xrightarrow{+} \bullet \xrightarrow{-} \bullet$

By Proposition 5.10(b), the g -polytope of B_{Γ_1} is given by the convex hull in \mathbb{R}^3 of all integer vectors appearing in the above table and described as follows.



It is isomorphic to the root polytope of type A_3 up to the linear isomorphism by $\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$.

(c) Let Γ_2 be the following Brauer odd-cycle having 3 edges.




Then $B_{\Gamma_2} \cong k \left[1 \xrightleftharpoons[c]{a} 3 \xrightleftharpoons[d]{b} 2 \right] / \langle ac, db, bdca - cabd \rangle$. Then $B_{\Gamma_2} = \frac{1}{3} \frac{2}{3} \frac{1}{1} \oplus \frac{2}{3} \frac{1}{3} \frac{2}{2} \oplus \frac{3}{1} \frac{1}{3} \frac{1}{2}$, and there are 18 elements in $2\text{-psilt}^1 B_{\Gamma_2}$ whose g -vectors are given by the following table.

$[W^\epsilon]$	$(1, 0, 0)$	$(0, 1, 0)$	$(0, 0, 1)$	$(1, -1, 0)$	$(1, 0, -1)$	$(0, 1, -1)$
W^ϵ	$\bullet \xrightarrow{+} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{+} \bullet$	$\bullet \cdots \bullet \xrightarrow{+} \bullet$	$\bullet \xrightarrow{+} \bullet \xrightarrow{-} \bullet \cdots \bullet$	$\bullet \xrightarrow{+} \bullet \xrightarrow{-} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{+} \bullet \xrightarrow{-} \bullet$
	$(1, 1, -1)$	$(2, 0, -1)$	$(0, 2, -1)$	$(-1, 0, 0)$	$(0, -1, 0)$	$(0, 0, -1)$
	$\bullet \xrightarrow{+} \bullet \xrightarrow{+} \bullet \cdots \bullet$	$\bullet \xrightarrow{+} \bullet \xrightarrow{+} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{+} \bullet \xrightarrow{+} \bullet$	$\bullet \xrightarrow{-} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{-} \bullet$	$\bullet \cdots \bullet \xrightarrow{-} \bullet$
	$(-1, 1, 0)$	$(-1, 0, 1)$	$(0, -1, 1)$	$(-1, -1, 1)$	$(-2, 0, 1)$	$(0, -2, 1)$
	$\bullet \xrightarrow{-} \bullet \xrightarrow{+} \bullet \cdots \bullet$	$\bullet \xrightarrow{-} \bullet \cdots \bullet \xrightarrow{+} \bullet$	$\bullet \cdots \bullet \xrightarrow{-} \bullet \xrightarrow{+} \bullet$	$\bullet \xrightarrow{-} \bullet \xrightarrow{-} \bullet \cdots \bullet$	$\bullet \xrightarrow{-} \bullet \xrightarrow{-} \bullet \cdots \bullet$	$\bullet \cdots \bullet \xrightarrow{-} \bullet \xrightarrow{+} \bullet$

Then $P(B_{\Gamma_2})$ is described in the left diagram of Figure 2, which is isomorphic to the root polytope of type C_3 up to linear isomorphism by $\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & -2 \\ 0 & 0 & 1 \end{pmatrix}$.

- (d) The following Brauer odd-cycle Γ_3 has the same underlying graph as Γ_2 in (c) but they are different as Brauer graphs.



Then, B_{Γ_3} is an algebra A in Example 4.4. There are 18 elements in $2\text{-psilt}^1 B_{\Gamma_3}$ whose g -vectors are given by the following table.

$[W^\epsilon]$	$(1, 0, 0)$	$(0, 1, 0)$	$(0, 0, 1)$	$(1, -1, 0)$	$(1, 0, -1)$	$(0, 1, -1)$
W^ϵ						
	$(1, 1, -1)$	$(2, 0, -1)$	$(0, 2, -1)$	$(-1, 0, 0)$	$(0, -1, 0)$	$(0, 0, -1)$
	$(-1, 1, 0)$	$(-1, 0, 1)$	$(0, -1, 1)$	$(-1, -1, 1)$	$(-2, 0, 1)$	$(0, -2, 1)$

From the tables in (c) and (d), we have

$$[\text{AW}(\Gamma_2)] = [\text{AW}(\Gamma_3)] \subseteq \mathbb{Z}^3.$$

It implies that B_{Γ_2} and B_{Γ_3} determine the same g -polytope, while they do not determine the same g -fan, see Figure 2. Notice that they are not derived equivalent.




FIGURE 2. The g -polytopes for (c) and (d)

9.3. Signed half-walks and root lattices. Let $\Gamma = (H, \sigma, \overline{})$ be a Brauer graph having n edges, which is either a Brauer tree or a Brauer odd-cycle. In this subsection, we associate the root system $\Phi(\Gamma)$ to Γ and show that $\Phi(\Gamma)$ is in bijection with the set $\text{AW}(\Gamma)$ of admissible signed walks of Γ (Proposition 9.16). Using this result, we prove Theorem 9.4.

Definition 9.15. Let V be the set of vertices of Γ and $\mathbb{R}V$ the vector space with basis V . Then

$$\dim \mathbb{R}V = \begin{cases} n + 1 & \text{if } \Gamma \text{ is a Brauer tree,} \\ n & \text{if } \Gamma \text{ is a Brauer odd-cycle.} \end{cases}$$




FIGURE 3. A bipartite orientation on the spanning tree.

We define the inner product on $\mathbb{R}V$ which makes V an orthonormal basis. We define the *root system* associated to Γ by

$$\Phi(\Gamma) := \begin{cases} \Phi_{A_n} = \{u - v \mid (u, v) \in V \times V, u \neq v\} & \text{if } \Gamma \text{ is a Brauer tree,} \\ \Phi_{C_n} = \{\pm u \pm v \mid (u, v) \in V \times V, u \neq v\} \cup \{\pm 2u \mid u \in V\} & \text{if } \Gamma \text{ is a Brauer odd-cycle.} \end{cases}$$

We denote the root lattice of $\Phi(\Gamma)$ by $L(\Gamma)$. The *root polytope* $P_{\Phi(\Gamma)}$ is defined as the convex hull of $\Phi(\Gamma)$ in $\mathbb{R} \otimes_{\mathbb{Z}} L(\Gamma)$.

Proposition 9.16. *Let Γ be a Brauer graph which is either a Brauer tree or a Brauer odd-cycle. Let E be the set of edges of Γ . Then, we have an isomorphism $\partial: \mathbb{Z}E \simeq L(\Gamma)$ which restricts to a bijection $[\text{AW}(\Gamma)] \simeq \Phi(\Gamma)$.*

$$\begin{array}{ccc} \partial: & \mathbb{Z}E & \xrightarrow{\sim} L(\Gamma) \\ & \cup & \cup \\ & [\text{AW}(\Gamma)] & \xrightarrow{\sim} \Phi(\Gamma). \end{array} \quad (9.6)$$

From now on, we prove Proposition 9.16. We need to fix a spanning tree Γ_{sp} of Γ , that is, Γ_{sp} is one of subtrees which include all vertices of Γ . Let H_{sp} be the set of half-edges of Γ_{sp} and H_{sp}^c the complement of H_{sp} . If Γ is a Brauer tree, then $H_{\text{sp}}^c = \emptyset$; Otherwise, H_{sp}^c consists of two half-edges which form an edge lying in the unique odd cycle of Γ .

Definition 9.17. An *orientation* of Γ_{sp} is a complete set \mathfrak{o} of representatives of $H_{\text{sp}}/\overline{(\cdot)}$ in H_{sp} . For such \mathfrak{o} , we say that a vertex v is a *source* (respectively, a *sink*) if there are no half-edges $h \in \mathfrak{o}$ such that $v = s(\overline{h})$ (resp., $v = s(h)$). A *bipartite orientation* is an orientation such that every vertex of Γ_{sp} is either a source or a sink.

Since Γ_{sp} is a tree, it has precisely two bipartite orientation which are related to each other by involution of half-edges. Take one of such an orientation \mathfrak{o} of Γ_{sp} . For a vertex $v \in V$, we set

$$\mathfrak{o}(v) := \begin{cases} 1 & \text{if } v \text{ is a source in } \Gamma_{\text{sp}}, \\ -1 & \text{if } v \text{ is a sink in } \Gamma_{\text{sp}}. \end{cases}$$

By definition, we have $\mathfrak{o}(s(h)) = -\mathfrak{o}(s(\overline{h}))$ for any $h \in H_{\text{sp}}$, while we have $\mathfrak{o}(s(h)) = \mathfrak{o}(s(\overline{h}))$ for the half-edges $h \in H_{\text{sp}}^c$ since Γ is a Brauer odd-cycle (if h exists) and h belongs to the odd cycle, see Figure 3. Consider the surjective homomorphism

$$\pi: \mathbb{Z}H \rightarrow \mathbb{Z}E \quad \text{given by} \quad h \mapsto \mathfrak{o}(s(h))[h] \quad \text{for all } h \in H,$$

and a homomorphism

$$\delta: \mathbb{Z}H \rightarrow \mathbb{Z}V \quad \text{given by} \quad h \mapsto \begin{cases} s(h) - s(\overline{h}) & \text{for all } h \in H_{\text{sp}}, \\ s(h) + s(\overline{h}) & \text{for all } h \in H_{\text{sp}}^c. \end{cases}$$

Proposition 9.18. *We have the following commutative diagram.*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \ker \pi & \longrightarrow & \mathbb{Z}H & \xrightarrow{\pi} & \mathbb{Z}E \longrightarrow 0 \\
 & & & & \downarrow \delta & \nearrow \partial & \\
 & & & & \mathbb{Z}V & &
 \end{array} \quad (\text{exact})$$

Moreover, for any $h \in H$, we have

$$\partial([h]) = \begin{cases} \mathfrak{o}(s(h))(s(h) - s(\bar{h})) & \text{if } h \in H_{\text{sp}}, \\ \mathfrak{o}(s(h))(s(h) + s(\bar{h})) & \text{if } h \in H_{\text{sp}}^c. \end{cases} \quad (9.7)$$

Proof. For the former assertion, it suffices to show $\delta(\ker \pi) = 0$. By definition, the kernel $\ker \pi$ is generated by elements of the forms $h + \bar{h}$ with $h \in H_{\text{sp}}$ and $h - \bar{h}$ with $h \in H_{\text{sp}}^c$. Then, we have

$$\begin{aligned} \delta(h + \bar{h}) &= (s(h) - s(\bar{h})) + (s(\bar{h}) - s(h)) = 0 \quad \text{for all } h \in H_{\text{sp}} \text{ and} \\ \delta(h - \bar{h}) &= (s(h) + s(\bar{h})) - (s(\bar{h}) + s(h)) = 0 \quad \text{for all } h \in H_{\text{sp}}^c. \end{aligned}$$

Thus, we get the former assertion. The latter one is clear. \square

Proposition 9.19. *The map ∂ restricts to an isomorphism $\partial: \mathbb{Z}E \simeq L(\Gamma)$.*

Proof. As we mentioned in Example 9.13, a basis E of $\mathbb{Z}E$ is included in $[\text{AW}(\Gamma)]$. We first assume that Γ is a Brauer tree and hence $\Phi(\Gamma) = \Phi_{A_n}$. By (9.7), E is bijectively sent to

$$\partial(E) = \{s(h) - s(\bar{h}) \mid h \in \mathfrak{o}\} \subseteq \Phi_{A_n},$$

which is clearly a basis of $L(\Gamma)$. Next, we assume that Γ is a Brauer odd-cycle and hence $\Phi(\Gamma) = \Phi_{C_n}$. We denote by $X = \{h_0, \bar{h}_0\}$ the unique edge which does not lie in Γ_{sp} . Then, E is bijectively sent to

$$\partial(E) \stackrel{(9.7)}{=} \{s(h) - s(\bar{h}) \mid h \in \mathfrak{o}\} \cup \{\mathfrak{o}(s(h_0))(s(h_0) + s(\bar{h}_0))\} \subseteq \Phi_{C_n}.$$

which is easily shown to be a basis of $L(\Gamma)$. \square

Now, we study signed walks of Γ . The following is basic.


Lemma 9.20. *Let Γ be a Brauer tree or a Brauer odd-cycle.*

- (a) *Let $w^\epsilon = (h_1, \dots, h_m; \epsilon)$ be a signed half-walk. For any edge X lying in the odd cycle in Γ , there is at most one $1 \leq i \leq m$ such that $h_i \in X$. For a signed half-walk w^ϵ of Γ , every half-edge lying in the odd cycle in Γ appears at most once in w^ϵ .*
- (b) *Every signed walk W^ϵ of Γ satisfies (NC0) and (NC1).*

Proof. (a) We only have to consider a Brauer odd-cycle Γ . We denote by Z the unique odd cycle in Γ , by ℓ the length of Z . Take an integers $1 \leq j < k \leq m$ such that $h_j, h_k \in X$ and there are no $j < i < k$ satisfying $h_i \in X$. Since Γ has no cycle except for Z , we have $h_j = h_k$ and $k - j = \ell$. Since ℓ is odd, we have $\epsilon(h_j) = \epsilon(h_k) = (-1)^{k-j}\epsilon(h_j) = (-1)^\ell\epsilon(h_j) = -\epsilon(h_j)$, a contradiction.

(b) The condition (NC1) is automatic for any signed walk.

Let $W^\epsilon = \{w^\epsilon = (h_1, \dots, h_m; \epsilon), \bar{w}^\epsilon\}$ be a signed walk with endpoints $u := s(h_1)$ and $v := s(\bar{h}_m)$. If $u \neq v$, the condition (NC0) is automatic. Assume that $u = v$. In this case, Γ is a Brauer odd-cycle. We denote by Z the unique odd cycle in Γ , by ℓ the length of Z . Since Γ has no cycles except for Z , using (a), we can write $w = pz\bar{p}$, where $p = (h_1, \dots, h_r)$ with $r \geq 0$ and $z = (h_{r+1}, \dots, h_{r+\ell})$ forms the cycle Z .



Since ℓ is odd, so is $m = 2r + \ell$. Thus, $\epsilon(h_m) = (-1)^{m-1}\epsilon(h_1) = \epsilon(h_1)$ as desired. \square

By Lemma 9.20(b), for a signed walk $W = \{w^\epsilon = (h_1, \dots, h_m; \epsilon), \overline{w}^\epsilon\}$, one can extend the signature ϵ to the endpoints $s(w) = s(h_1)$ and $s(\overline{w}) = s(\overline{h_m})$ of W by $\epsilon(s(w)) := \epsilon(h_1)$ and $\epsilon(s(\overline{w})) := \epsilon(h_m)$. Moreover, one can assign the sign $\mathfrak{o}\epsilon(s(w)) := \mathfrak{o}(s(w))\epsilon(s(w)) \in \{\pm 1\}$ on $s(w)$, and similarly $\mathfrak{o}\epsilon(s(\overline{w}))$ on $s(\overline{w})$.

The following result means that the element $\partial([W^\epsilon])$ is completely determined by the endpoints of W and the signature on them.

Lemma 9.21. *For a signed walk $W^\epsilon = \{w^\epsilon, \overline{w}^\epsilon\}$ of Γ , the following holds.*

$$\partial([W^\epsilon]) = \begin{cases} \mathfrak{o}\epsilon(s(w))(s(w) - s(\overline{w})) & \text{if } W \text{ is a walk of } \Gamma_{\text{sp}}, \\ \mathfrak{o}\epsilon(s(w))(s(w) + s(\overline{w})) & \text{else.} \end{cases} \quad (9.8)$$

Proof. Let $W^\epsilon = \{w^\epsilon = (h_1, \dots, h_m; \epsilon), \overline{w}^\epsilon\}$ be a signed walk. By (9.4) and (9.7), we have

$$\begin{aligned} [W^\epsilon] &= \sum_{i=1}^m \epsilon(h_i)[h_i] \in \mathbb{Z}E \quad \text{and} \\ \partial([h_i]) &= \begin{cases} \mathfrak{o}(s(h_i))(s(h_i) - s(h_{i+1})) & \text{if } h_i \in H_{\text{sp}}, \\ \mathfrak{o}(s(h_i))(s(h_i) + s(h_{i+1})) & \text{if } h_i \in H_{\text{sp}}^c, \end{cases} \end{aligned}$$

where $1 \leq i \leq m$ and $s(h_{m+1}) = s(\overline{h_m})$. If W is a walk of Γ_{sp} (that is, $h_i \in H_{\text{sp}}$ for all $1 \leq i \leq m$), we have

$$\begin{aligned} \partial([W^\epsilon]) &= \sum_{i=1}^m \epsilon(h_i)\partial([h_i]) = \mathfrak{o}\epsilon(s(h_1)) \sum_{i=1}^m (s(h_i) - s(h_{i+1})) \\ &= \mathfrak{o}\epsilon(s(h_1))(s(h_1) - s(\overline{h_m})) = \mathfrak{o}\epsilon(s(w))(s(w) - s(\overline{w})). \end{aligned}$$

Otherwise, by Lemma 9.20(a), there exists a unique integer $1 \leq j \leq m$ such that $h_j \in H_{\text{sp}}^c$. Then we have

$$\begin{aligned} \partial([W^\epsilon]) &= \sum_{i=1}^m \epsilon(h_i)\partial([h_i]) \\ &= \mathfrak{o}\epsilon(s(h_1)) \left(\sum_{i=1}^{j-1} (s(h_i) - s(h_{i+1})) + (s(h_j) + s(h_{j+1})) - \sum_{i=j+1}^m (s(h_i) - s(h_{i+1})) \right) \\ &= \mathfrak{o}\epsilon(s(h_1))(s(h_1) + s(\overline{h_m})) = \mathfrak{o}\epsilon(s(w))(s(w) + s(\overline{w})). \end{aligned}$$

Thus, we obtain the desired equalities (9.8). \square

Lemma 9.22. *Let T be a subtree of Γ . Then, the following hold.*

- (a) *Every signed walk of T is admissible on Γ .*
- (b) *For two vertices u, v of T with $u \neq v$, there exists a unique walk W having u, v as endpoints and it gives rise to admissible signed walks $W^{\pm\epsilon}$. Conversely, every admissible signed walks of T can be obtained in this way.*

Proof. (a) Let W^ϵ be a signed walk of T . By Lemma 9.20(b), W^ϵ satisfies the conditions (NC0) and (NC1). Since it has no proper maximal common subwalks and no intersecting vertices with itself, (NC2) and (NC3) are automatic. We conclude that W^ϵ is admissible on T , and also on Γ .

(b) Let u, v be distinct vertices of T . Since T is a tree, there exists a unique half-walk $w = (h_1, \dots, h_m)$ such that $s(h_1) = u$ and $s(\overline{h_m}) = v$ and it gives rise to a walk $W = \{w, \overline{w}\}$ having u, v as endpoints. In this case, h_1, \dots, h_m are pairwise distinct because T is a tree. Then one can find a signature $\pm\epsilon$ on W , and both of $W^{\pm\epsilon}$ are admissible by (a). Conversely, it is easy to see that every admissible signed walk of T can be obtained in this way. \square

Now, we are ready to prove Proposition 9.16.

Proof of Proposition 9.16. In the above notations, we show that the isomorphism

$$\partial: \mathbb{Z}E \xrightarrow{\sim} L(\Gamma)$$

in Proposition 9.19 restricts to a bijection

$$\partial|_{[\text{AW}(\Gamma)]}: [\text{AW}(\Gamma)] \xrightarrow{\sim} \Phi(\Gamma). \quad (9.9)$$

We first consider the case when Γ is a Brauer tree and hence $\Phi(\Gamma) = \Phi_A$. From (9.8), we have $\partial([\text{AW}(\Gamma)]) \subseteq \Phi_A$ since every signed walk of Γ is admissible and has distinct endpoints by Lemma 9.22(b). On the other hand, by Lemma 9.22(b), for two distinct vertices u, v of Γ , we have two admissible signed walks $W^{\pm\epsilon}$ having u, v as the endpoints and

$$\partial(\{W^\epsilon, W^{-\epsilon}\}) = \{\pm(u - v)\}.$$

Therefore, we get the assertion in this case.

Next, we assume that Γ is a Brauer odd-cycle and hence $\Phi(\Gamma) = \Phi_C$. We denote by Z the unique odd cycle in Γ , by ℓ the length of Z . For each vertex $v \in V$, we define a subtree T_v of Γ as follows: Consider a graph obtained from Γ by deleting all (half-)edges lying in Z and then T_v is defined to be its connected component containing the vertex v . We write \tilde{v} for the the unique common vertex of T_v and Z .

By (9.8), we have $\partial([W^\epsilon]) \in \Phi_C$ for any signed walk W^ϵ of Γ . In fact, for the endpoints u, v of W^ϵ , we have $\partial([W^\epsilon]) = \pm u \pm v$ if $u \neq v$, $\partial([W^\epsilon]) = \pm 2u$ by (NC0) otherwise. Thus, $\partial([\text{AW}(\Gamma)]) \subseteq \Phi_C$. In order to show the bijectivity, it is enough to show $\Phi_C \subseteq \partial([\text{AW}(\Gamma)])$.


(a) Firstly, we consider the elements of the form $\pm(u - v) \in \Phi_C$ with $u \neq v$. Let u, v be distinct vertices of Γ . Applying Lemma 9.22(b) to the spanning tree Γ_{sp} , there are two admissible signed walks $W^{\pm\epsilon}$ having u, v as endpoints. By (9.8), we have

$$\partial(\{W^\epsilon, W^{-\epsilon}\}) = \{\pm(u - v)\}.$$

(b) Secondly, we show that $\pm(u + v)$ belong to $\partial([\text{AW}(\Gamma)])$ for vertices $u \neq v$ with $T_u \neq T_v$. Recall from Lemma 9.22 that we have a unique half-walk w of Γ_{sp} such that $u = s(w)$ and $v = s(\overline{w})$. Since Γ contains no cycles except for Z , w can be written as $w = pzq$, where

- $p = (p_1, \dots, p_m)$ ($m \geq 0$) is a unique half-walk on T_u such that $u = s(p_1)$ and $\tilde{u} = s(\overline{p_m})$,
- $z := (z_1, \dots, z_j)$ ($1 \leq j < \ell$) is a half-walk consisting of half-edges appearing in Z such that $\tilde{u} = s(z_1)$ and $\tilde{v} = s(\overline{z_j})$ with $z_i \in H_{\text{sp}}$ for all $1 \leq i \leq j$, and
- $q = (q_1, \dots, q_r)$ ($r \geq 0$) is a unique half-walk on T_v such that $\tilde{v} = s(q_1)$ and $v = s(\overline{q_r})$.

For a half-walk z , there is a half-walk $z' = (z_{j+1}, \dots, z_\ell)$ such that $zz' = (z_1, \dots, z_j, z_{j+1}, \dots, z_\ell)$ forms the cycle Z . Since $s(\overline{z_\ell}) = \tilde{u}$ and $s(z_{j+1}) = \tilde{v}$, we obtain a half-walk $w' := pz'q$ and a walk $W' := \{w', \overline{w'}\}$ of Γ .



From our construction, W' is not a walk of Γ_{sp} but a walk of a certain subtree of Γ . By Lemma 9.22(b), it gives rise to two admissible signed walks $W'^{\pm\epsilon'}$. By (9.8), we have

$$\partial(\{W'^{\epsilon'}, W'^{-\epsilon'}\}) = \{\pm(u + v)\}.$$

(c) Thirdly, we consider vertices $u \neq v$ with $T_u = T_v$, i.e., $\tilde{u} = \tilde{v}$. We will construct $X, Y \in \text{AW}(\Gamma)$ such that


$$\partial(\{X, Y\}) = \{\pm(u + v)\}. \quad (9.10)$$

We may assume $u \neq \tilde{u}$ by replacing u and v if necessary. Then, one can find a half-walk $w = (h_1, \dots, h_m)$ of the form $w := pzq$, where

- $p = (p_1, \dots, p_m)$ ($m \geq 1$) is a unique half-walk on T_u such that $u = s(p_1)$ and $\tilde{u} = s(\overline{p_m})$,
- $z = (z_1, \dots, z_\ell)$ is a half-walk which forms the cycle Z so that $\tilde{u} = s(z_1) = s(\overline{z_\ell})$, and
- $q = (q_1, \dots, q_r)$ ($r \geq 0$) is a unique half-walk on T_u such that $\tilde{u} = s(q_1)$ and $v = s(\overline{q_r})$.


On the other hand, let w' be a half-walk of Γ defined by $w' := p\bar{z}q$. Now, we fix a signature ϵ on a walk $w \in W$. Then, it gives a signature on $w' \in W'$. By Lemma 9.20(b), all $W^{\pm\epsilon}, W'^{\pm\epsilon}$ satisfy (NC0) and (NC1). For the admissibility, we need to check the conditions (NC2) and (NC3). Depending on the structure of Γ , we have three cases (c-i)-(c-iii) as follows.

(c-i) We first consider the case when $r \geq 1$ and $\overline{p_m} = q_1$. In this case, (NC3) is automatic. For a signed walk W^ϵ , take an integer $k > 0$ such that $t_i := p_{m-k+i} = \overline{q_i}$ for all $i \in \{1, \dots, k\}$ but $\overline{p_{m-k}} \neq q_{k+1}$. Since $u \neq v$, at most one of $\overline{p_{m-k}}$ and q_{k+1} is the virtual edge attached to W^ϵ . Then, (t_1, \dots, t_k) gives a unique proper maximal common subwalk of W with itself. By definition, W^ϵ satisfies (NC2) if and only if the cyclic subordering around \tilde{u} and around $s := s(t_1)$ accounting the virtual edges is either



By the cyclic ordering around \tilde{u} , precisely one of W^ϵ and W'^ϵ satisfies (NC2) by taking in account of virtual edges attached to them. Similarly, precisely one of $W^{-\epsilon}$ and $W'^{-\epsilon}$ satisfies (NC2). As a consequence, we obtain the desired admissible signed walks X and Y satisfying (9.10).

(c-ii) Next, we assume that $r \geq 1$ and $\overline{p_m} \neq q_1$. In this case, the condition (NC2) is automatic. For a signed walk W^ϵ , \tilde{u} is the intersecting vertex of W with itself, whose neighbourhoods are $\{a, b\} = \{\overline{p_m}, z_1\}$ and $\{c, d\} = \{\overline{z_\ell}, q_1\}$. Recall that W^ϵ satisfies (NC3) if and only if the cyclic subordering around \tilde{u} accounting virtual edges and the signatures are either



From the cyclic ordering around \tilde{u} , precisely one of W^ϵ and W'^ϵ satisfies (NC3). Similarly, precisely one of $W^{-\epsilon}$ and $W'^{-\epsilon}$ satisfies (NC3). Then, we obtain the desired X and Y satisfying (9.10).

(c-iii) Lastly, we assume that $r = 0$, i.e., $v = \tilde{u}$. In this case, by letting $q_1 := z_{\ell+1}$ be the virtual edge attached to W^ϵ , a similar argument as in (c-ii) gives the desired X and Y satisfying (9.10).

(d) Finally, we show that $\pm 2u \in \partial([AW(\Gamma)])$ for any vertex $u \in V$. By the same way as a proof of Lemma 9.20(b), one can find a half-walk $w = (h_1, \dots, h_m)$ of the form $w = p\bar{z}\bar{p}$, where

- $p = (h_1, \dots, h_r)$ ($r \geq 0$) is a unique half-walk on T_u such that $u = s(h_1)$ and $\tilde{u} = s(\overline{h_r})$ and
- $z = (h_{r+1}, \dots, h_{r+\ell})$ is a half-walk which forms the cycle Z so that $\tilde{u} = s(h_{r+1}) = s(\overline{h_{r+\ell}})$.

Then, it gives rise to a walk W equipped with signatures $\pm\epsilon$. By Lemma 9.20(b), it satisfies the conditions (NC0) and (NC1). Since the endpoints of W are the same, the condition (NC2) is automatic. Now, we consider (NC3). This is automatic if $u \neq \tilde{u}$. Assume that $u = \tilde{u}$ and let $\{a, b\} = \{\overline{h_r}, h_{r+1}\}$ and $\{c, d\} = \{\overline{h_{r+\ell}}, h_{r+\ell+1}\}$ be neighbourhoods of \tilde{u} in Γ . Since two of a, b, c, d are virtual, the condition (NC3) is satisfied. Therefore, both of $W^{\pm\epsilon}$ are admissible. By (9.8), we have

$$\partial(\{W^\epsilon, W^{-\epsilon}\}) = \{\pm 2u\}.$$

By (a)-(d), we conclude that the map (9.9) is bijective. \square

We end this subsection with a proof of Theorem 9.4.

Proof of Theorem 9.4. Let Γ be a Brauer graph having n edges and $B := B_\Gamma$ the Brauer graph algebra of Γ .

(a) Thanks to Proposition 4.5(c), we can assume that a base field k is algebraically closed. In order to show that the algebra B is pairwise g -convex, we first consider a left mutation of B with respect to the indecomposable projective B -module P so that the exchange triangle is

$$P \rightarrow U \rightarrow P' \rightarrow P[1]. \quad (9.11)$$

The triangle (9.11) is explicitly described in [Ai, Section 6], in particular, the number of indecomposable direct summands of U is at most two.

Next, let $T \in 2\text{-silt} B$. Then T is a tilting complex since B is a symmetric algebra [AiI, Example 2.8]. We have a triangle equivalence

$$F: D^b(\text{mod } B) \xrightarrow{\sim} D^b(\text{mod } \text{End}_{D^b(\text{mod } B)}(T))$$

mapping T to $\text{End}_{D^b(\text{mod } B)}(T)$. By [AZ, Corollary 1.3], we have $\text{End}_{D^b(\text{mod } B)}(T) \cong B_{\Gamma'}$ for some Brauer graph Γ' having n edges. Since F sends an exchange triangle $X \rightarrow U' \rightarrow Y \rightarrow X[1]$ of T to an exchange triangle of $B_{\Gamma'}$, by applying the argument above to $B_{\Gamma'}$, the number of indecomposable direct summands of U' is at most two.

Consequently, B is pairwise g -convex.

We show the latter assertion. (i) \Leftrightarrow (iii) is [AAC, Theorem 6.7]. (i) \Leftrightarrow (ii) follows from Theorem 5.10(b).

(b) Assume that Γ is a Brauer tree or a Brauer odd-cycle. Combining (9.5) and (9.6), we obtain the following commutative diagram.

$$\begin{array}{ccccc} K_0(\text{proj } B_\Gamma) & \xrightarrow{\sim} & \mathbb{Z}E & \xrightarrow{\sim} & L(\Gamma) \\ \cup & & \cup & & \cup \\ [2\text{-psilt}^1 B_\Gamma] & \xrightarrow{\sim} & [\text{AW}(\Gamma)] & \xrightarrow{\sim} & \Phi(\Gamma). \end{array}$$

Thus, we obtain (9.1), which clearly gives rise to the desired isomorphism $P(B_\Gamma) \cong P_{\Phi(\Gamma)}$ of lattice polytopes. The last assertion follows from [ABHPS, Theorem 1]. \square

ACKNOWLEDGMENTS

T.A is supported by JSPS Grants-in-Aid for Scientific Research JP19J11408. A.H is supported by JSPS Grant-in-Aid for Scientists Research (C) 20K03513. O.I is supported by JSPS Grant-in-Aid for Scientific Research (B) 16H03923, (C) 18K3209 and (S) 15H05738. R.K is supported by JSPS Grant-in-Aid for Young Scientists (B) 17K14169. Y.M is supported by Grant-in-Aid for Scientific Research (C) 20K03539. Y.M would like to thank Haruhisa Enomoto and Hiroyuki Minamoto for helpful discussions

REFERENCES

- [AHKM1] T. Aoki, A. Higashitani, O. Iyama, R. Kase, Y. Mizuno, *Fans and polytopes in tilting theory II: Classification of g -fans of rank 2*, in preparation.
- [AHKM2] T. Aoki, A. Higashitani, O. Iyama, R. Kase, Y. Mizuno, *Fans and polytopes in tilting theory III: Classification of convex g -fans of rank 3*, in preparation.
- [Ada] T. Adachi, *Characterizing τ -tilting finite algebras with radical square zero*, Proc. Amer. Math. Soc. 144 (2016) 4673–4685.
- [AAC] T. Adachi, T. Aihara, A. Chan, *Classification of two-term tilting complexes over Brauer graph algebras*, Math. Z. 290 (2018), no. 1–2, 1–36.
- [AIR] T. Adachi, O. Iyama, I. Reiten, τ -tilting theory, Compos. Math. 150 (2014), no. 3, 415–452.
- [Adi] K. Adiprasito, *Combinatorial Lefschetz theorems beyond positivity*, arXiv:1812.10454.
- [Ai] T. Aihara, *Derived equivalences between symmetric special biserial algebras*, J. Pure Appl. Algebra 219 (2015), no. 5, 1800–1825.
- [AiI] T. Aihara, O. Iyama, *Silting mutation in triangulated categories*, J. Lond. Math. Soc. (2) 85 (2012), no. 3, 633–668.

- [AM] T. Aihara, Y. Mizuno, *Classifying tilting complexes over preprojective algebras of Dynkin type*, Algebra Number Theory 11 (2017), No. 6, 1287–1315.
- [AIRT] C. Amiot, O. Iyama, I. Reiten, G. Todorov, *Preprojective algebras and c -sortable words*, Proc. Lond. Math. Soc. (3) 104 (2012), no. 3, 513–539.
- [An] L. Angeleri Hügel, *Silting objects*, Bull. Lond. Math. Soc. 51 (2019), no. 4, 658–690.
- [AMV] L. Angeleri Hügel, F. Marks, J. Vitoria, *Silting modules*, Int. Math. Res. Not. IMRN 2016, no. 4, 1251–1284.
- [AZ] M. Antipov, A. Zvonareva, *Brauer graph algebras are closed under derived equivalence*, Math. Z. 301 (2022), no. 2, 1963–1981.
- [Ao] T. Aoki, *Classifying torsion classes for algebras with radical square zero via sign decomposition*, J. Algebra 610 (2022), 167–198.
- [AY] T. Aoki, T. Yurikusa, *Complete special biserial algebras are g -tame*, arXiv:2003.09797.
- [ABHPS] F. Ardila, M. Beck, S. Hocsten, J. Pfeifle, K. Seashore, *Root polytopes and growth series of root lattices*, SIAM J. Discrete Math. 25 (2011), no. 1, 360–378.
- [As1] S. Asai, *Semibricks*, Int. Math. Res. Not. IMRN 2020, no. 16, 4993–5054.
- [As2] S. Asai, *The wall-chamber structures of the real Grothendieck groups*, Adv. Math. 381 (2021), 107615.
- [AsI] S. Asai, O. Iyama, *Semistable torsion classes and canonical decompositions in Grothendieck groups*, arXiv:2112.14908.
- [AMN] H. Asashiba, Y. Mizuno, K. Nakashima, *Simplicial complexes and tilting theory for Brauer tree algebras*, J. Algebra 551 (2020), 119–153.
- [ASS] I. Assem, D. Simson, A. Skowroński, *Elements of the Representation Theory of Associative Algebras. Vol. 1*, London Mathematical Society Student Texts 65, Cambridge university press (2006).
- [ARS] M. Auslander, I. Reiten, S. O. Smalø, *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1997.
- [AS] M. Auslander, S. O. Smalø, *Preprojective modules over Artin algebras*, J. Algebra 66 (1980), no. 1, 61–122.
- [BCDMTY] V. Bazier-Matte, N. Chapelier-Laget, G. Douville, K. Mousavand, H. Thomas, and E. Yildirim, *ABHY associahedra and newton polytopes of F -polynomials for finite type cluster algebras*, arXiv:1808.09986.
- [BKT] P. Baumann, J. Kamnitzer, P. Tingley, *Affine Mirkovic-Vilonen polytopes*, Publ. Math. Inst. Hautes Etudes Sci. 120 (2014), 113–205.
- [BR] M. Beck, S. Robins, *Computing the continuous discretely. Integer-point enumeration in polyhedra*, Second edition. With illustrations by David Austin. Undergraduate Texts in Mathematics. Springer, New York, 2015.
- [BFMN] L. Bossinger, B. Frias-Medina, T. Magee, A. Najera Chavez, *Toric degenerations of cluster varieties and cluster duality*, Compos. Math. 156 (2020), no. 10, 2149–2206.
- [B] T. Bridgeland, *Scattering diagrams, Hall algebras and stability conditions*, Algebr. Geom. 4 (2017), no. 5, 523–561.
- [BY] T. Brüstle, D. Yang, *Ordered exchange graphs*, Advances in representation theory of algebras, 135–193, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2013.
- [BST] T. Brüstle, D. Smith, H. Treffinger, *Wall and chamber structure for finite-dimensional algebras*, Adv. Math. 354 (2019), 106746.
- [BH] W. Bruns, J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, Cambridge, 1993.
- [BIRS] A. B. Buan, O. Iyama, I. Reiten, J. Scott, *Cluster structures for 2-Calabi-Yau categories and unipotent groups*, Compos. Math. 145 (2009), 1035–1079.
- [BP] V. Buchstaber, T. Panov, *Toric topology*, Mathematical Surveys and Monographs, 204. American Mathematical Society, Providence, RI, 2015.
- [C] J. W. S. Cassels, *An introduction to the geometry of numbers*, Springer-Verlag, Berlin, 1997.
- [CKLP] G. Cerulli Irelli, Giovanni, B. Keller, D. Labardini-Fragoso, P. Plamondon, *Linear independence of cluster monomials for skew-symmetric cluster algebras*, Compos. Math. 149 (2013), no. 10, 1753–1764.
- [CFZ] F. Chapoton, S. Fomin, A. Zelevinsky, *Polytopal realizations of generalized associahedra*, Canad. Math. Bull. 45 (2002), no. 4, 537–566.
- [DIJ] L. Demonet, O. Iyama, G. Jasso, *τ -tilting finite algebras, bricks, and g -vectors*, Int. Math. Res. Not. IMRN 2019, no. 3, 852–892.
- [DIRRT] L. Demonet, O. Iyama, N. Reading, I. Reiten, H. Thomas, *Lattice theory of torsion classes*, to appear in Trans. Amer. Math. Soc., arXiv:1711.01785.
- [DDPW] B. Deng, J. Du, B. Parshall, J. Wang, *Finite dimensional algebras and quantum groups*, Mathematical Surveys and Monographs, 150. American Mathematical Society
- [DF] H. Derksen, J. Fei, *General presentations of algebras*, Adv. Math. 278 (2015), 210–237.
- [DWZ] H. Derksen, J. Weyman, A. Zelevinsky, *Quivers with potentials and their representations II: applications to cluster algebras*, J. Amer. Math. Soc. 23 (2010), no. 3, 749–790.
- [D] V. Drinfeld, *DG quotients of DG categories*, J. Algebra 272 (2004), no. 2, 643–691.
- [EJR] F. Eisele, G. Janssens, T. Raedschelders, *A reduction theorem for τ -rigid modules*, Math. Z. 290 (2018), no. 3–4, 1377–1413.
- [E] G. Ewald, *On the classification of toric Fano varieties*, Discrete Comput. Geom. 3 (1988), no. 1, 49–54.

- [Fe1] J. Fei, *Combinatorics of F -polynomials*, to appear in Int. Math. Res. Not., arXiv:1909.10151.
- [Fe2] J. Fei, *Tropical F -polynomials and General Presentations*, arXiv:1911.10513.
- [FT] S. Fomin, D. Thurston, *Cluster algebras and triangulated surfaces Part II: Lambda lengths*, Mem. Amer. Math. Soc. 255 (2018), no. 1223.
- [FZ1] S. Fomin, A. Zelevinsky, *Y -systems and generalized associahedra*, Ann. of Math. (2) 158 (2003), no. 3, 977–1018.
- [FZ2] S. Fomin, A. Zelevinsky, *Cluster algebras. II. Finite type classification*, Invent. Math. 154 (2003), no. 1, 63–121.
- [FZ3] S. Fomin, A. Zelevinsky, *Cluster algebras. IV. Coefficients*, Compos. Math. 143 (2007), no. 1, 112–164.
- [FG] C. Fu, S. Geng, *Tilting modules and support τ -tilting modules over preprojective algebras associated with symmetrizable Cartan matrices*, Algebr. Represent. Theory 22 (2019), no. 5, 1239–1260.
- [Fu] W. Fulton, *Introduction to Toric Varieties*, Ann of Math. Studies **131**, Princeton Univ. Press, 1993.
- [GLS] C. Geiss, B. Leclerc, J. Schröer, *Quivers with relations for symmetrizable Cartan matrices I: Foundations*, Invent. Math. 209 (2017), no. 1, 61–158.
- [GI] J. Grant, O. Iyama, *Higher preprojective algebras, Koszul algebras, and superpotentials*, Compos. Math. 156 (2020), no. 12, 2588–2627.
- [GHKK] M. Gross, P. Hacking, S. Sean M. Kontsevich, *Canonical bases for cluster algebras*, J. Amer. Math. Soc. 31 (2018), no. 2, 497–608.
- [H1] L. Hille, *On the volume of a tilting module*, Abh. Math. Sem. Univ. Hamburg 76 (2006), 261–277.
- [H2] L. Hille, *Tilting Modules over the Path Algebra of Type A, Polytopes, and Catalan Numbers*, Lie algebras and related topics, 91–101, Contemp. Math., 652, Amer. Math. Soc., Providence, RI, 2015.
- [HLT] C. Hohlweg, C. Lange, H. Thomas, *Permutahedra and generalized associahedra*, Adv. Math. 226 (2011), no. 1, 608–640.
- [HPS] C. Hohlweg, V. Pilaud, S. Stella, *Polytopal realizations of finite type g -vector fans*, Adv. Math. 328 (2018), 713–749.
- [INT] N. Ilten, A. Najera Chavez, H. Treffinger, *Deformation Theory for Finite Cluster Complexes*, arXiv: 2111.02566.
- [IT] C. Ingalls, H. Thomas, *Noncrossing partitions and representations of quivers*, Compos. Math. 145 (2009), no. 6, 1533–1562.
- [IR] O. Iyama, I. Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, Amer. J. Math. 130 (2008), no. 4, 1087–1149.
- [J] G. Jasso, *Reduction of τ -tilting modules and torsion pairs*, Int. Math. Res. Not. IMRN 2015, no. 16, 7190–7237.
- [IK] O. Iyama, Y. Kimura, *Classifying subcategories of modules over Noetherian algebras*, arXiv:2106.00469.
- [IRRT] O. Iyama, N. Reading, I. Reiten, H. Thomas, *Lattice structure of Weyl groups via representation theory of preprojective algebras*, Compos. Math. 154 (2018), no. 6, 1269–1305.
- [IY] O. Iyama, D. Yang, *Silting reduction and Calabi–Yau reduction of triangulated categories*, Trans. Amer. Math. Soc. 370 (2018), no. 11, 7861–7898.
- [Ka] R. Kase, *From support τ -tilting posets to algebras*, arXiv:1709.05049.
- [Ke1] B. Keller, *Deriving DG categories*, Ann. Sci. Ecole Norm. Sup. (4) 27 (1994), no. 1, 63–102.
- [Ke2] B. Keller, *On the cyclic homology of exact categories*, J. Pure Appl. Algebra 136 (1999), no. 1, 1–56.
- [KV] B. Keller, D. Vossieck, *Aisles in derived categories*, Deuxieme Contact Franco-Belge en Algebre (Faulx-les-Tombes, 1987). Bull. Soc. Math. Belg. Sér. A 40 (1988), no. 2, 239–253.
- [KeY] B. Keller, D. Yang, *Derived equivalences from mutations of quivers with potential*, Adv. Math. 226 (2011), no. 3, 2118–2168.
- [KM] Y. Kimura, Y. Mizuno, *Two-term tilting complexes for preprojective algebras of non-Dynkin type*, Communications in Algebra. <https://doi.org/10.1080/00927872.2021.1962337>.
- [Ki] A. D. King, *Moduli of representations of finite-dimensional algebras*, Quart. J. Math. Oxford Ser. (2) 45 (1994), no. 180, 515–530.
- [KY] S. Koenig, D. Yang, *Silting objects, simple-minded collections, t -structures and co- t -structures for finite-dimensional algebras*, Doc. Math. **19** (2014), 403–438.
- [KS] M. Kreuzer and H. Skarke, *Classification of polyhedra in three dimensions*, Adv. Theor. Math. Phys. 2 (1998), 853–871.
- [LZ] J. Lagarias and G. M. Ziegler, *Bounds for lattice polytopes containing a fixed number of interior points in a sublattice*, Canad. J. Math. 43 (1991), no. 5, 1022–1035.
- [L] T. Y. Lam *Lectures on modules and rings*, Graduate Texts in Mathematics, 189. Springer-Verlag.
- [M] Y. Mizuno, *Classifying τ -tilting modules over preprojective algebras of Dynkin type*, Math. Z. 277 (2014), no. 3–4, 665–690.
- [MY] Y. Mizuno, K. Yamaura, *Higher APR tilting preserves n -representation infiniteness*, J. Algebra 447 (2016), 56–73.
- [Nag] K. Nagao, *Donaldson-Thomas theory and cluster algebras*, Duke Math. J. 162 (2013), no. 7, 1313–1367.
- [Nak] T. Nakanishi, *Cluster Algebras and Scattering Diagrams, Part II. Cluster Patterns and Scattering Diagrams*, arXiv:2103.16309.

- [NZ] T. Nakanishi, A. Zelevinsky, *On tropical dualities in cluster algebras*, Algebraic groups and quantum groups, 217–226, Contemp. Math., 565, Amer. Math. Soc., Providence, RI, 2012.
- [O] M. Øbro, *Classification of smooth Fano polytopes*, Ph. D. Thesis, University of Aarhus.
- [PPPP] A. Padrol, Y. Palu, V. Pilaud, P. Plamondon, *Associahedra for finite type cluster algebras and minimal relations between g -vectors*, arXiv:1906.06861.
- [Pe] T. Petersen, *Eulerian numbers* Birkhäuser Advanced Texts: Basler Lehrbücher. Birkhäuser/Springer, New York, 2015.
- [Pl1] P. Plamondon, *Cluster algebras via cluster categories with infinite-dimensional morphism spaces*, Compos. Math. 147 (2011), no. 6, 1921–1954.
- [Pl2] P. Plamondon, *Generic bases for cluster algebras from the cluster category*, Int. Math. Res. Not. IMRN 2013, no. 10, 2368–2420.
- [PR] B. Poonen and F. Rodriguez-Villegas, *Lattice Polygons and the Number 12*, Amer. Math. Monthly 107 (2000), pp. 238 – 250.
- [Re] N. Reading, *Universal geometric cluster algebras*, Math. Z. 277 (2014), no. 1-2, 499–547.
- [ReS1] N. Reading, D. Speyer, *Cambrian fans*, J. Eur. Math. Soc. (JEMS) 11 (2009), no. 2, 407–447.
- [ReS2] N. Reading, D. Speyer, *Sortable elements in infinite Coxeter groups*, Trans. Amer. Math. Soc. 363 (2011), no. 2, 699–761.
- [RiS] C. Riedtmann, A. Schofield, *On a simplicial complex associated with tilting modules*, Comment. Math. Helv. 66 (1991), no. 1, 70–78.
- [Ri] C. M. Ringel, *Representations of K -species and bimodules*, J. Algebra 41 (1976), no. 2, 269–302.
- [Sa] H. Sato, *Towards the classification of higher-dimensional toric Fano varieties*, *Tohoku Math J.* **52** (2000), 383–413.
- [Sc] S. Schroll, *Brauer graph algebras: a survey on Brauer graph algebras, associated gentle algebras and their connections to cluster theory*, Homological methods, representation theory, and cluster algebras, 177–223, CRM Short Courses, Springer, Cham, 2018.
- [Sm] S. Smalø, *Torsion theories and tilting modules*, Bull. London Math. Soc. 16 (1984), no. 5, 518–522.
- [Sö] C. Söderberg, *Preprojective algebras of d -representation finite species with relations*, arXiv:2109.15187.
- [U] L. Unger, *Shellability of simplicial complexes arising in representation theory*, Adv. Math. 144 (1999), no. 2, 221–246.
- [V] R. H. Villarreal, *Monomial algebras*, Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, FL, 2015.
- [VK] V. E. Voskresenskii and A. A. Klyachko, *Toroidal Fano varieties and root systems*, *Math. USSR Izvestiya* **24** (1985), 221–244.
- [Y1] T. Yurikusa, *Wide subcategories are semistable*, Doc. Math. 23 (2018), 35–47.
- [Y2] T. Yurikusa, *Acyclic cluster algebras with dense g -vector fans*, arXiv:2107.13482.
- [Zi] G. M. Ziegler, *Lectures on polytopes*. Graduate Texts in Mathematics, 152. Springer-Verlag, New York, 1995.
- [Zh] Y. Zhang, *On mutation of τ -tilting modules*, Comm. Algebra 45 (2017), no. 6, 2726–2729.

GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, 1-5 YAMADAOKA, SUITA, OSAKA 565-0871, JAPAN

Email address: aoki-t@ist.osaka-u.ac.jp

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, 1-5 YAMADAOKA, SUITA, OSAKA 565-0871, JAPAN

Email address: higashitani@ist.osaka-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1 KOMABA MEGURO-KU TOKYO 153-8914, JAPAN

Email address: iyama@ms.u-tokyo.ac.jp

DEPARTMENT OF INFORMATION SCIENCE AND ENGINEERING, OKAYAMA UNIVERSITY OF SCIENCE, 1-1 RIDAICHO, KITA-KU, OKAYAMA 700-0005, JAPAN

Email address: r-kase@ous.ac.jp

FACULTY OF LIBERAL ARTS, SCIENCES AND GLOBAL EDUCATION / GRADUATE SCHOOL OF SCIENCE, OSAKA METROPOLITAN UNIVERSITY, 1-1 GAKUEN-CHO, NAKA-KU, SAKAI, OSAKA 599-8531, JAPAN

Email address: yuya.mizuno@omu.ac.jp