

NEW PROOFS OF SOME DOUBLE SUM ROGERS-RAMANUJAN TYPE IDENTITIES

LIUQUAN WANG

ABSTRACT. Recently, Rosengren utilized an integral method to prove a number of conjectural identities found by Kanade and Russell. Using this integral method, we give new proofs to some double sum identities of Rogers-Ramanujan type. These identities were earlier proved by approaches such as combinatorial arguments or by using q -difference equations. Our proofs are based on streamlined calculations, which relate these double sum identities to some known Rogers-Ramanujan type identities with single sums. Moreover, we prove a conjectural identity of Andrews and Uncu which was earlier confirmed by Chern.

1. INTRODUCTION AND MAIN RESULTS

The famous Rogers-Ramanujan identities state that

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n} = \frac{1}{(q, q^4; q^5)_{\infty}}, \quad \sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q; q)_n} = \frac{1}{(q^2, q^3; q^5)_{\infty}}. \quad (1.1)$$

Here and throughout this paper, we assume that $|q| < 1$ for convergence and use the standard q -series notation

$$(a; q)_0 := 1, \quad (a; q)_n := \prod_{k=0}^{n-1} (1 - aq^k), \quad (1.2)$$

$$(a_1, \dots, a_m; q)_n := (a_1; q)_n \cdots (a_m; q)_n, \quad n \in \mathbb{N} \cup \{\infty\}. \quad (1.3)$$

Despite their analytic forms given in (1.1), the Rogers-Ramanujan identities can also be formulated as combinatorial partition identities. The i -th ($i = 1, 2$) identity is equivalent to saying that the number of partitions of n such that the adjacent parts differ by least 2 and such that the smallest part is at least i is the same as the number of partitions of n into parts congruent to $\pm i$ modulo 5.

Since the appearances of (1.1), lots of similar identities have been found and they were called as Rogers-Ramanujan type identities. One of the famous works on this topic is Slater's list [26], which contains 130 such identities. For example, the following identities are in Slater's list [26, Eqs. (34), (36)]

$$\sum_{n=0}^{\infty} \frac{q^{n^2+2n}(-q; q^2)_n}{(q^2; q^2)_n} = \frac{1}{(q^3, q^4, q^5; q^8)_{\infty}}, \quad (1.4)$$

2010 *Mathematics Subject Classification.* 11P84, 33D15, 11B65.

Key words and phrases. Rogers-Ramanujan identities; sum-product identities; Partitions.

$$\sum_{n=0}^{\infty} \frac{q^{n^2}(-q; q^2)_n}{(q^2; q^2)_n} = \frac{1}{(q, q^4, q^7; q^8)_{\infty}}. \quad (1.5)$$

Both identities were earlier found by Ramanujan [5, Entries 1.17.11, 1.7.12, 4.2.15]. Göllnitz [15, Eqs. (2.22), (2.24)] found the following companion identities:

$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)}(-q^{-1}; q^2)_n}{(q^2; q^2)_n} = \frac{1}{(q, q^5, q^6; q^8)_{\infty}}, \quad (1.6)$$

$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)}(-q; q^2)_n}{(q^2; q^2)_n} = \frac{1}{(q^2, q^3, q^7; q^8)_{\infty}}. \quad (1.7)$$

There is now a huge list of Rogers-Ramanujan type identities, and it is evidently that more and more will be discovered in the future. We refer the reader to Sills' book [25] on this topic for more information.

One of the way to generalize the Rogers-Ramanujan identities is to consider multi-sum identities. For instance, the Andrews-Gordon identities state that

$$\sum_{n_1, \dots, n_{k-1} \geq 0} \frac{q^{N_1^2 + \dots + N_{k-1}^2 + N_1 + \dots + N_{k-1}}}{(q; q)_{n_1} \cdots (q; q)_{n_{k-1}}} = \frac{(q^i, q^{2k+1-i}, q^{2k+1}; q^{2k+1})_{\infty}}{(q; q)_{\infty}}, \quad (1.8)$$

where $1 \leq i \leq k$, $k \geq 2$ and $N_j = n_j + \dots + n_{k-1}$. This identity was given by Andrews [1], and is the analytic counterpart of Gordon's [16] combinatorial generalization of (1.1).

In recent years, multi-sum identities of the Rogers-Ramanujan type have drawn lots of attention. In particular, a number of recent works studied Rogers-Ramanujan type identities in the form

$$\text{finite sum of } \sum_{(i_1, \dots, i_k) \in S} \frac{(-1)^{t(i_1, \dots, i_k)} q^{Q(i_1, \dots, i_k)}}{(q^{n_1}; q^{n_1})_{i_1} \cdots (q^{n_k}; q^{n_k})_{i_k}} = \prod_{(a, n) \in P} (q^a; q^n)_{\infty}^{r(a, n)}. \quad (1.9)$$

Here $t(i_1, \dots, i_k)$ is an integer-valued function, $Q(i_1, \dots, i_k)$ is a rational polynomial in variables i_1, \dots, i_k , n_i ($i = 1, 2, \dots, k$) are positive integers with $\gcd(n_1, \dots, n_k) = 1$, P is a finite subset of \mathbb{Q}^2 and $r(a, n)$ are integer-valued functions.

We shall call any identity of the form (1.9) a Rogers-Ramanujan type identity with *index* (n_1, \dots, n_k) . Note that here we do not allow q -Pochhammer symbols to appear in the numerator in each summand. But as we can see from (1.4)–(1.7), this may be allowed if we discuss in a more general context.

Around 2019, Kanade and Russell [19] searched for Rogers-Ramanujan type identities related to level 2 characters of the affine Lie algebra $A_9^{(2)}$, and they conjectured a number of such identities. Let

$$F(u, v, w) := \sum_{i, j, k=0}^{\infty} \frac{(-1)^k q^{3k(k-1) + (i+2j+3k)(i+2j+3k-1)} u^i v^j w^k}{(q; q)_i (q^4; q^4)_j (q^6; q^6)_k}, \quad (1.10)$$

which is roughly speaking a triple sum of index $(1, 4, 6)$ according to our definition. Some of their conjectural identities are

$$F(q, q^6, q^9) = \frac{1}{(q, q^4, q^6, q^8, q^{11}; q^{12})_\infty}, \quad (1.11)$$

$$F(q, q, q^6) = \frac{1}{(q^3; q^4)_\infty (q, q^8; q^{12})_\infty}. \quad (1.12)$$

Five of their conjectural identities were proved by Bringmann, Jennings-Shaffer and Mahlburg [8]. In a recent work, Rosengren [24] presented a streamlined approach and proved all of the nine conjectural identities on $F(u, v, w)$.

There are two steps in Rosengren's approach. First, one uses an integral representation of the triple sum to reduce it to a single sum. Second, one needs to evaluate the single sum via q -series techniques. The difficulty in the second step varies a lot for different identities. We shall call this approach as the integral method. With this method, Rosengren also gave a new proof of the following identity with index $(1, 3)$:

$$\sum_{i,j \geq 0} \frac{q^{2i^2+6ij+6j^2}}{(q; q)_i (q^3; q^3)_j} = \frac{1}{(q^3; q^6)_\infty (q^2, q^{10}; q^{12})_\infty}. \quad (1.13)$$

As shown in [19] and [20], this identity is equivalent to a conjectural partition identity of Capparelli [10], which was first proved by Andrews [3].

There are more double sum Rogers-Ramanujan type identities in the literature and their proofs differ from each other. Some of them can be converted to a single sum in a straightforward way. For example, Takigiku and Tsuchioka [27] proved three identities with index $(1, 2)$ such as

$$\sum_{i,j \geq 0} \frac{(-1)^j q^{\binom{i}{2} + 2\binom{j}{2} + 2ij + i + j}}{(q; q)_i (q^2; q^2)_j} = \frac{1}{(q^2, q^3, q^4, q^{10}, q^{11}, q^{12}; q^{14})_\infty}. \quad (1.14)$$

Their original proofs of these identities use q -difference equations. But they also mentioned [27, Remark 4.5] that, by summing over j first using Euler's identity

$$\sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} z^n}{(q; q)_n} = (-z; q)_\infty, \quad (1.15)$$

the sum side in (1.14) can easily be shown equal to the left side of Slater's identity [26, Eq. (81)]

$$\sum_{i=0}^{\infty} \frac{q^{\frac{1}{2}(i^2+i)}}{(q; q)_i (q; q^2)_i} = \frac{(q, q^6, q^7; q^7)_\infty (q^5, q^9; q^{14})_\infty}{(q; q)_\infty (q; q^2)_\infty}, \quad (1.16)$$

and hence (1.14) holds.

The same trick was applied by Andrews in [4] and he found three more identities with index $(1, 2)$ such as

$$\sum_{i,j \geq 0} \frac{(-1)^j q^{(i+j)^2}}{(q; q)_i (q^2; q^2)_j} = \frac{1}{(q^2, q^4, q^{10}, q^{12}; q^{14})_\infty}. \quad (1.17)$$

We will focus on double sum Rogers-Ramanujan type identities in which the sum indexes cannot be separated or eliminated directly. Clearly (1.13) is such an identity. We can find some other such identities in the literature. In a recent work, Uncu and Zudilin [28] discovered two identities with index $(1, 2)$ (see (1.18) and (1.19) below). In the way of proving their identities, we find a companion identity (1.20) and we state them together as follows.

Theorem 1.1. *We have*

$$\sum_{i,j \geq 0} \frac{q^{i^2+2ij+2j^2}}{(q; q)_i (q^2; q^2)_j} = \frac{(q^3; q^3)_\infty^2}{(q; q)_\infty (q^6; q^6)_\infty}, \quad (1.18)$$

$$\sum_{i,j \geq 0} \frac{q^{i^2+2ij+2j^2+i+2j}}{(q; q)_i (q^2; q^2)_j} = \frac{(q^6; q^6)_\infty^2}{(q^2; q^2)_\infty (q^3; q^3)_\infty}, \quad (1.19)$$

$$\sum_{i,j \geq 0} \frac{q^{i^2+2ij+2j^2+j}}{(q; q)_i (q^2; q^2)_j} = (-q; q)_\infty. \quad (1.20)$$

Uncu and Zudilin [28] pointed out that, as explained by Ole Warnaar, these two identities are instances of Bressoud's identities from [7]. We will provide new proofs, which relate (1.18) to Ramanujan's identity [5, p. 87, Entry 4.2.11]:

$$\sum_{n=0}^{\infty} \frac{(-1)^n (q; q^2)_n q^{n(n+2)}}{(q^4; q^4)_n} = \frac{(q^3; q^3)_\infty (q^{12}; q^{12})_\infty}{(q^4; q^4)_\infty (q^6; q^6)_\infty}, \quad (1.21)$$

and relate (1.19) to the following identities of Ramanujan [5, pp. 88, 101, Entries 4.2.13 and 5.3.2]:

$$\sum_{n=0}^{\infty} \frac{(-q^2; q^2)_n q^{n(n+1)}}{(q; q)_{2n+1}} = \frac{(q^3; q^3)_\infty (q^{12}; q^{12})_\infty}{(q; q)_\infty (q^6; q^6)_\infty}, \quad (1.22)$$

$$\sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^{n^2}}{(q; q)_{2n}} = \frac{(q^6; q^6)_\infty^2}{(q; q)_\infty (q^{12}; q^{12})_\infty}. \quad (1.23)$$

The identity (1.20) appears to be new. We will see that it is a consequence of (1.15).

In 2021, through q -difference equations, Andrews and Uncu [6, Theorem 1.1] proved the following interesting identity with index $(1, 3)$.

Theorem 1.2.

$$\sum_{i,j \geq 0} \frac{(-1)^j q^{i^2+3ij+3j(3j+1)/2}}{(q; q)_i (q^3; q^3)_j} = \frac{1}{(q; q^3)_\infty}. \quad (1.24)$$

They also conjectured that [6, Conjecture 1.2]

$$\sum_{i,j \geq 0} \frac{(-1)^j q^{\frac{3j(3j+1)}{2}+i^2+3ij+i+j}}{(q; q)_i (q^3; q^3)_j} = \frac{1}{(q^2, q^3; q^6)_\infty}. \quad (1.25)$$

Let $\omega = e^{2\pi i/3}$ throughout this paper. We find that (1.24) can be proved through the following Rogers-Ramanujan type identity

$$\sum_{n=0}^{\infty} \frac{(-1)^n \omega^n q^{\frac{1}{2}n(n+1)}}{(q, \omega^2 q; q)_n} = \frac{1}{(q; q^3)_{\infty}}, \quad (1.26)$$

which we will prove in this paper. Furthermore, we find that in order to prove the conjectural identity (1.25), it suffices to show the following identity.

Theorem 1.3. *We have*

$$\sum_{n=0}^{\infty} \frac{(-1)^n q^{\frac{3}{2}n^2 + \frac{5}{2}n} (q; q^3)_n}{(q^9; q^9)_n} = \frac{(q^4; q^6)_{\infty} (q^{12}; q^{18})_{\infty}}{(q^5; q^6)_{\infty} (q^9; q^{18})_{\infty}}. \quad (1.27)$$

Remark 1. This theorem was posed as a conjecture in the first version of this paper (appeared as arXiv: 2203.15572v1), which contains essentially all materials of this paper except that Theorem 1.3 is unproven. After the first draft has been written, we heard from Shane Chern [12] that he has independently worked on the identity (1.25) and also discovered (1.27). He [13] has successfully proved (1.27) by proving the identity

$$\sum_{n=0}^{\infty} \frac{(a; q)_n (a^{-1} q^2; q^2)_n}{(a^2 q; q^2)_n (q^3; q^3)_n} (-1)^n a^n q^{n(n+1)/2} = \frac{(aq; q^2)_{\infty} (a^3 q^3; q^6)_{\infty}}{(a^2 q; q^2)_{\infty} (q^3; q^6)_{\infty}}. \quad (1.28)$$

The above theorem follows by replacing q with q^3 and setting $a = q$ in (1.28).

Note that Chern proved (1.28) by proving a finite sum version of it, we will give a different proof of Theorem 1.3 by pointing out that (1.28) is in fact a direct consequence of a cubic summation formula due to Rahman [22]. As a result, this yields another proof of the conjectural identity (1.25).

Theorem 1.4. *The identity (1.25) holds.*

Finally, we discuss a set of identities with index $(1, 4)$, which can be found in the work of Kurşungöz [20, Corollary 18].

Theorem 1.5. *We have*

$$\sum_{i,j \geq 0} \frac{q^{(3i^2-i)/2 + 4ij + 4j^2}}{(q; q)_i (q^4; q^4)_j} = \frac{1}{(q, q^4, q^7; q^8)_{\infty}}, \quad (1.29)$$

$$\sum_{i,j \geq 0} \frac{q^{(3i^2+3i)/2 + 4ij + 4j^2 + 4j}}{(q; q)_i (q^4; q^4)_j} = \frac{1}{(q^3, q^4, q^5; q^8)_{\infty}}, \quad (1.30)$$

$$\sum_{i,j \geq 0} \frac{q^{(3i^2+i)/2 + 4ij + 4j^2 + 2j} (1 + q^{2i+4j+1})}{(q; q)_i (q^4; q^4)_j} = \frac{1}{(q, q^5, q^6; q^8)_{\infty}}, \quad (1.31)$$

$$\sum_{i,j \geq 0} \frac{q^{(3i^2+i)/2 + 4ij + 4j^2 + 2j}}{(q; q)_i (q^4; q^4)_j} = \frac{1}{(q^2, q^3, q^7; q^8)_{\infty}}. \quad (1.32)$$

We will provide new proofs of these identities using the integral method, and by this we see that they are related to the identities (1.4)–(1.7).

The paper is organized as follows. In Section 2 we collect some useful q -series formulas which will be used to derive the identities. In Section 3 we present new proofs for Theorems 1.1, 1.2, 1.5 and give a proof for Theorem 1.4. Finally, in Section 4 we propose some conjectural identities with index (1, 2, 3, 4).

2. PRELIMINARIES

Besides (1.15), we also need another identity due to Euler

$$\sum_{n=0}^{\infty} \frac{z^n}{(q; q)_n} = \frac{1}{(z; q)_{\infty}}, \quad |z| < 1 \quad (2.1)$$

and the Jacobi triple product identity

$$(q, z, q/z; q)_{\infty} = \sum_{n=-\infty}^{\infty} (-1)^n q^{\binom{n}{2}} z^n. \quad (2.2)$$

They will be used frequently to express multi-sums as integrals of infinite products. Both (1.15) and (2.1) can be deduced from the q -binomial theorem: for $|q|, |z| < 1$,

$$\sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} z^n = \frac{(az; q)_{\infty}}{(z; q)_{\infty}}. \quad (2.3)$$

One of the key step in establishing Rogers-Ramanujan type identities by the integral method is to evaluate integrals of infinite q -products and express the result as infinite q -products. We will mainly use an identity from the book of Gasper and Rahman [14]. Before we state this result, we remark that the symbol “idem ($c_1; c_2, \dots, c_C$)” after an expression stands for the sum of the $(C - 1)$ expressions obtained from the preceding expression by interchanging c_1 with each c_k , $k = 2, 3, \dots, C$.

Lemma 2.1. (Cf. [14, Eq. (4.10.6)]) Suppose that

$$P(z) := \frac{(a_1 z, \dots, a_A z, b_1/z, \dots, b_B/z; q)_{\infty}}{(c_1 z, \dots, c_C z, d_1/z, \dots, d_D/z; q)_{\infty}}$$

has only simple poles. We have

$$\begin{aligned} \oint P(z) \frac{dz}{2\pi i z} &= \frac{(b_1 c_1, \dots, b_B c_1, a_1/c_1, \dots, a_A/c_1; q)_{\infty}}{(q, d_1 c_1, \dots, d_D c_1, c_2/c_1, \dots, c_C/c_1; q)_{\infty}} \\ &\times \sum_{n=0}^{\infty} \frac{(d_1 c_1, \dots, d_D c_1, q c_1/a_1, \dots, q c_1/a_A; q)_n}{(q, b_1 c_1, \dots, b_B c_1, q c_1/c_2, \dots, q c_1/c_C; q)_n} \\ &\times \left(-c_1 q^{(n+1)/2} \right)^{n(C-A)} \left(\frac{a_1 \cdots a_A}{c_1 \cdots c_C} \right)^n + \text{idem } (c_1; c_2, \dots, c_C) \end{aligned} \quad (2.4)$$

when $C > A$, or if $C = A$ and

$$\left| \frac{a_1 \cdots a_A}{c_1 \cdots c_C} \right| < 1. \quad (2.5)$$

Here the integration is over a positively oriented contour so that the poles of $(c_1 z, \dots, c_C z; q)_\infty^{-1}$ lie outside the contour, and the origin and poles of $(d_1/z, \dots, d_D/z; q)_\infty^{-1}$ lie inside the contour.

Heine's transformation formula [14, p. 13, Eq. (1.4.3)] states that for $|t| < 1$,

$$\sum_{n=0}^{\infty} \frac{(a, b; q)_n}{(c, q; q)_n} t^n = \frac{(abt/c; q)_\infty}{(t; q)_\infty} \sum_{n=0}^{\infty} \frac{(\frac{c}{a}, \frac{c}{b}; q)_n}{(c, q; q)_n} \left(\frac{abt}{c} \right)^n. \quad (2.6)$$

We also need Lebesgue's identity (see [2, p. 21, Cor. 2.7], for example)

$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)/2} (a; q)_n}{(q; q)_n} = (aq; q^2)_\infty (-q; q)_\infty. \quad (2.7)$$

We will occasionally use the notation $f_m := (q^m; q^m)_\infty$ when doing calculations. The following lemma will be used in some proofs.

Lemma 2.2. *We have*

$$\frac{f_3}{f_1^3} = \frac{f_4 f_6^3}{f_2^9 f_{12}^2} + 3q \frac{f_4^2 f_6 f_{12}^2}{f_2^7}, \quad (2.8)$$

$$\frac{f_1^3}{f_3} = \frac{f_4^3}{f_{12}} - 3q \frac{f_2^2 f_{12}^3}{f_4 f_6^2}. \quad (2.9)$$

Proofs of (2.8) and (2.9) can be found in [29] and [17], respectively.

3. PROOFS OF THE THEOREMS

In this section, we will present proofs of the main results stated in Section 1. Throughout this section, the contours of integrals will be chosen in the way as in Lemma 2.1.

Proof of Theorem 1.1. We shall consider

$$F_1(u, v; q) := \oint \frac{(uz, qz, 1/z, q; q)_\infty}{(vz, -vz; q)_\infty} \frac{dz}{2\pi iz}. \quad (3.1)$$

By (1.15) and (2.1) we see that

$$\begin{aligned} F_1(u, v; q) &= \oint \sum_{i=0}^{\infty} \frac{(-1)^i q^{\binom{i}{2}} u^i z^i}{(q; q)_i} \sum_{j=0}^{\infty} \frac{v^{2j} z^{2j}}{(q^2; q^2)_j} \sum_{k=-\infty}^{\infty} (-1)^k q^{\binom{k}{2}} z^{-k} \frac{dz}{2\pi iz} \\ &= \sum_{i, j \geq 0} \frac{u^i v^{2j} q^{i^2 + 2ij + 2j^2 - i - j}}{(q; q)_i (q^2; q^2)_j}. \end{aligned} \quad (3.2)$$

To prove (1.18), we choose $u = q$ and $v = q^{\frac{1}{2}}$. Setting

$$(A, B, C, D) = (2, 1, 2, 0), \quad a_1 = a_2 = q, \quad b_1 = 1, \quad (c_1, c_2) = (q^{1/2}, -q^{1/2})$$

in (2.4), we deduce that

$$F_1(q, q^{1/2}; q) = (q; q)_\infty \left(\frac{(q^{\frac{1}{2}}; q)_\infty^3}{(q, -1; q)_\infty} S_1(q) + \frac{(-q^{\frac{1}{2}}; q)_\infty^3}{(q, -1; q)_\infty} S_2(q) \right), \quad (3.3)$$

where

$$S_1(q) = \sum_{n=0}^{\infty} \frac{(q^{\frac{1}{2}}; q)_n (-q)^n}{(q, -q; q)_n}, \quad S_2(q) = \sum_{n=0}^{\infty} \frac{(-q^{\frac{1}{2}}; q)_n (-q)^n}{(q, -q; q)_n}.$$

By Heine's transformation (2.6) with $(q, a, b, c, t) \rightarrow (q^2, q, 0, -q^2, -q^2)$, we obtain

$$S_1(q^2) = \frac{1}{(-q^2; q^2)_{\infty}} \sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^{n^2+2n}}{(-q^2, q^2; q^2)_n} = \frac{f_2 f_6^2}{f_3 f_4^2}, \quad (3.4)$$

where the last equality follows from (1.21) with q replaced by $-q$.

Replacing q by $-q$ in the expression of $S_1(q^2)$ we get $S_2(q^2)$ and hence

$$S_2(q^2) = \frac{f_2 f_3 f_{12}}{f_4^2 f_6}. \quad (3.5)$$

Hence

$$\begin{aligned} F_1(q^2, q; q^2) &= (q^2; q^2)_{\infty} \left(\frac{(q; q^2)_{\infty}^3}{2(q^4; q^4)_{\infty}} S_1(q^2) + \frac{(-q; q^2)_{\infty}^3}{2(q^4; q^4)_{\infty}} S_2(q^2) \right) \\ &= \frac{1}{2} f_2 \left(\frac{f_1^3}{f_3} \cdot \frac{f_6^2}{f_2^2 f_4^3} + \frac{f_3}{f_1^3} \cdot \frac{f_2^7 f_{12}}{f_4^6 f_6} \right) \\ &= \frac{f_6^2}{f_2 f_{12}}. \end{aligned} \quad (3.6)$$

Here the last equality follows by substitution of (2.8) and (2.9) and simplification. Replacing q by $q^{\frac{1}{2}}$, we obtain (1.18).

To prove (1.19), we need to evaluate $F_1(q^2, q^{\frac{3}{2}}; q)$. But in this case Lemma 2.1 is not applicable since $C = A$ and the condition (2.5) does not hold. Instead, we shall use the integral

$$F_2(u, v; q) := \oint \frac{(vz, q^2 z, 1/z, q^2; q^2)_{\infty}}{(uz; q)_{\infty}} \frac{dz}{2\pi i z}. \quad (3.7)$$

By (1.15), (2.1) and (2.2) we have

$$\begin{aligned} F_2(u, v; q) &= \oint \sum_{i=0}^{\infty} \frac{u^i z^i}{(q; q)_i} \sum_{j=0}^{\infty} \frac{(-1)^j z^j v^j q^{j^2-j}}{(q^2; q^2)_j} \sum_{\ell=-\infty}^{\infty} (-1)^{\ell} q^{\ell^2-\ell} z^{-\ell} \frac{dz}{2\pi i z} \\ &= \sum_{i,j \geq 0} \frac{(-1)^i u^i v^j q^{i^2+2ij+2j^2-i-2j}}{(q; q)_i (q^2; q^2)_j}. \end{aligned} \quad (3.8)$$

Now we set $u = -q^2, v = q^4$. Replacing q by q^2 in (2.4) and letting

$$(A, B, C, D) = (2, 1, 2, 0), \quad (a_1, a_2) = (q^2, q^4), \quad b_1 = 1, \quad (c_1, c_2) = (-q^2, -q^3),$$

we deduce that

$$F_2(-q^2, q^4; q) = 2 \frac{(-q^2; q^2)_{\infty}^3}{(q; q)_{\infty}} T_1(q) - \frac{(-q; q^2)_{\infty}^3}{(q; q)_{\infty}} T_2(q), \quad (3.9)$$

where

$$T_1(q) = \sum_{n=0}^{\infty} \frac{(-1; q^2)_n q^n}{(q; q)_{2n}}, \quad T_2(q) = \sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^n}{(q; q)_{2n+1}}. \quad (3.10)$$

Now by Heine's transformation (2.6) with $(q, a, b, c, t) \rightarrow (q^2, -1, 0, q, q)$ and (1.23), we deduce that

$$T_1(q) = \frac{1}{(q; q^2)_{\infty}} \sum_{n=0}^{\infty} \frac{(-q; q^2)_n q^{n^2}}{(q, q^2; q^2)_n} = \frac{f_2 f_6^2}{f_1^2 f_{12}}. \quad (3.11)$$

Similarly, using (2.6) with $(q, a, b, c, t) \rightarrow (q^2, -q, 0, q^3, q)$ and (1.22), we deduce that

$$T_2(q) = \frac{1}{(q; q^2)_{\infty}} \sum_{n=0}^{\infty} \frac{(-q^2; q^2)_n q^{n^2+n}}{(q; q)_{2n+1}} = \frac{f_2 f_3 f_{12}}{f_1^2 f_6}. \quad (3.12)$$

Now substituting (3.11) and (3.12) into (3.9), we deduce that

$$\begin{aligned} F_2(-q^2, q^4; q) &= 2 \frac{f_4^3 f_6^2}{f_1^3 f_2^2 f_{12}} - \frac{f_2^7 f_3 f_{12}}{f_1^6 f_4^3 f_6} \\ &= \frac{1}{f_1^3} \left(2 \frac{f_4^3 f_6^2}{f_2^2 f_{12}} - \frac{f_2^7 f_{12}}{f_4^3 f_6} \cdot \frac{f_3}{f_1^3} \right) \quad (\text{use (2.8)}) \\ &= \frac{1}{f_1^3} \left(2 \frac{f_4^3 f_6^2}{f_2^2 f_{12}} - \frac{f_2^7 f_{12}}{f_4^3 f_6} \cdot \left(\frac{f_4^6 f_6^3}{f_2^9 f_{12}^2} + 3q \frac{f_4^2 f_6 f_{12}^2}{f_2^7} \right) \right) \\ &= \frac{f_6^2}{f_1^3 f_2^2} \left(\frac{f_4^3}{f_{12}} - 3q \frac{f_2^2 f_{12}^3}{f_4 f_6^2} \right) \quad (\text{by (2.9)}) \\ &= \frac{f_6^2}{f_2^2 f_3}. \end{aligned}$$

Finally, to prove (1.20), setting $u = v = q$ in (3.2), we see that the left side of (1.20) is

$$\begin{aligned} F_1(q, q; q) &= \oint \frac{(qz, 1/z, q; q)_{\infty}}{(-qz; q)_{\infty}} \frac{dz}{2\pi i z} \\ &= \oint \sum_{k=0}^{\infty} \frac{(-qz)^k}{(q; q)_k} \sum_{\ell=-\infty}^{\infty} (-1)^{\ell} q^{\binom{\ell}{2}} z^{-\ell} \frac{dz}{2\pi i z} \\ &= \sum_{k=0}^{\infty} \frac{q^{k(k+1)/2}}{(q; q)_k} = (-q; q)_{\infty}. \end{aligned}$$

Here the second and last equalities follow from (1.15) and (2.1). \square

Remark 2. For (1.20), we cannot use (2.4) to evaluate $F_1(q, q; q)$ directly since $A = C = 1$ and $|a_1/c_1| = 1$.

Proof of Theorem 1.2. We start with the integral

$$\begin{aligned} R(q) &:= \oint \sum_{i=0}^{\infty} \frac{q^{\frac{1}{2}(i^2+i)}(-1)^i z^i}{(q; q)_i} \sum_{j=0}^{\infty} \frac{q^{3j} z^{3j}}{(q^3; q^3)_j} \sum_{k=-\infty}^{\infty} (-1)^k q^{\frac{1}{2}(k^2-k)} z^{-k} \frac{dz}{2\pi i z} \\ &= \sum_{i,j \geq 0} \frac{(-1)^j q^{i^2+3ij+\frac{3}{2}j(3j+1)}}{(q; q)_i (q^3; q^3)_j}. \end{aligned} \quad (3.13)$$

By (1.15), (2.1) and (2.2), we can write

$$R(q) = \oint \frac{(qz, qz, 1/z, q; q)_{\infty}}{(q^3 z^3; q^3)_{\infty}} \frac{dz}{2\pi i z} = \oint \frac{(qz, 1/z, q; q)_{\infty}}{(\omega q z, \omega^2 q z; q)_{\infty}} \frac{dz}{2\pi i z}. \quad (3.14)$$

Now applying (2.4) with

$$(A, B, C, D) = (1, 1, 2, 0), \quad a_1 = q, \quad b_1 = 1, \quad (c_1, c_2) = (\omega q, \omega^2 q),$$

we deduce that

$$R(q) = \frac{(\omega^2; q)_{\infty}}{1 - \omega} R_1(q) + \frac{(\omega; q)_{\infty}}{1 - \omega^2} R_2(q), \quad (3.15)$$

where

$$R_1(q) = \sum_{n=0}^{\infty} \frac{(-1)^n \omega^n q^{\frac{1}{2}n(n+1)}}{(q, \omega^2 q; q)_n}, \quad (3.16)$$

$$R_2(q) = \sum_{n=0}^{\infty} \frac{(-1)^n \omega^{2n} q^{\frac{1}{2}n(n+1)}}{(q, \omega q; q)_n}. \quad (3.17)$$

From [21, Eq. (6.1.11)] we find

$$\sum_{n=0}^{\infty} \frac{q^{\frac{1}{2}n(n-1)}(-b)^n}{(a, q; q)_n} = (b; q)_{\infty} \sum_{n=0}^{\infty} \frac{q^{\frac{1}{2}(3n^2-3n)}(-ab)^n}{(q, a, b; q)_n}. \quad (3.18)$$

Setting $a = \omega^2 q$ and $b = \omega q$ in the above identity and using (1.15), we obtain

$$R_1(q) = (\omega q; q)_{\infty} \sum_{n=0}^{\infty} \frac{(-1)^n q^{\frac{1}{2}(3n^2+n)}}{(q^3; q^3)_n} = (\omega q; q)_{\infty} (q^2; q^3)_{\infty}. \quad (3.19)$$

Replacing ω by ω^2 , we obtain

$$R_2(q) = (\omega^2 q; q)_{\infty} (q^2; q^3)_{\infty}. \quad (3.20)$$

Substituting (3.19) and (3.20) into (3.15), we deduce that

$$R(q) = (q^2; q^3)_{\infty} (\omega q; q)_{\infty} (\omega^2 q; q)_{\infty} \left(\frac{1 - \omega^2}{1 - \omega} + \frac{1 - \omega}{1 - \omega^2} \right) = \frac{1}{(q; q^3)_{\infty}}.$$

□

Proof of Theorem 1.3. Recall the following formula found by Rahman [22]:

$$\sum_{n=0}^{\infty} \frac{(1 - a q^{5n})(a, b; q^2)_n (ab^2/q^3; q^3)_n (q^2/b; q)_n (a q^3/b; q^6)_n}{(1 - a)(q^3, a q^3/b; q^3)_n (q^5/b^2; q^2)_n (ab, ab q^2; q^4)_n} \left(-\frac{q^2}{b} \right)^n q^{\binom{n+1}{2}}$$

$$= \frac{(aq^2, q^3/b; q^2)_\infty (ab^2, q^9/b^3; q^6)_\infty}{(ab, q^5/b^2; q^2)_\infty (q^3, aq^6/b; q^6)_\infty}. \quad (3.21)$$

This formula also appears as [14, Exercise 3.29(i)] except that a typo has been corrected here.

Setting $a = 0$ in (3.21), we obtain

$$\sum_{n=0}^{\infty} \frac{(q^2/b; q)_n (b; q^2)_n}{(q^5/b^2; q^2)_n (q^3; q^3)_n} (-1)^n \left(-\frac{q^2}{b}\right)^n q^{n(n+1)/2} = \frac{(q^3/b; q^2)_\infty (q^9/b^3; q^6)_\infty}{(q^5/b^2; q^2)_\infty (q^3; q^6)_\infty}. \quad (3.22)$$

Setting $b = q$ we complete the proof of Theorem 1.3. In fact, replacing b by q^2/b we recover Chern's formula (1.28). \square

Proof of Theorem 1.4. We consider the series

$$\begin{aligned} \overline{R}(q) &:= \oint \sum_{i=0}^{\infty} \frac{q^{\frac{1}{2}(i^2+3i)} (-1)^i z^i}{(q; q)_i} \sum_{j=0}^{\infty} \frac{z^{3j} q^{4j}}{(q^3; q^3)_j} \sum_{k=-\infty}^{\infty} (-1)^k z^{-k} q^{\frac{1}{2}(k^2-k)} \frac{dz}{2\pi i z} \\ &= \sum_{i,j \geq 0} \frac{(-1)^j q^{i^2+3ij+\frac{9}{2}j^2+i+\frac{5}{2}j}}{(q; q)_i (q^3; q^3)_j}. \end{aligned} \quad (3.23)$$

Now using (1.15), (2.1) and (2.2) we deduce that

$$\overline{R}(q) = \oint \frac{(q^2 z, qz, 1/z, q; q)_\infty}{(q^4 z^3; q^3)_\infty} \frac{dz}{2\pi i z} = \oint \frac{(q^2 z, qz, 1/z, q; q)_\infty}{(q^{\frac{4}{3}} z, \omega q^{\frac{4}{3}} z, \omega^2 q^{\frac{4}{3}} z; q)_\infty} \frac{dz}{2\pi i z} \quad (3.24)$$

Now setting

$$(A, B, C, D) = (2, 1, 3, 0), \quad (a_1, a_2) = (q^2, q), \quad b_1 = 1, \quad (c_1, c_2, c_3) = (q^{\frac{4}{3}}, \omega q^{\frac{4}{3}}, \omega^2 q^{\frac{4}{3}})$$

in (2.4), we deduce that

$$\begin{aligned} \overline{R}(q) &= \frac{(q^{\frac{4}{3}}, q^{\frac{2}{3}}, q^{-\frac{1}{3}}; q)_\infty}{(\omega, \omega^2; q)_\infty} \overline{R}_1(q) + \frac{(\omega q^{\frac{4}{3}}, \omega^2 q^{\frac{2}{3}}, \omega^2 q^{-\frac{1}{3}}; q)_\infty}{(\omega^2, \omega; q)_\infty} \overline{R}_2(q) \\ &\quad + \frac{(\omega^2 q^{\frac{4}{3}}, \omega q^{\frac{2}{3}}, \omega q^{-\frac{1}{3}}; q)_\infty}{(\omega, \omega^2; q)_\infty} \overline{R}_3(q) \\ &= -\frac{1}{3} q^{-\frac{1}{3}} \frac{(q; q)_\infty}{(q^3; q^3)_\infty} \left((q^{\frac{1}{3}}, q^{\frac{2}{3}}, q^{\frac{2}{3}}; q)_\infty \overline{R}_1(q) + \omega^2 (\omega q^{\frac{1}{3}}, \omega^2 q^{\frac{2}{3}}, \omega^2 q^{\frac{2}{3}}; q)_\infty \overline{R}_2(q) \right. \\ &\quad \left. + \omega (\omega^2 q^{\frac{1}{3}}, \omega q^{\frac{2}{3}}, \omega q^{\frac{2}{3}}; q)_\infty \overline{R}_3(q) \right), \end{aligned} \quad (3.25)$$

where

$$\overline{R}_1(q) = \sum_{n=0}^{\infty} \frac{(-1)^n (q^{\frac{1}{3}}; q)_n q^{\frac{1}{2}n^2 + \frac{5}{6}n}}{(q, \omega q, \omega^2 q; q)_n}, \quad (3.26)$$

$$\overline{R}_2(q) = \sum_{n=0}^{\infty} \frac{(-1)^n (\omega q^{\frac{1}{3}}; q)_n \omega^n q^{\frac{1}{2}n^2 + \frac{5}{6}n}}{(q, \omega q, \omega^2 q; q)_n}, \quad (3.27)$$

$$\overline{R}_3(q) = \sum_{n=0}^{\infty} \frac{(-1)^n (\omega^2 q^{\frac{1}{3}}; q)_n \omega^{2n} q^{\frac{1}{2}n^2 + \frac{5}{6}n}}{(q, \omega q, \omega^2 q; q)_n}. \quad (3.28)$$

By Theorem 1.3 we have

$$\overline{R}_1(q^3) = \frac{(q^4; q^6)_{\infty} (q^{12}; q^{18})_{\infty}}{(q^5; q^6)_{\infty} (q^9; q^{18})_{\infty}}. \quad (3.29)$$

If we replace q by ωq or $\omega^2 q$ in the definition of $\overline{R}_1(q^3)$, then we get $\overline{R}_2(q^3)$ and $\overline{R}_3(q^3)$, respectively. Thus

$$\overline{R}_2(q^3) = \frac{(\omega q^4; q^6)_{\infty} (q^{12}; q^{18})_{\infty}}{(\omega^2 q^5; q^6)_{\infty} (q^9; q^{18})_{\infty}}, \quad (3.30)$$

$$\overline{R}_3(q^3) = \frac{(\omega^2 q^4; q^6)_{\infty} (q^{12}; q^{18})_{\infty}}{(\omega q^5; q^6)_{\infty} (q^9; q^{18})_{\infty}}. \quad (3.31)$$

Now replacing q by q^3 in (3.25) and substituting (3.29)–(3.31) into it, we obtain

$$\overline{R}(q^3) = -\frac{1}{3} \frac{(q^3; q^3)_{\infty} (q^{12}; q^{18})_{\infty}}{(q^9; q^9)_{\infty} (q^9; q^{18})_{\infty}} (W(q) + W(\omega q) + W(\omega^2 q)), \quad (3.32)$$

where

$$W(q) = q^{-1} (q, q^2, q^2; q^3)_{\infty} \frac{(q^4; q^6)_{\infty}}{(q^5; q^6)_{\infty}} = \frac{f_1 f_2}{q f_3 f_6}. \quad (3.33)$$

Recall the cubic continued fraction studied by Ramanujan (see page 366 of [23]):

$$\nu(q) := \frac{q^{1/3}}{1 + \frac{q + q^2}{1 + \frac{q^2 + q^4}{1 + \dots}}}. \quad (3.34)$$

Chan [11] proved that

$$\frac{1}{\nu(q^3)} - 1 - 2\nu(q^3) = \frac{(q; q)_{\infty} (q^2; q^2)_{\infty}}{q(q^9; q^9)_{\infty} (q^{18}; q^{18})_{\infty}}. \quad (3.35)$$

If we write the series expansion of $W(q)$ as $W(q) = \sum_{n=-1}^{\infty} a(n)q^n$, then by (3.32) and (3.35) we deduce that

$$\overline{R}(q^3) = -\frac{(q^3; q^3)_{\infty} (q^{12}; q^{18})_{\infty}}{(q^9; q^9)_{\infty} (q^9; q^{18})_{\infty}} \sum_{n=0}^{\infty} a(3n)q^{3n} = \frac{(q^{12}, q^{18}; q^{18})_{\infty}}{(q^9; q^{18})_{\infty} (q^6; q^6)_{\infty}}. \quad (3.36)$$

Replacing q by $q^{\frac{1}{3}}$, we obtain (1.25). □

Proof of Theorem 1.5. Let

$$F(u, v; q) := \sum_{i, j \geq 0} \frac{u^i v^j q^{\binom{i}{2} + 2\binom{i+2j}{2}}}{(q; q)_i (q^4; q^4)_j}. \quad (3.37)$$

By (1.15), (2.1) and (2.2), this double sum satisfies

$$\begin{aligned}
F(u, v; q) &= \oint \sum_{i=0}^{\infty} \frac{(-1)^i u^i z^i q^{\binom{i}{2}}}{(q; q)_i} \sum_{j=0}^{\infty} \frac{v^j z^{2j}}{(q^4; q^4)_j} \sum_{k=-\infty}^{\infty} (-1)^k q^{2\binom{k}{2}} z^{-k} \frac{dz}{2\pi i z} \\
&= \oint \frac{(uz; q)_{\infty} (q^2 z, 1/z, q^2; q^2)_{\infty}}{(vz^2; q^4)_{\infty}} \frac{dz}{2\pi i z} \\
&= \oint \frac{(uz, uqz, q^2 z, 1/z, q^2; q^2)_{\infty}}{(v^{\frac{1}{2}} z, -v^{\frac{1}{2}} z; q^2)_{\infty}} \frac{dz}{2\pi i z}.
\end{aligned} \tag{3.38}$$

Taking $(u, v) = (x, x^2)$, from (3.37) and (3.38) we have

$$\begin{aligned}
F(x, x^2; q) &= \sum_{i, j \geq 0} \frac{q^{\frac{3}{2}(i^2 - i) + 4ij + 4j^2 - 2j} x^{i+2j}}{(q; q)_i (q^4; q^4)_j} \\
&= \oint \frac{(xqz, q^2 z, 1/z, q^2; q^2)_{\infty}}{(-xz; q^2)_{\infty}} \frac{dz}{2\pi i z}.
\end{aligned} \tag{3.39}$$

By the q -binomial theorem (2.3), we deduce that

$$\begin{aligned}
F(x, x^2; q) &= \oint \sum_{n=0}^{\infty} \frac{(-q; q^2)_n (-xz)^n}{(q^2; q^2)_n} \sum_{k=-\infty}^{\infty} (-1)^k q^{k(k-1)} z^{-k} \frac{dz}{2\pi i z} \\
&= \sum_{n=0}^{\infty} \frac{q^{n^2 - n} (-q; q^2)_n x^n}{(q^2; q^2)_n}.
\end{aligned} \tag{3.40}$$

Setting $x = q$ in (3.40), we obtain by (1.5) that

$$F(q, q^2; q) = \sum_{n=0}^{\infty} \frac{q^{n^2} (-q; q^2)_n}{(q^2; q^2)_n} = \frac{1}{(q, q^4, q^7; q^8)_{\infty}}. \tag{3.41}$$

This proves (1.29).

Setting $x = q^3$ in (3.40), we obtain by (1.4) that

$$F(q^3, q^6; q) = \sum_{n=0}^{\infty} \frac{q^{n^2 + 2n} (-q; q^2)_n}{(q^2; q^2)_n} = \frac{1}{(q^3, q^4, q^5; q^8)_{\infty}}. \tag{3.42}$$

This proves (1.30).

By (3.40) and (2.7) with $(a, q) \rightarrow (-q^3, q^2)$, we see that the left side of (1.31) is

$$\begin{aligned}
F(q^2, q^4; q) + qF(q^4, q^8; q) &= \sum_{n=0}^{\infty} \frac{q^{n^2 + n} (-q; q^2)_n}{(q^2; q^2)_n} (1 + q^{2n+1}) \\
&= (1 + q) \sum_{n=0}^{\infty} \frac{q^{n(n+1)} (-q^3; q^2)_n}{(q^2; q^2)_n} \\
&= (-q; q^4)_{\infty} (-q^2; q^2)_{\infty} = \frac{1}{(q, q^5, q^6; q^8)_{\infty}}.
\end{aligned}$$

This proves (1.31).

Setting $x = q^2$ in (3.40), we obtain by (1.7) that

$$F(q^2, q^4; q) = \sum_{n=0}^{\infty} \frac{q^{n^2+n}(-q; q^2)_n}{(q^2; q^2)_n} = \frac{1}{(q^2, q^3, q^7; q^8)_{\infty}}. \quad (3.43)$$

This proves (1.32). \square

Remark 3. In the proof of (1.31) we have shown that

$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)}(-q; q^2)_{n+1}}{(q^2; q^2)_n} = \frac{1}{(q, q^5, q^6; q^8)_{\infty}}. \quad (3.44)$$

This can be compared with (1.6). They have different sum sides but the same product sides. It is easy to see that (1.6) is also a special case of (2.7) with q replaced by q^2 and $a = -q^{-1}$.

4. CONCLUDING REMARKS

So far we have seen that the integral method is quite useful in proving certain Rogers-Ramanujan type identities. It should be mentioned that this method has its limitations. When the sum side cannot be expressed in terms of a nice integral which is calculable, this method may not work.

Besides the triple sum identities including (1.11) and (1.12), Kanade and Russell [19] also discovered some conjectural quadruple sum identities. To state their conjectures, we shall define

$$F(u, v, w, t; q) := \sum_{i,j,k,\ell \geq 0} \frac{(-1)^{\ell} u^i v^j w^k t^{\ell} q^{\binom{i+2j+3k+4\ell}{2} + 2\ell^2 - 2\ell}}{(q; q)_i (q^2; q^2)_j (q^3; q^3)_k (q^4; q^4)_{\ell}}. \quad (4.1)$$

The equivalent version of the conjectures of Kanade and Russell [19, Eqs. (62)–(65)] are as follows:

$$F(q, q^3, q^6, q^8; q) = \frac{1}{(q, q^3, q^4, q^6, q^8, q^9, q^{11}; q^{12})_{\infty}}, \quad (4.2)$$

$$F(q^3, q^5, q^6, q^{12}; q) = \frac{1}{(q^3, q^4, q^5, q^6, q^7, q^8, q^9; q^{12})_{\infty}}, \quad (4.3)$$

$$F(q^2, q^3, q^5, q^8; q) = \frac{1}{(q^2, q^3, q^4, q^5, q^8, q^9, q^{11}; q^{12})_{\infty}} \quad (4.4)$$

and

$$\begin{aligned} & F(q^2, q^3, q^4, q^8; q) + qF(q^3, q^5, q^7, q^{10}; q) + q^3F(q^4, q^7, q^{14}, q^{10}; q) \\ & - q^7F(q^5, q^9, q^{18}, q^{13}; q) = \frac{1}{(q, q^3, q^4, q^7, q^8, q^9, q^{10}; q^{12})_{\infty}}. \end{aligned} \quad (4.5)$$

With the help of Maple, we searched for similar identities and find the following identities:

$$F(q, q^3, q^4, q^6; q) = \frac{1}{(q, q^3, q^4, q^6, q^7, q^{10}, q^{11}; q^{12})_{\infty}}, \quad (4.6)$$

$$F(q^2, q^5, q^5, q^{10}; q) = \frac{1}{(q^2, q^3, q^5, q^6, q^7, q^8, q^{11}; q^{12})_\infty}. \quad (4.7)$$

It turns out that they were already implicitly included as conjectures in the work of Kanade and Russell [19]. Indeed, (4.6) (resp. (4.7)) can be viewed as an analytic form of the conjectural partition identity I_5 (resp. I_6) in [18] (see also [19, Secs. 5.1.1, 5.1.2]). In fact, setting $x = 1$ in [19, Eq. (54)], we get (conjecturally) (4.7). One can treat the second line of [18, Eq. (45)] in a way similar to [19, Eqs. (52), (53)] and then get an expression similar to [19, Eq. (54)], and then we get (4.6) (conjecturally) after setting $x = 1$ in the resulting expression.

Furthermore, Kanade and Russell [19] also gave different analytical forms for the conjectural partitions identities I_5 and I_6 . After setting $x = 1$ in [19, Eqs. (47), (51)] (except for a typo there), we see that the conjectural partition identities I_5 and I_6 are equivalent to the following triple sum identities

$$\sum_{i,j,k \geq 0} \frac{q^{\binom{i+2j+3k}{2} + i + j^2 + 2j + 4k}}{(q; q)_i (q^2; q^2)_j (q^3; q^3)_k} = \frac{1}{(q, q^3, q^4, q^6, q^7, q^{10}, q^{11}; q^{12})_\infty}, \quad (4.8)$$

$$\sum_{i,j,k \geq 0} \frac{q^{\binom{i+2j+3k}{2} + 2i + j^2 + j + 4j + 5k}}{(q; q)_i (q^2; q^2)_j (q^3; q^3)_k} = \frac{1}{(q^2, q^3, q^5, q^6, q^7, q^8, q^{11}; q^{12})_\infty}. \quad (4.9)$$

They have already been proved by Bringmann, Jennings-Shaffer and Mahlburg [8]. Hence the partition identities I_5 and I_6 have been confirmed and so the conjectural identities (4.6) and (4.7) are also confirmed.

So far the conjectural identities (4.2)–(4.5) are still open. We believe that the identities (4.2)–(4.7) can all be proved via the integral method. In fact, we can use (1.15), (2.1) and (2.2) to express the sum side as

$$\begin{aligned} F(u, v, w, t; q) &= \oint \sum_{i=0}^{\infty} \frac{(-uz)^i}{(q; q)_i} \sum_{j=0}^{\infty} \frac{(vz^2)^j}{(q^2; q^2)_j} \sum_{k=0}^{\infty} \frac{(-wz^3)^k}{(q^3; q^3)_k} \sum_{\ell=0}^{\infty} \frac{q^{2\ell^2 - 2\ell} (-tz^4)^\ell}{(q^4; q^4)_\ell} \\ &\quad \times \sum_{s=-\infty}^{\infty} (-1)^s q^{\binom{s}{2}} z^{-s} \frac{dz}{2\pi iz} \\ &= \oint \frac{(tz^4; q^4)_\infty (qz, 1/z, q; q)_\infty}{(-uz; q)_\infty (vz^2; q^2)_\infty (-wz^3; q^3)_\infty} \frac{dz}{2\pi iz} \end{aligned} \quad (4.10)$$

$$= \oint \frac{(t^{\frac{1}{4}}z, -t^{\frac{1}{4}}z, it^{\frac{1}{4}}z, -it^{\frac{1}{4}}z, qz, 1/z, q; q)_\infty}{(-uz, v^{\frac{1}{2}}z, -v^{\frac{1}{2}}z, -w^{\frac{1}{3}}z, -\omega w^{\frac{1}{3}}z, -\omega^2 w^{\frac{1}{3}}z; q)_\infty} \frac{dz}{2\pi iz}. \quad (4.11)$$

It is still possible to evaluate this integral using Lemma 2.1 and hence give a way to prove the above conjectures of Kanade and Russell. We have made some progress towards these conjectures and plan to discuss this in a separate paper.

It is clear that the integral method can also be applied to discover new identities. In another paper [9], using this method, we will establish some new multi-sum Rogers-Ramanujan type identities with various indexes.

Acknowledgements. We thank Matthew Russell for pointing out the relations between (4.6)–(4.7) and the conjectural identities I_5 and I_6 in [18] and their discussions of these identities in Sections 5.1.1 and 5.1.2 in [19]. This work was supported by the National Natural Science Foundation of China (12171375).

REFERENCES

- [1] G.E. Andrews, On the General Rogers-Ramanujan Theorem. Providence, RI: Amer. Math. Soc., 1974.
- [2] G.E. Andrews, The Theory of Partitions, Addison-Wesley, 1976; Reissued Cambridge, 1998.
- [3] G.E. Andrews, Schur’s theorem, Capparelli’s conjecture and q -trinomial coefficients. In: G.E. Andrews, D.M. Bressoud, L.A. Parson (eds.) The Rademacher Legacy to Mathematics, pp. 141–154. American Mathematical Society, Providence (1994).
- [4] G.E. Andrews, Sequences in partitions, double q -series and the mock theta function $\rho_3(q)$, Algorithmic Combinatorics-Enumerative Combinatorics, Special Functions and Computer Algebra, Springer, Cham. Switzerland, (2020), pp. 25–46.
- [5] G.E. Andrews and B.C. Berndt, Ramanujan’s Lost Notebook, Part II, Springer 2009.
- [6] G.E. Andrews and A.K. Uncu, Sequences in overpartitions, arXiv:2111.15003v1.
- [7] D.M. Bressoud, A generalization of the Rogers-Ramanujan identities for all moduli, J. Combin. Theory Ser. A 27 (1979), 64–68.
- [8] K. Bringmann, C. Jennings-Shaffer and K. Mahlburg, Proofs and reductions of various conjectured partition identities of Kanade and Russell, J. Reine Angew. Math. 766 (2020), 109–135.
- [9] Z. Cao and L. Wang, Multi-sum Rogers-Ramanujan type identities, arXiv:2205.12786.
- [10] S. Capparelli, On some representations of twisted affine Lie algebras and combinatorial identities. J. Algebra 154 (1993), 335–355.
- [11] H.C. Chan, Ramanujan’s cubic continued fraction and a generalization of his “most beautiful identity”. Int. J. Number Theory 6 (2010), 673–680.
- [12] S. Chern, Private communication on March 29, 2022.
- [13] S. Chern, Asymmetric Rogers-Ramanujan type identities. I, The Andrews-Uncu Conjecture, submitted.
- [14] G. Gasper and M. Rahman, Basic Hypergeometric Series, 2nd Edition, Encyclopedia of Mathematics and Its Applications, Vol. 96, Cambridge University Press, 2004.
- [15] H. Göllnitz, Partitionen mit Differenzenbedingungen, J. Reine Angew. Math. 225 (1967) 154–190.
- [16] B. Gordon, A combinatorial generalization of the Rogers-Ramanujan identities, Amer. J. Math. 83 (1961), 393–399.
- [17] M.D. Hirschhorn, F. Garvan and J. Borwein, Cubic analogs of the Jacobian cubic theta function $\theta(z, q)$, Canad. J. Math. 45 (1993), 673–694.
- [18] S. Kanade and M.C. Russell, IdentityFinder and some new partition identities of Rogers-Ramanujan type, Exp. Math. 24 (2015), no. 4, 419–423.
- [19] S. Kanade and M.C. Russell, Staircases to analytic sum-sides for many new integer partition identities of Rogers-Ramanujan type. Electron. J. Combin. 26 (2019), 1–6.
- [20] K. Kurşungöz, Andrews-Gordon type series for Capparelli’s and Göllnitz-Gordon identities, J. Combin. Theory Ser. A 165 (2019), 117–138.
- [21] J.M. Laughlin, A.V. Sills and P. Zimmer, Rogers-Ramanujan-Slater type identities, Electron. J. Combin. 15 (2008), #DS15.
- [22] M. Rhaman, Some infinite integrals of q -Bessel functions. Ramanujan International Symposium on Analysis (Pune, 1987), 117–137, Macmillan of India, New Delhi, 1989.
- [23] S. Ramanujan, The Lost Notebook and Other Unpublished Papers. Narosa, New Delhi (1988).
- [24] H. Rosengren, Proofs of some partition identities conjectured by Kanade and Russell, Ramanujan J. <https://doi.org/10.1007/s11139-021-00389-9>

- [25] A.V. Sills, An Invitation to the Rogers-Ramanujan Identities, CRC Press (2018).
- [26] L.J. Slater, Further identities of the Rogers-Ramanujan type, Proc. Lond. Math. Soc. (2) 54 (1) (1952), 147–167.
- [27] M. Takigiku and S. Tsuchioka, A proof of conjectured partition identities of Nandi, arXiv:1910.12461, 2019.
- [28] A. Uncu and W. Zudilin, Reflecting (on) the modulo 9 Kanade-Russell (conjectural) identities, arXiv: 2106.02959v2.
- [29] E.X.W. Xia and O.X.M. Yao, Analogues of Ramanujan’s partition identities, Ramanujan J. 31 (2013), 373–396.

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN 430072, HUBEI,
PEOPLE’S REPUBLIC OF CHINA

Email address: wanglq@whu.edu.cn; mathlqwang@163.com