

Non-Lie subgroups in Lie groups over local fields of positive characteristic

Helge Glöckner¹

Abstract

By Cartan's Theorem, every closed subgroup H of a real (or p -adic) Lie group G is a Lie subgroup. For Lie groups over a local field \mathbb{K} of positive characteristic, the analogous conclusion is known to be wrong. We show more: There exists a \mathbb{K} -analytic Lie group G and a non-discrete, compact subgroup H such that, for every \mathbb{K} -analytic manifold M , every \mathbb{K} -analytic map $f: M \rightarrow G$ with $f(M) \subseteq H$ is locally constant. In particular, the set H does not admit a non-discrete \mathbb{K} -analytic manifold structure which makes the inclusion of H into G a \mathbb{K} -analytic map. We can achieve that, moreover, H does not admit a \mathbb{K} -analytic Lie group structure compatible with the topological group structure induced by G on H .

Classification: 22E20 (primary); 22E35, 22E50, 32P05 (secondary).

Key words: Lie group, local field, positive characteristic, Cartan's theorem, subgroup, Lie subgroup, submanifold, initial Lie subgroup, compatible Lie group structure

1 Introduction and statement of results

If \mathbb{K} is the field \mathbb{Q}_p of p -adic numbers or the field \mathbb{R} of real numbers, then Cartan's Theorem ensures that every closed subgroup H of a \mathbb{K} -analytic Lie group G is a submanifold and hence a Lie subgroup (see the corollary to Theorem 1 in [5, Part II, Chapter V, §9]); viz., for each $x \in H$, there exists a \mathbb{K} -analytic diffeomorphism ϕ from an open x -neighbourhood $U \subseteq G$ onto an open subset V of $E \cong \mathbb{K}^n$ such that

$$\phi(U \cap H) = V \cap F$$

for some vector subspace $F \subseteq E$. It is well known that the conclusion becomes invalid for Lie groups over the field \mathbb{C} of complex numbers or a local field \mathbb{K} of positive characteristic, i.e., a field $\mathbb{K} = \mathbb{F}((X))$ of formal Laurent

¹Supported by Deutsche Forschungsgemeinschaft, project GL 357/10-1.

series over a finite field \mathbb{F} (see [6] for basic information concerning such fields). In fact, \mathbb{R} is a closed subgroup of the 1-dimensional complex Lie group $(\mathbb{C}, +)$ which cannot be a Lie subgroup as it is neither open nor discrete. For the same reason, the closed subgroup

$$\mathbb{F}[[X^2]] := \left\{ \sum_{k=0}^{\infty} a_k X^{2k} : (a_k)_{k \in \mathbb{N}_0} \in \mathbb{F}^{\mathbb{N}_0} \right\}$$

is not a Lie subgroup of the 1-dimensional \mathbb{K} -analytic Lie group $(\mathbb{K}, +)$, if $\mathbb{K} = \mathbb{F}((X))$. Our goal is to describe subgroups of Lie groups over $\mathbb{K} = \mathbb{F}((X))$ with more special properties.

Recall that a Lie group over a locally compact field of characteristic 0 induces a Lie group structure on each subgroup (see Definition 3 in [2, Chapter III, §4.5]). In the example of $\mathbb{R} \subseteq \mathbb{C}$ just recalled, the induced structure is discrete, as observed in [2, Chapter III, §4.5].

We show that the non-discrete, compact subgroup

$$H := \left\{ \sum_{k=0}^{\infty} a_k X^{2^k} : (a_k)_{k \in \mathbb{N}_0} \in \mathbb{F}^{\mathbb{N}_0} \right\} \quad (1)$$

of $(\mathbb{F}((X)), +)$ has analogous properties. With terminology as in §2, we have:

Theorem 1.1. *Let $\mathbb{K} := \mathbb{F}((X))$ for a finite field \mathbb{F} . The subgroup H of the 1-dimensional \mathbb{K} -analytic Lie group $G := (\mathbb{K}, +)$ described in (1) has the following properties:*

- (a) *H is an initial Lie subgroup of G . The initial Lie subgroup topology on H is the discrete topology.*
- (b) *For each \mathbb{K} -analytic manifold structure on the set H making the inclusion map $H \rightarrow G$ a \mathbb{K} -analytic map, the topology on the manifold H must be discrete.*

In particular, H is not a submanifold of G .

In Section 2, we introduce certain *constancy-enforcing* subsets N of a \mathbb{K} -analytic manifold M . These can be regarded as initial submanifolds such that N as a manifold is endowed with the discrete topology (see Remark 2.5). Using this terminology, Theorem 1.1(a) requires that H be a constancy-enforcing subset of G . Part (b) follows from (a).

Note that H from (1) can be given a 1-dimensional \mathbb{K} -analytic Lie group structure making the bijection

$$\mathbb{F}[[X]] \rightarrow H, \quad \sum_{k=0}^{\infty} a_k X^k \rightarrow \sum_{k=0}^{\infty} a_k X^{2^k}$$

an isomorphism of \mathbb{K} -analytic Lie groups, where $\mathbb{F}[[X]] \subseteq \mathbb{F}((X))$ is the compact open subring of formal power series. This \mathbb{K} -analytic Lie group structure on H is compatible with the given compact topology on H , induced by G ; it does not make the inclusion map $H \rightarrow G$ a \mathbb{K} -analytic mapping, by (b).

In our second example, a manifold structure compatible with the induced topology does not exist. For $\mathbb{K} = \mathbb{F}((X))$, we consider the unit group $(\mathbb{K}^\times, \cdot)$ of non-zero field elements, which is a 1-dimensional \mathbb{K} -analytic Lie group. It is well known that \mathbb{K}^\times contains a compact open subgroup isomorphic to a countable power $(\mathbb{Z}_p)^\mathbb{N}$ of the compact group \mathbb{Z}_p of p -adic integers, where p is the characteristic of \mathbb{F} (cf. [6, Chapter II, §3, Proposition 10]). Hence \mathbb{K}^\times has a compact subgroup H which is isomorphic to \mathbb{Z}_p as a topological group. We can take H as before and $G := \mathbb{K}^\times$ in the following result.

Theorem 1.2. *Let G be a Lie group over $\mathbb{K} = \mathbb{F}((X))$ and $H \subseteq G$ be a compact subgroup such that $H \cong \mathbb{Z}_p$ as a topological group. Then H does not admit a \mathbb{K} -analytic Lie group structure compatible with the topology induced by G .*

The pathologies described in Theorem 1.1 and Theorem 1.2 can occur simultaneously, as the following example shows.

Example 1.3. Let p be a prime, \mathbb{F} be the field with p elements and $2 \leq \ell \in \mathbb{N}$ such that p and ℓ are coprime. Then the closed subgroup H generated by $1 + X^\ell$ in the multiplicative group $G := \mathbb{F}((X))^\times$ is isomorphic to \mathbb{Z}_p , whence it does not admit a Lie group structure over $\mathbb{K} := \mathbb{F}((X))$ compatible with its topology (by Theorem 1.2). Note that $H \subseteq \mathbb{F}[[X^\ell]]$. Thus H is a constancy-enforcing subset of G (and hence an initial Lie subgroup when endowed with the discrete topology), by the following theorem.

Theorem 1.4. *Let p be a prime, \mathbb{F} be the field of order p and $2 \leq \ell \in \mathbb{N}$ with p and ℓ coprime. Then $\mathbb{F}[[X^\ell]]$ is a constancy-enforcing subset of $\mathbb{F}((X))$.*

Remark 1.5. If H is as in Example 1.3, then also every subgroup $S \subseteq H$ is constancy-enforcing. Moreover, S does not admit a non-discrete \mathbb{K} -analytic

Lie group structure making the inclusion map $S \rightarrow G$ continuous. [Otherwise, S would have a non-trivial compact open subgroup K . Then K would be a non-trivial closed subgroup of $H \cong \mathbb{Z}_p$ and thus $K \cong \mathbb{Z}_p$, contrary to Theorem 1.2].

General conventions. We write $\mathbb{N} := \{1, 2, \dots\}$ and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

Acknowledgement. The author thanks C.R.E. Raja (Indian Statistical Institute, Bangalore) for questions and discussions which inspired the work. The referee's comments helped to improve the presentation.

2 Initial submanifolds and constancy-enforcing subsets

Recall that a topological field \mathbb{K} is called a *local field* if it is non-discrete and locally compact (see [6]). We fix an absolute value $|\cdot|$ on \mathbb{K} which defines its topology. For the basic theory of \mathbb{K} -analytic mappings between open subsets of finite-dimensional \mathbb{K} -vector spaces and the corresponding \mathbb{K} -analytic manifolds and Lie groups modeled on \mathbb{K}^m , we refer to [5] (cf. also [1] and [2]). The following well-known concept is useful.

Definition 2.1. Let M be a \mathbb{K} -analytic manifold over a local field \mathbb{K} . A subset $N \subseteq M$ is called an *initial submanifold* if there exists a \mathbb{K} -analytic manifold structure on N with the following property: For each \mathbb{K} -analytic manifold Z , each map $f: Z \rightarrow N$ is analytic if and only if $\iota_N \circ f: Z \rightarrow M$ is analytic, where $\iota_N: N \rightarrow M$, $x \mapsto x$ is the inclusion map.

Remark 2.2.

- (a) If $N \subseteq M$ is an initial submanifold, then $\iota_N: N \rightarrow M$ is analytic. In fact, $\text{id}_N: N \rightarrow N$ is analytic and hence also $\iota_N \circ \text{id}_N = \iota_N$.
- (b) The \mathbb{K} -analytic manifold structure on an initial submanifold is uniquely determined.

In fact, let us write N_1 and N_2 for N , endowed with \mathbb{K} -analytic manifold structures as described in Definition 2.1. Write ι_{N_j} for the inclusion map $N_j \rightarrow M$, for $j \in \{1, 2\}$. Let $f: N_1 \rightarrow N_2$ be the identity map, $x \mapsto x$. Since $\iota_{N_2} \circ f = \iota_{N_1}$ is analytic, the map f is analytic. Likewise, the analyticity of $\iota_{N_1} \circ f^{-1} = \iota_{N_2}$ implies that f^{-1} is analytic. Thus $f: N_1 \rightarrow N_2$ is an analytic diffeomorphism.

- (c) Since ι_N is \mathbb{K} -analytic and hence continuous, the topology on an initial submanifold (as a manifold) is finer than the topology induced by M . It can be properly finer.

Definition 2.3. If G is a \mathbb{K} -analytic Lie group, we call a subgroup $H \subseteq G$ an *initial Lie subgroup* if it is an initial submanifold.

If $m_H: H \times H \rightarrow H$ is the group multiplication of H and m_G the group multiplication of G , then $\iota_H \circ m_H = m_G \circ (\iota_H \times \iota_H)$ is \mathbb{K} -analytic, entailing that m_H is \mathbb{K} -analytic. Likewise, the map $H \rightarrow H$, $x \mapsto x^{-1}$ is \mathbb{K} -analytic, and thus H is a \mathbb{K} -analytic Lie group in its initial manifold structure.

We consider a related concept.

Definition 2.4. Let M be a \mathbb{K} -analytic manifold over a local field \mathbb{K} . We say that a subset $N \subseteq M$ is *constancy-enforcing* if, for each \mathbb{K} -analytic manifold Z , each \mathbb{K} -analytic map $f: Z \rightarrow M$ with image $f(Z) \subseteq N$ is locally constant.

Remark 2.5. Note that N is constancy-enforcing if and only if N is an initial submanifold of M and the topology underlying the initial submanifold structure is the discrete topology. For our ends, the shorter phrase is valuable.

Let us write $\mathbb{D} := \{z \in \mathbb{K} : |z| < 1\}$ for the open unit ball.

Lemma 2.6. A subset $N \subseteq M$ is constancy-enforcing if and only if each \mathbb{K} -analytic map $f: \mathbb{D} \rightarrow M$ with $f(\mathbb{D}) \subseteq N$ is constant on some 0-neighbourhood.

Proof. The necessity of the condition is clear. Conversely, assume that the condition is satisfied. It suffices to show that $N \cap U$ is constancy-enforcing for all U in an open cover of M . We may therefore assume that N is contained in the domain P of a \mathbb{K} -analytic diffeomorphism $\psi: P \rightarrow Q$ onto an open subset $Q \subseteq \mathbb{K}^m$. It then suffices to show that $\psi(N)$ is a constancy-enforcing subset of Q (or, equivalently, of \mathbb{K}^m). We may therefore assume that $M = \mathbb{K}^m$. Let Z be a \mathbb{K} -analytic manifold and $f: Z \rightarrow \mathbb{K}^m$ be a \mathbb{K} -analytic mapping such that $f(Z) \subseteq N$. Given $x \in Z$, there exists a \mathbb{K} -analytic diffeomorphism $\phi: U \rightarrow V$ from an open x -neighbourhood $U \subseteq Z$ onto an open subset $V \subseteq \mathbb{K}^n$, with $\phi(x) = 0$. It suffices to show that $f \circ \phi^{-1}$ is constant on some 0-neighbourhood. We may therefore assume that $Z = B_\varepsilon(0)$ is an open ball in \mathbb{K}^n for some norm $\|\cdot\|$ on \mathbb{K}^n and some $\varepsilon > 0$. After a translation in the

range, we may assume that $f(0) = 0$. We may also assume that

$$f(y) = \sum_{k=1}^{\infty} p_k(y)$$

pointwise for $y \in B_\varepsilon(0)$ with certain homogeneous polynomials $p_k: \mathbb{K}^n \rightarrow \mathbb{K}^m$ of degree k . For each $y \in B_\varepsilon(0)$, we obtain a \mathbb{K} -analytic map

$$h: \mathbb{D} \rightarrow \mathbb{K}^m, \quad z \mapsto f(z y) = \sum_{k=1}^{\infty} z^k p_k(y)$$

with $h(\mathbb{D}) \subseteq f(U) \subseteq N$. By hypothesis, h is constant on a 0-neighbourhood $V \subseteq \mathbb{D}$, and thus $h|_V = 0$. As a consequence, $p_k(y) = 0$ for all $k \in \mathbb{N}$ (see second lemma in [5, Part II, Chapter II], applied to power series in a single variable). Thus $p_k = 0$, as $y \in B_\varepsilon(0)$ was arbitrary and p_k is homogeneous. Hence $f = 0$. \square

3 Proof of Theorem 1.1

We shall deduce the theorem from the following lemma. Let $\mathbb{K} := \mathbb{F}((X))$ and $\mathbb{O} := \mathbb{F}[[X]]$. Then $\mathbb{D} := \{z \in \mathbb{K}: |z| < 1\} = X\mathbb{O}$.

Lemma 3.1 *H as in (1) is a constancy-enforcing subset of \mathbb{K} .*

Proof. By Lemma 2.6, we have to show that each \mathbb{K} -analytic map $f: \mathbb{D} \rightarrow \mathbb{K}$ with $f(\mathbb{D}) \subseteq H$ is constant on some 0-neighbourhood. After replacing f with $z \mapsto f(\lambda z)$ for a small non-zero element $\lambda \in \mathbb{D}$, we may assume that f is given globally by a convergent power series,

$$f(z) = \sum_{k=0}^{\infty} a_k z^k$$

with $a_k \in \mathbb{K}$. Then $a_0 = f(0) \in H$. Since H is a subgroup, it follows that $f(y) - a_0 \in H$ for all $y \in \mathbb{D}$. We may therefore assume without loss of generality that $a_0 = 0$. Pick $0 \neq w \in \mathbb{D}$. Since the series for $f(w)$ converges, the summands form a bounded sequence; thus

$$M := \sup\{|a_k| |w|^k: k \in \mathbb{N}\} < \infty.$$

Choose $\theta \in \mathbb{D}$ such that $0 < |\theta| \leq \frac{1}{M+1}$. Then $|\theta^k| \leq \frac{1}{M+1}$ for all $k \in \mathbb{N}$ and thus $|a_k(w\theta)^k| \leq 1$ for all $k \in \mathbb{N}$. After replacing f with $z \mapsto f(zw\theta)$, we may therefore assume that the coefficients of f satisfy

$$|a_k| \leq 1 \quad \text{for all } k \in \mathbb{N}.$$

We show that $f = 0$. If $f \neq 0$, then there exists a minimal $\ell \in \mathbb{N}$ such that $a_\ell \neq 0$; we shall derive a contradiction. We can write

$$a_\ell = bX^m$$

with $m \in \mathbb{N}_0$ and $b \in \mathbb{K}$ such that $|b| = 1$. For all $n \in \mathbb{N}$ such that $n \geq m+1$,

$$f(X^n) = a_\ell X^{\ell n} + R_n = bX^{\ell n+m} + R_n$$

holds with

$$R_n = \sum_{k=\ell+1}^{\infty} a_k X^{kn} \in X^{\ell n+m+1}\mathbb{O},$$

since $a_k \in \mathbb{O}$ and $kn \geq (\ell+1)n = \ell n + n \geq \ell n + m + 1$. Thus $f(X^n) \in H$ necessitates

$$\ell n + m \in \{2^j : j \in \mathbb{N}_0\},$$

for all $n \geq m+1$. This is absurd. \square

Proof of Theorem 1.1. (a) Let Z be a \mathbb{K} -analytic manifold and $f: Z \rightarrow H$ be a map. If $\iota_H \circ f$ is \mathbb{K} -analytic, then f is locally constant, since H is constancy-enforcing (by Lemma 3.1). Hence f is \mathbb{K} -analytic as a map to H , endowed with the discrete topology (and corresponding \mathbb{K} -analytic manifold structure modeled on \mathbb{K}^0). If, conversely, f is \mathbb{K} -analytic as a map to H with the discrete topology, then f is locally constant, whence also $\iota_H \circ f$ is locally constant and hence \mathbb{K} -analytic.

(b) Endow H with a \mathbb{K} -analytic manifold structure which turns ι_H into a \mathbb{K} -analytic map. Since H is constancy-enforcing, ι_H is locally constant. Since ι_H is injective, we deduce that H is discrete. \square

4 Proof of Theorem 1.2

The point is that \mathbb{Z}_p cannot be endowed with a \mathbb{K} -analytic Lie group structure compatible with its natural topology, for $\mathbb{K} = \mathbb{F}((X))$. This can be deduced from more elaborate general arguments (see [3, Theorem 13.23] or further developments in [4]). For the reader's convenience, we provide a direct proof. Suppose that \mathbb{Z}_p admits such a manifold structure, modeled on \mathbb{K}^n with $n \in \mathbb{N}$; we shall derive a contradiction. Write H for \mathbb{Z}_p with this structure. Let q be the order of the finite field \mathbb{F} ; thus $q = p^m$ for some $m \in \mathbb{N}$. Choose the absolute value $|\cdot|$ on \mathbb{K} such that $|X| = 1/q$. Let $\|\cdot\|$ be the maximum norm on \mathbb{K}^n given by $\|(z_1, \dots, z_n)\| := \max\{|z_1|, \dots, |z_n|\}$ and define

$$B_\varepsilon := \{z \in \mathbb{K}^n : \|z\| < \varepsilon\}, \quad \overline{B}_\varepsilon := \{z \in \mathbb{K}^n : \|z\| \leq \varepsilon\}$$

for $\varepsilon > 0$. As $\mathbb{D} = X\mathbb{O}$, we have $B_1 = X\mathbb{O}^n = X\overline{B}_1$. More generally,

$$B_{q^{-k}} = \overline{B}_{q^{-k-1}} = X^{k+1}\overline{B}_1 \quad \text{for each integer } k.$$

Considering balls as compact, open subgroups of $(\mathbb{K}^n, +)$, we obtain the index

$$[\overline{B}_1 : X\overline{B}_1] = [\mathbb{O} : X\mathbb{O}]^n = q^n.$$

Since $\mathbb{K}^n \rightarrow \mathbb{K}^n$, $y \mapsto X^k y$ is an automorphism of $(\mathbb{K}^n, +)$, we deduce that $[\overline{B}_{q^{-k}} : \overline{B}_{q^{-k-1}}] = [X^k \overline{B}_1 : X^k(X\overline{B}_1)] = q^n$ for each integer k and thus

$$[\overline{B}_{q^{-k}} : \overline{B}_{q^{-\ell}}] = (q^n)^{\ell-k} \quad \text{for all integers } \ell \geq k.$$

By [5, Part II, Chapter IV, §8, Theorem], H has an open subgroup U which is a standard group. Notably, there exists $r \in]0, 1]$ and an isomorphism $\phi: U \rightarrow B_r$ of \mathbb{K} -analytic Lie groups for a group multiplication $B_r \times B_r \rightarrow B_r$, $(x, y) \mapsto x * y$ on B_r with neutral element 0, such that B_ε is a normal subgroup of B_r for all $\varepsilon \in]0, r]$ (cf. §9 in the book chapter just cited). Moreover, the cosets of B_ε in $(B_r, *)$ and $(B_r, +)$ coincide, for all $\varepsilon \in]0, r]$ (see Theorem 1 in [5, Part II, Chapter IV, §9]). After shrinking r if necessary, we may assume that the components $f_1, \dots, f_n: B_r \rightarrow \mathbb{K}$ of the p th power map

$$f: B_r \rightarrow B_r, \quad z \mapsto z^p = z * \dots * z$$

in $(B_r, *)$ are given by power series

$$f_j(z) = \sum_{\alpha \in (\mathbb{N}_0)^n} a_{\alpha,j} z^\alpha$$

on B_r for all $j \in \{1, \dots, n\}$, with $\sum_{\alpha \in (\mathbb{N}_0)^n} |a_{\alpha,j}| r^{|\alpha|} < \infty$. We use multi-index notation here, as in [5]; notably, $|\alpha| := \alpha_1 + \dots + \alpha_n$ for $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{N}_0)^n$, $z^\alpha := z_1^{\alpha_1} \dots z_n^{\alpha_n} \in \mathbb{K}$ for $z = (z_1, \dots, z_n) \in \mathbb{K}^n$, and $a_{\alpha,j} \in \mathbb{K}$. Setting $a_\alpha := (a_{\alpha,1}, \dots, a_{\alpha,n}) \in \mathbb{K}^n$, we have $M := \sum_{\alpha \in (\mathbb{N}_0)^n} \|a_\alpha\| r^{|\alpha|} < \infty$ and

$$f(z) = \sum_{\alpha \in (\mathbb{N}_0)^n} a_\alpha z^\alpha \quad \text{in } \mathbb{K}^n \text{ for all } z \in B_r,$$

as the limit of the net of finite partial sums. Then $a_0 = 0$ as $f(0) = 0^p = 0$ and $a_\alpha = 0$ for all multi-indices α of length $|\alpha| = 1$. In fact, the power map f has derivative $Df(0) = p \text{id}_{\mathbb{K}^n} = 0 \in \text{End}_{\mathbb{K}}(\mathbb{K}^n)$ at 0 since \mathbb{K} has characteristic p (see [5, Part II, Chapter IV, §7]). For all $z \in B_r$ and $\alpha \in (\mathbb{N}_0)^n$ with $|\alpha| \geq 2$,

$$|z^\alpha| \leq (\|z\|)^{|\alpha|} = \left(\frac{\|z\|}{r}\right)^{|\alpha|} r^{|\alpha|} \leq \left(\frac{\|z\|}{r}\right)^2 r^{|\alpha|}$$

holds, whence

$$\|f(z)\| \leq \sum_{|\alpha| \geq 2} \|a_\alpha\| |z^\alpha| \leq \left(\frac{\|z\|}{r}\right)^2 \sum_{|\alpha| \geq 2} \|a_\alpha\| r^{|\alpha|} = C \|z\|^2$$

with $C := M/r^2$. Thus

$$f(B_\varepsilon) \subseteq B_{C\varepsilon^2}$$

for all $\varepsilon \in]0, r]$. Notably, $f(B_\varepsilon) \subseteq B_\varepsilon$ if $\varepsilon \in]0, r]$ such that $C\varepsilon \leq 1$. Choose $\ell \in \mathbb{N}$ so large that $Cq^{-\ell} \leq q^{-2}$ and $q^{-\ell} \leq r$. Then $C(q^{-\ell})^2 = Cq^{-\ell}q^{-\ell} \leq q^{-\ell-2}$ and thus, choosing $\varepsilon := q^{-\ell}$,

$$f(\overline{B}_{q^{-\ell-1}}) = f(B_{q^{-\ell}}) \subseteq B_{C(q^{-\ell})^2} \subseteq B_{q^{-\ell-2}} = \overline{B}_{q^{-\ell-3}},$$

whence

$$[\overline{B}_{q^{-\ell-1}} : f(\overline{B}_{q^{-\ell-1}})] \geq [\overline{B}_{q^{-\ell-1}} : \overline{B}_{q^{-\ell-3}}] = (q^n)^2.$$

But $W := \phi^{-1}(\overline{B}_{q^{-\ell-1}})$ is a closed subgroup of \mathbb{Z}_p and hence of the form $p^k \mathbb{Z}_p$ for some $k \in \mathbb{N}_0$. Since ϕ is a homomorphism of groups and f the p th power map, we have $\phi(pW) = f(\phi(W))$. Thus

$$p = [p^k \mathbb{Z}_p : p^{k+1} \mathbb{Z}_p] = [W, pW] = [\phi(W), f(\phi(W))] \geq (q^n)^2 \geq p^2,$$

contradiction. □

5 Proof of Theorem 1.4

Give $\mathbb{K} := \mathbb{F}((X))$ the absolute value taking X to $1/p$. By Lemma 2.6, we have to show that each \mathbb{K} -analytic map $f: \mathbb{D} \rightarrow \mathbb{K}$ with $f(\mathbb{D}) \subseteq H := \mathbb{F}[[X^\ell]]$ is constant on some 0-neighbourhood. As in the proof of Lemma 3.1, we may assume that $f(0) = 0$ and that, for suitable $a_k \in \mathbb{K}$ with $|a_k| \leq 1$,

$$(\forall z \in \mathbb{D}) \quad f(z) = \sum_{k=1}^{\infty} a_k z^k.$$

We show that $f = 0$. If $f \neq 0$, we get a contradiction: First, note that $f'(z) = 0$ for all $z \in \mathbb{D}$. In fact, if $f'(z_0) \neq 0$, the inverse function theorem would show that $f(\mathbb{D})$ is a neighbourhood of $f(z_0)$ in \mathbb{K} ; this is absurd, as $f(\mathbb{D})$ is a subset of H and $H = \mathbb{F}[[X^\ell]]$ has empty interior in \mathbb{K} . Hence

$$(\forall z \in \mathbb{D}) \quad \sum_{k=1}^{\infty} k a_k z^{k-1} = 0,$$

whence $a_k = 0$ for all $k \in \mathbb{N}$ such that $k \notin p\mathbb{N}$ (see the second lemma in [5, Part II, Chapter II]). As we suppose that $f \neq 0$, some coefficient of f is non-zero. Hence, there exists a maximal $n \in \mathbb{N}$ such that $a_k = 0$ for all $k \in \mathbb{N}$ such that $k \notin p^n\mathbb{N}$. Abbreviate $q := p^n$. By maximality of n , there exists $m \in \mathbb{N}$ with $a_m \neq 0$ such that $m \in q\mathbb{N}$ but $m \notin pq\mathbb{N}$ and thus $m = q\mu$ with $\mu \in \mathbb{N} \setminus p\mathbb{N}$. By the preceding,

$$(\forall z \in \mathbb{D}) \quad f(z) = \sum_{k=1}^{\infty} a_{qk} z^{qk}.$$

We now consider \mathbb{K} as a subfield of $\mathbb{L} := \mathbb{F}((Y)) \cong \mathbb{K}$ using the embedding

$$\sum_{j=j_0}^{\infty} c_j X^j \mapsto \sum_{j=j_0}^{\infty} c_j Y^{qj}$$

for $j_0 \in \mathbb{Z}$, $c_j \in \mathbb{F}$. Thus $\mathbb{K} = (\mathbb{L})^q$ is the image of \mathbb{L} under the q th power of the Frobenius endomorphism of \mathbb{L} . We give \mathbb{L} the absolute value (also denoted $|\cdot|$) which extends the given absolute value $|\cdot|$ on \mathbb{K} ; thus $|Y| = \sqrt[q]{1/p}$. For each $k \in \mathbb{N}$, there exists $b_k \in \mathbb{L}$ such that $(b_k)^q = a_{qk}$. Then $b_\mu \neq 0$ and

$$g: W \rightarrow \mathbb{L}, \quad w \mapsto \sum_{k=1}^{\infty} b_k w^k$$

is an \mathbb{L} -analytic function on $W := \{w \in \mathbb{L} : |w| < 1\}$ as all $|b_k| \leq 1$. Now

$$f(z) = (g(z))^q \quad \text{for all } z \in \mathbb{D},$$

by construction. Since $\mu b_\mu \neq 0$ holds in \mathbb{L} , the function $g': W \rightarrow \mathbb{L}$, $w \mapsto \sum_{k=1}^{\infty} k b_k w^{k-1}$ does not vanish identically. Let $\nu \in \mathbb{N}_0$ be minimal such that $(\nu + 1)b_{\nu+1} \neq 0$; then

$$g'(w) = w^\nu h(w)$$

with $h(w) := \sum_{k=\nu+1}^{\infty} k b_k w^{k-1-\nu}$ and $h(0) \neq 0$. There exists $r \in]0, 1]$ such that $h(w) \neq 0$ for all $w \in \mathbb{L}$ such that $|w| < r$. There exists $z_0 \in \mathbb{K} \setminus \{0\}$ such that $|z_0| < r$; then $g'(z_0) = (z_0)^\nu h(z_0) \neq 0$. Now

$$g(w) = g(z_0) + g'(z_0)(w - z_0) + (w - z_0)R(w)$$

for a continuous function $R: W \rightarrow \mathbb{L}$ such that $R(z_0) = 0$. Since $g'(z_0) = \sum_{k=1}^{\infty} k b_k (z_0)^{k-1}$ with $|k b_k| \leq 1$, we have $|g'(z_0)| \leq 1$. Thus, we can write $g'(z_0) = aY^\tau$ with $\tau \in \mathbb{N}_0$ and $a \in \mathbb{L}$ such that $|a| = 1$. Then $a = b + c$ with $b \in \mathbb{F}^\times$ and $c \in Y\mathbb{F}[[Y]]$. There is $s \in]0, 1]$ such that $|R(w)| < |g'(z_0)|$ and thus

$$g'(z_0) + R(w) = bY^\tau + \text{higher order terms}$$

for all $w \in W$ such that $|w - z_0| < s$. We choose $j \in \mathbb{N}$ so large that $(1/p)^j < s$. Then $|X^j| < s$ and $|X^{j+1}| < s$. Moreover,

$$H \ni f(z_0 + X^j) - f(z_0) = (g(z_0 + X^j) - g(z_0))^q = b^q X^{\tau+qj} + \text{higher order terms},$$

whence $\tau + qj \in \ell\mathbb{N}$. Likewise, $f(z_0 + X^{j+1}) - f(z_0) \in H$ implies that $\tau + q(j+1) \in \ell\mathbb{N}$. Hence $q \in \ell\mathbb{N}$, which is absurd. \square

References

- [1] N. Bourbaki, “Variétés différentielles et analytiques”, Fascicule de résultats, Paragraphes 1 à 7; Hermann, Paris, 1967.
- [2] N. Bourbaki, “Lie Groups and Lie Algebras”, Chapters 1–3, Springer, Berlin, 1989.
- [3] J. Dixon, M. du Sautoy, A. Mann and D. Segal, “Analytic Pro- p Groups”, Cambridge University Press, Cambridge, ²1999.
- [4] A. Jaikin-Zapirain and B. Klopsch, *Analytic groups over general pro- p domains*, J. Lond. Math. Soc. **76** (2007), 365–383.
- [5] J.-P. Serre, “Lie Algebras and Lie Groups”, Springer, Berlin, 1992.
- [6] A. Weil, “Basic Number Theory”, Springer, Berlin, ³1974.

Helge Glöckner, Universität Paderborn, Institut für Mathematik,
Warburger Str. 100, 33098 Paderborn, Germany; glockner@math.upb.de