

Scattering for the Equivariant $U(1)$ Problem

Nishanth Gudapati

Abstract

Extending our previous works on the Cauchy problem for the $2+1$ equivariant Einstein-wave map system, we prove that the linear part dominates the nonlinear part of the wave maps equation coupled to the full set of the Einstein equations, for small data. A key ingredient in the proof is a nonlinear Morawetz estimate for the fully coupled equivariant Einstein-wave maps. The $2 + 1$ dimensional Einstein-wave map system occurs naturally in the $3 + 1$ vacuum Einstein equations of general relativity.

Keywords. Critical Wave Maps, $U(1)$ Einstein equations

1 $2+1$ Equivariant Einstein-Wave Map System

The aim of this work is to complete the task undertaken in [8], on the scattering of the Cauchy problem of $2 + 1$ equivariant wave maps coupled to Einstein's equations for general relativity. Let (M, g) be a $2+1$ dimensional, globally hyperbolic, regular, Lorentzian spacetime what is foliated by 2 dimensional Riemannian hypersurfaces (Σ, q) such that the group $SO(2)$ acts through isometries with a nonempty fixed point set Γ . The fixed point set Γ forms a timelike boundary of the quotient $M/SO(2)$. In this setting, suppose (M, g) can be represented in double null coordinates (ξ, η, θ) as

N. Gudapati: Department of Mathematics, Clark University, 950 Main Street, Worcester, MA 01610; e-mail: ngudapati@clarku.edu

Mathematics Subject Classification (2010): Primary 83C40, Secondary 53Z05

$$g = -e^{2Z} d\xi d\eta + r^2 d\theta^2 \quad (1.1)$$

where $Z = Z(\xi, \eta)$ and $r = r(\xi, \eta)$ are functions only of (ξ, η) , $\theta \in [0, 2\pi)$ is the parameter corresponding to $SO(2)$ action on (Σ, q) , $r(\xi, \eta) \in [0, \infty)$ is the radius of the area of group orbits of $SO(2)$ so that the curve $\{r = 0\}$ corresponds to the points on Γ . ξ and η are the outgoing and incoming null coordinates respectively, such that

$$\xi = \eta, \quad \partial_\xi r + \frac{1}{2} = 0, \quad \partial_\eta r - \frac{1}{2} = 0, \quad \text{on } \Gamma. \quad (1.2)$$

Now consider the 2+1 dimensional equivariant Einstein-wave map system

$$\mathbf{E}_{\mu\nu} = \mathbf{T}_{\mu\nu} \quad (1.3a)$$

$$\square_{g(u)} u = \frac{k^2 f_u(u) f(u)}{r^2}, \quad (1.3b)$$

where \mathbf{E} is the Einstein tensor of (M, g)

$$\mathbf{E} := R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R_g,$$

with the Ricci tensor \mathbf{R} and the scalar curvature R_g respectively.

$$U : (M, g) \rightarrow (N, h)$$

is an equivariant wave map, where the target (N, h) is a surface of revolution such that

$$f(0) = 0, \quad f'(0) = 1 \quad \text{and} \quad \int_0^u f(s) ds \rightarrow \infty \quad \text{as} \quad u \rightarrow \infty \quad (1.4)$$

with the odd and smooth generating function f . The equivariant ansatz $U = (u, k\theta)$, where u is independent of the angular variable θ , reduces the wave maps equation

$$\square_g U^i + {}^{(h)}\Gamma_{jk}^i(U) g^{\mu\nu} \partial_\mu U^j \partial_\nu U^k = 0$$

to

$$\square_g u = \frac{k^2 f_u(u) f(u)}{r^2}, \quad (1.5)$$

as it is well known. The wave maps equation has an associated energy-momentum tensor given by

$$\mathbf{T}_{\mu\nu} := \langle \partial_\mu U, \partial_\nu U \rangle_{h(U)} - \frac{1}{2} g_{\mu\nu} \langle \partial^\sigma U, \partial_\sigma U \rangle_{h(U)}. \quad (1.6)$$

In the double null coordinate system introduced above, the Ricci tensor \mathbf{R} of (M, g) is

$$\begin{aligned} \mathbf{R}_{\xi\xi} &= r^{-1}(2Z_\xi r_\xi - r_{\xi\xi}), \\ \mathbf{R}_{\xi\eta} &= -(2Z_{\xi\eta} + r^{-1}r_{\xi\eta}), \\ \mathbf{R}_{\eta\eta} &= r^{-1}(2Z_\eta r_\eta - r_{\eta\eta}), \\ \mathbf{R}_{\theta\theta} &= 4r e^{-2Z} r_{\xi\eta}, \\ \mathbf{R}_{\xi\theta} &= 0 \text{ and} \\ \mathbf{R}_{\eta\theta} &= 0. \end{aligned}$$

The scalar curvature is

$$R_g = 8e^{-2Z} (\partial_\xi \partial_\eta Z + r^{-1} r_{\xi\eta}),$$

we have the Einstein tensor and energy-momentum tensor as

$$\begin{aligned} \mathbf{E}_{\xi\xi} &= r^{-1}(2\partial_\xi Z \partial_\xi r - \partial_\xi^2 r), \\ \mathbf{E}_{\xi\eta} &= r^{-1} \partial_\xi \partial_\eta r, \\ \mathbf{E}_{\eta\eta} &= r^{-1}(2\partial_\eta Z \partial_\eta r - \partial_\eta^2 r), \\ \mathbf{E}_{\theta\theta} &= -4r^2 e^{-2Z} \partial_\xi \partial_\eta Z, \\ \mathbf{E}_{\xi\theta} &= 0 \text{ and} \\ \mathbf{E}_{\eta\theta} &= 0 \end{aligned}$$

Furthermore,

$$\mathcal{L} = -4e^{-2Z} \partial_\xi u \partial_\eta u + \frac{f^2(u)}{r^2}$$

The components of $\mathbf{T}_{\mu\nu}$ are

$$\begin{aligned} \mathbf{T}_{\xi\xi} &= \partial_\xi u \partial_\xi u, \\ \mathbf{T}_{\eta\eta} &= \partial_\eta u \partial_\eta u, \\ \mathbf{T}_{\xi\eta} &= \frac{e^{2Z}}{4} \frac{f^2(u)}{r^2}, \\ \mathbf{T}_{\theta\theta} &= \frac{r^2}{2} e^{-2Z} \left(4\partial_\eta u \partial_\xi u + e^{2Z} \frac{f^2(u)}{r^2} \right). \end{aligned}$$

Suppose we introduce coordinates

$$T = \frac{\xi + \eta}{2}, \quad R = \frac{\xi - \eta}{2}$$

we can construct the ‘retarded time’ coordinates (η, R, θ) such that the metric g of M can be expressed as

$$g = -e^{2Z(\eta, R)} d\eta^2 - 2e^{2Z(\eta, R)} d\eta dR + r^2(\eta, R) d\theta^2. \quad (1.7)$$

Let us now calculate the relevant quantities in the retarded-time coordinates:

Ricci tensor \mathbf{R}

$$\begin{aligned} \mathbf{R}_{\eta\eta} &= r^{-1} (r \partial_R^2 Z - \partial_R r \partial_R Z + 2\partial_\eta Z \partial_\eta r - 2r \partial_{\eta R}^2 Z - \partial_\eta^2 r), \\ \mathbf{R}_{\eta R} &= r^{-1} (\partial_R Z \partial_R r + r \partial_R^2 Z - 2r \partial_{\eta R}^2 Z - \partial_{\eta R}^2 r), \\ \mathbf{R}_{RR} &= r^{-1} (2\partial_R Z \partial_R r - \partial_R^2 r), \\ \mathbf{R}_{\theta\theta} &= r e^{-2Z} (-\partial_R^2 r + 2\partial_{\eta R}^2 r), \\ \mathbf{R}_{\eta\theta} &= 0 \text{ and} \\ \mathbf{R}_{R\theta} &= 0. \end{aligned}$$

The scalar curvature can be represented as

$$R_g = -r^{-1} 2e^{-2Z} (r \partial_R^2 Z + \partial_R^2 r - 2r \partial_{\eta R}^2 Z - 2\partial_{\eta R}^2 r) \quad (1.8)$$

Consequently, the Einstein tensor \mathbf{E} is given by

$$\begin{aligned} \mathbf{E}_{\eta\eta} = & -r^{-1}(-\partial_R Z \partial_R r + \partial_R^2 r - \partial_{R\eta}^2 r) \\ & - r^{-1}(\partial_\eta r \partial_R Z + \partial_R Z \partial_\eta r - 2\partial_\eta Z \partial_\eta r - \partial_{\eta R}^2 r + \partial_\eta^2 r), \end{aligned} \quad (1.9a)$$

$$\mathbf{E}_{\eta R} = -r^{-1}(-\partial_R Z \partial_R r + \partial_R^2 r - \partial_{\eta R}^2 r), \quad (1.9b)$$

$$\mathbf{E}_{RR} = r^{-1}(2\partial_R Z \partial_R r - \partial_R^2 r), \quad (1.9c)$$

$$\mathbf{E}_{\theta\theta} = e^{-2Z} r^2 (\partial_R^2 Z - 2\partial_{\eta R}^2 Z), \quad (1.9d)$$

$$\mathbf{E}_{R\theta} = 0 \text{ and} \quad (1.9e)$$

$$\mathbf{E}_{\eta\theta} = 0. \quad (1.9f)$$

The energy-momentum tensor \mathbf{T} is given by

$$\mathbf{T}_{\mu\nu} := \langle \partial_\mu U, \partial_\nu U \rangle_h - \frac{1}{2} g_{\mu\nu} \mathcal{L}. \quad (1.10)$$

$$\begin{aligned} \mathcal{L} &= h_{ij} g^{\mu\nu} \partial_\mu U^i \partial_\nu U^j \\ &= -e^{2Z} (\partial_\eta u)^2 - 2e^{-2Z} \partial_\eta u \partial_R u + \frac{f^2(u)}{r^2} \end{aligned} \quad (1.11)$$

$$\mathbf{T}_{\eta\eta} = \frac{1}{2} \left((\partial_\eta u)^2 - 2\partial_\eta u \partial_R u + e^{2Z} \frac{f^2(u)}{r^2} \right), \quad (1.12a)$$

$$\mathbf{T}_{\eta R} = \frac{1}{2} \left(-(\partial_\eta u)^2 + \frac{f^2(u)}{r^2} \right), \quad (1.12b)$$

$$\mathbf{T}_{RR} = (\partial_R u)^2, \quad (1.12c)$$

$$\mathbf{T}_{\theta\theta} = \frac{r^2}{2} e^{-2Z} \left((\partial_\eta u)^2 + 2\partial_\eta u \partial_R u + e^{2Z} \frac{f^2(u)}{r^2} \right). \quad (1.12d)$$

If we introduce a time coordinate function $T := \eta + R$, we can represent the metric g also in terms of (T, R, θ) coordinates

$$ds^2 = e^{2Z(T,R)} (-dT^2 + dR^2) + r^2(T, R) d\theta^2.$$

The Ricci tensor in (T, R, θ) is

$$\mathbf{R}_{TT} = r^{-1} (\partial_R Z \partial_R r + r \partial_R^2 Z + \partial_T Z \partial_T r - r \partial_T^2 Z - \partial_T^2 r), \quad (1.13)$$

$$\mathbf{R}_{TR} = r^{-1} (\partial_R r \partial_T Z + \partial_R Z \partial_T r - \partial_{TR}^2 r), \quad (1.14)$$

$$\mathbf{R}_{RR} = -r^{-1} (-\partial_R Z \partial_R r + r \partial_R^2 Z + \partial_R^2 r - \partial_T Z \partial_T r - r \partial_T^2 Z), \quad (1.15)$$

$$\mathbf{R}_{\theta\theta} = r e^{-2Z} (-\partial_T^2 r + \partial_R^2 r), \quad (1.16)$$

$$\mathbf{R}_{T\theta} = 0 \text{ and} \quad (1.17)$$

$$\mathbf{R}_{R\theta} = 0. \quad (1.18)$$

The scalar curvature is given by

$$R_g = -2r^{-1} e^{-2Z} (r(-\partial_T^2 Z + \partial_R^2 Z) + (-\partial_T^2 r + \partial_R^2 r))$$

The Einstein tensor

$$\mathbf{E}_{TT} = r^{-1} (\partial_R Z \partial_R r + \partial_R^2 r - \partial_T Z \partial_T r), \quad (1.19)$$

$$\mathbf{E}_{TR} = r^{-1} (\partial_R r \partial_T Z + \partial_R Z \partial_T r - \partial_{TR}^2 r), \quad (1.20)$$

$$\mathbf{E}_{RR} = r^{-1} (\partial_R Z \partial_R r + \partial_T Z \partial_T r - \partial_T^2 r), \quad (1.21)$$

$$\mathbf{E}_{\theta\theta} = r^2 e^{-2Z} (-\partial_T^2 Z + \partial_R^2 Z), \quad (1.22)$$

$$\mathbf{E}_{T\theta} = 0 \text{ and} \quad (1.23)$$

$$\mathbf{E}_{R\theta} = 0. \quad (1.24)$$

Furthermore, we have

$$\mathcal{L} = -e^{-2Z} (\partial_T u)^2 + e^{-2Z} (\partial_R u)^2 + \frac{f^2(u)}{r^2}$$

Thus,

$$\mathbf{T}_{TT} = \frac{1}{2} \left((\partial_T u)^2 + (\partial_R u)^2 + e^{2Z} \frac{f^2(u)}{r^2} \right), \quad (1.25)$$

$$\mathbf{T}_{TR} = (\partial_T u \partial_R u), \quad (1.26)$$

$$\mathbf{T}_{RR} = \frac{1}{2} \left((\partial_T u)^2 + (\partial_R u)^2 - e^{2Z} \frac{f^2(u)}{r^2} \right), \quad (1.27)$$

$$\mathbf{T}_{\theta\theta} = \frac{1}{2} r^2 \left(-e^{-2Z} (\partial_T u)^2 + e^{-2Z} (\partial_R u)^2 + \frac{f^2(u)}{r^2} \right) \quad (1.28)$$

Initial Value Problem

We shall set up the initial value problem at a $T = 0$ Cauchy hypersurface (Σ_0, q_0) . The initial data of geometric part of the $2 + 1$ Einstein-wave map system consists of a Riemannian 2-manifold (Σ, q) and a symmetric 2-tensor k_0 . The Einstein's equations for general relativity are an overdetermined system, thus to formulate an initial value problem the following constraint equations are imposed on (Σ_0, q_0, k_0)

$$R_q + \text{tr}(k_0)^2 - \|k_0\|_q^2 = 2\mathbf{e} \quad (1.29a)$$

$$\nabla(q_0)_a (k_0)_b^a - \nabla(q_0)_b (k_0)_a^a = \mathbf{m}, \quad (1.29b)$$

where \mathbf{e} and \mathbf{m} are the energy and momentum densities respectively. It follows from a standard result that there exists a unique (upto an isometry) smooth, globally hyperbolic, future development (M, g, U) of the initial data (Σ) [4]. Subsequently, k is the second fundamental form of the embedding $\Sigma \hookrightarrow M$ and if \mathcal{N} is the unit normal of $\Sigma \hookrightarrow M$ the above system is

$$\mathbf{E}(\mathcal{N}, \mathcal{N}) = \mathbf{T}(\mathcal{N}, \mathcal{N}) \quad (1.30a)$$

$$\mathbf{E}(\mathcal{N}, e_1) = \mathbf{T}(\mathcal{N}, e_1) \quad (1.30b)$$

and

$$\mathbf{e} = \frac{1}{2} \left(\|\mathcal{N}(U)\|_h^2 + \|\mathcal{S}(U)\|_h^2 + \left\| \frac{1}{r} \partial_\theta(U) \right\|_h^2 \right) \quad (1.31a)$$

$$\mathbf{m} = \mathcal{N}(U) \cdot \mathcal{S}(U) \quad (1.31b)$$

with $\mathcal{N} := e^{-Z} \partial_T$ and $\mathcal{S} := e^{-Z} \partial_R$.

The initial data of the wave maps equation are

$$U_0 : (\Sigma, q) \rightarrow (N, h) \quad (1.32a)$$

$$U_1 : (\Sigma, q) \rightarrow T_{U_0}(N, h) \quad (1.32b)$$

Therefore the 5-tuple (Σ, q, K, U_0, U_1) that satisfies the constraint equations constitutes the initial data of the Einstein-wave map system. For our convenience, we introduce the following quantities on the initial Cauchy surface

$$u|_{\Sigma_0} = u_0, \quad \partial_T u|_{\Sigma_0} = u_1, \quad (1.33a)$$

$$Z|_{\Sigma_0} = Z_0, \quad \partial_T Z|_{\Sigma_0} = Z_1, \quad (1.33b)$$

$$r|_{\Sigma_0} = r_0, \quad \partial_T r|_{\Sigma_0} = r_1. \quad (1.33c)$$

with $r_1|_{\Gamma} = 0$ and $\partial_R r_0|_{\Gamma} = 1$.

Remark 1.1. Define a function v such that $u = Rv$, then the following results hold

1. v satisfies the 4+1 wave equation:

$$\left. \begin{array}{ll} {}^{4+1}\square v &= F(v) & \text{on } \mathbb{R}^{4+1} \\ v_0 = v(0, x) \text{ and } v_1 = \partial_T v(0, x) & & \text{on } \mathbb{R}^4 \end{array} \right\} \quad (1.34)$$

with

$$\begin{aligned} F(v) &= \left(e^{2Z} - 1 + \left(\frac{r}{R} \partial_\eta r + \frac{1}{2} \right) - \left(\frac{r}{R} \partial_\xi r - \frac{1}{2} \right) \right) \frac{v}{r^2} \\ &\quad + 2\partial_\xi v \partial_\eta \log\left(\frac{r}{R}\right) + 2\partial_\eta v \partial_\xi \log\left(\frac{r}{R}\right) + e^{2Z} \frac{R^2}{r^2} v^3 \zeta(Rv) \end{aligned} \quad (1.35)$$

2. If we define $E_0 := \int_{\Sigma_0} \mathbf{T}(\mathcal{N}, \mathcal{N}) \bar{\mu}_q = \int_{\Sigma_0} \mathbf{e} \bar{\mu}_q$, the energy of v satisfies the estimate

$$E_0 \geq \|u_0\|_{H^1(\mathbb{R}^2)}^2 + \|u_1\|_{L^2(\mathbb{R}^2)}^2 \quad (1.36)$$

$$\geq \|v_0\|_{H^1(\mathbb{R}^4)}^2 + \|v_1\|_{L^2(\mathbb{R}^4)}^2. \quad (1.37)$$

The results follow from the Hardy's inequality and assumptions on the target manifold (see [8]).

In the following, we shall define the notion of asymptotic flatness of the 2-metric (Σ, q)

Definition 1.2 (Asymptotic flatness and AV-mass). *Suppose (Σ, q, k) is a Riemannian 2-metric, then it is called asymptotically flat if*

1. *for some compact set Ω , \exists a global diffeomorphism $\Sigma \setminus \Omega \rightarrow \mathbb{R}^2 \setminus \mathbb{B}_1(0)$*
2. *The (Σ, q, k) admits the asymptotic expansion*

$$\begin{aligned} q &= \rho^{-m_{AV}} (\delta_{ab} + \mathcal{O}(\rho^{-1})) \\ k &= \mathcal{O}(\rho^{-2}), \quad \text{tr}(k) = \mathcal{O}(\rho^{m_{AV}-3}) \end{aligned}$$

and the energy momentum tensor

$$\mathbf{T}_{\mu\nu} = \mathcal{O}(\rho^{m_{AV}-3}) \quad (1.38)$$

in the asymptotic region, where δ_{ab} is the Euclidean metric and $\rho = |x|$. We define the quantity m_{AV} as the Ashtekar-Varadarajan mass of the Cauchy hypersurface (Σ, q, k) [3].

Recall the form of the 2-metric

$$q = e^{2Z} dR^2 + r^2 d\theta^2, \quad (1.39)$$

then a calculation shows that m_{AV} is given by

$$m_{AV} = 2(1 - e^{-\tilde{Z}_\infty}), \quad (\Sigma, q) \quad (1.40)$$

where

$$e^{2\tilde{Z}_\infty} := \lim_{R \rightarrow \infty} e^{2\tilde{Z}} = \lim_{R \rightarrow \infty} e^{2Z} \left(\frac{\partial r}{\partial R} \right)^{-2}, \quad \text{on } (\Sigma, q). \quad (1.41)$$

Furthermore, let us define the mass aspect function, that is closely related to the Hawking mass in 3+1 dimensional spacetimes

$$m = 1 + 4e^{-2Z} \partial_\xi r \partial_\eta r, \quad (1.42)$$

Review of Known Results

To put our results into context let first us review the results for the 2+1 equivariant Einstein-wave map system and the the literature on critical wave maps shall be discussed later. For the initial data described in the previous section, we have the following global existence result from [1], which also holds without the smallness restriction of the initial data in view of a non-concentration of energy argument [10].

Theorem 1.3 (Global regularity of equivariant Einstein-wave maps). *Let $E_0 < \epsilon^2$ for ϵ sufficiently small and let (M, g, u) be the maximal Cauchy development of an asymptotically flat, compactly supported, regular Cauchy data set for the 2 + 1 equivariant Einstein-wave map problem (1.3) with target (N, h) satisfying*

$$\int_0^s f(s') ds' \rightarrow \infty \quad \text{for } s \rightarrow \infty. \quad (1.43)$$

Then (M, g, u) is regular and causally geodesically complete.

The results in [11, 1, 8] and the current work are part of the larger program to understand the global behaviour of the 3 + 1 Einstein equations with one translational $U(1)$ symmetry. In global existence proof of [1], it was proved that, in future development of the initial data of the 2+1 equivariant Einstein-wave map system, the trapped surfaces cannot form. In particular, it was proved that

Theorem 1.4 (No trapped surfaces; Theorem 3.1 in [1]). *Suppose (M, g, U) is the regular, globally hyperbolic maximal development of the 2+1 equivariant Einstein-wave map system, define $\mathcal{Q} := M \setminus U(1)$ and*

$$\mathcal{R} := \{p \in \mathcal{Q} \quad \text{such that} \quad \partial_\xi r > 0, \partial_\eta r < 0\} \quad (1.44)$$

then,

$$1. \quad \mathcal{Q} = \mathcal{R}$$

$$2. \quad q \in \mathcal{Q},$$

$$0 \leq m(q) < m_\infty < 1 \quad (1.45)$$

where m_∞ is the spatial asymptotic limit of m on (Σ, q, k) .

Consider the system of coupled nonlinear wave equations occurring in the 2+1 Einstein-equivariant wave map system:

$$-\partial_T^2 Z + \partial_R^2 Z = -e^{2Z} \frac{f^2(u)}{8r^2} - \frac{1}{8} (-(\partial_T u)^2 + (\partial_R u)^2) \quad (1.46a)$$

$$-\partial_T^2 r + \partial_R^2 r = r\kappa(\partial_T u + \partial_R u)^2 \quad (1.46b)$$

$$\square_{g(u)} u = \frac{f(u)f_u(u)}{r^2}. \quad (1.46c)$$

In a previous work, we have established scattering for the following partially coupled Einstein-wave map equations, classified as Problems I and II below.

Problem I

Consider a function v such that

$$\left. \begin{array}{ll} {}^{4+1}\square v = F(v) & \text{on } \mathbb{R}^{4+1} \\ v_0 = v(0, x) \text{ and } v_1 = \partial_T v(0, x) & \text{on } \mathbb{R}^4 \end{array} \right\} \quad (1.47)$$

with

$$F(v) = \left(e^{2Z} - 1 + \left(\frac{r}{R} \partial_\eta r + \frac{1}{2} \right) - \left(\frac{r}{R} \partial_\xi r - \frac{1}{2} \right) \right) \frac{v}{r^2} + e^{2Z} \frac{R^2}{r^2} v^3 \zeta(Rv)$$

Problem II

Suppose the function v is such that

$$\left. \begin{array}{ll} {}^{4+1}\square v = F(v) & \text{on } \mathbb{R}^{4+1} \\ v_0 = v(0, x) \text{ and } v_1 = \partial_T v(0, x) & \text{on } \mathbb{R}^4 \end{array} \right\} \quad (1.48)$$

where

$$\begin{aligned} F(v) = & \left(\frac{1}{r} \partial_\eta r + \frac{1}{2R} \right) \partial_\xi v + \left(\frac{1}{r} \partial_\xi r - \frac{1}{2R} \right) \partial_\eta v \\ & + \left(\left(\frac{r}{R} \partial_\eta r + \frac{1}{2} \right) - \left(\frac{r}{R} \partial_\xi r - \frac{1}{2} \right) \right) \frac{v}{r^2} \\ & + \frac{R^2}{r^2} v^3 \zeta(Rv) \end{aligned} \quad (1.49)$$

and v is coupled to Einstein's equations (1.3) with $u = Rv$. We were able to prove scattering for both Problems I and II:

Theorem 1.5 (Theorem 2.11 and Theorem 3.2 in [8]). *Suppose v is a global solution of the partially linearized wave maps equations (1.47) or (1.48), with the energy $E_0 < \epsilon^2$, then v scatters forward and backward in time*

The coupling of the wave maps equation with the Einstein's equations introduces nonlocality into the problem, which causes significant difficulties in closing the estimates at critical regularity. It may be noted that the equations in Problem I and II are partially linearized approximations of the equation (1.46c), where they correspond to the linearization of the equation (1.46b) and equation (1.46a) respectively, which in turn correspond to partial decoupling of the wave maps equation (1.46c). The methods employed in the wave maps equation are based on a optimal control of the Morawetz and Strichartz estimates that allow us to control the bootstrap estimates at critical regularity. The nonlinear terms in the wave map equations in Problem I and II involve all the terms of the fully coupled wave maps equation, but in view of the nonlinear theory the equivalent results carry forward. In this work we are able to prove scattering for the fully coupled wave maps equation:

$$\left. \begin{array}{ll} {}^{4+1}\square v &= F(v) \quad \text{on } \mathbb{R}^{4+1} \\ v_0 = v(0, x) \quad \text{and} \quad v_1 = \partial_T v(0, x) &\quad \text{on } \mathbb{R}^4 \end{array} \right\} \quad (1.50)$$

$$\begin{aligned} F(v) = & \left(e^{2Z} - 1 + \left(\frac{r}{R} \partial_\eta r + \frac{1}{2} \right) - \left(\frac{r}{R} \partial_\xi r - \frac{1}{2} \right) \right) \frac{v}{r^2} \\ & + 2\partial_\xi v \partial_\eta \log \frac{r}{R} + 2\partial_\eta v \partial_\xi \log \left(\frac{r}{R} \right) + e^{2Z} \frac{R^2}{r^2} v^3 \zeta(Rv) \end{aligned} \quad (1.51)$$

Theorem 1.6. *Suppose $E_0 < \epsilon^2$, for ϵ sufficiently small and (M, g, U) is a global solution of the 2+1 equivariant Einstein-wave map system, then v scatters backward and forward in time in the energy topology*

In the proof of the above scattering theorem, we use the following nonlinear Morawetz estimate:

Theorem 1.7 (Nonlinear Morawetz estimate). *Suppose (M, g, U) is a globally*

regular solution of the 2+1 Einstein-equivariant wave map system, then

$$\int_{\mathbb{R}^{4+1}} \frac{v^2}{R^3} \bar{\mu}_{\check{g}} \leq \|v_0\|_{\dot{H}^1(\mathbb{R}^4)}^2 + \|v_1\|_{L^2(\mathbb{R}^4)}^2. \quad (1.52)$$

where $\bar{\mu}_{\check{g}}$ is the volume form of the 4 + 1 Minkowski space $(\mathbb{R}^{4+1}, \check{g})$, with

$$\check{g} = -dT^2 + dR^2 + R^2 d\omega_{\mathbb{S}^3}. \quad (1.53)$$

Subsequently, using the scattering result for v , we obtain scattering for the fully coupled equivariant wave maps:

Theorem 1.8 (Temporal Scattering). *Let $E_0 < \epsilon^2$ for ϵ sufficiently small and let (M, g, u) be the maximal geodesically complete Cauchy development of an asymptotically flat, compactly supported, regular Cauchy data set for the 2 + 1 equivariant Einstein-wave map problem (1.3) with target (N, h) satisfying*

$$\int_0^s f(s') ds' \rightarrow \infty \quad \text{when } s \rightarrow \infty. \quad (1.54)$$

Then the maximal development (M, g, U) is such that U converges to a linearized system of (1.3b) as $T \rightarrow \infty$ along future pointing the timelike curves.

We would like to point out that, both conceptually and technically, the wave map fields (and their conjugate momenta) is the dominant field that represents the true dynamical degrees of freedom of the 2 + 1 dimensional Einstein-wave map system.

The key ingredient in the proof is the non-linear Morawetz estimate (1.52) for the fully coupled Einstein-wave map system. An analogous result holds for the partially coupled equivariant Einstein-wave system (1.48) (Problem II), but (1.52) now holds for the fully coupled system. Likewise, a similar result holds for scattering backwards in time. It may be noted that a canonical notion of energy for the Einstein equations is defined relative to a chosen reference background. However, in view of the Remark 1.1, in this work we use a weaker notion of mass-energy that nevertheless gives scattering and sharp asymptotics relative to the same reference background.

In view of the fact that the Jacobians of the transition functions between the (T, R, θ) , double-null and retarded-time coordinates are constants, the scattering result of Theorem 1.8 can be adapted to establish equivalent results along the $\eta = \text{const.}$ in the asymptotic region.

Corollary 1.9. *Suppose (M, g, U) is the regular, globally hyperbolic, causally geodesically complete maximal development of the asymptotically flat initial data (Σ, q, k, U) with u compactly supported, then u admits the decay rate*

$$|u| \leq c(1 + \xi)^{-1/2}(1 + \eta)^{-1/2} \quad (1.55)$$

in the temporal asymptotic region $(T \rightarrow \infty)$ and

$$u = \mathcal{O}(R^{-1/2}) \quad (1.56)$$

as $R \rightarrow \infty$ in the $\eta = \text{const.}$ hypersurfaces.

Let us now discuss a few salient points about the 2+1 dimensional equivariant Einstein-wave map system.

Metric functions

The wave map U and its conjugate momentum form the dynamical degrees of freedom of the system and drive the dynamics of the whole system. However, in our quest to understand the asymptotic geometry of the maximal future development (M, g, U) of the system, the form of the Einstein tensor poses obstacles. For instance, consider the equation

$$\left. \begin{array}{ll} {}^{1+1}\square w &= e^x \\ w(0, x) = 0 \quad \text{and} \quad \partial_T w(0, x) = 0 & \end{array} \right\} \begin{array}{l} \text{on } \mathbb{R}^{1+1} \\ \text{on } \mathbb{R} \end{array} \quad (1.57)$$

The energy of w in the equation (1.57) goes out of control asymptotically as $T \rightarrow \infty$ even though we start with ‘small’ initial data. Thus we are motivated to analyze the asymptotic behaviour separately for metric functions to rule out such behaviour. As per our analysis, we expect that, as the maximal development (M, g, U) attains its scattering state near temporal infinity, it converges to the Minkowski space. In other words, as the maximal development attains its

scattering state near temporal infinity, the geometric mass (Ashtekar-Varadarajan) converges to zero. This topic, with the together the behaviour of the curvature, merits a dedicated discussion and shall be addressed in a separate work.

Sharp Asymptotics

We would like to remark that the scattering methods have an advantage in the characterization of the asymptotic geometry of the initial value problem of the system as the decay rate so obtained is sharp compared to the energy methods in 2+1 dimensions:

1. The asymptotic decay rate obtained is sharp

$$|u| \leq c(1 + \xi)^{-1/2}(1 + \eta)^{-1/2} \quad \text{in } \mathbb{R}^{2+1}, \quad (1.58)$$

in view of the fact that the geometric mass and spatial asymptotics are conserved globally, including the region where the system (M, g, U) attains its scattering state.

The small data problem and the target

We would also like to remark that our scattering result reaffirms the intuition the curvature of the target manifold (N, h) does not play a decisive role in the scattering in the small data context.

In the above, we mentioned several results for the 2+1 equivariant Einstein-wave system which hold in the asymptotic limits. In the context of Einstein's equations, an important aim of the initial value formulation is the characterization of the asymptotic geometry of the future development of the system. In a classic 1964 work, Penrose had formulated a notion of conformal infinity in the context of Einstein's equations for general relativity. From a PDE perspective, the work of Penrose gave a crucial answer for the question of what it means for solutions of the Einstein equations to exist 'globally'. Following the definition of Penrose, the conformal infinity of in our context is defined as follows:

Definition 1.10. *A 2+1 Lorentzian manifold (M, g) admits a conformal null infinity if \exists a 2+1 Lorentzian manifold $(\mathcal{M}, \mathbf{g})$ with boundary \mathcal{I} such that*

- \exists a diffeomorphism $\mathcal{M} \setminus \mathcal{I} \rightarrow M$
- \exists a smooth function Ω on $(\mathcal{M}, \mathbf{g})$, $\mathbf{g} = \Omega^2 g$ in the interior of \mathcal{M} such that $\Omega = 0$ on \mathcal{I} and $\nabla_\nu \Omega \neq 0$ at \mathcal{I}

Minkowski Spacetime \mathbb{R}^{2+1}

In our problem, the Minkowski spacetime \mathbb{R}^{2+1} plays an important role as an example of a spacetime that admits a conformal completion. Consider the Minkowski metric

$$ds^2 = -d\eta^2 - d\eta dR + R^2 d\theta^2 \quad (1.59)$$

in the retarded-time coordinates (η, R, θ) . Suppose we introduce the function $\rho = \frac{1}{r}$ and define the unphysical metric with $\Omega = \rho$ then \mathbb{R}^{2+1} admits a conformal infinity at $\rho = 0$. In this work we shall be interested in the geometry and behaviour of fields near the future null infinity, so this choice of the conformal factor to construct null infinity suffices. Likewise, the 4+1 dimensional Minkowski space admits a conformal null infinity similarly.

Wave maps from \mathbb{R}^{2+1} with symmetry have been studied in a series of landmark papers of Christodoulou-Tahvildar-Zadeh and Shatah [7, 6, 18, 17], where global regularity and asymptotics of initial value problems of equivariant and spherically symmetric wave maps were studied (see also [9]).

In [12] the Einstein-scalar wave system without symmetry was studied and stability of Minkowski space in exponential time (\sqrt{t}^{-1}) was established. The techniques in [12] were inspired from [16] which provided a simpler proof of the seminal and classic work on stability of Minkowski space [5]. The asymptotic behaviour of perturbations of the 3+1 Minkowski space in harmonic coordinates is studied in [15].

In a pioneering and influential series of works [21], global behaviour of wave maps in multiple dimensions and regularity levels has been studied. Local and global well-posedness of wave maps at various regularity levels are studied in [23, 22, 24]. Global regularity for small data for critical wave maps $U : \mathbb{R}^{2+1} \rightarrow$

\mathbb{H}^2 was proved in [13]. In a voluminous work [14], the important question of concentration compactness for wave maps $U : \mathbb{R}^{2+1} \rightarrow \mathbb{H}^2$ was thoroughly settled. Likewise, scattering for large data for critical wave maps with more general targets was settled in [19][20].

Notation

A wave operator ${}^{n+1}\square$ without a metric subscript refers to a flat-space wave operator in $n + 1$ dimensions. We shall use the Einstein summation convention throughout.

2 Asymptotics of a partially linearized Equivariant wave map system

As we already remarked, in a previous work we have proved the scattering for the following problem

$$\left. \begin{array}{ll} {}^{4+1}\square v &= F(v) \quad \text{on } \mathbb{R}^{4+1} \\ v_0 = v(0, x) \quad \text{and} \quad v_1 = \partial_T v(0, x) &\quad \text{on } \mathbb{R}^4 \end{array} \right\} \quad (2.1)$$

with

$$F(v) = \left(e^{2Z} - 1 + \left(\frac{r}{R} \partial_\eta r + \frac{1}{2} \right) - \left(\frac{r}{R} \partial_\xi r - \frac{1}{2} \right) \right) \frac{v}{r^2} + e^{2Z} \frac{R^2}{r^2} v^3 \zeta(Rv)$$

and coupled to the Einstein's equations (1.3a) with $u = Rv$. If we consider the special case of $r \equiv R$, we have the partial decoupling of system, resulting in the partially linearized wave maps equation

$${}^{4+1}\square v = (e^{2Z} - 1) \frac{v^2}{R^2} + e^{2Z} \zeta(rv) v^3 \quad (2.2)$$

It follows from the scattering theorem that for data that is sufficiently decaying at ι^0 , $\|u\|_{L^\infty} = \mathcal{O}(T^{-1})$ in the temporal asymptotic region. Let us now consider the partially linearized system in the Bondi ('retarded time') coordinates (η, R, θ)

$$g = -e^{2Z(\eta, R)} d\eta^2 - 2e^{2Z(\eta, R)} d\eta dR + R^2 d\theta^2 \quad (2.3)$$

with the usual behaviour on the axis and $\theta \in [0, 2\pi)$:

$$Z = 0, \quad \partial_R Z = 0, \quad \text{on } \Gamma.$$

We have the following equations from (1.9) and (1.12),

$$R^{-1} \partial_\eta Z = \frac{1}{2} \left((\partial_\eta u)^2 - 2 \partial_\eta u \partial_R u + e^{2Z} \frac{f^2(u)}{r^2} \right) \quad (2.4)$$

$$R^{-1} \partial_R Z = \frac{1}{2} (\partial_R u)^2 \quad (2.5)$$

Proposition 2.1. *Suppose the partially linearized wave maps equation admits the asymptotic expansions ([2], also compare with the rate (1.58))*

$$u(\eta, R) = f_0(\eta) R^{-1/2} + \sum_{n=1}^{\infty} f_n(\eta) R^n \quad (2.6)$$

$$\partial_\eta u(\eta, R) = \dot{f}_0(\eta) R^{-1/2} + \sum_{n=1}^{\infty} \dot{f}_1 R^n \quad (2.7)$$

$$\partial_R u(\eta, R) = -f'_0(\eta) R^{-3/2} + \sum_{n=1}^{\infty} f'_1(\eta) R^{n-1} \quad (2.8)$$

in the asymptotic region ($R \rightarrow \infty$) of $\eta = \text{const.}$ hypersurfaces then

1. $Z = \mathcal{O}(1)$, as $R \rightarrow \infty$ along $\eta = \text{const.}$ hypersurfaces
2. $Z \rightarrow 0$ asymptotically along the timelike curves inside the future null cones

Proof. 1. follows directly from the integration of (2.4).

2. Consider a curve with the parameter τ such that $\tau = \eta + (1 - \lambda)R$, for $\lambda > 1$, we have

$$\partial_\tau Z = R \left((\partial_\tau u)^2 + \left(\frac{1}{2(1-\lambda)} + \frac{2}{1-\lambda} + \frac{1}{(1-\lambda)^2} \right) (\partial_R u)^2 - 2 \partial_\tau u \partial_R u + e^{2Z} \frac{f^2(u)}{R^2} \right), \quad (2.9)$$

now integrating (2.9) in view of the fall of rates gives the result as $R \rightarrow \infty$ for fixed λ .

□

In order to obtain equivalent results of Proposition 2.1 for the fully coupled equivariant Einstein-wave map system we turn to the double null coordinates (ξ, η, θ) . In addition, we shall give a proof without the use of asymptotic expansions and analyticity.

Consider the asymptotic behaviour of the metric (Σ, q)

$$q_{ab} = R^{-m_{AV}}(dR^2 + R^2 d\theta^2)$$

In view of the asymptotic fall-off conditions along the $\eta = c$ lines we shall prove the mass-loss formula. Consider the $T = c$ slices of the system, then the Hamiltonian constraint $E(\mathcal{N}, \mathcal{N}) = T(\mathcal{N}, \mathcal{N})$, $N = e^{-Z} \partial_T$ can be calculated from (??), as

$$R^{-1} e^Z \partial_R Z = \frac{1}{2} \left((\partial_T u)^2 + (\partial_R u)^2 + e^{2Z} \frac{f^2(u)}{R^2} \right) e^Z \quad (2.10)$$

If

$$E(U) = \int_{\Sigma_0} (\partial_T u)^2 + (\partial_R u)^2 + e^{2Z} \frac{f^2(u)}{R^2} \bar{\mu}_q \quad (2.11)$$

$$\pi m_{AV} = 2\pi(1 - e^{Z_\infty}) = E(U) \quad (2.12)$$

where $Z_\infty := \lim_{R \rightarrow \infty} Z$ on (Σ, q) . Now then, we have the following

Remark 2.2 (Mass loss). *Suppose u , $\partial_R u$ and $\partial_\eta u$ admit the asymptotic expansions as in Proposition 2.1, and $Z(\eta, R)$ is a globally regular solution with $|e^{2Z} - 1| \leq c(E_0)$ then we have the following mass loss formula*

$$e^{Z(\eta, R)} = \left(1 - \frac{m_{AV}}{2}\right) - \int_\infty^\eta f_0^2(\eta) \bar{\mu}_\eta \quad (2.13)$$

which follows from the integration of (2.4) and noting that $\lim_{\eta \rightarrow -\infty} \lim_{R \rightarrow \infty} Z = Z_\infty$. In the case of the fully coupled 2+1 equivariant Einstein-wave map system we shall prove the scattering of the system, but the quantitative aspects asymptotic geometry are encoded in the rate at which the maximal development (M, g, U) of the system converges to its scattering state. A characterization of this rate and its close connection with the asymptotics of the Weyl tensor of the original 3 + 1 metric shall be the subject of a subsequent article.

3 Scattering for 2+1 Equivariant Einstein Wave Map System

Recall the 2+1 dimensional equivariant Einstein-wave system:

$$\mathbf{E}_{\mu\nu} = \mathbf{T}_{\mu\nu}, \quad \text{on } (M, g) \quad (3.1)$$

$$\square_{g(u)} u = \frac{f_u(u)f(u)}{r^2} \quad (3.2)$$

for the initial data $(\Sigma_0, q_0, K_0, u_0, u_1)$. Now consider the following nonlinear second-order hyperbolic partial differential equation ($n \neq 1$):

$$\left. \begin{aligned} L\psi &= N(\psi) && \text{on } (M, g) \\ \psi(0, x) = \psi_0 \text{ and } \partial_T \psi(0, x) = \psi_1 &&& \text{on } (\Sigma, q) \end{aligned} \right\} \quad (3.3)$$

where L is a linear operator and N is the nonlinearity. We say that the wave equation (3.6) exhibits temporal scattering if there exists a solution of a corresponding linearized equation ψ_S and an energy topology E_ψ such that

$$\left. \begin{aligned} L\psi_S &= 0 && \text{on } (M, g) \\ \psi_S(0, x) = (\psi_S)_0 \text{ and } \partial_T \psi_S(0, x) = (\psi_S)_1 &&& \text{on } (\Sigma, q) \end{aligned} \right\} \quad (3.4)$$

and

$$\|\psi - \psi_S\|_{E_\psi} \rightarrow 0, \quad \text{as } T \rightarrow \infty, \quad (3.5)$$

where $((\psi_S)_0, (\psi_S)_1)$ are the ‘induced’ initial data of (3.7). As we already remarked, we shall recast the wave maps equation as a 4+1 dimensional wave equation,

$$\left. \begin{aligned} {}^{4+1}\square v &= F(v) && \text{on } \mathbb{R}^{4+1} \\ v(0, x) = \psi_0 \text{ and } \partial_T v(0, x) = v_1 &&& \text{on } \mathbb{R}^4 \end{aligned} \right\} \quad (3.6)$$

where v is coupled to the 2+1 Einstein’s equations with $u = Rv$.

Following our previous definition, the equation (3.6) scatters in time if there exists a v_S such that

$$\left. \begin{aligned} {}^{4+1}\square v_S &= 0 && \text{on } \mathbb{R}^{4+1} \\ v_S(0, x) = (v_S)_0 \text{ and } \partial_T v_S(0, x) = (v_S)_1 &&& \text{on } \mathbb{R}^4 \end{aligned} \right\} \quad (3.7)$$

$$\|v - v_S\|_{E_v} \rightarrow 0 \quad \text{as } T \rightarrow \infty \quad (3.8)$$

where $(v_S)_0$ and $(v_S)_1$ are the scattering data

$$(v_S)_0 = \mathcal{F}^{-1} \left(\hat{v}_0 - \int_0^\infty \frac{\sin \sqrt{\Delta} s}{\sqrt{\Delta}} \hat{F}(s) ds \right) \quad (3.9)$$

$$(v_S)_1 = \mathcal{F}^{-1} \left(\hat{v}_1 - \int_0^\infty \cos(\sqrt{\Delta} s) \hat{F}(s) ds \right) \quad (3.10)$$

and the energy topology of the problem (3.6) is $\dot{H}(\mathbb{R}^4) \times L^2(\mathbb{R}^4)$. We shall use the scattering theory for v to make a scattering statement for (3.3) for u . Likewise, if we consider a 1D wave equation

$$\left. \begin{aligned} {}^{1+1}\square w &= g(w) && \text{on } (M, g) \\ w_\Gamma = w_0 \quad \text{and} \quad \partial_R w_\Gamma = w_1 &&& \text{on } \Gamma \end{aligned} \right\} \quad (3.11)$$

Suppose $S_w(\tau)(w_0, w_1)$ is the evolution operator for the wave equation where τ is the evolution parameter. In this work we shall set up the initial value problem also on the axis Γ and study the scattering for this asymptotics. In this case, we say that the solution for (3.11) if there exists a solution w_S corresponding to the linearization of (3.11) with respect to the full 2+1 equivariant Einstein-wave map system, such that

$$\|w - w_S\|_{L^\infty} \rightarrow 0, \quad \text{as } \tau \rightarrow \infty. \quad (3.12)$$

Lemma 3.1. *Suppose Z is a solution of the wave equation,*

$${}^{1+1}\square Z = \frac{f^2(u)}{r^2} - \partial_\xi u \partial_\eta u \quad (3.13)$$

then Z satisfies:

$$\begin{aligned} Z(T, R) = & c_1 \int_0^R v^2(s, T + R - s) s ds + c_2 \int_0^R v^2(s, T - R + s) s ds \\ & + c_3 R^2 v^2(R, T) + c_4 \int_0^R \int_{T-(R+s)}^{T+(R-s)} v^2 ds dT' + c_5 \int_0^R \int_{T-(R+s)}^{T+(R-s)} F(v) v s^2 ds dT'. \end{aligned} \quad (3.14)$$

for some constants $c_i, i = 1 \cdots 5$.

Proof. Now since $u = Rv$, by the product rule

$$-\frac{1}{2}(\partial_\eta u)(\partial_\xi u) = -\frac{1}{2}R^2(\partial_\eta v)(\partial_\xi v) - \frac{1}{2}R(\partial_\eta v)v - \frac{1}{2}R(\partial_\xi v)v - \frac{1}{2}v^2. \quad (3.15)$$

Making a change of variables,

$$\begin{aligned} \int_0^R \int_{T-(R-s)}^{T+(R-s)} s(\partial_\eta v)v dt ds &= \frac{1}{2} \int_{T-R}^{T+R} \int_{T-R}^\xi (\xi - \eta)(\partial_\eta v)v d\eta d\xi \\ &= \frac{1}{4} \int_{T-R}^{T+R} \int_{T-R}^\xi (\xi - \eta)\partial_\eta(v^2) d\eta d\xi. \end{aligned} \quad (3.16)$$

Integrating by parts,

$$= \frac{1}{4} \int_{T-R}^{T+R} (\xi - (T - R))v^2(T - R, \xi) d\xi - \frac{1}{4} \int_{T-R}^{T+R} \int_{T-R}^\xi v^2(\eta, \xi) d\eta d\xi. \quad (3.17)$$

By a similar calculation,

$$\begin{aligned} \int_0^R \int_{T-(R-s)}^{T+(R-s)} s(\partial_\xi v)v dt ds &= \frac{1}{2} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta)(\partial_\xi v)v d\xi d\eta \\ &= \frac{1}{4} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta)\partial_\xi(v^2) d\xi d\eta \end{aligned} \quad (3.18)$$

$$\begin{aligned} &= \frac{1}{4} \int_{T-R}^{T+R} (T + R - \eta)v^2(\eta, T + R) d\eta \\ &\quad - \frac{1}{4} \int_{T-R}^{T+R} \int_\eta^{T+R} v^2(\eta, \xi) d\xi d\eta. \end{aligned} \quad (3.19)$$

Next, integrating by parts,

$$\frac{1}{2} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta)^2(\partial_\xi v)(\partial_\eta v) d\xi d\eta = \frac{1}{2} \int_{T-R}^{T+R} (T + R - \eta)^2 v(\partial_\eta v)(T + R, \eta) d\eta \quad (3.20)$$

$$-\frac{1}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta)^2 v(\partial_{\xi\eta} v) d\xi d\eta - \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta) v(\partial_{\eta} v) d\xi d\eta. \quad (3.21)$$

Now by a change of variables,

$$\frac{1}{4} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta) v(\partial_{\eta} v) d\xi d\eta = (3.105). \quad (3.22)$$

Also, integrating by parts,

$$\begin{aligned} & \frac{1}{2} \int_{T-R}^{T+R} (T + R - \eta)^2 v(\partial_{\eta} v)(T + R, \eta) d\eta \\ &= -\frac{1}{4} (2R)^2 v^2(T + R, T - R) + \frac{1}{2} \int_{T-R}^{T+R} (T + R - \eta) v^2(T + R, \eta) d\eta. \end{aligned} \quad (3.23)$$

Finally, by (??),

$$\begin{aligned} -\frac{1}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta)^2 v(\partial_{\xi\eta} v) d\xi d\eta &= \frac{3}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta) v(\partial_R v) d\xi d\eta \\ &\quad - \frac{1}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta)^2 v F(v) d\xi d\eta. \end{aligned} \quad (3.24)$$

Now, making a change of variables,

$$\frac{3}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta) v(\partial_R v) d\xi d\eta = \frac{3}{2} \int_0^R \int_{T-(R+s)}^{T+(R-s)} v(\partial_s v) s ds dT'. \quad (3.25)$$

Integrating by parts,

$$\begin{aligned} &= \frac{3}{4} \int_0^R v^2(s, T + R - s) s ds + \frac{3}{4} \int_{T-R}^T v^2(s, T - R + s) s ds \\ &\quad - \frac{3}{2} \int_0^R \int_{T-(R+s)}^{T+(R-s)} v^2 ds dT'. \end{aligned} \quad (3.26)$$

Therefore we have

$$\begin{aligned} Z = & c_1 \int_0^R v^2(s, T + R - s) s ds + c_2 \int_0^R v^2(s, T - R + s) s ds \\ & + c_3 R^2 v^2(R, T) + c_4 \int_0^R \int_{T-(R+s)}^{T+(R-s)} v^2 ds dT' + c_5 \int_0^R \int_{T-(R+s)}^{T+(R-s)} F(v) v s^2 ds dT'. \end{aligned} \quad (3.27)$$

□

Consider the radial wave equation,

$${}^{4+1}\square v = F(v), \quad \text{on } (\mathbb{R}^{4+1}, \check{g}) \quad (3.28)$$

and the energy-momentum tensor that arises directly from the variational principle of (3.28)

$$\check{T}_{\mu\nu} := \nabla_\mu v \nabla_\nu v - \frac{1}{2} \check{g}_{\mu\nu} \nabla^\sigma v \nabla_\sigma v - \check{g}_{\mu\nu} \tilde{F}(v), \quad \mu, \nu, \sigma = 0, 1 \dots 4. \quad (3.29)$$

where $\tilde{F}(v)$ is the primitive of F with respect to v . The divergence of \check{T} is given by

$$\begin{aligned} \nabla^\nu \check{T}_{\mu\nu} &= (\partial^\nu \partial_\mu v) \partial_\nu v + \partial_\mu v {}^{4+1}\square v - \frac{1}{2} \check{g}_{\mu\nu} \partial^\nu (\partial^\sigma v \partial_\sigma v) - \check{g}_{\mu\nu} \partial^\nu \tilde{F}(v) \\ &= \partial_\mu v ({}^{4+1}\square v) - \check{g}_{\mu\nu} \partial^\nu \tilde{F}(v) \end{aligned} \quad (3.30)$$

Consider the vector field \mathfrak{X} such that its momentum is given by

$$J_{\mathfrak{X}} = \check{\mathbf{T}}(\mathfrak{X}) = \check{T}_\mu^\nu \mathfrak{X}^\mu \quad (3.31)$$

so that, after relabeling the indices for convenience, we have the identity

$$\nabla_\nu J_{\mathfrak{X}}^\nu = \frac{1}{2} {}^{(\mathfrak{X})} \pi_{\mu\nu} \check{T}^{\mu\nu} + \mathfrak{X}^\mu \nabla^\nu \check{T}_{\mu\nu}. \quad (3.32)$$

Define \check{e} and \check{m} such that

$$\check{e} + \tilde{F}(v) = \check{T}(\partial_T, \partial_T), \quad \check{m} := \check{T}(\partial_T, \partial_R). \quad (3.33)$$

Now consider a Morawetz multiplier vector $\mathfrak{X} := \mathfrak{F}(R)\partial_R$ so that the corresponding momentum is given by

$$\begin{aligned} J_{\mathfrak{X}} &= \check{\mathbf{T}}(\mathfrak{X}) \\ &= \mathfrak{F}(R) \left(-\check{m}\partial_T + (\check{e} + \tilde{F}(v))\partial_R \right) \end{aligned} \quad (3.34)$$

and its divergence

$$\nabla_{\nu} J_{\mathfrak{X}}^{\nu} = \frac{1}{2} \check{\mathbf{T}}^{\mu\nu}(\mathfrak{X}) \pi_{\mu\nu} + \mathfrak{X}^{\mu} \partial_{\mu} v^{(4+1)} \square v - \mathfrak{X}^{\mu} \check{g}_{\mu\nu} \partial^{\nu} \tilde{F}(v), \quad (3.35)$$

where the non-zero terms of deformation tensor

$${}^{(\mathfrak{X})}\pi_{\mu\nu} := \check{g}_{\sigma\nu} \partial_{\mu} \mathfrak{X}^{\sigma} + \check{g}_{\sigma\mu} \partial_{\nu} \mathfrak{X}^{\sigma} + \mathfrak{X}^{\sigma} \partial_{\sigma} \check{g}_{\mu\nu}$$

are given by

$$\begin{aligned} {}^{(\mathfrak{X})}\pi_{RR} &= 2g_{RR}\partial_R \mathfrak{F}(R), {}^{(\mathfrak{X})}\pi_{\theta\theta} = \frac{2}{R} g_{\theta\theta} \mathfrak{F}(R), \\ {}^{(\mathfrak{X})}\pi_{\phi\phi} &= \frac{2}{R} g_{\phi\phi} \mathfrak{F}(R), {}^{(\mathfrak{X})}\pi_{\psi\psi} = \frac{2}{R} g_{\psi\psi} \mathfrak{F}(R). \end{aligned}$$

Consequently a calculation shows that (3.35) can be represented as

$$\nabla_{\nu} J_{\mathfrak{X}}^{\nu} = \left(-\frac{6\mathfrak{F}(R)}{R} \check{\mathcal{L}} + \check{e} \partial_R \mathfrak{F}(R) \right) + \mathfrak{F}(R) \partial_R v F(v) - \mathfrak{F}(R) \partial^R \tilde{F}(v) \quad (3.36)$$

Now define the following lower-order momentum vector

$$J_1^{\nu}[v] := \kappa v \nabla^{\nu} v - \frac{1}{2} v^2 \nabla^{\nu} \kappa + \tilde{F}(v) \mathfrak{X}^{\nu}. \quad (3.37)$$

Its divergence is

$$\begin{aligned} \nabla_{\nu} J_1^{\nu} &= \kappa v^{4+1} \square v + \kappa \nabla^{\nu} v \nabla_{\nu} v + v \nabla^{\nu} v \nabla_{\nu} \kappa - (\square \kappa) \frac{v^2}{2} \\ &\quad - v \nabla^{\nu} \kappa \nabla_{\nu} v + \mathfrak{X}^{\nu} \partial_{\nu} \tilde{F}(v) + \tilde{F}(v) \nabla_{\nu} \mathfrak{X}^{\nu} \\ &= \kappa v^{4+1} \square v + \kappa \nabla^{\nu} v \nabla_{\nu} v - (\square \kappa) \frac{v^2}{2} + \mathfrak{X}^{\nu} \partial_{\nu} \tilde{F}(v) + \tilde{F}(v) \nabla_{\nu} \mathfrak{X}^{\nu} \\ &= \kappa v^{4+1} \square v + 2\kappa \check{\mathcal{L}} - (\square \kappa) \frac{v^2}{2} + \mathfrak{X}^{\nu} \partial_{\nu} \tilde{F}(v) + \tilde{F}(v) \nabla_{\nu} \mathfrak{X}^{\nu}. \end{aligned} \quad (3.38)$$

In addition, consider a timelike multiplier vector field $\mathfrak{T} = \partial_T$ such that the corresponding current is

$$\begin{aligned} J_T &:= \check{T}(\mathfrak{T}) \\ &= (-\check{e} - \tilde{F}(v))\partial_T + \check{m}\partial_R \end{aligned} \quad (3.39)$$

the divergence

$$\begin{aligned} \nabla_\nu J_T^\nu &= \frac{1}{2}{}^{(\mathfrak{T})}\pi_{\mu\nu}\check{T}^{\mu\nu} + (\mathfrak{T})^\mu\nabla^\nu\check{T}_{\mu\nu} \\ &= \partial_T v F(v) + \partial^T \tilde{F}(v) \end{aligned} \quad (3.40)$$

in view of ${}^{(\mathfrak{T})}\pi_{\mu\nu} \equiv 0$. Likewise, if we define a lower-order momentum vector field

$$J_2^\nu := \tilde{F}(v)\mathfrak{T}^\nu$$

then

$$\nabla_\nu J_2^\nu = \mathfrak{T}^\nu \partial_\nu \tilde{F}(v) + \tilde{F}(v)\nabla_\nu \mathfrak{T}^\nu. \quad (3.41)$$

Consider the volume 3-form of (M, g) :

$$\bar{\mu}_g = \frac{1}{2}re^{2Z}d\eta \wedge d\xi \wedge d\theta \quad (3.42)$$

Define the 2-forms $\bar{\mu}_\xi$ and $\bar{\mu}_\eta$ such that

$$d\xi \wedge \bar{\mu}_\xi := \bar{\mu}_g \quad (3.43)$$

$$d\eta \wedge \bar{\mu}_\eta := \bar{\mu}_g \quad (3.44)$$

so that we have

$$\bar{\mu}_\xi = -\frac{1}{2}re^{2Z}(d\eta \wedge d\theta) \quad (3.45)$$

$$\bar{\mu}_\eta = \frac{1}{2}re^{2Z}(d\xi \wedge d\theta) \quad (3.46)$$

explicitly. Suppose \vec{v} is smooth vector field defined on (M, g) then the flux through the $\eta = c$ and $\xi = c$ null hypersurfaces are

$$\text{Flux}^+(\vec{v}) := \int_{\{\eta=c\}} d\eta(\vec{v}) \bar{\mu}_\eta \quad (3.47)$$

$$\text{Flux}^-(\vec{v}) := \int_{\{\xi=c\}} d\xi(\vec{v}) \bar{\mu}_\xi \quad (3.48)$$

respectively. Now having fixed the orientation, let us now give the equivalent definitions for the corresponding 4+1 Minkowski metric $(\mathbb{R}^{4+1}, \check{g})$ that are consistent with the original spacetime (M, g) :

$$\check{\mu}_{\check{g}} = \frac{1}{2} \sqrt{-\check{g}} d\eta \wedge d\xi \wedge d\omega_{\mathbb{S}^3}. \quad (3.49)$$

Define $\check{\mu}_\eta$ and $\check{\mu}_\xi$ such that,

$$d\xi \wedge \check{\mu}_\xi := \check{\mu}_{\check{g}} \quad (3.50)$$

$$d\eta \wedge \check{\mu}_\eta := \check{\mu}_{\check{g}}, \quad (3.51)$$

and for a smooth vector field P in $(\mathbb{R}^{4+1}, \check{g})$ the fluxes through $\eta = c$ and $\xi = c$ hypersurfaces are

$$\text{Flux}^+(P) := \int_{\{\eta=c\}} d\eta(P) \bar{\mu}_\eta \quad (3.52)$$

$$\text{Flux}^-(P) := \int_{\{\xi=c\}} d\xi(P) \bar{\mu}_\xi. \quad (3.53)$$

If we consider the vector field J_T , we have

$$\text{Flux}^+(J_T) = \int_{\{\eta=c\}} -(\check{e} + \tilde{F}(v) + \check{m}) \check{\mu}_\eta \quad (3.54a)$$

$$\text{Flux}^-(J_T) = \int_{\{\xi=c\}} -(\check{e} + \tilde{F}(v) - \check{m}) \check{\mu}_\xi \quad (3.54b)$$

$$\text{Flux}^+(J_2) = \int_{\{\eta=c\}} (\tilde{F}(v)) \check{\mu}_\eta \quad (3.54c)$$

$$\text{Flux}^-(J_2) = \int_{\{\xi=c\}} -(\tilde{F}(v)) \check{\mu}_\xi \quad (3.54d)$$

Lemma 3.2. *Suppose (M, g, U) is a globally regular solution of the initial value problem of the Einstein-equivariant wave map system with $E_0 < \epsilon^2$, for ϵ sufficiently small, we have*

$$\text{Flux}^+(J) \geq 0, \quad (3.55a)$$

$$\text{Flux}^-(J) \leq 0, \quad (3.55b)$$

where $J := J_T + J_2$.

Theorem 3.3 (Nonlinear Morawetz Estimate). *Suppose v is a globally regular solution of the equation*

$$\left. \begin{array}{ll} {}^{4+1}\square v &= F(v) & \text{on } \mathbb{R}^{4+1} \\ v_0 = v(0, x) \text{ and } v_1 = \partial_T v(0, x) & & \text{on } \mathbb{R}^4 \end{array} \right\} \quad (3.56)$$

where

$$\begin{aligned} F(v) := & \left(e^{2Z} - 1 + \left(\frac{r}{R} \partial_\eta r + \frac{1}{2} \right) - \left(\frac{r}{R} \partial_\xi r - \frac{1}{2} \right) \right) \frac{v}{r^2} + 2\partial_\xi v \partial_\eta \log \left(\frac{r}{R} \right) \\ & + 2\partial_\eta v \partial_\xi \log \left(\frac{r}{R} \right) + e^{2Z} \frac{R^2}{r^2} v^3 \zeta(Rv) \end{aligned} \quad (3.57)$$

with Z and r coupled as in (1.3), then

$$\int_{\mathbb{R}^{4+1}} \frac{v^2}{|x|^3} \bar{\mu}_{\check{g}} \leq \|v_0\|_{H^1(\mathbb{R}^4)}^2 + \|v_1\|_{L^2(\mathbb{R}^4)}^2. \quad (3.58)$$

Proof. To prove the result, we shall use the Morawetz multiplier method we introduced above. Consider the multiplier \mathfrak{X} such that, $\mathfrak{F}(R) = \frac{1}{3}$, we have

$$J_{\mathfrak{X}} = \frac{1}{3} (-\check{m} \partial_T + (\check{e} + \tilde{F}(v)) \partial_R) \quad (3.59)$$

and

$$\nabla_\nu J_{\mathfrak{X}}^\nu = -\frac{2}{R} \check{\mathcal{L}} + \frac{1}{3} \partial_R v F(v) - \mathfrak{X}^\mu \check{g}_{\mu\nu} \partial^\nu \tilde{F}(v). \quad (3.60)$$

Likewise, if we consider the choice of $\kappa = \frac{1}{R}$, then

$${}^{4+1}\square\kappa = -\frac{1}{R^3} \quad (3.61)$$

$$J_\kappa = -\left(\frac{1}{R}v\partial_T v\right)\partial_T + \left(\frac{1}{R}v\partial_R v + \frac{v}{R^2} + \mathfrak{X}^R\tilde{F}(v)\right)\partial_R \quad (3.62)$$

$$\nabla_\nu J_\kappa^\nu = \frac{2}{R}\check{\mathcal{L}} + \frac{1}{R}vF(v) + \frac{1}{2}\frac{v^2}{R^3} + \mathfrak{X}^\nu\partial_\nu\tilde{F}(v) \quad (3.63)$$

Subsequently, if we consider the sum vector $J^\nu := J_\mathfrak{X}^\nu + J_\kappa^\nu$, then we have

$$\begin{aligned} \nabla_\nu J^\nu &= \nabla_\nu J_\mathfrak{X}^\nu + \nabla_\nu J_\kappa^\nu \\ &= \frac{1}{2}\frac{v^2}{R^3} + \frac{1}{3}\partial_R v F(v) + \frac{1}{R}vF(v) \end{aligned} \quad (3.64)$$

We have the following identity from the Stokes theorem, for $\check{\Sigma}_0$ and $\check{\Sigma}_T$,

$$\frac{1}{2}\int\frac{v^2}{R^3}\bar{\mu}_{\check{g}} = \int\nabla_\nu J_S^\nu\bar{\mu}_{\check{g}} = \int_{\check{\Sigma}_0}\langle\partial_T, J_S\rangle\bar{\mu}_{\check{q}} - \int_{\check{\Sigma}_T}\langle\partial_T, J_S\rangle\bar{\mu}_{\check{q}} \quad (3.65)$$

If we consider the boundary terms on $\check{\Sigma}_0$ and $\check{\Sigma}_T$

$$\check{g}_{\mu\nu}J_S^\mu(\partial_T)^\nu = \check{g}_{\mu\nu}J_\mathfrak{X}^\mu(\partial_T)^\nu + \check{g}_{\mu\nu}J_1^\mu(\partial_T)^\nu \quad (3.66)$$

where

$$\check{g}_{\mu\nu}J_\mathfrak{X}^\mu(\partial_T)^\nu = \check{g}_{TT}J_\mathfrak{X}^T(\partial_T)^T = \mathfrak{F}(R)\check{m} \quad (3.67)$$

and

$$\check{g}_{\mu\nu}J_1^\mu(\partial_T)^\nu = \check{g}_{TT}J_1^T(\partial_T)^T = -\frac{1}{|x|}v\partial_T v \quad (3.68)$$

it follows from the dominant energy condition and Hardy's inequality that

$$\int_{\check{\Sigma}_0}\langle\partial_T, J_S\rangle\bar{\mu}_{\check{q}}, \quad \int_{\check{\Sigma}_T}\langle\partial_T, J_S\rangle\bar{\mu}_{\check{q}} \leq \|v_0\|_{\dot{H}^1(\mathbb{R}^4)}^2 + \|v_1\|_{L^2(\mathbb{R}^4)}^2. \quad (3.69)$$

Thus,

$$\int_{\mathbb{R}^{4+1}} \frac{v^2}{R^3} \bar{\mu}_{\tilde{g}} \leq E(v) + \left| \int \int F(v) v R^2 dR dT \right| + \left| \int \int F(v) v_R R^3 dR dT \right|. \quad (3.70)$$

Therefore, to prove the theorem it suffices to absorb the last two terms in the right hand side of (3.70) into the left hand side of (3.70). The term

$$\int_{\mathbb{R}^{4+1}} \frac{F(v) v}{|x|} \bar{\mu}_{\tilde{g}} \quad (3.71)$$

is easier. Making a change of variables,

$$\int_{\mathbb{R}} \int_{\mathbb{R}} F(v) v R^2 dR dT = \frac{1}{4} \int_{\mathbb{R}} \int_{\mathbb{R}} F(v(\xi, \eta)) v(\xi, \eta) (\xi - \eta)^2 d\xi d\eta. \quad (3.72)$$

Now since

$$\left| \frac{R}{r} - 1 \right|, \left| \partial_{\eta} r - \frac{1}{2} \right|, \left| \partial_{\xi} r + \frac{1}{2} \right| \leq \epsilon(E(v)),$$

where $\epsilon(E(v)) \searrow 0$ as $E \searrow 0$, the Cauchy-Schwartz inequality

$$\begin{aligned} & \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \partial_{\eta} r - \frac{1}{2} \right| \left| \partial_{\xi} v(\xi, \eta) \right| |v(\xi, \eta)| |\xi - \eta| d\xi d\eta \\ & \leq \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \partial_{\eta} r - \frac{1}{2} \right| \left| \partial_{\xi} v(\xi, \eta) \right|^2 |\xi - \eta|^2 d\xi d\eta + \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \partial_{\eta} r - \frac{1}{2} \right| |v(\xi, \eta)|^2 d\xi d\eta. \end{aligned} \quad (3.73)$$

Now again by a change of variables

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \partial_{\eta} r - \frac{1}{2} \right| |v(\xi, \eta)|^2 d\xi d\eta \leq \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} |v(R, T)|^2 dR dT. \quad (3.74)$$

It is also straightforward to show that

$$\begin{aligned} & \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \partial_{\eta} r - \frac{1}{2} \right| |\partial_{\xi} v(\xi, \eta)|^2 d\xi d\eta \\ & \leq \left(\int_{\mathbb{R}} \left(\sup_{\xi} \left| \frac{1}{|\xi - \eta|} \right| \left| \partial_{\eta} r - \frac{1}{2}(\xi, \eta) \right| \right) d\eta \right) \cdot \sup_{\eta} \left(\int |\xi - \eta|^3 |\partial_{\xi} v(\xi, \eta)|^2 d\xi \right). \end{aligned} \quad (3.75)$$

By energy bounds on the boundary of a light cone, since $|\xi - \eta| = R$,

$$\int_{\mathbb{R}} |\xi - \eta|^3 |\partial_{\xi} v(\xi, \eta)|^2 d\xi \leq E(v). \quad (3.76)$$

Also by the fundamental theorem of calculus, the fact that $\partial_{\eta} r \Big|_{R=0} = \frac{1}{2}$, and $|\partial_{\xi} \partial_{\eta} r| \leq R|v|^2$,

$$\sup_{\xi} \frac{1}{|\xi - \eta|} \left| (\partial_{\eta} r - \frac{1}{2})(\xi, \eta) \right| \leq \int_{\mathbb{R}} |v(\xi, \eta)|^2 d\eta. \quad (3.77)$$

Therefore,

$$(3.73) \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} |v(\xi, \eta)|^2 d\xi d\eta. \quad (3.78)$$

By a similar calculation,

$$\begin{aligned} & \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{1}{R} \left| \partial_{\xi} r + \frac{1}{2} \right| \left| \partial_{\eta} v(R, T) \right| |v(R, T)| R^2 dR dT \\ & \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT + \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \end{aligned} \quad (3.79)$$

Also, since $|Z| \leq \epsilon(E(v))$,

$$\begin{aligned} & \int_{\mathbb{R}} \int_{\mathbb{R}} \left(e^{2Z} - 1 + \partial_{\eta} r + \frac{1}{2} - (\partial_{\xi} r - \frac{1}{2}) \right) \frac{v(R, T)^2}{R^2} R^2 dR dT \\ & \leq \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \end{aligned} \quad (3.80)$$

Therefore, combining (3.74), (3.78), and (3.80),

$$(3.132) \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT + \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \quad (3.81)$$

The bulk term

$$\int_{\mathbb{R}} F(v) \partial_R v \bar{\mu}_{\dot{g}} \quad (3.82)$$

is more difficult to control. Several components of this term are essentially the same as in (3.132). Indeed, as in (3.76) - (3.78),

$$\int_{\mathbb{R}^{4+1}} \frac{1}{R} \left| \partial_\eta r - \frac{1}{2} \right| \left| \partial_\xi v(R, T) \right| \left| \partial_R v(R, T) \right| \bar{\mu}_{\dot{g}} \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT, \quad (3.83)$$

and

$$\int_{\mathbb{R}^{4+1}} \frac{1}{R} \left| \partial_\xi r + \frac{1}{2} \right| \left| \partial_\eta v(R, T) \right| \left| \partial_R v(R, T) \right| \bar{\mu}_{\dot{g}} \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \quad (3.84)$$

Also,

$$\begin{aligned} & \int_{\mathbb{R}^{4+1}} \left| \partial_\eta r + \frac{1}{2} \right| \left| \frac{v(R, T)}{R^2} \right| |v_R(R, T)| \bar{\mu}_{\dot{g}} \\ & \leq \int_{\mathbb{R}^{4+1}} \frac{1}{R} \left| \partial_\eta r + \frac{1}{2} \right| v_R(R, T)^2 \bar{\mu}_{\dot{g}} + \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \partial_\eta r + \frac{1}{2} \right| v(R, T)^2 dR dT \\ & \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT + \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT, \end{aligned} \quad (3.85)$$

and

$$\begin{aligned} & \int_{\mathbb{R}^{4+1}} \left| \partial_\xi r - \frac{1}{2} \right| \left| \frac{v(R, T)}{R^2} \right| |v_R(R, T)| \bar{\mu}_{\dot{g}} \\ & \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT + \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \end{aligned} \quad (3.86)$$

It remains to compute

$$\int_{\mathbb{R}^{4+1}} (e^{2Z} - 1) \frac{v(R, T)}{R^2} v_R(R, T) \bar{\mu}_{\dot{g}}, \quad (3.87)$$

and

$$\int_{\mathbb{R}^{4+1}} e^{2Z} \frac{R^2}{r^2} v(R, T)^3 \zeta(Rv) v_R(R, T) \bar{\mu}_{\dot{g}}. \quad (3.88)$$

Making a power series expansion,

$$e^{2Z} = 1 + 2Z + 2Z^2 + \frac{4}{3}Z^3 + \dots \quad (3.89)$$

because $|Z| \leq \epsilon(E(v))$ it suffices to consider only $1 + 2Z$. The other terms follow in a similar manner. In view of Lemma 3.1, to estimate

$$2 \int \int Z(R, T) \frac{v(R, T)}{R^2} v_R(R, T) R^3 dR dT, \quad (3.90)$$

we shall split $Z(R, T) = Z_1(R, T) + Z_2(R, T)$, where $Z_1(R, T)$ has good integral properties and $Z_2(R, T)$ has a radial derivative with good integral properties. By equation (7d),

$$\partial_{\xi\eta}^2 Z = -\frac{1}{2}(\partial_\eta u)(\partial_\xi u) - \frac{e^{2Z}}{8} \frac{f^2(u)}{r^2}. \quad (3.91)$$

Then by Duhamel's principle and the fact that $Z = Z_R = 0$ on $R = 0$,

$$Z(R, T) = \int_0^R \int_{T-(R-s)}^{T+(R-s)} (\partial_{\xi\eta}^2 Z)(s, t) dt ds. \quad (3.92)$$

Now the term

$$- \int_0^R \int_{T-(R-s)}^{T+(R-s)} \frac{e^{2Z}}{8} \frac{f^2(sv)}{r^2} ds dt = Z_2^{(1)}(R, T) \quad (3.93)$$

may be safely placed in Z_2 . Indeed, integrating by parts,

$$\int_{\mathbb{R}} \int_{\mathbb{R}} Z_2^{(1)}(R, T) v(R, T) v_R(R, T) R dR dT \quad (3.94)$$

$$= -\frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} Z_2^{(1)}(R, T) v(R, T)^2 dR dT - \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \partial_R(Z_2^{(1)}(R, T)) v(R, T)^2 R dR dT. \quad (3.95)$$

Since $|Z_2^{(1)}(R, T)| \lesssim \int \int v(r, t)^2 dr dt$,

$$\frac{1}{2} \int \int Z_2^{(1)} v(R, T)^2 dR dT \leq \left(\int \int v(R', T')^2 dR' dT' \right)^2. \quad (3.96)$$

Meanwhile, we can directly compute

$$\begin{aligned} -\partial_R \left(\int_0^R \int_{T-(R-s)}^{T+(R-s)} \frac{e^{2Z}}{8} \frac{f^2(sv)}{r^2} ds dT' \right) &= - \int_0^R \frac{e^{2Z}}{8} \frac{f^2(sv)}{r^2} (s, T + (R-s)) ds \\ &\quad - \int_0^R \frac{e^{2Z}}{8} \frac{f^2(sv)}{r^2} (s, T - (R-s)) ds. \end{aligned} \quad (3.97)$$

Since $|Z|$ is small,

$$\sup_R \int_0^R \frac{e^{2Z}}{8} \frac{f^2(sv)}{r^2} (s, T + (R-s)) ds \leq \int v^2(\eta, \xi = R+T) d\eta, \quad (3.98)$$

and

$$\sup_R \int_0^R \frac{e^{2Z}}{8} \frac{f^2(sv)}{r^2} (s, T - (R-s)) ds \lesssim \int v^2(\eta = T-R, \xi) d\xi. \quad (3.99)$$

Then since Hardy's inequality implies

$$\int_0^\infty v^2(s, T + R - s) s ds \leq E(v), \quad (3.100)$$

and

$$\int_0^\infty v^2(s, T - R + s) s ds \leq E(v), \quad (3.101)$$

so therefore,

$$-\frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \partial_R (Z_2^{(1)}(R, T)) v(R, T)^2 R dR dT \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v^2(R, T) dR dT. \quad (3.102)$$

Now by the product rule, $u = Rv$,

$$-\frac{1}{2} (\partial_\xi u) (\partial_\eta u) = \frac{1}{2} v^2 + \frac{1}{2} Rv (\partial_\eta v) - \frac{1}{2} Rv (\partial_\xi v) - \frac{1}{2} R^2 (\partial_\xi v) (\partial_\eta v). \quad (3.103)$$

Similar to (3.93),

$$Z_2^{(2)}(R, T) = -\frac{1}{2} \int_0^R \int_{T-(R-s)}^{T+(R-s)} v^2(s, t) dr dt, \quad (3.104)$$

and can be analyzed in a manner similar to (3.94) - (3.102). Next, making a change of variables,

$$\begin{aligned} \int_0^R \int_{T-(R-s)}^{T+(R-s)} s(\partial_\eta v) v dt ds &= \frac{1}{2} \int_{T-R}^{T+R} \int_{T-R}^\xi (\xi - \eta)(\partial_\eta v) v d\eta d\xi \\ &= \frac{1}{4} \int_{T-R}^{T+R} \int_{T-R}^\xi (\xi - \eta) \partial_\eta (v^2) d\eta d\xi. \end{aligned} \quad (3.105)$$

Integrating by parts,

$$\begin{aligned} &= \frac{1}{4} \int_{T-R}^{T+R} (\xi - (T - R)) v^2(T - R, \xi) d\xi - \frac{1}{4} \int_{T-R}^{T+R} \int_{T-R}^\xi v^2(\eta, \xi) d\eta d\xi \\ &= \frac{1}{4} \int_0^R s \cdot v^2(s, (T - R) + s) ds + \frac{1}{4} \int_0^R \int_{T-(R-s)}^{T+(R-s)} s v^2(s, t) dt ds \\ &= Z_1^{(1)}(R, T) + Z_2^{(3)}(R, T). \end{aligned} \quad (3.106)$$

The contribution of $Z_2^{(3)}(R, T)$ may again be estimated as in (3.94) - (3.102). Now,

$$Z_1^{(1)}(R, T) = \frac{1}{4} \int_0^R s \cdot v^2(s, (T - R) + s) ds \leq R \int_{\eta=T-R} v^2(\xi, \eta) d\xi. \quad (3.107)$$

Then by (3.76) and (3.100),

$$\begin{aligned} &\int_{\mathbb{R}} \int_{\mathbb{R}} v_R(R, T) v(R, T) R Z_1^{(1)}(R, T) dR dT \\ &\leq \int \int v_R(\xi, \eta) R^2 v(\xi, \eta) \left(\int v^2(\xi, \eta = T - R) d\xi \right) d\xi d\eta \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v^2(R, T) dR dT. \end{aligned} \quad (3.108)$$

By a similar calculation,

$$\int_0^R \int_{T-(R-s)}^{T+(R-s)} s(\partial_\xi v)v dT' ds = \frac{1}{2} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta)(\partial_\xi v)v d\xi d\eta \quad (3.109)$$

$$\begin{aligned} &= \frac{1}{4} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta) \partial_\xi (v^2) d\xi d\eta \\ &= \frac{1}{4} \int_{T-R}^{T+R} (T + R - \eta) v^2(\eta, \xi = T + R) d\eta - \frac{1}{4} \int_{T-R}^{T+R} \int_\eta^{T+R} v^2(\eta, \xi) d\xi d\eta. \end{aligned} \quad (3.110)$$

Then set

$$Z_1^{(2)}(R, T) = \frac{1}{4} \int_{T-R}^{T+R} (T + R - \eta) v^2(\eta, \xi = T + R) d\eta \quad (3.111)$$

and

$$Z_2^{(4)}(R, T) = -\frac{1}{4} \int_{T-R}^{T+R} \int_\eta^{T+R} v^2(\eta, \xi) d\xi d\eta = -\frac{1}{4} \int_0^R \int_{T-(R-s)}^{T+(R-s)} v^2(s, T') dT' ds. \quad (3.112)$$

It only remains to compute

$$\frac{1}{2} \int_0^R \int_{T+(R-s)}^{T+(R-s)} s^2(\partial_\xi v(s, T'))(\partial_\eta v(s, T')) ds dT' = \frac{1}{2} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta)^2 (\partial_\xi v)(\partial_\eta v) d\xi d\eta. \quad (3.113)$$

Integrating by parts,

$$\frac{1}{2} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta)^2 (\partial_\xi v)(\partial_\eta v) d\xi d\eta = \frac{1}{2} \int_{T-R}^{T+R} (T + R - \eta)^2 v(\partial_\eta v)(T + R, \eta) d\eta \quad (3.114)$$

$$- \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta) v(\partial_\eta v) d\xi d\eta - \frac{1}{2} \int_{T-R}^{T+R} \int_\eta^{T+R} (\xi - \eta)^2 v(\partial_{\xi\eta} v) d\xi d\eta. \quad (3.115)$$

By the Cauchy - Schwartz inequality,

$$\begin{aligned}
& \frac{1}{2} \int_{T-R}^{T+R} (T+R-\eta)^2 v(\partial_\eta v)(T+R, \eta) d\eta = Z_2^{(3)}(R, T) \\
& = \frac{1}{2} \int_0^R s^2 v(s, T+R-s) (\partial_\eta v(s, T+R-s)) ds \\
& \leq \frac{R^{1/2}}{2} \left(\int s^3 (\partial_\eta v)^2(\xi = T+R, \eta) d\eta \right)^{1/2} \left(\int v(\xi = T+R, \eta)^2 d\eta \right)^{1/2}.
\end{aligned} \tag{3.116}$$

Again by (3.76) and (3.100),

$$\begin{aligned}
& \int_{\mathbb{R}} \int_{\mathbb{R}} Z_2^{(3)}(R, T) v_R(R, T) v(R, T) R dR dT \leq \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} (\xi - \eta)^{3/2} v_R(\xi, \eta) v(\xi, \eta) \\
& \leq E(v) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT.
\end{aligned} \tag{3.117}$$

Also observe that

$$\int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta) v(\partial_\eta v) d\xi d\eta = (3.105), \tag{3.118}$$

and may be computed in exactly the same manner, splitting into $Z_2^{(4)} + Z_1^{(5)}$. Now

$$\partial_{\xi\eta}^2 v = (\partial_{TT} - \partial_{RR})v. \tag{3.119}$$

Then

$$\begin{aligned}
& -\frac{1}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta)^2 v(\partial_{\xi\eta} v) d\xi d\eta = \frac{3}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta) v(\partial_R v) d\xi d\eta \\
& \quad - \frac{1}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta)^2 v F(v) d\xi d\eta.
\end{aligned} \tag{3.120}$$

Now, making a change of variables,

$$\frac{3}{2} \int_{T-R}^{T+R} \int_{\eta}^{T+R} (\xi - \eta) v(\partial_R v) d\xi d\eta = \frac{3}{2} \int_0^R \int_{T-(R+s)}^{T+(R-s)} v(\partial_s v) s ds dt, \quad (3.121)$$

and so integrating by parts,

$$\begin{aligned} &= \frac{3}{4} \int_0^R v^2(s, T+R-s) s ds + \frac{3}{4} \int_0^R v^2(s, T-R+s) s ds \\ &\quad - \frac{3}{2} \int_0^R \int_{T-(R+s)}^{T+(R-s)} v^2(s, T') ds dT' = Z_2^{(5)}(R, T) + Z_1^{(6)}(R, T). \end{aligned} \quad (3.122)$$

Next split $F(v) = F_1(v) + F_2(v)$ with

$$\begin{aligned} F_1(v) &= \frac{1}{R}(\partial_\eta r - \frac{1}{2})(\partial_\xi v) + \frac{1}{R}(\partial_\xi r + \frac{1}{2})(\partial_\eta v), \\ F_2(v) &= (e^{2Z} - 1 + \partial_\eta r + \frac{1}{2} - (\partial_\xi r - \frac{1}{2})) \frac{v}{R^2} + e^{2Z} \frac{R^2}{r^2} v^3 \zeta(Rv). \end{aligned} \quad (3.123)$$

By the radial Sobolev embedding theorem, the boundedness of $\zeta(Rv)$, and

$$\left| \partial_\xi r + \frac{1}{2} \right|, \left| \partial_\eta r - \frac{1}{2} \right| \quad \text{and} \quad |Z| \leq \epsilon(E(v)),$$

then

$$\int_0^R \int_{T-(R+s)}^{T+(R-s)} F_2(v) v s^2 ds dT' \leq \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v^2(R, T) dR dT. \quad (3.124)$$

Next, integrating by parts,

$$\int_0^R \int_{T-(R-s)}^{T+(R-s)} s(\partial_\eta r - \frac{1}{2}) v(\partial_\xi v) ds dT', \quad (3.125)$$

may be computed in a way that is very similar to (3.109), the only additional term is a term of the form

$$- \frac{1}{2} \int_0^R \int_{T-(R-s)}^{T+(R-s)} s(\partial_{\xi\eta} r) v^2 ds dT'. \quad (3.126)$$

Now since $|\partial_{\xi\eta}r| \leq \frac{\epsilon(E(v))}{r}$, this term may be bounded by

$$\int_0^R \int_{T-(R-s)}^{T+(R-s)} v^2(s, T') ds dT'. \quad (3.127)$$

A similar computation may be made for

$$\int_0^R \int_{T-(R-s)}^{T+(R-s)} s(\partial_{\xi}r - \frac{1}{2})v(\partial_{\eta}v) ds dT'. \quad (3.128)$$

Thus,

$$(3.87) \lesssim \epsilon(E(v)) \int \int v(R, T)^2 dR dT. \quad (3.129)$$

Many of the computations in (3.87) may also be used to estimate (3.88). Expanding $\zeta(z)$ for $|z| \leq \epsilon$, by the radial Sobolev embedding theorem which implies $R|v(R, T)| \leq \epsilon(E(v))$,

$$\zeta(Rv) = c_0 + c_1(Rv) + c_2(Rv)^2 + \dots \quad (3.130)$$

$$\begin{aligned} c_j \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{2+j}}{r^2} v^{3+j}(R, T) v_R(R, T) R^3 dR dT &= \frac{c_0}{4+j} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{5+j}}{r^2} \partial_R(v(R, T)^{4+j}) dR dT \\ &= -\frac{(5+j)c_j}{4+j} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{4+j}}{r^2} v(R, T)^{4+j} dR dT - \frac{c_j}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{5+j}}{r^3} (\partial_R r) v(R, T)^{4+j} dR dT. \end{aligned} \quad (3.131)$$

Therefore,

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^2}{r^2} v(R, T)^3 v_R(R, T) R^3 dR dT \lesssim \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \quad (3.132)$$

Next, we estimate

$$c_j \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{2+j}}{r^2} Z(R, T) v^{3+j}(R, T) v_R(R, T) R^3 dR dT. \quad (3.133)$$

Again splitting $Z(R, T) = Z_1(R, T) + Z_2(R, T)$,

$$\begin{aligned}
& \frac{c_0}{4+j} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{5+j}}{r^2} Z_1(R, T) v(R, T)^{3+j} \partial_R v(R, T) dR dT \\
&= \frac{c_0}{4+j} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{5+j}}{r^2} Z_1(R, T) \partial_R (v(R, T)^{4+j}) dR dT \\
&= -\frac{(5+j)c_j}{4+j} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{4+j}}{r^2} Z_1(R, T) v(R, T)^{4+j} dR dT \\
&\quad -\frac{c_j}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{5+j}}{r^3} Z_1(R, T) (\partial_R r) v(R, T)^{4+j} dR dT \\
&\quad -\frac{c_j}{4+j} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{5+j}}{r^2} (\partial_R Z_1(R, T)) v(R, T)^{4+j} dR dT.
\end{aligned} \tag{3.134}$$

Then by the radial Sobolev embedding theorem, properties of r , and the analysis of (3.90) implies that

$$(3.134) \leq \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \tag{3.135}$$

The radial Sobolev embedding theorem, properties of r , and the analysis of (3.90) also implies that

$$\frac{c_0}{4+j} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{R^{5+j}}{r^2} Z_2(R, T) v(R, T)^{3+j} \partial_R v(R, T) dR dT \leq \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \tag{3.136}$$

Therefore, we have now proved

$$\int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT \leq E(v) + \epsilon(E(v)) \int_{\mathbb{R}} \int_{\mathbb{R}} v(R, T)^2 dR dT. \tag{3.137}$$

This completes the proof of Theorem 3.3. \square

\square

Theorem 3.4. *If v is a global solution of (1.50) for $E_0 < \epsilon^2$, then*

$$v(T, x) \rightarrow \bar{v}(T, x) + o(1) \tag{3.138}$$

where \bar{v} is a solution of linearized (1.46c)

$$^{4+1}\square v = 0 - \partial_T^2 \bar{v} + \partial_R^2 \bar{v} + \frac{3}{R} \partial_R \bar{v} = 0 \quad (3.139)$$

which in turn is equivalent to linearized (1.46c)

$$-\partial_T^2 \bar{u} + \partial_R^2 \bar{u} + \frac{1}{R} \partial_R \bar{u} - \frac{1}{R^2} \bar{u} = 0 \quad (3.140)$$

Proof. Consider a large time T_0 and let $v = \bar{v} + v_S$, such that \bar{v} and v_S are solutions of

$$\left. \begin{aligned} ^{4+1}\square \bar{v} &= F(v) && \text{on } \mathbb{R}^{4+1} \\ \bar{v}_0 = \bar{v}(T_0, x) = 0 \quad \text{and} \quad \bar{v}_1 = \partial_T \bar{v}(T_0, x) = 0 &&& \text{on } \mathbb{R}^4 \end{aligned} \right\} \quad (3.141)$$

where the forcing term

$$\begin{aligned} F(v) := & \left(e^{2Z} - 1 + \left(\frac{r}{R} \partial_\eta r + \frac{1}{2} \right) - \left(\frac{r}{R} \partial_\xi r - \frac{1}{2} \right) \right) \frac{v}{r^2} + 2 \partial_\xi v \partial_\eta \log \left(\frac{r}{R} \right) \\ & + 2 \partial_\eta v \partial_\xi \log \left(\frac{r}{R} \right) + e^{2Z} \frac{R^2}{r^2} v^3 \zeta(Rv) \end{aligned} \quad (3.142)$$

and

$$\left. \begin{aligned} ^{4+1}\square v_S &= 0 && \text{on } \mathbb{R}^{4+1} \\ (v_S)_0 = v(T_0, x) \quad \text{and} \quad (v_S)_1 = \partial_T v(T_0, x) &&& \text{on } \mathbb{R}^4 \end{aligned} \right\} \quad (3.143)$$

so that $v(T) = \bar{v}(T) + S(T - T_0)(v_0, v_1)$

where $S(T)(v_0, v_1)$ is the linear evolution operator of the wave equation (3.143).

It follows from the triangle inequality that $\bar{E}(\bar{v}) \leq E(v)$, where

$$\bar{E} = \|\partial_T \bar{v}\|_{L^2(\mathbb{R}^4)}^2 + \|\nabla \bar{v}\|_{L^2(\mathbb{R}^4)}^2 + \frac{1}{2} \|\bar{v}\|_{L^4(\mathbb{R}^4)}^4 \quad (3.144)$$

Consider the energy momentum tensor corresponding to the variational principle of (3.141):

$$\check{T}(\bar{v})_{\mu\nu} = \nabla_\mu \bar{v} \nabla_\nu \bar{v} - \frac{1}{2} \check{g}_{\mu\nu} \nabla^\sigma \bar{v} \nabla_\sigma \bar{v} - \check{g}_{\mu\nu} \mathcal{F}(\bar{v}) \quad (3.145)$$

where $\bar{F}(\bar{v})$ is the primitive of $F(v)$ with respect to a variation of \bar{v} . Now consider the vector fields $\bar{J} := \mathfrak{T}$ and $J_2 := \tilde{F}(v) \mathfrak{T}^\nu$ and from the Stokes theorem, we have

$$\int_{\Sigma_{T_1}} \check{e} \bar{\mu}_q - \int_{\Sigma_{T_2}} \check{e} \bar{\mu}_q = \int \partial_T v F(v) \bar{\mu}_{\check{g}} \quad (3.146)$$

which corresponds to the instantaneous identity

$$\frac{\partial}{\partial T} \left(\frac{1}{2} \langle \partial_T \bar{v}, \partial_T \bar{v} \rangle + \frac{1}{2} \langle \nabla \bar{v}, \nabla \bar{v} \rangle \right) = \langle \partial_T \bar{v}, F(v) \rangle \quad (3.147)$$

where $\langle X, Y \rangle = \int X \cdot Y \bar{\mu}_{\check{g}}$. Now consider the terms on the right hand side of (3.147) consecutively,

1. Firstly consider the terms

$$\int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_{\eta} r + \frac{1}{2}) \partial_{\xi} v \partial_T \bar{v} \bar{\mu}_{\check{g}} \text{ and } \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_{\xi} r - \frac{1}{2}) \partial_{\eta} v \partial_T \bar{v} \bar{\mu}_{\check{g}} :$$

we have

$$\int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_{\eta} r + \frac{1}{2}) \partial_{\xi} v \partial_T \bar{v} R^3 dR dT \quad (3.148)$$

$$= \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_{\eta} r + \frac{1}{2}) \partial_{\xi} v \partial_T \bar{v} R^3 d\xi \eta \quad (3.149)$$

$$\leq \int_{\eta_0} (\sup_{R>0} \frac{1}{R} \int_0^R f^2(v) s ds) (\int (dv)^2 d\xi) d\eta \quad (3.150)$$

going back to (T, R, θ)

$$\leq E(v) \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}} \quad (3.151)$$

analogously, we have

$$\int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_{\xi} r - \frac{1}{2}) (\partial_{\eta} v) (\partial_T v) \bar{\mu}_{\check{g}} \leq E(v) \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}} \quad (3.152)$$

2. Next, the terms

$$\int_{T_0}^{\infty} \int_{\mathbb{R}^4} (\partial_{\eta} r + \frac{1}{2}) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\check{g}} \text{ and } \int_{T_0}^{\infty} \int_{\mathbb{R}^4} (\partial_{\xi} r - \frac{1}{2}) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\check{g}}$$

using Cauchy-Schwarz

$$\begin{aligned}
& \int_{T_0}^{\infty} \int_{\mathbb{R}^4} (\partial_{\eta} r + \frac{1}{2}) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\check{g}} \\
& \leq \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_{\eta} r + \frac{1}{2}) (\partial_T \bar{v})^2 \bar{\mu}_{\check{g}} + \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R^3} (\partial_{\eta} r + \frac{1}{2}) v^2 \bar{\mu}_{\check{g}} \\
& \leq E(v) \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}} + \epsilon \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}}
\end{aligned} \tag{3.153}$$

likewise, we have

$$\int_{T_0}^{\infty} \int_{\mathbb{R}^4} (\partial_{\xi} r - \frac{1}{2}) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\check{g}} \leq E(v) \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}} + \epsilon \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}}$$

3. Controlling the term

$$\int_{T_0}^{\infty} \int_{\mathbb{R}^4} (e^{2Z} - 1) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\check{g}} \tag{3.154}$$

is similar,

$$\begin{aligned}
\int_{T_0}^{\infty} \int_{\mathbb{R}^4} (e^{2Z} - 1) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\check{g}} & \leq \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (e^{2Z} - 1) \bar{\mu}_{\check{g}} + \int_{T_0}^{\infty} \int_{\mathbb{R}^4} (e^{2Z} - 1) \frac{v^2}{R^3} \bar{\mu}_{\check{g}} \\
& \leq E(v) \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}} + \epsilon \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}}.
\end{aligned} \tag{3.155}$$

In view of the fact that

$$\lim_{T \rightarrow \infty} \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{v^2}{R^3} \bar{\mu}_{\check{g}} = 0$$

we now have

$$\lim_{T_0 \rightarrow \infty} \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_\eta r + \frac{1}{2}) \partial_\xi v \partial_T \bar{v} \bar{\mu}_{\tilde{g}} = 0 \quad (3.156a)$$

$$\lim_{T_0 \rightarrow \infty} \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_\xi r - \frac{1}{2}) \partial_\eta v \partial_T \bar{v} \bar{\mu}_{\tilde{g}} = 0 \quad (3.156b)$$

$$\lim_{T_0 \rightarrow \infty} \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_\eta r + \frac{1}{2}) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\tilde{g}} = 0 \quad (3.156c)$$

$$\lim_{T_0 \rightarrow \infty} \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{1}{R} (\partial_\xi r - \frac{1}{2}) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\tilde{g}} = 0 \quad (3.156d)$$

$$\lim_{T_0 \rightarrow \infty} \int_{T_0}^{\infty} \int_{\mathbb{R}^4} (e^{2Z} - 1) \frac{v}{R^2} \partial_T \bar{v} \bar{\mu}_{\tilde{g}} = 0 \quad (3.156e)$$

from the dominated convergence theorem.

4. Now consider the term

$$\int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{R^2}{r^2} e^{2Z} v^3 \zeta(Rv) \partial_T \bar{v} \bar{\mu}_{\tilde{g}}$$

again expand out $\zeta(R\tilde{v})$:

$$\zeta(Rv) = c_0 + c_1 Rv + c_2 (Rv)^2 + \dots \quad (3.157)$$

Then the leading order term of

$$c_j \int_{T_0}^{\infty} \int_{\mathbb{R}^4} \frac{R^{2+j}}{r^2} e^{2Z} v^{3+j} \partial_T \bar{v} \bar{\mu}_{\tilde{g}} \quad (3.158)$$

transforms as

$$c_0 \int \frac{R^2}{r^2} \bar{v}^3 \bar{v}_T R^3 \bar{\mu}_{\tilde{g}} = \left(-\frac{5c_0}{4} \int \frac{R^2}{r^2} \bar{v}^4 \bar{\mu}_{\tilde{g}} \right) - \frac{c_0}{2} \int \frac{R^2}{r^3} \bar{v}^4 (\partial_T r) \bar{\mu}_{\tilde{g}} \quad (3.159)$$

Now by the radial Sobolev embedding theorem, combined with the Morawetz estimates,

$$\int_{T \geq T_0} \int \frac{R^2}{r^3} \bar{v}^4 (\partial_T r) \bar{\mu}_{\tilde{g}} \rightarrow 0 \quad (3.160)$$

as $T_0 \rightarrow \infty$. Next, using the radial Strichartz estimate

$$\left\| |x|^{1/2} S(t)((v_S)_0, (v_S)_1) \right\|_{L_t^2 L_x^\infty} \leq \|(v_S)_0\|_{\dot{H}^1} + \|(v_S)_1\|_{L^2}, \quad (3.161)$$

so

$$\begin{aligned} \int_{T \geq T_0} \int \frac{R^2}{r^2} v^2 v_S \bar{v}_T R^3 dR dT &\leq \left(\int_{T \geq T_0} \int v^2 dR dT \right)^{1/2} \\ &\quad \cdot \|R^{1/2} v_S\|_{L_T^2 L_x^\infty} \|\bar{v}_T\|_{L_T^\infty L_x^2} \|Rv\|_{L_{T,x}^\infty}. \end{aligned} \quad (3.162)$$

If

$$E(\bar{v}(T)) = \|\bar{v}_T\|_{L^2}^2 + \|\nabla \bar{v}(T)\|_{L^2}^2 + \frac{1}{2} \|\bar{v}(T)\|_{L^4}^4, \quad (3.163)$$

then the Sobolev embedding theorem implies that for small energy, the RHS of eq. 3.162

$$\leq \left(\sup_{T \geq T_0} E(\bar{v}(T)) \right)^{1/2} E(v) \left(\int_{T \geq T_0} \int v^2 dR dT \right)^{1/2}. \quad (3.164)$$

Also,

$$\begin{aligned} \int_{T \geq T_0} \int \frac{R^2}{r^2} v \cdot v_S \cdot \bar{v} \bar{v}_T R^3 dR dT &\leq \left(\int_{T \geq T_0} \int v^2 dR dT \right)^{1/2} \\ &\quad \cdot \|R^{1/2} v_S\|_{L_T^2 L_x^\infty} \|\bar{v}_T\|_{L_T^\infty L_x^2} \|R\bar{v}\|_{L_{T,x}^\infty} \\ &\leq E(v) \left(\sup_{T \geq T_0} E(\bar{v}(T)) \right). \end{aligned} \quad (3.165)$$

Finally,

$$\begin{aligned}
\int_{T \geq T_0} \int \frac{R^2}{r^2} v_S \bar{v}^2 \bar{v}_T R^3 dR dT &\leq \left\| |x|^{1/2} v_S \right\|_{L_T^2 L_x^\infty} \|v_T\|_{L_T^\infty L_x^2} \\
&\quad \cdot \left(\int v + (v_S)^2 dR dT \right)^{1/2} \| |x| \bar{v} \|_{L_{T,x}^\infty} \\
&\leq E(v) \left(\sup_{T \geq T_0} E(\bar{v})(T) \right).
\end{aligned} \tag{3.166}$$

Now consider,

$$c_j \int_{T_0} \int_{\mathbb{R}^4} \frac{R^{2+j}}{r^2} Z(R, T) v^{3+j} \partial_T \bar{v} \bar{\mu}_{\bar{g}} \tag{3.167}$$

Analogously, we have

$$c_j \int_{T_0} \int_{\mathbb{R}^4} \frac{R^{2+j}}{r^2} Z(R, T) v^{3+j} \partial_T \bar{v} \bar{\mu}_{\bar{g}} \leq c(E_0) \int v^2 dR dT \tag{3.168}$$

□

We would now like to remark on the decay rates for our scattering problem in comparison with the decay rates for the linear wave equation.

Remark 3.5. Suppose $u : \mathbb{R}^{2+1} \rightarrow \mathbb{R}$ satisfies the linear wave equation

$$\left. \begin{aligned}
{}^{2+1}\square v &= 0 && \text{on } \mathbb{R}^{2+1} \\
u_0 = u(0, x) \text{ and } u_1 = \partial_T u(0, x) &&& \text{on } \mathbb{R}^2
\end{aligned} \right\} \tag{3.169}$$

such that the smooth initial data (u_0, u_1) is compactly supported or sufficiently rapidly decaying, then u admits the decay rate:

$$|u| \leq (1 + \xi)^{-1/2} (1 + \eta)^{-1/2}. \tag{3.170}$$

This decay rate has to be contrasted with the decay rate from the (standard) energy estimates:

$$\|u\| \leq (1+t)^{n-1/2}(1+|t-x|)^{1/2}. \quad (3.171)$$

In the following we shall prove that this decay rate is also consistent with the decay rate for the fully coupled 2+1 equivariant Einstein-wave map system.

Corollary 3.6. *Suppose (M, g, U) is the smooth, globally hyperbolic, geodesically complete maximal development of the smooth, asymptotically flat initial data $(\Sigma_0, q_0, k_0, U_0)$, with U_0 compactly supported, then u admits the decay rate*

$$|u| \leq (1+\xi)^{-1/2}(1+\eta)^{-1/2} \quad (3.172)$$

in the temporal asymptotic region $(T \rightarrow \infty)$ and

$$|u| \leq \mathcal{O}(R^{-1/2}) \quad (3.173)$$

as $R \rightarrow \infty$ along the $\eta = \text{const.}$ hypersurfaces.

Proof. Now let us turn to the scattering for our problem, we have

$$\|v - v_S\| \rightarrow 0$$

this implies

$$\|u - \bar{u}\|_{L^\infty} \rightarrow 0, \quad T \rightarrow \infty, \text{ where } \bar{u} = r\bar{v}.$$

Note that if \bar{v} satisfies $\square v = 0$ then \bar{u} satisfies

$$\square u_S - \frac{u_S}{R^2} = 0. \quad (3.174)$$

Following [7], if we introduce the following ‘projection’ formulas

$$\begin{aligned} u_1 &= u_S \cos \theta \\ u_2 &= u_S \sin \theta \end{aligned} \quad (3.175)$$

so that $u = u_1 \cos \theta + u_2 \sin \theta$ and

$${}^{2+1}\square u_1 = 0, \quad \mathbb{R}^{2+1} \quad (3.176)$$

$${}^{2+1}\square u_2 = 0, \quad \mathbb{R}^{2+1} \quad (3.177)$$

If we turn to the 2+1 decay rates, we have

$$\begin{aligned} u_1 &\leq (1 + \xi)^{-1/2}(1 + \eta)^{-1/2} \\ u_2 &\leq (1 + \xi)^{-1/2}(1 + \eta)^{-1/2} \end{aligned} \quad (3.178)$$

which is view of the previous equations gives

$$u_S \leq (1 + \xi)^{-1/2}(1 + \eta)^{-1/2} \quad (3.179)$$

□

References

- [1] L. Andersson, N. Gudapati, and J. Szeftel, *Global regularity for 2+1 dimensional Einstein-wave map system*, Ann. PDE **3** (2017), no. 13.
- [2] A. Ashtekar, J. Bicak, and B. Schmidt, *Behaviour of Einstein-Rosen waves at null infinity*, Phys. Rev. D. **55** (1997), no. 2, 687–694.
- [3] A. Ashtekar and M. Varadarajan, *Striking property of the gravitational Hamiltonian*, Phys. Rev. D **50** (1994), no. 8, 4944–4956.
- [4] Y. Choquet-Bruhat and R. Geroch, *Global aspects of the Cauchy problem in general relativity*, Commun. math. Phys. **14** (1969), 329–335.
- [5] D. Christodoulou and S. Klainerman, *The global nonlinear stability of the minkowski space (pms-41)*, Princeton University Press (1994).
- [6] D. Christodoulou and A.S. Tahvildar-Zadeh, *On the asymptotic behavior of spherically symmetric wave maps*, Duke Math. J. **71** (1993), no. 1, 31–69.
- [7] ———, *On the regularity of spherically symmetric wave maps*, Comm. Pure Appl. Math. **46** (1993), no. 7, 1041–1091.
- [8] B. Dodson and N. Gudapati, *On scattering for small data of 2+1 dimensional equivariant Einstein-wave map system*, Ann. Henri. Poincaré **18** (2017), no. 9, 3097–3142.

- [9] M. Grillakis, *Classical solutions for the equivariant wave map in 1+2 dimensions*, preprint (1991).
- [10] N. Gudapati, *The Cauchy problem for energy critical self-gravitating wave maps*, Dissertation (FU Berlin) (2013).
- [11] ———, *The Cauchy problem for energy critical self-gravitating wave maps*, Dissertation (FU Berlin) (2013).
- [12] C. Huneau, *Stability in exponential time of Minkowski space–time with a translation space-like Killing field*, Ann. PDE **2** (2016), no. 7.
- [13] J. Krieger, *Global regularity of wave maps from \mathbb{R}^{2+1} to \mathbb{H}^2* , Comm. Math. Phys. **250** (2004), no. 4, 507–580.
- [14] J. Krieger and W. Schlag, *Concentration compactness for critical wave maps*, AMS Monographs in Mathematics (2012).
- [15] H. Lindblad, *On the asymptotic behavior of solutions of einstein vacuum equations in wave coordinates*, Commun. Math. Phys. **353** (2017), no. 1, 135–184.
- [16] H. Lindblad and I. Rodnianski, *The global stability of Minkowski space-time in harmonic gauge*, Ann. Math. **171** (2010), no. 3, 1401–1477.
- [17] J. Shatah and A.S. Tahvildar-Zadeh, *Regularity of harmonic maps from the Minkowski space into rotationally symmetric manifolds*, Comm. Pure Appl. Math. **45** (1992), 1041–1091.
- [18] ———, *Cauchy problem for equivariant wave maps*, Comm. Pure Appl. Math. **47** (1994), no. 5, 719–754.
- [19] J. Sterbenz and D. Tataru, *Energy dispersed large data wave maps in 2+1 dimensions*, Commun. Math. Phys. **298** (2010), 139–230.
- [20] ———, *Regularity of wave maps in dimension 2+1*, Commun. Math. Phys. **298** (2010), 231–264.

- [21] T. Tao, *Global regularity of wave maps i to viii*, arXiv (2001-09).
- [22] D. Tataru, *Local and global results for wave maps I*, Comm. PDE **23** (1998), no. 9-10, 1781–1793.
- [23] ———, *On global existence and scattering for the wave maps equation*, Amer. J. Math. **123** (2001), no. 3, 385–423.
- [24] ———, *Rough solutions for the wave maps equation*, Amer. J. Math. **127** (2005), no. 2, 293–377.