

GENERALIZED NUMERICAL RADIUS INEQUALITIES FOR SCHATTEN P-NORMS

JANA HAMZA AND HASSAN ISSA

ABSTRACT. In this paper, we present various inequalities for generalized numerical radius of 2×2 block matrices for Schatten p-norm. Moreover, we give a refinement of the triangle inequality for the Schatten p-generalized numerical radius.

1. INTRODUCTION

Consider the space $B(H)$ of bounded linear operators over a Hilbert space H . For $A \in B(H)$, the numerical radius, the usual operator norm, and the Schatten p-norm, are denoted by $\omega(A)$, $\|A\|$ and $\|A\|_p$ respectively.

A norm $\|\cdot\|$ on $B(H)$ is said to be unitarily invariant if $\|UAV\| = \|A\|$, where $A \in B(H)$ and $U, V \in B(H)$ being unitary, and weakly unitarily invariant if $\|UAU^*\| = \|A\|$ where $A \in B(H)$ and $U \in B(H)$ being unitary. It is known that $\|A\|_p$ is unitarily invariant.

We should note that if $B(H)$ is the space of $n \times n$ complex matrices, $M_n(\mathbb{C})$, then for $A, B \in M_n(\mathbb{C})$ we have

$$\|A \oplus A^*\| = \|A \oplus A\| \quad (1.1)$$

and

$$\|A \oplus B\| = \left\| \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right\| \quad (\text{see [11]}). \quad (1.2)$$

Moreover, we have

$$\|A \oplus B\|_p = (\|A\|_p^p + \|B\|_p^p)^{\frac{1}{p}}, \quad (1.3)$$

2010 *Mathematics Subject Classification.* 15B57.

Key words and phrases. numerical radius, Schatten p- norm, generalized Schatten p-numerical radius, inequality.

and

$$\|A \oplus A\|_p = 2^{\frac{1}{p}} \|A\|_p. \quad (1.4)$$

The numerical radius $\omega(\cdot)$ is defined by $\omega(A) = \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta} A)\|$, where $A \in B(H)$. Due to its importance, the numerical radius has been generalized several times, and their last was given by Abu-Omar and Kittaneh [2] in 2019, in which they generalized it on $B(H)$. This generalization is denoted by $\omega_N(\cdot)$, and is defined by $\omega_N(A) = \sup_{\theta \in \mathbb{R}} N(Re(e^{i\theta} A))$. It was proved by the authors in [2] that $\omega_N(\cdot)$ generalizes the numerical radius $\omega(\cdot)$ if N is the usual operator norm.

In this paper, we are interested in studying the space of $n \times n$ complex matrices, $M_n(\mathbb{C})$. We should note that $\omega_N(\cdot)$ has the following two important properties, see [1].

Property 1.1. *The following properties hold:*

- a) *The norm $\omega_N(\cdot)$ is self adjoint.*
- b) *If the norm $N(\cdot)$ is weakly unitarily invariant, then so is $\omega_N(\cdot)$.*

In [6], Bhatia and Kittaneh were able to prove the following theorem that relates the Schatten p-norm of an $n \times n$ block matrix by that of its block entries.

Theorem 1.2. *Let $T \in M_n(\mathbb{C})$ such that $T = [T_{ij}]$, $1 \leq i, j \leq n$ and $1 \leq p \leq \infty$ then*

$$n^{2-p} \|T\|_p^p \leq \sum_{i,j=1}^n \|T_{ij}\|_p^p \leq \|T\|_p^p, \quad (1.5)$$

for $2 \leq p \leq \infty$; and

$$\|T\|_p^p \leq \sum_{i,j=1}^n \|T_{ij}\|_p^p \leq n^{2-p} \|T\|_p^p, \quad (1.6)$$

for $1 \leq p \leq 2$.

Motivated by the results of Bhatia and Kittaneh in [6], and those of Aldalabih and Kittaneh in [3], we aim in this paper to prove some generalized numerical radius inequalities for partitioned general 2×2 block matrices considering the case when N is taken to be the Schatten p-norm. We denote this norm by $\omega_p(\cdot)$ and call it the Schatten p-generalized numerical radius. We emphasize on finding such inequalities

for the off-diagonal part of block matrices. We also provide an application of this norm in which we give a refinement of the triangle inequality for the Schatten p-generalized numerical radius. The following Lemma was proved by the authors in [1], and will be used in our work.

Lemma 1.3. *Let $A, B \in M_n(\mathbb{C})$, then*

$$\omega_p \left(\begin{bmatrix} 0 & B \\ B & 0 \end{bmatrix} \right) = 2^{\frac{1}{p}} \omega_p(B)$$

for all p .

2. GENERAL 2×2 BLOCK MATRICES INEQUALITIES

In this section we give bounds for the generalized Schatten p-numerical radius of general 2×2 block matrices. We give emphasis for 2×2 block diagonal matrices. Most of the results in this section, generalize those presented in [3].

Lemma 2.1. *Let $A, B \in M_n(\mathbb{C})$ then for all p , we have*

$$\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) = 2^{\frac{1}{p}-1} \sup_{\theta \in \mathbb{R}} \|e^{i\theta} A + e^{-i\theta} B^*\|_p.$$

Proof. By equations (1.1), (1.2) and (1.4), we have

$$\begin{aligned} \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \frac{1}{2} \sup_{\theta \in \mathbb{R}} \left\| \begin{bmatrix} 0 & e^{i\theta} A + e^{-i\theta} B^* \\ e^{i\theta} B + e^{-i\theta} A^* & 0 \end{bmatrix} \right\|_p \\ &= \frac{1}{2} \sup_{\theta \in \mathbb{R}} \left\| \begin{bmatrix} e^{i\theta} A + e^{-i\theta} B^* & 0 \\ 0 & (e^{i\theta} A + e^{-i\theta} B^*)^* \end{bmatrix} \right\|_p \\ &= \frac{1}{2} \sup_{\theta \in \mathbb{R}} \left\| \begin{bmatrix} e^{i\theta} A + e^{-i\theta} B^* & 0 \\ 0 & e^{i\theta} A + e^{-i\theta} B^* \end{bmatrix} \right\|_p \\ &= \frac{1}{2} \sup_{\theta \in \mathbb{R}} 2^{\frac{1}{p}} \|e^{i\theta} A + e^{-i\theta} B^*\|_p \\ &= 2^{\frac{1}{p}-1} \sup_{\theta \in \mathbb{R}} \|e^{i\theta} A + e^{-i\theta} B^*\|_p. \end{aligned}$$

□

Proposition 2.2. *Let $A, B \in M_n(\mathbb{C})$, then the following inequality holds for all p*

$$\omega_p \left(\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \right) \leq (\omega_p^p(A) + \omega_p^p(B))^{\frac{1}{p}}.$$

Proof. We have

$$\begin{aligned}
\omega_p \left(\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \right) &= \sup_{\theta \in \mathbb{R}} \left\| \operatorname{Re} \left(e^{i\theta} \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \right) \right\|_p \\
&= \sup_{\theta \in \mathbb{R}} \left\| \begin{bmatrix} \operatorname{Re}(e^{i\theta} A) & 0 \\ 0 & \operatorname{Re}(e^{i\theta} B) \end{bmatrix} \right\|_p \\
&= \sup_{\theta \in \mathbb{R}} \left(\left\| \operatorname{Re}(e^{i\theta} A) \right\|_p^p + \left\| \operatorname{Re}(e^{i\theta} B) \right\|_p^p \right)^{\frac{1}{p}} \quad (\text{by equation (1.3)}) \\
&\leq (\omega_p^p(A) + \omega_p^p(B))^{\frac{1}{p}}
\end{aligned}$$

as required. $\square$

Theorem 2.3. *Let $A = [A_{ij}]$ be a 2×2 block matrix, then*

$$\omega_p^p(A) \leq \frac{1}{2^{p-2}} \sum_{i,j=1}^2 \omega_p^p(a_{ij}) \quad (2.1)$$

for $2 \leq p \leq \infty$, and

$$\omega_p^p(A) \leq \sum_{i,j=1}^2 \omega_p^p(a_{ij}) \quad (2.2)$$

for $1 \leq p \leq 2$, where

$$a_{ij} = \begin{cases} A_{ij} & i = j \\ 2^{-\frac{1}{p}} \begin{bmatrix} 0 & A_{ij} \\ A_{ji} & 0 \end{bmatrix} & i \neq j. \end{cases}$$

Proof. We have

$$\left\| \operatorname{Re}(e^{i\theta} A) \right\|_p = \left\| \begin{bmatrix} \operatorname{Re}(e^{i\theta} A_{11}) & \frac{1}{2}(e^{i\theta} A_{12} + e^{-i\theta} A_{21}^*) \\ \frac{1}{2}(e^{i\theta} A_{21} + e^{-i\theta} A_{12}^*) & \operatorname{Re}(e^{i\theta} A_{22}) \end{bmatrix} \right\|_p,$$

then by inequality (1.5) and Lemma 2.1, we get

$$\begin{aligned}
\|Re(e^{i\theta} A)\|_p^p &\leq \frac{1}{2^{p-2}} \sum_{i,j=1}^2 \|(Re(e^{i\theta} A))_{ij}\|_p^p \\
&= \frac{1}{2^{p-2}} \left(\sum_{i=j} \|(Re(e^{i\theta} A_{ij}))\|_p^p + \sum_{i \neq j} \left(\frac{1}{2} \|e^{i\theta} A_{ij} + e^{-i\theta} A_{ji}^*\|_p \right)^p \right) \\
&\leq \frac{1}{2^{p-2}} \left(\sum_{i=j} \omega_p^p(A_{ij}) + \sum_{i \neq j} \left(2^{-\frac{1}{p}} \omega_p \left(\begin{bmatrix} 0 & A_{ij} \\ A_{ji} & 0 \end{bmatrix} \right) \right)^p \right) \\
&= \frac{1}{2^{p-2}} \sum_{i=j} \omega_p^p(a_{ij}).
\end{aligned}$$

Then

$$\begin{aligned}
\omega_p^p(A) &= \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta} A)\|_p^p \\
&\leq \frac{1}{2^{p-2}} \sum_{i=j} \omega_p^p(a_{ij})
\end{aligned}$$

for $2 \leq p \leq \infty$. The second inequality is proved in a similar manner, using Lemma 2.1, and inequality (1.6). $\square$

3. OFF-DIAGONAL 2×2 BLOCK MATRICES INEQUALITIES

In this section, our interest was finding inequalities for ω_p of the off-diagonal 2×2 block matrices. The following lemma is useful in our work.

Lemma 3.1. *Let $A, B \in M_n(\mathbb{C})$, then*

$$\begin{aligned}
\text{a) } \omega_p \left(\begin{bmatrix} 0 & A \\ e^{i\theta} B & 0 \end{bmatrix} \right) &= \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \text{ for all } \theta \in \mathbb{R}. \\
\text{b) } \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(\begin{bmatrix} 0 & B \\ A & 0 \end{bmatrix} \right).
\end{aligned}$$

for all p .

Proof. Let $U = \begin{bmatrix} I & 0 \\ 0 & e^{i\frac{\theta}{2}}I \end{bmatrix}$, then U is unitary. Then by Property 1.1 we have

$$\begin{aligned} \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(U \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} U^* \right) \\ &= \omega_p \left(e^{-i\frac{\theta}{2}} \begin{bmatrix} 0 & A \\ e^{i\theta}B & 0 \end{bmatrix} \right) \\ &= \omega_p \left(\begin{bmatrix} 0 & A \\ e^{i\theta}B & 0 \end{bmatrix} \right) \end{aligned}$$

which ends the proof of (a). Now to prove the equality (b), consider $U = \begin{bmatrix} 0 & I \\ I & 0 \end{bmatrix}$, then U is unitary.

Then by Property 1.1 we have

$$\begin{aligned} \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(U \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} U^* \right) \\ &= \omega_p \left(\begin{bmatrix} 0 & B \\ A & 0 \end{bmatrix} \right) \end{aligned}$$

which ends the proof. $\square$

The next theorem gives upper and lower bounds for $\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right)$ in terms of $\omega_p(A+B)$ and $\omega_p(A-B)$.

Theorem 3.2. *Let $A, B \in M_n(\mathbb{C})$, then*

$$\frac{\max(\omega_p(A+B), \omega_p(A-B))}{2^{1-\frac{1}{p}}} \leq \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq \frac{\omega_p(A+B) + \omega_p(A-B)}{2^{1-\frac{1}{p}}}$$

for all p .

Proof. By Lemma 1.3, we have

$$\begin{aligned}
2^{\frac{1}{p}}\omega_p(A+B) &= \omega_p \left(\begin{bmatrix} 0 & A+B \\ A+B & 0 \end{bmatrix} \right) \\
&= \omega_p \left(\begin{bmatrix} 0 & B \\ A & 0 \end{bmatrix} + \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \\
&\leq \omega_p \left(\begin{bmatrix} 0 & B \\ A & 0 \end{bmatrix} \right) + \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \quad (\text{by triangle inequality}). \\
&= 2\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \quad (\text{by Lemma 3.1}),
\end{aligned}$$

then

$$\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \geq \frac{1}{2^{1-\frac{1}{p}}} \omega_p(A+B), \quad (3.1)$$

replacing B by $-B$ in inequality (3.1), we get

$$\omega_p \left(\begin{bmatrix} 0 & A \\ -B & 0 \end{bmatrix} \right) \geq \frac{1}{2^{1-\frac{1}{p}}} \omega_p(A-B),$$

taking $\theta = \pi$ in Lemma 3.1, we get

$$\begin{aligned}
\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(\begin{bmatrix} 0 & A \\ -B & 0 \end{bmatrix} \right) \\
&\geq \frac{1}{2^{1-\frac{1}{p}}} \omega_p(A-B),
\end{aligned} \quad (3.2)$$

therefore, by the estimations (3.1) and (3.2), we get

$$\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \geq \frac{\max(\omega_p(A+B), \omega_p(A-B))}{2^{1-\frac{1}{p}}}.$$

Now for the second inequality, consider $U = \frac{1}{\sqrt{2}} \begin{bmatrix} I & -I \\ I & I \end{bmatrix}$, where I is the $n \times n$ identity matrix, then U is unitary, and thus by Property 1.1, we get

$$\begin{aligned}
\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(U \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} U^* \right) \\
&= \frac{1}{2} \omega_p \left(\begin{bmatrix} -(A+B) & A-B \\ -(A-B) & A+B \end{bmatrix} \right) \\
&\leq \frac{1}{2} \left(\omega_p \left(\begin{bmatrix} -(A+B) & 0 \\ 0 & A+B \end{bmatrix} \right) + \omega_p \left(\begin{bmatrix} 0 & A-B \\ -(A-B) & 0 \end{bmatrix} \right) \right) \\
&\quad (\text{by triangle inequality}) \\
&\leq \frac{1}{2} \left(2^{\frac{1}{p}} \omega_p(A+B) + 2^{\frac{1}{p}} \omega_p(A-B) \right) \\
&\quad (\text{by Proposition 2.2, Lemma 3.1 and Lemma 1.3}).
\end{aligned}$$

as required. $\square$

Corollary 3.3. *Let $T \in M_n(\mathbb{C})$ such that $T = A + iB$, where $A = \text{Re}(T)$ and $B = \text{Im}(T)$, then*

$$\frac{\omega_p(T)}{2} \leq \frac{1}{2^{\frac{1}{p}}} \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq \omega_p(T).$$

for all p .

Proof. Replacing B by iB in theorem 3.2, we get

$$\frac{\max(\omega_p(A + iB), \omega_p(A - iB))}{2^{1-\frac{1}{p}}} \leq \omega_p \left(\begin{bmatrix} 0 & A \\ iB & 0 \end{bmatrix} \right) \leq \frac{\omega_p(A + iB) + \omega_p(A - iB)}{2^{1-\frac{1}{p}}},$$

then,

$$\frac{\max(\omega_p(T), \omega_p(T^*))}{2^{1-\frac{1}{p}}} \leq \omega_p \left(\begin{bmatrix} 0 & A \\ iB & 0 \end{bmatrix} \right) \leq \frac{\omega_p(T) + \omega_p(T^*)}{2^{1-\frac{1}{p}}}.$$

However, $\omega_p(T) = \omega_p(T^*)$, then

$$\frac{\omega_p(T)}{2^{1-\frac{1}{p}}} \leq \omega_p \left(\begin{bmatrix} 0 & A \\ iB & 0 \end{bmatrix} \right) \leq \frac{\omega_p(T)}{2^{-\frac{1}{p}}}.$$

Take $\theta = \frac{\pi}{2}$ in Lemma 3.1, then

$$\frac{\omega_p(T)}{2^{1-\frac{1}{p}}} \leq \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq \frac{\omega_p(T)}{2^{-\frac{1}{p}}}. \quad (3.3)$$

Multiply (3.3) by $2^{-\frac{1}{p}}$, then

$$\frac{\omega_p(T)}{2} \leq \frac{1}{2^{\frac{1}{p}}} \omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq \omega_p(T).$$

□

Remark 3.1. Let $A \in M_n(\mathbb{C})$, then we have

$$e^{i(\theta-\frac{\pi}{2})} = -ie^{i\theta}, \text{ and } e^{-i(\theta-\frac{\pi}{2})} = ie^{-i\theta}.$$

Therefore,

$$\begin{aligned} \operatorname{Re}(e^{i(\theta-\frac{\pi}{2})} A) &= \frac{e^{i(\theta-\frac{\pi}{2})} A + e^{-i(\theta-\frac{\pi}{2})} A^*}{2} \\ &= \frac{-ie^{i\theta} A + ie^{-i\theta} A^*}{2} \\ &= \frac{e^{i\theta} A - e^{-i\theta} A^*}{2i} \\ &= \operatorname{Im}(e^{i\theta} A). \end{aligned}$$

Therefore,

$$\begin{aligned} \omega_p(A) &= \sup_{\alpha \in \mathbb{R}} \left\| \operatorname{Re}(e^{i\alpha} A) \right\|_p \\ &= \sup_{\theta \in \mathbb{R}} \left\| \operatorname{Re}(e^{i(\theta-\frac{\pi}{2})} A) \right\|_p \\ &= \sup_{\theta \in \mathbb{R}} \left\| \operatorname{Im}(e^{i\theta} A) \right\|_p. \end{aligned}$$

See [8]

Remark 3.2. Let $X \in M_n(\mathbb{C})$, and $2 \leq p < \infty$ then

$$\begin{aligned}
\omega_p \left(\begin{bmatrix} X & X \\ -X & -X \end{bmatrix} \right) &= \sup_{\theta \in \mathbb{R}} \left\| \begin{bmatrix} \operatorname{Re}(e^{i\theta} X) & \operatorname{Im}(e^{i\theta} X) \\ -\operatorname{Im}(e^{i\theta} X) & -\operatorname{Re}(e^{i\theta} X) \end{bmatrix} \right\|_p \\
&\leq \frac{1}{2^{\frac{2}{p}-1}} \sup_{\theta \in \mathbb{R}} \left(2 \|\operatorname{Re}(e^{i\theta} X)\|_p^p + 2 \|\operatorname{Im}(e^{i\theta} X)\|_p^p \right)^{\frac{1}{p}} \\
&\quad (\text{by inequality (1.5)}) \\
&= \frac{1}{2^{\frac{2}{p}-1}} 2^{\frac{2}{p}} \omega_p(X) \quad (\text{by Remark 3.1}) \\
&= 2\omega_p(X).
\end{aligned}$$

Theorem 3.4. Let $A, B \in M_n(\mathbb{C})$, then

$$\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq 2^{\frac{1}{p}} \min(\omega_p(A), \omega_p(B)) + \min(\omega_p(A+B), \omega_p(A-B)).$$

for $2 \leq p < \infty$.

Proof. Consider $U = \frac{1}{\sqrt{2}} \begin{bmatrix} I & I \\ -I & I \end{bmatrix}$, where I is the $n \times n$ identity matrix, then U is unitary, we have

$$\begin{aligned}
\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(U \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} U^* \right) \\
&= \frac{1}{2} \omega_p \left(\begin{bmatrix} A+B & A-B \\ -(A-B) & -(A+B) \end{bmatrix} \right) \\
&= \frac{1}{2} \omega_p \left(\begin{bmatrix} A+B & A+B \\ -(A+B) & -(A+B) \end{bmatrix} + \begin{bmatrix} 0 & -2B \\ 2B & 0 \end{bmatrix} \right) \\
&\leq \frac{1}{2} \left(\omega_p \left(\begin{bmatrix} A+B & A+B \\ -(A+B) & -(A+B) \end{bmatrix} \right) + \omega_p \left(\begin{bmatrix} 0 & -2B \\ 2B & 0 \end{bmatrix} \right) \right).
\end{aligned}$$

$$\begin{aligned}
& \text{(by triangle inequality)} \\
& \leq \frac{1}{2} \left(2\omega_p(A+B) + 2^{\frac{1}{p}}\omega_p(2B) \right) \text{ (by Remark 3.2, Lemma 3.1 and Lemma 1.3)} \\
& = \omega_p(A+B) + 2^{\frac{1}{p}}\omega_p(B).
\end{aligned} \tag{3.4}$$

Replacing B by $-B$ in (3.4), we get

$$\begin{aligned}
\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(\begin{bmatrix} 0 & A \\ -B & 0 \end{bmatrix} \right) \quad \text{(taking } \theta = \pi \text{ in Lemma 3.1)} \\
&\leq \omega_p(A-B) + 2^{\frac{1}{p}}\omega_p(B).
\end{aligned} \tag{3.5}$$

Then by (3.4) and (3.5), we get

$$\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq 2^{\frac{1}{p}}\omega_p(B) + \min(\omega_p(A+B), \omega_p(A-B)). \tag{3.6}$$

Interchanging A and B in (3.6), we get

$$\begin{aligned}
\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) &= \omega_p \left(\begin{bmatrix} 0 & B \\ A & 0 \end{bmatrix} \right) \quad \text{(by Lemma 3.1)} \\
&\leq 2^{\frac{1}{p}}\omega_p(A) + \min(\omega_p(A+B), \omega_p(A-B)).
\end{aligned} \tag{3.7}$$

Therefore, by (3.6) and (3.7), we have

$$\omega_p \left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \right) \leq 2^{\frac{1}{p}} \min(\omega_p(A), \omega_p(B)) + \min(\omega_p(A+B), \omega_p(A-B))$$

as required. □

4. AN APPLICATION

In this section we present an application, which is a refinement of the triangle inequality for the generalized Schatten p -numerical radius. The following remark is presented by "Yamazaki" in [14]

Remark 4.1. *Let $T \in M_n(\mathbb{C})$, we have*

$$\omega_p(T) = \sup_{\alpha^2 + \beta^2 = 1} \|\alpha \operatorname{Re}(T) + \beta \operatorname{Im}(T)\|_p.$$

for all p . Then

$$\|T + T^*\|_p \leq 2\omega_p(T).$$

Proof. We have

$$\begin{aligned}
\omega_p(T) &= \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta}T)\|_p \\
&= \frac{1}{2} \sup_{\theta \in \mathbb{R}} \|e^{i\theta}T + e^{-i\theta}T^*\|_p \\
&= \sup_{\theta \in \mathbb{R}} \|\cos(\theta)T + i\sin(\theta)T + \cos(\theta)T^* - i\sin(\theta)T^*\|_p \\
&= \sup_{\theta \in \mathbb{R}} \|\cos(\theta)Re(T) - \sin(\theta)Im(T)\|_p \\
&= \sup_{\alpha^2 + \beta^2 = 1} \|\alpha Re(T) + \beta Im(T)\|_p.
\end{aligned}$$

Take $\theta = 2\pi$, then

$$\begin{aligned}
\omega_p(T) &\geq \|\cos(2\pi)Re(T) - \sin(2\pi)Im(T)\|_p \\
&= \frac{1}{2} \|T + T^*\|_p
\end{aligned}$$

so, $\|T + T^*\|_p \leq 2\omega_p(T)$. □

The next theorem is a refinement of the triangle inequality for the generalized Schatten p-numerical radius.

Theorem 4.1. *Let $A, B \in M_n(\mathbb{C})$, then*

$$\|A + B\|_p \leq 2^{1-\frac{1}{p}} \omega_p \left(\begin{bmatrix} 0 & A \\ B^* & 0 \end{bmatrix} \right) \leq \|A\|_p + \|B\|_p.$$

for all p .

Proof. Let $T = \begin{bmatrix} 0 & A \\ B^* & 0 \end{bmatrix}$. We have

$$\begin{aligned}
\left\| \begin{bmatrix} 0 & A+B \\ A^*+B^* & 0 \end{bmatrix} \right\|_p^p &= \|(A+B) \oplus (A^*+B^*)\|_p^p && \text{(by equation (1.2))} \\
&= \|(A+B) \oplus (A+B)\|_p^p && \text{(by equation (1.1))} \\
&= 2 \|A+B\|_p^p. && \text{(by equation (1.4))} \tag{4.1}
\end{aligned}$$

Then,

$$\begin{aligned}
2 \|A + B\|_p^p &= \left\| \begin{bmatrix} 0 & A + B \\ A^* + B^* & 0 \end{bmatrix} \right\|_p^p \\
&= \|T + T^*\|_p^p \\
&\leq 2^p \omega_p^p(T) \quad (\text{by Remark 4.1}) \\
&= 2^p \omega_p^p \left(\begin{bmatrix} 0 & A \\ B^* & 0 \end{bmatrix} \right).
\end{aligned}$$

Thus,

$$\|A + B\|_p \leq 2^{1-\frac{1}{p}} \omega_p \left(\begin{bmatrix} 0 & A \\ B^* & 0 \end{bmatrix} \right).$$

For the second inequality, we have

$$\begin{aligned}
\omega_p(T) &= \sup_{\theta \in \mathbb{R}} \|Re(e^{i\theta} T)\|_p \\
&= \frac{1}{2} \sup_{\theta \in \mathbb{R}} \left\| \begin{bmatrix} 0 & e^{i\theta} A + e^{-i\theta} B \\ e^{-i\theta} A^* + e^{i\theta} B^* & 0 \end{bmatrix} \right\|_p \\
&= \frac{1}{2^{1-\frac{1}{p}}} \sup_{\theta \in \mathbb{R}} \|e^{i\theta} A + e^{-i\theta} B\|_p \quad (\text{by same argument as (4.1)}) \\
&\leq \frac{1}{2^{1-\frac{1}{p}}} (\|A\|_p + \|B\|_p). \quad (\text{by triangle inequality})
\end{aligned}$$

Therefore,

$$\|A + B\|_p \leq 2^{1-\frac{1}{p}} \omega_p \left(\begin{bmatrix} 0 & A \\ B^* & 0 \end{bmatrix} \right) \leq \|A\|_p + \|B\|_p.$$

□

REFERENCES

- [1] H. Abbas, S. Harb, and H. Issa. Inequalities for the generalized numerical radius, *arXiv: 2004.09955* (2020).
- [2] A. Abu-Omar and F. Kittaneh. A generalization of the numerical radius. *Linear Algebra Appl.* 569 (2019), pp. 323-334.
- [3] A. Aldalabih and F. Kittaneh. Hilbert-Schmidt numerical radius inequalities for operator matrices. *Linear Algebra Appl.* 581 (2019), pp. 72-84.
- [4] D. AL-Saafin, *Spectral Radius, Numerical Radius and Unitarily Invariant Norm Inequalities in Hilbert Space*, Master's dissertation, Zarqa University, Jordan, 2016.
- [5] R. Bhatia. *Matrix Analysis*, Grad. Texts in Math., Springer-Verlag, New York, 1997.

- [6] R. Bhatia and F. Kittaneh. Norm inequalities for partitioned operators and an application. *Math. Ann.* 287 (1990), pp. 719-726.
- [7] T. Bottazzi and C. Conde. Generalized numerical radius and related inequalities, *arXiv: 1909.09243* (2019).
- [8] R. Horn and C. Johnson. *Matrix Analysis*. 2nd ed. Cambridge: Cambridge University Press, 2013.
- [9] F. Hiai and D. Petz. *Introduction to Matrix Analysis and Applications*, Universitext, Springer, Switzerland, 2014.
- [10] G.J.O. Jameson. Some inequalities for $(a + b)^p$ and $(a + b)^p + (a - b)^p$. *Math. Gazette* 98 (2014), pp. 96-103.
- [11] F. Kittaneh. Norm inequalities for sums and differences of positive operators. *Linear Algebra and its Applications* 383 (2004), pp. 85-91.
- [12] F. Kittaneh. Norm inequalities for certain operator sums. *Journal of Functional Analysis* 143 (1997), pp. 337-348.
- [13] F. Kittaneh. A numerical radius inequality and an estimate for the numerical radius of the Frobenius companion matrix, *Studia Math.* 158(1) (2003), pp. 11-17.
- [14] T. Yamazaki. Upper and lower bounds of numerical radius and equality condition. 2007.
- [15] F. Zhang. *Matrix Theory*, Springer Science and Business Media, 2011.

JANA HAMZA

DEPARTMENT OF MATHEMATICS, FACULTY OF ARTS AND SCIENCES, LEBANESE INTERNATIONAL UNIVERSITY, BEKAA, LEBANON

Email address: 21610074@students.liu.edu.lb

HASSAN ISSA

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, LEBANESE UNIVERSITY, BEIRUT, LEBANON

Email address: hassan.issa@mathematik.uni-goettingen.de