A MODIFIED NONLINEAR CONJUGATE GRADIENT ALGORITHM FOR FUNCTIONS WITH NON-LIPSCHITZ CONTINUOUS GRADIENT

A PREPRINT

Bingjie Li Department of Statistics & Data Science National University of Singapore, Singapore bjlistat@nus.edu.sg Tianhao Ni School of Mathematical Science Zhejiang University, China 12035009@zju.edu.cn

Schenyue Zhang Nanjing Center for Applied Mathematics, China School of Mathematical Science, Zhejiang University, China zyzhang@zju.edu.cn

April 19, 2022

ABSTRACT

In this paper, we propose a modified nonlinear conjugate gradient (NCG) method for functions with a non-Lipschitz continuous gradient. First, we present a new formula for the conjugate coefficient β_k in NCG, conducting a search direction that provides an adequate function decrease. We can derive that our NCG algorithm guarantees strongly convergent for continuous differential functions without Lipschitz continuous gradient. Second, we present a simple interpolation approach that could automatically achieve shrinkage, generating a step length satisfying the standard Wolfe conditions in each step. Our framework considerably broadens the applicability of NCG and preserves the superior numerical performance of the PRP-type methods.

Keywords nonlinear conjugate gradient · line search · strongly convergent · non-Lipschitz gradient

1 Introduction

Consider the unconstrained optimization problem:

$$\min f(x) \quad x \in \mathbb{R}^n,\tag{1}$$

where f(x) is continuously differentiable. The nonlinear conjugate gradient (NCG) method provides an iterative scheme for minimizing f(x) via the two steps: starting at a point x_0 and setting k = 0 and $d_0 = -\nabla f(x_0)$, the NCG updates the current point x_k via a line search along the direction d_k as that

$$x_{k+1} = x_k + \alpha_k d_k,\tag{2}$$

and then updates the direction d_k in a linear combination of the previous direction d_k and the current gradient $g_{k+1} = \nabla f(x_{k+1})$ for next search,

$$d_{k+1} = -g_{k+1} + \beta_k d_k. \tag{3}$$

In each iteration step, the step length α_k and conjugate coefficient β_k determine the convergence behavior of the NCG. The sequence is called globally convergent if $\liminf_{k\to\infty} ||g_k|| = 0$, and called strongly convergent if $\lim_{k\to\infty} ||g_k|| = 0$.

1.1 The step length

Basically, given a search direction d_k , a step length α_k is chosen to yield a smaller value of the objective function f at the updated point x_{k+1} . The ideal line search set the α_k that minimizes $f(x_k + \alpha d_k)$,

$$\alpha_k = \arg\min_{\alpha} f(x_k + \alpha d_k).$$

Since *f* is nonlinear and not convex in many application, the ideal search may cost much. As a substitute, an inexact search approach is commonly used. In an inexact line search, the step length α_k is chosen to satisfy Armijo-Goldstein Condition [1]

$$f(x_k + \alpha_k d_k) \le f(x_k) + \rho \,\alpha_k \langle g_k, d_k \rangle \tag{4}$$

$$f(x_k + \alpha_k d_k) \ge f(x_k) + (1 - \rho) \alpha_k \langle g_k, d_k \rangle, \tag{5}$$

or the standard Wolfe-Powell conditions [2]

$$f(x_k + \alpha_k d_k) \le f(x_k) + \rho \alpha_k \langle g_k, d_k \rangle \tag{6}$$

$$\langle \nabla f(x_k + \alpha_k d_k), d_k \rangle \ge \sigma \langle g_k, d_k \rangle,$$
(7)

or the strong Wolfe-Powell conditions[3]

$$f(x_k + \alpha_k d_k) \le f(x_k) + \rho \alpha_k \langle g_k, d_k \rangle$$

$$|\langle \nabla f(x_k + \alpha_k d_k), d_k \rangle| \le -\sigma \langle g_k, d_k \rangle.$$
(8)
(9)

Here, the parameters ρ and σ are positive and $\rho < \sigma < 1$. Generally, α_k can be obtained via bisection [4] or interpolation [5], or combination of the two approaches [6].

1.2 The search direction

In nonlinear conjugate gradient methods, the behavior of d_{k+1} is determined by β_k . There are many approaches in the literature to setting β_k . Classical formulas for β_k are called Fletcher-Reeves (FR) [7], Hestenes-Stiefel (HS) [8], Polak-Ribiere-Polyak (PRP) [9]. They are given by

$$\boldsymbol{\beta}_k^{\text{FR}} = \frac{\|\boldsymbol{g}_{k+1}\|_2^2}{\|\boldsymbol{g}_k\|_2^2}, \quad \boldsymbol{\beta}_k^{\text{HS}} = \frac{\langle \boldsymbol{g}_{k+1}, \boldsymbol{y}_k \rangle}{\langle \boldsymbol{d}_k, \boldsymbol{y}_k \rangle}, \quad \boldsymbol{\beta}_k^{\text{PRP}} = \frac{\langle \boldsymbol{g}_{k+1}, \boldsymbol{y}_k \rangle}{\|\boldsymbol{g}_k\|_2^2}$$

where $y_k = g_{k+1} - g_k$. In practice, the PRP method outperforms others in many optimization problems because it can immediately recover after generating a tiny step. However, the PRP method only guarantees global convergence for strictly convex functions, limiting its applicability. To improve it, Gilbert and Nocedal [10] modified the PRP method by setting

$$\boldsymbol{\beta}_{k}^{\text{PRP+}} = \max\{\boldsymbol{\beta}_{k}^{\text{PRP}}, \boldsymbol{0}\},\tag{10}$$

and showed that this modification of the PRP method, called PRP+, is globally convergent if the search direction is sufficient descending and the step length satisfies the standard Wolfe conditions.

In recent years, a variety of new nonlinear conjugate gradient methods have been proposed to find a search direction satisfying the descending condition $\langle d_{k+1}, g_{k+1} \rangle < 0$ or the sufficient descending condition $\langle d_{k+1}, g_{k+1} \rangle < -c ||g_{k+1}||^2$, where *c* is a positive number. In [11], Dai and Yuan proposed a formula with

$$\beta_k^{\rm DY} = \frac{\langle g_{k+1}, d_{k+1} \rangle}{\langle g_k, d_k \rangle},\tag{11}$$

and it provides a descending direction. In [12], Hager and Zhang modified HS method to

$$\beta_{k}^{\text{HZ}} = \max\{\beta_{k}^{\text{HS}} - \frac{2\|y_{k}\|^{2} \langle g_{k+1}, d_{k} \rangle}{\langle d_{k}, y_{k} \rangle^{2}}, -\frac{1}{\|d_{k}\|\min\{\eta, \|g_{k}\|\}}\},\tag{12}$$

where $\eta > 0$ is a constant. Similar modification on PRP method was proposed by Yuan [13], that is,

$$\beta_{k}^{\text{PRP-Y}} = \max\{\beta_{k}^{\text{PRP}} - \frac{\nu \|y_{k}\|^{2}}{\|g_{k}\|^{4}} \langle g_{k+1}, d_{k} \rangle, 0\} \quad \nu > \frac{1}{4}.$$
(13)

Both β_k^{HZ} and β_k^{MPRP} provide a sufficient descent direction.

The convergence of the above nonlinear conjugate gradient methods requires the gradient g(x) of the objective function f(x) to be Lipschitz continuous. That is, there exists a constant L > 0 such that

$$|g(x) - g(y)|| \le L||x - y||, \quad \text{for all} \quad x, y \in \mathbb{R}^n.$$
(14)

This requirement of the Lipschitz continuous gradient limits the application of nonlinear conjugate gradient methods when faced with complicated practical problems.

1.3 Our contribution

In this paper, we propose a modified nonlinear conjugate gradient method, which does not require the gradient g(x) to be Lipschitz continuous. The novelty of our approach comes from two aspects:

- We propose a new formula for β_k , called MPRP, obtaining an adequate descending direction. The strong convergence of our approach is guaranteed even though f(x) is just a continuous differential function with a non-Lipschitz gradient.
- We suggest a more straightforward line search method for a step length that satisfies the standard Wolfe conditions in finite iterations. In practice, it works very well. The line search iteration terminates within one or two iterations generally in our experiments.

This paper is organized as follows. In Section 2, we propose our new formula for β_k , and the line search approach is given in Section 3. We discuss the convergence of our method in Section 4. The numerical experiment is also given in Section 5, to show the performance of our approach. At last, we end the article with a conclusion in Section 6.

2 The new formula for β_k

Besides the Lipschitz continuity of the gradient, the convergence of the PRP-Y method [13] requires that the step length $\{\alpha_k\}$ have a positive lower bound. This requirement is difficult to guarantee in practice. To weaken the conditions required for the convergence of the PRP-Y method, we provide a formula for β_k as

$$\beta_{k} = \min\left\{\frac{\langle g_{k+1}, g_{k+1} - g_{k} - \frac{v \|g_{k+1} - g_{k}\|^{2}}{\|g_{k}\|^{2}} d_{k}\rangle_{+}}{\|g_{k}\|^{2}}, \frac{\kappa \|g_{k+1}\|_{2}}{\|d_{k}\|_{2}}\right\}.$$
(15)

where $v > \frac{1}{4}$ as in (13) and $\kappa > 0$. The modification can guarantee a stronger sufficient descent condition than that of the PRP-Y method. That is,

Lemma 2.1. Let β_k be defined by (15) and $\mu = \frac{4\nu - 1}{4\nu(1+\kappa)}$. Then

$$\langle d_{k+1}, g_{k+1} \rangle \le -\mu \| d_{k+1} \|_2 \| g_{k+1} \|_2.$$
 (16)

Proof. Let $\tilde{\beta}_k = \beta_k^{\text{PRP}}$ for short, and let $\tilde{d}_{k+1} = -g_{k+1} + \tilde{\beta}_k d_k$. We rewrite

$$\beta_k = \rho_k \beta_k, \quad d_{k+1} = \rho_k d_{k+1} + (\rho_k - 1)g_{k+1}$$

with a scale $\rho_k \in [0, 1]$ since $\beta_k \leq \tilde{\beta}_k$. At first, we require the inequality

$$\tilde{\beta}_k \langle d_k, g_{k+1} \rangle \le \frac{1}{4\nu} \|g_{k+1}\|^2, \tag{17}$$

concluded by the definition $\tilde{\beta}_k = \frac{\langle g_{k+1}, g_{k+1} - g_k \rangle}{\|g_k\|_2^2} - \frac{v\|g_{k+1} - g_k\|^2}{\|g_k\|^4} \langle g_{k+1}, d_k \rangle$. It gives

$$egin{aligned} & ilde{eta}_k \langle d_k, g_{k+1}
angle &= rac{\langle g_{k+1}, g_{k+1} - g_k
angle}{\|g_k\|_2^2} \langle g_{k+1}, d_k
angle - rac{m{v} \|g_{k+1} - g_k\|^2}{\|g_k\|_4^4} \langle g_{k+1}, d_k
angle^2 \ &= \Big\langle g_{k+1}, rac{\langle g_{k+1}, d_k
angle}{\|g_k\|_2^2} (g_{k+1} - g_k) \Big
angle - rac{m{v} \langle g_{k+1}, d_k
angle^2}{\|g_k\|_4^4} \|g_{k+1} - g_k\|^2 \end{aligned}$$

Let $q_k = \frac{\langle g_{k+1}, d_k \rangle}{\|g_k\|_2^2} (g_{k+1} - g_k)$ for simplicity. Then

ļ

$$\tilde{\beta}_k \langle d_k, g_{k+1} \rangle = \langle g_{k+1}, q_k \rangle - \mathbf{v} \| q_k \|^2 = \frac{\| g_{k+1} \|^2}{4\mathbf{v}} - \| (2\sqrt{\mathbf{v}})^{-1} g_{k+1} - \sqrt{\mathbf{v}} q_k \|^2.$$

Therefore, (17) is true. Following it, we get that $\langle \tilde{d}_{k+1}, g_{k+1} \rangle \leq (\frac{1}{4v} - 1) ||g_{k+1}||^2$ and

$$\begin{aligned} \langle d_{k+1}, g_{k+1} \rangle &= \rho_k \langle \tilde{d}_{k+1}, g_{k+1} \rangle + (\rho_k - 1) \|g_{k+1}\|^2 \\ &\leq \left(\rho_k (\frac{1}{4\nu} - 1) + (\rho_k - 1) \right) \|g_{k+1}\|^2 \leq \frac{1 - 4\nu}{4\nu} \|g_{k+1}\|^2. \end{aligned}$$

Here we have used $\rho_k \leq 1$. On the other hand, since $|\beta_k| \leq \frac{\kappa ||g_{k+1}||}{||d_k||}$, we also have that

$$\|d_{k+1}\| = \|-g_{k+1} + \beta_k d_k\| \le \|g_{k+1}\| + |\beta_k| \|d_k\| \le (1+\kappa) \|g_{k+1}\|.$$

Therefore, (16) holds since $||g_{k+1}||_2^2 \ge \frac{1}{1+\kappa} ||d_{k+1}||_2 ||g_{k+1}||_2$ and $\frac{1-4\nu}{4\nu} < 0$.

We call a search direction d_{k+1} an adequate descending direction if it satisfies (16). Obvious, the gradient itself satisfies (16) with $\mu = 1$. To the best of our knowledge, (15) is the first conjugate gradient method that provides an adequate descending direction.

3 A simple interpolation line search approach

For a general continuously differentiable f(x), the interpolation method does not guarantee capturing required α_k satisfying the standard Wolfe conditions since it asks for a three times continuously differentiable [5]. One can get α_k by the combination method [6] that is more efficient than the bisection approach [4]. In [6], the bisection is combined with the interpolation in a bit complicated way for interval shrinking. Here we give a simpler approach for determining α_k satisfying the weak Wolfe-Powell conditions.

Theoretically, at a current point $x = x_k$ with the conjugate direction $d = d_k$, the required inexact line search $\alpha = \alpha_k$ satisfying the weak Wolfe-Powell conditions (6-7) can be chosen as

$$\alpha^* = \sup \left\{ \hat{\alpha} : \text{the Wolfe-Powell condition (6) holds over } (0, \hat{\alpha}) \right\}.$$
(18)

It exists, is positive, and satisfies (6-7). To verify this claim, let's consider the function

$$g(\alpha) = f(x) + \rho \alpha \langle \nabla f(x), d \rangle - f(x + \alpha d).$$

Clearly, (6) is equivalent to $g(\alpha) \ge 0$, and meanwhile, (7) holds if $g'(\alpha) \le 0$. By the definition and the continuity of f, (6) is true for $0 < \alpha \le \alpha^*$. The supremum in (18) implies that $g(\alpha^*) = 0$ and $g'(\alpha^*) \le 0$. Hence, (7) is also satisfied for $\alpha = \alpha^*$. Practically, there is a relative large sub-interval of $(0, \alpha^*]$ in which both (6) and (7) are true. For instance, if $\hat{\alpha} \in (0, \alpha^*]$ is the largest point such that $g(\alpha)$ is a local maximum, then $g'(\alpha) \le 0$ in $[\hat{\alpha}, \alpha^*]$. Therefore, (6-7) hold for $\alpha \in [\hat{\alpha}, \alpha^*]$.

An ideal choice of α is the minimizer α_{\min} of $f(x + \alpha d)$ over $(0, \alpha^*]$ since it decreases f as small as possible, while both (6) and (7) are still satisfied. In this subsection, we give a simple rule for pursuing α_{\min} via a quadratic interpolation to $f(x + \alpha d)$, assuming f is continuously differentiable. It generates a nested and shrunk interval sequence containing the required α . The pursuing terminates as soon as a point satisfying (6-7) is found.

Initially, we set $\alpha'_0 = 0$ that satisfies (6) but (7), and choose a relatively large $\alpha''_0 > 0$ that does not satisfy (6). A simple choice of α''_0 will be given later. Starting with $[\alpha'_0, \alpha''_0]$, we generate a sequence of intervals $[\alpha'_0, \alpha''_0]$ iteratively such that each α'_ℓ satisfies (6) but α''_ℓ does not, and meanwhile, α'_ℓ doesn't satisfy (7). That is, for $x'_\ell = x + \alpha'_\ell d$ and $x''_\ell = x + \alpha''_\ell d$

$$f(x_{\ell}') \le f(x) + \rho \alpha_{\ell}'(g, d), \ f(x_{\ell}'') > f(x) + \rho \alpha_{\ell}''(g, d), \ \langle \nabla f(x_{\ell}'), d \rangle < \sigma \langle g, d \rangle,$$
(19)

where $g = \nabla f(x)$. The third inequality above implies that $\langle \nabla f(x'_{\ell}), d \rangle < 0$. Furthermore, by the first two inequalities in (19), we have that

$$f(x_{\ell}'') > f(x_{\ell}') + \rho(\alpha_{\ell}'' - \alpha_{\ell}') \langle g, d \rangle > f(x_{\ell}') + \frac{\rho}{\sigma} (\alpha_{\ell}'' - \alpha_{\ell}') \langle \nabla f(x_{\ell}'), d \rangle.$$

$$(20)$$

In the current interval, we consider a quadratic function $q(\alpha)$ with interpolation conditions

$$q(\boldsymbol{\alpha}'_{\ell}) = f(x'_{\ell}), \quad q'(\boldsymbol{\alpha}'_{\ell}) = \langle \nabla f(x'_{\ell}), d \rangle, \quad q(\boldsymbol{\alpha}''_{\ell}) = f(x''_{\ell}),$$

It can be represented as

$$\begin{split} q(\boldsymbol{\alpha}) &= f(x'_{\ell}) + (\boldsymbol{\alpha} - \boldsymbol{\alpha}'_{\ell}) \langle \nabla f(x'_{\ell}), d \rangle \\ &+ \left(f(x''_{\ell}) - f(x'_{\ell}) - (\boldsymbol{\alpha}''_{\ell} - \boldsymbol{\alpha}'_{\ell}) \langle \nabla f(x'_{\ell}), d \rangle \right) \frac{(\boldsymbol{\alpha} - \boldsymbol{\alpha}'_{\ell})^2}{(\boldsymbol{\alpha}''_{\ell} - \boldsymbol{\alpha}'_{\ell})^2} \end{split}$$

with the minimizer $c_{\ell} = \arg \min_{\alpha} q(\alpha)$ given by

$$c_{\ell} = \alpha_{\ell}' + \frac{\alpha_{\ell}'' - \alpha_{\ell}'}{2} \frac{-(\alpha_{\ell}'' - \alpha_{\ell}') \langle \nabla f(x_{\ell}'), d \rangle}{f(x_{\ell}'') - f(x_{\ell}') - (\alpha_{\ell}'' - \alpha_{\ell}') \langle \nabla f(x_{\ell}), d \rangle} > \alpha_{\ell}'.$$

$$(21)$$

By the Mean-Value Theorem for derivatives and the second inequality in (20),

$$0 < (1 - M_{\ell})^{-1} \le \left(1 - \frac{\langle \nabla f(\bar{x}_{\ell}), d \rangle}{\langle \nabla f(x'_{\ell}), D_k \rangle}\right)^{-1} = \frac{-(\alpha''_{\ell} - \alpha'_{\ell}) \langle \nabla f(x'_{\ell}), D_k \rangle}{f(x''_{\ell}) - f(x'_{\ell}) - (\alpha''_{\ell} - \alpha'_{\ell}) \langle \nabla f(x'_{\ell}), D_k \rangle} < \frac{\sigma}{\sigma - \rho},$$
(22)

Algorithm 1 An inexact line search satisfying the weak Wolfe-Powell conditions

Require: point *x*, f = f(x), $g = \nabla f(x)$, direction *d*, and parameters σ , ρ .

Ensure: α satisfying (6-7) within accuracy ε , $x := x + \alpha d$, f(x), and $g = \nabla f(x)$.

- 1: Set $\alpha' = 0$, x' = x, f' = f, g' = g, $v = \rho \langle g, d \rangle$. Find the smallest integer $p \ge 1$ such that (6) does not hold for $\alpha = \eta 2^p$, and set $\alpha'' = \eta 2^p$.
- 2: Repeat the following iteration until terminating.
- 3: Compute *c* as (21), \tilde{c} as (24), and $\tilde{f} = f(\tilde{x})$ at $\tilde{x} = x + \tilde{c}d$.
- 4: If $\tilde{f} > f + \tilde{c}v$, update $(\alpha'', f(x''))$ by $(\tilde{c}, f(\tilde{x}))$ and go to Step 3.
- 5: Compute $\tilde{g} = \nabla f(\tilde{x})$. If $\langle \tilde{g}, d \rangle \ge \sigma \langle g, d \rangle$, set $x = \tilde{x}$, $f = \tilde{f}$, $g = \tilde{g}$, and terminate.
- 6: Otherwise, update α', f', g' by $\tilde{c}, \tilde{f}, \tilde{g}$, respectively.

7: End iteration

where $\bar{x}_{\ell} = x + \bar{\alpha}_{\ell} d$ with $\bar{\alpha}_{\ell} \in [\alpha_{\ell}', \alpha_{\ell}'']$ and

$$M_{\ell} = \min_{\alpha \in [\alpha'_{\ell}, \alpha''_{\ell}]} \frac{\langle \nabla f(x + \alpha d), d \rangle}{\langle \nabla f(x'_{\ell}), d \rangle} \leq \frac{\langle \nabla f(\bar{x}_{\ell}), d \rangle}{\langle \nabla f(x'_{\ell}), d \rangle} < \frac{\rho}{\sigma}.$$

Hence, if $0 < 2\rho < \sigma$, we have that

$$\alpha_{\ell}' < \alpha_{\ell}' + \frac{1}{2(1-M_{\ell})}(\alpha_{\ell}'' - \alpha_{\ell}') < c_{\ell} < \alpha_{\ell}' + \frac{\sigma}{2(\sigma-\rho)}(\alpha_{\ell}'' - \alpha_{\ell}') < \alpha_{\ell}''.$$

$$\tag{23}$$

We may shrink $[\alpha'_{\ell}, \alpha''_{\ell}]$ to $[c_{\ell}, \alpha''_{\ell}]$ or $[\alpha'_{\ell}, c_{\ell}]$, if $\alpha = c_{\ell}$ satisfies (6) or does not. However, if (6) is satisfied, the interval length is $\alpha''_{\ell} - c_{\ell} \leq \frac{1-2M_{\ell}}{2-2M_{\ell}}(\alpha''_{\ell} - \alpha'_{\ell})$. When $M_{\ell} < 0$ and $|M_{\ell}|$ is large, $\frac{1-2M_{\ell}}{2-2M_{\ell}} \approx 1$. The interval shrinking is inefficient in this case. To avoid this phenomenon, we slightly modify c_{ℓ} as that with $\eta = \frac{\sigma}{2(\sigma-\rho)}$

$$\tilde{c}_{\ell} = \max\left\{c_{\ell}, \, \eta \, \alpha_{\ell}' + (1 - \eta) \, \alpha_{\ell}''\right\} \in (\alpha_{\ell}', \alpha_{\ell}'').$$
(24)

Since $\tilde{c}_{\ell} \geq \eta \alpha'_{\ell} + (1 - \eta) \alpha''_{\ell}$ and $c_{\ell} < \alpha'_{\ell} + \eta (\alpha''_{\ell} - \alpha'_{\ell})$ by (23), we get

$$\alpha_\ell'' - \tilde{c}_\ell \leq \eta(\alpha_\ell'' - \alpha_\ell'), \quad \tilde{c}_\ell - \alpha_\ell' \leq \max\left\{\eta, 1 - \eta\right\}(\alpha_\ell'' - \alpha_\ell') = \eta(\alpha_\ell'' - \alpha_\ell').$$

The last equality holds since $\eta > 1/2$. Hence, if the Wolfe-Powell conditions (6-7) are satisfied for $\alpha = \tilde{c}_{\ell}$, we get the required $\alpha_k = \tilde{c}_{\ell}$. Otherwise, shrink $[\alpha'_{\ell}, \alpha''_{\ell}]$ as

$$[\alpha_{\ell+1}', \alpha_{\ell+1}''] = \begin{cases} [\alpha_{\ell}', \tilde{c}_{\ell}], & \text{if (6) does not hold for } \alpha = \tilde{c}_{\ell}; \\ [\tilde{c}_{\ell}, \alpha_{\ell}''], & \text{otherwise.} \end{cases}$$
(25)

The interval length is significantly decreased as $0 < \alpha''_{\ell+1} - \alpha'_{\ell+1} \le \eta(\alpha''_{\ell} - \alpha'_{\ell})$, where $\eta < 1$ since $2\rho < \sigma$. Hence, $\alpha''_{\ell} - \alpha'_{\ell} \to 0$ as $\ell \to \infty$.

A good choice of α_0'' helps to pursue the minimizer α_{\min} . Motivated by the above analysis on the estimation of the shrinking rate η_ℓ , we suggest the experiential setting

$$\alpha_0^{\prime\prime} = \min\left\{\alpha = 2^p \eta : (6) \text{ is not satisfied for } \alpha = 2^p \eta \text{ with integer } p \ge 0\right\}.$$
(26)

Algorithm 1 gives the details of the procedure for determining an inexact line search α_k , given x_k , f_k , g_k , the conjugate direction d_k .

4 Convergence of the Algorithm

Combining the formula (15) and line search Algorithm 1, we are able to provide our modified PRP-type (MPRP) nonlinear conjugate gradient method, as shown in Algorithm 2. To show the convergence of the MPRP method, we first prove that the line search Algorithm 1 will converge to a step length that satisfies the standard Wolfe condition.

Lemma 4.1. If *f* is lower bounded and continuously differentiable, an $\alpha = \tilde{c}_{\ell^*}$ satisfying (6-7) can be obtained within *a finite iterations of (25) if* $0 < 2\rho < \sigma < 1$.

Algorithm 2 The modified PRP-type (MPRP) nonlinear conjugated gradient method

Require: initial point *x*, parameters ε , ρ , σ , *v*, κ , and k_{\max}^{NCG} .

Ensure: an approximate solution x_* of $\min_x f(x)$ with the given accuracy

- 1: Compute f = f(x), $g = \nabla f(x)$, and set d = -g.
- 2: For $k = 1, \cdots, k_{\text{max}}^{\text{NCG}}$
- 3: Update (x, f, g) by Algorithm 1 with searching direct *d*.
- 4: If $||g||_{\infty} < \varepsilon$, then set $x^* = x$ and terminate the iteration.
- 5: Otherwise, compute β by (15) and update $d := -g + \beta d$.
- 6: End for

Proof. If (6-7) do not hold for all \tilde{c}_{ℓ} , the updating rule (25) yields a sequence of nested intervals $\{[\alpha'_{\ell}, \alpha''_{\ell}]\}$. Since $0 < 2\rho < \sigma < 1$, the intervals tend to a single point α_* and both $\{x'_{\ell}\}$ and $\{x''_{\ell}\}$ tend to $x_* = x + \alpha_* d$. Hence, by (20) and the Taylor extension of $f(x + \alpha d)$ at $\alpha = \alpha_*$, we get

$$\langle \nabla f(x_*), d \rangle = \lim_{\ell \to \infty} \frac{f(x_{\ell}') - f(x_{\ell}')}{\alpha_{\ell}'' - \alpha_{\ell}'} \ge \rho \langle \nabla f(x), d \rangle > \sigma \langle \nabla f(x), d \rangle$$
(27)

since $\langle \nabla f(x), d \rangle < 0$ and $\rho < \sigma$. However, by (19), $\langle \nabla f(x_*), d \rangle \leq \sigma \langle \nabla f(x), d \rangle$, a contradiction with (27).

Because the search direction of MPRP is adequate descending, the proof of convergence of the algorithm is simple, similar to the proof of the steepest descent method. We have

Theorem 4.2. Assume that f is lower bounded and continuously derivative. If the inexact line search $\{\alpha_k\}$ satisfies the weak Wolfe-Powell condition (6-7) and

$$\langle d_k, g_k \rangle \le -\mu \| d_k \|_2 \| g_k \|_2$$
 (28)

for a constant $\mu > 0$, then the NCG converges: $\{f(x_k)\}$ is monotone decreasing and converges, and $\nabla f(x_k) \rightarrow 0$.

Proof. We assume $g_k = \nabla f(x_k) \neq 0$ for each k without loss of generalities, and let $s_k = \alpha_k d_k$. The condition (28) becomes $\langle g_k, s_k \rangle \leq -\mu ||g_k|| ||s_k|| \leq 0$. Hence, the Wolfe-Powell condition (6) gives the monotone decreasing of $\{f(x_k)\}$,

$$f(x_{k+1}) - f(x_k) \leq \rho \langle g_k, s_k \rangle \leq -\rho \mu \|g_k\| \|s_k\| \leq 0,$$

and $\{f(x_k)\}$ is convergent since *f* itself is lower bounded. We also conclude from the convergence and the above inequality that $||g_k|| ||s_k|| \to 0$.

We further show that $||g_k|| \to 0$. Otherwise, there is a subsequence $\{||g_{k_i}||\}$ that has a positive lower bound. The lower bound implies that $||s_{k_i}|| \to 0$ since we also have $||g_{k_i}|| ||s_{k_i}|| \to 0$. Note that $s_{k_i} = x_{k_i+1} - x_{k_i}$ is also the gap vector between x_{k_i} and x_{k_i+1} , $f(x_{k_i})$ and $f(x_{k_i+1})$ can be represented each other in terms of s_{k_i} via the Taylor extensions

$$f(x_{k_i+1}) = f(x_{k_i}) + \langle g_{k_i}, g_{k_i} \rangle + o(||g_{k_i}||),$$

$$f(x_{k_i}) = f(x_{k_i+1}) - \langle g_{k_i+1}, g_{k_i} \rangle + o(||g_{k_i}||).$$

Clearly, the each other implies that $\langle g_{k_i}, g_{k_i} \rangle - \langle g_{k_i+1}, g_{k_i} \rangle = o(||g_{k_i}||).$

Turn back to the Wolfe-Powell condition (7). Since it gives $\langle g_{k_i+1}, s_{k_i} \rangle \ge \sigma \langle g_{k_i}, s_{k_i} \rangle$,

$$o(\|s_{k_i}\|) = \langle g_{k_i}, s_{k_i} \rangle - \langle g_{k_i+1}, s_{k_i} \rangle \le (1-\sigma) \langle g_{k_i}, s_{k_i} \rangle \le -(1-\sigma) \mu \|g_{k_i}\| \|s_{k_i}\|.$$

Here we have used the inequality $\langle g_{k_i}, g_{k_i} \rangle \leq -\mu ||g_{k_i}|| ||s_{k_i}||$ from the condition (28) and $\sigma < 1$. Hence, $(1 - \sigma)\mu ||g_{k_i}|| \leq -\frac{o(||s_{k_i}||)}{||s_{k_i}||}$, and

$$0 \leq (1 - \sigma) \mu \liminf \|g_{k_i}\| \leq 0.$$

It implies $(1 - \sigma)\mu \leq 0$, a contradiction, since $\liminf ||g_{k_i}|| > 0$ by assumption.

5 Numerical Experiments

In this section, we show the performance of our MPRP in Algorithm 2, compared with three other PRP-type NCG methods: the classical PRP method [9], the PRP+ method [10] and the PRP-Y method [13]. The numerical experiments are divided into 3 parts. In the first part, we test these PRP-type methods on 84 unconstrained optimization problems with Lipschitz continuous gradient from [14]. The second part aims to show the enhancements brought by our line search method, compared with the bisection line search method. In the third part, we adopt our algorithm on a regression problem with an objective function whose gradient is non-Lipschitz continuous. The following parameters were adopted in our implementation

$$v = 0.8, \kappa = 10, \rho = 0.1, \sigma = 0.4.$$

We set the termination criterion as $\|\nabla f(x)\|_{\infty} \le 10^{-5}$ and the maximum number of iterations as 20000. All compared algorithms are executed on the Windows system in a PC with Intel Core i5-8250U CPU@1.80GHz and 8GB RAM.

We adopt the commonly used performance profile of Dolan and Moré proposed in [15], to display the performance of compared NCG methods in terms of CPU time and the number of iterations. Take the CPU time as an example, let *S* and *P* be the set of solvers and problems, respectively, and denote $n_s = |S|$ and $n_p = |P|$. For each solver *s* and problem *p*, let $t_{p,s}$ be the computing time requiring by solver *s* to solve problem *p*. For each problem *p*, define the performance ratio as $r_{p,s} = \frac{t_{p,s}}{\min\{t_{p,s},s\in S\}}$ and the ratio $r_{p,s} \ge 1$ obviously for all *p* and *s*. If a solver fails to solve a problem, the ratio $r_{p,s}$ is set to a large enough positive number *M* that larger than $r_{p,s}$ of problem *p* that can be solved by solver *s*. Finally, the performance profile is defined by

$$\rho(\tau) = \frac{1}{n_p} |\{p \in P : r_{p,s} \le \tau\}|$$

. The performance profile of the number of iterations is similar.

5.1 The performance of compared NCG methods on tested functions

This subsection shows the performance of 4 tested NCG algorithms on 84 unconstrained optimization problems drawn from [14]. The performance profile of CPU time and number of iterations is illustrated in Figure 1.



Figure 1: Performance profile of PRP,PRP+,PRP-Y and MPRP with CPU Time and Number of iteration

Among the four algorithms, MPRP and PRP-Y have significant advantages over the PRP and PRP+ method in the CPU Time and the number of iterations. However, PRP-Y performs slightly better than our MPRP. This is due to two reasons. For one thing, MPRP controls the angle between the search direction and the negative gradient, weakening the effect of the conjugate direction. Hence, the zig-zag phenomenon may occur in a small part of optimization problems. For another thing, MPRP requires an additional computation of $||d_k||$ at each step, which also increases the CPU time slightly.



Figure 2: Performance profile of different line search methods on PRP,PRP+,PRP-Y and MPRP with CPU Time and Number of iterations

5.2 Line searches via bisection and interpolation

In this subsection, we test the performance of our interpolation line search in Algorithm 1, compared with the classical bisection line search [4]. The two line search approaches are used for the 4 tested NCG methods, on the 84 unconstrained optimization problems in [14]. The Comparison is shown on CPU time and the number of iterations in Figure 2.

As shown in Figure 2, the interpolation line search performs much better than the bisection line search in most problems regardless of the NCG methods we adopt. It has been shown that our interpolation line search method saves a significant amount of computation both in terms of the number of iterations and the CPU time.

5.3 Performance of NCG methods on a function with non-Lipschitz continuous gradient

In this subsection, we consider a linear regression model with a regular term as follows:

$$\min_{x} \frac{1}{2} \|Ax - b\|_{2}^{2} + \frac{\lambda}{2} \|x\|_{p}^{p},$$
(29)

where $||x||_p^p = |x_1|^p + ... + |x_n|^p$. The model (29) becomes the lasso regression model or the ridge regression model, if one select p = 1, or p = 2, respectively. Here we set p = 1.5 so that $f(x) = \frac{1}{2} ||Ax - b||_2^2 + \frac{\lambda}{2} ||x||_p^p$ is continuous differential with a non-Lipschitz gradient, and test the compared NCG methods on it.

In our experiment, we set A as a random matrix in $\mathbb{R}^{10\times 50}$ with entries drawn from the uniform distribution in [0, 1], and b = Au with u is sparse with 10% non-zero entries drawn from the standard normal distribution. We also set $\lambda = 0.01$. We test the four algorithms 10 times, each with different random seeds.

Table 1: The performance of four NCG methods on (29) with p = 1.5.

Method	PRP	PRP+	PRP-Y	MPRP
CPU Time	1.13	1.01	0.81	0.81
Iterations	504.5	447.9	422.9	419.4

Table 1 shows the average CPU time and the average number of iterations for the 10 runs. It can be seen that the performance of PRP-Y and MPRP is slightly better than that of the other two algorithms. Surprisingly, PRP, PRP+ and PRP-Y all converge on this problem, even though they do not theoretically have a guarantee of convergence. One conjecture is that these algorithms skip the non-Lipschitz region of the gradient and converge to a stationary point with a neighbor where the gradient is Lipschitz continuous.

6 Conclusion

This paper proposed a modified nonlinear conjugate gradient method for continuous differential function. The strong convergence is guaranteed without the condition of the Lipschitz continuous gradient. Furthermore, a simpler but more efficient interpolation Wolfe line search method is also introduced. The numerical results demonstrate the feasibility of the new NCG method and the new line search method. However, although in theory our algorithm gains a greater range of applicability, this advantage does not manifest itself numerically. The reasons behind this are worth further investigation.

Acknowledgments

The work was supported by NSFC project 11971430.

References

- [1] Larry Armijo. Minimization of functions having lipschitz continuous first partial derivatives. *Pacific Journal of mathematics*, 16(1):1–3, 1966.
- [2] Philip Wolfe. Convergence conditions for ascent methods. SIAM review, 11(2):226–235, 1969.
- [3] Philip Wolfe. Convergence conditions for ascent methods. ii: Some corrections. *SIAM review*, 13(2):185–188, 1971.
- [4] Jorge J Moré and Danny C Sorensen. Newton's method. Technical report, Argonne National Lab., IL (USA), 1982.
- [5] Wenyu Sun and Ya-Xiang Yuan. *Optimization theory and methods: nonlinear programming*, volume 1. Springer Science & Business Media, 2006.
- [6] Roger Fletcher. Practical methods of optimization. John Wiley & Sons, 2013.
- [7] Reeves Fletcher and Colin M Reeves. Function minimization by conjugate gradients. *The computer journal*, 7(2):149–154, 1964.
- [8] Magnus R Hestenes and Eduard Stiefel. Methods of conjugate gradients for solving. *Journal of research of the National Bureau of Standards*, 49(6):409, 1952.
- [9] Boris Teodorovich Polyak. The conjugate gradient method in extremal problems. USSR Computational Mathematics and Mathematical Physics, 9(4):94–112, 1969.
- [10] Jean Charles Gilbert and Jorge Nocedal. Global convergence properties of conjugate gradient methods for optimization. *SIAM Journal on optimization*, 2(1):21–42, 1992.
- [11] Yu-Hong Dai and Yaxiang Yuan. A nonlinear conjugate gradient method with a strong global convergence property. *SIAM Journal on optimization*, 10(1):177–182, 1999.
- [12] William W Hager and Hongchao Zhang. A new conjugate gradient method with guaranteed descent and an efficient line search. *SIAM Journal on optimization*, 16(1):170–192, 2005.
- [13] Gonglin Yuan. Modified nonlinear conjugate gradient methods with sufficient descent property for large-scale optimization problems. *Optimization Letters*, 3(1):11–21, 2009.
- [14] Neculai Andrei. An unconstrained optimization test functions collection. *Adv. Model. Optim*, 10(1):147–161, 2008.
- [15] Elizabeth D Dolan and Jorge J Moré. Benchmarking optimization software with performance profiles. *Mathematical programming*, 91(2):201–213, 2002.