Comment on "Scattering of light by a parity-time-symmetric dipole beyond the first Born approximation"

Farhang Loran¹, Ali Mostafazadeh², Sema Seymen³, and O. Teoman Turgut³

¹Department of Physics, Isfahan University of Technology, Isfahan 84156-83111, Iran

²Departments of Mathematics and Physics, Koç University, 34450 Sariyer, Istanbul, Turkey

³Department of Physics, Boğaziçi University, 34342 Bebek, Istanbul, Turkey

In [J. A. Rebouças and P. A. Brandão, Phys. Rev. A **104**, 063514 (2021)] the authors compute the scattering amplitude for a \mathcal{PT} -symmetric double-delta-function potential in three dimensions by invoking the far-zone approximation and summing the resulting Born series. We show that the analysis of this paper suffers from a basic error. Therefore its results are inconclusive. We give an exact closed-form expression for the scattering amplitude of this potential.

The authors of [1] consider the scattering problem for a \mathcal{PT} -symmetric double-delta-function potential in threedimensions,

$$p(\mathbf{r}) := \mathfrak{z}_1 \delta(\mathbf{r} - \mathbf{r}_0) + \mathfrak{z}_2 \delta(\mathbf{r} + \mathbf{r}_0), \qquad (1)$$

where $\mathfrak{z}_1 = \mathfrak{z}_2^* = -\alpha k^2 (\sigma + i\gamma)$ and α, σ , and γ are real parameters, k is the wavenumber, and $\pm \mathbf{r}_0$ are the positions of the point scatterers. They substitute the Born series $u(\mathbf{r}) = \sum_{n=0}^{\infty} u_n(\mathbf{r}) \alpha^n$ in the Lippmann-Schwinger equation to show that

$$u_n(\mathbf{r}) = k^2 \int_{\mathbb{R}^3} \chi(\mathbf{r}') G(|\mathbf{r} - \mathbf{r}'|) u_{n-1}(\mathbf{r}') d^3 r', \quad n \ge 1, \ (2)$$

where $\chi(\mathbf{r}') := v(\mathbf{r})/\alpha k^2$, and $G(x) := e^{ikx}/4\pi x$. Then they let $r := |\mathbf{r}|, r' := |\mathbf{r}'|, \hat{s} := \mathbf{r}/r$, denote the direction of the incident wave vector by \hat{a} , and use $u_0(\mathbf{r}) = e^{ik\hat{a}\cdot\mathbf{r}}$ and the far-zone (f.z.) approximation,

$$G(|\mathbf{r} - \mathbf{r}'|) \sim \frac{e^{ikr}}{4\pi r} e^{-ik\hat{s}\cdot\mathbf{r}'} \quad \text{for} \quad r \gg r', \qquad (3)$$

in (2) to obtain the recursion relation,

$$u_n(\mathbf{r}) = \frac{k^2 e^{ikr}}{4\pi r} [(\sigma + i\gamma)u_{n-1}(\mathbf{r}_0)e^{-ik\hat{s}\cdot\mathbf{r}_0} + (\sigma - i\gamma)u_{n-1}(-\mathbf{r}_0)e^{ik\hat{s}\cdot\mathbf{r}_0}], \quad n \ge 1.$$
(4)

The results of [1] rely on the authors' solution of this relation. But as we explain below, there is a basic error in their analysis. To determine $u_1(\mathbf{r})$, they substitute (3) in (2) and set n = 1. Because (3) holds whenever $r \gg r'$, this gives an approximate expression $u_1^{\text{f.z.}}(\mathbf{r})$ for $u_1(\mathbf{r})$ which is valid in the f.z., i.e., $u_1(\mathbf{r}) \sim u_1^{\text{f.z.}}(\mathbf{r})$ for $r \to \infty$. Repeating the same procedure for n = 2, they express $u_2(\mathbf{r})$ in terms of $u_1(\pm \mathbf{r}_0)$ which they calculate by substituting $\pm \mathbf{r}_0$ for \mathbf{r} in $u_1^{\text{f.z.}}(\mathbf{r})$, i.e., set $u_1(\pm \mathbf{r}_0) \sim$ $u_1^{\text{f.z.}}(\pm \mathbf{r}_0)$. This is inadmissible, because $|\mathbf{r}_0|$ does not tend to infinity. In general, the iterative solution of (4) given in the appendix of [1] is unacceptable, because this equation holds for $r \to \infty$. Therefore, one cannot use it to determine $u_n(\pm \mathbf{r}_0)$ even approximately.

There is actually no need for invoking the f.z. approximation in treating this problem [2, 3]. Ref. [4] gives the exact solution of the scattering problem for the multidelta-function potentials,

$$v(\mathbf{r}) := \sum_{n=1}^{N} \mathfrak{z}_n \delta(\mathbf{r} - \mathbf{a}_n), \tag{5}$$

in two dimensions, where \mathfrak{z}_n are real or complex coupling constants, and \mathbf{a}_n are the positions of the point scatterers. The analysis of [4] has a straightforward extension to three dimensions. To see this, first we use the notation of Ref. [1] to express the Lippmann-Schwinger equation for the potential (5) in the form

$$u(\mathbf{r}) = u_0(\mathbf{r}) - \sum_{n=1}^N \mathfrak{z}_n G(|\mathbf{r} - \mathbf{a}_n|) u(\mathbf{a}_n).$$
(6)

Setting $\mathbf{r} = \mathbf{a}_m$ in this equation, we find a system of linear equations for $u(\mathbf{a}_n)$. Because $G(0) = \infty$, the matrix of coefficients of this system has divergent diagonal entries. Therefore, we regularize G(x) and perform a couplingconstant renormalization to remove the singularities. We can do this by a cut-off renormalization or dimensional regularization as outlined in [5] or other renormalization schemes [6]. In this way, we can set $\mathbf{r} = \mathbf{a}_m$ in (6) to arrive at

$$\sum_{n=1}^{N} A_{mn} X_n = e^{ik\mathbf{a}_m \cdot \hat{a}},\tag{7}$$

where

$$A_{mn} := \begin{cases} \tilde{\mathfrak{z}}_n^{-1} + \frac{ik}{4\pi} & \text{for } m = n, \\ G(|\mathbf{a}_m - \mathbf{a}_n|) & \text{for } m \neq n, \end{cases}$$
(8)

 $\tilde{\mathfrak{z}}_n$ are the renormalized coupling constants, and $X_n := \tilde{\mathfrak{z}}_n u(\mathbf{a}_n)$. Solving (7) for X_n , substituting the result in (6), and noting that the scattering amplitude $\tilde{u}_s(\mathbf{r})$ is given by

$$u(\mathbf{r}) \to u_0(\mathbf{r}) + \tilde{u}_s(\mathbf{r}) \frac{e^{ikr}}{r} \quad \text{for} \quad r \to \infty,$$

we find

$$\tilde{u}_s(\mathbf{r}) = -\frac{1}{4\pi} \sum_{m,n=1}^N A_{mn}^{-1} e^{ik(\mathbf{a}_n \cdot \hat{a} - \mathbf{a}_m \cdot \hat{s})}, \qquad (9)$$

where A_{mn}^{-1} are the entries of the inverse of the $N \times N$ matrix $\mathbf{A} = [A_{mn}]$, and we have also made use of (3).

For the \mathcal{PT} -symmetric double-delta-function potential (1), N = 2, $\tilde{\mathfrak{z}}_1 = \tilde{\mathfrak{z}}_2^* = -k^2 \tilde{\alpha}(\tilde{\sigma} + i\tilde{\gamma})$, $\tilde{\alpha}, \tilde{\sigma}, \tilde{\gamma}$ are real renormalized parameters, and $\mathbf{a}_1 = -\mathbf{a}_2 = \mathbf{r}_0$. Subtituting these relations and Eq. (8) in Eq. (9), we obtain

$$\tilde{u}_s(\mathbf{r}) = \frac{1}{2\pi D} \left[\frac{\mathfrak{s}\cos\xi_- - \mathfrak{g}\sin\xi_-}{k^2(\mathfrak{s}^2 + \mathfrak{g}^2)} + \frac{e^{2ikr_0}\cos\xi_+ - 2ikr_0\cos\xi_-}{8\pi r_0} \right]$$

where

$$D := \det \mathbf{A} = \frac{2\pi - ik^3\mathfrak{s}}{2\pi k^4(\mathfrak{s}^2 + \mathfrak{g}^2)} - \frac{4k^2r_0^2 + e^{4ikr_0}}{64\pi^2r_0^2},$$

- J. A. Rebouças and P. A. Brandão, "Scattering of light by a parity-time-symmetric dipole beyond the first Born approximation," Phys. Rev. A 104, 063514 (2021).
- [2] Yu. N. Demkov and V. N. Ostrovskii, Zero-Range Potentials and Their Applications in Atomic Physics (Plenum press, New York, 1988).
- [3] S. Albeverio, F. Gesztesy, R. Hoegh-Krohn, and H. Holden, *Solvable Models in Quantum Mechanics* (American Mathematical Society, Providence, RI, 2005).
- [4] H. V. Bui and A. Mostafazadeh, "Geometric scattering of

 $\mathfrak{s} := \tilde{\alpha}\tilde{\sigma}, \mathfrak{g} := \tilde{\alpha}\tilde{\gamma}, r_0 := |\mathbf{r}_0|, \text{ and } \xi_{\pm} := k\mathbf{r}_0 \cdot (\hat{a} \pm \hat{s}).$ Notice that the parameters \mathfrak{s} and \mathfrak{g} enter our calculations after we renormalize the bare coupling constants $\alpha\sigma$ and $\alpha\gamma$. Therefore, they may depend on other physical parameters of the problem.

Acknowledgements: This work has been supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK) in the framework of the Project No. 120F061 and by the Turkish Academy of Sciences (TÜBA).

a scalar particle moving on a curved surface in the presence of point defects" Ann. Phys. (NY) **407**, 228 (2019).

- [5] R. Jackiw, "Delta-function potentials in two- and threedimensional quantum mechanics," in: *M.A.B. Beg Memorial Volume*, eds. A. All and P. Hoodbhoy (World Scientific, Singapore, 1991).
- [6] B. Altunkaynak, F. Erman, and O. T. Turgut, "Finitely many Dirac-delta interactions on Riemannian manifolds," J. Math. Phys. 47, 082110 (2006).