Chemotaxis systems with singular sensitivity and logistic source: Boundedness, persistence, absorbing set, and entire solutions

Halil Ibrahim Kurt and Wenxian Shen Department of Mathematics and Statistics Auburn University Auburn University, AL 36849 U.S.A.

Abstract

This paper deals with the following parabolic-elliptic chemotaxis system with singular sensitivity and logistic source,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (\frac{u}{v} \nabla v) + u(a(t, x) - b(t, x)u), & x \in \Omega, \\ 0 = \Delta v - \mu v + \nu u, & x \in \Omega, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial\Omega, \end{cases}$$
(0.1)

where $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain, a(t, x) and b(t, x) are positive smooth functions, and χ , μ and ν are positive constants. In recent years, a lot of attention has been drawn to the question of whether logistic kinetics prevents finite-time blow-up in various chemotaxis models. In the very recent paper [25], we proved that for given nonnegative initial function $0 \neq u_0 \in C^0(\bar{\Omega})$ and $s \in \mathbb{R}$, (0.1) has a unique globally defined classical solution $(u(t,x;s,u_0), v(t,x;s,u_0))$ with $u(s,x;s,u_0) = u_0(x)$, provided that $a_{\inf} = \inf_{t \in \mathbb{R}, x \in \Omega} a(t,x)$ is large relative to χ and u_0 is not small.

In this paper, we further investigate qualitative properties of globally defined positive solutions of (0.1) under the assumption that a_{\inf} is large relative to χ and u_0 is not small. Among others, we provide some concrete estimates for $\int_{\Omega} u^{-p}$ and $\int_{\Omega} u^q$ for some p > 0 and $q > \max\{2, N\}$ and prove that any globally defined positive solution is bounded above and below eventually by some positive constants independent of its initial functions. We prove the existence of a "rectangular" type bounded invariant set (in L^q) which eventually attracts all the globally defined positive solutions. We also prove that (0.1) has a positive entire classical solution ($u^*(t, x), v^*(t, x)$), which is periodic in t if a(t, x) and b(t, x) are periodic in t and is independent of t if a(t, x) and b(t, x) are independent of t.

Key words. Parabolic-elliptic chemotaxis system, logistic source, singular sensitivity, global boundedness, absorbing set, entire positive solution, pointwise persistence, stationary positive solution, periodic positive solution.

1 Introduction and Main Results

Chemotaxis systems, also known as Keller-Segel systems, have been widely studied since the pioneering works [22, 23] by Keller and Segel at the beginning of 1970s on the mathematical modeling of the aggregation process of Dictyostelium discoideum. The current paper is devoted to the study of the asymptotic dynamics of the following parabolic-elliptic chemotaxis system,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (\frac{u}{v} \nabla v) + u(a(t, x) - b(t, x)u), & x \in \Omega, \\ 0 = \Delta v - \mu v + \nu u, & x \in \Omega, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial\Omega, \end{cases}$$
(1.1)

where $\Omega \subset \mathbb{R}^N$ is a smooth bounded domain, a(t, x) and b(t, x) are nonnegative smooth functions, and χ, μ and ν are positive constants. Biologically, (1.1) describes the evolution of a biological process in which cells (with density u) move towards higher concentrations of a chemical substance with density v produced by cells themselves. In (1.1), the cross-diffusion term $-\chi \nabla \cdot (\frac{u}{v} \nabla v)$ reflects the chemotactic movement and $\frac{\chi}{v}$ is referred to as chemotaxis sensitivity; the reaction term u(a(t, x) - b(t, x)u) represents the cell kinetic mechanism and is referred to as logistic source; $\mu > 0$ denotes the degradation rate of the chemical substance, and $\nu > 0$ is the rate at which the mobile species produces the chemical substance. It is seen that the chemotaxis sensitivity $\frac{\chi}{v}$ is singular near v = 0, reflecting an inhibition of chemotactic migration at high signal concentrations. Such a sensitivity was first proposed in [24] due to the Weber–Fechner law of stimulus perception. The time and space dependence of the logistic source reflects the heterogeneity of the underlying environment of the chemotaxis system.

Since the pioneering works of Keller and Segel ([22, 23, 24]), considerable efforts have been devoted to identifying the effects of the cross-diffusion and the kinetic term on the blow-up or global boundedness of solutions of (1.1). For example, consider (1.1) without logistic source (i.e. $a(x,t) = b(x,t) \equiv 0$) and $\mu = \nu = 1$. When Ω being a ball, it is shown in [30] that the classical radially symmetric positive solutions are global and bounded when $\chi > 0$ and N = 2, or $\chi < \frac{2}{N-2}$ and $N \ge 3$, and there exist radial blow-up solutions if $\chi > \frac{2N}{N-2}$ and $N \ge 3$. Without the requirement for symmetry, Biler in [3] proved the global existence of positive solutions when $\chi \le 1$ and N = 2, or $\chi < \frac{2}{N}$ and $N \ge 2$. Fujie, Winkler, and Yokota in [13] proved the boundedness of globally defined positive solutions when $\chi < \frac{2}{N}$ and $N \ge 2$. More recently, Fujie and Senba in [11] proved the global existence of classical positive solutions for the case of N = 2 for any $\chi > 0$. The existence of finite-time blow-up is then completely ruled out for any $\chi > 0$ in the case N = 2. In [4], global existence of weak solutions is proved if $0 < \chi < \frac{N}{N-2}$.

Consider (1.1) with a(t, x), b(t, x) > 0. Central questions include whether the logistic source prevents the occurrence of finite-time blow-up in (1.1) (i.e. any positive solution exists globally); if so, whether the logistic source prevents the occurrence of infinite-time blow-up in (1.1) (i.e. any globally defined positive solution is bounded), and what is the long time behavior of globally defined bounded positive solutions, etc. To recall the existing results related to these central questions, we first make the following standing assumption on a(t, x) and b(t, x): (H) a(t, x) and b(t, x) are continuous in $x \in \overline{\Omega}$ uniformly with respect to $t \in \mathbb{R}$, Hölder continuous in $t \in \mathbb{R}$ with exponent $0 < \gamma_0 < 1$ uniformly with respect to $x \in \overline{\Omega}$, i.e., there is $B_1 > 0$ such that

$$|a(t,x) - a(s,x)| \le B_1 |t-s|^{\gamma_0}, \quad |b(t,x) - b(s,x)| \le B_1 |t-s|^{\gamma_0} \quad \forall t,s \in \mathbb{R}, \ x \in \overline{\Omega},$$

and there are positive constants α , B_2 such that

$$\alpha \le a(t,x) \le B_2, \quad \alpha \le b(t,x) \le B_2$$

Put

$$a_{\inf} = \inf_{x \in \bar{\Omega}, t \in \mathbb{R}} a(t, x), \ b_{\inf} = \inf_{x \in \bar{\Omega}, t \in \mathbb{R}} b(t, x), \ a_{\sup} = \sup_{x \in \bar{\Omega}, t \in \mathbb{R}} a(t, x), \ b_{\sup} = \sup_{x \in \bar{\Omega}, t \in \mathbb{R}} b(t, x).$$

We consider the classical solutions (u(t,x), v(t,x)) of (1.1) with initial functions $u_0(x)$ satisfying

$$u_0 \in C^0(\overline{\Omega}), \quad u_0 \ge 0, \quad \text{and} \quad \int_{\Omega} u_0 > 0.$$
 (1.2)

Definition 1.1. For given $s \in \mathbb{R}$ and u_0 satisfying (1.2), we say (u(t, x), v(t, x)) is a classical solution of (1.1) on (s, s + T) for some $T \in (0, \infty]$ with initial condition $u(s, x) = u_0(x)$ if

$$u(\cdot, \cdot) \in C([s, s+T) \times \overline{\Omega}) \cap C^{1,2}((s, s+T) \times \overline{\Omega}), \quad v(\cdot, \cdot) \in C^{0,2}((s, s+T) \times \overline{\Omega}),$$
$$\lim_{t \to s+} \|u(t, \cdot; s, u_0) - u_0(\cdot)\|_{C^0(\overline{\Omega})} = 0, \tag{1.3}$$

and (u(t,x), v(t,x)) satisfies (1.1) for all $(t,x) \in (s, s+T) \times \Omega$.

Sometime, we may assume

$$a_{\inf} > \begin{cases} \frac{\mu\chi^2}{4}, & \text{if } 0 < \chi \le 2, \\ \mu(\chi - 1), & \text{if } \chi > 2. \end{cases}$$
(1.4)

The following proposition is on the existence and uniqueness of the classical solutions of (1.1) with given initial function u_0 satisfying (1.2) and follows from the arguments in [12, Lemma 2.2].

Proposition 1.1. (Local existence) For any $s \in \mathbb{R}$ and u_0 satisfying (1.2), there is $T_{\max}(s, u_0) \in (0, \infty]$ such that the system (1.1) possesses a unique classical solution, denoted by $(u(t, x; s, u_0), v(t, x; s, u_0))$, on $(s, T_{\max}(s, u_0))$ with initial condition $u(s, x; s, u_0) = u_0(x)$. Furthermore, if $T_{\max}(s, u_0) < \infty$, then

$$\limsup_{t \nearrow T_{\max}(s,u_0)} \|u(t,\cdot;s,u_0)\|_{C^0(\bar{\Omega})} = \infty \quad or \quad \liminf_{t \nearrow T_{\max}(s,u_0)} \inf_{x \in \Omega} v(t,\cdot;s,u_0) = 0.$$

Note that, by the assumption (**H**), it is not difficult to prove that $\inf_{s \in \mathbb{R}} T_{\max}(s, u_0) > 0$.

We say that finite-time blow-up occurs in (1.1) if for some $s \in \mathbb{R}$ and u_0 satisfying (1.2), $T_{\max}(s, u_0) < \infty$, and infinite-time blow-up occurs if for some $s \in \mathbb{R}$ and u_0 satisfying (1.2), $T_{\max}(s, u_0) = \infty$ and

$$\limsup_{t \to \infty} \|u(t, \cdot; s, u_0)\|_{C^0(\bar{\Omega})} = \infty \quad or \quad \liminf_{t \to \infty} \inf_{x \in \Omega} v(t, \cdot; s, u_0) = 0.$$

Throughout the rest of this section, we assume that $u_0(x)$ satisfies (1.2) and $(u(t, x), v(t, x)) := (u(t, x; s, u_0), v(t, x; s, u_0))$ is the unique classical solution of (1.1) on $(s, T_{\max}(s, u_0))$ with the initial condition $u(s, x; s, u_0) = u_0(x)$. We now recall some existing results related to the central questions mentioned in the above.

When N = 2 and $a(t, x) \equiv a$, $b(t, x) \equiv b$ are positive constants, the following have been proved:

- (a) For any u_0 satisfying (1.2) and $s \in \mathbb{R}$, $T_{\max}(s, u_0) = \infty$, that is, finite-time blow-up does not occur (see [12, Theorem 1.1]).
- (b) If (1.4) holds, then for any u_0 satisfying (1.2) and $s \in \mathbb{R}$, $\sup_{t \ge s} ||u(t, \cdot; s, u_0)||_{\infty} < \infty$, that is, infinite-time blow-up does not occur under the assumption (1.4) ([12, Theorem 1.2]).
- (c) The constant solution $(\frac{a}{b}, \frac{\nu}{\mu}\frac{a}{b})$ is exponentially stable under the assumption (1.4) and some other assumptions (see [6, Theorem 1]).

The authors of the current paper studied the global existence of classical solutions of (1.1) for general $N \ge 1$ and time and space dependent functions a(t, x), b(t, x) in [25]. To recall some results proved in [25], we first state some additional conditions on initial data u_0 and on the coefficients in (1.1). Here is an additional condition on the initial function u_0 :

$$u_0 \text{ satisfies (1.2)} \text{ and } \exists \tau_0 > 0 \text{ s.t. } \int_{\Omega} u^{-1}(s + \tau_0, x; s, u_0) \le \frac{b_{\sup} |\Omega| \max\{1, \frac{1}{\chi}\}}{a_{\inf} - a_{\chi,\mu}} \quad \forall s \in \mathbb{R}.$$
 (1.2)'

By [25, Proposition 1.3] (see also Lemma 3.2), for any $q \ge 3$ and $q-1 \le k < 2q-2$, there exist positive constants M(k,q) > 0 and $M^*(k,q) > 0$ such that for any given $s \in \mathbb{R}$ and u_0 satisfying (1.2),

$$\int_{\Omega} \frac{|\nabla v(x,t;s,u_0)|^{2q}}{v^{k+1}(x,t;s,u_0)} \le M(k,q) \int_{\Omega} \frac{u^q(x,t;s,u_0)}{v^{k-q+1}(x,t;s,u_0)} + M^*(k,q) \int_{\Omega} v^{2q-k-1}(x,t;s,u_0)$$

for all $t \in (s, T_{\max}(s, u_0))$. The following is an additional condition on the parameters:

$$a_{\inf} > a_{\chi,\mu} + \frac{b_{\sup} |\Omega| (p_N - 1) (C_n^*)^{\frac{1}{p_N + 1}} \max\{\chi, \chi^2\}}{4b_{\inf} \delta_0},$$
(1.5)'

where δ_0 is as in Lemma 2.3, $p_N = \max\{2, N\}$, $C_N^* = M(p_N, p_N + 1)$, and

$$a_{\chi,\mu} := 2(\chi + 2 - 2\sqrt{\chi + 1})\mu.$$

Note that if $\int_{\Omega} u_0^{-1}(x) dx \leq \frac{b_{\sup} |\Omega| \max\{1, \frac{1}{\chi}\}}{a_{\inf} - a_{\chi,\mu}}$, then (1.2)' holds. The condition (1.2)' indicates that u_0 is not small, which prevents v becomes too small as time evolutes and is a natural assumption. Note also that $a_{\chi,\mu}$ in (1.5)' satisfies

$$a_{\chi,\mu} \le \begin{cases} \frac{\mu\chi^2}{2}, & \text{if } 0 < \chi \le 2\\ 2\mu(\chi - 1), & \text{if } \chi > 2. \end{cases}$$

The condition (1.5)' indicates that $a(\cdot, \cdot)$ is large relative to the chemotaxis sensitivity coefficient χ , which also prevents v becomes too small as time evolutes, and is a natural condition. Assuming that (1.2)' and (1.5)' hold, among others, the following are proved in the recent paper [25]:

(i) (Global existence) For any $s \in \mathbb{R}$,

$$T_{\max}(s, u_0) = \infty \tag{1.5}$$

(see [25, Theorem 1.2(3)]).

(ii) (Boundedness of $\int_{\Omega} u^q$) There is q > N such that for any $s \in \mathbb{R}$,

$$\sup_{t \ge s} \int_{\Omega} u^q(t, x; s, u_0) dx < \infty$$
(1.6)

(see [25, Theorem 1.1(3)]).

(iii) (Boundedness of $\int_{\Omega} u^{-p}$) There is p > 0 such that for any $s \in \mathbb{R}$,

$$\limsup_{t \to \infty} \int_{\Omega} u^{-p}(t, x; s, u_0) dx < \infty$$
(1.7)

(see [25, Lemma 3.4]).

(iv) (Mass persistence) For any $s \in \mathbb{R}$,

$$\inf_{t \ge s} \int_{\Omega} u(t, x; s, u_0) dx > 0, \quad \inf_{t \ge s, x \in \Omega} v(t, x; s, u_0) > 0$$
(1.8)

(see [25, Proposition 1.2(2)]).

(v) (Boundedness of $||u||_{\infty}$) For any $s \in \mathbb{R}$,

$$\sup_{t \ge s} \|u(t, \cdot; s, u_0)\|_{\infty} < \infty \tag{1.9}$$

(see [25, Theorem 1.2(3)]).

We remark that (1.9) implies (1.6). But (1.9) is proved in [25] by using (1.6) and (1.8), and (1.8) is proved in [25] by using (1.7). Hence the boundedness of $\int_{\Omega} u^q$ in (1.6) and the boundedness of $\int_{\Omega} u^{-p}$ in (1.7) play essential roles in the proofs of the main results in [25].

The results (a)-(c) and (i)-(v) provide some answers to the central questions mentioned in the above. For example, results (i)-(v) imply that logistic kinetics prevents the occurrence of finite-time blow-up provided that the intrinsic growth rate function a(t,x) is large relative to the chemotaxis sensitivity and the initial condition u_0 is not small, which is an interesting biological phenomenon. However, there are still many interesting questions associated to those central questions. For example, whether the ultimate upper bound of $\int_{\Omega} u^q$, $\int_{\Omega} u^{-p}$, and $||u||_{\infty}$ in (ii), (iii), and (v), respectively, are independent of the initial functions; whether the mass persistence in (iv) (i.e. $\int_{t\geq s} \int_{\Omega} u(t,x;s,u_0)dx > 0$) can be replaced by the pointwise persistence (i.e. $\liminf_{t-s\to\infty} \inf_{x\in\Omega} u(t,x;s,u_0) > 0$); whether (1.1) has bounded positive entire solutions (i.e. bounded positive solutions (u(t,x), v(t,x)) which are defined for all $t \in \mathbb{R}$); whether (i)-(v) hold without the assumptions (1.2)' and (1.5)'.

In the current paper, we will further investigate those central questions mentioned in the above. We state our main results in next subsection.

1.1 Main results and remarks

Assume that (1.5)' holds. For any $s \in \mathbb{R}$, $\tau \geq s$, u_0 satisfying (1.2)', and p > 0, let

$$m^*(\tau, s, u_0) = \max\left\{\int_{\Omega} u(\tau, x; s, u_0) dx, \frac{a_{\sup}}{b_{\inf}} |\Omega|\right\}$$

and

$$\tilde{M}_1(p,\tau,s,u_0) = pb_{\sup}|1-p|\left(m^*(\tau,s,u_0) - \frac{a_{\sup}}{b_{\inf}}|\Omega|\right).$$

Note that

$$u(t,x;s,u_0), v(t,x;s,u_0) > 0 \quad \forall x \in \Omega, \ t \in (s,\infty)$$

A positive entire solution of (1.1) is a solution (u(t,x), v(t,x)) which is defined for all $t \in \mathbb{R}$ and $\inf_{t \in \mathbb{R}, x \in \Omega} u(t,x) > 0$.

The first main result of the current paper is on the boundedness of $\int_{\Omega} u^{-p}$ and $\int_{\Omega} u^{q}$ of the globally defined classical solutions of (1.1).

Theorem 1.1 (Boundedness of $\int_{\Omega} u^{-p}$ and $\int_{\Omega} u^{q}$). Assume that (1.5)' holds. Then the following hold.

(1) Let
$$p = 1$$
, $\gamma = a_{\inf} - a_{\chi,\mu}$, $M_1(p) = \frac{b_{\sup}|\Omega|}{a_{\inf} - a_{\chi,\mu}}$. Then for any u_0 satisfying $(1.2)'$, $s \in \mathbb{R}$ and $\tau > s$,

$$\int_{\Omega} u^{-p}(t,x;s,u_0) dx \le e^{-\gamma(t-\tau)} \int_{\Omega} u^{-p}(\tau,x;s,u_0) dx + M_1(p) + \tilde{M}_1(p,u_0,\tau) \quad \forall t \ge \tau$$

and

$$\int_{\Omega} u^{-p}(t,x;s,u_0) dx \le \max\left\{\int_{\Omega} u^{-p}(\tau,x;s,u_0) dx, M_1(p) + \tilde{M}_1(p,\tau,s,u_0)\right\} \quad \forall \ t \ge \tau,$$
(1.10)

as well as

$$\lim_{t-s \to \infty} \int_{\Omega} u^{-p}(t,x;s,u_0) \le M_1(p).$$
(1.11)

(2) Let p > 0 be as in (1). There are $q > \max\{2, N\}$ and $M_2(p,q) > 0$ such that for any $s \in \mathbb{R}$, u_0 satisfying (1.2)', and $\tau > s$, there is $\tilde{M}_2(p,q,\tau,s,u_0)$ such that

$$\int_{\Omega} u^{q}(t,x;s,u_{0})dx \le e^{-(t-\tau)} \int_{\Omega} u^{q}(\tau,x;s,u_{0})dx + M_{2}(p,q) + \tilde{M}_{2}(p,q,\tau,s,u_{0}) \quad \forall t \ge \tau,$$

and

$$\int_{\Omega} u^q(t,x;s,u_0) dx \le \max\left\{\int_{\Omega} u^q(\tau,x;s,u_0) dx, M_2(p,q) + \tilde{M}_2(p,q,\tau,s,u_0)\right\} \quad \forall t \ge \tau,$$

as well as

$$\limsup_{t-s\to\infty}\int_{\Omega}u^q(t,x;s,u_0)ds\leq M_2(p,q).$$

- **Remark 1.1.** (1) Theorem 1.1(1) improves boundedness result of $\int_{\Omega} u^{-p}(t, x; s, u_0) dx$ in [25, Lemma 3.4] (see (iii) in the above) in the following two aspects. First, it provides some concrete estimates for $\int_{\Omega} u^{-p}(t, x; s, u_0) dx$. Second, it provides an ultimate upper bound independent of u_0 for $\int_{\Omega} u^{-p}(t, x; s, u_0) dx$.
 - (2) Theorem 1.1(2) improves the boundedness result of $\int_{\Omega} u^q(t,x;s,u_0)dx$ in [25, Theorem 1.1(2)] (see (ii) in the above) in the following two aspects. First, it provides some concrete estimates for $\int_{\Omega} u^q(t,x;s,u_0)dx$. Second, it provides an ultimate upper bound independent of u_0 for $\int_{\Omega} u^q(t,x;s,u_0)dx$.
 - (3) The inequality (1.11) provides some improvement of (1.8) proved in [25]. To be more precise, let p be as in Theorem 1.1(1). By the Hölder's inequality, for any $s \in \mathbb{R}$ and u_0 satisfying (1.2)', we have

$$\begin{aligned} |\Omega| &= \int_{\Omega} u^{\frac{p}{p+1}}(t,x;s,u_0) u^{-\frac{p}{p+1}}(t,x;s,u_0) dx \\ &\leq \left(\int_{\Omega} u(t,x;s,u_0) dx \right)^{\frac{p}{p+1}} \left(\int_{\Omega} u^{-p}(t,x;s,u_0) dx \right)^{\frac{1}{p+1}} \quad \forall t > s. \end{aligned}$$

Then by (1.11),

$$\lim_{t-s \to \infty} \int_{\Omega} u(t,x;s,u_0) dx \ge \frac{|\Omega|^{\frac{p+1}{p}}}{(2M_1(p))^{\frac{1}{p}}},\tag{1.12}$$

which provides an ultimate lower bound independent of u_0 for $\int_{\Omega} u(t, x; s, u_0) dx$ and hence improves (1.8) proved in [25].

(4) The upper bounds of $\int_{\Omega} u^{-p}(t,x;s,u_0)dx$ and $\int_{\Omega} u^q(t,x;s,u_0)dx$ obtained in Theorem 1.1 provide some useful tool for the study of the asymptotic behavior of globally defined positive solutions of (1.1) (see Theorems 1.2-1.4 in the following).

The second main result of this paper is on the existence of globally absorbing sets, which provides some insight on the asymptotic behavior of globally defined positive solutions of (1.1).

Theorem 1.2 (Globally absorbing rectangle). Assume that (1.5)' holds. Let $q > \max\{2, N\}$ and p > 0 be as in Theorem 1.1. There are $M_0^* > 0$, $M_1^*, M_2^* > 0$ such that the following hold.

(1) The set

$$\mathcal{E} = \left\{ u \in C^{0}(\bar{\Omega}) \, | \, u \ge 0, \, \int_{\Omega} u(x) dx \le M_{0}^{*}, \int_{\Omega} u^{-p}(x) dx \le M_{1}^{*}, \, \int_{\Omega} u^{q}(x) dx \le M_{2}^{*} \right\} (1.13)$$

is an invariant set of (1.1) in the sense that for any $u_0 \in \mathcal{E}$ and $s \in \mathbb{R}$, u_0 satisfies (1.2)' and $u(t, \cdot; s, u_0) \in \mathcal{E}$ for all $t \geq s$. Moreover, for any $0 < \theta < 1 - \frac{2N}{q}$ and $\tau > 0$, there is $M_3^*(\theta, \tau) > 0$ such that for any $u_0 \in \mathcal{E}$ and $s \in \mathbb{R}$,

$$\|u(t,\cdot;s,u_0)\|_{C^{\theta}(\bar{\Omega})} \le M_3^*(\theta,\tau) \quad \forall t \ge s+\tau.$$

$$(1.14)$$

(2) The set \mathcal{E} is globally absorbing in the sense that for any u_0 satisfying (1.2)' and $s \in \mathbb{R}$,

$$\begin{cases} \limsup_{t-s\to\infty} \int_{\Omega} u(t,x;s,u_0) \le M_0^* \\ \limsup_{t-s\to\infty} \int_{\Omega} u^{-p}(t,x;s,u_0) dx \le M_1^* \\ \limsup_{t-s\to\infty} \int_{\Omega} u^q(t,x;s,u_0) dx \le M_2^*. \end{cases}$$
(1.15)

Moreover, for any $0 < \theta < 1 - \frac{2N}{q}$, there is $M_4^*(\theta) > 0$ such that for any u_0 satisfying (1.2)', there is $T(u_0) > 0$ such that

$$\|u(t,\cdot;s,u_0)\|_{C^{\theta}(\bar{\Omega})} \le M_4^*(\theta) \quad \forall s \in \mathbb{R}, \ t \ge s + T(u_0).$$

$$(1.16)$$

Remark 1.2. (1) Assume that (1.5)' holds. (1.16) provides some improvement of (1.9) proved in [25]. In fact, by (1.16), for any $s \in \mathbb{R}$ and u_0 satisfying (1.2)',

$$\|u(t,\cdot;s,u_0)\|_{\infty} \le M_4^*(\theta) \quad \forall t-s \gg 1,$$

$$(1.17)$$

which provides an ultimate upper bound independent of u_0 for $||u(t, \cdot; s, u_0)||_{\infty}$ and hence improves (1.9) proved in [25].

(2) Theorem 1.2(2) implies that \mathcal{E} is both pullback absorbing and forward absorbing in the sense that for any u_0 satisfying (1.2)',

$$\begin{cases} \limsup_{s \to -\infty} \int_{\Omega} u(t, x; s, u_0) dx \leq M_0^* & \forall t \in \mathbb{R} \\ \limsup_{s \to -\infty} \int_{\Omega} u^{-p}(t, x; s, u_0) dx \leq M_1^* & \forall t \in \mathbb{R} \\ \limsup_{s \to -\infty} \int_{\Omega} u^q(t, x; s, u_0) dx \leq M_2^* & \forall t \in \mathbb{R}, \end{cases}$$

and

$$\begin{cases} \limsup_{t \to \infty} \int_{\Omega} u(t, x; s, u_0) \leq M_0^* & \forall \ s \in \mathbb{R} \\ \limsup_{t \to \infty} \int_{\Omega} u^{-p}(t, x; s, u_0) dx \leq M_1^* & \forall \ s \in \mathbb{R} \\ \limsup_{t \to \infty} \int_{\Omega} u^q(t, x; s, u_0) dx \leq M_2^* & \forall \ s \in \mathbb{R}, \end{cases}$$

respectively.

(3) Note that the existence of bounded absorbing sets and eventual initial-independent boundedness of positive solutions are strongly related. It should be pointed out that the existence of absorbing sets and eventual boundedness of positive solutions have been studied in various logistic chemotaxis models. For example, in [39], Winkler studied the existence of bounded absorbing sets in L[∞] for the following parabolic-elliptic chemotaxis model with regular sensitivity and logistic type source,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + g(u), & x \in \Omega \\ 0 = \Delta v - v + u, & x \in \Omega \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial \Omega, \end{cases}$$

where $g(u) = Au - Bu^{\alpha}$ for some A, B > 0 and $\alpha > 1$. We refer the reader to [34], [35] and references therein for the study of eventual boundedness of positive solutions of the following parabolic-parabolic chemotaxis model,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + g(u), & x \in \Omega \\ v_t = \Delta v - v + u, & x \in \Omega \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial \Omega, \end{cases}$$

where g(u) is a logistic type nonlinear function.

The following theorem is on the pointwise persistence, which is strongly related to (1.12) and provides some further insight on the asymptotic behavior of globally defined positive solutions of (1.1).

Theorem 1.3 (Uniform pointwise persistence). Assume that (1.5)' holds. There is $m^* > 0$ such that for any u_0 satisfying (1.2)' and $s \in \mathbb{R}$,

$$\liminf_{t-s \to \infty} \inf_{x \in \Omega} u(t, x; s, u_0) \ge m^*.$$
(1.18)

Remark 1.3. The pointwise persistence (1.18) is proved for the first time for chemotaxis models with singular sensitivity and logistic source, and implies the mass persistence. We point out that mass persistence for chemotaxis systems with regular chemotaxis sensitivity and logistic source was studied in [32] and pointwise persistence was studied in [20, 21].

By (1.17) and (1.18), under the assumption (1.5)', any globally defined positive solution with initial condition u_0 satisfying (1.2)' is eventually bounded above and below by some positive constants independent of its initial condition. It is natural to ask whether globally defined positive solutions converge to some positive entire solution. When $a(t,x) \equiv a$ and $b(t,x) \equiv b$, it is clear that $(\frac{a}{b}, \frac{\nu}{\mu} \frac{a}{b})$ is a positive entire solution of (1.1). When a(t,x) and b(t,x) depend on t and x, it is not clear at all whether (1.1) has positive entire solutions. The last main result of the current paper is on the existence of positive entire solutions of (1.1).

Theorem 1.4 (Existence of positive entire solutions). Suppose that (1.5)' holds. Let \mathcal{E} be as in Theorem 1.2. Then the followings hold.

- (1) There is a positive entire solution $(u^*(t,x), v^*(t,x))$ of (1.1) with $u^*(t, \cdot) \in \mathcal{E}$ for any $t \in \mathbb{R}$.
- (2) If there is T > 0 such that a(t + T, x) = a(t, x) and b(t + T, x) = b(t, x), then (1.1) has a positive T-periodic solution $(u, v) = (u^*(t, x), v^*(t, x))$ with $u^*(t, \cdot) \in \mathcal{E}$ for any $t \in \mathbb{R}$.
- (3) If $a(t,x) \equiv a(x)$ and $b(t,x) \equiv b(x)$, then (1.1) has a positive stationary solution $(u,v) = (u^*(x), v^*(x))$ with $u^*(\cdot) \in \mathcal{E}$.

Remark 1.4. As stated in (c), when N = 2 and $a(t,x) \equiv a$, $b(t,x) \equiv b$, the constant entire solution $(\frac{a}{b}, \frac{\nu}{a}\frac{a}{b})$ is exponentially stable under the assumption (1.4) and some other assumption

(see [6, Theorem 1]). By the arguments of [6, Theorem 1], it can also be proved that, for general $N \ge 1$, $(\frac{a}{b}, \frac{\nu}{\mu} \frac{a}{b})$ is exponentially stable under the assumption (1.4) and some other assumption. But when a(t, x), b(t, x) are not constant functions, the arguments of [6, Theorem 1] are difficult to apply. We plan to study the stability of positive entire solutions of (1.1) somewhere else.

It should be pointed out that a considerable amount of research has also been carried out toward the finite-time blow-up or global boundedness of solutions of the following parabolicparabolic chemotaxis model with singular sensitivity,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (\frac{u}{v} \nabla v) + u(a - bu), & x \in \Omega, \\ v_t = \Delta v - v + u, & x \in \Omega, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial \Omega. \end{cases}$$
(1.19)

For example, for the case that a = b = 0, it is proved that if $0 < \chi < \sqrt{\frac{2}{N}}$, then for any initial data $u_0 \in C^0(\bar{\Omega})$ and $v_0 \in W^{1,\infty}(\Omega)$ with $u_0 \ge 0$ and $v_0 > 0$ in $\bar{\Omega}$, there exists a global-in-time classical solution of (1.19) (see [37]), and moreover, the global-in-time classical solution is bounded (see [9]). When Ω is a smooth, bounded, convex two-dimensional domain, it is shown in [26] that there is $\chi_0 > 1$ such that for any initial data $u_0 \in C^0(\bar{\Omega})$ and $v_0 \in W^{1,\infty}(\Omega)$ with $u_0 \ge 0$ and $v_0 > 0$ in $\bar{\Omega}$, (1.19) has a global bounded solution for $\chi \in (0, \chi_0)$, which implies that $0 < \chi < 1$ is not critical for the global existence of (1.19) on two-dimensional domains. See [10, 27, 37] for more results.

For the case that a, b > 0, it is proved that if N = 2, then for any initial data $0 \le u_0 \in L^2(\Omega)$ and $0 < v_0 \in H^{1+\epsilon_0}(\Omega)$ with $\inf_{x\in\Omega} v_0(x) > 0$ and $\epsilon_0 \in (0, \frac{1}{2})$, (1.19) has a global classical solution (see [1, 40]), and if (1.4) holds with $\mu = 1$ and $a_{\inf} = a$, then globally defined positive solutions of (1.19) are bounded (see [40]) and the constant solution $(\frac{a}{b}, \frac{a}{b})$ is exponentially stable (see [42, Theorem 1.1]). See [41] for the existence of weak solutions of (1.19) with a, b > 0 in the case $N \ge 3$. However, there is little study on the global existence of classical solutions of (1.19) when $N \ge 3$.

We also point out that a considerable amount of research has been carried out toward the finite-time blow-up or global boundedness of solutions of the following chemotaxis model with regular chemotaxis sensitivity and logistic source,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + u(a - bu), & x \in \Omega, \\ \tau v_t = \Delta v - \mu v + \nu u, & x \in \Omega, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial \Omega. \end{cases}$$
(1.20)

For example, when $\tau = 0$ and $\mu = \nu = 1$, it is proved in [33] that, if $N \leq 2$ or $b > \frac{N-2}{N}\chi$, then for every nonnegative initial data $u_0 \in C^0(\bar{\Omega})$, (1.20) possesses a global bounded classical solution which is unique. It should be pointed out that, when a = b = 0 and $N \geq 2$, finite-time blow-up of positive solutions occurs under some condition on the mass and the moment of the initial data (see [17], [18], [28], [29]). Hence the finite time blow-up phenomena in (1.20) is suppressed to some extent by the logistic source. But it remains open whether in any space dimensional setting, for every nonnegative initial data $u_0 \in C^0(\overline{\Omega})$ (1.20) possesses a unique global classical solution for every $\chi > 0$ and every b > 0. It should be pointed out that finite-time blow-up occurs in various variants of (1.20), for example, it occurs in (1.20) with $\tau = 0$, with the logistic source being replaced by logistic-type superlinear degradation (see [31, 38]), and/or with the second equation being replaced by the following one

$$0 = \Delta v - \frac{1}{|\Omega|} \int_{\Omega} u(\cdot, t) + u, \quad x \in \Omega$$

(see [5, 7, 8, 36]).

The rest of this paper is organized in the following way. In section 2, we present some lemmas to be used in later sections. In section 3, we study the boundedness of $\int_{\Omega} u^{-p}$ and $\int_{\Omega} u^{q}$ of globally defined positive solutions of (1.1) and prove the Theorem 1.1. In section 4, we investigate the existence of absorbing invariant sets and the ultimate upper and lower bounds of globally defined positive solutions and prove Theorems 1.2 and 1.3. In section 5, we explore the existence of positive entire solutions of (1.1) and prove the Theorem 1.4.

2 Preliminaries

In this section, we present some lemmas to be used in later sections. Throughout this section, $\mu > 0$ and $\nu > 0$ are fixed. C denotes some generic constant which is independent of solutions, but may not be the same at different places.

First, for convenience, we present two lemmas on fixed point theorems.

Lemma 2.1. If F is a closed subset of a Banach space X, G is a subset of a Banach space Y, $T_y: F \to F, y$ in G is a uniform contraction on F and T_yx is continuous in $y \in G$ for each fixed x in F, then the unique fixed point g(y) of T_y, y in G, is continuous in y.

Proof. See [15, Theorem 3.2 in Chapter 0].

Lemma 2.2. Let G be a closed convex set in a Banach space X and let T be a continuous mapping of G into itself such that the image TG is precompact. Then T has a fixed point.

Proof. It follows from [14, Corollary 11.2].

Next, we present two lemmas on the lower and upper bounds of the solutions of

$$\begin{cases} \Delta v - \mu v + \nu u = 0, & x \in \Omega, \\ \frac{\partial v}{\partial n} = 0, & x \in \partial \Omega. \end{cases}$$
(2.1)

For given $u \in L^p(\Omega)$, let $v(\cdot; u)$ be the solution of (2.1).

Lemma 2.3. Suppose that $u \in C^0(\overline{\Omega})$ is nonnegative and $\int_{\Omega} u > 0$. Then

$$v(x;u) \ge \delta_0 \int_{\Omega} u > 0$$
 in Ω

for some positive constants δ_0 depending only on Ω .

Proof. It follows from the arguments of [13, Lemma 2.1].

Lemma 2.4. For any p > 1, there exists $C_p > 0$ such that

$$\max\left\{\|v(\cdot;u)\|_{L^{p}(\Omega)}, \|\nabla v(\cdot;u)\|_{L^{p}(\Omega)}\right\} \leq C_{p}\|u(\cdot)\|_{L^{p}(\Omega)} \quad \forall u \in L^{p}(\Omega).$$

Proof. It follows from L^p -estimates for elliptic equations (see [2, Theorem 12.1]).

Now, we present some properties of the semigroup generated by $-\Delta + \mu I$ complemented with Neumann boundary condition on $L^p(\Omega)$. For given $1 , let <math>X_p = L^p(\Omega)$ and $A_p = -\Delta + \mu I$ with

$$D(A_p) = \left\{ u \in W^{2,p}(\Omega) \, | \, \frac{\partial u}{\partial n} = 0 \quad \text{on } \partial \Omega \right\}.$$

It is well known that A_p is a sectorial operator in X_p and thus generates an analytic semigroup $(e^{-A_p t})_{t>0}$ in X_p (see, for example, [2, Theorem 13.4]). Moreover $0 \in \rho(A_p)$ and

 $||e^{-A_p t}u||_{X_p} \le e^{-\mu t} ||u||_{X_p}$ for $t \ge 0$ and $u \in X_p$.

Let $X_p^{\alpha} = D(A_p^{\alpha})$ equipped with the graph norm $||u||_{\alpha,p} := ||u||_{X_p^{\alpha}} = ||A_p^{\alpha}u||_{L^p}$.

Lemma 2.5. (i) Let $p \in (1, \infty)$. For each $\beta \ge 0$, there is $C_{p,\beta} > 0$ such that for some $\gamma > 0$,

$$||A_p^{\beta}e^{-A_pt}||_{L^p(\Omega)} \le C_{p,\beta}t^{-\beta}e^{-\gamma t} \text{ for } t > 0.$$

(ii) If $m \in \{0,1\}$ and $q \in [p,\infty]$ are such that $m - \frac{N}{q} < 2\beta - \frac{N}{p}$, then $X_n^\beta \hookrightarrow W^{m,q}(\Omega).$

(iii) If $2\beta - \frac{N}{p} > \theta \ge 0$, then

 $X_p^\beta \hookrightarrow C^\theta(\Omega).$

Proof. (i) It follows from [16, Theorem 1.4.3].

(ii) It follows from [16, Theorem 1.6.1].

(iii) It also follows from [16, Theorem 1.6.1].

Lemma 2.6. Let $\beta \ge 0$, $p \in (1, \infty)$. Then for any $\epsilon > 0$ there exists $C_{p,\beta,\epsilon} > 0$ such that for any $w \in C_0^{\infty}(\Omega)$ we have

$$\|A_p^{\beta}e^{-tA_p}\nabla \cdot w\|_{L^p(\Omega)} \le C_{p,\beta,\epsilon}t^{-\beta-\frac{1}{2}-\epsilon}e^{-\gamma t}\|w\|_{L^p(\Omega)} \quad \text{for all } t > 0 \text{ and some } \gamma > 0.$$
(2.2)

Accordingly, for all t > 0 the operator $A_p^{\beta} e^{-tA_p} \nabla \cdot$ admits a unique extension to all of $L^p(\Omega)$ which is again denoted by $A_p^{\beta} e^{-tA_p} \nabla \cdot$ and satisfies (2.2) for all \mathbb{R}^n -valued $w \in L^p(\Omega)$.

Proof. It follows from [19, Lemma 2.1].

Finally, we present some basic properties of solutions of (1.1). In the rest of this paper, if no confusion occurs, we put $A = A_p$ for some 1 .

Lemma 2.7. Assume (1.5)'. For any given $u_0 \in C^0(\overline{\Omega})$ satisfying (1.2)' and $s \in \mathbb{R}$, there is a unique classical solution $(u(t, x; s, u_0), v(t, x; s, v_0))$ of (1.1) on (s, ∞) with initial condition $u(s, x; s, u_0) = u_0(x)$ (i.e. $T_{\max}(s, u_0) = \infty$). Moreover, $(u(t, x; s, u_0), v(t, x; s, u_0))$ satisfies

$$u(t, \cdot; s, u_0) = e^{-A(t-s)} u_0 - \chi \int_s^t e^{-A(t-\tau)} \nabla \cdot \left(\frac{u(\tau, \cdot; s, u_0)}{v(\tau, \cdot; s, u_0)} \nabla v(\tau, \cdot; s, u_0) \right) d\tau + \int_s^t e^{-A(t-\tau)} u(\tau, \cdot; s, u_0) \left[\mu + a(\tau, \cdot) - b(\tau, \cdot) u(\tau, \cdot; s, u_0) \right] d\tau$$
(2.3)

for any t > s, and if $\inf_{x \in \Omega} u_0(x) > 0$, then for any p > 0,

$$\lim_{t \to s+} \int_{\Omega} u^{-p}(t,x;s,u_0) dx = \int_{\Omega} u_0^{-p}(x) dx.$$
(2.4)

Proof. First, by the arguments of [12, Lemma 2.2] and [25, Theorem 1.2], for any given $u_0 \in C^0(\overline{\Omega})$ satisfying (1.2) and $s \in \mathbb{R}$, there is a unique classical solution $(u(t, x; s, u_0), v(t, x; s, v_0))$ of (1.1) on (s, ∞) with initial condition $u(s, x; s, u_0) = u_0(x)$ and $(u(t, \cdot; s, u_0), v(t, \cdot; s, v_0))$ satisfies (2.3).

Next, if $\inf_{x\in\bar{\Omega}} u_0(x) > 0$, by (1.3),

$$\lim_{t \to s+} u^{-p}(t, x; s, u_0) = u_0^{-p}(x) \quad \text{uniformly in } x \in \bar{\Omega}.$$

This implies (2.4) holds. The lemma is thus proved.

To indicate the dependence of the classical solution $(u(t, x; s, u_0), v(t, x; s, u_0))$ of (1.1) on a(t, x) and b(t, x), we may put

$$(u(t,x;s,u_0,a,b),v(t,x;s,u_0,a,b)) = (u(t,x;s,u_0),v(t,x;s,u_0)).$$

Lemma 2.8. Fix $s \in \mathbb{R}$. Let $u_n, u_0 \in C^0(\overline{\Omega})$ with $u_n(x) \ge 0$ and $a_n(t, x), b_n(t, x)$ satisfy (H). If u_0 satisfies (1.2)', $\lim_{n\to\infty} ||u_n - u_0||_{\infty} = 0$, and

$$\lim_{n \to \infty} \sup_{x \in \Omega} |a_n(t, x) - a_0(t, x)| = 0, \quad \lim_{n \to \infty} \sup_{x \in \Omega} |b_n(t, x) - b_0(t, x)| = 0$$

locally uniformly in $t \in \mathbb{R}$ for some $a_0(t, x)$ and $b_0(t, x)$, then $a_0(t, x), b_0(t, x)$ satisfy (**H**), there is K > 0 such that u_n satisfies (1.2) for $n \geq K$, and

$$\lim_{n \to \infty} \|u(t, \cdot; s, u_n, a_n, b_n) - u(t, \cdot; s, u_0, a_0, b_0)\|_{\infty} = 0 \text{ uniformly in } t \in [s, s+T]$$
(2.5)

for any T > 0.

Proof. This lemma is about the continuity of solutions of (1.1) with respect to initials and the coefficients in the equations. Note that solutions of (1.1) satisfy the integral equation (2.3). It then suffices to prove the continuity of solutions of (2.3) with respect to initials and the coefficients in the equations.

First of all, it is clear that $a_0(t, x)$ and $b_0(t, x)$ satisfy **(H)**. Assume that $u_0 \in C^0(\overline{\Omega})$ satisfies (1.2). For given r > 0, define

$$\mathcal{B}(u_0, r) = \{ u \in C^0(\Omega) \, | \, \|u - u_0\|_{\infty} \le r \}.$$

Fix $0 < r \ll 1$ such that

$$\int_{\Omega} \tilde{u}_0(x) dx \ge \frac{2}{3} \int_{\Omega} u_0(x) dx \quad \text{for any } \tilde{u}_0 \in \mathcal{B}(u_0, r).$$
(2.6)

Since $\lim_{n\to\infty} \|u_n - u_0\|_{\infty} = 0$, there is K > 0 such that $u_n \in \mathcal{B}(u_0, r)$ and $\int_{\Omega} u_n(x) dx \ge \frac{2}{3} \int_{\Omega} u_0(x) dx > 0$ for $n \ge K$. This implies that u_n satisfies **(H)** for $n \ge K$. Without loss of generality, we may assume that $u_n \in \mathcal{B}(u_0, r)$ for all $n \ge 1$. Then $\int_{\Omega} u_n(x) dx > 0$ for all $n \ge 1$. By Lemma 2.7, $u(t, x; s, u_n, a_n, b_n)$ and $v(t, x; s, u_0, a_0, b_0)$ are defined for $t \ge s$.

Next, we note that it suffices to prove that (2.5) holds for $0 < T \ll 1$. To prove this, set

$$\varepsilon := \frac{\delta_0}{2} \int_{\Omega} u_0(x) dx > 0,$$

where δ_0 is as in Lemma 2.3. For given T > 0 and $R > r + ||u_0||_{C^0(\overline{\Omega})}$, we define the Banach space

$$\mathcal{X}_T = C^0([s, s+T], C^0(\bar{\Omega}))$$

equipped with the norm

$$||u||_{\mathcal{X}_T} = \max_{s \le t \le s+T} ||u(t,x)||_{C^0(\bar{\Omega})}.$$

Set

$$\mathcal{S}(T) = \left\{ u \in \mathcal{X}_T : \|u\|_{\mathcal{X}_T} \le R, \text{ and } \nu A^{-1} u \ge \varepsilon \text{ for all } t \in [s, s+T] \right\},\$$

and

$$\mathcal{Y}(T) = \{a(\cdot, \cdot) \in \mathcal{X}_T : |a(t, x) - a(t', x)| \le B_1 | t - t' |^{\gamma}, \ \alpha \le a(t, x) \le B_2 \ \forall t, t' \in [s, s + T], \ x \in \bar{\Omega}\},\$$

where B_1, B_2, α, γ are as in (**H**). It is clear that $\mathcal{S}(T)$ and $\mathcal{Y}(T)$ are closed subsets of the Banach space \mathcal{X}_T .

For $\tilde{u}_0 \in \mathcal{B}(u_0, r)$ and $(a, b) \in \mathcal{Y}(T) \times \mathcal{Y}(T)$, we define

$$(\mathcal{M}(\tilde{u}_0, a, b)u)(t, \cdot) = e^{-A(t-s)}\tilde{u}_0 - \chi \int_s^t e^{-A(t-\tau)} \nabla \cdot \left(\frac{u(\tau, \cdot)}{v(\tau, \cdot)} \nabla v(\tau, \cdot)\right) d\tau + \int_s^t e^{-A(t-\tau)} u(\tau, \cdot) \left[\mu + a(\tau, \cdot) - b(\tau, \cdot)u(\tau, \cdot)\right] d\tau,$$

where $u \in \mathcal{S}(T)$ and $v(\tau, \cdot) = \nu A^{-1}u(\tau, \cdot)$ for $\tau \in [s, s + T]$. It is not difficult to prove that $\mathcal{M}(\tilde{u}_0, a, b)u \in \mathcal{X}_T$ for any $u \in \mathcal{S}(T)$.

We claim that $\mathcal{M}(\tilde{u}_0, a, b)$ maps $\mathcal{S}(T)$ into itself for $0 < T \ll 1$. To see this, let p, β , and $\epsilon > 0$ be such that N < p, $\frac{N}{2p} < \beta < \frac{1}{2}$, and $\epsilon \in (0, \frac{1}{2} - \beta)$. Let $C_p, C_{p,\beta}$, and $C_{p,\beta,\epsilon}$ be as in by Lemma 2.4, Lemma 2.5, and Lemma 2.6, respectively. Then for any $\tilde{u}_0 \in \mathcal{B}(u_0, r)$ and $u \in \mathcal{S}(T)$, we have

$$\|(\mathcal{M}(\tilde{u}_0, a, b)u)(t, \cdot)\|_{C^0(\bar{\Omega})}$$

$$\leq \|e^{-A(t-s)}\tilde{u}_{0}\|_{C^{0}(\bar{\Omega})} + \chi C_{p,\beta} \int_{s}^{t} \|A^{\beta}e^{-A(t-\tau)}\nabla \cdot \left(\frac{u(\tau,\cdot)}{v(\tau,\cdot)}\nabla v(\tau,\cdot)\right)\|_{L^{p}(\Omega)}d\tau \\ + C_{p,\beta} \int_{s}^{t} \|A^{\beta}e^{-A(t-\tau)}u(\tau,\cdot)[\mu + a(\tau,\cdot) - b(\tau,\cdot)u(\tau,\cdot)]\|_{L^{p}(\Omega)}d\tau \\ \leq \|\tilde{u}_{0}\|_{C^{0}(\bar{\Omega})} + \frac{\chi}{\varepsilon}C_{p,\beta}C_{p,\beta,\epsilon}C_{p}|\Omega|^{1/p}R^{2}\int_{s}^{t}(t-\tau)^{-\beta-\frac{1}{2}-\epsilon}d\tau \\ + C_{p,\beta}^{2}|\Omega|^{1/p}(R\mu + RB_{2} + R^{2}B_{2})\int_{s}^{t}(t-\tau)^{-\beta}d\tau \\ \leq \|u_{0}\|_{C^{0}(\bar{\Omega})} + r + \frac{\chi}{\varepsilon}C_{p,\beta}, C_{p,\beta,\epsilon}C_{p}|\Omega|^{1/p}R^{2}T^{\frac{1}{2}-\beta-\epsilon} + C_{p,\beta}^{2}|\Omega|^{1/p}R(\mu+B_{2}+RB_{2})T^{1-\beta}$$

for all $t \in [s, s+T]$. We then have $\|\mathcal{M}(u_0, a, b)u\|_{\mathcal{X}_T} \leq R$ if $T \in (0, 1)$ is suitably small such that

$$T \le \left(\frac{R - r - \|u_0\|_{C^0(\bar{\Omega})}}{\frac{\chi}{\varepsilon} C_{p,\beta} C_{p,\beta,\epsilon} C_p |\Omega|^{1/p} R^2 + C_{p,\beta}^2 |\Omega|^{1/p} R(\mu + B_2 + RB_2)}\right)^{\frac{1}{\frac{1}{2} - \beta - \epsilon}}.$$
(2.7)

Note that

$$\begin{split} \int_{\Omega} (e^{-A(t-s)}\tilde{u}_0)(t,x)dx &= e^{-(t-s)} \int_{\Omega} e^{(t-s)\Delta} \tilde{u}_0(x)dx \ge e^{-T} \int_{\Omega} \tilde{u}_0(x)dx, \\ \int_{\Omega} \left[\int_s^t e^{-A(t-\tau)} \nabla \cdot \left(\frac{u}{v} \nabla v\right) d\tau \right] dx &= e^{-(t-s)} \int_s^t \left[\int_{\Omega} e^{(t-\tau)\Delta} \nabla \cdot \left(\frac{u}{v} \nabla v\right) dx \right] d\tau \\ &= e^{-(t-s)} \int_s^t \left[\int_{\Omega} \nabla \cdot \left(\frac{u}{v} \nabla v\right) dx \right] d\tau \\ &= e^{-(t-s)} \int_s^t 0 d\tau = 0, \end{split}$$

and

$$\begin{split} \int_{\Omega} \left[\int_{s}^{t} e^{-A(t-\tau)} u \big(\mu + a(\tau, \cdot) - b(\tau, \cdot) u \big) d\tau \right] dx &= e^{-(t-s)} \int_{s}^{t} \left[\int_{\Omega} e^{(t-\tau)\Delta} u \big(\mu + a(\tau, \cdot) - b(\tau, \cdot) u \big) dx \right] d\tau \\ &= e^{-(t-s)} \int_{s}^{t} \left[\int_{\Omega} (\mu + a(\tau, \cdot)) u - b(\tau, \cdot) u^{2} dx \right] d\tau \\ &\geq -B_{2} |\Omega| R^{2} T e^{-T}. \end{split}$$

It then follows that

$$\begin{split} \int_{\Omega} (\mathcal{M}(\tilde{u}_0, a, b)u)(t, x) dx &= \int_{\Omega} e^{-A(t-s)} \tilde{u}_0(x) dx + \int_{\Omega} \left[\int_s^t e^{-A(t-\tau)} \nabla \cdot \left(\frac{u(\tau, x)}{v(\tau, x)} \nabla v(\tau, x) \right) d\tau \right] dx \\ &+ \int_{\Omega} \left[\int_s^t e^{-A(t-\tau)} u(\tau, x) \left(\mu + a(\tau, x) - b(\tau, x)u(\tau, x) \right) d\tau \right] dx \\ &\geq e^{-T} \int_{\Omega} \tilde{u}_0(x) dx - e^{-T} B_2 |\Omega| R^2 T \quad \forall t \in [s, s+T]. \end{split}$$

This together with (2.6) implies that, if

$$0 < \frac{6T}{4 - 3e^T} < \frac{1}{B_2 |\Omega| R^2} \int_{\Omega} u_0(x) dx,$$
(2.8)

then

$$\int_{\Omega} (\mathcal{M}(\tilde{u}_0, a, b)u)(t, x) dx \ge \frac{1}{2} \int_{\Omega} u_0(x) dx \quad \forall \, \tilde{u}_0 \in \mathcal{B}(u_0, r), \ (a, b) \in \mathcal{Y}_0(T), \ u \in \mathcal{S}(T).$$

Hence

$$(\nu A^{-1}\mathcal{M}(\tilde{u}_0, a, b)u)(t, x) \ge \delta_0 \int_{\Omega} \mathcal{M}(\tilde{u}_0, a, b)u(x, t)dx \ge \frac{\delta_0}{2} \int_{\Omega} u_0(x)dx = \varepsilon$$

for all $t \in [s, s + T]$. Therefore, the claim holds with any T > 0 satisfies (2.7) and (2.8).

We then prove that the mapping $\mathcal{M}(\tilde{u}_0, a, b)$ is a uniform contraction on $\mathcal{S}(T)$ for $0 < T \ll 1$. Let p, β, ϵ , and $C_p, C_{p,\beta}, C_{p,\beta,\epsilon}$ be as in the above. By similar arguments as in the above, for given $u, w \in \mathcal{S}(T)$ with $v := \nu A^{-1}u, \bar{v} := \nu A^{-1}w$, we have

$$\begin{split} \|(\mathcal{M}(\tilde{u}_{0},a,b)u)(t,\cdot) - (\mathcal{M}(\tilde{u}_{0},a,b)w)(t,\cdot)\|_{C^{0}(\bar{\Omega})} \\ &\leq \chi C_{p,\beta} \int_{s}^{t} \|A^{\beta}e^{-A(t-\tau)}\nabla \cdot \left(\frac{u(\tau,\cdot)}{v(\tau,\cdot)}\nabla v(\tau,\cdot) - \frac{w(\tau,\cdot)}{\bar{v}(\tau,\cdot)}\nabla \bar{v}(\tau,\cdot)\right)\|_{L^{p}(\Omega)}d\tau \\ &+ C_{p,\beta} \int_{s}^{t} \|A^{\beta}e^{-A(t-\tau)}(u(\tau,\cdot) - w(\tau,\cdot))[\mu + a(\tau,\cdot) - b(\tau,\cdot)(u(\tau,\cdot) + w(\tau,\cdot))]\|_{L^{p}(\Omega)}d\tau \\ &\leq \frac{\chi C_{p,\beta}C_{p,\beta,\epsilon}C_{p}|\Omega|^{1/p}}{\varepsilon} \int_{s}^{t} (t-\tau)^{-\beta-\frac{1}{2}-\epsilon}\|u(\tau) - w(\tau)\|_{C^{0}(\Omega)} \\ &\quad \cdot \left(\|u(\tau)\|_{C^{0}(\Omega)} + \|w(\tau)\|_{C^{0}(\Omega)} + \frac{C_{2p}}{\varepsilon}\|w(\tau)\|_{C^{0}(\Omega)}^{2}\right)d\tau \\ &\quad + C_{p,\beta}^{2}|\Omega|^{1/p} \int_{s}^{t} (t-\tau)^{-\beta}\|u(\tau) - w(\tau)\|_{C^{0}(\Omega)} \left(\mu + B_{2} + B_{2}(\|u(\tau)\|_{C^{0}(\Omega)} + \|w(\tau)\|_{C^{0}(\Omega)})\right)d\tau \\ &\leq T^{\frac{1}{2}-\beta-\epsilon} \left(\frac{\chi C_{p,\beta}C_{p,\beta,\epsilon}C_{p}|\Omega|^{1/2}}{\varepsilon^{2}}(2\varepsilon R + C_{2p}R^{2}) + C_{p,\beta}^{2}|\Omega|^{1/p}(\mu + B_{2} + 2RB_{2})\right)\|u-w\|_{\mathcal{X}} \tag{2.9}$$

for all $t \in [s, s + T]$. It then follows that $\mathcal{M}(\tilde{u}_0, a, b)$ is a uniform contraction on $\mathcal{S}(T)$ for any T > 0 satisfying (2.7), (2.8), and

$$T < \left[\frac{\chi C_{p,\beta} C_{p,\beta,\epsilon} C_p |\Omega|^{1/p}}{\varepsilon^2} (2\varepsilon R + C_{2p} R^2) + C_{p,\beta}^2 |\Omega|^{1/p} (\mu + B_2 + 2RB_2)\right]^{\frac{-1}{\frac{1}{2} - \beta - \epsilon}}.$$
 (2.10)

We now prove $\mathcal{M}(\tilde{u}_0, a, b)$ is uniformly continuous in $\tilde{u}_0 \in \mathcal{B}(u_0, r)$ and $(a, b) \in \mathcal{Y}(T) \times \mathcal{Y}(T)$. Note that for any $\tilde{u}_1, \tilde{u}_2 \in \mathcal{B}(u_0, r), (a_1, b_1), (a_2, b_2) \in \mathcal{Y}(T) \times \mathcal{Y}(T)$, and $u \in \mathcal{S}(T)$,

$$\mathcal{M}(\tilde{u}_2, a_2, b_2)u - \mathcal{M}(\tilde{u}_1, a_1, b_1)u = [\mathcal{M}(\tilde{u}_2, a_2, b_2)u - \mathcal{M}(\tilde{u}_1, a_2, b_2)u] \\ + [\mathcal{M}(\tilde{u}_1, a_2, b_2)u - \mathcal{M}(\tilde{u}_1, a_1, b_2)u] \\ + [\mathcal{M}(\tilde{u}_1, a_1, b_2)u - \mathcal{M}(\tilde{u}_1, a_1, b_1)u].$$

It then suffices to prove that $\mathcal{M}(\tilde{u}_0, a, b)$ is uniformly continuous in $\tilde{u}_0 \in \mathcal{B}(u_0, r)$ (resp. uniformly continuous in $a \in \mathcal{Y}(T)$, uniformly continuous in $b \in \mathcal{Y}(T)$). Observe that, for any $\tilde{u}_n, \tilde{u}_0 \in \mathcal{Y}(T)$

 $\mathcal{B}(u_0, r), (a, b) \in \mathcal{Y}_0(T) \times \mathcal{Y}_0(T), \text{ and } u \in \mathcal{S}(T), \text{ by comparison principles for parabolic equations,}$ we have

$$\|(\mathcal{M}(\tilde{u}_n, a, b)u)(t, \cdot) - (\mathcal{M}(\tilde{u}_0, a, b)u)(t, \cdot)\|_{C^0(\bar{\Omega})} = \|e^{-(t-s)A}(\tilde{u}_n - \tilde{u}_0)\|_{C^0(\bar{\Omega})} \le \|\tilde{u}_n - \tilde{u}_0\|_{C^0(\bar{\Omega})}.$$

Thus, the mapping $\mathcal{M}(\tilde{u}_0, a, b)$ is uniformly continuous in $\tilde{u}_0 \in \mathcal{B}(u_0, r)$. For any $\tilde{u}_0 \in \mathcal{B}(u_0, r)$, $a_n, a_0, b \in \mathcal{Y}(T)$, and $u \in \mathcal{S}(T)$,

$$\begin{aligned} \|(\mathcal{M}(\tilde{u}_{0},a_{n},b)u)(t,\cdot) - (\mathcal{M}(\tilde{u}_{0},a_{0},b)u)(t,\cdot)\|_{C^{0}(\bar{\Omega})} &= \|\int_{s}^{t} e^{-A(t-\tau)}u(\tau,\cdot)[a_{n}(\tau,\cdot) - a_{0}(\tau,\cdot)]d\tau\|_{C^{0}(\bar{\Omega})} \\ &\leq R\|a_{n} - a_{0}\|_{\mathcal{X}_{T}}T. \end{aligned}$$

This implies that the mapping $\mathcal{M}(\tilde{u}_0, a, b)$ is uniformly continuous in $a \in \mathcal{Y}(T)$. For any $\tilde{u}_0 \in \mathcal{B}(u_0, r), a, b_n, b_0 \in \mathcal{Y}(T)$, and $u \in \mathcal{S}(T)$,

$$\begin{aligned} \|(\mathcal{M}(\tilde{u}_{0},a,b_{n})u)(t,\cdot) - (\mathcal{M}(\tilde{u}_{0},a,b_{0})u)(t,\cdot)\|_{C^{0}(\bar{\Omega})} &= \|\int_{s}^{t} e^{-A(t-\tau)}u^{2}(\tau,\cdot)[b_{n}(\tau,\cdot) - b_{0}(\tau,\cdot)]d\tau\|_{C^{0}(\bar{\Omega})} \\ &\leq R^{2}\|b_{n} - b_{0}\|_{\mathcal{X}_{T}}T. \end{aligned}$$

Therefore, the mapping $\mathcal{M}(\tilde{u}_0, a, b)$ is also uniformly continuous in $b \in \mathcal{Y}(T)$.

Finally, Let T > 0 satisfy (2.7), (2.8), and (2.10). By Lemmas 2.1 and 2.7, we conclude that $\mathcal{M}(\tilde{u}_0, a, b)$ has a unique fixed point $u \in \mathcal{S}(T)$ fulfilling $(\mathcal{M}(\tilde{u}_0, a, b)u)(t, \cdot) = u(t, \cdot; s, \tilde{u}_0, a, b)$ for $t \in [s, s + T]$, and we also have that $u(t, \cdot; s, \tilde{u}_0, a, b) \in C^0(\bar{\Omega})$ is continuous in $\tilde{u}_0 \in \mathcal{B}(u_0, r)$ and $(a, b) \in \mathcal{Y}(T)$ uniformly in $t \in [s, s + T]$. The lemma is thus proved.

3 Boundedness of positive solutions

In this section, we investigate the boundedness of positive solutions of (1.1), and prove Theorem 1.1. Throughout this section, we assume that (1.5)' holds.

For given $s \in \mathbb{R}$ and u_0 satisfying (1.2), put $(u(t,x), v(t,x)) = (u(t,x;s,u_0), v(t,x;s,u_0))$ for $t \geq s$ and $x \in \overline{\Omega}$. Observe that, for any $1 < q < \infty$,

$$\begin{split} \frac{1}{q} \int_{\Omega} u_t^q(t,x) dx &= -(q-1) \int_{\Omega} u^{q-2}(t,x) |\nabla u(t,x)|^2 dx - \chi(q-1) \int_{\Omega} \frac{u^{q-1}(t,x)}{v(t,x)} \nabla u(t,x) \cdot \nabla v(t,x) dx \\ &+ \int_{\Omega} a(t,x) u^q(t,x) - \int_{\Omega} b(t,x) u^{q+1}(t,x) dx \quad \forall t > s, \end{split}$$

and for any p > 0,

$$\frac{1}{p} \cdot \frac{d}{dt} \int_{\Omega} u^{-p}(t,x) = -(p+1) \int_{\Omega} u^{-p-2}(t,x) |\nabla u(t,x)|^2 + (p+1)\chi \int_{\Omega} \frac{u^{-p-1}(t,x)}{v(t,x)} \nabla u(t,x) \cdot \nabla v(t,x) - \int_{\Omega} a(t,x) u^{-p}(t,x) + \int_{\Omega} b(t,x) u^{-p+1}(t,x) \quad \forall t > s.$$
(3.1)

In the rest of this section, we may omit (t, x) inside the integrals if no confusion occurs.

We first present some lemmas.

Lemma 3.1. For any $s \in \mathbb{R}$ and u_0 satisfying (1.2)',

$$\int_{\Omega} u(t,x)dx \le m^*(\tau,s,u_0) = \max\left\{\int_{\Omega} u(\tau,x)dx, \frac{a_{\sup}}{b_{\inf}}|\Omega|\right\} \quad \forall \ t > \tau \ge s$$

and

$$\limsup_{t-s\to\infty}\int u(t,x)dx \le \frac{a_{\sup}}{b_{\inf}}|\Omega|,$$

where $|\Omega|$ is the Lebesgue measure of Ω .

Proof. By integrating the first equation in (1.1) with respect to x, we get that

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} u &= \int_{\Omega} \Delta u - \chi \int_{\Omega} \nabla \cdot \left(\frac{u}{v} \nabla v\right) + \int_{\Omega} a(x,t)u - b(x,t)u^2 \\ &= \int_{\Omega} a(x,t)u(x,t)dx - \int_{\Omega} b(x,t)u^2(x,t)dx \\ &\leq a_{\sup} \int_{\Omega} u - \frac{b_{\inf}}{|\Omega|} \Big(\int_{\Omega} u\Big)^2. \end{aligned}$$

This together with comparison principle for scalar ODEs implies that

$$\int_{\Omega} u(t, x) dx \le \max \left\{ \int_{\Omega} u(\tau, x) dx, \frac{a_{\sup}}{b_{\inf}} |\Omega| \right\} \quad \forall \ t > \tau \ge s$$

and

$$\limsup_{t-s\to\infty}\int u(t,x)dx \le \frac{a_{\sup}}{b_{\inf}}|\Omega|$$

The lemma is thus proved.

Lemma 3.2. Let $q \ge 3$ and $q - 1 \le k < 2q - 2$. There exist positive constants M(q,k) > 0 and $M^*(q,k) > 0$ such that

$$\int_{\Omega} \frac{|\nabla v|^{2q}}{v^{k+1}} \le M(k,q) \int_{\Omega} \frac{u^q}{v^{k-q+1}} + M^*(k,q) \int_{\Omega} v^{2q-k-1}$$

for any $s \in \mathbb{R}$, $u_0 \in C^0(\overline{\Omega})$ satisfying (1.2)', and $t \in (s, \infty)$.

Proof. It follows from [25, Proposition 1.3].

Lemma 3.3. For any p > 0, $s \in \mathbb{R}$ and $u_0 \in C^0(\overline{\Omega})$ satisfying (1.2)',

$$\int_{\Omega} u \ge |\Omega|^{\frac{p+1}{p}} \left(\int_{\Omega} u^{-p} \right)^{-\frac{1}{p}} \quad \forall t > s.$$

Proof. For any given p > 0, by Hölder's inequality, we have

$$|\Omega| = \int_{\Omega} u^{\frac{p}{p+1}} u^{-\frac{p}{p+1}} \le \left(\int_{\Omega} u\right)^{\frac{p}{p+1}} \left(\int_{\Omega} u^{-p}\right)^{\frac{1}{p+1}} \quad \forall t > s.$$

This implies that

$$\int_{\Omega} u \ge |\Omega|^{\frac{p+1}{p}} \left(\int_{\Omega} u^{-p} \right)^{-\frac{1}{p}} \quad \forall t > s.$$

The lemma is thus proved.

Lemma 3.4. Let p > 0. Then for every $\beta > 0$, we have

$$\begin{split} (p+1)\chi\int_{\Omega}\frac{u^{-p-1}}{v}\nabla u\cdot\nabla v &\leq (p+1)\int_{\Omega}u^{-p-2}|\nabla u|^2 + \frac{(p+1)\beta\mu}{p}\int_{\Omega}u^{-p} \\ &+ \left[\frac{(p+1)(\chi-\beta)^2}{4} - \frac{(p+1)\beta}{p}\right]\int_{\Omega}u^{-p}\frac{|\nabla v|^2}{v^2} \end{split}$$

for all $t \in (s, \infty)$.

Proof. This is Lemma [25, Lemma 3.3].

Lemma 3.5. Let R > 0 be such that

$$R > \begin{cases} \frac{\mu\chi^2}{4}, & \text{if } 0 < \chi \leq 2\\ \mu(\chi - 1), & \text{if } \chi > 2. \end{cases}$$

Then there is $\beta > 0$, $\beta \neq \chi$ such that

$$\frac{(p+1)\beta\mu}{p} - R < 0,$$

where p is given by

$$p = \frac{4\beta}{(\chi - \beta)^2}.$$

Proof. This is [25, Lemma 3.1].

We now prove Theorem 1.1.

Proof of Theorem 1.1. (1) It follows from the arguments of [25, Theorem 1.1(3)]. To be more precise, let $\beta = \chi + 2 - 2\sqrt{\chi + 1}$. Then

$$p := \frac{4\beta}{(\chi - \beta)^2} = \frac{4(\chi + 2 - 2\sqrt{\chi + 1})}{(-2 + 2\sqrt{\chi + 1})^2} = 1 \implies \frac{(\chi - \beta)^2}{4} - \beta = 0,$$

and

$$\frac{(p+1)\beta\mu}{p} = 2\beta\mu = 2(\chi + 2 - 2\sqrt{\chi + 1})\mu = a_{\chi,\mu} < a_{\inf}$$

By (3.1),

$$\frac{d}{dt} \int_{\Omega} u^{-1} \leq -2 \int_{\Omega} u^{-3} |\nabla u|^2 + 2\chi \int_{\Omega} \frac{u^{-2}}{v} \nabla u \cdot \nabla v - a_{\inf} \int_{\Omega} u^{-1} + b_{\sup} |\Omega|$$

for all t > s. By Lemma 3.4, we have that

$$\frac{d}{dt} \int_{\Omega} u^{-1} \leq 2 \left[\frac{(\chi - \beta)^2}{4} - \beta \right] \int_{\Omega} u^{-1} \frac{|\nabla v|^2}{v^2} + \left[2\beta\mu - a_{\inf} \right] \int_{\Omega} u^{-1} + b_{\sup} |\Omega|$$
$$= -(a_{\inf} - a_{\chi,\mu}) \int_{\Omega} u^{-1} + b_{\sup} |\Omega|, \quad \forall t > \tau > s.$$

r	-	ч.
L		н
L		
-		-

It then follows from comparison principle for scalar ODEs that

$$\int_{\Omega} u^{-1}(t,x;s,u_0) dx \le e^{-(a_{\inf} - a_{\chi,\mu})(t-\tau)} \int_{\Omega} u^{-1}(\tau,x;s,u_0) dx + \frac{b_{\sup}|\Omega|}{a_{\inf} - a_{\chi,\mu}}$$

and

$$\int_{\Omega} u^{-1}(t,x;s,u_0) dx \le \max\left\{\int_{\Omega} u^{-1}(\tau,x;s,u_0) dx, \frac{b_{\sup}|\Omega|}{a_{\inf} - a_{\chi,\mu}}\right\}$$
(3.2)

for any $t > \tau > s$. Moreover,

$$\lim_{t-s\to\infty}\int_{\Omega}u^{-1}(t,x;s,u_0)dx = \frac{b_{\sup}|\Omega|}{a_{\inf}-a_{\chi,\mu}}.$$

Theorem 1.1(1) thus follows.

(2) It also follows from the arguments of [25, Theorem 1.1(3)]. To be more precise, by Lemmas 2.3 and 3.3, we have

$$v(t,x) \ge \delta_0 \int_{\Omega} u \ge \delta_0 |\Omega|^2 \left(\int_{\Omega} u^{-1} \right)^{-1} \quad \forall t > s.$$

By (3.2) and the assumption (1.2)'

$$\int_{\Omega} u \ge \frac{(a_{\inf} - a_{\chi,\mu}) \min\{1,\chi\}}{b_{\sup}|\Omega|}, \quad v \ge \frac{\delta_0(a_{\inf} - a_{\chi,\mu}) \min\{1,\chi\}|\Omega|}{b_{\sup}} \quad \text{for all } t \in [s + \tau_0, \infty).$$

By the arguments of [25, Theorem 1.1 (3)], for any $q > \max\{2, N\}$ and any $\varepsilon > 0$, there is $C(\varepsilon, q) > 0$ such that

$$\frac{1}{q} \cdot \frac{d}{dt} \int_{\Omega} u^{q} \leq \frac{b_{\sup} |\Omega| (q-1) \left\{ (C^{*})^{\frac{1}{q+1}} + \varepsilon \right\} \max\{\chi, \chi^{2}\}}{4\delta_{0}(a_{\inf} - a_{\chi,\mu})} \int_{\Omega} u^{q+1} + \frac{b_{\sup} |\Omega| (q-1) C(\varepsilon, p) \max\{\chi, \chi^{2}\}}{4\delta_{0}(a_{\inf} - a_{\chi,\mu})} \left(\int_{\Omega} u \right)^{q+1} + a_{\sup} \int_{\Omega} u^{q} - b_{\inf} \int_{\Omega} u^{q+1} \ \forall t \in (s + \tau_{0}, \infty).$$

By the assumption (1.5)', there are $q > \max\{2, n\}$ and $0 < \varepsilon = \varepsilon(q) \ll 1$ such that

$$\frac{b_{\sup}|\Omega|(q-1)\left\{ (C^*(n,\nu))^{\frac{1}{q+1}} + \varepsilon \right\} \max\{\chi,\chi^2\}}{4\delta_0(a_{\inf} - a_{\chi,\mu})} < b_{\inf}.$$

Note that

$$\int_{\Omega} u^{q+1} \ge \frac{1}{|\Omega|} \left(\int_{\Omega} u^{q} \right)^{\frac{q+1}{p}} \quad \text{for all } t \in (s, \infty).$$

Therefore, there is $b_1 > 0$ such that

$$\frac{1}{q} \cdot \frac{d}{dt} \int_{\Omega} u^{q} \leq \frac{b_{\sup} |\Omega| (q-1) C(\varepsilon, q) \max\{\chi, \chi^{2}\}}{4\delta_{0}(a_{\inf} - a_{\chi, \mu})} \left(\int_{\Omega} u\right)^{q+1} + a_{\sup} \int_{\Omega} u^{q} - b_{1} \left(\int_{\Omega} u^{q}\right)^{\frac{q+1}{p}} \quad \text{for all } t \in (s + \tau_{0}, \infty).$$

This together with Lemma 3.1 implies that there are $M_2(1,q)$ and $\tilde{M}_2(1,q,\tau,s,u_0)$ such that

$$\begin{cases} \frac{d}{dt} \int_{\Omega} u^{q} \leq -\int_{\Omega} u^{q} + M_{2}(1,q) & \forall t \geq s + \tau_{0} \\ \frac{d}{dt} \int_{\Omega} u^{q} \leq -\int_{\Omega} u^{q} + M_{2}(1,q) + \tilde{M}_{2}(1,q,\tau,s,u_{0}) & \forall t \geq \tau > s. \end{cases}$$
(3.3)

Theorem 1.1(2) then follows.

4 Globally attracting invariant rectangle and pointwise persistence

In this section, we investigate the existence of globally attracting invariant rectangle for (1.1) and pointwise persistence, and prove Theorems 1.2 and 1.3.

We first prove a lemma.

Lemma 4.1. Assume that (1.5)' holds. Let p > 0 be as in Theorem 1.1(1). There are $M_1^* > 0$ and $\tilde{M}_1^* > 0$ such that for every $u_0 \in C^0(\bar{\Omega})$ satisfying (1.2)' and $\int_{\Omega} u_0(x) dx \leq \frac{a_{\sup}}{b_{\inf}} |\Omega|$, $s \in \mathbb{R}$, and $\tau > 0$,

$$\int_{\Omega} u^{-p}(t,x;s,u_0)dx \le \max\left\{\int_{\Omega} u^{-p}(s+\tau,x;s,u_0)dx, M_1^*\right\} \quad \text{for all } s+\tau < t < \infty$$
(4.1)

and

$$\limsup_{t-s\to\infty} \int_{\Omega} u^{-p}(t,x;s,u_0) dx \le M_1^*, \tag{4.2}$$

$$\liminf_{t-s\to\infty} \int_{\Omega} u(t,x;s,u_0) dx \ge \tilde{M}_1^*, \quad \liminf_{t-s\to\infty} \inf_{x\in\Omega} v(t,x;s,u_0) \ge \delta_0 \tilde{M}_1^*.$$
(4.3)

In addition, if $\inf_{x\in\Omega} u_0(x) > 0$ and $\int_{\Omega} u_0^{-p}(x) dx \leq M_1^*$, then

$$\int_{\Omega} u^{-p}(t, x; s, u_0) dx \le M_1^* \quad \forall \ s < t < \infty$$

$$\tag{4.4}$$

and

$$\int_{\Omega} u(t,x;s,u_0) \ge \tilde{M}_1^*, \quad \inf_{x \in \Omega} v(t,x;s,u_0) \ge \delta_0 \tilde{M}_1^* \quad \forall t \in [s,\infty).$$

$$(4.5)$$

Proof. First, let $M_1^* = M_1(p)$. Then, by Lemma 3.1,

$$\int_{\Omega} u(\tau, x; s, u_0) dx \le \frac{a_{\sup}}{b_{\inf}} |\Omega| \quad \forall \ \tau \ge s.$$

Hence

$$m^*(\tau, s, u_0) = \frac{a_{\text{sup}}}{b_{\text{inf}}} \quad \text{and} \quad \tilde{M}_1(p, \tau, s, u_0) = 0 \quad \forall \ \tau \ge s.$$

$$(4.6)$$

This together with (1.10) and (1.11) implies (4.1) and (4.2) hold.

Next, by Hölder inequality, we have

$$\begin{aligned} |\Omega| &= \int_{\Omega} u^{\frac{p}{p+1}}(t,x;s,u_0) u^{-\frac{p}{p+1}}(t,x;s,u_0) dx \\ &\leq \left(\int_{\Omega} u(t,x;s,u_0) dx\right)^{\frac{p}{p+1}} \left(\int_{\Omega} u^{-p}(t,x;s,u_0) dx\right)^{\frac{1}{p+1}} \quad \forall t > s. \end{aligned}$$
(4.7)

This together with (4.2) and Lemma 2.3 implies that (4.3) holds with $\tilde{M}_1^* = \frac{|\Omega|^{\frac{p+1}{p}}}{(M_1^*)^{\frac{1}{p}}}$.

Now, assume $\inf_{x\in\Omega} u_0(x) > 0$ and $\int_{\Omega} u_0^{-p}(x) dx \le M_1^*$. By (2.4),

$$\lim_{\tau \to 0} \int_{\Omega} u^{-p}(s+\tau, x; s, u_0) dx = \int_{\Omega} u_0^{-p}(x) dx \le M_1^*.$$

This together with (4.1) implies (4.4). (4.5) then follows from (4.4), (4.7), and Lemma 2.3. The lemma is thus proved.

Next, we prove Theorem 1.2.

Proof of Theorem 1.2. Let p and $M_1(p)$ be as in Theorem 1.1(1), and q and $M_2(p,q)$ be as in Theorem 1.1(2). Let $M_0^* = \frac{a_{\sup}}{b_{\inf}} |\Omega|$, $M_1^* = M_1(p)$, and $M_2^* = M_2(p,q)$. In the following, we prove that (1) and (2) in Theorem 1.2 hold with these p, q, M_1^*, M_2^* and some $M_3^*(\tau, \theta), M_4^*(\theta)$.

(1) We first prove that the set \mathcal{E} is invariant.

First of all, for any $u_0 \in \mathcal{E}$,

$$|\Omega|^{2} = \left(\int_{\Omega} u_{0}^{-p/2}(x)u_{0}^{p/2}(x)dx\right)^{2} \le \int_{\Omega} u_{0}^{-p}(x)dx \cdot \int_{\Omega} u_{0}^{p}(x)dx \le M_{1}^{*}\int_{\Omega} u_{0}^{p}(x)dx.$$

This implies that $\int_{\Omega} u_0^p(x) dx > 0$ and then $\int_{\Omega} u_0(x) dx > 0$. Recall that p = 1 and

$$M_1^* = M_1(p) = \frac{b_{\sup}|\Omega|}{a_{\inf} - a_{\chi,\mu}} \le \frac{b_{\sup}|\Omega| \max\{1, \frac{1}{\chi}\}}{a_{\inf} - a_{\chi,\mu}},$$

Hence u_0 satisfies (1.2)' with $\tau_0 = 0$.

For any $s \in \mathbb{R}$ and $u_0 \in \mathcal{E}$, by Lemma 3.1,

$$\int_{\Omega} u(t,x;s,u_0)dx \le \max\left\{\int_{\Omega} u_0(x)dx, M_0^*\right\} = M_0^* \quad \forall t \ge s.$$

$$(4.8)$$

By Lemma 4.1, there holds

$$\int_{\Omega} u^{-p}(t,x;s,u_0)dx \le M_1^* \quad \forall t > s.$$

$$\tag{4.9}$$

By (1.10), (3.3) with $\tau_0 = 0$, (4.6), and (4.9),

$$\int_{\Omega} u^q(t,x;s,u_0) dx \le \max\left\{\int_{\Omega} u^q(\tau,x;s,u_0) dx, M_2^*\right\} \quad \forall t > \tau > s.$$

Letting $\tau \to s^+$, we have

$$\int_{\Omega} u^q(t,x;s,u_0) dx \le M_2^* \quad \forall t \ge s.$$
(4.10)

It then follows from (4.8), (4.9) and (4.10) that the set \mathcal{E} is invariant.

Next, we prove (1.14) holds. To this end, for any given $0 < \theta < 1 - \frac{2N}{q}$, choose $\beta \in (0, \frac{1}{2})$ such that

$$2\beta - \frac{2N}{q} > \theta.$$

By Lemma 2.5, there is K_1 such that for any $u_0 \in \mathcal{E}$ and $s \in \mathbb{R}$,

$$\|u(t,\cdot;s,u_0)\|_{C^{\theta}(\bar{\Omega})} \le K_1 \|u(t,\cdot;s,u_0)\|_{X^{\beta}_{q/2}} \quad \forall t > s.$$
(4.11)

Note that for any $u_0 \in \mathcal{E}$ and $s \in \mathbb{R}$,

$$\begin{aligned} \|u(t,\cdot;s,u_{0})\|_{X_{q/2}^{\beta}} &\leq \|A^{\beta}e^{-A(t-s)}u_{0}\|_{L^{q/2}(\Omega)} \\ &+ \chi \int_{s}^{t} \left\|A^{\beta}e^{-A(t-\lambda)}\nabla \cdot \left(\frac{u(\lambda,\cdot;s,u_{0})}{v(\lambda,\cdot;s,u_{0})}\nabla v(\lambda,\cdot;s,u_{0})\right)\right\|_{L^{q/2}(\Omega)} d\lambda \\ &+ \int_{s}^{t} \|A^{\beta}e^{-A(t-\lambda)}\left((1+a(\lambda,\cdot))u(\lambda,\cdot;s,u_{0})-b(\lambda)u^{2}(\lambda,\cdot;s,u_{0})\right)\|_{L^{q/2}(\Omega)} d\lambda, \end{aligned}$$

$$(4.12)$$

where $A = A_{q/2}$. By Lemma 2.5 again, there is K_2 such that

$$\|A^{\beta}e^{-A(t-s)}u_0\|_{L^{q/2}(\Omega)} \le K_2(t-s)^{-\beta(t-s)}e^{-\gamma(t-s)}\|u_0\|_{L^{q/2}} \quad \forall t > s.$$
(4.13)

By Lemma 2.6, there is K_3 such that

$$\begin{split} \chi \int_{s}^{t} \left\| A^{\beta} e^{-A(t-\lambda)} \nabla \cdot \left(\frac{u(\lambda, \cdot; s, u_{0})}{v(\lambda, \cdot; s, u_{0})} \nabla v(\lambda, \cdot; s, u_{0}) \right) \right\|_{L^{q/2}(\Omega)} d\lambda \\ &\leq K_{3} \chi \int_{s}^{t} (t-\lambda)^{-\beta-\frac{1}{2}-\epsilon} e^{-\gamma(t-\lambda)} \left\| \frac{u(\lambda, \cdot; s, u_{0})}{v(\lambda, \cdot; s, u_{0})} \cdot \nabla v(\lambda, \cdot; s, u_{0}) \right\|_{L^{q/2}(\Omega)} d\lambda \\ &\leq \frac{K_{3} \chi}{\tilde{M}_{1}^{*}} \int_{s}^{t} (t-\lambda)^{-\beta-\frac{1}{2}-\epsilon} e^{-\gamma(t-\lambda)} \left\| u(\lambda, \cdot; s, u_{0}) \cdot \nabla v(\lambda, \cdot; s, u_{0}) \right\|_{L^{q/2}(\Omega)} d\lambda \quad \text{(by Lemma 4.1)} \\ &\leq \frac{K_{3} \chi}{\tilde{M}_{1}^{*}} \int_{s}^{t} (t-\lambda)^{-\beta-\frac{1}{2}-\epsilon} e^{-\gamma(t-\lambda)} \left\| u(\lambda, \cdot; s, u_{0}) \right\|_{L^{q}(\Omega)} \left\| \nabla v(\lambda, \cdot; s, u_{0}) \right\|_{L^{q}(\Omega)} d\lambda \quad \text{(by Hölder inequality)} \\ &\leq \frac{CK_{3} \chi (M_{2}^{*})^{2/q}}{\tilde{M}_{1}^{*}} \int_{s}^{t} (t-\lambda)^{-\beta-\frac{1}{2}-\epsilon} e^{-\gamma(t-\lambda)} d\lambda \quad \text{(by Lemma 2.4 and Theorem 1.1)}. \end{aligned}$$

By Theorem 1.1, there is $K_4 > 0$ such that

$$\int_{s}^{t} \|A^{\beta}e^{-A(t-\lambda)} \left((1+a(\lambda,\cdot))u(\lambda,\cdot;s,u_{0}) - b(\lambda)u^{2}(\lambda,\cdot;s,u_{0}) \right)\|_{L^{q/2}(\Omega)} d\lambda$$

$$\leq \int_{s}^{t} (t-\lambda)^{-\beta}e^{-\gamma(t-\lambda)} \left(|\Omega|^{2/q} + a_{\max} \|u(\lambda,\cdot;su_{0})\|_{L^{q/2}} + b_{\max} \|u^{2}(\lambda,\cdot;s,u_{0})\|_{L^{q/2}} \right) d\lambda$$

$$\leq K_{4} \int_{s}^{t} (t-\lambda)^{-\beta}e^{-\gamma(t-\lambda)} d\lambda.$$
(4.15)

By (4.11)-(4.15), for any $\tau > 0$ and $0 < \theta < 1 - \frac{2N}{q}$, there is $M_3^*(\theta, \tau) > 0$ such that (1.14) holds. (1) is thus proved.

(2) First of all, it is not difficult to see that (1.15) follows directly from Lemma 3.1 and Theorem 1.1. It then suffices to prove (1.16).

For any u_0 satisfying (1.2)', by (1.15), there is $T_0(u_0) > 0$ such that for any $s \in \mathbb{R}$,

$$\int_{\Omega} u^{-p}(t,x;s,u_0) dx \le 2M_1^* \quad \text{and} \quad \int_{\Omega} u^q(t,x;s,u_0) dx \le 2M_2^* \quad \forall t \ge s + T_0(u_0).$$

Observe that

$$(u(t,x;s,u_0),v(t,x;s,u_0)) = (u(t,x;\tilde{s},\tilde{u}_0),v(t,x;\tilde{s},\tilde{u}_0)) \quad \forall t \ge \tilde{s},$$
(4.16)

where $\tilde{s} = s + T_0(u_0)$ and $\tilde{u}_0(x) = u(s + T_0(u_0), x; s, u_0)$. Hence

$$\int_{\Omega} u^{-p}(t,x;\tilde{s},\tilde{u}_0)dx \le 2M_1^* \quad \text{and} \quad \int_{\Omega} u^q(t,x;\tilde{s},\tilde{u}_0)dx \le 2M_2^* \quad \forall t \ge \tilde{s}.$$

By the arguments in the proof of (1.14), there is $M_4^*(\theta)$ such that

$$\|u(t,\cdot;\tilde{s},\tilde{u}_0)\|_{C^{\theta}(\bar{\Omega})} \le M_4^*(\theta) \quad \forall t \ge \tilde{s}+1.$$

This together with (4.16) implies that (1.16) holds with $T(u_0) = T_0(u_0) + 1$.

We now prove Theorem 1.3. In the rest of this section, to indicate the dependence of $(u(t, x; s, u_0), v(t, x; s, u_0))$ on a(t, x), b(t, x), we put

$$(u(t,x;s,u_0,a,b),v(t,x;s,u_0,a,b)) = (u(t,x;s,u_0),v(t,x;s,u_0)).$$

Proof of Theorem 1.3. We prove this theorem by contradiction. Assume that there is no $m^* > 0$ such that (1.18) holds. Then for any $n \in \mathbb{N}$, there is u_n satisfying (1.2)' such that

$$\liminf_{t-s\to\infty} \inf_{x\in\Omega} u(t,x;s,u_n,a,b) \le m_n := \frac{1}{n}.$$
(4.17)

Let $p, q, \theta, M_1^*, M_2^*$, and $M_4^*(\theta)$ be as in Theorem 1.2. By Theorem 1.2 and Lemma 4.1, there is $T_n > 0$ such that

$$\|u(t, \cdot; s, u_n, a, b)\|_{C^{\theta}(\bar{\Omega})} \le M_4^*(\theta) \quad \forall t - s \ge T_n$$

$$(4.18)$$

and

$$\int_{\Omega} u(t,x;s,u_n,a,b)dx \ge \frac{\tilde{M}_1^*}{2} \quad \forall t-s \ge T_n.$$
(4.19)

By (4.17), there are $t_n, s_n \in \mathbb{R}$ with $t_n - s_n \ge T_n + 1$ and $x_n \in \overline{\Omega}$ such that

$$u(t_n, x_n; s_n, u_n, a, b) \le \frac{2}{n}.$$
 (4.20)

Let

$$a_n(t,x) = a(t+t_n-1,x), \quad b_n(t,x) = b(t+t_n-1,x).$$

Observe that

$$u(t_n, x; s, u_n, a, b) = u(t_n, x; t_n - 1, u(t_n - 1, \cdot; s_n, u_n, a, b), a, b)$$

= $u(1, x; 0, u(t_n - 1, \cdot; s_n, u_n, a, b), a_n, b_n).$ (4.21)

By (4.18) and **(H)**, without loss of generality, we may assume that there are $u_0^*, u_1^* \in C(\bar{\Omega})$ and $a^*(t, x), b^*(t, x)$ satisfying **(H)** such that

$$\lim_{n \to \infty} u(t_n - 1, x; s_n, u_n, a, b) = u_0^*(x), \quad \lim_{n \to \infty} u(t_n, x; s_n, u_n, a, b) = u_1^*(x)$$

uniformly in $x \in \overline{\Omega}$, and

$$\lim_{n \to \infty} a_n(t, x) = a^*(t, x), \quad \lim_{n \to \infty} b_n(t, x) = b^*(t, x)$$

uniformly in $x \in \overline{\Omega}$ and locally uniformly in $t \in \mathbb{R}$. It then follows from Lemma 2.8 and (4.21) that

$$u_1^*(x) = u(1, x; 0, u_0^*, a^*, b^*).$$
(4.22)

By (4.18), (4.19), and the Dominated Convergence Theorem,

$$\int_{\Omega} u_0^*(x) = \lim_{n \to \infty} \int_{\Omega} u(t_n - 1, x; s_n, u_n, a, b) dx \ge \frac{\tilde{M}_1^*}{2}.$$

Then by (4.22) and the strong maximum principle for parabolic equations,

$$\inf_{x\in\bar{\Omega}}u_1^*(x)>0.$$

This together with (4.20) implies that

$$\frac{2}{n} \ge \inf_{x \in \Omega} u(t_n, x; s_n, u_n, a, b) \ge \frac{1}{2} \inf_{x \in \Omega} u_1^*(x) \quad \forall n \gg 1,$$

which is a contradiction. Therefore, there is $m^* > 0$ such that for any u_0 satisfying (1.2),

$$\liminf_{t-s\to\infty}\inf_{x\in\Omega}u(t,x;s,u_0,a,b)\geq m^*.$$

The theorem is thus proved.

5 Existence of positive entire solutions

In this section, we study the existence of positive entire solutions and prove Theorem 1.4.

We first prove a lemma.

Lemma 5.1. Let p > 0, q > 2N, and $M_1^*, M_2^* > 0$ be as in Theorem 1.2. The set

$$\mathcal{E} = \left\{ u \in C^{0}(\bar{\Omega}) \, | \, u \ge 0, \, \int_{\Omega} u(x) \le M_{0}^{*}, \, \int_{\Omega} u^{-p}(x) dx \le M_{1}^{*}, \, \int_{\Omega} u^{q}(x) \le M_{2}^{*} \right\}$$

is a convex, closed subset of $C^0(\overline{\Omega})$.

-		-
L		
L		
L		

Proof. First, we prove the set \mathcal{E} is convex. Note that $g(u) = u^r$ is a convex function on $(0, \infty)$, where r = 1, -p or q. Hence, for any $\lambda \in [0, 1]$ and $u_*, u^* > 0$, we have

$$g(\lambda u_* + (1 - \lambda)u^*) \le \lambda g(u_*) + (1 - \lambda)g(u^*).$$
 (5.1)

For any $u_1, u_2 \in \mathcal{E}$, we have $\int_{\Omega} u_1^{-p}(x) dx \leq M_1^*$, $\int_{\Omega} u_2^{-p}(x) dx \leq M_1^*$, and $u_1(x) > 0, u_2(x) > 0$ for a.e. $x \in \Omega$. By (5.1), we have that

$$g(\lambda u_1(x) + (1-\lambda)u_2(x)) \le \lambda g(u_1(x)) + (1-\lambda)g(u_2(x)) \quad \text{for} \quad a.e. \ x \in \Omega.$$

This implies that

$$\int_{\Omega} g(\lambda u_1(x) + (1-\lambda)u_2(x))dx \le \lambda \int_{\Omega} g(u_1(x))dx + (1-\lambda) \int_{\Omega} g(u_2(x))dx.$$

This implies that for any $\lambda \in [0, 1]$ and $u_1, u_2 \in \mathcal{E}$, $\lambda u_1 + (1 - \lambda)u_2 \in \mathcal{E}$. Thus, the set \mathcal{E} is convex.

Next, we prove the set \mathcal{E} is closed. Suppose that $u_n \in \mathcal{E}$, $u \in C^0(\overline{\Omega})$, and $u_n \to u$ in $C^0(\overline{\Omega})$. Then $u(x) \ge 0$ for any $x \in \overline{\Omega}$, and for any $\varepsilon > 0$, there is N_{ε} such that for any $n \ge N_{\varepsilon}$, there holds

$$|u_n(x) - u(x)| < \varepsilon \quad \forall \, x \in \bar{\Omega}.$$

This implies that

$$0 \le u_n(x) = u_n(x) - u(x) + u(x) \le \varepsilon + u(x) \quad \forall n \ge N_{\varepsilon}, \ x \in \overline{\Omega}.$$
(5.2)

For any $\varepsilon > 0$, let

$$\Omega_{\varepsilon} = \{ x \in \Omega \, | \, u(x) \le \varepsilon \}.$$

Let

$$\Omega_0 = \{ x \in \Omega \, | \, u(x) = 0 \}.$$

By (5.2),

$$0 \le u_n(x) \le 2\varepsilon \quad \forall x \in \Omega_{\varepsilon}.$$

Since $\int_{\Omega} u_n^{-p}(x) dx \leq M_1^*$, we have that

$$|\Omega_{\epsilon}| = \int_{\Omega} u_n^{-p} \cdot u_n^p \le \int_{\Omega} u_n^{-p} \cdot (2\epsilon)^p \le (2\epsilon)^p \int_{\Omega} u_n^{-p} \le (2\epsilon)^p M_1^*$$

This implies that

$$|\Omega_0| = 0.$$

Let g(u) be as in the above. Then by Fatou's lemma, we have

$$\int_{\Omega} g(u(x))dx = \int_{\Omega \setminus \Omega_0} g(u(x))dx = \int_{\Omega \setminus \Omega_0} \lim_{n \to \infty} g(u_n(x))dx \le \lim_{n \to \infty} \inf \int_{\Omega \setminus \Omega_0} g(u_n(x))dx.$$

This implies that

$$\int_{\Omega} u(x)dx \le M_0^*, \ \int_{\Omega} u^{-p}(x)dx \le M_1^*, \ \int_{\Omega} u^q(x)dx \le M_2^*$$

Hence $u \in \mathcal{E}$ and the set \mathcal{E} is closed.

We now prove Theorem 1.4.

Proof of Theorem 1.4. (1) First of all, fix $u_0 \in \mathcal{E}$, where \mathcal{E} as in (1.13). By Theorem 1.2, $u(t, \cdot; s, u_0) \in \mathcal{E}$ for any t > s.

Next, for any $n \in \mathbb{N}$, define $u_n(x) = u(0, x; -n, u_0)$. Fix $\theta \in (0, \frac{2N}{q})$. By Theorem 1.2 again, there is $\tilde{M}_3^* > 0$ such that

$$\|u(t,\cdot;-n,u_n)\|_{C^{\theta}} = \sup_{x\in\Omega} |u(t,x;-n,u_0)| + \sup \frac{|u(t,x;-n,u_0) - u(t,y;-n,u_0)|}{|x-y|^{\theta}} \le \tilde{M}_3^*$$
(5.3)

for all $t \ge -n+1$ and $n \ge 1$. This implies that the sequence $\{u_n(\cdot)\}$ is uniformly bounded and equicontinuous on $\overline{\Omega}$. By the Arzelá-Ascoli Theorem and Lemma 5.1, there are $n_k, u_0^* \in \mathcal{E}$ such that $u_{n_k} \to u_0^*$ in $C^0(\overline{\Omega})$ as $n_k \to \infty$.

Note that

$$u(\cdot, t; -n_k, u_0) = u(\cdot, t; 0, u(\cdot, 0; -n_k, u_0)) = u(\cdot, t; 0, u_{n_k}) \quad \forall t > 0.$$

By Lemma 2.8, for any t > 0,

$$\lim_{n_k \to \infty} u(\cdot, t; -n_k, u_0) = \lim_{n_k \to \infty} u(\cdot, t; 0, u_{n_k}) = u(\cdot, t; 0, u_0^*) \quad \text{in} \quad C^0(\bar{\Omega})$$

Since $v(\cdot,t;0,u_0^*) = A^{-1}u(\cdot,t;0,u_0^*)$, we also have that for any t > 0,

$$\lim_{n_k \to \infty} v(\cdot, t; -n_k, u_0) = \lim_{n_k \to \infty} A^{-1} u(\cdot, t; -n_k, u_0) = v(\cdot, t; 0, u_0^*) \quad \text{in} \quad C^0(\bar{\Omega}).$$

We now prove that $u(\cdot, t; 0, u_0^*)$ has a backward extension on $(-\infty, 0)$. By (5.3) and the Arzelá-Ascoli Theorem, without loss of generality, we may assume that for any $m \in \mathbb{N}$, there is $u_m^*(\cdot) \in \mathcal{E}$ such that $u(\cdot, -m; -n_k, u_0) \to u_m^*(\cdot)$ in $C^0(\overline{\Omega})$. By Lemma 2.8 again,

$$\lim_{n_k \to \infty} u(\cdot, t; -n_k, u_0) = \lim_{n_k \to \infty} u(\cdot, t; -m, u(\cdot, -m; -n_k, u_0)) = u(\cdot, t; -m, u_m^*)$$

and for t > -m. Note that $u_0^* = u(\cdot, 0; -m, u_m^*)$. This implies that $u^*(x, t; 0, u_0^*)$ has a backward extension up to t = -m. Letting $m \to \infty$ yields that $u^*(x, t)$ has a backward extension on $(-\infty, 0)$.

Finally, let us denote $u^*(t, x) = u^*(t, x; 0, u_0^*)$, hence $v^*(t, x) = A^{-1}u^*(t, x)$. Then $(u^*(t, x), v^*(t, x))$ is an entire solution of (1.1) and $u^*(t, \cdot) \in \mathcal{E}$ for any $t \in \mathbb{R}$. (1) is thus proved.

(2) Assume that there exists T > 0 such that a(t + T, x) = a(t, x) and b(t + T, x) = b(t, x). First, define the map $\mathcal{T}(T) : \mathcal{E} \to C^0(\overline{\Omega})$ as follows:

$$\mathcal{T}(T)u_0 = u(T, \cdot; 0, u_0) \quad \forall \, u_0 \in \mathcal{E}.$$

By Theorem 1.2, $\mathcal{T}(T)$ maps \mathcal{E} into \mathcal{E} . By Lemma 2.7 and Theorem 1.2, $\mathcal{T}(T) : \mathcal{E} \to \mathcal{E}$ is continuous and relatively compact in $C^0(\bar{\Omega})$. By Lemma 5.1, \mathcal{E} is a convex closed subset of $C^0(\bar{\Omega})$. Then, by Schauder fixed point theorem (see Lemma 2.2), one can find $u^T \in \mathcal{E}$ such that $\mathcal{T}(T)u^T = u^T$, that is, $u(\cdot, T; s, u^T) = u^T(\cdot)$. Thus,

$$u(t+T, \cdot; 0, u^{T}) = u(t, \cdot; T, u(T, \cdot; 0, u^{T})) = u(t, \cdot; 0, u^{T}).$$

Hence, $u(t, \cdot; 0, u^T)$ is periodic with period T. Moreover, the second equation of (1.3) and the uniqueness of solutions of

$$\begin{cases} -\Delta v + v = u(t, \cdot; 0, u^T), & x \in \Omega \\ \frac{\partial v}{\partial n} = 0, & x \in \partial \Omega, \end{cases}$$

it is obtained that $v(t, \cdot; 0, u^T) = (I - \Delta)^{-1}u(t, \cdot; 0, u^T)$ is periodic with period T, which concludes that $(u(t, \cdot; 0, u^T), v(t, \cdot; 0, u^T))$ is a positive periodic solution of (1.3) with $u(t, \cdot; 0, u^T) \in \mathcal{E}$ for any $t \in \mathbb{R}$. (2) is thus proved.

(3) Assume that a(t,x) = a(x) and b(t,x) = b(x). Observe that every T > 0 is a period for a(t,x) and b(t,x). By (2), for fixed T > 0, there exists $u^T \in \mathcal{E}$ such that $(u(t,\cdot;0,u^T), (I - \Delta)^{-1}u(t,\cdot;0,u^T))$ is a positive periodic solution of (1.1) with period T.

Let $T_n = \frac{1}{n}$ and $u_n(x) = u^{T_n}(x)$. Then $u_n(x) = u(kT_n, x; 0, u_n)$ for all $k \in \mathbb{N}$. By Theorem 1.2, $\{u_n(\cdot)\}$ is a uniformly bounded and equi-continuous sequence in $C^0(\bar{\Omega})$. Then by the Arzelá-Ascoli theorem, there exist a subsequence $\{u_{n_k}\}$ and $u^* \in \mathcal{E}$ such that $u_{n_k} \to u^*$ in $C^0(\bar{\Omega})$ as $n_k \to \infty$.

We claim that

$$u(\cdot, t; 0, u^*) = u^*(\cdot) \text{ for all } t \ge 0.$$
 (5.4)

For any t > 0 and $\epsilon > 0$, let $\tau_{n_k} \in [0, T_{n_k})$ be such that

$$\frac{t - \tau_{n_k}}{T_{n_k}} \in \mathbb{Z}^+$$

Then

$$\tau_{n_k} \to 0$$
 and $k \to \infty$.

By Lemma 2.7,

$$|u(\tau_{n_k}, \cdot; 0, u^*) - u^*(\cdot)||_{\infty} < \epsilon \quad \forall k \gg 1.$$

By Lemma 2.8,

$$\|u(\tau,\cdot;0,u^*) - u(\tau,\cdot;0,u_{n_k})\|_{\infty} < \epsilon \quad \forall \tau \in [0,t] \quad \text{and} \quad \forall k \gg 1.$$

It then follows that

$$\begin{aligned} |u(t,x;0,u^*) - u^*(x)| &\leq |u(t,x;0,u^*) - u(t,x;0,u_{n_k})| + |u(t,x;0,u_{n_k}) - u^*(x)| \quad (\text{choose } k \gg 1) \\ &= |u(t,x;0,u^*) - u(t,x;0,u_{n_k})| + |u(\tau_{n_k},x;0,u_{n_k}) - u^*(x)| \\ &\leq |u(t,x;0,u^*) - u(t,x;0,u_{n_k})| + |u(\tau_{n_k},x;0,u_{n_k}) - u(\tau_{n_k},x;0,u^*)| \\ &+ |u(\tau_{n_k},x;0,u^*) - u^*(x)| \\ &\leq 3\epsilon \quad \forall x \in \bar{\Omega}. \end{aligned}$$

Letting $\epsilon \to 0$ entails (5.4). This shows that the solution pair $(u(\cdot,t;0,u^*),v(\cdot,t;0,u^*))$ with $v(\cdot,t;0,u^*) = (I - \Delta)^{-1}u(\cdot,t;0,u^*)$ is a steady state solution of (1.1) with $u^*(\cdot) \in \mathcal{E}$. (3) is thus proved.

References

- M. Aida, K. Osaka, T. Tsujikawa, A. Yagi, and M. Mimura, Chemotaxis and growth system with sigular sensitivity function, *Nonliear Anal. Real World Appl.* 6 (2005), 323-336.
- H. Amann, Existence and regularity for semilinear parabolic evolution equations, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 11 (1984), no. 4, 593-676.
- [3] P. Biler, Global solutions to some parabolic-elliptic systems of chemotaxis, Advanced Mathematics and Applications, 9 (1999), 347-359.
- [4] T. Black, Global generalized solutions to a parabolic-elliptic Keller-Segel system with singular sensitivity, *Discrete Contin. Dyn. Syst. Ser. S* 13 (2020) 119-137.
- [5] T. Black, M. Fuest, and J. Lankeit, Relaxed parameter conditions for chemotactic collapse in logistic-type parabolic-elliptic Keller-Segel systems, Z. Angew. Math. Phys. 72 (2021), no. 3, Paper No. 96, 23 pp.
- [6] J. Cao, W. Wang, and H. Yu, Asymptotic behavior of solutions to two-dimensional chemotaxis system with logistic source and singular sensitivity, J. Math. Anal. Appl. 436 (2016), no. 1, 382-392.
- [7] M. Fuest, Finite-time blow-up in a two-dimensional Keller-Segel system with an environmental dependent logistic source, Nonlinear Anal. Real World Appl., 52 (2020), 103022, 14 pp.
- [8] M. Fuest, Approaching optimality in blow-up results for Keller-Segel systems with logistictype dampening, NoDEA Nonlinear Differential Equations Appl. 28 (2021), no. 2, Paper No. 16, 17 pp.
- [9] K. Fujie, Boundedness in a fully parabolic chemotaxis system with singular sensitivity, J. Math. Anal. Appl., 424 (2015), 675-684.
- [10] K. Fujie and T. Senba, Global existence and boundedness of radial solutions to a two dimensional fully parabolic chemotaxis system with general sensitivity, *Nonlinearity* 29 (2016), 2417-2450.
- [11] K. Fujie and T. Senba, Global existence and boundedness in a parabolic-elliptic Keller-Segel system with general sensitivity, *Discrete Contin. Dyn. Syst. Ser. B*, **21(1)** (2016), 81-102.
- [12] K. Fujie, M. Winkler, and T. Yokota, Blow-up prevention by logistic sources in a parabolicelliptic Keller-Segel system with singular sensitivity, *Nonlinear Anal.* **109** (2014), 56-71.
- [13] K. Fujie, M. Winkler, and T. Yokota. Boundedness of solutions to parabolic-elliptic Keller-Segel systems with signal dependent sensitivity, *Math. Methods Appl. Sci.*, **38(6)** (2015), 1212-1224.

- [14] D. Gilbarg, N. Trudinger, Elliptic Partial Differential Equations of Second Order. Springer, Berlin, (1998) ISBN 3-540-41160-7.
- [15] J. K. Hale, Ordinary Differential Equations, Wiley Interscience, New York (1969). Second Edition, Robert E. Krieger Publ. Co., Malabar, Florida.
- [16] D. Henry, Geometric Theory of Semilinear Parabolic Equations, Springer, Berlin, Heidelberg, New York, 1977.
- [17] M.A. Herrero, E. Medina, and J.J.L. Velzquez, Singularity patterns in a chemotaxis model, Math. Ann., 306 (1996), 583-623.
- [18] M.A. Herrero and J.J.L. Velzquez, Finite-time aggregation into a single point in a reactiondiffusion system, *Nonlinearity*, **10** (1997), 1739-1754.
- [19] D. Horstmann and M. Winkler, Boundedness vs.blow-up in a chemotaxis system, J. Differential Equations, 215 (2005), 52-107.
- [20] T.B. Issa and W. Shen, Persistence, coexistence and extinction in two species chemotaxis models on bounded heterogeneous environments, J. Dynam. Differential Equations 31 (2019), no. 4, 1839-1871.
- [21] T.B. Issa and W. Shen, Pointwise persistence in full chemotaxis models with logistic source on bounded heterogeneous environments, J. Math. Anal. Appl. 490 (2020), no. 1, 124204, 30 pp.
- [22] E.F. Keller and L.A. Segel, Initiation of slime mold aggregation viewed as an instability, J. Theoret. Biol. 26 (1970) 399-415.
- [23] E.F. Keller and L.A. Segel, Model for chemotaxis, J. Theoret. Biol., **30** (1971), 225-234.
- [24] E.F. Keller and L.A. Segel, Traveling bans of chemotactic bacteria: a theoretical analysis. J. Theor. Biol. 30 (1971), 377–380.
- [25] H. I. Kurt and W. Shen, Finite-time blow-up prevention by logistic source in chemotaxis models with singular sensitivity in any dimensional setting, https://arxiv.org/abs/2008.01887.
- [26] J. Lankeit, A new approach toward boundedness in a two-dimensional parabolic chemotaxis system with singular sensitivity, *Math. Methods Appl. Sci.*, **39** (2016), no. 3, 394-404.
- [27] J. Lankeit and M. Winkler, A generalized solution concept for the Keller-Segel system with logarithmic sensitivity: global solvability for large nonradial data, *Nonlinear Differential Equations Appl.* 24 (2017), no. 4, Art. 49, 33 pp.
- [28] T. Nagai, Blowup of nonradial solutions to parabolic-elliptic systems modeling chemotaxis in two-dimensional domains, J. Inequal. Appl., 6 (2001), 37-55.

- [29] T. Nagai and T. Senba, Global existence and blow-up of radial solutions to a parabolic-elliptic system of chemotaxis, Adv. Math. Sci. Appl., 8 (1998), 145-156.
- [30] T. Nagai and T. Senba, Global existence and blow-up of radial solutions to a parabolic-elliptic system of chemotaxis, *Advanced Mathematics and Applications*, 8 (1998), 145-156.
- [31] Y. Tanaka and T. Yokota, Blow-up in a parabolic–elliptic Keller–Segel system with densitydependent sublinear sensitivity and logistic source, *Math Meth Appl Sci.*, 43 (2020), 7372– 7396.
- [32] Y. Tao and M. Winkler, Persistence of mass in a chemotaxis system with logistic source, J. Differential Equations, 259 (2015), no. 11, 6142-6161.
- [33] J.I. Tello and W. Winkler M, A chemotaxis system with logistic source, Common Partial Diff. Eq., 32 (2007), 849-877.
- [34] G. Viglialoro, Boundedness properties of very weak solutions to a fully parabolic chemotaxissystem with logistic source, *Nonlinear Anal. Real World Appl.* **34** (2017), 520-535.
- [35] G. Viglialoro and T. E. Woolley, Eventual smoothness and asymptotic behaviour of solutions to a chemotaxis system perturbed by a logistic growth, *Discrete Contin. Dyn. Syst. Ser. B* 23 (2018), no. 8, 3023-3045.
- [36] M. Winkler, Blow-up in a higher-dimensional chemotaxis system despite logistic growth restriction, *Journal of Mathematical Analysis and Applications*, **384** (2011), 261-272.
- [37] M. Winkler, Global solutions in a fully parabolic chemotaxis system with singular sensitivity, Math. Methods Appl. Sci. 34 (2011), 176-190.
- [38] Winkler, M. Finite-time blow-up in low-dimensional Keller-Segel systems with logistic-type superlinear degradation, Z. Angew. Math. Phys., 69, 40 (2018), no. 2, Paper No. 69, 40 pp.
- [39] M. Winkler, Chemotaxis with logistic source: very weak global solutions and their boundedness properties, J. Math. Anal. Appl. 348 (2008), no. 2, 708-729.
- [40] X. Zhao and S. Zheng, Global boundedness to a chemotaxis system with singular sensitivity and logistic source, Z. Angew. Math. Phys. 68 (2017), no. 1, Art. 2, 13 pp.
- [41] X. Zhao and S. Zheng, Global existence and boundedness of solutions to a chemotaxis system with singular sensitivity and logistic-type source, J. Differential Equations 267 (2019), no. 2, 826-865.
- [42] P. Zheng, C. Mu, R. Willie, and Z. Hu, Global asymptotic stability of steady states in a chemotaxis-growth system with singular sensitivity, *Comput. Math. Appl.* 75 (2018), no. 5, 1667-1675.