Intersections and Distinct Intersections in Cross-intersecting Families

Peter Frankl¹, Jian Wang²

¹Rényi Institute, Budapest, Hungary

²Department of Mathematics Taiyuan University of Technology Taiyuan 030024, P. R. China

E-mail: ¹peter.frankl@gmail.com, ²wangjian01@tyut.edu.cn

Abstract

Let \mathcal{F}, \mathcal{G} be two cross-intersecting families of k-subsets of $\{1, 2, \ldots, n\}$. Let $\mathcal{F} \wedge \mathcal{G}$, $\mathcal{I}(\mathcal{F}, \mathcal{G})$ denote the families of all intersections $F \cap G$ with $F \in \mathcal{F}, G \in \mathcal{G}$, and all distinct intersections $F \cap G$ with $F \neq G, F \in \mathcal{F}, G \in \mathcal{G}$, respectively. For a fixed $T \subset \{1, 2, \ldots, n\}$, let \mathcal{S}_T be the family of all k-subsets of $\{1, 2, \ldots, n\}$ containing T. In the present paper, we show that $|\mathcal{F} \wedge \mathcal{G}|$ is maximized when $\mathcal{F} = \mathcal{G} = \mathcal{S}_{\{1\}}$ for $n \geq 2k^2 + 8k$, while surprisingly $|\mathcal{I}(\mathcal{F}, \mathcal{G})|$ is maximized when $\mathcal{F} = \mathcal{S}_{\{1,2\}} \cup \mathcal{S}_{\{1,4,5\}} \cup \mathcal{S}_{\{2,3,6\}}$ and $\mathcal{G} = \mathcal{S}_{\{1,3\}} \cup \mathcal{S}_{\{2,4\}} \cup \mathcal{S}_{\{1,4,6\}} \cup \mathcal{S}_{\{2,3,5\}}$ for $n \geq 100k^2$. The maximum number of distinct intersections in a t-intersecting family is determined for $n \geq 3(t+2)^3k^2$ as well.

1 Introduction

Let n, k be positive integers and let $[n] = \{1, 2, ..., n\}$ denote the standard *n*-element set. Let $\binom{[n]}{k}$ denote the collection of all *k*-subsets of [n]. Subsets of $\binom{[n]}{k}$ are called *k*-uniform hypergraphs or *k*-graphs for short. A *k*-graph \mathcal{F} is called *intersecting* if $F \cap F' \neq \emptyset$ for all $F, F' \in \mathcal{F}$. For a fixed set $T \subset [n]$, define the *T*-star \mathcal{S}_T by $\mathcal{S}_T = \{S \in \binom{[n]}{k}: T \subset S\}$. We often write $\mathcal{S}_p, \mathcal{S}_{pq}$ and \mathcal{S}_{pqr} for $\mathcal{S}_{\{p\}}, \mathcal{S}_{\{p,q\}}$ and $\mathcal{S}_{\{p,q,r\}}$, respectively. One of the most fundamental theorems in extremal set theory is the following:

Erdős-Ko-Rado Theorem ([1]). Suppose that $n \ge 2k$ and $\mathcal{F} \subset {\binom{[n]}{k}}$ is intersecting. Then

(1)
$$|\mathcal{F}| \le \binom{n-1}{k-1}$$

Hilton and Milner [6] proved that S_1 is the only family that achieves equality in (1) up to isomorphism for n > 2k.

Two families $\mathcal{F}, \mathcal{G} \subset {[n] \choose k}$ are called *cross-intersecting* if any two sets $F \in \mathcal{F}, G \in \mathcal{G}$ have non-empty intersection. If $\mathcal{A} \subset {[n] \choose k}$ is intersecting, then $\mathcal{F} = \mathcal{A}, \mathcal{G} = \mathcal{A}$ are cross-intersecting. Therefore the following result is a strengthening of (1).

Theorem 1.1 ([7]). Suppose that $n \ge 2k$ and $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ are cross-intersecting. Then

(2)
$$|\mathcal{F}||\mathcal{G}| \le {\binom{n-1}{k-1}}^2$$

Let us introduce the central notion of the present paper.

Definition 1.2. For $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ define

$$\mathcal{F} \land \mathcal{G} = \{F \cap G \colon F \in \mathcal{F}, G \in \mathcal{G}\} \text{ and } \mathcal{I}(\mathcal{F}, \mathcal{G}) = \{F \cap G \colon F \in \mathcal{F}, G \in \mathcal{G}, F \neq G\}.$$

Clearly $\mathcal{F} \wedge \mathcal{G} = (\mathcal{F} \cap \mathcal{G}) \cup \mathcal{I}(\mathcal{F}, \mathcal{G})$. For $\mathcal{F} = \mathcal{G}$, we often write $\mathcal{I}(\mathcal{F})$ instead of $\mathcal{I}(\mathcal{F}, \mathcal{F})$.

The first result of the present paper shows another extremal property of the full star.

Theorem 1.3. Suppose that $n \ge 2k^2 + 8k$, $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ are cross-intersecting. Then

(3)
$$|\mathcal{F} \wedge \mathcal{G}| \le \sum_{0 \le i \le k-1} \binom{n-1}{i}$$

where equality holds if and only if $\mathcal{F} = \mathcal{G} = \mathcal{S}_1$ up to isomorphism.

Corollary 1.4. Suppose that $n \ge 2k^2 + 8k$, $\mathcal{F} \subset {\binom{[n]}{k}}$ is intersecting. Then

$$|\mathcal{F} \wedge \mathcal{F}| \leq \sum_{0 \leq i \leq k-1} \binom{n-1}{i}$$

where equality holds if and only if $\mathcal{F} = \mathcal{S}_1$ up to isomorphism.

One would expect that both Theorem 1.3 and Corollary 1.4 hold for n > ck for some absolute constant c. Unfortunately, we could not prove it. We can demonstrate the same results for $n > c'k^2/\log k$ with a more complicated proof.

Let us now consider the probably more natural quantity $|\mathcal{I}(\mathcal{F},\mathcal{G})|$, namely the case that intersections of identical sets are not counted. Quite surprisingly the pairs of families maximizing $|\mathcal{I}(\mathcal{F},\mathcal{G})|$ is rather peculiar. The fact that we can prove the optimality of such a pair shows the strength of our methods.

Let us define the two families

$$\mathcal{A}_1 = \mathcal{S}_{12} \cup \mathcal{S}_{34} \cup \mathcal{S}_{145} \cup \mathcal{S}_{236} \text{ and } \mathcal{A}_2 = \mathcal{S}_{13} \cup \mathcal{S}_{24} \cup \mathcal{S}_{146} \cup \mathcal{S}_{235}.$$

One can check that $\mathcal{A}_1, \mathcal{A}_2$ are cross-intersecting.

Proposition 1.5.

$$|\mathcal{I}(\mathcal{A}_{1},\mathcal{A}_{2})| = 4 \sum_{0 \le i \le k-2} \binom{n-4}{i} + 6 \sum_{0 \le i \le k-3} \binom{n-4}{i} + 4 \sum_{0 \le i \le k-4} \binom{n-4}{i} + \sum_{0 \le i \le k-5} \binom{n-4}{i} + 2 \sum_{i \le i \le k-3} \binom{n-6}{i} + \sum_{0 \le i \le k-4} \binom{n-6}{i}.$$

Proof. For any $A_1 \in \mathcal{A}_1$ and $A_2 \in \mathcal{A}_2$, there are $\binom{4}{1} \sum_{\substack{0 \leq i \leq k-2 \\ i}} \binom{n-4}{i}$ distinct intersections for $|A_1 \cap A_2 \cap \{1, 2, 3, 4\}| = 1$. There are $\binom{4}{2} \sum_{\substack{0 \leq i \leq k-3 \\ 0 \leq i \leq k-3}} \binom{n-4}{i}$ distinct intersections for $|A_1 \cap A_2 \cap \{1, 2, 3, 4\}| = 2$. There are $\binom{4}{3} \sum_{\substack{0 \leq i \leq k-4 \\ i}} \binom{n-4}{i}$ distinct intersections for $|A_1 \cap A_2 \cap \{1, 2, 3, 4\}| = 4$. There are $2 \sum_{\substack{0 \leq i \leq k-3 \\ 0 \leq i \leq k-3}} \binom{n-6}{i}$ distinct intersections for $|A_1 \cap A_2 \cap \{1, 2, 3, 4\}| = 0$ and $|A_1 \cap A_2 \cap \{5, 6\}| = 1$. There are $\sum_{\substack{0 \leq i \leq k-4 \\ i}} \binom{n-6}{i}$ distinct intersections for $|A_1 \cap A_2 \cap \{1, 2, 3, 4\}| = 0$ and $|A_1 \cap A_2 \cap \{5, 6\}| = 2$. Thus the proposition follows.

Our main result shows that $|\mathcal{I}(\mathcal{F}, \mathcal{G})|$ is maximized by $\mathcal{A}_1, \mathcal{A}_2$ over all cross-intersecting families $\mathcal{F}, \mathcal{G} \subset {[n] \choose k}$ for $n \geq 100k^2$.

Theorem 1.6. If $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ are cross-intersecting families and $n \geq 100k^2$, then $|\mathcal{I}(\mathcal{F}, \mathcal{G})| \leq |\mathcal{I}(\mathcal{A}_1, \mathcal{A}_2)|$.

Let $n \ge k > t$. A family $\mathcal{F} \subset {\binom{[n]}{k}}$ is called *t*-intersecting if any two members of it intersect in at least *t* elements. Note that for $n \le 2k - t$ the whole set ${\binom{[n]}{k}}$ is *t*-intersecting. Thus we always assume that n > 2k - t when considering extremal problems for *t*-intersecting families.

Define

$$\mathcal{A}(n,k,t) = \left\{ A \in \binom{[n]}{k} : |A \cap [t+2]| \ge t+1 \right\}.$$

This family was first defined in [2] and it is easily seen to be *t*-intersecting.

Proposition 1.7.

$$|\mathcal{I}(\mathcal{A}(n,k,t))| = \binom{t+2}{t} \sum_{0 \le i \le k-t-1} \binom{n-t-2}{i} + \binom{t+2}{t+1} \sum_{0 \le i \le k-t-2} \binom{n-t-2}{i} + \sum_{0 \le i \le k-t-3} \binom{n-t-2}{i}.$$
(5)

Proof. For any $A_1, A_2 \in \mathcal{A}(n, k, t)$, we have $|A_1 \cap A_2 \cap [t+2]| \ge t$. Note that $|A_i \cap [t+2]| \ge t+1$ for i = 1, 2. There are $\binom{t+2}{t} \sum_{\substack{0 \le i \le k-t-1 \\ i}} \binom{n-t-2}{i}$ distinct intersections for $|A_1 \cap A_2 \cap [t+2]| = t+1$. 2]| = t. There are $\binom{t+2}{t+1} \sum_{\substack{0 \le i \le k-t-2 \\ i}} \binom{n-t-2}{i}$ distinct intersections for $|A_1 \cap A_2 \cap [t+2]| = t+1$. There are $\sum_{\substack{0 \le i \le k-t-2 \\ i}} \binom{n-t-2}{i}$ distinct intersections for $|A_1 \cap A_2 \cap [t+2]| = t+2$. Thus the proposition follows.

Our third result shows that $|\mathcal{I}(\mathcal{F})|$ is maximized by $\mathcal{A}(n,k,t)$ over all intersecting families $\mathcal{F} \subset {[n] \choose k}$ for $n \geq 3(t+2)^3 k^2$.

Theorem 1.8. If $\mathcal{F} \subset {[n] \choose k}$ is a t-intersecting family and $n \geq 3(t+2)^3k^2$, then $\mathcal{I}(\mathcal{F}) \leq |\mathcal{I}(\mathcal{A}(n,k,t))|$.

We should mention that this result was proved for the case t = 1 in [5].

Let us list some notions and results that we need for the proofs. Define the family of *t*-transversals of $\mathcal{F} \subset {[n] \choose k}$:

$$\mathcal{T}_t(\mathcal{F}) = \{T \subset [n] \colon |T| \le k, |T \cap F| \ge t \text{ for all } F \in \mathcal{F}\}.$$

Clearly, if \mathcal{F} is *t*-intersecting then $\mathcal{F} \subset \mathcal{T}_t(\mathcal{F})$ and vice versa. The *t*-covering number $\tau_t(\mathcal{F})$ is defined as follows:

$$\tau_t(\mathcal{F}) = \min\{|T| \colon |T \cap F| \ge t \text{ for all } F \in \mathcal{F}\}.$$

For t = 1, we often write $\mathcal{T}(\mathcal{F}), \tau(\mathcal{F})$ instead of $\mathcal{T}_1(\mathcal{F}), \tau_1(\mathcal{F})$, respectively. If \mathcal{F}, \mathcal{G} are cross-intersecting, then clearly $\mathcal{F} \subset \mathcal{T}(\mathcal{G})$ and $\mathcal{G} \subset \mathcal{T}(\mathcal{F})$.

Let us recall the following common notations:

$$\mathcal{F}(i) = \{F \setminus \{i\} \colon i \in F \in \mathcal{F}\}, \qquad \mathcal{F}(\overline{i}) = \{F \in \mathcal{F} : i \notin F\}.$$

Note that $|\mathcal{F}| = |\mathcal{F}(i)| + |\mathcal{F}(\overline{i})|$.

Define $\nu(\mathcal{F})$, the matching number of \mathcal{F} as the maximum number of pairwise disjoint edges in \mathcal{F} . Note that $\nu(\mathcal{F}) = 1$ iff \mathcal{F} is intersecting. We need the following inequality generalising the Erdős-Ko-Rado Theorem.

Proposition 1.9 ([3]). Suppose that $\mathcal{F} \subset {\binom{[n]}{k}}$ then

(6)
$$|\mathcal{F}| \le \nu(\mathcal{F}) \binom{n-1}{k-1}.$$

An intersecting family \mathcal{F} is called *non-trivial* if $\bigcap_{F \in \mathcal{F}} F = \emptyset$. We also need the following stability theorem concerning the Erdős-Ko-Rado Theorem.

Hilton-Milner Theorem ([6]). If n > 2k and $\mathcal{F} \subset {\binom{[n]}{k}}$ is non-trivial intersecting, then

(7)
$$|\mathcal{F}| \le \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1.$$

Let us list some inequalities that will be used frequently in the proof.

Proposition 1.10. Let n, k, ℓ, t, p be positive integers with $k > \ell$, k > t and n > 2k + p. Then

(8)
$$\binom{n}{k} \leq \frac{n-p}{n-p(k+1)} \binom{n-p}{k}$$

(9)
$$\sum_{0 \le i \le k-\ell} \binom{n-t}{i} \le \frac{n-t-p}{n-t-pk} \sum_{0 \le i \le k-\ell} \binom{n-t-p}{i},$$

(10)
$$\sum_{0 \le i \le k-\ell-1} \binom{n-t}{i} \le \frac{k}{n-t-k} \sum_{0 \le i \le k-\ell} \binom{n-t}{i},$$

(11)
$$\qquad for \ \ell \ge t+1, \ \sum_{t \le j \le \ell} \binom{\ell}{j} \ge \frac{1}{2t+2} \sum_{t \le j \le \ell+1} \binom{\ell+1}{j}.$$

Proof. Note that

$$\frac{\binom{n-p}{k}}{\binom{n}{k}} = \frac{(n-k)(n-k-1)\cdots(n-k-p+1)}{n(n-1)\cdots(n-p+1)} \ge \left(1-\frac{k}{n-p}\right)^p \ge 1-\frac{pk}{n-p}.$$

Then (8) holds. By (8), we have for i < k

$$\binom{n-t}{i} \le \frac{n-t-p}{n-t-p(i+1)} \binom{n-t-p}{i} \le \frac{n-t-p}{n-t-pk} \binom{n-t-p}{i},$$

and thereby (9) follows. Since

$$\binom{n-t}{i-1} / \binom{n-t}{i} = \frac{i}{n-t-i+1} \le \frac{k}{n-t-k},$$

we obtain (10).

For $\ell \geq 2t$, since

$$\sum_{t \le j \le \ell} {\ell \choose j} \ge 2^{\ell-1} \text{ and } \sum_{t \le j \le \ell+1} {\ell+1 \choose j} \le 2^{\ell+1}$$

we see that

$$\frac{\sum\limits_{t \le j \le \ell} {\ell \choose j}}{\sum\limits_{t \le j \le \ell+1} {\ell+1 \choose j}} \ge \frac{1}{4}.$$

For $t+1 \leq \ell \leq 2t$,

$$\begin{split} \sum_{t \leq j \leq \ell} \binom{\ell}{j} &\geq \sum_{t \leq j \leq \ell} \frac{\ell + 1 - j}{\ell + 1} \binom{\ell + 1}{j} \\ &\geq \sum_{t \leq j \leq \ell - 1} \frac{\ell + 1 - j}{\ell + 1} \binom{\ell + 1}{j} + \frac{1}{\ell + 1} \binom{\ell + 1}{\ell} \\ &\geq \frac{1}{\ell + 1} \sum_{t \leq j \leq \ell - 1} \binom{\ell + 1}{j} + \frac{1}{\ell + 2} \left(\binom{\ell + 1}{\ell} + \binom{\ell + 1}{\ell + 1} \right) \\ &> \frac{1}{2t + 2} \sum_{t \leq j \leq \ell + 1} \binom{\ell + 1}{j}. \end{split}$$

Thus (11) holds.

2 Intersections in cross-intersecting families

In this section, we determine the maximum size of $\mathcal{F} \wedge \mathcal{G}$ over all cross-intersecting families $\mathcal{F}, \mathcal{G} \subset {[n] \choose k}$. We also determine the maximum size of $(\mathcal{F}_1 \wedge \mathcal{G}_1) \cup (\mathcal{F}_2 \wedge \mathcal{G}_2)$ over all families $\mathcal{F}_1, \mathcal{F}_2, \mathcal{G}_1, \mathcal{G}_2 \subset {[n] \choose k}$ with $\mathcal{F}_1, \mathcal{G}_1$ being cross-intersecting and $\mathcal{F}_2, \mathcal{G}_2$ being cross-intersecting. This result will be used in Section 3.

First we prove a key proposition to the proof of Theorem 1.3.

Proposition 2.1. Let $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ be cross-intersecting families and set $\mathcal{H} = \mathcal{I}(\mathcal{F}, \mathcal{G}) \cap {\binom{[n]}{k-1}}$. Then $\nu(\mathcal{H}) \leq 4$.

Proof. Suppose that $F_i \cap G_i = D_i$ are pairwise disjoint (k-1)-sets, $0 \le i \le 4$. Define x_i, y_i by $F_i = D_i \cup \{x_i\}, G_i = D_i \cup \{y_i\}$ and note that $x_i \ne y_i$. There are altogether 5×4 conditions $F_i \cap G_j \ne \emptyset$ to satisfy. Each of them is assured by either of the following three relations: $x_i \in D_j, y_j \in D_i, x_i = y_j$. From the first two types there are at most one for each x_i and y_j . Altogether at most 5+5=10. If no multiple equalities (e.g. $x_1 = y_2 = y_3$)

exist, we get only at most 5 more relations and 10 + 5 < 20. Thus there must be places of coincidence, say by symmetry that of the form $x_i = x_{i'}$. Thus, again by symmetry, we may assume that $x_i \notin D_0$ for $0 \le i \le 4$. Note that $y_0 \in D_i$ holds for at most one value of *i*. Without loss of generality assume $y_0 \notin D_i$, $1 \le i \le 3$. By $F_i \cap G_0 \neq \emptyset$, $y_0 = x_i$, i = 1, 2, 3. Look at y_1 . By symmetry assume $y_1 \notin D_2$. Now $G_1 \cap F_2 \neq \emptyset$ implies $y_1 = x_2$. Hence $y_1 = x_1$, a contradiction.

Let D_1, D_2, D_3, D_4 be pairwise disjoint (k-1)-sets. Pick an element $d_i \in D_i$, i = 1, 2, 3. Define x_i, y_i by $x_1 = x_2 = y_4 = d_3$, $x_3 = y_1 = d_2$ and $x_4 = y_2 = y_3 = d_1$. Setting $F_i = D_i \cup \{x_i\}, G_i = D_j \cup \{y_j\}$. One can check easily that $F_i \cap G_j \neq \emptyset$ for $1 \le i \ne j \le 4$. This example shows that Proposition 2.1 is best possible.

Proof of Theorem 1.3. We distinguish two cases. First we suppose that

(12)
$$|\mathcal{F} \cap \mathcal{G}| > \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1.$$

Since \mathcal{F}, \mathcal{G} are cross-intersecting, $\mathcal{F} \cap \mathcal{G}$ is intersecting. By (7) and (12), without loss of generality, we assume that $1 \in F$ for all $F \in \mathcal{F} \cap \mathcal{G}$. We claim that $1 \in H$ for all $H \in \mathcal{F} \cup \mathcal{G}$. Indeed, if $1 \notin H \in \mathcal{F} \cup \mathcal{G}$ then $H \cap F \neq \emptyset$ for $F \in \mathcal{F} \cap \mathcal{G}$ yields

$$|\mathcal{F} \cap \mathcal{G}| \le {\binom{n-1}{k-1}} - {\binom{n-k-1}{k-1}}$$

contradicting (12). We proved that $1 \in H$ for all $H \in \mathcal{F} \cup \mathcal{G}$ and thereby (3) holds.

Suppose next that (12) does not hold. By Proposition 2.1 and (6), we have for $n \ge 5k$,

$$\left| \mathcal{I}(\mathcal{F},\mathcal{G}) \cap \binom{[n]}{k-1} \right| \le 4 \binom{n-1}{k-2} \stackrel{(8)}{\le} \frac{4(n-2)}{n-k} \binom{n-2}{k-2} \le 5 \binom{n-2}{k-2}$$

Since the remaining sets in $\mathcal{I}(\mathcal{F},\mathcal{G})$ are of size at most k-2, we have

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| \le 5\binom{n-2}{k-2} + \sum_{0 \le i \le k-2} \binom{n}{i}.$$

Moreover,

$$|\mathcal{F} \cap \mathcal{G}| \le \binom{n-1}{k-1} - \binom{n-k-1}{k-1} + 1 \le k \binom{n-2}{k-2}.$$

Thus, for $n \ge 2k + 1$ we have

$$\begin{aligned} |\mathcal{F} \wedge \mathcal{G}| &\leq (k+5)\binom{n-2}{k-2} + \sum_{0 \leq i \leq k-2} \binom{n}{i} \\ &\stackrel{(10)}{\leq} \frac{(k+5)(k-1)}{n-1}\binom{n-1}{k-1} + \frac{k}{n-k} \sum_{0 \leq i \leq k-1} \binom{n}{i} \\ &\stackrel{(9)}{\leq} \frac{(k+5)(k-1)}{n-1}\binom{n-1}{k-1} + \frac{k}{n-k} \frac{n-1}{n-k} \sum_{0 \leq i \leq k-1} \binom{n-1}{i}. \end{aligned}$$

Note that $n \ge 2k^2 + 8k$ implies

$$\frac{(k+5)(k-1)}{n-1} \le \frac{1}{2}$$

and

$$\frac{k}{n-k}\frac{n-1}{n-k} < \frac{k}{n-k}\left(1+\frac{k}{n-k}\right) < \frac{k}{2k^2+7k}\left(1+\frac{k}{2k^2}\right) = \frac{2k+1}{2k(2k+7)} \le \frac{1}{2}.$$

Thus,

1

$$|\mathcal{F} \wedge \mathcal{G}| \leq \frac{1}{2} \binom{n-1}{k-1} + \frac{1}{2} \sum_{0 \leq i \leq k-1} \binom{n-1}{i} < \sum_{0 \leq i \leq k-1} \binom{n-1}{i}.$$

Lemma 2.2. Suppose that $n \ge 2k^2 + 9k$, $\mathcal{F}_1, \mathcal{G}_1 \subset {\binom{[n]}{k}}$ are cross-intersecting and $\mathcal{F}_2, \mathcal{G}_2 \subset {\binom{[n]}{k}}$ are cross-intersecting. Then

(13)
$$|(\mathcal{F}_1 \wedge \mathcal{G}_1) \cup (\mathcal{F}_2 \wedge \mathcal{G}_2)| \le 2 \sum_{0 \le i \le k-1} \binom{n-2}{i} + \sum_{0 \le i \le k-2} \binom{n-2}{i}.$$

with equality holding if and only if $\mathcal{F}_1 = \mathcal{G}_1 = \mathcal{S}_1$ and $\mathcal{F}_2 = \mathcal{G}_2 = \mathcal{S}_2$ up to isomorphism. Proof. By Theorem 1.3, for j = 1, 2

$$|\mathcal{F}_j \wedge \mathcal{G}_j| \le \sum_{0 \le i \le k-1} \binom{n-1}{i}.$$

By Proposition 2.1 and (6), for j = 1, 2

$$\left| \mathcal{I}(\mathcal{F}_j, \mathcal{G}_j) \cap \binom{[n]}{k-1} \right| \le 4\binom{n-1}{k-2} \stackrel{(8)}{\le} \frac{4(n-2)}{n-k} \binom{n-2}{k-2} \le 5\binom{n-2}{k-2},$$

Since the remaining sets in $\mathcal{I}(\mathcal{F}_j, \mathcal{G}_j)$ are of size at most k-2, for j=1,2

$$|\mathcal{I}(\mathcal{F}_j, \mathcal{G}_j)| \le 5 \binom{n-2}{k-2} + \sum_{0 \le i \le k-2} \binom{n}{i}.$$

If $|\mathcal{F}_{j} \cap \mathcal{G}_{j}| \leq {\binom{n-1}{k-1}} - {\binom{n-k-1}{k-1}} \leq k {\binom{n-2}{k-2}}$ for some $j \in \{1, 2\}$, then for $n \geq 2k+2$ $|(\mathcal{F}_{1} \wedge \mathcal{G}_{1}) \cup (\mathcal{F}_{2} \wedge \mathcal{G}_{2})|$ $\leq |\mathcal{F}_{1} \wedge \mathcal{G}_{1}| + |\mathcal{F}_{2} \wedge \mathcal{G}_{2}|$ $\leq k {\binom{n-2}{k-2}} + 5 {\binom{n-2}{k-2}} + \sum_{0 \leq i \leq k-2} {\binom{n}{i}} + \sum_{0 \leq i \leq k-1} {\binom{n-1}{i}}$ $\stackrel{(9)}{\leq} \frac{(k+5)(k-1)}{n-k} {\binom{n-2}{k-1}} + \frac{n-2}{n-2k} \sum_{0 \leq i \leq k-2} {\binom{n-2}{i}} + \frac{n-2}{n-1-k} \sum_{0 \leq i \leq k-1} {\binom{n-2}{i}}.$

Note that $n \ge 2k^2 + 9k \ge 10k$ implies

$$\frac{(k+5)(k-1)}{n-k} \le \frac{1}{2}, \ \frac{n-2}{n-2k} \le \frac{5}{4} \text{ and } \frac{n-2}{n-1-k} \le \frac{5}{4}.$$

Thus,

$$|(\mathcal{F}_{1} \wedge \mathcal{G}_{1}) \cup (\mathcal{F}_{2} \wedge \mathcal{G}_{2})| \le \frac{1}{2} \binom{n-2}{k-1} + \frac{5}{4} \sum_{0 \le i \le k-2} \binom{n-2}{i} + \frac{5}{4} \sum_{0 \le i \le k-1} \binom{n-2}{i} \le 2 \sum_{0 \le i \le k-1} \binom{n-2}{i} + \sum_{0 \le i \le k-2} \binom{n-2}{i}.$$

Thus we may assume that $|\mathcal{F}_j \cap \mathcal{G}_j| \ge {\binom{n-1}{k-1}} - {\binom{n-k-1}{k-1}} + 1$ for each j = 1, 2. By (7), both $\mathcal{F}_1 \cap \mathcal{G}_1$ and $\mathcal{F}_2 \cap \mathcal{G}_2$ are trivial intersecting families. By the same argument as in Theorem 1.3, we see that there exist x, y such that $\mathcal{F}_1 \cup \mathcal{G}_1 \subset \mathcal{S}_x$ and $\mathcal{F}_2 \cup \mathcal{G}_2 \subset \mathcal{S}_y$. If $x \ne y$, then we are done. If x = y, then

$$\begin{aligned} |(\mathcal{F}_1 \wedge \mathcal{G}_1) \cup (\mathcal{F}_2 \wedge \mathcal{G}_2)| &\leq |\mathcal{S}_x \wedge \mathcal{S}_x| \\ &= \sum_{0 \leq i \leq k-1} \binom{n-1}{i} \\ &\leq 2 \sum_{0 \leq i \leq k-1} \binom{n-2}{i} + \sum_{0 \leq i \leq k-2} \binom{n-2}{i}. \end{aligned}$$

3 Distinct intersections in cross-intersecting families

In this section, we determine the maximum number of distinct intersections in crossintersecting families.

For the proof, we need the following notion of basis. Two cross-intersecting families \mathcal{F}, \mathcal{G} are called *saturated* if any cross-intersecting families $\tilde{\mathcal{F}}, \tilde{\mathcal{G}}$ with $\mathcal{F} \subset \tilde{\mathcal{F}}, \mathcal{G} \subset \tilde{\mathcal{G}}$ have $\mathcal{F} = \tilde{\mathcal{F}}$ and $\mathcal{G} = \tilde{\mathcal{G}}$. Since $\mathcal{F} \subset \tilde{\mathcal{F}}$ and $\mathcal{G} \subset \tilde{\mathcal{G}}$ imply $\mathcal{I}(\mathcal{F}, \mathcal{G}) \subset \mathcal{I}(\tilde{\mathcal{F}}, \tilde{\mathcal{G}})$, we may always assume that \mathcal{F}, \mathcal{G} are saturated when maximizing the size of $\mathcal{I}(\mathcal{F}, \mathcal{G})$. Let $\mathcal{B}(\mathcal{F})$ be the family of minimal (for containment) sets in $\mathcal{T}(\mathcal{G})$ and let $\mathcal{B}(\mathcal{G})$ be the family of minimal sets in $\mathcal{T}(\mathcal{F})$. Let us prove some properties of the basis.

Lemma 3.1. Suppose that $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ are saturated cross-intersecting families. Then (i) and (ii) hold.

(i) Both $\mathcal{B}(\mathcal{F})$ and $\mathcal{B}(\mathcal{G})$ are antichains, and $\mathcal{B}(\mathcal{F}), \mathcal{B}(\mathcal{G})$ are cross-intersecting,

(*ii*)
$$\mathcal{F} = \left\{ F \in \binom{[n]}{k} : \exists B \in \mathcal{B}(\mathcal{F}), B \subset F \right\}$$
 and $\mathcal{G} = \left\{ G \in \binom{[n]}{k} : \exists B \in \mathcal{B}(\mathcal{G}), B \subset G \right\}.$

Proof. (i) Clearly, $\mathcal{B}(\mathcal{F})$ and $\mathcal{B}(\mathcal{G})$ are both anti-chains. Suppose for contradiction that $B \in \mathcal{B}(\mathcal{F}), B' \in \mathcal{B}(\mathcal{G})$ but $B \cap B' = \emptyset$. If |B| = |B'| = k, then $B \in \mathcal{F}, B' \in \mathcal{G}$ follows from saturatedness, a contradiction. If |B| < k, then there exists $F \supset B$ such that |F| = k and $|F \cap B'| = |B \cap B'| = 0$. By definition $F \in \mathcal{T}(\mathcal{G})$. Since \mathcal{F}, \mathcal{G} are saturated, we see that $F \in \mathcal{F}$. But this contradicts the assumption that B' is a transversal of \mathcal{F} . Since \mathcal{F}, \mathcal{G} are saturated, (ii) is immediate from the definition of $\mathcal{B}(\mathcal{F})$ and $\mathcal{B}(\mathcal{G})$.

Let $r(\mathcal{B}) = \max\{|B|: B \in \mathcal{B}\}$ and $s(\mathcal{B}) = \min\{|B|: B \in \mathcal{B}\}$. For any ℓ with $s(\mathcal{B}) \leq \ell \leq r(\mathcal{B})$, define

$$\mathcal{B}^{(\ell)} = \{B \in \mathcal{B} \colon |B| = \ell\} \text{ and } \mathcal{B}^{(\leq \ell)} = \bigcup_{i=s(\mathcal{B})}^{\ell} \mathcal{B}^{(i)}$$

It is easy to see that $s(\mathcal{B}(\mathcal{G})) = \tau(\mathcal{F})$.

By a branching process, we establish an upper bound on the size of the basis.

Lemma 3.2. Suppose that $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ are saturated cross-intersecting families. Let $\mathcal{B}_1 = \mathcal{B}(\mathcal{F})$ and $\mathcal{B}_2 = \mathcal{B}(\mathcal{G})$. For each i = 1, 2, if $s(\mathcal{B}_i) \geq 2$ and $\tau(\mathcal{B}_i^{(\leq r_i)}) \geq 2$ then

(14)
$$\sum_{r_i \le \ell \le k} \ell^{-2} k^{-\ell+2} |\mathcal{B}_{3-i}^{(\ell)}| \le 1.$$

Proof. By symmetry, it is sufficient to prove the lemma only for i = 1. For the proof we use a branching process. During the proof a sequence $S = (x_1, x_2, \ldots, x_\ell)$ is an ordered sequence of distinct elements of [n] and we use \hat{S} to denote the underlying unordered set $\{x_1, x_2, \ldots, x_\ell\}$. At the beginning, we assign weight 1 to the empty sequence S_{\emptyset} . At the first stage, we choose $B_{1,1} \in \mathcal{B}_1$ with $|B_{1,1}| = s(\mathcal{B}_1)$. For any vertex $x_1 \in B_{1,1}$, define one sequence (x_1) and assign the weight $s(\mathcal{B}_1)^{-1}$ to it.

At the second stage, since $\tau(\mathcal{B}_1^{(\leq r_1)}) \geq 2$, for each sequence $S = (x_1)$ we may choose $B_{1,2} \in \mathcal{B}_1^{(\leq r_1)}$ such that $x_1 \notin B_{1,2}$. Then we replace $S = (x_1)$ by $|B_{1,2}|$ sequences of the form (x_1, y) with $y \in B_{1,2}$ and weight $\frac{w(S)}{|B_{1,2}|}$.

In each subsequent stage, we pick a sequence $S = (x_1, \ldots, x_p)$ and denote its weight by w(S). If $\hat{S} \cap B_1 \neq \emptyset$ holds for all $B_1 \in \mathcal{B}_1$ then we do nothing. Otherwise we pick $B_1 \in \mathcal{B}_1$ satisfying $\hat{S} \cap B_1 = \emptyset$ and replace S by the $|B_1|$ sequences (x_1, \ldots, x_p, y) with $y \in B_1$ and assign weight $\frac{w(S)}{|B_1|}$ to each of them. Clearly, the total weight is always 1.

We continue until $S \cap B_1 \neq \emptyset$ for all sequences and all $B_1 \in \mathcal{B}_1$. Since [n] is finite, each sequence has length at most n and eventually the process stops. Let S be the collection of sequences that survived in the end of the branching process and let $S^{(\ell)}$ be the collection of sequences in S with length ℓ .

Claim 1. To each $B_2 \in \mathcal{B}_2^{(\ell)}$ with $\ell \geq r_1$ there is some sequence $S \in \mathcal{S}^{(\ell)}$ with $\widehat{S} = B_2$.

Proof. Let us suppose the contrary and let $S = (x_1, \ldots, x_p)$ be a sequence of maximal length that occurred at some stage of the branching process satisfying $\widehat{S} \subseteq B_2$. Since $\mathcal{B}_1, \mathcal{B}_2$ are cross-intersecting, $B_{1,1} \cap B_2 \neq \emptyset$, implying that $p \geq 1$. Since \widehat{S} is a proper subset of B_2 and $B_2 \in \mathcal{B}_2 = \mathcal{B}(\mathcal{G})$, it follows that $\widehat{S} \notin \mathcal{B}(\mathcal{G}) \subset \mathcal{T}(\mathcal{F})$. Thereby there exists $F \in \mathcal{F}$ with $\widehat{S} \cap F = \emptyset$. In view of Lemma 3.1 (ii), we can find $B'_1 \in \mathcal{B}_1$ such that $\widehat{S} \cap B'_1 = \emptyset$. Thus at some point we picked S and some $\widetilde{B}_1 \in \mathcal{B}_1$ with $\widehat{S} \cap \widetilde{B}_1 = \emptyset$. Since $\mathcal{B}_1, \mathcal{B}_2$ are cross-intersecting, $B_2 \cap \widetilde{B}_1 \neq \emptyset$. Consequently, for each $y \in B_2 \cap \widetilde{B}_1$ the sequence (x_1, \ldots, x_p, y) occurred in the branching process. This contradicts the maximality of p. Hence there is an S at some stage satisfying $\widehat{S} = B_2$. Since $\mathcal{B}_1, \mathcal{B}_2$ are cross-intersecting, $\widehat{S} \cap B'_1 = B_2 \cap B'_1 \neq \emptyset$ for all $B'_1 \in \mathcal{B}_1$. Thus $\widehat{S} \in \mathcal{S}$ and the claim holds.

By Claim 1, we see that $|\mathcal{B}_{2}^{(\ell)}| \leq |\mathcal{S}^{(\ell)}|$ for all $\ell \geq r_{1}$. Let $S = (x_{1}, \ldots, x_{\ell}) \in \mathcal{S}^{(\ell)}$ and let $S_{i} = (x_{1}, \ldots, x_{i})$ for $i = 1, \ldots, \ell$. At the first stage, $w(S_{1}) = 1/s(\mathcal{B}_{1})$. Assume that $B_{1,i}$ is the selected set when replacing S_{i-1} in the branching process for $i = 2, \ldots, \ell$. Clearly, $x_{i} \in B_{i}, B_{1,2} \in \mathcal{B}_{1}^{(\leq r_{1})}$ and

$$w(S) = \frac{1}{s(\mathcal{B}_1)} \prod_{i=2}^{\ell} \frac{1}{|B_{1,i}|}$$

Note that $s(\mathcal{B}_1) \leq \ell$, $|B_{1,2}| = r_1 \leq \ell$ and $|B_{1,i}| \leq k$ for $i \geq 3$. It follows that

$$w(S) \ge \left(\ell^2 k^{\ell-2}\right)^{-1} = \ell^{-2} k^{-\ell+2}.$$

Thus,

$$\sum_{r_1 \le \ell \le k} \ell^{-2} k^{-\ell+2} |\mathcal{B}_2^{(\ell)}| \le \sum_{r_1 \le \ell \le k} \sum_{S \in \mathcal{S}^{(\ell)}} w(S) \le \sum_{S \in \mathcal{S}} w(S) = 1.$$

For the proof of Theorem 1.6, we also need the following lemma.

Lemma 3.3. Suppose that $\mathcal{F} \subset {\binom{[n]}{k}}, \mathcal{G} \subset {\binom{[n]}{k-1}}$ are cross-intersecting. Then

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| \le 2\binom{n-1}{k-2} + (2k+1)\binom{n-1}{k-3} + \sum_{0 \le i \le k-3} \binom{n}{i}.$$

Proof. Let $\mathcal{H}_1 = \mathcal{I}(\mathcal{F}, \mathcal{G}) \cap {[n] \choose k-1}$. We claim that $\nu(\mathcal{H}_1) \leq 2$. Otherwise, let $G_i = F_i \cap G_i$, i = 1, 2, 3, be three pairwise disjoint members in \mathcal{H}_1 with $F_i \in \mathcal{F}, G_i \in \mathcal{G}$. Define x_i by $F_i \setminus G_i = \{x_i\}$. By symmetry we may assume that $x_1 \notin G_3$. Then F_1, G_3 are disjoint, contradicting the fact that \mathcal{F}, \mathcal{G} are cross-intersecting. Thus $\nu(\mathcal{H}_1) \leq 2$.

If $\nu(\mathcal{H}_1) \leq 1$, then (1) implies $|\mathcal{H}_1| \leq \binom{n-1}{k-2}$. Since the remaining sets in $\mathcal{I}(\mathcal{F}, \mathcal{G})$ are all of size at most k-2, it follows that

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| \le {\binom{n-1}{k-2}} + \sum_{0 \le i \le k-2} {\binom{n}{i}}.$$

If $\nu(\mathcal{H}_1) = 2$, let $G_1 = F_1 \cap G_1, G_2 = F_2 \cap G_2$ be two disjoint members in \mathcal{H}_1 and let $\mathcal{H}_2 = \mathcal{I}(\mathcal{F}, \mathcal{G}) \cap {\binom{[n] \setminus (F_1 \cup F_2)}{k-2}}$. We claim that \mathcal{H}_2 is intersecting. Suppose not, let $D_3 = F_3 \cap G_3, D_4 = F_4 \cap G_4$ be two disjoint members in \mathcal{H}_2 . Define x_i by $F_i \setminus G_i = \{x_i\}$ for i = 1, 2 and define x_i, y_i, z_i by $F_i \setminus D_i = \{x_i, y_i\}, G_i \setminus D_i = \{z_i\}$ for i = 3, 4. Since $F_3 \cap G_1 \neq \emptyset$ and $F_3 \cap G_2 \neq \emptyset$, by symmetry we may assume that $x_3 \in G_1$ and $y_3 \in G_2$. Similarly, assume that $x_4 \in G_1$ and $y_4 \in G_2$. Since $F_1 \cap G_3 \neq \emptyset$ and $F_2 \cap G_3 \neq \emptyset$, we see that $z_3 \in F_1 \cap F_2$. It follows that $x_1 = x_2 = z_3$. Similarly we have $x_1 = x_2 = z_4$. But then F_3, G_4 are disjoint, contradicting the fact that \mathcal{F}, \mathcal{G} are cross-intersecting. Thus \mathcal{H}_2 is intersecting. By (1) we have

$$\left|\mathcal{I}(\mathcal{F},\mathcal{G})\cap \binom{[n]}{k-2}\right| \leq |F_1\cup F_2|\binom{n-1}{k-3} + \binom{n-2k}{k-3} \leq (2k+1)\binom{n-1}{k-3}.$$

By (6) we obtain that

$$\left| \mathcal{I}(\mathcal{F},\mathcal{G}) \cap {[n] \choose k-1} \right| \leq 2 {n-1 \choose k-2}.$$

Hence

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| \le 2\binom{n-1}{k-2} + (2k+1)\binom{n-1}{k-3} + \sum_{0 \le i \le k-3} \binom{n}{i}.$$

Corollary 3.4. Let $\mathcal{F}, \mathcal{G} \subset {\binom{[n]}{k}}$ be cross-intersecting families. If \mathcal{G} is a star, then

(15)
$$\mathcal{I}(\mathcal{F},\mathcal{G}) \le 2\sum_{0 \le i \le k-2} \binom{n-1}{i} + \binom{n-2}{k-2} + (2k+1)\binom{n-2}{k-3}.$$

Proof. Assume without loss of generality that \mathcal{F} and \mathcal{G} are saturated. Since \mathcal{G} is a star, we may assume that $\mathcal{G} \subset \mathcal{S}_1$. Then $\{1\} \in \mathcal{T}(\mathcal{G})$ whence $\{1\} \in \mathcal{B}(\mathcal{F})$. By Lemma 3.1 (ii) $\mathcal{S}_1 \subset \mathcal{F}$. Note that $\mathcal{F}(\bar{1}) \subset {\binom{[2,n]}{k}}, \mathcal{G}(1) \subset {\binom{[2,n]}{k-1}}$ are cross-intersecting. By Lemma 3.3, we have

$$\begin{aligned} |\mathcal{I}(\mathcal{F}(\bar{1}),\mathcal{G}(1))| &\leq 2\binom{n-2}{k-2} + (2k+1)\binom{n-2}{k-3} + \sum_{0 \leq i \leq k-3} \binom{n-1}{i} \\ &\leq \sum_{0 \leq i \leq k-2} \binom{n-1}{i} + \binom{n-2}{k-2} + (2k+1)\binom{n-2}{k-3}. \end{aligned}$$

Thus,

$$\begin{aligned} |\mathcal{I}(\mathcal{F},\mathcal{G})| &\leq |\mathcal{I}(\mathcal{S}_1,\mathcal{S}_1)| + |\mathcal{I}(\mathcal{F}(\bar{1}),\mathcal{G}(1))| \\ &\leq 2\sum_{0\leq i\leq k-2} \binom{n-1}{i} + \binom{n-2}{k-2} + (2k+1)\binom{n-2}{k-3}. \end{aligned}$$

Now we are in position to prove the main theorem.

Proof of Theorem 1.6. Let $\mathcal{B}_1 = \mathcal{B}(\mathcal{F}), \mathcal{B}_2 = \mathcal{B}(\mathcal{G})$ and let $s_1 = s(\mathcal{B}_1), s_2 = s(\mathcal{B}_2)$. Suppose first that $\min\{s_1, s_2\} = 1$. By symmetry let $s_2 = 1$, then \mathcal{G} is a star. By (15) and $n \ge 2k+3$, we have

$$\begin{aligned} |\mathcal{I}(\mathcal{F},\mathcal{G})| \\ &\leq 2 \sum_{0 \leq i \leq k-2} \binom{n-1}{i} + \binom{n-2}{k-2} + (2k+1)\binom{n-2}{k-3} \\ &\stackrel{(8)}{\leq} 2 \sum_{0 \leq i \leq k-2} \binom{n-1}{i} + \frac{n-2-2}{n-2-2(k-1)}\binom{n-4}{k-2} + \frac{(2k+1)(k-2)}{n-1}\binom{n-1}{k-2} \\ &\stackrel{(8)}{\leq} 2 \sum_{0 \leq i \leq k-2} \binom{n-1}{i} + \frac{n-4}{n-2k}\binom{n-4}{k-2} + \frac{(2k+1)(k-2)}{n-1} \frac{n-1-3}{n-1-3(k-1)}\binom{n-4}{k-2} \\ &\stackrel{(9)}{\leq} \frac{2(n-4)}{n-1-3k} \sum_{0 \leq i \leq k-2} \binom{n-4}{i} + \frac{n-4}{n-2k}\binom{n-4}{k-2} + \frac{(2k+1)(k-2)(n-4)}{(n-1)(n-3k)}\binom{n-2}{k-3}. \end{aligned}$$

Note that $n \ge 63k$ implies

$$\frac{2(n-4)}{n-1-3k} \le \frac{21}{10}, \ \frac{n-4}{n-2k} \le \frac{11}{10}$$

and $n \ge 44k^2$ implies

$$\frac{(2k+1)(k-2)}{n-1} \le \frac{1}{22}, \ \frac{n-4}{n-3k} \le \frac{11}{10}.$$

Thus,

(16)
$$|\mathcal{I}(\mathcal{F},\mathcal{G})| \leq \frac{21}{10} \sum_{0 \leq i \leq k-2} \binom{n-4}{i} + \frac{11}{10} \binom{n-4}{k-2} + \frac{1}{20} \binom{n-4}{k-2} < \frac{13}{4} \sum_{0 \leq i \leq k-2} \binom{n-4}{i} < |\mathcal{I}(\mathcal{A}_1,\mathcal{A}_2)|.$$

Thus, we may assume that $s_1, s_2 \ge 2$. Let us partition \mathcal{F} into $\mathcal{F}^{(s_1)} \cup \ldots \cup \mathcal{F}^{(k)}$ where $F \in \mathcal{F}^{(\ell)}$ if max{ $|B|: B \in \mathcal{B}_1, B \subset F$ } = ℓ . Similarly, partition \mathcal{G} into $\mathcal{G}^{(s_2)} \cup \ldots \cup \mathcal{G}^{(k)}$ where $G \in \mathcal{G}^{(\ell)}$ if $\max\{|B|: B \in \mathcal{B}_2, B \subset G\} = \ell$. Fix an $F \in \mathcal{F}^{(\ell)}$ with $B_1 \subset F, B_1 \in \mathcal{B}_1^{(\ell)}$. For an arbitrary $G \in \mathcal{G}$, we have

$$F \cap G = (B_1 \cap G) \cup ((F \setminus B_1) \cap G)$$

where $B_1 \cap G \neq \emptyset$ and $|(F \setminus B_1) \cap G| \leq |F \setminus B_1| = k - \ell$. It follows that for $s_1 \leq \ell \leq k$

(17)
$$|\mathcal{I}(\mathcal{F}^{(\ell)},\mathcal{G})| \le |\mathcal{B}_1^{(\ell)}| (2^\ell - 1) \sum_{0 \le i \le k-\ell} \binom{n-1}{i}.$$

Similarly, for $s_2 \leq \ell \leq k$

(18)
$$|\mathcal{I}(\mathcal{F}, \mathcal{G}^{(\ell)})| \le |\mathcal{B}_2^{(\ell)}| (2^\ell - 1) \sum_{0 \le i \le k-\ell} \binom{n-1}{i}$$

Let α be the smallest integer such that $\tau(\mathcal{B}_1^{(\leq \alpha)}) \geq 2$ and let β be the smallest integer such that $\tau(\mathcal{B}_2^{(\leq \beta)}) \geq 2$. By symmetry, we may assume that $\alpha \geq \beta$. We distinguish three cases.

Case 1. $\beta \geq 3$. Let $\mathcal{F}' = \mathcal{F}^{(s_1)} \cup \cdots \cup \mathcal{F}^{(\beta-1)}$. Note that \mathcal{F}' and \mathcal{G} are cross-intersecting and \mathcal{F}' is a star. By (15) and (16), we have

(19)

$$|\mathcal{I}(\mathcal{F}',\mathcal{G})| \le 2\sum_{0\le i\le k-2} \binom{n-1}{i} + \binom{n-2}{k-2} + (2k+1)\binom{n-2}{k-3} < \frac{13}{4}\sum_{0\le i\le k-2} \binom{n-4}{i}.$$

Define

$$f(n,k,\ell) = 2^{\ell} \ell^2 k^{\ell-2} \sum_{0 \le i \le k-\ell} \binom{n-1}{i}.$$

and let

$$\lambda_{\ell} = \ell^{-2} k^{-\ell+2} |\mathcal{B}_1^{(\ell)}|.$$

By (17), we see

$$\sum_{\beta \le \ell \le k} |\mathcal{I}(\mathcal{F}^{(\ell)}, \mathcal{G})| = \sum_{\beta \le \ell \le k} \lambda_{\ell} f(n, k, \ell).$$

Since

$$\frac{f(n,k,\ell)}{f(n,k,\ell+1)} = \frac{\ell^2}{2k(\ell+1)^2} \cdot \frac{\sum_{0 \le i \le k-\ell} \binom{n-1}{i}}{\sum_{0 \le i \le k-\ell-1} \binom{n-1}{i}} \stackrel{(9)}{\ge} \frac{(n-1-k)\ell^2}{2k^2(\ell+1)^2} \ge 1 \text{ for } n \ge 5k^2,$$

 $f(n, k, \ell)$ is decreasing as a function of ℓ . Moreover, by (14) we have

$$\sum_{\beta \leq \ell \leq k} \lambda_\ell \leq 1$$

Hence,

(20)
$$\sum_{\beta \le \ell \le k} |\mathcal{I}(\mathcal{F}^{(\ell)}, \mathcal{G})| \le f(n, k, \beta) \le f(n, k, 3) = 72k \sum_{0 \le i \le k-3} \binom{n-1}{i}.$$

Using (9) and (10), for $n \ge 2k + 3$ we have

$$\sum_{\beta \le \ell \le k} |\mathcal{I}(\mathcal{F}^{(\ell)}, \mathcal{G})| \le 72k \sum_{0 \le i \le k-3} \binom{n-1}{i}$$

$$\stackrel{(10)}{\le} \frac{72k^2}{n-1-k} \sum_{0 \le i \le k-2} \binom{n-1}{i}$$

$$\stackrel{(9)}{\le} \frac{72k^2}{n-1-k} \frac{n-1-3}{n-1-3k} \sum_{0 \le i \le k-2} \binom{n-4}{i}$$

Since $n \ge 100k^2 \ge 100k$, we infer

$$\frac{72k^2}{n-1-k} \le \frac{8}{11}$$
 and $\frac{n-1-3}{n-1-3k} \le \frac{33}{32}$

It follows that for $\beta \geq 3$

(21)
$$\sum_{\beta \le \ell \le k} |\mathcal{I}(\mathcal{F}^{(\ell)}, \mathcal{G})| \le \frac{3}{4} \sum_{0 \le i \le k-2} \binom{n-4}{i}$$

Using (19) and (21), we have

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| \leq |\mathcal{I}(\mathcal{F}',\mathcal{G})| + \sum_{\beta \leq \ell \leq k} |\mathcal{I}(\mathcal{F}^{(\ell)},\mathcal{G})| \leq 4 \sum_{0 \leq i \leq k-2} \binom{n-4}{i} < |\mathcal{I}(\mathcal{A}_1,\mathcal{A}_2)|.$$

Case 2. $\beta = 2$ and $\alpha > 2$.

By (21) we have

$$\sum_{3 \le \ell \le k} |\mathcal{I}(\mathcal{F}^{(\ell)}, \mathcal{G})| \le \frac{3}{4} \sum_{0 \le i \le k-2} \binom{n-4}{i}.$$

Since $\alpha > 2$, it follows that $\mathcal{F}^{(2)}$ is a star. By (15) and (16), we have

$$|\mathcal{I}(\mathcal{F}^{(2)},\mathcal{G})| < \frac{13}{4} \sum_{0 \le i \le k-2} \binom{n-4}{i}.$$

Thus,

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| \le |\mathcal{I}(\mathcal{F}^{(2)},\mathcal{G})| + \sum_{3 \le \ell \le k} |\mathcal{I}(\mathcal{F}^{(\ell)},\mathcal{G})| \le 4 \sum_{0 \le i \le k-2} \binom{n-4}{i} < |\mathcal{I}(\mathcal{A}_1,\mathcal{A}_2)|.$$

Case 3. $\beta = \alpha = 2$.

Case 3. $\beta = \alpha = 2$. Since $\mathcal{B}_1^{(2)}, \mathcal{B}_2^{(2)}$ are cross-intersecting, we see that $\nu(\mathcal{B}_1^{(2)}) \leq 2$ and $\nu(\mathcal{B}_2^{(2)}) \leq 2$. More-over, $\beta = \alpha = 2$ implies $\tau(\mathcal{B}_1^{(2)}) \geq 2$ and $\tau(\mathcal{B}_2^{(2)}) \geq 2$. It follows that $\mathcal{B}_1^{(2)}, \mathcal{B}_2^{(2)}$ are either both triangles or both subgraphs of K_4 with a matching of size two. **Case 3.1.** $\mathcal{B}_1^{(2)}, \mathcal{B}_2^{(2)}$ are both triangles.

Without loss of generality, assume that $\mathcal{B}_{1}^{(2)} = \mathcal{B}_{2}^{(2)} = \{(1,2), (1,3), (2,3)\}$. By saturatedness, we have

$$\mathcal{F} = \mathcal{G} = \mathcal{A}_3 = \left\{ A \in \binom{[n]}{k} : |A \cap \{1, 2, 3\}| \ge 2 \right\}.$$

Therefore,

$$\begin{aligned} |\mathcal{I}(\mathcal{F},\mathcal{G})| &= |\mathcal{I}(\mathcal{A}_{3},\mathcal{A}_{3})| \\ &= 3 \sum_{0 \le i \le k-2} \binom{n-3}{i} + 3 \sum_{0 \le i \le k-3} \binom{n-3}{i} + \sum_{0 \le i \le k-4} \binom{n-3}{i} \\ &\stackrel{(9)}{=} \frac{n-4}{n-3-k} \left(3 \sum_{0 \le i \le k-2} \binom{n-4}{i} + 3 \sum_{0 \le i \le k-3} \binom{n-4}{i} + \sum_{0 \le i \le k-4} \binom{n-4}{i} \right). \end{aligned}$$

Since $n \ge 13k$ implies $\frac{n-4}{n-3-k} \le \frac{13}{12}$, we obtain that

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| < \frac{13}{4} \sum_{0 \le i \le k-2} \binom{n-4}{i} + \frac{13}{4} \sum_{0 \le i \le k-3} \binom{n-4}{i} + 2 \sum_{0 \le i \le k-4} \binom{n-4}{i} < |\mathcal{I}(\mathcal{A}_1,\mathcal{A}_2)| \le 1$$

Case 3.2. $\mathcal{B}_1^{(2)}, \mathcal{B}_2^{(2)}$ are both subgraphs of K_4 with a matching of size two. By symmetry, we may assume that $(1,3), (2,4) \in \mathcal{B}_1^{(2)}$ and $(1,2), (3,4) \in \mathcal{B}_2^{(2)}$. We

further assume that $|\mathcal{B}_{1}^{(2)}| \geq |\mathcal{B}_{2}^{(2)}|$. **Case 3.2.1.** $\mathcal{B}_{1}^{(2)} = \{(1,3), (2,4), (1,4), (2,3)\}$ and $\mathcal{B}_{2}^{(2)} = \{(1,2), (3,4)\}$. Since \mathcal{F}, \mathcal{G} are saturated, we have $\mathcal{F} = \mathcal{F}^{(2)}$ and $\mathcal{G} = \mathcal{G}^{(2)}$. Thus,

$$\begin{aligned} |\mathcal{I}(\mathcal{F},\mathcal{G})| &= 4\sum_{0 \le i \le k-2} \binom{n-4}{i} + 6\sum_{0 \le i \le k-3} \binom{n-4}{i} + 4\sum_{0 \le i \le k-4} \binom{n-4}{i} \\ &+ \sum_{0 \le i \le k-5} \binom{n-4}{i} \end{aligned}$$

 $< |\mathcal{I}(\mathcal{A}_1, \mathcal{A}_2)|.$

Case 3.2.2. $\mathcal{B}_1^{(2)} = \{(1,3), (2,4), (1,4)\}$ and $\mathcal{B}_2^{(2)} = \{(1,2), (3,4), (1,4)\}.$ Since \mathcal{F}, \mathcal{G} are saturated, we have $\mathcal{F} = \mathcal{F}^{(2)}$ and $\mathcal{G} = \mathcal{G}^{(2)}$. Thus,

$$\begin{aligned} |\mathcal{I}(\mathcal{F},\mathcal{G})| &= 4 \sum_{0 \le i \le k-2} \binom{n-4}{i} + 6 \sum_{0 \le i \le k-3} \binom{n-4}{i} + 4 \sum_{0 \le i \le k-4} \binom{n-4}{i} \\ &+ \sum_{0 \le i \le k-5} \binom{n-4}{i} \end{aligned}$$

 $< |\mathcal{I}(\mathcal{A}_1, \mathcal{A}_2)|.$

Case 3.2.3. $\mathcal{B}_{1}^{(2)} = \{(1,3), (2,4)\}$ and $\mathcal{B}_{2}^{(2)} = \{(1,2), (3,4)\}.$ By Lemma 3.1, we have $\mathcal{S}_{13} \cup \mathcal{S}_{24} \subset \mathcal{F}$ and $\mathcal{S}_{12} \cup \mathcal{S}_{34} \subset \mathcal{G}$. Let $\mathcal{F}' = \mathcal{F} \setminus (\mathcal{S}_{13} \cup \mathcal{S}_{24})$ and $\mathcal{G}' = \mathcal{G} \setminus (\mathcal{S}_{12} \cup \mathcal{S}_{34}).$ Since $\mathcal{B}_1^{(2)}, \mathcal{G}'$ are cross-intersecting, $G \cap \{1,3\} \neq \emptyset$ and $G \cap \{2,4\} \neq \emptyset$ for all $G \in \mathcal{G}'$. Moreover, $G \notin \mathcal{S}_{12} \cup \mathcal{S}_{34}$. It follows that $G \cap [4] = \{1, 4\}$ or $G \cap [4] = \{2, 3\}$ for all $G \in \mathcal{G}'$. Similarly, $F \cap [4] = \{1, 4\}$ or $F \cap [4] = \{2, 3\}$ for all $F \in \mathcal{F}'$. Let

$$\mathcal{F}'_{14} = \{F \colon F \in \mathcal{F}', F \cap [4] = \{1,4\}\}, \ \mathcal{F}'_{23} = \{F \colon F \in \mathcal{F}', F \cap [4] = \{2,3\}\}$$

and

$$\mathcal{G}'_{14} = \{ G \colon G \in \mathcal{G}', G \cap [4] = \{1, 4\} \}, \ \mathcal{G}'_{23} = \{ G \colon G \in \mathcal{G}', G \cap [4] = \{2, 3\} \}.$$

Since $\mathcal{F}'_{14}, \mathcal{G}'_{23}$ are cross-intersecting and $\mathcal{F}'_{23}, \mathcal{G}'_{14}$ are cross-intersecting, by (13) we have

$$|(\mathcal{F}'_{14} \land \mathcal{G}'_{23}) \cup (\mathcal{F}'_{23} \land \mathcal{G}'_{14})| \le 2\sum_{0 \le i \le k-3} \binom{n-6}{i} + \sum_{0 \le i \le k-4} \binom{n-6}{i}$$

Note that $\mathcal{I}(\mathcal{F}'_{14}, \mathcal{G} \setminus \mathcal{G}'_{23}) \subset \mathcal{I}(\mathcal{S}_{13} \cup \mathcal{S}_{24}, \mathcal{S}_{12} \cup \mathcal{S}_{34})$ and $\mathcal{I}(\mathcal{F}'_{23}, \mathcal{G} \setminus \mathcal{G}'_{14}) \subset \mathcal{I}(\mathcal{S}_{13} \cup \mathcal{S}_{24}, \mathcal{S}_{12} \cup \mathcal{S}_{24})$ \mathcal{S}_{34}). Thus,

$$\begin{aligned} |\mathcal{I}(\mathcal{F},\mathcal{G})| &= |\mathcal{I}(\mathcal{S}_{13} \cup \mathcal{S}_{24}, \mathcal{S}_{12} \cup \mathcal{S}_{34})| + |(\mathcal{F}'_{14} \wedge \mathcal{G}'_{23}) \cup (\mathcal{F}'_{23} \wedge \mathcal{G}'_{14})| \\ &\leq 4 \sum_{0 \leq i \leq k-2} \binom{n-4}{i} + 6 \sum_{0 \leq i \leq k-3} \binom{n-4}{i} + 4 \sum_{0 \leq i \leq k-4} \binom{n-4}{i} \\ &+ \sum_{0 \leq i \leq k-5} \binom{n-4}{i} + 2 \sum_{0 \leq i \leq k-3} \binom{n-6}{i} + \sum_{0 \leq i \leq k-4} \binom{n-6}{i} \\ &= |\mathcal{I}(\mathcal{A}_1, \mathcal{A}_2)|. \end{aligned}$$

Case 3.2.4. $\mathcal{B}_1^{(2)} = \{(1,3), (2,4), (1,4)\}$ and $\mathcal{B}_2^{(2)} = \{(1,2), (3,4)\}.$

By Lemma 3.1 (ii), we have $S_{13} \cup S_{24} \cup S_{14} \subset \mathcal{F}$ and $S_{12} \cup S_{34} \subset \mathcal{G}$. Let $\mathcal{F}' = \mathcal{F} \setminus (S_{13} \cup S_{24} \cup S_{14})$ and $\mathcal{G}' = \mathcal{G} \setminus (S_{12} \cup S_{34})$. Since $\mathcal{B}_1^{(2)}, \mathcal{G}'$ are cross-intersecting, $G \cap [4] = \{1, 4\}$ for all $G \in \mathcal{G}'$. Similarly, $F \cap [4] = \{2, 3\}$ for all $F \in \mathcal{F}'$. Since $\mathcal{F}', \mathcal{G}'$ are cross-intersecting, by (3) we have

$$|\mathcal{F}'_{23} \wedge \mathcal{G}'_{14}| \le \sum_{0 \le i \le k-3} \binom{n-5}{i} < 2\sum_{0 \le i \le k-3} \binom{n-6}{i} + \sum_{0 \le i \le k-4} \binom{n-6}{i}.$$

Note that $\mathcal{I}(\mathcal{F}'_{23}, \mathcal{G} \setminus \mathcal{G}'_{14}) \subset \mathcal{I}(\mathcal{S}_{13} \cup \mathcal{S}_{24}, \mathcal{S}_{12} \cup \mathcal{S}_{34})$. Thus,

$$|\mathcal{I}(\mathcal{F},\mathcal{G})| = |\mathcal{I}(\mathcal{S}_{13} \cup \mathcal{S}_{24}, \mathcal{S}_{12} \cup \mathcal{S}_{34})| + |\mathcal{F}'_{23} \wedge \mathcal{G}'_{14}| \le |\mathcal{I}(\mathcal{A}_1, \mathcal{A}_2)|.$$

4 Distinct intersections in a *t*-intersecting family

In this section, we determine the maximum number of distinct intersections in a t-intersecting family.

Since $\mathcal{F} \subset \tilde{\mathcal{F}}$ implies $\mathcal{I}(\mathcal{F}) \subset \mathcal{I}(\tilde{\mathcal{F}})$, we may always assume that \mathcal{F} is saturated. Let $\mathcal{B} = \mathcal{B}_t(\mathcal{F})$ be the family of minimal (for containment) sets in $\mathcal{T}_t(\mathcal{F})$.

Lemma 4.1. Suppose that $\mathcal{F} \subset {\binom{[n]}{k}}$ is a saturated t-intersecting family. Then (i) and (ii) hold.

(i) \mathcal{B} is a t-intersecting antichain,

(*ii*)
$$\mathcal{F} = \left\{ H \in {[n] \choose k} : \exists B \in \mathcal{B}, B \subset H \right\}.$$

Proof. (i) Clearly, \mathcal{B} is an anti-chain. Suppose for contradiction that $B, B' \in \mathcal{B}$ but $|B \cap B'| < t$. If |B| = |B'| = k, then $B, B' \in \mathcal{F}$ as \mathcal{F} is saturated, a contradiction. If |B'| < k, then there exists $F' \supset B'$ such that |F'| = k and $|F' \cap B| = |B' \cap B| < t$. By definition $F' \in \mathcal{T}_t(\mathcal{F})$. Since \mathcal{F} is saturated, we see that $F' \in \mathcal{F}$. But this contradicts the assumption that B is a t-transversal. Since \mathcal{F} is saturated, (ii) is immediate from the definition of \mathcal{B} .

Let $r(\mathcal{B}) = \max\{|B|: B \in \mathcal{B}\}$ and $s(\mathcal{B}) = \min\{|B|: B \in \mathcal{B}\}$. For any ℓ with $s(\mathcal{B}) \leq \ell \leq r(\mathcal{B})$ define

$$\mathcal{B}^{(\ell)} = \{B \in \mathcal{B} \colon |B| = \ell\} \text{ and } \mathcal{B}^{(\leq \ell)} = \bigcup_{i=s(\mathcal{B})}^{\ell} \mathcal{B}^{(i)}$$

It is easy to see that $s(\mathcal{B}_t(\mathcal{F})) = \tau_t(\mathcal{F}).$

Lemma 4.2. Suppose that $\mathcal{F} \subset {\binom{[n]}{k}}$ is a saturated t-intersecting family and $\mathcal{B} = \mathcal{B}_t(\mathcal{F})$. If $s(\mathcal{B}) \ge t+1$ and $\tau_t(\mathcal{B}^{(\le r)}) \ge t+1$, then

(22)
$$\sum_{r \le \ell \le k} \left(\binom{\ell}{t} \ell k^{\ell-t-1} \right)^{-1} |\mathcal{B}^{(\ell)}| \le 1.$$

Proof. For the proof we use a branching process. During the proof a sequence $S = (x_1, x_2, \ldots, x_\ell)$ is an ordered sequence of distinct elements of [n] and we use \widehat{S} to denote the underlying unordered set $\{x_1, x_2, \ldots, x_\ell\}$. At the beginning, we assign weight 1 to the empty sequence S_{\emptyset} . At the first stage, we choose $B_1 \in \mathcal{B}_t$ with $|B_1| = s(\mathcal{B}) \ge t + 1$. For any t-subset $\{x_1, \ldots, x_t\} \subset B_1$, define one sequence (x_1, \ldots, x_t) and assign the weight $\binom{s(\mathcal{B})}{t}^{-1}$ to it.

At the second stage, since $\tau_t(\mathcal{B}^{(\leq r)}) \geq t+1$, for each t-sequence $S = (x_1, \ldots, x_t)$ we may choose $B \in \mathcal{B}^{(\leq r)}$ such that $|\widehat{S} \cap B| < t$. Then we replace $S = (x_1, \ldots, x_t)$ by $|B \setminus \widehat{S}|$ (t+1)-sequences of the form (x_1, \ldots, x_t, y) with $y \in B \setminus \widehat{S}$ and weight $\frac{w(S)}{|B \setminus \widehat{S}|}$.

In each subsequent stage, we pick a sequence $S = (x_1, \ldots, x_p)$ and denote its weight by w(S). If $|\widehat{S} \cap B| \ge t$ for all $B \in \mathcal{B}$ then we do nothing. Otherwise we pick $B \in \mathcal{B}$ satisfying $|\widehat{S} \cap B| < t$ and replace S by the $|B \setminus \widehat{S}|$ sequences (x_1, \ldots, x_p, y) with $y \in B \setminus \widehat{S}$ and assign weight $\frac{w(S)}{|B \setminus \widehat{S}|}$ to each of them. clearly, the total weight is always 1.

We continue until $|S \cap B| \ge t$ for all sequences and all $B \in \mathcal{B}$. Since [n] is finite, each sequence has length at most n and eventually the process stops. Let S be the collection of sequences that survived in the end of the branching process and let $S^{(\ell)}$ be the collection of sequences in S with length ℓ .

Claim 2. To each $B \in \mathcal{B}^{(\ell)}$ with $\ell \geq r$ there is some sequence $S \in \mathcal{S}^{(\ell)}$ with $\widehat{S} = B$.

Proof. Let us suppose the contrary and let $S = (x_1, \ldots, x_p)$ be a sequence of maximal length that occurred at some stage of the branching process satisfying $\widehat{S} \subseteq B$. Since \mathcal{B} is t-intersecting, $|B \cap B_1| \ge t$, implying that $p \ge t$. Since \widehat{S} is a proper subset of B, there exists $F \in \mathcal{F}$ with $|\widehat{S} \cap F| < t$. In view of Lemma 4.1 (ii) we can find $B' \in \mathcal{B}$ such that $|\widehat{S} \cap B'| < t$. Thus at some point we picked S and some $\widetilde{B} \in \mathcal{B}$ with $|\widehat{S} \cap \widetilde{B}| < t$. Since \mathcal{B} is t-intersecting, $|B \cap \widetilde{B}| \ge t$. Consequently, for each $y \in B \cap (\widetilde{B} \setminus \widehat{S})$ the sequence (x_1, \ldots, x_p, y) occurred in the branching process. This contradicts the maximality of p. Hence there is an S at some stage satisfying $\widehat{S} = B$. Since \mathcal{B} is t-intersecting, $|\widehat{S} \cap B'| \ge t$ for all $B' \in \mathcal{B}$. Thus $\widehat{S} \in S$ and the claim holds.

By Claim 2, we see that $|\mathcal{B}^{(\ell)}| \leq |\mathcal{S}^{(\ell)}|$. Let $S = (x_1, \ldots, x_\ell) \in \mathcal{S}^{(\ell)}$ and let $S_i = (x_1, \ldots, x_i)$ for $i = 1, \ldots, \ell$. At the first stage, $w(S_t) = 1/\binom{s(\mathcal{B})}{t}$. Assume that B_i is the selected set when replacing S_{i-1} in the branching process for $i = t + 1, \ldots, \ell$. Clearly, $x_i \in B_i$, $B_{t+1} \in \mathcal{B}^{(\leq r)}$ and

$$w(S) = \frac{1}{\binom{s(\mathcal{B})}{t}} \prod_{i=t+1}^{\ell} \frac{1}{|B_i \setminus \widehat{S_{i-1}}|}$$

Note that $s(\mathcal{B}) \leq r \leq \ell$, $|B_{t+1} \setminus \widehat{S}_t| \leq \ell$ and $|B_i \setminus \widehat{S_{i-1}}| \leq k$ for $i \geq t+2$. It follows that

$$w(S) \ge \left(\binom{\ell}{t}\ell k^{\ell-t-1}\right)^{-1}$$

Thus,

$$\sum_{r \le \ell \le k} \left(\binom{\ell}{t} \ell k^{\ell-t-1} \right)^{-1} |\mathcal{B}^{(\ell)}| \le \sum_{r \le \ell \le k} \sum_{S \in \mathcal{S}^{(\ell)}} w(S) \sum_{S \in \mathcal{S}} w(S) = 1.$$

Lemma 4.3. Suppose that $\tau_t(\mathcal{B}^{(t+1)} \ge t+1)$. Then $\mathcal{F} = \mathcal{A}(n, k, t)$.

Proof. Choose $B_1, B_2 \in \mathcal{B}^{(t+1)}$ and assume by symmetry that $B_i = [t] \cup \{t+i\}$ for i = 1, 2. Since $\tau_t(\mathcal{B}^{(t+1)}) \ge t+1$, we may choose $B_3 \in \mathcal{B}^{(t+1)}$ satisfying $[t] \nsubseteq B_3$. Now $|B_2 \cap B_i| \ge t$ implies $\{t+1, t+2\} \subset B_3$. Using $|B_3| = t+1$, by symmetry we may assume that $B_3 = [t+2] \setminus \{t\}$. Now take an arbitrary $F \in \mathcal{F}$. It is clear that $|F \cap B_i| \ge t$ can only hold for all $1 \le i \le 3$ if $|F \cap [t+2]| \ge t+1$. That is $\mathcal{F} \subset \mathcal{A}(n,k,t)$. Since \mathcal{F} is saturated, $\mathcal{F} = \mathcal{A}(n,k,t)$.

Proof of Theorem 1.8. By (5) and (9), we have

$$\begin{aligned} |\mathcal{I}(\mathcal{S}_{[t]})| &= \sum_{0 \le i \le k-t-1} \binom{n-t}{i} \\ \stackrel{(9)}{\le} \frac{n-t-2}{n-t-2k} \sum_{0 \le i \le k-t-1} \binom{n-t-2}{i} \\ &< \binom{t+2}{t} \sum_{0 \le i \le k-t-1} \binom{n-t-2}{i} \\ &< |\mathcal{I}(\mathcal{A}(n,k,t))|. \end{aligned}$$

Thus, we may assume that $s = s(\mathcal{B}) \ge t + 1$. Let us partition \mathcal{F} into $\mathcal{F}^{(s)} \cup \ldots \cup \mathcal{F}^{(k)}$ where $F \in \mathcal{F}^{(\ell)}$ if $\max\{|B|: B \in \mathcal{B}, B \subset F\} = \ell$. Set

$$\mathcal{I}_{\ell} = \left\{ F \cap F' \colon F \in \mathcal{F}^{(\ell)}, F' \in \mathcal{F}^{(s)} \cup \ldots \cup \mathcal{F}^{(\ell)} \right\}.$$

Then

$$|\mathcal{I}(\mathcal{F})| \le \sum_{s \le \ell \le k} |\mathcal{I}_{\ell}|$$

The point is that for $F \in \mathcal{F}^{(\ell)}$ and $B \subset F, B \in \mathcal{B}^{\ell}$ for an arbitrary $F' \in \mathcal{F}$,

$$F \cap F' = (B \cap F') \cup ((F \setminus B) \cap F').$$

Note that $|B \cap F'| \ge t$ and $|(F \setminus B) \cap F'| \le |F \setminus B| = k - \ell$. It follows that for $s \le \ell \le k$

(23)
$$|\mathcal{I}_{\ell}| \leq \left(\sum_{t \leq j \leq \ell} \binom{\ell}{j}\right) |\mathcal{B}^{(\ell)}| \sum_{0 \leq i \leq k-\ell} \binom{n-t}{i}.$$

Let α be the smallest integer such that $\tau_t(\mathcal{B}^{(\leq \alpha)}) \geq t + 1$. The family $\mathcal{F}' = \bigcup_{i=l}^{\alpha-1} \mathcal{F}^{(i)}$ is a trivial *t*-intersecting family. By (9), we have for $n \geq 5k$

(24)
$$\left| \bigcup_{i=s}^{\alpha-1} \mathcal{I}_i \right| \leq |\mathcal{I}(\mathcal{S}_{[t]})| = \sum_{0 \leq i \leq k-t-1} \binom{n-t}{i}$$
$$\stackrel{(9)}{\leq} \frac{n-t-2}{n-t-2k} \sum_{0 \leq i \leq k-t-1} \binom{n-t-2}{i}$$
$$\leq 2 \sum_{0 \leq i \leq k-t-1} \binom{n-t-2}{i}.$$

If $\alpha = s = t + 1$, then $\mathcal{B}^{(t+1)}$ is a *t*-intersecting (t+1)-uniform family with *t*-covering number t+1. By Lemma 4.3, $\mathcal{F} = \mathcal{A}(n,k,t)$ and there is nothing to prove. Thus we may assume that $\alpha \geq t+2$.

Define

$$f(n,k,\ell) = \left(\sum_{t \le j \le \ell} \binom{\ell}{j}\right) \binom{\ell}{t} \ell k^{\ell-t-1} \sum_{0 \le i \le k-\ell} \binom{n-t}{i}$$

and let

$$\lambda_{\ell} = \left(\binom{\ell}{t} \ell k^{\ell-t-1} \right)^{-1} |\mathcal{B}^{(\ell)}|.$$

Then by (23)

(25)
$$\sum_{\alpha \le \ell \le k} |\mathcal{I}_{\ell}| = \sum_{\alpha \le \ell \le k} \lambda_{\ell} \cdot f(n, k, \ell).$$

By (10) and (11), we have

$$\frac{f(n,k,\ell)}{f(n,k,\ell+1)} = \frac{\sum_{t \le j \le \ell} {\ell \choose j}}{\sum_{t \le j \le \ell+1} {\ell+1 \choose j}} \cdot \frac{{\ell \choose t} \ell k^{\ell-t-1}}{{\ell+1 \choose t} (\ell+1)k^{\ell-t}} \cdot \frac{\sum_{0 \le i \le k-\ell} {n-t \choose i}}{\sum_{0 \le i \le k-\ell-1} {n-t \choose i}}$$
$$\ge \frac{1}{2(t+1)} \cdot \frac{(\ell+1-t)\ell}{(\ell+1)^2k} \cdot \frac{n-t-k}{k}.$$

By $\ell \ge t+1 \ge 3$, we have

$$\frac{\ell+1-t}{\ell+1} \cdot \frac{\ell}{\ell+1} \ge \frac{2}{t+2} \cdot \frac{3}{4} \ge \frac{3}{2(t+2)}.$$

Then by $n \ge \frac{4}{3}(t+2)^2k^2$

$$\frac{f(n,k,\ell)}{f(n,k,\ell+1)} \geq \frac{3(n-t-k)}{4(t+1)(t+2)k^2} \geq 1.$$

Hence $f(n, k, \ell)$ is decreasing as a function of ℓ . Moreover, (22) implies $\sum_{\alpha \le \ell \le k} \lambda_{\ell} \le 1$. From (25) we see

$$\sum_{\alpha \leq \ell \leq k} |\mathcal{I}_{\ell}| \leq f(n,k,\alpha) \leq f(n,k,t+2).$$

Therefore,

$$\sum_{\alpha \le \ell \le k} |\mathcal{I}_{\ell}| \le \left(\binom{t+2}{t} + \binom{t+2}{t+1} + \binom{t+2}{t+2} \right) \binom{t+2}{t} (t+2)k \sum_{0 \le i \le k-t-2} \binom{n-t}{i}$$

$$\stackrel{(10)}{\le} \frac{(t+2)^2(t+1)(t^2+5t+8)k}{4} \cdot \frac{k}{n-t-k} \sum_{0 \le i \le k-t-1} \binom{n-t}{i}$$

$$\stackrel{(9)}{\le} \frac{(t+2)^2(t+1)(t+2)(t+4)k^2(n-t-2)}{4(n-t-k)(n-t-2k)} \sum_{0 \le i \le k-t-1} \binom{n-t-2}{i}.$$

Note that $n \ge 5k$ implies

$$\frac{n-t-2}{n-t-2k} \le 2$$

and $n \ge 3(t+2)^3 k^2$, $t \ge 2$ imply

$$\frac{(t+2)^3(t+1)(t+4)k^2}{4(n-t-k)} \le \frac{1}{2} \left(\binom{t+2}{2} - 2 \right).$$

It follows that

(26)
$$\sum_{\alpha \le \ell \le k} |\mathcal{I}_{\ell}| \le \left(\binom{t+2}{2} - 2 \right) \sum_{0 \le i \le k-t-1} \binom{n-t-2}{i}.$$

By (24) and (26), we obtain that

$$|\mathcal{I}(\mathcal{F})| \le \left| \bigcup_{i=s}^{\alpha-1} \mathcal{I}_i \right| + \sum_{\alpha \le \ell \le k} |\mathcal{I}_\ell| < |\mathcal{I}(\mathcal{A}(n,k,t))|$$

concluding the proof of the theorem.

5 Further problems and results

In their seminal paper [1] Erdős, Ko and Rado actually proved their main result for antichains. Namely, instead of considering k-graphs $\mathcal{F} \subset {[n] \choose k}$ they suppose that \mathcal{F} is an antichain of rank k, that is $|F| \leq k$ for all $F \in \mathcal{F}$. The reason that this tendency has all but disappeared from recent research is that a t-intersecting antichain \mathcal{F} of rank k which is not k-uniform can always be replaced by a t-intersecting family $\tilde{\mathcal{F}} \subset {[n] \choose k}$ with $|\tilde{\mathcal{F}}| > |\mathcal{F}|$. The way to do is to apply an operation on antichains discovered already by Sperner [8].

For a family $\mathcal{A} \subset {[n] \choose a}$ define its shade $\sigma^+(\mathcal{A})$ by

$$\sigma^+(\mathcal{A}) = \left\{ B \in \binom{[n]}{a+1} : \exists A \in \mathcal{A}, A \subset B \right\}$$

Sperner [8] proved that for a < n/2, $|\sigma^+(\mathcal{A})| \ge |\mathcal{A}|$ with strict inequality unless $a = \frac{n-1}{2}$ and $\mathcal{A} = \binom{[n]}{\frac{n-1}{2}}$. Let $\mathcal{F} \subset 2^{[n]}$ be a *t*-intersecting antichain of rank $k, n \ge 2k - t$. Suppose that $a = \min\{|F|: F \in \mathcal{F}\}$ and a < k. Define

$$\mathcal{F}^{(a)} = \{F \in \mathcal{F} : |\mathcal{F}| = a\} \text{ and } \tilde{\mathcal{F}} = (F \setminus \mathcal{F}^{(a)}) \cup \sigma^+(\mathcal{F}^{(a)})$$

Then not only is $\tilde{\mathcal{F}}$ a *t*-intersecting antichain of rank k with $|\tilde{\mathcal{F}}| > |\mathcal{F}|$ but $\mathcal{I}(\tilde{\mathcal{F}}) \supset \mathcal{I}(\mathcal{F})$ can be checked easily as well. This shows that it was reasonable to restrict our attention to k-uniform families.

However there is a related, very natural problem.

Problem 5.1. Determine or estimate $\max |\mathcal{I}(\mathcal{A})|$ over all antichain $\mathcal{A} \subset 2^{[n]}$.

Example 5.2. Let $\ell \leq \frac{n}{2}$ and define $\mathcal{A} = {\binom{[n]}{n-\ell}}$. Clearly,

$$\mathcal{I}(\mathcal{A}) = \{ B \subset [n] \colon n - 2\ell \le |B| < n - \ell \}.$$

Choosing $\ell = \lfloor n/3 \rfloor$, we have

$$|\mathcal{I}(\mathcal{A})| = 2^n - \sum_{0 \le i \le \lfloor \frac{n}{3} \rfloor} {\binom{n}{i}} - \sum_{0 \le j \le n-2 \lfloor \frac{n}{3} \rfloor} {\binom{n}{j}}.$$

Proposition 5.3. If $\mathcal{A} \subset 2^{[n]}$ is an antichain, then $|\mathcal{I}(\mathcal{A})| < 2^n - \sqrt{2}^n$.

Proof. Note that

$$|\mathcal{I}(\mathcal{A})| \le \binom{|\mathcal{A}|}{2}.$$

Consequently, if $|\mathcal{A}| \leq \sqrt{2}^n$ then $|\mathcal{I}(\mathcal{A})| < 2^n/2 < 2^n - \sqrt{2}^n$. Thus we can assume $|\mathcal{A}| > \sqrt{2}^n$. Since $\mathcal{A} \cap \mathcal{I}(\mathcal{A}) = \emptyset$, we have

$$|\mathcal{I}(\mathcal{A})| \le 2^n - |\mathcal{A}| < 2^n - \sqrt{2}^n.$$

Two families \mathcal{A}, \mathcal{B} are called *cross-Sperner* if $A \not\subset B$ and $B \not\subset A$ hold for all $A \in \mathcal{A}$, $B \in \mathcal{B}$. Set

$$\mathcal{I}(\mathcal{A},\mathcal{B}) = \{A \cap B \colon A \in \mathcal{A}, B \in \mathcal{B}\}.$$

Define

$$m(n) = \max\{|\mathcal{I}(\mathcal{A}, \mathcal{B})| : \mathcal{A}, \mathcal{B} \subset 2^{[n]} \text{ are cross-Sperner}\}$$

Example 5.4. Let $[n] = X \cup Y$ be a partition. Define

$$\mathcal{A} = \{ A \cup Y \colon A \subsetneq X \}, \ \mathcal{B} = \{ X \cup B \colon B \subsetneq Y \}.$$

Then

$$\mathcal{I}(\mathcal{A},\mathcal{B}) = \{A \cup B \colon A \subsetneq X, B \subsetneq Y\}$$

and

$$|\mathcal{I}(\mathcal{A}, \mathcal{B})| = 2^n - 2^{|X|} - 2^{|Y|} + 1.$$

Theorem 5.5. $m(n) = 2^n - 2 \cdot 2^{n/2} + 1$ holds for n = 2d even.

Proof. The lower bound comes from the example with $|X| = \lfloor n/2 \rfloor$, $|Y| = \lceil n/2 \rceil$. Note that for $A, A' \in \mathcal{A}, B, B' \in \mathcal{B}$ the cross-Sperner property implies $A \not\subset A' \cap B', B \not\subset A' \cap B'$. In particular,

$$\mathcal{A} \cap \mathcal{I}(\mathcal{A}, \mathcal{B}) = \emptyset = \mathcal{B} \cap \mathcal{I}(\mathcal{A}, \mathcal{B}).$$

Cross-Sperner property implies $\mathcal{A} \cap \mathcal{B} = \emptyset$ and $[n] \notin \mathcal{A} \cup \mathcal{B} \cup \mathcal{I}(\mathcal{A}, \mathcal{B})$. Thus

$$|\mathcal{A}| + |\mathcal{B}| + |\mathcal{I}(\mathcal{A}, \mathcal{B})| \le 2^n - 1$$

or equivalently

(27)
$$|\mathcal{I}(\mathcal{A},\mathcal{B})| \le 2^n - |\mathcal{A}| - |\mathcal{B}| - 1.$$

Obviously,

(28)
$$|\mathcal{I}(\mathcal{A},\mathcal{B})| \le |\mathcal{A}| \cdot |\mathcal{B}|.$$

Suppose that n = 2d (even). If $|A| + |B| \ge 2(2^d - 1)$, then (27) implies

$$|\mathcal{I}(\mathcal{A},\mathcal{B})| \le 2^n - 2 \cdot 2^d + 1.$$

If $\frac{|A|+|B|}{2} \le 2^d - 1$ then the inequality between arithmetic and geometric mean yields via (28):

$$|\mathcal{I}(\mathcal{A},\mathcal{B})| \le \left(2^d - 1\right)^2 = 2^n - 2 \cdot 2^d + 1.$$

However the proof only gives $\mathcal{I}(\mathcal{A}, \mathcal{B}) \leq 2^n - 2 \cdot 2^{n/2} + 1$ for n = 2d + 1.

Problem 5.6. For n = 2d + 1, does $m(n) = 2^n - 2^{d+1} - 2^d + 1$ hold?

References

- P. Erdős, C. Ko, R. Rado, Intersection theorems for systems of finite sets, Quart. J. Math. Oxford Ser. 12 (1961), 313–320.
- [2] P. Frankl, The Erdős-Ko-Rado theorem is true for n = ckt, Coll. Math. Soc. J. Bolyai 18 (1978), 365–375.
- [3] P. Frankl, The shifting technique in extremal set theory, Surveys in Combinatorics 123 (1987), 81–110.
- [4] P. Frankl, Improved bounds for Erdős' matching conjecture, J. Combin. Theory, Ser. A 120 (2013), 1068–1072.
- [5] P. Frankl, A. Kupavskii, S. Kiselev, On the maximum number of distinct intersections in an intersecting family, Discrete Math. 345 (2022), 112757. https://doi.org/10.1016/j.disc.2021.112757.
- [6] A.J.W. Hilton, E.C. Milner, Some intersection theorems for systems of finite sets, Q. J. Math. 18 (1) (1967), 369–384.
- [7] L. Pyber, A new generalization of the Erdős-Ko-Rado theorem, J. Combin. Theory Ser. A 43 (1986), 85–90.
- [8] E. Sperner, Ein Satz über Untermengen einer endlichen Menger, Math. Zeitschrift 27 (1928), 544–548.