Proportional Allocation of Indivisible Goods up to the Least Valued Good on Average^{*}

Yusuke Kobayashi[†] Ryoga Mahara[‡]

Abstract

We study the problem of fairly allocating a set of indivisible goods to multiple agents and focus on the proportionality, which is one of the classical fairness notions. Since proportional allocations do not always exist when goods are indivisible, approximate concepts of proportionality have been considered in the previous work. Among them, proportionality up to the maximin good (PROPm) has been the best approximate notion of proportionality that can be achieved for all instances [5]. In this paper, we introduce the notion of *proportionality up to the least valued good on average* (PROPavg), which is a stronger notion than PROPm, and show that a PROPavg allocation always exists for all instances and can be computed in polynomial time. Our results establish PROPavg as a notable non-trivial fairness notion that can be achieved for all instances. Our proof is constructive, and based on a new technique that generalizes the cut-and-choose protocol and uses a recursive technique.

^{*}An earlier version of this work was presented in ISAAC 2022 [21].

[†]Research Institute for Mathematical Sciences, Kyoto University, Japan. E-mail: yusuke@kurims.kyotou.ac.jp

[‡]Department of Mathematical Informatics, University of Tokyo. E-mail: mahara@mist.i.u-tokyo.ac.jp

1 Introduction

1.1 Proportional Allocation of Indivisible Goods

We study the problem of fairly allocating a set of indivisible goods to multiple agents under additive valuations. Fair division of indivisible goods is a fundamental and well-studied problem in Economics and Computer Science. We are given a set M of m indivisible goods and a set N of n agents with individual valuations. Under additive valuations, each agent $i \in N$ has a value $v_i(\{g\}) \ge 0$ for each good g and her value for a bundle S of goods is equal to the sum of the values of each good $g \in S$, i.e., $v_i(S) = \sum_{g \in S} v_i(\{g\})$. An indivisible good can not be split among multiple agents and this causes finding a fair division to be a difficult task.

One of the standard notions of fairness is proportionality (PROP). Let an allocation $X = (X_1, X_2, \ldots, X_n)$ denote a partition of M to N into n bundles such that X_i is allocated to agent i. An allocation X is proportional if $v_i(X_i) \ge \frac{1}{n}v_i(M)$ holds for each agent i. In other words, in a proportional allocation, every agent receives a set of goods whose value is at least 1/n fraction of the value of the entire set. Unfortunately, proportional allocations do not always exist when goods are indivisible. For instance, when allocating a single indivisible good to more than one agents it is impossible to achieve any proportional allocation. Thus, several relaxations of proportionality such as PROP1, PROPx, and PROPm have been considered in the previous work.

Each of these notions requires that each agent $i \in N$ receives value at least $\frac{1}{n}v_i(M) - d_i(X)$, where $d_i(X)$ is a nonnegative real number appropriately defined for each notion. Proportionality up to the largest valued good (PROP1) is a relaxation of proportionality that was introduced by Conitzer et al. [16]. PROP1 requires $d_i(X)$ to be the largest value that agent *i* has for any good allocated to other agents, i.e., $d_i(X) = \max_{k \in N \setminus \{i\}} \max_{g \in X_k} v_i(\{g\})$. It is shown in [16] that there always exists a Pareto optimal¹ allocation that satisfies PROP1. Moreover, Aziz et al. [3] presented a polynomial-time algorithm that finds a PROP1 and Pareto optimal allocation even in the presence of chores, i.e., some items can have negative value.

Another relaxation is proportionality up to the least valued good (PROPx), which is much stronger than PROP1. PROPx requires $d_i(X)$ to be the least value that agent *i* has for any good allocated to other agents, i.e., $d_i(X) = \min_{k \in N \setminus \{i\}} \min_{g \in X_k} v_i(\{g\})$. Moulin [26] gave an example for which no PROPx allocation exists, and Aziz et al. [3] gave a simpler example.

Recently, Baklanov et al. [4] introduced proportionality up to the maximin good (PROPm). PROPm requires $d_i(X) = \max_{k \in N \setminus \{i\}} \min_{g \in X_k} v_i(\{g\})$, which shows that PROPm is the notion between PROP1 and PROPx. It is shown in [4] that a PROPm allocation always exists for instances with at most five agents, and later Baklanov et al. [5] showed that there always

¹An allocation $X = (X_1, \ldots, X_n)$ is Pareto optimal if there is no allocation $Y = (Y_1, \ldots, Y_n)$ such that $v_i(Y_i) \ge v_i(X_i)$ for any agent *i*, and there exists an agent *j* such that $v_j(Y_j) > v_j(X_j)$.

exists a PROPm allocation for any instance and it can be computed in polynomial time. To the best of our knowledge, PROPm has been the best approximate notion of proportionality that is shown to exist for all instances.

However, PROPm is not a good enough relaxation of proportionality in some cases. For example, suppose that there exists a good $g \in M$ for which every agent has value at least 1/n fraction of the value of M. Then allocating g to some agent i and allocating all the goods in $M \setminus \{g\}$ to another agent achieves a PROPm allocation, whereas it will be better to allocate $M \setminus \{g\}$ to $N \setminus \{i\}$ in a fair manner (see Example 1). This motivates the study of better relaxations of proportionality than PROPm.

1.2 Our Contribution

In this paper, we introduce proportionality up to the least valued good on average (PROPavg), a new relaxation of proportionality, and show that there always exists a PROPavg allocation for all instances and can be computed in polynomial time. PROPavg requires $d_i(X)$ to be the average of minimum values that agent *i* has for any good allocated to other agents, i.e., $d_i(X) = \frac{1}{n-1} \sum_{k \in N \setminus \{i\}} \min_{g \in X_k} v_i(\{g\})$. It is easy to see that PROPavg implies PROPm. Note that a similar and slightly stronger notion was introduced by Baklanov et al. [4] with the name of Average-EFX (Avg-EFX), where $d_i(X) = \frac{1}{n} \sum_{k \in N \setminus \{i\}} \min_{g \in X_k} v_i(\{g\})$. It remains open whether an Avg-EFX allocation always exists even in the case of four agents. The following example demonstrates that PROPavg (or Avg-EFX) is a reasonable relaxation of proportionality compared to PROPm.

Example 1. Suppose that $N = \{1, 2, 3\}$, $M = \{g_1, g_2, g_3, g_4\}$, and each agent has an identical additive valuation defined as follows: $v(\{g_1\}) = 10, v(\{g_2\}) = v(\{g_3\}) = 7, v(\{g_4\}) = 6$.

Table 1 summarizes whether several allocations satisfy or not each fairness notions in this instance. As $v(\{g_1\}) \ge 10$, the allocation $(\{g_1, g_2, g_3, g_4\}, \emptyset, \emptyset)$ satisfies PROP1 even though agent 2 and 3 receive no good. Similarly, the allocation $(\{g_1\}, \{g_2, g_3, g_4\}, \emptyset)$ satisfies PROPm even though agent 3 receives no good. It is easy to see that every agent has to receive at least one good in any PROPavg allocation in this instance.

Table 1: The relationship between several allocations and fairness notions in Example 1. The symbol " \checkmark " indicates that the corresponding allocation satisfies the corresponding fairness. The symbol " \checkmark " indicates that the corresponding allocation does not satisfy the corresponding fairness. See Section 1.4 for the definition of EFX.

	EFX	PROPavg	PROPm	PROP1
$(\{g_1\},\{g_2,g_4\},\{g_3\})$	1	✓	1	1
$(\{g_1\},\{g_2,g_3\},\{g_4\})$	X	✓	1	1
$(\{g_1\},\{g_2,g_3,g_4\},\emptyset)$	X	X	1	1
$(\{g_1,g_2,g_3,g_4\},\emptyset,\emptyset)$	X	×	×	1

	$d_i(X)$	Does it always exist?	Time complexity		
PROPx	$\min_{k \in N \setminus \{i\}} \min_{g \in X_k} v_i(\{g\})$	No [3,26]	_		
Avg-EFX	$\frac{1}{n} \sum_{k \in N \setminus \{i\}} \min_{g \in X_k} v_i(\{g\})$	Open	Open		
PROPavg	$\frac{1}{n-1}\sum_{k\in N\setminus\{i\}}\min_{g\in X_k}v_i(\{g\})$	Yes (our result)	Polynomial time (our result)		
PROPm	$\max_{k \in N \setminus \{i\}} \min_{g \in X_k} v_i(\{g\})$	Yes $[5]$	Polynomial time		
PROP1	$\max_{k \in N \setminus \{i\}} \max_{g \in X_k} v_i(\{g\})$	Yes $[3]$	Polynomial time		

Table 2: Relaxations of Proportionality

The main contribution of this paper is the following theorem which extends the results on PROPm allocations shown by Baklanov et al. [5].

Theorem 2. There always exists a **PROPavg** allocation when each agent has a non-negative additive valuation. Furthermore, it can be computed in polynomial time.

Known results on relaxations of proportionality are summarized in Table 2.

1.3 Our Techniques

Our algorithm can be seen as a generalization of *cut-and-choose* protocol, which is a wellknown procedure to fairly allocate resources between two agents. In the cut-and-choose protocol, one agent partitions resources into two bundles for her valuation, and then the other agent chooses the best bundle of the two for her valuation. We generalize this protocol from two agents to n agents in the following way: some n - 1 agents partition the goods into n bundles, and then the remaining agent chooses the best bundle among them for her valuation. To apply this protocol, it suffices to show that there exists a partition of the goods into n bundles such that no matter which bundle the remaining agent chooses, the remaining n - 1 bundles can be allocated to the first n - 1 agents fairly.

In our algorithm, we find such a partition by using an auxiliary graph called *PROPavg-graph*. A formal definition of the PROPavg-graph is given in Section 3. In Section 4, we show the existence of a PROPavg allocation by constructing a pseudo-polynomial time algorithm for finding it. In Section 5, by improving our algorithm in Section 4, we show how to find a PROPavg allocation in polynomial time.

Let us emphasize that introducing the PROPavg-graph is a key technical ingredient in this paper. It is also worth noting that Hall's marriage theorem [20], a classical and famous result in discrete mathematics, plays an important role in our argument.

1.4 Related Work

Fair division of divisible resources is a classical topic starting from the 1940's [29] and has a long history in multiple fields such as Economics, Social Choice Theory, and Computer




Figure 1: The relationship among some fairness notions and the existence results for these fairness notions. EF, PROP, and PROPx do not always exist, whereas PROPavg, PROPm, EF1, and PROP1 always exist when each agent has non-negative additive valuation. It is not known whether EFX or Avg-EFX always exist or not for all instances.

Science [8,9,25,28]. On the other hand, fair division of indivisible items has actively studied in recent years (see, e.g., [2] for a recent survey).

In the context of fair division, besides proportionality, envy-freeness (EF) is another wellstudied notion of fairness. An allocation is called *envy-free* if for each agent, she receives a set of goods for which she has value at least value of the set of goods any other agent receives. As in the proportionality case, envy-free allocations do not always exist when goods are indivisible, and several relaxations of envy-freeness have been considered. Among them, a notable one is *envy-freeness up to one good* ($\mathsf{EF1}$) [10]. It is known that there always exists an EF1 allocation, and it can be computed in polynomial time [22]. Another notable relaxation is envy-freeness up to any good (EFX) [12]. An allocation $X = (X_1, \ldots, X_n)$ is called EFX if for any pair of agents $i, j \in N$, $v_i(X_i) \ge v_i(X_j) - m_i(X_j)$, where $m_i(X_j)$ is the value of the least valuable good for agent i in X_i . It is one of the major open problems in fair division whether EFX allocations always exist or not. As mentioned in [4], it is easy to see that EFX implies Avg-EFX. As with EFX, it is not known whether Avg-EFX allocations always exist for instances with four or more agents. The relationship among notions mentioned above and the existence results are summarized in Figure 1. There have been several studies on the existence of an EFX allocation for restricted cases. Plaut and Roughgarden [27] showed that an EFX allocation always exists for instances with two agents even when each agent can have more general valuations than additive valuations. Chaudhury et al. [13] showed that an EFX allocation always exists for instances with three agents. It is not known whether EFX allocations always exist even in the case of four agents having additive valuations. There are several studies of the cases with restricted valuations. For example, there always exists an EFX allocation when valuations are identical [27], two types [23, 24], binary [6, 17], or

bi-valued [1].

Another approach related to EFX is a setting to allow unallocated goods, which is known as *EFX-with-charity*. Obviously, without any constraints, the problem is trivial: leaving all goods unallocated results in an envy-free allocation. Thus, the goal here is to find allocations with better guarantees. For additive valuations, Caragiannis et al. [11] showed that there exists an EFX allocation with some unallocated goods where every agent receives at least half the value of her bundle in a maximum *Nash social welfare* allocation². For normalized and monotone valuations, Chaudhury et al. [15] showed that there exist an EFX allocation and a set of unallocated goods U such that every agent has value for her own bundle at least value for U, and |U| < n. Berger et al. [7] showed that the number of the unallocated goods can be decreased to n - 2, and to just one for the case of four agents having nice cancelable valuations, which are more general than additive valuations. Mahara [24] showed that the number of the unallocated goods can be decreased to n - 2 for normalized and monotone valuations, which are more general than nice cancelable valuations. For additive valuations, Chaudhury et al. [14] presented a polynomial-time algorithm for finding an approximate EFX allocation with at most a sublinear number of unallocated goods and high Nash social welfare.

2 Preliminaries

Let $N = \{1, \ldots, n\}$ be a set of n agents and M be a set of m goods. We assume that goods are indivisible: a good can not be split among multiple agents. Each agent $i \in N$ has a nonnegative valuation $v_i : 2^M \to \mathbb{R}_{\geq 0}$, where 2^M is the power set of M. We assume that each valuation v_i is *additive*, i.e., $v_i(S) = \sum_{g \in S} v_i(\{g\})$ for any $S \subseteq M$. Note that since valuations are non-negative and additive, they have to be *normalized*: $v_i(\emptyset) = 0$ and *monotone*: $S \subseteq T$ implies $v_i(S) \leq v_i(T)$ for any $S, T \subseteq M$. For ease of explanation, we normalize the valuations so that $v_i(M) = 1$ for all $i \in N$.

To simplify notation, we denote $\{1, \ldots, k\}$ by [k] for any positive integer k, write $v_i(g)$ instead of $v_i(\{g\})$ for $g \in M$, and use $S \setminus g$ and $S \cup g$ instead of $S \setminus \{g\}$ and $S \cup \{g\}$, respectively.

We say that $X = (X_1, X_2, ..., X_n)$ is an allocation of M to N if it is a partition of Minto n disjoint subsets such that each set is indexed by $i \in N$. Each X_i is the set of goods given to agent i, which we call a bundle. It is simply called an allocation to N if M is clear from context. For $i \in N$ and $S \subseteq M$, let $m_i(S)$ denote the value of the least valuable good for agent i in S, that is, $m_i(S) = \min_{g \in S} \{v_i(g)\}$ if $S \neq \emptyset$ and $m_i(\emptyset) = 0$. For an allocation $X = (X_1, X_2, ..., X_n)$ to N, we say that an agent i is *PROPavg-satisfied* by X if

$$v_i(X_i) + \frac{1}{n-1} \sum_{k \in [n] \setminus i} m_i(X_k) \ge \frac{1}{n},$$

²This is an allocation that maximizes $\prod_{i=1}^{n} v_i(X_i)$.

where we recall that $v_i(M) = 1$. In other words, agent *i* receives a set of goods for which she has value at least 1/n fraction of her total value minus the average of minimum value of the set of goods any other agent receives. An allocation X is called *PROPavg* if every agent $i \in N$ is **PROPavg**-satisfied by X.

Let G = (V, E) be a graph. For $S \subseteq V$, let $\Gamma_G(S) = \{v \in V \setminus S \mid (s, v) \in E \text{ for some } s \in S\}$ denote the set of neighbors of S in G. For $v \in V$, let G - v denote the graph obtained from G by deleting v. A *perfect matching* in G is a set of pairwise disjoint edges of G covering all the vertices of G.

3 Key Ingredient: **PROPavg-**Graph

In order to prove Theorem 2, we give an algorithm for finding a PROPavg allocation. As described in Section 1.3, our algorithm is a generalization of the cut-and-choose protocol that consists of the following three steps.

- 1. We partition the goods into n bundles without assigning them to agents.
- 2. A specified agent, say n, chooses the best bundle for her valuation.
- 3. We determine an assignment of the remaining bundles to the agents in $N \setminus n$.

The partition given in the first step is represented by an allocation of M to a newly introduced set of size n, say V_2 , and the assignment in the third step is represented by a matching in an auxiliary bipartite graph, which we call *PROPavg-graph*. In this section, we define the **PROPavg-graph** and its desired properties.

Let V_2 be a set of n elements and fix a specified element $r \in V_2$. We say that $X = (X_u)_{u \in V_2}$ is an allocation to V_2 if it is a partition of M into n disjoint subsets such that each set is indexed by an element in V_2 , that is, $\bigcup_{u \in V_2} X_u = M$ and $X_u \cap X_{u'} = \emptyset$ for distinct $u, u' \in V_2$. For an allocation $X = (X_u)_{u \in V_2}$ to V_2 , we define a bipartite graph $G_X = (V_1, V_2; E)$ called *PROPavg-graph* as follows. The vertex set consists of $V_1 = N \setminus n$ and V_2 , and the edge set Eis defined by

$$(i,u) \in E \iff v_i(X_u) + \frac{1}{n-1} \sum_{u' \in V_2 \setminus \{r,u\}} m_i(X_{u'}) \ge \frac{1}{n}$$

for $i \in V_1$ and $u \in V_2$. It should be emphasized that the summation is taken over $V_2 \setminus \{r, u\}$, i.e., $m_i(X_r)$ is not counted, in the above definition, which is crucial in our argument. The following lemma shows that the PROPavg-graph is closely related to the definition of PROPavg-satisfaction.

Lemma 3. Suppose that $G_X = (V_1, V_2; E)$ is the PROPavg-graph for an allocation $X = (X_u)_{u \in V_2}$ to V_2 . Let σ be a bijection from N to V_2 and define an allocation $Y = (Y_1, \ldots, Y_n)$ to N by $Y_i = X_{\sigma(i)}$ for $i \in N$. For $i^* \in V_1$, if $(i^*, \sigma(i^*)) \in E$, then i^* is PROPavg-satisfied by Y.




Figure 2: An example of (a) represents a PROPavg-graph G_X corresponding to X that satisfies (P1) but does not satisfy (P2). An example of (b) represents a PROPavg-graph G_X corresponding to X that satisfies (P2).

Proof. Let $u^* = \sigma(i^*)$ and suppose that $(i^*, u^*) \in E$. We directly obtain

$$v_{i^*}(Y_{i^*}) + \frac{1}{n-1} \sum_{j \in [n] \setminus i^*} m_{i^*}(Y_j) \ge v_{i^*}(X_{u^*}) + \frac{1}{n-1} \sum_{u \in V_2 \setminus \{u^*, r\}} m_{i^*}(X_u) \ge \frac{1}{n}$$

where the first inequality follows from the definition of Y and $m_{i^*}(X_r) \ge 0$, and the second inequality follows from $(i^*, u^*) \in E$.

For an allocation $X = (X_u)_{u \in V_2}$ to V_2 , we consider the following two properties, which play an important role in our argument.

(P1) $G_X - r$ has a perfect matching.

(P2) For any $u \in V_2$, $G_X - u$ has a perfect matching.

Obviously, property (P2) is stronger than property (P1). Roughly speaking, our goal is to find an allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P2). As shown later, we find an allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P2) while keeping (P1). Examples of a PROPavg-graph G_X corresponding to X satisfying (P1) or (P2) are described in Figure 2. We can rephrase these conditions by using the following classical theorem known as Hall's marriage theorem in discrete mathematics.

Theorem 4 (Hall's marriage theorem [20]). Suppose that G = (A, B; E) is a bipartite graph with |A| = |B|. Then, G has a perfect matching if and only if $|S| \le |\Gamma_G(S)|$ for any $S \subseteq A$.

The property (P1) is equivalent to $|S| \leq |\Gamma_{G_X-r}(S)|$ for any $S \subseteq V_1$ by this theorem. The property (P2) is equivalent to $|S| \leq |\Gamma_{G_X-u}(S)|$ for any $u \in V_2$ and $S \subseteq V_1$ by Hall's marriage theorem. By simple observation, we can obtain another characterization of property (P2).

Lemma 5. Let $X = (X_u)_{u \in V_2}$ be an allocation to V_2 . Then, X satisfies (P2) if and only if $|S| + 1 \leq |\Gamma_{G_X}(S)|$ for any non-empty subset $S \subseteq V_1$.

Proof. By Hall's marriage theorem, it is sufficient to show that the following two conditions are equivalent:

- (i) $|S| \leq |\Gamma_{G_X-u}(S)|$ for any $u \in V_2$ and $S \subseteq V_1$, and
- (ii) $|S| + 1 \le |\Gamma_{G_X}(S)|$ for any non-empty subset $S \subseteq V_1$.

Suppose that (i) holds. Let S be a nonempty subset of V_1 . Since (i) implies that $|\Gamma_{G_X}(S)| \geq |S| \geq 1$, we obtain $\Gamma_{G_X}(S) \neq \emptyset$. Let $u \in \Gamma_{G_X}(S)$. By (i) again, we obtain $|\Gamma_{G_X}(S)| = |\Gamma_{G_X-u}(S)| + 1 \geq |S| + 1$. This shows (ii).

Conversely, suppose that (ii) holds. Let $u \in V_2$ and let $S \subseteq V_1$. If $S = \emptyset$, then it clearly holds that $|S| \leq |\Gamma_{G_X-u}(S)|$. If $S \neq \emptyset$, then we have $|S| + 1 \leq |\Gamma_{G_X}(S)| \leq |\Gamma_{G_X-u}(S)| + 1$, which implies that $|S| \leq |\Gamma_{G_X-u}(S)|$. This shows (i).

4 Existence of a **PROPavg** Allocation

In this section, we show that there always exists a PROPavg allocation by constructing a pseudo-polynomial time algorithm for finding it. Improvements in time complexity are discussed in Section 5. Our algorithm begins with obtaining an initial allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P1). Unless X satisfies (P2), we appropriately choose a good in $\bigcup_{u \in V_2 \setminus r} X_u$ and move it to X_r while keeping (P1). Finally, we get an allocation $X^* = (X_u^*)_{u \in V_2}$ to V_2 satisfying (P2). As we will see later, we can obtain a PROPavg allocation to N from X^* .

4.1 Our Algorithm

In order to obtain an initial allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P1), we use the following previous result about *EFX-with-charity*.

Theorem 6 (Chaudhury et al. [15]). For normalized and monotone valuations, there always exists an allocation $X = (X_1, \ldots, X_n)$ of $M \setminus U$ to N, where U is a set of unallocated goods, such that

- X is EFX, that is, $v_i(X_i) + m_i(X_j) \ge v_i(X_j)$ for any pair of agents $i, j \in N$,
- $v_i(X_i) \ge v_i(U)$ for any agent $i \in N$, and
- |U| < n.

The following lemma shows that by applying Theorem 6 to agents $N \setminus n$, we can obtain an initial allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P1).

Lemma 7. There exists an allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P1).

Proof. By applying Theorem 6 to agents $N \setminus n$, we can obtain an allocation $Y = (Y_1, \ldots, Y_{n-1})$ of $M \setminus U$ to $N \setminus n$, where U is a set of unallocated goods, satisfying the conditions in Theorem 6. Let $V_2 = \{r, u_1, \ldots, u_{n-1}\}$ and define an allocation $X = (X_u)_{u \in V_2}$ to V_2 as $X_{u_j} = Y_j$ for $j \in [n-1]$ and $X_r = U$. Let $G_X = (V_1, V_2; E)$ be the PROPavg-graph for X. We show that X satisfies (P1).

Fix any agent $i \in V_1$. We have $v_i(X_{u_i}) + m_i(X_{u_j}) \ge v_i(X_{u_j})$ for any $j \in [n-1] \setminus i$ since Y is EFX and $X_{u_j} = Y_j$. We also have $v_i(X_{u_i}) = v_i(Y_i) \ge v_i(U) = v_i(X_r)$ and a trivial inequality $v_i(X_{u_i}) \ge v_i(X_{u_i})$. By summing up these inequalities, we obtain $n \cdot v_i(X_{u_i}) + \sum_{j \in [n-1] \setminus i} m_i(X_{u_j}) \ge \sum_{u \in V_2} v_i(X_u)$. Since $\sum_{u \in V_2} v_i(X_u) = v_i(M) = 1$, this shows that $v_i(X_{u_i}) + \frac{1}{n} \sum_{j \in [n-1] \setminus i} m_i(X_{u_j}) \ge \frac{1}{n}$, and hence $(i, u_i) \in E$. Therefore, $G_X - r$ has a perfect matching $\{(i, u_i) \mid i \in [n-1]\}$, which implies that X satisfies (P1).

The following lemma shows that if we obtain an allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P2), then there exists a PROPavg allocation to N.

Lemma 8. Suppose that $X = (X_u)_{u \in V_2}$ is an allocation to V_2 satisfying (P2). Then, we can construct a **PROPavg** allocation to N.

Proof. Let $X = (X_u)_{u \in V_2}$ be an allocation to V_2 satisfying (P2). First, agent n chooses the best bundle X_{u^*} for her valuation among $\{X_u \mid u \in V_2\}$ (if there is more than one such bundle, choose one arbitrarily). Since X satisfies (P2), there exists a perfect matching A in $G_X - u^*$. For each agent $i \in V_1(=N \setminus n)$, the bundle that matches i in A is allocated to i. By Lemma 3, agent i is PROPavg-satisfied for each agent $i \in V_1$. Furthermore, since we have $v_n(X_{u^*}) = \max_{u \in V_2} v_n(X_u) \geq \frac{1}{n}$, agent n is also PROPavg-satisfied. Therefore, the obtained allocation is a PROPavg allocation to N.

The following proposition shows how we update an allocation in each iteration, whose proof is given in Section 4.2.

Proposition 9. Suppose that $X = (X_u)_{u \in V_2}$ is an allocation to V_2 that satisfies (P1) but does not satisfy (P2). Then, there exists another allocation $X' = (X'_u)_{u \in V_2}$ to V_2 satisfying (P1) such that $|X'_r| = |X_r| + 1$.

As we will see in Section 4.2, an allocation X' in Proposition 9 is obtained by moving some appropriate item $g \in \bigcup_{u \in V_2 \setminus r} X_u$ to X_r . If Proposition 9 holds, then we can show that there always exists a PROPavg allocation when each agent has a non-negative additive valuation as follows. See also Algorithm 1.

By Lemma 7, we first obtain an initial allocation $X = (X_u)_{u \in V_2}$ to V_2 satisfying (P1). By Proposition 9, unless X satisfies (P2), we can increase $|X_r|$ by one while keeping the condition (P1). Since $|X_r| \leq |M|$, this procedure terminates in at most m steps, and we finally obtain an allocation X^* to V_2 satisfying (P2). Therefore, there exists a PROPavg allocation to N by Lemma 8. Algorithm 1 Algorithm for finding a PROPavg allocation

Input: agents N, goods M, and valuation v_i for each $i \in N$

Output: a PROPavg allocation to N

- 1: Apply Lemma 7 to obtain an allocation X to V_2 satisfying (P1).
- 2: while X does not satisfy (P2) do
- 3: Apply Proposition 9 to X and obtain another allocation X' to V_2 .
- 4: $X \leftarrow X'$.
- 5: Apply Lemma 8 to obtain a PROPavg allocation to N.

4.2 **Proof of Proposition 9**

Let $X = (X_u)_{u \in V_2}$ be an allocation to V_2 . For $u^* \in V_2 \setminus r$ and $g \in X_{u^*}$, we say that an allocation $X' = (X'_u)_{u \in V_2}$ to V_2 is obtained from X by moving g to X_r if

$$X'_{u} = \begin{cases} X_{r} \cup g & \text{if } u = r, \\ X_{u^{*}} \setminus g & \text{if } u = u^{*}, \\ X_{u} & \text{otherwise.} \end{cases}$$

The following lemma guarantees that if there exists an agent $i \in V_1$ such that $(i, r) \notin E$ in the PROPavg-graph $G_X = (V_1, V_2; E)$, then we can move some good in $\bigcup_{u \in V_2 \setminus r} X_u$ to X_r so that the edges incident to i do not disappear. This lemma is crucial in the proof of Proposition 9.

Lemma 10. Let $X = (X_u)_{u \in V_2}$ be an allocation to V_2 and let $i \in V_1$ be an agent such that $(i, r) \notin E$ in the PROPavg-graph $G_X = (V_1, V_2; E)$. Then, there exist $u^* \in V_2$ and $g \in X_{u^*}$ such that $(i, u^*) \in E$, $|X_{u^*}| \ge 2$, and the following property holds: if an allocation X' to V_2 is obtained from X by moving g to X_r , then the corresponding PROPavg-graph $G_{X'}$ has an edge (i, u^*) .

Proof. We first show that $X_u \neq \emptyset$ for any $u \in V_2$ with $(i, u) \in E$. Indeed, if $X_u = \emptyset$, then we have

$$v_i(X_r) + \frac{1}{n-1} \sum_{u' \in V_2 \setminus r} m_i(X_{u'}) \ge v_i(X_u) + \frac{1}{n-1} \sum_{u' \in V_2 \setminus \{r,u\}} m_i(X_{u'}) \ge \frac{1}{n}$$

where the first inequality follows from $v_i(X_u) = m_i(X_u) = 0$ and the second inequality follows from $(i, u) \in E$. This contradicts $(i, r) \notin E$.

To derive a contradiction, assume that u^* and g satisfying the conditions in Lemma 10 do not exist. Then, we have the following claim.

Claim 11. For any $u \in V_2$ with $(i, u) \in E$, we obtain

$$v_i(X_u) - m_i(X_u) + \frac{1}{n-1} \sum_{\substack{u' \in V_2 \setminus \{r, u\}:\\(i, u') \in E}} m_i(X_{u'}) < \frac{1}{n}.$$
(1)

Proof of the Claim. Fix $u \in V_2$ with $(i, u) \in E$. Let g be a good in X_u that minimizes $v_i(g)$, where we note that $X_u \neq \emptyset$ as described above. Then, $v_i(g) = m_i(X_u)$. Define $X' = (X'_{u'})_{u' \in V_2}$ as the allocation to V_2 that is obtained from X by moving g to X_r . Let $G_{X'} = (V_1, V_2; E')$ be the PROPavg-graph corresponding to X'. Since u and g do not satisfy the conditions in Lemma 10 by our assumption, we have $(i, u) \notin E'$ or $|X_u| = 1$.

If $(i, u) \in E'$, then we obtain $|X_u| = 1$, and hence

$$v_{i}(X_{r}) + \frac{1}{n-1} \sum_{u' \in V_{2} \setminus r} m_{i}(X_{u'}) \ge \frac{1}{n-1} \sum_{u' \in V_{2} \setminus \{r,u\}} m_{i}(X_{u'})$$
$$= v_{i}(X'_{u}) + \frac{1}{n-1} \sum_{u' \in V_{2} \setminus \{r,u\}} m_{i}(X'_{u'})$$
$$\ge \frac{1}{n},$$

where the equality follows from $v_i(X'_u) = v_i(\emptyset) = 0$ and the last inequality follows from $(i, u) \in E'$. This contradicts $(i, r) \notin E$.

Thus, it holds that $(i, u) \notin E'$. Since $v_i(X_u) - m_i(X_u) = v_i(X'_u)$, we obtain

$$v_{i}(X_{u}) - m_{i}(X_{u}) + \frac{1}{n-1} \sum_{\substack{u' \in V_{2} \setminus \{r, u\}:\\(i, u') \in E}} m_{i}(X_{u'})$$

$$\leq v_{i}(X_{u}) - m_{i}(X_{u}) + \frac{1}{n-1} \sum_{\substack{u' \in V_{2} \setminus \{r, u\}}} m_{i}(X_{u'})$$

$$= v_{i}(X'_{u}) + \frac{1}{n-1} \sum_{\substack{u' \in V_{2} \setminus \{r, u\}}} m_{i}(X'_{u'})$$

$$< \frac{1}{n},$$

where the last inequality follows from $(i, u) \notin E'$.

By summing up inequality (1) for each $u \in V_2$ with $(i, u) \in E$, we obtain the following inequality:

$$\sum_{\substack{u \in V_2: \\ (i,u) \in E}} v_i(X_u) + \left(-1 + \frac{l-1}{n-1}\right) \sum_{\substack{u' \in V_2 \setminus r: \\ (i,u') \in E}} m_i(X_{u'}) < \frac{l}{n},\tag{2}$$

where $l = |\{u \in V_2 \mid (i, u) \in E\}|.$

On the other hand, for any $u \in V_2$ with $(i, u) \notin E$, we have

$$v_i(X_u) + \frac{1}{n-1} \sum_{\substack{u' \in V_2 \setminus r: \\ (i,u') \in E}} m_i(X_{u'}) \le v_i(X_u) + \frac{1}{n-1} \sum_{u' \in V_2 \setminus \{r,u\}} m_i(X_{u'}) < \frac{1}{n},$$
(3)

where the both inequalities follow from $(i, u) \notin E$. Summing up inequality (3) for each $u \in V_2$ with $(i, u) \notin E$, we obtain

$$\sum_{\substack{u \in V_2:\\(i,u) \notin E}} v_i(X_u) + \left(\frac{n-l}{n-1}\right) \sum_{\substack{u' \in V_2 \setminus r:\\(i,u') \in E}} m_i(X_{u'}) < \frac{n-l}{n},\tag{4}$$

where we note that $|\{u \in V_2 \mid (i, u) \notin E\}| = n - l.$

By taking the sum of inequalities (2) and (4), we obtain

$$\sum_{\substack{u \in V_2: \\ i,u) \in E}} v_i(X_u) + \sum_{\substack{u \in V_2: \\ (i,u) \notin E}} v_i(X_u) < 1,$$

which contradicts $\sum_{u \in V_2} v_i(X_u) = 1.$

Therefore, there exist $u^* \in V_2$ and $g \in X_{u^*}$ satisfying the conditions in Lemma 10. \Box

We are now ready to prove Proposition 9.

Proof of Proposition 9. Suppose that $X = (X_u)_{u \in V_2}$ is an allocation to V_2 that satisfies (P1) but does not satisfy (P2). Let $G_X = (V_1, V_2; E)$ be the PROPavg-graph corresponding to X. Since X does not satisfy (P2), there exists a non-empty set $S \subseteq V_1$ such that $|S| + 1 > |\Gamma_{G_X}(S)|$ by Lemma 5. Among such sets, let $S^* \subseteq V_1$ be an inclusion-wise minimal one. By the integrality of $|\Gamma_{G_X}(S^*)|$ and $|S^*|$, this means that $|S^*| \ge |\Gamma_{G_X}(S^*)|$ and $|S|+1 \le |\Gamma_{G_X}(S)|$ for any non-empty proper subset $S \subsetneq S^*$. We now show some properties of S^* .

Claim 12. For any $i \in S^*$, it holds that $(i, r) \notin E$.

Proof of the claim. Since X satisfies (P1), we have $|S^*| \leq |\Gamma_{G_X-r}(S^*)|$ by Hall's marriage theorem. Hence, we obtain $|S^*| \leq |\Gamma_{G_X-r}(S^*)| \leq |\Gamma_{G_X}(S^*)| \leq |S^*|$, where the last inequality follows from the definition of S^* . This shows that all the above inequalities are tight. Since $|\Gamma_{G_X-r}(S^*)| = |\Gamma_{G_X}(S^*)|$, we obtain $r \notin \Gamma_{G_X}(S^*)$, that is, $(i,r) \notin E$ for any $i \in S^*$.

Claim 13. For any non-empty proper subset $S \subsetneq S^*$, it holds that $|S| + 1 \leq |\Gamma_{G_X - r}(S)|$.

Proof of the claim. Let S be a non-empty proper subset of S^* . Since $(i, r) \notin E$ for any $i \in S \subsetneq S^*$ by Claim 12, we have that $\Gamma_{G_X}(S) = \Gamma_{G_X-r}(S)$. Hence, we obtain $|S| + 1 \leq |\Gamma_{G_X}(S)| = |\Gamma_{G_X-r}(S)|$, where the inequality follows from the minimality of S^* . \Box

Claim 14. For any $i \in S^*$ and $u \in \Gamma_{G_X}(S^*)$ with $(i, u) \in E$, $G_X - r$ has a perfect matching in which i matches u.

Proof of the claim. Fix any $i \in S^*$ and $u \in \Gamma_{G_X}(S^*)$ with $(i, u) \in E$. Note that $r \notin \Gamma_{G_X}(S^*)$ by Claim 12, and hence $u \neq r$.

Since X satisfies (P1), $G_X - r$ has a perfect matching A. In A, it is obvious that every vertex in S^* is matched to a vertex in $\Gamma_{G_X-r}(S^*)$. Conversely, every vertex in $\Gamma_{G_X-r}(S^*)$




Figure 3: A PROPavg-graph G_X corresponding to X in the proof of Proposition 9. A nonempty minimal subset S^* of V_1 such that $|S^*| + 1 > |\Gamma_{G_X}(S^*)|$ is represented by vertices in the above dotted rectangle.

is matched to a vertex in S^* as $|S^*| = |\Gamma_{G_X - r}(S^*)|$ (see the proof of Claim 12). Thus, by removing the edges between S^* and $\Gamma_{G_X}(S^*)$ from A, we obtain a matching $A_1 \subseteq A$ that exactly covers $V_1 \setminus S^*$ and $V_2 \setminus (\Gamma_{G_X}(S^*) \cup \{r\})$.

Let G'_X be the subgraph of G_X induced by $(S^* \setminus i) \cup (\Gamma_{G_X}(S^*) \setminus u)$. We now show that G'_X has a perfect matching. Consider any $S \subseteq S^* \setminus i$. If $S = \emptyset$, then it clearly holds that $|S| \leq |\Gamma_{G'_X}(S)|$. If $S \neq \emptyset$, then $|S| + 1 \leq |\Gamma_{G_X - r}(S)| \leq |\Gamma_{G'_X}(S) \cup u| = |\Gamma_{G'_X}(S)| + 1$, where the first inequality is by Claim 13. Therefore, $|S| \leq |\Gamma_{G'_X}(S)|$ holds for any $S \subseteq S^* \setminus i$, and hence G'_X has a perfect matching A_2 by Hall's marriage theorem.

Then, $A_1 \cup A_2 \cup \{(i, u)\}$ is a desired perfect matching in $G_X - r$.

Fix any agent $i^* \in S^*$. Since $(i^*, r) \notin E$ by Claim 12, by applying Lemma 10 to agent i^* , we obtain $u^* \in V_2$ and $g \in X_{u^*}$ satisfying the conditions in Lemma 10. See also Figure 3. Let $X' = (X'_u)_{u \in V_2}$ be the allocation to V_2 obtained from X by moving g to X_r and let $G_{X'} = (V_1, V_2; E')$ be the PROPavg-graph corresponding to X'. Then, the conditions in Lemma 10 show that $(i^*, u^*) \in E \cap E'$ and $|X_{u^*}| \geq 2$. We also see that E' satisfies the following.

Claim 15. For any $i \in V_1$ and $u \in V_2 \setminus u^*$, if $(i, u) \in E$ then $(i, u) \in E'$.

Proof of the claim. Since $|X_{u^*}| \geq 2$, we have $m_i(X'_{u^*}) = m_i(X_{u^*} \setminus g) \geq m_i(X_{u^*})$ for any agent $i \in V_1$. Hence, for any $i \in V_1$ and $u \in V_2 \setminus u^*$ with $(i, u) \in E$, we obtain

$$v_i(X'_u) + \frac{1}{n-1} \sum_{u' \in V_2 \setminus \{r,u\}} m_i(X'_{u'}) \ge v_i(X_u) + \frac{1}{n-1} \sum_{u' \in V_2 \setminus \{r,u\}} m_i(X_{u'}) \ge \frac{1}{n},$$

which shows that $(i, u) \in E'$.

By Claim 14 and $(i^*, u^*) \in E$, there exists a perfect matching A in $G_X - r$ in which i^* matches u^* . Then, by Claim 15 and $(i^*, u^*) \in E'$, we see that $A \subseteq E'$, that is, A is a perfect matching also in $G_{X'} - r$. Therefore, X' satisfies (P1). Since $|X'_r| = |X_r| + 1$ clearly holds by definition, X' satisfies the conditions in Proposition 9.

5 Finding a **PROPavg** Allocation in Polynomial Time

In this section, we show how to find a PROPavg allocation in polynomial time. As mentioned in Section 4, Algorithm 1 runs in pseudo-polynomial time. This is because we can not guarantee the polynomial solvability in line 1 of Algorithm 1. We can see that the other parts of Algorithm 1 run in polynomial time as follows. In line 2, we can check (P2) in polynomial time by applying a maximum matching algorithm for each $G_X - u$. In line 3, it suffices to find a good $g \in \bigcup_{u \in V_2 \setminus r} X_u$ such that (P1) is kept after moving g. Since (P1) can be checked in polynomial time, this can be done in polynomial time by considering all g in a brute-force way. Finally, line 5 is executed in polynomial time by Lemma 8. Note that we can speed up lines 2 and 3 by using the DM-decomposition of G_X [18,19], but we do not go into details, because we only focus on the polynomial solvability.

Let us now consider how to find an initial allocation X to V_2 satisfying (P1) in polynomial time. Our idea is to use a recursive algorithm. That is, we use a PROPavg allocation of M to n-1 agents as an initial allocation X to V_2 satisfying (P1). Indeed, if it holds that $v_i(g) \leq \frac{1}{n}$ for any agent $i \in N$ and any good $g \in M$, then we can show that a PROPavg allocation of M to n-1 agents satisfies (P1) as follows.

Lemma 16. Suppose that for any agent $i \in N$ and any good $g \in M$, we have $v_i(g) \leq \frac{1}{n}$. Let (X_1, \ldots, X_{n-1}) be a **PROPavg** allocation for $N \setminus n$. Then, $X = (X_1, \ldots, X_{n-1}, X_n)$ is an allocation to $V_2 = [n]$ satisfying (P1), where $X_n = \emptyset$ and the specific element $r \in V_2$ is equal to n.

Proof. Let $G_X = (V_1, V_2; E)$ be the PROPavg-graph corresponding to X. It is enough to

show that $(i, X_i) \in E$ for any $i \in [n-1]$. Fix any $i \in [n-1]$. We obtain that

$$v_{i}(X_{i}) \geq \frac{1}{n-1} - \frac{1}{n-2} \sum_{j \in [n-1] \setminus i} m_{i}(X_{j})$$

$$= \frac{1}{n} - \frac{1}{n-1} \sum_{j \in [n-1] \setminus i} m_{i}(X_{j})$$

$$+ \underbrace{\frac{1}{n-1} \left(\frac{1}{n} - \frac{1}{n-2} \sum_{j \in [n-1] \setminus i} m_{i}(X_{j}) \right)}_{\geq 0}$$

$$\geq \frac{1}{n} - \frac{1}{n-1} \sum_{j \in [n-1] \setminus i} m_{i}(X_{j}),$$

where the first inequality follows from the assumption that (X_1, \ldots, X_{n-1}) is a PROPavg allocation and the second inequality follows from the assumption that $v_i(g) \leq \frac{1}{n}$ for any $i \in N$ and $g \in M$. This implies that $(i, X_i) \in E$ and thus X is an allocation to $V_2 = [n]$ satisfying (P1).

Unfortunately, the argument in Lemma 16 does not work without the assumption that $v_i(g) \leq \frac{1}{n}$ for any $i \in N$ and $g \in M$. To elude this difficulty, our algorithm applies preprocessing. This preprocessing allocates g to i and remove i and g from our instance as long as there exists an agent i and a good g such that $v_i(g) \geq \frac{1}{n}$. See Algorithm 2 for the entire algorithm.

If this preprocessing removes at least one agent from our instance, then our algorithm recursively computes a PROPavg allocation for the remaining agents and goods, and returns the overall allocation together with the removed agents. In order to verify that the returned allocation is a PROPavg allocation for n agents, we need a refined condition (see line 7 of Algorithm 2).

Otherwise, our algorithm recursively computes a PROPavg allocation for n-1 agents. Since $v_i(g) < \frac{1}{n}$ holds for any agent *i* and good *g*, we can use this allocation as an initial allocation to V_2 satisfying (P1) by Lemma 16. The rest of our algorithm finds an allocation to V_2 satisfying (P2) and returns a PROPavg allocation as in Algorithm 1.

In the remaining part of this section, we show the correctness and the polynomial solvability of Algorithm 2. The following lemma shows that if the preprocessing removes at least one agent from our instance, then the algorithm returns a legal PROPavg allocation for N.

Lemma 17. In line 14 of Algorithm 2, $X = (X_1, \ldots, X_{|N|})$ is a PROPavg allocation to N.

Proof. Fix any $i \in N$. We show that i is **PROPavg**-satisfied by X.

Case 1: $i \in N_2$

Algorithm 2 Algorithm for finding a PROPavg allocation in polynomial time

1: procedure PROPAVG $(N, M, \{v_i\}_{i \in N})$ if |N| = 1 then 2:3: return X = (M)4: else $N_1 \leftarrow N, N_2 \leftarrow \emptyset$ 5:6: $M_1 \leftarrow M, M_2 \leftarrow \emptyset$ while $\exists i \in N_1$ and $\exists g \in M_1$ s.t. $v_i(g) \geq \frac{v_i(M)}{|N|} - \frac{1}{|N|-1} \sum_{j \in N_2} m_i(X_j)$ do 7: $X_i \leftarrow \{g\}$ 8: $N_1 \leftarrow N_1 \setminus i, N_2 \leftarrow N_2 \cup i$ 9: $M_1 \leftarrow M_1 \setminus g, M_2 \leftarrow M_2 \cup g$ 10: Let $N_1 = \{1, \ldots, l\}$ and $N_2 = \{l + 1, \ldots, |N|\}$, renumbering if necessary. 11: if $|N_2| \geq 1$ then 12:13: $(X_1,\ldots,X_l) \leftarrow \operatorname{PROPAVG}(N_1, M_1, \{v_i\}_{i \in N_1})$ return $X = (X_1, \ldots, X_{|N|})$ 14:else 15: $(X_1, \ldots, X_{n-1}) \leftarrow \operatorname{PROPAVG}(N \setminus n, M, \{v_i\}_{i \in N \setminus n})$ $\triangleright N_1 = N, M_1 = M$ 16:Apply Lemma 16 to obtain an allocation $X = (X_1, \ldots, X_n)$ satisfying (P1). 17:while X does not satisfy (P2) do 18:Apply Proposition 9 to X and obtain another allocation X' to V_2 . 19: $X \leftarrow X'$. 20:Apply Lemma 8 to obtain a PROPavg allocation $X = (X_1, \ldots, X_{|N|})$ to N. 21:22: return $X = (X_1, \ldots, X_{|N|})$

In this case, agent i receives exactly one good in the while statement. By the while condition, we have

$$v_i(X_i) \ge \frac{1}{n} - \frac{1}{n-1} \sum_{j \in N_2} m_i(X_j)$$
$$\ge \frac{1}{n} - \frac{1}{n-1} \sum_{j \in N \setminus i} m_i(X_j).$$

Thus, i is PROPavg-satisfied by X.

Case 2: $i \in N_1$ and l = 1

In this case, we have

$$v_{i}(X_{i}) = v_{i}(M_{1}) = v_{i}(M) - \sum_{j \in N_{2}} m_{i}(X_{j})$$

$$= \left(\frac{1}{n} - \frac{1}{n-1} \sum_{j \in N_{2}} m_{i}(X_{j})\right) + \underbrace{\left(\frac{n-1}{n} - \frac{n-2}{n-1} \sum_{j \in N_{2}} m_{i}(X_{j})\right)}_{\geq 0}$$

$$\geq \frac{1}{n} - \frac{1}{n-1} \sum_{j \in N_{2}} m_{i}(X_{j})$$

$$= \frac{1}{n} - \frac{1}{n-1} \sum_{j \in N \setminus i} m_{i}(X_{j}),$$

where the inequality follows from $\frac{n-1}{n} \ge \frac{n-2}{n-1} \ge \frac{n-2}{n-1} \sum_{j \in N_2} m_i(X_j)$. Case 3: $i \in N_1$ and $l \ge 2$

Since (X_1, \ldots, X_l) is a **PROPavg** allocation of M_1 to N_1 , we have

$$v_{i}(X_{i}) \geq \frac{v_{i}(M_{1})}{l} - \frac{1}{l-1} \sum_{j \in N_{1} \setminus i} m_{i}(X_{j})$$

= $\frac{1}{l} - \frac{1}{l} \sum_{j \in N_{2}} m_{i}(X_{j}) - \frac{1}{l-1} \sum_{j \in N_{1} \setminus i} m_{i}(X_{j}).$ (5)

In line 13 of Algorithm 2, the while condition in line 7 does not hold for agent i. Thus, it holds that

$$m_i(X_j) < \frac{1}{n} - \frac{1}{n-1} \sum_{j' \in N_2} m_i(X_{j'})$$
 (6)

for any $j \in N_1 \setminus i$. Summing up inequality (6) for each $j \in N_1 \setminus i$, we obtain

$$\sum_{j \in N_1 \setminus i} m_i(X_j) < \frac{l-1}{n} - \frac{l-1}{n-1} \sum_{j \in N_2} m_i(X_j).$$
(7)

By multiplying inequality (7) by $\frac{n-l}{l(l-1)} > 0$ and rearranging, we have

$$0 > -\frac{n-l}{ln} + \frac{n-l}{l(n-1)} \sum_{j \in N_2} m_i(X_j) + \frac{n-l}{l(l-1)} \sum_{j \in N_1 \setminus i} m_i(X_j).$$
(8)

Summing up inequalities (5) and (8), we have

$$v_i(X_i) > \frac{1}{n} + \left(-\frac{1}{l} + \frac{n-l}{l(n-1)}\right) \sum_{j \in N_2} m_i(X_j) + \left(-\frac{1}{l-1} + \frac{n-l}{l(l-1)}\right) \sum_{j \in N_1 \setminus i} m_i(X_j).$$
(9)

Since $2 \leq l < n$, by direct calculation, we have

$$-\frac{1}{l} + \frac{n-l}{l(n-1)} + \frac{1}{n-1} = \frac{1}{l(n-1)} \ge 0$$

and

$$\begin{aligned} -\frac{1}{l-1} + \frac{n-l}{l(l-1)} + \frac{1}{n-1} &= \frac{1}{l(l-1)(n-1)} \left(-l(n-1) + (n-l)(n-1) + l(l-1) \right) \\ &= \frac{(n-l)(n-l-1)}{l(l-1)(n-1)} \\ &\ge 0. \end{aligned}$$

Applying these inequalities to inequality (9), we finally obtain

$$v_i(X_i) > \frac{1}{n} - \frac{1}{n-1} \left(\sum_{j \in N_2} m_i(X_j) + \sum_{j \in N_1 \setminus i} m_i(X_j) \right).$$

= $\frac{1}{n} - \frac{1}{n-1} \sum_{j \in N \setminus i} m_i(X_j),$

which implies that i is PROPavg-satisfied by X.

Therefore, X is a PROPavg allocation to N in line 16 of Algorithm 2.

We finally give the proof of Theorem 2 by showing that Algorithm 2 is a polynomial time algorithm to find a PROPavg allocation.

Proof of Theorem 2. We first show the correctness of Algorithm 2. If |N| = 1, our algorithm obviously returns a PROPavg allocation in line 3. Assume that $|N| \ge 2$. If $|N_2| \ge 1$, it returns a PROPavg allocation in line 14 by Lemma 17. Otherwise, since the while condition in line 7 does not hold for any agent in N_1 , $v_i(g) < \frac{1}{n}$ holds for any agent $i \in N_1$ and good $g \in M_1$ in line 16. Thus, $X = (X_1, \ldots, X_n)$ satisfies (P1) by Lemma 16, where X_n is an empty set. The rest of the algorithm finds an allocation to V_2 satisfying (P2) and returns a PROPavg allocation as in Algorithm 1. Therefore, Algorithm 2 returns a PROPavg allocation in all cases.

We finally show that Algorithm 2 completes in time polynomial in the number of agents and items. Let T(n, m) be the worst case time complexity of Algorithm 2 when |N| = n and |M| = m. Clearly, T(1, m) = O(m). We can check the while condition in line 7 and execute the body of the while loop in polynomial time of n and m. In addition, as mentioned at the beginning of Section 5, Lines 17 to 22 can be executed in polynomial time of n and m. Thus, T(n,m) can be expressed as

$$T(n,m) = O(\text{poly}(n,m)) + \max\{\max_{\substack{1 \le n' \le n-1 \\ 1 \le m' \le m-1}} T(n',m'), T(n-1,m)\}.$$

Therefore, T(n, m) is polynomially bounded in n and m.

6 Conclusion

In this paper, we have introduced PROPavg, which is a stronger notion than PROPm, and shown that a PROPavg allocation always exists and can be computed in polynomial time when each agent has a non-negative additive valuation. In order to devise our algorithm, we have developed a new technique that generalizes the cut-and-choose protocol. This technique is interesting by itself and seems to have a potential for further applications. In fact, we can define a bipartite graph like the PROPavg-graph for another fairness notion, and our argument works if we obtain an allocation satisfying a (P2)-like condition. We expect that this technique will be used in other contexts as well.

There are still several future work. Whether Avg-EFX, which is a stronger notion than PROPavg exists for four or more agents is an interesting open problem. Another direction is to consider weighted approximate proportionality. In the weighted case, each agent i has a non-negative weight α_i , where $\alpha_1 + \cdots + \alpha_n = 1$. The goal is to find an allocation X such that $v_i(X_i) \geq \frac{\alpha_i}{n} - d_i(X)$ for each agent $i \in N$. Whether weighted PROPavg allocation exists is also an interesting problem.

Acknowledgments

This work was partially supported by the joint project of Kyoto University and Toyota Motor Corporation, titled "Advanced Mathematical Science for Mobility Society", and by JSPS, KAKENHI grant number JP19H05485, Japan.

References

- Georgios Amanatidis, Georgios Birmpas, Aris Filos-Ratsikas, Alexandros Hollender, and Alexandros A Voudouris. Maximum Nash welfare and other stories about EFX. *Theoretical Computer Science*, 863:69–85, 2021.
- [2] Georgios Amanatidis, Georgios Birmpas, Aris Filos-Ratsikas, and Alexandros A. Voudouris. Fair division of indivisible goods: A survey. In Proceedings of the 31st International Joint Conference on Artificial Intelligence (IJCAI), pages 5385–5393, 2022.
- [3] Haris Aziz, Hervé Moulin, and Fedor Sandomirskiy. A polynomial-time algorithm for computing a Pareto optimal and almost proportional allocation. Operations Research Letters, 48(5):573–578, 2020.
- [4] Artem Baklanov, Pranav Garimidi, Vasilis Gkatzelis, and Daniel Schoepflin. Achieving proportionality up to the maximin item with indivisible goods. In *Proceedings of the* AAAI Conference on Artificial Intelligence, volume 35, pages 5143–5150, 2021.

- [5] Artem Baklanov, Pranav Garimidi, Vasilis Gkatzelis, and Daniel Schoepflin. PROPm allocations of indivisible goods to multiple agents. In *Proceedings of the 30th International Joint Conference on Artificial Intelligence (IJCAI)*, pages 24–30, 2021.
- [6] Siddharth Barman, Sanath Kumar Krishnamurthy, and Rohit Vaish. Greedy algorithms for maximizing Nash social welfare. In Proceedings of the 17th International Conference on Autonomous Agents and MultiAgent Systems (AAMAS), pages 7–13, 2018.
- [7] Ben Berger, Avi Cohen, Michal Feldman, and Amos Fiat. Almost full EFX exists for four agents. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 36, pages 4826–4833, 2022.
- [8] Steven J Brams, Steven John Brams, and Alan D Taylor. *Fair Division: From cake-cutting to dispute resolution*. Cambridge University Press, 1996.
- [9] Felix Brandt, Vincent Conitzer, Ulle Endriss, Jérôme Lang, and Ariel D Procaccia. Handbook of computational social choice. Cambridge University Press, 2016.
- [10] Eric Budish. The combinatorial assignment problem: Approximate competitive equilibrium from equal incomes. *Journal of Political Economy*, 119(6):1061–1103, 2011.
- [11] Ioannis Caragiannis, Nick Gravin, and Xin Huang. Envy-freeness up to any item with high Nash welfare: The virtue of donating items. In *Proceedings of the 20th ACM Conference on Economics and Computation (EC)*, pages 527–545, 2019.
- [12] Ioannis Caragiannis, David Kurokawa, Hervé Moulin, Ariel D. Procaccia, Nisarg Shah, and Junxing Wang. The unreasonable fairness of maximum Nash welfare. ACM Transactions on Economics and Computation, 7(3):1–32, 2019.
- [13] Bhaskar Ray Chaudhury, Jugal Garg, and Kurt Mehlhorn. EFX exists for three agents. In Proceedings of the 21st ACM Conference on Economics and Computation (EC), pages 1–19, 2020.
- [14] Bhaskar Ray Chaudhury, Jugal Garg, Kurt Mehlhorn, Ruta Mehta, and Pranabendu Misra. Improving EFX guarantees through rainbow cycle number. In Proceedings of the 22nd ACM Conference on Economics and Computation (EC), pages 310–311, 2021.
- [15] Bhaskar Ray Chaudhury, Telikepalli Kavitha, Kurt Mehlhorn, and Alkmini Sgouritsa. A little charity guarantees almost envy-freeness. SIAM Journal on Computing, 50(4):1336– 1358, 2021.
- [16] Vincent Conitzer, Rupert Freeman, and Nisarg Shah. Fair public decision making. In Proceedings of the 18th ACM Conference on Economics and Computation (EC), pages 629–646, 2017.

- [17] Andreas Darmann and Joachim Schauer. Maximizing Nash product social welfare in allocating indivisible goods. *European Journal of Operational Research*, 247(2):548–559, 2015.
- [18] Andrew L Dulmage. A structure theory of bipartite graphs of finite exterior dimension. The Transactions of the Royal Society of Canada, Section III, 53:1–13, 1959.
- [19] Andrew L Dulmage and Nathan S Mendelsohn. Coverings of bipartite graphs. Canadian Journal of Mathematics, 10:517–534, 1958.
- [20] P Hall. On representatives of subsets. Journal of the London Mathematical Society, 1(1):26–30, 1935.
- [21] Yusuke Kobayashi and Ryoga Mahara. Proportional Allocation of Indivisible Goods up to the Least Valued Good on Average. In 33rd International Symposium on Algorithms and Computation (ISAAC), pages 55:1–55:13, 2022.
- [22] Richard J. Lipton, Evangelos Markakis, Elchanan Mossel, and Amin Saberi. On approximately fair allocations of indivisible goods. In *Proceedings of the 5th ACM Conference* on *Electronic Commerce (EC)*, pages 125–131, 2004.
- [23] Ryoga Mahara. Existence of EFX for two additive valuations. arXiv preprint arXiv:2008.08798, 2020.
- [24] Ryoga Mahara. Extension of additive valuations to general valuations on the existence of EFX. In Proceedings of the 29th Annual European Symposium on Algorithms (ESA), pages 66:1–15, 2021.
- [25] Hervé Moulin. Fair division and collective welfare. MIT press, 2004.
- [26] Hervé Moulin. Fair division in the internet age. Annual Review of Economics, 11:407– 441, 2019.
- [27] Benjamin Plaut and Tim Roughgarden. Almost envy-freeness with general valuations. SIAM Journal on Discrete Mathematics, 34(2):1039–1068, 2020.
- [28] Jack Robertson and William Webb. Cake-cutting algorithms: Be fair if you can, 1998.
- [29] Hugo Steinhaus. The problem of fair division. *Econometrica*, 16(1):101–104, 1948.