

# On quasi-nodal spheres and the spin Hall effect: the case of $\text{YH}_3$ and $\text{CaTe}$ .

Rafael González-Hernández,<sup>1,2,\*</sup> Carlos Pinilla,<sup>1,3,†</sup> and Bernardo Uribe<sup>4,5,‡</sup>

<sup>1</sup>*Departamento de Física y Geociencias, Universidad del Norte,*

*Km. 5 Vía Antigua Puerto Colombia, Barranquilla 080020, Colombia*

<sup>2</sup>*Institut für Physik, Johannes Gutenberg Universität Mainz, D-55099 Mainz, Germany*

<sup>3</sup>*School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, United Kingdom*

<sup>4</sup>*Departamento de Matemáticas y Estadística, Universidad del Norte,*

*Km. 5 Vía Antigua Puerto Colombia, Barranquilla 080020, Colombia*

<sup>5</sup>*Max Planck Institut fuer Mathematik, Vivatsgasse 7, 53115, Bonn, Germany*

(Dated: September 26, 2022)

Band inversion is a known feature in a wide range of topological insulators characterized by a change of orbital type around a high symmetry point close to the Fermi level. In some cases of band inversion in topological insulators, the existence of quasi-nodal spheres has been detected, and the change of orbital type is shown to be concentrated along these spheres in momentum space. In order to understand this phenomenon, we develop a local effective four-fold Hamiltonian which models the band inversion and reproduces the quasi-nodal sphere. This model shows that the signal of the spin Hall conductivity, as well as the change of orbital type, are both localized on the quasi-nodal sphere, and moreover, that these two indicators characterize the topological nature of the material. Using K-theoretical methods we show that the change of orbital type parametrized by an odd clutching function is equivalent to the strong Fu-Kane-Mele invariant. We corroborate these results with *ab-initio* calculations for the materials  $\text{YH}_3$  and  $\text{CaTe}$  where in both cases the signal of the spin Hall conductivity is localized on the quasi-nodal spheres in momentum space. We conclude that a non-trivial spin Hall conductivity localized on the points of change of orbital type is a good indicator for Topological Insulation.

## INTRODUCTION

The incursion of topological invariants in condensed matter physics has led to an enhanced classification of quantum materials [1–4]. Among the insulators, the topological ones (TI) have attracted significant attention due to the presence of conducting surface states and because they show efficient spin transport properties [5–8]. The presence of these unique surface states, together with the topological Fu-Kane-Mele invariant, have been used to characterize these topological insulator phases from normal insulators systems [9–11].

The topological order of some insulators may be alternatively deduced from the change of orbital type on the last valence bands around a time-reversal invariant point (TRIM). This feature is known in the literature as “Band inversion” and has been extensively used to classify TIs [12–14]. Band inversion is assumed to be induced from band splitting due to strong spin-orbit coupling (SOC) interaction from heavy elements in materials [15–19]. In these systems, SOC can have a significant impact on the band structure and induces an opening gap between the conduction and valence bands thus changing the orbital type. This band inversion may be coupled with a change in the inversion symmetry eigenvalues at the appropriate TRIM. It was this precise feature that led to the establishment of the Fu-Kane-Mele invariant in topological insulators with inversion symmetry as  $\text{Bi}_2\text{Se}_3$ ,  $\text{Bi}_2\text{Te}_3$  and  $\text{Sb}_2\text{Te}_3$  [20, 21]. However, some materials show that the band inversion can appear spontaneously, or may be induced by strain, even without SOC interaction [13, 22].

This phenomenon can be observed in  $\alpha\text{-Sn}$ ,  $\text{HgTe}$ , several half-Heusler and chalcopyrite semiconductors, where the band inversion is presented at a particular TRIM and independent of the SOC interaction [23–27]

The type of Band inversion which is of interest for this work appears whenever there is a hybridization of the energy bands, and the change of orbital type is concentrated on the  $\mathbf{k}$ -states where the energy gap is opened. This form of band inversion is present, among others, on the trihydrides materials and it has been previously reported in references [28, 29]. On these materials a small energy gap is present and the  $\mathbf{k}$ -states with a small energy gap form a 2-dimensional sphere. In [29] these spheres were called “Pseudo Dirac nodal spheres” (PDNS) and here we have denoted them “Quasi-nodal spheres”. In the pioneer work by Wang et al. [29], the possible formation of PDNS in crystal structures is reported on different symmetry point groups through the study of band crossings with pairs of 1D irreducible representations. It is also found that the PDNS phase is robust against the SOC effect, in particular for the  $M\text{H}_3$  (with  $M=\text{Y}, \text{Ho}, \text{Tb}, \text{Nd}$ ) and  $\text{Si}_3\text{N}_2$  materials. In addition, the prediction of a realizable topological state with exotic transport properties is suggested for these PDNS prototypes [29]. The analysis of the topological nature and the spin transport properties of the PDNS (or quasi-nodal spheres) remained an open question and it has been one of the motivations of the present work.

In this work, we study the mechanism underlying the band inversion occurring through hybridization as it happens on the trihydrides. We argue that the change of or-

bital type is concentrated on the quasi-nodal sphere, and moreover, that the Fu-Kane-Mele invariant characterizing the TI property of the material can be deduced from the non-triviality of the spin Hall conductivity (SHC) and the localization of its signal on the quasi-nodal sphere. This fact explains the 2-dimensional nature of the  $\mathbf{k}$ -points where hybridization is happening and shows how the Fu-Kane-Mele invariant relates to the SHC and the change of orbital type.

The paper contains two main parts, the theoretical analysis and the material realization. In the first part, we analyze the mechanism of band inversion by using a four-fold effective Hamiltonian for a system where the orbital character change is present, the quasi-nodal sphere is induced, and both the signal of the SHC and the orbital type change are concentrated on the quasi-nodal sphere. Here we show explicitly how the valence states of the effective Hamiltonian define an element in the appropriate K-theory, and how this K-theoretical element matches the K-theoretical version of the strong Fu-Kane-Mele invariant. In the second part, we investigate two possible materials, yttrium trihydride ( $\text{YH}_3$ ) and calcium telluride ( $\text{CaTe}$ ), which show an efficient spin/change conversion generated from the band inversion on the quasi-nodal sphere.

## BAND INVERSION

Let us consider a Hamiltonian with time reversal symmetry  $\mathbb{T}$  and inversion symmetry  $\mathcal{I}$  where spin orbit coupling is taken into account. Band inversion could be understood as the change of orbital character on the last valence bands around a time reversal invariant point (TRIM). This change of character, whenever coupled with a change of eigenvalue of the inversion operator, determines the strong topological insulator nature of the material.

In the case of interest, the change of orbital type is occurring on a sphere of points where the energy gap is small. This 2-dimensional sphere in momentum space with small energy gap (comparable to room temperature  $\sim 25$  meV) has been coined **Quasi-nodal Sphere**, generalizing the concept of quasi-nodal line that was presented by the first and third authors in [30].

In what follows we argue that band inversion due to hybridization produces a quasi-nodal sphere on momentum space where orbital characters are mixed, and moreover, where the Fu-Kane-Mele invariant can be extracted. We first define a local model for the low energy effective Hamiltonian inducing band inversion due to hybridization, we calculate its quasi-nodal sphere together with a  $2 \times 2$  matrix information on each point obtained from the change of orbital type on the valence bands, and finally, we relate these parametrized matrices to the topological invariants obtained through K-theory.

## Model Hamiltonian

Take the Pauli matrices  $\tau_i$  and  $\sigma_j$  in orbital and spin coordinates respectively, and consider the Hamiltonian

$$H(\mathbf{k}) = M(\mathbf{k})\tau_3\sigma_0 + A(\mathbf{k})\tau_1\sigma_3 + B(\mathbf{k})\tau_2\sigma_0 \quad (1)$$

where

$$M(\mathbf{k}) = D_1 - m_1 k_z^2 - n_1(k_x^2 + k_y^2), \quad (2)$$

$$A(\mathbf{k}) = D_2 k_z + E_2 k_z^3 + F_2(3k_x^2 k_y - k_y^3), \quad (3)$$

$$B(\mathbf{k}) = D_3(k_x^3 - k_x k_y^2). \quad (4)$$

This Hamiltonian preserves time reversal, inversion and a three-fold rotation with matrices

$$\mathbb{T} = i\tau_0\sigma_2\mathbb{K}, \quad \mathcal{I} = -\tau_3\sigma_0, \quad \text{and} \quad C_3 = e^{i\tau_0\sigma_3\frac{\pi}{3}} \quad (5)$$

respectively. Here  $C_3$  is a three-fold rotation around the  $z$ -axis. The bands are double-degenerate due to the presence of both time reversal and inversion, and the energies of the bands are:

$$\pm E(\mathbf{k}) = \pm \sqrt{M(\mathbf{k})^2 + A(\mathbf{k})^2 + B(\mathbf{k})^2}. \quad (6)$$

Tuning up the constants to locally model the band inversion present in  $\text{YH}_3$  at the point  $\Gamma$ , we set the coefficients to the following values:

$$\begin{aligned} D_1 &= 0.2, m_1 = n_1 = 18, \\ D_2 &= 0.1, E_2 = F_2 = D_3 = 5. \end{aligned} \quad (7)$$

The energy bands of the Hamiltonian are presented in Fig. 1 a) and b) together with their projections on the first orbital. There is no symmetry protecting energy crossings and therefore the eigenstates hybridize producing an energy gap. The change of orbital type is concentrated along the points in momentum space whose energy gap is small, and they define a 2-dimensional sphere as can be seen in Fig. 1 d). This 2-dimensional sphere is a quasi-nodal sphere and it is necessary for the band inversion that induces the Hamiltonian. Note that the inversion operator  $\mathcal{I} = -\tau_3\sigma_0$  is coupled with the orbital types since it acts by  $-1$  on the first orbital and by  $+1$  on the second orbital (see Fig. 1 a). The interdependence of the inversion operator with the orbit type is what permits to distinguish the TI nature of the material by the change of eigenvalues of the inversion. Nevertheless, the TI nature of the material is kept while inversion is broken, thus the change of orbital type, together with the SHC is what allows to distinguish its topological nature. From Fig. 1 d) we see that the valence bands are concentrated on the second orbital inside the quasi-nodal sphere while concentrated on the first orbital outside of it.

The change of orbital type of the valence bands along the quasi-nodal sphere permits to define a  $2 \times 2$  unitary complex matrix for each point on the nodal-sphere. This




FIG. 1. Energy bands of the Hamiltonian of Eqn. (1) with projection to the first orbital. a). Top panel: valence and conduction bands on  $k_x$  coordinate axis with projection on the first orbital. The eigenvalues of the inversion operator at  $(0, 0, 0)$  on the valence bands are  $+1$  and on the conduction bands are  $-1$ . Bottom panel: sum of the spin Berry curvatures of the two valence bands  $\Omega_{xy}^z$ . b) Top panel: close up view of the energy bands where the change of orbital type is concentrated. Bottom panel: close up view of the sum of the spin Berry curvatures  $\Omega_{xy}^z$ . The signal of the the spin Berry curvature is concentrated where the change of orbital type occurs. c) Valence band on the plane  $k_z = 0$  with projection on the first orbital. d)  $\mathbf{k}$ -states with mixed orbital type (more than 3% of each orbital type,  $k_y \leq 0$ ) equivalent to the quasi-nodal sphere with energy gap of less than 0.05eV. The color represents the magnitude of the projection on the first orbital. Inside the quasi-nodal sphere the states are concentrated on the first orbital, while outside are concentrated on the second orbital.

assignment produces a map from the sphere  $S^2$  to the Lie group of special unitary matrices  $SU(2)$  which is equivariant with respect to the inversion operator. Here the inversion operator acts by the antipodal action on the sphere and by multiplication by  $-1$  on  $SU(2)$ . This map can be understood as the clutching map (in the words of Atiyah [31, pp.20]) that defines the rank 2 complex vector bundle over the 3-dimensional sphere which produces the non-trivial Fu-Kane-Mele invariant. Let us postpone the construction of the Fu-Kane-Mele invariant on the 3-dimensional sphere to the next section and let us show how the matrices are defined.

Denote by  $\varphi_{\pm}^i(\mathbf{k})$ ,  $i \in \{1, 2\}$ , the four eigenvectors of the Hamiltonian of Eqn. (1) with the fixed parameters described in Eqns. (7) satisfying the equation:

$$H(\mathbf{k})\varphi_{\pm}^i(\mathbf{k}) = \pm E(\mathbf{k})\varphi_{\pm}^i(\mathbf{k}). \quad (8)$$

Note that the valence states  $\varphi_{-}^i(\mathbf{k})$  are well defined for all  $\mathbf{k}$  since the system is gapped, and therefore we may take the vector projections of each valence state to the first and second orbital respectively. The eigenstates have four coordinates

$$\varphi_{-}^i = (\varphi_{-,1}^i, \varphi_{-,2}^i, \varphi_{-,3}^i, \varphi_{-,4}^i) \in \mathbb{C}^4 \quad (9)$$

and the projection on the first orbital takes the first two coordinates and the projection on the second takes the last two. Denoting  $pr_j$  the projection on the  $j$ -th orbital we have:

$$pr_1\varphi_{-}^i = (\varphi_{-,1}^i, \varphi_{-,2}^i), \quad (10)$$

$$pr_2\varphi_{-}^i = (\varphi_{-,3}^i, \varphi_{-,4}^i). \quad (11)$$

Along the quasi-nodal sphere both projections are non-trivial (we may take the sphere of radius  $\|\mathbf{k}\| = 0.11$ , see Fig. (1) c)), and therefore we may define the following matrix coefficients:

$$A_{ij}(\mathbf{k}) = (-1)^i \left\langle \frac{pr_1\varphi_{-}^i(\mathbf{k})}{\|pr_1\varphi_{-}^i(\mathbf{k})\|} \left| \frac{pr_2\varphi_{-}^j(\mathbf{k})}{\|pr_2\varphi_{-}^j(\mathbf{k})\|} \right. \right\rangle \quad (12)$$

for  $i, j \in \{1, 2\}$ . Since the composition  $\mathbb{T}\mathcal{I}$  commutes with the Hamiltonian, we may consider  $\varphi_{-}^2$  to be the Kramer's pair of  $\varphi_{-}^1$  satisfying the equation

$$\varphi_{-}^2 = \mathbb{T}\mathcal{I}\varphi_{-}^1. \quad (13)$$

From Eqns. (5) we know that  $\mathbb{T}\mathcal{I} = -i\tau_3\sigma_2\mathbb{K}$  and therefore Eqn. (13) implies the following equality:

$$(\varphi_{-,1}^2, \varphi_{-,2}^2, \varphi_{-,3}^2, \varphi_{-,4}^2) = (-\bar{\varphi}_{-,2}^1, \bar{\varphi}_{-,1}^1, -\bar{\varphi}_{-,4}^1, \bar{\varphi}_{-,3}^1). \quad (14)$$

Replacing Eqn. (14) on the definition of the matrix coefficients of Eqn. (12) we see that the matrix  $A$  is unitary with  $A_{11} = \bar{A}_{22}$ ,  $A_{12} = -\bar{A}_{21}$ , and its determinant is 1. Hence the matrix  $A$  belongs to the Lie group  $SU(2)$  of special unitary matrices.

If we take  $S_r^2 = \{\mathbf{k}: \|\mathbf{k}\| = r\}$  to be the quasi-nodal sphere of the system ( $r = 0.11$  for the Hamiltonian with parameters in Eqns. (7)), we obtain a map

$$A : S_r^2 \rightarrow SU(2), \quad \mathbf{k} \mapsto A(\mathbf{k}). \quad (15)$$

The explicit form of the inversion operator  $\mathcal{I} = -\tau_3\sigma_0$  implies the following equation:

$$pr_j\varphi_{-}^i(\mathbf{k}) = (-1)^j pr_j\varphi_{-}^i(-\mathbf{k}), \quad j \in \{1, 2\}, \quad (16)$$

which makes the map  $A$  equivariant with respect to the inversion action, i.e.

$$A(-\mathbf{k}) = -A(\mathbf{k}). \quad (17)$$

Any equivariant map such as  $A$  produces the Fu-Kane-Mele invariant since any two maps satisfying Eqn. (17) are homotopic [32, Lem. 3.27]. In particular the map  $A$  is homotopic to the map  $C : S_r^2 \rightarrow SU(2)$ ,

$$C(\mathbf{k}) = \begin{pmatrix} ik_z & k_x + ik_y \\ -k_x + ik_y & -ik_z \end{pmatrix}, \quad (18)$$

which is a simple clutching function that defines the rank 2 bundle with non-trivial Fu-Kane-Mele invariant.

Another important feature of the quasi-nodal sphere of this Hamiltonian is the fact that the signal for the spin Hall conductivity  $\Omega_{xy}^z$  localizes around it. In Fig. 1 d) we have plotted the points on which the orbital type is mixed, i.e.  $\mathbf{k}$ -points with more than 3% on each orbital

$$0.03 < ||pr_1\varphi_-^1(\mathbf{k})|| < 0.97, \quad (19)$$

and we notice that these mixed states define the quasi-nodal sphere. From Fig. 1 b) we notice that the signal of the SHC is localized on the  $\mathbf{k}$ -points with mixed orbital type. This feature could be appreciated on the materials  $\text{YH}_3$  and  $\text{CaTe}$  on Fig. 3 d), f), g) and Fig. 4 d), f), g) respectively.

The topological invariant that the Hamiltonian of Eqn. (1) defines can be better understood with the help of K-theory. This is the subject of the next section.

### K-theory, clutching functions and sewing matrices

Restrict the Hamiltonian to a ball  $B$  centered at the origin which includes the quasi-nodal sphere. The valence eigenstates define a rank 2 complex vector bundle over  $B$  which incorporates the clutching function  $A$ . This bundle possesses the topological information to capture the Fu-Kane-Mele invariant, but unfortunately it does not trivialize on the boundary of  $B$  in order to define an appropriate element in K-theory (the Hamiltonian of Eqn. (1) provided only a local model and it was not defined over a compact space). To overcome this issue we will construct a rank 2 complex vector bundle over the sphere  $S^3$  which models the structure of the valence eigenstates of the Hamiltonian of Eqn. (1) and we will show that this vector bundle incorporates the Fu-Kane-Mele invariant for K-theory. This bundle will model the K-theoretical properties of the valence states, and will permit us to calculate its topological invariants.

Consider the 3-dimensional sphere

$$S^3 = \{(t, k_1, k_2, k_3) | t^2 + ||k||^2 = 1\}, \quad (20)$$

where  $S^3 \setminus \{(-1, \mathbf{0})\}$  could be thought as the stereographic projection of the ball  $B$  and  $S^3$  its one-point

compactification. Take  $\rho : [-1, 1] \rightarrow [0, 1]$  a partition of unity on the interval  $[-1, 1]$  with  $\rho|_{[-1, -1+\varepsilon]} = 0$  and  $\rho|_{(1-\varepsilon, 1]} = 1$  for small  $\varepsilon$ . Define the rank 2 vector bundle  $E \subset S^3 \times (\mathbb{C}^2)^2$  by the equation

$$E = \{[(t, \mathbf{k}), (\rho(t)u, (1 - \rho(t))C(\mathbf{k})u)] \mid u \in \mathbb{C}^2\}, \quad (21)$$

and define the actions of  $\mathcal{I}$  and  $\mathbb{T}$  as in Eqns. (5) by the following formulas:

$$\mathcal{I} \cdot ((t, \mathbf{k}), (u, v)) = ((t, -\mathbf{k}), (-u, v)), \quad (22)$$

$$\mathbb{T} \cdot ((t, \mathbf{k}), (u, v)) = ((t, -\mathbf{k}), (\mathbb{J}u, \mathbb{J}v)). \quad (23)$$

Here  $\mathbb{J} = i\tau_2\mathbb{K}$  commutes with the matrices in  $SU(2)$  and therefore the action of  $\mathbb{T}$  is well-defined. Moreover, the partition of unity  $\rho$  could be understood as the projection map on the first orbital while  $(1 - \rho)$  as the projection on the second.

When the action of  $\mathcal{I}$  is disregarded, the relative bundle  $[E] - [S^3 \times \mathbb{C}^2]$  generates the only non-trivial class on the appropriate relative K-theory groups. Choosing as base point  $(-1, \mathbf{0})$  and following the notation of [32, App. C], we have that

$$[E] - [S^3 \times \mathbb{C}^2] \in \widetilde{KQ}^0(S^3) \cong \mathbb{Z}/2, \quad (24)$$

and therefore the bundle  $E$  induces the strong Fu-Kane-Mele invariant. Here  $KQ^{n+4} \cong KR^n$  where  $KR^*$  is Atiyah's real K-theory [33], and

$$\widetilde{KQ}^0(S^3) \cong \widetilde{KR}^4(S^3) \cong KR^7(*) \cong KO^{-1} \cong \mathbb{Z}/2. \quad (25)$$

Taking into account the involution  $\mathcal{I}$ , the relative bundle  $[E] - [S^3 \times \mathbb{C}^2]$  gives the generator of the relative equivariant symplectic K-theory groups:

$$[E] - [S^3 \times \mathbb{C}^2] \in \widetilde{KSp}_{(\mathcal{I})}^0(S^3) \cong \mathbb{Z}. \quad (26)$$

Here the composition  $\mathcal{I}\mathbb{T}$  defines the quaternionic structure on the fibers of bundle and  $\mathcal{I}$  induces a  $\mathbb{Z}/2$ -equivariant quaternionic action on the bundle. The groups  $\widetilde{KSp}_{(\mathcal{I})}^0(S^3)$  are the reduced  $\mathbb{Z}/2$ -equivariant symplectic K-theory groups of the sphere  $S^3$  [34], cf. [35]. The restriction map to the point  $(1, \mathbf{0})$  counts the number of quaternionic  $\mathbb{Z}/2$ -representations with  $\pm 1$ -eigenvalues:

$$\widetilde{KSp}_{(\mathcal{I})}^0(S^3) \rightarrow \mathbb{Z}_{-1} \oplus \mathbb{Z}_{+1} \cong KSp_{(\mathcal{I})}^0(\{(1, \mathbf{0})\}), \quad (27)$$

$$E \mapsto 1 \oplus 0, \quad (28)$$

where  $\mathbb{Z}_{-1}$  counts the non-trivial ones and  $\mathbb{Z}_{+1}$  the trivial ones. Since this restriction map is injective, the topological class of the bundle can be determined by the eigenvalues of  $\mathcal{I}$  on the fixed point set of the action. Since  $E$  has the non-trivial quaternionic  $\mathbb{Z}/2$ -representation on the fixed point  $(1, \mathbf{0})$  and the trivial one on  $(-1, \mathbf{0})$ , the strong Fu-Kane-Mele invariant can be determined by the




FIG. 2. Spin Berry curvature  $\Omega_{xy}^z$  at the Fermi level with respect to  $k_x$  of the Hamiltonian of Eqn. (1). The energy gap is widened by setting the structural constants  $E_2, F_2, D_3$  of the Hamiltonian to the following values: a)  $E_2 = F_2 = D_3 = 250$  b)  $E_2 = F_2 = D_3 = 50$  and c)  $E_2 = F_2 = D_3 = 5$ . The values of c) are the original values from Eqn. (7). The signal of the SHC is concentrated on the region where the orbital type changes. Note that despite the fact that the difference of energies appears on the denominator of the Kubo formula of the SHC (see Eqn. (36)), the signal of the spin Berry curvature localizes whenever the change of orbit type occurs.

parity of the number of pairs of complex  $-1$  eigenvalues on the fixed points of the  $\mathcal{I}$  action; this is the main result of Fu et al. [9].

The restriction of the bundle  $E$  to the sphere of points with  $t = 0$ , permits us to define the transformation from the first to the second orbital, and this transformation is precisely the function  $C$  presented in Eqn. (18). This clutching function is shown in [32, Cor. 4.1] to induce the Fu-Kane-Mele invariant.

The bundle  $E$  models the topological structure of the valence states of the Hamiltonian of Eqn. (1) and it induces the topological generators in both the equivariant symplectic K-theory and the quaternionic K-theory of involution spaces. Hence the bundle  $E$  models the topological nature of the band inversion and it induces the strong Fu-Kane-Mele invariant [9]. The change of eigenvalues of the inversion operator on the fixed points on the sphere  $S^3$  is only possible whenever the clutching function  $C$  is defined on the sphere  $S^2$  and it is odd (Eqn. (17)).

The relation of the clutching functions with the SHC can be seen in Fig. 2. The signal of the SHC concentrates on the points where the change of orbital type occurs despite the size of the energy gap. We can therefore conclude that not only the  $\mathbf{k}$ -points with mixed orbital type make the quasi-nodal sphere, but moreover, that the signal of the spin Hall conductivity localize on this quasi-nodal sphere.

We can go one step further and we may calculate the sewing matrices of the inversion and the time reversal operator. For this we need to take a trivialization of the bundle  $E$  (since any rank 2 complex bundle over  $S^3$  is trivializable), and we may write the inversion and the

time reversal operator on the new basis. Now we have  $E = S^3 \times \mathbb{C}^2$  and the actions become

$$\mathcal{I} \cdot ((t, \mathbf{k}), u) = ((t, -\mathbf{k}), (t\text{Id} + C(\mathbf{k}))u), \quad (29)$$

$$\mathbb{T} \cdot ((t, \mathbf{k}), u) = ((t, -\mathbf{k}), (t\text{Id} + C(\mathbf{k}))\mathbb{J}u). \quad (30)$$

For  $(e_1, e_2)$  the canonical basis on  $\mathbb{C}^2$ , the sewing matrices for the operator  $P$  on  $E$  are defined by the equation

$$G(P)_{ij}(t, \mathbf{k}) = \langle e_i | P(t, \mathbf{k}) | e_j \rangle. \quad (31)$$

For the inversion and the time reversal operator the sewing matrices can be described as follows

$$G(I)(t, \mathbf{k}) = t\text{Id} + C(\mathbf{k}), \quad (32)$$

$$G(\mathbb{T})(t, \mathbf{k}) = (t\text{Id} + C(\mathbf{k}))\mathbb{J}. \quad (33)$$

In both cases the sewing matrices define maps from  $S^3$  to  $SU(2)$  whose degree is  $\pm 1$ . From [36–38] we know that the parity of the degree of those maps recovers the Chern-Simons axion coupling term

$$\theta = \frac{1}{4\pi} \int_{\text{BZ}} dk^3 \epsilon^{\alpha\beta\gamma} \text{Tr} \left( \mathcal{A}_\alpha \partial_\beta \mathcal{A}_\gamma - \frac{2i}{3} \mathcal{A}_\alpha \mathcal{A}_\beta \mathcal{A}_\gamma \right). \quad (34)$$

Since the parity of the degrees of both  $G(I)$  and  $G(\mathbb{T})$  is odd, then we know that  $\theta = \pi$ . Hence we see the topological nature of the trivial complex bundle  $E$  whenever the action of either  $I$  or  $\mathbb{T}$  are defined as in Eqns. (29), (30).

Note that whenever  $t = 0$ , both the clutching function of the bundle of Eqn. (21) and the sewing matrices of the inversion operator agree. The topological nature of the bundle could be theoretically deduced then from either one. Unfortunately both approaches are not well suited for computational calculations. On the one hand the definition of the clutching functions is not gauge invariant, and on the other, the calculation of the Chern-Simons coupling term has been until now elusive for real materials [39]. We hope in the future new methods are developed to calculate both.

## MATERIAL REALIZATION

### Yttrium trihydride

The yttrium trihydride ( $\text{YH}_3$ ) compound can crystallize in a hexagonal structure with space group P-3c1 #165 [41], as shown in Fig. 3 a). This space group contains 12 symmetry operations which can be generated by the rotation  $C_{3z}$ , the screw rotation  $S_{2(x+y)}$ , the inversion  $\mathcal{I}$  and the time-reversal  $\mathbb{T}$  symmetry operations. The presence of both  $\mathcal{I}$  and  $\mathbb{T}$  symmetries indicate that energy




FIG. 3. a) Ball stick model of the hexagonal crystal structure of  $\text{YH}_3$  material [40]. b) Hexagonal Brillouin zone indicating the high-symmetry points. c) Position of electronic states in reciprocal space with energy band gap of less than 0.01 eV. d) Projected band structure with H-s orbitals along the high symmetry lines, where the symbols + and - indicate the inversion symmetry eigenvalues of +1 and -1 respectively at  $\Gamma$  point. e) Spin Hall conductivity (in  $\hbar/e/S/\text{cm}$ ). f) Total density of states (DOS) as a function of Fermi energy. g) Spin Berry curvature  $\Omega_{xy}^z(\mathbf{k})$  [ $\text{\AA}^2$ ] along high symmetry lines added over the valence band.  $\Omega_{xy}^z(\mathbf{k})$  signal is concentrated where the change of orbital character occurs. It is noted a large SHC in the band gap where there are no electronic states. Spin-orbit coupling is included in all the calculations and the Fermi level is set to zero.

bands are double degenerate along the full Brillouin zone (BZ). The BZ for this material is an hexagonal unit cell in reciprocal space with high-symmetry points as indicated in Fig. 3 b).

Based on *ab-initio* calculations, we have obtained the electronic band structure of the  $\text{YH}_3$  material and we have confirmed the appearance of the quasi-nodal sphere centered at the  $\Gamma$  point. The band structure along the high-symmetry lines, including the spin-orbit coupling interaction, is shown in Fig. 3 d). It should be noted a hybridization band gap between the valence and conduction bands for all  $\mathbf{k}$ -paths that connect the  $\Gamma$  point with any other high symmetry point. The hybridization band gap is originated by the SOC interaction and it is extended along the quasi-nodal sphere in reciprocal space. Symmetry analysis indicates that no crystal symmetry protects a possible nodal sphere or line whenever SOC is included. To confirm the existence of the quasi-nodal sphere, we carried out a systematic search of the  $\mathbf{k}$ -points in BZ where almost zero band gaps between the occupied and unoccupied bands are located. To visualize the quasi-nodal sphere, we have plotted the  $\mathbf{k}$ -points in the BZ with energy band gap of less than 0.01 eV in Fig. 3 c).

The orbital-resolved H-s projection is also shown in the band structure of Fig. 3 d). The figure shows that these orbitals dominate the valence band close to the Fermi level. In addition, a band inversion (change of the orbital character) is noted around the  $\Gamma$  point, which is occurring on the surface of the quasi-nodal sphere, as it was predicted by the  $\mathbf{k}\cdot\mathbf{p}$  model presented in the previous section.

We have also calculated the inversion symmetry eigenvalues on all TRIMs and we have observed the change of eigenvalues from the valence to the conduction bands at the  $\Gamma$  point (symbols + and - in Fig. 3 d). This change of eigenvalues indicates that the material can be considered a topological insulator with a tiny band gap spread along the quasi-nodal sphere. We have corroborated the topological insulator property of this material by using the Wilson loop method in order to find the  $\mathbb{Z}_2$  invariants for the six  $k_i=0$  planes in the BZ. Our calculations show that for the  $k_i=0$  planes the invariant is 1 while for the  $k_i=\pi$  planes the invariant is -1 (here  $i=x, y, z$ ). These results indicate the existence of the strong  $\mathbb{Z}_2$  Fu-Kane-Mele topological index in the  $\text{YH}_3$  material.

As it is well known, topological insulators are materials that can provide a platform to reach a large spin Hall



conductivity (SHC) signal [7, 8]. For the case of  $\text{YH}_3$ , we have plotted the SHC as a function of Fermi level in Fig. 3 e). In Fig. 3 g) we found a strong SHC response at the Fermi level, well distributed all around the quasi-nodal sphere. Note that the shape of the signal of the SHC along  $\Gamma$ -M matches the one of the model Hamiltonian presented in Fig. 1. a) This particular feature has been also observed in 2D topological insulators, where the  $\mathbf{k}$ -resolved SHC signal is distributed around the  $\Gamma$  point [42]. Furthermore, it is important to highlight that a strong SHC signal is observed in the absence of energy states at the Fermi level, as it can be observed in the total density of states calculation in Fig. 3 f). These results indicate that hydride materials can be promissory materials to efficient spin/charge conversion by means of the SHE [43].

### Calcium telluride

Calcium telluride (CaTe) is another material with quasi-nodal sphere realization due to band inversion. CaTe is a non-magnetic material whose space group is Pm-3m #221 and has 48 symmetry operations. This symmetry group can be generated by the operations:  $C_{2z}$ ,  $C_{2y}$ ,  $C_{2(x+y)}$ ,  $C_{3(x+y+z)}$ ,  $\mathcal{I}$  and  $\mathcal{T}$ . The cubic crystal structure and Brillouin zone for the CaTe compound are shown in Fig. 4 a) and Fig. 4 b), respectively. In Fig. 4 d) we present the electronic band structure with the orbital-projected Ca- $p$  states. We note a band character inversion between the valence and conduction bands around the M point. The symbols + and - in Fig. 4 d) indicate a reversal of the inversion symmetry eigenvalues at the M point, which indicates a topological response on this compound.

We have also found a Dirac point (DP) along the M-R  $\mathbf{k}$ -path that can be observed in the band structure of Fig. 4 e). However, a tiny band gap is observed along any  $\mathbf{k}$ -path starting from M when the SOC is taken into account. The stability of the DP is confirmed by the projection of the  $C_{4x}$  rotation symmetry eigenvalues at the two bands that generate the DP as shown in Fig. 4 e). The eigenvalues of  $C_{4x}$  are the fourth roots of  $-1$  and by Kramer's rule they come in pairs:  $(e^{i\frac{\pi}{2}}, e^{-i\frac{\pi}{2}})$  and  $(e^{i\frac{3\pi}{2}}, e^{-i\frac{3\pi}{2}})$ . The trace of the associated matrix permits us to detect the type of corepresentation at the high symmetry line and these traces are  $\sqrt{2}$  and  $-\sqrt{2}$  respectively. Whenever the traces differ, it means that the bands have different corepresentations of the group generated by  $C_{4x}$  and  $\mathcal{IT}$  and therefore the bands cannot hybridize [44]. In this particular case the traces of the operator  $C_{4x}$  are different and therefore the DP is protected by the four-fold rotation symmetry.

The procedure for finding the quasi-nodal sphere was also applied in the CaTe case. The quasi-nodal spheres

are centered at the M points, and the signal of the  $\mathbf{k}$ -resolved spin Berry curvature peaks at the  $\mathbf{k}$ -points where the quasi-nodal sphere is located as shown in Fig. 4 g). We found that the SHC response of the quasi-nodal sphere arises from the band inversion around the M point, as illustrated in Fig. 1 b) obtained from the  $\mathbf{k} \cdot \mathbf{p}$  Hamiltonian model of Eqn. (1). In the case of CaTe this feature is noticed, namely that the regions near the quasi-nodal sphere (M-point centered) retain locally large  $\mathbf{k}$ -resolved SHC (see Fig. 4 g)).

Topological insulators share a strong spin Hall conductivity, however, the presence of a Dirac point is not always an indication of SHC. We have found a large signal of the SHC at the Fermi level for the CaTe as shown in 4 f), where the spin Hall conductivity as a function of Fermi energy is presented. We can notice that the SHC signal is strong and coincides with the energy range of the quasi-nodal sphere and the DP. However, it is not clear whether the signal of the SHC comes from the quasi-nodal sphere, the DP or both. In order to distinguish the contribution to the spin Hall conductivity by the DP and the quasi-nodal sphere we have applied a small (3%) diagonal distortion on the cubic CaTe. This strain deformation transforms CaTe from the cubic (Pm-3m) to the rhombohedral (R-3m) structure. The deformation preserves the symmetry generators:  $C_{2(x+y)}$ ,  $C_{3(x+y+z)}$ ,  $\mathcal{I}$  and  $\mathcal{T}$ , but breaks the  $C_{4x}$  symmetry. The breaking of the  $C_{4x}$  symmetry leads to a band gap and a subsequent absence of the DP is noticed (see Fig. 5 e).

The rhombohedral structure of CaTe preserves the band inversion, the reversal of the inversion symmetry eigenvalues, and the quasi-nodal sphere around the M point. For this rhombohedral phase, it is noted a topological insulator behavior with a small band gap, similar to the  $\text{YH}_3$  case and the four-states Hamiltonian model of Eqn. (1). The calculation shows a non-zero SHC in the band gap, which emerges from the band inversion at the quasi-nodal sphere as shown in Fig. 5 f) and corroborated by the  $\mathbf{k} \cdot \mathbf{p}$  model (Fig. 1 b).

The CaTe crystalline phase displayed in Fig. 4 a) has been experimentally known to be dynamically stable for pressures above 33 GPa[45]. This has also been corroborated by our *ab-initio* calculations (see figures 5 a) and 5 c). We can see that the DP along the M-R  $\mathbf{k}$ -path is preserved even when an isotropic pressure is applied (see figures 5 b) and 5 d)). This result is expected since the  $C_{4x}$  symmetry, band inversion and quasi-nodal sphere are not affected by an isotropic compression.

Our results suggest that CaTe and  $\text{YH}_3$  could both display large spin Hall angles for practical applications since they show small longitudinal conductivities (low density of states) at the Fermi level. The SHC signal is concentrated at the hybridized band regions, as it was also found by Lau et al. [46].




FIG. 4. a) Crystal structure [40] and b) cubic Brillouin zone indicating the high-symmetry points of the CaTe material (cubic phase). c) Total density of states (DOS) as a function of Fermi energy. d) Projected band structure with Ca- $p$  orbitals along the high symmetry lines, where the symbols + and - denote the inversion symmetry eigenvalues at M point, +1 and -1, respectively. e)  $C_{4z}$  symmetry eigenvalues resolved at the electronic bands along the X-M-R  $k$ -path. f) Spin Hall conductivity (in  $\hbar/e/S/\text{cm}$ ) as a function of Fermi energy. It is noted a strong SHC signal at the Fermi level, which is attributed to the quasi-nodal sphere and the Dirac point. g) Spin Berry curvature  $\Omega_{xy}^z(k)$  [ $\text{\AA}^2$ ] along high symmetry lines added over the valence band.  $\Omega_{xy}^z(k)$  signal is concentrated where the change of orbital character occurs. Spin-orbit coupling is included in all the calculations and the Fermi level is set as zero.

### Computational methods

For  $\text{YH}_3$  and CaTe materials, we have carried out *ab-initio* calculations within the density-functional theory with an uniform  $\mathbf{k}$ -mesh in the Brillouin zone of  $13 \times 13 \times 11$  and  $14 \times 14 \times 14$ , respectively. The energy cut-off was taken to be 520 eV and the exchange-correlation contribution to the total energy treated within the PBE parametrization [47], as implemented in the Vienna *ab-initio* simulation package (VASP) [48]. Projected band structures were plotted using the `pyprocar` program [49]. The symmetry eigenvalues along the high symmetry lines were calculated using the `irrep` code [50]. To estimate the spin transport properties, we employed the `wannier90` code [51, 52] to build a Wannier Hamiltonian. The intrinsic spin Hall conductivity (SHC) was evaluated by integrating the spin Berry curvature on a dense  $240 \times 240 \times 240$  grid in the first Brillouin zone, using the `linres` code [53]. In this model, the SHC can be written as:

$$\sigma_{xy}^z = -\frac{e^2}{\hbar} \sum_n \int_{BZ} \frac{dk^3}{(2\pi)^3} f_n(k) \Omega_{n,xy}^z(k), \quad (35)$$

where  $f_n(k)$  is the Fermi-Dirac distribution and  $\Omega_{n,xy}^z(k)$  is the spin Berry curvature for the  $n$ th band. The spin Berry curvature for the  $n$ th band can be calculated using the Kubo formula:

$$\Omega_{n,xy}^z(k) = -2\hbar^2 \text{Im} \sum_{m \neq n} \frac{\langle n, k | \hat{j}_x^z | m, k \rangle \langle m, k | \hat{v}_y | n, k \rangle}{(\epsilon_{n,k} - \epsilon_{m,k})^2}, \quad (36)$$

where  $|n, k\rangle$  are the Bloch functions,  $k$  is the Bloch wave vector,  $\epsilon_{n,k}$  is the electronic band energy,  $\hat{v}_i$  is the velocity operator in the  $i$  direction and  $\hat{j}_x^z = \frac{1}{2} \{\hat{v}_x, \hat{s}_z\}$  is the spin current operator.

Vibrational frequencies have been calculated within the perturbation theory method as implemented in VASP using the `phonopy` code [54] to obtain the force constants and vibrational frequencies. In this case, we have used a  $2 \times 2 \times 2$  supercell with a  $8 \times 8 \times 8$   $\mathbf{q}$ -mesh to guarantee convergence up to  $0.1 \text{ cm}^{-1}$ .






FIG. 5. Top four panels: dynamical stability of cubic CaTe under high pressure. a) and c) correspond to phonon dispersion curves at 30 GPa and 50 GPa respectively. Electronic band structure at the same given pressures are contained in panels b) and d) with preserved band inversion at  $\Gamma$  point. The colors indicate the eigenvalues of the four-fold rotation  $C_{4x}$ . Bottom two panels: CaTe with a small ( $\epsilon=3\%$ ) diagonal distortion, thus breaking the four-fold symmetry. e) Electronic band structure with gapless phase; note that there is no DP due to the broken four-fold symmetry. f) Energy resolved spin Hall conductivity. The SHC arises mainly from the electronic bands that generate the quasi-nodal sphere and it survives despite the absence of the DP.

## CONCLUSIONS

Band inversion in topological insulators induced by hybridization of the energy bands produce quasi-nodal spheres. The orbital type change on the bands occurs on the quasi-nodal sphere and the matrix information on how this change is happening encodes the strong Fu-Kane-Mele invariant. The signal of the SHC is localized where the orbital type changes, and therefore the signal is strong close to the Fermi level, where hybridization occurs.

The appearance of quasi-nodal spheres on materials with band inversion due to hybridization is therefore enforced. The change of orbital type must occur in a 2-dimensional sphere, otherwise the Fu-Kane-Mele invariant is forced to be trivial.

The band inversion on  $\text{YH}_3$  and CaTe is confirmed and the SHC is calculated in both materials. The signal is localized on the quasi-nodal sphere and, by removing the Dirac point of CaTe by strain, it is shown that in both

cases the signal of the SHC is localized on the quasi-nodal spheres where the orbital type changes. This result is expected to hold in materials with a similar type of band inversion as the trihydrides ( $\text{XH}_3$ ,  $\text{X} = \text{Y, Gd, Ho, Tb, Nd}$ ) and  $\text{Si}_3\text{N}_2$ .

## ACKNOWLEDGMENTS

The first author gratefully acknowledges the computing time granted on the supercomputer Mogon at Johannes Gutenberg University Mainz (hpc.uni-mainz.de). The second author thanks the Royal Society of the UK for an International Collaboration Award and the Science and Technology Facilities Council of the UK for computing time on SCARF. The third author acknowledges the support of CONACYT through project CB-2017-2018-A1-S-30345-F-3125. The first and the third authors thank the continuous support of the Alexander Von Humboldt Foundation, Germany.

\* rhernandezj@uninorte.edu.co

† ccpinilla@uninorte.edu.co

‡ bjongbloed@uninorte.edu.co, uribe@mpim-bonn.mpg.de

- [1] D. Castelvecchi, *Nature* **547**, 272 (2017), ISSN 1476-4687, URL <https://doi.org/10.1038/547272a>.
- [2] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, *Phys. Rev. B* **78**, 195125 (2008), URL <https://link.aps.org/doi/10.1103/PhysRevB.78.195125>.
- [3] B. Q. Lv, T. Qian, and H. Ding, *Rev. Mod. Phys.* **93**, 025002 (2021), URL <https://link.aps.org/doi/10.1103/RevModPhys.93.025002>.
- [4] A. Bansil, H. Lin, and T. Das, *Rev. Mod. Phys.* **88**, 021004 (2016), URL <https://link.aps.org/doi/10.1103/RevModPhys.88.021004>.
- [5] C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 226801 (2005), URL <https://link.aps.org/doi/10.1103/PhysRevLett.95.226801>.
- [6] B. A. Bernevig and S.-C. Zhang, *Phys. Rev. Lett.* **96**, 106802 (2006), URL <https://link.aps.org/doi/10.1103/PhysRevLett.96.106802>.
- [7] X.-L. Qi and S.-C. Zhang, *Physics Today* **63**, 33 (2010), <https://doi.org/10.1063/1.3293411>, URL <https://doi.org/10.1063/1.3293411>.
- [8] J. Sinova, S. O. Valenzuela, J. Wunderlich, C. H. Back, and T. Jungwirth, *Rev. Mod. Phys.* **87**, 1213 (2015), URL <https://link.aps.org/doi/10.1103/RevModPhys.87.1213>.
- [9] L. Fu, C. L. Kane, and E. J. Mele, *Phys. Rev. Lett.* **98**, 106803 (2007), URL <https://link.aps.org/doi/10.1103/PhysRevLett.98.106803>.
- [10] A. A. Soluyanov and D. Vanderbilt, *Phys. Rev. B* **83**, 235401 (2011), URL <https://link.aps.org/doi/10.1103/PhysRevB.83.235401>.
- [11] C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 146802 (2005), URL <https://link.aps.org/doi/10.1103/PhysRevLett.95.146802>.

- [12] A. Bansil, H. Lin, and T. Das, *Rev. Mod. Phys.* **88**, 021004 (2016), URL <https://link.aps.org/doi/10.1103/RevModPhys.88.021004>.
- [13] Z. Zhu, Y. Cheng, and U. Schwingenschlöggl, *Phys. Rev. B* **85**, 235401 (2012), URL <https://link.aps.org/doi/10.1103/PhysRevB.85.235401>.
- [14] Y. Ando, *Journal of the Physical Society of Japan* **82**, 102001 (2013), <https://doi.org/10.7566/JPSJ.82.102001>, URL <https://doi.org/10.7566/JPSJ.82.102001>.
- [15] A. Manchon, H. C. Koo, J. Nitta, S. M. Frolov, and R. A. Duine, *Nature Materials* **14**, 871 (2015), URL <https://doi.org/10.1038/nmat4360>.
- [16] A. J. Browne, A. Krajewska, and A. S. Gibbs, *J. Mater. Chem. C* **9**, 11640 (2021), URL <http://dx.doi.org/10.1039/D1TC02070F>.
- [17] P. Dziawa, B. J. Kowalski, K. Dybko, R. Buczko, A. Szczerbakow, M. Szot, E. Lusakowska, T. Balasubramanian, B. M. Wojek, M. H. Berntsen, et al., *Nature Materials* **11**, 1023 (2012), URL <https://doi.org/10.1038/nmat3449>.
- [18] Y. Zhang, K. He, C.-Z. Chang, C.-L. Song, L.-L. Wang, X. Chen, J.-F. Jia, Z. Fang, X. Dai, W.-Y. Shan, et al., *Nature Physics* **6**, 584 (2010), URL <https://doi.org/10.1038/nphys1689>.
- [19] W. Witczak-Krempa, G. Chen, Y. B. Kim, and L. Balents, *Annual Review of Condensed Matter Physics* **5**, 57 (2014), <https://doi.org/10.1146/annurev-conmatphys-020911-125138>, URL <https://doi.org/10.1146/annurev-conmatphys-020911-125138>.
- [20] S. Chege, P. Ning'i, J. Sifuna, and G. O. Amolo, *AIP Advances* **10**, 095018 (2020), URL <https://doi.org/10.1063/5.0022525>.
- [21] H. Zhang, C.-X. Liu, X.-L. Qi, X. Dai, Z. Fang, and S.-C. Zhang, *Nature Physics* **5**, 438 (2009), URL <https://doi.org/10.1038/nphys1270>.
- [22] M. Zhao, X. Zhang, and L. Li, *Scientific Reports* **5**, 16108 (2015), ISSN 2045-2322, URL <https://doi.org/10.1038/srep16108>.
- [23] H. Lin, L. A. Wray, Y. Xia, S. Xu, S. Jia, R. J. Cava, A. Bansil, and M. Z. Hasan, *Nature Materials* **9**, 546 (2010), URL <https://doi.org/10.1038/nmat2771>.
- [24] S. Chadov, X. Qi, J. Kübler, G. H. Fecher, C. Felser, and S. C. Zhang, *Nature Materials* **9**, 541 (2010), URL <https://doi.org/10.1038/nmat2770>.
- [25] D. Xiao, Y. Yao, W. Feng, J. Wen, W. Zhu, X.-Q. Chen, G. M. Stocks, and Z. Zhang, *Phys. Rev. Lett.* **105**, 096404 (2010), URL <https://link.aps.org/doi/10.1103/PhysRevLett.105.096404>.
- [26] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, *Science* **314**, 1757 (2006), <https://www.science.org/doi/pdf/10.1126/science.1133734>, URL <https://www.science.org/doi/abs/10.1126/science.1133734>.
- [27] W. Feng, D. Xiao, J. Ding, and Y. Yao, *Phys. Rev. Lett.* **106**, 016402 (2011), URL <https://link.aps.org/doi/10.1103/PhysRevLett.106.016402>.
- [28] D. Shao, T. Chen, Q. Gu, Z. Guo, P. Lu, J. Sun, L. Sheng, and D. Xing, *Scientific Reports* **8**, 1467 (2018), URL <https://doi.org/10.1038/s41598-018-19870-5>.
- [29] J. Wang, Y. Liu, K.-H. Jin, X. Sui, L. Zhang, W. Duan, F. Liu, and B. Huang, *Phys. Rev. B* **98**, 201112 (2018), URL <https://link.aps.org/doi/10.1103/PhysRevB.98.201112>.
- [30] R. González-Hernández, E. Tuiran, and B. Uribe, *Phys. Rev. B* **104**, 205128 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevB.104.205128>.
- [31] M. F. Atiyah, *K-theory* (W. A. Benjamin, Inc., New York-Amsterdam, 1967), lecture notes by D. W. Anderson.
- [32] G. De Nittis and K. Gomi, *Comm. Math. Phys.* **339**, 1 (2015), ISSN 0010-3616, URL <https://doi.org/10.1007/s00220-015-2390-0>.
- [33] M. F. Atiyah, *Quart. J. Math. Oxford Ser. (2)* **17**, 367 (1966), ISSN 0033-5606, URL <https://doi.org/10.1093/qmath/17.1.367>.
- [34] G. Luke and A. S. Mishchenko, *Vector bundles and their applications*, vol. 447 of *Mathematics and its Applications* (Kluwer Academic Publishers, Dordrecht, 1998), ISBN 0-7923-5154-1, URL <https://doi.org/10.1007/978-1-4757-6923-4>.
- [35] J. L. Dupont, *Math. Scand.* **24**, 27 (1969), ISSN 0025-5521, URL <https://doi.org/10.7146/math.scand.a-10918>.
- [36] J. Ahn and B.-J. Yang, *Phys. Rev. B* **99**, 235125 (2019), URL <https://link.aps.org/doi/10.1103/PhysRevB.99.235125>.
- [37] Z. Wang, X.-L. Qi, and S.-C. Zhang, *New Journal of Physics* **12**, 065007 (2010), URL <https://doi.org/10.1088/1367-2630/12/6/065007>.
- [38] T. L. Hughes, E. Prodan, and B. A. Bernevig, *Phys. Rev. B* **83**, 245132 (2011), URL <https://link.aps.org/doi/10.1103/PhysRevB.83.245132>.
- [39] J. Liu and D. Vanderbilt, *Phys. Rev. B* **92**, 245138 (2015), URL <https://link.aps.org/doi/10.1103/PhysRevB.92.245138>.
- [40] K. Momma and F. Izumi, *Journal of Applied Crystallography* **44**, 1272 (2011), URL <https://doi.org/10.1107/S0021889811038970>.
- [41] P. Rizo-Acosta, F. Cuevas, and M. Latroche, *J. Mater. Chem. A* **7**, 23064 (2019), URL <http://dx.doi.org/10.1039/C9TA05440E>.
- [42] S.-j. Zhang, W.-x. Ji, C.-w. Zhang, P. Li, and P.-j. Wang, *Scientific Reports* **7**, 45923 (2017), URL <https://doi.org/10.1038/srep45923>.
- [43] X. Zuo, Q. Gao, X. Sui, X. Xu, X. Jiang, L. Han, H. Li, D. Li, D. Liu, B. Huang, et al., *Phys. Rev. B* **103**, 125159 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevB.103.125159>.
- [44] R. González-Hernández, E. Tuiran, and B. Uribe, *Phys. Rev. B* **103**, 235143 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevB.103.235143>.
- [45] H. Luo, R. G. Greene, K. Ghandehari, T. Li, and A. L. Ruoff, *Phys. Rev. B* **50**, 16232 (1994), URL <https://link.aps.org/doi/10.1103/PhysRevB.50.16232>.
- [46] Y.-C. Lau, H. Lee, G. Qu, K. Nakamura, and M. Hayashi, *Phys. Rev. B* **99**, 064410 (2019), URL <https://link.aps.org/doi/10.1103/PhysRevB.99.064410>.
- [47] J. P. Perdew, K. Burke, and M. Ernzerhof, *Phys. Rev. Lett.* **77**, 3865 (1996), ISSN 0031-9007, URL <http://link.aps.org/doi/10.1103/PhysRevLett.77.3865>.
- [48] G. Kresse and J. Furthmüller, *Phys. Rev. B* **54**, 11169 (1996), URL <https://link.aps.org/doi/10.1103/PhysRevB.54.11169>.
- [49] U. Herath, P. Tavazde, X. He, E. Bousquet, S. Singh, F. Muñoz, and A. H. Romero, *Computer Physics Communications* **251**, 107080 (2020), ISSN 0010-4655, URL <http://www.sciencedirect.com/science/>

- article/pii/S0010465519303935.
- [50] M. Iraola, J. L. Mañes, B. Bradlyn, M. K. Horton, T. Neupert, M. G. Vergniory, and S. S. Tsirkin, *Computer Physics Communications* **272**, 108226 (2022), ISSN 0010-4655, URL <https://www.sciencedirect.com/science/article/pii/S0010465521003386>.
- [51] A. A. Mostofi, J. R. Yates, G. Pizzi, Y.-S. Lee, I. Souza, D. Vanderbilt, and N. Marzari, *Computer Physics Communications* **185**, 2309 (2014), ISSN 0010-4655, URL <http://www.sciencedirect.com/science/article/pii/S001046551400157X>.
- [52] G. Pizzi, V. Vitale, R. Arita, S. Blügel, F. Freimuth, G. Géranton, M. Gibertini, D. Gresch, C. Johnson, T. Koretsune, et al., *Journal of Physics: Condensed Matter* **32**, 165902 (2020), URL <https://doi.org/10.1088/1361-648x/ab51ff>.
- [53] J. Železný, <https://bitbucket.org/zeleznyj/wannier-linear-response/wiki/Home>.
- [54] A. Togo and I. Tanaka, *Scr. Mater.* **108**, 1 (2015).