METZLER/ZETA CORRESPONDENCE

Yusuke IDE

Department of Mathematics, College of Humanities and Sciences, Nihon University Sakura-josui, Setagaya-ku, Tokyo 156-8550, JAPAN

> Takashi KOMATSU Math. Research Institute Calc for Industry, Minami, Hiroshima, 732-0816, JAPAN e-mail: ta.komatsu@sunmath-calc.co.jp

Norio KONNO Department of Applied Mathematics, Faculty of Engineering, Yokohama National University Hodogaya, Yokohama 240-8501, JAPAN

e-mail: konno-norio-bt@ynu.ac.jp

Iwao SATO Oyama National College of Technology Oyama, Tochigi 323-0806, JAPAN e-mail: isato@oyama-ct.ac.jp

July 4, 2022

Abstract

We present an explicit formula for the determinant on the Metzler matrix of a digraph D. Furthermore, we introduce a walk-type zeta function with respect to this Metzler matrix of the symmetric digraph of a finite torus, and express its limit formula by using the integral expression.

2000 Mathematical Subject Classification: 05C50, 15A15. **Key words and phrases** : digraph, Metzler matrix, zeta function, torus

The contact author for correspondence: Iwao Sato Oyama National College of Technology, Oyama, Tochigi 323-0806, JAPAN E-mail: isato@oyama-ct.ac.jp

1 Introduction

Ihara [7] defined the Ihara zeta functions of graphs, and showed that the reciprocals of the Ihara zeta functions of regular graphs are explicit polynomials. The Ihara zeta function of a regular graph G associated with a unitary representation of the fundamental group of G was developed by Sunada [21, 22]. Hashimoto [6] treated multivariable zeta functions of bipartite graphs. Furthermore, Hashimoto [6] gave a determinant expression for the Ihara zeta function of a general graph by using the edge matrix. Bass [2] generalized Ihara's result on the Ihara zeta function of a regular graph to an irregular graph G. Stark and Terras [20] gave an elementary proof of Bass' theorem, and discussed three different zeta functions of any graph. Various proofs of Bass' theorem were given by Kotani and Sunada [18], and Foata and Zeilberger [5].

Recently, there were exciting developments between quantum walk [1, 8, 9, 16, 23] on a graph and the Ihara zeta function of a graph. We investigated a new class of zeta functions for many kinds of walks including the quantum walk (QW) and the random walk (RW) on a graph by a series of "Zeta Correspondence" of our previous work [10, 11, 12, 13, 14, 15, 17]. In Walk/Zeta Correspondence [11], a walk-type zeta function was defined without use of the determinant expressions of zeta function of a graph G, and various properties of walk-type zeta functions of RW, correlated random walk (CRW) and QW on G were studied. Also, their limit formulas by using integral expressions were presented.

The Ihara zeta function has many applications in both pure and applied mathematics, including for instance dynamical systems, spectral graph theory and complex network analysis. In complex network analysis, Masuda, Preciado and Ogura [19] studied the stochastic susceptible-infected-susceptible model (SIS model) of epidemic processes on finite directed and weighted networks, and presented a new lower bound on the exponential rate by using the leading eigenvalue of some matrix composed from the digraph version of the edge matrix. The edge matrix appeared in the determinant expressions for the Ihara zeta function. Furthermore, the above matrix is a kind of the Metzler matrix.

In this paper, we present an explicit formula for a determinant on the Metzler matrix of a digraph D. Moreover, we introduce a walk-type zeta function with respect to this Metzler matrix of the symmetric digraph of a finite torus, and express its limit formula by using integral expression.

This paper is organized as follows: In Section 2, we give a short review about the Ihara zeta function of a graph. In Section 3, we review "Walk/Zeta Correspondence" studied in [11]. In Section 4, we state a new lower bound on the exponential rate by Masuda, Preciado and Ogura [19]. In Section 5, we present an explicit formula for the determinant on the above Metzler matrix of a digraph D. In Section 6, we give an explicit formula for the characteristic polynomial of the above Metzler matrix of the symmetric digraph of a graph, and obtain its spectrum. In Section 7, we introduce a walk-type zeta function with respect to this Metzler matrix of the symmetric digraph of a finite torus, and express its limit formula by using integral expression.

2 The Ihara zeta function of a graph

Graphs and digraphs treated here are finite. Let G be a connected graph and $D(G) = \{(u, v), (v, u) \mid uv \in E(G)\}$ the arc set of the symmetric digraph corresponding to G. For $e = (u, v) \in D(G)$, set u = o(e) and v = t(e). Furthermore, let $e^{-1} = (v, u)$ be the *inverse* of e = (u, v).

A path P of length n in G is a sequence $P = (e_1, \dots, e_n)$ of n arcs such that $e_i \in D(G)$, $t(e_i) = o(e_{i+1})(1 \le i \le n-1)$. If $e_i = (v_{i-1}, v_i)$ for $i = 1, \dots, n$, then we write $P = (v_0, v_1, \dots, v_{n-1}, v_n)$. Set |P| = n, $o(P) = o(e_1)$ and $t(P) = t(e_n)$. Also, P is called an (o(P), t(P))-path. We say that a path $P = (e_1, \dots, e_n)$ has a backtracking if $e_{i+1}^{-1} = e_i$ for some i $(1 \le i \le n-1)$. A (v, w)-path is called a *v*-cycle (or *v*-closed path) if v = w. The inverse cycle of a cycle $C = (e_1, \dots, e_n)$ is the cycle $C^{-1} = (e_n^{-1}, \dots, e_1^{-1})$.

We introduce an equivalence relation between cycles. Two cycles $C_1 = (e_1, \dots, e_m)$ and $C_2 = (f_1, \dots, f_m)$ are called *equivalent* if $f_j = e_{j+k}$ for all j. The inverse cycle of C is not equivalent to C if $|C| \ge 3$. Let [C] be the equivalence class which contains a cycle C. Let B^r be the cycle obtained by going r times around a cycle B. Such a cycle is called a *multiple* of B. A cycle C is *reduced* if both C and C^2 have no backtracking. Furthermore, a cycle C is *prime* if it is not a multiple of a strictly smaller cycle. Note that each equivalence class of prime, reduced cycles of a graph G corresponds to a unique conjugacy class of the fundamental group $\pi_1(G, v)$ of G at a vertex v of G.

The *Ihara*(-Selberg) zeta function of G is defined by

$$\mathbf{Z}(G, u) = \prod_{[C]} (1 - u^{|C|})^{-1},$$

where [C] runs over all equivalence classes of prime, reduced cycles of G.

Let G be a connected graph with N vertices and M edges. Then two $2M \times 2M$ matrices $\mathbf{B} = \mathbf{B}(G) = (\mathbf{B}_{ef})_{e,f \in D(G)}$ and $\mathbf{J}_0 = \mathbf{J}_0(G) = (\mathbf{J}_{ef})_{e,f \in D(G)}$ are defined as follows:

$$\mathbf{B}_{ef} = \begin{cases} 1 & \text{if } t(e) = o(f), \\ 0 & \text{otherwise,} \end{cases} \quad \mathbf{J}_{ef} = \begin{cases} 1 & \text{if } f = e^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

Note that

$$\mathbf{J}_0 = \mathbf{B} \circ \mathbf{B}^T,$$

where \mathbf{B}^T is the transpose of \mathbf{B} , and the *Schur/Hadamard product* $\mathbf{A} \circ \mathbf{B}$ of two matrices \mathbf{A} and \mathbf{B} is defined by

$$(\mathbf{A} \circ \mathbf{B})_{ij} = \mathbf{A}_{ij} \cdot \mathbf{B}_{ij}.$$

The matrix $\mathbf{B} - \mathbf{J}_0$ is called the *edge matrix* of *G*.

Theorem 1 (Ihara; Hashimoto; Bass) Let G be a connected graph with N vertices and M edges. Then the reciprocal of the Ihara zeta function of G is given by

$$\mathbf{Z}(G, u)^{-1} = \det(\mathbf{I}_{2M} - u(\mathbf{B} - \mathbf{J}_0)) = (1 - u^2)^{M-N} \det(\mathbf{I}_N - u\mathbf{A}(G) + u^2(\mathbf{D}_G - \mathbf{I}_N)),$$

where $\mathbf{D}_G = (d_{ij})$ is the diagonal matrix with $d_{ii} = \deg_G v_i$ $(V(G) = \{v_1, \cdots, v_N\})$.

3 Walk/Zeta Correspondence on torus

We state the Walk/Zeta Correspondence on a finite torus. At first, we give the definition of the 2*d*-state discrete-time walk on T_N^d . The discrete-time walk is defined by using a *shift* operator and a coin matrix which will be mentioned below.

Let $f: V(T_N^d) \longrightarrow \mathbb{C}^{2d}$. For j = 1, 2, ..., d and $\mathbf{x} = (x_1, ..., x_d) \in V(T_N^d)$, the shift operator τ_j is defined by

$$(\tau_j f)(\mathbf{x}) = f(\mathbf{x} - \mathbf{e}_j),$$

where $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_d\}$ denotes the standard basis of \mathbb{R}^d .

Let $A = (a_{ij})_{i,j=1,2,...,2d}$ be a $2d \times 2d$ matrix with $a_{ij} \in \mathbb{C}$ for i, j = 1, 2, ..., 2d. We call A the *coin matrix*. If $a_{ij} \in [0,1]$ and $\sum_{i=1}^{2d} a_{ij} = 1$ for any j = 1, 2, ..., 2d, then the walk is a CRW. In particular, when $a_{i1} = a_{i2} = \cdots = a_{i2d}$ for any i = 1, 2, ..., 2d, this CRW becomes a RW. If A is unitary, then the walk is a QW. So our class of walks contains RWs, CRWs, and QWs as special models.

To describe the evolution of the walk, we decompose the $2d \times 2d$ coin matrix A as

$$A = \sum_{j=1}^{2d} P_j A_j$$

where P_j denotes the orthogonal projection onto the one-dimensional subspace $\mathbb{C}\eta_j$ in \mathbb{C}^{2d} . Here $\{\eta_1, \eta_2, \ldots, \eta_{2d}\}$ denotes a standard basis on \mathbb{C}^{2d} .

The discrete-time walk associated with the coin matrix A on T_N^d is determined by the $2dN^d \times 2dN^d$ matrix

$$M_A = \sum_{j=1}^d \left(P_{2j-1} A \tau_j^{-1} + P_{2j} A \tau_j \right).$$
(1)

Let $\mathbb{Z}_{\geq} = \mathbb{Z} \cup \{0\}$. Then the state at time $n \in \mathbb{Z}_{\geq}$ and location $\mathbf{x} \in V(T_N^d)$ can be expressed by a 2*d*-dimensional vector:

$$\Psi_n(\mathbf{x}) = egin{bmatrix} \Psi_n^1(\mathbf{x}) \ \Psi_n^2(\mathbf{x}) \ dots \ \Psi_n^{2d}(\mathbf{x}) \end{bmatrix} \in \mathbb{C}^{2d}.$$

For $\Psi_n: V(T_N^d) \longrightarrow \mathbb{C}^{2d}$ $(n \in \mathbb{Z}_{\geq})$, from (1), the evolution of the walk is defined by

$$\Psi_{n+1}(\mathbf{x}) \equiv (M_A \Psi_n)(\mathbf{x}) = \sum_{j=1}^d \left(P_{2j-1} A \Psi_n(\mathbf{x} + \mathbf{e}_j) + P_{2j} A \Psi_n(\mathbf{x} - \mathbf{e}_j) \right).$$
(2)

Now, we define the *walk-type zeta function* by

$$\overline{\zeta}\left(A, T_N^d, u\right) = \det\left(I_{2dN^d} - uM_A\right)^{-1/N^d}.$$
(3)

In general, for a $d_c \times d_c$ coin matrix A, we put

$$\overline{\zeta}(A, T_N^d, u) = \det\left(I_{d_cN^d} - uM_A\right)^{-1/N^d}.$$

Komatsu et al. [11] obtained the following result.

Theorem 2 (Komatsu et al. [11])

$$\overline{\zeta} \left(A, T_N^d, u \right)^{-1} = \exp\left[\frac{1}{N^d} \sum_{\widetilde{\mathbf{k}} \in \widetilde{\mathbb{K}}_N^d} \log\left\{ \det\left(F(\widetilde{\mathbf{k}}, u) \right) \right\} \right],$$
$$\lim_{N \to \infty} \overline{\zeta} \left(A, T_N^d, u \right)^{-1} = \exp\left[\int_{[0, 2\pi)^d} \log\left\{ \det\left(F\left(\Theta^{(d)}, u\right) \right) \right\} d\Theta_{unif}^{(d)} \right],$$

where $\tilde{\mathbb{K}}_N = \{0, 2\pi/N, \ldots, 2\pi(N-1)/N\}, \Theta^{(d)} = (\theta_1, \theta_2, \ldots, \theta_d) (\in [0, 2\pi)^d)$ and $d\Theta_{unif}^{(d)}$ denotes the uniform measure on $[0, 2\pi)^d$, that is,

$$d\Theta_{unif}^{(d)} = \frac{d\theta_1}{2\pi} \cdots \frac{d\theta_d}{2\pi}.$$

Furthermore,

$$F(\mathbf{w},u) = I_{2d} - u\widehat{M}_A(\mathbf{w}) \quad and \quad \widehat{M}_A(\mathbf{w}) = \sum_{j=1}^d \left(e^{iw_j} P_{2j-1}A + e^{-iw_j} P_{2j}A \right),$$

with $\mathbf{w} = (w_1, w_2, \dots, w_d) \in \mathbb{R}^d$.

4 A lower bound on the exponential rate in the SIS model

At first, we state the stochastic SIS model on a digraph. Let D = (V(D), A(D)) be a connected digraph with N vertices and M arcs, where V(D) and A(D) are the set of vertices and arcs of D, respectively. At any given continuous time $t \ge 0$, each vertex is in one of two possible states, namely, susceptible or infected. An infected vertex v stochastically transits to the susceptible state at a constant instantaneous rate of $\delta_v > 0$, which is called the *recovery rate* of v. Whenever v is susceptible and its infected in-neighbor w is infected, then w stochastically and independently infects v at a constant instantaneous rate of $\beta_{(w,v)}$. The value $\beta_{(w,v)}$ is called the *infection rate*.

Let $p_v(t)$ $(v \in V(D))$ be the probability that v is infected at time t. Then the *decay rate* of the SIS model is defined by

$$\gamma = -\limsup_{t \to \infty} \frac{\log \sum_{v \in V(D)} p_v(t)}{t},$$

where all vertices are assumed to be infected at t = 0.

Next, let $V(D) = \{v_1, \ldots, v_N\}$ and $A(D) = \{e_1, \ldots, e_M\}$, and let an $N \times M$ matrix C be the *incidence matrix* of D. Furthermore, we define the *non-backtracking matrix* $H = (H_{ef})_{e,f \in A(D)}$ of D as follows:

$$H_{ef} = \begin{cases} 1 & \text{if } t(e) = o(f) \text{ and } f \neq e^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

The matrix H is a digraph version of the edge matrix of a graph.

A real matrix A is called *nonnegative*, denoted $A \ge \mathbf{0}$, if all entries of A are nonnegative. For two matrices A and B of same size, we say that $A \le B$ if $B-A \ge \mathbf{0}$. A square matrix A is called *Metzler* if all its off-diagonal entries are nonnegative [3]. For the Metzler matrix A, the maximum real part of the eigenvalues of A is denoted by $\lambda_{\max}(A)$ (see [3]). Let $\mathbf{M}_1 \oplus \cdots \oplus \mathbf{M}_s$ be the block diagonal sum of square matrices $\mathbf{M}_1, \cdots, \mathbf{M}_s$. If $\mathbf{M}_1 = \mathbf{M}_2 = \cdots = \mathbf{M}_s = \mathbf{M}$, then we write $s \circ \mathbf{M} = \mathbf{M}_1 \oplus \cdots \oplus \mathbf{M}_s$. Then an $(N+M) \times (N+M)$ A is defined as follows:

$$\mathcal{A} = \begin{bmatrix} -\mathbf{D} & C_+ B' \\ D'_2 C_-^T & H^T B' - B' - D'_1 - D'_2 \end{bmatrix},$$

where

$$C_{+} = \max(C, \mathbf{0}), \ C_{-} = \max(-C, \mathbf{0})$$

and

$$\mathbf{D} = (\delta_{v_1}) \otimes \cdots \otimes (\delta_{v_N}),$$

$$D'_1 = (\delta_{o(e_1)}) \otimes \cdots \otimes (\delta_{o(e_M)}),$$

$$D'_2 = (\delta_{t(e_1)}) \otimes \cdots \otimes (\delta_{t(e_m)}),$$

$$B' = (\beta_{e_1}) \otimes \cdots \otimes (\beta_{e_m}).$$

Two matrices C_+ and C_- denote the positive and negative parts of the incidence matrix C, respectively. Note that \mathcal{A} is Metzler.

Masuda, Preciado and Ogura [19] presented a new lower bound on the exponential rate by using the leading eigenvalue of \mathcal{A} .

Theorem 3 (Masuda, Preciado and Ogura [19]) A lower bound on the decay rate γ is given as follows:

$$\gamma \geq -\lambda_{\max}(\mathcal{A}).$$

Furthermore, we use the Weinstein-Aronszajn identity (see [4]).

Theorem 4 (the Weinstein-Aronszajn identity) If **A** and **B** are an $r \times s$ and an $s \times r$ matrix, respectively, then we have

$$\det(\mathbf{I}_r - \mathbf{AB}) = \det(\mathbf{I}_s - \mathbf{BA}).$$

5 A determinant formula for the matrix \mathcal{A} of a digraph

At first, we state the properties for various matrices composing the Metzler matrix \mathcal{A} .

Let *D* be a connected graph with *N* vertices v_1, \dots, v_N and *M* arcs, and A(D) the arc set of *D*. Furthermore, let two $N \times N$ matrices $B = (B_{uv})$ and $\mathbf{D} = \mathbf{D}(D) = (D_{uv})$ be given as follows:

$$B_{uv} = \begin{cases} \beta_{uv} & \text{if } (u,v) \in A(D), \\ 0 & \text{otherwise,} \end{cases} \quad D_{uv} = \begin{cases} \delta_u & \text{if } u = v, \\ 0 & \text{otherwise} \end{cases}$$

where $\beta_{uv} > 0$, $\delta_u > 0$ for each $e = (u, v) \in A(D)$ and $u \in V(D)$. Set $\beta_e = \beta_{uv}$ for $e = (u, v) \in A(D)$.

The *incidence matrix* $C = (C_{ue})$ of D is the $N \times M$ matrix given as follows:

$$C_{eu} = \begin{cases} 1 & \text{if } t(e) = u, \\ -1 & \text{if } o(e) = u. \\ 0 & \text{otherwise.} \end{cases}$$

Furthermore, two matrices C_+ and C_- are defined as follows:

$$C_{+} = \max(C, \mathbf{0}), \ C_{-} = \max(-C, \mathbf{0}).$$

Then we have

$$(C_{+})_{eu} = \begin{cases} 1 & \text{if } t(e) = u, \\ 0 & \text{otherwise,} \end{cases} \quad (C_{-})_{eu} = \begin{cases} 1 & \text{if } o(e) = u, \\ 0 & \text{otherwise.} \end{cases}$$

That is, C_+ and C_- are the terminus-arc incidence matrix and the origin-arc incidence matrix of D, respectively. For brevity, we set

$$K = C_+, \ L = C_-.$$

Three $M \times M$ diagonal matrices $B' = (B'_{ef}), D_1 = (D^{(1)}_{ef})$ and $D'_2 = (D^{(2)}_{ef})$ are given by

$$B'_{ef} = \begin{cases} \beta_e & \text{if } e = f, \\ 0 & \text{otherwise}, \end{cases} \quad D^{(1)}_{ef} = \begin{cases} \delta_{o(e)} & \text{if } e = f, \\ 0 & \text{otherwise}, \end{cases} \quad D^{(2)}_{ef} = \begin{cases} \delta_{t(e)} & \text{if } e = f, \\ 0 & \text{otherwise}, \end{cases}$$

Then KB' and D'_2L^T are the following matrices:

$$(KB')_{ue} = \begin{cases} \beta_e & \text{if } t(e) = u, \\ 0 & \text{otherwise,} \end{cases} \quad (D'_2 L^T)_{ue} = \begin{cases} \delta_{t(e)} & \text{if } o(e) = u, \\ 0 & \text{otherwise.} \end{cases}$$

Next, let

$$M_0 = |\{e \in A(D) \mid e^{-1} \notin A(D)\}|, \ M_1 = |\{e \in A(D) \mid e^{-1} \in A(D)\}|/2.$$

Note that $M = M_0 + 2M_1$. Furthermore, let

$$A(D) = \{e_1, \dots, e_{M_0}, f_1, \dots, f_{M_1}, f_1^{-1}, \dots, f_{M_1}^{-1}\}.$$

We consider the matrices under this order. The matrix $M \times M$ matrix J is defined as follows:

$$J = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_M \\ \mathbf{0} & \mathbf{I}_M & \mathbf{0} \end{bmatrix}.$$

Then the following result holds.

Proposition 1 The non-backtracking matrix H of D is expressed as follows:

$$H = K^T L - J.$$

Here, we set $F = K^T L$. Then we have

$$H = F - J, \ H^T = LK^T - J = F^T - J.$$

The non-backtracking matrix of a digraph is a digraph version of the edge matrix of a graph. Furthermore, the Metzler matrix $\mathcal{A} = \mathcal{A}(D)$ is written as follows:

$$\mathcal{A} = \begin{bmatrix} -\mathbf{D} & KB' \\ D'_2 L^T & H^T B' - B' - D'_1 - D'_2 \end{bmatrix}.$$

 Set

$$E = B' + D_1' + D_2'.$$

Then the following result holds for det $(I_{N+M} - u\mathcal{A})$. The Kronecker product $\mathbf{A} \otimes \mathbf{B}$ of matrices \mathbf{A} and \mathbf{B} is considered as the matrix \mathbf{A} having the element a_{ij} replaced by the matrix $a_{ij}\mathbf{B}$.

Theorem 5 Let D be a connected digraph with N vertices and M arcs. Furthermore, let

 $A(D) = \{e_1, \dots, e_{M_0}, f_1, \dots, f_{M_1}, f_1^{-1}, \dots, f_{M_1}^{-1}\},\$

where $e_i^{-1} \notin A(D)$ ($\leq i \leq M_0$). Then

$$\det(I_{n+M} - u\mathcal{A}) = \prod_{j=1}^{M_0} (1 + \gamma_{e_j} u) \prod_{k=1}^{M_1} G_k \det(I_N + uD - uA_{nsym}^T - uA_{sym} + u^2 \tilde{D}),$$

where $\gamma_e = \beta_e + \delta_{o(e)} + \delta_{t(e)}$ for each $e \in A(D)$, and

$$G_k = 1 + (\gamma_{f_k} + \gamma_{f_k^{-1}})u + u^2(\gamma_{f_k}\gamma_{f_k^{-1}} - \beta_{f_k}\beta_{f_k^{-1}}) \ (1 \le k \le M_1).$$

Proof. At first, we have

$$\det(I_{N+M} - u\mathcal{A})$$

$$= \det \begin{bmatrix} I_N + u\mathbf{D} & -uKB' \\ -uD'_2L^T & I_M - u(H^TB' - E) \end{bmatrix}$$

$$\times \det \begin{bmatrix} I_N & u(I_N + u\mathbf{D})^{-1}KB' \\ \mathbf{0} & I_M \end{bmatrix}$$

$$= \det \begin{bmatrix} I_N + u\mathbf{D} & \mathbf{0} \\ -uD'_2L^T & I_M - u(H^TB' - E) - u^2D'_2L^T(I_N + u\mathbf{D})^{-1}KB' \end{bmatrix}.$$

By Proposition 1, we get

$$det(I_{N+M} - u\mathcal{A}) = det(I_N + u\mathbf{D}) det(I_M - u(H^TB' - E) - u^2D'_2L^T(I_N + u\mathbf{D})^{-1}KB') = det(I_N + u\mathbf{D}) det(I_M + uE - u(L^TKB' - JB') - u^2D'_2L^T(I_N + u\mathbf{D})^{-1}KB') = det(I_N + u\mathbf{D}) det(I_M + u(JB' + E) - u(L^T + uD'_2L^T(I_N + u\mathbf{D})^{-1})KB') = det(I_N + u\mathbf{D}) det(I_M - u(L^T + uD'_2L^T(I_N + u\mathbf{D})^{-1})KB'(I_M + u(JB' + E)^{-1}) \times det(I_M + u(JB' + E)).$$

From Theorem 4, we obtain

$$det(I_{N+M} - u\mathcal{A}) = det(I_N + u\mathbf{D}) det(I_N - uKB'(I_M + u(JB' + E))^{-1}(L^T + uD'_2L^T(I_N + u\mathbf{D})^{-1})) \times det(I_M + u(JB' + E)) = det((I_N + u\mathbf{D}) - uKB'(I_M + u(JB' + E))^{-1}(L^T(I_N + u\mathbf{D}) + uD'_2L^T)) \times det(I_M + u(JB' + E)) = det(I_N + u\mathbf{D} - uKB'(I_M + u(JB' + E))^{-1}(L^T + u(L^T\mathbf{D} + D'_2L^T)) \times det(I_M + u(JB' + E)).$$

Next, we consider $\det(I_M + u(JB' + E))$. Let three diagonal matrices B_0 , B_1 and B_2 be given as follows:

$$(B_0)_{e_i e_i} = \beta_{e_i}, \ (B_1)_{f_j f_j} = \beta_{f_j}, \ (B_2)_{f_j^{-1} f_j^{-1}} = \beta_{f_j^{-1}} \ (1 \le i \le M_0; 1 \le j \le M_1).$$

Then we get

$$(I_{N+M}+J)B' = \begin{bmatrix} B_0 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & B_1 & B_2 \\ \mathbf{0} & B_1 & B_2 \end{bmatrix}.$$

Thus, we have

$$JB' + E = \begin{bmatrix} F_0 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & F_1 & B_2 \\ \mathbf{0} & B_1 & F_2 \end{bmatrix},$$

where F_0 , F_1 and F_2 are $M_0 \times M_0$, $M_1 \times M_1$ and $M_1 \times M_1$ diagonal matrices such that

$$(F_0)_{e_i e_i} = \gamma_{e_i}, \ (F_1)_{f_j f_j} = \gamma_{f_j}, \ (F_2)_{f_j^{-1} f_j^{-1}} = \gamma_{f_j^{-1}} \ (1 \le i \le M_0, 1 \le j \le M_1).$$

Therefore,

$$\begin{aligned} \det(I + u(JB' + E)) \\ &= \det \begin{bmatrix} I + uF_0 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & I + uF_1 & uB_2 \\ \mathbf{0} & uB_1 & I + uF_2 \end{bmatrix} \\ &= \det(I + uF_0) \det \begin{bmatrix} I + uF_1 & uB_2 \\ uB_1 & I + uF_2 \end{bmatrix} \det \begin{bmatrix} I & -u(I + uF_1)^{-1}B_2 \\ \mathbf{0} & I \end{bmatrix} \\ &= \det(I + uF_0) \det \begin{bmatrix} I + uF_1 & \mathbf{0} \\ uB_1 & I + uF_2 - u^2B_1(I + uF_1)^{-1}B_2 \end{bmatrix} \\ &= \det(I + uF_0) \det(I + uF_1) \det(I + uF_2 - u^2B_1(I + uF_1)^{-1}B_2). \end{aligned}$$

But, we get

$$B_1(i+uF_1)^{-1}B_2 = \bigoplus_{j=1}^{M_1} \left(\frac{\beta_{f_j}\beta_{f_j}^{-1}}{1+\gamma_{f_j}u}\right).$$

Thus, we have

$$\det(I + u(JB' + E)) = \det(I + uF_0) \det((I + uF_1)(I + uF_2) - u^2(I + uF_1)B_1(I + uF_1)^{-1}B_2).$$

Since

$$(I+uF_1)B_1(I+uF_1)^{-1}B_2 = \bigoplus_{j=1}^{M_1} \left(\beta_{f_j}\beta_{f_j^{-1}}\right),$$

$$\begin{aligned} \det(I + u(JB' + E)) \\ &= \det(I + uF_0) \det((I + uF_1)(I + uF_2) - u^2 \oplus_{j=1}^{M_1} (\beta_{f_j} \beta_{f_j^{-1}})) \\ &= \prod_{j=1}^{M_0} (1 + \gamma_{e_j} u) \prod_{k=1}^{M_1} (1 + \gamma_{f_k} u)(1 + \gamma_{f_k^{-1}} u) - \beta_{f_k} \beta_{f_k^{-1}}) \\ &= \prod_{j=1}^{M_0} (1 + \gamma_{e_j} u) \prod_{k=1}^{M_1} (1 + (\gamma_{f_k} + \gamma_{f_k^{-1}}) u + u^2(\gamma_{f_k} \gamma_{f_k^{-1}} - \beta_{f_k} \beta_{f_k^{-1}})). \end{aligned}$$

Now, we consider $(I_M + u(JB' + E))^{-1}$ by the Schur complement on the inverse matrix (see [24]). If the suitable matrices are nonsingular, then the inverse matrix U^{-1} of a square matrix

$$U = \begin{bmatrix} A & B \\ C & D \end{bmatrix},$$
$$U^{-1} = \begin{bmatrix} (A - BD^{-1}C)^{-1} & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}.$$

Substituting $A = I + F_1 u$, $B = uB_2$, $C = uB_1$ and $D = I + F_2 u$, we have

$$(A - BD^{-1}C)^{-1} = \bigoplus_{j=1}^{M_1} \left(\frac{1 + \gamma_{f_k^{-1}u}}{G_k}\right),$$
$$(D - CA^{-1}B)^{-1} = \bigoplus_{j=1}^{M_1} \left(\frac{1 + \gamma_{f_k}u}{G_k}\right),$$
$$A^{-1}B(D - CA^{-1}B)^{-1} = \bigoplus_{j=1}^{M_1} \left(\frac{u\beta_{f_k^{-1}}}{G_k}\right),$$
$$(D - CA^{-1}B)^{-1}CA^{-1} = \bigoplus_{j=1}^{M_1} \left(\frac{u\beta_{f_k}}{G_k}\right),$$

that is,

$$\begin{bmatrix} I+F_1u & uB_2\\ uB_1 & I+F_2u \end{bmatrix}^{-1} = \begin{bmatrix} \oplus_{j=1}^{M_1} \left(\frac{1+\gamma_{f_k}^{-1}u}{G_k}\right) & \oplus_{j=1}^{M_1} \left(\frac{u\beta_{f_k}^{-1}}{G_k}\right)\\ \oplus_{j=1}^{M_1} \left(\frac{u\beta_{f_k}}{G_k}\right) & \oplus_{j=1}^{M_1} \left(\frac{1+\gamma_{f_k}u}{G_k}\right) \end{bmatrix}.$$

Thus,

$$(I_M + u(JB' + E))^{-1}$$

$$= \begin{bmatrix} \oplus_{j=1}^{M_0} \left(\frac{1}{1+\gamma_{e_j}^{-1}u}\right) & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \oplus_{j=1}^{M_1} \left(\frac{1+\gamma_{f_k}^{-1}u}{G_k}\right) & \oplus_{j=1}^{M_1} \left(\frac{u\beta_{f_k}^{-1}}{G_k}\right) \\ \mathbf{0} & \oplus_{j=1}^{M_1} \left(\frac{u\beta_{f_k}}{G_k}\right) & \oplus_{j=1}^{M_1} \left(\frac{1+\gamma_{f_k}u}{G_k}\right) \end{bmatrix}.$$

Now, let

$$M = KB'(I_M + u(JB' + E))^{-1}(L^T + u(L^T\mathbf{D} + D'_2L^T)).$$

Furthermore, put

$$\epsilon_e = \delta_{o(e)} + \delta_{t(e)}$$
 for $e \in A(D)$.

Then we get

$$(L^T + u(L^T \mathbf{D} + D'_2 L^T))_{eu} = \begin{cases} 1 + u\epsilon_e & \text{if } o(e) = u, \\ 0 & \text{otherwise.} \end{cases}$$

Thus, the entries of M are given as follows:

$$M_{uv} = \begin{cases} \beta_{e^{-1}}(1 + \epsilon_{e^{-1}}u)/(1 + \gamma_{e^{-1}}u) & \text{if } e = (u, v) \in A(D) \text{ and } e^{-1} \notin A(D), \\ \beta_{e^{-1}}(1 + \gamma_{e}u)(1 + \epsilon_{e^{-1}}u)/G_{e} & \text{if } e = (u, v), e^{-1} \in A(D), \\ -u \sum_{o(e)=u} \beta_{e}\beta_{e^{-1}}(1 + \epsilon_{e}u)/G_{e} & \text{if } u = v, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, it follows that

$$\det(I_{N+M} - u\mathcal{A}) = \prod_{j=1}^{M_0} (1 + \gamma_{e_j} u) \prod_{k=1}^{M_1} G_k \det(I_N + uD - uM).$$

Now, we define three $N \times N$ matrices A_{nsym} , A_{sym} and \tilde{D} as follows:

$$(A_{nsym})_{uv} = \begin{cases} (\beta_{e-1}(1+\epsilon_{e^{-1}}u)/(1+\gamma_{e^{-1}}u)) & \text{if } e = (u,v) \in A(D) \text{ and } e^{-1} \notin A(D), \\ 0 & \text{otherwise}, \end{cases}$$
$$(A_{sym})_{uv} = \begin{cases} (\beta_{e-1}(1+\gamma_{e}u)(1+\epsilon_{e^{-1}}u)/G_{e}) & \text{if } e = (u,v), e^{-1} \in A(D), \\ 0 & \text{otherwise}, \end{cases}$$
$$(\tilde{D})_{uv} = \begin{cases} -u \sum_{o(e)=u} (\beta_{e}\beta_{e-1}(1+\epsilon_{e}u)/G_{e}) & \text{if } u = v, \\ 0 & \text{otherwise}. \end{cases}$$

Hence,

$$\det(I_{n+M} - u\mathcal{A}) = \prod_{j=1}^{M_0} (1 + \gamma_{e_j} u) \prod_{k=1}^{M_1} G_k \det(I_N + uD - uA_{nsym} - uA_{sym}^T + u^2 \tilde{D}).$$

6 The spectrum for the Metzler matrix of the symmetric digraph of a graph

Let D be a connected digraph with N vertices and M arcs, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in A(D)$ and $u \in V(D)$. Then we have

$$\gamma_e = \beta + 2\delta - \gamma \ (const), \ \epsilon_e = 2\delta = \epsilon \ (const) \ for \ each \ e \in A(D)$$

and

$$G_j = (1 + (\beta + 2\delta)u)^2 - \beta^2 u^2 = 1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2 = G \quad (1 \le j \le M_1).$$

Thus, we have

$$A_{nsym} = \frac{\beta(1+\epsilon u)}{1+\gamma u} A_0, \quad A_{sym} = \frac{\beta(1+\epsilon u)(1+\gamma u)}{G} A_1,$$
$$\tilde{D} = \frac{\beta^2(1+\epsilon u)}{G} D_0,$$

where three matrices A_0 , A_1 and D_0 are given as follows:

$$(A_0)_{uv} = \begin{cases} 1 & \text{if } (u,v) \in A(D) \text{ and } (v,u) \notin A(D), \\ 0 & \text{otherwise,} \end{cases} \qquad (A_1)_{uv} = \begin{cases} 1 & \text{if } (u,v), (v,u) \in A(D), \\ 0 & \text{otherwise,} \end{cases}$$
$$(D_0)_{uv} = \begin{cases} \deg^+ u & \text{if } u = v, \\ 0 & \text{otherwise.} \end{cases}$$

Here, deg $+ u = |\{e \in A(D) \mid o(e) = u\}|$. Thus, we have

$$\begin{aligned} \det(I_{n+M} - u\mathcal{A}) &= (1 + \gamma u)^{M_0} G^{M_1} \\ &\times \det(I_N + \delta u I_N - u \frac{\beta(1 + \epsilon u)}{1 + \gamma u} A_0^T - u \frac{\beta(1 + \gamma u)(1 + \epsilon u)}{G} A_1 + u^2 \frac{\beta^2(1 + \epsilon u)}{G} D_0) \\ &= (1 + \gamma u)^{M_0 - N} G^{M_1 - N} \\ &\times \det((1 + \delta u)(1 + \gamma u)(1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2) I_N \\ &- \beta u(1 + \epsilon u)(1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2) A_0^T - u\beta(1 + \gamma u)^2(1 + \epsilon u) A_1 + \beta^2(1 + \gamma u)(1 + \epsilon u) D_0) \\ &= (1 + \gamma u)^{M_0 - N} (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{M_1 - N} \\ &\times \det((1 + \delta u)(1 + (\beta + 2\delta)u)(1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2) I_N \\ &- (1 + 2\delta u)\beta u((1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2) A_0^T + (1 + (\beta + 2\delta)u)^2 A_1) - \beta(1 + (\beta + 2\delta)u) D_0). \end{aligned}$$

Therefore, we obtain

Corollary 1 Let D be a connected graph with N vertices v_1, \dots, v_N and M arcs, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in A(D)$ and $u \in V(D)$. Then

$$det(I_{n+M} - u\mathcal{A}) = (1 + \gamma u)^{M_0 - N} (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{M_1 - N} \\ \times det((1 + \delta u)(1 + (\beta + 2\delta)u)(1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)I_N \\ - (1 + 2\delta u)\beta u((1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)A_0^T + (1 + (\beta + 2\delta)u)^2A_1) - \beta(1 + (\beta + 2\delta)u)D_0).$$

Next, let D = G be a connected (undirected) graph with N vertices and M edges, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in D(G)$ and $u \in V(G)$. Then we have

$$M_0 = 0, \ M_1 = M, \ |D(G)| = 2M.$$

Furthermore, we get

$$A_0 = \mathbf{0}, \ A_1 = \mathbf{A}(G) \ and \ D_0 = \mathbf{D}_G.$$

Thus,

Corollary 2 Let D = G be a connected (undirected) graph with N vertices and M edges, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in D(G)$ and $u \in V(G)$. Then

$$det(I_{n+2M} - u\mathcal{A}) = (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{M-N}$$

× det((1 + δu)(1 + 2(β + 2 δ)u + 4 $\delta(\delta$ + 1)u^2)I_N
- (1 + 2 δu) βu (1 + (β + 2 δ)u) \mathbf{A} (G) + β^2 (1 + 2 δu) $u^2\mathbf{D}_G$).

Finally, let G be a connected d-regular graph with N vertices and M edges, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in D(G)$ and $u \in V(G)$. By Corollary 2, we have

$$det(I_{n+2M} - u\mathcal{A}) = (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{M-N} \\ \times det((1 + \delta u)(1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)I_N \\ - (1 + 2\delta u)\beta u(1 + (\beta + 2\delta)u)\mathbf{A}(G) + d\beta^2(1 + 2\delta u)u^2I_N) \\ = (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{M-N} \\ \times det(\{1 + (2\beta + 5\delta)u + (2\delta(\beta + 4\delta + 2) + d\beta^2)u^2 + 2\delta(2\delta(\delta + 1) + d\beta^2)u^3\}I_N \\ - \beta(1 + 2\delta u)(1 + (\beta + 2\delta)u)\mathbf{A}(G)).$$

Therefore,

Corollary 3 Let D = G be a connected d-regular graph with N vertices and M edges, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in D(G)$ and $u \in V(G)$. Then

$$\det(I_{n+2M} - u\mathcal{A}) = (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{M-N} \\ \times \det(\{1 + (2\beta + 5\delta)u + (2\delta(\beta + 4\delta + 2) + d\beta^2)u^2 + 2\delta(2\delta(\delta + 1) + d\beta^2)u^3\}I_N \\ - \beta(1 + 2\delta u)(1 + (\beta + 2\delta)u)\mathbf{A}(G)).$$

From Corollary 3, the following result holds.

Corollary 4 Let D = G be a connected d-regular graph with N vertices and M edges, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in D(G)$ and $u \in V(G)$. Then

$$\det(\lambda I_{n+2M} - \mathcal{A}) = (\lambda^2 + 2(\beta + 2\delta)\lambda + 4\delta(\delta + 1))^{M-N}$$

$$\times \det(\{\lambda^3 + (2\beta + 5\delta)\lambda^2 + (2\delta(\beta + 4\delta + 2) + d\beta^2)\lambda + 2\delta(2\delta(\delta + 1) + d\beta^2)\}I_N$$

$$-\beta(\lambda + 2\delta)(\lambda + (\beta + 2\delta))\mathbf{A}(G)).$$

Substituting $u = 1/\lambda$ to Corollary 4, the spectra for \mathcal{A} of a regular graph are obtained.

Corollary 5 Let D = G be a connected d-regular graph with N vertices and M edges, and let $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in D(G)$ and $u \in V(G)$. Then the eigenvalues of A are given as follows:

1. 3N eigenvalues: solutions of

$$\lambda^{3} + (2\beta - \beta\mu + 5\delta)\lambda^{2} + (2\delta(\beta + 4\delta + 2) + d\beta^{2} - \beta(\beta + 4\delta)\mu)\lambda + 2\delta(2\delta(\delta + 1) + d\beta^{2} - \beta(\beta + 2\delta)\mu) = 0, \quad \mu \in \operatorname{Spec}(\mathbf{A}(G)),$$

2. 2(M-N) eigenvalues: $\lambda = -(\beta + 2\delta) \pm \sqrt{\beta^2 + 4\delta(\beta - 1)}$.

Proof. By Corollary 4, we have

$$\det(\lambda I_{n+2M} - \mathcal{A}) = (\lambda^2 + 2(\beta + 2\delta)\lambda + 4\delta(\delta + 1))^{M-N}$$
$$\times \prod_{\mu \in Spec(\mathbf{A}(G))} \{\lambda^3 + (2\beta - \beta\mu + 5\delta)\lambda^2 + (2\delta(\beta + 4\delta) + 2) + d\beta^2$$
$$-\beta(\beta + 4\delta)\mu)\lambda + 2\delta(2\delta(\delta + 1) + d\beta^2 - \beta(\beta + 2\delta)\mu)\}.$$

Solving $\lambda^2 + 2(\beta + 2\delta)\lambda + 4\delta(\delta + 1) = 0$, we obtain

$$\lambda = -(\beta + 2\delta) \pm \sqrt{\beta^2 + 4\delta(\beta - 1)}.$$

The result follows. \Box

7 A walk-type zeta function with respect to the Metzler matrix of the symmetric digraph of a finite torus

As an application, we present the Walk/Zeta Correspondence (see [11]) for the Metzler matrix of the symmetric digraph of the *d*-dimensional torus $G = T_N^d$. For a natural number $N \geq 2$, let $G = T_N^d$ be the *d*-dimensional finite torus (graph). Furthermore, assume that $\beta_e = \beta$, $\delta_u = \delta$ for each $e \in D(T_N^d)$ and $u \in V(T_N^d)$. Then we have

$$n = |V(T_N^d)| = N^d, \ m = |E(T_N^d)| = dN^d.$$

Note that T_N^d is a 2*d*-regular graph.

Similarly to the Walk/Zeta Correspondence of [11], we introduce a zeta function for the Metzler matrix \mathcal{A} of the symmetric digraph of $G = T_N^d$ as follows:

$$\overline{\zeta}(\mathcal{A}, T_N^d, u) = \det(\mathbf{I}_{(2d+1)N^d} - u\mathcal{A})^{-1/N^d}.$$

Let $\mathbf{A} = \mathbf{A}(G) = \mathbf{A}(T_N^d)$. It is known that

Spec
$$(T_N^d) = \left\{ 2\sum_{j=1}^d \cos\left(\frac{2\pi k_j}{N}\right) \mid k_j = 0, 1, \dots, N-1 \ (1 \le j \le d) \right\}.$$

By Theorem 5 and Corollary 3, we have

$$\begin{split} \overline{\zeta}(\mathcal{A}, T_N^d, u)^{-1} &= \det(\mathbf{I}_{(2d+1)N^d} - u\mathcal{A})^{1/N^d} \\ &= \{(1+2(\beta+2\delta)u + 4\delta(\delta+1)u^2)^{(2d-1)N^d} \times \det(\{1+(2\beta+5\delta)u + (2\delta(\beta+4\delta+2)+2d\beta^2)u^2 + 2\delta(2\delta(\delta+1)+2d\beta^2)u^3\}I_{N^d} - \beta(1+2\delta u)(1+(\beta+2\delta)u)\mathbf{A}(T_N^d))\}^{1/N^d} \\ &= (1+2(\beta+2\delta)u + 4\delta(\delta+1)u^2)^{2d-1}\prod_{k_1=0}^{N-1} \cdots \prod_{k_d=0}^{N-1}\{\{1+(2\beta+5\delta)u + 2(\delta(\beta+4\delta+2)+d\beta^2)u^2 + 4\delta(\delta+1) + d\beta^2)u^3\} \\ &\quad + 2(\delta(\beta+4\delta+2) + d\beta^2)u^2 + 4\delta(\delta+1) + d\beta^2)u^3\} \\ &\quad - 2\beta(1+2\delta u)(1+(\beta+2\delta)u)\sum_{j=1}^d \cos\left(\frac{2\pi k_j}{N}\right)\}^{1/N^d} \\ &= (1+2(\beta+2\delta)u + 4\delta(\delta+1)u^2)^{2d-1} \\ &\quad \times \exp(\frac{1}{N^d}\sum_{k_1=0}^{N-1} \cdots \sum_{k_d=0}^{N-1} \log\{\{1+(2\beta+5\delta)u + 2(\delta(\beta+4\delta+2) + d\beta^2)u^2 + 4\delta(\delta+1) + d\beta^2)u^3\} \\ &\quad - 2\beta(1+2\delta u)(1+(\beta+2\delta)u)\sum_{j=1}^d \cos\left(\frac{2\pi k_j}{N}\right)\}). \end{split}$$

When $N \to \infty$, we get

$$\begin{split} &\lim_{N \to \infty} \overline{\zeta} (\mathcal{A}, T_N^d, u)^{-1} \\ &= (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{2d - 1} \\ &\times \exp(\int_0^{2\pi} \cdots \int_0^{2\pi} \log\{\{1 + (2\beta + 5\delta)u + 2(\delta(\beta + 4\delta + 2) + d\beta^2)u^2 + 4\delta(\delta + 1) + d\beta^2)u^3\} \\ &- 2\beta(1 + 2\delta u)(1 + (\beta + 2\delta)u) \sum_{j=1}^d \cos\theta_j \frac{d\theta_1}{2\pi} \cdots \frac{d\theta_d}{2\pi}\}). \end{split}$$

Therefore, we obtain

Theorem 6 Let $G = T_N^d$ be the d-dimensional finite torus. Then

$$\begin{split} \overline{\zeta}(\mathcal{A}, T_N^d, u)^{-1} &= (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{2d-1} \\ &\times \exp\left[\frac{1}{N^d} \sum_{k_1=0}^{N-1} \cdots \sum_{k_d=0}^{N-1} \log\left\{\{1 + (2\beta + 5\delta)u + 2(\delta(\beta + 4\delta + 2) + d\beta^2)u^2 \\ &+ 4\delta(\delta + 1) + d\beta^2)u^3\}I_{N^d} - 2\beta(1 + 2\delta u)(1 + (\beta + 2\delta)u)\sum_{j=1}^d \cos\left(\frac{2\pi k_j}{N}\right)\}\right] \end{split}$$

and

$$\begin{split} \lim_{N \to \infty} \overline{\zeta} (\mathcal{A}, T_N^d, u)^{-1} \\ &= (1 + 2(\beta + 2\delta)u + 4\delta(\delta + 1)u^2)^{2d - 1} \\ &\times \exp\Big[\int_0^{2\pi} \cdots \int_0^{2\pi} \log\Big\{\{1 + (2\beta + 5\delta)u + 2(\delta(\beta + 4\delta + 2) + d\beta^2)u^2 + 4\delta(\delta + 1) + d\beta^2)u^3\} \\ &- 2\beta(1 + 2\delta u)(1 + (\beta + 2\delta)u)\sum_{j=1}^d \cos\theta_j \frac{d\theta_1}{2\pi} \cdots \frac{d\theta_d}{2\pi}\Big\}\Big]. \end{split}$$

References

- A. Ambainis, Quantum walks and their algorithmic applications, Int. J. Quantum Inf. 1 (2003), 507–518.
- [2] H. Bass, The Ihara-Selberg zeta function of a tree lattice, Internat. J. Math. 3 (1992), 717-797.
- [3] L. Farina and S. Rinaldi, Positive Linear Systems-Theory and Applications, John Wiley & Sons, Inc. New York, NY, 2000.
- [4] H. Flanders, The elementary divisors of AB and BA, Proc. Am. Math. Soc. 2 (1951), 871-874.
- [5] D. Foata and D. Zeilberger, A combinatorial proof of Bass's evaluations of the Ihara-Selberg zeta function for graphs, Trans. Amer. Math. Soc. 351 (1999), 2257-2274.
- [6] K. Hashimoto, Zeta Functions of Finite Graphs and Representations of p-Adic Groups, Adv. Stud. Pure Math. Vol. 15, Academic Press, New York, 1989, pp. 211-280.
- [7] Y. Ihara, On discrete subgroups of the two by two projective linear group over *p*-adic fields, J. Math. Soc. Japan 18 (1966), 219-235.
- [8] J. Kempe, Quantum random walks an introductory overview, Contemporary Physics 44 (2003), 307–327.
- [9] V. Kendon, Decoherence in quantum walks a review, Math. Struct. in Comp. Sci. 17 (2007), 1169–1220.
- [10] T. Komatsu, N. Konno and I. Sato, Grover/Zeta Correspondence based on the Konno-Sato theorem, Quantum Inf. Process. 20, 268 (2021).
- [11] T. Komatsu, N. Konno and I. Sato, Walk/Zeta Correspondence, arXiv:2104.10287 (2021).

- [12] T. Komatsu, N. Konno and I. Sato, IPS/Zeta Correspondence, Quantum Inf. Comput. 22 (2022), 251–269.
- [13] T. Komatsu, N. Konno and I. Sato, Vertex-Face/Zeta Correspondence, J. Algebraic Combin. (in press), arXiv:2107.03300 (2021).
- [14] T. Komatsu, N. Konno and I. Sato, CTM/Zeta Correspondence, Quantum Stud.: Math. Found. 9 (2022), 165–173.
- [15] T. Komatsu, N. Konno, I. Sato and S. Tamura, A Generalized Grover/Zeta Correspondence, arXiv:2201.03973 (2022).
- [16] N. Konno, Quantum Walks, In: Lecture Notes in Mathematics: Vol.1954, pp.309–452, Springer-Verlag, Heidelberg (2008).
- [17] N. Konno and S. Tamura, Walk/Zeta Correspondence for quantum and correlated random walks, Yokohama Math. J. 67 (2021), 125–152.
- [18] M. Kotani and T. Sunada, Zeta functions of finite graphs, J. Math. Sci. U. Tokyo 7 (2000) 7-25.
- [19] N. Masuda, V. M. Preciado and M. Ogura, Analysis of the susceptible-infectedsusceptible epidemic dynamics in networks via the non-backtracking matrix, IMA Journal of Applied Mathematics 85 (2020), 214-230.
- [20] H. M. Stark and A. A. Terras, Zeta functions of finite graphs and coverings, Adv. Math. 121 (1996), 124-165.
- [21] T. Sunada, L-Functions in Geometry and Some Applications, in Lecture Notes in Math., Vol. 1201, Springer-Verlag, New York, 1986, pp. 266-284.
- [22] T. Sunada, Fundamental Groups and Laplacians (in Japanese), Kinokuniya, Tokyo, 1988.
- [23] S. E. Venegas-Andraca, Quantum walks: a comprehensive review, Quantum Inf. Process. 11 (2012), 1015-1106.
- [24] F. Zhang, The Schur Complement and Its Applications, Springer 2005.