

# RATE OF MOMENT CONVERGENCE IN THE CENTRAL LIMIT THEOREM FOR ELEPHANT RANDOM WALKS

MASAFUMI HAYASHI, SO OSHIRO, AND MASATO TAKEI

ABSTRACT. The one-dimensional elephant random walk is a typical model of discrete-time random walk with step-reinforcement, and is introduced by Schütz and Trimper (2004). It has a parameter  $\alpha \in (-1, 1)$ : The case  $\alpha = 0$  corresponds to the simple symmetric random walk, and when  $\alpha > 0$  (resp.  $\alpha < 0$ ), the mean displacement of the walker at time  $n$  grows (resp. vanishes) like  $n^\alpha$ . The walk admits a phase transition at  $\alpha = 1/2$  from the diffusive behavior to the superdiffusive behavior. In this paper, we study the rate of the moment convergence in the central limit theorem for the position of the walker when  $-1 < \alpha \leq 1/2$ . We find a crossover phenomenon in the rate of convergence of the  $2m$ -th moments with  $m = 2, 3, \dots$  inside the diffusive regime  $-1 < \alpha < 1/2$ .

## 1. INTRODUCTION

The elephant random walk, introduced by Schütz and Trimper [11], is one of the simplest models of step-reinforced random walks: Let  $p, q \in (0, 1)$ .

- The first step  $X_1$  of the walker is  $+1$  with probability  $q$ , and  $-1$  with probability  $1 - q$ .
- For each  $n \in \mathbb{N} := \{1, 2, \dots\}$ , let  $U_n$  be uniformly distributed on  $\{1, \dots, n\}$ , and

$$X_{n+1} = \begin{cases} X_{U_n} & \text{with probability } p, \\ -X_{U_n} & \text{with probability } 1 - p. \end{cases}$$

Each of choices in the above procedure is made independently. The sequence  $\{X_i\}$  generates a one-dimensional random walk  $\{S_n\}$  by

$$S_0 := 0, \quad \text{and} \quad S_n = \sum_{i=1}^n X_i \quad \text{for } n \in \mathbb{N}.$$

We begin with a brief review of fundamental results. The new parameters defined by

$$\alpha := 2p - 1 \in (-1, 1) \quad \text{and} \quad \beta := E[X_1] = 2q - 1 \in (-1, 1)$$

will be convenient later. Let  $\mathcal{F}_n$  be the  $\sigma$ -algebra generated by  $X_1, \dots, X_n$ . For  $n \in \mathbb{N}$ , the conditional distribution of  $X_{n+1}$  given the history up to time

---

M.H. is partially supported by JSPS KAKENHI Grant Numbers JP19H01791 and JP21K03272. M.T. is partially supported by JSPS KAKENHI Grant Numbers JP19H01793 and JP19K03514.

$n$  is

$$\begin{aligned} & P(X_{n+1} = \pm 1 \mid \mathcal{F}_n) \\ &= \frac{\#\{i = 1, \dots, n : X_i = \pm 1\}}{n} \cdot p + \frac{\#\{i = 1, \dots, n : X_i = \mp 1\}}{n} \cdot (1-p) \\ &= \frac{n \pm S_n}{2n} \cdot p + \frac{n \mp S_n}{2n} \cdot (1-p) = \frac{1}{2} \left( 1 \pm \alpha \cdot \frac{S_n}{n} \right). \end{aligned} \quad (1.1)$$

The conditional expectations of  $X_{n+1}$  and  $S_{n+1}$  are

$$E[X_{n+1} \mid \mathcal{F}_n] = \alpha \cdot \frac{S_n}{n}, \quad (1.2)$$

and

$$E[S_{n+1} \mid \mathcal{F}_n] = \left( 1 + \frac{\alpha}{n} \right) S_n. \quad (1.3)$$

We set

$$a_n := \prod_{j=1}^{n-1} \left( 1 + \frac{\alpha}{j} \right) = \prod_{j=1}^{n-1} \frac{j + \alpha}{j} = \frac{\Gamma(n + \alpha)}{\Gamma(n)\Gamma(1 + \alpha)}. \quad (1.4)$$

Stirling's formula yields that for  $a, b \in \mathbb{R}$ ,

$$\frac{\Gamma(n + a)}{\Gamma(n + b)} \sim n^{a-b} \quad (n \rightarrow \infty), \quad (1.5)$$

and

$$a_n \sim \frac{n^\alpha}{\Gamma(1 + \alpha)} \quad (n \rightarrow \infty), \quad (1.6)$$

where  $x_n \sim y_n$  means that  $x_n/y_n$  converges to 1 as  $n \rightarrow \infty$ .

Let

$$M_n := \frac{S_n}{a_n} \quad \text{for } n \in \mathbb{N}.$$

Then  $\{M_n\}$  is a martingale with respect to  $\{\mathcal{F}_n\}$ . As  $E[M_1] = E[S_1] = \beta$ , we have

$$E[S_n] = \beta a_n \quad \text{for any } n \in \mathbb{N}. \quad (1.7)$$

By (1.6), the mean displacement is asymptotically of order  $n^\alpha$ , provided that  $\beta \neq 0$ .

By a further calculation (cf. page 12 of [2]),

$$E[(S_n)^2] = \begin{cases} \frac{n}{1-2\alpha} + \frac{1}{2\alpha-1} \cdot \frac{\Gamma(n+2\alpha)}{\Gamma(n)\Gamma(2\alpha)} & (\alpha \neq 1/2), \\ n \sum_{\ell=1}^n \frac{1}{\ell} & (\alpha = 1/2). \end{cases} \quad (1.8)$$

Here we regard

$$\frac{1}{\Gamma(s)} = 0 \quad \text{for } s = 0, -1, -2, \dots \quad (1.9)$$

From (1.5), the asymptotic behavior of the mean square displacement is as follows:

$$E[(S_n)^2] \sim \begin{cases} \frac{n}{1-2\alpha} & (-1 < \alpha < 1/2), \\ n \log n & (\alpha = 1/2), \\ \frac{n^{2\alpha}}{(2\alpha-1)\Gamma(2\alpha)} & (1/2 < \alpha < 1). \end{cases} \quad (1.10)$$

Several limit theorems are obtained in [1, 2, 5, 6, 9]. Here we summarize central limit theorems for  $\{S_n\}$ . When  $-1 < \alpha < 1/2$ , the elephant random walk is diffusive, and the fluctuation is Gaussian:

$$\frac{S_n}{\sqrt{n/(1-2\alpha)}} \xrightarrow{d} N(0, 1) \quad (n \rightarrow \infty). \quad (1.11)$$

where  $\xrightarrow{d}$  denotes the convergence in distribution, and  $N(0, 1)$  is the standard normal distribution. When  $\alpha = 1/2$ , the walk is marginally superdiffusive, but still

$$\frac{S_n}{\sqrt{n \log n}} \xrightarrow{d} N(0, 1) \quad (n \rightarrow \infty). \quad (1.12)$$

On the other hand, if  $\alpha > 1/2$ , then  $\{M_n\}$  is an  $L^2$ -bounded martingale, and there exists a random variable  $M_\infty$  such that  $P(M_\infty \neq 0) > 0$  and

$$M_n = \frac{S_n}{a_n} \rightarrow M_\infty \quad \text{a.s. and in } L^2 \text{ as } n \rightarrow \infty. \quad (1.13)$$

Although  $M_\infty$  is non-Gaussian,

$$\frac{S_n - M_\infty \cdot a_n}{\sqrt{n/(2\alpha-1)}} \xrightarrow{d} N(0, 1) \quad (n \rightarrow \infty). \quad (1.14)$$

## 2. MAIN RESULTS

We denote by  $\mu_k$  the  $k$ -th moment of the standard normal distribution:

$$\mu_k := \begin{cases} 0 & (k = 2m-1), \\ (k-1)!! & (k = 2m), \end{cases} \quad (m \in \mathbb{N}).$$

Then, in view of (1.11) and (1.12), we can expect that

$$\lim_{n \rightarrow \infty} E \left[ \left( \frac{S_n}{\sqrt{n/(1-2\alpha)}} \right)^k \right] = \mu_k \quad (k \in \mathbb{N}) \quad (2.1)$$

holds true if  $-1 < \alpha < 1/2$ , and

$$\lim_{n \rightarrow \infty} E \left[ \left( \frac{S_n}{\sqrt{n \log n}} \right)^k \right] = \mu_k \quad (k \in \mathbb{N}) \quad (2.2)$$

if  $\alpha = 1/2$ . The aim of this paper is not only to show that the above equalities (2.1) and (2.2) are valid, but also to investigate the exact rate of these convergence.

Our first result is about odd order moments.

**Theorem 2.1.** Assume that  $\beta = 2q - 1 \neq 0$ .

(i) If  $-1 < \alpha < 1/2$ , then we have, as  $n \rightarrow \infty$ ,

$$E \left[ \left( \frac{S_n}{\sqrt{n/(1-2\alpha)}} \right)^{2m-1} \right] \sim \frac{\beta\sqrt{1-2\alpha}}{\Gamma(1+\alpha)} \cdot (2m-1)!! \cdot n^{-(1-2\alpha)/2}. \quad (2.3)$$

(ii) If  $\alpha = 1/2$ , then we have, as  $n \rightarrow \infty$ ,

$$E \left[ \left( \frac{S_n}{\sqrt{n \log n}} \right)^{2m-1} \right] \sim \frac{2\beta}{\sqrt{\pi}} \cdot (2m-1)!! \cdot (\log n)^{-1/2}. \quad (2.4)$$

**Remark 2.2.** It follows from (1.1) that if  $\beta = 0$ , then  $\{S_n\}$  has the same law as the process  $\{-S_n\}$ . Hence we have  $E[(S_n)^{2m-1}] = 0$  for any  $n \in \mathbb{N}$ .

The next result shows that for nonpositive  $\alpha$ , the exponent in the rate of convergence of the  $2m$ -th moments with  $m \geq 2$  is different from that of the second moment.

**Theorem 2.3.** The following three assertions hold true:

(i) If  $-1 < \alpha < 0$  and  $\alpha \neq -1/2$ , then

$$E \left[ \left( \frac{S_n}{\sqrt{n/(1-2\alpha)}} \right)^2 \right] - 1 \sim -\frac{1}{\Gamma(2\alpha)} \cdot n^{-(1-2\alpha)} \quad (n \rightarrow \infty). \quad (2.5)$$

The left hand side of (2.5) is identically zero if  $\alpha = 0$  and  $n \geq 1$ , or if  $\alpha = -1/2$  and  $n \geq 2$ . On the other hand, for any  $-1 < \alpha \leq 0$  and  $m = 2, 3, \dots$ , we have that, as  $n \rightarrow \infty$ ,

$$\frac{1}{(2m-1)!!} E \left[ \left( \frac{S_n}{\sqrt{n/(1-2\alpha)}} \right)^{2m} \right] - 1 \sim \frac{m(m-1)}{2} \cdot c(\alpha) \cdot n^{-1}, \quad (2.6)$$

where

$$c(\alpha) = -\frac{2(2\alpha^2 + 1)}{3(1-4\alpha)}. \quad (2.7)$$

(ii) If  $0 < \alpha < 1/2$ , then we have that, as  $n \rightarrow \infty$ ,

$$\frac{1}{(2m-1)!!} E \left[ \left( \frac{S_n}{\sqrt{n/(1-2\alpha)}} \right)^{2m} \right] - 1 \sim -\frac{m}{\Gamma(2\alpha)} \cdot n^{-(1-2\alpha)}, \quad (2.8)$$


holds for any  $m \in \mathbb{N}$ .

(iii) If  $\alpha = 1/2$ , then we have that, as  $n \rightarrow \infty$ ,

$$\frac{1}{(2m-1)!!} E \left[ \left( \frac{S_n}{\sqrt{n \log n}} \right)^{2m} \right] - 1 \sim m\gamma \cdot (\log n)^{-1}. \quad (2.9)$$

holds for any  $m \in \mathbb{N}$ . Here  $\gamma$  is Euler's constant.

Our results are summarized in Figure 1.

FIGURE 1.  $\alpha$ -dependence of the convergence rate.

### 3. LITERATURE REVIEW

Let us review literatures closely related to the current paper.

The rate of convergence in the central limit theorem for  $\{S_n\}$  when  $-1 < \alpha \leq 1/2$  is studied in Fan, Hu, and Ma [7]. Let

$$\sigma_n := a_n \left\{ \sum_{i=1}^n \frac{1}{(a_i)^2} \right\}^{1/2},$$

for  $n \in \mathbb{N}$ . By (1.6) and Lemma A.1 in Appendix,

$$\sigma_n \sim \begin{cases} \sqrt{n/(1-2\alpha)} & (-1 < \alpha < 1/2), \\ \sqrt{n \log n} & (\alpha = 1/2), \end{cases} \quad (n \rightarrow \infty).$$

Let  $Z$  be a random variable with the standard normal distribution, and for another random variable  $X$ , we define

$$D(X) := \sup_{x \in \mathbb{R}} |P(X \leq x) - P(Z \leq x)|.$$

In [7], the following Berry–Esseen type bound is established:

$$D\left(\frac{S_n}{\sigma_n}\right) \leq \begin{cases} \frac{C \log n}{\sqrt{n}} & (-1 < \alpha \leq 0), \\ \frac{C \log n}{n^{(1-2\alpha)/2}} & (0 \leq \alpha < 1/2), \\ \frac{C \log \log n}{\sqrt{\log n}} & (\alpha = 1/2). \end{cases}$$

There is a crossover phenomenon in the *upper bounds* at  $\alpha = 0$ . A similar result on Wasserstein-1 distance is obtained by Ma, El Machkouri, and Fan [10]. Our main results give more precise information about such transition.

Bercu, Chabanol, and Ruch [3] derives the asymptotic behavior of the  $k$ -th moment  $E[(S_n)^k]$  of  $S_n$  when  $\alpha > 1/2$ . In [3], it is shown that  $\{M_n\}$  is an  $L^k$ -bounded martingale for any  $k \in \mathbb{N}$ , which implies that

$$E[(M_n)^k] \rightarrow E[(M_\infty)^k] =: C_k \quad (n \rightarrow \infty).$$

If  $k$  is an odd integer and  $\beta = 0$ , then  $C_k = 0$  (see Remark 2.2). Except this case, the constant  $C_k$  is positive and we have

$$E[(S_n)^k] \sim C_k (a_n)^k \quad (n \rightarrow \infty).$$

The main purpose in [3] is to establish new hypergeometric identities by calculating  $E[(M_\infty)^k]$  in two different ways. One of interesting future problems is to investigate the rate of moment convergence corresponding to (1.14).

We remark that  $c(\alpha)$  defined by (2.7) attains its maximum at  $\alpha = -1/2$ . Bertoin [4] shows that there is a transition at  $\alpha = -1/2$  about the recurrence property of the elephant random walk. Define

$$R := \inf\{n > 0 : S_n = 0\}.$$

As a consequence of the law of the iterated logarithm (see [6] and [2]) we can see that  $P(R < +\infty) = 1$  if  $-1 < \alpha \leq 1/2$ . On the other hand, it follows from (1.13) that  $P(R = +\infty) > 0$  if  $\alpha > 1/2$ . It is proved in [4] that

$$E[R] \begin{cases} < +\infty & (-1 < \alpha < -1/2), \\ = +\infty & (-1/2 \leq \alpha \leq 1/2). \end{cases}$$

#### 4. PROOF OF THEOREM 2.1

Let  $m, n \in \mathbb{N}$ . We start with the following recursion, which is a consequence of (4.3) in [3]:

$$E[(S_{n+1})^{2m-1}] = \sum_{\ell=1}^m \left\{ \binom{2m-1}{2\ell-1} + \frac{\alpha}{n} \binom{2m-1}{2\ell-2} \right\} E[(S_n)^{2\ell-1}]. \quad (4.1)$$

We set

$$\begin{aligned} M_n^{(2m-1)} &:= E[(S_n)^{2m-1}], \\ f_n^{(2m-1)} &:= \sum_{\ell=1}^{m-1} \left\{ \binom{2m-1}{2\ell-1} + \frac{\alpha}{n} \binom{2m-1}{2\ell-2} \right\} M_n^{(2\ell-1)}, \\ g_n^{(2m-1)} &:= 1 + \frac{(2m-1)\alpha}{n}. \end{aligned} \quad (4.2)$$

Then the recursion (4.1) can be put into the following form: For  $n \in \mathbb{N}$ ,

$$M_{n+1}^{(2m-1)} = f_n^{(2m-1)} + g_n^{(2m-1)} M_n^{(2m-1)}. \quad (4.3)$$

It is convenient to define

$$f_0^{(2m-1)} := M_1^{(2m-1)} = E[(X_1)^{2m-1}] = \beta \quad \text{and} \quad g_0^{(2m-1)} := 0.$$

To solve (4.3), we use the following two lemmas.

**Lemma 4.1** (cf. Schütz and Trimper [11], (12) and (13)). Let  $j_0 \in \mathbb{N}$ . The general solution  $\{x_n\}_{n=j_0+1, j_0+2, \dots}$  to the recursion

$$\begin{cases} x_{j_0+1} = f_{j_0}, \\ x_{n+1} = f_n + g_n \cdot x_n \quad (n = j_0 + 1, j_0 + 2, \dots) \end{cases}$$

is given by

$$x_n = x_{j_0+1} \prod_{k=j_0+1}^{n-1} g_k + \sum_{j=j_0+1}^{n-1} f_j \cdot \prod_{k=j+1}^{n-1} g_k \quad (n = j_0 + 1, j_0 + 2, \dots).$$

**Lemma 4.2.** For  $\delta \in \mathbb{R}$ , we define

$$j_0(\delta) := \begin{cases} k & \text{if } \delta = -k \text{ for some } k \in \mathbb{N}, \\ 0 & \text{otherwise,} \end{cases} \quad (4.4)$$

and

$$A_\delta := \begin{cases} k! & \text{if } \gamma = -k \text{ for some } k \in \mathbb{N}, \\ \frac{1}{\Gamma(1 + \delta)} & \text{otherwise.} \end{cases}$$

Then we have

$$\prod_{j=j_0(\delta)+1}^{n-1} \left(1 + \frac{\delta}{j}\right) \sim A_\delta \cdot n^\delta \quad (n \rightarrow \infty).$$

*Proof.* If  $\delta$  is not a negative integer, then for any  $n > 1$ ,

$$\prod_{j=j_0(\delta)+1}^{n-1} \left(1 + \frac{\delta}{j}\right) = \prod_{j=1}^{n-1} \frac{j + \delta}{j} = \frac{\Gamma(n + \delta)}{\Gamma(n)\Gamma(1 + \delta)} \quad (4.5)$$

If  $\delta = -k$  for some  $k \in \mathbb{N}$ , then for any  $n > k + 1$ ,

$$\prod_{j=j_0(\delta)+1}^{n-1} \left(1 + \frac{\delta}{j}\right) = \prod_{j=k+1}^{n-1} \frac{j - k}{j} = \frac{\Gamma(n - k) \times k!}{\Gamma(n)}. \quad (4.6)$$

We obtain the desired result by (1.5).  $\square$

We set  $j_0 = j_0((2m - 1)\alpha)$ . Note that

$$g_{j_0}^{(2m-1)} = 0, \quad \text{and} \quad g_n^{(2m-1)} \neq 0 \quad \text{for all } n > j_0.$$

By Lemma 4.1,  $\{M_n^{(2m-1)}\}$  satisfies

$$M_n^{(2m-1)} = M_{j_0+1}^{(2m-1)} \bar{g}_n^{(2m-1)} + \bar{g}_n^{(2m-1)} \sum_{j=j_0+1}^{n-1} \frac{f_j^{(2m-1)}}{\bar{g}_{j+1}^{(2m-1)}}, \quad (4.7)$$

where

$$\bar{g}_n^{(2m-1)} := \prod_{k=j_0+1}^{n-1} g_k^{(2m-1)} \quad (n = j_0 + 1, j_0 + 2, \dots). \quad (4.8)$$

From Lemma 4.2, we have

$$\bar{g}_n^{(2m-1)} \sim A_{(2m-1)\alpha} n^{(2m-1)\alpha} \quad (n \rightarrow \infty). \quad (4.9)$$

We put

$$F_n^{(2m-1)} := \bar{g}_n^{(2m-1)} \sum_{j=j_0+1}^{n-1} \frac{f_j^{(2m-1)}}{\bar{g}_{j+1}^{(2m-1)}}. \quad (4.10)$$

**4.1. Proof of (2.3) in Theorem 2.1.** Assume that  $-1 < \alpha < 1/2$  and  $\beta \neq 0$ . We define

$$C_\ell = C_\ell(\alpha, \beta) := \frac{(2\ell - 1)!!}{(1 - 2\alpha)^{\ell-1}} \cdot \frac{\beta}{\Gamma(1 + \alpha)} \quad (\ell \in \mathbb{N}).$$

It suffices to show that

$$M_n^{(2\ell-1)} \sim C_\ell n^{\ell-1+\alpha} \quad (n \rightarrow \infty), \quad (4.11)$$

for  $\ell \in \mathbb{N}$ , by induction. We note that (4.11) with  $\ell = 1$  follows from (1.7) and (1.6). Assume that  $m > 1$  and (4.11) holds true for  $\ell = 1, \dots, m-1$ . Then we have that, as  $n \rightarrow \infty$ ,

$$f_n^{(2m-1)} \sim \binom{2m-1}{2m-3} M_n^{(2m-3)} \sim (2m-1)(m-1) C_{m-1} n^{m-2+\alpha}.$$

By (4.9) and Lemma A.1 in Appendix, we have

$$\begin{aligned} F_n^{(2m-1)} &\sim A_{(2m-1)\alpha} n^{(2m-1)\alpha} \sum_{j=j_0+1}^{n-1} \frac{(2m-1)(m-1) C_{m-1} j^{m-2+\alpha}}{A_{(2m-1)\alpha} (j+1)^{(2m-1)\alpha}} \\ &\sim (2m-1)(m-1) C_{m-1} \cdot n^{(2m-1)\alpha} \sum_{j=j_0+1}^{n-1} j^{m-2+\alpha-(2m-1)\alpha} \\ &\sim \frac{(2m-1)(m-1)}{m-1+\alpha-(2m-1)\alpha} \cdot C_{m-1} n^{m-1+\alpha} \\ &= \frac{2m-1}{1-2\alpha} \cdot C_{m-1} n^{m-1+\alpha} = C_m n^{m-1+\alpha}, \end{aligned} \quad (4.12)$$

as  $n \rightarrow \infty$ . Since  $m > 1$  and  $-1 < \alpha < 1/2$ , we can see that  $m-1+\alpha > (2m-1)\alpha$ . From (4.7), (4.9) and (4.12),

$$M_n^{(2m-1)} \sim F_n^{(2m-1)} \sim C_m n^{m-1+\alpha} \quad (n \rightarrow \infty).$$

This completes the proof.  $\square$

**4.2. Proof of (2.4) in Theorem 2.1.** Here we assume that  $\alpha = 1/2$  and  $\beta \neq 0$ . It suffices to show that for  $\ell \in \mathbb{N}$ ,

$$M_n^{(2\ell-1)} \sim C'_\ell \cdot n^{\ell-1/2} (\log n)^{\ell-1} \quad (n \rightarrow \infty) \quad (4.13)$$

holds true. Here constants  $C'_\ell$  are given by

$$C'_\ell = C'_\ell(\beta) := \frac{2\beta}{\sqrt{\pi}} \cdot (2\ell - 1)!!.$$

We can see that (1.7) and (1.6) imply (4.13) with  $\ell = 1$ . Assume that  $m > 1$  and (4.13) holds true for  $\ell = 1, \dots, m-1$ . Then we have that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} f_n^{(2m-1)} &\sim \binom{2m-1}{2m-3} M_n^{(2m-3)} \\ &\sim (2m-1)(m-1) C'_{m-1} n^{m-3/2} (\log n)^{m-2}. \end{aligned}$$



By (4.9) and Lemma A.1, we have, as  $n \rightarrow \infty$ ,

$$\begin{aligned}
F_n^{(2m-1)} &\sim A_{m-1/2} n^{m-1/2} \sum_{j=1}^{n-1} \frac{(2m-1)(m-1)C'_{m-1} j^{m-3/2} (\log j)^{m-2}}{A_{m-1/2} (j+1)^{m-1/2}} \\
&\sim (2m-1)(m-1)C'_{m-1} n^{m-1/2} \sum_{j=1}^{n-1} \frac{(\log j)^{m-2}}{j} \\
&\sim (2m-1)C'_{m-1} n^{m-1/2} (\log n)^{m-1} \\
&= C'_m n^{m-1/2} (\log n)^{m-1}. \tag{4.14}
\end{aligned}$$

Here we used (A.1). As  $m > 1$ , it follows from (4.7), (4.9) and (4.14) that

$$M_n^{(2m-1)} \sim F_n^{(2m-1)} \sim C'_m n^{m-1/2} (\log n)^{m-1} \quad (n \rightarrow \infty).$$

This completes the proof.  $\square$

## 5. PROOF OF THEOREM 2.3 (i) AND (ii)

Let  $m, n \in \mathbb{N}$ . As a counterpart of (4.1) for odd order moments, we use the following recursion for even order moments, which can be derived from (4.2) in [3]:

$$E[(S_{n+1})^{2m}] = 1 + \sum_{\ell=1}^m \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} E[(S_n)^{2\ell}]. \tag{5.1}$$

Assume that  $-1 < \alpha < 1/2$ . For  $n, m \in \mathbb{N}$ , we put

$$M_n^{(2m)} := \frac{1}{(2m-1)!!} E \left[ \left( \frac{S_n}{\sqrt{n/(1-2\alpha)}} \right)^{2m} \right] - 1.$$

The asymptotic behavior of  $M_n^{(2)}$  can be obtained from (1.8) and (1.5). For the sake of completeness, we give a short proof for  $-1 < \alpha < 1/2$  ( $\alpha = 1/2$  will be treated in Section 6 below). Using (5.1), one can show that  $\{M_n^{(2)}\}$  satisfies the following recursion:

$$M_1^{(2)} = -2\alpha, \quad M_{n+1}^{(2)} = \frac{n}{n+1} \left( 1 + \frac{2\alpha}{n} \right) M_n^{(2)} \quad (n \in \mathbb{N}).$$

If  $-1 < \alpha < 1/2$  and  $\alpha \neq 0, -1/2$ , then

$$M_n^{(2)} = \frac{M_1^{(2)}}{n} \prod_{j=1}^{n-1} \left( 1 + \frac{2\alpha}{j} \right) = -\frac{\Gamma(n+2\alpha)}{\Gamma(n+1)\Gamma(2\alpha)} \sim -\frac{n^{-(1-2\alpha)}}{\Gamma(2\alpha)}, \tag{5.2}$$

as  $n \rightarrow \infty$ , while if  $\alpha = 0$  and  $n \geq 1$ , or if  $\alpha = -1/2$  and  $n \geq 2$ , then

$$M_n^{(2)} = 0. \tag{5.3}$$

Putting

$$s_n^{(2m)} = \left( \frac{n}{1-2\alpha} \right)^m (2m-1)!!,$$

we have  $E[(S_n)^{2m}] = s_n^{(2m)}(M_n^{(2m)} + 1)$ . Hence (5.1) can be put into the following form:

$$\begin{aligned} M_{n+1}^{(2m)} &= \frac{1}{s_{n+1}^{(2m)}} \left( 1 + \sum_{\ell=1}^m \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} s_n^{(2\ell)} M_n^{(2\ell)} \right. \\ &\quad \left. + \sum_{\ell=1}^m \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} s_n^{(2\ell)} \right) - 1 \\ &= \left( \frac{n}{n+1} \right)^m \left( 1 + \frac{2m\alpha}{n} \right) M_n^{(2m)} \\ &\quad + \frac{1}{s_{n+1}^{(2m)}} \sum_{\ell=1}^{m-1} \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} s_n^{(2\ell)} M_n^{(2\ell)} \\ &\quad + \frac{1}{s_{n+1}^{(2m)}} \left\{ 1 + \sum_{\ell=1}^m \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} s_n^{(2\ell)} \right\} - 1. \end{aligned}$$

Hence, by putting

$$\begin{aligned} f_n^{(2m)} &:= \frac{1}{s_{n+1}^{(2m)}} \sum_{\ell=1}^{m-1} \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} s_n^{(2\ell)} M_n^{(2\ell)}, \\ g_n^{(2m)} &:= \left( \frac{n}{n+1} \right)^m \left( 1 + \frac{2m\alpha}{n} \right), \\ h_n^{(2m)} &:= \frac{1}{s_{n+1}^{(2m)}} \left\{ 1 + \sum_{\ell=1}^m \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} s_n^{(2\ell)} \right\} - 1, \end{aligned} \tag{5.4}$$

we have

$$M_{n+1}^{(2m)} = f_n^{(2m)} + h_n^{(2m)} + g_n^{(2m)} M_n^{(2m)}. \tag{5.5}$$

By Lemma 4.1, the recursion (5.5) is solved by

$$M_n^{(2m)} = M_{j_0+1}^{(2m)} \bar{g}_n^{(2m)} + F_n^{(2m)} + H_n^{(2m)}. \tag{5.6}$$

Here  $j_0 = j_0(2m\alpha)$  is defined by (4.4), and we put

$$\bar{g}_n^{(2m)} := \prod_{j=j_0+1}^{n-1} g_j^{(2m)} = \left( \frac{j_0+1}{n} \right)^m \prod_{j=j_0+1}^{n-1} \left( 1 + \frac{2m\alpha}{j} \right), \tag{5.7}$$

and

$$F_n^{(2m)} := \bar{g}_n^{(2m)} \sum_{j=j_0+1}^{n-1} \frac{f_j^{(2m)}}{\bar{g}_{j+1}^{(2m)}}, \quad H_n^{(2m)} := \bar{g}_n^{(2m)} \sum_{j=j_0+1}^{n-1} \frac{h_j^{(2m)}}{\bar{g}_{j+1}^{(2m)}}. \tag{5.8}$$

Applying Lemma 4.2, we have

$$\bar{g}_n^{(2m)} \sim (j_0+1)^m A_{2m\alpha} n^{-m(1-2\alpha)}, \tag{5.9}$$

as  $n \rightarrow \infty$ .

Next, let us investigate the asymptotic behavior of  $H_n^{(2m)}$ .

**Lemma 5.1.** Assume that  $m \geq 2$ .

(i) If  $-1 < \alpha \leq 0$ , then

$$H_n^{(2m)} \sim \frac{(1-4\alpha)c_{2m}}{m(1-2\alpha)-1} \cdot n^{-1} \quad (n \rightarrow \infty). \quad (5.10)$$

Here  $c_{2m} = \frac{m(m-1)}{2}c(\alpha)$ , and  $c(\alpha)$  is the constant defined by (2.7).

(ii) If  $0 < \alpha < 1/2$ , then  $H_n^{(2m)} = o(n^{-(1-2\alpha)})$  as  $n \rightarrow \infty$ .

*Proof.* We put

$$\begin{aligned} I_n &:= \frac{1}{s_{n+1}^{(2m)}} \left\{ \left(1 + \frac{2m\alpha}{n}\right) s_n^{(2m)} + \binom{2m}{2m-2} s_n^{(2m-2)} \right\} - 1, \\ J_n &:= \frac{1}{s_{n+1}^{(2m)}} \cdot \frac{\alpha}{n} \binom{2m}{2m-3} s_n^{(2m-2)}, \\ K_n &:= \frac{1}{s_{n+1}^{(2m)}} \left( 1 + \sum_{\ell=1}^{m-2} \left\{ \binom{2m}{2\ell} + \frac{\alpha}{n} \binom{2m}{2\ell-1} \right\} s_n^{(2\ell)} \right). \end{aligned}$$

Then  $h_n^{(2m)}$  can be written as follows:

$$h_n^{(2m)} = I_n + J_n + K_n.$$

We first note that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} I_n &= \left(\frac{n}{n+1}\right)^m \left(1 + \frac{2m\alpha}{n}\right) + m \cdot \frac{1-2\alpha}{n+1} \left(\frac{n}{n+1}\right)^{m-1} - 1 \\ &= -\frac{(n+1)^m - n^m - mn^{m-1}}{(n+1)^m} \sim -\frac{m(m-1)}{2} \cdot n^{-2}. \end{aligned}$$

One can see that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} J_n &\sim \frac{2m(m-1)}{3} \cdot \alpha(1-2\alpha) \cdot n^{-2}, \\ K_n &\sim \binom{2m}{2m-4} \frac{s_n^{(2m-4)}}{s_{n+1}^{(2m)}} \sim \frac{m(m-1)}{6} \cdot (1-2\alpha)^2 \cdot n^{-2}. \end{aligned}$$

Thus we obtain

$$\begin{aligned} h_n^{(2m)} &\sim \left(-\frac{1}{2} + \frac{2\alpha(1-2\alpha)}{3} + \frac{(1-2\alpha)^2}{6}\right) m(m-1) \cdot n^{-2} \\ &= -\frac{m(m-1)}{3} \cdot (2\alpha^2 + 1) \cdot n^{-2}. \end{aligned} \quad (5.11)$$

If  $-1 < \alpha \leq 0$  and  $m \geq 2$ , then  $-2 + m(1-2\alpha) \geq 0$ . Using (5.9), (5.11), and Lemma A.1, we have that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} H_n^{(2m)} &\sim -\frac{m(m-1)}{3} \cdot (2\alpha^2 + 1) \cdot n^{-m(1-2\alpha)} \sum_{j=1}^{n-1} j^{-2+m(1-2\alpha)} \\ &\sim -\frac{m(m-1)}{3} \cdot \frac{2\alpha^2 + 1}{m(1-2\alpha) - 1} \cdot n^{-1} = \frac{(1-4\alpha)c_{2m}}{m(1-2\alpha) - 1} \cdot n^{-1}. \end{aligned}$$

This proves (i). If  $0 < \alpha < 1/2$  and  $m \geq 2$ , then as  $n \rightarrow \infty$ ,

$$H_n^{(2m)} = \begin{cases} O(n^{-m(1-2\alpha)}) & \text{if } -2 + m(1-2\alpha) < -1, \\ O(n^{-m(1-2\alpha)} \log n) & \text{if } -2 + m(1-2\alpha) = -1, \\ O(n^{-1}) & \text{if } -2 + m(1-2\alpha) > -1, \end{cases}$$

which implies (ii).  $\square$

**5.1. Proof of (2.8) in Theorem 2.3.** Here we suppose that  $0 < \alpha < 1/2$ . By induction with respect to  $m$ , we will prove (2.8). The case  $m = 1$  is proved in (5.2). Let  $m \geq 2$  be an integer. For  $\ell = 1, \dots, m-1$ , suppose that, as  $n \rightarrow \infty$ ,

$$M_n^{(2\ell)} \sim -\frac{\ell}{\Gamma(2\alpha)} n^{-(1-2\alpha)},$$

holds true. By the hypothesis of the induction, we have

$$f_n^{(2m)} \sim \binom{2m}{2m-2} \frac{s_n^{(2m-2)}}{s_{n+1}^{(2m)}} M_n^{(2m-2)} \sim -(1-2\alpha) \frac{m(m-1)}{\Gamma(2\alpha)} \cdot n^{-2(1-\alpha)}$$

as  $n \rightarrow \infty$ . By (5.9) and Lemma A.1, we have

$$\begin{aligned} F_n^{(2m)} &\sim -(1-2\alpha) \frac{m(m-1)}{\Gamma(2\alpha)} n^{-m(1-2\alpha)} \sum_{j=1}^{n-1} j^{-2(1-\alpha)+m(1-2\alpha)} \\ &\sim -\frac{m}{\Gamma(2\alpha)} n^{-(1-2\alpha)}, \end{aligned} \quad (5.12)$$

as  $n \rightarrow \infty$ . Here we used the fact that

$$-2(1-\alpha) + m(1-2\alpha) \geq -2(1-\alpha) + 2(1-2\alpha) = -2\alpha > -1.$$

In view of (5.6), (5.9) and Lemma 5.1 (ii), we have that, as  $n \rightarrow \infty$ ,  $M_n^{(2m)} \sim F_n^{(2m)}$ . This completes the proof.  $\square$

**5.2. Proof of (2.6) in Theorem 2.3.** Suppose that  $-1 < \alpha \leq 0$ . By induction with respect to  $m \geq 2$ , we will prove (2.6). By (5.4),

$$f_n^{(4)} = \frac{s_n^{(2)}}{s_{n+1}^{(4)}} \left\{ \binom{4}{2} + \frac{\alpha}{n} \binom{4}{1} \right\} M_n^{(2)}.$$

By (5.3),

$$f_n^{(4)} = F_n^{(4)} = 0 \quad \text{for } n \geq 2, \text{ if } \alpha = 0 \text{ or } -1/2.$$

If  $-1 < \alpha < 0$  and  $\alpha \neq -1/2$ , then (5.2) and the same computation leading (5.12) yield that

$$F_n^{(4)} \sim -\frac{2}{\Gamma(2\alpha)} n^{-(1-2\alpha)} \quad (n \rightarrow \infty).$$

Taking this, (5.6), (5.9) and Lemma 5.1 (i) into consideration, we have that, as  $n \rightarrow \infty$ ,

$$M_n^{(4)} \sim H_n^{(4)} \sim \frac{(1-4\alpha)c_4}{2(1-2\alpha)-1} \cdot n^{-1} = c_4 n^{-1}.$$

Therefore (2.6) with  $m = 2$  holds true.

Now, let  $m \geq 3$  be an integer, and assume that for  $\ell = 2, \dots, m-1$ ,

$$M_n^{(2\ell)} \sim c_{2\ell} n^{-1} \quad (n \rightarrow \infty)$$

holds true. The hypothesis of the induction yields that, as  $n \rightarrow \infty$ ,

$$f_n^{(2m)} \sim \binom{2m}{2m-2} \frac{s_n^{(2m-2)}}{s_{n+1}^{(2m)}} M_n^{(2m-2)} \sim (1-2\alpha)m c_{2m-2} n^{-2}.$$

Using (5.9) and Lemma A.1 again, we have that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} F_n^{(2m)} &\sim (1-2\alpha)m c_{2m-2} n^{-m(1-2\alpha)} \sum_{j=j_0+1}^{n-1} j^{-2+m(1-2\alpha)} \\ &\sim \frac{(1-2\alpha)m c_{2m-2}}{m(1-2\alpha)-1} \cdot n^{-1} = \frac{(1-2\alpha)(m-2)}{m(1-2\alpha)-1} \cdot c_{2m} n^{-1}. \end{aligned}$$

By (5.6), (5.9) and Lemma 5.1 (ii), we have that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} M_n^{(2m)} &\sim H_n^{(2m)} + F_n^{(2m)} \\ &\sim \left\{ \frac{1-4\alpha}{m(1-2\alpha)-1} + \frac{(1-2\alpha)(m-2)}{m(1-2\alpha)-1} \right\} c_{2m} n^{-1} = c_{2m} n^{-1}. \end{aligned}$$

Thus we obtain the desired result.  $\square$

## 6. PROOF OF THEOREM 2.3 (iii)

Here we suppose that  $\alpha = 1/2$ . Then (5.1) can be put into the following form:

$$E[(S_{n+1})^{2m}] = 1 + \sum_{\ell=1}^m \left\{ \binom{2m}{2\ell} + \frac{1}{2n} \binom{2m}{2\ell-1} \right\} E[(S_n)^{2\ell}]. \quad (6.1)$$

We define

$$L_n^{(2m)} := E[(S_n)^{2m}] \Big/ \binom{n+m-1}{n-1} = \frac{(n-1)! m!}{(n+m-1)!} E[(S_n)^{2m}].$$

Then, from (6.1), one can see that  $\{L_n^{(2m)}\}_{n=1,2,\dots}$  satisfies

$$L_{n+1}^{(2m)} - L_n^{(2m)} = \binom{n+m}{n}^{-1} \left( 1 + \sum_{\ell=1}^{m-1} \left\{ \binom{2m}{2\ell} + \frac{1}{2n} \binom{2m}{2\ell-1} \right\} E[(S_n)^{2\ell}] \right).$$

Thus, we obtain

$$\begin{aligned} L_n^{(2m)} &= 1 + \sum_{j=1}^{n-1} \binom{j+m}{j}^{-1} \\ &\quad + \sum_{j=1}^{n-1} \sum_{\ell=1}^{m-1} \left\{ \binom{2m}{2\ell} + \frac{1}{2j} \binom{2m}{2\ell-1} \right\} \binom{j+m}{j}^{-1} E[(S_j)^{2\ell}]. \end{aligned} \quad (6.2)$$

Here we have used the fact that  $L_1^{(2m)} = 1$ . Now, we put

$$M_n^{(2m)} := \frac{1}{(2m-1)!!} E \left[ \left( \frac{S_n}{\sqrt{n \log n}} \right)^{2m} \right] - 1,$$

$$t_n^{(2m)} := (n \log n)^m \cdot (2m-1)!! \left/ \binom{n+m-1}{n-1} \right.$$

Then we have  $L_n^{(2m)} = t_n^{(2m)}(M_n^{(2m)} + 1)$ . Hence, (6.2) can be put into the following form:

$$M_n^{(2m)} = I_n^{(2m)} + J_n^{(2m)} + K_n^{(2m)}, \quad (6.3)$$

where

$$I_n^{(2m)} := \frac{1}{t_n^{(2m)}} \left( 1 + \sum_{j=1}^{n-1} \binom{j+m}{j}^{-1} \right),$$

$$J_n^{(2m)} := \frac{1}{t_n^{(2m)}} \sum_{j=1}^{n-1} \sum_{\ell=1}^{m-1} \left\{ \binom{2m}{2\ell} + \frac{1}{2j} \binom{2m}{2\ell-1} \right\} \binom{j+m}{j}^{-1} \binom{j+\ell-1}{j-1} t_j^{(2\ell)} - 1$$

$$K_n^{(2m)} := \frac{1}{t_n^{(2m)}} \sum_{j=1}^{n-1} \sum_{\ell=1}^{m-1} \left\{ \binom{2m}{2\ell} + \frac{1}{2j} \binom{2m}{2\ell-1} \right\} \binom{j+m}{j}^{-1} \binom{j+\ell-1}{j-1} t_j^{(2\ell)} M_j^{(2\ell)}.$$
(6.4)

We first investigate the asymptotic behavior of  $I_n^{(2m)}$  and  $J_n^{(2m)}$ .

**Lemma 6.1.** For any  $m \geq 2$ , we have that, as  $n \rightarrow \infty$ ,

$$I_n^{(2m)} = O((\log n)^{-m}), \quad (6.5)$$

$$J_n^{(2m)} = O((\log n)^{-m}). \quad (6.6)$$

*Proof.* To prove (6.5), note that

$$\binom{j+m}{j} \sim \frac{j^m}{m!} \quad (j \rightarrow \infty), \quad (6.7)$$

$$t_n^{(2m)} \sim m! \cdot (2m-1)!! \cdot (\log n)^m \quad (n \rightarrow \infty). \quad (6.8)$$

As  $m \geq 2$ , we have

$$I_n^{(2m)} \sim \frac{1}{t_n^{(2m)}} \left( 1 + \sum_{j=1}^{\infty} \binom{j+m}{j}^{-1} \right) = O((\log n)^{-m}).$$

Next, we will show (6.6). For  $1 \leq \ell \leq m-1$ , we have that, as  $j \rightarrow \infty$ ,

$$d_j := \binom{j+m}{j}^{-1} \binom{j+\ell-1}{j-1} t_j^{(2\ell)} \sim m! \cdot (2\ell-1)!! \cdot j^{-(m-\ell)} (\log j)^\ell.$$

This implies that, for  $1 \leq \ell \leq m-2$ ,

$$\frac{1}{t_n^{(2m)}} \sum_{j=1}^{n-1} d_j = O \left( \frac{1}{t_n^{(2m)}} \right) = O((\log n)^{-2m}),$$

and also, for  $1 \leq \ell \leq m - 1$ ,

$$\frac{1}{t_n^{(2m)}} \sum_{j=1}^{n-1} \frac{1}{2j} \cdot d_j = O\left(\frac{1}{t_n^{(2m)}}\right) = O((\log n)^{-2m}).$$

Therefore we obtain

$$\begin{aligned} J_n^{(2m)} &= \frac{1}{t_n^{(2m)}} \sum_{j=1}^{n-1} \binom{2m}{2m-2} \binom{j+m}{j}^{-1} \binom{j+m-2}{j-1} t_j^{(2m-2)} - 1 \\ &\quad + O((\log n)^{-2m}). \end{aligned} \quad (6.9)$$

The first term in the right hand side of (6.9) is

$$\frac{(n+m-1)!}{n^m(n-1)!} \cdot \frac{m}{(\log n)^m} \sum_{j=1}^{n-1} \frac{j^m j!}{(j+m)!} \cdot \frac{(\log j)^{m-1}}{j} - 1. \quad (6.10)$$

Note that

$$\begin{aligned} c_n &:= \frac{(n+m-1)!}{n^m(n-1)!} = \prod_{k=1}^{m-1} \left(1 + \frac{k}{n}\right) = 1 + O\left(\frac{1}{n}\right) \quad (n \rightarrow \infty), \\ c'_n &:= \frac{n^m n!}{(n+m)!} = \prod_{k=1}^m \left(1 + \frac{k}{n}\right)^{-1} = 1 + O\left(\frac{1}{n}\right) \quad (n \rightarrow \infty). \end{aligned} \quad (6.11)$$

a Hence (6.11) and (A.1) in Appendix imply that (6.10) is

$$\begin{aligned} &c_n \frac{m}{(\log n)^m} \sum_{j=1}^{n-1} c'_j \frac{(\log j)^{m-1}}{j} - 1 \\ &= c_n \frac{m}{(\log n)^m} \sum_{j=1}^{n-1} (c'_j - 1) \frac{(\log j)^{m-1}}{j} + (c_n - 1) \frac{m}{(\log n)^m} \sum_{j=1}^{n-1} \frac{(\log j)^{m-1}}{j} \\ &\quad + \frac{m}{(\log n)^m} \sum_{j=1}^{n-1} \frac{(\log j)^{m-1}}{j} - 1 \\ &= O((\log n)^{-m}), \end{aligned}$$

as  $n \rightarrow \infty$ . Here we use the fact that  $\sum_{j=1}^{\infty} (c'_j - 1) \frac{(\log j)^{m-1}}{j}$  converges. Now, taking (6.9) into consideration, one can see that, as  $n \rightarrow \infty$ ,

$$J_n^{(2m)} = O((\log n)^{-m}).$$

Thus we obtain the desired result.  $\square$

Now, by induction with respect to  $m$ , we will prove (2.9) in Theorem 2.3. By (6.2), we have

$$L_n^{(2)} = 1 + \sum_{j=1}^{n-1} \frac{1}{j+1} = \sum_{j=1}^n \frac{1}{j},$$

and

$$M_n^{(2)} = \frac{L_n^{(2)}}{t_n^{(2)}} - 1 = \frac{1}{\log n} \left( \sum_{j=1}^n \frac{1}{j} - \log n \right) \sim \gamma (\log n)^{-1} \quad (n \rightarrow \infty).$$

Let  $m \geq 2$  be fixed and suppose that for any  $1 \leq \ell \leq m-1$

$$M_n^{(2^\ell)} \sim \ell \gamma (\log n)^{-1} \quad (n \rightarrow \infty)$$

holds true. Then we have that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} K_n^{(2m)} &\sim \frac{1}{t_n^{(2m)}} \sum_{j=1}^{n-1} \binom{2m}{2m-2} \binom{j+m}{j}^{-1} \binom{j+m-2}{j-1} t_j^{(2m-2)} M_j^{(2m-2)} \\ &\sim \frac{m(m-1)\gamma}{(\log n)^m} \sum_{j=1}^{n-1} \frac{(\log j)^{m-2}}{j} \\ &\sim m\gamma (\log n)^{-1}. \end{aligned}$$

Here we used (A.1). In view of Lemma 6.1,  $M_n^{(2m)} \sim K_n^{(2m)}$  as  $n \rightarrow \infty$ . This completes the proof of (2.9) in Theorem 2.3.  $\square$

#### APPENDIX A. LEMMAS FROM CALCULUS

**Lemma A.1** (Stolz-Cesàro: See e.g. Knopp [8], p.34). If a real sequence  $\{a_n\}$  and a positive sequence  $\{b_n\}$  satisfy

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L \in \mathbb{R} \cup \{\pm\infty\}, \quad \text{and} \quad \sum_{n=1}^{\infty} b_n = +\infty,$$

then

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n a_k}{\sum_{k=1}^n b_k} = L.$$

**Lemma A.2** (cf. Knopp [8], p.64). Let  $f(t)$  be a positive function on  $[1, \infty)$ , which is non-increasing on  $[n_0, \infty)$  for some  $n_0 \in \mathbb{N}$ . For each  $n \in \mathbb{N}$ , we set

$$S_n := \sum_{j=1}^n f(j) \quad \text{and} \quad I_n := \int_1^n f(t) dt.$$

If  $I_n \rightarrow +\infty$  as  $n \rightarrow \infty$ , then

$$\frac{S_n}{I_n} = 1 + O((I_n)^{-1}) \quad \text{as } n \rightarrow \infty.$$

*Proof.* For  $n \geq n_0$ , let

$$S'_n := \sum_{j=n_0}^n f(j) \quad \text{and} \quad I'_n := \int_{n_0}^n f(t) dt.$$

As in p.64 of [8], we can see that  $0 \leq S'_n - I'_n \leq f(n_0)$  for any  $n \geq n_0$ . Since

$$\frac{S_n}{I_n} - 1 = \frac{S_{n_0-1} - I_{n_0}}{I_n} + \frac{S'_n - I'_n}{I_n} \quad (n \geq n_0),$$

we obtain the conclusion.  $\square$



Let  $m \geq 1$ . The function

$$f(t) = \frac{(\log t)^{m-1}}{t} \quad (t \geq 1)$$

is non-increasing on  $[e^{m-1}, \infty)$ . As

$$I_n = \int_1^n f(t) dt = \frac{(\log n)^m}{m},$$

Lemma A.2 implies that

$$\frac{m}{(\log n)^m} \sum_{j=1}^n \frac{(\log j)^{m-1}}{j} = 1 + O((\log n)^{-m}) \quad (n \rightarrow \infty). \quad (\text{A.1})$$

#### REFERENCES

- [1] Baur, E. and Bertoin, J. (2016). Elephant random walks and their connection to Pólya-type urns, *Phys. Rev. E*, **94**, 052134
- [2] Bercu, B. (2018). A martingale approach for the elephant random walk, *J. Phys. A: Math. Theor.*, **51**, 015201
- [3] Bercu, B., Chabanol, M.-L., and Ruch, J.-J. (2019). Hypergeometric identities arising from the elephant random walk, *J. Math. Anal. Appl.*, **480**, 123360
- [4] Bertoin, J. (2021). Counting the zeros of an elephant random walk, arXiv:2105.09569
- [5] Coletti, C. F., Gava, R. J., and Schütz, G. M. (2017a). Central limit theorem for the elephant random walk, *J. Math. Phys.*, **58**, 053303
- [6] Coletti, C. F., Gava, R. J., and Schütz, G. M. (2017b). A strong invariance principle for the elephant random walk, *J. Stat. Mech.*, 123207
- [7] Fan, X., Hu, H., and Ma, X. (2021). Cramér moderate deviations for the elephant random walk, *J. Stat. Mech.*, 023402
- [8] Knopp, K. (1956). Infinite sequences and series, Dover Publications.
- [9] Kubota, N. and Takei, M. (2019). Gaussian fluctuation for superdiffusive elephant random walks, *J. Statist. Phys.*, **177**, 1157–1171.
- [10] Ma, X., El Machkouri, M., and Fan, X. (2022). On Wasserstein-1 distance in the central limit theorem for elephant random walk, *J. Math. Phys.*, **63**, 013301
- [11] Schütz, G. M. and Trimper, S. (2004). Elephants can always remember: Exact long-range memory effects in a non-Markovian random walk, *Phys. Rev. E*, **70**, 045101

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF THE RYUKYU, OKINAWA, JAPAN

*Email address:* hayashi@math.u-ryukyu.ac.jp

DEPARTMENT OF MATHEMATICAL SCIENCES, GRADUATE SCHOOL OF ENGINEERING AND SCIENCE, UNIVERSITY OF THE RYUKYUS, OKINAWA, JAPAN

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF ENGINEERING, YOKOHAMA NATIONAL UNIVERSITY, YOKOHAMA, JAPAN

*Email address:* takei-masato-fx@ynu.ac.jp