

# ON THE TOPOLOGY OF GAUSSIAN RANDOM ZERO SETS

ZHENGJIANG LIN

ABSTRACT. We study the asymptotic laws for the number, Betti numbers, and diffeomorphism types of connected components of zero sets of real smooth Gaussian random fields defined on either a Euclidean space or a Riemannian manifold without boundary, where the random zero sets almost surely consist of submanifolds of high codimensions.

## CONTENTS

|  |    |
|--|----|
| 1. Introduction  | 1  |
| 1.1. Main Results  | 2  |
| 2. Random Fields on Cubes  | 8  |
| 3. Proof of Theorem 1.1 and Theorem 1.2  | 16 |
| 4. Parametric Ensembles and Proof of Theorem 1.3                                       | 21 |
| Appendix A. Quantitative Bulinskaya’s Lemma for Gaussian Random Fields                 | 35 |
| Appendix B. Stability Lemmas   | 36 |
| Appendix C. Conditions for Ergodicity in Theorem 1.1 and Theorem 3.1                   | 38 |
| Appendix D. Conditions for Positive Limiting Constants in Theorem 1.1 and Theorem 3.1  | 40 |
| Appendix E. The Two-Point Conditions (A3) in Definition 2.1 and (B4) in Definition 4.2 | 48 |
| Appendix F. Curvatures Computations  | 51 |
| References   | 54 |

## 1. INTRODUCTION

In the past twenty years, the study of statistics related to the level sets, in particular zero sets, of random fields is a rising area with many interesting but challenging problems. And indeed, the study of them can be dated back even to early twentieth century. The zeros of random polynomials might be first studied by Boch and Pólya [BP32]. Then, Kac [K43] studied asymptotic laws for the expected number of real zeros of random polynomials of degree  $d$  as  $d \rightarrow \infty$ .

It is very natural that one wants to do similar things in multivariate cases. But now a branch appears. Unlike univariate case where there is no big difference between “numbers”

---

*Date:* May 3, 2022.

and “measures” for discrete points, for multivariate cases, one can either consider the Hausdorff measures (volumes) or the number of connected components of the level sets. Different views have led different ways to tackle these related problems.

In this manuscript, we study the numbers and possible topological behaviors of connected components of those random zero sets. Before our work, one of previous great breakthroughs was made by Nazarov and Sodin in [NS09], where the authors first studied the asymptotic behaviors for numbers of connected components of zero sets of random functions from Gaussian ensembles of spherical harmonics. After that, in [S16] and [NS16], they developed more tools which are very useful in studying other Gaussian ensembles, for example trigonometric ensemble and Kostlan’s ensemble. In Example 4.3, we give the construction of Kostlan’s ensemble, and readers can see [S16] and [NS16] for more concrete examples. Then, in [SW19], Sarnak and Wigman developed more results in diffeomorphism types and tree structures of connected components of zero sets of those random functions. There are many other related remarkable works concerning the asymptotic topological behaviors, see for example [GW14], [GW16],[CS19], [W21], and references therein. Also, in recent years, one possible reason of interests is the percolation model raised by Bogomolny and Schmit [BS02]. There are also some connections to a statistical version of the Hilbert’s sixteenth problem, see for example [LL15].

Our main results are generalizations for parts of brilliant works built up in [NS16], [W21], and [SW19], from zero sets of Gaussian random functions to zero sets of Gaussian random fields, which are random submanifolds of higher codimensions. And some of our results are parallel to those great works in [GW14] and [GW16], but are adapted to some more general situations. In our model, a special case is when random fields consist of several independent random functions although we do not need such a strong assumption in the main parts. But to understand our results here better, readers should better keep this simplest model in mind first. The main tools we use are some statistical estimates for curvature integrals over zero sets and the well-known Kac–Rice formula, see for example Section 2 of [CH20] or Chapter 6 of [AW09]. More specifically, we consider the integrals involving second fundamental forms and mean curvatures, which appeared in Fenchel–Borsuk–Willmore and Chern–Lashof inequalities, see [CL57], [CL58], and [C71]. The statistics involving curvature integral is a very natural idea, see for example [L16], where the author considered curvature integrals in Chern–Gauss–Bonnet’s theorem. But there is a new difficult problem in our cases that we need to calculate a highly singular integral due to the calculations of second fundamental forms and mean curvatures. Apart from this difficulty, we also need to build up other related technical estimates in our model. See for example our proofs for Theorem 2.3, Lemma 2.6, and Theorem 4.7 for more discussions.

We also believe that there are many other statistical estimates for curvature integrals which can imply more integral geometric properties for level sets. But as far as we know, this direction is still in a rising status and there must be many applications for future development.

## 1.1. Main Results.

Let  $(\Omega, \mathfrak{S}, \mathcal{P})$  be a probability space and we use  $\mathcal{E}(\cdot)$  to denote expectations. We consider  $C^{3-}$ -smooth Gaussian fields  $F : \mathbb{R}^n \times \Omega \rightarrow \mathbb{R}^m$ ,  $m \in (0, n)$ . Here,  $C^{3-} \equiv \bigcap_{0 < \beta < 1} C^{2+\beta}$ . This smoothness assumption is not so essential and readers can take infinitely smooth random fields at first, while our techniques only use derivatives up to the second order.

Let  $Z(F) \equiv F^{-1}(0)$  be the random zero set of  $F$ . Almost surely, 0 is not a critical value of  $F$ , which follows from either the Bulinskaya's lemma, see for example Proposition 6.12 of [AW09], or our Lemma A.1 for a quantitative version. Hence, almost surely,  $Z(F)$  consists of  $(n-m)$ -dimensional regular submanifolds without boundaries, either bounded or unbounded.

By  $N(R; F)$ , we denote the number of connected components of  $Z(F)$  fully contained in the open cube  $C_R \equiv (-R, R)^n$  of  $\mathbb{R}^n$ . We use cubes here for some technical reasons proving (3.3) in Theorem 3.1, but of course one can replace open cubes by open convex polyhedrons or open balls.

From a topological point of view, we can also regard  $N(R; F)$  as the sum of 0-th Betti numbers for connected components of  $Z(F)$  fully contained in  $C_R$ . Hence, similarly, for each  $l = 0, 1, \dots, n-m$ , we define  $\beta_l(R; F)$  as the sum of  $l$ -th Betti numbers over  $\mathbb{R}$  for connected components of  $Z(F)$  fully contained in  $C_R$ , i.e.,

$$(1.1) \quad \beta_l(R; F) \equiv \sum_{\gamma \subset Z(F) \cap C_R} \beta_l(\gamma),$$

where the summation counts closed components  $\gamma$  fully contained in  $C_R$  and  $\beta_l(\gamma)$  is the  $l$ -th Betti number over  $\mathbb{R}$  of  $\gamma$ . By Poincaré duality, we know that  $\beta_l(R; F) = \beta_{n-m-l}(R; F)$  for each  $l$ . We will still use the notation  $N(R; F)$  in most of cases, as our theorems for  $N(R; F)$  hold in a more complete way, while those theorems for Betti numbers are direct once the proofs for  $N(R; F)$  are given.

Also, we let  $H(n-m)$  be the set of diffeomorphism types of closed  $(n-m)$ -dimensional manifolds that can be embedded into  $\mathbb{R}^n$ .  $H(n-m)$  is a countable set by Cheeger's finiteness theorem. For any diffeomorphism type  $H \in H(n-m)$ , let  $N(R; F, H)$  be the number of connected components of  $Z(F)$  lying entirely in  $C_R$ , and  $C^1$ -diffeomorphic to  $H$ .

As we mentioned before, our results work for the cases of independent Gaussian random functions, i.e., if we denote  $F = (f_1, \dots, f_m)$ , then  $f_{i_1}(x)$  and  $f_{i_2}(y)$  are independent as two Gaussian variables for all  $x, y \in \mathbb{R}^n$  if  $i_1 \neq i_2$ . But of course our methods and results adapt to more general settings.

We say that  $F$  is stationary if for any  $x, y, z \in \mathbb{R}^n$ ,  $(F(x), F(y))$  and  $(F(x+z), F(y+z))$  have the same distribution.

**Theorem 1.1.** *Assume that  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $m \in (0, n)$ , is a  $C^{3-}$ -smooth centred stationary Gaussian random field such that the joint distribution of  $(F(0), \nabla F(0))$  is non-degenerate.*

(1) *There exists a number  $\nu = \nu_F \geq 0$  so that*

$$(1.2) \quad \mathcal{E}(N(R; F)) = \nu \cdot |C_R| + o(R^n),$$

*as  $R \rightarrow +\infty$ . Here  $|C_R|$  is the volume of  $C_R$ .*

(2) If the translation action of  $\mathbb{R}^n$  is ergodic, then

$$(1.3) \quad \lim_{R \rightarrow \infty} \frac{N(R; F)}{|C_R|} = \nu \text{ almost surely} \quad \text{and} \quad \lim_{R \rightarrow \infty} \mathcal{E} \left| \frac{N(R; F)}{|C_R|} - \nu \right| = 0.$$

Moreover, for any  $H \in H(n - m)$ , (1) and (2) hold true if one replaces  $N(R; F)$  by  $N(R; F, H)$  and replaces the number  $\nu_F$  by a  $\nu_{F, H} \geq 0$ .

(3)

$$(1.4) \quad \nu_F = \sum_{H \in H(n-m)} \nu_{F, H}.$$

See more details in Section 3 about the translation action of  $\mathbb{R}^n$  and the definition of ergodicity. Another natural question is when those  $\nu_F$  and  $\nu_{F, H}$  are positive. In the special case when  $F$  consists of independent  $\{f_i\}_{i=1}^m$ , if for each  $i = 1, \dots, m$ , the spectral measure  $\rho_i$  associated to  $f_i$  has a nonempty interior in its compact support, then one can get that  $\nu_F > 0$  and  $\nu_{F, H} > 0$  for some  $H$ . This condition on spectral measure is exactly the one appeared in [NS16], [SW19], and [W21] when  $m = 1$ . And we will also discuss some weaker assumptions in Appendix D. We want to emphasize another thing that, even under some strong assumptions on  $F$ , it may not be true that  $\nu_{F, H} > 0$  for all  $H \in H(n - m)$  when  $m > 1$ . See our Remark 1.7 for more discussions later.

For Betti numbers  $\beta_l(R; F)$ , we have the following counterparts, which, when  $m = 1$ , were built up as Theorem 1.2 in [W21].

**Theorem 1.2.** *If  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $m \in (0, n)$ , is a  $C^{3-}$ -smooth centred stationary Gaussian random field such that the joint distribution of  $(F(0), \nabla F(0))$  is non-degenerate. Then, for each  $l = 0, 1, \dots, n - m$ , we have the following results.*

(1) There exists a number  $\nu_l = \nu_{l; F} \geq 0$  so that

$$(1.5) \quad \mathcal{E}(\beta_l(R; F)) = \nu_l \cdot |C_R| + o(R^n),$$

as  $R \rightarrow +\infty$ .

(2) If the translation action of  $\mathbb{R}^n$  is ergodic, then

$$(1.6) \quad \lim_{R \rightarrow \infty} \mathcal{E} \left| \frac{\beta_l(R; F)}{|C_R|} - \nu_l \right| = 0.$$

Under the assumptions we discussed after Theorem 1.1 or in Appendix D, these  $\nu_l$  are positive numbers. On the other hand, we believe that with tools in our manuscript, in particular, our  $L^1$ -estimate Theorem 2.8 and  $L^q$ -estimate Theorem 4.8, these constants  $\nu_{l; F}$  will depend on the field  $F$  in some continuous sense. One can see [KW18] for the case of random plane waves when  $n = 2$  and  $m = 1$ .

Next, we are going to consider parametric ensembles of Gaussian random fields defined on either open sets of  $\mathbb{R}^n$  or more generally, manifolds without boundary. Let  $\{F_L\}_{L \in \mathcal{L}}$  be a family of Gaussian fields defined on an open subset  $U \subset \mathbb{R}^n$ , where the index set  $L$  attains a discrete set  $\mathcal{L} \subset \mathbb{R}$ . One can also regard our  $F_L$  here as  $(f_{L,1}, \dots, f_{L,m})$  for independent  $f_{L,i}$

with each  $\{f_{L,i}\}_{L \in \mathcal{L}}$  being an ensemble of Gaussian random functions, for example Kostlan's ensemble or the ensemble of spherical harmonics. And  $L \in \mathcal{L}$  will be the parameters in the corresponding ensembles. In Section 4, we give the concrete example of Kostlan's ensemble as our Example 4.3. See more concrete examples in [S16] and [NS16]. Actually, the Gaussian fields defined on manifolds we are going to discuss are very generic. One can see [CH20] and references therein for more discussions.

**Theorem 1.3.** *Suppose that  $\{F_L\}_{L \in \mathcal{L}}$  is tame on an open set  $U \subset \mathbb{R}^n$ . Then, there is a measurable locally bounded function  $x \mapsto \bar{\nu}(x)$  on  $U$ , such that for every sequence of connected components counting measure  $n_L$  of  $F_L$  and for every  $\varphi \in C_c(U)$ , we have that*

$$(1.7) \quad \lim_{L \rightarrow \infty} \mathcal{E}^* \left[ \left| \frac{1}{L^n} \int_U \varphi(x) dn_L(x) - \int_U \varphi(x) \bar{\nu}(x) dx \right| \right] = 0.$$

Let us explain some notations and terminologies in Theorem 1.3. The definition of  $\{F_L\}_{L \in \mathcal{L}}$  being tame is in Definition 4.2. The definition is similar to the one in Theorem 2 of [NS16] for random functions cases, but we add one more thing, which is the assumption (B4) in Definition 4.2, and also, the assumption (A3) in Definition 2.1. The (A3) and (B4) assumptions are about the two-point covariance kernel, which are added to help us prove the two  $L^q$ -estimates, i.e., our Theorem 4.7 and Theorem 4.8. And these two assumptions are natural in most of the cases. For example,  $F_L$  consists of independent  $f_{L,i}$  with each  $\{f_{L,i}\}_{L \in \mathcal{L}}$  being Kostlan's ensemble or those generic Gaussian ensembles in [CH20].

And also, similar to the settings in Theorem 2 of [NS16], we say that a Borel measure  $n_L$  is a connected components counting measure of  $F_L$  if  $\text{spt}(n_L) \subset Z(F_L)$  and the  $n_L$ -mass of each component is 1. Readers can just regard them as numbers of connected components fully contained in a compact set of  $U$  for simplicity.

Then, one can also state a manifold version of Theorem 1.3. Suppose that  $X$  is an  $n$ -dimensional  $C^3$ -manifold without boundary and may not be compact. We say that a parametric Gaussian ensemble  $\{F_L\}_{L \in \mathcal{L}}$  is tame if for every  $C^3$ -chart  $(U, \pi)$ ,  $\pi : U \subset \mathbb{R}^n \rightarrow X$ ,  $\{F_L \circ \pi\}_{L \in \mathcal{L}}$  is tame on  $U$ . This definition does not depend on the particular choice of charts. See more discussions in section 9 of [NS16].

**Theorem 1.4.** *Suppose that  $\{F_L\}_{L \in \mathcal{L}}$  is tame on  $X$ , then the followings hold.*

- (1) *There is a locally finite Borel non-negative measure  $n_\infty$  on  $X$  such that for every choice of connected components counting measures  $n_L$  of  $F_L$  and every  $\varphi \in C_c(X)$ , we have that*

$$(1.8) \quad \lim_{L \rightarrow \infty} \mathcal{E}^* \left[ \left| \frac{1}{L^n} \int_X \varphi(x) dn_L(x) - \int_X \varphi(x) dn_\infty(x) \right| \right] = 0.$$

- (2) *For every chart  $\pi : U \rightarrow X$ , the measure  $n_\infty$  coincides on  $\pi(U)$  with the pushforward  $\pi_*(\bar{\nu}_\pi \cdot \mathcal{H}^n)$ , where  $\bar{\nu}_\pi$  is the point intensity associated with the ensemble  $\{F_L \circ \pi\}_{L \in \mathcal{L}}$ , and  $\mathcal{H}^n$  is the  $n$ -dimensional Hausdorff measure.*

**Remark 1.5.** For any  $H \in H(n - m)$ , Theorem 1.3 also holds true if one replaces  $\nu_{F_x}$  by  $\nu_{F_x, H}$  and replaces counting measure  $n_L$  by an  $H$ -counting measure  $n_{L, H}$ . And similarly,

Theorem 1.4 holds true if one replaces  $n_L$  by an  $n_{L,H}$  and  $n_\infty$  by an  $n_{\infty,H}$ . The idea is that we only count those components of the diffeomorphism type  $H$ .

**Remark 1.6.** For a closed manifold  $X$ , one can also constructs diffeomorphism types counting measures on  $H(n-m)$  by similar ways in [SW19]. For example, define  $N(X; F_L) \equiv \int_X 1 \cdot dn_L(x)$  and  $N(X; F_L, H) \equiv \int_X 1 \cdot dn_{L,H}(x)$ , we then let

$$(1.9) \quad \mu_L \equiv \frac{1}{N(X; F_L)} \cdot \sum_{H \in H(n-m)} N(X; F_L, H) \cdot \delta_H,$$

be a random probability measure on  $H(n-m)$ . Then, follow the same arguments proving Theorem 1.1 of [SW19], one can show that there is a limiting probability measure (in discrepancy sense) on  $H(n-m)$  if  $n_\infty(X) > 0$ , which is

$$(1.10) \quad \mu_\infty \equiv \frac{1}{n_\infty(X)} \cdot \sum_{H \in H(n-m)} n_{\infty,H}(X) \cdot \delta_H.$$

That is, for any  $\epsilon > 0$ ,

$$(1.11) \quad \lim_{L \rightarrow \infty} \mathcal{P} \left( \sup_{A \in H(n-m)} |\mu_L(A) - \mu_\infty(A)| > \epsilon \right) = 0.$$

**Remark 1.7.** Moreover, one can also use countable classifications thinner than  $H(n-m)$ . Say, let  $\mathcal{C}$  with elements  $c$  be the  $C^1$ -isotopy classes for  $(n-m)$ -dimensional submanifolds embedded in  $\mathbb{R}^n$ , which means that for any two  $(n-m)$ -dimensional manifolds  $M_1, M_2$  embedded in  $\mathbb{R}^n$ , if  $M_1$  is  $C^1$ -isotopic to  $M_2$ , then they are in the same class  $c \in \mathcal{C}$ . And  $\mathcal{C}$  is also a countable set. In this setting, our results including Theorem 1.1, Theorem 1.3, Theorem 1.4, and Remark 1.6, will also hold true. One interesting situation is that, for example, when  $n = 3$  and  $m = 2$ ,  $H(1)$  only contains the diffeomorphism type of  $\mathbb{S}^1$ , but we can consider  $\mathcal{C}$  as all possible  $C^1$ -knots structures in  $\mathbb{R}^3$ , which are also countable. The Remark 1.6 then holds true if one replaces  $H(1)$  and  $H$  by  $\mathcal{C}$  and knots  $c \in \mathcal{C}$ .

On the other hand, for either diffeomorphism types or  $C^1$ -isotopy classes, although we can write an infinite summation, for example, our Remark 1.6 or our (3) in Theorem 1.1, there are some  $\nu_{F,H}$  always being zero. This is a topological drawback, and the reason is that it may not be correct that for any closed  $(n-m)$ -dimensional submanifold  $M \subset \mathbb{R}^n$ , there is a  $G \in C^1(\mathbb{R}^n, \mathbb{R}^m)$  such that  $M$  is a non-degenerate connected component of  $Z(G)$ , which means that  $\nabla G$  is of full rank on  $M$ . But if the normal bundle of  $M$  in  $\mathbb{R}^n$  is trivial, then it is possible to find such a  $G$ . And then, follow steps in Appendix D, one can show that under some assumptions on  $F$ , we can get that  $\nu_{F,H(M)} > 0$ , where  $H(M)$  can be either the diffeomorphism type or the  $C^1$ -isotopy class of  $M$ . In particular, knots or even links always have trivial normal bundles in any  $\mathbb{R}^n$  for  $n \geq 3$  because they are embeddings of  $\mathbb{S}^1$ , whose vector bundles are either orientable trivial bundles or a direct sum of orientable trivial bundles and one Möbius band.

Our final results are for asymptotic laws of Betti numbers, which are new even in the case when  $m = 1$  under the more general parametric ensembles settings. They are also similar

and parallel to some of the works in [GW14] and [GW16] in some specific situations. For each  $l = 0, \dots, n - m$ , we use  $\beta_l(F_L)$  to denote the summation of  $l$ -th Betti numbers over  $\mathbb{R}$  for all connected components of  $Z(F_L)$  in the manifold  $X$ .

**Theorem 1.8.** *Assume that  $\{F_L\}_{L \in \mathcal{L}}$  is tame on  $X$  and  $X$  is a closed manifold. Then for each  $l = 0, \dots, n - m$ , there is a nonnegative measurable and bounded function  $\bar{\nu}_l(x)$  on  $X$ , and a positive constant  $C_l$  depending on the uniform smoothness and nondegeneracy of  $\{F_L\}_{L \in \mathcal{L}}$ , such that*

$$(1.12) \quad \int_X \bar{\nu}_l(x) \, dx \leq \liminf_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(F_L))}{L^n} \leq \limsup_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(F_L))}{L^n} \leq C_l \cdot |X|,$$

where  $|X|$  is the volume of  $X$ .

The precise definitions for uniform smoothness and nondegeneracy are stated in our Definition 4.1, where we say that  $\{F_L\}_{L \in \mathcal{L}}$  is uniformly controllable.

We organize the proofs as the followings. In Section 2, we will show  $L^1$ -estimates for the number and Betti numbers of connected components fully contained in  $C_R$ , i.e. upper bounds for  $\mathcal{E}(N(R; F))/R^n$  and  $\mathcal{E}(\beta_l(R; F))/R^n$  independent of  $R$ , which are our Theorem 2.3 and Theorem 2.8. This part contains the core ideas of this manuscript so we suggest readers to read them first before reading other sections.

And in Section 3, we will restate Theorem 1.1 as Theorem 3.1 and prove it. The proof for Theorem 1.2 is similar and we will mention necessary modifications at the end of Section 3. The ergodicity issue of translation actions and the positivity issue of  $\nu$  related to Theorem 1.1 and Theorem 3.1 will be included in Appendix C and Appendix D.

In Section 4, we restate Theorem 1.3 as Theorem 4.4 and prove it, which also needs two lemmas similar to those included in [NS16]. One in Appendix A is a quantitative Bulinskaya's lemma for random fields. And another is a stability lemma in Appendix B. The proof of Theorem 4.4 also needs an upper bound for  $\mathcal{E}((N(R; F))^q)/R^{nq}$  independent of  $R$  for some  $q = q(n, m) > 1$ , which is our Theorem 4.7. The proof of Theorem 1.4 is exactly the same as Theorem 3 in [NS16] once Theorem 1.3 is completed, so we omit the details in this manuscript. And finally, we restate Theorem 1.8 as Theorem 4.9 and prove it.

The principal novel results are the two technical  $L^1$  and  $L^q$  estimates we mentioned above, see our Theorem 2.3, Theorem 2.8, Theorem 4.7, and Theorem 4.8, for which we use geometric inequalities and consider the integrals involving curvatures. In all of these proofs, we will use  $|\cdot|$  or  $\|\cdot\|$  to denote norms of vectors or volumes of a set when there is no ambiguity. We only use  $\|\cdot\|_{C^{1+\beta}}$  or  $\|\cdot\|_{C^2}$ , i.e.  $\|\cdot\|$  with subindices, to denote norms related to smoothness. We will also use the notations  $v = (v_i)$  (or  $A = (A_{ij})$ ) to denote vectors (or matrices) with elements  $v_i$  (or  $A_{ij}$ ). When we do computations, we will use, for example,  $C(n, m)$ , to denote constants depending on  $n, m$ , but they are not necessarily same.

**Acknowledgements** The author would like to thank his advisor Professor Fang-Hua Lin for his endless encouragement and inspiring discussions during the Covid pandemic. In

the finalizing stage, the author would also like to thank Professor Paul Bourgade for very helpful discussions and suggestions. Professor Yuri Bakhtin recommended [GS04] and Ao Sun recommended [CL58] to the author and these references helped the author a lot. Finally, the author wants to thank Ao Sun and Professor Chao Li for their consistent encouragement in the past year.

## 2. RANDOM FIELDS ON CUBES

We denote  $C_R \equiv (-R, R)^n$  as open cubes with side lengths  $2R$  for  $R > 0$  in  $\mathbb{R}^n$ . Suppose that  $F : C_{R+1} \rightarrow \mathbb{R}^m$ ,  $m \in (0, n)$ , is a continuous Gaussian random field, and denote that  $F(x) = (f_1(x), \dots, f_m(x))$ . Then, the  $(x, y)$ -covariance kernel (two-point covariance kernel),

$$(2.1) \quad K_{i_1 i_2}(x, y) = \mathcal{E}(f_{i_1}(x)f_{i_2}(y)), \quad i_1, i_2 = 1, 2, \dots, m,$$

is an  $m \times m$  matrix.

**Definition 2.1.** We say that  $F : C_{R+1} \rightarrow \mathbb{R}^m$  satisfies  $(R; M, k_1)$ -assumptions if on  $C_{R+1}$ ,

- (A1)  $F$  is centred, i.e.,  $\mathcal{E}(f_i(x)) = 0$  for  $i = 1, \dots, m$ . And the joint distribution of  $F(x)$  and its Jacobian  $\nabla F(x)$ , i.e.,  $(f_1(x), \dots, f_m(x), \partial_1 f_1(x), \partial_2 f_1(x), \dots, \partial_n f_m(x))$ , is a non-degenerate Gaussian vector with a uniform lower bound for all  $x \in C_{R+1}$ , i.e.,

$$(2.2) \quad \mathcal{E} \left( \left| \sum_{i=1}^m \eta_i \cdot f_i(x) + \sum_{i=1}^m \sum_{j=1}^n \xi_{ij} \cdot \partial_j f_i(x) \right|^2 \right) \geq k_1 > 0,$$

for all  $x \in C_{R+1}$ , and for any  $\eta = (\eta_i) \in \mathbb{R}^m$ ,  $\xi = (\xi_{ij}) \in \mathbb{R}^{nm}$ , with  $\|\eta\|^2 + \|\xi\|^2 = 1$ .

- (A2)  $F$  is almost surely  $C^{3-} \equiv \cap_{0 < \beta < 1} C^{2+\beta}$  in  $C_{R+1}$ :

$$(2.3) \quad \max_{0 \leq |\alpha| \leq 3} \sup_{x \in C_{R+1}} |D_x^\alpha D_y^\alpha K_{i_1 i_2}(x, y)|_{y=x} \leq M, \quad i_1, i_2 = 1, 2, \dots, m.$$

Furthermore, we say that  $F$  satisfies  $(R; M, k_1, k_2)$ -assumptions if on  $C_{R+1}$ ,  $F$  also satisfies the following two-point nondegeneracy assumption (A3) for some  $k_2 > 0$ .

- (A3) For every pair of distinct points  $x, y \in C_R$ , the Gaussian vector  $(F(x), F(y))$  has a nondegenerate distribution with the following restrictions:

$$(2.4) \quad p_{(F(x), F(y))}(0, 0) \leq \begin{cases} \frac{k_2}{\|x-y\|^m}, & \text{if } \|x-y\| \leq 1, \\ k_2, & \text{if } \|x-y\| > 1, \end{cases}$$

where  $p_{(F(x), F(y))}(0, 0)$  is the density of the joint distribution  $(F(x), F(y))$  at  $(0, 0)$ .

**Remark 2.2.** The condition (A3), and the later condition (B4) in Definition 4.2, will only be used when we prove Theorem 4.7, Theorem 4.4, and Theorem 4.9 in Section 4. The upper bound  $\frac{k_2}{\|x-y\|^m}$  is natural when  $x$  and  $y$  are close to each other, which actually follows from the condition (A1). In our Appendix E, we will explain this and show that (A1) and (A2) would imply (2.4) only under the condition that  $(F(x), F(y))$  has a nondegenerate distribution.



**Theorem 2.3.** *Suppose that  $F : C_{R+1} \rightarrow \mathbb{R}^m$ , satisfies  $(R; M, k_1)$ -assumptions on  $C_{R+1}$ . Then, there is a constant  $D_1 = D_1(n, m, M, k_1) > 0$ , such that*

$$(2.5) \quad \mathcal{E}(N(R; F)) \leq D_1 \cdot |C_R|.$$

Notice that Theorem 2.3 also holds true for any  $r \in (0, R)$  with the the same  $D_1$ .

We need two theorems before we prove Theorem 2.3. The first is the following Kac–Rice theorem, see for example Remark 10 in Section 2 of [CH20] and Theorem 6.10 in Chapter 6 of [AW09]. We restate it here in consistent with our Definition 2.1.

**Theorem 2.4** (Kac-Rice Theorem with Integrand). *Let  $W : C_{R+1} \rightarrow \mathbb{R}^{n_1}$  be a continuous Gaussian random field such that  $(F, W)$  is also a Gaussian field on  $C_{R+1}$ . And suppose that  $h = h(x, w) : \mathbb{R}^n \times \mathbb{R}^{n_1} \rightarrow \mathbb{R}$  is a positive continuous bounded function.*

*Take  $s = 1$  or  $s = 2$ . For  $s = 1$ , we assume that  $F$  satisfies the  $(R; M, k_1)$ -assumptions. For  $s = 2$ , we assume that  $F$  satisfies the  $(R; M, k_1, k_2)$ -assumptions.*

*Then, since  $F : C_{R+1} \rightarrow \mathbb{R}^m$  and  $m \in (0, n)$ , we have that*

$$(2.6) \quad \mathcal{E} \left[ \left( \int_{\{F=0\} \cap B} h(u, W(u)) d\mathcal{H}^{n-m}(u) \right)^s \right] = \int_{B^s} J_{s,F}(h; u_1, \dots, u_s) p_{(F(u_1), \dots, F(u_s))}(0) du_1 \dots du_s$$

for every Borel measurable set  $B \subset C_{R+1}$ , where

$$(2.7) \quad J_{s,F}(h; u_1, \dots, u_s) = \mathcal{E} \left[ \prod_{t=1}^s h(u_t, W(u_t)) \sqrt{\det[(\nabla F(u_t))^T (\nabla F(u_t))]} \Big| F(u_t) = 0, t = 1, \dots, s \right]$$

and  $p_{(F(u_1), \dots, F(u_s))}(0)$  is the density of  $(F(u_t))_{t=1}^s$  at  $(0, \dots, 0)$ ,  $(\nabla F)^T (\nabla F)$  is the  $m \times m$  matrix  $(\nabla f_{i_1} \cdot \nabla f_{i_2})_{m \times m}$ , and  $\mathcal{H}^{n-m}$  is the  $(n - m)$ -dimensional Hausdorff measure, i.e., surface measure for submanifolds in  $\mathbb{R}^n$ .

Our second theorem is to link the number of connected components to some curvature integrals over  $Z(F)$ . Let me cite Theorem 3 in [C71] and readers can see more references therein. We will give more intuitions of geometric inequalities when we state Theorem 2.8 and Theorem 2.9 later.

**Theorem 2.5** (Fenchel–Borsuk–Willmore–Chern–Lashof Inequality). *Let  $M$  be a connected compact  $(n - m)$ -dimensional submanifold of  $\mathbb{R}^n$  without boundary. Then,*

$$(2.8) \quad |\mathbb{S}^{n-m}| \leq \int_M \left\| \frac{\mathbf{H}}{n - m} \right\|^{n-m} d\mathcal{H}^{n-m},$$

where  $|\mathbb{S}^{n-m}|$  is the volume of  $(n - m)$ -dimensional unit sphere and  $\mathbf{H}$  is the mean curvature vector of  $M \subset \mathbb{R}^n$ . The equality in (2.8) holds if and only if when  $n - m > 1$ ,  $M$  is embedded as a hypersphere in an  $(n - m + 1)$ -dimensional linear subspace of  $\mathbb{R}^n$ ; when  $n - m = 1$ ,  $M$  is embedded as a simple convex closed curve in a 2-dimensional linear subspace of  $\mathbb{R}^n$ .

If we already know that all components of  $Z(F)$  contained in  $C_R$  are regular submanifolds, we can then apply this Theorem 2.5, and see that there is a positive constant  $C = C(m, n)$  such that

$$(2.9) \quad \begin{aligned} N(R; F) &\leq C \cdot \sum_{M \subset F^{-1}(0) \cap C_R, \text{disjoint}} \int_M \|\mathbf{H}\|^{n-m} d\mathcal{H}^{n-m} \\ &\leq C \cdot \int_{F^{-1}(0) \cap C_R} \|\mathbf{H}\|^{n-m} d\mathcal{H}^{n-m}. \end{aligned}$$

And then, we can use Theorem 2.4 to compute the expectation of the mean curvature integral. Fortunately, the regularity follows from Bulinskaya's lemma, see for example Proposition 6.12 of [AW09]. In our Appendix A, we will give a quantitative version for Bulinskaya's lemma, which is similar to Lemma 7 in [NS16]. Now, we can use (2.9) to compute our expectation and prove Theorem 2.3.

*Proof of Theorem 2.3.* The general computations for  $\|\mathbf{H}\|$  are included in Appendix F and Theorem F.1. Let us recall the results and rewrite them adapting to our current settings. We can give an expression of  $\|\mathbf{H}\|$  by terms and derivatives of  $F$ . If we set  $\varphi : M \rightarrow \mathbb{R}^n$  as an embedding from a regular connected component  $M$  of  $Z(F)$ ,  $(g_{ij}) \equiv (\partial_i \varphi \cdot \partial_j \varphi)$  and  $(g^{ij}) \equiv (g_{ij})^{-1}$ ,  $A = (A_{\alpha\beta}) \equiv (\nabla f_\alpha \cdot \nabla f_\beta) = ((\nabla F)^T (\nabla F))$ , and  $A^{\alpha\beta}$  denotes  $(A^{-1})_{\alpha\beta}$ , where  $v_1 \cdot v_2$  is the standard inner product for two vectors  $v_1, v_2 \in \mathbb{R}^n$ . Then, we have the following inequalities ( $\|\cdot\|$  are all vectors norms or matrices norms),

$$(2.10) \quad \begin{aligned} \|\mathbf{H}\|^2 &= \sum_{1 \leq \alpha, \beta \leq m} \sum_{1 \leq i, j, k, l \leq n-m} (\nabla^2 f_\alpha)(\partial_i \varphi, \partial_j \varphi) g^{ij} A^{\alpha\beta} (\nabla^2 f_\beta)(\partial_k \varphi, \partial_l \varphi) g^{kl} \\ &\leq C(n, m) \cdot \|A^{-1}\| \cdot \|\nabla^2 F\|^2 \\ &= C(n, m) \cdot |\det A|^{-1} \cdot \|\text{adj} A\| \cdot \|\nabla^2 F\|^2 \\ &\leq C(n, m) \cdot |\det A|^{-1} \cdot \|\nabla F\|^{2(m-1)} \cdot \|\nabla^2 F\|^2, \end{aligned}$$

where  $\text{adj} A$  is the adjugate matrix associated with  $A$ . Here,  $C(n, m)$  are different positive constants but they only depend on  $n, m$ .

According to Theorem 2.4, we now let

$$(2.11) \quad W = (\nabla F, \nabla^2 F),$$

which satisfies that  $(F(u), W(u))$  is Gaussian for every  $u \in C_{R+1}$ . And according to terms in (2.9) and (2.10), we set

$$(2.12) \quad h(x, W) \equiv h(W) \equiv (C(n, m))^{\frac{(n-m)}{2}} \cdot |\det A|^{-\frac{(n-m)}{2}} \cdot \|\nabla F\|^{(n-m)(m-1)} \cdot \|\nabla^2 F\|^{(n-m)}.$$

Here, we regard  $W$  as a vector with elements  $(\nabla F, \nabla^2 F)$  and regard  $h$  as a function of this vector  $W$  and independent of  $x$ . To be consistent with the boundedness assumption in our Theorem 2.4 (see also Remark 10 in Section 2 of [CH20] and Theorem 6.10 in [AW09]), we need to modify  $h$  to make it bounded. We denote the closed singular set of  $h$  as  $\mathcal{S}$ , which

is of Hausdorff codimension 1 in the vector space of  $W$  because  $\det A$  is a polynomial with elements in  $\nabla F$ . So, we can construct a monotone family of positive continuous cut-off functions  $\varphi_Q(W)$ , such that  $\varphi_Q \rightarrow 1$  as  $Q \rightarrow \infty$ , and  $\varphi_{Q_1}(W) < \varphi_{Q_2}(W)$  when  $Q_1 < Q_2$ , and  $\varphi_Q(W) = 0$  when  $\text{dist}(W, \mathcal{S}) < 1/Q$  or  $\|W\| > Q$ . We then set

$$(2.13) \quad h_Q(x, W) \equiv h_Q(W) \equiv \varphi_Q(W)h(W), \quad Q > 0,$$

which is a monotone family of positive continuous bounded functions.

We can then see that, by monotone convergence theorem,

$$(2.14) \quad \begin{aligned} & \mathcal{E} \left( \int_{F^{-1}(0) \cap C_R} (\|\mathbf{H}(u)\|^{(n-m)}) d\mathcal{H}^{n-m}(u) \right) \leq \mathcal{E} \left( \int_{F^{-1}(0) \cap C_R} h(u, W(u)) d\mathcal{H}^{n-m}(u) \right) \\ & = \mathcal{E} \left( \lim_{Q \rightarrow \infty} \int_{F^{-1}(0) \cap C_R} h_Q(u, W(u)) d\mathcal{H}^{n-m}(u) \right) \\ & = \lim_{Q \rightarrow \infty} \mathcal{E} \left( \int_{F^{-1}(0) \cap C_R} h_Q(u, W(u)) d\mathcal{H}^{n-m}(u) \right) \\ & = \lim_{Q \rightarrow \infty} \int_{C_R} J_{1,F}(h_Q; u) \cdot p_{F(u)}(0) du \\ & = \lim_{Q \rightarrow \infty} \int_{C_R} \mathcal{E} [h_Q(u, W(u)) \cdot |\det A|^{1/2} \mid F(u) = 0] \cdot p_{F(u)}(0) du \\ & = \int_{C_R} \mathcal{E} [h(u, W(u)) \cdot |\det A|^{1/2} \mid F(u) = 0] \cdot p_{F(u)}(0) du. \end{aligned}$$

Since  $F$  satisfies (A1) in Definition 2.1,  $p_{F(u)}(0) \leq C(n, m, k_1)$  for any  $u \in C_R$ . To estimate the conditional expectation, we first notice that there are terms including  $\nabla F$  and  $\nabla^2 F$ , which we are going to separate using the inequality  $ab \leq \frac{k}{k+1} a^{\frac{k}{k+1}} + \frac{1}{k+1} b^{k+1} \leq a^{\frac{k}{k+1}} + b^{k+1}$  for some positive integer  $k = k(m, n)$  to be determined later. The conditional expectation term in (2.14) then equals to

$$(2.15) \quad \begin{aligned} & \mathcal{E} [|\det A(u)|^{\frac{1-(n-m)}{2}} \cdot \|\nabla F(u)\|^{(n-m)(m-1)} \cdot \|\nabla^2 F(u)\|^{(n-m)} \mid F(u) = 0] \\ & \leq \mathcal{E} \left[ \left( |\det A(u)|^{\frac{1-(n-m)}{2}} \cdot \|\nabla F(u)\|^{(n-m)(m-1)} \right)^{\frac{k+1}{k}} + \|\nabla^2 F(u)\|^{(n-m)(k+1)} \mid F(u) = 0 \right]. \end{aligned}$$

We now need to bound

$$(2.16) \quad \mathcal{E} \left[ \left( |\det A(u)|^{\frac{1-(n-m)}{2}} \cdot \|\nabla F(u)\|^{(n-m)(m-1)} \right)^{\frac{k+1}{k}} \mid F(u) = 0 \right]$$

and

$$(2.17) \quad \mathcal{E} \left[ \|\nabla^2 F(u)\|^{(n-m)(k+1)} \mid F(u) = 0 \right]$$

separately.

The estimate for (2.17) is direct. For each  $u \in C_R$ , and each  $\partial_{j_1 j_2}^2 f_i(u)$ , one needs to find  $a_{j_1 j_2, i}^t(u)$  such that for each  $s = 1, \dots, m$ ,

$$(2.18) \quad \mathcal{E} \left[ \left( \partial_{j_1 j_2}^2 f_i(u) + \sum_{t=1}^m a_{j_1 j_2, i}^t f_t(u) \right) \cdot f_s(u) \right] = 0.$$

Since  $F$  satisfies (A1) in Definition 2.1, the matrix  $(\mathcal{E}(f_t(u) \cdot f_s(u)))_{m \times m}$  is always invertible and has uniformly lowerly bounded determinant value for all  $u \in C_R$ , and then those  $a_{j_1 j_2, i}^t$  are solvable and there is a constant  $C = C(n, m, M, k_1)$  such that

$$(2.19) \quad |a_{j_1 j_2, i}^t| \leq C.$$

And hence, there is another constant  $C = C(n, m, M, k_1, k) = C(n, m, M, k_1)$  (since  $k$  only depend on  $n, m$  and will be determined later) such that

$$(2.20) \quad \begin{aligned} & \mathcal{E} \left[ |\partial_{j_1 j_2}^2 f_i(u)|^{(n-m)(k+1)} \mid F(u) = 0 \right] \\ &= \mathcal{E} \left[ \left| \partial_{j_1 j_2}^2 f_i(u) + \sum_{t=1}^m a_{j_1 j_2, i}^t f_t(u) \right|^{(n-m)(k+1)} \right] \leq C. \end{aligned}$$

The estimate for (2.16) needs more ingredients. First, notice that since  $(F, \nabla F)$  satisfies (A1) in Definition 2.1, then for each  $u \in C_R$ , and each  $\partial_j f_i(u)$ , we need to find  $a_{j, i}^t(u)$  such that for each  $s = 1, \dots, m$ ,

$$(2.21) \quad \mathcal{E} \left[ \left( \partial_j f_i(u) + \sum_{t=1}^m a_{j, i}^t f_t(u) \right) \cdot f_s(u) \right] = 0.$$

By similar reasons estimating those  $a_{j_1 j_2, i}^t$  in (2.18),  $a_{j, i}^t(u)$  are also solvable and there is a constant  $C = C(n, m, M, k_1)$  such that

$$(2.22) \quad |a_{j, i}^t| \leq C.$$

Next, we replace each  $\partial_j f_i(u)$  in (2.16) by  $(\partial_j f_i)^\# \equiv \partial_j f_i(u) + \sum_{t=1}^m a_{j, i}^t f_t(u)$ , and also define  $(\nabla F(u))^\#$  as replacing elements  $\partial_j f_i$  in  $\nabla F$  by  $(\partial_j f_i)^\#$ . Since the joint distribution  $(F, \nabla F)$  is non-degenerate, the Gaussian vector  $(\nabla F(u))^\#$  is also non-degenerate.

We can divide the non-degenerate  $nm$ -dimensional Gaussian vector  $(\nabla F(u))^\#$  into  $m$   $n$ -dimensional vectors:

$$(2.23) \quad (\nabla F(u))^\# = (v_1(u), v_2(u), \dots, v_m(u)),$$

where for each  $t = 1, \dots, m$ ,

$$(2.24) \quad v_t(u) = ((\nabla f_t(u))^\#)^T = ((\partial_1 f_t(u))^\#, (\partial_2 f_t(u))^\#, \dots, (\partial_n f_t(u))^\#).$$

With these notations, we also set  $(A(u))^\# = (v_{t_1} \cdot v_{t_2})_{m \times m}$ , and we see that (2.16) equals to

$$(2.25) \quad \mathcal{E} \left[ \left( \left| \det(A(u))^\# \right|^{\frac{1-(n-m)}{2}} \cdot \|(\nabla F(u))^\#\|^{(n-m)(m-1)} \right)^{\frac{k+1}{k}} \right],$$

and notice that the determinant of the covariance kernel for  $(\nabla F(u))^\#$  is lowerly bounded by some constant  $C = C(n, m, M, k_1)$  as in (A1) and (A2) in Definition 2.1, the (2.25) then reduces to estimating the following integral:

$$(2.26) \quad C(n, m, M, k_1) \int_{(\mathbb{R}^n)^{\otimes m}} |\det(v_{t_1} \cdot v_{t_2})|^\alpha \cdot \|v_1\|^2 + \dots + \|v_m\|^2 |^\beta \cdot e^{-c(n, m, M, k_1)(\|v_1\|^2 + \dots + \|v_m\|^2)} dv_1 \cdots dv_m,$$

where  $\alpha = \frac{k+1}{k} \cdot \frac{1-(n-m)}{2}$ , which is very close to but more negative than  $-\frac{n-m-1}{2}$  since  $k$  will be chosen to be large enough, and  $\beta = \frac{k+1}{2k} \cdot (n-m)(m-1)$ , and we use  $\|\cdot\|$  to denote vector norms temporarily. And  $c(n, m, M, k_1)$  is a positive constant related to the covariance kernel for  $(\nabla F(u))^\#$ . We can simplify the integral in (2.26) further and then get

$$(2.27) \quad \left( \int_{(\mathbb{S}^{n-1})^{\otimes m}} |\det(w_{t_1} \cdot w_{t_2})|^\alpha d\sigma(w_1) \cdots d\sigma(w_m) \right) \cdot \int_{(\mathbb{R}_+)^{\otimes m}} |r_1|^{2\alpha+n-1} \cdots |r_m|^{2\alpha+n-1} \|r_1\|^2 + \dots + \|r_m\|^2 |^\beta \cdot e^{-c(n, m, M, k_1)(\|r_1\|^2 + \dots + \|r_m\|^2)} dr_1 \cdots dr_m,$$

where we use spherical coordinates and  $\sigma(\cdot)$  is the surface measure on the unit sphere  $\mathbb{S}^{n-1}$ . Notice that  $0 < 2\alpha + n - 1 < n - 1$  since  $m \geq 1$  and  $\alpha = -\frac{n-m-1}{2} - \epsilon$  for some small  $\epsilon = \epsilon(m, n) > 0$  to be determined later. Hence, the second multiplier is bounded by some constant  $C = C(n, m, M, k_1)$ . The rest is to consider the singular integral

$$(2.28) \quad \int_{(\mathbb{S}^{n-1})^{\otimes m}} |\det(w_{t_1} \cdot w_{t_2})|^\alpha d\sigma(w_1) \cdots d\sigma(w_m)$$

for some  $\alpha < 0$ .

Currently, our  $\alpha$  is close to  $-\frac{n-m-1}{2}$  but not less than  $-\frac{n-m}{2}$ . But indeed, we can prove a stronger statement and the sharp  $\alpha$  is actually  $-\frac{n-m+1}{2}$ ! Hence, one can just choose any  $\epsilon = \epsilon(m, n) \in (0, 1/2]$ , so that  $k$  is well-chosen in the previous settings and only depend on  $m, n$ . The sharp value for  $\alpha$  will be used when we prove the quantitative Bulinskaya's lemma, our Lemma A.1 in Appendix A.

**Lemma 2.6.** *Let  $n > m > 0$ . For any  $s_0 \in (0, 1)$  and  $\alpha \in (-\frac{n-m+1}{2}s_0, 0)$ , there is a positive constant  $C = C(s_0, n, m)$  such that*

$$(2.29) \quad \int_{(\mathbb{S}^{n-1})^{\otimes m}} |\det(w_{t_1} \cdot w_{t_2})|^\alpha d\sigma(w_1) \cdots d\sigma(w_m) \leq C.$$

*Proof of Lemma 2.6.* If  $m = 1$ , there is nothing to prove. For  $m \geq 2$ , we fix  $\{e_1, \dots, e_n\}$  as the standard orthonormal frame of  $\mathbb{R}^n$ . Notice that  $|\det(w_{t_1} \cdot w_{t_2})|$  is invariant under actions of  $O(n)$ , hence, the left hand side of (2.29) equals to

$$(2.30) \quad |\mathbb{S}^{n-1}| \cdot \int_{(\mathbb{S}^{n-1})^{\otimes (m-1)}} |\det(A(e_1))|^\alpha d\sigma(w_2) \cdots d\sigma(w_m),$$

where  $A(e_1)$  has the form

$$(2.31) \quad A(e_1) = \begin{bmatrix} 1 & e_1 \cdot w_2 & \cdots & e_1 \cdot w_n \\ w_2 \cdot e_1 & 1 & \cdots & w_2 \cdot w_n \\ \vdots & \vdots & \cdots & \vdots \\ w_n \cdot e_1 & w_n \cdot w_2 & \cdots & 1 \end{bmatrix}.$$

By elementary row operations, we see that  $A(e_1)$  has the same determinant value as the matrix

$$(2.32) \quad A'(e_1) = \begin{bmatrix} 1 & e_1 \cdot w_2 & \cdots & e_1 \cdot w_n \\ 0 & & & \\ \vdots & & & \\ 0 & & \left[ w_{t_1} \cdot w_{t_2} - (e_1 \cdot w_{t_1})(e_1 \cdot w_{t_2}) \right]_{(m-1) \times (m-1)} & \end{bmatrix}.$$

Then, we can parametrize  $\mathbb{S}^{n-1}$  by  $B^{n-1}$ , the unit disc of dimension  $n-1$ . That is, for each  $t = 2, \dots, m$ , almost all  $w_t \in \mathbb{S}^{n-1}$  can be written in coordinates of the form  $(\sqrt{1 - |w'_t|^2}, w'_t)$  or  $(-\sqrt{1 - |w'_t|^2}, w'_t)$  for  $w'_t = (w_t(2), \dots, w_t(n)) \in B^{n-1}$ . And then  $w_{t_1} \cdot w_{t_2} - (e_1 \cdot w_{t_1})(e_1 \cdot w_{t_2}) = \langle w'_{t_1}, w'_{t_2} \rangle_{\mathbb{R}^{n-1}}$ , where  $\langle \cdot, \cdot \rangle_{\mathbb{R}^{n-1}}$  is the standard inner product in  $\mathbb{R}^{n-1}$ . By the area formula, we can rewrite (2.30) as

$$(2.33) \quad \begin{aligned} & |\mathbb{S}^{n-1}| \cdot 2^{m-1} \cdot \int_{(B^{n-1})^{\otimes (m-1)}} |\det(\langle w'_{t_1}, w'_{t_2} \rangle_{\mathbb{R}^{n-1}})|^\alpha \cdot \prod_{t=2}^m \frac{1}{\sqrt{1 - |w'_t|^2}} dw'_2 \cdots dw'_m \\ & = |\mathbb{S}^{n-1}| \cdot 2^{m-1} \cdot \int_{(\mathbb{S}^{n-2})^{\otimes (m-1)}} |\det(\langle \tilde{w}_{t_1}, \tilde{w}_{t_2} \rangle_{\mathbb{R}^{n-1}})|^\alpha d\tilde{\sigma}(\tilde{w}_2) \cdots d\tilde{\sigma}(\tilde{w}_m) \\ & \quad \cdot \int_{(0,1)^{\otimes (m-1)}} \prod_{t=2}^m \frac{r_t^{2\alpha+n-2}}{\sqrt{1 - |r_t|^2}} dr_2 \cdots dr_m. \end{aligned}$$

Here  $\tilde{\sigma}(\cdot)$  is the surface measure of  $\mathbb{S}^{n-2}$ . Notice that since  $n > m \geq 2$ , we have that  $-(n-m+1)s_0 + n - 2 \geq -(n-1)s_0 + n - 2 = (n-1)(1-s_0) - 1 \geq 2(1-s_0) - 1 = 1 - 2s_0$ . Hence,

$$(2.34) \quad \int_0^1 r^{2\alpha+n-2} dr \leq \int_0^1 r^{-(n-m+1)s_0+n-2} dr \leq \frac{1}{2(1-s_0)}.$$

If  $m = 2$ , we finish our proof as in (2.33), the determinant integral is just 1. If  $m > 2$ , then we run the same process for the pair  $n-2, m-1$  with the condition that  $n-2 \geq m-1 \geq 2$ . Continue doing this process, we finish the proof.  $\square$

With this Lemma 2.6, we then bound (2.16) by a constant  $C = C(n, m, M, k_1) > 0$ . Combine this constant with the constant  $C = C(n, m, M, k_1) > 0$  bounding (2.17), we then use (2.9) and finish our proofs for Theorem 2.3.

□

**Remark 2.7.** The sharp constant in Theorem 2.5 does not influence our proofs for Theorem 2.3 as we only need an upper bound. On the other hand, one does not expect that we can calculate the accurate expectation for the number of connected components in such a way. For example, when  $n = 3$  and  $m = 2$ ,  $Z(F)$  consists of many closed curves, which we call knots. And then, the Fáry–Milnor’s theorem says that if a closed curve is not an unknot, then the integral on the right hand side of (2.8) is strictly greater than  $4\pi$ , while in this case  $|\mathbb{S}^1| = 2\pi$ . And by our later Theorem 3.1 (also stated as Theorem 1.1 previously), all types of knots will appear with positive probabilities in limiting cases under some assumptions. So, even an accurate calculation of curvature integrals will not tell us the accurate value of the limiting expectation for the number of connected components.

Next, for Betti numbers, we can build up similar upper bounds as in Theorem 2.3.

**Theorem 2.8.** *Suppose that  $F : C_{R+1} \rightarrow \mathbb{R}^m$ , satisfies  $(R; M, k_1)$ -assumptions on  $C_{R+1}$ . Then, there is a constant  $\tilde{D}_1 = \tilde{D}_1(n, m, M, k_1) > 0$  such that*

$$(2.35) \quad \sum_{l=0}^{n-m} \mathcal{E}(\beta_l(R; F)) \leq \tilde{D}_1 \cdot |C_R|.$$

The proof is exactly the same as Theorem 2.3 once we replace Theorem 2.5 by Chern–Lashof inequalities in [CL57] and [CL58], which connect the topology of a closed manifold to its mean curvature related curvature integrals. On the other hand, there are some other works and techniques in differential geometry which lead similar topological controls from the study of well-known vanishing theorems. See for example, [B88] and references therein, where a De Giorgi–Nash–Moser iteration scheme also leads to a control on Betti numbers. But the forms shown in [CL57] and [CL58] are cleaner and well adapt to our applications here. The author also proved them independently when he studied optimal constants problems in geometric inequalities, but then learnt that these things were studied well by very great geometers many years ago. The following inequality is exactly the Theorem 1 in [CL58] but we rewrite it a little to be adapted to our notations.

**Theorem 2.9** (Chern–Lashof Inequality). *If  $M$  is a connected compact  $(n - m)$ -dimensional submanifold of  $\mathbb{R}^n$  without boundary, then,*

$$(2.36) \quad \sum_{l=0}^{n-m} \beta_l(M) \leq \frac{1}{|\mathbb{S}^{n-1}|} \cdot \int_M \int_{T_x^\perp M \cap \{|y|=1\}} |\det(\langle \text{II}(x), y \rangle_{\mathbb{R}^n})| d\sigma^{m-1}(y) d\mathcal{H}^{n-m}(x).$$

In (2.36),  $|\mathbb{S}^{n-1}|$  is the volume of  $(n - 1)$ -dimensional unit sphere,  $\beta_l(M)$  is the  $l$ -th Betti number of  $M$  over  $\mathbb{R}$ ,  $\text{II}(x)$  is the second fundamental form of  $M \subset \mathbb{R}^n$  at  $x$ , and  $y$  ranges over unit vectors on the fiber  $T_x^\perp M$  of normal bundle  $T^\perp M$ ,  $\langle \cdot, \cdot \rangle_{\mathbb{R}^n}$  is the Euclidean inner product in  $\mathbb{R}^n$ , and  $\sigma^{m-1}$  is the surface measure of  $\mathbb{S}^{m-1} \subset T_x^\perp M$ .

Then, it is easy to see that  $|\det(\langle \mathbb{H}(x), y \rangle)| \leq C(n, m) \cdot \|\mathbb{H}(x)\|^{n-m}$ . To estimate the norm of  $\mathbb{H}(x)$ , we can get from Theorem F.1 that, similar in estimating  $\|\mathbf{H}\|^2$  in (2.10),

$$\begin{aligned}
(2.37) \quad \|\mathbb{H}\|^2 &\leq C(n, m) \cdot \|A^{-1}\| \cdot \|\nabla^2 F\|^2 \\
&= C(n, m) \cdot |\det A|^{-1} \cdot \|\text{adj} A\| \cdot \|\nabla^2 F\|^2 \\
&\leq C(n, m) \cdot |\det A|^{-1} \cdot \|\nabla F\|^{2(m-1)} \cdot \|\nabla^2 F\|^2,
\end{aligned}$$

for some positive constants  $C(n, m)$ . We just use the same  $h$  and  $h_Q$  as shown in the proof of Theorem 2.3 and then can get a proof for Theorem 2.8.

### 3. PROOF OF THEOREM 1.1 AND THEOREM 1.2

In this section, we assume that  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $m \in (0, n)$ , is a centred stationary Gaussian random field satisfying (A1) and (A2) in Definition 2.1 at the origin  $x = 0$ , which is equivalent to say that  $F$  satisfies the  $(R; M, k_1)$ -assumptions on at the origin  $x = 0$ . Also, since  $F$  is stationary, these assumptions hold at any point  $x \in \mathbb{R}^n$ .

We will focus on Theorem 1.1 first and explain the necessary modifications in proving Theorem 1.2 after the proof of Theorem 1.1. We first explain some definitions in Theorem 1.1 and give some basic settings similar to Section 6 in [NS16]. We denote  $\mathfrak{B}(C^1(\mathbb{R}^n, \mathbb{R}^m))$  as the Borel  $\sigma$ -algebra generated by open sets in  $C^1(\mathbb{R}^n, \mathbb{R}^m)$ , and  $\gamma_F = F_*(\mathcal{P})$  with  $\mathcal{P}$  the probability measure in the background probability space  $(\Omega, \mathfrak{S}, \mathcal{P})$ . By Bulinskaya's lemma (see either Proposition 6.12 of [AW09] or our Lemma A.1), we see that we only need to consider a Borel subset of  $C^1(\mathbb{R}^n, \mathbb{R}^m)$ :

$$(3.1) \quad C_*^1(\mathbb{R}^n, \mathbb{R}^m) \equiv \{G \in C^1(\mathbb{R}^n, \mathbb{R}^m) \mid \nabla G \text{ is of full rank on } Z(G)\},$$

since  $\gamma_F(C^1(\mathbb{R}^n, \mathbb{R}^m) \setminus C_*^1(\mathbb{R}^n, \mathbb{R}^m)) = 0$ .

The action of  $\mathbb{R}^n$  on  $(C_*^1(\mathbb{R}^n, \mathbb{R}^m), \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m)), \gamma_F)$  is by shifts  $\tau_v : G(x) \mapsto G(x + v)$ , which is measure preserving since  $F$  is stationary. We say that the action of  $\mathbb{R}^n$  is *ergodic* if for every set  $A \subset \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m))$  satisfying  $\gamma_F((\tau_v)A\Delta A) = 0$  for all  $v \in \mathbb{R}^n$ , then either  $\gamma_F(A) = 0$  or  $\gamma_F(A) = 1$ . Here  $\Delta$  is the symmetric difference.

So, we can restate Theorem 1.1 in the following way.

**Theorem 3.1.** *Assume that  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $m \in (0, n)$ , is a centred stationary Gaussian random field satisfying the  $(R; M, k_1)$ -assumptions at the origin  $x = 0$ .*

(1) *There exists a number  $\nu = \nu_F \geq 0$  so that*

$$(3.2) \quad \mathcal{E}(N(R; F)) = \nu \cdot |C_R| + o_{R \rightarrow \infty}(R^n).$$

(2) *If the translation action of  $\mathbb{R}^n$  on  $(C_*^1(\mathbb{R}^n, \mathbb{R}^m), \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m)), \gamma_F)$  is ergodic, then*

$$(3.3) \quad \lim_{R \rightarrow \infty} \frac{N(R; F)}{|C_R|} = \nu \text{ almost surely} \quad \text{and} \quad \lim_{R \rightarrow \infty} \mathcal{E} \left| \frac{N(R; F)}{|C_R|} - \nu \right| = 0.$$

Moreover, for any  $H \in H(n - m)$ , (1) and (2) hold true if one replaces  $N(R; F)$  by  $N(R; F, H)$  and replaces the number  $\nu_F$  by a  $\nu_{F, H} \geq 0$ .



(3)

$$(3.4) \quad \nu_F = \sum_{H \in H(n-m)} \nu_{F,H}.$$

**Remark 3.2.** When  $m = 1$ , two main assumptions in [NS16] are about when the shifting action of  $\mathbb{R}^n$  is ergodic and when  $\nu_F > 0$ . Similar assumptions were made in [SW19] for  $\nu_{F,H} > 0$ . For general  $m > 1$ , one may need more ingredients. But at least, if the random fields  $F$  consists of independent random functions, i.e.,  $F = (f_1, \dots, f_m)$  and  $f_{i_1}$  is independent of  $f_{i_2}$  when  $i_1 \neq i_2$ , we can denote the spectral measure of  $f_i$  as  $\rho_i$ , and assume that  $\rho_i$  has no atoms and its compact support has nonempty interior. These two assumptions are exactly those in [NS16] and [SW19], under which we can get the ergodicity and positivity. In Appendix C and Appendix D, we discuss the ergodicity and positivity without the independence assumption, but add other assumptions to obtain the ergodicity and positivity respectively.

To prove Theorem 3.1, we define  $N(x, R; F)$  as the number of all connected components of  $Z(F)$  fully contained in the open cube  $C_R(x) = C_R + x$ , shifted from  $C_R$  by  $x$ , of  $\mathbb{R}^n$ , and  $N^*(x, R; F)$  to be the number of all connected components of  $Z(F)$  that intersect the closed cube  $\overline{C_R(x)}$ . For any  $H \in H(n-m)$ , we can also define  $N(x, R; F, H)$  and  $N^*(x, R; F, H)$  in a similar way.

So, for any  $G \in C_*^1(\mathbb{R}^n, \mathbb{R}^m)$ ,  $N^*(R; G) - N(R; G) \leq \mathfrak{N}_\#(R; G)$  for

$$(3.5) \quad \mathfrak{N}_\#(R; G) = \begin{cases} \sum_{k=i}^{n-1} \mathfrak{N}_k(R; G), & \text{if } Z(G) \text{ transversally intersect and only intersect with} \\ & i\text{-th skeletons to } (n-1)\text{-th skeletons of } \partial C_R \\ & \text{for some } i \in \{m, \dots, n-1\}, \\ +\infty, & \text{otherwise. And we denote this set as } \text{Degen}(R). \end{cases}$$

Here,  $\mathfrak{N}_k(R; G)$  is the number of connected components of  $Z(G)$  which intersect with skeletons of dimension  $k$  but do not intersect with skeletons in dimensions  $\leq k$ .

Notice that the measurability of  $G \mapsto N(R; G)$  for fixed  $R$  follows from its lower semicontinuity on  $C_*^1(\mathbb{R}^n, \mathbb{R}^m)$ . For the measurability of  $\mathfrak{N}_\#(R; G)$ , one just notices that transversal intersection is stable under small  $C^1$ -perturbations, and hence  $C_*^1(\mathbb{R}^n, \mathbb{R}^m) \setminus \text{Degen}(R)$  is an open subset of  $C_*^1(\mathbb{R}^n, \mathbb{R}^m)$ . By applying Bulinskaya's Lemma on each skeleton of  $\partial C_R$ , we see that  $\gamma_F(\text{Degen}(R)) = 0$ . The measurability of  $\mathfrak{N}_\#(R; G)$  then follows again from the lower semicontinuity on  $C_*^1(\mathbb{R}^n, \mathbb{R}^m) \setminus \text{Degen}(R)$ .

For the expectation of  $\mathcal{E}(\mathfrak{N}_k(R; F))$ , one just restricts the Gaussian random field  $F$  on each  $k$ -th skeleton of  $\partial C_R$ , which is a  $k$ -dimensional open cube in the corresponding  $k$ -dimensional linear subspace of  $\mathbb{R}^n$ . Such a restriction still keeps the conditions (A1) and (A2) in Definition 2.1. Hence, by Theorem 2.3, there is a constant  $C = C(k, m, M, k_1) > 0$  such that  $\mathcal{E}(\mathfrak{N}_k(R; F)) \leq C \cdot R^k$ . And hence, there is a constant  $C = C(n, m, M, k_1) > 0$  such that  $\mathcal{E}(\mathfrak{N}_\#(R; F)) \leq C \cdot R^{n-1}$ . This observation is crucial in proving Theorem 3.1.

We then need an Integral–Geometric Sandwich lemma, see for example Lemma 1 in [NS16] or Lemma 1 in [S16].

**Lemma 3.3.** *For every  $r \in (0, R)$ , we have that*

$$(3.6) \quad \frac{1}{|C_r|} \int_{C_{R-r}} N(x, r; F) \, dx \leq N(R; F) \leq \frac{1}{|C_r|} \int_{C_{R+r}} N^*(x, r; F) \, dx.$$

The proof of this Lemma is exactly the same as the one in [NS16]. For any  $H \in H(n-m)$ , similar inequalities hold true if one replaces  $N(x, r; F)$  (resp.,  $N(R; F)$ ,  $N^*(x, r; F)$ ) by  $N(x, r; F, H)$  (resp.,  $N(R; F, H)$ ,  $N^*(x, r; F, H)$ ). See for example Lemma 3.7 in [SW19].

*Proof of (1) in Theorem 3.1.* We only prove (1) for  $N(R; F)$  and the proof for  $N(R; F, H)$  is same. Set

$$(3.7) \quad \nu = \limsup_{R \rightarrow \infty} \frac{\mathcal{E}(N(R; F))}{R^n},$$

which is bounded by a constant  $D_1 = D_1(n, m, M, k_1) > 0$  by Theorem 2.3. So for each  $\epsilon > 0$ , we can choose an  $r > 0$  such that

$$(3.8) \quad \frac{\mathcal{E}(N(r; F))}{r^n} \geq \nu - \epsilon.$$

By Lemma 3.3, we have that for all  $R > r$ ,

$$(3.9) \quad \mathcal{E}(N(R; F)) \geq \frac{1}{|C_r|} \int_{C_{R-r}} \mathcal{E}(N(x, r; F)) \, dx = \frac{|C_{R-r}|}{|C_r|} \cdot \mathcal{E}(N(r; F)),$$

since  $F$  is stationary. Hence,

$$(3.10) \quad \liminf_{R \rightarrow \infty} \frac{\mathcal{E}(N(R; F))}{R^n} \geq \frac{\mathcal{E}(N(r; F))}{r^n} \geq \nu - \epsilon.$$

Let  $\epsilon \rightarrow 0$ , we then get (3.2). □

*Proof of (2) in Theorem 3.1.* We only prove (2) for  $N(R; F)$  and the proof for  $N(R; F, H)$  is same. We will also use the Wiener’s ergodic theorem on page 28 of [NS16] since we have the ergodicity assumption. Hence, we set the two random variables:

$$(3.11) \quad \Phi(R, r) \equiv \frac{1}{|C_r|} \int_{C_{R-r}} N(x, r; F) \, dx = \frac{1}{|C_r|} \int_{C_{R-r}} N(r; \tau_x(F)) \, dx,$$

and

$$(3.12) \quad \Psi(R, r) \equiv \frac{1}{|C_r|} \int_{C_{R+r}} \mathfrak{N}_\#(x, r; F) \, dx = \frac{1}{|C_r|} \int_{C_{R+r}} \mathfrak{N}_\#(r; \tau_x(F)) \, dx.$$

By Lemma 3.3, we see that  $\Phi(R, r) \leq N(R; F) \leq \Phi(R + 2r, r) + \Psi(R, r)$ . Also, the Wiener’s ergodic theorem gives that, for fixed  $r$ ,

$$(3.13) \quad \lim_{R \rightarrow \infty} \frac{\Phi(R, r)}{|C_{R-r}|} = \frac{\mathcal{E}(N(r; F))}{|C_r|}, \text{ and } \lim_{R \rightarrow \infty} \frac{\Psi(R, r)}{|C_{R+r}|} = \frac{\mathcal{E}(\mathfrak{N}_\#(r; F))}{|C_r|},$$

a.s. and in  $L^1(\mathcal{P})$ . Equivalently,

$$(3.14) \quad \lim_{R \rightarrow \infty} \frac{\Phi(R, r)}{|C_R|} = \frac{\mathcal{E}(N(r; F))}{|C_r|}, \text{ and } \lim_{R \rightarrow \infty} \frac{\Psi(R, r)}{|C_R|} = \frac{\mathcal{E}(\mathfrak{N}_\#(r; F))}{|C_r|},$$

a.s. and in  $L^1(\mathcal{P})$ . Notice that

$$(3.15) \quad \begin{aligned} \left| \frac{N(R; F)}{|C_R|} - \nu \right| &\leq \left| \frac{N(R; F)}{|C_R|} - \frac{\Phi(R, r)}{|C_R|} \right| + \left| \frac{\Phi(R, r)}{|C_R|} - \frac{\mathcal{E}(N(r; F))}{|C_r|} \right| + \left| \frac{\mathcal{E}(N(r; F))}{|C_r|} - \nu \right| \\ &= \left| \frac{N(R; F)}{|C_R|} - \frac{\Phi(R, r)}{|C_R|} \right| + \left| \frac{\Phi(R, r)}{|C_R|} - \frac{\mathcal{E}(N(r; F))}{|C_r|} \right| + \left| \frac{\mathcal{E}(N(r; F))}{|C_r|} - \nu \right|. \end{aligned}$$

Hence, for the convergence in mean part, after taking expectations on both sides and letting  $R \rightarrow \infty$ , by (3.2),

$$(3.16) \quad \begin{aligned} 0 &\leq \limsup_{R \rightarrow \infty} \mathcal{E} \left[ \left| \frac{N(R; F)}{|C_R|} - \nu \right| \right] \\ &\leq \nu - \frac{\mathcal{E}(N(r; F))}{|C_r|} + \left| \frac{\mathcal{E}(N(r; F))}{|C_r|} - \nu \right|. \end{aligned}$$

Then, let  $r \rightarrow \infty$ , by (3.2) again, we get the convergence in mean part of (3.3).

For the almost sure convergence part, we see that we can further reduce (3.15) and get

$$(3.17) \quad \begin{aligned} \left| \frac{N(R; F)}{|C_R|} - \nu \right| &\leq \frac{\Phi(R + 2r, r) + \Psi(R, r)}{|C_R|} - \frac{\Phi(R, r)}{|C_R|} + \left| \frac{\Phi(R, r)}{|C_R|} - \frac{\mathcal{E}(N(r; F))}{|C_r|} \right| \\ &\quad + \left| \frac{\mathcal{E}(N(r; F))}{|C_r|} - \nu \right|. \end{aligned}$$

So, a.s.,

$$(3.18) \quad 0 \leq \limsup_{R \rightarrow \infty} \left| \frac{N(R; F)}{|C_R|} - \nu \right| \leq \frac{\mathcal{E}(\mathfrak{N}_\#(r; F))}{|C_r|} + \left| \frac{\mathcal{E}(N(r; F))}{|C_r|} - \nu \right|.$$

By the discussions before Lemma 3.3, we see that  $\mathcal{E}(\mathfrak{N}_\#(r; F)) \leq C \cdot r^{n-1}$  for some  $C = C(n, m, M, k_1) > 0$ . Then, let  $r \rightarrow \infty$  and use (3.2) again, we get the almost sure convergence part of (3.3).  $\square$

*Proof of (3) in Theorem 3.1.* By definition, for any  $F \in C_*^1(\mathbb{R}^n, \mathbb{R}^m)$  and any  $R > 1$ ,

$$(3.19) \quad N(R; F) = \sum_{H \in H(n-m)} N(R; F, H).$$

For a subset  $A \subset H(n-m)$ , we define  $N(R; F, A)$  to be the number of connected components of  $Z(F)$  lying entirely in  $C_R$  with the diffeomorphism classes lying in  $A$ . Hence, for any finite  $A \subset H(n-m)$ ,

$$(3.20) \quad N(R; F) \geq \sum_{H \in A} N(R; F, H).$$

Take expectations on both sides and divide both sides by  $|C_R|$ , then let  $R \rightarrow \infty$ , by (1) of Theorem 3.1, we see that  $\nu_F \geq \sum_{H \in A} \nu_{F,H}$ . Since  $A$  is arbitrary, we see that

$$(3.21) \quad \nu_F \geq \sum_{H \in H(n-m)} \nu_{F,H}.$$

On the other hand, for any  $H \in H(n-m)$ , by Lemma 3.3 applied for  $N(R; F, H)$ , since  $F$  is stationary,

$$(3.22) \quad \frac{|C_{R-r}|}{|C_r|} \cdot \mathcal{E}(N(r; F, H)) = \frac{1}{|C_r|} \int_{C_{R-r}} \mathcal{E}(N(x, r; F, H)) dx \leq \mathcal{E}(N(R; F, H)),$$

when  $0 < r < R$ . Divide both sides by  $|C_R|$  and let  $R \rightarrow \infty$ , we see that for any  $r > 0$ ,

$$(3.23) \quad \frac{\mathcal{E}(N(r; F, H))}{|C_r|} \leq \nu_{F,H}.$$

Hence, by monotone convergence theorem,

$$(3.24) \quad \frac{\mathcal{E}(N(r; F))}{|C_r|} = \mathcal{E} \left( \sum_{H \in H(n-m)} \frac{N(r; F, H)}{|C_r|} \right) = \sum_{H \in H(n-m)} \frac{\mathcal{E}(N(r; F, H))}{|C_r|} \leq \sum_{H \in H(n-m)} \nu_{F,H}.$$

Let  $r \rightarrow \infty$ , the left hand side becomes  $\nu_F$  and we then finish the proof.  $\square$

**Remark 3.4.** One can also prove part (3) in Theorem 3.1 by analyzing Cheeger's finiteness theorem quantitatively like the proof of Theorem 4.2 in [SW19], which will give readers more understanding of the geometry of random zero sets.

Then, for Theorem 1.2, the proof needs the following Integral–Geometric Sandwich type inequality like Lemma 3.3, which was proved as Lemma 3.1 in [W21].

**Lemma 3.5.** *For each  $l = 0, 1, \dots, n-m$  and every  $r \in (0, R)$ , we have that*

$$(3.25) \quad \frac{1}{|C_r|} \int_{C_{R-r}} \beta_l(x, r; F) dx \leq \beta_l(R; F),$$

where  $\beta_l(x, r; F)$  is the sum of  $l$ -th Betti numbers over  $\mathbb{R}$  of all connected components of  $Z(F)$  fully contained in the open cube  $C_r(x)$ .

Combine Lemma 3.5 and Theorem 2.8, a same proof as the one for Theorem 3.1 will conclude Theorem 1.2. The reason that we do not have an almost sure convergence for Betti numbers is because we cannot similarly define a meaningful  $\mathfrak{N}_\#(r; F)$  and get a bound like  $\mathcal{E}(\mathfrak{N}_\#(r; F)) \leq C \cdot r^{n-1}$ .

## 4. PARAMETRIC ENSEMBLES AND PROOF OF THEOREM 1.3

In this section, we are going to handle the version of parametric ensembles similar to [NS16]. Let us first give some settings compatible with our random fields cases.

Let  $\{F_L\}_{L \in \mathcal{L}}$  be a family of Gaussian field ensemble defined on an open subset  $U \subset \mathbb{R}^n$ , where the index set  $L$  attains a discrete set  $\mathcal{L} \subset \mathbb{R}$ . We set the covariance matrix associated to  $F_L = (f_{L,1}, \dots, f_{L,m})$  as

$$(4.1) \quad (K_L)_{i_1 i_2}(x, y) = \mathcal{E}(f_{L,i_1}(x)f_{L,i_2}(y)), \quad i_1, i_2 = 1, 2, \dots, m,$$

and the scaled version for  $F_{x,L}(u) \equiv F_L(x + L^{-1}u)$  as

$$(4.2) \quad (K_{x,L})_{i_1 i_2}(u, v) = \mathcal{E}((F_{x,L})_{i_1}(u)(F_{x,L})_{i_2}(v)) = (K_L)_{i_1 i_2}(x + L^{-1}u, x + L^{-1}v).$$

**Definition 4.1.** We say that a family of Gaussian random fields  $\{F_L\}_{L \in \mathcal{L}}$  is locally uniformly controllable, if for every compact subset  $Q \subset U$ , there are positive constants  $M = M(Q), k_1 = k_1(Q)$  such that

(B1) For each  $x \in Q$ ,  $F_L(x)$  is centred and the joint distribution of  $(F_L(x), \nabla F_L(x))$  is a non-degenerate Gaussian vector with a rescaled uniformly lower bound on the covariance kernel, i.e.,

$$(4.3) \quad \liminf_{L \rightarrow \infty} \inf_{x \in Q} \inf_{(\eta, \xi) \in \mathbb{S}^{m+mn-1}} \mathcal{E} \left( \left| \sum_{i=1}^m \eta_i \cdot f_{L,i}(x) + L^{-1} \cdot \sum_{i=1}^m \sum_{j=1}^n \xi_{ij} \cdot \partial_j f_{L,i}(x) \right|^2 \right) \geq k_1 > 0,$$

where  $\eta = (\eta_i) \in \mathbb{R}^m$ ,  $\xi = (\xi_{ij}) \in \mathbb{R}^{nm}$  with  $\|\eta\|^2 + \|\xi\|^2 = 1$ .

(B2) We have the local uniform  $C^{3-}$ -smoothness:

$$(4.4) \quad \limsup_{L \rightarrow \infty} \max_{0 \leq |\alpha| \leq 3} \sup_{x \in Q} L^{-2|\alpha|} |D_x^\alpha D_y^\alpha (K_L)_{i_1 i_2}(x, y)|_{y=x} \leq M, \quad i_1, i_2 = 1, 2, \dots, m.$$

For an  $x \in U$ , if there exists a continuous stationary Gaussian field  $F_x : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , such that for every finite point set  $\mathcal{U} \subset \mathbb{R}^n$ , the Gaussian vectors  $(F_{x,L})|_{\mathcal{U}}$  converge to  $(F_x)|_{\mathcal{U}}$  in distribution, then we call  $F_x$  the translation invariant local limit of  $\{F_L\}_{L \in \mathcal{L}}$  at  $x$ . And hence,

$$(4.5) \quad \lim_{L \rightarrow \infty} K_{x,L}(u, v) = K_x(u - v) \quad \text{for any } (u, v) \in \mathbb{R}^n \times \mathbb{R}^n,$$

where  $K_x$  is the covariance kernel of  $F_x$ . On the other hand, since we have the uniform smoothness assumption (B2), the existence of such an  $F_x$  is equivalent to the existence of the limiting covariance kernel  $K_x$ . See more discussions in Appendix A.12 of [NS16].

**Definition 4.2.** For a family of locally uniformly controllable Gaussian fields  $\{F_L\}_{L \in \mathcal{L}}$  defined on  $U \subset \mathbb{R}^n$ , we say that  $\{F_L\}_{L \in \mathcal{L}}$  is tame if there exists a Borel subset  $U' \subset U$  of full Lebesgue measure such that, for all  $x \in U'$ ,

(B3)  $\{F_L\}_{L \in \mathcal{L}}$  has a translation invariant local limit  $F_x$  at  $x$ , which satisfies that the action of  $\mathbb{R}^n$  on  $(C_*^1(\mathbb{R}^n, \mathbb{R}^m), \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m)), \gamma_{F_x})$  is ergodic.

(B4) There is a  $k_2 = k_2(x)$ , and for every  $R > 0$ , there is an  $L_0 = L_0(R, x) > 0$ , such that when  $L > L_0$ ,  $F_{x,L}(u)$  satisfies (A3) in Definition 2.1, i.e., if  $x \in Q \cap U'$  for some compact set  $Q \subset U$ , then when  $L > L_0$ ,  $F_{x,L}(u)$  satisfies  $(R; M(Q), k_1(Q), k_2(x))$ -assumptions on  $C_{R+1}$ .

As mentioned in our introduction part, for an  $n$ -dimensional  $C^3$ -manifold  $X$  without boundary, we say that a parametric Gaussian ensemble  $\{F_L\}_{L \in \mathcal{L}}$  is tame if for every  $C^3$ -chart  $(U, \pi)$ ,  $\pi : U \subset \mathbb{R}^n \rightarrow X$ ,  $\{F_L \circ \pi\}_{L \in \mathcal{L}}$  is tame on  $U$ . And this definition does not depend on the choice of charts, which was discussed in section 9 of [NS16] when  $m = 1$ .

We then give an example, the Kostlan's ensemble, as a tame ensemble defined on a manifold when  $m = 1$ . For  $m > 1$ , one can take, for example,  $F_L = (f_{L,1}, \dots, f_{L,m})$  consisting of independent  $f_{L,i}$  with each  $f_{L,i}$  being independent Kostlan's ensemble (or more generally, different tame Gaussian functions ensembles) defined on the same manifold. For more examples, one can see [S16] and [NS16]. And apart from these concrete examples, in [CH20] and references therein, one will see a large family of Gaussian functions ensembles satisfying our definitions of tame ensembles and these definitions are actually very generic.

**Example 4.3** (Kostlan's ensemble). Consider the linear space of real homogeneous polynomials of degree  $d$  with coordinates  $(x_0, \dots, x_n) \in \mathbb{R}^{n+1}$ , which is endowed with the inner product

$$(4.6) \quad \langle P_d, Q_d \rangle = \sum_{|J|=d} \binom{d}{J}^{-1} p_J q_J,$$

where

$$(4.7) \quad J = (j_0, \dots, j_n), |J| = \sum_{s=0}^n j_s, \binom{d}{J} = \frac{d!}{j_0! \cdot \dots \cdot j_n!}$$

and

$$(4.8) \quad P_d(x) = \sum_{|J|=d} p_J x^J, \quad Q_d(x) = \sum_{|J|=d} q_J x^J, \quad x^J = x_0^{j_0} \cdot \dots \cdot x_n^{j_n}.$$

This inner product, up to a positive constant  $C = C(n, d)$ , equal to the inner product on the Bargmann–Fock space [B61], i.e., on the subspace of analytic functions on  $\mathbb{C}^{n+1}$  where the inner product

$$(4.9) \quad \langle f, g \rangle_{BK} \equiv C(n, d) \int_{\mathbb{C}^{n+1}} f(z) \overline{g(z)} e^{-\|z\|^2} d \text{Vol}(z)$$

is well-defined, one has that  $\langle P_d, Q_d \rangle = \langle P_d, Q_d \rangle_{BK}$  after the extension of  $P_d$  and  $Q_d$  to  $\mathbb{C}^{n+1}$ . Hence, we get an orthonormal basis

$$(4.10) \quad \left\{ \sqrt{\binom{d}{J}} x^J \right\}_{|J|=d}$$

on the linear space of real homogeneous polynomials of degree  $d$ , with respect to the inner product we defined above. We then randomize them by random  $P_d(x)$ , i.e.,

$$(4.11) \quad P_d(x) = \sum_{|J|=d} \sqrt{\binom{d}{J}} a_J x^J,$$

and  $\{a_J\}_{|J|=d}$  are i.i.d. standard Gaussian random variables. The zero sets of  $P_d(x)$  make sense as hypersurfaces on either  $\mathbb{S}^n$  or  $\mathbb{R}P^n$ , the real projective space of dimension  $n$ . View  $\{P_d(x)\}_{d \in \mathbb{N}}$  as a family of Gaussian random functions on  $\mathbb{S}^n$ , the two-point covariance kernel then is

$$(4.12) \quad \mathcal{E}(P_d(x)P_d(y)) = (\langle x, y \rangle_{\mathbb{R}^{n+1}})^d = (\cos(\theta(x, y)))^d,$$

where  $\langle x, y \rangle_{\mathbb{R}^{n+1}}$  is the standard inner product and  $\theta(x, y)$  is the angle between  $x, y$  as vectors in  $\mathbb{R}^{n+1}$ . Fix an  $x \in \mathbb{S}^n$ , let  $\exp_x : \mathbb{R}^n = T_x \mathbb{S}^n \rightarrow \mathbb{S}^n$  be the exponential map at  $x$ . We consider

$$(4.13) \quad P_{x,d}(u) \equiv P_d(\exp_x(d^{-1/2} \cdot u))$$

and then

$$(4.14) \quad K_{x,d}(u, v) \equiv \mathcal{E}(P_{x,d}(u)P_{x,d}(v)) = (\langle \exp_x(d^{-1/2} \cdot u), \exp_x(d^{-1/2} \cdot v) \rangle_{\mathbb{R}^{n+1}})^d.$$

As  $d \rightarrow \infty$ ,  $K_{x,d}(u, v)$ , together with its partial derivatives of any finite order, locally uniformly converges to a stationary Gaussian function  $R(u)$  defined on  $\mathbb{R}^n$ , whose associated covariance kernel is

$$(4.15) \quad K_R(u - v) = e^{-\|u-v\|^2/2}.$$

One can verify that this Kostlan's ensemble satisfies all of our definitions about tame ensembles. The way is that,  $K_R$  will satisfy all definitions for some universal constants only depending on  $n$ , and one use the uniform convergence from  $K_{x,d}$  to  $K_R$  to show that these  $K_{x,d}$  satisfy our definitions uniformly.

Now, We focus on  $\{F_L\}_{L \in \mathcal{L}}$  defined on an open subset  $U \subset \mathbb{R}^n$  again. Assume that  $\{F_L\}_{L \in \mathcal{L}}$  is tame and  $U'$  is the full measure set in Definition 4.2. For each  $x \in U'$ , we define  $\bar{\nu}(x) \equiv \nu_{F_x}$ , where we obtain each  $\nu_{F_x}$  from Theorem 3.1. We say that a Borel measure  $n_L$  is a connected components counting measure of  $F_L$  if  $\text{spt}(n_L) \subset Z(F_L)$  and the  $n_L$ -mass of each component is 1. Now, we can restate our Theorem 1.3 as the following.

**Theorem 4.4.** *Suppose that  $\{F_L\}_{L \in \mathcal{L}}$  is tame on an open set  $U \subset \mathbb{R}^n$ . Then,*

- (1) *The function  $x \mapsto \bar{\nu}(x)$  is measurable and locally bounded in  $U$ .*
- (2) *For every sequence of connected components counting measure  $n_L$  of  $F_L$  and for every  $\varphi \in C_c(U)$ , we have that*

$$(4.16) \quad \lim_{L \rightarrow \infty} \mathcal{E}^* \left[ \left| \frac{1}{L^n} \int_U \varphi(x) dn_L(x) - \int_U \varphi(x) \bar{\nu}(x) dx \right| \right] = 0.$$

Consider an  $x \in U'$  with the translation invariant limit  $F_x : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , which is centred and stationary. We see, for example by Appendix A.12 of [NS16], that  $F_x(u)$  satisfies (A1), (A2), and (A3) in Definition 2.1 at the origin  $u = 0$  for the same  $M(Q)$  and  $k_1(Q)$  if  $x \in Q$  for a compact set  $Q \subset U$ , and for  $k_2(x)$  in (B4) of Definition 4.2. We then have the local double limit lemma since  $F_x$  also satisfies the ergodicity assumption (B3) of Definition 4.2.

**Lemma 4.5.** *Suppose that  $\{F_L\}_{L \in \mathcal{L}}$  is locally uniformly controllable and  $F_x$  is its translation invariant local limit at a fixed  $x \in U$ . Also suppose that for  $F_x$ , the action of  $\mathbb{R}^n$  on  $(C_*^1(\mathbb{R}^n, \mathbb{R}^m), \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m)), \gamma_{F_x})$  is ergodic. Then, for any  $\epsilon > 0$ ,*

$$(4.17) \quad \lim_{R \rightarrow \infty} \limsup_{L \rightarrow \infty} \mathcal{P} \left( \left| \frac{N(R; F_{x,L})}{|C_R|} - \nu_{F_x} \right| > \epsilon \right) = 0.$$

The proof of this lemma is exactly the same as Lemma 12 in [NS16], Theorem 5 in [S16], and Proposition 4.1 in [W21]. But in the proof, one needs to replace some lemmas for random functions to those for random fields, which are our Lemma A.1 and Lemma B.1. After this replacement, one will find the proof and so we omit this proof.

**Remark 4.6.** For each  $l = 0, 1, \dots, n - m$ , Lemma 4.5 holds true if we replace  $N(R; F_{x,L})$  by  $\beta_l(R; F_{x,L})$  and  $\nu_{F_x}$  by  $\nu_{l;F_x}$ . And for any  $H \in H(n - m)$ , Lemma 4.5 also holds true if one replaces  $N(R; F_{x,L})$  by  $N(R; F_{x,L}, H)$  and replaces  $\nu_{F_x}$  by  $\nu_{F_x, H}$ . The proofs for both then need the  $C^1$ -stability result as in Remark B.2, one can see, for example Theorem 1.5 of [W21] or Theorem 6.2 of [SW19], which are the corresponding cases when  $m = 1$ .

So far, the general settings like Definition 2.1 and Definition 4.1 are similar to those in [NS16]. But remember that we also made (A3) and (B4) assumptions which do not play roles yet. These are key assumptions we need to make in order to prove global type theorems similar to Theorem 2 in [NS16] or Theorem 4 in [S16], which are our Theorem 1.3, Theorem 1.4, and Theorem 1.8. First, we can get a stronger theorem than Theorem 2.3 with the help of assumption (A3) in Definition 2.1.

**Theorem 4.7.** *Suppose that  $F : C_{R+1} \rightarrow \mathbb{R}^m$ , satisfies  $(R; M, k_1, k_2)$ -assumptions on  $C_{R+1}$ , then there is a constant  $D_2 = D_2(n, m, M, k_1, k_2) > 0$  and a constant  $q = q(n, m) > 1$ , such that for  $R > 1$ ,*

$$(4.18) \quad \mathcal{E}((N(R; F))^q) \leq D_2 \cdot |C_R|^q.$$

Before we give the proof, let us remark that the same result holds for Betti numbers.

**Theorem 4.8.** *Suppose that  $F : C_{R+1} \rightarrow \mathbb{R}^m$ , satisfies  $(R; M, k_1, k_2)$ -assumptions on  $C_{R+1}$ , then there is a constant  $\tilde{D}_2 = \tilde{D}_2(n, m, M, k_1, k_2) > 0$  and a constant  $q = q(n, m) > 1$ , such that for  $R > 1$ ,*

$$(4.19) \quad \sum_{l=0}^{n-m} \mathcal{E}((\beta_l(R; F))^q) \leq \tilde{D}_2 \cdot |C_R|^q.$$



The necessary modification is again using Chern–Lashof inequality in Theorem 2.9. Hence, we only give the proof for Theorem 4.7.

*Proof of Theorem 4.7.* For some  $q = q(n, m) > 1$  but close to 1 and a positive integer  $k$  to be determined later, by (2.9), we have that for some positive constant  $C = C(n, m)$ ,

$$\begin{aligned}
(4.20) \quad (N(R; F))^q &\leq C(n, m) \cdot \left( \int_{F^{-1}(0) \cap C_R} \|\mathbf{H}\|^{n-m} d\mathcal{H}^{n-m} \right)^q \\
&\leq C \cdot \left( \int_{F^{-1}(0) \cap C_R} \|\mathbf{H}\|^{(n-m)q} d\mathcal{H}^{n-m} \right) \cdot (\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R))^{q-1} \\
&= C \cdot \left( \int_{F^{-1}(0) \cap C_R} \|\mathbf{H}\|^{(n-m)q} d\mathcal{H}^{n-m} \right) \cdot \left( \frac{\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R)}{|C_R|} \right)^{q-1} \cdot |C_R|^{q-1} \\
&= C \cdot |C_R|^{q-1} \int_{F^{-1}(0) \cap C_R} (\|\mathbf{H}\|^{\frac{(n-m)q(k+1)}{k}})^{\frac{k}{k+1}} \left( \left( \frac{\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R)}{|C_R|} \right)^{(k+1)(q-1)} \right)^{\frac{1}{k+1}} d\mathcal{H}^{n-m} \\
&\leq \frac{C}{k+1} |C_R|^{q-1} \int_{F^{-1}(0) \cap C_R} k (\|\mathbf{H}\|^{\frac{(n-m)q(k+1)}{k}}) + \left( \left( \frac{\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R)}{|C_R|} \right)^{(k+1)(q-1)} \right) d\mathcal{H}^{n-m} \\
&\leq C \cdot |C_R|^{q-1} \int_{F^{-1}(0) \cap C_R} (\|\mathbf{H}\|^{\frac{(n-m)q(k+1)}{k}}) d\mathcal{H}^{n-m} \\
&\quad + C \cdot |C_R|^{-k(q-1)} (\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R))^{1+(k+1)(q-1)}.
\end{aligned}$$

We set  $k = \frac{2-q}{q-1}$  and  $q = q(m, n)$  to be determined later, then we derive from the above that

$$(4.21) \quad (N(R; F))^q \leq C \cdot |C_R|^{q-1} \int_{F^{-1}(0) \cap C_R} (\|\mathbf{H}\|^{\frac{(n-m)q}{2-q}}) d\mathcal{H}^{n-m} + C \cdot |C_R|^{q-2} (\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R))^2.$$

Notice that when  $q$  is greater than 1 and close to 1,  $q/(2-q)$  is also greater than 1 and close to 1. To simplify the notation, we let  $q/(2-q) = \theta > 1$ .

In order to estimate  $\mathcal{E}((N(R; F))^q)$ , by (4.21), we need to estimate

$$(4.22) \quad \mathcal{E} \left( \int_{F^{-1}(0) \cap C_R} (\|\mathbf{H}\|^{(n-m)\theta}) d\mathcal{H}^{n-m} \right)$$

and

$$(4.23) \quad \mathcal{E} \left( (\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R))^2 \right).$$

Since  $\theta$  is close to 1, the estimate for (4.22) is exactly the same as Theorem 2.3 and in particular the same as estimating (2.16). The  $\theta$  is then allowed to be greater than 1 but close to 1 because of the sharp constant  $\alpha$  shown in Lemma 2.6. So we can choose a  $\theta = (n, m) > 1$

and hence  $q = q(n, m) > 1$  for which  $k + 1 = \frac{1}{q-1}$  is a positive integer. And we also get that (4.22) is bounded by some  $C = C(n, m, M, k_1) > 0$ .

For estimating (4.23), by the Kac–Rice Theorem 2.4, we have that

$$(4.24) \quad \mathcal{E} \left( (\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R))^2 \right) = \int_{C_R \times C_R} J_{2,F}(1; u_1, u_2) p_{(F(u_1), F(u_2))}(0) du_1 du_2,$$

where  $J_{2,F}(h; u_1, u_2)$  has the expression

$$(4.25) \quad J_{2,F} = \mathcal{E} \left[ \sqrt{\det[(\nabla F(u_1))^T (\nabla F(u_1))]} \sqrt{\det[(\nabla F(u_2))^T (\nabla F(u_2))]} \Big| F(u_1) = F(u_2) = 0 \right].$$

We have an estimate for  $p_{(F(u_1), F(u_2))}(0)$  from the (A3) assumption in Definition 2.1. For this  $J_{2,F}$  term, we prove that it has a positive upper bound  $C = C(n, m, M, k_1, k_2) < \infty$ . By the inequality  $2ab \leq a^2 + b^2$  and symmetry, we only need to estimate

$$(4.26) \quad \mathcal{E} \left[ \det[(\nabla F(u_1))^T (\nabla F(u_1))] \mid F(u_1) = F(u_2) = 0 \right],$$

for  $u_1, u_2 \in C_R$  and  $u_1 \neq u_2$ . One can just use Proposition 7 or Lemma 9 of [CH20] since by Theorem E.1, the assumptions there are satisfied. For completeness, let us include this part of proof in consistent with our current settings. By similar tricks in (2.18), fix  $i \in \{1, \dots, m\}$  and  $j \in \{1, \dots, n\}$ , we need to first find  $a_{j,i}^t$  and  $b_{j,i}^t$  such that for each  $s = 1, \dots, m$ ,

$$(4.27) \quad \mathcal{E} \left[ (\partial_j f_i(u_1) + \sum_{t=1}^m a_{j,i}^t f_t(u_1) + b_{j,i}^t f_t(u_2)) \cdot f_s(u_1) \right] = 0,$$

and

$$(4.28) \quad \mathcal{E} \left[ (\partial_j f_i(u_1) + \sum_{t=1}^m a_{j,i}^t f_t(u_1) + b_{j,i}^t f_t(u_2)) \cdot f_s(u_2) \right] = 0.$$

To simplify the notations, we let  $v = \partial_j f_i(u_1)$ ,  $A = (a_{j,i}^1, \dots, a_{j,i}^m)$ ,  $B = (b_{j,i}^1, \dots, b_{j,i}^m)$ ,  $\mathcal{E}(F(u_1)v) = (\mathcal{E}(f_1(u_1)v), \dots, \mathcal{E}(f_m(u_1)v))$ ,  $\mathcal{E}(F(u_2)v) = (\mathcal{E}(f_1(u_2)v), \dots, \mathcal{E}(f_m(u_2)v))$ . Set  $\text{Cov}(x, y)$  as the  $m \times m$  matrix with elements  $(\text{Cov}(x, y))_{i_1 i_2} = \mathcal{E}(f_{i_1}(x) f_{i_2}(y))$  and  $\Lambda(x, y)$  as the covariance kernel of the joint distribution  $(F(x), F(y))$ , i.e.,

$$(4.29) \quad \Lambda(x, y) = \begin{bmatrix} \text{Cov}(x, x) & \text{Cov}(x, y) \\ \text{Cov}(y, x) & \text{Cov}(y, y) \end{bmatrix}.$$

$\Lambda(x, y)$  is invertible when  $x \neq y$  by the (A3) assumption. And hence, the conditions (4.27) and (4.28) are then solvable and we get

$$(4.30) \quad (A, B) = -(\mathcal{E}(F(u_1)v), \mathcal{E}(F(u_2)v)) \cdot (\Lambda(u_1, u_2))^{-1}.$$

Notice that

$$(4.31) \quad \det[(\nabla F(u_1))^T (\nabla F(u_1))] \leq C(n, m) \cdot \|\nabla F\|^{2m},$$

we see that the upper bound for (4.26) reduces to the upper bound for

$$(4.32) \quad \mathcal{E}(v + A \cdot F(u_1) + B \cdot F(u_2))^{2m}.$$

Since it is the  $2m$ -th moment of a centred Gaussian variable, we only need to estimate its variance. By (4.30), we have that

$$(4.33) \quad \mathcal{E}(A \cdot F(u_1) + B \cdot F(u_2))^2 = (\mathcal{E}(F(u_1)v), \mathcal{E}(F(u_2)v)) \cdot (\Lambda(u_1, u_2))^{-1} \cdot (\mathcal{E}(F(u_1)v), \mathcal{E}(F(u_2)v))^T.$$

We will use Theorem E.1 and consider the cases  $|u_1 - u_2| \leq \delta$  and  $|u_1 - u_2| > \delta$  separately, where  $\delta$  is chosen in Theorem E.1. When  $|u_1 - u_2| \leq \delta$ , by elementary row operations,

$$(4.34) \quad \Lambda(u_1, u_2) = \begin{bmatrix} \text{Id}_m & 0 \\ \text{Cov}(u_2, u_1)\text{Cov}(u_1, u_1)^{-1} & \text{Id}_m \end{bmatrix} \cdot \begin{bmatrix} \text{Cov}(u_1, u_1) & 0 \\ 0 & \text{Cov}(u_2, u_2) - \text{Cov}(u_2, u_1)\text{Cov}(u_1, u_1)^{-1}\text{Cov}(u_1, u_2) \end{bmatrix} \cdot \begin{bmatrix} \text{Id}_m & \text{Cov}(u_1, u_1)^{-1}\text{Cov}(u_1, u_2) \\ 0 & \text{Id}_m \end{bmatrix}.$$

Hence,

$$(4.35) \quad \begin{aligned} \mathcal{E}(AF(u_1) + BF(u_2))^2 &= (\mathcal{E}(F(u_1)v), \mathcal{E}(F(u_2)v) - \mathcal{E}(F(u_1)v)\text{Cov}(u_1, u_1)^{-1}\text{Cov}(u_1, u_2)) \\ &\cdot \begin{bmatrix} \text{Cov}(u_1, u_1) & 0 \\ 0 & \text{Cov}(u_2, u_2) - \text{Cov}(u_2, u_1)\text{Cov}(u_1, u_1)^{-1}\text{Cov}(u_1, u_2) \end{bmatrix}^{-1} \\ &\cdot (\mathcal{E}(F(u_1)v), \mathcal{E}(F(u_2)v) - \mathcal{E}(F(u_1)v)\text{Cov}(u_1, u_1)^{-1}\text{Cov}(u_1, u_2))^T \end{aligned}$$

The first  $m \times m$  block is good to be bounded since by (A1) of Definition 2.1,  $\text{Cov}(u_1, u_1)$  is quantitatively non-degenerate. For the second  $m \times m$  block, by Theorem E.1, the smallest eigenvalue of  $\text{Cov}(u_2, u_2) - \text{Cov}(u_2, u_1)\text{Cov}(u_1, u_1)^{-1}\text{Cov}(u_1, u_2)$  is lowerly bounded by  $(k_1/2) \cdot |u_1 - u_2|^2$ , while by mean value theorem,

$$(4.36) \quad |\mathcal{E}(F(u_2)v) - \mathcal{E}(F(u_1)v)\text{Cov}(u_1, u_1)^{-1}\text{Cov}(u_1, u_2)|^2 \leq C(n, m, M) \cdot |u_1 - u_2|^2.$$

Hence, when  $|u_1 - u_2| \leq \delta$ ,

$$(4.37) \quad \mathcal{E}(AF(u_1) + BF(u_2))^2 \leq C(n, m, M, k_1).$$

In the case that  $|u_1 - u_2| > \delta$ , by (A3) of Definition 2.1, there is a positive constant  $c = c(n, m, M, k_1, k_2)$  such that

$$(4.38) \quad \det(\Lambda(u_1, u_2)) \geq c.$$

Since the norm of  $(\mathcal{E}(F(u_1)v), \mathcal{E}(F(u_2)v))$  is bounded by some positive  $C = C(n, m, M)$ , we get that

$$(4.39) \quad \mathcal{E}(AF(u_1) + BF(u_2))^2 \leq C(n, m, M, k_1, k_2).$$

Combine with the fact that  $\mathcal{E}(v^2) \leq C(n, m, M)$ , we can then prove that

$$(4.40) \quad \mathcal{E}(v + A \cdot F(u_1) + B \cdot F(u_2))^2 \leq C(n, m, M, k_1, k_2).$$

And hence, by the previous arguments,

$$(4.41) \quad J_{2,F} \leq C(n, m, M, k_1, k_2).$$

Combine this with (A4) in Definition 2.1, i.e.

$$(4.42) \quad p_{(F(u_1), F(u_2))}(0) \leq \begin{cases} \frac{k_2}{|u_1 - u_2|^m}, & \text{if } |u_1 - u_2| \leq 1, \\ k_2, & \text{if } |u_1 - u_2| > 1, \end{cases}$$

one can get that, for  $R > 1$ ,

$$(4.43) \quad \begin{aligned} \mathcal{E}\left(\left(\mathcal{H}^{n-m}(F^{-1}(0) \cap C_R)\right)^2\right) &\leq C(n, m, M, k_1, k_2) \cdot R^n \cdot \left(\int_0^1 r^{n-1-m} dr + \int_1^{2R} r^{n-1} dr\right) \\ &\leq C(n, m, M, k_1, k_2) \cdot R^{2n}. \end{aligned}$$

Now, combine estimates for (4.22) and (4.23), we can estimate  $\mathcal{E}((N(R; F))^q)$  for some  $q = q(n, m) > 1$  but close to 1 by (4.21), and get that for all  $R > 1$ ,

$$(4.44) \quad \mathcal{E}((N(R; F))^q) \leq C(n, m, M, k_1, k_2) \cdot R^q.$$

□

In Theorem 4.7, we need to impose an additional condition (A3) to insure the nondegeneracy of  $F$  at two different points, and then we can get  $L^q$  bounds for the number and Betti numbers of connected components. In order to get a uniform control on the family  $\{F_L\}_{L \in \mathcal{L}}$ , we need to impose the tame conditions in Definition 4.2, with which we can get the proof for our Theorem 4.4. Notice that, if  $\{F_L\}_{L \in \mathcal{L}}$  satisfies (B4) for some  $k_2(x)$  at  $x \in U$  in Definition 4.2, then its translation invariant limit  $F_x$  also satisfies (A3) in Definition 2.1 for  $k_2(x)$  and any  $R > 1$ .

*Proof of (1) in Theorem 4.4.* Assume that  $x \in Q \cap U'$  for some compact subset  $Q \subset U$ , then  $F_x$  satisfies  $(R; M(Q), k_1(Q), k_2(x))$ -assumptions on  $C_{R+1}$  for each  $R > 0$ . By Theorem 2.3,

$$(4.45) \quad \bar{\nu}(x) = \lim_{R \rightarrow \infty} \frac{\mathcal{E}(N(R; F))}{|C_R|} \leq D_1(n, m, M(Q), k_1(Q)).$$

The measurability is directly if one consider the function

$$(4.46) \quad \nu_{R,L}(x, w) \equiv \frac{N(R; F_{x,L})}{|C_R|},$$

defined on  $U_{-(R+1)/L} \times \Omega'$  for  $U_{-r} \equiv \{z \in U \mid \text{dist}(z, \partial U) > r\}$  and  $\Omega' = \{w \in \Omega \mid F_L \in C_*^1(\mathbb{R}^n, \mathbb{R}^m) \text{ for all } L\}$ . Notice that  $\mathcal{P}(\Omega \setminus \Omega') = 0$  by Bulinskaya's lemma. The measurability

of  $\nu_{R,L}(x, w)$  follows from the measurability of compositions of lower semicontinuous and measurable maps.

We then claim that for any fixed  $x \in Q \cap U'$ ,

$$(4.47) \quad \lim_{R \rightarrow \infty} \limsup_{L \rightarrow \infty} \mathcal{E}(|\nu_{R,L}(x, \cdot) - \bar{\nu}(x)|) = 0,$$

which is a stronger version than Lemma 4.5. And the measurability of  $\bar{\nu}(x)$  follows from the Fubini–Tonelli’s theorem. First, we notice that for each  $x \in Q \cap U'$  and each  $R > 0$ , when  $L > L_0(R, x)$ , Theorem 4.7 gives that

$$(4.48) \quad \mathcal{E}((\nu_{R,L}(x, \cdot))^q) \leq D_2$$

for some  $q = q(n, m)$  and  $D_2 = D_2(n, m, M(Q), k_1(Q), k_2(x))$ . On the other hand, given any  $\epsilon > 0$ , Lemma 4.5 gives that

$$(4.49) \quad \lim_{R \rightarrow \infty} \limsup_{L \rightarrow \infty} \mathcal{P}(\Omega_\epsilon) = 0,$$

where  $\Omega_\epsilon \equiv \{w \in \Omega' \mid |\nu_{R,L}(x, w) - \bar{\nu}(x)| > \epsilon\}$ . Hence, when  $L > L_0(R, x)$ , by Hölder inequality,

$$(4.50) \quad 0 \leq \mathcal{E}(|\nu_{R,L}(x, \cdot) - \bar{\nu}(x)|) \leq \epsilon + (\mathcal{P}(\Omega_\epsilon))^{1-\frac{1}{q}} \cdot C(n, m, D_1, D_2).$$

Since  $D_1, D_2$  are independent of  $R, L$ , we take  $L \rightarrow \infty$  first, then  $R \rightarrow \infty$ , and finally let  $\epsilon \rightarrow 0$ , we finish the proof of the claim.  $\square$

*Proof of (2) in Theorem 4.4.* The proof is exactly the same as Section 8 in [NS16]. For completeness, we include this adapting to our settings here. We can just assume that  $\varphi \geq 0$  and set  $Q = \text{spt}(\varphi)$ . Fix an arbitrary  $\delta \in (0, 1)$  such that  $Q_{+4\delta} \subset U$  and we denote  $Q_1 = Q_{+\delta}$ ,  $Q_2 = Q_{+2\delta}$ , where  $Q_{+\delta} \equiv \{z \in \mathbb{R}^n \mid \text{dist}(z, Q) \leq \delta\}$ . For any  $x \in Q_1$ , let  $\varphi_-(x) \equiv \inf_{C_\delta(x)} \varphi$  and  $\varphi_+(x) \equiv \sup_{C_\delta(x)} \varphi$ , where we recall that  $C_\delta(x)$  is the shift from the open cube  $C_\delta(0)$  to the center  $x$ . Then, for parameters  $T, R, L$ , with  $1 < T < R < \delta L$ ,

$$(4.51) \quad \begin{aligned} \int_{Q_1} \varphi_-(x) \nu_{R,L}(x, w) dx &\leq \int_{Q_1} \varphi_-(x) \frac{n_L(C_{R/L}(x))}{|C_R|} dx \\ &\leq \int_{Q_1} \int_{C_{R/L}(x)} \frac{\varphi(y)}{|C_R|} dn_L(y) dx = \frac{1}{L^n} \int_U \varphi(y) dn_L(y) \\ &\leq \int_{Q_1} \varphi_+(x) \frac{n_L(C_{R/L}(x))}{|C_R|} dx. \end{aligned}$$

Then, we cover  $Q_2$  by  $C(n) \cdot |Q_2| \cdot (L/T)^n$  open cubes with side lengths  $2 \cdot (T/L)$ , which we denote as  $\{C_j\}$ . For those components of  $Z(F_L) \cap C_{R/L}(x)$  that does not intersect with the boundaries of any  $C_j$ , each of them is fully contained in one  $C_j$ , and hence is fully contained in  $C_{(R+T)/L}(x)$ . For those components of  $Z(F_L)$  that intersect with at least one  $C_j$ , the number of them are bounded by  $\sum_j \mathfrak{N}_\#(C_j; F_L)$ . Remember, from Bulinskaya’s lemma and

the discussions after the definition of  $\mathfrak{N}_\#(R; G)$  in (3.5), for each  $j$ , it is in probability 1 that  $\mathfrak{N}_\#(C_j; F_L)$  is finite and  $\mathcal{E}(\mathfrak{N}_\#(C_j; F_L)) \leq C(n, m, M(Q), k_1(Q)) \cdot (T)^{n-1}$ . Hence,

$$(4.52) \quad \int_{Q_1} \varphi_+(x) \frac{n_L(C_{R/L}(x))}{|C_R|} dx \leq \left( \frac{R+T}{R} \right)^n \cdot \int_{Q_1} \varphi_+(x) \nu_{R+T,L}(x, w) dx \\ + (\sup_U \varphi) \cdot L^{-n} \cdot \left( \sum_j \mathfrak{N}_\#(C_j; F_L) \right).$$

For the  $\int_U \varphi(x) \bar{\nu}(x) dx$  term, set  $\omega_\varphi(\cdot)$  as the modulus of continuity of  $\varphi$ ,

$$(4.53) \quad \int_U \varphi(x) \bar{\nu}(x) dx \geq \left( \frac{R+T}{R} \right)^n \cdot \int_{Q_1} \varphi_+(x) \bar{\nu}(x) dx - C(n) \cdot (T/R) \cdot (\sup_U \varphi) \cdot \|\bar{\nu}\|_{L^\infty(Q_1)} \cdot |Q_1| \\ - \omega_\varphi(\delta) \cdot \|\bar{\nu}\|_{L^\infty(Q_1)} \cdot |Q_1|.$$

Combine with (4.51) and (4.52),

$$(4.54) \quad \frac{1}{L^n} \int_U \varphi(y) dn_L(y) - \int_U \varphi(x) \bar{\nu}(x) dx \leq \left( \frac{R+T}{R} \right)^n \cdot (\sup_U \varphi) \cdot \int_{Q_1} |\nu_{R+T,L}(x, w) - \bar{\nu}(x)| dx \\ + C(n) \cdot (T/R) \cdot (\sup_U \varphi) \cdot \|\bar{\nu}\|_{L^\infty(Q_1)} \cdot |Q_1| + \omega_\varphi(\delta) \cdot \|\bar{\nu}\|_{L^\infty(Q_1)} \cdot |Q_1| \\ + (\sup_U \varphi) \cdot L^{-n} \cdot \left( \sum_j \mathfrak{N}_\#(C_j; F_L) \right).$$

and

$$(4.55) \quad \frac{1}{L^n} \int_U \varphi(y) dn_L(y) - \int_U \varphi(x) \bar{\nu}(x) dx \geq -(\sup_U \varphi) \cdot \int_{Q_1} |\nu_{R,L}(x, w) - \bar{\nu}(x)| dx \\ - \omega_\varphi(\delta) \cdot \|\bar{\nu}\|_{L^\infty(Q_1)} \cdot |Q_1|.$$

Hence,

$$(4.56) \quad \mathcal{E}^* \left[ \left| \frac{1}{L^n} \int_U \varphi(x) dn_L(x) - \int_U \varphi(x) \bar{\nu}(x) dx \right| \right] \leq 2^n \cdot (\sup_U \varphi) \cdot \int_{Q_1} \mathcal{E}(|\nu_{R+T,L}(x, \cdot) - \bar{\nu}(x)|) dx \\ + (\sup_U \varphi) \cdot \int_{Q_1} \mathcal{E}(|\nu_{R,L}(x, w) - \bar{\nu}(x)|) dx + (\sup_U \varphi) \cdot L^{-n} \cdot \left( \sum_j \mathcal{E}(\mathfrak{N}_\#(C_j; F_L)) \right) \\ + C(n) \cdot (T/R) \cdot (\sup_U \varphi) \cdot \|\bar{\nu}\|_{L^\infty(Q_1)} \cdot |Q_1| + \omega_\varphi(\delta) \cdot \|\bar{\nu}\|_{L^\infty(Q_1)} \cdot |Q_1|.$$

Notice that, by (B1) and (B2) in Definition 4.1, for fixed  $R > T > 1$ , there is a uniform  $L_1 = L_1(R) > 0$ , such that when  $L > L_1$ ,

$$(4.57) \quad \mathcal{E}(|\nu_{R+T,L}(x, w) - \bar{\nu}(x)|) \leq D_1(n, m, M(Q), k_1(Q))$$

for every  $x \in Q_1$  by Theorem 2.3. And hence, the function

$$(4.58) \quad \eta_R(x) \equiv \limsup_{L \rightarrow \infty} \mathcal{E}(|\nu_{R,L}(x, w) - \bar{\nu}(x)|)$$

has the same upper bound  $D_1$ . Recall that the claim (4.47) in proving part (1) of Theorem 4.4 shows that  $\lim_{R \rightarrow \infty} \eta_R(x) = 0$ . Hence, apply  $\limsup_{L \rightarrow \infty}$  first and then  $\lim_{R \rightarrow \infty}$ , the first two terms in (4.56) go to 0 by dominated convergence theorem. The fourth term will also go to 0 since  $T < R$  is still fixed at this step. As previously mentioned, since  $\{F_L\}_{L \in \mathcal{L}}$  satisfies (B1) and (B2) in Definition 4.1, when  $L > L_1(R)$ , for each  $j$ ,  $\mathcal{E}(\mathfrak{N}_\#(C_j; F_L)) \leq C(n, m, M(Q), k_1(Q)) \cdot (T)^{n-1}$ . And hence, the third term is bounded by  $(\sup_U \varphi) \cdot C(n, m, M(Q), k_1(Q)) \cdot T^{-1}$  for any fixed  $T < R < L$ . We now let  $T \rightarrow \infty$  and finally let  $\delta \rightarrow \infty$  to see that the right hand side of (4.56) goes to 0. We finish the proof.  $\square$

The proof for Theorem 1.4 is the same as the proof of Theorem 3 in [NS16] after we finished Theorem 4.4 as above. So we omit the details.

Instead, let us explore more on the Betti numbers behaviors of tame parametric ensembles  $\{F_L\}_{L \in \mathcal{L}}$  defined on an  $n$ -dimensional closed  $C^3$ -manifold  $X$ . Recall that we say that a parametric Gaussian ensemble  $\{F_L\}_{L \in \mathcal{L}}$  is tame if for every  $C^3$ -chart  $\pi : U \subset \mathbb{R}^n \rightarrow X$ ,  $\{F_L \circ \pi\}_{L \in \mathcal{L}}$  is tame on  $U$  as in the Definition 4.2.

For each  $l = 0, \dots, n - m$ , we use  $\beta_l(F_L)$  to denote the summation of  $l$ -th Betti numbers over  $\mathbb{R}$  for all connected components of  $Z(F_L)$  in  $X$ . For each  $x \in X$ , let  $F_x$  be the translation invariant local limit of  $\{F_L\}$  at  $x$ . And by Theorem 1.2, we get a limiting constant  $\nu_{l, F_x}$ , and we then define a function on  $X$  by  $\bar{\nu}_l(x) = \nu_{l, F_x}$ . The following theorem also holds if  $X$  is noncompact and without boundary, but one needs to modify the definitions of  $\beta_l(F_L)$  by summation of Betti numbers for those connected components fully contained in a geodesic ball of finite radius in  $X$ .

**Theorem 4.9.** *Assume that  $\{F_L\}_{L \in \mathcal{L}}$  is tame on  $X$  and  $X$  is closed. Then for each  $l = 0, \dots, n - m$ , the function  $x \mapsto \bar{\nu}_l(x)$  is measurable and bounded on  $X$ , and there is a positive constant  $C_l = C_l(n, m, M, k_1)$ , such that*

$$(4.59) \quad \int_X \bar{\nu}_l(x) d\text{Vol}(x) \leq \liminf_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(F_L))}{L^n} \leq \limsup_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(F_L))}{L^n} \leq C_l \cdot |X|,$$

where  $|X|$  is the volume of  $X$ , and the two positive constants  $M = M(X)$  and  $k_1 = k_1(X)$  are chosen in the Definition 4.1 since  $X$  is closed.

*Proof of the lower bound in Theorem 4.9.* The proof is quite similar to Theorem 4.4. For the measurability and local boundedness of  $\bar{\nu}_l(x)$ , the proof is exactly the same as part (1) of Theorem 4.4 because one just needs to verify these properties on local charts. Since  $X$  is closed,  $\bar{\nu}_l$  is actually bounded on  $X$ . And one thing we can similarly get is the following. Choose a local chart  $\pi : U \rightarrow X$  for some bounded open set  $U \subset \mathbb{R}^n$ . For  $x \in \pi(U)$  and  $y = \pi^{-1}(x)$ , we define a random function

$$(4.60) \quad \nu_{l;R,L}(y, w) \equiv \frac{\beta_l(R; (F \circ \pi)_{y,L})}{|C_R|}.$$

Here  $(F \circ \pi)_{y,L}$  is the pull-back of  $F_L$  from  $\pi(U)$  to  $U$  and rescaled, i.e.,

$$(4.61) \quad (F \circ \pi)_{y,L}(u) \equiv F_L(\pi(y + L^{-1}u)),$$

for  $u \in \mathbb{R}^n$  with  $y + L^{-1}u \in U$ . Hence,  $\beta_l(R; (F \circ \pi)_{y,L}) = \beta_l(C_{R/L}(y); F_L \circ \pi)$ .

Then, using Theorem 4.8, we can similarly get, as in the claim (4.47) in proving part (1) of Theorem 4.4, that

$$(4.62) \quad \lim_{R \rightarrow \infty} \limsup_{L \rightarrow \infty} \mathcal{E}(|\nu_{l;R,L}(y, \cdot) - \bar{\nu}_l(x) \sqrt{\det g_y}|) = 0.$$

Here  $g_y$  is the metric tensor of  $X$  at  $x = \pi(y)$ , which is to be compatible with change of coordinates to add this determinant term here, because we used  $|C_R|$  in standard  $\mathbb{R}^n$  volume to define  $\nu_{l;R,L}(y, w)$ . For more discussions on change of coordinates, see Section 9 of [NS16].

Recall that  $\beta_l(R; (F \circ \pi)_{y,L})$  only counts those connected components fully contained in the cube  $C_R$ . We set parameters  $R, L$  with  $1 < R < \delta L$  for some  $\delta \in (0, 1)$  and let  $Q \subset U$  be a compact subset such that  $Q_{+\delta} \subset U$  but  $Q_{+2\delta} \not\subset U$ . We have that

$$(4.63) \quad \begin{aligned} \int_Q \nu_{l;R,L}(y, w) dy &= \int_Q \frac{\beta_l(C_{R/L}(y); F_L \circ \pi)}{|C_R|} dy = \frac{1}{|C_R|} \int_Q \sum_{\gamma \subset Z(F_L \circ \pi)} \mathbf{1}_{\gamma \subset C_{R/L}(y)} \cdot \beta_l(\gamma) dy \\ &\leq \frac{1}{|C_R|} \sum_{\gamma \subset Z(F_L \circ \pi) \cap U} \beta_l(\gamma) \int_Q \mathbf{1}_{\gamma \subset C_{R/L}(y)} dy \\ &\leq \frac{|C_{R/L}|}{|C_R|} \sum_{\gamma \subset Z(F_L \circ \pi) \cap U} \beta_l(\gamma) = \frac{\beta_l(U; F_L \circ \pi)}{L^n} = \frac{\beta_l(\pi(U); F_L)}{L^n}. \end{aligned}$$

Here, we use  $\beta_l(\pi(U); F_L)$  to denote the summation of  $l$ -th Betti numbers for all connected components of  $Z(F_L)$  fully contained in the open set  $\pi(U)$ . The last inequality is because

$$(4.64) \quad \int_Q \mathbf{1}_{\gamma \subset C_{R/L}(y)} dy = |\{y \in Q \mid y \in \cap_{z \in \gamma} C_{R/L}(z)\}| \leq |C_{R/L}|.$$

And hence,

$$(4.65) \quad \begin{aligned} \int_Q \bar{\nu}_l(\pi(y)) \sqrt{\det g_y} dy &\leq \int_Q \mathcal{E}(\nu_{l;R,L}(y, w)) dy + \int_Q \mathcal{E}(|\nu_{l;R,L}(y, \cdot) - \bar{\nu}_l(x) \sqrt{\det g_y}|) dy \\ &\leq \frac{\mathcal{E}(\beta_l(\pi(U); F_L))}{L^n} + \int_Q \mathcal{E}(|\nu_{l;R,L}(y, \cdot) - \bar{\nu}_l(x) \sqrt{\det g_y}|) dy. \end{aligned}$$

Use (4.62), and take limits  $\lim_{R \rightarrow \infty} \liminf_{L \rightarrow \infty}$  at both sides, and apply dominated convergence theorem on the second term of the right hand side, which is dominated by the constant in Theorem 2.8, we then get that

$$(4.66) \quad \int_{\pi(Q)} \bar{\nu}_l(x) d\text{Vol}(x) = \int_Q \bar{\nu}_l(\pi(y)) \sqrt{\det g_y} dy \leq \liminf_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(\pi(U); F_L))}{L^n}.$$



Since  $\delta$  and  $Q$  were chosen arbitrarily, we see that

$$(4.67) \quad \int_{\pi(U)} \bar{v}_l(x) \, dx \leq \liminf_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(\pi(U); F_L))}{L^n}.$$

Finally, consider a triangulation of  $X$  with finite partitions  $\{\pi_j(U_j)\}_{j=1}^J$  such that each  $(U_j, \pi_j)$  is a local chart on  $X$ . Then,

$$(4.68) \quad \int_X \bar{v}_l(x) \, dx = \sum_{j=1}^J \int_{\pi(U_j)} \bar{v}_l(x) \, dx \leq \liminf_{L \rightarrow \infty} \sum_{j=1}^J \frac{\mathcal{E}(\beta_l(\pi(U); F_L))}{L^n} \leq \liminf_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(X; F_L))}{L^n}.$$

□

For the upper bound, the methods in proving Theorem 4.4 do not work, because one cannot drop those giant components which intersect with boundaries of small cubes  $\{C_j\}$  now. And hence we do not know whether limits exist for Betti numbers cases. In fact, those giant components exist in many known concrete examples. Readers can see the discussions in page 6 of [W21] and references therein.

Here, we use the Chern–Lashof inequality in Theorem 2.9 to calculate a global upper bound directly.

*Proof of the upper bound in Theorem 4.9.* For the upper bound, it even holds for uniformly controllable Gaussian fields  $\{F_L\}_{L \in \mathcal{L}}$ , i.e.,  $\{F_L\}_{L \in \mathcal{L}}$  only satisfies Definition 4.1. Since  $X$  is a  $C^3$ -manifold, let  $\Phi : X \rightarrow \mathbb{R}^{n+q}$  for some  $q > 0$  be an isometric embedding according to Nash's embedding theorem. Then, we can regard  $X$  as a submanifold of  $\mathbb{R}^{n+q}$  and  $Z(F_L) = F_L^{-1}(0)$  as finitely many closed submanifolds of  $\mathbb{R}^{n+q}$ . According to Theorem 2.9, similar to the proofs of Theorem 2.3 and Theorem 2.8 (see (2.9)), one can get that

$$(4.69) \quad \beta_l(X; F_L) \leq C(n, m) \cdot \int_{Z(F_L)} \|\Pi^{Z(F_L)}(x)\|^{n-m} \, d\mathcal{H}^{n-m}(x).$$

Here, the second fundamental form is the one for  $Z(F_L)$  as submanifolds in  $\mathbb{R}^{n+q}$ . And then, by Theorem F.1,

$$(4.70) \quad \beta_l(X; F_L) \leq C(n, m) \int_{Z(F_L)} C(n, m, X) + \|A_L^{-1}\|^{\frac{(n-m)}{2}} \cdot \|(\nabla^X)^2 F_L\|^{(n-m)} \, d\mathcal{H}^{n-m}(x),$$

where  $C(n, m, X)$  is a constant depending on  $X$  and the embedding map  $\Phi$ , and  $A_L$  is an  $m \times m$  matrix, whose elements are  $(A_L)_{\alpha\beta} = \langle (\nabla^X f_{L,\alpha}), (\nabla^X f_{L,\beta}) \rangle_X$ . We also use  $\nabla^X$  to denote gradients on  $X$ ,  $(\nabla^X)^2$  to denote Hessians on  $X$ , and  $\langle \cdot, \cdot \rangle_X$  the Riemannian metric on  $X$ . One can use the similar trick in (2.10) and hence can give an upper bound for  $\|A_L^{-1}\|$  by  $|\det A_L|^{-1} \cdot \|\nabla^X F\|^{2(m-1)}$ . And hence, again, we set the Gaussian field as

$$(4.71) \quad W_L = (\nabla^X F_L, (\nabla^X)^2 F_L)$$

in the Kac–Rice formula Theorem 2.4, and set

(4.72)

$$h(x, W_L(x)) = C(n, m, X) + |\det A_L(x)|^{-\frac{(n-m)}{2}} \|\nabla^X F_L(x)\|^{(n-m)(m-1)} \|(\nabla^X)^2 F_L(x)\|^{(n-m)}.$$

Hence,

$$(4.73) \quad \begin{aligned} \mathcal{E}(\beta_i(X; F_L)) &\leq \mathcal{E} \left( \int_{Z(F_L)} h(x, W_L(x)) d\mathcal{H}^{n-m}(x) \right) \\ &= \int_X \mathcal{E} [h(x, W_L(x)) \cdot |\det A_L(x)|^{1/2} \mid F_L(x) = 0] \cdot p_{F_L(x)}(0) d\text{Vol}(x). \end{aligned}$$

The term  $p_{F_L(x)}(0)$  is bounded by some constant  $C = C(n, m, k_1, M) > 0$  according to (B1) and (B2) in Definition 4.1 for positive  $M = M(X)$  and  $k_1 = k_1(X)$ . So, let us tackle the expectation at each  $x \in X$ . For a local chart  $(U, \pi)$  around  $x \in \pi(U)$ , after choosing local normal coordinates  $(u_1, \dots, u_n)$  around  $x = 0$ , the  $\nabla^X$  and  $(\nabla^X)^2$  are just the usual gradient and the usual Hessian at  $x = 0$ . And then, rewrite them by the rescaled Gaussian fields we defined at the beginning of Section 4, i.e.,  $F_{x,L}(u) = F_L \circ \pi(x + L^{-1}u)$ ,  $\nabla^X F_L(x) = L \cdot \nabla F_{x,L}(0)$ , and  $(\nabla^X)^2 F_L(x) = L^2 \cdot (\nabla^2 F_{x,L}(0))$ . And also, the elements of  $A_L$  then become  $(A_L)_{\alpha\beta} = L^2 \cdot \langle (\nabla(F_{x,L})_{\alpha}(0)), (\nabla(F_{x,L})_{\beta}(0)) \rangle_{\mathbb{R}^n} \equiv L^2 \cdot (A_{x,L})_{\alpha\beta}$ , where  $\langle \cdot, \cdot \rangle_{\mathbb{R}^n}$  is the standard inner product in  $\mathbb{R}^n$  formed by coordinates  $(u_1, \dots, u_n)$ . Then,

(4.74)

$$\begin{aligned} &|\det A_L(x)|^{-\frac{(n-m)}{2}} \|\nabla^X F_L(x)\|^{(n-m)(m-1)} \|(\nabla^X)^2 F_L(x)\|^{(n-m)} \\ &= L^{-m(n-m)} |\det A_{x,L}(0)|^{-\frac{(n-m)}{2}} L^{(n-m)(m-1)} \|\nabla F_{x,L}(0)\|^{(n-m)(m-1)} L^{2(n-m)} \|(\nabla)^2 F_{x,L}(0)\|^{(n-m)} \\ &= L^{(n-m)} |\det A_{x,L}(0)|^{-\frac{(n-m)}{2}} \|\nabla F_{x,L}(0)\|^{(n-m)(m-1)} \|(\nabla)^2 F_{x,L}(0)\|^{(n-m)} \\ &\equiv L^{(n-m)} \cdot P(F_{x,L})(0), \end{aligned}$$

for  $P(F_{x,L})(0)$  denoting the long terms in order to simplify the notation in this proof. And we can then rewrite the expectation term at  $x$  as

$$(4.75) \quad \begin{aligned} &\mathcal{E} [h(x, W_L(x)) \cdot |\det A_L(x)|^{1/2} \mid F_L(x) = 0] \\ &= L^m \cdot C(n, m, X) \cdot \mathcal{E} [|\det A_{x,L}(0)|^{1/2} \mid F_{x,L}(0) = 0] \\ &\quad + L^n \cdot \mathcal{E} [P(F_{x,L})(0) \cdot |\det A_{x,L}(0)|^{1/2} \mid F_{x,L}(0) = 0]. \end{aligned}$$

Recall that  $F_L \circ \pi$  satisfies (B1) and (B2) in Definition 4.1 for  $M = M(X)$ ,  $k_1 = k_1(X)$ , when  $L$  is large. We then see, from the same proof in Theorem 2.3,

$$(4.76) \quad \mathcal{E} [|\det A_{x,L}(0)|^{1/2} \mid F_{x,L}(0) = 0] \leq C(n, m, M, k_1),$$

and

$$(4.77) \quad \mathcal{E} [P(F_{x,L})(0) \cdot |\det A_{x,L}(0)|^{1/2} \mid F_{x,L}(0) = 0] \leq C(n, m, M, k_1),$$

for some positive constants  $C(n, m, M, k_1)$ . Since  $n > m$ , the  $L^n$  term is the much larger term. And hence,

$$(4.78) \quad \limsup_{L \rightarrow \infty} \frac{\mathcal{E}(\beta_l(F_L))}{L^n} \leq \limsup_{L \rightarrow \infty} \frac{C(n, m, M, k_1)}{L^n} \int_X (L^m \cdot C(n, m, X) + L^n) d\mathcal{H}^n(x) = C_l \cdot |X|,$$

for  $C_l = C_l(n, m, M, k_1)$  denoting the constant  $C(n, m, M, k_1)$ , which is independent of the constant  $C(n, m, X)$  and hence independent of the arbitrarily chosen embedding  $\Phi$ .  $\square$

#### APPENDIX A. QUANTITATIVE BULINSKAYA'S LEMMA FOR GAUSSIAN RANDOM FIELDS

In this appendix, we assume that our Gaussian random field  $F : C_{R+1} \rightarrow \mathbb{R}^m$  satisfies  $(R; M, k_1)$ -assumptions on  $C_{R+1}$ . But only  $C^2$ -smoothness for  $F$  is enough for our proof here. We use similar methods as in [NS16] to prove the following lemma. First, let  $\|\cdot\|$  denote the norm of a vector and we define

$$(A.1) \quad \lambda(F(x)) \equiv \min_{w \in \mathbb{S}^{m-1}} \|\nabla(F \cdot w)\|(x).$$

It is easy to see that  $\lambda(F(x))^2$  is the smallest eigenvalue of the matrix  $(\nabla f_{i_1} \cdot \nabla f_{i_2})_{m \times m}$  at  $x$ .

**Lemma A.1.** *Given  $\delta > 0$ , there exists a  $\tau = \tau(\delta, R, n, m, M, k_1) \in (0, 1/2)$ , such that*

$$(A.2) \quad \mathcal{P}\left(\min_{x \in \bar{C}_R} \max\{\|F(x)\|, \lambda(F(x))\} < \tau\right) < \delta$$

*Proof.* Denote  $\Omega_\tau$  as the event that

$$(A.3) \quad \{\exists y \in \bar{C}_R, \text{ s.t. } \|F(y)\| < \tau, \lambda(F(y)) < \tau\}.$$

We put  $W = 1 + \|F\|_{C^{1+\beta}(C_{R+1/2})}$  with  $\beta \in (0, 1)$  to be specified later. Here  $\|F\|_{C^{1+\beta}(C_{R+1/2})}$  is the  $C^{1+\beta}$ -norm for  $F$  on the cube  $C_{R+1/2}$ . Then, if  $\Omega_\tau$  occurs for a  $y \in \bar{C}_R$ , we suppose that  $\|\nabla(F \cdot w_y)\|(y) < \tau$  for a corresponding fixed  $w_y \in \mathbb{S}^{m-1}$ . Then, for  $x$  in the ball  $B(y, \tau)$ ,

$$(A.4) \quad \|F(x)\| \leq \tau + \tau \cdot \|F\|_{C^{1+\beta}(C_{R+1/2})} = W\tau,$$

and

$$(A.5) \quad \lambda(F(x)) \leq \|\nabla(F \cdot w_y)\|(x) \leq \tau + \tau^\beta \cdot \|F\|_{C^{1+\beta}(C_{R+1/2})} \leq W\tau^\beta.$$

Since each element of the matrix  $(\nabla F(x))^T(\nabla F(x)) = (\nabla f_{i_1} \cdot \nabla f_{i_2})_{m \times m}$  has an upper bound  $W^2$ , the largest eigenvalue of it is also bounded by  $C(n, m) \cdot W^2$ . We can further get that

$$(A.6) \quad |\det((\nabla F(x))^T(\nabla F(x)))| \leq C(n, m) \cdot (\lambda(F(x)))^2 \cdot W^{2(m-1)} \leq C(n, m) \cdot W^{2m} \tau^{2\beta}.$$

So, set  $\alpha_1 = m + (n - m + 1)m$ ,  $\alpha_2 = m + (n - m + 1)\beta$ , then for some  $\eta \in (0, 1)$ , we define a function  $\Phi_\eta(x)$  for  $x \in C_{R+1}$  and get that

$$(A.7) \quad \Phi_\eta(x) \equiv \|F(x)\|^{-m\eta} \cdot |\det((\nabla F(x))^T(\nabla F(x)))|^{-\frac{n-m+1}{2}\eta} \geq C(n, m) \cdot W^{-\alpha_1\eta} \cdot \tau^{-\alpha_2\eta},$$

where the inequality is for  $x \in B(y, \tau)$ . So, whenever  $\Omega_\tau$  happens, we have that

$$(A.8) \quad \int_{C_{R+1}} \Phi_\eta(x) \geq \int_{B(y, \tau)} \Phi_\eta(x) \geq C(n, m) \cdot W^{-\alpha_1 \eta} \cdot \tau^{n - \alpha_2 \eta}.$$

Notice that  $n - \alpha_2 \eta = n(1 - \eta\beta) - m\eta(1 - \beta) - \beta\eta$ . Hence, we can choose  $\beta = \beta(n, m) \in (0, 1)$ ,  $\eta = \eta(n, m) \in (0, 1)$ , which are close to 1, such that  $n - \alpha_2 \eta < -\epsilon$  for some small positive  $\epsilon = \epsilon(n, m) \in (0, 1/2)$ . Hence,

$$(A.9) \quad \begin{aligned} \mathcal{P}(\Omega_\tau) &\leq C(n, m) \cdot \tau^{\alpha_2 \eta - n} \cdot \mathcal{E} \left[ W^{\alpha_1 \eta} \cdot \int_{C_{R+1}} \Phi_\eta(x) \right] \\ &\leq C(n, m) \cdot \tau^\epsilon \cdot \left( \mathcal{E}(W^{p\alpha_1 \eta}) \right)^{\frac{1}{p}} \cdot \left( \mathcal{E} \left( \int_{C_{R+1}} \Phi_\eta(x) \right)^q \right)^{\frac{1}{q}} \\ &\leq C(n, m) \cdot \tau^\epsilon \cdot |C_{R+1}|^{1 - \frac{1}{q}} \cdot \left( \mathcal{E}(W^{p\alpha_1 \eta}) \right)^{\frac{1}{p}} \cdot \left( \int_{C_{R+1}} \mathcal{E}(\Phi_\eta^q(x)) \right)^{\frac{1}{q}} \end{aligned}$$

for some  $p = p(n, m) > 1$ ,  $q = q(n, m) > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$  and  $q \cdot \eta < 1$ .

For the term  $\mathcal{E}(W^{p\alpha_1 \eta})$ , an upper bound in  $n, m, M, k_1$  follows from the Kolmogorov's theorem, see for example Appendix A.9 and Appendix A.11 in [NS16]. Next, since for each  $x \in C_{R+1}$ , the joint distribution of  $(F(x), \nabla F(x))$  satisfies (A1) in Definition 2.1, then similar to (2.26), we can write

$$(A.10) \quad \begin{aligned} \mathcal{E}(\Phi_\eta^q(x)) &= \mathcal{E} \left( \|F(x)\|^{-m\eta q} \cdot \left| \det((\nabla F(x))^T (\nabla F(x))) \right|^{-\frac{n-m+1}{2}\eta q} \right), \\ &\leq C(n, m, M, k_1) \cdot \int_{\mathbb{R}^m} \|v_0\|^{-m\eta q} \cdot e^{-c(n, m, M, k_1)(\|v_0\|^2)} dv_0 \\ &\quad \cdot \int_{(\mathbb{R}^n)^{\otimes m}} |\det(v_{t_1} \cdot v_{t_2})|^{-\frac{n-m+1}{2}\eta q} \cdot e^{-c(n, m, M, k_1)(\|v_1\|^2 + \dots + \|v_m\|^2)} dv_1 \cdots dv_m, \end{aligned}$$

where we use  $v_0$  to denote the components of  $F(x) = (f_1(x), \dots, f_m(x))$ , and  $(v_1, \dots, v_m)$  to denote components of  $\nabla F(x) = ((\nabla f_1(x))^T, \dots, (\nabla f_m(x))^T)$ , and  $(v_{t_1} \cdot v_{t_2})$  to denote the matrix with elements  $v_{t_1} \cdot v_{t_2}$ . Since  $\eta q < 1$  by our choice, the boundedness of the above two terms by some constants  $C = C(n, m, M, k_1) > 0$  (We do not need  $k_2$ -dependence here) are deduced previously in the proof of Theorem 2.3. See the process proving (2.16), (2.27) and Lemma 2.6. □

## APPENDIX B. STABILITY LEMMAS

We fix the  $\beta = \beta(m, n) \in (0, 1)$  chosen in the proof of Lemma A.1.

**Lemma B.1.** *Fix  $\alpha > 0$ ,  $A > 0$ . Assume that  $G(x) \in C^{1+\beta}(C_{R+1}, \mathbb{R}^m)$  for some  $m \in (0, n)$  and  $\|G\|_{C^{1+\beta}(C_{R+1})} < A$ . If for each  $x \in C_{R+1}$ , either  $\|G(x)\| > \alpha$  or  $\lambda(G(x)) > \alpha$  (see the*

definition for  $\lambda(G(x))$  in Lemma A.1), then, there is a positive constant  $\delta = \delta(n, m, \alpha, A) > 0$ , such that for any continuous vector field  $\tilde{G}(x) \in C(C_{R+1}, \mathbb{R}^m)$  with  $\|\tilde{G}(x) - G(x)\|_{C(C_{R+1})} < \delta$ , then  $N(R+1; \tilde{G}) \geq N(R; G)$ .

*Proof.* We will actually prove the following: Under the assumptions in the lemma, for each component  $\gamma$  of  $Z(G) = G^{-1}(0)$  inside  $C_R$ , there is a component  $\tilde{\gamma}$  of  $Z(\tilde{G})$  such that  $\tilde{\gamma}$  is contained in a connected component of  $\{\|G\| < \alpha\}$ . And for different components  $\gamma$  of  $Z(G)$  fully contained in  $C_R$ , they are contained in different connected components of  $\{\|G\| < \alpha\}$ .

We may write  $G(x) = (g_1(x), \dots, g_m(x))$ . Then,

$$(B.1) \quad \nabla(\|G(x)\|^2) = 2(g_1(x)\nabla g_1(x) + \dots + g_m(x)\nabla g_m(x)).$$

Whenever  $\|G(x)\| \in (0, \alpha)$ , since  $\lambda(G(x)) > \alpha > 0$ , we know that  $\nabla(\|G(x)\|^2) \neq 0$ . Hence, each component of  $\{\|G\| = a\}$  for  $a \in (0, \alpha)$  which evolves from a closed  $\gamma$  in  $C_R$  is a closed regular hypersurface. Hence, different components  $\gamma$  of  $Z(G)$  inside  $C_R$  are contained in different connected components of  $\{\|G\| < \alpha\}$ , which we denote as  $\mathfrak{C}(\gamma)$ . Notice that since  $\lambda(G(x)) > \alpha$  in  $\mathfrak{C}(\gamma)$ ,  $\mathfrak{C}(\gamma)$  is contained in the 1-neighbourhood of  $\gamma$ , and hence in  $C_{R+1}$ .

Now, for each  $\gamma$  of  $Z(G)$ , we choose an arbitrary point  $x \in \gamma$ . We consider an  $m$ -dimensional disc  $D^m(x, r)$  contained in the normal vector space of  $\gamma$  at  $x$  for some radius  $r$  chosen later, i.e.,  $D^m(x, r) \subset (T_x\gamma)^\perp \subset \mathbb{R}^n$ . Since  $\lambda(G(x)) > \alpha$ , there is an  $r = r(n, m, \alpha, A) \in (0, 1)$  such that for all  $y \in D^m(x, r)$ ,

$$(B.2) \quad \lambda^\perp(G(y)) \equiv \min_{w \in \mathbb{S}^{m-1}} \|\nabla^{(T_x\gamma)^\perp}(G \cdot w)\|(y) > \frac{\alpha}{2},$$

where  $\nabla^{(T_x\gamma)^\perp}g$  is the projection of  $\nabla g$  onto  $(T_x\gamma)^\perp$  (Also notice that  $\lambda^\perp(G(x)) = \lambda(G(x))$ ). By the inverse function theorem, we can regard  $G$  as a coordinates chart on  $D^m(x, r)$ . Then, we see that  $\{\|G\| \leq \frac{\alpha r}{4}\} \cap (T_x\gamma)^\perp$  is contained in  $D^m(x, r)$  and can be regarded as an  $m$ -dimensional closed ball under the chart  $G$ .

Now, we choose  $\delta = \frac{\alpha r}{8}$  and take an arbitrary  $\tilde{G}$  such that  $\|\tilde{G} - G\|_{C(C_{R+1})} < \delta$ . If we regard  $G - \tilde{G}$  as a map defined on  $\{\|G\| \leq \frac{\alpha r}{4}\} \cap (T_x\gamma)^\perp$ , i.e.,  $Id - \tilde{G} \circ G^{-1}$  is a map from  $G[\{\|G\| \leq \frac{\alpha r}{4}\} \cap (T_x\gamma)^\perp]$  to  $G[\{\|G\| \leq \frac{\alpha r}{8}\} \cap (T_x\gamma)^\perp]$ . By Brouwer's fixed point theorem under the chart  $G$ , there is a point  $y \in \{\|G\| \leq \frac{\alpha r}{4}\} \cap (T_x\gamma)^\perp$  such that  $G(y) - \tilde{G}(y) = G(y)$ , equivalently,  $\tilde{G}(y) = 0$ . Since  $\|\tilde{G}\| > \frac{\alpha}{2}$  on the boundary of the cooresponding  $\mathfrak{C}(\gamma)$ , we see that  $Z(\tilde{G})$  has at least one connected component inside  $\mathfrak{C}(\gamma)$ . Since each  $\mathfrak{C}(\gamma)$  does not intersect with the others, this finishes the proof.  $\square$

**Remark B.2.** I hope that one can directly choose  $\delta = \alpha$  as in Lemma 3 of [NS16], while our Lemma here is enough for our proofs of Lemma 4.5. On the other hand, if we choose  $\tilde{G} \in C^1(C_{R+1}, \mathbb{R}^m)$  and assume that  $\|\tilde{G}(x) - G(x)\|_{C^1(C_{R+1})}$  is small enough (depending on  $n, m, A, \alpha$ ), one can prove that there is a  $\tilde{\gamma}$  which is isotopic to corresponding  $\gamma$  via a family of  $C^1$ -embeddings and these embeddings are also contained in the corresponding  $\mathfrak{C}(\gamma)$ . For this  $C^1$ -stability result and the construction for the family of  $C^1$ -embeddings in  $\mathbb{R}^n$ , one can see,

for example Proposition 6.8 of [SW19] and Thom's isotopy theorem, which is the case when  $m = 1$ . General  $m > 1$  cases are similar and hence we omit the details. The general idea is considering  $G_t(x) = G(x) + t \cdot (\tilde{G}(x) - G(x))$ ,  $t \in [0, 1]$ . Since now  $\|\tilde{G}(x) - G(x)\|_{C^1(C_{R+1})}$  is small, then first, for each  $t$ ,  $Z(G_t)$  is regular. Then, one can replace the Brouwer's fixed point theorem by contraction mapping theorem in our previous proof and get the uniqueness of  $y_t$  in each  $(T_x\gamma)^\perp$  such that  $G_t(y_t) = 0$ . This process actually builds up the  $C^1$ -isotopy.

### APPENDIX C. CONDITIONS FOR ERGODICITY IN THEOREM 1.1 AND THEOREM 3.1

In this appendix, we give a sufficient condition to guarantee the ergodicity of translations on  $(C_*^1(\mathbb{R}^n, \mathbb{R}^m), \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m)), \gamma_F)$ . Here  $\mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m))$  is the Borel  $\sigma$ -algebra generated by all open sets in  $C_*^1(\mathbb{R}^n, \mathbb{R}^m)$ . Recall that  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a centred stationary Gaussian random field satisfying (A1) and (A2) in Definition 2.1 at the origin  $x = 0$ . And we write  $F(x) = (f_1(x), \dots, f_m(x))$ ,

$$(C.1) \quad k_{i_1 i_2}(x - y) \equiv K_{i_1 i_2}(x, y) = \mathcal{E}(f_{i_1}(x)f_{i_2}(y)), \quad i_1, i_2 = 1, 2, \dots, m,$$

is an  $m \times m$  matrix. Notice that, since  $F$  is stationary,  $\mathcal{E}(f_{i_1}(x)f_{i_2}(y)) = \mathcal{E}(f_{i_1}(x - y)f_{i_2}(0))$  while  $k_{i_2 i_1}(x - y) = \mathcal{E}(f_{i_2}(x)f_{i_1}(y)) = \mathcal{E}(f_{i_2}(0)f_{i_1}(y - x)) = k_{i_1 i_2}(y - x)$ . Hence,  $k_{i_2 i_1}(x) = k_{i_1 i_2}(-x)$  for all  $x \in \mathbb{R}^n$ .

**Theorem C.1.** *If*

$$(C.2) \quad \lim_{R \rightarrow \infty} \sum_{1 \leq i_1, i_2 \leq m} \frac{1}{|B_R|} \int_{B_R} (k_{i_1 i_2}(x))^2 dx = 0,$$

*then the translations actions of  $\mathbb{R}^n$  on  $(C_*^1(\mathbb{R}^n, \mathbb{R}^m), \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m)), \gamma_F)$  is ergodic.*

This is the Fomin–Grenander–Maruyama's theorem. The proof is exactly the same as the one in Appendix B of [NS16]. For completeness, we prove it for our random fields version.

*Proof.* Take an  $A \in \mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m))$  satisfying  $\gamma_F(\tau_v(A)\Delta A) = 0$  for every  $v \in \mathbb{R}^n$ , we need to show that  $\gamma_F(A)$  is either 0 or 1. Here  $(\tau_v G)(x) = G(x + v)$  for any  $v \in \mathbb{R}^n$  and  $G \in C_*^1(\mathbb{R}^n, \mathbb{R}^m)$ .

Notice that  $\mathfrak{B}(C_*^1(\mathbb{R}^n, \mathbb{R}^m))$  is generated by evaluations sets  $I(x; a, b)$  for  $x \in \mathbb{R}^n$  and  $a \prec b \in \mathbb{R}^m$  ( $a \prec b$  means that  $a_i < b_i$  for any  $i = 1, \dots, m$ ):

$$(C.3) \quad I(x; a, b) = \{G \in C_*^1(\mathbb{R}^n, \mathbb{R}^m) \mid a_i \leq G_i(x) < b_i \text{ for all } i = 1, \dots, m\},$$

where  $G_i(x)$  is the  $i$ -th component of  $G(x)$ . Hence, for any  $\epsilon > 0$ , we can take finitely many points  $x_1, \dots, x_k \in \mathbb{R}^n$  and a Borel set  $B \subset \mathbb{R}^{mk}$  so that  $\gamma_F(A\Delta P) < \epsilon$  for

$$(C.4) \quad P = P(x_1, \dots, x_k; B) \equiv \{G \in C_*^1(\mathbb{R}^n, \mathbb{R}^m) \mid (G(x_1), \dots, G(x_k)) \in B\}.$$

We can also assume that the distribution of the Gaussian vector  $(F(x_1), \dots, F(x_k))$  is non-degenerate (Otherwise, one just use the maximal linear basis of this Gaussian vector.). So,

we can get

$$(C.5) \quad \gamma_F(P) = (2\pi)^{-\frac{mk}{2}} (\det \Lambda)^{-\frac{1}{2}} \int_B e^{-\frac{1}{2} \langle \Lambda^{-1} t, t \rangle} dt,$$

where  $\Lambda_{j_1 i_1, j_2 i_2} = \mathcal{E}(f_{i_1}(x_{j_1}) f_{i_2}(x_{j_2})) = k_{i_1 i_2}(x_{j_1} - x_{j_2})$  is the covariance matrix for the Gaussian vector  $(F(x_1), \dots, F(x_k))$  and it is an  $mk \times mk$  matrix, and  $\langle \cdot, \cdot \rangle$  is the inner product for vectors in  $\mathbb{R}^{mk}$ .

Since  $\tau_v P = P(x_1 - v, \dots, x_k - v; B)$ , we have

$$(C.6) \quad P \cap \tau_v P = P(x_1, \dots, x_k, x_1 - v, \dots, x_k - v; B \times B).$$

If we know that  $(F(x_1), \dots, F(x_k), F(x_1 - v), \dots, F(x_k - v))$  is also non-degenerate, we can write

$$(C.7) \quad \gamma_F(P \cap \tau_v P) = (2\pi)^{-mk} (\det \tilde{\Lambda})^{-\frac{1}{2}} \int_{B \times B} e^{-\frac{1}{2} \tilde{\Lambda}^{-1} t \cdot t} dt,$$

where

$$(C.8) \quad \tilde{\Lambda} = \begin{bmatrix} \Lambda & \Theta(v) \\ \Theta^T(v) & \Lambda \end{bmatrix}$$

with  $\Theta_{j_1 i_1, j_2 i_2} = \mathcal{E}(f_{i_1}(x_{j_1}) f_{i_2}(x_{j_2} - v)) = k_{i_1 i_2}(x_{j_1} - x_{j_2} + v)$ . We will prove that we can choose a sequence  $v_l \in \mathbb{R}^n$  so that  $\|\Theta(v_l)\| \rightarrow 0$  as  $l \rightarrow \infty$ , and hence we can not only get the nondegeneracy for (C.8) and then (C.7), but also conclude that

$$(C.9) \quad \lim_{l \rightarrow \infty} \gamma_F(P \cap \tau_{v_l} P) = (\gamma_F(P))^2.$$

And then

$$(C.10) \quad \gamma_F(A) = \limsup_{l \rightarrow \infty} \gamma_F(A \cap \tau_{v_l} A) \leq \limsup_{l \rightarrow \infty} \gamma_F(P \cap \tau_{v_l} P) + 2\epsilon \leq (\gamma_F(A) + \epsilon)^2 + 2\epsilon.$$

Let  $\epsilon \rightarrow 0$ , we see that  $\gamma_F(A)$  is 0 or 1 as  $\gamma_F$  is a probability measure.

Now, the reason we can choose such a sequence  $v_l$  is because of our assumption. Notice that if we denote  $T = \max_{j_1, j_2} |x_{j_1} - x_{j_2}|$ , then for any  $R > 0$ , we have that

$$(C.11) \quad \begin{aligned} \frac{1}{|B_R|} \int_{B_R} \|\Theta(v)\|^2 dv &\leq \frac{C(m, k)}{|B_R|} \sum_{i_1, i_2, j_1, j_2} \int_{B_R} (k_{i_1 i_2}(x_{j_1} - x_{j_2} + v))^2 dv \\ &\leq \frac{C(m, k)}{|B_R|} \sum_{1 \leq i_1, i_2 \leq m} \int_{B_{R+T}} (k_{i_1 i_2})^2(v) dv, \end{aligned}$$

which goes to 0 as  $R \rightarrow \infty$  by our assumption. We then choose  $v_l$  by mean value theorem.  $\square$

**Remark C.2.** In the case that  $m = 1$ , by Wiener's lemma, our assumption is equivalent to say that the spectral measure, i.e., the Fourier transform, of the covariance function  $k(x - y) = \mathcal{E}(f(x)f(y))$  has no atoms. This is mentioned in Appendix B of [NS16] in proving this ergodicity theorem. In our random fields cases, if we assume that  $F$  consists of independent centred stationary Gaussian random functions, i.e., each  $f_{i_1}(x)$  is independent

of  $f_{i_2}(y)$  for  $i_2 \neq i_1$  and any  $x, y \in \mathbb{R}^n$ , and we make the assumption that for each  $f_{i_1}$ , the corresponding spectral measure  $\rho_{i_1}$  does not have atoms, then we can also get (C.2) by Wiener's lemma and hence get our ergodicity theorem. Also, for  $m = 1$  cases, it is mentioned in [NS16] that the ergodicity is actually equivalent to spectral measures having no atoms, which is actually the full version of Fomin–Grenander–Maruyama's theorem.

#### APPENDIX D. CONDITIONS FOR POSITIVE LIMITING CONSTANTS IN THEOREM 1.1 AND THEOREM 3.1

In this appendix, similar to Appendix C, we let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a centred stationary Gaussian random field satisfying (A1) and (A2) in Definition 2.1 at the origin  $x = 0$ . Our main goal in this appendix is to build up the following fact under some further assumptions on  $F$ . For any bounded open set  $B \subset \mathbb{R}^n$ , any  $G \in C^1(\mathbb{R}^n, \mathbb{R}^m)$ , and any  $\epsilon > 0$ ,

$$(D.1) \quad \mathcal{P}(\|F - G\|_{C^1(B)} < \epsilon) > 0.$$

For this purpose, we need to recall some basic facts for Gaussian fields first. One can see Chapter 4 of [GS04] together with Appendix A of [NS16] as references for the following discussions.

We write  $F(x) = (f_1(x), \dots, f_m(x))$ , and then

$$(D.2) \quad k_{i_1 i_2}(x - y) \equiv K_{i_1 i_2}(x, y) = \mathcal{E}(f_{i_1}(x)f_{i_2}(y)), \quad i_1, i_2 = 1, 2, \dots, m,$$

is an  $m \times m$  matrix. Since  $F$  is stationary,  $k_{i_2 i_1}(x) = k_{i_1 i_2}(-x)$  for all  $x \in \mathbb{R}^n$ . Also,  $k$  has a positive property in the following sense. Let  $x_1, \dots, x_r \in \mathbb{R}^n$  and  $a_1, \dots, a_r \in \mathbb{C}$ , then

$$(D.3) \quad \sum_{1 \leq t, s \leq r} k(x_t - x_s) a_t \bar{a}_s \quad \text{is positive semi-definite in } \text{End}_{\mathbb{C}}(\mathbb{R}^m).$$

This is because for any  $\eta = (\eta_1, \dots, \eta_m)^T \in \mathbb{C}^m$ ,

$$(D.4) \quad \begin{aligned} \eta^T \cdot \left( \sum_{1 \leq t, s \leq r} k(x_t - x_s) a_t \bar{a}_s \right) \cdot \bar{\eta} &= \sum_{1 \leq i_1, i_2 \leq m} \sum_{1 \leq t, s \leq r} \mathcal{E} \left( a_t \eta_{i_1} f_{i_1}(x_t) \cdot \overline{a_s \eta_{i_2} f_{i_2}(x_s)} \right) \\ &= \mathcal{E} \left| \sum_{i_1=1}^m \sum_{t=1}^r a_t \eta_{i_1} f_{i_1}(x_t) \right|^2 \geq 0. \end{aligned}$$

Then, by Bochner–Herglotz's theorem, the Fourier transform of  $k$ , which we denote as  $\rho = (\rho_{i_1 i_2})$ , is a finite positive semi-definite  $\text{End}_{\mathbb{C}}(\mathbb{R}^m)$ -valued measure on  $\mathbb{R}^n$  and satisfies that

$$(D.5) \quad \mathcal{E}(f_{i_1}(x)f_{i_2}(y)) = k_{i_1 i_2}(x - y) = \int_{\mathbb{R}^n} e^{i\langle x-y, \xi \rangle} d\rho_{i_1 i_2}(\xi).$$

Since  $k(x) = (k(-x))^T$ , we see that  $\rho = \rho^* \equiv (\bar{\rho})^T$  and  $\rho(\xi) = (\rho(-\xi))^T$ . Indeed,  $F$  has the following expression.

$$(D.6) \quad F(x) = \int_{\mathbb{R}^n} e^{i\langle x, \xi \rangle} dZ(\xi),$$



for a complex vector-valued orthogonal random measure  $Z(\cdot)$ , which satisfies that for any Borel sets  $U_1, U_2 \subset \mathbb{R}^n$  with  $\rho(U_1)$  and  $\rho(U_2)$  finite, we have that  $\mathcal{E}(Z(U_1)) = \mathcal{E}(Z(U_2)) = 0$  and  $\mathcal{E}(Z_{i_1}(U_1)\overline{Z_{i_2}(U_2)}) = \rho_{i_1 i_2}(U_1 \cap U_2)$  for  $i_1, i_2 = 1, \dots, m$ . Notice that this also implies that  $|\rho_{i_1 i_2}(U_1 \cap U_2)|^2 \leq |\rho_{i_1 i_1}(U_1)| \cdot |\rho_{i_2 i_2}(U_2)|$  by Cauchy–Schwarz inequality. Hence, for any Borel set  $U \subset \mathbb{R}^n$ , if  $\rho_{i_1 i_2}(U) \neq 0$ , then  $\rho_{i_1 i_1}(U) \neq 0$  and  $\rho_{i_2 i_2}(U) \neq 0$ .

We define the space of square-summable Hermitian fields as

$$(D.7) \quad L_H^2(\rho) \equiv \{G : \mathbb{R}^n \rightarrow \mathbb{C}^m \mid G \in L^2(\rho), \text{ and } G(x) = \overline{G(-x)} \text{ for all } x \in \mathbb{R}^n\},$$

where for  $G = (g_1, \dots, g_m)^T \in L^2(\rho)$ ,

$$(D.8) \quad \|G\|_{L^2(\rho)}^2 \equiv \sum_{1 \leq i_1, i_2 \leq m} \int_{\mathbb{R}^n} g_{i_1}(\xi) \overline{g_{i_2}(\xi)} d\rho_{i_1 i_2}(\xi) < \infty.$$

And we also define the reproducing kernel Hilbert spaces associated with  $\rho$  (or equivalently, associated with  $k$ ), which are

$$(D.9) \quad \mathcal{H}(\rho) \equiv \mathcal{F}_\rho(L^2(\rho)), \quad \mathcal{H}_0(\rho) \equiv \mathcal{F}_\rho(L_H^2(\rho)),$$

where for  $G = (g_1, \dots, g_m)^T \in L^2(\rho)$  and for each  $i_2 = 1, \dots, m$ ,

$$(D.10) \quad \hat{g}_{i_2}(x) \equiv (\mathcal{F}_\rho(G)(x))_{i_2} = \sum_{i_1=1}^m \int_{\mathbb{R}^n} e^{-i\langle x, \xi \rangle} g_{i_1}(\xi) d\rho_{i_1 i_2}(\xi).$$

We denote  $\hat{G} = \mathcal{F}_\rho(G) = (\hat{g}_1, \dots, \hat{g}_m)^T$ . The inner product of  $\mathcal{H}(\rho)$  is defined by

$$(D.11) \quad \langle \widehat{G}_1, \widehat{G}_2 \rangle_{\mathcal{H}(\rho)} \equiv \langle G_1, G_2 \rangle_{L^2(\rho)}.$$

Define a map  $\Phi : L^2(\Omega, \mathfrak{S}, \mathcal{P}) \rightarrow C^1(\mathbb{R}^n, \mathbb{R}^m)$ ,  $h \mapsto \Phi[h](x)$ , and for each  $t = 1, \dots, m$ , the  $t$ -th component of  $\Phi[h]$ ,  $(\Phi[h])_t$ , is given by  $\mathcal{E}(hf_t(x))$ , the covariance of  $h$  and  $f_t(x)$ . Notice that although our random variables in  $L^2(\Omega, \mathfrak{S}, \mathcal{P})$  and  $F$  are real-valued, we still write  $\overline{(\cdot)}$  to be consistent. Then, the image of  $\Phi$ ,  $\mathcal{H}(F) \equiv \Phi(L^2(\Omega, \mathfrak{S}, \mathcal{P}))$ , is a Hilbert space with induced inner product

$$(D.12) \quad \langle \Phi[h_1], \Phi[h_2] \rangle_{\mathcal{H}(F)} \equiv \mathcal{E}(h_1 \overline{h_2}) = \langle h_1, h_2 \rangle_{L^2(\Omega, \mathfrak{S}, \mathcal{P})},$$

for any  $h_1, h_2 \in H(F)$ , where  $H(F) \subset L^2(\Omega, \mathfrak{S}, \mathcal{P})$  is the closed  $\mathbb{R}$ -linear span of all  $(F(x) \cdot w)$  for  $x \in \mathbb{R}^n$  and  $w \in \mathbb{R}^m$ , which is exactly the orthogonal complement of  $\text{Ker}(\Phi)$ . Let  $\{h_j\}$  be a countable orthonormal basis of  $H(F)$  and set  $e_j = \Phi[h_j] \in C^1(\mathbb{R}^n, \mathbb{R}^m)$ , we see that for any  $w \in \mathbb{R}^m$ , and  $x \in \mathbb{R}^n$  fixed,

$$(D.13) \quad F(x) \cdot w = \sum_j \mathcal{E}((F(x) \cdot w)h_j)h_j = \sum_j (\Phi[h_j](x) \cdot w)h_j,$$

and equivalently,

$$(D.14) \quad F(x) = \sum_j e_j(x)h_j.$$

For any  $\tilde{G} \in \mathcal{H}(F)$ , one can write a series convergent in  $\mathcal{H}(F)$ ,

$$(D.15) \quad \tilde{G}(x) = \sum_j e_j(x) \langle \tilde{G}, e_j \rangle_{\mathcal{H}(F)}.$$

Since we made the assumption (A2) in Definition 2.1 that  $F$  is a  $C^{3-}$ -smooth Gaussian field, the convergence (D.15) is actually, at least, in locally  $C^1$ -sense. And then we can get that for any  $\epsilon > 0$  and any bounded open set  $B \subset \mathbb{R}^n$ ,

$$(D.16) \quad \mathcal{P}(\|F - \tilde{G}\|_{C^1(B)} > \epsilon) > 0.$$

On the other hand, there is a way to link the inner product  $\langle \cdot, \cdot \rangle_{\mathcal{H}(\rho)}$  and  $\langle \cdot, \cdot \rangle_{\mathcal{H}(F)}$  by the following isometry map, which shows that these two spaces,  $\mathcal{H}_0(\rho)$  and  $\mathcal{H}(F)$ , and their inner products are actually the same. For any  $G = (g_1, \dots, g_m)^T \in L_H^2(\rho)$ , let

$$(D.17) \quad I(G) \equiv \int_{\mathbb{R}^n} G(\xi) \cdot dZ(\xi) = \sum_{i=1}^m \int_{\mathbb{R}^n} g_i(\xi) dZ_i(\xi),$$

then

$$(D.18) \quad \begin{aligned} \|I(G)\|_{L^2(\Omega, \mathfrak{S}, \mathcal{P})}^2 &= \mathcal{E} \left( \int_{\mathbb{R}^n} G(\xi_1) \cdot dZ(\xi_1) \int_{\mathbb{R}^n} \overline{G(\xi_2)} \cdot d\overline{Z(\xi_2)} \right) \\ &= \sum_{1 \leq i_1, i_2 \leq m} \int_{\mathbb{R}^n} g_{i_1}(\xi) \overline{g_{i_2}(\xi)} d\rho_{i_1 i_2}(\xi) = \|G\|_{L^2(\rho)}^2. \end{aligned}$$

Notice that  $H(F) \subset I(L_H^2(\rho))$ . Actually, one can even show that  $H(F) = I(L_H^2(\rho))$ . This is because that  $\{e^{i\langle x, \xi \rangle}\}_{x \in \mathbb{R}^n}$  is a dense family as  $\xi$ -functions in  $L_{\rho_{tt}}^2(\mathbb{R}^n)$  for each  $t = 1, \dots, m$ , which means that the linear combinations of this family can approximate smooth functions with compact support, and one then compares (D.6) and (D.17). Then, one can check that

$$(D.19) \quad \begin{aligned} (\Phi[I(G)](x))_{i_2} &= \mathcal{E}(I(G) \overline{f_{i_2}(x)}) = \mathcal{E} \left( \int_{\mathbb{R}^n} G(y) \cdot dZ(y) \int_{\mathbb{R}^n} e^{-i\langle x, \xi \rangle} d\overline{Z_{i_2}(\xi)} \right) \\ &= \sum_{i_1=1}^m \int_{\mathbb{R}^n} e^{-i\langle x, \xi \rangle} g_{i_1}(\xi) d\rho_{i_1 i_2}(\xi) = (\mathcal{F}_\rho(G)(x))_{i_2}, \end{aligned}$$

and then

$$(D.20) \quad \begin{aligned} \langle \widehat{G}_1, \widehat{G}_2 \rangle_{\mathcal{H}(\rho)} &= \langle G_1, G_2 \rangle_{L^2(\rho)} = \langle I(G_1), I(G_2) \rangle_{L^2(\Omega, \mathfrak{S}, \mathcal{P})} \\ &= \langle \Phi[I(G_1)], \Phi[I(G_2)] \rangle_{\mathcal{H}(F)} = \langle \widehat{G}_1, \widehat{G}_2 \rangle_{\mathcal{H}(F)}. \end{aligned}$$

For similar reason, one can show that  $\mathcal{H}(F) = \mathcal{H}_0(\rho)$ .

We define a subset  $\Sigma_\rho$  of  $C^1(\mathbb{R}^n, \mathbb{R}^m)$  as the following. For each element  $H(x) \in \Sigma_\rho$ , write  $H(x)$  as  $(h_1(x), \dots, h_m(x))^T$ , and then for each  $t = 1, \dots, m$ ,  $h_t(x)$  has the form

$$(D.21) \quad h_t(x) = \sum_{p=1}^q \sum_{s=1}^m \left( e^{-i\langle x, \xi_p \rangle} g_s(\xi_p) \rho_{st}(V_p) + e^{i\langle x, \xi_p \rangle} g_s(-\xi_p) \rho_{st}(-V_p) \right),$$

where  $V_p$  are Borel sets,  $\xi_p \in V_p$ , and  $G = (g_1, \dots, g_m)^T \in L_H^2(\rho)$ . We emphasize that these  $V_p, \xi_p$  are the same for different  $t = 1, \dots, m$ . This  $H(x)$  is actually a Riemann sum of  $\mathcal{F}_\rho(G)$  in (D.10) if  $G$  is also continuous. Also, notice that  $h_t(x)$  is real, because  $g_s(-\xi_p) = \overline{g_s(\xi_p)}$  and  $\rho_{st}(-V_p) = \overline{\rho_{st}(V_p)}$ .

Our aim is to prove that under some conditions, one can show that  $\Sigma_\rho$ , and hence  $\mathcal{H}_0(\rho)$ , is  $C^1$ -dense in  $C^1(B, \mathbb{R}^m)$  for any bounded open set  $B \subset \mathbb{R}^n$  and use (D.16) to get an approximation by  $F(x)$  with positive probability. This strategy was sketched in the Lemma 5.5 and Proposition 5.2 in [SW19].

We define that the  $\mathbb{C}^m$ -valued orthogonal random measure  $Z(\cdot) = (Z_1(\cdot), \dots, Z_m(\cdot))^T$  in (D.6) is *m-balls absolute non-degenerate* if there are  $m$  open balls  $\{B_{r_t}(y_t)\}_{t=1}^m$  such that for each  $t = 1, \dots, m$ , and for any nonempty open subset  $U \subset B_{r_t}(y_t)$ ,

$$(D.22) \quad Z_t(U) \notin \text{Span}_{\mathbb{C}}\{Z_1(U), \dots, Z_{t-1}(U), Z_{t+1}(U), \dots, Z_m(U)\},$$

which is equivalent to say that there are complex constants  $c_{t1}(U), \dots, c_{tm}(U)$ , such that

$$(D.23) \quad \mathcal{E} \left( \left( \sum_{i=1}^m c_{ti}(U) Z_i(U) \right) \overline{Z_s(U)} \right) = \sum_{i=1}^m c_{ti}(U) \rho_{is}(U) = \delta_{ts}.$$

This also implies that  $B_{r_t}(y_t)$  is in the interior of the compact support of  $\rho_{tt}$ . Moreover, we requires that there are absolute positive constants  $C_1(Z)$  and  $C_2(Z)$  such that for any open subset  $V \subset \cup_i B_{r_i}(y_i)$ ,  $|\rho_{st}|(V) \leq C_1(Z) \cdot |\rho_{st}(V)|$  for any  $s, t$ ; and for any  $s, t$  and  $U \subset B_{r_t}(y_t)$ ,  $\sum_{i=1}^m |c_{ti}(U) \rho_{is}(U)| \leq C_2(Z)$ .

**Example D.1.** For random fields with independent components, i.e.,  $Z_s(\cdot)$  is independent of  $Z_t(\cdot)$  when  $s \neq t$ , this *m-balls absolute non-degenerate* condition is equivalent to say that for each  $t$ , the interior of the compact support of  $\rho_{tt}$  is nonempty. In this case, each  $\rho_{tt}$  is also a nonnegative measure and one chooses  $B_{r_t}(y_t)$  contained in the interior of the compact support of  $\rho_{tt}$ , and  $c_{ti}(U) = \delta_{ti} \cdot \rho_{tt}(U)^{-1}$  for all  $t$  and  $i$ . And then  $C_1(Z)$  and  $C_2(Z)$  can be chosen as 1 automatically.

In general, the constant  $C_2(Z)$  is quite tricky to get. One may use the following weaker but more explicit assumption.

**Example D.2.** Assume that there is an open ball  $B \subset \mathbb{R}^n$ , and a positive constant  $\lambda$ , such that for any open subset  $U \subset B$ ,

$$(D.24) \quad \lambda^{-1} \cdot |U| \cdot \text{Id}_m \leq \rho(U) \leq \lambda \cdot |U| \cdot \text{Id}_m,$$

where  $\rho(U)$  is the  $m \times m$  matrix with elements  $\rho_{i_1 i_2}(U)$ , and for matrix  $A \geq B$ , we mean that the smallest eigenvalue of  $A - B$  is nonnegative. And  $|U|$  is the volume of  $U$  in  $\mathbb{R}^n$ . One can also replace  $|\cdot|$  by any other measure which is mutually absolutely continuous with respect to  $|\cdot|$ . With this nondegenerate assumption, one can choose  $c(U)$  as the inverse of  $\rho(U)$  and set a  $C_2(Z)$  only depending on  $\lambda$ . And if we also have some  $C_1(Z)$  satisfying the definition, we can get that the random measure  $Z(\cdot)$  satisfies this *m-balls absolute non-degenerate* condition.

According to the above definitions and examples, assume that we can get those two universal positive constants  $C_1(Z)$  and  $C_2(Z)$  for the  $m$ -balls *absolute non-degenerate* condition, we then have the following theorem.

**Theorem D.3.** *If  $Z(\cdot)$  is  $m$ -balls absolute non-degenerate as defined above, then  $\mathcal{H}_0(\rho)$  is  $C^1$ -dense in  $C^1(B, \mathbb{R}^m)$  for any bounded open set  $B \subset \mathbb{R}^n$ .*

*Proof.* By the assumption, denote  $B_{r_t}(y_t)$  by  $V^t$ . We assume that  $\{V^t\}_{t=1}^m$  are disjoint, otherwise we choose smaller open balls of them. Recall that  $\rho(x) = \rho^*(x) = (\rho(-x))^T$ . And we can also assume that  $V^t \cap (-V^t) = \emptyset$ , otherwise we choose smaller balls again. The following strategies were sketched in Lemma 5.5 and Proposition 5.2 in [SW19]. For completeness, we include them here.

Now, fix any  $p(x) = (p_1(x), \dots, p_m(x))^T$  such that for each  $t = 1, \dots, m$ ,  $p_t(x)$  is a real polynomial in  $x_1, \dots, x_n$ . Then, for any  $\epsilon > 0$  and any bounded open set  $B$ , we need to find a  $G \in L^2_H(\rho)$  such that

$$(D.25) \quad \|(\mathcal{F}_\rho(G)(x))_t - p_t(x)\|_{C^1(B)} < \epsilon.$$

We also fix such  $\epsilon$  and  $B$  and then proceed.

Let  $\phi(\xi)$  be a smooth, nonnegative function supported in  $B_1(0)$  such that  $\int_{\mathbb{R}^n} \phi(\xi) d\xi = 1$  and  $\phi(\xi_1) = \phi(\xi_2)$  if  $\|\xi_1\| = \|\xi_2\|$ . For any  $\delta_t \in (0, 1)$  and any multi-index  $\beta_t$ , we see that the function

$$(D.26) \quad \phi_{\delta_t, \beta_t}(\xi) \equiv \frac{\partial^{\beta_t}}{\partial \xi^{\beta_t}} \left( \frac{1}{(r_t \delta_t)^n} \phi\left(\frac{\xi - y_t}{r_t \delta_t}\right) \right)$$

is supported in  $V^t = B_{r_t}(y_t)$ . Now, we do the Fourier transform on  $(-i)^{\beta_t} \phi_{\delta_t, \beta_t}(\xi)$ , and then get a function

$$(D.27) \quad \begin{aligned} \tilde{g}_{\delta_t, \beta_t}(x) &\equiv (-i)^{\beta_t} \cdot \int_{\mathbb{R}^n} (\phi_{\delta_t, \beta_t}(\xi)) e^{-i\langle x, \xi \rangle} d\xi \\ &= x^{\beta_t} \cdot \int_{\mathbb{R}^n} \frac{1}{(r_t \delta_t)^n} \phi\left(\frac{\xi - y_t}{r_t \delta_t}\right) e^{-i\langle x, \xi \rangle} d\xi \\ &\rightarrow x^{\beta_t} \cdot e^{-i\langle x, y_t \rangle}, \end{aligned}$$

as  $\delta_t \rightarrow 0^+$ . Notice that, for any bounded open set  $B \subset \mathbb{R}^n$ , the convergence of (D.27) is in  $C^1(B, \mathbb{C})$  as functions in  $x$ . Here, for  $g \in C^1(B, \mathbb{C})$ , we mean that  $g = u + iv$  for  $u, v \in C^1(B, \mathbb{R})$ , and the  $C^1$ -norm is induced by  $C^1$ -norms on  $u, v$  components.

We first consider a polynomial with complex coefficient, which we denote as  $q_\epsilon(x)$ , such that  $\|q_\epsilon(x) - e^{i\langle x, y_t \rangle}\|_{C^1(B)} \leq \epsilon$ . Then by (D.27), one can find a smooth complex-valued function  $\phi_t(\xi)$  supported in  $V^t$  such that the Fourier transform of  $\phi_t$ , i.e.,  $\mathcal{F}(\phi_t)(x) \equiv \int_{\mathbb{R}^n} \phi_t(\xi) e^{-i\langle x, \xi \rangle} d\xi$ , satisfies that

$$(D.28) \quad \|\mathcal{F}(\phi_t)(x) - p_t(x) q_\epsilon(x) e^{-i\langle x, y_t \rangle}\|_{C^1(B)} \leq \epsilon.$$

And hence

$$(D.29) \quad \|\mathcal{F}(\phi_t)(x) - p_t(x)\|_{C^1(B)} \leq C(\|p_t\|_{C^1(B)}, \|y_t\|) \cdot \epsilon,$$

where  $C(\|p_t\|_{C^1(B)}, \|y_t\|)$  is a positive constant depending on the  $C^1$ -norm of  $p_t$  on  $B$  and the length of  $y_t$ . We then choose  $\varphi_t(\xi) = (\phi_t(\xi) + \overline{\phi_t(-\xi)})/2$ , whose Fourier transform is the real part of  $\mathcal{F}(\phi_t)(x)$  and (D.29) still holds for  $\mathcal{F}(\varphi_t)(x)$ . Notice that  $\varphi_t(-\xi) = \overline{\varphi_t(\xi)}$  for any  $\xi \in \mathbb{R}^n$  and is supported in  $V^t \cup (-V^t)$ .

Now, approximate  $\mathcal{F}(\varphi_t)(x)$  in  $C^1(B)$  by a Riemann sum form function, i.e., for the chosen  $\epsilon > 0$ , we have an

$$(D.30) \quad R(\varphi_t)(x) = \frac{1}{2} \sum_{p=1}^{q_t} \left( e^{-i\langle x, \xi_p^t \rangle} \phi_t(\xi_p^t) |V_p^t| + e^{i\langle x, \xi_p^t \rangle} \overline{\phi_t(\xi_p^t)} |V_p^t| \right),$$

such that  $\|\mathcal{F}(\varphi_t)(x) - R(\varphi_t)(x)\|_{C^1(B)} \leq \epsilon$ , where  $\{V_p^t\}_{p=1}^{q_t}$  is a disjoint open subdivision of  $V^t$  and each  $\xi_p^t$  is a point in  $V_p^t$ . Also, this subdivision is so narrow such that for any  $\xi \in V_p^t$  and any  $x \in B$ ,

$$(D.31) \quad |e^{-i\langle x, \xi_p^t \rangle} - e^{-i\langle x, \xi \rangle}| \leq \epsilon \cdot \left( \sum_{s=1}^m \|\phi_s\|_{L^1(\mathbb{R}^n)} \right)^{-1}, \quad |\xi_p e^{-i\langle x, \xi_p^t \rangle} - \xi e^{-i\langle x, \xi \rangle}| \leq \epsilon \cdot \left( \sum_{s=1}^m \|\phi_s\|_{L^1(\mathbb{R}^n)} \right)^{-1},$$

and

$$(D.32) \quad \sum_{l=1}^m \sum_{V_p^l} |\phi_l(\xi_p^l)| \cdot |V_p^l| \leq 2 \cdot \left( \sum_{l=1}^m \|\phi_l\|_{L^1(\mathbb{R}^n)} \right).$$

We then collect all  $\{V_p^t\}$  for all  $t = 1, \dots, m$ , and get a family of disjoint open sets  $\{V_p\}$  and their corresponding  $\{\xi_p^s\}$ . Notice that  $\phi_t(\xi_p^s) = 0$  for  $s \neq t$ . We denote  $q = \sum_{t=1}^m q_t$ .

Compare  $R(\varphi_t)(x)$  and functions in  $\Sigma_\rho$  of the form (D.21), we consider a function  $G = (g_1, \dots, g_m)^T \in L_H^2(\rho)$  such that  $G(x) = 0$  if  $x$  is not contained in any  $V_p$  or  $-V_p$ . And for each  $g_s$  and each  $V_p^t$ , let  $g_s$  be the constant  $c_{ts}(V_p^t)(\phi_t(\xi_p^t)|V_p^t|)$  on  $V_p^t$  and  $c_{ts}(V_p^t)(\overline{\phi_t(\xi_p^t)}|V_p^t|)$  on  $(-V_p^t)$ , where the constants  $c_{ts}(V_p^t)$  are from (D.23). This  $G$  is well-defined and in  $L_H^2(\rho)$ .

By the transform (D.21), we see that for each  $t = 1, \dots, m$ ,

$$(D.33) \quad \begin{aligned} (H(G)(x))_t &\equiv \sum_{p=1}^q \sum_{s=1}^m \left( e^{-i\langle x, \xi_p \rangle} g_s(\xi_p) \rho_{st}(V_p) + e^{i\langle x, \xi_p \rangle} g_s(-\xi_p) \rho_{st}(-V_p) \right) \\ &= 2\text{Re} \sum_{l=1}^m \sum_{V_p^l} \sum_{s=1}^m e^{-i\langle x, \xi_p^l \rangle} g_s(\xi_p^l) \rho_{st}(V_p^l) = 2\text{Re} \sum_{l=1}^m \sum_{V_p^l} \sum_{s=1}^m e^{-i\langle x, \xi_p^l \rangle} c_{ts}(V_p^l) \rho_{st}(V_p^l) (\phi_l(\xi_p^l) |V_p^l|) \\ &= 2\text{Re} \sum_{l=1}^m \sum_{V_p^l} e^{-i\langle x, \xi_p^l \rangle} \delta_{lt} (\phi_l(\xi_p^l) |V_p^l|) = 2\text{Re} \sum_{V_p^t} e^{-i\langle x, \xi_p^t \rangle} (\phi_t(\xi_p^t) |V_p^t|) = 2R(\varphi_t)(x), \end{aligned}$$

where  $\text{Re}(\cdot)$  means the real part of a complex number. And hence, we see that  $R(\varphi_t)(x)$  is actually expressed by  $G/2$  in transform (D.21).

Then, we compare  $H(G)$  and the  $\rho$ -Fourier transform of  $G$ . Recall that the subdivision  $\{V_p\}$  satisfies that for any  $\xi \in V_p$ , both  $|e^{-i\langle x, \xi_p \rangle} - e^{-i\langle x, \xi \rangle}|$  and  $|\xi_p e^{-i\langle x, \xi_p \rangle} - \xi e^{-i\langle x, \xi \rangle}|$  are bounded by the chosen  $\epsilon$  and a constant in (D.31). And the absolute non-degeneracy of  $Z(\cdot)$  assumption is that  $|\rho_{st}(U) \leq C_1(Z) \cdot |\rho_{st}(U)|$  for any  $s, t$ . Hence, by the choice of the subdivision, and notice that  $G$  is a constant on each  $V_p$ ,

$$\begin{aligned}
& \frac{1}{2} |(\mathcal{F}_\rho(G)(x))_t - (H(G)(x))_t| \\
&= \left| \text{Re} \sum_{p=1}^q \sum_{s=1}^m \int_{V_p} e^{-i\langle x, \xi \rangle} g_s(\xi) - e^{-i\langle x, \xi_p \rangle} g_s(\xi_p) d\rho_{st}(\xi) \right| \\
&= \left| \text{Re} \sum_{p=1}^q \sum_{s=1}^m g_s(\xi_p) \int_{V_p} e^{-i\langle x, \xi \rangle} - e^{-i\langle x, \xi_p \rangle} d\rho_{st}(\xi) \right| \\
&\leq \epsilon \cdot \left( \sum_{s=1}^m \|\phi_s\|_{L^1(\mathbb{R}^n)} \right)^{-1} \cdot \sum_{p=1}^q \sum_{s=1}^m |g_s(\xi_p)| |\rho_{st}(V_p)| \\
&\leq \epsilon \cdot \left( \sum_{s=1}^m \|\phi_s\|_{L^1(\mathbb{R}^n)} \right)^{-1} \cdot C_1(Z) \sum_{p=1}^q \sum_{s=1}^m |g_s(\xi_p) \rho_{st}(V_p)| \\
&= \epsilon \cdot \left( \sum_{s=1}^m \|\phi_s\|_{L^1(\mathbb{R}^n)} \right)^{-1} \cdot C_1(Z) \sum_{l=1}^m \sum_{V_p^l} \sum_{s=1}^m |c_{ls}(V_p^l) \rho_{st}(V_p^l)| \cdot |\phi_l(\xi_p^l)| \cdot |V_p^l| \\
&\leq \epsilon \cdot \left( \sum_{s=1}^m \|\phi_s\|_{L^1(\mathbb{R}^n)} \right)^{-1} \cdot C_1(Z) C_2(Z) \sum_{l=1}^m \sum_{V_p^l} \cdot |\phi_l(\xi_p^l)| \cdot |V_p^l| \\
&\leq 2\epsilon \cdot C_1(Z) C_2(Z).
\end{aligned} \tag{D.34}$$

One can do similar estimates on  $|\nabla[(\mathcal{F}_\rho(G)(x))_t - (H(G)(x))_t]|$ . Hence, there is a positive constant  $C(C_1(Z), C_2(Z))$  such that

$$\|(\mathcal{F}_\rho(G)(x))_t - (H(G)(x))_t\|_{C^1(B)} \leq C(C_1(Z), C_2(Z)) \cdot \epsilon. \tag{D.35}$$

And then

$$\|(\mathcal{F}_\rho(G)(x))_t - p_t(x)\|_{C^1(B)} \leq C(\|p_t\|_{C^1(B)}, |y_t|, C_1(Z), C_2(Z)) \cdot \epsilon. \tag{D.36}$$

Since  $C^1(B, \mathbb{R}^m)$  can be  $C^1$ -approximated by polynomial vectors  $p(x)$ , we finish the proof.  $\square$

For the positivity of the limiting constant  $\nu_F$  in Theorem 3.1, notice that by Lemma 3.3, we have that for all  $R > 0$ ,

$$\frac{\mathcal{E}(N(R; F))}{|C_R|} \leq \nu_F. \tag{D.37}$$

Hence, we only need to show that there is some large  $R = R_0$  such that  $\mathcal{P}(N(R_0; F) > 0) > 0$ .

According to (D.16) (also see Proposition 5.2 of [SW19]), together with our Theorem D.3 and  $\mathcal{H}_0(\rho) = \mathcal{H}(F)$ , for any  $\epsilon > 0$  and any  $G \in C^1(\mathbb{R}^n, \mathbb{R}^m)$ ,

$$(D.38) \quad \mathcal{P}(\|F - G\|_{C^1(B_{10}(0))} < \epsilon) > 0,$$

where  $B_{10}(0)$  is the ball with center 0 and radius 10 in  $\mathbb{R}^n$ . We consider

$$(D.39) \quad G = (g_1, \dots, g_m) \equiv (x_1, \dots, x_{m-1}, 1 - |x_m|^2 - |x_{m+1}|^2 - \dots - |x_n|^2),$$

so that  $Z(G)$  is a unit  $\mathbb{S}^{n-m}$  contained in the linear subspace of coordinates  $x_m, \dots, x_n$ . Hence, there is an  $\epsilon = \epsilon(n, m)$  chosen in Lemma B.1 such that

$$(D.40) \quad \mathcal{P}(N(10; F) \geq 1) \geq \mathcal{P}(\|F - G\|_{C^1(C_{B_{10}(0)})} < \epsilon) > 0.$$

This shows the positivity of  $\nu_F$ .

Moreover, under our assumptions, by Remark B.2, the component  $\gamma$  of  $Z(F)$  in  $B_{10}(0)$  is  $C^1$ -isotopic to the component of  $Z(G)$ , which is  $\mathbb{S}^{n-m}$  for this  $G$ . But actually, one can replace  $G$  here by any  $G \in C^1(B_{10}(0), \mathbb{R}^m)$  such that  $Z(G) \cap B_1(0) \neq \emptyset$  and 0 is a regular value for  $G$ . For similar reason, one can show that for any diffeomorphism type  $H$  (or  $C^1$ -isotopic type  $H$ ) which can be realized as a regular connected component of some  $Z(G)$  in  $B_1(0)$ , we always have that

$$(D.41) \quad \mathcal{P}(N(10; F, H) \geq 1) \geq \mathcal{P}(\|F - G\|_{C^1(C_{B_{10}(0)})} < \epsilon) > 0.$$

**Remark D.4.** We can also show the positive probability for those  $\nu_F, \nu_{F,H}$  in the case that  $F$  consists of independent  $f_i, i = 1, \dots, m$ , and each  $f_i$  is a frequency 1 random wave on  $\mathbb{R}^n$ , i.e.,

$$(D.42) \quad \mathcal{E}(f_{i_1}(u)f_{i_2}(v)) = \delta_{i_1 i_2} \cdot \int_{\mathbb{S}^{n-1}} e^{i\langle u-v, \xi \rangle} d\mathcal{H}^{n-1}(\xi),$$

and solves

$$(D.43) \quad \Delta f_i + f_i = 0.$$

In this case, one can use the similar proof for Theorem 1 in [CS19], which is in their section 3. The rough idea is like the following. We consider an arbitrary  $(n-m)$ -dimensional connected closed submanifold  $\gamma$  of  $\mathbb{R}^{n+q}$  fully contained in  $B_1(0)$ , and  $\gamma$  is also realized as a regular connected component for zero sets of some  $C^1$ -smooth function  $G$ , i.e.,  $\gamma \subset Z(G) \cap B_1(0)$ , and  $\nabla G$  is of full rank on  $\gamma$ . Our goal is to modify this  $\gamma$  such that it is  $C^1$ -isotopic to another  $\tilde{\gamma}$  which can be expressed as the transversal intersection of  $m$  closed analytic hypersurfaces, which also enclose  $m$  bounded domains with the same first Dirichlet eigenvalue.

Denote that  $G = (g_1, \dots, g_m)$  and assume that 0 is a regular value of each  $g_i$ , otherwise one just replace each  $g_i$  by some  $g_i + \epsilon$  with  $\epsilon$  small, and get a new component  $C^1$ -isotopic to the original  $\gamma$ . We do not know whether each  $Z(g_i)$  is a closed hypersurface yet. But since, at least,  $Z(g_i)$  is a regular hypersurface for each  $i$ , one can replace each  $g_i$  by  $\tilde{g}_i \equiv g_i(x)^2 + \epsilon_i |x|^2 - \delta_i^2$  for very small  $\epsilon_i$  and very small  $\delta_i$ , such that 0 is a regular value of  $\tilde{g}_i$  by Sard's theorem. Notice that  $Z(\tilde{g}_i) \subset B_{\delta_i/\epsilon_i}(0)$ , hence, we get a new component  $\tilde{\gamma}$  realized as

a part of the intersection of those  $Z(\tilde{g}_i)$  with each  $Z(\tilde{g}_i)$  bounded, and  $\tilde{\gamma}$  is also  $C^1$ -isotopic to  $\gamma$  since  $\epsilon_i$  and  $\delta_i$  are small. We can also assume that each  $\tilde{g}_i$  is  $C^\infty$ -smooth.

Assume that  $\tilde{\gamma} = \cap_i c_i$ , for each  $c_i$  a bounded connected component of  $Z(\tilde{g}_i)$ , which can also be regarded as a closed hypersurface. Let  $A_i$  be the bounded component of  $\mathbb{R}^n \setminus c_i$ . Now, for each  $A_i$ , one can make smooth connected sum with long and thin necks to some large balls  $B_i$  and avoid touching  $\tilde{\gamma}$ . We do these connected sums and get new bounded sets  $\mathcal{A}_i$ , such that all these  $\{\mathcal{A}_i\}_{i=1}^m$  have the same first Dirichlet eigenvalue  $\lambda$ . This same first Dirichlet eigenvalue property is possible, because the first Dirichlet eigenvalue for a given domain is always larger than the first eigenvalues of domains which contain it, and we can adjust the sizes of balls  $B_i$  to let  $\mathcal{A}_i$  have the same first eigenvalue. And  $\tilde{\gamma}$  now becomes a part of the transversal intersection of those  $\mathcal{A}_i$ .

For each  $\mathcal{A}_i$ , we assume that  $\partial\mathcal{A}_i$  is realized as a component of the regular zero of another smooth function  $\bar{g}_i$ . Approximate each  $\bar{g}_i$  by another analytic function  $\mathbf{g}_i$  defined in a neighbourhood of  $\partial\mathcal{A}_i$ , we see that the set  $\{\mathbf{g}_i = 0\}$  has a bounded component  $\tilde{c}_i$  which is very close to  $\partial\mathcal{A}_i$ , and these  $\tilde{c}_i$  form another transversal intersection  $\tilde{\gamma}_1$ , which is  $C^1$ -isotopic to  $\tilde{\gamma}$ . The existence of such an analytic  $\mathbf{g}_i$  is by the same arguments as proving Theorem 1 in page 7 of [CS19], where they used Whitney's approximation Theorem.

The new problem here is that  $\tilde{\mathcal{A}}_i$ , the bounded component of  $\mathbb{R}^n \setminus \tilde{c}_i$ , may not have the same first Dirichlet eigenvalue. To fix this, since  $\mathcal{A}_i$  have the same first Dirichlet eigenvalue, and each  $\mathcal{A}_i$  is very close to  $\tilde{\mathcal{A}}_i$ , we may scale  $\tilde{\mathcal{A}}_i$  a little, or equivalently, replace  $\mathbf{g}_i(x)$  by  $\mathbf{g}_i(\eta_i x)$  for some  $\eta_i$  close to 1, and this small scaling will not affect the transversal intersection  $\tilde{\gamma}_1$  up to another  $C^1$ -isotopy.

To summarize, we get a  $\tilde{\gamma}_1$  in the same  $C^1$ -isotopy class, which can also be expressed as the transversal intersection of analytic hypersurfaces  $\tilde{c}_i = \partial\tilde{\mathcal{A}}_i$ , where all these  $\tilde{\mathcal{A}}_i$  are bounded and have the same first Dirichlet eigenvalue.

After these, the remaining arguments are exactly the same as the proof in page 8 of [CS19].

#### APPENDIX E. THE TWO-POINT CONDITIONS (A3) IN DEFINITION 2.1 AND (B4) IN DEFINITION 4.2

As we mentioned previously, we can use (A1) and (A2) to deduce (A3) in Definition 2.1. Let us formulize it more specifically. The idea is inspired by Proposition 6 and Proposition 7 of [CH20]. Set the two-point  $m \times m$  covariance matrix  $\text{Cov}(x, y)$  by

$$(E.1) \quad (\text{Cov}(x, y))_{i_1 i_2} \equiv \mathcal{E}(f_{i_1}(x) f_{i_2}(y)), \quad i_1, i_2 = 1, \dots, m.$$

**Theorem E.1.** *Suppose that  $F : C_{R+1} \rightarrow \mathbb{R}^m$  satisfies  $(R; M, k_1)$ -assumptions in Definition 2.1, then there is a  $\delta = \delta(n, m, M, k_1) \in (0, 1)$  such that when  $x \in C_R$  and  $\|x - y\| \leq \delta$ , we have that for any  $\eta \in \mathbb{S}^{m-1}$ ,*

$$(E.2) \quad \eta^T \cdot (\text{Cov}(y, y) - \text{Cov}(y, x) \text{Cov}(x, x)^{-1} \text{Cov}(x, y)) \cdot \eta \geq \frac{k_1}{2} \cdot \|y - x\|^2.$$



In particular, there is a  $k_2 = k_2(n, m, M, k_1) > 0$  such that when  $x \in C_R$  and  $\|x - y\| < \delta$ , then

$$(E.3) \quad p_{(F(x), F(y))}(0, 0) \leq \frac{k_2}{\|x - y\|^m},$$

which also implies that the joint distribution for  $(F(x), F(y))$  is non-degenerate if  $\|x - y\| \leq \delta$ .

*Proof.* Notice that (E.3) follows from (E.2) by elementary row operations in calculating the determinant of the covariance kernel for the joint distribution  $(F(x), F(y))$ . To prove (E.2) we fix an arbitrary  $x \in C_R$ . For any  $w = (w_1, \dots, w_n)^T \in \mathbb{S}^{n-1}$ , we define an  $m \times m$  symmetric matrix-valued function on  $[-1, 1] \subset \mathbb{R}$ :

$$(E.4) \quad k_w(t) \equiv \text{Cov}(x + t \cdot w, x + t \cdot w) - \text{Cov}(x + t \cdot w, x) \text{Cov}(x, x)^{-1} \text{Cov}(x, x + t \cdot w).$$

Since  $F$  satisfies (A1) in Definition 2.1,  $\text{Cov}(x, x)$  is invertible and hence  $k_w(t)$  is well defined on  $[-1, 1]$ . It is easy to see that  $k_w(0) = 0$  and  $k'_w(0) = 0$ . Then,

$$(E.5) \quad \begin{aligned} (k''_w(t))_{i_1 i_2} &= \mathcal{E}((\nabla_{w,w}^2 f_{i_1})(x + tw) f_{i_2}(x + tw)) + \mathcal{E}(f_{i_1}(x + tw) (\nabla_{w,w}^2 f_{i_2})(x + tw)) \\ &\quad + 2\mathcal{E}((\nabla_w f_{i_1})(x + tw) (\nabla_w f_{i_2})(x + tw)) \\ &\quad - \sum_{p,q} \mathcal{E}((\nabla_{w,w}^2 f_{i_1})(x + tw) f_p(x)) (\text{Cov}(x, x)^{-1})_{pq} (\text{Cov}(x, x + t \cdot w))_{qi_2} \\ &\quad - \sum_{p,q} (\text{Cov}(x + t \cdot w, x))_{i_1 p} (\text{Cov}(x, x)^{-1})_{pq} \mathcal{E}(f_q(x) (\nabla_{w,w}^2 f_{i_2})(x + tw)) \\ &\quad - 2 \sum_{p,q} \mathcal{E}((\nabla_w f_{i_1})(x + tw) f_p(x)) \cdot (\text{Cov}(x, x)^{-1})_{pq} \cdot \mathcal{E}(f_q(x) (\nabla_w f_{i_2})(x + tw)). \end{aligned}$$

By Cauchy's mean value theorem, we have that for any  $i_1, i_2 = 1, \dots, m$ , there is a  $\xi_{i_1 i_2}$  between 0 and  $t$ , such that

$$(E.6) \quad (k_w(t))_{i_1 i_2} = \frac{t^2}{2} \cdot (k''_w(\xi_{i_1 i_2}))_{i_1 i_2}.$$

Furthermore, notice that  $F$  is  $C^3$ -smooth, by assumption (A2), we have that

$$(E.7) \quad |(k''_w(\xi_{i_1 i_2}))_{i_1 i_2} - (k''_w(0))_{i_1 i_2}| \leq \sqrt{\xi_{i_1 i_2}} \cdot C_1 \leq \sqrt{t} \cdot C_1$$

for some positive constant  $C_1 = C_1(n, m, M)$ . We set the remaining term as the symmetric matrix  $R$  with elements  $R_{i_1 i_2} \equiv t^{-1/2} \cdot [(k''_w(\xi_{i_1 i_2}))_{i_1 i_2} - (k''_w(0))_{i_1 i_2}]$ , so that

$$(E.8) \quad k_w(t) = \frac{t^2}{2} (k''_w(0) + \sqrt{t} R).$$

For the expression of  $k''_w(0)$ , we set  $t = 0$  in (E.5) and then get

$$(E.9) \quad \begin{aligned} (k''_w(0))_{i_1 i_2} &= 2[\mathcal{E}((\nabla_w f_{i_1})(x) (\nabla_w f_{i_2})(x)) \\ &\quad - \sum_{p,q} \mathcal{E}((\nabla_w f_{i_1})(x) f_p(x)) \cdot (\text{Cov}(x, x)^{-1})_{pq} \cdot \mathcal{E}(f_q(x) (\nabla_w f_{i_2})(x))]. \end{aligned}$$

For any  $\eta = (\eta_1, \dots, \eta_m)^T \in \mathbb{S}^{m-1}$ , we let

$$(E.10) \quad v_\eta = \sum_{i=1}^m \eta_i ((\nabla_w f_i)(x)) = \sum_{i=1}^m \sum_{j=1}^n \eta_i w_j ((\partial_j f_i)(x)).$$

Notice that  $\text{Cov}(x, x)$  is a positive definite matrix, so it can be written as the square of another positive definite matrix, i.e.  $\text{Cov}(x, x) = B^2$  and  $B$  is positive definite. Set the Gaussian vector  $G = (g_1, \dots, g_m) \equiv F(x)^T \cdot B^{-1}$  so that  $\mathcal{E}(g_{i_1} g_{i_2}) = \delta_{i_1 i_2}$ . And then

$$(E.11) \quad \begin{aligned} \frac{1}{2} \eta^T \cdot k_w''(0) \cdot \eta &= \mathcal{E}(v_\eta^2) - \sum_p \mathcal{E}(v_\eta g_p) \cdot \mathcal{E}(g_p v_\eta) \\ &= \mathcal{E} \left[ \left( v_\eta - \sum_p g_p \cdot \mathcal{E}(v_\eta g_p) \right)^2 \right]. \end{aligned}$$

Notice that  $v_\eta$  is a linear combination of  $\nabla F(x)$  terms while  $\sum_p g_p \cdot \mathcal{E}(v_\eta g_p)$  is a linear combination of  $F(x)$  terms. Since the joint distribution of  $(F(x), \nabla F(x))$  is non-degenerate and by (A1) of Definition 2.1, we see that

$$(E.12) \quad \mathcal{E} \left[ \left( v_\eta - \sum_p g_p \cdot \mathcal{E}(v_\eta g_p) \right)^2 \right] \geq k_1 \cdot \left| \sum_{i=1}^m \sum_{j=1}^n (\eta_i w_j)^2 \right| = k_1,$$

where for the inequality, we only collect the coefficients for  $\nabla F(x)$  terms. Hence, by (E.8), there is a  $\delta = \delta(n, m, M, k_1) \in (0, 1)$  such that when  $|t| \leq \delta$ , we have that

$$(E.13) \quad \eta^T \cdot k_w(t) \cdot \eta = \frac{t^2}{2} \eta^T \cdot (k_w''(0) + \sqrt{t}R) \cdot \eta \geq \frac{t^2}{2} \cdot k_1.$$

□

**Remark E.2.** The Theorem E.1 gives the upper bound for the density  $p_{(F(x), F(y))}(0, 0)$  when  $\|x - y\| < \delta$  but  $x \neq y$ . If we know that  $(F(x), F(y))$  is always non-degenerate when  $x \neq y$  and  $x, y \in C_{R+1}$ , then condition (A3) in Definition 2.1 will hold automatically, and the  $k_2$  depends on the covariance kernel for the joint distribution of  $(F(x), F(y))$  when  $\|x - y\| > \delta$ .

For the condition (B4), in most of applications or concrete examples like Kostlan's ensemble or spherical harmonics ensembles, first, the convergence for kernels  $\{K_{x,L}\}_{L \in \mathcal{L}}$  is in  $C^k$ , i.e., for all  $R > 0$ ,

$$(E.14) \quad \lim_{L \rightarrow \infty} \|K_{x,L}(u, v) - K_x(u - v)\|_{C^k(C_{R+1} \times C_{R+1})} = 0.$$

In this case, the conditions (B1), (B2) and (B4) will hold if for all compact set  $Q \subset U$ , and  $x \in Q$ ,  $K_x(u - v)$  satisfies a uniform  $(R; M(Q), k_1(Q), k_2(x))$ -assumptions. In particular, for (B4), one way to verify it is that if we know for all  $x \in U$  and  $R > 0$ ,

$$(E.15) \quad \lim_{L \rightarrow \infty} \sup_{u, v \in C_{R+1}} |K_{x,L}(u, v) - K_x(u - v)| = 0,$$

and  $K_x$ , or equivalently  $F_x$ , satisfies (A3) in Definition 2.1, then when  $\|u - v\| > \delta$ , the quantitative two-point nondegeneracy for  $K_{x,L}$ , or equivalently  $F_{x,L}$ , holds from this uniform convergence, while when  $\|u - v\| \leq \delta$ , one can just apply Theorem E.1.

## APPENDIX F. CURVATURES COMPUTATIONS

In this section, we give some preliminary computations for curvatures involved in the two geometric inequalities, Theorem 2.5 and Theorem 2.9. More precisely, we compute second fundamental forms and mean curvatures for submanifolds obtained from zero sets.

Let  $X \subset \mathbb{R}^{n+q}$  be an  $n$ -dimensional  $C^3$ -manifold embedded in  $\mathbb{R}^{n+q}$  for some  $q \geq 0$ . A special case is when  $q = 0$  and  $X = \mathbb{R}^n$ . And we let  $M$  be an  $(n - m)$ -dimensional manifold embedded in  $X$ , and hence  $M$  is also a submanifold of  $\mathbb{R}^{n+q}$ . We assume that  $M$  is realized as a regular part of the zero set of a non-degenerate  $m$ -dimensional  $C^2$ -vector field  $F$  on  $X$ , i.e., for  $F = (f_1, \dots, f_m)$ ,  $M$  is a connected component of  $F^{-1}(0) = Z(F)$ , and on  $M$ ,  $\nabla^X F$ , the gradient of  $F$  with respect to the Riemannian connection on  $X$ , is of full rank.

Choose a point  $p_0 \in M$ , assume that  $p = (t_1, \dots, t_{n-m})$  is a local coordinate on  $M$  around  $p_0 = (0, \dots, 0)$ . Denote the embedding from  $M$  into  $X$  as  $\varphi$ , and choose  $x = (x_1, \dots, x_n)$  as a local coordinate on  $X$  around  $x_0 = \varphi(p_0)$ . So, one may write  $\varphi = (\varphi^1, \dots, \varphi^n)$ . We denote the embedding from  $X$  into  $\mathbb{R}^{n+q}$  to be  $\Phi$ . If we view all these objects as subsets in  $\mathbb{R}^{n+q}$ , then  $p_0$ ,  $x_0 = \varphi(p_0)$ , and  $y_0 = \Phi(x_0) = \Phi(\varphi(p_0))$  are actually the same point.

Our aim in this section is to compute the second fundamental form and mean curvature of  $M$  as a submanifold of  $\mathbb{R}^{n+q}$  at the point  $y_0 = \Phi(\varphi(p_0))$ . Since we only consider the value of tensors at a fixed point, for simplicity, let us assume that  $(t_1, \dots, t_{n-m})$  and  $(x_1, \dots, x_n)$  are local normal coordinates on  $M$  and  $X$  at the points  $p_0, x_0 = \varphi(p_0)$  respectively. That is, at  $x_0$ ,

$$(F.1) \quad \langle \partial_{x_l} \Phi, \partial_{x_k} \Phi \rangle_{\mathbb{R}^{n+q}} = \delta_{lk},$$

for  $\langle \cdot, \cdot \rangle_{\mathbb{R}^{n+q}}$  the standard inner product in  $\mathbb{R}^{n+q}$ , together with that

$$(F.2) \quad \partial_{x_l x_k}^2 \Phi \in T_{y_0}^\perp X, \quad \forall l, k = 1, \dots, n,$$

where  $T_{y_0}^\perp X$  is the normal vector space of  $X$  at  $y_0 = \Phi(x_0)$  in  $\mathbb{R}^{n+q}$ . And at  $p_0$ ,

$$(F.3) \quad \sum_l (\partial_{t_i} \varphi^l)(\partial_{t_j} \varphi^l) = \delta_{ij},$$

together with that

$$(F.4) \quad \sum_l (\partial_{t_i t_j}^2 \varphi^l)(\partial_{t_s} \varphi^l) = 0, \quad \forall i, j, s = 1, \dots, n - m.$$

We first consider the second fundamental form  $\mathbb{II} = \mathbb{II}^M$  of  $M$  as a submanifold of  $\mathbb{R}^{n+q}$  at  $y_0 = \Phi(\varphi(p_0))$ , which is defined as

$$(F.5) \quad \langle \mathbb{II}_{ij}, \mathbf{n} \rangle \equiv \left\langle \frac{\partial^2 (\Phi \circ \varphi)}{\partial t_i \partial t_j} (p_0), \mathbf{n} \right\rangle_{\mathbb{R}^{n+q}},$$

where  $\mathbf{n} \in T_{y_0}^\perp M$  is a unit normal vector of  $M$  at  $y_0$  in  $\mathbb{R}^{n+q}$ . Let us compute these derivatives.

$$(F.6) \quad \partial_{t_j}(\Phi \circ \varphi) = \sum_l (\partial_{x_l} \Phi)(\partial_{t_j} \varphi^l),$$

and

$$(F.7) \quad \partial_{t_i t_j}^2(\Phi \circ \varphi) = \sum_{l,k} (\partial_{x_l x_k}^2 \Phi)(\partial_{t_j} \varphi^l)(\partial_{t_i} \varphi^k) + \sum_l (\partial_{x_l} \Phi)(\partial_{t_i t_j}^2 \varphi^l).$$

If  $\mathbf{n} \in T_{y_0}^\perp X$ , then  $\mathbf{n} \perp (\partial_{x_l} \Phi)$ , and hence

$$(F.8) \quad |\langle \mathbf{II}_{ij}, \mathbf{n} \rangle| = \left| \sum_{l,k} \langle (\partial_{x_l x_k}^2 \Phi), \mathbf{n} \rangle (\partial_{t_j} \varphi^l)(\partial_{t_i} \varphi^k) \right| \leq \| \mathbf{II}^X(y_0) \|.$$

The last inequality is because of (F.3), which says that  $\partial_{t_i} \varphi$  and  $\partial_{t_j} \varphi$  are two unit vectors, and hence the value is bounded by the norm of the matrix, which is  $\| \mathbf{II}^X(y_0) \|$ , the norm of the second fundamental form of  $X$  as a submanifold of  $\mathbb{R}^{n+q}$  at  $y_0$ . This is a constant only depend on  $X$  and the embedding  $\Phi$ , so we denote an upper bound of this constant by  $C(X)$ .

If  $\mathbf{n} \in T_{y_0} X \cap T_{y_0}^\perp M$ , i.e., vectors tangent to  $X$  but normal to  $M$  at  $y_0$  in  $\mathbb{R}^{n+q}$ , we have that

$$(F.9) \quad |\langle \mathbf{II}_{ij}, \mathbf{n} \rangle| = \left| \sum_l \langle (\partial_{x_l} \Phi), \mathbf{n} \rangle (\partial_{t_i t_j}^2 \varphi^l) \right|,$$

which is because of (F.2),  $\partial_{x_l x_k}^2 \Phi \perp \mathbf{n}$ . Now, let us use the fact that  $M$  is realized as part of  $Z(F)$  to do more calculations and estimate (F.9). Since  $f_\alpha(\varphi) = 0$  for  $\alpha = 1, \dots, m$ , we differentiate these equations and get

$$(F.10) \quad \sum_l (\partial_{x_l} f_\alpha)(\partial_{t_j} \varphi^l) = 0,$$

and

$$(F.11) \quad \sum_{l,k} (\partial_{x_l x_k}^2 f_\alpha)(\partial_{t_j} \varphi^l)(\partial_{t_i} \varphi^k) + \sum_l (\partial_{x_l} f_\alpha)(\partial_{t_i t_j}^2 \varphi^l) = 0.$$

Since  $\nabla^X F$  is of full rank on  $M$ , we see that these  $\nabla^X f_\alpha \equiv \sum_l (\partial_{x_l} f_\alpha)(\partial_{x_l} \Phi)$ , actually form a basis of  $T_{y_0} X \cap T_{y_0}^\perp M$ , and hence we take an  $m \times m$  symmetric matrix  $T = (T^{\alpha\beta})$  such that these

$$(F.12) \quad \mathbf{n}_\alpha = \sum_\beta T^{\alpha\beta} \sum_l (\partial_{x_l} f_\beta)(\partial_{x_l} \Phi)$$

are orthonormal, i.e.,  $\langle \mathbf{n}_{\alpha_1}, \mathbf{n}_{\alpha_2} \rangle = \delta_{\alpha_1 \alpha_2}$ . On the other hand,

$$(F.13) \quad \langle \mathbf{n}_{\alpha_1}, \mathbf{n}_{\alpha_2} \rangle = \sum_{\beta_1, \beta_2} T^{\alpha_1 \beta_1} T^{\alpha_2 \beta_2} \sum_l (\partial_{x_l} f_{\beta_1})(\partial_{x_l} f_{\beta_2}),$$

where if one uses matrices notations, the right hand side is  $TAT^T$ , for  $T^T$  the transpose of  $T$ , and  $A$  is the matrix consists of inner products of  $f_\beta$ , i.e.,  $A = (A_{\beta_1\beta_2}) = (\sum_l (\partial_{x_l} f_{\beta_1})(\partial_{x_l} f_{\beta_2}))$ . And hence,

$$(F.14) \quad T^T T = A^{-1}.$$

Then, according to (F.1), (F.9), and (F.11),

$$(F.15) \quad |\langle \mathbb{I}_{ij}, \mathbf{n}_\alpha \rangle| = \left| \sum_\beta T^{\alpha\beta} \sum_l (\partial_{x_l} f_\beta)(\partial_{t_j}^2 \varphi^l) \right| = \left| \sum_\beta T^{\alpha\beta} \sum_{l,k} (\partial_{x_l x_k}^2 f_\beta)(\partial_{t_j} \varphi^l)(\partial_{t_i} \varphi^k) \right|,$$

and then

$$(F.16) \quad \begin{aligned} \sum_\alpha |\langle \mathbb{I}_{ij}, \mathbf{n}_\alpha \rangle|^2 &= \left| \sum_\beta T^{\alpha\beta} \sum_{l,k} (\partial_{x_l x_k}^2 f_\beta)(\partial_{t_j} \varphi^l)(\partial_{t_i} \varphi^k) \right|^2 \\ &= \sum_\alpha \sum_{\beta_1, \beta_2} T^{\alpha\beta_1} T^{\alpha\beta_2} \left[ \sum_{l_1, k_1} (\partial_{x_{l_1} x_{k_1}}^2 f_{\beta_1})(\partial_{t_j} \varphi^{l_1})(\partial_{t_i} \varphi^{k_1}) \right] \left[ \sum_{l_2, k_2} (\partial_{x_{l_2} x_{k_2}}^2 f_{\beta_2})(\partial_{t_j} \varphi^{l_2})(\partial_{t_i} \varphi^{k_2}) \right] \\ &= \sum_{\beta_1, \beta_2} (A^{-1})_{\beta_1 \beta_2} \left[ \sum_{l_1, k_1} (\partial_{x_{l_1} x_{k_1}}^2 f_{\beta_1})(\partial_{t_j} \varphi^{l_1})(\partial_{t_i} \varphi^{k_1}) \right] \left[ \sum_{l_2, k_2} (\partial_{x_{l_2} x_{k_2}}^2 f_{\beta_2})(\partial_{t_j} \varphi^{l_2})(\partial_{t_i} \varphi^{k_2}) \right]. \end{aligned}$$

For an upper bound, one just notice again that by (F.3) that those  $\partial_{t_j} \varphi$  are unit vectors, and then

$$(F.17) \quad \sum_\alpha |\langle \mathbb{I}_{ij}, \mathbf{n}_\alpha \rangle|^2 \leq C(n, m) \cdot \|A^{-1}\| \cdot \|(\nabla^X)^2 F\|^2,$$

for some positive constant  $C(n, m)$ , where  $(\nabla^X)^2 F$  is the Hessian of  $F$  and those  $\|\cdot\|$  are matrices norms.

We denote  $\{\mathbf{n}_\mu\}$  as an orthonormal basis of  $T_{y_0}^\perp X$ , then the norm square of the second fundamental form at  $y_0$  is defined as

$$(F.18) \quad \|\mathbb{I}(y_0)\|^2 \equiv \sum_{i,j} \left( \sum_\alpha |\langle \mathbb{I}_{ij}, \mathbf{n}_\alpha \rangle|^2 + \sum_\mu |\langle \mathbb{I}_{ij}, \mathbf{n}_\mu \rangle|^2 \right).$$

We then summarize our results previously as the following theorem.

**Theorem F.1.** *The second fundamental form  $\mathbb{I}^M$  of  $M$  as a submanifold of  $\mathbb{R}^{n+q}$  satisfies that*

$$(F.19) \quad \|\mathbb{I}^M\|^2 \leq C(n, m, X) + C(n, m) \cdot \|A^{-1}\| \cdot \|(\nabla^X)^2 F\|^2,$$

where the matrix  $A$  consists of inner products of  $\nabla^X f_\alpha$  for  $\alpha = 1, \dots, m$ , and  $(\nabla^X)^2 F$  is the Hessian. The positive constant  $C(n, m, X)$  depends on  $X$  and the embedding  $\Phi : X \rightarrow \mathbb{R}^{n+q}$ . In particular, when  $X = \mathbb{R}^n$ , one can set  $C(n, m, X) = 0$ .

We remark that the mean curvature  $\mathbf{H}$  at  $y_0$  is defined as

$$(F.20) \quad \langle \mathbf{H}(y_0), \mathbf{n} \rangle \equiv \sum_i \langle \Pi_{ii}, \mathbf{n} \rangle,$$

for  $\mathbf{n} \in T_{y_0}^\perp M$ . And hence

$$(F.21) \quad \|\mathbf{H}(y_0)\|^2 \equiv \sum_\alpha \left| \sum_i \langle \Pi_{ii}, \mathbf{n}_\alpha \rangle \right|^2 + \sum_\mu \left| \sum_i \langle \Pi_{ii}, \mathbf{n}_\mu \rangle \right|^2 \leq C(n, m) \cdot \|\Pi^M(y_0)\|^2,$$

for some positive constant  $C(n, m)$ .

## REFERENCES

- [AW09] J. Azaïs and M. Wschebor. *Level sets and extrema of random processes and fields*. John Wiley & Sons, 2009.
- [B61] V. Bargmann. *On a Hilbert space of analytic functions and an associated integral transform part I*. Communications on pure and applied mathematics 14.3 (1961): 187-214.
- [B88] P. H. Bérard. *From vanishing theorems to estimating theorems: the Bochner technique revisited*. Bulletin of the American Mathematical Society 19.2 (1988): 371-406.
- [BP32] A. Bloch and G. Pólya. *On the roots of certain algebraic equations*. Proceedings of the London Mathematical Society 2.1 (1932): 102-114.
- [BS02] E. Bogomolny and C. Schmit. *Percolation model for nodal domains of chaotic wave functions*. Physical Review Letters 88.11 (2002): 114102.
- [C71] B. Y. Chen. *On a theorem of Fenchel-Borsuk-Willmore-Chern-Lashof*. Mathematische Annalen 194, no. 1 (1971): 19-26.
- [CH20] Y. Canzani and B. Hanin. *Local universality for zeros and critical points of monochromatic random waves*. Communications in Mathematical Physics 378, no. 3 (2020): 1677-1712.
- [CL57] S. S. Chern and R. K. Lashof. *On the total curvature of immersed manifolds*. American journal of mathematics 79.2 (1957): 306-318.
- [CL58] S. S. Chern and R. K. Lashof. *On the total curvature of immersed manifolds, II*. Michigan Math. J. 5.1 (1958): 5-12.
- [CS19] Y. Canzani and P. Sarnak. *Topology and nesting of the zero set components of monochromatic random waves*. Communications on Pure and Applied Mathematics 72.2 (2019): 343-374.
- [GS04] I. I. Gikhman and A. V. Skorokhod. *The Theory of Stochastic Processes, I*, in: Classics in Mathematics, Springer, Berlin, 2004, Translated from the Russian, by S. Kotz., Corrected Printing of the First Edition.
- [GW14] D. Gayetand and J. Y. Welschinger. *Lower estimates for the expected Betti numbers of random real hypersurfaces*. Journal of the London Mathematical Society 90.1 (2014): 105-120.
- [GW16] D. Gayetand and J. Y. Welschinger. *Betti numbers of random real hypersurfaces and determinants of random symmetric matrices*. Journal of the European Mathematical Society 18.4 (2016): 733-772.
- [K43] M. Kac. *On the average number of real roots of a random algebraic equation*. Bulletin of the American Mathematical Society 49.4 (1943): 314-320.
- [KW18] P. Kurlberg and I. Wigman. *Variation of the Nazarov-Sodin constant for random plane waves and arithmetic random waves*. Advances in Mathematics 330 (2018): 516-552.
- [L16] T. Letendre. *Expected volume and Euler characteristic of random submanifolds*. Journal of Functional Analysis 270.8 (2016): 3047-3110.
- [LL15] A. Lerario and E. Lundberg. *Statistics on Hilbert's 16th problem*. International Mathematics Research Notices 2015.12 (2015): 4293-4321.

- [NS09] F. Nazarov and M. Sodin. *On the number of nodal domains of random spherical harmonics*. American Journal of Mathematics 131.5 (2009): 1337-1357.
- [NS16] F. Nazarov and M. Sodin. *Asymptotic Laws for the Spatial Distribution and the Number of Connected Components of Zero Sets of Gaussian Random Functions*. Journal of Mathematical Physics, Analysis, Geometry 12, no. 3 (2016): 205-278.
- [S16] M. Sodin. *Lectures on random nodal portraits*. In Probability and statistical physics in St. Petersburg, volume 91 of Proc. Sympos. Pure Math., pages 395-422. Amer. Math.Soc., Providence, RI, 2016.
- [SW19] P. Sarnak and I. Wigman. *Topologies of Nodal Sets of Random Band-Limited Functions*. Communications on pure and applied mathematics 72.2 (2019): 275-342.
- [W21] I. Wigman. *On the expected Betti numbers of the nodal set of random fields*. Analysis & PDE 14, no. 6 (2021): 1797-1816.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, 251 MERCER STREET,  
NEW YORK, NY10012, USA

*Email address:* malin@nyu.edu