Lattice Minors and Eulerian Posets

William Gustafson

May 4, 2022

Abstract

Deletion and contraction operations on lattices enriched with a generating set were introduced by the author in [10]; and from these a notion of minors. When applied to the lattice of flats of a graph, these correspond to the same operations applied to the graph when the vertices are labeled and the edges unlabeled, followed by simplifying the resultant graph. We introduce a partial ordering of the minors of a given generator enriched lattice which we call the minor poset. We show that the minor poset of any generator enriched lattice is isomorphic to the face poset of a regular CW sphere, and in particular, is Eulerian. Furthermore, inequalities for the coefficients of cd-indices of minor posets are established. We also give a forbidden minor characterization of the generator enriched lattices whose minor poset is itself a lattice. Minor posets admit a decomposition into a disjoint union of Boolean algebras. We use this decomposition to derive formulas for the rank generating function of minor posets associated to geometric lattices with generating set consisting of the join irreducibles, and for minor posets associated to a class generalizing distributive lattices.

1 Introduction

Let G be a finite graph with labeled vertices and unlabeled edges. The minors of G, that is, graphs obtained by deleting and contracting edges of G, inherit a vertex labeling. These vertex labeled minors of G may be partially ordered by defining $H_1 \leq H_2$ when H_1 is a minor of H_2 . When G admits a planar embedding where all the vertices lie on a circle, this partial order on the simple minors of G arises as a lower interval in the well-studied uncrossing poset; see [11, 15, 16]. In [11] Hersh and Kenyon show the uncrossing poset is isomorphic to the face poset of a regular CW complex. Hence, the poset of minors of a graph admitting a planar embedding with all vertices on a circle is the face poset of a regular CW sphere, and in particular, is Eulerian.

It is natural to ask whether this result holds for larger classes of graphs, or even if this construction may be extended to matroids. Given a graph, the minors of the lattice of flats (defined in Section 2.1), are in bijection with the vertex labeled minors of the graph ([10, Theorem 3.9]). This result generalizes to the setting of polymatroids ([10, Theorem 3.11]). In this work we begin the study of minor posets of generator enriched lattices, which generalize the notion of posets of simple vertex labeled minors discussed above.

In Section 2 generator enriched lattices and the notions of deletion and contraction on generator enriched lattices are defined, and some basic results are discussed. In Section 3 the minor poset is introduced. The class of generator enriched lattices for which the minor poset is itself a lattice is characterized in terms of five forbidden minors in Proposition 3.3.2 and Theorem 3.3.6. A decomposition of the minor poset into Boolean algebras is given in Theorem 3.4.1. This decomposition is used to give expressions for the rank generating function of minor posets of geometric lattices with generating set consisting of the join irreducibles and of lattices with generating set consisting of the join irreducibles. See Theorems 3.4.3 and 3.4.5.

In Section 4 the minor poset of any generator enriched lattice is shown to be isomorphic to the face poset of a regular CW sphere; see Section 4.3.3. This is done via a construction using the zipping operation of Reading [18] in Theorem 4.3.1. Using this construction, in Corollaries 4.3.4 and 4.3.5 we derive inequalities between the coefficients of **cd**-indices of minor posets when there is a structure preserving surjection between the associated generator enriched lattices. In particular, we show that the **cd**-index of the n-dimensional cube is the coefficientwise maximum of **cd**-indices of minor posets of rank n + 1.

We discuss some open questions and future work in Section 5.

2 Lattice minors

2.1 Definitions

We begin by defining a few basic notions, which were introduced by the author in [10].

Definition 2.1.1. A generator enriched lattice is a pair (L, G) such that L is a finite lattice and $G \subseteq L \setminus \{\widehat{0}\}$ generates the lattice L via the join operation.

Given a generator enriched lattice (L, G) the elements of G will be referred to as generators of (L, G). Necessarily G must include the set of join irreducibles of L, which we denote by $\operatorname{irr}(L)$. When $G = \operatorname{irr}(L)$ the generator enriched lattice (L, G) is said to be minimally generated. Note that the generating set G may be empty. In this case the lattice L consists of a single element, say x, and we denote this generator enriched lattice as (x, \emptyset) as opposed to $(\{x\}, \emptyset)$.




Figure 1: In (a) is the Hasse diagram of a lattice L with $irr(L) = \{g, h, i\}$, and in (b) is the diagram of the associated minimally generated lattice (L, irr(L)). In (c) is the Hasse diagram of the Boolean algebra B_2 , which is also the diagram of the minimally generated lattice $(B_2, \{j, k\})$, and in (d) is the diagram of the generator enriched lattice $(B_2, \{j, k, \widehat{1}\})$.

Given a lattice L, let $H \subseteq L \setminus \{\widehat{0}\}$ and let $z \in L$ be an element such that z < h for all $h \in H$. Define the generator enriched lattice with generating set H and minimal element z to be

$$\begin{split} \langle H|z\rangle &= \left(\left\{z\vee\bigvee_{x\in X}x:X\subseteq H\right\},H\right)\\ &= \left(\left\{z\right\}\cup\left\{\bigvee_{x\in X}x:\emptyset\neq X\subseteq H\right\},H\right). \end{split}$$

Usually when listing H explicitly the set brackets will be omitted.

Generator enriched lattices will be depicted by diagrams analogous to Cayley graphs for groups. The diagram of a generator enriched lattice (L,G) is a directed graph with vertex set L. The edges are all pairs $(\ell,\ell\vee g)$ for $\ell\in L$ and $g\in G$ provided $\ell\neq\ell\vee g$. Similar to Hasse diagrams all diagrams of generator enriched lattices will be depicted with edges directed upwards. The diagram determines the generator enriched lattice. The minimal element $\widehat{0}$ is the unique source vertex and the generating set consists of all elements incident to $\widehat{0}$. Furthermore, the order relation is read off just as from a Hasse diagram, the relation $\ell_1 \leq \ell_2$ holds if and only if there is a directed path in the diagram from ℓ_1 to ℓ_2 . Figure 1 shows examples of diagrams of generator enriched lattices.

The structure preserving maps between generator enriched lattices are called strong maps.

Definition 2.1.2. Let (L,G) and (K,H) be two generator enriched lattices. A strong map is a map $f: L \to K$ that is join-preserving and satisfies $f(G) \subseteq H \cup \{\widehat{0}_K\}$. We use the notation $f: (L,G) \to (K,H)$ to denote a strong map f from (L,G) to (K,H).

A strong map $f:(L,G)\to (K,H)$ is said to be *injective* when it is injective as a map on the underlying lattices, and *surjective* when $f(G\cup\{\widehat{0}_L\})=H\cup\{\widehat{0}_K\}$. Two generator enriched lattices are said to be *isomorphic* when there is a strong bijection between them.

If both L and K are minimally generated geometric lattices then the definition of a strong map from L to K given here agrees with the usual definition of a strong map between simple matroids as introduced by Higgs in [12, pp. 1].

Let B_n denote the Boolean algebra of rank n, that is, the set of all subsets of $[n] = \{1, \ldots, n\}$ ordered by inclusion. For any generator enriched lattice (L, G) with n generators there is a strong surjection from $(B_n, \operatorname{irr}(B_n))$ onto (L, G). To construct such a map, label the elements of the generating set as $G = \{g_1, \ldots, g_n\}$. Let $\theta : B_n \to L$ be the map defined for $X \subseteq [n]$ by

$$\theta(X) = \bigvee_{x \in X} g_x.$$

This map θ will be referred to as the canonical strong map onto (L, G).

Definition 2.1.3. Given a generator enriched lattice (L,G) and $I \subseteq G$ the deletion by I is the generator enriched lattice

$$(L,G)\setminus I=\langle G\setminus I|\widehat{0}\rangle.$$

Set $i_0 = \bigvee_{i \in I} i$ and $J = \{g \vee i_0 : g \in G\} \setminus \{i_0\}$. The contraction of (L, G) by I is the generator enriched lattice

$$(L,G)/I = \langle J|i_0\rangle.$$

The restriction of (L,G) by I is the generator enriched lattice

$$(L,G)|_{I}=(L,G)\setminus (G\setminus I)=\langle G\cap I|\widehat{0}\rangle.$$

It will be convenient at times to instead index deletion and contractions by subsets of [n] or by elements of L. To this end given a labeling $G = \{g_1, \ldots, g_n\}$ of the generating set define deletions and contractions of (L, G) by $X \subseteq [n]$ as:

$$(L,G) \setminus X = (L,G) \setminus \{g_x : x \in X\},\$$

$$(L,G)/X = (L,G)/\{g_x : x \in X\}.$$

Deletions and contractions of (L,G) by an element $\ell \in L$ are defined by as:

$$(L,G) \setminus \ell = (L,G) \setminus \{g \in G : g \le \ell\},\$$
$$(L,G)/\ell = (L,G)/\{g \in G : g \le \ell\}.$$

Finally a minor of (L, G) is any generator enriched lattice that is the result of some sequence of deletions and contractions applied to (L, G). See Figure 2 for examples.

Remark 2.1.4. By definition the underlying lattice of a minor of (L, G) is a join subsemilattice of L. In general the underlying lattice of a minor of (L, G) may not be a sublattice of L. For example, consider the partition lattice Π_4 with minimal generating set

$$\operatorname{irr}(\Pi_4) = \{12/3/4, \ 13/2/4, \ 14/2/3, \ 1/23/4, \ 1/24/3, \ 1/2/34\}.$$

Deleting the atom 13/2/4 results in a minor that is not a sublattice of Π_4 . In said minor the meet of 123/4 and 134/2 is the minimal partition 1/2/3/4 as opposed to 13/2/4 when computed in L.




Figure 2: In (a) is the diagram of the face lattice of the square as a minimally generated lattice. In (b)-(d) are several minors.

Remark 2.1.5. Any interval of L is the underlying lattice of a minor of (L,G). If $a \leq b$ in L then the minor $(K,H) = (L/a)|_b$ has underlying lattice K = [a,b] of L. The example given in Remark 2.1.4 shows the converse is false, that in general not all minors of (L,G) have as underlying lattice an interval of L.

Example 2.1.6. The deletion and contraction operations of generator enriched lattices do not in general commute. For example, consider the generator enriched lattice (L,G) depicted in Figure 3. The contraction (L,G)/2 has a single generator $\widehat{1} = g_3 \vee g_1 = g_2 \vee g_3$ labeled both by 1 and 3. Thus, the minor $((L,G)/2) \setminus 1$ has a single element g_2 , while the minor $((L,G) \setminus 1)/2$ has a single generator, namely $\widehat{1} = g_3 \vee g_2$.




Figure 3: A generator enriched lattice (L, G), where $G = \{g_1, g_2, g_3\}$, for which deletions and contractions do not commute, along with the relevant minors.

2.2 Basic results

We record a few basic results for minors of generator enriched lattices. The proofs can be found in [10, Lemma 3.6, Lemma 3.7, Lemma 3.8].

Lemma 2.2.1 (Gustafson). Any minor of a generator enriched lattice (L,G) may be expressed as the result of a contraction followed by a deletion. Namely, a minor (K,H) of (L,G) may be expressed as $(K,H) = ((L,G)/\widehat{0}_K)|_H$.

Lemma 2.2.2 (Gustafson). For any generator enriched lattice (L,G) the minors are precisely generator enriched lattices of the form $\langle \ell \vee g_1, \ldots, \ell \vee g_k | \ell \rangle$ for $\ell \in L$ and $\{g_1, \ldots, g_k\} \subseteq G$ such that $g_j \not\leq \ell$ for $1 \leq j \leq k$.

Lemma 2.2.3 (Gustafson). If L is a geometric lattice then the minors of $(L, \operatorname{irr}(L))$ are the generator enriched lattices of the form $\langle \ell_1, \ldots, \ell_k | \ell \rangle$ such that $\ell_i \succ \ell \in L$ for $1 \leq i \leq k$. In particular, every minor of $(L, \operatorname{irr}(L))$ is minimally generated and geometric.

We present here a few results which give an alternate description of minors for minimally generated distributive lattices. Recall that the fundamental theorem of finite distributive lattices states that every finite distributive lattice L is isomorphic to the lattice of lower order ideals of the subposet irr(L) of L [5, Theorem 3, Chapter VIII].

Definition 2.2.4. Let P be a poset. An order minor of P is a pair (I, J) of disjoint subsets of P such that J is a lower order ideal of P.

When $(L, \operatorname{irr}(L))$ is the minimally generated lattice of lower order ideals of a poset P there is an implicit bijection between P and the generating set $\operatorname{irr}(L)$. Through this bijection deletions and contractions of $(L, \operatorname{irr}(L))$ may be indexed by subsets of P, just as deletions and contractions indexed by subsets of [n] are defined. The following bijection between order minors of the poset P and minors of the generator enriched lattice $(L, \operatorname{irr}(L))$ was given in $[10, \operatorname{Proposition} 3.14]$.

Proposition 2.2.5 (Gustafson). Let L be the lattice of lower order ideals of a poset P. The order minors of P and the minors of (L, irr(L)) are in bijection via the map

$$(I,J)\mapsto ((L,\operatorname{irr}(L))|_{I\cup J})/J.$$

Not only do the order minors of a poset index the minors of the minimally generated lattice of lower order ideals, the order minor also describes the isomorphism type of the corresponding lattice minor ([10, Proposition 3.15]).

Proposition 2.2.6 (Gustafson). Let P be a poset and L the lattice of lower order ideals of P, and let (I, J) be an order minor of P. The minor $((L, irr(L))|_{I \cup J})/J$ of (L, irr(L)) is isomorphic to the minimally generated lattice of lower order ideals of I.




Figure 4: The minor poset of a generator enriched lattice. The elements are depicted via their diagrams.

3 The minor poset

In this section a partial order structure on the set of minors of a given generator enriched lattice called the minor poset is studied. We begin with some basic results for minor posets in 3.1. In Section 3.2 we discuss a few operations on generator enriched lattices and their effects on the minor poset. In Section 3.3 a characterization is given of the generator enriched lattices for which the minor poset is itself a lattice. In Section 3.4 a decomposition theorem is presented which is used to derive expressions for the rank generating function of minor posets in special cases.

3.1 Basic results

Definition 3.1.1. Let (L,G) be a generator enriched lattice. The minor poset, denoted M(L,G), is the poset consisting of a unique minimal element \emptyset and the minors of (L,G) and with order relation defined by $(K_1,H_1) \leq (K_2,H_2)$ when (K_1,H_1) is a minor of (K_2,H_2) .

As an immediate observation, the lower interval $[\emptyset, (K, H)]$ in the minor poset M(L, G) is the minor poset M(K, H). See Figure 4 for an example of a minor poset.

Recall a ranked poset is said to be *thin* if all length 2 intervals are isomorphic to the Boolean algebra B_2 . The following lemma is used to show minor posets are thin and graded.

Lemma 3.1.2. Let (L,G) be a generator enriched lattice and let (K_1, H_1) and (K_2, H_2) be minors of (L,G) such that $(K_1, H_1) < (K_2, H_2)$ in the minor poset M(L,G). If $|H_2| - |H_1| = 2$ then the interval $[(K_1, H_1), (K_2, H_2)]$ of M(L,G) is isomorphic to the Boolean algebra B_2 .

Proof. The minor (K_1, H_1) may be presented as $((K_2, H_2)/I) \setminus J$ for some sets of generators I and J. We proceed by considering the different possibilities for I and J. When I is empty the set J must contain two elements, say j_1 and j_2 . Then $(K_1, H_2) = (K_2, H_2) \setminus \{j_1, j_2\}$ and the open interval $((K_1, H_1), (K_2, H_2))$ consists of the two minors $(K_2, H_2) \setminus \{j_1\}$ and $(K_2, H_2) \setminus \{j_2\}$.

Now consider the case where J is empty. In this case I may consist of either one element or two elements. First suppose $I = \{i_1, i_2\}$. As a further subcase assume $i_1 \not< i_2$ and $i_1 \not> i_2$. Since by assumption $|H_2| - |H_1| = 2$, every generator j of (K_1, H_1) corresponds to a unique generator i of (K_2, H_2) such that $i \lor i_1 \lor i_2 = j$. Due to this uniqueness no deletion of (K_2, H_2) has (K_1, H_1) as a minor. If $(K, H) = (K_2, H_2) \setminus \{j'\}$ then $j' \lor i_1 \lor i_2$ is not an element of $(K, H)/\{i_1, i_2\}$ but is an element of K_1 . By similar reasoning no contraction of (K_2, H_2) other than $(K_2, H_2)/i_1$ and $(K_2, H_2)/i_2$ as well as (K_1, H_1) itself has (K_1, H_1) as a minor. This establishes the open interval $((K_1, H_1), (K_2, H_2))$ consists solely of the minors $(K_2, H_2)/i_1$ and $(K_2, H_2)/i_2$.

Now return to the case $I = \{i_1, i_2\}$ and suppose $i_1 < i_2$. In this case $i_1 \lor i_2 = i_2$ so $(K_2, H_2)/\{i_1, i_2\} = (K_2, H_2)/\{i_2\}$. By a similar argument as used in the previous subcase each generator j of (K_1, H_1) corresponds to a unique generator j' of (K_2, H_2) such that $j' \lor i_2 = j$. Due to this uniqueness no deletion of (K_2, H_2) other than $(K_2, H_2) \setminus \{i_1\}$ contains (K_1, H_1) as a minor. Similarly, no contraction of (K_2, H_2) with the exceptions of $(K_2, H_2)/i_1$ and (K_1, H_1) itself contain (K_1, H_1) as a minor. Thus, the open interval $((K_1, H_1), (K_2, H_2))$ consists solely of the minors $(K_2, H_2)/i_1$ and $(K_2, H_2) \setminus i_1$.

Suppose $I = \{i\}$ and $J = \emptyset$. If there exists $i' \in H_2$ such that i' < i then we may take $I = \{i, i'\}$ which falls under a previous case. Otherwise, there is one generator $j \in H_1$ that corresponds to two generators $j_1, j_2 \in H_2$. All other generators in H_1 correspond to a unique generator in H_2 . Thus in this case

$$((K_1, H_1), (K_2, H_2)) = \{(K_2, H_2) \setminus \{j_1\}, (K_2, H_2) \setminus \{j_2\}\}.$$

What remains is the case when both I and J are singletons. Suppose $I = \{i\}$ and $J = \{i \lor j\}$ for some generator j of (K_2, H_2) . Once again since $|H_2| - |H_1| = 2$, every generator of (K_1, H_1) corresponds to a unique generator of (K_2, H_2) . Hence for all generators j_1, j_2 of (K_2, H_2) the joins $i \lor j_1$ and $i \lor j_2$ are distinct. Hence, the open interval $((K_1, H_1), (K_2, H_2))$ consists of the minors $(K_2, H_2) \setminus j$ and $(K_2, H_2)/i$.

Lemma 3.1.3. For any generator enriched lattice (L, G) the minor poset M(L, G) is graded by rk(K, H) = |H| + 1 and is thin.

Proof. First observe the atoms of the minor poset M(L,G) are the minors of (L,G) that contain one element. A generator enriched lattice with a single element has no generators, and thus has no minors. Thus, such elements cover the minimal element \emptyset of M(L,G). On the other hand if a minor of (L,G) covers the minimal element \emptyset of M(L,G), then by definition it must have no minors. A generator enriched lattice with no minors must have no generators and thus consists of a single element.

By Lemma 3.1.2 whenever $(K_1, H_1) \prec (K_2, H_2)$ the difference $|H_2| - |H_1|$ must be less than 2. Clearly this difference is positive, so it must be equal to 1. Since every atom has zero generators every saturated chain from the minimal element \emptyset to a minor (K, H) must have the same length, namely |H| + 1.

Having proven the minor poset M(L,G) is graded by rk(K,H) = |H| + 1, it follows from Lemma 3.1.2 that the poset M(L,G) is thin.

The order relation of the minor poset of a generator enriched lattice satisfying the join irreducible lift property is reduces to inclusion.

Definition 3.1.4. A generator enriched lattice (L, G) is said to lift join irreducibles if for all $\ell \in L$ and $i \in G$ the element $i \vee \ell$ is join irreducible in $[\ell, \widehat{1}]$.

Taking $\ell = \widehat{0}$ in the above definition it is seen a generator enriched lattice that lifts join irreducibles must be minimally generated. This property may be equivalently stated as every minor of (L, G) is minimally generated. In particular, a minimally generated distributive lattice has the join irreducible lift property by Proposition 2.2.6, as do minimally generated geometric lattices by Lemma 2.2.3. Figure 2 shows that the face lattice of the square is an example of a minimally generated lattice that does not lift join irreducibles.

Proposition 3.1.5. If (L,G) is a generator enriched lattice with the join irreducible lift property, and (K_1, H_1) and (K_2, H_2) are minors of (L,G) then $(K_1, H_1) \leq (K_2, H_2)$ if and only if $K_1 \subseteq K_2$.

Proof. If $(K_1, H_1) \leq (K_2, H_2)$, that is, (K_1, H_1) is a minor of (K_2, H_2) , then clearly $K_1 \subseteq K_2$. Now assume conversely $K_1 \subseteq K_2$. Set $k_1 = \widehat{0}_{K_1}$ and $k_2 = \widehat{0}_{K_2}$. Let $M_1 = [k_1, \widehat{1}]$ and $M_2 = [k_2, \widehat{1}]$. Since (L, G) has the join irreducible lift property, $H_1 = \operatorname{irr}(K_1) \subseteq \operatorname{irr}(M_1)$. Furthermore,

$$\operatorname{irr}(M_1) = \{i \vee k_1 : i \in \operatorname{irr}(L)\} = \{i \vee k_1 : i \in \operatorname{irr}(M_2)\}.$$

By assumption $K_1 \subseteq K_2$. Thus $\operatorname{irr}(K_1) \subseteq \operatorname{irr}(M_1) \cap K_2$. Irreducibility in M_1 implies irreducibility in $M_1 \cap K_2$. Thus $\operatorname{irr}(K_1) \subseteq \operatorname{irr}(M_1 \cap K_2)$, and (K_1, H_1) is a deletion of

$$(M_1 \cap K_2, \operatorname{irr}(M_1 \cap K_2)) = (K_2, H_2)/k_1.$$

Therefore $(K_1, H_1) \leq (K_2, H_2)$ in M(L, G).

At this point the isomorphism type of the minor posets of Boolean algebras and chains are readily determined.

Proposition 3.1.6. The minor poset $M(B_n, irr(B_n))$ of the Boolean algebra B_n with minimal generating set is isomorphic to the face lattice of the n-dimensional cube.

Proof. Recall the face lattice of the n-dimensional cube is isomorphic to the poset of intervals of B_n with a unique minimal element $\widehat{0}$ appended [21, Chapter 3, Exercise 177]. It is easy to see from Lemma 2.2.3 that the minors of $(B_n, \operatorname{irr}(B_n))$ are exactly the intervals of B_n . Proposition 3.1.5 implies the order relations are the same.

Proposition 3.1.7. The minor poset of the length n chain is isomorphic to the rank n + 1 Boolean algebra.

Proof. In a chain every element except the minimal element is join irreducible and must be a generator. As a result, every subset of a chain is a minor. Proposition 3.1.5 implies the minors are ordered by inclusion. \Box

Minor posets of minimally generated distributive lattices can be described in terms of order minors.

Proposition 3.1.8. Let (L, irr(L)) be the minimally generated lattice of lower order ideals of a poset P. The minor poset M(L, irr(L)) is isomorphic to the poset of order ideals of P ordered via $(I_1, J_1) \leq (I_2, J_2)$ when $I_1 \subseteq I_2$ and the maximal elements of J_1 are contained in $I_2 \cup J_2$.

Proof. Recall from Proposition 2.2.5 that an order minor (I, J) corresponds to the lattice minor $((L, \operatorname{irr}(L))|_{I \cup J})/J$. Let

$$(K_1, \operatorname{irr}(K_1)) = ((L, \operatorname{irr}(L))|_{I_1 \cup J_1})/J_1,$$

and

$$(K_2, \operatorname{irr}(K_2)) = ((L, \operatorname{irr}(L))|_{I_2 \cup J_2})/J_2.$$

We have $(K_1, \operatorname{irr}(K_1)) \leq (K_2, \operatorname{irr}(K_2))$ if and only if $\widehat{0}_{K_1} \in K_2$ and

$$\operatorname{irr}(K_1) \subseteq \{i \vee \widehat{0}_{K_1} : i \in \operatorname{irr}(K_2)\}.$$

The former occurs precisely when the maximal elements of J_1 are contained in $I_2 \cup J_2$. The latter condition holds if and only if $I_1 \subseteq I_2$.

See Figure 5 for an example.

The lowest three ranks of minor posets are easily described. In particular, from Lemma 2.2.2 the rank 1 elements of M(L, G) are in bijection with the elements of L and the rank 2 elements




Figure 5: The poset of order minors of a 3 element poset P. An order minor (I, J) is depicted by coloring the elements of $I \cup J$ black, circling the elements of J and coloring the elements of $P \setminus (I \cup J)$ white.

of M(L, G) are in bijection with the edges of the diagram of (L, G). Somewhat more can be said for the rank 2 and 3 minors when (L, G) is minimally generated and geometric. The following lemma will be used for this purpose.

Lemma 3.1.9. A generator enriched lattice is minimally generated and geometric if and only if it has no minors isomorphic to the length 2 chain.

Proof. The class of minimally generated geometric lattices is closed under taking minors and the length 2 chain is not geometric. Conversely, consider a generator enriched lattice (L,G) that is not a minimally generated geometric lattice. First consider the case in which (L,G) has some generator g that is not an atom. There must be some atom a with a < g. Observe the minor $L|_{\{a,g\}}$ is isomorphic to the length 2 chain. Now suppose L is not upper semimodular. Let $x,y \in L$ such that $x \wedge y \prec x$ and $x \wedge y \prec y$ but $x \not\prec x \vee y$. Since y covers $x \wedge y$ it is a generator in the contraction $(L,G)/(x \wedge y)$. This implies $x \vee y$ is a generator of the minor $(L,G)/x = ((L,G)/(x \wedge y))/x$. Since $x \not\prec x \vee y$ the minor (L,G)/x has a generator that is not an atom. By the preceding case (L,G)/x has a minor isomorphic to the length 2 chain, which is a minor of (L,G) as well.

Proposition 3.1.10. The diagram of a generator enriched lattice (L,G) is equal to the Hasse diagram of L if and only if (L,G) is minimally generated and geometric.

Proof. First assume (L, G) is minimally generated and geometric. A minor (L, G) with one generator is of the form $\langle b|a\rangle$ for $a, b \in L$ with $a \prec b$. Thus all edges of the diagram are edges of the Hasse diagram of L. Since the converse always holds the two graphs are equal.

Now let (L, G) be a generator enriched lattice that either is not minimally generated or is not geometric. By Lemma 3.1.9 the generator enriched lattice (L, G) must have a minor isomorphic to the length 2 chain. The diagram of a minor of (L, G) is a subgraph of the diagram of (L, G), so there must be an edge in the diagram of (L, G) that is not an edge of the Hasse diagram of L.

Proposition 3.1.11. For any generator enriched lattice (L, G) the lower intervals $[\widehat{0}, (K, H)]$ where $\operatorname{rk}(K, H) = 3$ of the minor poset $\operatorname{M}(L, G)$ are each isomorphic to the Boolean algebra B_3 or the face lattice Q_2 of a square. Moreover, all intervals $[\widehat{0}, (K, H)]$ with $\operatorname{rk}(K, H) = 3$ are isomorphic to Q_2 if and only if (L, G) is minimally generated and geometric.

Proof. The only possible isomorphism types of a generator enriched lattice with 2 generators are the minimally generated Boolean algebra $(B_2, \text{irr}(B_2))$ and the length 2 chain. The minor poset of the Boolean algebra B_2 is isomorphic to the face lattice of a square while the minor poset of the length 2 chain is isomorphic to the Boolean algebra B_3 . The second statement follows from Lemma 3.1.9.

3.2 Operations on generator enriched lattices

In this subsection we discuss several operations on generator enriched lattices and how these affect the associated minor posets. First we establish the following lemma which allows us to induce order-preserving maps on minor posets from strong maps. Incidentally, Theorem 4.3.1 is a considerable strengthening of this lemma in the particular case of strong surjections.

Lemma 3.2.1. Let $f:(L,G) \to (K,H)$ be a strong map between generator enriched lattices. The induced map $F: M(L,G) \to M(K,H)$ defined by

$$F(\langle I|z\rangle) = \langle f(I)|f(z)\rangle$$

is order-preserving.

Proof. Let (L_1, G_1) and (L_2, G_2) be minors of (L, G) such that $(L_1, G_1) \leq (L_2, G_2)$. Set $(K_1, H_1) = F(L_1, G_1)$ and $(K_2, H_2) = F(L_2, G_2)$. Since (L_1, G_1) is a minor of (L_2, G_2) the generating set G_1 is a subset of the set $\{g \vee \widehat{0}_{L_1} : g \in G_2\} \setminus \{\widehat{0}_{L_1}\}$. Applying the map F the set H_1 is a subset of the set

$$\{f(g) \lor f(\widehat{0}_{L_1}) : g \in G_2\} \setminus \{f(\widehat{0}_{L_1})\} = \{h \lor \widehat{0}_{K_1} : h \in H_2\} \setminus \{\widehat{0}_{K_1}\}.$$

This establishes that if the contraction $(K_2, H_2)/\widehat{0}_{K_1}$ is defined then $(K_1, H_1) \leq (K_2, H_2)$. For the contraction to be defined the element $\widehat{0}_{K_1}$ must be able to be expressed as a join of elements from H_2 . The element $\widehat{0}_{L_1}$ can be expressed as a join $\widehat{0}_{L_1} = g_1 \vee \cdots \vee g_r$ for some generators $g_i \in G_2$ for $i = 1, \ldots, r$. Applying the map f we have $\widehat{0}_{K_1} = f(g_1) \vee \cdots \vee f(g_r)$, the elements $f(g_i)$ are contained in $H_2 \cup \{\widehat{0}_{K_2}\}$. Possibly for some i we have $f(g_i) = \widehat{0}_{K_2}$, removing all such elements results in an expression of $\widehat{0}_{K_1}$ as a join of elements from H_2 . Therefore $(K_1, H_1) \leq (K_2, H_2)$ and the induced map F is order-preserving.

We first examine the Cartesian product of generator enriched lattices. Given two generator enriched lattices (L, G) and (K, H) define the Cartesian product $(L, G) \times (K, H)$ to be the generator enriched lattice

$$(L,G)\times (K,H)=(L\times K,(G\times \{\widehat{0}_K\})\cup (\{\widehat{0}_L\}\times K)).$$

This operation behaves nicely on the minor posets, it corresponds to the diamond product. Recall that for two posets P and Q the diamond product $P \diamond Q$ is defined to be the poset $((P \setminus \{\widehat{0}_P\}) \times (Q \setminus \{\widehat{0}_Q\})) \cup \{\widehat{0}\}$. The pyramid and prism operations are defined on a poset P as $Pyr(P) = P \times B_1$ and $Prism(P) = P \diamond B_2$. We define Pyr on generator enriched lattices in the same manner as $Pyr(L, G) = (L, G) \times (B_1, irr(B_1))$. For a discussion of the diamond product, see [9, end of Section 2].

Proposition 3.2.2. For any generator enriched lattices (L,G) and (K,H) we have

$$M((L,G)\times (K,H))\cong M(L,G)\diamond M(K,H).$$

In particular, $M(Pyr(L, G)) \cong Prism(M(L, G))$.

Proof. Let $\pi_1: (L,G) \times (K,H) \to (L,G)$ and $\pi_2: (L,G) \times (K,H) \to (K,H)$ be the projection maps. By Lemma 3.2.1 these induce order-preserving maps $\overline{\pi}_1$ and $\overline{\pi}_2$ between the minor posets. Neither of these maps send a minor to the minimal element \emptyset , so we have an order-preserving map $\phi: M((L,G) \times (K,H)) \to M(L,G) \diamond M(K,H)$ defined on minors by $\phi(M,I) = (\overline{\pi}_1(M,I), \overline{\pi}_2(M,I))$.

The inverse of ϕ is the map ψ defined for $((L',G'),(K',H'))\in (\mathrm{M}(L,G)\diamond \mathrm{M}(K,H))\setminus\{\emptyset\}$ by

$$\psi((L', G'), (K', H')) = \langle (G' \times \{\widehat{0}_{K'}\}) \cup (\{\widehat{0}_{L'}\} \times H') | (\widehat{0}_{L'}, \widehat{0}_{K'}) \rangle.$$

To see that the image under ψ is indeed a minor, let $G'' \subseteq G$ such that $G' = \{g \lor \widehat{0}_{L'} : g \in G''\}$. Similarly define H''. The generating set of the image under ψ can be described as

$$\{(g,\widehat{0}_K)\vee(\widehat{0}_{L'},\widehat{0}_{K'}):g\in G''\}\cup\{(\widehat{0}_L,h)\vee(\widehat{0}_{L'},\widehat{0}_{K'}):h\in H''\}.$$

Lemma 2.2.2 implies the image is indeed a minor of the Cartesian product.

Observe that for $I \subseteq G'$ we have

$$\psi((L',G')\setminus I,(K',H')) = \psi((L',G'),(K',H'))\setminus (I\times\{\widehat{0}'_K\}),$$

and

$$\psi((L',G')/I,(K',H')) = \psi((L',G'),(K',H'))/(I \times \{\widehat{0}_K'\}).$$

A similar statement holds for deletions and contractions of (K', H'). Thus the map ψ is order-preserving.

We now consider the minor poset of the result of adjoining a new maximal element to a generator enriched lattice. Given a generator enriched lattice (L, G) let \widehat{L} denote the lattice obtained from L by adjoining a new maximal element $m > \widehat{1}_L$ and let $\widehat{G} = G \cup \{m\}$.

Proposition 3.2.3. For any generator enriched lattice (L,G) we have

$$M(\widehat{L}, \widehat{G}) \cong Pyr(M(L, G)).$$

Proof. Let m be the maximal element of \widehat{L} . Observe since m is join irreducible and maximal in \widehat{L} , for any minor $(K,H) \neq (m,\emptyset)$ of $(\widehat{L},\widehat{G})$, both $(K \cup \{m\}, H \cup \{m\})$ and $(K \setminus \{m\}, H \setminus \{m\})$ are minors of (L,G). Define a map $\phi : M(\widehat{L},\widehat{G}) \to \operatorname{Pyr}(M(L,G))$ by setting $\phi(\emptyset) = (\emptyset,\widehat{0})$, and for minors (K,H) of (L,G) setting

$$\phi(K,H) = \begin{cases} (K,H,\widehat{0}) & \text{if } m \not\in K, \\ (K \setminus \{m\}, H \setminus \{m\},\widehat{1}) & \text{if } m \in H, \\ (\emptyset,\widehat{1}) & \text{if } K = \{m\}, \ H = \emptyset. \end{cases}$$

The map ϕ is order-preserving. Define $\psi : \operatorname{Pyr}(\operatorname{M}(L,G)) \to \operatorname{M}(\widehat{L},\widehat{G})$ by setting $\psi(\emptyset,\widehat{0}) = \emptyset$ and $\psi(\emptyset,\widehat{1}) = (m,\emptyset)$ and setting

$$\psi(K, H, \varepsilon) = \begin{cases} (K, H) & \text{if } \varepsilon = \widehat{0}, \\ (K \cup \{m\}, H \cup \{m\}) & \text{if } \varepsilon = \widehat{1}. \end{cases}$$

Clearly the map ψ is order-preserving and is the inverse of ϕ .

3.3 The lattice property for the poset of minors

In this subsection we characterize the generator enriched lattices where the associated minor poset is itself a lattice. This characterization (Theorem 3.3.6 and Proposition 3.3.2) is in terms of five forbidden minors.

To prove the characterization the following concept of a lattice with no parallels is required. This property is weaker than the property that the minor poset is a lattice, and is crucial to the proof of Theorem 3.3.6.

Definition 3.3.1. A generator enriched lattice (L,G) has no parallels if for any element $\ell \in L$ and generators $g,h \in G$ whenever $g \lor \ell \neq \ell$ then $g \lor \ell \neq h \lor \ell$. If $g \lor \ell = h \lor \ell \neq \ell$ holds the generator enriched lattice (L,G) is said to have a parallel.




Figure 6: The forbidden minors for the no parallels property.

A parallel in a minimally generated geometric lattice corresponds to a nontrivial parallel class in a loopless contraction of the associated simple matroid. It is straightforward to show any minimally generated distributive lattice has no parallels ([10, Lemma 3.13]).

Proposition 3.3.2. A generator enriched lattice has no parallels if and only if it has no minor isomorphic to one of the four generator enriched lattices whose diagrams are depicted in Figure 6.

Proof. It is clear from the definition that the class of generator enriched lattices with no parallels is closed under taking minors. Observe that each of the four depicted generator enriched lattices has a parallel formed by the elements labeled g, h and ℓ . Thus any generator enriched lattice with a minor isomorphic to one of the four generator enriched lattices depicted in Figure 6 has a parallel.

Let (L,G) be a generator enriched lattice with a parallel. By definition there exists an element $\ell \in L$ and generators $g_1, g_2 \in G$ such that $g_1 \vee \ell = g_2 \vee \ell \neq \ell$. Choose ℓ' to be minimal among such ℓ , that is, $g_1 \vee \ell' = g_2 \vee \ell' \neq \ell'$ and for all $\ell < \ell'$ the elements $g_1 \vee \ell$ and $g_2 \vee \ell$ are distinct. Choose an element ℓ_0 of L such that there is a generator h of (L,G) with $h \vee \ell_0 = \ell'$. Now consider the contraction $(L,G)/\ell_0$. The elements $g_1 \vee \ell_0$, $g_2 \vee \ell_0$ and ℓ' are distinct and each is a generator of the contraction $(L,G)/\ell_0$. Let $(K,H) = ((L,G)/\ell_0)|_{\{g_1 \vee \ell_0, g_2 \vee \ell_0, \ell'\}}$. The generator enriched lattice (K,H) has a parallel and three generators. It is readily checked that the only possibilities for (K,H) are the four generator enriched lattices depicted in Figure 6.

Figure 7 gives an example of a generator enriched lattice which is not distributive and has no parallels. The same example also shows a lattice with no parallels need not be semimodular. On the other hand the generator enriched lattice shown in Figure 6 (a) is an example of a minimally generated modular lattice which has a parallel.

Recall a generator enriched lattice (L,G) is said to lift join irreducibles if for any generator g and element $\ell \in L$ the element $g \vee \ell$ is join irreducible in the interval $[\ell, \widehat{1}]$. Equivalently a generator enriched lattice lifts join irreducibles whenever every minor is minimally generated. The following forbidden minor characterization shows this property is weaker than the property of no parallels.



Figure 7: The diagram of a generator enriched lattice with no parallels which is not distributive.

Lemma 3.3.3. A generator enriched lattice lifts join irreducibles if and only if it has no minors isomorphic to the generator enriched lattice depicted in Figure 6 (d).

Proof. The generator enriched lattice depicted in Figure 6 (d) has underlying lattice isomorphic to B_2 and has three generators. Since this generator enriched lattice is not minimally generated, it does not lift join irreducibles. Since the join irreducible lift property is closed under taking minors, no generator enriched lattice with a minor isomorphic to B_2 with three generators lifts join irreducibles.

Let (L, G) be a generator enriched lattice that does not lift join irreducibles. There is some minor (K, H) of (L, G) that is not minimally generated. Let h be a generator of (K, H) that is not join irreducible. The element h may be expressed as the join of join-irreducibles of K, say

$$h = i_1 \vee \cdots \vee i_r$$
.

Assume this set $I = \{i_1, \ldots, i_r\}$ is minimal in the sense that for any proper subset of I the join is not equal to h. Consider the contraction $(K', H') = (K, H)/(i_1 \vee \cdots \vee i_{r-2})$. Note that the element h is equal to $h \vee i_1 \vee \cdots \vee i_{r-2}$ hence h is a generator of (K', H'). By the minimality of I the generators $a = i_1 \vee \cdots \vee i_{r-1}$ and $b = i_1 \vee \cdots \vee i_{r-2} \vee i_r$ of (K', H') are not equal to h. Furthermore, a and b must be incomparable since $a \vee b = h$. This shows the minor $(K', H')|_{\{a,b,h\}}$ of (L, G), which has three generators, has underlying lattice isomorphic to B_2 .

The following lemma gives a third equivalent definition of the no parallels property. This is needed to prove Theorem 3.3.6.

Lemma 3.3.4. Let (L,G) be a generator enriched lattice and let $\theta: B_{|G|} \to L$ be the canonical strong surjection. The generator enriched lattice (L,G) has no parallels if and only if for every $\ell \in L$ the set $\theta^{-1}(\ell)$ has a unique minimal element in the Boolean algebra.

Proof. Label the elements of G as g_1, \ldots, g_n . Assume (L, G) has a parallel, that is, for some element $\ell \in L$ and $i, j \in [n]$ we have $g_i \vee \ell = g_j \vee \ell \neq \ell$. Choose some $X \subseteq [n]$ such that $\theta(X) = \ell$. Observe that $\theta(X \cup \{i\}) = \theta(X \cup \{j\}) = g_i \vee \ell$. For any sets $A \subseteq B$ that are elements of the fiber $\theta^{-1}(g_i \vee \ell)$, we have $[A, B] \subseteq \theta^{-1}(g_i \vee \ell)$. Since the preimage set contains $X \cup \{i\}$ and $X \cup \{j\}$ but not the intersection X, it does not have a unique minimal element.

To show the converse, suppose for some $\ell \in L$ the preimage $\theta^{-1}(\ell)$ has two minimal elements X and Y. If either X or Y is of cardinality 1 then ℓ is a generator of (L,G). Furthermore, the existence of a second minimal element of the preimage $\theta^{-1}(\ell)$ implies ℓ is join reducible in L. Thus in this case (L,G) does not lift join irreducibles, hence (L,G) has a parallel. Now assume $|X| \geq 2$ and $|Y| \geq 2$. Choose some $x \in X \setminus Y$ and set $X' = X \setminus \{x\}$. Since X is minimal in $\theta^{-1}(\ell)$ the image $\theta(X')$ is not equal to ℓ . Consider the contraction (L,G)/X'. Since $\theta(X') \vee \theta(\{x\}) = \ell$ the element ℓ is a generator of (L,G)/X'. On the other hand, $\ell = \theta(Y) = \theta(X') \vee \theta(Y)$ so

$$\ell = \theta(X') \lor \bigvee_{y \in Y} \theta(\{y\}).$$

Either $\theta(X') \vee \theta(\{y\}) = \ell$ for some $y \in Y$ or ℓ is not join irreducible in (L, G)/X'. The first case shows (L, G) has a parallel and the second shows that (L, G) does not lift join irreducibles. In either case (L, G) has a parallel.

The above lemma can be restated as: a generator enriched lattice has no parallels if and only if every element has a unique minimal expression as a join of generators. Dilworth characterized lattices satisfying this condition as lower semi-modular lattices in which every modular sublattice is distributive; see [7, Theorem 1.1].

The following lemma characterizes the join operation in the minor poset for any generator enriched lattice.

Lemma 3.3.5. Let (L,G) be a generator enriched lattice, let (K_1, H_1) and (K_2, H_2) be minors of (L,G) and set $\ell_0 = \widehat{0}_{K_1} \wedge \widehat{0}_{K_2}$. The join $(K_1, H_1) \vee (K_2, H_2)$ in the minor poset M(L,G) exists if and only if the following three conditions hold:

- 1. Let θ be the canonical strong map onto $(L,G)/\ell_0$. The fibers $\theta^{-1}(\widehat{0}_{K_1})$ and $\theta^{-1}(\widehat{0}_{K_2})$ each have a unique minimal element, say X_1 and X_2 respectively.
- 2. For each generator h of (K_1, H_1) there is a unique generator g of $(L, G)/\ell_0$ with $g \vee \widehat{0}_{K_1} = h$, and similarly for (K_2, H_2) .
- 3. Any minor (K, H) of (L, G) such that $(K, H) \geq (K_1, H_1)$ and $(K, H) \geq (K_2, H_2)$ contains the element ℓ_0 .

Let I be the set

$$I = \{ \theta(\{x\}) : x \in X_1 \cup X_2 \}$$

$$\cup \{ g \vee \ell_0 : g \in G \text{ and } g \vee \widehat{0}_{K_1} \in H_1 \}$$

$$\cup \{ g \vee \ell_0 : g \in G \text{ and } g \vee \widehat{0}_{K_2} \in H_2 \}.$$

If the above conditions are satisfied then the join $(K_1, H_1) \vee (K_2, H_2)$ is the minor $((L, G)/\ell_0)|_I$.



Figure 8: A generator enriched lattice whose minor poset is not a lattice. The other four obstructions appear in Figure 6.

Proof. Assume the join $(K_1, H_1) \vee (K_2, H_2)$ exists in M(L, G). Both (K_1, H_1) and (K_2, H_2) are a minor of the contraction $(L, G)/\ell_0$, hence the join $(K_1, H_1) \vee (K_2, H_2)$ is a minor of $(L, G)/\ell_0$ as well. Let z be the minimal element of the underlying lattice of $(K_1, H_1) \vee (K_2, H_2)$. We have that $z \geq \ell_0$. On the other hand $\widehat{0}_{K_1} \geq z$ and $\widehat{0}_{K_2} \geq z$ so $z \leq \ell_0$. Therefore $z = \ell_0$. The join $(K_1, H_1) \vee (K_2, H_2)$ is thus a deletion of the minor $(L, G)/\ell_0$, say $(K_1, H_1) \vee (K_2, H_2) = ((L, G)/\ell_0)|_J$ for some set J of generators. The set J must be the unique minimal set with the properties $\widehat{0}_{K_1}$ can be expressed as a join of elements in J, the set H_1 is included in the set $\{j \vee \widehat{0}_{K_1} : j \in J\}$, and the corresponding statements hold for (K_2, H_2) . For if J' is another set with these properties then $((L, G)/\ell_0)|_{J'}$ is greater than or equal to both (K_1, H_1) and (K_2, H_2) ; thus $((L, G)/\ell_0)|_J \leq ((L, G)/\ell_0)|_{J'}$ and $J \subseteq J'$. The existence of the set J implies Conditions 1 and 2 hold. Condition 3 holds since the join $(K_1, H_1) \vee (K_2, H_2)$ contains the element ℓ_0 .

Now suppose Conditions 1 through 3 are satisfied. Let I be the set defined in the statement and let $(K_0, H_0) = ((L, G)/\ell_0)|_I$. Consider a minor (K, H) of (L, G) with $(K, H) \geq (K_1, H_1)$ and $(K, H) \geq (K_2, H_2)$. By Condition 3 the minor (K, H) contains ℓ_0 . The contraction $(K, H)/\ell_0$ satisfies $(K, H)/\ell_0 \geq (K_1, H_1)$ and $(K, H)/\ell_0 \geq (K_2, H_2)$. Thus the set

$$\{g \vee \widehat{0}_{K_1} : g \text{ is a generator of } (K, H)/\ell_0)\}$$

includes H_1 and similarly for (K_2, H_2) . Furthermore $\widehat{0}_{K_1} \in (L, G)/\ell_0$ and $\widehat{0}_{K_2} \in (L, G)/\ell_0$ so by condition 1 the generating set of $(L, G)/\ell_0$ includes $\{\theta(x) : x \in X_1 \cup X_2\}$. By condition 2 the generating set of $(L, G)/\ell_0$ and H include I, hence $(K, H) \geq ((L, G)/\ell_0)|_{I}$. Therefore $((L, G)/\ell_0)|_{I}$ is the join $(K_1, H_1) \vee (K_2, H_2)$.

Theorem 3.3.6. For any generator enriched lattice (L,G) the poset of minors M(L,G) is a lattice if and only if (L,G) has no parallels and no minor isomorphic to the generator enriched lattice whose diagram is depicted in Figure 8.

Proof. Since for any minor (K, H) of (L, G) the minor poset M(K, H) is a lower interval in M(L, G), the property that the minor poset is a lattice is closed under taking minors. It is readily checked that for each of the five generator enriched lattices depicted in Figures 6 and 8 the minor poset is not a lattice.

Let (L,G) be a generator enriched lattice with no parallels and such that there are two minors (K_1,H_1) and (K_2,H_2) of (L,G) for which the join $(K_1,H_1) \vee (K_2,H_2)$ in M(L,G) does not exist. Set $\ell_1 = \widehat{0}_{K_1}$, $\ell_2 = \widehat{0}_{K_2}$ and $\ell_0 = \ell_1 \wedge \ell_2$. It will be shown (L,G) must have a minor isomorphic to the generator enriched lattice in Figure 8. Since (L,G) has no parallels conditions 1 and 2 in Lemma 3.3.5 must be satisfied; condition 1 follows from Lemma 3.3.4 and condition 2 follows from the definition of the no parallels property. Thus condition 3 must fail, that is, there is some minor (K,H) with $(K,H) \geq (K_1,H_1)$ and $(K,H) \geq (K_2,H_2)$, but where ℓ_0 is not an element of K. In particular this implies the element ℓ_0 is neither ℓ_1 nor ℓ_2 .

Let θ be the canonical map from $B_{|G|}$ onto (L,G). The fiber $\theta^{-1}(\ell_0)$ contains a unique minimal element, say X. Let I_0 be the θ images of all singletons included in X, that is,

$$I_0 = \{\theta(\{x\}) : x \in X\}.$$

Similarly define I_1 , I_2 and I for ℓ_1 , ℓ_2 and $\widehat{0}_K$ respectively. By Lemma 3.3.4 and choice of I_0 the join of a set of generators is greater than or equal to ℓ_0 only if said set includes I_0 . A similar statement holds for I_1 and I_2 . Since $\ell_0 \notin K$ the set

$$I \cup \{g \in G : g \vee \widehat{0}_K \in H\}$$

does not include I_0 . On the other hand since ℓ_1 and ℓ_2 are elements of K the sets I_1 and I_2 are included in the above set. Thus, there is some $i_0 \in I_0 \setminus (I_1 \cup I_2)$. Recall ℓ_0 is neither ℓ_1 nor ℓ_2 . Thus $\ell_1 \not\leq \ell_2 = \ell_2 \vee \ell_0$ so I_1 cannot be included in $I_0 \cup I_2$. Let $i_1 \in I_1 \setminus (I_0 \cup I_2)$. Similarly, there is some element $i_2 \in I_2 \setminus (I_0 \cup I_1)$.

Let i be the join of all elements in the set

$$J = I_0 \cup I_1 \cup I_2 \setminus \{i_0, i_1, i_2\}.$$

Since i is the join of the elements of J, and this set includes none of I_0 , I_1 or I_2 the element i is not greater than or equal to any of i_0 , i_1 or i_2 . Thus, the elements $i \vee i_0$, $i \vee i_1$, and $i \vee i_2$ are each distinct from i. Furthermore, they are generators of the contraction (L, G)/i and by the no parallels assumption must be distinct. We claim the minor $((L, G)/i)|_{\{i_0\vee i, i_1\vee i, i_2\vee i\}}$ is isomorphic to the generator enriched lattice depicted in Figure 8. Note that $i_j \vee i = \ell_j \vee i$ for j = 0, 1, 2. Since taking the join with i is an order-preserving map, we have $i_0 \vee i \leq i_1 \vee i$ and $i_0 \vee i \leq i_2 \vee i$. It remains to show that $i_1 \vee i$ and $i_2 \vee i$ are incomparable. The element $i_2 \vee i$ is the join of the elements of $I_2 \cup J$ which does not contain i_1 hence does not include I_1 . Thus $i_2 \vee i \not\geq i_1$ hence $i_2 \vee i \not\geq i_1 \vee i$. Similarly $i_1 \vee i \not\geq i_2 \vee i$.

The four generator enriched lattices depicted in Figure 6 and the generator enriched lattice depicted in Figure 8 together form a forbidden minor characterization of the generator enriched lattices for which the minor poset is a lattice. Since all five of these generator enriched lattices have three generators the following corollary is immediate.

Corollary 3.3.7. For any generator enriched lattice (L, G) the minor poset M(L, G) is a lattice if and only if every interval $[\emptyset, (K, H)]$ such that $\operatorname{rk}(K, H) = 4$ is a lattice.

Recall that the minors of a minimally distributive lattice are in bijection with the order minors of the poset P of join irreducibles. Furthermore, the minors are all minimally generated and distributive and are isomorphic to the lattice of lower order ideals of a subposet of P; see Proposition 2.2.6. Since any minimally generated distributive lattice has no parallels, the forbidden minor characterization reduces to the following characterization for distributive lattices. In particular this indicates that minimally generated distributive lattices whose minor poset is a lattice are quite sparse.

Corollary 3.3.8. Let L be a finite distributive lattice. The minor poset M(L, irr(L)) is a lattice if and only if the poset irr(L) contains no triple x, y, z such that $x, y \ge z$ and x and y are incomparable.

3.4 A decomposition theorem

In this subsection we describe a decomposition of minor posets into a disjoint union of Boolean algebras. This decomposition is leveraged to derive formulas for the rank generating function of minor posets of minimally generated geometric lattices, and of generator enriched lattices with no parallels.

To state the decomposition theorem some notation is needed. Let (L, G) be a generator enriched lattice and let $\ell \in L$. Define $M(L, G, \ell)$ to be the subposet of M(L, G) consisting of the minors of (L, G) for which the minimal element is ℓ .

Theorem 3.4.1. Let (L,G) be a generator enriched lattice. The poset $M(L,G) \setminus \{\emptyset\}$ decomposes as a disjoint union of the subposets $M(L,G,\ell)$, that is,

$$M(L,G) \setminus \{\emptyset\} = \bigcup_{\ell \in L} M(L,G,\ell).$$

Furthermore, the subposet $M(L,G,\ell)$ is the interval $[(\ell,\emptyset),(L,G)/\ell]$ of M(L,G), and each such interval is isomorphic to a Boolean algebra.

Proof. Clearly the union is disjoint and consists of all elements of the minor $M(L,G)\setminus\{\emptyset\}$. Fix $\ell\in L$ and $(K,H)\in M(L,G,\ell)$. Deleting all generators from (K,H) results in the minor (ℓ,\emptyset) consisting of only the minimal element ℓ of K. Thus $(K,H)\geq (\ell,\emptyset)$. By Lemma 2.2.1 the minor (K,H) can be expressed as $((L,G)/\ell)\setminus I$ for some set I. Thus $(K,H)\leq (L,G)/\ell$ and $M(L,G,\ell)\subseteq [(\ell,\emptyset),(L,G)/\ell]$. The minimal element of any minor of $(L,G)/\ell$ is greater than or equal to ℓ . On the other hand if (K,H) is a minor of (L,G) such that (ℓ,\emptyset) is a minor of (K,H), then $\widehat{0}_K\leq \ell$. Thus $[(\ell,\emptyset),(L,G)/\ell]\subseteq M(L,G,\ell)$ so equality holds.

Since the contraction operation changes the minimal element of a generator enriched lattice, all relations $(K_1, H_1) \leq (K_2, H_2)$ in the interval $M(L, G, \ell)$ must be induced by deletions. Thus, $(K_1, H_1) \leq (K_2, H_2)$ in $M(L, G, \ell)$ if and only if $H_1 \subseteq H_2$. Therefore $M(L, G, \ell)$ is isomorphic to the Boolean algebra $B_{\text{rk}((L,G)/\ell)-1}$.




Figure 9: The Boolean decomposition of the minor poset depicted in Figure 4.

See Figure 9 for an example.

For a ranked poset P we denote the rank generating function as F(P;q). The rank generating function of the Boolean algebra B_n is $F(B_n;q) = \sum_{X\subseteq [n]} q^{|X|} = (1+q)^n$. Since the minimal element of each interval $M(L,G,\ell)$ is rank 1 in M(L,G), the decomposition theorem above implies the following.

Lemma 3.4.2. Let (L,G) be a generator enriched lattice and for $\ell \in L$ set

$$\alpha(\ell) = |\{g \vee \ell : g \in G\} \setminus \{\ell\}|.$$

The rank generating function of the minor poset M(L,G) is given by

$$F(M(L,G);q) = 1 + q \sum_{\ell \in L} (1+q)^{\alpha(\ell)}.$$

We now proceed to derive more compact formulas for minimally generated geometric lattices and generator enriched lattices with no parallels. The formula for the first of these is in terms of the incidence algebra of the lattice. See [21, Section 3.6] for a more thorough introduction to incidence algebras.

Recall the incidence algebra of a finite poset P is the set of all maps $f: Int(P) \to \mathbb{C}$, where Int(P) denotes the set of intervals of P. The incidence algebra is equipped with a product, known as the convolution:

$$(f * g)(x,y) = \sum_{x \le z \le y} f(x,z)g(z,y).$$

The identity is the map δ defined by $\delta(x,x) = 1$ for all $x \in P$ and $\delta(x,y) = 0$ for all $x \neq y$ in P. The zeta function $\zeta : \text{Int}(P) \to \mathbb{C}$ is defined by $\zeta(x,y) = 1$. In order to state the rank generating function formula an exponentiation operation is needed. This is the operation defined by

$$f^g(x,y) = \prod_{x \le z \le y} f(x,z)^{g(z,y)}.$$

We also need the map κ which encodes cover relations, defined by $\kappa(x,y)=1$ when $x \prec y$ and $\kappa(x,y)=0$ otherwise.

Theorem 3.4.3. Let L be a geometric lattice. The rank generating function of the minor poset M(L, irr(L)) is

$$F(M(L, irr(L)); q) = 1 + q(\zeta * (\zeta + q\kappa)^{\zeta})(\widehat{0}_L, \widehat{1}_L).$$

Proof. Since the lattice L is geometric,

$$\{i \lor \ell : i \in \operatorname{irr}(L)\} \setminus \{\ell\} = \{\ell' \in L : \ell' \succ \ell\}.$$

Applying this substitution to the sum from Lemma 3.4.2 and expanding the term inside the sum gives

$$\sum_{\ell \in L} (1+q)^{|\{\ell' \in L: \ell' \succ \ell\}|} = \sum_{\ell \in L} \prod_{\ell' \ge \ell} \begin{cases} 1+q & \text{if } \ell' \succ \ell, \\ 1 & \text{if } \ell' \ne \ell, \end{cases}$$

$$= \sum_{\ell \in L} \prod_{\ell' \ge \ell} (\zeta + q\kappa)(\ell, \ell')$$

$$= \sum_{\ell \in L} \prod_{\ell' \ge \ell} (\zeta + q\kappa)^{\zeta(\ell', \widehat{1}_L)}(\ell, \ell')$$

$$= \sum_{\ell \in L} (\zeta + q\kappa)^{\zeta}(\ell, \widehat{1})$$

$$= (\zeta * (\zeta + q\kappa)^{\zeta})(\widehat{0}, \widehat{1}).$$

For generator enriched lattices with no parallels, we give an expression for the rank generating function of M(L, G) in terms of the rank generating function of the dual lattice L^* . First we establish that a lattice with no parallels is indeed ranked. In fact, we give a description of the rank function which gives another characterization of lattices with no parallels.

Proposition 3.4.4. A generator enriched lattice (L, G) has no parallels if and only if the lattice L is graded and the rank function is given by $rk(\ell) = |\{g \in G : g \leq \ell\}|$.

Proof. For $\ell \in L$ let $r(\ell) = |\{g \in G : g \leq \ell\}|$. First assume (L, G) has a parallel, say for $g, h \in G$ and $\ell \in L$ we have $\ell \vee g = \ell \vee h \neq \ell$. If $x \prec \ell \vee g$ then $g, h \not\leq x$. Thus we must have $r(\ell \vee g) - r(x) \geq 2$ so r cannot be the rank function of L.

Now assume (L,G) has no parallels. For any atom $a \in L$ clearly r(a) = 1, so it will suffice to show that whenever $x \prec y$ we have r(y) - r(x) = 1. Since $x \prec y$ for any $g \in G$ such that $g \leq y$ but $g \not\leq x$ we have $y = x \vee g$. If $y = x \vee g = x \vee h$ then since (L,G) has no parallels we have g = h. Thus there is exactly one such generator and r(y) - r(x) = 1. \square

Theorem 3.4.5. If L is a lattice with no parallels then the rank generating function of the poset of minors M(L, irr(L)) is given by

$$F(M(L, irr(L)); q) = 1 + qF(L^*; 1 + q).$$

Proof. Since $(L, \operatorname{irr}(L))$ has no parallels $|\{g \vee \ell : g \in G\} \setminus \{\ell\}| = |\{i \in \operatorname{irr}(L) : i \not\leq \ell\}|$. On the other hand, the rank of the element $\ell \in L$ is given by

$$rk(\ell) = |\{i \in irr(L) : i \le \ell\}|.$$

The cardinality $|\{i \in \operatorname{irr}(L) : i \not\leq \ell\}|$ is the corank of ℓ , that is, $\operatorname{rk}(\widehat{1}) - \operatorname{rk}(\ell)$. This difference is equal to the rank of ℓ in the dual lattice L^* . Substituting the rank of ℓ in L^* for the exponent in Lemma 3.4.2 results in the desired expression.

4 The zipping construction

In this section a construction is given for strong minor posets using the zipping operation introduced by Reading in [18]. This construction implies that any minor poset is isomorphic to the face poset of a regular CW sphere, and in particular is Eulerian. The construction also yields inequalities for the **cd**-indices of minor posets.

4.1 Background

To begin with some background on Eulerian posets, the zipping operation and PL-topology as related to posets is recalled. The *order complex* $\Delta(P)$ of a poset P is the simplicial complex consisting of all chains in P. The *proper part* of a poset P with $\widehat{0}$ and $\widehat{1}$ is the subposet $P \setminus \{\widehat{0}, \widehat{1}\}$.

A simplicial complex Δ is a *PL-sphere* when there is a piecewise linear homeomorphism from Δ to the boundary of a simplex. A poset is said to be a PL-sphere when the order complex is a PL-sphere. A basic fact of PL-topology is that the proper part of the face lattice of any polytope is a PL-sphere.

The link of a face X in a simplicial complex Δ is the subcomplex

$$\operatorname{link}_{\Delta}(X) = \{Y \in \Delta : Y \cap X = \emptyset \text{ and } Y \cup X \in \Delta\}.$$

Lemma 4.1.1 (Hudson [13, Corollary 1.16]). Given a simplicial complex Δ that is a PL-sphere, for any face X of Δ the link link $_{\Delta}(X)$ is a PL-sphere.

Let P be a poset with $\widehat{0}$ and $\widehat{1}$ where the proper part is a PL-sphere and let [x,y] be an interval of P. The order complex $\Delta((x,y))$ is the link $\operatorname{link}_{\Delta(P)}(C)$ where C is the union of a maximal chain in $[\widehat{0},x]$ and a maximal chain in $[y,\widehat{1}]$. By Lemma 4.1.1 the order complex $\Delta((x,y))$ is a PL-sphere. In summation, if P is a poset such that the proper part is a PL-sphere then the proper part of any closed interval of P is a PL-sphere as well.

The following definition and theorem due to Björner characterize face posets of regular CW complexes.

Definition 4.1.2 (Björner [6, Definition 2.1]). A poset P is a CW poset if the following three conditions hold:

- 1. P has a unique minimal element $\hat{0}$ and a unique maximal element $\hat{1}$.
- 2. $|P| \ge 3$.
- 3. $\Delta((\widehat{0}, p))$ is homeomorphic to a sphere for all elements p in the open interval $(\widehat{0}, \widehat{1})$.

The definition we give differs slightly from the one given in [6] since in the present context face posets are adjoined with a maximal element which does not correspond to a cell.

Theorem 4.1.3 (Björner [6, Proposition 3.1]). A poset is a CW poset if and only if it is isomorphic to the face poset of a regular CW complex.

A graded poset P with $\widehat{0}$ and $\widehat{1}$ is said to be Eulerian if for all x < y in P,

$$\sum_{x \le z \le y} (-1)^{\operatorname{rk}(z)} = 0.$$

Equivalently the Möbius function of P satisfies $\mu(x,y)=(-1)^{\mathrm{rk}(y)-\mathrm{rk}(x)}$. Face lattices of polytopes are the motivating examples of Eulerian posets. More generally face posets of regular CW spheres are Eulerian posets.

The zipping operation was introduced by Reading in [18, Section 4] as a tool to study Bruhat intervals.

Definition 4.1.4. Let P be a poset and let $x, y, z \in P$. The triple x, y, z is said to form a zipper if the following three conditions hold:

(i) z covers only the two elements x and y.

- (ii) $z = x \vee y$.
- (iii) $\{p \in P : p < x\} = \{p \in P : p < y\}.$

Let P be a poset with a zipper $x, y \prec z$. The poset zip(P, z) has underlying set obtained from P by replacing the elements x, y and z with a new element w. The order relation of zip(P, z) is defined by the following three conditions that hold for all p and q in $P\setminus\{x, y, z\}$.

- (a) $p \le w$ in zip(P, z) if and only if $p \le z$ in P.
- (b) $w \le p$ in zip(P, z) if and only if $x \le p$ or $y \le p$ in P.
- (c) $p \le q$ in zip(P, z) if and only if $p \le q$ in P.

When P and zip(P,z) are graded we have $rk_{zip(P,z)}(w) = rk_P(z) - 1$.

Let Q be a poset obtained from P by a sequence of zipping operations and let $\pi: P \to Q$ be the projection map induced by the zipping operations. The relation $q_1 \leq q_2$ in Q holds if and only if there exist $p_1 \leq p_2$ in P such that $\pi(p_1) = q_1$ and $\pi(q_1) = q_2$.

A graded poset is said to be *thin* if all length two intervals are isomorphic to the Boolean algebra B_2 .

Remark 4.1.5. If P is a thin poset then $x, y \prec z$ form a zipper whenever conditions (i) and (ii) in Definition 4.1.4 are satisfied. Condition (iii) in the same definition follows from thinness and condition (i).

Proposition 4.1.6 (Reading [18, Proposition 4.4]). If P is a graded and thin poset, with a zipper $x, y \prec z$, then the poset zip(P, z) is graded and thin as well.

Theorem 4.1.7 (Reading [18, Theorem 4.7]). If the proper part of P is a PL-sphere and $x, y \prec z$ form a zipper, then so is the proper part of zip(P, z).

The definition of the **cd**-index is briefly presented here. For a further discussion of the **cd**-index see the survey by Bayer [1].

Let P be a graded poset of rank n+1 with $\widehat{0}$ and $\widehat{1}$. Let \mathbf{a} and \mathbf{b} be noncommuting variables of degree 1. For each chain $C = \{\widehat{0} < x_1 < \cdots < x_k < \widehat{1}\}$ in P define a weight $w(C) = w_1 \cdots w_n$ by

$$w_i = \begin{cases} \mathbf{b} & \text{if there is a rank } i \text{ element in } C, \\ \mathbf{a} - \mathbf{b} & \text{otherwise.} \end{cases}$$

The ab-index of P is the polynomial

$$\Psi(P) = \sum_{C} w(C),$$

where the sum above is over all chains $C = \{\widehat{0} < x_1 < \dots < x_k < \widehat{1}\}$. Define noncommutative variables $\mathbf{c} = \mathbf{a} + \mathbf{b}$ and $\mathbf{d} = \mathbf{ab} + \mathbf{ba}$ of degree 1 and 2, respectively. If the \mathbf{ab} -index of P can be expressed in terms of \mathbf{c} and \mathbf{d} , the resulting polynomial is called the \mathbf{cd} -index of P and is also denoted $\Psi(P)$. Not every poset has a \mathbf{cd} -index, but every Eulerian poset has a \mathbf{cd} -index [2, Theorem 4].

In order to establish certain cd -index inequalities for minor posets we will need a condition for nonnegativity of the cd -index. The Gorenstein* condition is such a condition. A poset P whose proper part is P' is said to be $\operatorname{Gorenstein}^*$ when the order complex $\Delta(P')$ is a real homology sphere. This means the homology groups of the order complex and of every link in the order complex are isomorphic to those of a sphere of the same dimension. See [8, Section 2.1] for a precise definition of Gorenstein* posets. A Gorenstein* poset is necessarily Eulerian. Lemma 4.1.1 implies that any poset P whose proper part P' is a PL-sphere is a Gorenstein* poset.

Many results on nonnegativity of **cd**-indices led up to the nonnegativity result for Gorenstein* posets. Purtill showed that the coefficients of the **cd**-index are nonnegative for polytopes of dimension at most 5 ([17, Proposition 7.11]), and polytopes with simplicial facets ([17, Corollary 7.8]). More generally Purtill showed that any CL-shellable Eulerian poset whose proper upper intervals are all Boolean algebras has a **cd**-index with nonnegative coefficients in [17, Corollary 7.4]. Stanley showed that the **cd**-index of the face poset of any S-shellable regular CW sphere, a class which includes all polytopes, has nonnegative coefficients in [20, Theorem 2.2], and also that the **cd**-index of any Gorenstein* poset such that all lower intervals are Boolean algebras has nonnegative coefficients in [20, Corollary 3.1]. Reading established the nonnegativity of certain coefficients of **cd**-indices of all Gorenstein* posets in [19, Theorem 3]. We will require nonnegativity of the **cd**-index in the most general case as stated below. This was conjectured by Stanley in [20, Conjecture 2.1]

Theorem 4.1.8 (Karu [14, Theorem 1.3]). If P is a Gorenstein* poset then the coefficients of the $\operatorname{\mathbf{cd}}$ -index $\Psi(P)$ are nonnegative.

There has also been interest in inequalities relating **cd**-indices of different posets, in particular, **cd**-index inequalities imply flag vector inequalities. Billera, Ehrenborg and Readdy showed that the set of **cd**-indices of lattices of regions of oriented matroids is minimized by the **cd**-index of the cross polytope ([4, Corollary 7.5]), and in particular that the set of **cd**-indices of zonotopes is minimized by the **cd**-index of the cube ([4, Corollary 7.6]). Billera and Ehrenborg gave certain inequalities between the **cd**-index of a polytope and the **cd**-index of a face ([3, Theorem 5.1]), a particular case being $\Psi(P) \geq \Psi(\text{Pyr}(F))$ where F is a facet of the polytope P ([3, Corollary 5.2]). This result was then used to show that the set of **cd**-indices of polytopes is minimized by the **cd**-index of the simplex, that is, the **cd**-index of the Boolean algebra ([3, Theorem 5.3]). Billera and Ehrenborg also showed that the **cd**-indices of the cyclic polytopes coefficientwise maximize **cd**-indices of polytopes in [3, Theorem 6.5]. Generalizing the lower bound for polytopes, Ehrenborg and Karu showed that **cd**-index of the Boolean algebra is the minimum among all Gorenstein* lattices.

Reading gave a formula for how zipping operations affect the **cd**-index, we will use this to derive inequalities between **cd**-indices of minor posets in Section 4.3.

Theorem 4.1.9. If P is an Eulerian poset and $x, y \prec z$ form a zipper, then the poset zip(P, z) is Eulerian as well.

(a) (Reading [18, Proposition 4.5, Theorem 4.6]) If $z \neq \widehat{1}_P$ then

$$\Psi(\operatorname{zip}(P,z)) = \Psi(P) - \Psi([\widehat{0},x]_P) \cdot \mathbf{d} \cdot \Psi([z,\widehat{1}]_P).$$

(b) (Stanley [20, Lemma 1.1]) If $z = \widehat{1}_P$ then

$$\Psi(\operatorname{zip}(P,z)) = \Psi(P) \cdot \mathbf{c}.$$

4.2 Factoring strong surjections

In this subsection we provide a process to factor any strong surjection into strong surjections that only identify two elements. For the maps appearing in this factorization we give a description of the fibers of the induced map between the minor posets which is needed for the proof of Theorem 4.3.1.

Definition 4.2.1. Let $\mathcal{E}(L,G)$ denote the poset consisting of edges of the diagram of a generator enriched lattice (L,G), that is, pairs $(\ell,\ell\vee g)$ for $\ell\in L$ and $g\in G$ such that $\ell\vee g\neq \ell$. The order relation of $\mathcal{E}(L,G)$ is defined by $(\ell,\ell\vee g)\leq (a\vee\ell,a\vee\ell\vee g)$ for $a\in L$.

An oriented edge in the diagram of a generator enriched lattice (L, G) is determined by its vertices. For notational convenience the elements of $\mathcal{E}(L, G)$ will be denoted as unordered pairs.

Definition 4.2.2. Given an equivalence relation ϕ on a generator enriched lattice (L, G), an edge of ϕ is an edge $(\ell, \ell \vee g)$ of (L, G) such that $\ell \equiv \ell \vee g(\phi)$. Let $\mathcal{E}(\phi)$ denote the set of edges of ϕ . The equivalence relation ϕ is said to be connected if for all $a \equiv b(\phi)$ there is a sequence $a = c_0, c_1, \ldots, c_k = b$ such that for $1 \le i \le k$ the pair $\{c_{i-1}, c_i\}$ is an edge of ϕ .

Note that a connected relation is determined by its set of edges.

Lemma 4.2.3. Let ϕ be an equivalence relation on a generator enriched lattice (L, G). The relation ϕ is join-preserving if and only if (i) it is connected and (ii) the set of edges $\mathcal{E}(\phi)$ forms an upper order ideal in the poset $\mathcal{E}(L, G)$ of edges of (L, G).

Proof. Let ϕ be a join preserving equivalence relation on (L, G). To show ϕ is connected let $a, b \in (L, G)$ with $a \equiv b(\phi)$. Then $a \equiv a \lor b \equiv b(\phi)$. Choose some sequence g_1, \ldots, g_r of

generators of (L, G) such that $a < a \lor g_1 < \cdots < a \lor (g_1 \lor \cdots \lor g_r) = a \lor b$. Each term of the sequence must be congruent to a since said terms lie in the interval $[a, a \lor b]$. Thus each pair of subsequent terms in the sequence is an edge of ϕ . There exists a similarly defined sequence from b to $a \lor b$. Concatenating these two sequences gives a sequence from a to b consisting of edges of ϕ . Therefore ϕ is connected.

In order to show the set $\mathcal{E}(\phi)$ forms an upper order ideal of the poset $\mathcal{E}(L,G)$, let $\{a,b\}$ be an edge of ϕ and let $\ell \in L$. Since $a \equiv b(\phi)$ and ϕ is join-preserving $a \lor \ell \equiv b \lor \ell(\phi)$. Hence if $a \lor \ell \neq b \lor \ell$ then $\{a \lor \ell, b \lor \ell\}$ is an edge of ϕ . Thus $\mathcal{E}(\phi)$ is an upper order ideal of $\mathcal{E}(L,G)$.

Conversely, consider a connected equivalence relation ϕ on L such that the set of edges $\mathcal{E}(\phi)$ forms an upper order ideal in the poset $\mathcal{E}(L,G)$ of edges of (L,G). Let $a,b\in L$ such that $a\equiv b(\phi)$. Since ϕ is connected there is a sequence $a=c_0,c_1,\ldots,c_k=b$ such that for $1\leq i\leq k$ the pair $\{c_{i-1},c_i\}$ is an edge of ϕ . Taking the join with any $\ell\in L$ results in a sequence $a\vee\ell=c_0\vee\ell,c_1\vee\ell,\ldots,c_k\vee\ell=b\vee\ell$. For $1\leq i\leq k$ either $c_{i-1}\vee\ell=c_i\vee\ell$ or $\{c_{i-1}\vee\ell,c_i\vee\ell\}$ is an edge of L. In the second case since $\mathcal{E}(\phi)$ is an upper order ideal, the edge $\{c_{i-1},c_i\}\in\mathcal{E}(\phi)$ implies that $\{c_{i-1}\vee\ell,c_i\vee\ell\}\in\mathcal{E}(\phi)$. Removing repeated terms results in a sequence from $a\vee\ell$ to $b\vee\ell$ consisting of edges of ϕ . Thus $a\vee\ell\equiv b\vee\ell(\phi)$, hence ϕ is a join preserving equivalence relation.

Lemma 4.2.4. Let (L,G) and (K,H) be generator enriched lattices and $f:(L,G) \to (K,H)$ be a strong map. If the map f has a single nontrivial fiber $\{x,y\}$ in L, then the element x is only covered by y or vice versa.

Proof. Since f(x) = f(y) it must be that $f(x \lor y) = f(x) \lor f(y) = f(x)$. Since $f^{-1}(x) = \{x, y\}$ the element $x \lor y$ is either x or y. Without loss of generality we may assume x < y. In fact $x \prec y$, since if $x \le z \le y$ then $f(x) \le f(z) \le f(y) = f(x)$ so f(z) = f(x). Hence z must be equal to x or to y. On the other hand, if z > x then $f(z) = f(x \lor z) = f(y \lor z)$. The map f is invertible when restricted to $L \setminus \{x\}$, so this implies $y \lor z = z$ hence $y \le z$. Therefore x is only covered by y.

Lemma 4.2.5. Let (L,G) and (K,H) be generator enriched lattices. Any strong surjection $f:(L,G)\to (K,H)$ can be factored as $f=f_r\circ\cdots\circ f_1$ where each map f_i is a strong surjection that identifies only two elements.

Proof. Consider the map f as an equivalence relation on L defined by $a \equiv b(f)$ when f(a) = f(b). By Lemma 4.2.3 this equivalence relation is connected and the edges form an upper order ideal in the poset $\mathcal{E}(L,G)$ of edges of (L,G). Choose a linear extension of $\mathcal{E}(L,G)$. Order the edges of f as $\{x_1,y_1\},\ldots,\{x_r,y_r\}$ by the chosen linear extension. Define equivalence relations f_i for $1 \leq i \leq r$ by letting f_i be the transitive closure of the relation defined by $x_j \equiv y_j$ for $1 \leq j \leq i$. By definition f_i is connected and the edges of f_i form an upper order ideal in $\mathcal{E}(L,G)$ so f_i is join-preserving. Thus, each relation f_i defines a generator enriched lattice, namely the quotient $(L,G)/f_i$. Furthermore, for each i either $(L,G)/f_{i+1} = (L,G)/f_i$ or $(L,G)/f_{i+1}$ is obtained from $(L,G)/f_i$ by

identifying two elements, namely $f_i(x_{i+1})$ and $f_i(y_{i+1})$. Thus, the strong map f factors as the product $f_r \circ \cdots \circ f_1$ of the quotient maps. Removing any instances of identity maps gives the desired factorization.

Recall from Lemma 3.2.1 that a strong map $f:(L,G)\to (K,H)$ induces an order-preserving map $F: \mathcal{M}(L,G)\to \mathcal{M}(K,H)$ defined by

$$F(\langle I|z\rangle) = \langle f(I)|f(z)\rangle.$$

Lemma 4.2.6. Let $f:(L,G) \to (K,H)$ be a strong map between generator enriched lattices. If the map f has a single nontrivial fiber $x \prec y$ then the induced map $F: M(L,G) \to M(K,H)$ has nontrivial fibers of the form

$$\{(M,I),(M_x,I_x),(M_y,I_y)\},\$$

where

$$x, y \in I \cup \{\widehat{0}_M\},$$

$$(M_x, I_x) = (M, I) \setminus \{y\},$$

$$(M_y, I_y) = \begin{cases} (M, I) \setminus \{x\} & \text{if } x \in I, \\ (M, I)/\{y\} & \text{if } x = \widehat{0}_M. \end{cases}$$

Proof. Clearly such a triple $\{(M,I),(M_x,I_x),(M_y,I_y)\}$ consists of a single fiber of F, and given any minor (M,I) with $x,y\in I\cup\{\widehat{0}_M\}$ there is such a triple. Furthermore, the fiber containing (M,I) is precisely the set $\{(M,I),(M_x,I_x),(M_y,I_y)\}$. Now suppose (M_1,I_1) and (M_2,I_2) are minors of (L,G) in the same fiber of F such that neither $I_1\cup\{\widehat{0}_{M_1}\}$ nor $I_2\cup\{\widehat{0}_{M_2}\}$ contains both x and y. By assumption $f(\widehat{0}_{M_1})=f(\widehat{0}_{M_2})$. Either $\widehat{0}_{M_1}=\widehat{0}_{M_2}$ or one is x and the other y. Consider the case $\widehat{0}_{M_1}=\widehat{0}_{M_2}$. The equality $F(M_1,I_1)=F(M_2,I_2)$ holds, thus $f(I_1)=f(I_2)$. Hence, these sets differ by exchanging x and y. There is a minor (M,I) of (L,G) with $I=I_1\cup I_2$ and $\widehat{0}_M=\widehat{0}_{M_1}$. This minor satisfies $F(M,I)=F(M_1,I_1)=F(M_2,I_2)$ and (M_1,I_1) and (M_2,I_2) are the minors $(M,I)\setminus\{x\}$ and $(M,I)\setminus\{y\}$.

Now consider the case where $\widehat{0}_{M_1} \neq \widehat{0}_{M_2}$. Since $f(\widehat{0}_{M_1}) = f(\widehat{0}_{M_2})$ one must equal x and the other must equal y. Say $\widehat{0}_{M_1} = x$. Since $x \prec y$ the element y is a generator of the contraction (L,G)/x. Consequently, there is a minor (M,I) of (L,G) with $\widehat{0}_M = x$ and $I = I_1 \cup \{y\}$. Furthermore, $(M_1,I_1) = (M,I) \setminus \{y\}$ and $(M_2,I_2) = (M,I)/\{y\}$.

4.3 The main theorem

Theorem 4.3.1. Let (L,G) and (K,H) be generator enriched lattices such that there is a strong surjection from (L,G) onto (K,H). The minor poset M(K,H) can be obtained from M(L,G) via a sequence of zipping operations.

Proof. By Lemma 4.2.5 we may assume f has a single nontrivial fiber $\{x \prec y\}$. Let $F: \mathrm{M}(L,G) \to \mathrm{M}(K,H)$ be the map induced by f. By Lemma 4.2.6 the nontrivial fibers of F are of the form $\{(M,I),(M_x,I_x),(M_y,I_y)\}$ in which $(M_x,I_x)=(M,I)\setminus\{y\}$ and $(M_y,I_y)=(M,I)\setminus\{y\}$ when $x\neq \widehat{0}_M$ and $(M_y,I_y)=(M,I)/\{y\}$ when $x=\widehat{0}_M$. Let Z be the set of maximal elements of these nontrivial fibers of F. Choose some total ordering $<_{\mathrm{zip}}$ of Z with the property that if (M_1,I_1) and (M_2,I_2) are elements of Z such that $\mathrm{rk}(M_1,I_1)<_{\mathrm{rk}}(M_2,I_2)$ then $(M_1,I_1)<_{\mathrm{zip}}(M_2,I_2)$. The poset $\mathrm{M}(K,H)$ will be obtained from $\mathrm{M}(L,G)$ by identifying each of the triples $\{(M,I),(M_x,I_x),(M_y,I_y)\}$ as described above. It will be shown these identifications may be made by zipping each fiber containing an element of Z with respect to the order $<_{\mathrm{zip}}$.

Let $(M,I) \in Z$ and let (M_x,I_x) and (M_y,I_y) be the other elements of the same fiber as (M,I) as before. Let P be the poset obtained from M(L,G) by zipping elements of Z in increasing order with respect to $<_{\text{zip}}$ up to but not including the step of zipping the element (M,I). The minor poset M(L,G) is graded and thin by Lemma 3.1.3, so Proposition 4.1.6 implies that P is graded and thin as well. Let $\pi: M(L,G) \to P$ be the projection map induced by the zipping operations. By Remark 4.1.5 to show the triple $\pi(M_x,I_x), \pi(M_y,I_y), \pi(M,I)$ forms a zipper in P it suffices to show that $\pi(M,I)$ only covers the elements $\pi(M_x,I_x)$ and $\pi(M_y,I_y)$, and show that the join $\pi(M_x,I_x) \vee \pi(M_y,I_y)$ in P is $\pi(M,I)$.

We claim any minor (M',I') other than (M_x,I_x) or (M_y,I_y) covered by (M,I) satisfies $x,y\in I'\cup\{\widehat{0}_{M'}\}$. This claim is obvious when (M',I') is a deletion of (M,I). Since x is only covered by y, for any $z\in L$ either $z\vee x=z\vee y$ or $z\leq x$. Thus contracting (M,I) by an element $\ell\neq x$ either fixes both elements x and y or removes more than one generator. We conclude any minor $(M',I')\prec (M,I)$ satisfies $x,y\in I'\cup\{\widehat{0}_{M'}\}$. By construction such a minor (M',I') was the maximal element of a zipper in the construction of P and thus $\operatorname{rk}_P(\pi(M',I'))<\operatorname{rk}_{M(L,G)}(M',I')$. Hence the element $\pi(M',I')$ is not covered by $\pi(M,I)$. Therefore $\pi(M,I)$ only covers the elements $\pi(M_x,I_x)$ and $\pi(M_y,I_y)$ in the poset P.

It remains to show $\pi(M,I) = \pi(M_x,I_x) \vee \pi(M_y,I_y)$ in P. To this end, we first observe $(M,I) = (M_x,I_x) \vee (M_y,I_y)$ in M(L,G). From the definitions of (M_x,I_x) and (M_y,I_y) either $\widehat{0}_{M_x} = \widehat{0}_{M_y}$ or $\widehat{0}_{M_x} = x$. In the case the minimal element $\widehat{0}_{M_x} = x$ it is only covered by $\widehat{0}_{M_y} = y$. With these relations in mind it is straightforward to check via Lemma 3.3.5 that $(M,I) = (M_x,I_x) \vee (M_y,I_y)$ in M(L,G).

Now consider an element $p \in P$ that is an upper bound for the elements $\pi(M_x, I_x)$ and $\pi(M_y, I_y)$. By construction, the fibers $\pi^{-1} \circ \pi(M_x, I_x)$ and $\pi^{-1} \circ \pi(M_y, I_y)$ are trivial. The fact that $p > \pi(M_x, I_x)$ implies there is some minor (N_x, J_x) of (L, G) such that $(N_x, J_x) > (M_x, I_x)$ and $\pi(N_x, J_x) = p$. Similarly, there is a minor (N_y, J_y) of (L, G) such that $(N_y, J_y) > (M_y, I_y)$ and $\pi(N_y, J_y) = p$. If $(N_y, J_y) = (N_x, J_x)$ then $(N_x, J_x) \geq (M, I)$, hence $p \geq \pi(M, I)$ as desired. It remains to show this equality must hold. Since $p > \pi(M_x, I_x)$ we have that $\operatorname{rk}_P(p) \geq \operatorname{rk}_P(\pi(M_x, I_x)) + 1 = \operatorname{rk}(M, I)$. By construction no minors of (L, G) of rank greater than $\operatorname{rk}(M, I)$ were the maximal element of




Figure 10: An example of the zipping construction.

a zipper in the construction of P, so $\pi^{-1}(p)$ consists of minors with rank at most $\operatorname{rk}(M,I)$. If the fiber $\pi^{-1}(p)$ was not trivial then since P was constructed via zipping operations we have $\operatorname{rk}_P(p) \leq \operatorname{rk}(M,I) - 1 = \operatorname{rk}_P(\pi(M_x,I_x))$. Since $p > \pi(M_x,I_x)$ this cannot be the case. Hence, the fiber $\pi^{-1}(p)$ is trivial and the equality $(N_x,J_x) = (N_y,J_y)$ holds, hence $p \geq \pi(M,I)$. Therefore $\pi(M,I)$ is the join of $\pi(M_x,I_x)$ and $\pi(M_y,I_y)$ in P. This establishes the triple $\pi(M_x,I_x), \pi(M_y,I_y), \pi(M,I)$ forms a zipper in P and completes the proof.

Figure 10 depicts an example of the zipping construction.

Corollary 4.3.2. For any generator enriched lattice (L,G) the minor poset M(L,G) is Eulerian, and if $G \neq \emptyset$ the proper part of M(L,G) is a PL-sphere.

Proof. Let n = |G|. Proposition 3.1.6 says the minor poset $M(B_n, \operatorname{irr}(B_n))$ is isomorphic to the face lattice Q_n of the n-dimensional cube. In particular, this poset is Eulerian and the proper part is a PL-sphere. There is a strong surjection from $(B_n, \operatorname{irr}(B_n))$ onto (L, G), namely the canonical strong map. Thus, Theorem 4.3.1 implies the minor poset M(L, G) can be obtained from the poset Q_n via zipping operations. Theorem 4.1.9 and Theorem 4.1.7 imply the minor poset M(L, G) is Eulerian and that if $G \neq \emptyset$ then its proper part is a PL-sphere.

Corollary 4.3.3. For any generator enriched lattice (L,G) with $G \neq \emptyset$ the minor poset M(L,G) is isomorphic to the face poset of a regular CW sphere.

Proof. Since lower intervals of a minor poset are themselves minor posets, Section 4.3.2 along with Theorem 4.1.3 implies the result.

Recall the rank 1 elements of the minor poset M(L, G) are in bijection with the elements of (L, G) and the rank 2 elements are in bijection with the edges of the diagram of (L, G). The

regular CW complex whose face poset is isomorphic to M(L,G) has a 1-skeleton isomorphic to the diagram of (L,G). In particular, by Proposition 3.1.10 when (L,G) is minimally generated and geometric, the 1-skeleton is isomorphic to the Hasse diagram of L. In any case, when |G| = 3 the associated CW complex can be constructed simply by embedding the diagram of (L,G) into the 2-sphere.

Corollary 4.3.4. Let (L, G) be a generator enriched lattice with n generators. The following inequalities hold coefficientwise among the $\operatorname{\mathbf{cd}}$ -indices:

$$0 \le \Psi(\mathcal{M}(L,G)) \le \Psi(Q_n).$$

Proof. The left-hand inequality is implied by Section 4.3.2 and Theorem 4.1.8. Theorem 4.1.8 also applies to the face lattice of the n-dimensional cube, to the intermediate posets in the zipping construction of M(L,G) and to all intervals of these posets; as the proper parts of these posets are PL-spheres. Theorem 4.1.9 implies the **cd**-index of the minor poset M(L,G) may be obtained from the **cd**-index of the face lattice of the n-dimensional cube by subtracting terms which all have nonnegative coefficients.

Corollary 4.3.5. Let (L,G) and (K,H) be generator enriched lattices such that there is a strong surjection from (L,G) onto (K,H). The following inequality of $\operatorname{\mathbf{cd}}$ -indices is satisfied coefficientwise:

$$\Psi(\mathcal{M}(K,H)) \cdot \mathbf{c}^{|G|-|H|} \le \Psi(\mathcal{M}(L,G)). \tag{4.1}$$

Proof. First consider the case where |G| = |H|. By Theorem 4.3.1 we have a sequence of zipping operations that takes the minor poset M(L,G) to M(K,H). The minor poset M(L,G) has proper part a PL-sphere, hence so does every intermediate poset resulting from the sequence of zipping operations. By Theorem 4.1.8 every intermediate poset, and every interval of every intermediate poset, has a **cd**-index with nonnegative coefficients. By assumption rk(M(L,G)) = rk(M(K,H)) so no zipping operation involves the maximal element. Thus, Theorem 4.1.9 (a) implies each zipping operation corresponds, on the level of **cd**-indices, to subtracting off some **cd**-polynomial with nonnegative coefficients. Therefore, we have $\Psi(M(K,H)) \leq \Psi(M(L,G))$ coefficientwise when |G| = |H|.

Now consider the case where |G| = |H| + 1. Let $f: (L, G) \to (K, H)$ be a strong surjection. Factor the map f as in Lemma 4.2.5, and then group maps which do not decrease the number of generators. This results in a factorization $f = f_3 \circ f_2 \circ f_1$, for strong surjections

$$f_1: (L,G) \to (M,I),$$

 $f_2: (M,I) \to (N,J),$
 $f_3: (N,J) \to (K,H),$

such that |G| = |I| = |J| + 1. By the previous case we have $\Psi(M(M, I)) \leq \Psi(M(L, G))$ and $\Psi(M(K, H)) \leq \Psi(M(N, J))$ coefficientwise. Consider the zipping sequence induced by the map f_2 . Since |I| - |J| = 1 we have $\operatorname{rk}(M(M, I)) - \operatorname{rk}(M(N, J)) = 1$, thus there is

one zipping operation that involves the maximal element. By the construction given in Theorem 4.3.1 this is the final zipping operation. Let P be the result of the zipping operations applied to $\mathrm{M}(M,I)$ with the exception of the final zipping operation. The zipping operations used to construct P all correspond, on the level of \mathbf{cd} -indices, to subtracting off a \mathbf{cd} -polynomial with nonnegative coefficients. Thus, we have $\Psi(P) \leq \Psi(\mathrm{M}(M,I))$. The minor poset $\mathrm{M}(N,J)$ is constructed from P by zipping the maximal element $\widehat{1}_P$. By Theorem 4.1.9(b) we have $\Psi(P) = \Psi(\mathrm{M}(N,J)) \cdot \mathbf{c}$. We have established the following inequalities:

$$\Psi(\mathcal{M}(K,H)) \cdot \mathbf{c} \le \Psi(\mathcal{M}(N,J)) \cdot \mathbf{c} \le \Psi(\mathcal{M}(M,I)) \le \Psi(\mathcal{M}(L,G)).$$

Finally, consider the case where |G| - |H| > 1. Factor the map f as in Lemma 4.2.5 and group maps so that each group ends with a map decreasing the number of generators by one. This results in a factorization $f = f_r \circ \cdots \circ f_1$ consisting of strong surjections $f_i: (M_i, I_i) \to (M_{i+1}, I_{i+1})$ such that

$$|I_i| - |I_{i+1}| = 1,$$

 $(M_1, I_1) = (L, G),$
 $(M_{r+1}, I_{r+1}) = (K, H).$

The previous case shows $\Psi(M(M_{i+1}, I_{i+1})) \cdot \mathbf{c} \leq \Psi(M(M_i, I_i))$ for i = 1, ..., r. Setting r = |G| - |H|, it follows that $\Psi(M(K, H)) \cdot \mathbf{c}^r \leq \Psi(M(L, G))$.

Remark 4.3.6. It would be too much to expect the converse of the above corollary to hold. That is, to expect cd -index inequalities for minor posets to imply the existence of strong surjections between the associated generator enriched lattices. Indeed, the converse is false. As a counterexample, let (L, G) be the generator enriched lattice depicted in Figure 6 (b) and let (K, H) be the generator enriched lattice depicted in Figure 6 (a). There is no strong surjection from (L, G) onto (K, H), but the inequality is satisfied since

$$\Psi(\mathcal{M}(L,G)) = \mathbf{c}^3 + 2\mathbf{cd} + 3\mathbf{dc},$$

$$\Psi(\mathcal{M}(K,H)) = \mathbf{c}^3 + \mathbf{cd} + 3\mathbf{dc}.$$

To end this section we consider a special case of inequality (4.1). Namely, it is shown any generator enriched lattice with no parallels admits a strong surjection onto the chain with the same number of generators. Thus, the set of **cd**-indices of minor posets of generator enriched lattices with no parallels and n generators is minimized by the **cd**-index of the Boolean algebra B_{n+1} . This result bares some similarity to the theorem due to Ehrenborg and Karu that says the coefficientwise minimum of **cd**-indices of Gorenstein* posets is achieved by the Boolean algebra ([8, Corollary 1.3]). The bound for minor posets of generator enriched lattices with no parallels is not a special case of Ehrenborg and Karu's theorem, Figure 8 depicts a generator enriched lattice with no parallels whose minor poset is not a lattice. On the other hand, any minor poset that is a lattice must be the minor poset of a generator enriched lattice with no parallels by Theorem 3.3.6.

Lemma 4.3.7. If (L,G) is a generator enriched lattice with n generators and no parallels then there is a strong surjection onto the length n chain.

Proof. Construct a linear extension g_1, \ldots, g_n of the set G of generators as follows. For g_1 choose any element minimal in the subposet G of L. Consider the contraction $(L, G)/g_1$. By the no parallels assumption, and the fact that g_1 is minimal, the contraction $(L, G)/g_1$ must have n-1 generators. Next choose any minimal generator g of $(L, G)/g_1$. The element g corresponds to a unique generator g_2 of (L, G) satisfying $g_2 \vee g_1 = g$. Now the process repeats considering the contraction $(L, G)/\{g_1, g_2\}$. This process ends with an ordering g_1, \ldots, g_n of G. This ordering has the property that for i < j

$$q_1 \vee \cdots \vee q_i \not> q_i \vee (q_1 \vee \cdots \vee q_{i-1}).$$

In particular, the ordering is a linear extension of the poset G.

Label the elements of the length n chain C_n as $0 < 1 < \cdots < n$. Define a map

$$f:(L,G)\to (C_n,\mathrm{irr}(C_n))$$

by

$$f(\ell) = \max(\{0\} \cup \{i : g_i \le \ell\}).$$

To show f is join-preserving, let $a, b \in L$ and suppose $f(a) \vee f(b) = i$. Let j > i, by construction $g_j \not\leq g_1 \vee \cdots \vee g_i \geq a \vee b$. Thus $g_j \not\leq a \vee b$, and $f(a \vee b) < j$. Hence, $f(a \vee b) \leq f(a) \vee f(b)$. Clearly $f(a \vee b) \geq f(a) \vee f(b)$ so $f(a \vee b) = f(a) \vee f(b)$. Since the ordering g_1, \ldots, g_n is a linear extension of the subposet G of L, the image $f(g_i)$ is i, hence f is a surjection.

Corollary 4.3.8. The class of \mathbf{cd} -indices of minor posets of generator enriched lattices with n generators and no parallels is coefficientwise minimized by the \mathbf{cd} -index of the rank n+1 Boolean algebra.

Proof. Lemma 4.3.7 and Section 4.3.4 imply

$$\Psi(B_{n+1}) = \Psi(\mathcal{M}(C_n, \operatorname{irr}(C_n))) \le \Psi(\mathcal{M}(L, G)).$$

5 Concluding remarks

The original motivation for studying minor posets was an effort to describe the structure of the uncrossing poset. The uncrossing poset of rank $\binom{n}{2}$ is the poset on complete matchings of [2n] where the order relations correspond to resolving crossings in certain diagrams,

adjoined with a unique minimal element $\widehat{0}$. Certain lower intervals in the uncrossing poset are isomorphic to the minor poset of the lattice of flats of a graph derived from a diagram associated to the pairing. The structure of the entire uncrossing poset is more nuanced, the author aims to describe the uncrossing poset as a generalized minor poset in future work.

The minor poset of a generator enriched lattice is isomorphic to the face poset of a CW sphere, and such a CW sphere can be realized with piecewise linear cells. Another open question the author plans to study is whether there is some combinatorially significant realization of such CW complexes with piecewise linear cells. Certain polytopes defined from the canonical strong map appear to be a promising candidate.

Stanley posed the broad open question of finding a direct combinatorial interpretation of the coefficients of the **cd**-index for posets with a nonnegative **cd**-index in [20]. Is there a direct combinatorial interpretation of the coefficients of the **cd**-index of the minor poset of a given generator enriched lattice? A related question is whether the **cd**-index of the minor poset can be calculated directly from the generator enriched lattice in some manner.

A further question about the **cd**-indices of minor posets is whether one can derive tight bounds for the coefficients of **cd**-indices of minor posets over interesting classes of generator enriched lattices. Section 4.3.4 and Section 4.3.8 provide a tight upper and lower bound respectively for the class of generator enriched lattices with no parallels and n generators. Another class of generator enriched lattices for which this question may be tractable is minimally generated geometric lattices with n generators; and as a refinement, minimally generated geometric lattice of rank at most r with n generators. Any minimally generated geometric lattice of rank at most r is the image under a strong map of the lattice of flats of the uniform matroid of rank r with ground set [n]. On the other end, every geometric lattice with n generators admits a strong surjection onto the lattice of flats of the uniform matroid of rank 2 with ground set [n]. Thus, calculating **cd**-indices of minor posets for uniform matroids will produce tight bounds.

6 Acknowledgments

The author thanks Richard Ehrenborg and Margaret Readdy for extensive comments.

References

- [1] Margaret M. Bayer. The cd-index: a survey. In Polytopes and discrete geometry, volume 764 of Contemp. Math., pages 1–19. Amer. Math. Soc., Providence, RI, 2021.
- [2] Margaret M. Bayer and Andrew Klapper. A new index for polytopes. *Discrete Comput. Geom.*, 6(1):33–47, 1991.

- [3] Louis J. Billera and Richard Ehrenborg. Monotonicity of the cd-index for polytopes. *Math. Z.*, 233(3):421–441, 2000.
- [4] Louis J. Billera, Richard Ehrenborg, and Margaret Readdy. The c-2d-index of oriented matroids. J. Combin. Theory Ser. A, 80(1):79–105, 1997.
- [5] Garrett Birkhoff. Lattice theory. American Mathematical Society Colloquium Publications, Vol. XXV. American Mathematical Society, Providence, R.I., third edition, 1967.
- [6] A. Björner. Posets, regular CW complexes and Bruhat order. European J. Combin., 5(1):7–16, 1984.
- [7] R. P. Dilworth. Lattices with unique irreducible decompositions. *Ann. of Math.* (2), 41:771–777, 1940.
- [8] Richard Ehrenborg and Kalle Karu. Decomposition theorem for the **cd**-index of Gorenstein posets. J. Algebraic Combin., 26(2):225–251, 2007.
- [9] Richard Ehrenborg and Margaret Readdy. Coproducts and the *cd*-index. *J. Algebraic Combin.*, 8(3):273–299, 1998.
- [10] W. Gustafson. Polymatroids, closure operators and lattices. Submitted, arXiv:2112.04600.
- [11] Patricia Hersh and Richard Kenyon. Shellability of face posets of electrical networks and the CW poset property. Adv. in Appl. Math., 127:Paper No. 102178, 37, 2021.
- [12] D. A. Higgs. Strong maps of geometries. J. Combinatorial Theory, 5:185–191, 1968.
- [13] J. F. P. Hudson. *Piecewise linear topology*. W. A. Benjamin, Inc., New York-Amsterdam, 1969. University of Chicago Lecture Notes prepared with the assistance of J. L. Shaneson and J. Lees.
- [14] Kalle Karu. The cd-index of fans and posets. Compos. Math., 142(3):701–718, 2006.
- [15] Thomas Lam. The uncrossing partial order on matchings is Eulerian. *J. Combin. Theory Ser. A*, 135:105–111, 2015.
- [16] Thomas Lam. Electroid varieties and a compactification of the space of electrical networks. Adv. Math., 338:549–600, 2018.
- [17] Mark Purtill. André permutations, lexicographic shellability and the *cd*-index of a convex polytope. *Trans. Amer. Math. Soc.*, 338(1):77–104, 1993.
- [18] Nathan Reading. The cd-index of Bruhat intervals. *Electron. J. Combin.*, 11(1):Research Paper 74, 25, 2004.
- [19] Nathan Reading. Non-negative cd-coefficients of Gorenstein* posets. *Discrete Math.*, 274(1-3):323–329, 2004.

- [20] Richard P. Stanley. Flag f-vectors and the cd-index. $Math.\ Z.,\ 216(3):483-499,\ 1994.$
- [21] Richard P. Stanley. *Enumerative combinatorics. Volume 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2012.

William Gustafson

University of Kentucky Department of Mathematics Lexington KY 40506-0027 USA

william.gustafson@uky.edu