

LONG GEODESICS IN THE SPACE OF KÄHLER METRICS.

BO BERNDTSSON

ABSTRACT. We give some remarks on geodesics in the space of Kähler metrics that are defined for all time. Such curves are conjecturally induced by holomorphic vector fields, and we show that this is indeed so for regular geodesics, whereas the question for generalized geodesics is still open (as far as we know). We also give a result about the derivative of such geodesics which implies a variant of a theorem of Atiyah and Guillemin-Sternberg on convexity of the image of certain moment maps.

Dedicated to László Lempert, friend and collaborator.

1. INTRODUCTION

Let (X, ω) be a compact Kähler manifold of complex dimension n . We denote by

$$\mathcal{H}_\omega = \{u \in C^\infty; i\partial\bar{\partial}u + \omega > 0\}$$

the Mabuchi space of potentials of Kähler metrics in the same cohomology class as ω , [14]. A *complex geodesic* in this space is a complex curve $\tau \in U \rightarrow u_\tau \in \mathcal{H}_\omega$, where U is a domain in \mathbb{C} , such that $i\partial\bar{\partial}_{\tau x}u_\tau + \omega \geq 0$ and

$$(i\partial\bar{\partial}_{\tau x}u_\tau + \omega)^{n+1} = 0.$$

Here the subscript of the $\partial\bar{\partial}$ operator indicates that it is to be taken with respect to the variables τ, x jointly. If u_τ does not depend on the imaginary part of τ , this means that $u_t = u_\tau$ is a geodesic for the Riemannian structure on the Mabuchi space, see [16], [10]

For a complex geodesic

$$i\partial\bar{\partial}_{\tau x}u_\tau$$

is a $(1, 1)$ -form on $U \times X$, where U is an open set in \mathbb{C} , which we usually take to be a strip, if u_τ is independent of s . We also note that $i\partial\bar{\partial}_{\tau x}u_\tau + \omega \geq 0$ on $U \times X$ implies that u_τ is subharmonic in τ for x fixed, so u_t is a convex function of t if u_τ is independent of s .

A priori, u_τ should be smooth and strictly ω -plurisubharmonic for τ fixed, but as is customary we will allow also less regular functions (albeit always bounded), satisfying $i\partial\bar{\partial}_x u_\tau + \omega \geq 0$. Such functions are called generalized geodesics, and sometimes we will refer to the bona fide geodesics as 'regular' geodesics.

In this note we will be mainly interested in 'long' geodesics, i. e. geodesics u_t defined for all $-\infty < t < \infty$. Assuming also that u_t is of class C^1 with respect to t , we denote by \dot{u}_t the derivative of u_t with respect to t and put

$$A_t := \{\dot{u}_t(x); x \in X\}$$

and

$$B_x := \{\dot{u}_t(x); -\infty < t < \infty\}.$$

Thus A_t is the range of the velocity vector for t fixed $\dot{u}_t(x)$ as x varies, whereas B_x is the range when t varies and x is fixed. Our first result is as follows:

Theorem 1.1. *Let u_t be a geodesic of class C^1 defined for all t in \mathbb{R} that is strictly ω -plurisubharmonic for each t . Then $A_t = \overline{B_x}$ for all t and all x outside a pluripolar set in X .*

Our assumptions on the geodesic are very strong, but there is at least one class of examples: Let V be a holomorphic vector field with flow $F_\tau(x)$. This is defined for all $\tau \in \mathbb{C}$ (since X is compact), and we can define

$$\omega_\tau = F_\tau^*(\omega).$$

This a complex curve of Kähler forms on X , defined for all τ and ω_τ always lies in the same cohomology class as ω in $H^{1,1}(X)$ since F_τ is homotopic to F_0 ; the identity map. By the $\partial\bar{\partial}$ -lemma for Kähler manifolds we can write

$$\omega_\tau = i\partial\bar{\partial}u_\tau + \omega$$

for some functions u_τ , uniquely determined up to a function $f(\tau)$. Thus we get a curve in \mathcal{H}_ω , but the function $f(\tau)$ cannot always be chosen so that u_τ is a (complex) geodesic. This is, however, possible if V satisfies the cohomological condition that the contraction of the Kähler form with V is $\bar{\partial}$ -exact,

$$V \lrcorner \omega = \bar{\partial}h$$

for some function h . (It is always $\bar{\partial}$ -closed since V is holomorphic.) This condition is in particular fulfilled if the imaginary part of V is Hamiltonian for the symplectic form ω , and this in turn guarantees also that u_τ can be chosen to depend only on the real part of τ - since ω_τ then is independent of the imaginary part of τ . Hence all holomorphic vector fields whose imaginary parts are Hamiltonian furnish examples to which Theorem 1.1 applies.

The theorem has some features of an ergodic theorem, saying that averages over space equal averages over time; only here we don't really have averages, but images of maps. Moreover, the theorem says something about the real part of V , whereas it is the imaginary part that is measure preserving. A natural question in this context is whether there is an abstract counterpart of this result for measure-preserving maps that can in some sense be complexified.

We next move to an even more special situation – geodesics with multidimensional time. By this we mean functions u_t , defined for t in an open subset of \mathbb{R}^k , and whose restriction to any real line is a (one-variable) geodesic. More precisely, we have functions u_τ , with τ in a tube domain U in \mathbb{C}^k that are ω -plurisubharmonic on $U \times X$ and depend only on t , and the restriction of u_t to any real line is a geodesic. We can then define subsets A_t and B_x of \mathbb{R}^k as before, replacing \dot{u}_t by the gradient of u_t with respect to t . By essentially the same argument as in the one dimensional case and the Hahn-Banach theorem we then show that the same statement holds in the multidimensional case as well (Theorem 2.2). Since the gradient image of a convex function on \mathbb{R}^k is always convex, this gives us the corollary that the sets $A_t =: A$ all coincide, and that A is convex.

Just as in the one-dimensional case the main examples come from holomorphic vector fields whose imaginary parts are Hamiltonian, we can apply the higher dimensional case to commuting k -tuples of vector fields whose imaginary parts are Hamiltonian. As is well known, the gradient of u_t with respect to t (for, say, $t = 0$) is then the moment map of the Hamiltonian action, and

we get that the image of the moment map is convex. This is strongly related to the complex case of the celebrated theorem of Atiyah and Guillemin-Sternberg, see [2], [13].

In the next section we will give the proof of Theorem 1.1 and its multidimensional analog. In the section after that we will discuss when a holomorphic vector field induces a geodesic in the space of Kähler metrics, and give proofs of our claims above. Most likely, this material is well known, but I have not been able to find a reference. In a final section we will discuss if there are other examples of geodesics that are defined for all time $t \in \mathbb{R}$ than the ones induced by holomorphic vector fields, and we will prove that in case the geodesic is regular there are not. This is based on an analysis of the Mabuchi K-energy along such geodesics which may have an independent interest. Briefly, what we will show is the following.

It is well known that the K-energy is convex along geodesics, and that its second derivative can vanish identically only if the geodesic is induced by a holomorphic vector field. For regular geodesics, we will prove that, unless the second derivative vanishes identically, the metric on U (the strip where the geodesic is defined) defined by

$$\frac{d^2}{dt^2} K(u_t) d\tau \otimes d\bar{\tau}$$

has strictly negative curvature, bounded from above by a negative constant depending only on the volume of X and the dimension. This is clearly impossible if u_t is defined for all time, since we would then have a metric of strictly negative curvature on \mathbb{C} . Therefore the second derivative vanishes identically, which implies that the geodesic is induced by a holomorphic vector field.

This last part of the paper depends heavily on a result of Dan Burns on Monge-Ampère foliations. It also follows from recent work of Wan and Wang, [17], who, among many other things, give a direct proof of a curvature estimate from which our Theorem 4.2 follows. It seems to be an interesting problem if Theorem 4.2 also holds for generalized geodesics (in which case it is known that the K-energy is still convex).

Finally I'd like to thank Robert Berman, Sébastien Boucksom and Tamás Darvas for helpful comments, and the referee for spotting several errors and points that needed to be clarified.

2. THEOREM 1.1 AND ITS MULTIDIMENSIONAL ANALOG.

An important part in our proof is played by a construction from [5] of a certain measure on \mathbb{R} defined by a geodesic of class C^1 . The main observation is that if u_t is such a geodesic defined for $a < t < b$ (i. e. not necessarily defined for all times), and f is a continuous function on \mathbb{R} , then the integrals

$$\int_X f(\dot{u}_t) \omega_t^n / n! =: I(f)$$

do not depend on t . Thus they define a measure on \mathbb{R} , μ , the push forward of $\omega_t^n / n!$ under the map $\dot{u}_t : X \rightarrow \mathbb{R}$, and the main point is that this is the same for all t . In case the geodesic is induced by a holomorphic vector field V with Hamiltonian imaginary part, it is well known (see also the next section), that \dot{u}_0 is the Hamiltonian of $\text{Im}(V)$, and in that case μ is the measure studied by Duistermaat and Heckman, [11], in the case when the flow of $\text{Im}(V)$ is periodic.

If we assume that $\omega_t > 0$ for all t , this implies that the supremum norm on \dot{u}_t does not depend on t , since it is the right endpoint of the support of μ . It has been proved by Darvas, [9], that this

actually holds even without the positivity assumption on ω_t , and in a suitable formulation even without the C^1 -assumption on the geodesic.

In our case, when u_t is defined for all t an immediate consequence of this is that

$$u_t \leq u_0 + C|t|,$$

for some constant, so we have at most linear growth. We can therefore define a function on X

$$g(x) = \lim_{t \rightarrow \infty} u_t(x)/t;$$

the convexity in t implies that the limit exists. Being the limit of an essentially increasing sequence of functions that are $(1/t)\omega$ -plurisubharmonic, the upper semicontinuous regularization of g , g^* , is plurisubharmonic on X . Since X is compact, g^* is therefore constant, and it follows that g itself is constant outside a pluripolar set.

Now we recall the definitions from the introduction

$$A_t := \{\dot{u}_t(x); x \in X\}$$

and

$$B_x := \{\dot{u}_t(x); -\infty < t < \infty\}.$$

Since X is connected they are both intervals, and we want to prove that for most x they coincide, after taking the closure of B_x . So, fix an x_0 . Since the push-forward measure μ is independent of t and A_t is its support, $A_t = A$ does not depend on t . Moreover, $\dot{u}_t(x_0) \in A_t = A$ for all t , so it follows that $B_{x_0} \subset A$.

For the reverse inclusion, let x_0 be such that $g(x_0) = g^*(x_0)$. We use that for any x , $g(x) = \lim_{t \rightarrow \infty} \dot{u}_t(x)$, and \dot{u}_t is increasing in t . Since $g \leq g^* =: C$ everywhere it follows that

$$\dot{u}_t(x) \leq C = \lim_{t \rightarrow \infty} \dot{u}_t(x_0),$$

and $\lim_{t \rightarrow \infty} \dot{u}_t(x_0)$ is the right endpoint of the interval B_{x_0} . Replacing the geodesic u_t by u_{-t} , we find that $A \subset \overline{B_{x_0}}$, possibly after assuming that x_0 does not belong to another pluripolar set. This completes the proof of Theorem 1.1.

We next turn to the multidimensional case and the first step is a multidimensional variant of the 'Duistermaat-Heckman'-measure μ that we used above. We will consider functions $u_t(x)$ defined for x in X and for t in a convex domain in \mathbb{R}^k . As before we sometimes extend them to $\tau = t + is \in \mathbb{C}^k$ by putting $u_\tau := u_t$. We say that u_t is a multidimensional geodesic in \mathcal{H}_ω if u_τ is ω -plurisubharmonic as a function of (τ, x) and the restriction of u_t to any line in \mathbb{R}^k is a geodesic in \mathcal{H}_ω .

Theorem 2.1. *Let u_t be a multidimensional geodesic in \mathcal{H}_ω , of class C^2 in t . Let f be a continuous function on \mathbb{R}^k , and denote by $\partial_t u_t$ the gradient of u_t with respect to t . Then the integrals*

$$\int_X f(\partial_t u_t) \omega_t^n / n!$$

do not depend on t . In other words, the push-forward measures of $\omega_t^n / n!$ under the maps $\partial_t u_t$ do not depend on t and give a well defined measure $d\mu$ on \mathbb{R}^k .

The proof is essentially the same as in one variable and consists simply in differentiating with respect to t - we may of course assume that f is of class C^1 in the proof. Recall that the geodesic equation in one variable can be written

$$\ddot{u}_{tt} - |\bar{\partial}\dot{u}_t|_t^2 = 0,$$

where \ddot{u}_{tt} is the second derivative and $|\cdot|_t$ means the (pointwise) norm with respect to the Kähler metric ω_t . This means that if in the multidimensional case u_t is geodesic along each line, then

$$\ddot{u}_{ij} - \langle \bar{\partial}\dot{u}_i, \bar{\partial}\dot{u}_j \rangle_t = 0$$

for all (ij) , where we omit the t in the subscripts to make room for the variables with respect to which we differentiate. Differentiating the integral with respect to t_i we get

$$\int_X \sum_j f'_j \ddot{u}_{ij} \omega_t^n / n! + \int_X f i \bar{\partial} \bar{\partial} \dot{u}_i \wedge \omega_t^{n-1} / (n-1)!.$$

Applying Stokes' theorem to the second integral we see that this vanishes, which completes the proof of Theorem 2.1.

Now we define the sets A_t and B_x as before, replacing \dot{u}_t by the gradient with respect to t .

$$(2.1) \quad A_t := \{\partial_t u_t(x); x \in X\}$$

and

$$(2.2) \quad B_x := \{\partial_t u_t(x); -\infty < t < \infty\}.$$

Exactly as in one variable it follows from Theorem 2.1 that the sets $A_t =: A$ are all identical since they are equal to the support of $d\mu$, and that the closure of B_x is always a subset of A . Moreover, the closure of B_x is always convex, since B_x is the gradient image of a convex function. This is a standard fact that can be found e.g. in [12]. What remains is to prove that the closure of B_x fills out all of A .

For this we consider points λ on the unit sphere in \mathbb{R}^k and the geodesics

$$s \rightarrow \phi_{s\lambda} =: \Phi_s^\lambda.$$

The derivative is

$$\dot{\Phi}^\lambda = \lambda \cdot \partial_t \phi|_{s\lambda}.$$

Then

$$B_x^\lambda = \{\dot{\Phi}_s^\lambda(x), s \in \mathbb{R}\} = \{\lambda \cdot \xi; \xi \in B_x\},$$

and

$$A_s^\lambda = \{\dot{\Phi}_s^\lambda(x), x \in X\}.$$

By the result in one variable we have $A_t^\lambda = A^\lambda = B_x^\lambda$ for all x outside some pluripolar set, depending on λ . This means precisely that the orthogonal projection of A onto the line $\{s\lambda\}$ equals the projection of B_x . If we restrict attention to a countable dense set of λ 's this holds for x outside of a single pluripolar set in X . This is enough to show that A is included in the closure of B_x for all x outside this set. Indeed, if there were a point of A outside $\overline{B_x}$, we could find a hyperplane separating $\overline{B_x}$ from that point (here we are using that $\overline{B_x}$ is convex). We can

also assume this hyperplane is defined by a point λ in the countable dense set, and then get a contradiction to the one-dimensional result.

Summing up we have proved

Theorem 2.2. *Let u_t be a multidimensional geodesic in \mathcal{H}_ω , defined for t in all of \mathbb{R}^k . Then the sets A_t defined in (2.1) are all equal; $A_t =: A$, and they coincide with the closure of B_x for all x outside a pluripolar set. As a consequence, the set A is convex.*

3. HOLOMORPHIC VECTOR FIELDS THAT INDUCE GEODESICS.

Let V be a (nontrivial) holomorphic vector field on X , and let F_τ be its flow for (complex) time τ . Put $\omega_\tau = F_\tau^*(\omega)$. The question we discuss in this section is when there is a complex geodesic u_τ in \mathcal{H}_ω such that

$$\omega_\tau = i\partial\bar{\partial}u_\tau + \omega.$$

We first note that since F_τ is homotopic to the identity, ω_τ is always cohomologous to ω in $H^{1,1}(X)$. By the $\partial\bar{\partial}$ -lemma for Kähler manifolds this implies that we can always solve, for each τ

$$(3.1) \quad i\partial\bar{\partial}u_\tau = \omega_\tau - \omega,$$

and it is easy to see that we can make u_τ depend smoothly on τ . The solutions u_τ are uniquely determined up to constants for each τ , so as a function of τ we have uniqueness up to the addition of a function $f(\tau)$. The question is if this function can be chosen so that u_τ is a geodesic, i. e. that it is ω -plurisubharmonic on $\mathbb{C} \times X$ and satisfies

$$(i\partial\bar{\partial}_{\tau x}u_\tau + \omega)^{n+1} = 0.$$

Let u_τ be an arbitrary solution to (3.1), and let

$$\Omega := i\partial\bar{\partial}_{\tau x}u_\tau + \omega.$$

Let $\alpha_\tau := \bar{\partial}_x \dot{u}_\tau$, where $\dot{u}_\tau = \partial u_\tau / \partial \tau$. This is a $(0, 1)$ -form on X for each τ and it depends smoothly on τ . A direct computation shows that

$$(3.2) \quad \Omega = \omega_\tau + \ddot{u}_{\tau\bar{\tau}} i d\tau \wedge d\bar{\tau} + i(\bar{\alpha}_\tau \wedge d\bar{\tau} + d\tau \wedge \alpha_\tau),$$

and

$$\Omega^{n+1} / (n+1)! = (\ddot{u}_{\tau\bar{\tau}} - |\bar{\partial}_x \dot{u}_\tau|_\tau^2) i d\tau \wedge d\bar{\tau} \wedge \omega_\tau^n / n!.$$

Here $\ddot{u}_{\tau\bar{\tau}} = \partial^2 u_\tau / \partial \tau \partial \bar{\tau}$. We will use the simplifying notation

$$\ddot{u}_{\tau\bar{\tau}} - |\bar{\partial}_x \dot{u}_\tau|_\tau^2 =: c(u_\tau).$$

From this formula we see that if we change u_τ by adding a function $f(\tau)$, we change $c(u)$ by adding $f''(\tau)$. Hence we are able to choose u_τ so that it satisfies the HCMAE if and only if $c(u_\tau)$ depends only on τ , or equivalently if $\bar{\partial}_x c(u_\tau) = 0$. We will prove

Theorem 3.1. *Assume the holomorphic vector field V satisfies*

$$(3.3) \quad V \lrcorner \omega = i\bar{\partial}h$$

for some function h . Then there is a geodesic u_τ such that

$$F_\tau^*(\omega) = i\partial\bar{\partial}u_\tau + \omega.$$

Proof. Since V is invariant under the flow F_τ , it follows from (3.3) that

$$(3.4) \quad V \rfloor \omega_\tau = i\bar{\partial}h \circ F_\tau.$$

On the other hand, by Cartan's 'magic formula' for the Lie derivative

$$\partial V \rfloor \omega_\tau = \frac{\partial}{\partial \tau} \omega_\tau = i\partial\bar{\partial}\dot{u}_\tau.$$

Hence

$$\partial\bar{\partial}_x(h \circ F_\tau - \dot{u}_\tau) = 0,$$

so $h \circ F_\tau - \dot{u}_\tau$ is constant on X for each τ . By (3.4) this gives

$$(3.5) \quad V \rfloor \omega_\tau = i\bar{\partial}\dot{u}_\tau.$$

Next we apply $\partial/\partial\bar{\tau}$ to this and get

$$V \rfloor i\partial\bar{\partial}\dot{u}_\tau = i\bar{\partial}\ddot{u}_{\tau\bar{\tau}}.$$

This gives that

$$(3.6) \quad \bar{\partial}(V \rfloor \partial\dot{u}_\tau - \ddot{u}_{\tau\bar{\tau}}) = 0.$$

But taking the conjugate of (3.5) we get that

$$V \rfloor \partial\dot{u}_\tau = V \rfloor \bar{V} \rfloor \omega_\tau = |V|_\tau^2 = |\bar{\partial}\dot{u}_\tau|_\tau^2.$$

Hence (3.5) says precisely that $\bar{\partial}c(u_\tau) = 0$, so we are done. \square

Remark: If condition (3.3) is satisfied for a certain ω , it also holds for any other Kähler form $\omega' = \omega + i\partial\bar{\partial}v$ in the same cohomology class, with h replaced by $h + V(v)$. In particular it also holds for $G^*(\omega)$ where G is the time-1 flow of any holomorphic vector field. \square

Let us note that even though the geodesic is not unique, it can be chosen in a canonical way. It is clear that two geodesics differ only by a function of τ and this function has to be linear. Now consider the Aubin-Yau energy of u_τ ,

$$\mathcal{E}(u_\tau)$$

defined by $\mathcal{E}(u_0) = 0$ and

$$\partial\mathcal{E}(u_\tau)/\partial\tau = \int_X \dot{u}_\tau \omega_\tau^n / n!.$$

It is classical that this function is linear along geodesics (this follows also from the one-dimensional case of Theorem 2.1, since this implies that the derivative is constant), so by subtracting a suitable linear function of τ from the geodesic we can make it vanish. This is the 'canonical' choice, and it has the merit of removing all ambiguity from the definition. Theorem 3.1 can be restated as saying that if V satisfies condition (3.3), and for each τ , u_τ solves equation (3.1) and has energy zero, then u_τ is a geodesic.

We also remark that the cohomological condition (3.3) is not necessary for a geodesic to exist. A simple example is when X is the standard torus; the quotient of \mathbb{C} by the integer lattice \mathbb{Z}^2 . Take ω to be $idz \wedge d\bar{z}$ on the torus, and $V = \partial/\partial z$. Then $V \rfloor \omega = id\bar{z}$, which is not $\bar{\partial}$ -exact on the torus. But, $\omega_\tau = \omega$ for all τ , so $u_\tau = 0$ is a geodesic satisfying our conditions.

On the other hand, a geodesic as in Theorem 3.1 does not always exist. For this we take the same torus as above and the same vector field, but choose instead

$$\omega = \xi(x)idz \wedge d\bar{z}.$$

Then $\omega_\tau = \omega_t$ is independent of the imaginary part of τ and periodic in t , but not constant, if ξ is not constant, so there can be no geodesic. Indeed, the 'canonical' geodesic would then also be periodic in t , which by convexity would imply that it is constant in t . This example was communicated to me by Tamás Darvas.

Assume next that the imaginary part of V is Hamiltonian for the symplectic form ω , i. e. that

$$(V - \bar{V}) \lrcorner \omega = idH,$$

for a real valued function H ; the Hamiltonian. Identifying terms of the same bidegree, we see that $V \lrcorner \omega = i\bar{\partial}H$, so V satisfies the condition in Theorem 3.1 and therefore induces a geodesic, u_τ . In the course of the proof of Theorem 3.1 we saw also that

$$V \lrcorner \omega = i\bar{\partial}\dot{u}_0.$$

Hence \dot{u}_0 is a Hamiltonian for the imaginary part of V and the symplectic form ω , and by (3.5), \dot{u}_t is a Hamiltonian for $\text{Im}(V)$ for the symplectic form ω_t .

We next turn to multidimensional geodesics. If V is any holomorphic vector field on X , we denote by $F^V \in \text{Aut}(X)$ its time-1 flow. It is clear that if V and W are commuting holomorphic fields, then

$$F^{V+W} = F^V \circ F^W,$$

since $F^{t(V+W)}$ and $F^{tV} \circ F^{tW}$ satisfy the same ODE. Let V_1, \dots, V_k be commuting holomorphic vector fields satisfying the cohomological condition of Theorem 3.1. Put

$$F^{(\tau_1, \dots, \tau_k)} := F^{\tau_1 V_1 + \dots + \tau_k V_k},$$

$$\omega_{(\tau_1, \dots, \tau_k)} := (F^{(\tau_1, \dots, \tau_k)})^*(\omega).$$

and solve

$$i\partial\bar{\partial}u_{(\tau_1, \dots, \tau_k)} = \omega_{(\tau_1, \dots, \tau_k)} - \omega$$

with $u_{(\tau_1, \dots, \tau_k)}$ having energy zero. We claim that the restriction of $u_{(\tau_1, \dots, \tau_k)}$ to any complex line is a complex geodesic. Indeed, this is clear since if the line is $\{a + \tau b\}$ then

$$\omega_{a+\tau b} = (F^{\tau b})^*(F^a)^*(\omega)$$

satisfies the hypothesis of Theorem 3.1 (by the remark immediately after its proof). In particular, this means that if a and b lie in \mathbb{R}^k , then $u_{a+\tau b}$ is a real geodesic if the imaginary parts of V_j are all Hamiltonian. In other words, u_{t_1, \dots, t_k} is a multidimensional geodesic.

Moreover its gradient satisfies

$$\partial_t u_t|_0 =: (H_1, \dots, H_k),$$

where H_i is a Hamiltonian for V_i . In other words, the gradient of u_t at $t = 0$ is a moment map for V . Thus Theorem 2.2 has the following corollary, which is a special case of the results of Atiyah, [2], and Guillemin-Sternberg, [13].

Corollary 3.2. *Let $V = (V_1, \dots, V_k)$ be a k -tuple of commuting holomorphic vector fields, whose imaginary parts are Hamiltonian for the symplectic (Kähler) form ω . Then the image of its moment map is a convex subset of \mathbb{R}^k .*

4. GEODESICS THAT DEFINE HOLOMORPHIC VECTOR FIELDS

The main objective of this section is to prove

Theorem 4.1. *Let u_τ be a regular geodesic in \mathcal{H}_ω defined for all $\tau \in \mathbb{C}$. Assume moreover that u_τ does not depend on $\text{Im}(\tau)$. Then there is a holomorphic vector field on X that induces u_τ in the sense of the previous section.*

In the proof of the theorem we will use the Mabuchi K-energy, $K : \mathcal{H}_\omega \rightarrow \mathbb{R}$, see [15]. It is a classical fact that if u_τ is a regular complex geodesic, then $K(u_\tau)$ is a subharmonic function of τ (this was later shown to also hold for generalized geodesics with enough regularity in [4] and [8]). Moreover, if K is harmonic along a regular geodesic, then the geodesic is induced by a holomorphic vector field, [10]. This vector field may be time dependent, i. e. its coefficients may depend on τ , but it is a holomorphic function of τ , see e.g. [6]. If the geodesic is independent of the imaginary part of τ , the same thing goes for the vector field, which must therefore be independent of τ , and induce the geodesic in the sense of the previous section.

Theorem 4.2. *Let u_τ be a regular geodesic defined for τ in a domain U in \mathbb{C} . Let*

$$\Theta := \frac{\partial^2}{\partial \tau \partial \bar{\tau}} K(u_\tau) i d\tau \wedge d\bar{\tau},$$

where K is the Mabuchi K-energy. Then either Θ is identically equal to zero, or Θ defines a (possibly singular) metric on U with negative curvature, bounded from above by $-(2/nV(X))$ where $V(X)$ is the volume of X for the Kähler metric ω .

To explain the statement of the theorem, recall that if $e^v i d\tau \wedge d\bar{\tau}$ is the Kähler form of a metric on a domain in \mathbb{C} , then its curvature form is

$$-i\partial\bar{\partial}v.$$

We say that a possibly singular metric has curvature less than $-C$ if its curvature form is bounded from above by $-C$ times the metric form, or in other words if v is subharmonic and satisfies

$$i\partial\bar{\partial}v \geq C e^v i d\tau \wedge d\bar{\tau}.$$

Since there are no metrics on \mathbb{C} of strictly negative curvature, it is clear that Theorem 4.1 follows from Theorem 4.2. For a general domain U , we can also compare Θ to Ahlfors' ultrahyperbolic metric, [1] and get the following corollary.

Corollary 4.3. *For a regular geodesic, defined in a domain in \mathbb{C} , the Laplacian of the K-energy is bounded from above by $n/2V(X)$ times the ultrahyperbolic metric of the domain. In particular, for a regular geodesic ray, u_t (defined for $t > 0$)*

$$\frac{d^2}{dt^2} K(u_t) \leq (2n/V(X)) t^{-2}.$$

Thus we get uniform bounds of the second derivative of the K-energy, depending only on the dimension and the volume of X . It seems it would be interesting to know if this holds also for generalized geodesics.

The proof of Theorem 4.2 depends heavily on a remarkable result of Burns, [7]. A convenient starting point for our discussion is the formula for the $i\partial\bar{\partial}$ of the K-energy from [4]

$$(4.1) \quad i\partial\bar{\partial}K(u_\tau) = p_*((i\partial\bar{\partial}\log(\Omega^n \wedge id\tau \wedge d\bar{\tau}) \wedge \Omega^n/n!).$$

This needs to be explained a bit: First, as above,

$$\Omega = i\partial\bar{\partial}_{\tau x}u_\tau + \omega.$$

Second

$$\Omega^n \wedge id\tau \wedge d\bar{\tau}$$

is a form of top degree on $\mathbb{C} \times X$ and when we take its logarithm we mean that we take the logarithm of its density with respect to $c_nv \wedge \bar{v}$ where v is a nonvanishing holomorphic form of maximal degree. This depends on the choice of v , so the density is not well defined, but the $\partial\bar{\partial}$ of the logarithm does not depend on the choice of v . Finally, p is the projection map from $\mathbb{C} \times X$ to \mathbb{C} . In the sequel we will put

$$\theta := i\partial\bar{\partial}\log(\Omega^n \wedge id\tau \wedge d\bar{\tau}).$$

Since u_τ is a regular geodesic it is smooth and strictly ω -plurisubharmonic on X for each τ . This means that Ω is strictly positive on $\{\tau\} \times X$. Since $\Omega^{n+1} = 0$, Ω defines a foliation by holomorphic graphs of functions

$$f^x(\tau) : \mathbb{C} \rightarrow X, \quad f^x(0) = x,$$

for each x in X , along which Ω vanishes. Denote by $Y_x = \{(\tau, f^x(\tau))\}$ the graph of f^x .

Proposition 4.4. *Let $[Y_x]$ be the current of integration on Y_x . Then*

$$\Omega^n/n! = \int_X [Y_x]\omega^n/n!.$$

In the proof of the proposition we will use a well known lemma.

Lemma 4.5. *Let $f_\tau(x) := f^x(\tau)$. Then*

$$f_\tau^*(\omega_\tau) = \omega_0 = \omega.$$

Proof. Define first a time dependent vector field V_τ on X by

$$V_\tau \rfloor \omega_\tau = i\bar{\partial}\dot{u}_\tau,$$

where, as before $\dot{u}_\tau = \partial u_\tau / \partial \tau$. Then, let

$$\mathcal{V} := \partial / \partial \tau - V_\tau$$

(a vector field on $\mathbb{C} \times X$). It follows from formula (3.2) that $\mathcal{V} \rfloor \Omega = 0$. Hence \mathcal{V} is tangent to the foliation defined by Ω , so

$$\frac{\partial f_\tau}{\partial \tau} = -V_\tau.$$

This gives

$$\frac{\partial f_\tau^*(\omega_\tau)}{\partial \tau} = f_\tau^*(-V_\tau(\omega_\tau) + \dot{\omega}_\tau) = f_\tau^*(-di\bar{\partial}\dot{u}_\tau - i\partial\bar{\partial}\dot{u}_\tau) = 0.$$

Therefore $f_\tau^*(\omega_\tau)$ is constant; hence equal to ω since f_0 is the identity map. \square

We can now continue with the proof of the proposition. The statement means that if Θ is a form of bidegree $(1, 1)$, then

$$(4.2) \quad \int \Theta \wedge \Omega^n/n! = \int_X \left(\int_{Y_x} \Theta \right) \omega^n/n!.$$

Parametrizing the leaf Y_x by the map $\tau \rightarrow (\tau, f_\tau(x)) =: F_\tau(x)$ we have

$$\int_{Y_x} \Theta = \int_U \Theta(\mathcal{V}, \bar{\mathcal{V}})(F_\tau(x)) id\tau \wedge d\bar{\tau}.$$

On the other hand we have the pointwise formula

$$\Theta \wedge \Omega^n/n! = \Theta(\mathcal{V}, \bar{\mathcal{V}})\Omega^n/n! \wedge id\tau \wedge d\bar{\tau} = \Theta(\mathcal{V}, \bar{\mathcal{V}})\omega_\tau^n/n! \wedge id\tau \wedge d\bar{\tau},$$

which can be seen by contracting the identity

$$\Theta \wedge \Omega^n/n! id\tau \wedge d\bar{\tau} = 0$$

by first $\bar{\mathcal{V}}$ and then \mathcal{V} , and using that $\mathcal{V}|_\Omega = 0$ and $\mathcal{V}|_d\tau = 1$.

Hence we get in view of the lemma that

$$\begin{aligned} \int_X \left(\int_{Y_x} \Theta \right) \omega^n/n! &= \int_X \left(\int F_\tau^*(\Theta(\mathcal{V}, \bar{\mathcal{V}})\omega_\tau^n/n!) \right) id\tau \wedge d\bar{\tau} = \\ &= \int d\tau \wedge d\bar{\tau} \int \Theta(\mathcal{V}, \bar{\mathcal{V}})\omega_\tau^n/n! = \int \Theta \wedge \Omega^n/n!. \end{aligned}$$

This completes the proof of the proposition.

By the proposition and formula (4.1) we have

$$(4.3) \quad i\partial\bar{\partial}K(u_\tau) = \int_X p_*(\theta \wedge [Y_x])\omega^n/n!.$$

But

$$(4.4) \quad p_*(\theta \wedge [Y_x]) = G_x^*(\theta),$$

where $G_x : \mathbb{C} \rightarrow \mathbb{C} \times X$ is the parametrization of the leaf Y_x , so (4.3) expresses $i\partial\bar{\partial}K(u_\tau)$ as a superposition of the forms $\theta_x := G_x^*(\theta)$ with respect to the measure $\omega^n/n!$ on X .

This is where Burns' result enters the picture. In an earlier work by Bedford and Burns, [3], it was proved that θ is a positive form on each leaf. This means that the Kähler volume forms ω_τ^n are log-subharmonic along each leaf. They also found an explicit formula for $i\partial\bar{\partial}\log \omega_\tau^n$. Then, in [7], Burns proved that if we interpret this form as the Kähler form of a metric on Y_x , the Gaussian curvature of this metric is negative and bounded from above by $-2/n$. Thus the metric defined by $i\partial\bar{\partial}K(u_\tau)$ is the superposition of metrics of metrics with curvature less than $-2/n$ with respect to a positive measure of total mass equal to the volume of X for the metric defined by ω . Theorem (4.2) therefore follows from the following

Proposition 4.6. *Let g_α be (possibly singular) metrics on a domain in \mathbb{C} with curvature bounded from above by a negative constant $-a$. Let*

$$g := \int_{\alpha} g_{\alpha} d\nu(\alpha),$$

where $d\nu$ is a positive measure of total mass C . Then the curvature of g is bounded from above by $-a/C$.

Proof. Recall that the hypothesis on the curvature of g_α means that

$$\Delta \log g_\alpha \geq a g_\alpha,$$

where $\Delta = \partial^2 / \partial \tau \partial \bar{\tau}$. Equivalently

$$(4.5) \quad \Delta g_\alpha \geq a g_\alpha^2 + \frac{|\partial g_\alpha|^2}{g_\alpha}.$$

By Cauchy's inequality

$$|\partial g|^2 \leq \int \frac{|\partial g_\alpha|^2}{g_\alpha} d\nu(\alpha) \int g_\alpha d\nu(\alpha) = g \int \frac{|\partial g_\alpha|^2}{g_\alpha} d\nu(\alpha),$$

and

$$g^2 \leq C \int g_\alpha^2 d\nu(\alpha).$$

Thus the claim follows by integrating (4.5). □

REFERENCES

- [1] Ahlfors, L. Conformal invariants. Topics in geometric function theory. AMS, 2010.
- [2] Atiyah, M. Convexity and commuting Hamiltonians. Bull. London Math. Soc, 14 (1982), 1-15.
- [3] Bedford, E., Burns, D. Holomorphic mappings of annuli in \mathbb{C}^n and the associated extremal function. Ann Scuola Normali Sup di Pisa, 6, 3 (1979), pp. 381-414.
- [4] Berman, R., Berndtsson, B. . Convexity of the K-energy on the space of Kähler metrics and uniqueness of extremal metrics. Journal of the American Mathematical society, 30, (2017), pp. 1165-1196.
- [5] Berndtsson, B. Probability Measures Associated to Geodesics in the Space of Kähler Metrics. In: Hitrik M., Tamarkin D., Tsygan B., Zelditch S. (eds) Algebraic and Analytic Microlocal Analysis. AAMA 2013. Springer Proceedings in Mathematics & Statistics, vol 269. Springer, Cham. (ArXiv 0907.1806)
- [6] Berndtsson, B. Positivity of direct image bundles and convexity on the space of Kähler metrics. J. Differential Geom. 81 (2009), 457-482.
- [7] Burns, D. Curvatures of Monge-Ampère foliations and parabolic manifolds. Annals of Mathematics 115 (1982) , pp. 349-373.
- [8] Chen, XX., Li, L., Paun, M. Approximation of weak geodesics and subharmonicity of Mabuchi energy. Ann Fac Sci Toulouse, 25(5) (2016), pp. 935-957.
- [9] Darvas, Weak geodesic rays in the space of Kähler potentials and the class $\mathcal{E}(X, \omega)$. J. Inst. Math. Jussieu (2017) 16(4), 837-850
- [10] Donaldson, S. Symmetric spaces, Kähler geometry and Hamiltonian dynamics, Amer. Math. Soc. Transl. Ser. 2, vol. 196, Amer. Math. Soc., Providence RI, 1999, 13-33.
- [11] Duistermaat, J., Heckman, H., On the Variation in the Cohomology of the Symplectic Form of the Reduced Phase Space, Invent Math, 69 (1982), 259-268.

- [12] Gromov, M. Convex sets and Kähler manifolds. *Advances in J. Differential Geom.*, F. Tricerri ed., World Sci., Singapore, (1990), 1-38.
- [13] Guillemin, V., Sternberg, S. Convexity properties of the moment mapping, *Invent Math* 67, 491-513 (1982).
- [14] Mabuchi, T., Some symplectic geometry on compact Kähler manifolds I, *Osaka J. Math.* 24, 1987, 227-252.
- [15] Mabuchi, T., K-energy-maps integrating Futaki invariants. *Tohoku Math. J.* 38 (1986), 575-593.
- [16] Semmes, S.; Complex Monge-Ampère and symplectic manifolds, *Amer. J. Math.* 114 (1992), 495-550.
- [17] Wan X. , Wang, X., Poisson–Kähler fibration I: curvature of the base manifold . [arXiv:1908.03955](https://arxiv.org/abs/1908.03955).