

Integrable Zhaidary equations: reductions and gauge equivalence

Zh. Sagidullayeva^{1*}, K. Yesmakhanova^{1†}, R. Myrzakulov^{1‡}, Zh. Myrzakulova^{1§},
N. Serikbayev^{1¶}, G. Nugmanova^{1||}, A. Sergazina^{1**} and K. Yerzhanov^{1††}

¹*Ratbay Myrzakulov Eurasian International Centre for Theoretical Physics,
Astana, 010009, Kazakhstan*

Abstract

The present work addresses the study and characterization of the integrability of some spin systems (ISS) in 1+1 dimensions. Lax representations for these ISS are successfully obtained. The gauge equivalent counterparts of these integrable ISS are presented. Finally, we consider Zhanbota transcendents and some integrable Zhanbota equations. In particular, the gauge equivalence between some Zhanbota equations and the six Painleve equations is established.

Key words: Integrable equations, nonlinear Schrödinger equation, Heisenberg ferromagnet equation, Landau-Lifshitz equation, soliton equations, gauge equivalence, Zhanbota transcendents, derivative nonlinear Schrödinger equation, Zhanbota equations, spin systems, Painleve equations.

Contents

1	Introduction	2
2	Zhaidary equation	3
3	Gauge equivalent counterpart	4
4	Integrable reductions	5
4.1	Kuralay equation	5
4.2	Shynaray equation	6
5	Integrable generalizations	7
5.1	Nurshuak equation	7

*Email: zrmurzakulova@gmail.com

†Email: krmurzakulova@gmail.com

‡Email: rmyrzakulov@gmail.com

§Email: zhrmyrzakulova@gmail.com

¶Email: ns.serikbayev@gmail.com

||Email: nugmanovagn@gmail.com

**Email: asergazina@gmail.com

††Email: yerzhanovkk@gmail.com

5.2	Aizhan equation	8
5.3	Zhanbota equation	9
5.4	Akbota equation	11
6	Integrable spin systems in 1+0 dimensions: Zhanbota transcendents and Zhanbota equations	12
6.1	Zhanbota-III equation	12
6.2	Zhanbota-IV equation	13
6.3	Other Zhanbota equations and their gauge equivalent Painleve equations	14
7	Conclusion	14
8	Acknowledgments	14

1 Introduction

The Heisenberg ferromagnet equation (HFE)

$$iS_t + \frac{1}{2}[S, S_{xx}] = 0 \quad (1.1)$$

is one of the most fundamental integrable equations in soliton theory. It is well known that some integrable nonlinear evolution equations are related, often in an unexpected way, to each other by means of so-called gauge equivalences. For example, the gauge partner of the HFE is the following nonlinear Schrödinger equation (NLSE) [1]-[2]

$$iq_t + q_{xx} + 2\epsilon|q|^2q = 0. \quad (1.2)$$

Gauge equivalences between integrable nonlinear equations can be exploited to gain insight into either equation from its gauge partner, for instance by transforming solutions of one equation to solutions of the other. Formally the system can be cast into two gauge equivalent zero curvature conditions for the two sets of Lax operators. Returning to the HFE (1.1) and NLSE (1.2) we can note that they play important role in modern nonlinear physics and mathematics. They admit several integrable generalizations in 1+1 and 2+1 dimensions (see, e.g. [3]-[7] and references therein). In this paper, for example among the integrable generalizations of HFE and NLSE, we are interested in the study of their derivative versions. In particular, for the NLSE (1.2) there exist the following three celebrated derivative nonlinear Schrödinger equations (DNLSE) of this kind, the DNLSE-I, DNLSE-II and the DNLSE-III (see, e.g. [11] and references therein):

i) the Kaup-Newell equation (KNE) or DNLSE-I [8]

$$iq_t - q_{xx} - i(|q|^2q)_x = 0, \quad (1.3)$$

ii) the Chen-Lee-Liu equation (CLLE) or DNLSE-II [9]

$$iq_t - q_{xx} - i|q|^2q_x = 0, \quad (1.4)$$

iii) the Gerdjikov-Ivanov equation (GIE) or DNLSE-III [10]

$$iq_t - q_{xx} + iq^2\bar{q}_x - 0.5|q|^4q = 0. \quad (1.5)$$

Through the $U(1)$ - gauge transformation, these three DNLS can be related to each other. In fact, if $q(x, t)$ is a solution of the KNE (1.3), it is easy to find that the new field $p(x, t)$ defined as (see, e.g. [11] and references therein)

$$p(x, t) = q(x, t)e^{0.5\delta\zeta(x, t)}, \quad (1.6)$$

with

$$\zeta_x = |q|^2, \quad \zeta_t = i(q\bar{q}_x - \bar{q}q_x) + 1.5|q|^4, \quad \delta = \text{const} \quad (1.7)$$

satisfies the CLLE (1.4) for $\delta = 1$, and the GIE (1.5) for $\delta = 2$.

In the modern mathematics and physics, gauge transformations constitute an useful tool to link integrable nonlinear evolution equations (INLEE) in soliton theory, since they provide some transformations (like Darboux, Bäcklund) between those INLEE as well as the relation of their associated linear problems (Lax representations) (see, e.g. [12]-[37] and references therein). In this paper, we exploit this gauge transformation property to construct the gauge equivalent counterpart of the Zhaidary equation and to find its integrable reductions and generalizations. In particular, the gauge equivalence between some Zhanbota equations and the six Painleve equations is established.

This paper is organized as follows. In section 2 we study the Zhaidary equation (ZE). In section 3, we construct the gauge equivalent partner of the ZE. The section 4, provides the particular reductions of the ZE and their gauge equivalents and gives the associated new Lax operators in the explicit form. Some integrable generalizations of the ZE is presented in section 5. Finally in Section 6, we consider Zhanbota transcendents and integrable Zhanbota equations. Our conclusions are stated in section 7.

2 Zhaidary equation

The anisotropic Zhaidary-I equation (Z-IE) is given by [50]-[43], [48]

$$(1 + 2\beta(c\beta + d))\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} - u\mathbf{S}_x + 4cw\mathbf{S}_x - \mathbf{S} \wedge J\mathbf{S} = 0, \quad (2.1)$$

$$u_x + \frac{1}{2}(\mathbf{S}_x^2)_t = 0, \quad (2.2)$$

$$w_x + \frac{1}{4(2\beta c + d)^2}(\mathbf{S}_x^2)_t = 0, \quad (2.3)$$

where $\mathbf{S} = (S_1, S_2, S_3)$ is the unit spin vector, $\mathbf{S}^2 = 1$, $J = \text{diag}(J_1, J_2, J_3)$ ($J_1 \leq J_2 \leq J_3$), u and v are scalar functions (potentials). The isotropic Z-IE has the form

$$(1 + 2\beta(c\beta + d))\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} - u\mathbf{S}_x - 2\beta(c\beta + d)\mathbf{S}_t + 4cw\mathbf{S}_x = 0, \quad (2.4)$$

$$u_x + \frac{1}{2}(\mathbf{S}_x^2)_t = 0, \quad (2.5)$$

$$w_x + \frac{1}{4(2\beta c + d)^2}(\mathbf{S}_x^2)_t = 0. \quad (2.6)$$

The Z-IE (2.4)-(2.6) we sometime call the spin Zhaidary equation. The isotropic Z-IE is completely integrable that can be solved by the inverse scattering transformation method (IST). It possesses all the basic characters of integrable equations. The corresponding Lax representation (LR) has the form

$$\Psi_x = U_1\Psi, \quad (2.7)$$

$$\Psi_t = V_1\Psi. \quad (2.8)$$

Here

$$U_1 = [ic(\lambda^2 - \beta^2) + id(\lambda - \beta)]S + \frac{c(\lambda - \beta)}{2c\beta + d}SS_x, \quad (2.9)$$

$$V_1 = \frac{1}{1 - 2c\lambda^2 - 2d\lambda}\{[2c(\lambda^2 - \beta^2) + 2d(\lambda - \beta)]B + \lambda^2 F_2 + \lambda F_1 + F_0\}, \quad (2.10)$$

where

$$F_2 = -4ic^2wS, \quad F_1 = -4icdwS - \frac{4c^2}{2c\beta + d}wSS_x - \frac{ic}{2c\beta + d}S\{(SS_x)_t - [SS_x, B]\}, \quad (2.11)$$

$$F_0 = -\beta F_1 - \beta^2 F_2, \quad B = 0.25([S, S_t] + 2iuS), \quad S = \mathbf{S} \cdot \sigma. \quad (2.12)$$

If $\beta = 0$, the Z-IE takes the form

$$\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} - u\mathbf{S}_x + 4cw\mathbf{S}_x = 0, \quad (2.13)$$

$$u_x + \frac{1}{2}(\mathbf{S}_x^2)_t = 0, \quad (2.14)$$

$$w_x + \frac{1}{4d^2}(\mathbf{S}_x^2)_t = 0, \quad (2.15)$$

or

$$\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} + (2cd^{-2} - 1)u\mathbf{S}_x = 0, \quad (2.16)$$

$$u_x + \frac{1}{2}(\mathbf{S}_x^2)_t = 0. \quad (2.17)$$

3 Gauge equivalent counterpart

Let us find the gauge equivalent counterpart of the Z-IE (2.4)-(2.6). For this aim, consider the gauge transformation

$$\Phi = g\Psi, \quad g = \Phi|_{\lambda=\beta}, \quad S = g^{-1}\sigma_3g. \quad (3.1)$$

Then the new function Φ satisfies the following set of the linear equations

$$\Phi_x = U_2\Phi, \quad (3.2)$$

$$\Phi_t = V_2\Phi, \quad (3.3)$$

where

$$U_2 = [i(c\lambda^2 + d\lambda)\sigma_3 + (2c\lambda + d)Q], \quad (3.4)$$

$$V_2 = \frac{1}{1 - 2c\lambda^2 - 2d\lambda}(\lambda^2 B_2 + \lambda B_1 + B_0). \quad (3.5)$$

Here

$$B_2 = -4ic\sigma_3, \quad B_1 = -4icdv\sigma_3 - 2ic\sigma_3Q_t - 8c^2vQ, \quad B_0 = \frac{d}{2c}B_1 - \frac{d^2}{4c^2}B_2, \quad (3.6)$$

and

$$Q = \begin{pmatrix} 0 & q \\ r & 0 \end{pmatrix}, \quad r = \epsilon\bar{q}, \quad \epsilon = \pm 1. \quad (3.7)$$

The compatibility condition

$$U_{2t} - V_{2x} + [U_2, V_2] = 0 \quad (3.8)$$

gives the following equation

$$iq_t - q_{xt} + 4ic(vq)_x - 2d^2vq = 0, \quad (3.9)$$

$$ir_t + r_{xt} + 4ic(vr)_x + 2d^2vr = 0, \quad (3.10)$$

$$v_x - (rq)_t = 0, \quad (3.11)$$

which is called the Zhaidary-II equation (Z-IIIE).

4 Integrable reductions

The Zhaidary equations (ZE) admit several integrable reductions. Let us present these particular cases.

4.1 Kuralay equation

First, we assume that $c = 0$. Then from the Z-IE (2.4)-(2.6) we obtain the following Kuralay-I equation (K-IE) [38]-[41], [47]:

$$(1 + 2\beta d)\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} - u\mathbf{S}_x = 0, \quad (4.1)$$

$$u_x + \frac{1}{2}(\mathbf{S}_x^2)_t = 0. \quad (4.2)$$

If $\beta = 0$, this K-IE takes the form

$$\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} - u\mathbf{S}_x = 0, \quad (4.3)$$

$$u_x + \frac{1}{2}(\mathbf{S}_x^2)_t = 0. \quad (4.4)$$

The K-IE is integrable by IST method. The corresponding LR looks like

$$\Psi_x = U_3 \Psi, \quad (4.5)$$

$$\Psi_t = V_3 \Psi. \quad (4.6)$$

Here

$$U_3 = id(\lambda - \beta)S, \quad S = \begin{pmatrix} S_3 & S^- \\ S^+ & -S_3 \end{pmatrix}, \quad S^2 = I, \quad S^\pm = S_1 \pm iS_2, \quad (4.7)$$

$$V_3 = \frac{1}{1 - 2d\lambda} \{2d(\lambda - \beta)]B + \lambda^2 F_2 + \lambda F_1 + F_0\}, \quad (4.8)$$

where

$$F_2 = 0, \quad F_1 = 0, \quad F_0 = 0, \quad B = 0.25([S, S_t] + 2iuS) = 0.25Z. \quad (4.9)$$

Let us now we present the gauge equivalent counterpart of the K-IE (4.3)-(4.4). As usual, we consider the gauge transformation

$$\Phi = g\Psi, \quad g = \Phi|_{\lambda=\beta}, \quad S = g^{-1}\sigma_3 g. \quad (4.10)$$

Then the new function Φ satisfies the following set of the linear equations

$$\Phi_x = U_4 \Phi, \quad (4.11)$$

$$\Phi_t = V_4 \Phi, \quad (4.12)$$

where

$$U_4 = [id\lambda\sigma_3 + dQ, \quad (4.13)$$

$$V_4 = \frac{1}{1 - 2d\lambda}(\lambda^2 B_2 + \lambda B_1 + B_0). \quad (4.14)$$

Here

$$B_2 = 0, \quad B_1 = 0, \quad B_0 = \frac{d}{2}[-4idv\sigma_3 - 2i\sigma_3 Q_t - 8cvQ,], \quad (4.15)$$

and

$$Q = \begin{pmatrix} 0 & q \\ r & 0 \end{pmatrix}. \quad (4.16)$$

The compatibility condition

$$U_{4t} - V_{4x} + [U_4, V_4] = 0 \quad (4.17)$$

is equivalent to the following Kuralay-II equation (K-IIIE):

$$iq_t - q_{xt} - 2d^2vq = 0, \quad (4.18)$$

$$ir_t + r_{xt} + 2d^2vr = 0, \quad (4.19)$$

$$v_x - (rq)_t = 0. \quad (4.20)$$

4.2 Shynaray equation

Our next particular case is when $d = 0$. Then from the Z-IE (2.1)-(2.3) we get the following Shynaray-I equation (S-IE):

$$(1 + 2c\beta^2)\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} - u\mathbf{S}_x + 4cw\mathbf{S}_x = 0, \quad (4.21)$$

$$u_x + \frac{1}{2}(\mathbf{S}_x^2)_t = 0, \quad (4.22)$$

$$w_x + \frac{1}{16c^2\beta^2}(\mathbf{S}_x^2)_t = 0, \quad (4.23)$$

As the particular case of the integrable equation, the S-IE is also integrable by the IST method. The corresponding LR is given by

$$\Psi_x = U_5\Psi, \quad (4.24)$$

$$\Psi_t = V_5\Psi. \quad (4.25)$$

Here

$$U_5 = ic(\lambda^2 - \beta^2)S + \frac{\lambda - \beta}{2\beta}SS_x, \quad (4.26)$$

$$V_5 = \frac{1}{1 - 2c\lambda^2}\{2c(\lambda^2 - \beta^2)B + \lambda^2F_2 + \lambda F_1 + F_0\}, \quad (4.27)$$

where

$$F_2 = -4ic^2wS, \quad F_1 = -\frac{4c^2}{2c\beta}wSS_x - \frac{ic}{2c\beta}S\{(SS_x)_t - [SS_x, B]\}, \quad (4.28)$$

$$F_0 = -\beta F_1 - \beta^2 F_2, \quad B = 0.25([S, S_t] + 2iuS), \quad S = \mathbf{S} \cdot \sigma. \quad (4.29)$$

Let us find the gauge equivalent counterpart of the S-IE (4.21)-(4.23). Consider the gauge transformation

$$\Phi = g\Psi, \quad g = \Phi|_{\lambda=\beta}, \quad S = g^{-1}\sigma_3g. \quad (4.30)$$

Then the function Φ satisfies the following set of the linear equations

$$\Phi_x = U_6\Phi, \quad (4.31)$$

$$\Phi_t = V_6\Phi, \quad (4.32)$$

where

$$U_6 = ic\lambda^2\sigma_3 + 2c\lambda Q, \quad (4.33)$$

$$V_6 = \frac{1}{1-2c\lambda^2}(\lambda^2 B_2 + \lambda B_1 + B_0). \quad (4.34)$$

Here

$$B_2 = -4ic\sigma_3, \quad B_1 = -2ic\sigma_3 Q_t - 8c^2 v Q, \quad B_0 = 0. \quad (4.35)$$

Hence, the compatibility condition of the equations (4.31)-(4.32)

$$U_{6t} - V_{6x} + [U_6, V_6] = 0 \quad (4.36)$$

gives the following nonlinear evolution equation

$$iq_t - q_{xt} + 4ic(vq)_x = 0, \quad (4.37)$$

$$ir_t + r_{xt} + 4ic(vr)_x = 0, \quad (4.38)$$

$$v_x - (rq)_t = 0. \quad (4.39)$$

It is the Shynaray-II equation (S-IIIE).

5 Integrable generalizations

In the previous section we have presented some integrable reductions of the ZE (2.4)-(2.6). It is interesting to note that this ZE also admits several integrable extensions/generalizations. Now we are going to present some of these integrable generalizations of the ZE.

5.1 Nurshuak equation

One of such integrable generalizations of the ZE (2.4)-(2.6) is the following Nurshuak-I equation (N-IE):

$$iS_t + 2\epsilon_1 Z_x + i\epsilon_2(S_{xt} + [S_x, Z])_x + (wS)_x + \frac{1}{\omega}[S, W] = 0, \quad (5.1)$$

$$u_x - \frac{i}{4}tr(S \times [S_x, S_t]) = 0, \quad (5.2)$$

$$w_x - \frac{i}{4}\epsilon_2[tr(S_x^2)]_t = 0, \quad (5.3)$$

$$iW_x + \omega[S, W] = 0, \quad (5.4)$$

where

$$Z = \frac{1}{4}([S, S_t] + 2iuS). \quad (5.5)$$

As integrable equation, the N-IE admits the following LR:

$$\Psi_x = U_7 \Psi, \quad (5.6)$$

$$\Psi_t = V_7 \Psi. \quad (5.7)$$

Here

$$U_7 = -i\lambda S, \quad (5.8)$$

$$V_7 = \frac{1}{1-2\epsilon_1\lambda-4\epsilon_2\lambda^2}\{(2\epsilon_1\lambda+4\epsilon_2\lambda^2)Z + \lambda V_1 + \frac{i}{\lambda+\omega}W - \frac{i}{\omega}W\}, \quad (5.9)$$

where

$$V_1 = wS + i\epsilon_2(S_{xt} + [S_x, Z]), \quad (5.10)$$

$$W = \begin{pmatrix} W_3 & W^- \\ W^+ & -W_3 \end{pmatrix}, \quad S = \begin{pmatrix} S_3 & S^- \\ S^+ & -S_3 \end{pmatrix}, \quad S^\pm = S_1 \pm iS_2. \quad (5.11)$$

The compatibility condition $\Psi_{xt} = \Psi_{tx}$ gives the N-IE (5.1)–(5.4). The gauge partner of the N-IE (5.1)–(5.4) is the following Nurshuak-II equation (N-IIIE)

$$iq_t + \epsilon_1 q_{xt} + i\epsilon_2 q_{xxt} - vq + (wq)_x - 2ip = 0, \quad (5.12)$$

$$ir_t - \epsilon_1 r_{xt} + i\epsilon_2 r_{xxt} + vr + (wr)_x - 2ik = 0, \quad (5.13)$$

$$v_x + 2\epsilon_1(rq)_t - 2i\epsilon_2(r_{xt}q - rq_{xt}) = 0, \quad (5.14)$$

$$w_x - 2i\epsilon_2(rq)_t = 0, \quad (5.15)$$

$$p_x - 2i\omega p - 2\eta q = 0, \quad (5.16)$$

$$k_x + 2i\omega k - 2\eta r = 0, \quad (5.17)$$

$$\eta_x + rp + kq = 0, \quad (5.18)$$

where $r = \delta_1 \bar{q}$, $k = \delta_2 \bar{p}$, $\delta_j = \pm 1$. As the gauge equivalent of the integrable N-IE, this N-IIIE (5.12) – (5.18) is also integrable. Its LR reads as

$$\Psi_x = U_8 \Psi, \quad (5.19)$$

$$\Psi_t = V_8 \Psi. \quad (5.20)$$

Here

$$U_8 = -i\lambda\sigma_3 + A_0, \quad (5.21)$$

$$V_8 = \frac{1}{1 - (2\epsilon_1\lambda + 4\epsilon_2\lambda^2)} \{ \lambda B_1 + B_0 + \frac{i}{\lambda + \omega} B_{-1} \}, \quad (5.22)$$

where

$$B_1 = w\sigma_3 + 2i\epsilon_2\sigma_3 A_{0t}, \quad A_0 = \begin{pmatrix} 0 & q \\ -r & 0 \end{pmatrix}, \quad (5.23)$$

$$B_0 = -\frac{i}{2}v\sigma_3 + \begin{pmatrix} 0 & i\epsilon_1 q_t - \epsilon_2 q_{xt} + iwq \\ i\epsilon_1 r_t + \epsilon_2 r_{xt} - iwr & 0 \end{pmatrix}, \quad (5.24)$$

$$B_{-1} = \begin{pmatrix} \eta & -p \\ -k & -\eta \end{pmatrix}. \quad (5.25)$$

5.2 Aizhan equation

The Aizhan-I equation is given by

$$iS_t + i\epsilon_2(S_{xt} + [S_x, Z])_x + (wS)_x + \frac{1}{\omega}[S, W] = 0, \quad (5.26)$$

$$u_x - \frac{i}{4}tr(S \times [S_x, S_t]) = 0, \quad (5.27)$$

$$w_x - \frac{i}{4}\epsilon_2[tr(S_x^2)]_t = 0, \quad (5.28)$$

$$iW_x + \omega[S, W] = 0. \quad (5.29)$$

The Aizhan-I equation is one of integrable generalizations of the Z-IE. The Lax representation of the Aizhan-I equation has the form

$$\Psi_x = U_9 \Psi, \quad (5.30)$$

$$\Psi_t = V_9 \Psi, \quad (5.31)$$

where

$$U_9 = -i\lambda S, \quad (5.32)$$

$$V_9 = \frac{1}{1-4\epsilon_2\lambda^2} \{4\epsilon_2\lambda^2 Z + \lambda V_1 + \frac{i}{\lambda+\omega} W - \frac{i}{\omega} W\} \quad (5.33)$$

with

$$V_1 = wS + i\epsilon_2(S_{xt} + [S_x, Z]), \quad (5.34)$$

$$W = \begin{pmatrix} W_3 & W^- \\ W^+ & -W_3 \end{pmatrix}. \quad (5.35)$$

The gauge equivalent counterpart of the Aizhan-I equation (5.26)-(5.29) is the following Aizhan-II equation

$$iq_t + i\epsilon_2 q_{xxt} - vq + (wq)_x - 2ip = 0, \quad (5.36)$$

$$ir_t + i\epsilon_2 r_{xxt} + vr + (wr)_x - 2ik = 0, \quad (5.37)$$

$$v_x - 2i\epsilon_2(r_{xt}q - rq_{xt}) = 0, \quad (5.38)$$

$$w_x - 2i\epsilon_2(rq)_t = 0, \quad (5.39)$$

$$p_x - 2i\omega p - 2\eta q = 0, \quad (5.40)$$

$$k_x + 2i\omega k - 2\eta r = 0, \quad (5.41)$$

$$\eta_x + rp + kq = 0. \quad (5.42)$$

This Aizhan-II equation is integrable. The corresponding Lax representation is given by

$$\Phi_x = U_{10}\Phi, \quad (5.43)$$

$$\Phi_t = V_{10}\Phi, \quad (5.44)$$

where

$$U_{10} = -i\lambda\sigma_3 + A_0, \quad (5.45)$$

$$V_{10} = \frac{1}{1-4\epsilon_2\lambda^2} \{\lambda B_1 + B_0 + \frac{i}{\lambda+\omega} B_{-1}\}. \quad (5.46)$$

Here

$$B_1 = w\sigma_3 + 2i\epsilon_2\sigma_3 A_{0t}, \quad (5.47)$$

$$A_0 = \begin{pmatrix} 0 & q \\ -r & 0 \end{pmatrix}, \quad (5.48)$$

$$B_0 = -\frac{i}{2}v\sigma_3 + \begin{pmatrix} 0 & -\epsilon_2 q_{xt} + iwq \\ \epsilon_2 r_{xt} - iwr & 0 \end{pmatrix}, \quad (5.49)$$

$$B_{-1} = \begin{pmatrix} \eta & -p \\ -k & -\eta \end{pmatrix}. \quad (5.50)$$

5.3 Zhanbota equation

In this section, we consider the Zhanbota-I equation. It has the form [44]

$$\mathbf{S}_t - \mathbf{S} \wedge \mathbf{S}_{xt} - u\mathbf{S}_x - \frac{1}{\omega} \mathbf{S} \wedge \mathbf{W} = 0, \quad (5.51)$$

$$u_x + \mathbf{S} \times (\mathbf{S}_x \wedge \mathbf{S}_t) = 0, \quad (5.52)$$

$$\mathbf{W}_x - \omega \mathbf{S} \wedge \mathbf{W} = 0. \quad (5.53)$$

The Zhanbota-I equation is also one of integrable generalization of the Zhaidary-I equation. The matrix form of the Zhanbota-I equation reads as

$$iS_t + \frac{1}{2}[S, S_{xt}] + iuS_x + \frac{1}{\omega}[S, W] = 0, \quad (5.54)$$

$$u_x - \frac{i}{4}tr(S[S_x, S_t]) = 0, \quad (5.55)$$

$$iW_x + \omega[S, W] = 0, \quad (5.56)$$

where, $S = S_i\sigma_i$, $W = W_i\sigma_i$, ($i = 1, 2, 3$) and ω is a constant parameter. The $\mathbf{W} = (W_1, W_2, W_3)$ is the vector potential. The Zhanbota-I equation possesses the following Lax representation:

$$\Psi_x = U_{11}\Psi, \quad (5.57)$$

$$\Psi_t = V_{11}\Phi. \quad (5.58)$$

Here, the matrix operators U_{11} and V_{11} have the forms

$$U_{11} = -i\lambda S, \quad (5.59)$$

$$V_{11} = \frac{1}{1-2\lambda}\{\lambda V_1 + \frac{i}{\lambda+\omega}W - \frac{i}{\omega}W\}, \quad (5.60)$$

where

$$V_1 = 2Z = \frac{1}{2}([S, S_t] + 2iuS), \quad (5.61)$$

$$W = \begin{pmatrix} W_3 & W^- \\ W^+ & -W_3 \end{pmatrix}. \quad (5.62)$$

Let us find the gauge equivalent counterpart of the Zhanbota-I equation (5.51) - (5.53). It is not difficult to verify that the gauge equivalent counterpart of the Zhanbota-I equation is the following Zhanbota-II equation:

$$q_t + \frac{\kappa}{2i}q_{xt} + ivq - 2p = 0, \quad (5.63)$$

$$r_t - \frac{\kappa}{2i}r_{xt} - ivr - 2k = 0, \quad (5.64)$$

$$v_x + \frac{\kappa}{2}(rq)_t = 0, \quad (5.65)$$

$$p_x - 2i\omega p - 2\eta q = 0, \quad (5.66)$$

$$k_x + 2i\omega k - 2\eta r = 0, \quad (5.67)$$

$$\eta_x + rp + kq = 0, \quad (5.68)$$

where q, r, p, k are some complex functions; v, η are real potential functions and κ is a constant parameter. The Zhanbota-II equation is integrable. The Lax representation of the Zhanbota-II equations (5.63) - (5.68) reads as

$$\Phi_x = U_{12}\Phi, \quad (5.69)$$

$$\Phi_t = V_{12}\Phi, \quad (5.70)$$

where

$$U_{12} = -i\lambda\sigma_3 + A_0, \quad (5.71)$$

$$V_{12} = \frac{1}{1-\kappa\lambda}\{B_0 + \frac{i}{\lambda+\omega}B_{-1}\}. \quad (5.72)$$

Here

$$A_0 = \begin{pmatrix} 0 & q \\ -r & 0 \end{pmatrix}, \quad (5.73)$$

$$B_0 = -\frac{i}{2}v\sigma_3 - \frac{\kappa}{2i} \begin{pmatrix} 0 & q_y \\ r_y & 0 \end{pmatrix}, \quad (5.74)$$

$$B_{-1} = \begin{pmatrix} \eta & -p \\ -k & -\eta \end{pmatrix}. \quad (5.75)$$

Next, we consider the reduction $r = \delta_1 \bar{q}$, $k = \delta_2 \bar{p}$ with $\kappa = 2$, $\delta_j = \pm 1$, where the bar means the complex conjugate. Then, the Zhanbota-II equation (5.63)–(5.68) takes the following more compact form

$$iq_t + q_{xt} - vq - 2ip = 0, \quad (5.76)$$

$$v_x + 2\delta_1(|q|^2)_t = 0, \quad (5.77)$$

$$p_x - 2i\omega p - 2\eta q = 0, \quad (5.78)$$

$$\eta_x + (\delta_1 \bar{q}p + \delta_2 \bar{p}q) = 0. \quad (5.79)$$

Note that it is nothing but the Zhanbota-II equation.

5.4 Akbota equation

In this subsection, we consider the following Akbota-I equation [45]–[46], [47]

$$\mathbf{S}_t - \mathbf{S} \wedge (\alpha S_{xx} + \beta \mathbf{S}_{xt}) - u \mathbf{S}_x = 0, \quad (5.80)$$

$$u_x + \mathbf{S} \times (\mathbf{S}_x \wedge \mathbf{S}_t) = 0. \quad (5.81)$$

This Akbota-I equation is one of the integrable generalizations of the Zhaidary-I equation. It has the following Lax representation

$$\Psi_x - U_{13}\Psi = 0, \quad (5.82)$$

$$\Psi_t - V_{13}\Psi = 0, \quad (5.83)$$

where

$$U_{13} = \frac{i}{2}\lambda S, \quad V_{13} = \frac{1}{1-\lambda\beta} \left\{ \alpha \left(\frac{1}{2}i\lambda^2 S + \frac{1}{4}[S, S_x] \right) + \beta \lambda Z \right\}. \quad (5.84)$$

Note that the gauge equivalent counterpart of the Akbota-I equation is the following Akbota-II equation

$$iq_t + \alpha q_{xx} + \beta q_{xt} + vq = 0, \quad (5.85)$$

$$v_x - 2[\alpha(|q|^2)_x + \beta(|q|^2)_t] = 0. \quad (5.86)$$

The Lax representation of this Akbota-II equation is given by

$$\Phi_x = U_{14}\Phi, \quad (5.87)$$

$$\Phi_t = V_{14}\Phi, \quad (5.88)$$

where

$$U_{14} = \frac{i\lambda}{2}\sigma_3 + Q, \quad Q = \begin{pmatrix} 0 & \bar{q} \\ q & 0 \end{pmatrix}, \quad V_{14} = \frac{1}{1-\lambda\beta} \left\{ \frac{i\lambda^2}{2}\alpha\sigma_3 + \alpha\lambda Q + V_0 \right\} \quad (5.89)$$

with

$$V_0 = \begin{pmatrix} \alpha i|q|^2 + i\beta \partial_x^{-1}|q|_t^2 & -i\beta \bar{q}_t - i\alpha \bar{q}_x \\ i\beta q_y + \alpha i q_x & -[\alpha i|q|^2 + i\beta \partial_x^{-1}|q|_t^2] \end{pmatrix}. \quad (5.90)$$

6 Integrable spin systems in 1+0 dimensions: Zhanbota transcendents and Zhanbota equations

Interesting subclass of the integrable spin systems is integrable Zhanbota equations in 1+1 and 1+0 dimensions. These integrable Zhanbota equations are called as:

the Zhanbota-I equation,
the Zhanbota-II equation,
the Zhanbota-III equation,
the Zhanbota-IV equation,
the Zhanbota-V equation,
the Zhanbota-VI equation,
the Zhanbota-VII equation,
the Zhanbota-VIII equation

etc. As integrable equations, these Zhanbota equations admit Lax representations, infinite number conservation laws, Hamiltonian structures and so on. Among of these Zhanbota equations, six Zhanbota equations, namely, the Zhanbota-III, Zhanbota-IV, Zhanbota-V, Zhanbota-VI, Zhanbota-VII and Zhanbota-VIII equations have the especial property. Zhanbota transcendents are solutions of these six Zhanbota equations in the complex plane. Note that these six Zhanbota equations are not generally solvable in terms of elementary functions. It is interesting to note that these six Zhanbota equations correspond to the famous six Painleve equations. This remarkable property of these six Zhanbota equations we listed in the following Table 1:

Table 1. Some integrable Zhanbota equations and their gauge equivalent counterparts.

No	Name	Equivalent counterpart	References
1	Zhanbota-III equation	Painleve-I equation	[2205.02073]
2	Zhanbota-IV equation	Painleve-II equation	[2205.02073]
3	Zhanbota-V equation	Painleve-III equation	[2205.02073]
4	Zhanbota-VI equation	Painleve-IV equation	[2205.02073]
5	Zhanbota-VII equation	Painleve-V equation	[2205.02073]
6	Zhanbota-VIII equation	Painleve-VI equation	[2205.02073]

Table 1. In this table we present (listed) some integrable (1+0)-dimensional Zhanbota equations (that is in 1+0 dimensions) as well as their gauge equivalent counterparts. In this section, we want to present some of these six Zhanbota equations.

6.1 Zhanbota-III equation

One of interesting integrable spin systems in 1+0 dimensions is the Zhanbota-III equation. It has the form [48]-[50]

$$S_{xx} + \frac{1}{2}tr(S_x^2)S - \frac{1}{2}(\ln(tr(S_x^2)))_x S_x + (\beta + \epsilon\sqrt{tr(S_x^2)})[S, S_x] = 0, \quad (6.1)$$

where $\epsilon = \pm 1$ and β is some complex constant. This Zhanbota-III equation is integrable. The corresponding Lax representation is given by

$$\Psi_x = U_{III}\Psi, \quad (6.2)$$

$$\Psi_\lambda = V_{III}\Psi, \quad (6.3)$$

where

$$U_{III} = \left(\lambda - \beta + \frac{q_{III}}{\lambda} - \frac{q_{III}}{\beta} \right) S + \frac{\beta}{4} \left(\frac{1}{\lambda} - \frac{1}{\beta} \right) [S, S_x], \quad (6.4)$$

$$V_{III} = (4\lambda^4 + 2q_{III}^2 + x)S + \frac{\beta}{2q_{III}} \left(2\lambda q_{IIIx} + \frac{1}{2\lambda} \right) S_x + \frac{\beta}{4q_{III}} (4\lambda^2 q_{III} + 2q_{III}^2 + x) [S, S_x] \quad (6.5)$$

with

$$q_{III} = \pm 0.5\beta\sqrt{\mathbf{S}_x^2}, \quad \mathbf{S}_x^2 = 0.5\text{tr}(S_x^2), \quad q_{III}^2 = 0.25\beta^2\mathbf{S}_x^2 = \frac{\beta^2}{8}\text{tr}(S_x^2). \quad (6.6)$$

The compatibility condition of the linear equations (6.2)-(6.3) $\Psi_{x\lambda} = \Psi_{\lambda x}$ that is

$$U_{III\lambda} - V_{IIIx} + [U_{III}, V_{III}] = 0 \quad (6.7)$$

is equivalent to the Zhanbota-III equation (6.1). It is interesting to find the gauge equivalent counterpart of the Zhanbota-III equation. As it follows from the Table 1, the gauge equivalent counterpart of the Zhanbota-III equation is the Painleve-I equation. To prove it, consider the following well known representation of the spin matrix S:

$$S = \Phi^{-1}\sigma_3\Phi, \quad (6.8)$$

where $\Phi(x, \lambda)$ is a complex matrix. Substituting this formula to the Zhanbota-III equation (6.1), we obtain the following set of two linear equations

$$\Phi_x = U'_{III}\Phi, \quad (6.9)$$

$$\Phi_\lambda = V'_{III}\Phi, \quad (6.10)$$

where

$$U'_{III} = \left(\lambda + \frac{q_{III}}{\lambda} \right) \sigma_3 - \frac{iq_{III}}{\lambda} \sigma_2, \quad (6.11)$$

$$V'_{III} = (4\lambda^4 + 2q_{III}^2 + x)\sigma_3 - i(4\lambda^2 q_{III} + 2q_{III}^2 + x)\sigma_2 - (2\lambda q_{IIIx} + \frac{1}{2\lambda})\sigma_1. \quad (6.12)$$

From the compatibility condition of the two linear equations (6.9)-(6.10) that is from $\Phi_{x\lambda} = \Phi_{\lambda x}$, we obtain

$$U'_{III\lambda} - V'_{IIIx} + [U'_{III}, V'_{III}] = 0. \quad (6.13)$$

This equation gives us the following famous Painleve-I equation

$$q_{IIIxx} - 6q_{III}^2 - x = 0. \quad (6.14)$$

Thus we have proved that the Zhanbota-III equation (6.1) and the Painleve-I equation (6.14) is gauge equivalent to each other. Note that takes place the following formulas:

$$\Psi = g^{-1}\Phi, \quad g = \Phi|_{\lambda=\beta}. \quad (6.15)$$

6.2 Zhanbota-IV equation

Another example of integrable spin systems in 1+0 dimensions is the Zhanbota-IV equation. The Zhanbota-IV equation reads as [48]-[50]

$$S_{xx} + \frac{1}{2}\text{tr}(S_x^2)S - \frac{1}{2} \left(\ln \left(-\frac{\text{tr}(S_x^2)}{8} \right) \right)_x S_x - i\beta[S, S_x] = 0, \quad (6.16)$$

where β is an complex constant. As integrable equation, this Zhanbota-IV equation admits the Lax representation, infinite number of conservation laws and so on. For example, its Lax representation has the form

$$\Psi_x = U_{IV}\Psi, \quad (6.17)$$

$$\Psi_\lambda = V_{IV}\Psi, \quad (6.18)$$

where

$$U_{IV} = -i(\lambda - \beta)S, \quad (6.19)$$

$$V_{IV} = -i(4\lambda^2 + 2q_{IV}^2 + x)S + \frac{iq_{IVx}}{q_{IV}}S_x + \left(\lambda - \frac{\alpha}{4\lambda q_{IV}}\right)[S, S_x]. \quad (6.20)$$

In this case

$$q_{IV} = \pm i0.25\sqrt{\mathbf{S}_x^2}, \quad \mathbf{S}_x^2 = 0.5tr(S_x^2), \quad q_{IV}^2 = -0.25\mathbf{S}_x^2 = -\frac{1}{8}tr(S_x^2). \quad (6.21)$$

From the compatibility condition of the linear equations (6.9)-(6.10) $\Psi_{x\lambda} = \Psi_{\lambda x}$:

$$U_{IV\lambda} - V_{IVx} + [U_{IV}, V_{IV}] = 0 \quad (6.22)$$

we obtain the Zhanbota-IV equation (6.8). As in the previous subsection, we can prove that the gauge equivalent counterpart of the Zhanbota-IV equation (6.16) is the following Painleve-II equation:

$$q_{IVxx} - 2q_{IV}^3 - xq_{IV} - \alpha = 0. \quad (6.23)$$

6.3 Other Zhanbota equations and their gauge equivalent Painleve equations

As in the previous subsections, we can prove that the Zhanbota-V, Zhanbota-VI, Zhanbota-VII and Zhanbota-VIII equations are gauge equivalent to the Painleve-III, Painleve-IV, Painleve-V and Painleve-VI equations, respectively. The corresponding results we will publish in the other place.

7 Conclusion

In this paper, we have shown that the Zhaidary equation is integrable by the IST method. Its gauge equivalent equation is constructed. Some integrable particular cases (reductions) is studied. Also several integrable generalizations of the ZE are presented. For these reductions and generalizations their gauge partners and their Lax representations are found. Various issues are worthy of further exploration. In particular, we hope to investigate the discrete, nonlocal and fractional integrable versions of the ZE, its reductions and generalizations as well as solutions and geometry of the above presented integrable equations in the future. For example, the ZE and related integrable equations admit a geometric interpretation that directly relates the curvature and torsion of a vector field to any their solution. We will demonstrate that these geometrical relations and interpretations can be extended to the extended discrete, nonlocal and fractional integrable versions of the ZE. Finally, we consider Zhanbota transcendents and integrable Zhanbota equations. In particular, the gauge equivalence between some Zhanbota equations and the famous six Painleve equations is established.

8 Acknowledgments

This work was supported by the Ministry of Education and Science of Kazakhstan, Grant AP08857372.

References

- [1] M. Lakshmanan. Phys. Lett. A, **64**, 53-54 (1977)
- [2] V. E. Zakharov, L. A. Takhtajan, Theor. Math. Phys., **38**, 17-23 (1979)
- [3] Y. Ishimori. *Multi-vortex solutions of a two-dimensional nonlinear wave equation*, Prog. Theor. Phys., **72**, 33-37 (1984)
- [4] Chen Chi, Zhou Zi-Xiang. *Darboux Transformation and Exact Solutions of the Myrzakulov-I Equation*, Chin. Phys. Lett., **26**, N8, 080504 (2009)
- [5] Chen Hai, Zhou Zi-Xiang. *Darboux Transformation with a Double Spectral Parameter for the Myrzakulov-I Equation*, Chin. Phys. Lett., **31**, N12, 120504 (2014)
- [6] Hai Chen, Zi-Xiang Zhou. *Global explicit solutions with n double spectral parameters for the Myrzakulov-I equation*, Modern Physics Letters B, **30**, N29, 1650358 (2016)
- [7] Hai-Rong Wang, Rui Guo. *Soliton, breather and rogue wave solutions for the Myrzakulov-Lakshmanan-IV equation*, Optik, **242**, 166353 (2021)
- [8] D. Kaup and A. C. Newell. *An exact solution for a derivative nonlinear Schrödinger equation*, J. Math. Phys. **19**, 798-801 (1978)
- [9] H. H. Chen, Y. C. Lee and C. S. Liu. *Integrability of Nonlinear Hamiltonian Systems by Inverse Scattering Method*, Phys. Scr., **20**, 490-492 (1979)
- [10] V. Gerdjikov and I. Ivanov. *A quadratic pencil of general type and nonlinear evolution equations. II. Hierarchies of Hamiltonian structures*, Bulg. J. Phys. **10**, 130-143 (1983)
- [11] Paz Albares. *Integrability and rational soliton solutions for gauge invariant derivative nonlinear Schrödinger equations*, [arXiv:2102.12183]
- [12] Li-Yuan Ma, Shou-Feng Shen, Zuo-Nong Zhu. *Integrable nonlocal complex mKdV equation: soliton solution and gauge equivalence*, [arXiv:1612.06723]
- [13] Li-Yuan Ma, Zuo-Nong Zhu. *Nonlocal nonlinear Schrödinger equation and its discrete version: soliton solutions and gauge equivalence*, [arXiv:1503.06909]
- [14] Julia Cen, Francisco Correa, Andreas Fring. *Nonlocal gauge equivalence: Hirota versus extended continuous Heisenberg and Landau-Lifschitz equation*, [arXiv:1910.07272]
- [15] V. S. Gerdjikov, G. G. Grahovski, N. A. Kostov. *On N -wave Type Systems and Their Gauge Equivalent*, [arxiv:0111027]
- [16] A. Kundu. *Landau-Lifshitz and higher-order nonlinear systems gauge generated from nonlinear Schrödinger-type equations*, J. Math. Phys., **25**, 3433-3438 (1984)
- [17] M. Wadati and K. Sogo. *Gauge Transformations in Soliton Theory*, J. Phys. Soc. Jpn., **52**, 394-398 (1983)
- [18] R. Myrzakulov, A. Danlybaeva and G. Nugmanova. Theor. and Math. Phys., V.118, 13, P. 441-451 (1999).
- [19] R. Myrzakulov, G. Mamyrbekova, G. Nugmanova, M. Lakshmanan. Symmetry, **7**(3), 1352-1375 (2015).
- [20] Z. S. Yersultanova, M. Zhassylbayeva, K. Yesmakhanova, G. Nugmanova, R. Myrzakulov. *Darboux transformation and exact solutions of the integrable Heisenberg ferromagnetic equation with self-consistent potentials*, Int. Jour. Geom. Meth. Mod. Phys., **13**, N1, 1550134 (2016).

- [21] R. Myrzakulov, G. N. Nugmanova, R. N. Syzdykova. *Gauge equivalence between (2+1)-dimensional continuous Heisenberg ferromagnetic models and nonlinear Schrödinger-type equations*, J. Phys. A: Math. Gen., **31**, N47, 9535-9545 (1998)
- [22] R. Myrzakulov, S. Vijayalakshmi, R. Syzdykova, M. Lakshmanan, *On the simplest (2+1) dimensional integrable spin systems and their equivalent nonlinear Schrödinger equations*, J. Math. Phys., **39**, 2122-2139 (1998).
- [23] V.G. Makhankov, R. Myrzakulov. *σ -Model Representation of the Yajima-Oikawa Equation System*, Preprint P5-84-719, JINR, Dubna, 1984
- [24] V. G. Makhankov, R. Myrzakulov, O. K. Pashaev. *Gauge Equivalence, Supersymmetry and Classical Solutions of the ospu(1,1/1) Heisenberg Model and the Nonlinear Schrödinger Equation*, Letters in Mathematical Physics, **16**, 83-92 (1988).
- [25] R. Myrzakulov, S. Vijayalakshmi, G.N. Nugmanova, M. Lakshmanan. *A (2+1)-dimensional integrable spin model: Geometrical and gauge equivalent counterpart, solitons and localized coherent structures*, Phys. Lett. A, **233**, 391 (1997). [arXiv:solv-int/9704005]
- [26] Nevin Ertug Gürbüz, R. Myrzakulov, Z. Myrzakulova. *Three anholonomy densities for three formulations with anholonomic coordinates with hybrid frame in R_1^3* , Optik, 169161 (2022)
- [27] R. Myrzakulov, O. K. Pashaev, Kh. T. Kholmurodov. *Particle-Like Excitations in Many Component Magnon-Phonon Systems*, Physica Scripta, **33**, N4, 378 (1986)
- [28] R. Myrzakulov, M. Lakshmanan, S. Vijayalakshmi, A. Danlybaeva, J. Math. Phys., **39**, 3765-3771 (1998).
- [29] R. Myrzakulov, G. Nugmanova, R. Syzdykova, Journal of Physics A: Mathematical & Theoretical, **31** 147, 9535-9545 (1998).
- [30] R. Myrzakulov, M. Daniel, R. Amuda, Physica A., **234**, 13-4, 715-724 (1997).
- [31] S.C. Anco, R. Myrzakulov, Journal of Geometry and Physics, v.60, 1576-1603 (2010).
- [32] R. Myrzakulov, G. K. Mamyrbekova, G. N. Nugmanova, K. Yesmakhanova, M. Lakshmanan. Physics Letters A, **378**, 30-31, 2118-2123 (2014).
- [33] Z. Myrzakulova, G. Nugmanova, K. Yesmakhanova, N. Serikbayev, R. Myrzakulov. *Integrable generalized Heisenberg ferromagnet equations with self-consistent potentials and related Yajima-Oikawa type equations*, DOI: 10.13140/RG.2.2.19991.85923
- [34] L. Martina, Kur. Myrzakul, R. Myrzakulov, G. Soliani, Journal of Mathematical Physics, V.42, 13, P.1397-1417 (2001).
- [35] A. Myrzakul, G. Nugmanova, N. Serikbayev, R. Myrzakulov. *Surfaces and Curves Induced by Nonlinear Schrödinger-Type Equations and Their Spin Systems*, Symmetry, **13**, N10, 1827 (2021)
- [36] A. Myrzakul and R. Myrzakulov. *Integrable geometric flows of interacting curves/surfaces, multilayer spin systems and the vector nonlinear Schrödinger equation*, International Journal of Geometric Methods in Modern Physics, **14**, N10, 1750136 (2017)
- [37] A. Myrzakul and R. Myrzakulov. *Integrable motion of two interacting curves, spin systems and the Manakov system*, International Journal of Geometric Methods in Modern Physics, **14**, N7, 1750115 (2017)
- [38] Z. Sagidullayeva, G. Nugmanova, R. Myrzakulov and N. Serikbayev. *Integrable Kuralay equations: geometry, solutions and generalizations*, Symmetry, 14(7), 1374 (2022). [arXiv:2206.05348]

- [39] Waqas Ali Faridi, Muhammad Abu Bakar, Zhaidary Myrzakulova, Ratbay Myrzakulov, Ali Akgul, Sayed M. El Din. *The formation of solitary wave solutions and their propagation for Kuralay equation*, Results in Physics, **52**, 106774 (2023)
- [40] Hai-Yang Kong, Rui Guo. *Breathers, rogue waves and semi-rational solutions for a Heisenberg ferromagnet-type equation*. Modern Physics Letters B, **37**, No20, 2350053 (2023). <https://doi.org/10.1142/S0217984923500537>
- [41] Asim Zafar, Muhammad Raheel, Mohamed R. Ali, Zhaidary Myrzakulova, Ahmet Bekir, Ratbay Myrzakulov. *Exact Solutions of M-Fractional Kuralay Equation via Three Analytical Schemes*. Symmetry, **15**, N10, 1862 (2023).
- [42] Z. Myrzakulova, G. Nugmanova, K. Yesmakhanova, N. Serikbayev, R. Myrzakulov. *Integrable generalized Heisenberg ferromagnet equations with self-consistent potentials and related Yajima-Oikawa type equations*
- [43] Z. Myrzakulova, G. Nugmanova, K. Yesmakhanova, R. Myrzakulov. *Integrable motion of anisotropic space curves and surfaces induced by the Landau-Lifshitz equation*, [arXiv:2202.00748]
- [44] Qin-Ling Liu, Hui-Qin Hao, Rui Guo. *Multiple nonlinear wave solutions of a generalized Heisenberg ferromagnet model and their interactions*, Modern Physics Letters B, <https://doi.org/10.1142/S0217984924500349>
- [45] Hai-Yang Kong, Rui Guo. *Dynamic behaviors of novel nonlinear wave solutions for the Akbota equation*, Optik, **282**, 170863 (2023). <https://doi.org/10.1016/j.ijleo.2023.170863>
- [46] Thilagarajah Mathanaranjan, Ratbay Myrzakulov. *Conservation laws, soliton solutions and stability analysis for the Akbota equation*, DOI:10.13140/RG.2.2.32430.08006
- [47] Kuralay Yesmakhanova, Zhaidary Myrzakulova, Ratbay Myrzakulov, Gulgasyl Nugmanova. *Soliton surfaces induced by the Kuralay, Kairat and Akbota equations*, DOI: 10.13140/RG.2.2.16410.41928 . Conference: 15th Symposium on Integrable Systems, Bialystok, Poland, June 29-30, 2023
- [48] Zh. Sagidullayeva, K. Yesmakhanova, R. Myrzakulov. Zh. Myrzakulova, N. Serikbayev, G. Nugmanova, A. Sergazina, K. Yerzhanov. *Integrable Zhaidary equations: reductions and gauge equivalence*, [DOI: 10.13140/RG.2.2.31425.74082]
- [49] A.Sergazina, Z.Myrzakulova, K.Yesmakhanova, R.Myrzakulov, G.Nugmanova. *Integrable spin systems in 2+1, 1+1 and 1+0 dimensions*, [DOI:10.13140/RG.2.2.36416.30724]
- [50] Zh. Sagidullayeva, K. Yesmakhanova, R. Myrzakulov, Zh. Myrzakulova, N. Serikbayev, G. Nugmanova, A. Sergazina, K. Yerzhanov. *Integrable Zhaidary equations: reductions and gauge equivalence*, [arXiv:2205.02073]