

Nonequilibrium symmetry-protected topological order: emergence of semilocal Gibbs ensembles

Maurizio Fagotti,* Vanja Marić,* and Lenart Zadnik*
Université Paris-Saclay, CNRS, LPTMS, 91405, Orsay, France

We consider nonequilibrium time evolution in quantum spin chains after a global quench. We show that global symmetries can invalidate the standard picture of local relaxation to a (generalised) Gibbs ensemble and provide a solution to the problem. We introduce in particular a family of statistical ensembles, which we dub “semilocal (generalised) Gibbs ensembles”. The issue arises when the Hamiltonian possesses conservation laws that are not (pseudo)local but act as such in the symmetry-restricted space where time evolution occurs. Because of them, the stationary state emerging at infinite time can exhibit exceptional features. We focus on a specific example with a spin-flip symmetry, which is the commonest global symmetry encountered in spin-1/2 chains. Among the exceptional properties, we find that, at late times, the excess of entropy of a spin block triggered by a local perturbation in the initial state grows logarithmically with the subsystem’s length. We establish a connection with symmetry-protected topological order in equilibrium at zero temperature and study the melting of the order induced either by a (symmetry-breaking) rotation of the initial state or by an increase of the temperature.

CONTENTS

	20		20
I. Introduction	2	V. Signatures of semilocal order	20
II. Overview and results	3	A. Reduced density matrices across different theories of local observables	20
A. Local observables in spin chains	3	1. Restricted duality transformation	20
1. Quasilocality	3	2. Dual reduced density matrices	21
2. Semilocality	3	B. Excess of entropy	22
3. Cluster decomposition	4	1. Example: Dual generalised XY model	23
4. Semilocal <i>vs.</i> symmetry-protected topological order	4	2. Free-fermion techniques for the entropies	24
B. Conservation laws with semilocal densities	5	3. Results	25
1. Breakdown of maximum-entropy descriptions	5	C. Melting of the order	26
2. Semilocal (generalised) Gibbs ensemble	6	1. Slow relaxation after weak symmetry breaking	26
C. Signatures of semilocal order	6	2. Melting at low finite temperatures	27
1. Excess of entropy	7	VI. Discussion	28
2. Melting of the order	7	Open problems	28
III. Long time limit: state of the art	8	Acknowledgments	29
A. Background	8	A. Convergence of the iTEBD data	29
B. Maximum-entropy descriptions	9	B. $\mathbb{Z}_2 \times \mathbb{Z}_2$ duality	29
1. Example: XY model	10	C. Zero-temperature phases of the dual XY model	30
2. Example: Dual XY model	11	D. Tilted initial state	30
C. Failure of Gibbs and generalised Gibbs ensemble in symmetric systems	12	1. Tilted state $ \not\mathcal{A}_\theta\rangle_\sigma$	30
IV. Beyond (generalised) Gibbs ensembles	13	2. Decay of correlations in the state $ \not\mathcal{A}_\theta\rangle_\sigma$	31
A. Kramers-Wannier duality	13	E. Excess of entropy	32
B. Semilocality and hidden symmetry breaking	14	1. Bounds on Renyi entropies	32
C. Semilocal (generalised) Gibbs ensembles	16	2. Correlation matrix after the quench	32
1. Example: Dual XY model revisited	16	3. Asymptotic results for Renyi entropies	33
2. Example: generic model revisited	18	4. Exact Procedure	34
3. Canonical and non-canonical descriptions	18	F. Melting of the order and scaling limits	35
4. Remark	19	1. Effective density matrix	35
		2. Semilocal order parameter	37
		References	37

* The authors have contributed equally.

I. INTRODUCTION

There is no topological order in one-dimensional systems in equilibrium at zero temperature; the ground states of gapped spin-chain Hamiltonians are all equivalent to trivial product states [1]. When restricting to systems with a given symmetry, however, it becomes possible to distinguish different phases. On the one hand there are standard disordered and ordered Landau phases characterised by symmetry breaking; on the other hand, there can be nontrivial symmetry-protected topological phases, in which the order is manifested in excitations, such as the celebrated edge modes [2–7], or in exceptional entanglement properties [8].

Focusing on symmetry-protected topological order, perhaps the most prominent feature that is accessible in the bulk of the system is the so-called string order [9–11]. It refers to the existence of sequences of bounded operators with arbitrarily large support (e.g., strings of spins) whose expectation values remain nonzero in the limit of infinite support. String order was shown not to survive an increase in the temperature [12], and most results pertaining to its fate when the system is not kept in equilibrium [12–20] seem to point at its melting.

This comes of no surprise for the physical community that has been working on relaxation in nonequilibrium systems. Specifically, there has been a lot of progress in predicting the behaviour of local observables in integrable and generic systems composed of a macroscopic number of degrees of freedom [21–28]. It is now well accepted that, in an isolated many-body system, local relaxation occurs: finite regions of the full system relax because the rest of the system acts as an effective bath for them. At late times, local observables can be effectively described by statistical ensembles incorporating conservation laws with certain locality properties [29, 30], the simplest example being a conserved operator with a local density. In the end the effective stationary state can be thought of as a thermal state of an effective Hamiltonian. Since Ref. [12] ruled out string order at finite temperature, there is little if any hope to keep string order out of equilibrium.

In this work we reconsider symmetry-protected topological order from the perspective of local relaxation after a global quench in a quantum spin chain. We show that string order does not always melt down and explain a mechanism behind its persistence. We study the corresponding exotic nonequilibrium states emerging at late times. Embracing Occam’s razor for the description of such novel states of matter forces us to a fundamental change of view. We question the meaning of locality in a quantum spin chain and build on the traditional representation of local observables; we reformulate the theory in order for it to become consistent with the recent hints pointing at the emergence of symmetry-protected topological order after global quantum quenches [31].

The incriminated systems possess hidden symmetries that can not be associated with local conservation laws.

In infinite chains the corresponding charges lie outside of the theory defined on the local operators and their quasilocal completion, in turn invalidating standard descriptions in terms of maximum-entropy statistical ensembles. This is observed both in generic and integrable systems. In the generic case, the persistence of topological order impairs local relaxation to an effective thermal state [32], together with related results such as the celebrated eigenstate thermalisation hypothesis [32–35]. In integrable systems, instead, we experience the failure of the generalised Gibbs ensemble [36], the latter not being able to describe the infinite-time limit even when defined in its more refined version that includes every pseudolocal conservation law [29].

We overcome this problem by introducing two novel statistical ensembles: the *G-semilocal Gibbs ensemble* and the *G-semilocal generalised Gibbs ensemble*, which live in an extension of the theory, characterised by the symmetry G . These ensembles are able to capture string order in a setting where the absence of string order is the norm: they are the defining feature of a novel phase of nonequilibrium matter. This stands in stark contrast with the situation in equilibrium, in which string order is not sufficient to characterise a topological phase: even a trivial product state exhibits string order.

Purely for the sake of simplicity, we focus on the commonest symmetry in spin-1/2 chains — the invariance under a spin flip (a \mathbb{Z}_2 symmetry). We show the failure of maximum-entropy statistical descriptions both in generic and in integrable systems and report quantitative results for the simplest model we know to exhibit a \mathbb{Z}_2 -semilocal generalised Gibbs ensemble: the dual XY model, which, to the best of our knowledge, has been introduced in Ref. [37]. We elaborate on the observations of Ref. [31] about the macroscopic everlasting effects of a local perturbation like a spin flip in the nonequilibrium time evolution of systems described by Hamiltonians such as that of the dual XY model. While a similar phenomenology had been observed before in the case of symmetry-breaking perturbations of ground states [38, 39], and excited states in a jammed sector [40, 41], the question of whether or not these situations are just different facets of the same phenomenon remains open.

We take inspiration from Refs. [42, 43], on one side, and Ref. [44], on the other, and investigate the late-time effect of the perturbation on the Rényi entropies

$$S_\alpha[A] = \frac{1}{1-\alpha} \log \text{tr}(\rho_A^\alpha) \quad (1.1)$$

of spin blocks A (ρ_A is the reduced density matrix of A). We show that a symmetry-breaking localised perturbation results in a logarithmic correction to the standard extensive behaviour:

$$S_\alpha[A] \xrightarrow{t \rightarrow \infty} a_\alpha |A| + b_\alpha \log |A| + \mathcal{O}(|A|^0), \quad (1.2)$$

where $|A|$ is the subsystem’s length. In the examples considered the prefactor b_α of the logarithm is computed

analytically. Strictly speaking, b_α is neither quantised nor universal, but it is nevertheless nonzero and depends on only few system details.

Finally, we address the question of the instability of the symmetry-protected topological phase from the constructive point of view of the separation of time scales [45–48]. Specifically, we consider two ways of breaking the relevant symmetry: applying a weak symmetry-breaking transformation to the initial state or heating it up to a finite low temperature. As expected, string order melts down, but it does so over a time scale that is much longer than the times that can be reached, for example, in numerical simulations based on tensor networks.

II. OVERVIEW AND RESULTS

We consider time evolution after a global quench in an infinite spin-chain system that, in almost the entire paper, is assumed to be translationally invariant. That is to say, we investigate systems prepared in the ground state or in a thermal state of a given pre-quench Hamiltonian $\mathbf{H}_0 = \sum_\ell \mathbf{h}_{0,\ell}$ and then time-evolved with a different post-quench Hamiltonian $\mathbf{H} = \sum_\ell \mathbf{h}_\ell$, e.g.,

$$\begin{aligned} |\Psi(t)\rangle &= e^{-i\mathbf{H}t} |\Psi(0)\rangle, \\ (\mathbf{H}_0 |\Psi(0)\rangle &= E_{\text{GS}} |\Psi(0)\rangle), \end{aligned} \quad (2.1)$$

where E_{GS} is the ground state energy of \mathbf{H}_0 . We restrict ourselves to local Hamiltonian densities $\mathbf{h}_{0,\ell}$, \mathbf{h}_ℓ (see discussion below). The focus is on systems that are invariant under a global (sometimes called “on-site”) symmetry characterised by some global unitary operator \mathbf{U} , such that $\mathbf{U}\mathbf{h}_\ell\mathbf{U}^\dagger = \mathbf{h}_\ell$ and $\mathbf{U}\mathbf{h}_{0,\ell}\mathbf{U}^\dagger = \mathbf{h}_{0,\ell}$.

We have three main goals:

1. Show that the established picture of local relaxation to a Gibbs or a generalised Gibbs ensemble can fail in symmetric systems;
2. Uncover the reasons behind the failure of the standard descriptions and define a novel family of statistical ensembles able to capture the stationary values of local operators;
3. Identify signatures of the exotic nonequilibrium phases.

We warn the reader that, in order to achieve our objectives, we need to distinguish the notion of “local observable” from that of “local operator”. The latter is for us only a representation of the former, as will be clarified later.

We now provide an informal overview and contextualisation of our results.

A. Local observables in spin chains

In quantum mechanics the concept of locality is connected with the algebra of the operators representing the

observables. In practical terms, an observable is local if it can be associated with a position in such a way that its measurement does not affect the measurement of any other sufficiently distant local observable. We wrote “sufficiently distant” because in a spin chain local observables are not point-like objects but have a range corresponding to the size of the finite subsystem affected by their measurement. Generically local observables are represented by local operators of the form $\mathbf{O}_A \otimes \mathbf{I}_{\bar{A}}$, where \mathbf{O}_A has support in some finite connected subsystem A of the lattice, \bar{A} denoting its complement. The range of the corresponding local observable is then $|A|$.

1. Quasilocality

Time evolution does not preserve the locality of an operator: if the Hamiltonian is not exceptionally simple, the Heisenberg representation of an operator that is local at time $t = 0$ does not have a finite range at any $t \neq 0$. This problem can be resolved by weakening the definition of locality so as to include also observables represented as limits of sequences of local operators ordered by their range. For example $\sum_{n=1}^{\infty} e^{-n} \sigma_{\ell-n}^z \sigma_{\ell+n}^z$ is not local but is quasilocal: it can be well approximated by the truncated sums, and hence the limit is well defined. For local Hamiltonians it is actually sufficient to restrict ourselves to the strong form of quasilocality in which operators have exponentially decaying tails. A careful formalisation of this intuitive idea eventually results in the definitions of Refs. [30, 49–52].

Time evolution with local Hamiltonians preserves strong quasilocality [53, 54], but quantum mechanics is still a non-relativistic theory: causality, usually expressed as a commutation of quasilocal operators at space-like distances, does not hold. Nonetheless a weaker form of causality still applies: Lieb and Robinson proved that the commutator of quasilocal operators becomes exponentially small at space-like distances, providing a bound that plays the role of the speed of light in relativistic systems [53].

2. Semilocality

Exponentially localised operators are usually regarded as the quintessential representation of local observables in spin chains. Still, there are systems with local observables that escape the boundaries of quasilocality. Let us consider for example a quantum quench between two Hamiltonians with local densities invariant under a spin flip \mathcal{P}^z :

$$\begin{aligned} \mathbf{h}_j &= \mathcal{P}^z[\mathbf{h}_j] \equiv \lim_{n \rightarrow \infty} \left[\prod_{\ell=-n}^n \sigma_\ell^z \right] \mathbf{h}_j \left[\prod_{\ell'=-n}^n \sigma_{\ell'}^z \right], \\ \mathbf{h}_{0,j} &= \mathcal{P}^z[\mathbf{h}_{0,j}], \end{aligned} \quad (2.2)$$




FIG. 1. Panel (a): odd operator $\mathbf{O}_o(x)$ ($\mathbf{O}_o(y)$) with support to the left (right) of the site j commutes (anticommutes) with the string $\Pi^z(j)$. For simplicity we assume $x \ll j \ll y$. Panel (b): even operator with support that does not include the site j commutes with the string $\Pi^z(j)$. Restricting to the space of even operators, the strings behave as local objects.

where σ_ℓ^z denotes the operator acting like the Pauli z matrix on site ℓ and like the identity elsewhere. Suppose that the initial state is spin-flip symmetric. Thus, the expectation value of any local observable \mathbf{O}_o which is odd under spin flip, i.e., $\mathcal{P}^z[\mathbf{O}_o] = -\mathbf{O}_o$, vanishes. The symmetry of the Hamiltonian moreover implies that this property holds true at any time. The system is completely characterised by operators \mathbf{O}_e that are even under spin flip, i.e., $\mathcal{P}^z[\mathbf{O}_e] = \mathbf{O}_e$; without loss of information, we can restrict the space of operators to the even ones.

Such a restriction has however a subtle by-product: in the restricted space there are observables that are not represented by local operators but that nevertheless behave as such. In this specific case the prototype of such an observable is represented by an operator $\Pi^z(j)$ that plays the role of a product of Pauli matrices σ^z extending from site j to infinity; see Section IV A for a proper definition. As done in Ref. [31], we call $\Pi^z(j)$ semilocal to remember that its action is local only in the restricted space of even operators — see Fig. 1. This example with a \mathbb{Z}_2 (spin-flip) symmetry will be our testing ground, but the idea can be applied to other symmetries: if every subsystem has a density matrix that is written at any time in terms of a restricted set of local operators,

an alternative representation of local observables can be obtained by supplementing the restricted set with operators that are not local but behave as such. As in the \mathbb{Z}_2 case, we call them “semilocal”.

Duality. We expect quite generally that, once extended to include also semilocal operators, the restricted space becomes isomorphic to the original space of quasilocal operators. In the \mathbb{Z}_2 case, this is manifested in the existence of a *duality transformation* that preserves the locality of the Hamiltonian and maps even semilocal operators into local ones that are odd under a spin flip (see Section IV A). Analogous duality transformations exist also when the system exhibits other symmetries — see, e.g., Refs. [55, 56].

3. Cluster decomposition

Thermal states and ground states of local Hamiltonians have clustering properties. That is to say, the expectation value of the product of two local operators far away from each other factorises:

$$\lim_{y \rightarrow \infty} (\langle \mathbf{O}_1(x) \mathbf{O}_2(x+y) \rangle - \langle \mathbf{O}_1(x) \rangle \langle \mathbf{O}_2(x+y) \rangle) = 0. \quad (2.3)$$

Cluster decomposition is intimately connected with the phenomenon of symmetry breaking in ordered phases: generic nonsymmetric perturbations break the symmetry of the ground state in such a way that the state always ends up satisfying clustering [57]. If our system is symmetric, semilocal operators represent local observables as well, therefore we require that

a symmetric physical state satisfies clustering also for semilocal operators.

This additional requirement allows us to completely break the barrier between semilocality and locality: there is no physical property distinguishing semilocal operators from local ones. In view of this, local, (strongly) quasilocal, and semilocal operators can all represent local observables. Incidentally, we mention that this equivalence becomes natural when taking the continuum limit close to a critical point, where all the aforementioned operators become represented by local fields.

In the previous example of the \mathbb{Z}_2 symmetry, clustering fixes the expectation value of the semilocal operators up to an overall sign. Section IV B will show that the auxiliary sign is determined by how the symmetry is broken in the dual representation.

4. Semilocal vs. symmetry-protected topological order

We are considering noncritical spin-chain Hamiltonians with discrete global symmetries. If, in equilibrium at zero temperature, some of the symmetries remain unbroken, the state can exhibit so-called symmetry-protected topological order [58]. Because of the symmetry of the Hamiltonian, we can find string operators \mathbf{O}_A that act differently from the identity everywhere within a connected region A and commute with the energy densities \mathbf{h}_ℓ with support inside A . The order is manifested in the fact that there are operators of that kind with a nonzero expectation value in the limit $|A| \rightarrow \infty$. This is usually referred to as “string order” [55].

We reinterpret string order as the existence of an alternative representation of local observables including semilocal operators with a nonzero expectation value.

As far as we can see, indeed, it is always possible to reinterpret $\langle \mathbf{O}_A \rangle$ as the correlation between two semilocal operators positioned at the boundaries of A . Clustering of

semilocal operators together with the nonvanishing value of $\langle \mathbf{O}_A \rangle$ in the limit $|A| \rightarrow \infty$ imply that the expectation values of those semilocal operators are different from zero. In analogy with the local order parameters characterising a phase with a spontaneously broken symmetry, we call them semilocal order parameters.

The reader can understand this as an alternative way of presenting the results of Refs. [55, 56], which relate symmetry-protected topological order to a standard Landau phase (with a broken symmetry) in a dual representation. For example, in the \mathbb{Z}_2 case $\lim_{n \rightarrow \infty} \langle \prod_{j=\ell-n}^{\ell+n} \sigma_j^z \rangle$ is the natural string-order parameter and we have $\prod_{j=\ell-n}^{\ell+n} \sigma_j^z = \Pi^z(\ell-n)\Pi^z(\ell+n+1)$.

Since we have not carefully investigated the equivalence between semilocal order and symmetry-protected topological order, we will refrain from using the latter terminology when there is a risk that our claims could not hold in full generality. We note however that the basic property of symmetry-protected topological order is satisfied: semilocal order is preserved if the initial state is perturbed without breaking the symmetry associated with the order. For example, in the case of a \mathbb{Z}_2 -symmetric initial state $|\Psi(0)\rangle$ the transformed state $e^{i\mathbf{W}}|\Psi(0)\rangle$ retains the symmetry for any spin-flip invariant operator \mathbf{W} with strongly quasilocal density. The collection of such states obtained by all possible symmetry-preserving perturbations forms a symmetry-protected phase [1]. All these states are characterised by a nonvanishing expectation value of a semilocal operator.

B. Conservation laws with semilocal densities

Most spin-flip invariant Hamiltonians with local densities do not admit dynamics in which the expectation values of semilocal operators remain nonzero also at late times. This can be readily understood if, like in the \mathbb{Z}_2 case, there is a duality transformation mapping semilocal operators into local ones that break the symmetry. In the dual representation their expectation value remains nonzero only if there are conserved charges that break the symmetry of the dual Hamiltonian.

Conversely, for every symmetric Hamiltonian with at least one local charge that doesn't respect the symmetry, there is a dual Hamiltonian with at least one conservation law with a semilocal density. In the specific case of spin-flip symmetric Hamiltonians, the simplest examples of systems of this kind have dual Hamiltonians that are spin-flip invariant in all the directions, like the Hamiltonian of the Heisenberg model. In particular,

any local Hamiltonian with a spin-flip symmetry and a $U(1)$ charge that breaks spin flip is dual to a model with a semilocal charge.

The prototypical example of the latter has the following Hamiltonian

$$\mathbf{H} = \sum_{\ell \in \mathbb{Z}} \sigma_{\ell-1}^x (\mathbf{1} - \sigma_{\ell}^z) \sigma_{\ell+1}^x + \mathbf{W}_{\ell}[\{\sigma^z\}], \quad (2.4)$$

where \mathbf{W}_{ℓ} can be any local operator written only in terms of σ^z . To the best of our knowledge, for generic \mathbf{W}_{ℓ} the model does not have conservation laws with quasilocal densities. On the other hand, just using the algebra satisfied by $\Pi^z(j)$, the reader can verify that \mathbf{H} has the following conservation law with a semilocal density:

$$\mathbf{Q} = \sum_{\ell \in \mathbb{Z}} \Pi^z(\ell). \quad (2.5)$$

Following Refs. [30, 52] literally, we would exclude this conservation law from the description of the long-time behaviour of the expectation values of local operators. Indeed $\Pi^z(\ell)$ cannot be written as a limit of a sequence of local operators and does not even have a nonzero infinite-temperature overlap with a local operator. Is this charge relevant after a global quench?

1. Breakdown of maximum-entropy descriptions

After a quantum quench of a global parameter at zero temperature, the local properties of the state approach those of an effective stationary state. The standard picture is that the latter is completely characterised by (pseudo)local integrals of motion carrying information about the initial state, and it can be represented by the maximum-entropy ensemble satisfying those constraints — see Section III B. As such, the emerging state, which is called Gibbs or generalised Gibbs ensemble, is expected to resemble an equilibrium state at finite temperature. We remind the reader that finite-temperature states in one dimension do not generally exhibit topological order; this was proven, in particular, for global onsite symmetries [12]. Because of the similarity between (generalised) Gibbs ensembles and thermal states, it is reasonable to expect that, in such maximum-entropy descriptions, the two-point functions of semilocal operators approach zero at large distances.

Let us now re-examine global quenches in symmetric systems in the light of our previous considerations about semilocality. If the state before the quench is not symmetric (i.e., its density matrix is not even), semilocal operators do not represent local observables, exactly as in global quenches without global symmetries. We expect, therefore, the emergence of a standard (generalised) Gibbs ensemble; we will consider an explicit example in Section III B 1.

The situation changes if the initial state exhibits semilocal order. It is simple to show that

semilocal charges enable the possibility to keep memory of semilocal order,

that is to say, they allow a symmetry-protected topological order phase to survive the long-time limit after a quantum quench.

In the \mathbb{Z}_2 case, this can be readily proved considering the fluctuations of a semilocal charge, let it be $\sum_{j \in \mathbb{Z}} \mathbf{q}_j^{\text{sl}}$,

with a nonzero expectation value at the initial time. We indeed have

$$\lim_{t \rightarrow \infty} \frac{1}{|A|^2} \sum_{j,n \in A} \langle \Psi(t) | \mathbf{q}_j^{\text{sl}} \mathbf{q}_n^{\text{sl}} | \Psi(t) \rangle \geq \langle \Psi(0) | \mathbf{q}^{\text{sl}} | \Psi(0) \rangle^2 \quad (2.6)$$

for every subsystem A , where we used the non-negativity of fluctuations and $\langle \Psi(t) | \mathbf{q}^{\text{sl}} | \Psi(t) \rangle = \langle \Psi(0) | \mathbf{q}^{\text{sl}} | \Psi(0) \rangle$. In the limit of large $|A|$ the left-hand side of Eq. (2.6) can only be nonzero if the state at infinite time exhibits string order, specifically, if $\lim_{n \rightarrow \infty} \langle \mathbf{q}_j^{\text{sl}} \mathbf{q}_{j+n}^{\text{sl}} \rangle \neq 0$. Thus, a nonzero initial expectation value $\langle \Psi(0) | \mathbf{q}^{\text{sl}} | \Psi(0) \rangle \neq 0$ implies survival of the string order in the limit of infinite time. Being blind to a string order, a (generalised) Gibbs ensemble constructed with the pseudolocal charges of the model (see Section III B for the definition of pseudolocal-ity) can not describe such a limit. For an explicit example of the breakdown of the maximum-entropy descriptions in integrable and generic systems we refer the reader to Section III C.

2. Semilocal (generalised) Gibbs ensemble

The charges responsible for the persistent order can be accommodated in a minimal extension of the standard theory with quasilocal operators, which must now include also semilocal operators. In such an extended space we define two novel maximum-entropy statistical ensembles: the G -semilocal Gibbs ensemble and the G -semilocal generalised Gibbs ensemble, where G refers to the symmetry exploited to enlarge the theory (e.g., the \mathbb{Z}_2 symmetry) — see Section IV C. The former emerges in generic systems, in which there is a finite number of local and semilocal conservation laws; the latter incorporates instead the richer structure of integrable systems, where there are infinitely many pseudolocal and semilocal charges. They are supposed to capture the infinite-time limit as the corresponding Gibbs and generalised Gibbs ensembles do in non-symmetric systems.

We posit an additional step: the semilocal ensemble should be projected back onto a theory of local observables. Indeed, not all operators in the extended theory can represent local observables (they do not satisfy the causality principle sketched in Section II A) and the choice of which of them can is ambiguous: in the spin-flip example ($G = \mathbb{Z}_2$), odd operators and even operators with half-infinite strings do not commute at infinite distance but they both, separately, can supplement the even quasilocal operators to represent local observables — see Fig. 2.

That said, local observables are typically represented by quasilocal operators (excluding therefore, in the spin-flip case, operators with half-infinite strings). In such a theory

persistent semilocal order manifests itself in the fact that the projected ensemble does




FIG. 2. Quasilocal theory (red line) and semilocal theory (blue dashed line). Each of them contains operators that are able to represent local observables, provided that the operators of the other theory are excluded (even semilocal and odd quasilocal operators can not simultaneously represent local observables — see Fig. 1).

not maximise the entropy constrained by the quasilocal integrals of motion.

Note however that, in all the cases we envisage, there is also an alternative representation of local observables in which the projected ensemble is a maximum-entropy state as in Ref. [59].

In conclusion, symmetric systems allow for multiple representations of local observables and only a part of them, which we will refer to as “canonical” theories, include the densities of all the operators that are conserved in the extended space — see Section IV C 3.

C. Signatures of semilocal order

The fact that the two-point function of semilocal operators does not vanish in the limit of infinite distance is arguably the most evident signature of semilocal order in nonequilibrium states (see Section V C for a study of a semilocal order parameter in the dual XY model). In this respect, we only add a remark:

the semilocal integrals of motion are semilocal order parameters with the additional property of being stationary.

In fact, this is only part of the story. That a symmetric system can be described in alternative ways (e.g., by a quasilocal vs. by a semilocal theory) can have striking consequences. In particular, perturbations that generally have insignificant effects can trigger macroscopic changes. Two phenomena of this kind are described in the following.

1. Excess of entropy

Ref. [31] has recently shown that, in a symmetric quench, a localised perturbation to the initial state affects the stationary values of local operators. Here we make the next step, studying the effect of the perturbation on the entanglement properties of large subsystems. We parametrise the perturbation by an odd transformation \mathbf{U}_ℓ that connects different symmetry sectors and is localised around some position ℓ (in our specific example we will consider $\mathbf{U}_\ell = \sigma_\ell^x$): $|\Psi(0)\rangle \rightarrow \mathbf{U}_\ell |\Psi(0)\rangle$. The reader can think of \mathbf{U}_ℓ as of the result of a projective measurement of a local operator — see also Ref. [40]. Since the perturbation is local, it has a limited effect on the initial state. In particular, it does not affect at all the entanglement entropies of subsystems that enclose the support of \mathbf{U}_ℓ completely.

After a global quench the entropies of spin blocks grow in time until reaching an extensive value [60]. If there are no semilocal charges, the effect of \mathbf{U}_ℓ approaches zero in the limit of infinite time. On the other hand, semilocal charges keep memory of the perturbation and even the entanglement entropies of large subsystems remain affected. We consider in particular the *excess of entropy*, which was recently studied in states at zero temperature in which a symmetry is spontaneously broken [43]. It is defined as the increase in the entanglement entropies produced by the local perturbation

$$\begin{aligned} \Delta_{\mathbf{U}_\ell} S_A(t) = & S[\text{tr}_A(e^{-i\mathbf{H}t} \mathbf{U}_\ell |\Psi(0)\rangle \langle \Psi(0)| \mathbf{U}_\ell^\dagger e^{i\mathbf{H}t})] \\ & - S[\text{tr}_A(e^{-i\mathbf{H}t} |\Psi(0)\rangle \langle \Psi(0)| e^{i\mathbf{H}t})], \end{aligned} \quad (2.7)$$

where $S[\rho]$ can be any functional of the density matrix ρ that measures the entanglement between A and the rest. We will focus on the Rényi entropies $S[\rho] = [1/(1 - \alpha)] \log \text{tr}(\rho^\alpha)$. Remarkably,

$\Delta_{\mathbf{U}_\ell} S_A(t)$ does not approach zero in the limit of infinite time; it becomes proportional to the logarithm of the subsystem's length

$$\Delta_{\mathbf{U}_\ell} S_A(t) \propto \log |A|. \quad (2.8)$$

In Section VB we compute the prefactor analytically in systems that are dual to noninteracting spin models. Despite appearing universal when considering the dual XY model, which is our favourite testing ground, we argue that it is not. The prefactor depends indeed on non-universal details that are accidentally irrelevant in that model.

While the importance of the prefactor could be questioned, the logarithm growth of the excess of entropy is, to the best of our understanding, a striking exceptional property of systems with semilocal conservation laws.

2. Melting of the order

In generic systems after global quenches the energy is the only information about the initial state that survives the limit of infinite time; the stationary values of local operators are then captured by effective Gibbs ensembles. Since finite-temperature phase transitions in spin chains described by local Hamiltonians are exceptional if not forbidden [61], the norm is that the stationary expectation values of local observables are smooth functions of the initial state.

The stationary expectation values of local operators are expected to be described by an effective Gibbs ensemble even in the presence of semilocal conservation laws, provided that the initial conditions are generic. Indeed, semilocal charges are relevant only for symmetric initial states; one could then argue that the systems we are considering require a fine-tuning and hence are physically irrelevant.

This unremarkable picture is however a consequence of a naive physical interpretation of asymptotic behaviour. In reality the limit of infinite time is an effective description of times sufficiently larger than the relaxation time [62]. The latter diverges in the limit in which the symmetry in the initial state becomes exact but it is still finite for an exactly symmetric initial state. This happens because the limit of exact symmetry does not commute with the limit of infinite time. Consequently, for any given arbitrarily large time, the discrepancy between the expectation value of local observable and its infinite time limit becomes larger and larger the closer the system is to the symmetric point (see, e.g., Fig. 5 in Section III C). Such a slow relaxation can be understood only taking into account the existence of semilocal charges at the symmetric point.

In Section VC we consider two simple ways of breaking the symmetry in the initial state: a global rotation and an increase of the temperature. In both cases we find a

nontrivial scaling behaviour in the limit where the time is large and comparable with the correlation length of semilocal operators.

The latter, indeed, diverges in the limit of exact symmetry (while being finite at the symmetric point): the limit corresponds to recovering the symmetry without allowing semilocal operators to have a nonzero expectation value, finally leading to the breakdown of cluster decomposition.

Organisation of the rest of the paper:

Section III: It reviews some established results on relaxation after global quenches in quantum spin chains. Section III A presents the general picture in a descriptive way, whereas Section III B goes into more details about the late-time description. Two simple

examples supporting the validity of the picture are exhibited. Section III C points out when the standard description fails, exhibiting also a suspected counterexample in a generic system.

Section IV: It is the core of the paper, where the failure of the standard picture is explained and a theory able to overcome those problems is developed from scratch. Since the issue has deep roots in the mathematical structure underlying the representation of local observables in infinite spin chains, the section is more abstract than the rest of the paper. Section IV A describes a Kramers-Wannier duality transformation introducing ingredients that will be used in the following. Section IV B explains how a state can have distinct representations in different theories of local observables. Section IV C formalises the structure revealed in the previous section and defines the semilocal statistical ensembles able to capture the infinite time limit.

Section V: It describes some effects of persistent semilocal order. Section V A shows that the notion of subsystem changes across different theories of local observables, culminating in a formula that expresses a reduced density matrix in a theory in terms of reduced density matrices of a different theory. Section V B studies the excess of Rényi entropy induced by a local perturbation. The asymptotic behaviour of the excess of entropy is computed analytically in an exactly solvable model, exhibiting an unusual log behaviour that is interpreted as a consequence of semilocal order. Section V C shows that, despite semilocal order being fragile under symmetry-breaking perturbations, it leaves clear marks in the time evolution of the expectation value of local operators.

Section VI: It collects additional comments and a list of open problems.

III. LONG TIME LIMIT: STATE OF THE ART

This section presents some established results on quench dynamics in translationally invariant systems described by Hamiltonians with densities \mathbf{h}_ℓ that are *local*. As reviewed in Section III B, this means that each density \mathbf{h}_ℓ acts non-trivially on a finite connected subsystem A_ℓ of the spin chain, while on the rest of the system \bar{A}_ℓ it acts as an identity.

If the reader is familiar with relaxation in isolated quantum many-body systems, they can skip Section III A. If they are also familiar with the concept of pseudolocality, they can also skip Section III B. We advise however to read Section III C, as it shows when the standard descriptions of relaxation fail.

A. Background

Generally only a tiny part of the excited states of the Hamiltonian gives a significant contribution to the expectation values of local observables after a global quench. Even if tiny with respect to the total Hilbert space, this part is however still exponentially large with respect to the volume of the system (this can be a-posteriori understood by observing that the entanglement entropy of subsystems becomes extensive after global quenches [63]). Consequently, studies of global quenches are challenging both numerically and analytically.

Quantum quenches in translationally invariant many-body systems have been intensively investigated in the last two decades, leading to the following picture:

- In finite systems, consistently with the quantum recurrence theorem [64], there is no relaxation. The distributions of the expectation values over time, however, are highly peaked at values that can be described by the so-called *diagonal ensemble* [32].
- In the thermodynamic limit, if the initial state has clustering properties and the Hamiltonian is local, relaxation is the norm, and the expectation values of local observables can be described by *Gibbs* or *generalised Gibbs ensembles* (GGE) [25].
- Generally, the diagonal ensemble and the (generalised) Gibbs ensemble are locally equivalent [28]. They belong to a family of stationary states with the same local properties, the (generalised) Gibbs ensemble being the state keeping the least amount of information about the initial state. This equivalence class is sometimes called *macrostate* [25].
- Eigenstate thermalisation hypothesis [32–35] and its generalisation to integrable systems [65] point at the fact that the aforementioned family of stationary states includes excited states, usually referred to as *representative states* [66].

The diagonal ensemble can be obtained by killing the non-stationary part of the time-evolving density matrix in a basis diagonalising the Hamiltonian (i.e., only the (block)-diagonal part of the density matrix remains). In practice this can only be done in sufficiently small systems. (Generalised) Gibbs ensembles and representative states are more suitable for analytical investigations. In integrable systems, in particular, they can be directly used to compute expectation values. Generalised Gibbs ensembles rely on the identification of the smallest set of conserved operators characterising the stationary properties of local observables. Representative states instead require the knowledge of the overlaps between the initial state and a generic excited state.

Contrary to classical systems, every quantum system has a number of conservation laws in involution equal to its dimension (namely, the projectors on the eigenstates). In the thermodynamic limit, however, relaxation

is a property of a restricted class of observables, therefore that exponentially large (in volume) set of conservation laws is redundant, like in turn is redundant the description in terms of the diagonal ensemble.

The first studies on local relaxation after global quenches focused on non-interacting systems of fermions and bosons [36]. In those cases the Wick's theorem is sufficient to conclude that the mode occupation numbers provide a sufficiently large set of conserved operators. Having in mind interacting models, the attention had then moved to reinterpreting the known results in a way that could be easily generalised in the presence of interactions.

The importance of locality of the conservation laws [67] was then pointed out. It sets a clear (in translationally invariant systems) division between integrable and generic systems. Such a locality principle was strongly questioned after some discrepancies between theory and numerics in the Heisenberg XXZ spin-1/2 chain were observed. Using the so-called “quench action” method [27], based on representative states, it was shown that the generalised Gibbs ensemble constructed with local charges in the XXZ spin-1/2 chain was inadequate [68, 69]. These problems have been finally resolved with the discovery of new families of conservation laws [51, 70–72] which had been previously overlooked. Their peculiarity is that their densities are not strictly local but exhibit exponential tails. Locality, now including also the new kind of operators, called pseudolocal [30], had resurrected. Some technical discussions apart [73–75], the new picture was positively received, and after the axiomatic definition of generalised Gibbs ensemble put forward in Ref. [52] the question of the relevance of conservation laws has been practically archived.

B. Maximum-entropy descriptions

Conserved quantities constrain the dynamics of the system and prevent the loss of information about the initial state. An important question of the past decade has been, which of these quantities enter in the ensemble ρ_∞ that describes local relaxation according to

$$\lim_{t \rightarrow \infty} \langle \Psi(t) | \mathbf{O} | \Psi(t) \rangle = \text{tr}(\rho_\infty \mathbf{O}), \quad \forall \mathbf{O} \text{ local.} \quad (3.1)$$

We remind the reader again that by “local” we mean that \mathbf{O} acts non-trivially only on a finite connected subsystem A , while on its complement \bar{A} it acts as an identity. We focus here on representations of ρ_∞ through maximum-entropy ensembles [59]. In our specific case they are called *Gibbs* or *generalised Gibbs*, depending on whether the system is generic or integrable.

The first assumption behind such a description is that, at late times, the entanglement entropy

$$S[\rho] = -\text{tr}[\rho \log \rho] \quad (3.2)$$

of subsystems becomes an extensive quantity. One can then look for a representation of ρ_∞ in terms of a state

with an extensive entropy. Thermal ensembles are the typical examples of such states. Imagine then to use a thermal ensemble as a testing description of the stationary expectation values (cf. Ref. [76]). Specifically, we define it in such a way as to capture the energy of the state. We wonder whether the presence of an additional conservation law $\mathbf{K} = \sum_j \mathbf{k}_j$ could change the expectation value of local operators. Guided by statistical physics, we use the Ansatz

$$\rho_\infty = \frac{e^{\mathbf{Q}}}{\text{tr}(e^{\mathbf{Q}})} \quad (3.3)$$

where \mathbf{Q} was originally assumed to be proportional to \mathbf{H} and now is questioned whether to include or not an additional linear dependence on \mathbf{K} . In the absence of \mathbf{K} , $\mathbf{Q} = -\beta \mathbf{H}$ maximises the entropy under the constraint given by the energy per site $\langle \mathbf{h}_j \rangle$ in the initial state. In order to be relevant, the introduction of \mathbf{K} should result in

- (i) a change in the expectation value of local operators;
- (ii) an extensive reduction of the entropy.

While (i) is self-explanatory, (ii) is subtler and can be seen as a condition allowing us to use the principle of maximum entropy [59].

To be more quantitative, the effect of including \mathbf{K} in the ensemble can be interpreted as the result of a smooth variation $\mathbf{Q} = -\beta \mathbf{H} \mapsto \mathbf{Q}(\lambda) = -\beta(\lambda) \mathbf{H} + \lambda \mathbf{K}$ that brings the original ensemble into the new one. The variation of the expectation value of a local operator \mathbf{O} under a small change of λ is

$$\delta \langle \mathbf{O} \rangle_\lambda = \langle \delta \mathbf{Q}(\lambda), \mathbf{O} \rangle_\lambda, \quad (3.4)$$

where $\delta \mathbf{Q}(\lambda) = \mathbf{Q}'(\lambda) \delta \lambda$, and the right-hand side is the Kubo-Mori inner product. In this specific case ($[\delta \mathbf{Q}, \mathbf{Q}] = 0$) the latter is reduced to the connected correlation function $\langle \mathbf{A}, \mathbf{B} \rangle = \langle \mathbf{A} \mathbf{B} \rangle - \langle \mathbf{A} \rangle \langle \mathbf{B} \rangle$; the subscript λ indicates that the expectation values should be taken in the state $\rho_\infty(\lambda)$.

Since the energy density is fixed by the initial state, the variation should preserve it, hence

$$\langle \delta \mathbf{Q}(\lambda), \mathbf{h}_j \rangle_\lambda = \delta \lambda \langle -\beta'(\lambda) \mathbf{H} + \mathbf{K}, \mathbf{h}_j \rangle_\lambda = 0. \quad (3.5)$$

Condition (i) then requires the existence of a local operator \mathbf{O} satisfying [77]

$$0 < \langle \mathbf{K}, \mathbf{O} \rangle_\lambda < \infty. \quad (3.6)$$

On the other hand, the variation of the entropy per unit site is $\delta \mathfrak{s}[\rho_\infty(\lambda)] = -\langle \delta \mathbf{Q}(\lambda), \mathbf{q}_j(\lambda) \rangle_\lambda$, where $\mathbf{q}_j(\lambda)$ is the density of $\mathbf{Q}(\lambda)$ [78]. Imposing the energy constraint (3.5) we then obtain

$$\delta \mathfrak{s}[\rho_\infty(\lambda)] = -\frac{1}{2} \delta(\lambda^2) \langle \mathbf{Q}'(\lambda), \mathbf{q}'_j(\lambda) \rangle_\lambda, \quad (3.7)$$

which is non-positive along a path with increasing $|\lambda|$. Explicitly it reads $\delta\mathfrak{s}[\rho_\infty(\lambda)] = -\frac{1}{2}\delta(\lambda^2)[\langle\mathbf{K}, \mathbf{k}_j\rangle_\lambda - (\beta'(\lambda))^2 \langle\mathbf{H}, \mathbf{h}_j\rangle_\lambda]$, where \mathbf{k}_j is the density of \mathbf{K} . In order to apply the maximum entropy principle, we need the entropy to be a smooth function of the Lagrange multiplier λ , which requires [79]

$$0 < \langle\mathbf{K}, \mathbf{k}_j\rangle_\lambda < \infty. \quad (3.8)$$

This is usually referred to as ‘‘extensivity condition’’ and represents (ii). The generalisation to systems with more conservation laws is straightforward and results in the same equations.

As a matter of fact, in the most formal works dealing with the emergence of maximum-entropy ensembles after global quenches (see, e.g., Ref. [52]), an additional assumption is made:

- (*) the density \mathbf{k}_j can be obtained as a limit of a sequence of local operators.

From our simplified explanation, it might not be evident where this extra condition comes from. Explaining it goes beyond the mathematical rigour of this work; we only mention that (*) has a role in making sense to the thermodynamic limit. We will see however that, in order to describe the stationary values in symmetric systems with maximum-entropy ensembles, we will be forced to relax (*).

Conditions (3.6), (3.8), and (*) define what is now known as *pseudolocality*. Charges that satisfy them include translationally invariant sums $\mathbf{Q} = \sum_\ell \mathbf{q}_\ell$, where the density has the form $\mathbf{q}_\ell = \sum_{r=1}^\infty \mathbf{q}_\ell(r)$ with $\{\mathbf{q}_\ell(r)\}$ being a sequence of local operators that act on subsystems of increasing size r and are exponentially suppressed in r , e.g., $\langle\mathbf{Q}, \mathbf{q}_\ell(r)\rangle \leq C e^{-r/\xi}$, for some $C, \xi > 0$ [30, 51].

In the following we consider two examples of maximum-entropy descriptions in integrable exactly solvable models. Note that, since the systems are integrable, the corresponding maximum-entropy ensembles will have to incorporate the constraints of infinitely many pseudolocal integrals of motion.

1. Example: XY model

To illustrate the validity of the generalised Gibbs ensemble description we consider quench dynamics in the XY model, whose time evolution is generated by the Hamiltonian

$$\mathbf{H} = \sum_{\ell \in \mathbb{Z}} J_x \sigma_\ell^x \sigma_{\ell+1}^x + J_y \sigma_\ell^y \sigma_{\ell+1}^y. \quad (3.9)$$

This is a very well known model [80]. We mention that the ground state is noncritical for $|J_x| \neq |J_y|$. The critical

line separates two ordered phases where spin flip symmetry is broken (for $J_x > J_y$ the spins acquire a nonzero x component, whereas for $J_y > J_x$ they acquire a nonzero y component).

Let the system be prepared in the initial state

$$|\Psi(0)\rangle = |\nearrow_\theta\rangle := \bigotimes_{\ell \in \mathbb{Z}} \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} \end{pmatrix}, \quad (3.10)$$

which is the ground state of the Hamiltonian

$$\mathbf{H}_0 = - \sum_{\ell \in \mathbb{Z}} \cos \left(\frac{\theta}{2}\right) \sigma_\ell^z + \sin \left(\frac{\theta}{2}\right) \sigma_\ell^x. \quad (3.11)$$

For $\theta \notin \{0, \pi\}$ the initial state breaks the \mathbb{Z}_2 symmetry \mathcal{P}^z of the post-quench Hamiltonian \mathbf{H} (see Eq. (2.2)). The latter can be conveniently rewritten in terms of Majorana fermions $\mathbf{a}_\ell^{x,y} = (\prod_{j < \ell} \sigma_j^z) \sigma_\ell^{x,y}$ as [81]

$$\mathbf{H} = \frac{1}{4} \sum_{\ell, n \in \mathbb{Z}} (\mathbf{a}_\ell^x \ \mathbf{a}_\ell^y) \mathcal{H}_{\ell, n} \begin{pmatrix} \mathbf{a}_n^x \\ \mathbf{a}_n^y \end{pmatrix}, \quad (3.12)$$

where $\mathcal{H}_{\ell, n} = \int \frac{dk}{2\pi} e^{i(\ell-n)k} \mathcal{H}(k)$ is the Fourier transform of the 2×2 matrix

$$\mathcal{H}(k) = 2(J_x - J_y) \sin k \sigma^x - 2(J_x + J_y) \cos k \sigma^y. \quad (3.13)$$

On account of the Toeplitz structure of \mathcal{H} , $\mathcal{H}(k)$ is also referred to as the symbol of the Hamiltonian (see, e.g., Ref. [82]). The symbol generates the evolution of the two-point correlations of Majorana fermions according to

$$\Gamma_t(k) = e^{-i\mathcal{H}(k)t} \Gamma(k) e^{i\mathcal{H}(k)t}, \quad (3.14)$$

where $\Gamma(k)$ is the symbol of the two-point correlation matrix

$$\Gamma_{\ell, n} = \delta_{\ell, n} I - \left\langle \begin{pmatrix} \mathbf{a}_\ell^x \\ \mathbf{a}_\ell^y \end{pmatrix} \begin{pmatrix} \mathbf{a}_n^x & \mathbf{a}_n^y \end{pmatrix} \right\rangle, \quad (3.15)$$

in the initial state. For example, in the ferromagnetic initial state (3.10) with $\theta = 0$ one has $\Gamma_{\ell, n} = \delta_{\ell, n} \sigma^y$ and $\Gamma(k) = \sigma^y$.

Under some very mild assumptions on the dispersion relation which are practically always satisfied (for example, the dispersion relation should not be flat), at large time the symbol of the correlation matrix relaxes to its time-averaged value $\bar{\Gamma}(k) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \Gamma_t(k)$. In our specific case we find

$$\bar{\Gamma}(k) = \frac{(\cos^2\theta + 1)\cos^2 k - 2\cos\theta \cos k + \frac{J_x - J_y}{J_x + J_y} \sin^2\theta \sin^2 k}{2\cos\theta \cos^2 k - (\cos^2\theta + 1)\cos k} \begin{pmatrix} 0 & \frac{-i(J_x + J_y)\cos k}{(J_x + J_y)\cos k + i(J_x - J_y)\sin k} \\ \frac{i(J_x + J_y)\cos k}{(J_x + J_y)\cos k - i(J_x - J_y)\sin k} & 0 \end{pmatrix}. \quad (3.16)$$




FIG. 3. Relaxation of $\langle \sigma_\ell^z \rangle$ towards $\langle \sigma_\ell^z \rangle_{\text{GGE}} \approx 0.466$ (0.717) (dashed lines) in the XY model, computed from Eq. (3.16). The dots represent the results of the iTEBD simulation, while the solid curve is the exact time evolution using Eq. (3.14). Parameters in Eqs. (3.9) and (3.10) are $J_x = 1$, $J_y = 2$, and $\theta = 0.9$ (0.3). The iTEBD evolution uses a second order Trotter scheme with two-site quantum gates, time step $\delta t = 0.01$, Schmidt values cutoff 10^{-6} and maximal allowed bond dimension $M_{\text{max}} = 800$.

In a non-interacting model, Ref. [83] established that (3.16) is actually equivalent to a symbol of the two-point correlation matrix in a generalised Gibbs ensemble, i.e., $\bar{\Gamma}(k) \equiv \Gamma_{\text{GGE}}(k)$. Under some assumptions it was later proven [84, 85] that, if the Hamiltonian generating the time evolution is translationally invariant, and if the initial state has clustering properties, the asymptotic state of the system is Gaussian, whence higher-order correlations can be accessed using Wick's theorem. These conditions are satisfied by Hamiltonian (3.9) and our state (3.10). The time-averaged correlation matrix (3.16) thus determines not only the two-point correlations of Majorana fermions at late times (e.g. $\langle \sigma_\ell^z \rangle$, shown in Fig. 3), but also higher-order correlations, as demonstrated in Fig. 4, which shows the relaxation of $\langle \sigma_\ell^x \rangle$ (a string of Majorana fermions).

2. Example: Dual XY model

We now consider a global quench from the same initial state $|\nearrow_\theta\rangle$ in the model described by the Hamiltonian

$$\mathbf{H} = \sum_{\ell \in \mathbb{Z}} \sigma_{\ell-1}^x (J_x \mathbf{I} - J_y \sigma_\ell^z) \sigma_{\ell+1}^x. \quad (3.17)$$




FIG. 4. Relaxation of $\langle \sigma_\ell^x \rangle$ towards $\langle \sigma_\ell^x \rangle_{\text{GGE}} \approx 0$ (dashed lines) in the XY model. Parameters are the same as in Fig. 3.

To the best of our knowledge, this model has not been studied much. We mention that the ground state is non-critical for $|J_x| \neq |J_y|$. As shown in Appendix C, for $|J_y| < |J_x|$ the ground state is in a Landau phase where both spin flip symmetries over odd and even sites are broken, for $|J_y| > |J_x|$, instead, it is in a nontrivial $\mathbb{Z}_2 \times \mathbb{Z}_2$ protected topological phase (see, e.g., Ref. [86] for $J_x = 0$).

Hamiltonian (3.17) can be mapped into the one of the quantum XY model, given in Eq. (3.9), by means of a Kramers-Wannier duality transformation, which will be described in detail in Section IV A. After the duality transformation the Hamiltonian (3.17) is therefore quadratic in terms of Majorana fermions: the time evolution of the n -point correlation functions is determined only by the n -point correlation functions at the initial time. As reviewed in the previous section, only the 2-point functions survive the limit of infinite time (the asymptotic state is Gaussian). The correlation matrix of the generalised Gibbs ensemble can thus easily be determined from the initial correlation matrix (which consists of the initial expectation values of all local operators mapped by the duality transformation into operators that are quadratic in Majorana fermions). This allows one to obtain the stationary values of local operators with minimal effort. Since however we are considering this model as a representative of a larger class of systems for which this procedure would not apply, we describe also an alternative approach.

Specifically, one could infer that this is an integrable model from the fact that a boost operator \mathbf{B} exists and




FIG. 5. Numerical (iTEBD) time evolution of $\langle \sigma_\ell^z \rangle$ and GGE predictions ($\langle \sigma_\ell^z \rangle_{\text{GGE}} \approx 0.038, 0.0012, \text{ and } 0.000078$ for $\theta = 0.9, 0.3, \text{ and } 0.15$, respectively) in the dual XY model. Relaxation towards the GGE prediction is slower for smaller θ . Parameters of the Hamiltonian (3.17) are $J_x = 1, J_y = 2$. The iTEBD uses second order Trotter scheme with four-site quantum gates, $\delta t = 0.01$, Schmidt values cutoff 10^{-6} , and maximum bond dimension $M_{\text{max}} = 1000$. The data converges up to times $t \sim 4.5$ (see Appendix A).

takes the standard form

$$\mathbf{B} = \sum_{\ell \in \mathbb{Z}} \ell \mathbf{h}_\ell, \quad (3.18)$$

where $\mathbf{h}_\ell = \sigma_{\ell-1}^x (J_x \mathbf{I} - J_y \sigma_\ell^z) \sigma_{\ell+1}^x$. That is to say, we can construct a tower of conserved operators \mathbf{Q}_n in the following manner:

$$\mathbf{Q}_{n+1} = i[\mathbf{B}, \mathbf{Q}_n], \quad \mathbf{Q}_1 = \mathbf{H}. \quad (3.19)$$

We are then in a position to either guess the corresponding Lax operator (which, in this model, was obtained in Ref. [87]) and exploit the integrable structure to compute the GGE expectation values [88], or to construct a truncated generalised Gibbs ensemble [89] using the most local charges. For generic θ all these methods converge to the same values.

Relaxation of local observables in the dual XY model is shown in Figs. 5 and 6. For $\theta \notin \{0, \pi\}$ all local observables eventually relax to their respective GGE predictions. Note, however, that for some local observables the time scale on which relaxation happens tends to increase when θ decreases — see Fig. 5.

C. Failure of Gibbs and generalised Gibbs ensemble in symmetric systems

The Hamiltonians of the XY and the dual XY model have a \mathbb{Z}_2 symmetry: $\mathcal{P}^z[\mathbf{H}] = 0$. If also the initial state is even under \mathcal{P}^z (e.g., $|\mathcal{Z}_\theta\rangle\langle\mathcal{Z}_\theta|$ for $\theta \in \{0, \pi\}$), the entire system is symmetric. Hence, only the even conservation laws are expected to contribute to the stationary behaviour of the local observables.




FIG. 6. Relaxation of $\sigma_{\ell-1}^x \sigma_{\ell+1}^x$ towards the GGE prediction ($\langle \sigma_{\ell-1}^x \sigma_{\ell+1}^x \rangle_{\text{GGE}} \approx 0.022808$ (0.005646) for $\theta = 0.3$ (0.15), respectively) in the dual XY model. Parameters are the same as in Fig. 5, except for the maximal bond dimension: here, $M_{\text{max}} = 1000$ ($\theta = 0.3$), $M_{\text{max}} = 600$ ($\theta = 0.15$).




FIG. 7. Relaxation of $\langle \sigma_\ell^z \rangle$ and $\langle \sigma_\ell^z \sigma_{\ell+1}^z \rangle$ towards the respective GGE predictions (dashed lines) in the XY model. Parameters are $J_x = 1, J_y = 2$, and $\theta = 0$. The colored solid lines correspond to exact calculation using Eq. (3.14), while the dots represent the iTEBD results. The inset shows the $1/\sqrt{t}$ decay of oscillations (the differences between the local maxima of oscillations and the GGE predictions are plotted versus $1/\sqrt{t}$). The iTEBD uses second order Trotter scheme with a two-site update rule, $\delta t = 0.01$, Schmidt values cutoff 10^{-5} , and maximal bond dimension $M_{\text{max}} = 400$.

Using the even-parity local charges in each model reproduces the generalised Gibbs ensembles described in the previous examples. As demonstrated in Fig. 7, the GGE description is correct in the XY model for $\theta = 0$. In contrast, in the dual XY model the prediction for the expectation value of σ_ℓ^z is zero, while its time evolution approaches a finite value $\langle \sigma_\ell^z \rangle_\infty \approx 0.5625$ — see




FIG. 8. Numerical evolution of σ_ℓ^z (points) and the GGE prediction $\langle \sigma_\ell^z \rangle_{\text{GGE}} = 0$ (blue line) in the dual XY model, for $\theta = 0$, $J_x = 1$ and $J_y = 2$. The time evolution relaxes towards the value $\langle \sigma_\ell^z \rangle \approx 0.5625$. The iTEBD parameters are the same as in Fig. 5.

Fig. 8 [90]. The generalised Gibbs ensemble using the even local charges of the dual XY model thus *fails* to capture the asymptotic behaviour of σ_ℓ^z .

Normally this would be indicative of having overlooked some pseudolocal integrals of motion. Here, however, the discrepancy has a different origin (which will be explained in Section IV) and the failure of the generalised Gibbs ensemble acquires a more fundamental meaning.

Finally, as a concrete example of a generic system, we consider Hamiltonian (2.4) with

$$\mathbf{W}_\ell[\{\sigma^z\}] = w_1 \sigma_\ell^z + \sum_{n=1}^{\infty} w_{2,n} \sigma_\ell^z \sigma_{\ell+n}^z \quad (3.20)$$

and initial state

$$|\Psi(0)\rangle = e^{i\frac{\varphi}{2} \sum_\ell \sigma_\ell^x \sigma_{\ell+1}^x} |\uparrow\rangle, \quad (3.21)$$

where $|\uparrow\rangle$ means that all spins are up. To the best of our knowledge, there is no (pseudo)local operator commuting with this Hamiltonian. The expectation value of the energy density reads $\langle \mathbf{h}_\ell \rangle_t = 1 + (w_1 + w_{2,1} - 1) \cos^2 \varphi + (\sum_{n=2}^{\infty} w_{2,n}) \cos^4 \varphi$. We now provide numerical evidence that the state does not thermalise by comparing the time evolution of $\langle \sigma_\ell^z \rangle$ starting from two initial states with the same energy. To that aim, we choose the Hamiltonian's parameters in such a way that $0 < (1 - w_1 - w_{2,1}) / (\sum_{n=2}^{\infty} w_{2,n}) < 1$, so that the energy is the same for initial states with either $\cos^2 \varphi = 0$ or $\cos^2 \varphi = (1 - w_1 - w_{2,1}) / (\sum_{n=2}^{\infty} w_{2,n})$. Fig. 9 shows quite clearly that the infinite-time limit of $\langle \sigma_\ell^z \rangle$ depends on the initial state (at the same energy), in contrast with the eigenstate thermalisation hypothesis and the conjecture of local thermalisation.




FIG. 9. Numerical evolution of σ_ℓ^z in the generic model given by Eqs. (2.4) and (3.20). Parameters are $J_x = J_y = w_{2,2} = 1$, and $w_1 = 0.6$. The iTEBD parameters are the same as in Fig. 5, except for the maximum bond dimension M_{max} (here various bond dimensions are used, yielding comparable results).

IV. BEYOND (GENERALISED) GIBBS ENSEMBLES

We aim at obtaining a canonical description of the macrostate describing the infinite-time limit after symmetric quenches in systems such as the dual XY model in which the standard (generalised) Gibbs ensemble fails — see Fig. 8. Since the limit of infinite time makes sense only in the thermodynamic limit, we make an effort to consider infinite chains directly.

Bearing in mind the duality correspondence highlighted in Refs. [55, 56] between symmetry-protected topological phases and Landau phases, we start this section describing a duality transformation. The most important property that we want to stress since the beginning is that duality transformations can be used to map algebras of operators representing local observables into one another. This will prove essential in unveiling the importance of semilocal integrals of motion on the relaxation of local observables.

A. Kramers-Wannier duality

Let us denote by \mathfrak{A}_{ql} the (C^*) algebra of quasilocal operators in an infinite spin-1/2 chain. This algebra includes local operators and limits of their sequences that converge in the operator norm. It is generated by the local spin operators, which act like Pauli matrices on site j and like the identity elsewhere; we will denote them either by σ_j^α or by τ_j^α .

We consider a duality transformation $\mathcal{D}_{\mathbb{Z}_2}$ that differs from the standard Kramers-Wannier duality, responsible for the self-duality of the transverse-field Ising model,

just in an additional rotation:

$$\boldsymbol{\tau}_j^x = \boldsymbol{\sigma}_{j-1}^x \boldsymbol{\sigma}_j^x, \quad \boldsymbol{\tau}_j^z \boldsymbol{\tau}_{j+1}^z = \boldsymbol{\sigma}_j^z. \quad (4.1)$$

In order to define it properly in the infinite chain, we exploit the same trick as Ref. [91], which needed to define a Jordan-Wigner transformation directly in infinite systems.

In particular, we define $\mathbf{T}_{\sigma,s}^\alpha$, where $\alpha \in \{x, z\}$, $s = \pm$, and $\sigma \leftrightarrow \tau$, as the Hermitian operators satisfying

$$[\mathbf{T}_{\sigma,s}^\alpha]^2 = \mathbf{I}, \quad \mathbf{T}_{\sigma,s}^\alpha \mathbf{O} \mathbf{T}_{\sigma,s}^\alpha = \lim_{n \rightarrow \infty} \left[\prod_{\ell=0}^n \boldsymbol{\sigma}_{s\ell}^\alpha \right] \mathbf{O} \left[\prod_{\ell=0}^n \boldsymbol{\sigma}_{s\ell}^\alpha \right] \quad (4.2)$$

for all quasilocal operators $\mathbf{O} \in \mathfrak{A}_{\text{ql}}$, extended then by linearity as $\mathbf{T}_{\sigma,s}^\alpha (\mathbf{O}_1 + \mathbf{O}_2) \mathbf{T}_{\sigma,s}^\alpha = \mathbf{T}_{\sigma,s}^\alpha \mathbf{O}_1 \mathbf{T}_{\sigma,s}^\alpha + (\mathbf{T}_{\sigma,s}^\alpha \mathbf{O}_2 \mathbf{T}_{\sigma,s}^\alpha) \mathbf{T}_{\sigma,s}^\alpha$ for $\mathbf{O}_1, \mathbf{O}_2 \in \mathfrak{A}_{\text{ql}}$ (the role of such an extension will be discussed in Section IV C). We also define the auxiliary operators

$$\mathbf{\Pi}_{\sigma,+}^z(j) = \mathbf{T}_{\sigma,+}^z \cdot \begin{cases} \boldsymbol{\sigma}_j^z \boldsymbol{\sigma}_{j+1}^z \cdots \boldsymbol{\sigma}_{-1}^z & j < 0 \\ \mathbf{I} & j = 0 \\ \boldsymbol{\sigma}_0^z \boldsymbol{\sigma}_1^z \cdots \boldsymbol{\sigma}_{j-1}^z & j > 0. \end{cases} \quad (4.3)$$

and

$$\mathbf{\Pi}_{\tau,-}^x(j) = \mathbf{T}_{\tau,-}^x \cdot \begin{cases} \boldsymbol{\tau}_{j+1}^x \boldsymbol{\tau}_{j+2}^x \cdots \boldsymbol{\tau}_0^x & j < 0 \\ \mathbf{I} & j = 0 \\ \boldsymbol{\tau}_1^x \boldsymbol{\tau}_2^x \cdots \boldsymbol{\tau}_j^x & j > 0, \end{cases} \quad (4.4)$$

These operators act like a spin flip on the right or left of site j included and like the identity elsewhere. The rotated Kramers-Wannier duality is then an algebra homomorphism mapping the algebra of operators generated by $\{\boldsymbol{\tau}, \mathbf{T}_{\tau,-}^x\}$ to the one generated by $\{\boldsymbol{\sigma}, \mathbf{T}_{\sigma,+}^z\}$, both algebras being extensions of \mathfrak{A}_{ql} . It reads

$$\begin{aligned} \mathbf{\Pi}_{\tau,-}^x(j) &= \boldsymbol{\sigma}_j^x, \\ \boldsymbol{\tau}_j^y &= \boldsymbol{\sigma}_{j-1}^x \boldsymbol{\sigma}_j^y \mathbf{\Pi}_{\sigma,+}^z(j+1), \\ \boldsymbol{\tau}_j^z &= \mathbf{\Pi}_{\sigma,+}^z(j). \end{aligned} \quad (4.5)$$

We stress that, while $\mathbf{T}_{\sigma,\pm}^z$ resembles a string of $\boldsymbol{\sigma}_j^z$ with j extending up to $\pm\infty$, the sequence $\{\prod_{\ell=0}^n \boldsymbol{\sigma}_{\pm\ell}^z\}$ does not converge in the operator norm as $n \rightarrow \infty$. This is why we could not avoid extending the algebra of the quasilocal operators.

The duality transformation (4.5) mixes the local space with some inaccessible degrees of freedom and only a subset of the operators remain local under the mapping — see Table I. In the table we also consider parities under spin flips. Specifically, even (odd) quasilocal operators obey $\mathcal{P}_\nu^\alpha[\mathbf{O}] = \mathbf{O}$ ($\mathcal{P}_\nu^\alpha[\mathbf{O}] = -\mathbf{O}$), where (α, ν) corresponds to either (x, τ) or (z, σ) — cf. Eq. (2.2)

$$\mathcal{P}_\nu^\alpha[\mathbf{O}] = \lim_{n \rightarrow \infty} \mathbf{\Pi}_{\nu,s\nu}^\alpha(-n) \mathbf{\Pi}_{\nu,s\nu}^\alpha(n) \mathbf{O} \mathbf{\Pi}_{\nu,s\nu}^\alpha(-n) \mathbf{\Pi}_{\nu,s\nu}^\alpha(n), \quad (4.6)$$

\mathcal{D}	LE_z	LO_z	$S_z E_z$	$S_z O_z$
LE_x	•			
LO_x			•	
$S_x E_x$		•		
$S_x O_x$				•

TABLE I. The duality transformation \mathcal{D} . L stands for “local”, S_α for “semilocal” with a string along the axis α , E_α and O_α for “even” and “odd”, respectively, with respect to spin flip along the axis α . The transformation preserves only even locality and odd semilocality.

and $s_{\sigma(\tau)} = +(-)$. The transformations \mathcal{P}_ν^α can be extended by linearity to operators $\mathbf{O}_1 + \mathbf{O}_2 \mathbf{T}_{\nu,s\nu}^\alpha$.

Finally, we report the inverse transformation $\mathcal{D}_{\mathbb{Z}_2}^{-1}$, which maps the algebra of observables generated by $\{\boldsymbol{\sigma}, \mathbf{T}_{\sigma,+}^z\}$ to the one generated by $\{\boldsymbol{\tau}, \mathbf{T}_{\tau,-}^x\}$:

$$\begin{aligned} \boldsymbol{\sigma}_j^x &= \mathbf{\Pi}_{\tau,-}^x(j), \\ \boldsymbol{\sigma}_j^y &= \mathbf{\Pi}_{\tau,-}^x(j-1) \boldsymbol{\tau}_j^y \boldsymbol{\tau}_{j+1}^z, \\ \mathbf{\Pi}_{\sigma,+}^z(j) &= \boldsymbol{\tau}_j^z. \end{aligned} \quad (4.7)$$

B. Semilocality and hidden symmetry breaking

When presenting the duality transformation $\mathcal{D}_{\mathbb{Z}_2}$ we were forced to extend the algebra of operators. Not all operators of the extended algebra can however be associated with local observables. A theory of local observables, indeed, requires some basic notion of locality:

- (a) the theory is generated by operators that, in the limit of infinite distance, commute with one another

$$\lim_{|x-y| \rightarrow \infty} [\mathbf{O}_1(x), \mathbf{O}_2(y)] = 0; \quad (4.8)$$

- (b) the state has clustering properties for the aforementioned operators

$$\langle \mathbf{O}_1(x) \mathbf{O}_2(y) \rangle \xrightarrow{|x-y| \rightarrow \infty} \langle \mathbf{O}_1(x) \rangle \langle \mathbf{O}_2(y) \rangle. \quad (4.9)$$

After an inspection of Eq. (4.5), we realise that condition (a) forces us to exclude either operators with strings or operators that are odd under spin flip (that is, under \mathcal{P}_τ^x or \mathcal{P}_σ^z). In either of the cases the remaining operators can represent local observables. This can be understood by noting that a state can not simultaneously have clustering properties, i.e., condition (b), and a nonzero expectation value of two operators that anticommute at an infinite distance:

$$\begin{aligned} \langle \mathbf{O}_1(x) \rangle \langle \mathbf{O}_2(y) \rangle &\xleftarrow{|x-y| \rightarrow \infty} \langle \mathbf{O}_1(x) \mathbf{O}_2(y) \rangle = \\ &= -\langle \mathbf{O}_2(y) \mathbf{O}_1(x) \rangle \xrightarrow{|x-y| \rightarrow \infty} -\langle \mathbf{O}_1(x) \rangle \langle \mathbf{O}_2(y) \rangle. \end{aligned} \quad (4.10)$$

Duality transformations such as $\mathcal{D}_{\mathbb{Z}_2}$ help identify theories in which local observables are not necessarily represented by local operators. When that happens we call the corresponding operators “semilocal”. In the following we will refer to the theory consisting of only quasilocal operators as “quasilocal theory”, in contrast to “semilocal theories”, which include also semilocal operators.

Specifically, in the \mathbb{Z}_2 case we identify two theories of local observables:

- Quasilocal Theory: It consists of quasilocal operators;
- Even \mathbb{Z}_2 -Semilocal Theory: It consists of even quasilocal operators and operators with even quasilocal “heads” and half-infinite “tails” (see (4.27) for an example).

Semilocal operators represent information that was lost in the thermodynamic limit, e.g., anything related to the boundaries of the system. This becomes evident when considering their expectation values. For example, let us take the equilibrium state

$$\rho^{(\sigma)}(\beta) = \frac{e^{-\beta \mathbf{H}^{(\sigma)}}}{Z_\beta}, \quad (4.11)$$

where $\mathbf{H}^{(\sigma)} = \sum_\ell \sigma_\ell^z$, the upper index referring to the representation of spins through σ_ℓ^α . Being even, i.e., $\mathcal{P}_\sigma^z[\rho^{(\sigma)}(\beta)] = \rho^{(\sigma)}(\beta)$, this state can be described both by the quasilocal theory and by the even \mathbb{Z}_2 -semilocal one, the latter being generated by the even local operators and the semilocal operator $\mathbf{T}_{\sigma,+}^z$. Indeed, the Hamiltonian $\mathbf{H}^{(\sigma)}$ through which the state is defined belongs to both theories — see Fig. 2. Since it is gapped with a nondegenerate ground state, one would be tempted to say that the limits $\beta \rightarrow \pm\infty$ of Eq. (4.11) are given by the states $|\uparrow\rangle$ and $|\downarrow\rangle$, respectively; one would also conclude that they are one-site shift invariant. Such a conclusion is correct in the quasilocal theory, in which we have

$$\lim_{\beta \rightarrow \infty} \text{tr}[\rho(\beta)\mathbf{O}] = \langle \uparrow | \mathbf{O} | \uparrow \rangle_\sigma, \quad \forall \mathbf{O} \in \mathfrak{A}_{\text{ql}}^{(\sigma)}. \quad (4.12)$$

We added the subscript σ in $|\uparrow\rangle_\sigma$ to stress that the state can be represented by all spins up when considering operators in $\mathfrak{A}_{\text{ql}}^{(\sigma)}$.

Instead, in the even \mathbb{Z}_2 -semilocal theory semilocal operators, such as $\mathbf{\Pi}_{\sigma,+}^z(\ell)$, do not satisfy Eq. (4.12): they are affected by what is not in the bulk of the system. We denote this uncertainty by $|\uparrow \bullet\rangle_{\sigma,\text{sl}}$ and $|\downarrow \bullet\rangle_{\sigma,\text{sl}}$, where \bullet represents our ignorance of what is not in the bulk. Since we are already in the thermodynamic limit, this independent degree of freedom risks to appear very abstract. In order to partially overcome this problem, it is convenient to map semilocal operators into local ones through a duality transformation. In our specific case this is achieved by $\mathcal{D}_{\mathbb{Z}_2}^{-1}$. As in Section IV A, we denote the transformed spins by τ_ℓ^α ; the reader can think of the τ representation

as of a shortcut for operators that could be semilocal in the σ representation (the system has not changed). We then have — see Eq. (4.7) —

$$\rho^{(\tau)}(\beta) = \frac{e^{-\beta \mathbf{H}^{(\tau)}}}{Z_\beta}, \quad (4.13)$$

with $\mathbf{H}^{(\tau)} = \sum_\ell \tau_\ell^z \tau_{\ell+1}^z$, which is even under \mathcal{P}_τ^x . The limits $\beta \rightarrow \pm\infty$ now exhibit a completely different phenomenology that uncovers the role of \bullet . Specifically, cluster decomposition requires spin-flip symmetry \mathcal{P}_τ^x to be spontaneously broken [57, 92]:

$$\lim_{\beta \rightarrow \infty} \text{tr}[\rho(\beta)\mathbf{O}] = \begin{cases} \langle \uparrow | \mathbf{O} | \uparrow \rangle_\tau \\ \langle \downarrow | \mathbf{O} | \downarrow \rangle_\tau \end{cases} \quad \forall \mathbf{O} \in \mathfrak{A}_{\text{ql}}^{(\tau)}. \quad (4.14)$$

(Note that, in the limit $\beta \rightarrow -\infty$, the state breaks also one-site shift invariance and becomes Néel or anti-Néel.) Semilocal operators, which have vanishing expectation value at any finite temperature (they are odd under \mathcal{P}_τ^x while the state is even), acquire a nonzero expectation value at zero temperature due to spontaneous hidden symmetry breaking.

We are now in a position to quantify \bullet . We show, in particular, that the ambiguity hidden behind \bullet is just a global sign. To that aim, let us consider a triplet of semilocal operators $\mathbf{O}^{(n_j)}(x_j)$ ($j \in \{1, 2, 3\}$) representing local observables at arbitrarily large distances $|x_2 - x_1|$, $|x_3 - x_2|$ and $|x_3 - x_1|$. Using clustering we have

$$\langle \mathbf{O}^{(n_2)}(x_2) \rangle = s^{(n_2)}(x_2) \lim_{|x_j - x_k| \rightarrow \infty} \sqrt{\frac{\langle \mathbf{O}^{(n_2)}(x_2) \mathbf{O}^{(n_3)}(x_3) \rangle \langle \mathbf{O}^{(n_1)}(x_1) \mathbf{O}^{(n_2)}(x_2) \rangle}{\langle \mathbf{O}^{(n_1)}(x_1) \mathbf{O}^{(n_3)}(x_3) \rangle}}, \quad (4.15)$$

which tells us that we can determine the expectation value of $\mathbf{O}^{(n_2)}(x_2)$ up to a sign $s^{(n_2)}(x_2)$ from the expectation values of local operators ($\mathbf{O}^{(n_j)}(x_j)\mathbf{O}^{(n_k)}(x_k)$ are indeed local).

Let us now consider whatever semilocal operator $\mathbf{O}^{(n)}(x)$ with $|x - x_2| \rightarrow \infty$. Using again clustering we immediately obtain $s^{(n)}(x) = s^{(n_2)}(x_2) \text{sgn}(\langle \mathbf{O}^{(n)}(x) \mathbf{O}^{(n_2)}(x_2) \rangle)$, and hence a single global sign $s^{(n_2)}(x_2)$ fixes the expectation value of every semilocal operator. We can choose, for example, $s^{(n_2)}(x_2)$ to be the sign of the expectation value of the fundamental semilocal operator $\mathbf{T}_{\sigma,+}^z$. We can then indicate the state in the \mathbb{Z}_2 -semilocal theory by

$$|\uparrow; \text{sgn}[\langle \mathbf{T}_{\sigma,+}^z \rangle]\rangle_{\sigma,\text{sl}} \quad \text{or} \quad |\downarrow; \text{sgn}[\langle \mathbf{T}_{\sigma,+}^z \rangle]\rangle_{\sigma,\text{sl}}. \quad (4.16)$$

Note that the symmetry breaking states $|\uparrow, \pm\rangle_{\sigma,\text{sl}}$ and $|\downarrow, \pm\rangle_{\sigma,\text{sl}}$ in the semilocal theory are stable under symmetry-breaking perturbations, which can now also be semilocal. This is the hidden symmetry breaking studied in Refs. [55, 93, 94]. It is hidden because no local operator can distinguish between $|\uparrow, +\rangle_{\sigma,\text{sl}}$ and $|\uparrow, -\rangle_{\sigma,\text{sl}}$ (or $|\downarrow, +\rangle_{\sigma,\text{sl}}$ and $|\downarrow, -\rangle_{\sigma,\text{sl}}$).

More generally, a pure state that can be written as

$$\begin{aligned} |\Psi_0\rangle^{(\sigma)} &= \mathbf{V}(1) |\uparrow; s\rangle_{\sigma, \text{sl}} \quad \text{or} \quad |\Psi_0\rangle^{(\sigma)} = \mathbf{V}(1) |\downarrow; s\rangle_{\sigma, \text{sl}}, \\ i\partial_\tau \mathbf{V}(\tau) &= \mathbf{W}(\tau) \mathbf{V}(\tau), \quad \mathbf{V}(0) = \mathbf{I}, \end{aligned} \quad (4.17)$$

where s is the sign ambiguity and $\mathbf{W}(\tau)$ an extensive translationally invariant Hermitian operator with a \mathcal{P}_σ^z -invariant local density [95] can be described by the even \mathbb{Z}_2 -semilocal theory.

C. Semilocal (generalised) Gibbs ensembles

In equilibrium at zero temperature the interplay between the symmetries of the (ground) state and of the Hamiltonian is sufficient to discriminate the symmetry-protected topological phases [1]. After a global quench that is not sufficient anymore: conserved operators constrain the dynamics as well as the Hamiltonian, therefore it is reasonable to expect that also their group of symmetry becomes important.

We have already reviewed that not every conservation law affects the late-time behaviour of local operators. In generic systems it was shown that the discriminating criterion is pseudolocality, which, in view of condition (*) in Section III B, is related to the fact that the operators we are interested in form the algebra of quasilocal operators \mathfrak{A}_{ql} . In the presence of a symmetry of the system (i.e., initial state and Hamiltonian), such as the \mathbb{Z}_2 symmetry taken as an example in this paper, only a subset of operators is relevant: by symmetry the rest of them have zero expectation values. In the \mathbb{Z}_2 case, the relevant operators form the subalgebra $\mathfrak{A}_{\text{ql}}^+ \subset \mathfrak{A}_{\text{ql}}$ of the quasilocal operators that are even under spin flip. We remark that the full algebra \mathfrak{A}_{ql} can be obtained as an extension of the even subalgebra, generated by multiplying the latter by a single element, which can be whatever invertible local odd operator. Specifically, denoting the latter by \mathbf{O} , one has

$$\mathfrak{A}_{\text{ql}} = \mathfrak{A}_{\text{ql}}^+ \oplus (\mathfrak{A}_{\text{ql}}^+ \mathbf{O}). \quad (4.18)$$

Because of the symmetry, however, this is not the only extension giving rise to an algebra of operators that represent local observables. We have indeed shown that also the semilocal operator \mathbf{T}_+^z represents a local observable (which can be shifted by means of local operators to become $\mathbf{\Pi}_+^z(\ell)$, for any ℓ — Eq. (4.3)), therefore we can use it to extend $\mathfrak{A}_{\text{ql}}^+$ into another algebra associated with local observables, say

$$\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^+ = \mathfrak{A}_{\text{ql}}^+ \oplus (\mathfrak{A}_{\text{ql}}^+ \mathbf{T}_+^z). \quad (4.19)$$

This provides a more abstract definition of the semilocal theories informally introduced in the previous section: they are associated with different extensions of the symmetric subalgebra of quasilocal operators through operators that are not local but still represent local observables.

We are now in a position to generalise the requisite of pseudolocality (i.e. Eqs. (3.6), (3.8)) for the relevance of a conservation law to semilocal theories. To that aim, let us call \mathbf{T}_i , with $i = 1, \dots, n$, the operators that, in a given semilocal theory, are added to the symmetric subalgebra $\mathfrak{A}_{\text{ql}}^+$ to form a representation of local observables. Such an algebra is supposed to be closed both under time evolution and under shifts of lattice sites. Without loss of generality we can assume that \mathbf{T}_i commute with every local operator in the symmetric subalgebra $\mathfrak{A}_{\text{ql}}^+$ as long as the latter's support is far enough from position 0. A conservation law is then relevant if its density around position 0 reads $\mathbf{q}_0^{(0)} + \sum_{i=1}^n \mathbf{q}_0^{(i)} \mathbf{T}_i$, with $\mathbf{q}_0^{(i)} \in \mathfrak{A}_{\text{ql}}^+$ being even (strongly) quasilocal operators. We will refer to it as a “semilocal conservation law”.

If there are several semilocal theories, it could not be obvious which theories include all the relevant conservation laws. We propose then to extend the entire algebra of quasilocal operators by adding all the semilocal operators that generate the various semilocal theories. In the \mathbb{Z}_2 case this corresponds to considering the algebra

$$\mathfrak{A}_{\mathbb{Z}_2\text{-sl}} = \mathfrak{A}_{\text{ql}} \oplus (\mathfrak{A}_{\text{ql}} \mathbf{T}_+^z), \quad (4.20)$$

which we have already encountered when we defined the rotated Kramers-Wannier transformation (4.5).

We conjecture that in the extended space there exists a maximum-entropy representation of the macrostate emerging at infinite time after the global quench. We call it G -semilocal (generalised) Gibbs ensemble, where G is the symmetry used to extend the algebra (e.g., $G = \mathbb{Z}_2$ in most cases considered here). Extending the algebra allows us to recover a simple description of the late-time stationary values.

There is however an inconvenience: in the extended theory there are operators that do not represent local observables. In fact, except for the operators in the subalgebra that is common to all theories, there is no unambiguous way to associate local observables to operators. For example, in the \mathbb{Z}_2 case, both \mathfrak{A}_{ql} and $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^+$, which are subalgebras of $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}$, are generated by operators representing local observables, but the operators do not coincide. While the even local operators form a common subset $\mathfrak{A}_{\text{ql}}^+$ of both algebras (see Fig. 10) and thus enter the description of the local observables in both theories, it is less clear how to choose between the even semilocal or odd local operators ($\mathfrak{A}_{\text{ql}}^+ \mathbf{T}_+^z$ or $\mathfrak{A}_{\text{ql}}^+ \sigma_0^x$, respectively). We will expand on this in Section IV C 3. Before that, we provide an explicit example of a model with semilocal conservation laws.

1. Example: Dual XY model revisited

Contrary to the transverse-field Ising model, the quantum XY model

$$\mathbf{H}^{(\tau)} = \sum_{\ell \in \mathbb{Z}} J_x \tau_\ell^x \tau_{\ell+1}^x + J_y \tau_\ell^y \tau_{\ell+1}^y \quad (4.21)$$




FIG. 10. Representation of local observables in a \mathbb{Z}_2 -symmetric theory. Quasilocal operators (left column, red line) can be extended to semilocal ones through multiplication by \mathbf{T}_+^z . The full semilocal theory (black frame) can be projected either onto the quasilocal theory (red frame), or even semilocal theory (blue dashed frame): both can represent local observables.

is not self-dual under the Kramers-Wannier transformation. The duality transformation (4.5) maps its Hamiltonian into the one of the dual XY model

$$\mathbf{H}^{(\sigma)} = \sum_{\ell \in \mathbb{Z}} \sigma_{\ell-1}^x (J_x \mathbf{I} - J_y \sigma_{\ell}^z) \sigma_{\ell+1}^x. \quad (4.22)$$

$$\begin{aligned} \mathbf{q}_{\ell}^{(2,+;\tau)} &= J_x \tau_{\ell}^x \tau_{\ell+1}^x + J_y \tau_{\ell}^y \tau_{\ell+1}^y, & \mathbf{q}_{\ell}^{(3,+;\tau)} &= (J_x \tau_{\ell}^x \tau_{\ell+2}^x + J_y \tau_{\ell}^y \tau_{\ell+2}^y) \tau_{\ell+1}^z - (J_x + J_y) \tau_{\ell+1}^z, \\ \mathbf{q}_{\ell}^{(n,+;\tau)} &= (J_x \tau_{\ell}^x \tau_{\ell+n-1}^x + J_y \tau_{\ell}^y \tau_{\ell+n-1}^y) \prod_{j=1}^{n-2} \tau_{\ell+j}^z + (J_x \tau_{\ell}^y \tau_{\ell+n-3}^y + J_y \tau_{\ell}^x \tau_{\ell+n-3}^x) \prod_{j=1}^{n-4} \tau_{\ell+j}^z, & \text{for } n > 3, \\ \mathbf{q}_{\ell}^{(n,-;\tau)} &= (\tau_{\ell}^x \tau_{\ell+n-1}^y - \tau_{\ell}^y \tau_{\ell+n-1}^x) \prod_{j=1}^{n-2} \tau_{\ell+j}^z, & \text{for } n \geq 2. \end{aligned} \quad (4.26)$$

For the product we use the standard convention with $\prod_{j=\ell}^{\ell-1} \tau_j^{\alpha} = \mathbf{I}$. The upper indices n and \pm denote, respectively, the number of sites the charge's local density acts upon and the charge's parity under spatial reflection. Note that the reflection-odd charges $\mathbf{Q}^{(n,-;\tau)}$ do not depend on the coupling constants J_x and J_y .

Besides the abelian charges with densities (4.26) the XY model possesses also local charges of a staggered form $\sum_{\ell} (-1)^{\ell} \mathbf{w}_{\ell}$ that do not commute with $\mathbf{Q}^{(n,\pm;\tau)}$ [82]. Since the expectation value of any staggered operator is

We denote the corresponding local Hamiltonian densities by $\mathbf{h}_{\ell}^{(\tau)} = J_x \tau_{\ell}^x \tau_{\ell+1}^x + J_y \tau_{\ell}^y \tau_{\ell+1}^y$, and $\mathbf{h}_{\ell}^{(\sigma)} = \sigma_{\ell-1}^x (J_x \mathbf{I} - J_y \sigma_{\ell}^z) \sigma_{\ell+1}^x$, respectively.

Under the duality transformation $\mathcal{D}_{\mathbb{Z}_2}$, the spin-flip symmetry $\mathcal{P}_{\tau}^x[\mathbf{h}_{\ell}^{(\tau)}] = \mathbf{h}_{\ell}^{(\sigma)}$, for any ℓ , becomes the following invariance of the dual XY model's local densities:

$$\lim_{n \rightarrow \infty} [\sigma_{-n}^x \sigma_n^x] \mathbf{h}_{\ell}^{(\sigma)} [\sigma_{-n}^x \sigma_n^x] = \mathbf{h}_{\ell}^{(\sigma)}, \quad \forall \ell. \quad (4.23)$$

While this invariance holds trivially for (quasi)local densities, the same is not true for the semilocal operators. Considering, for example, half-infinite strings $\mathbf{\Pi}_{\sigma,+}^z(\ell)$, we have

$$\lim_{n \rightarrow \infty} [\sigma_{-n}^x \sigma_n^x] \mathbf{\Pi}_{\sigma,+}^z(\ell) [\sigma_{-n}^x \sigma_n^x] = -\mathbf{\Pi}_{\sigma,+}^z(\ell), \quad (4.24)$$

i.e., $\mathbf{\Pi}_{\sigma,+}^z(\ell)$ are odd under the operation that is dual to \mathcal{P}_{τ}^x . Indeed, recall that $\mathcal{D}_{\mathbb{Z}_2}$ maps all operators that are odd under \mathcal{P}_{τ}^x into semilocal operators, as described in Table I. With this in mind, let us now consider the charges of the XY model and its dual counterpart.

The quantum XY model is non-abelian integrable [82]. That is to say, the Hamiltonian commutes with infinitely many pseudolocal operators, not necessarily commuting with one another. Its translationally invariant local charges are

$$\mathbf{Q}^{(n,\pm;\tau)} = \sum_{\ell \in \mathbb{Z}} \mathbf{q}_{\ell}^{(n,\pm;\tau)}, \quad (4.25)$$

where the local densities (including the Hamiltonian's one $\mathbf{h}_{\ell}^{(\tau)} \equiv \mathbf{q}_{\ell}^{(2,+;\tau)}$) read (see, e.g., [96])

zero in a translationally invariant state, these additional nonabelian charges are irrelevant for our discussion and we will therefore not report their explicit form here.

Consider now the densities (4.26) after the duality transformation. According to Table I the local densities $\{\mathbf{q}_{\ell}^{(2n,+;\tau)}, \mathbf{q}_{\ell}^{(2n+1,-;\tau)}\}_{n \in \mathbb{N}}$ remain local even in the σ representation, since they are even under \mathcal{P}_{τ}^x . Instead $\{\mathbf{q}_{\ell}^{(2n+1,+;\tau)}, \mathbf{q}_{\ell}^{(2n,-;\tau)}\}_{n \in \mathbb{N}}$ are odd under \mathcal{P}_{τ}^x and are mapped into operators with half-infinite strings. Hence,

the set $\{\mathbf{Q}^{(2n+1,+;\tau)}, \mathbf{Q}^{(2n,-;\tau)}\}_{n \in \mathbb{N}}$ is mapped into a set of semilocal charges in the dual XY model. For example, the density of the charge $\mathbf{Q}^{(3,+;\tau)}$ becomes

$$\mathbf{q}_\ell^{(3,+;\sigma)} = [\sigma_{\ell-1}^x (J_x \sigma_\ell^x \sigma_{\ell+1}^x + J_y \sigma_\ell^y \sigma_{\ell+1}^y) \sigma_{\ell+2}^x - (J_x + J_y) \mathbf{I}] \times \mathbf{\Pi}_{\sigma,+}^z(\ell+1). \quad (4.27)$$

In the symmetric quench we considered in Section III C this is a special charge: it is the only one with a nonzero expectation value in the state $|\uparrow\rangle_\sigma$. This is because its density is the only one among $\mathbf{q}_\ell^{(n,\pm;\sigma)}$ containing a term that consists solely of σ_j^z matrices.

2. Example: generic model revisited

Let us reconsider the generic model of Section III C — cf. Eqs (2.4) and (3.20) — which is described by the Hamiltonian

$$\mathbf{H} = \sum_{\ell \in \mathbb{Z}} \sigma_{\ell-1}^x (\mathbf{1} - \sigma_\ell^z) \sigma_{\ell+1}^x + w_1 \sigma_\ell^z + \sum_{n=1}^{\infty} w_{2,n} \sigma_\ell^z \sigma_{\ell+n}^z. \quad (4.28)$$

As anticipated in Section II B, this Hamiltonian has a semilocal conservation law, namely

$$\mathbf{Q} = \sum_{\ell \in \mathbb{Z}} \mathbf{\Pi}^z(\ell). \quad (4.29)$$

The expectation value of this semilocal charge is defined through clustering (i.e., using Eq. (2.3)), which yields $\langle \mathbf{\Pi}^z(j) \rangle_t^2 = \cos^2 \varphi$ for its density. The latter provides a positive lower bound for the string order parameter $\lim_{n \rightarrow \infty} \langle \prod_{j=-n}^n \sigma_n^z \rangle$ at infinite time through Eq. (2.6). According to Ref. [12], a thermal state can not exhibit string order, therefore we can immediately conclude that, for $\cos \varphi \neq 0$, the state at infinite time is not thermal. This is consistent with the behaviour shown in Fig. 9.

3. Canonical and non-canonical descriptions

A priori we do not see any reason to choose one theory of local observables over the other. If however we imagine the theoretical system as an idealisation of an experiment and the experimental apparatus as something that goes beyond the system under investigation, a theory could be somehow selected by how the apparatus was designed. For example, in the \mathbb{Z}_2 case, if the experimental apparatus is able to preserve spin-flip symmetry with high accuracy, the even \mathbb{Z}_2 semilocal theory could become a better framework where to study the effect of noise (even under spin flip) or other blind spots of the experiment. The question then becomes, how the G -semilocal (generalised) Gibbs ensemble is represented within the chosen theory. To answer it, the ensemble should be projected onto the subalgebra associated with that theory. There are two possibilities:

1. the G -semilocal (generalised) Gibbs ensemble is part of the subalgebra, hence the projected ensemble coincides with the maximum-entropy ensemble in the theory. We call such theories “canonical”;
2. the G -semilocal (generalised) Gibbs ensemble includes also conserved operators outside the theory; the theory is termed “non-canonical”.

For example, a \mathbb{Z}_2 -semilocal (generalised) Gibbs ensemble $\rho_{\mathbb{Z}_2\text{-sl}}$ constructed from an even local charge \mathbf{Q}_{ql}^+ and an even semilocal charge $\mathbf{Q}_{\mathbb{Z}_2\text{-sl}}^+$ can be written as

$$\rho_{\mathbb{Z}_2\text{-sl}} \propto e^{-\mathbf{Q}_{\text{ql}}^+ - \mathbf{Q}_{\mathbb{Z}_2\text{-sl}}^+}. \quad (4.30)$$

It coincides with its projection onto the theory represented by $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^+$, i.e.,

$$\rho_{\mathbb{Z}_2\text{-sl}} = \rho_{\mathbb{Z}_2\text{-sl}} \Big|_{\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^+}. \quad (4.31)$$

In the quasilocal theory associated with \mathfrak{A}_{ql} it instead takes a different form. Specifically, only the first term in

$$\rho_{\mathbb{Z}_2\text{-sl}} \propto e^{-\mathbf{Q}_{\text{ql}}^+ \cosh \mathbf{Q}_{\mathbb{Z}_2\text{-sl}}^+ - e^{-\mathbf{Q}_{\text{ql}}^+} \sinh \mathbf{Q}_{\mathbb{Z}_2\text{-sl}}^+} \quad (4.32)$$

survives the projection onto the quasilocal algebra (the second one consists of odd powers of $\mathbf{Q}_{\mathbb{Z}_2\text{-sl}}$ and thus contains strings). The projected ensemble

$$\rho_{\mathbb{Z}_2\text{-sl}} \Big|_{\mathfrak{A}_{\text{ql}}} \propto e^{-\mathbf{Q}_{\text{ql}}^+ \cosh \mathbf{Q}_{\mathbb{Z}_2\text{-sl}}^+} \quad (4.33)$$

does not maximize the von Neumann entropy (3.2) constrained by the pseudolocal charges, making the theory non-canonical [97].

Returning to the problem of defining and classifying nonequilibrium phases after global quenches, the way local observables are represented in a canonical theory constitutes a fundamental distinction, which goes much beyond the differences associated with the multiplicity and the symmetries of the conservation laws. It was observed in Ref. [65] that the generalised eigenstate thermalisation hypothesis (gETH), according to which all eigenstates with the same local integrals of motion are locally equivalent, could be sufficient to prove that a maximum-entropy ensemble description is possible (the criticism raised in Ref. [98] is resolved once pseudolocal integrals of motion are taken into account). In non-canonical theories we claim that gETH fails. As a matter of fact, the following much weaker assumption fails: in the limit of infinite time the only information retained from the initial state is encoded in the (pseudo)local conserved operators.

We have already shown an example of the breakdown of such a key property in the previous section, when we tried to describe the infinite-time limit after a symmetric quench in the dual XY model through the maximum-entropy ensemble of a non-canonical theory, which in the specific case was the (quasilocal) generalised Gibbs ensemble. That ensemble was unable to describe even a local observable such as σ_ℓ^z — Fig. 8. As expected, instead,

the stationary values are described by the \mathbb{Z}_2 -semilocal generalised Gibbs ensemble (4.30). In that specific case, the relaxation to the \mathbb{Z}_2 -semilocal GGE is a trivial consequence of the established result that noninteracting systems relax to generalised Gibbs ensembles. More generally, if a duality transformation between a canonical theory and the quasilocal theory is known, proving relaxation to the (generalised) Gibbs ensemble in the canonical theory becomes equivalent to proving relaxation of non-symmetric states in the standard quasilocal one.

Non-symmetric states. So far we have assumed that the entire system (initial state and Hamiltonian) is symmetric. On the other hand, we defined the semilocal charges in an extended algebra, which includes also non-symmetric operators. It is then natural to wonder whether a semilocal charge could make sense also with non-symmetric initial states. To that aim, we consider again the state $|\Psi\rangle^{(\sigma)} = |\nearrow_\theta\rangle_\sigma$ and $\theta \notin \{0, \pi\}$. As discussed in Section IV B, we must be careful about the interpretation of $|\Psi\rangle$ when we extend the theory so as to include also semilocal operators. It is again convenient to map even semilocality into odd locality. Appendix D 1 proves

$$|\Psi\rangle^{(\tau)} = |\beta_\theta\rangle_\tau := \frac{\sum_{\underline{s}} e^{-\frac{\beta\theta}{2} E(\underline{s})} |\underline{s}\rangle_\tau}{\sum_{\underline{s}} e^{-\beta\theta E(\underline{s})}}, \quad (4.34)$$

where $\beta_\theta = -\log \tan(\theta/2)$, $\underline{s} = (\dots, s_{-1}, s_0, s_1, \dots)$, with $s_j \in \{-1, 1\}$ denoting the eigenvalues of the spin operator τ_j^z , and $E(\underline{s})$ is the energy of the classical Ising model $E(\underline{s}) = -\sum_j s_j s_{j+1}$.

One can show that $|\beta\rangle_\tau$ has clustering properties for any finite β , i.e., for $\theta \notin \{0, \pi\}$ (see Appendix D 2). Importantly, despite the tilted state not being symmetric, the corresponding state in the dual representation, i.e., $|\beta_\theta\rangle_\tau$, is even under \mathcal{P}_τ^x . More generally we can understand this by interpreting $|\Psi\rangle$ as the ground state of a symmetric Hamiltonian in which the symmetry is spontaneously broken at zero temperature in the σ representation (a simple Hamiltonian with this symmetry-breaking ground state is known [99]: the quantum XY model in a transverse field, with density $\mathbf{h}_\ell = J_x \sigma_\ell^x \sigma_{\ell+1}^x + J_y \sigma_\ell^y \sigma_{\ell+1}^y - 2\sqrt{J_x J_y} \sigma_\ell^z$, with $J_y/J_x = \cos^2 \theta$). It turns out that the symmetry in the dual Hamiltonian remains unbroken and the ground state is symmetric. This implies that even semilocal operators have zero expectation values in the original system, and hence the expectation values of all semilocal charges vanish in $|\Psi\rangle$. On the one hand, this justifies taking into account semilocal charges also with non-symmetric initial states; on the other hands, it shows their irrelevance (as the corresponding integrals of motion vanish).

As a matter of fact, the naive approach of interpreting half-infinite strings as infinite products of Pauli matrices gives, in this case, the correct result: since $|\nearrow_\theta\rangle$ is a product state, for $\theta \notin \{0, \pi\}$ one has $|\langle \nearrow_\theta | \sigma_j^z | \nearrow_\theta \rangle| < 1$

for all j , so the strings are exponentially suppressed. The two approaches give the same result because the sequence of the expectation values of $\{\mathbf{O}_0 \Pi_{\ell=0}^n \sigma_\ell^z\}$ converges (to zero) as $n \rightarrow \infty$ for any operator \mathbf{O}_0 quasilocalised around 0. This actually allows one to go even further and conclude that also the expectation values of odd semilocal operators vanish.

4. Remark

We conclude with a clarification. So far we have presented (3.17) as a Hamiltonian with a \mathbb{Z}_2 symmetry, but, in fact, it exhibits a $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry. Indeed, the energy density is also even under

$$\mathcal{P}_e^z[\mathbf{h}_j] = \lim_{n \rightarrow \infty} \left[\prod_{\ell=-n}^n \sigma_{2\ell}^z \right] \mathbf{h}_j \left[\prod_{\ell=-n}^n \sigma_{2\ell}^z \right]. \quad (4.35)$$

The same comment applies to the symmetric initial state studied in the example of Section III C, namely, $|\uparrow\rangle$. The extended algebra is now obtained by supplementing the local operators that are even under both spin flips by two (commuting) semilocal operators $\mathbf{T}_{2,0}^z$ and $\mathbf{T}_{2,1}^z$, which play the role of products of Pauli matrices σ^z over even or odd sites, respectively, extending from site 0 or 1 to infinity (see Appendix B for a proper definition).

Let $\mathfrak{A}_{\text{ql}}^{+\bullet}$, $\mathfrak{A}_{\text{ql}}^{\bullet+}$, $\mathfrak{A}_{\text{ql}}^{\pm\pm}$, $\mathfrak{A}_{\text{ql}}^{++}$ be the subalgebras of quasilocal operators even under \mathcal{P}_e^z , $\mathcal{P}_o^z := \mathcal{P}_e^z \circ \mathcal{P}^z$, $\mathcal{P}^z (= \mathcal{P}_e^z \circ \mathcal{P}_o^z)$, both \mathcal{P}^z and \mathcal{P}_e^z , respectively ($\mathfrak{A}_{\text{ql}}^{\pm\pm}$ was previously denoted by $\mathfrak{A}_{\text{ql}}^+$, since only the symmetry \mathcal{P}^z was relevant). We identify five subalgebras generated by operators representing local observables:

- \mathfrak{A}_{ql} : this is the standard algebra of quasilocal operators, given by $\mathfrak{A}_{\text{ql}}^{\pm\pm} \oplus (\mathfrak{A}_{\text{ql}}^{\pm\pm} \mathbf{O})$, with \mathbf{O} an invertible local odd operator;
- $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^{+\bullet}$: this is the \mathbb{Z}_2 -semilocal algebra even under spin flip on even sites, and it is given by $\mathfrak{A}_{\text{ql}}^{+\bullet} \oplus (\mathfrak{A}_{\text{ql}}^{+\bullet} \mathbf{T}_{2,0}^z)$;
- $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^{\bullet+}$: this is the \mathbb{Z}_2 -semilocal algebra even under spin flip on odd sites, and it is given by $\mathfrak{A}_{\text{ql}}^{\bullet+} \oplus (\mathfrak{A}_{\text{ql}}^{\bullet+} \mathbf{T}_{2,1}^z)$;
- $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^{\pm\pm}$: this is the \mathbb{Z}_2 -semilocal algebra even under spin flip, and it is given by $\mathfrak{A}_{\text{ql}}^{\pm\pm} \oplus (\mathfrak{A}_{\text{ql}}^{\pm\pm} \mathbf{T}_{2,0}^z \mathbf{T}_{2,1}^z)$;
- $\mathfrak{A}_{\mathbb{Z}_2 \times \mathbb{Z}_2\text{-sl}}^{++}$: this is the $\mathbb{Z}_2 \times \mathbb{Z}_2$ -semilocal algebra even both under spin flip on odd sites and under spin flip on even sites, and it is given by $\mathfrak{A}_{\text{ql}}^{++} \oplus (\mathfrak{A}_{\text{ql}}^{++} \mathbf{T}_{2,0}^z) \oplus (\mathfrak{A}_{\text{ql}}^{++} \mathbf{T}_{2,1}^z) \oplus (\mathfrak{A}_{\text{ql}}^{++} \mathbf{T}_{2,0}^z \mathbf{T}_{2,1}^z)$.

It turns out that the $\mathbb{Z}_2 \times \mathbb{Z}_2$ -semilocal generalised Gibbs ensemble emerging after a quench from $|\uparrow\rangle$ belongs to the intersection of $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^{\pm\pm}$ and $\mathfrak{A}_{\mathbb{Z}_2 \times \mathbb{Z}_2\text{-sl}}^{++}$ [100]. The latter two algebras represent therefore canonical theories. On the other hand, \mathfrak{A}_{ql} , $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^{+\bullet}$, and $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^{\bullet+}$ are associated

with non-canonical theories. There, the stationary state capturing the infinite time expectation values is not a maximum-entropy statistical ensemble. Once projected onto the quasilocal theory, the $\mathbb{Z}_2 \times \mathbb{Z}_2$ -semilocal generalised Gibbs ensemble has the form (4.33) with $\mathbf{Q}_{\text{ql}} = 0$ (the form of the ensemble is specific to the initial state considered, i.e., $|\uparrow\rangle$). Since it is even both under \mathcal{P}_e^z and \mathcal{P}_o^z , the $\mathbb{Z}_2 \times \mathbb{Z}_2$ -semilocal generalised Gibbs ensemble belongs to the intersection of \mathfrak{A}_{ql} , $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^+$, and $\mathfrak{A}_{\mathbb{Z}_2\text{-sl}}^+$: the three theories share the same projected ensemble. This makes the algebraic structure associated with the $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry of the dual XY model redundant, and the reader can now understand why we have completely overlooked this larger symmetry group when describing the model and its conservation laws.

The remaining question is: can we realise that the state does not locally relax to the maximum-entropy state without comparing the predictions from the maximum-entropy state? This is the subject of the next two sections.

V. SIGNATURES OF SEMILOCAL ORDER

A way to assess whether a theory is canonical or not is by perturbing the initial state with an operator that represents a local observable but does not belong to the common subalgebra. Specifically, we consider a unitary transformation with finite support connecting different symmetry sectors, such as σ_r^x in the spin-flip case [31]. Such perturbations are semilocal in the eyes of charges belonging to a different theory, which will therefore be affected in a nonlocal way. If the quasilocal theory is not canonical, we find it reasonable to expect signatures of such a nonlocality in the entanglement of subsystems.

Being the first investigation of this kind in nonequilibrium symmetry-protected topological order phases, we focus on the entanglement entropies of connected blocks of spins A , which in the rest of the paper will be identified with the following set of sites

$$A \equiv \{1, \dots, \ell\}. \quad (5.1)$$

To that aim, we need to construct reduced density matrices of finite subsystems. Reduced density matrices describe the expectation values of local observables represented by local operators with support in the subsystem. In a spin-1/2 chain they can be expanded in an orthogonal basis of Hermitian matrices O_A ($\text{tr}[O_A O'_A] = \delta_{O_A, O'_A} \text{tr}[\mathbb{I}_A]$) representing local operators \mathbf{O}_A with support in the subsystem A

$$\rho_A(\Psi) = \frac{1}{\text{tr}[\mathbb{I}_A]} \sum_{O_A} \langle \Psi | \mathbf{O}_A | \Psi \rangle O_A, \quad (5.2)$$

where \mathbf{O}_A is the operator that acts like O_A in the subsystem and like the identity elsewhere. Reduced density

matrices are therefore embedded in \mathfrak{A}_{ql} : investigating reduced density matrices implicitly selects the quasilocal theory of local observables. The Rényi entanglement entropies are then defined as

$$S_\alpha(\ell, \Psi) \equiv \frac{1}{1-\alpha} \log \text{tr}[\rho_A^\alpha(\Psi)], \quad (5.3)$$

which include the von Neumann entropy “ S_1 ”, also known as entanglement entropy, as the limit $\alpha \rightarrow 1^+$ (this limit makes sense because the support of the distribution of eigenvalues of the density matrix is bounded, and hence the Rényi entropies with $\alpha = 2, 3, \dots$ characterise the distribution completely [101]).

A. Reduced density matrices across different theories of local observables

In a symmetric system $\rho_A(\Psi)$ is embedded in the subalgebra common to all the theories of local observables, which in the \mathbb{Z}_2 case is $\mathfrak{A}_{\text{ql}}^+$ (the sum in Eq. (5.2) can be restricted to matrices associated with operators that are even under spin flip). Thus, $\rho_A(\Psi)$ is also represented in the other theories of local observables, although, there, its interpretation as a subsystem’s reduced density matrix breaks down. The simplest way to understand the representation of a reduced density matrix in a semilocal theory is by mapping semilocal operators into local operators through a duality transformation. This is discussed below, where we specify the action of the duality transformation on the reduced density matrices and show how to estimate the entanglement entropy in the semilocal theory using the dual reduced density matrices.

1. Restricted duality transformation

We remind the reader that, when restricted to the even quasilocal subalgebra, the duality transformation defined in Eq. (4.5) maps local operators into local operators (see Table I). This allows us to use the rotated Kramer-Wannier duality transformation also in finite subsystems. For a subsystem \tilde{A} of $\ell + 1$ sites

$$\tilde{A} = \{1, \dots, \ell + 1\}, \quad (5.4)$$

we define $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$ as follows

$$\begin{aligned} \mathbb{I}^{\otimes(j-1)} \otimes \sigma^z \otimes \mathbb{I}^{\otimes(\ell+1-j)} &\mapsto \mathbb{I}^{\otimes(j-1)} \otimes (\tau^z)^{\otimes 2} \otimes \mathbb{I}^{\otimes(\ell-j)} \\ \mathbb{I}^{\otimes(j'-1)} \otimes (\sigma^x)^{\otimes 2} \otimes \mathbb{I}^{\otimes(\ell-j')} &\mapsto \mathbb{I}^{\otimes j'} \otimes \tau^x \otimes \mathbb{I}^{\otimes(\ell-j')}, \end{aligned} \quad (5.5)$$

where $j = 1, \dots, \ell$ and $j' = 1, \dots, \ell - 1$. Note that we have given up the bold notation because σ^α are not spin operators in an infinite system but Pauli matrices spanning the local 2-dimensional Hilbert space.

Since the matrices on the left hand side of Eq. (5.5) generate all matrices representing even operators with support in A — (5.1) (acting therefore as the identity

on site $\ell + 1$), it is convenient to consider an alternative representation of $S_\alpha(\ell, \Psi)$ (cf. Eq. (5.3)):

$$S_\alpha(\ell, \Psi) = \frac{\log \text{tr}[(\rho_A(\Psi) \otimes \frac{\mathbb{I}}{2})^\alpha]}{1 - \alpha} - \log 2. \quad (5.6)$$

Remark. Similarly to $\mathcal{D}_{\mathbb{Z}_2}^{-1}$, also $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$ is an algebra homomorphism, although it is defined only in a subspace of the full matrix algebra associated with the subsystem. Specifically, let us introduce the following:

$\mathfrak{A}_A^{(\sigma)}$: the full matrix algebra generated by the tensor products of Pauli matrices $\otimes_{j \in A} \sigma_j^{\alpha_j}$, with $\alpha_j = 0, x, y, z$, in the subsystem consisting of the sites $j \in A$ (and analogously for $\mathfrak{A}_A^{(\tau)}$);

$\mathfrak{A}_A^{+(\sigma)}$: the restriction of $\mathfrak{A}_A^{(\sigma)}$ to matrices commuting with $\otimes_{j \in A} \sigma_j^z$, which we refer to as even;

$\mathfrak{A}_A^{+(\tau)}$: the restriction of $\mathfrak{A}_A^{(\tau)}$ to matrices commuting with $\otimes_{j \in \tilde{A}} \tau_j^x$, which we refer to as even;

$\mathfrak{A}_B \otimes \mathbb{I}_C$: (with $B \cap C = \emptyset$) the matrix subalgebra of $\mathfrak{A}_{B \cup C}$ consisting of the matrices in \mathfrak{A}_B extended with the identity in C , to act on sites $j \in B \cup C$.

We recognise $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$, defined in Eq. (5.5), as the *trace-preserving* isomorphism that acts like (4.1) in a restricted space of matrices with support on \tilde{A} :

$$\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell) : \mathfrak{A}_A^{+(\sigma)} \otimes \mathbb{I}_{\ell+1} \rightarrow \tilde{\mathfrak{A}}_A^{+(\tau)} \subset \mathfrak{A}_A^{+(\tau)}. \quad (5.7)$$

Here the image $\tilde{\mathfrak{A}}_A^{+(\tau)}$, which is a subalgebra of $\mathfrak{A}_A^{+(\tau)}$, is isomorphic to $\mathfrak{A}_A^{+(\sigma)}$ despite consisting also of operators whose range extends up to $\ell + 1$.

Since we are considering a mapping that spoils the notion of a spatial subsystem, it is convenient to consider a generalisation of reduced density matrices which, instead of describing the expectation values of operators with support in a given region, describes the expectation values of operators that are represented by matrices belonging to a given algebra. Specifically, we introduce the following notation

$$\hat{P}_{\mathfrak{A}}(\Psi) = \frac{1}{\text{tr}[\mathbb{I}_{\mathfrak{A}}]} \sum_{O \in \mathfrak{A}} \langle \Psi | \mathbf{O} | \Psi \rangle O, \quad (5.8)$$

where $\mathbb{I}_{\mathfrak{A}}$ is the identity in \mathfrak{A} and the sum is over an orthogonal basis of Hermitian matrices, such that $\text{tr}[OO'] = \delta_{OO'}$ (we have always in mind operators written as tensor products of Pauli matrices).

If the reduced density matrix $\rho_A(\Psi)$ — (5.2) of the subsystem is even, the matrix $\rho_A(\Psi) \otimes \frac{\mathbb{I}}{2}$, which appears in Eq. (5.6), belongs to $\mathfrak{A}_A^{+(\sigma)} \otimes \mathbb{I}_{\ell+1}$,

$$\rho_A(\Psi) \otimes \frac{\mathbb{I}}{2} = \hat{P}_{\mathfrak{A}_A^{+(\sigma)} \otimes \mathbb{I}_{\ell+1}}(\Psi), \quad (5.9)$$

and hence it is in the domain of $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$.

Since this mapping preserves the trace, $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$ does not affect the value of functionals of $\rho_A(\Psi)$ such as the Rényi entropies (5.6). In light of Eq. (5.7), $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$ maps $\rho_A^{(\sigma)}(\Psi) \otimes \frac{\mathbb{I}}{2}$ into the projection of $\rho_A^{(\tau)}(\Psi)$ on the subalgebra $\tilde{\mathfrak{A}}_A^{+(\tau)}$

$$\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell) : \rho_A^{(\sigma)}(\Psi) \otimes \frac{\mathbb{I}}{2} \mapsto \rho_A^{(\tau)}(\Psi)|_{\tilde{\mathfrak{A}}_A^{+(\tau)}}. \quad (5.10)$$

We will see in a moment how to carry out this projection.

2. Dual reduced density matrices

By construction, the operators $\mathbf{O}^{(\sigma)}$ that have support in A and are represented by matrices $O^{(\sigma)} \otimes \mathbb{I}$ in the domain of $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$ are mapped by the rotated Kramers-Wannier duality transformation (4.5) into operators $\tilde{\mathbf{O}}$ that have support in \tilde{A} and are represented by $\tilde{O} = \mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)[O^{(\sigma)} \otimes \mathbb{I}]$. The following equality holds for them:

$$\langle \Psi | \mathbf{O} | \Psi \rangle^{(\sigma)} = \langle \Psi | \tilde{\mathbf{O}} | \Psi \rangle^{(\tau)}. \quad (5.11)$$

Thus, the (extended) reduced density matrix $\rho_A^{(\sigma)}(\Psi) \otimes \frac{\mathbb{I}}{2}$ is mapped into

$$\rho_A^{(\sigma)}(\Psi) \otimes \frac{\mathbb{I}}{2} \mapsto 2^{-\ell-1} \sum_{\substack{O \in \mathfrak{A}_A^{+(\sigma)} \\ \tilde{O} = \mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)[O \otimes \mathbb{I}]}} \langle \Psi | \tilde{\mathbf{O}} | \Psi \rangle^{(\tau)} \tilde{O}, \quad (5.12)$$

which can be compactly written as

$$\hat{P}_{\mathfrak{A}_A^{+(\sigma)} \otimes \mathbb{I}_{\ell+1}} \mapsto \hat{P}_{\tilde{\mathfrak{A}}_A^{+(\tau)}}. \quad (5.13)$$

While the right-hand side of Eq. (5.12) resembles

$$\rho_A^{(\tau)}(\Psi) := \hat{P}_{\tilde{\mathfrak{A}}_A^{+(\tau)}} = 2^{-\ell-1} \sum_{\tilde{O} \in \mathfrak{A}_A^{+(\tau)}} \langle \Psi | \tilde{\mathbf{O}} | \Psi \rangle^{(\tau)} \tilde{O}, \quad (5.14)$$

the sum in Eq. (5.12) does not cover the full matrix algebra of the subsystem ($\tilde{\mathfrak{A}}_A^{+(\tau)}$ is strictly contained in $\mathfrak{A}_A^{(\tau)}$). In order to obtain Eq. (5.12) from $\rho_A^{(\tau)}(\Psi)$ we must therefore project out all the operators that are not in the image of $\mathcal{D}_{\mathbb{Z}_2}^{-1}(\ell)$. This can be done by identifying the symmetries characterising $\tilde{\mathfrak{A}}_A^{+(\tau)}$:

1. Only operators even under spin flip \mathcal{P}_τ^x are represented in $\tilde{\mathfrak{A}}_A^{+(\tau)}$;
2. Operators $\tilde{\mathbf{O}}$ that are represented on the right hand side of (5.12) commute both with τ_1^z and $\tau_{\ell+1}^z$ — see Eq. (5.5).

These \mathbb{Z}_2 symmetries can be enforced by averaging over the corresponding (Hermitian) flip matrices P_i as follows

$$\hat{P}_{\mathfrak{A}_A^{+(\tau)}}(\Psi) = \bar{\rho}_A^{(\tau)}(\Psi) := \sum_{j_{1,2,3}=0}^1 \frac{P_3^{j_3} P_2^{j_2} P_1^{j_1} \rho_A^{(\tau)}(\Psi) P_1^{j_1} P_2^{j_2} P_3^{j_3}}{8},$$

$$P_1 = (\tau^x)^{\otimes(\ell+1)}, \quad P_2 = \tau^z \otimes \mathbb{I}^{\otimes \ell}, \quad P_3 = \mathbb{I}^{\otimes \ell} \otimes \tau^z$$
(5.15)

(see Fig. 11 for a graphical representation).




FIG. 11. A graphical representation of the mapping (5.12) when the subsystem A consists of a single site. Operators $P_{1,2,3}$ are colored and underlined. They anticommute with the operators in the correspondingly colored circles: their effect is to remove the undesired terms in the density matrix (5.14).

To sum up, we have started with enlarging the subsystem by one site $\rho_A(\Psi) \rightarrow \rho_A(\Psi) \otimes \frac{\mathbb{I}}{2}$; this has allowed us to apply the duality transformation (5.5), which alters the meaning of spatial interval. We have then found a representation of $\rho_A(\Psi) \otimes \frac{\mathbb{I}}{2}$ in terms of reduced density matrices in the semilocal theory with the new notion of spatial subsystem. The final object, $\bar{\rho}_A^{(\tau)}(\Psi)$ is an average of density matrices selecting the part that is symmetric under three spin-flip transformations. Finally, the entanglement entropies can be expressed in terms of $\bar{\rho}_A^{(\tau)}(\Psi)$ by applying Eq. (5.6):

$$S_\alpha(\ell, \Psi) = \frac{\log \text{tr}[(\bar{\rho}_A^{(\tau)}(\Psi))^\alpha]}{1 - \alpha} - \log 2, \quad (5.16)$$

where $\log 2$ cancels the effect of the spurious double degeneracy introduced when the subsystem was enlarged.

Bounds on the entropies of the dual RDMS.

We show here that, up to $\mathcal{O}(1)$ corrections, the averaged density matrix $\bar{\rho}_A^{(\tau)}(\Psi)$ can be replaced by $\rho_A^{(\tau)}(\Psi)$ in (5.16). Since no special property of the density matrix and of the \mathbb{Z}_2 symmetry we average over is required, we consider a generic density matrix ρ (Hermitian positive semidefinite operator of trace one) and the symmetry generated by a Hermitian involution P ; Eq. (5.15) is just a subsequent application of such an average.

We define the symmetric density matrix

$$\bar{\rho} = \frac{\rho + P\rho P}{2}, \quad (5.17)$$

and we call \bar{S}_α its Rényi entropies (S_α indicates instead the entropies of ρ).

Since $\bar{\rho}$ can be interpreted as the sum of two density matrices weighted with the probability distribution $\{1/2, 1/2\}$, general properties of the von Neumann entropy (see Section 12.2 of Ref. [102]) imply

$$S_1 \leq \bar{S}_1 \leq S_1 + \log 2 \quad (5.18)$$

For instance, the lower bound follows directly from the concavity of the von Neumann entropy. We obtain an analogous result for the Rényi entropies, which satisfy

$$S_\alpha \leq \bar{S}_\alpha \leq S_\alpha + \frac{\alpha \log 2}{\alpha - 1}; \quad (5.19)$$

see Appendix E 1. Returning to Eq. (5.16), these bounds imply

$$S_\alpha(\ell, \Psi) = S_\alpha^{(\tau)}(\ell + 1, \Psi) + \mathcal{O}(\ell^0), \quad (5.20)$$

where we have defined the entropy of the subsystem \tilde{A} in the dual representation

$$S_\alpha^{(\tau)}(\ell + 1, \Psi) = \frac{\log \text{tr}[(\rho_A^{(\tau)}(\Psi))^\alpha]}{1 - \alpha}. \quad (5.21)$$

B. Excess of entropy

The protocol. We consider a system invariant under \mathcal{P}^z and compare two quench protocols:

- (I) nonequilibrium time evolution of a translationally invariant state $|\Psi(t)\rangle = e^{-i\mathbf{H}t} |\Psi(0)\rangle$;
- (II) nonequilibrium time evolution of the same state as in (I) perturbed by a local(ised) unitary operator $|\Psi_r(t)\rangle = e^{-i\mathbf{H}t} \sigma_r^x |\Psi(0)\rangle$.

The question is: do the Rényi entropies of subsystems discriminate between the two protocols in the limit of infinite time? Inspired by Refs. [42, 43], we define the excess of entropy as

$$\Delta_r S_\alpha(\ell, t) = S_\alpha(\ell, \Psi_r(t)) - S_\alpha(\ell, \Psi(t)), \quad (5.22)$$

where the subsystem of length ℓ is associated with the sites $\{1, \dots, \ell\}$, that is to say, the reduced density matrix to be investigated reads

$$\rho_A(\Psi) = \frac{1}{2^\ell} \sum_{\substack{\{\alpha\}_\ell \\ \alpha_j=0,x,y,z}} \langle \Psi | \prod_{j=1}^{\ell} \sigma_j^{\alpha_j} | \Psi \rangle \bigotimes_{j=1}^{\ell} \sigma_j^{\alpha_j}. \quad (5.23)$$

Here Ψ can be either $\Psi_r(t)$ or $\Psi(t)$.

In Section IV C we have discussed the limit of infinite time in the quench protocol (I), showing that, in the quasilocal theory, the state locally relaxes to a projected \mathbb{Z}_2 -semilocal generalised Gibbs ensemble. In order to compute the excess of entropy, we should also take the limit of infinite time in the protocol (II), which goes however beyond our assumption of translational invariance. Although a theory to describe the infinite-time limit in inhomogeneous systems with semilocal charges is still missing, this problem can be solved in the dual representation, where the spin flip is mapped into a topological excitation resulting in a domain-wall initial state [31]. Specifically, if the initial state is $|\uparrow\rangle_\sigma$, it follows from the first equation of (4.7) that after flipping the r -th spin the state in the dual representation becomes $|\cdots \downarrow_{r-1} \downarrow_r \uparrow_{r+1} \uparrow_{r+2} \cdots\rangle_\tau$ (or $|\cdots \uparrow_{r-1} \uparrow_r \downarrow_{r+1} \downarrow_{r+2} \cdots\rangle_\tau$, depending on how the symmetry is broken). The problem then moves to identifying the stationary state describing the limit of infinite time after a quench from a domain-wall state in a canonical theory, which is a much more studied situation [103–108]. The drawback is that we need to express the observable under investigation in the dual representation; for the Rényi entropies this corresponds to using Eq. (5.16).

The main effect of the presence of semilocal charges (namely, odd charges in the dual representation) is that the stationary states emerging in the protocols (I) and (II) are macroscopically different [109]. As far as we can see, however, the two macrostates have generally the same entropy per unit length, therefore the excess of entropy is expected to be subextensive. In the examples that we consider, the excess of entropy still grows with ℓ , therefore we can also replace Eq. (5.16) by Eq. (5.20) without affecting the asymptotic behaviour. This important simplification allows us to express the excess of entropy in the dual representation as

$$\lim_{t \rightarrow \infty} \Delta_r S_\alpha(\ell, t) = S_\alpha^{(\tau)}(\ell, \rho_{\text{NESS}}^{(\tau)}) - S_\alpha^{(\tau)}(\ell, \rho_\infty^{(\tau)}) + O(1), \quad (5.24)$$

where

$$\begin{aligned} \rho_{\text{NESS}}^{(\tau)} &= \lim_{B \rightarrow \infty} \lim_{t \rightarrow \infty} \text{tr}_{\overline{B}} [e^{-i\mathbf{H}^{(\tau)}t} (|\downarrow\uparrow\rangle\langle\downarrow\uparrow|)_\tau e^{i\mathbf{H}^{(\tau)}t}] \\ \rho_\infty^{(\tau)} &= \lim_{B \rightarrow \infty} \lim_{t \rightarrow \infty} \text{tr}_{\overline{B}} [e^{-i\mathbf{H}^{(\tau)}t} (|\uparrow\rangle\langle\uparrow|)_\tau e^{i\mathbf{H}^{(\tau)}t}]. \end{aligned} \quad (5.25)$$

In order to gain some intuition about the behaviour of the excess of entropy, we follow the suggestion of Ref. [31] and consider the simplified case in which $\mathbf{H}^{(\tau)}$ commutes with \mathbf{S}^z (and hence $|\uparrow\rangle_\tau$ is an eigenstate). We also assume that $|\uparrow\rangle_\tau$ could be interpreted as the vacuum of stable excitations, whose number is linear in \mathbf{S}^z (this characterises a class of integrable systems). In this situation the nonequilibrium stationary state $\rho_{\text{NESS}}^{(\tau)}$ can be interpreted as a Fermi sea or split Fermi seas [110], which can be described by a conformal field theory (see also Ref. [111]). Since in a conformal field theory the Rényi entropies have a logarithmic dependence on the subsystem's length [112], we can expect the excess of entropy to grow logarithmically in ℓ . Although not a-priori evident (but somehow consistent with the observations of Ref. [31]), the log behaviour survives a symmetry-preserving local unitary transformation of the initial state, which transforms this protocol into a genuine global quench. This will be derived in the next section, in which we will investigate systems that are dual to \mathbb{Z}_2 -symmetric generalised XY models [113].

1. Example: Dual generalised XY model

In this section we work out the excess of entropy in a generalisation of the dual XY model that preserves the noninteracting structure and the existence of a family of semilocal conservation laws. We do not focus only on the dual XY model because, as will be clear later, the dual XY model is rather special and one could read the outcome of the investigation as an indication of a universality that instead is not present. In its most general form the Hamiltonian reads

$$\mathbf{H} = \sum_{\alpha \in \{x,y\}} \sum_{n=1}^{\infty} J_\alpha^{(n)} \sum_{j \in \mathbb{Z}} \mathbf{h}_j^{(\alpha,n)}, \quad (5.26)$$

where the coupling constants $J_\alpha^{(n)}$ are assumed to approach zero in the limit $n \rightarrow \infty$ exponentially fast in n . The operators $\mathbf{h}_j^{(\alpha,n)}$ are Hermitian and local, with support including j and extending over $n+1$ sites; their explicit form in the σ representation is not essential for the discussion but we report it for the sake of completeness:

$$\begin{aligned}
\mathbf{h}_j^{(x,1)} &= \sigma_{j-1}^x \sigma_{j+1}^x &= \tau_j^x \tau_{j+1}^x \\
\mathbf{h}_j^{(y,1)} &= -\sigma_{j-1}^x \sigma_j^z \sigma_{j+1}^x &= \tau_j^y \tau_{j+1}^y \\
\mathbf{h}_j^{(\alpha,n)} &= \sigma_{j-1}^x \sigma_j^\alpha \left\{ \begin{array}{ll} \prod_{l=1}^{\frac{n+1}{2}-\alpha} \sigma_{j+2(l-1)+\alpha}^z \sigma_{j+n-1}^\alpha & n \text{ odd} \\ \prod_{l=1}^{\frac{n}{2}-1} \sigma_{j+2(l-1)+\alpha}^z \sigma_{j+n-1}^{3-\alpha} & n \text{ even} \end{array} \right\} \sigma_{j+n}^x &= \tau_j^\alpha \prod_{l=1}^{n-1} \tau_l^z \left\{ \begin{array}{ll} \tau_{j+n}^\alpha & n \text{ odd} \\ \tau_{j+n}^{3-\alpha} & n \text{ even} \end{array} \right\} & n > 1,
\end{aligned} \tag{5.27}$$

where $\alpha \in \{x \equiv 1, y \equiv 2\}$ and we abused the notation by using the letter or the number depending on what is convenient in each case.

The dual XY model corresponds to $J_\alpha^{(n)} = \delta_{n1} J_\alpha$. More generally, the model with Hamiltonian (5.26) is the dual of a generalised XY model (introduced by Suzuki in Ref. [113]) with $\mathcal{P}_\tau^x[\mathbf{H}] = \mathbf{H}$ and $\mathcal{P}_\tau^z[\mathbf{H}] = \mathbf{H}$. In the explicit examples, we will only consider the effect of having also $J_\alpha^{(2)}$ and $J_\alpha^{(4)}$ different from zero (besides $J_\alpha^{(1)}$). However, since it is not convenient to write the explicit dependence on the coupling constants, we will still treat the dual generalised XY model in (almost) full generality.

When written in the τ representation, the Hamiltonian has the form (3.12) with $\mathbf{a}_\ell^{x,y} = (\prod_{j<\ell} \tau_j^z) \tau_\ell^{x,y}$. As done in Section III B 1, it is convenient to express the matrices $\mathcal{H}_{\ell,n}$ characterising the Hamiltonian in terms of the symbol $\mathcal{H}(p)$ as $\mathcal{H}_{\ell,n} = \int \frac{dk}{2\pi} e^{i(\ell-n)p} \mathcal{H}(p)$, where

$$\mathcal{H}^\dagger(p) = \mathcal{H}(p) = -\mathcal{H}^T(-p). \tag{5.28}$$

The symbol of \mathbf{H} — (5.26) — has the additional property of being a smooth matrix-valued function of p satisfying

$$\mathcal{H}(p + \pi) = \sigma^z \mathcal{H}(p) \sigma^z. \tag{5.29}$$

Under close scrutiny, we realise that this condition represents evenness under \mathcal{P}_τ^x , which is required in order for the Hamiltonian density to remain (strongly) quasilocally also in the σ representation. For the sake of simplicity, we are going to restrict ourselves to reflection symmetric systems ($J_x^{(2n)} = J_y^{(2n)}$), whose symbol has the additional property

$$(J_x^{(2n)} = J_y^{(2n)} \quad \forall n) \quad \Rightarrow \quad \text{tr}[\mathcal{H}(p)] = 0. \tag{5.30}$$

Incidentally, the dispersion relation of the quasiparticle excitations, which can be identified with the positive eigenvalues of the symbol, can be written as

$$\varepsilon(p) = \sqrt{\frac{\text{tr}[\mathcal{H}(p)^2]}{2}} \tag{5.31}$$

As mentioned in Section III B 1, the symbol of the Hamiltonian generates the time evolution of the symbol of the correlation matrix (3.14) and of the operators (in the Heisenberg representation) with a quadratic fermionic representation.

2. Free-fermion techniques for the entropies

Since \mathbf{H} is noninteracting in the dual representation, if the initial state $|\Psi(0)\rangle^{(\tau)}$ is Gaussian, Wick's theorem implies that every expectation value can be written in terms

of the correlation matrix $\Gamma_{t=0}$ capturing the expectation values of quadratic operators. As long as the subsystem is connected, this statement holds true even if both the operators and the correlation matrix are restricted to the subsystem. As a result, the density matrices $\rho_{\tilde{A}}^{(\tau)}(\Psi(t))$ and $\rho_{\tilde{A}}^{(\tau)}(\Psi_\tau(t))$ are Gaussian at any time, that is to say

$$\rho_{\tilde{A}}^{(\tau)}(\Psi) \propto \exp\left(\frac{1}{4} \sum_{m,n=1}^{\ell+1} \vec{a}_m^T \mathcal{W}_{m,n}(\Psi) \vec{a}_n\right) \tag{5.32}$$

with $\vec{a}_m = (a_m^x \ a_m^y)^T$, $a_m^{x,y}$ being the Majorana (Jordan-Wigner) fermions defined in $\tilde{A} = \{1, \dots, \ell + 1\}$:

$$a_m^{x,y} = (\tau^z)^{\otimes(m-1)} \otimes \tau^{x,y} \otimes \mathbb{I}^{(\ell+1-m)}. \tag{5.33}$$

The Gaussian structure is very useful also to compute the Rényi entropies, which can be expressed in terms of the correlation matrix as [114, 115]

$$S_\alpha^{(\tau)}(\ell+1; \Psi) = \frac{\log \det \left[\left(\frac{I + \Gamma_{\tilde{A}}}{2} \right)^\alpha + \left(\frac{I - \Gamma_{\tilde{A}}}{2} \right)^\alpha \right]}{2(1 - \alpha)}, \tag{5.34}$$

where $\Gamma_{\tilde{A}}$ is the correlation matrix of $|\Psi\rangle$ restricted to \tilde{A} . Therefore, using the approximation (5.24), as $\ell \rightarrow \infty$, the unbounded part of the excess of entropy can be expressed in terms of the correlation matrices of $\rho_{\text{NESS}}^{(\tau)}$ and $\rho_\infty^{(\tau)}$. They are reported in the following, together with the correlation matrix of the initial state, $\Gamma_0(p)$, to which they are related.

Correlation matrix of $|\Psi(0)\rangle^{(\tau)}$. In the numerical investigations we have focussed on the initial state $|\uparrow\rangle_\tau$, which is one of the two representations of $|\uparrow\rangle_\sigma$ with the hidden symmetry broken — see Section IV B. Its correlations matrix has the symbol $\Gamma_0(p) = \sigma^y$.

Correlation matrix of $\rho_\infty^{(\tau)}$. We have already reviewed how to obtain the correlation matrix of $\rho_\infty^{(\tau)}$ after a global quench — see Section III B 1. In our specific case with reflection symmetry (and one-site shift invariance), the symbol of the correlation matrix of $\rho_\infty^{(\tau)}$ can be written as

$$\Gamma_\infty(p) = \frac{\text{tr}[\Gamma_0(p) \mathcal{H}(p)] \mathcal{H}(p)}{2\varepsilon^2(p)}, \tag{5.35}$$

where we used that the only matrix commuting with $\mathcal{H}(p)$ different from the identity (which is excluded because the initial state is reflection symmetric and hence $\text{tr}[\Gamma_0(p)] = 0$ — see also (5.30)) is proportional to $\mathcal{H}(p)$. From $\Gamma_\infty(p)$ we can then obtain $\Gamma_{\tilde{A};\infty}$ by Fourier transforming $\Gamma_\infty(p)$ and restricting the indices to the subsystem \tilde{A} .

Correlation matrix of $\rho_{\text{NESS}}^{(\tau)}$. We can not apply the same technique for the correlation matrix of $\rho_{\text{NESS}}^{(\tau)}$: the time evolved domain wall $|\Psi_r(t)\rangle^{(\tau)}$ is not translationally invariant, so the symbol is not even defined. This problem can be exactly solved by redefining the symbol so as to incorporate the inhomogeneities of the Gaussian state [103]. As generally expected in bipartitioning protocols in integrable systems, however, $|\Psi_r(t)\rangle^{(\tau)}$ locally relaxes along rays at fixed $\zeta = (x-r)/t$ [103, 116]. It is then easier (and asymptotically correct in the limit $t \rightarrow \infty$) to associate a translationally invariant macrostate with each ray. Being translationally invariant, the macrostate is characterised by a symbol as defined before. Finally, a generalised hydrodynamic equation provides the connection between the macrostates at different rays [117, 118], and one can easily extract the symbol associated with the ray $\zeta = 0$, which corresponds to the infinite time limit that we indicated with $\rho_{\text{NESS}}^{(\tau)}$. Specifically, Appendix E 2 shows

$$\Gamma_{\text{NESS}}(p) = -\text{sgn}[v(p)] \frac{\text{tr}[\Gamma_0(p)\mathcal{H}(p)]\mathbb{I}}{2\varepsilon(p)}, \quad (5.36)$$

where $v(p) = \varepsilon'(p)$ is the velocity of the quasiparticle excitations.

We now have all the ingredients to work out Eq. (5.24) and to obtain analytical asymptotic results for large ℓ . In the next paragraph we show how, for $\alpha = 2, 3, \dots$, one can also obtain exact numerical results.

Exact numerical evaluation of the entropies. In order to evaluate the Rényi entropies exactly, we use Eq. (5.16), which is written in terms of the averaged density matrix $\bar{\rho}_A^{(\tau)}(\Psi)$ — (5.15). The method is explained in details in Appendix E 4 and is based on the realisation that each term of the sum in Eq. (5.15) is Gaussian. The Rényi entropies are expressed as traces of powers of $\bar{\rho}_A^{(\tau)}(\Psi)$, which are linear combinations of Gaussians. The trace of a product of Gaussians can be expressed as the square root of a determinant of a $2(\ell+1) \times 2(\ell+1)$ matrix [119, 120] and can therefore be computed efficiently. There is only a subtlety related to the sign ambiguity of the square root. For a product of two Gaussians (which appears in the expression for the second Rényi entropy) the sign problem is resolved by using the property that the trace of a product of two positive semidefinite Hermitian operators is non-negative [121]. For higher-order




FIG. 12. Time evolution of the excess of the second Rényi entropy, generated in the XY model (3.9) and in the dual XY model (3.17). For large times the excess in the XY model reaches a zero value, while in the dual XY model the its asymptotic value is nonzero. It is in agreement with the result obtained via the method used also in Fig. 13 (here the latter results are presented as dashed lines). The spin flip occurs in the middle of the subsystem A ($r = 5, |A| = 10$).




FIG. 13. Exact results for the excess of the second Rényi entropy after the quench, for different values of model's parameters. The excess grows logarithmically with the subsystem size (the x -axis is in the log scale). The analytical results for the coefficients b_α (i.e., the slopes), given by Eq. (5.39), are in agreement with the numerical data and indicated in the plot.

Rényi entropies the sign is fixed in a less straightforward way, as discussed in Ref. [119].

3. Results

Both with and without the local spin flip in the initial state, the quench is global and the Rényi entropies grow

linearly in time until they saturate to an extensive value. In noninteracting models this was established in Refs. [63, 103, 122]; it was later generalised to interacting integrable models in Refs. [123–125]. Less is known in regards to the subleading behaviour.

For the quench from the initial state $|\uparrow\rangle_\tau$ we find

$$\lim_{t \rightarrow \infty} S_\alpha^{(\tau)}(\ell + 1; \Psi(t)) = a_\alpha \ell + \mathcal{O}(1), \quad (5.37)$$

with positive constants a_α , whose values can be determined analytically using the integral representation for the entropies [114] and the Szegő-Widom theorem [126, 127]. Their values are reported in Appendix E 3.

For the quench from the domain-wall state (the dual representation of the state $\sigma_r^x |\uparrow\rangle_\sigma$) we instead find a subleading term that is logarithmic in ℓ :

$$\lim_{t \rightarrow \infty} S_\alpha^{(\tau)}(\ell; \Psi_r(t)) = a_\alpha \ell + b_\alpha \log \ell + \mathcal{O}(1), \quad (5.38)$$

with positive constants b_α . This result is obtained using the asymptotic formula for the determinant of a block Toeplitz matrix with piecewise-continuous symbols, conjectured and checked numerically in Refs. [128, 129]. The values of b_α are related to the discontinuities of the symbol (5.36), which coincide with the zeroes of the velocity $v(p)$ of the quasiparticle excitations. For instance, for $\alpha = 2$ we have

$$b_2 = \frac{2}{\pi^2} \sum_{p \in (-\pi, \pi]: v(p)=0} \text{Arg}^2 \left(1 + i \frac{\text{tr}[\Gamma(p; 0)\mathcal{H}(p)]}{2\varepsilon(p)} \right), \quad (5.39)$$

where Arg denotes (the principal value of) the argument, and the sum is over the zeroes of $v(p)$. For general α the expression can be found in Appendix E 3.

In the symmetric generalised XY model there are always zeroes of $v(p)$ at the momenta $p = -\pi/2, 0, \pi/2, \pi$, and additional ones can appear depending on the coupling constants $J_\alpha^{(n)}$. In the special case of the XY model these four zeros are the only ones and we find

$$b_\alpha^{(\text{XY})} = \frac{1}{6} \left(1 + \frac{1}{\alpha} \right). \quad (5.40)$$

For more general coupling constants $J_\alpha^{(n)}$ this expression is only a lower bound, i.e., $b_\alpha \geq b_\alpha^{(\text{XY})}$.

From Eqs. (5.38) and (5.39) it follows that the excess (5.22) of Rényi entropies grows logarithmically with ℓ :

$$\lim_{t \rightarrow \infty} \Delta_r S_\alpha(\ell, t) = b_\alpha \log \ell + \mathcal{O}(1). \quad (5.41)$$

This prediction is compared against exact numerical data for $\alpha = 2$: Fig. 12 shows the time evolution of $\Delta_r S_\alpha(\ell, t)$, while in Fig. 13 its dependence on ℓ is presented.

Finally, the von Neumann entropy is obtained by taking the limit $\alpha \rightarrow 1^+$; in the dual XY model it is given by $b_1^{(\text{XY})} = 1/3$.

In the next section we conclude our preliminary study of the signatures of semilocal order after global quenches by showing that the order melts down over large time scales even when the initial state is not symmetric.

C. Melting of the order

Just as a continuous phase transition is characterised by the behaviour of the system when the critical point is approached, so is the symmetry protected topological order when the symmetry is only approximate. We are familiar with an example of this: in Fig. 5 we showed that, after a quench from the tilted state $|\nearrow_\theta\rangle$, the time scale on which local observables relax towards the predictions of the (generalised) Gibbs ensemble grows as θ approaches 0. In the limit $\theta \rightarrow 0$ the relaxation time diverges: the canonical form of a quasilocal ensemble, valid for any finite θ , yields a wrong prediction when $\theta = 0$. At the symmetric point, instead, the correct prediction is given by the \mathbb{Z}_2 -semilocal ensemble, and the corresponding relaxation time is again finite. This implies that the limits $t \rightarrow \infty$ and $\theta \rightarrow 0$ do not commute (note that the existence of noncommuting limits is one of the hallmarks of spontaneous symmetry breaking [57] as well as of prethermalisation/prerelaxation behaviours after global quenches [45–48]).

Signatures of semilocal order thus exist also in states that weakly break the symmetry with which the order is associated. In particular, we envisage the existence of a scaling limit with $t \rightarrow \infty$ and $\theta \rightarrow 0$, in which local observables exhibit some form of prerelaxation. This scaling limit is discussed in Section V C 1.

Symmetric initial states exhibiting semilocal order can be regarded as ground states of prequench Hamiltonians \mathbf{H}_0 (see Section IV B). Since string order does not survive nonzero temperatures [12], the prerelaxation behaviour mentioned above is expected also for sufficiently low, but nonzero temperatures. We will see in Section V C 2 that the melting of the order arising at small nonzero temperatures has essentially the same origin as the prerelaxation occurring for small symmetry-breaking unitary perturbations of the initial state.

1. Slow relaxation after weak symmetry breaking

Being interested in the time scales on which semilocal order disappears after a weak symmetry breaking, we consider an initial state prepared close to the symmetric point, i.e., $|\nearrow_\theta\rangle$ for small $\theta > 0$.

Since $\langle \nearrow_\theta | \sigma_\ell^y | \nearrow_\theta \rangle = 0$ and $\langle \nearrow_\theta | \sigma_\ell^x | \nearrow_\theta \rangle = \sin \theta \rightarrow 0$ as $\theta \rightarrow 0$, the local operators to consider in order to explore the melting of the order consist solely of products of σ^z . This includes, in particular, the strings $\prod_{\ell=0}^{n-1} \sigma_{j+\ell}^z$,

which, when evaluated in a particular state, can be regarded as truncations of a semilocal order parameter, corresponding to the limit $n \rightarrow \infty$. They satisfy

$$\langle \not\rightarrow_\theta | \prod_{\ell=0}^{n-1} \sigma_{j+\ell}^z | \not\rightarrow_\theta \rangle = \cos^n \theta. \quad (5.42)$$

The inverse duality transformation (4.7) maps operators composed solely of σ_j^z into operators consisting of an even number of τ_j^z . Specifically, the strings appearing in Eq. (5.42) are mapped into $\tau_j^z \tau_{j+n}^z$, which are the 4-fermion operators $-\mathbf{a}_j^x \mathbf{a}_j^y \mathbf{a}_{j+n}^x \mathbf{a}_{j+n}^y$. To describe the expectation values of such operators we introduce an effective density matrix ρ_{eff} that is only asked to capture the expectation value of 4-fermion operators \mathbf{O} :

$$\langle \not\rightarrow_\theta | \mathbf{O}^{(\sigma)}(t) | \not\rightarrow_\theta \rangle = \text{tr}[\rho_{\text{eff}}(t) \mathbf{O}], \quad (5.43)$$

where $\mathbf{O}^{(\sigma)}(t) = e^{i\mathbf{H}^{(\sigma)}t} \mathbf{O} e^{-i\mathbf{H}^{(\sigma)}t}$ is the time evolution of \mathbf{O} in the Heisenberg picture, and $\mathbf{H}^{(\sigma)}$ is the dual XY model's Hamiltonian, given in Eq. (3.17).

For each t we can expand $\mathbf{O}^{(\sigma)}(t)$ in the operator basis $\{\mathbf{e}_\alpha\}$ as $\mathbf{O}^{(\sigma)}(t) = \sum_\alpha (\text{tr}[\mathbf{O}(t) \mathbf{e}_\alpha] / \text{tr} \mathbf{I}) \mathbf{e}_\alpha$, where \mathbf{e}_α can be chosen, for example, to be strings of Pauli matrices of various lengths. The time evolution preserves the number of fermions, so the expansion of $\mathbf{O}^{(\sigma)}(t)$ retains only the basis elements \mathbf{e}_α corresponding to 4-fermion operators. Moreover, since in Eq. (5.43) we project onto the tilted initial state, we can keep only the basis elements

for which $\langle \not\rightarrow_\theta | \mathbf{e}_\alpha | \not\rightarrow_\theta \rangle \neq 0$. The latter again correspond to strings of σ_j^z , whence we obtain

$$\begin{aligned} \langle \not\rightarrow_\theta | \mathbf{O}^{(\sigma)}(t) | \not\rightarrow_\theta \rangle &= \sum_{j \in \mathbb{Z}} \sum_{n \geq 1} \langle \not\rightarrow_\theta | \prod_{\ell=0}^{n-1} \sigma_{j+\ell}^z | \not\rightarrow_\theta \rangle \\ &\times \frac{\text{tr}[\mathbf{O}^{(\tau)}(t) \tau_j^z \tau_{j+n}^z]}{\text{tr} \mathbf{I}}. \end{aligned} \quad (5.44)$$

The effective density matrix for small θ is thus [130]

$$\rho_{\text{eff}}(t) \sim e^{-i\mathbf{H}^{(\tau)}t} \left[\frac{1}{\text{tr} \mathbf{I}} \sum_{j \in \mathbb{Z}} \sum_{n \geq 1} e^{-n\theta^2/2} \tau_j^z \tau_{j+n}^z \right] e^{i\mathbf{H}^{(\tau)}t}, \quad (5.45)$$

where we used Eq. (5.42) with its right-hand side rewritten, for small θ , as $\cos^n \theta = e^{-n\theta^2/2} [1 + \mathcal{O}(\theta^4)]$.

The right-hand side of Eq. (5.45) is in τ representation: $\mathbf{H}^{(\tau)}$ corresponds to an XY Hamiltonian (3.9), so the standard free-fermionic techniques apply for calculation of the expectation values — see Appendix F. For example, the truncated string order parameter $Z_n = \langle \prod_{\ell=0}^{n-1} \sigma_{j+\ell}^z \rangle$, which, in the fermionic language, corresponds to $-\langle \mathbf{a}_1^x \mathbf{a}_1^y \mathbf{a}_{n+1}^x \mathbf{a}_{n+1}^y \rangle$, can be efficiently computed by invoking Wick's theorem: the full expression derived in the scaling limit $t \rightarrow \infty$, $\theta \rightarrow 0$, at fixed finite $0 < \theta^2 t < \infty$ for any $n \geq 1$ is reported in Appendix F. It can be simplified in the limit $n \rightarrow \infty$, where it becomes

$$Z_\infty = \frac{1}{2} \int \frac{d^2 k}{(2\pi)^2} (e^{-|v(k_1) - v(k_2)| \frac{\theta^2 t}{2}} + e^{-|v(k_1) + v(k_2)| \frac{\theta^2 t}{2}}) \frac{(J_x + J_y)^4 \cos^2 k_1 \cos^2 k_2}{[J_x^2 + J_y^2 + 2J_x J_y \cos(2k_1)][J_x^2 + J_y^2 + 2J_x J_y \cos(2k_2)]}. \quad (5.46)$$

From this expression one can read off the time scale ξ on which the order melts: $\xi \sim \theta^{-2}$. In fact this time scale is inherent to the behaviour of the effective density matrix ρ_{eff} (see Appendix F) and is therefore exhibited also by the time evolution of Z_n for finite n . Fig. 14 corroborates it numerically for $n = 1$, i.e., in the case of $Z_1 = \langle \sigma_\ell^z \rangle$.

We warn the reader that setting $\theta^2 t = 0$ in Eq. (5.46) does not correspond to the expectation value of the string order parameter at the initial time, it is rather the starting value from which the preresolution behaviour ensues. This is because the derivation is based on asymptotics for large $t > 0$ and small $\theta > 0$. Despite this, setting $\theta^2 t = 0$ in Eq. (5.46) correctly reproduces the large time value of the string order parameter for $\theta = 0$.

2. Melting at low finite temperatures

The simplest example of a finite-temperature initial state reads

$$\rho(0) = \frac{e^{\beta \sum_\ell \sigma_\ell^z}}{\text{tr}[e^{\beta \sum_\ell \sigma_\ell^z}]}. \quad (5.47)$$

In the limit $\beta \rightarrow \infty$ this thermal state reproduces the symmetric state with semilocal order, as considered in Section IV B. For any finite β $\rho(0)$ contains only terms composed of products of σ_j^z . Hence, focusing again only on 4-fermion operators, the only relevant operator with a nonzero expectation value is again $\mathbf{a}_\ell^x \mathbf{a}_\ell^y \mathbf{a}_{\ell+n}^x \mathbf{a}_{\ell+n}^y$, i.e., a fermionic representation of a string of Pauli matrices σ_j^z . This time its expectation value reads

$$-\langle \mathbf{a}_\ell^x \mathbf{a}_\ell^y \mathbf{a}_{\ell+n}^x \mathbf{a}_{\ell+n}^y \rangle = \tanh^n \beta. \quad (5.48)$$




FIG. 14. Numerically computed (iTEBD) expectation value $\langle \sigma_\ell^z \rangle$ plotted versus $\theta^2 t$ is consistent with a scaling prediction derived in Appendix F. In the limit $\theta \rightarrow 0$ the scaling prediction yields $\langle \sigma_\ell^z \rangle \approx 0.5626$, which is also the GGE prediction for the expectation value of $\langle \sigma_\ell^z \sigma_{\ell+1}^z \rangle$ in the XY model prepared in the initial state $|\uparrow\rangle$ — see Fig. 7. The prediction coming from a GGE with the standard pseudolocal charges of the dual XY model is instead zero. The inset is in logarithmic scale.

For large β and small θ , this is equivalent to setting the rotation angle in the tilted initial state $|\nearrow_\theta\rangle$ to be

$$\theta_\beta \approx 2e^{-\beta}. \quad (5.49)$$

Having established a connection with the tilted initial state, we can retrace the steps in the previous section and observe that the nontrivial scaling limit is now $t \rightarrow \infty$, $\beta \rightarrow \infty$, at fixed finite $4e^{-2\beta}t$. The time scale ξ on which the order melts at a low, but finite temperature, after the system has been prepared in the state $\rho(0)$, is thus $\xi \sim \theta_\beta^{-2} = (e^\beta/2)^2$.

Finally, we observe a similarity with the findings of Ref. [47], in which an analogous scaling behaviour was observed at low temperature in a phase in which a symmetry is spontaneously broken at zero temperature. The reader can understand this connection as a manifestation of the duality between the quasilocal theory and the even \mathbb{Z}_2 -semilocal theory, the latter being characterised by a spontaneous symmetry breaking at zero temperature — cf. (4.7).

VI. DISCUSSION

We have shown that symmetry-protected topological phases of matter can emerge also at infinite times after

quenches of global Hamiltonian parameters in isolated many-body quantum systems. We have traced this phenomenon back to the existence of conserved operators (semilocal charges) that do not belong to the natural theory in which local observables are represented by local operators and their quasilocal completion. From that perspective, the topological character of the system is manifest: different theories of local observables are associated with different concepts of locality for the operators; being outside the quasilocal theory, semilocal charges can keep memory of information that is not contained in the bulk of the system.

We aimed at reconciling the observation that semilocal charges affect the stationary values of local operators with the established belief that those values are solely determined by the local conserved quantities. This has led us to introduce statistical ensembles sensitive to string (semilocal) order, which we called semilocal (generalised) Gibbs ensembles.

Finally, we have reported signatures of semilocal order in the entropies of subsystems and in the time evolution of systems that are not exactly symmetric. In particular, we have found that a perturbation with a finite support affects the entanglement entropies of subsystems with an exceptional leading correction scaling as the logarithm of the subsystem’s length.

Open problems

- The description we have proposed relies on translational invariance. If we break it, the state can be globally non-symmetric still exhibiting the local symmetries characterising semilocal order, in the sense that the two-point functions of semilocal operators could approach nonzero values at intermediate distances. In addition, since the expectation values of semilocal charges are also affected by local inhomogeneities, the latter could look relevant despite the Lieb-Robinson bounds ruling out their importance at space-like distances. A theory that could capture time evolution of inhomogeneous states goes beyond the purposes of this work and will be addressed in separate investigations.
- We define the G -semilocal (generalised) Gibbs ensemble in an extended space that includes operators that do not represent local observables. Having realised the existence of an ambiguity in what should be called “local observable” in the extended space, we propose to project back the ensemble onto a theory of local observables. Taking into account that similar situations seem to occur in quantum field theories, where fields that are semilocal with respect to each other are part of the theory, we wonder whether such a projection is really required. Perhaps a more rigorous treatment along the lines of the algebraic formulation of a local quantum field

theory [131] could clarify this issue. Incidentally, we wonder whether the finding of Ref. [132] could be connected with the picture presented here.

- We show that, in the presence of semilocal order, the excess of entropy triggered by a local perturbation of the initial state develops a logarithmic dependence on the subsystem’s length. We expect that even the entanglement properties of the symmetric translationally invariant system (without any perturbation) should exhibit exceptional properties, but, at the moment, this is still an open question.
- Despite our description applying to more general situations, the explicit examples that we consider are not interacting. This choice has a twofold motivation: on the one hand, we aim at providing the cleanest examples exhibiting the exotic phenomenology of semilocal order after global quenches; on the other hand, interactions could introduce complications that need clues from simpler models in order to be solved. Whether the interplay between interactions and semilocality could enable additional phenomena is an open problem.
- We discuss semilocal order both in generic and in integrable systems. Extending the theory to non-abelian integrable structures remains an intriguing development, in which semilocal dynamical symmetries could play an important role [133].

ACKNOWLEDGMENTS

We thank Kemal Bidzhiev, Saverio Bocini, Mario Colura, Fabian Essler, Ivan Kukuljan, Leonardo Mazza, Spyros Sotiriadis, and Gabor Takacs for useful discussions.

This work was supported by the European Research Council under the Starting Grant No. 805252 LoCo-Macro.

Appendix A: Convergence of the iTEBD data

Here we estimate the times t up to which the results of the iTEBD time evolution in Fig. 5 are virtually indistinguishable from those of the exact calculation. By “virtually indistinguishable” we mean, that the difference between the numerically computed and the exact expectation value $\langle \sigma_\ell^z \rangle$ at any particular time should be smaller than some accuracy, which we choose to be 0.01 in Fig. 5.

We compare the data from calculations with maximum bond dimensions that are increasing exponentially up to the largest one, $M_{\max} = 1000$ (bond dimension is expected to increase exponentially with t). The results of the calculation with $M_{\max} = 1000$ are assumed to match




FIG. 15. Differences between the results of the calculation with $M_{\max} = 1000$ and smaller maximum bond dimensions. Red circles denote when the differences become noticeable: the corresponding times grow linearly.

the exact time evolution longer than those at smaller M_{\max} , which is why we take them as a reference. Fig. 15 shows the differences $\langle \sigma_\ell^z \rangle_{1000} - \langle \sigma_\ell^z \rangle_{M_{\max}}$, where the subscript denotes the maximum bond dimension in the calculation. From the times at which these differences become virtually discernible (i.e., larger than 0.01), we estimate that the calculations with $M_{\max} = 1000$ can be trusted up to times $t \sim 4.5$.

Appendix B: $\mathbb{Z}_2 \times \mathbb{Z}_2$ duality

We report here what could be considered as a double application of the rotated Kramers-Wannier duality transformation. We denote it by $\mathcal{D}_{\mathbb{Z}_2 \times \mathbb{Z}_2}$ and it is an algebra homomorphism mapping the algebra of observables generated by $\{\tau, \mathbf{T}_{\tau; -2,0}^x, \mathbf{T}_{\tau; -2,1}^x\}$ into the one generated by $\{\sigma, \mathbf{T}_{\sigma; 2,0}^z, \mathbf{T}_{\sigma; 2,1}^z\}$. Here again σ_j^α and τ_j^α act like Pauli matrices on site j and like the identity elsewhere. The additional operators $\mathbf{T}_{\sigma; 2s,y}^\alpha$ are instead defined by the following conditions

$$\begin{aligned} [\mathbf{T}_{\sigma; 2s,y}^\alpha]^2 &= \mathbf{I}, & \mathbf{T}_{\sigma; 2s,1}^\alpha \mathbf{T}_{\sigma; 2s,0}^\alpha &= \mathbf{T}_{\sigma; 2s,0}^\alpha \mathbf{T}_{\sigma; 2s,1}^\alpha = \mathbf{T}_{\sigma; s}^\alpha, \\ \mathbf{T}_{\sigma; 2s,y}^\alpha \mathbf{O} \mathbf{T}_{\sigma; 2s,y}^\alpha &= \lim_{n \rightarrow \infty} \left[\prod_{\ell=0}^n \sigma_{s(2\ell+y)}^\alpha \right] \mathbf{O} \left[\prod_{\ell=0}^n \sigma_{s(2\ell+y)}^\alpha \right] \end{aligned} \quad (\text{B1})$$

for all local operators \mathbf{O} , extended then by linearity as in the case of $\mathcal{D}_{\mathbb{Z}_2}$, in such a way that $\mathbf{O}_1 + \mathbf{O}_2 \mathbf{T}_{\sigma; 2s,0}^\alpha + \mathbf{O}_3 \mathbf{T}_{\sigma; 2s,1}^\alpha + \mathbf{O}_4 \mathbf{T}_{\sigma; s}^\alpha$ is transformed by the adjoint action of $\mathbf{T}_{\sigma; 2s,y}^\alpha$ into an analogous operator in which the local operators \mathbf{O}_j are replaced by $\mathbf{T}_{\sigma; 2s,y}^\alpha \mathbf{O}_j \mathbf{T}_{\sigma; 2s,y}^\alpha$. Explicitly, the duality transformation $\mathcal{D}_{\mathbb{Z}_2 \times \mathbb{Z}_2}$ reads

$$\begin{aligned} \mathbf{\Pi}_{\tau, -2}^x(j) &= \sigma_j^x, \\ \tau_j^y &= \sigma_{j-2}^x \sigma_j^y \mathbf{\Pi}_{\sigma, +2}^z(j+2), \\ \tau_j^z &= \mathbf{\Pi}_{\sigma, +2}^z(j), \end{aligned} \quad (\text{B2})$$

where $\mathbf{\Pi}_{\tau,-2}^x(j)$ and $\mathbf{\Pi}_{\sigma,+2}^z(j)$ are defined as $\mathbf{\Pi}_{\sigma,s}^\alpha(j)$ in the sublattice of sites with the same parity of j . This is obtained using $\mathbf{T}_{\sigma;2s,j \bmod 2}^\alpha$ instead of $\mathbf{T}_{\sigma;s}^\alpha$. In the bulk the transformation can be expressed in the more standard form

$$\tau_j^x = \sigma_{j-2}^x \sigma_j^x \quad \tau_j^z \tau_{j+2}^z = \sigma_j^z, \quad (\text{B3})$$

which corresponds to two independent rotated Kramers-Wannier dualities on the sublattices consisting of the sites labelled by numbers with the same parity.

Finally, the inverse duality $\mathcal{D}_{\mathbb{Z}_2 \times \mathbb{Z}_2}^{-1}$ reads

$$\begin{aligned} \sigma_j^x &= \mathbf{\Pi}_{\tau,-2}^x(j), \\ \sigma_j^y &= \mathbf{\Pi}_{\tau,-2}^x(j-2) \tau_j^y \tau_{j+2}^z, \\ \mathbf{\Pi}_{\sigma,+2}^z(j) &= \tau_j^z. \end{aligned} \quad (\text{B4})$$

Appendix C: Zero-temperature phases of the dual XY model

Since the (rotated) Kramers-Wannier transformation maps the Hamiltonian (3.17) of the dual XY model into the Hamiltonian (3.9) of the XY model, we can immediately infer that the model is noncritical for $|J_x| \neq |J_y|$ and critical for $|J_x| = |J_y|$.

For $|J_y| < |J_x|$ it should thus be possible to connect smoothly the ground state(s) of the model to the one(s) at the classical point $J_y = 0$, above which there is a finite energy gap from the first excited state(s) in the thermodynamic limit. It is easy to see that at the classical point the ground states exhibit magnetic order with translational invariance under a shift by one or several sites in the bulk, the simplest being the ferromagnetic ground state $\otimes_j |\leftarrow\rangle_j$ for $J_x < 0$ (and the one related by the spin flip symmetry). A symmetry breaking field [57] can then select a ground state with a particular pattern of the magnetic order and we are in the standard Landau phase. Note that, besides symmetry breaking, there could be other ambiguities in the zero temperature limit. For example, for $J_y > 0$, imposing periodic boundary conditions on a chain with an odd number of sites frustrates the system, with consequences on the properties of the state that survive the thermodynamic limit [134–136].

It is simple to show that the Hamiltonian of the dual XY model with $|J_y| < |J_x|$ can be mapped through a symmetry-preserving local unitary transformation, into an operator, with strongly quasilocal densities, that commutes with

$$\mathbf{H}(J_x, 0) = J_x \sum_{\ell} \sigma_{\ell-1}^x \sigma_{\ell+1}^x \quad (\text{C1})$$

and shares the same ground states of $\mathbf{H}(J_x, 0)$. The latter states completely break the $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetry of the Hamiltonian (see Section IV C 4), belonging therefore to a standard ordered phase à la Landau. Specifically and just for the sake of example, the symmetry

breaking states of $\mathbf{H}(J_x, 0)$ with clustering properties are mapped into the analogues with $J_y \neq 0$ (and $|J_y| < |J_x|$) by the transformation $e^{-i\mathbf{W}}$, where \mathbf{W} is a dual generalised XY model — Section V B 1 and Appendix E 2 — characterised by the symbol

$$W(p) = -\frac{1}{2} \arctan \frac{2J_y \tan p}{J_x + J_y + (J_x - J_y) \tan^2 p} \sigma^z. \quad (\text{C2})$$

Analogously, there is a transformation mapping the Hamiltonian of the dual XY model with $|J_y| < |J_x|$ into an operator, with strongly quasilocal densities, that commutes with

$$\mathbf{H}(0, J_y) = -J_y \sum_{\ell} \sigma_{\ell-1}^x \sigma_{\ell}^z \sigma_{\ell+1}^x \quad (\text{C3})$$

and shares the same ground state of $\mathbf{H}(0, J_y)$. As discussed in Ref. [86], the latter Hamiltonian has the ground state in a nontrivial $\mathbb{Z}_2 \times \mathbb{Z}_2$ protected topological phase, which therefore describes the entire parameter space $|J_y| > |J_x|$.

Appendix D: Tilted initial state

1. Tilted state $|\nearrow_{\theta}\rangle_{\sigma}$

The mapping of the tilted initial state (3.10) is perhaps most conveniently performed on a finite lattice of size L . To this end we report the finite-system inverse duality transformation

$$\begin{aligned} \sigma_j^x &= \mathbf{\Pi}_{\tau,-}^x(j), \\ \sigma_j^y &= \begin{cases} \tau_1^x \tau_2^z & j = 1 \\ \mathbf{\Pi}_{\tau,-}^x(j-1) \tau_j^y \tau_{j+1}^z & 1 < j < L \\ -\tau_L^z & j = L, \end{cases} \\ \sigma_j^z &= \begin{cases} \tau_j^z \tau_{j+1}^z & 1 \leq j < L \\ \mathbf{\Pi}_{\tau}^x \tau_1^z \tau_L^z & j = L, \end{cases} \end{aligned} \quad (\text{D1})$$

where

$$\mathbf{\Pi}_{\tau,-}^x(j) = -\tau_1^y \prod_{\ell=2}^j \tau_{\ell}^x, \quad \mathbf{\Pi}_{\tau}^x = \prod_{\ell=1}^L \tau_{\ell}^x. \quad (\text{D2})$$

The state

$$|\nearrow_{\theta}\rangle_{\sigma} \equiv \bigotimes_{j=1}^L \left(\cos \frac{\theta}{2} \right), \quad (\text{D3})$$

is an eigenstate of all $\eta_j^{(\sigma)}(\theta) = \cos \theta \sigma_j^z + \sin \theta \sigma_j^x$ for $j = 1, \dots, L$, with eigenvalue 1. Applying the inverse duality transformation on a finite system (D1), we have

$$\eta_j^{(\tau)}(\theta) = \begin{cases} \cos \theta \tau_j^z \tau_{j+1}^z + \sin \theta \mathbf{\Pi}_{\tau,-}^x(j) & 1 \leq j < L \\ \cos \theta \mathbf{\Pi}_{\tau}^x \tau_1^z \tau_L^z - i \sin \theta \mathbf{\Pi}_{\tau}^x \tau_1^z & j = L. \end{cases} \quad (\text{D4})$$

In the basis of all τ_j^z (their eigenvalues being denoted by $s_j \in \{-1, 1\}$) the tilted state reads

$$|\nearrow_\theta\rangle_\sigma = \sum_{\underline{s} \in \{-1, 1\}^{\times L}} c(\underline{s}) |\underline{s}\rangle_\tau. \quad (\text{D5})$$

Applying the operator (D4) we now obtain relations

$$\begin{aligned} \frac{c(-s_1, \dots, -s_j, s_{j+1}, \dots, s_L)}{c(\underline{s})} &= \\ &= \begin{cases} \frac{1-s_j s_{j+1} \cos \theta}{i s_1 \sin \theta} & 1 \leq j < L \\ s_1 s_L e^{-i\theta s_L} & j = L, \end{cases} \end{aligned} \quad (\text{D6})$$

which are solved by

$$c(\underline{s}) = \frac{1}{\sqrt{Z}} e^{i\frac{\pi}{4}(s_1 - s_L) + i\frac{\theta}{2}s_L} e^{-\frac{\beta}{2}E(\underline{s})}. \quad (\text{D7})$$

Here $E(\underline{s}) = -\sum_{j=1}^{L-1} s_j s_{j+1}$, $\beta = -\log \tan(\theta/2)$, and

$$Z = \sum_{\underline{s} \in \{-1, 1\}^{\times L}} e^{-\beta E(\underline{s})}. \quad (\text{D8})$$

We recognize that the squared absolute value $|c(\underline{s})|^2$ yields the canonical ensemble probability distribution for the one-dimensional classical Ising model with free boundary conditions: β plays the role of an effective inverse temperature, while Z is the corresponding partition function. In this regard a nonzero angle θ corresponds to a nonzero temperature.

Finally, the partition function (D8) can be evaluated using the transfer matrix formalism (see, e.g., Ref. [137]). We define transfer matrix

$$T = \frac{1 + \sigma^x}{2} + \frac{1 - \sigma^x}{2} \cos \theta \quad (\text{D9})$$

with elements $T_{s,s'} = (1/2)e^{\beta s s'} \sin \theta$, $s, s' \in \{-1, 1\}$, and use it to rewrite the partition function:

$$Z = \left(\frac{2}{\sin \theta} \right)^{L-1} \sum_{s_1, s_L \in \{-1, 1\}} [T^{L-1}]_{s_1, s_L}. \quad (\text{D10})$$

The powers of the transfer matrix are readily obtained, i.e., $T^n = (1 + \sigma^x)/2 + [(1 - \sigma^x)/2] \cos^n \theta$, whence

$$Z = 2 \left(\frac{2}{\sin \theta} \right)^{L-1}. \quad (\text{D11})$$

2. Decay of correlations in the state $|\nearrow_\theta\rangle_\sigma$

The transfer matrix (D9) enables an efficient method for calculation of local operators that act in the bulk of the spin chain, i.e., far from the boundaries, which will eventually be sent to infinity as the thermodynamic limit is taken. The expectation value of a local operator \mathbf{O} acting on the sites $j, \dots, j+r-1$, where $r \geq 1$ is its range, reads

$$\langle \mathbf{O} \rangle = \frac{1}{Z} \sum_{\underline{s}, \underline{s}'} \langle \underline{s} | \mathbf{O} | \underline{s}' \rangle e^{\frac{\beta}{2}[E(\underline{s}) + E(\underline{s}')]}. \quad (\text{D12})$$

We have used $s_1 = s'_1$ and $s_L = s'_L$ to cancel the phases coming from the coefficients (D7). This is possible since spins in positions 1 and L are not affected by the action of \mathbf{O} :

$$\begin{aligned} \langle \underline{s} | \mathbf{O} | \underline{s}' \rangle &= \prod_{\ell=1}^{j-1} \delta_{s_\ell, s'_\ell} \prod_{m=j+r}^L \delta_{s_m, s'_m} \\ &\times \langle s_j, \dots, s_{j+r-1} | \mathbf{O} | s'_j, \dots, s'_{j+r-1} \rangle. \end{aligned} \quad (\text{D13})$$

In fact, this formula allows us to simplify Eq. (D12) even further. Defining a 2×2 matrix with elements

$$\begin{aligned} O_{s_{j-1}, s_{j+r}} &= \left(\frac{\sin \theta}{2} \right)^{r+1} \sum_{\substack{s_j, \dots, s_{j+r-1}, \\ s'_j, \dots, s'_{j+r-1}}} \langle s_j, \dots, s_{j+r-1} | \mathbf{O} | s'_j, \dots, s'_{j+r-1} \rangle \\ &\times \exp \left\{ \beta \left[s_{j-1}(s_j + s'_j) + s_{j+r}(s_{j+r-1} + s'_{j+r-1}) + \sum_{\ell=j}^{j+r-2} (s_\ell s_{\ell+1} + s'_\ell s'_{\ell+1}) \right] \right\}, \end{aligned} \quad (\text{D14})$$

we can rewrite the expectation value of \mathbf{O} as

$$\langle \mathbf{O} \rangle = \frac{1}{2} \sum_{s_1, s_L} [T^{j-2} O T^{L-j-r}]_{s_1, s_L}. \quad (\text{D15})$$

Assuming $\theta \notin \{0, \pi\}$, the thermodynamic limit suppresses the powers of $\cos \theta$, originating in the second term

of Eq. (D9). We finally obtain

$$\langle \mathbf{O} \rangle = \frac{1}{2} \sum_{s_1, s_L} \left[\frac{1 + \sigma^x}{2} O \frac{1 + \sigma^x}{2} \right]_{s_1, s_L} = \frac{1}{2} (1 \ 1) O \begin{pmatrix} 1 \\ 1 \end{pmatrix}. \quad (\text{D16})$$

Consider now two operators \mathbf{O}_1 and \mathbf{O}_2 , acting on re-

gions that are separated for d sites. Using the same argument as above, the thermodynamic limit now yields

$$\langle \mathbf{O}_1 \mathbf{O}_2 \rangle = \left[\frac{1}{2} (1 \ 1) O_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right] \left[\frac{1}{2} (1 \ 1) O_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right] + \mathcal{O}(\cos^d \theta), \quad (\text{D17})$$

implying

$$\langle \mathbf{O}_1 \mathbf{O}_2 \rangle - \langle \mathbf{O}_1 \rangle \langle \mathbf{O}_2 \rangle = \mathcal{O}(\cos^d \theta) \quad (\text{D18})$$

for the connected 2-point correlation function. The exponential decay of the correlations between any distant local operators in the τ -representation signifies clustering properties of the tilted state $|\mathcal{N}_\theta\rangle_\sigma$ for even local and even semilocal operators in the σ -representation.

Note that, for diagonal local operators \mathbf{O} (which are written solely in terms of $\{\tau_j^z\}$), Eq. (D12) becomes

$$\langle \mathbf{O} \rangle = \frac{\sum_{\underline{s}} \langle \underline{s} | \mathbf{O} | \underline{s} \rangle e^{\beta E(\underline{s})}}{\sum_{\underline{s}} e^{\beta E(\underline{s})}}, \quad (\text{D19})$$

i.e., a thermal expectation value. In the infinite system such operators include, for example, $\mathbf{\Pi}_{\sigma,+}^z(j)$ (the latter is mapped into τ_j^z by the inverse duality transformation (B4)). Eq. (D18) can then simply be seen as a consequence of the absence of finite-temperature phase transitions in the one-dimensional classical Ising model.

Appendix E: Excess of entropy

1. Bounds on Renyi entropies

Here we prove Eq. (5.19) by showing

$$2^{-\alpha} \text{tr}(\rho^\alpha) \leq \text{tr}(\bar{\rho}^\alpha) \leq \text{tr}(\rho^\alpha). \quad (\text{E1})$$

Taking the logarithm and multiplying the inequalities by $1/(1-\alpha)$ yields Eq. (5.19).

We prove both inequalities in Eq. (E1) using the inequality of Ault [138, 139], which says that, given arbitrary $n \times n$ complex matrices A_1, A_2, \dots, A_m , one has

$$\text{tr}\left(\frac{B + B^\dagger}{2}\right) \leq \frac{1}{m} \text{tr}\left(\sum_{j=1}^m (A_j A_j^\dagger)^{\frac{m}{2}}\right) \quad (\text{E2})$$

for $B = A_1 A_2 \dots A_m$.

To show the upper bound in (E1) we write

$$\text{tr} \bar{\rho}^\alpha = \text{tr}\left(\frac{\rho \bar{\rho}^{\alpha-1} + \text{h.c.}}{2}\right), \quad (\text{E3})$$

which follows from the definition (5.17) of $\bar{\rho}$, using the cyclic property of the trace and the invariance of $\bar{\rho}$ under the Hermitian involution P . The inequality of Ault for $A_1 = \rho$, $A_2 = A_3 = \dots = A_\alpha = \bar{\rho}$ then yields

$$\text{tr} \bar{\rho}^\alpha \leq \frac{1}{\alpha} [\text{tr}(\rho^\alpha) + (\alpha - 1) \text{tr}(\bar{\rho}^\alpha)]. \quad (\text{E4})$$

Rearranging the terms proves the second inequality in Eq. (E1).

To prove the lower bound in Eq. (E1) we introduce projectors $P_\pm = (I \pm P)/2$ and write

$$\bar{\rho} = P_+ \rho P_+ + P_- \rho P_-. \quad (\text{E5})$$

Note that the operators $P_+ \rho P_+$ and $P_- \rho P_-$ are Hermitian and positive semidefinite. Since $P_+ P_- = 0$ it follows

$$\bar{\rho}^\alpha = (P_+ \rho P_+)^\alpha + (P_- \rho P_-)^\alpha. \quad (\text{E6})$$

Since $\text{tr}[(\rho P_s)^\alpha] = \text{tr}[(P_s \rho P_s)^\alpha]$, we then see

$$0 \leq \text{tr}[(\rho P_s)^\alpha] \leq \text{tr} \bar{\rho}^\alpha, \quad s = \pm 1. \quad (\text{E7})$$

On the other hand using $P_+ + P_- = I$ we have

$$\text{tr} \rho^\alpha = \sum_{s_1, \dots, s_\alpha = \pm 1} \text{tr}(P_{s_1} \rho P_{s_2} \rho \dots P_{s_\alpha} \rho), \quad (\text{E8})$$

which can be rewritten, using the cyclic property of the trace, as

$$\text{tr} \rho^\alpha = \sum_{s_1, \dots, s_\alpha = \pm 1} \text{tr} \left[\frac{(\sqrt{\rho} P_{s_1} \sqrt{\rho}) \dots (\sqrt{\rho} P_{s_\alpha} \sqrt{\rho}) + \text{h.c.}}{2} \right]. \quad (\text{E9})$$

Here, $\sqrt{\rho}$ denotes the square root of the positive semidefinite operator ρ . Applying the inequality of Ault with $A_j = \sqrt{\rho} P_{s_j} \sqrt{\rho}$ for $j = 1, \dots, \alpha$ now yields

$$\text{tr} \rho^\alpha \leq \sum_{s_1, \dots, s_\alpha = \pm 1} \frac{1}{\alpha} \sum_{j=1}^{\alpha} \text{tr}[(\rho P_{s_j})^\alpha]. \quad (\text{E10})$$

Finally, using (E7) we obtain

$$\text{tr} \rho^\alpha \leq 2^\alpha \text{tr} \bar{\rho}^\alpha, \quad (\text{E11})$$

which proves the lower bound in Eq. (E1).

2. Correlation matrix after the quench

At time t the correlation matrix symbol for the initial states $|\uparrow\rangle_\tau$ and $|\downarrow\rangle_\tau$ reads, respectively,

$$\Gamma(p) = \pm e^{-i\mathcal{H}(p)t} \sigma^y e^{i\mathcal{H}(p)t}, \quad (\text{E12})$$

where $\mathcal{H}(p) = f_0(p)I + f_1(p)\sigma^x + f_2(p)\sigma^y + f_3(p)\sigma^z$ is the symbol of the Hamiltonian. For the generalized XY model we have explicitly

$$f_0(p) = 2 \sum_{k=1}^{\infty} (J_x^{(2k)} - J_y^{(2k)}) \sin(2kp), \quad (\text{E13})$$

$$f_1(p) = 2 \sum_{k=0}^{\infty} (J_x^{(2k+1)} - J_y^{(2k+1)}) \sin[(2k+1)p], \quad (\text{E14})$$

$$f_2(p) = -2 \sum_{k=0}^{\infty} (J_x^{(2k+1)} + J_y^{(2k+1)}) \cos[(2k+1)p], \quad (\text{E15})$$

$$f_3(p) = -2 \sum_{k=1}^{\infty} (J_x^{(2k)} + J_y^{(2k)}) \sin(2kp), \quad (\text{E16})$$

and focusing on the reflection symmetric system ($J_x^{(2k)} = J_y^{(2k)} \forall k$) we have the simplification $f_0(p) = 0$. For $t \rightarrow \infty$ due to dephasing the symbol can be effectively replaced by its time average, equal to

$$\Gamma(p) = \pm \frac{f_2(p)}{\varepsilon^2(p)} \mathcal{H}(p), \quad (\text{E17})$$

which gives Eq. (5.35) in the main text.

Let us determine the root density $\rho(p)$ starting from this expression, since we are going to need it to derive eq. (5.36). It can be shown [103] that for translationally invariant and reflection symmetric free fermionic models the root density $\rho(p)$ is related to the correlation matrix symbol $\Gamma(p)$ through the relation

$$V(p)\Gamma(p)V^\dagger(p) = 4\pi\rho_o(p)I + (4\pi\rho_e(p) - 1)\sigma^y + 4\pi\psi_R(p)\sigma^z - 4\pi\psi_I(p)\sigma^x. \quad (\text{E18})$$

Here $\rho_o(p)$ and $\rho_e(p)$ are the odd and the even part of $\rho(p)$ respectively, $\psi(p) = \psi_R(p) + i\psi_I(p)$ is a field related to the off-diagonal elements of the density matrix, that can be neglected in the limit $t \rightarrow \infty$, while $V(p)$ is a unitary transformation related to Bogoliubov rotation. The action of the latter on the symbol of the Hamiltonian reads

$$V(p)\mathcal{H}(p)V^\dagger(p) = \varepsilon(p)\sigma^y. \quad (\text{E19})$$

In the limit $t \rightarrow \infty$ we have thus simply

$$\Gamma(p) = 4\pi\rho_o(p)I + (4\pi\rho_e(p) - 1) \frac{\mathcal{H}(p)}{\varepsilon(p)}, \quad (\text{E20})$$

so, comparing with (E17), we find

$$4\pi\rho(p) - 1 = \pm \frac{f_2(p)}{\varepsilon(p)}, \quad (\text{E21})$$

where we have $\rho(p) = \rho_e(p)$ because the state is reflection symmetric.

When the initial state is the domain wall state $|\dots \downarrow\downarrow_r \uparrow\uparrow \dots\rangle_\tau$, as in bipartitioning protocols in general, the system reaches stationarity along rays $\zeta = x/t$, where x is the distance from the junction [103, 116]. Along a given ray one can still describe the observables by the root density $\rho_\zeta(p)$ or the correlation matrix symbol $\Gamma_\zeta(p)$, through (E20), which now holds along the ray. The late-time dynamics is described by the hydrodynamic equation [103, 117, 118]

$$\partial_t \rho_\zeta(p) + v(p) \partial_x \rho_\zeta(p) = 0, \quad (\text{E22})$$

where $v(p) = \varepsilon'(p)$ is the velocity of excitations. The equation implies that $\rho_\zeta(p)$ depends only on the difference $x - v(p)t = t[\zeta - v(p)]$.

The subsystem A corresponds to $\zeta = 0$, since the domain wall is placed inside. For $t \rightarrow \infty$ and $\zeta = 0$ we have $x - v(p)t \rightarrow -\text{sgn}[v(p)]\infty$. However, the same value of

$x - v(p)t$ is achieved for $\zeta \rightarrow -\text{sgn}[v(p)]\infty$ and $t \rightarrow \infty$. The latter corresponds simply to the evolution starting from the state $|\uparrow\rangle_\tau$ or $|\downarrow\rangle_\tau$, depending on the sign of the velocity. From (E21) we conclude that the zero ray root density is given by

$$4\pi\rho_{\zeta=0}(p) - 1 = -\text{sgn}[v(p)] \frac{f_2(p)}{\varepsilon(p)}. \quad (\text{E23})$$

Finally, from (E20) we find the correlation matrix symbol for the subsystem A ,

$$\Gamma_{\zeta=0}(p) = -\text{sgn}[v(p)] \frac{f_2(p)}{\varepsilon(p)} I. \quad (\text{E24})$$

3. Asymptotic results for Renyi entropies

The Renyi entropies can be expressed as the integral [114]

$$S_\alpha^{(\tau)}(\ell, \Psi) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi i} \oint_{C_\varepsilon} F_\alpha(z) \frac{d}{dz} \log \det [zI - \Gamma] dz, \quad (\text{E25})$$

where

$$F_\alpha(z) = \frac{1}{2(1-\alpha)} \log \left[\left(\frac{1+z}{2} \right)^\alpha + \left(\frac{1-z}{2} \right)^\alpha \right] \quad (\text{E26})$$

and the integration curve C_ε goes around the interval $[-1, 1]$ on the real axis, at distance ε from it. The information on the state $|\Psi(t)\rangle$ is, of course, encoded in the correlation matrix Γ .

A block Toeplitz matrix $T_\ell(\mathcal{M})$ with a 2×2 matrix symbol $\mathcal{M}(p)$ is defined by the elements

$$[T_\ell(\mathcal{M})]_{2j+m, 2l+n} = \int_{-\pi}^{\pi} \mathcal{M}(p) e^{ip(j-l)} \frac{dp}{2\pi} \quad (\text{E27})$$

for $0 \leq j, l \leq \ell - 1$, $0 \leq m, n \leq 1$. For a piecewise continuous symbol, such that $\det \mathcal{M}(p)$ is non-zero and has zero winding number, and jump discontinuities at points p_r for $r = 1, 2, \dots, R$, we use the asymptotic formula for large ℓ ,

$$\begin{aligned} & \log \det T_\ell(\mathcal{M}) \\ &= \int_{-\pi}^{\pi} \log \det \mathcal{M}(p) \frac{dp}{2\pi} + \log(\ell) \sum_{r=1}^R \mathcal{B}_r + O(1). \end{aligned} \quad (\text{E28})$$

Without discontinuities, the second term is not present. The leading term grows linearly with ℓ and its value comes from the Szegő-Widom theorem [126, 127]. It has been conjectured and checked numerically in [128, 129] that with discontinuities there is a subdominant term that grows logarithmically with ℓ . Denoting by $\mu_{r,j}^\pm$ for $j = 1, 2$ the eigenvalues of the limits $\lim_{p \rightarrow p_r^\pm} \mathcal{M}(p)$, the conjecture says that the logarithmic coefficients \mathcal{B}_r are given by

$$\mathcal{B}_r = \frac{1}{4\pi^2} \sum_{j=1}^2 \left(\log \frac{\mu_{r,j}^-}{\mu_{r,j}^+} \right)^2. \quad (\text{E29})$$

Note that the formula generalizes straightforwardly to a symbol of arbitrary size.

The asymptotic results for the Renyi entropies are obtained by using the asymptotic formula (E28) to compute the determinant of the Toeplitz matrix $zI - \Gamma$ in (E25). The latter is associated to the symbol $\mathcal{M}(p) = zI - \Gamma(p)$, where $\Gamma(p)$ is the symbol of the correlation matrix, given by (5.35) for a quench from the state of all spin up and by (5.36) for a quench from a domain wall state. In this way we find that the leading term both in (5.37) and in (5.38) is given by

$$a_\alpha = \frac{1}{1-\alpha} \int_{-\pi}^{\pi} \log \left[\left(\frac{1 + \frac{f_2(p)}{\varepsilon(p)}}{2} \right)^\alpha + \left(\frac{1 - \frac{f_2(p)}{\varepsilon(p)}}{2} \right)^\alpha \right] \frac{dp}{2\pi}, \quad (\text{E30})$$

where $f_2(p) = \text{tr}[\sigma^y \mathcal{H}(p)]$ for the generalized XY model is given by eq. (E15).

The discontinuities $p_1, p_2, \dots, p_R \in (-\pi, \pi]$ of the symbol (5.36) are the zeroes of the velocity, i.e. points for which $v(p_r) = 0$. From the asymptotic formula (E28) the logarithmic coefficient in (5.38) is given by

$$b_\alpha = \lim_{\varepsilon \rightarrow 0^+} \sum_{r=1}^R \frac{1}{2\pi i} \oint_{C_\varepsilon} F_\alpha(z) \frac{d}{dz} \log \mathcal{B}_r(z) dz, \quad (\text{E31})$$

where

$$\mathcal{B}_r(z) = \frac{1}{2\pi^2} \left(\log \frac{z - \frac{f_2(p_r)}{\varepsilon(p_r)}}{z + \frac{f_2(p_r)}{\varepsilon(p_r)}} \right)^2. \quad (\text{E32})$$

To evaluate the integral we use the partial integration,

$$b_\alpha = - \lim_{\varepsilon \rightarrow 0^+} \sum_{r=1}^R \frac{1}{2\pi i} \oint_{C_\varepsilon} \frac{d}{dz} [F_\alpha(z)] \mathcal{B}_r(z) dz, \quad (\text{E33})$$

and deform the contour around the poles of

$$\frac{d}{dz} F_\alpha(z) = \frac{\alpha}{2(1-\alpha)} \frac{(1+z)^{\alpha-1} - (1-z)^{\alpha-1}}{(1+z)^\alpha + (1-z)^\alpha}, \quad (\text{E34})$$

which are given by

$$z_j = i \tan \left[\frac{(2j-1)\pi}{2\alpha} \right], \quad j = 1, 2, \dots, \alpha, \quad j \neq \frac{\alpha+1}{2}. \quad (\text{E35})$$

As $z \rightarrow z_j$ we have

$$\frac{d}{dz} F_\alpha(z) = \frac{1}{2(1-\alpha)(z-z_j)} (1 + O(z-z_j)) \quad (\text{E36})$$

so we obtain

$$b_\alpha = - \frac{1}{4\pi^2(\alpha-1)} \times \sum_{r=1}^R \sum_{\substack{j=1,2,\dots,\alpha \\ j \neq \frac{\alpha+1}{2}}} \left(\log \frac{i \tan \left[\frac{(2j-1)\pi}{2\alpha} \right] - \frac{f_2(p)}{\varepsilon(p)}}{i \tan \left[\frac{(2j-1)\pi}{2\alpha} \right] + \frac{f_2(p)}{\varepsilon(p)}} \right)^2. \quad (\text{E37})$$

Since the term inside the logarithm is just a phase factor, the squared logarithm is real and negative. Because of the prefactor every term in the sum brings thus a positive contribution to b_α . For instance, for $\alpha = 2$ this expression simplifies to eq. (5.39) in the main text.

The velocity of excitations is given by

$$v(p) = \frac{\sum_{j=1}^3 f_j(p) f'_j(p)}{\varepsilon(p)}. \quad (\text{E38})$$

For the XY chain the velocity has four zeroes, given by $p_1 = -\pi/2$, $p_2 = 0$, $p_3 = \pi/2$, $p_4 = \pi$. Since $f_2(p) = 0$ for $p = p_1, p_3$, these two points do not contribute in (E37). As a matter of fact, since $f_2(p) = 0$ there is no discontinuity in the symbol at these points. For $p = p_2, p_4$ we have $f_2(p)/\varepsilon(p) \in \{-1, 1\}$ so the contribution of these two points is the same. From (E37) we find the logarithmic coefficient (5.40). A general model under consideration contains the aforementioned four zeroes of the velocity but it can contain also additional ones. Therefore, in general the logarithmic coefficient for the XY chain is the lower bound. Namely, $b_\alpha \geq b_\alpha^{(XY)}$.

4. Exact Procedure

In this section we show how to compute the second Renyi entropy exactly for systems mappable to free fermions, starting from eq. (5.15). Let us denote shortly $\rho_\tau \equiv \rho_A^{(\tau)}(\Psi)$ and $\bar{\rho} \equiv \bar{\rho}_A^{(\tau)}(\Psi)$.

The density matrix ρ_τ can be expressed as a Gaussian in Majorana fermions (5.33), following the standard techniques [115, 119, 140] based on Wick theorem [141]. Namely, defining the $2(\ell+1) \times 2(\ell+1)$ correlation matrix with the elements

$$\Gamma_{2j+\alpha, 2l+\beta} = \delta_{j,l} \delta_{\alpha,\beta} - \langle \Psi(t) | \mathbf{a}_j^\alpha \mathbf{a}_l^\beta | \Psi(t) \rangle, \quad (\text{E39})$$

$$0 \leq j, l \leq \ell, \quad 1 \leq \alpha, \beta \leq 2,$$

(with the usual identification $x \equiv 1, y \equiv 1$) and the matrix function

$$W(\Gamma) = \log[(I + \Gamma)(I - \Gamma)^{-1}] \quad (\text{E40})$$

we have

$$\rho_\tau = \frac{1}{\mathcal{Z}(\Gamma)} \exp \left(\frac{\vec{a}^T W(\Gamma) \vec{a}}{4} \right), \quad (\text{E41})$$

where $\vec{a} = (a_1^x, a_1^y, a_2^x, a_2^y, \dots, a_{\ell+1}^y)$. The normalization is given by the function

$$\mathcal{Z}(\Gamma) = \pm \left[\det \left(\frac{I - \Gamma}{2} \right) \right]^{-\frac{1}{2}}, \quad (\text{E42})$$

which can be computed from the formula (E47) that we introduce later. There is a sign ambiguity in the formula (E42), which can be avoided in computing the second Renyi entropy.

The effects of the transformations P_i on the Gaussian in eq. (5.15) are completely determined by their effects on the two-point products of Majorana fermions, i.e. by $P_i a_l^{x,y} a_l^{x,y} P_i$. We thus find

$$P_i \rho_\tau P_i = \frac{1}{\mathcal{Z}(\mathcal{F}_i(\Gamma))} \exp\left(\frac{\vec{a}^T W(\mathcal{F}_i(\Gamma)) \vec{a}}{4}\right), \quad (\text{E43})$$

where we define the functions \mathcal{F}_i by their action on the matrix elements of the correlation matrix:

$$\Gamma_{j,l} \xrightarrow{\mathcal{F}_1} (-1)^{\delta_{j,1} + \delta_{j,2} + \delta_{l,1} + \delta_{l,2}} \Gamma_{j,l} \quad (\text{E44})$$

$$\Gamma_{j,l} \xrightarrow{\mathcal{F}_2} (-1)^{\delta_{j,2\ell+1} + \delta_{j,2\ell+2} + \delta_{l,2\ell+1} + \delta_{l,2\ell+2}} \Gamma_{j,l} \quad (\text{E45})$$

$$\Gamma_{j,l} \xrightarrow{\mathcal{F}_3} (-1)^{\lfloor \frac{j}{2} \rfloor + \lfloor \frac{l}{2} \rfloor} \Gamma_{j,l}. \quad (\text{E46})$$

Of course, for products of different transformations, such as in $P_i P_j \rho_\tau P_j P_i$, we just have to consider the compositions such as $\mathcal{F}_i(\mathcal{F}_j(\cdot))$.

We have thus expressed the density matrix as a sum of Gaussian fermionic operators, similarly to the problem of the entanglement entropy of two disjoint blocks in the XY chain [119]. The Renyi entropies can thus be computed by exploring the formula for the trace of a product of Gaussians, given by [120]

$$\begin{aligned} \text{tr} \left(e^{\frac{\vec{a}^T W_1 \vec{a}}{4}} e^{\frac{\vec{a}^T W_2 \vec{a}}{4}} \dots e^{\frac{\vec{a}^T W_n \vec{a}}{4}} \right) \\ = \pm \sqrt{\det(I + e^{W_1} e^{W_2} \dots e^{W_n})}, \end{aligned} \quad (\text{E47})$$

for antisymmetric matrices W_1, W_2, \dots, W_n . The formula has a sign ambiguity, well known in such problems, that has to be resolved in some independent way. The normalization (E42) follows from formula (E47) with $n = 1$.

In computing the second Renyi entropy it is easy to avoid the sign ambiguity, based on fact that the trace of a product of two positive semidefinite Hermitian operators is non-negative [121]. Now, from the invariance $P_j \bar{\rho} P_j = \bar{\rho}$, for $j = 1, 2, 3$, that follows from (5.15), we get the simplification

$$\text{tr}[\bar{\rho}^2] = \text{tr}[\rho_\tau \bar{\rho}]. \quad (\text{E48})$$

Using (5.15) again, we can write this as a sum of eight non-negative terms

$$\begin{aligned} \text{tr}[\bar{\rho}^2] = \\ 2^{-3} \sum_{j_1, j_2, j_3=0,1} \text{tr} \left[\rho_\tau \left(P_3^{j_3} P_2^{j_2} P_1^{j_1} \rho_\tau P_1^{j_1} P_2^{j_2} P_3^{j_3} \right) \right]. \end{aligned} \quad (\text{E49})$$

Each of the eight terms can be evaluated using (E43) and the formula (E47) for the trace of a product of Gaussians. Using the identity

$$\frac{\det(I + e^{W(\Gamma_1)} e^{W(\Gamma_2)})}{\det(I + e^{W(\Gamma_1)}) \det(I + e^{W(\Gamma_2)})} = \det\left(\frac{I + \Gamma_1 \Gamma_2}{2}\right), \quad (\text{E50})$$

and eq. (5.16) we get then the second Renyi entropy,

$$\begin{aligned} S_2(\ell, \Psi) = 2 \log 2 - \log \left[\mathcal{G}(\Gamma, \Gamma) \right. \\ \left. + \sum_{i=1}^3 \mathcal{G}(\Gamma, \mathcal{F}_i(\Gamma)) + \sum_{1 \leq i < j \leq 3} \mathcal{G}(\Gamma, \mathcal{F}_i(\mathcal{F}_j(\Gamma))) \right. \\ \left. + \mathcal{G}(\Gamma, \mathcal{F}_1(\mathcal{F}_2(\mathcal{F}_3(\Gamma)))) \right], \end{aligned} \quad (\text{E51})$$

where we have defined the function

$$\mathcal{G}(\Gamma_1, \Gamma_2) = \sqrt{\det\left(\frac{I + \Gamma_1 \Gamma_2}{2}\right)}. \quad (\text{E52})$$

Appendix F: Melting of the order and scaling limits

This section provides details on the calculation of expectation values of local observables (e.g., strings of σ_j^z) in a time-evolved weakly tilted state $|\mathcal{A}_\theta\rangle$. The first part of the section discusses the effective density matrix given in Eq. (5.45), from which expectation values of semilocal order parameters can be calculated; the latter are discussed in the second part.

1. Effective density matrix

In the fermionic representation we have

$$\begin{aligned} e^{-i\mathbf{H}(\tau)t} \tau_j^z e^{i\mathbf{H}(\tau)t} = \frac{1}{2} \sum_{j_1, j_2} \int_{-\pi}^{\pi} \frac{d^2 k}{(2\pi)^2} e^{i[k_1(j_1-j) + k_2(j-j_2)]} \\ \times \vec{\mathbf{a}}_{j_1}^T \cdot \left[e^{-i\mathcal{H}(k_1)t} \sigma_y e^{i\mathcal{H}(k_2)t} \right] \cdot \vec{\mathbf{a}}_{j_2}, \end{aligned} \quad (\text{F1})$$

where $\vec{\mathbf{a}}_j = (\mathbf{a}_j^x \ \mathbf{a}_j^y)^T$ and $\mathcal{H}(k)$ is the symbol of the Hamiltonian, defined in Eq. (3.13). Using this representation in the expression (5.45) for the effective density matrix, performing the sums over j and n , and changing the integration variables as $k_1 \rightarrow k_2 - k_1$, $k_2 \rightarrow k_2$, $k_3 \rightarrow k_3$, $k_4 \rightarrow k_3 - k_4$, we obtain

$$\begin{aligned} \rho_{\text{eff}}(t) \sim \frac{1}{4 \text{tr} \mathbf{I}} \sum_{j_1, 2, 3, 4} \int \frac{d^3 k}{(2\pi)^3} \frac{e^{i[(j_4-j_1)k_1 + (j_1-j_2)k_2 + (j_3-j_4)k_3]}}{e^{\frac{\theta^2}{2} + ik_1} - 1} \\ \times \vec{\mathbf{a}}_{j_1}^T \cdot [e^{-i\mathcal{H}(k_2-k_1)t} \sigma_y e^{i\mathcal{H}(k_2)t}] \cdot \vec{\mathbf{a}}_{j_2} \\ \times \vec{\mathbf{a}}_{j_3}^T \cdot [e^{-i\mathcal{H}(k_3)t} \sigma_y e^{i\mathcal{H}(k_3-k_1)t}] \cdot \vec{\mathbf{a}}_{j_4}, \end{aligned} \quad (\text{F2})$$

for the effective density matrix.

Being interested in the expectation value of local operators at large times after a weak symmetry breaking, we will assume that t and θ are the only large and small

parameter, respectively, in Eq. (F2). We can assume this because local operator \mathbf{O} in $\text{tr}[\rho_{\text{eff}}(t)\mathbf{O}]$ is characterised by indices j_1, j_2, j_3 , and j_4 , confined to some finite interval. In order to obtain the asymptotic behaviour of the integral in Eq. (F2) one now has to consider the pole in k_1 . The first step consists of deforming the integration path in k_1 into a piecewise linear curve with lines parallel to the axes, in such a way that the horizontal lines give exponentially decaying contributions.




FIG. 16. The integration path over the interval $[-\pi, \pi]$ on the real axis is deformed depending on the sign of $v(k_3 - k_1^{\text{R}})s_2 - v(k_2 - k_1^{\text{R}})s_1$, to make the exponent in (F6) negative on the horizontal lines. Panel (a): A contour deformed without crossing the pole. Panel (b): Deforming the contour across the pole, its contribution has to be included.

We start from the relation

$$e^{-i\mathcal{H}(k_2 - k_1)t} = \sum_{s_1 = \pm 1} P_{s_1}(k_2 - k_1) e^{-i\varepsilon(k_2 - k_1)s_1 t}, \quad (\text{F3})$$

where $P_s(k)$ are the projectors on the eigenstates of $\mathcal{H}(k)$ with eigenvalues $\pm\varepsilon(k)$:

$$P_s(k) = \frac{1}{2} \left(I + s \frac{\mathcal{H}(k)}{\varepsilon(k)} \right). \quad (\text{F4})$$

Denoting the real and imaginary part of k_1 by $k_1^{\text{R}}, k_1^{\text{I}}$ respectively, assuming the imaginary part k_1^{I} to be small,

and expanding in it, we obtain

$$\begin{aligned} e^{-i\mathcal{H}(k_2 - k_1)t} &\sim \sum_{s_1} P_{s_1}(k_2 - k_1) e^{-i\varepsilon(k_2 - k_1^{\text{R}})s_1 t - k_1^{\text{I}} v(k_2 - k_1^{\text{R}})s_1 t}, \\ e^{i\mathcal{H}(k_3 - k_1)t} &\sim \sum_{s_2} P_{s_2}(k_3 - k_1) e^{i\varepsilon(k_3 - k_1^{\text{R}})s_2 t + k_1^{\text{I}} v(k_3 - k_1^{\text{R}})s_2 t}. \end{aligned} \quad (\text{F5})$$

where $v(k) := \varepsilon'(k)$. The expressions in each line of Eq. (F5) are multiplied in Eq. (F2). In particular, the real parts of each exponential in Eq. (F5) together yield a factor

$$e^{k_1^{\text{I}} [v(k_3 - k_1^{\text{R}})s_2 - v(k_2 - k_1^{\text{R}})s_1] t}. \quad (\text{F6})$$

Assuming $\theta^2 t$ to be fixed, we deform the integration path in k_1 so that $\theta^2 \ll |k_1^{\text{I}}| \ll 1$ and we choose the sign of k_1^{I} so that the factor multiplying t in Eq. (F6) is always negative and the horizontal lines in the contour are exponentially suppressed (the choice of the integration contour thus depends also on s_1, s_2). The contours that we work with are therefore determined by the sign of $v(k_3 - k_1^{\text{R}})s_2 - v(k_2 - k_1^{\text{R}})s_1$, and are presented in Fig. 16.

The pole is at $k_1^{\text{R}} = 0$ and there are two possible placements of the obtained contour with respect to the pole, depending on the exponent in eq. (F6):

1. For $v(k_3)s_2 - v(k_2)s_1 > 0$ we have $k_1^{\text{I}} < 0$ on the deformed contour when $k_1^{\text{R}} = 0$. In this case we do not have a pole contribution — see Fig. 16(a).
2. For $v(k_3)s_2 - v(k_2)s_1 < 0$ we have $k_1^{\text{I}} > 0$ on the deformed contour when $k_1^{\text{R}} = 0$. In such a case the contour has to be extended across the pole — see Fig. 16(b). In order for the integral along the extended contour to reproduce the correct result, the pole contribution has to be subtracted.

In the first case the integral is exponentially suppressed along the entire contour and it yields zero. In the second case only the subtracted pole contribution remains of the integral. This can be described simply by including the factor $\theta_H(v(k_2)s_1 - v(k_3)s_2)$ in the integral, where θ_H is the Heaviside step function. Assuming $\theta^2(\theta^2 t) \rightarrow 0$, we then include into Eq. (F2) the pole contribution

$$\begin{aligned} e^{-i\mathcal{H}(k_2 - i\frac{\theta^2}{2})t} &= \sum_{s_1 = \pm 1} P_{s_1}(k_2) e^{-i\mathcal{H}(k_2)s_1 t - v(k_2)\frac{\theta^2}{2}s_1 t} \\ &\quad \times (1 + \mathcal{O}(\theta^2)), \end{aligned} \quad (\text{F7})$$

and an analogous one for $\exp[i\mathcal{H}(k_3 - i\theta^2/2)t]$, to get

$$\begin{aligned} \rho_{\text{eff}}(t) &\sim \frac{1}{4 \text{tr} \mathbf{I}} \sum_{\substack{j_1, j_2, \\ j_3, j_4}} \int \frac{dk_2 dk_3}{(2\pi)^3} \sum_{s_1, s_2 = \pm 1} e^{i[(j_1 - j_2)k_2 + (j_3 - j_4)k_3]} \\ &\quad \times e^{-\frac{\theta^2}{2} t (v(k_2)s_1 - v(k_3)s_2)} \theta_H(v(k_2)s_1 - v(k_3)s_2) \\ &\quad \times \bar{\mathbf{a}}_{j_1}^T \cdot [P_{s_1}(k_2) \Gamma_t(k_2)] \cdot \bar{\mathbf{a}}_{j_2} \\ &\quad \times \bar{\mathbf{a}}_{j_3}^T \cdot [P_{s_2}(k_3) \Gamma_t(k_3)] \cdot \bar{\mathbf{a}}_{j_4}. \end{aligned} \quad (\text{F8})$$

Here we denoted $\Gamma_t(k) = e^{-i\mathcal{H}(k)t} \sigma^y e^{i\mathcal{H}(k)t}$. In the limit $t \rightarrow \infty$ we can replace the latter by its time average

$$\bar{\Gamma}(k) = -\frac{2(J_x + J_y) \cos k}{\varepsilon^2(k)} \mathcal{H}(k), \quad (\text{F9})$$

ending up with

$$\begin{aligned} \rho_{\text{eff}}(t) &\sim \frac{1}{\text{tr} \mathbf{I}} \int \frac{d^2 k}{(2\pi)^2} \frac{(J_x + J_y)^2 \cos k_1 \cos k_2}{\varepsilon(k_1) \varepsilon(k_2)} \\ &\times \sum_{s=\pm 1} s e^{-|v(k_1) - sv(k_2)| \frac{\theta^2 t}{2}} \\ &\times \sum_{j_1, j_2} e^{i(j_1 - j_2)k_1} \vec{\mathbf{a}}_{j_1}^T \cdot P_{\text{sgn}[v(k_1) - sv(k_2)]}(k_1) \cdot \vec{\mathbf{a}}_{j_2} \\ &\times \sum_{j_3, j_4} e^{i(j_3 - j_4)k_2} \vec{\mathbf{a}}_{j_3}^T \cdot P_{\text{sgn}[sv(k_1) - v(k_2)]}(k_2) \cdot \vec{\mathbf{a}}_{j_4}. \end{aligned} \quad (\text{F10})$$

For sufficiently large $\theta^2 t$ this generically decays as $1/(\theta^2 t)$.

2. Semilocal order parameter

Having obtained the effective reduced density matrix, we are now ready to investigate the expectation values of the string $\prod_{\ell=0}^{n-1} \sigma_{j+\ell}^z$, which corresponds to a product of four Majorana fermions. For $n = 1$ this is the simplest local operator sensitive to the semilocal order [31], while in the limit $n \rightarrow \infty$ it can be regarded as a string-order parameter.

Let us then consider $Z_n = \langle \prod_{\ell=0}^{n-1} \sigma_{j+\ell}^z \rangle = -\langle \mathbf{a}_1^x \mathbf{a}_1^y \mathbf{a}_{n+1}^x \mathbf{a}_{n+1}^y \rangle$. This can be computed from Eq. (F10) using Wick's theorem:

$$\begin{aligned} Z_n &= \int \frac{d^2 k}{(2\pi)^2} \frac{2(J_x + J_y)^2 \cos k_1 \cos k_2}{\varepsilon(k_1) \varepsilon(k_2)} \sum_{s, s'=\pm 1} s e^{-|v(k_1) - sv(k_2)| \frac{\theta^2 t}{2}} \sin^2 \left[n \frac{k_1 + s' k_2}{2} \right] \\ &\times \left(-s' + 4s \frac{(J_x + J_y)^2 \cos k_1 \cos k_2 + s'(J_x - J_y)^2 \sin k_1 \sin k_2}{\varepsilon(k_1) \varepsilon(k_2)} \right). \end{aligned} \quad (\text{F11})$$

This expression simplifies in the limit $n \rightarrow \infty$ (bear in mind that this limit has been taken after the limit of large time and small angle at fixed $\theta^2 t$), in which we

obtain Eq. (5.46).

-
- [1] X. Chen, Z.-C. Gu, and X.-G. Wen, Phys. Rev. B **83**, 035107 (2011).
 - [2] X. G. Wen, Phys. Rev. Lett. **64**, 2206 (1990).
 - [3] X. G. Wen, Phys. Rev. B **43**, 11025 (1991).
 - [4] X. G. Wen, Phys. Rev. Lett. **66**, 802 (1991).
 - [5] X. Chen, Z.-X. Liu, and X.-G. Wen, Phys. Rev. B **84**, 235141 (2011).
 - [6] P. Fendley, Journal of Physics A: Mathematical and Theoretical **49**, 30LT01 (2016).
 - [7] T. Scaffidi, D. E. Parker, and R. Vasseur, Phys. Rev. X **7**, 041048 (2017).
 - [8] F. Pollmann, A. M. Turner, E. Berg, and M. Oshikawa, Phys. Rev. B **81**, 064439 (2010).
 - [9] M. den Nijs and K. Rommelse, Phys. Rev. B **40**, 4709 (1989).
 - [10] F. Pollmann and A. M. Turner, Phys. Rev. B **86**, 125441 (2012).
 - [11] C. de Groot, A. Turzillo, and N. Schuch, Symmetry protected topological order in open quantum systems (2021).
 - [12] S. Roberts, B. Yoshida, A. Kubica, and S. D. Bartlett, Phys. Rev. A **96**, 022306 (2017).
 - [13] D. I. Tsomokos, A. Hamma, W. Zhang, S. Haas, and R. Fazio, Phys. Rev. A **80**, 060302 (2009).
 - [14] A. Rahmani and C. Chamon, Phys. Rev. B **82**, 134303 (2010).
 - [15] G. B. Halász and A. Hamma, Phys. Rev. Lett. **110**, 170605 (2013).
 - [16] L. Mazza, D. Rossini, M. Endres, and R. Fazio, Phys. Rev. B **90**, 020301 (2014).
 - [17] P. Wang, W. Yi, and G. Xianlong, New Journal of Physics **17**, 013029 (2015).
 - [18] M. Calvanese Strinati, L. Mazza, M. Endres, D. Rossini, and R. Fazio, Phys. Rev. B **94**, 024302 (2016).
 - [19] Y. Zeng, A. Hamma, and H. Fan, Phys. Rev. B **94**, 125104 (2016).
 - [20] L. Rademaker, SciPost Phys. **7**, 71 (2019).
 - [21] A. Polkovnikov, K. Sengupta, A. Silva, and M. Venugatlattore, Rev. Mod. Phys. **83**, 863 (2011).
 - [22] J. Eisert, M. Friesdorf, and C. Gogolin, Nature Phys **11**, 124 (2015).
 - [23] C. Gogolin and J. Eisert, Rep. Prog. Phys. **79**, 056001 (2016).
 - [24] P. Calabrese and J. Cardy, J. Stat. Mech. **2016**, 064003 (2016).
 - [25] F. H. L. Essler and M. Fagotti, J. Stat. Mech. **2016**,

- 064002 (2016).
- [26] M. A. Cazalilla and M.-C. Chung, *Journal of Statistical Mechanics: Theory and Experiment* **2016**, 064004 (2016).
- [27] J.-S. Caux, *Journal of Statistical Mechanics: Theory and Experiment* **2016**, 064006 (2016).
- [28] L. Vidmar and M. Rigol, *J. Stat. Mech.* **2016**, 064007 (2016).
- [29] E. Ilievski, J. De Nardis, B. Wouters, J.-S. Caux, F. H. L. Essler, and T. Prosen, *Phys. Rev. Lett.* **115**, 157201 (2015).
- [30] E. Ilievski, M. Medenjak, T. Prosen, and L. Zadnik, *J. Stat. Mech.* **2016**, 064008 (2016).
- [31] M. Fagotti, *Phys. Rev. Lett.* **128**, 110602 (2022).
- [32] M. Rigol, V. Dunjko, and M. Olshanii, *Nature* **452**, 854 (2008).
- [33] J. M. Deutsch, *Phys. Rev. A* **43**, 2046 (1991).
- [34] M. Srednicki, *Phys. Rev. E* **50**, 888 (1994).
- [35] J. M. Deutsch, *Reports on Progress in Physics* **81**, 082001 (2018).
- [36] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, *Phys. Rev. Lett.* **98**, 050405 (2007).
- [37] L. Zadnik and M. Fagotti, *SciPost Phys. Core* **4**, 10 (2021).
- [38] V. Eisler and F. Maislinger, *Phys. Rev. B* **98**, 161117 (2018).
- [39] V. Eisler and F. Maislinger, *SciPost Phys.* **8**, 37 (2020).
- [40] K. Bidzhiev, M. Fagotti, and L. Zadnik, *arXiv:2108.03230* (2021).
- [41] L. Zadnik, S. Bocini, K. Bidzhiev, and M. Fagotti, *arXiv:2111.06325* (2021).
- [42] M. Gruber and V. Eisler, *SciPost Phys.* **10**, 5 (2021).
- [43] V. Eisler, *J. Phys. A: Math. Theor.* **54**, 424002 (2021).
- [44] A. Kitaev and J. Preskill, *Phys. Rev. Lett.* **96**, 110404 (2006).
- [45] B. Bertini, F. H. L. Essler, S. Groha, and N. J. Robinson, *Phys. Rev. Lett.* **115**, 180601 (2015).
- [46] B. Bertini and M. Fagotti, *Journal of Statistical Mechanics: Theory and Experiment* **2015**, P07012 (2015).
- [47] V. Alba and M. Fagotti, *Phys. Rev. Lett.* **119**, 010601 (2017).
- [48] J. Durnin, M. J. Bhaseen, and B. Doyon, *Phys. Rev. Lett.* **127**, 130601 (2021).
- [49] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 1 & 2*, 2nd ed. (Berlin: Springer-Verlag, 1997).
- [50] E. Ilievski and T. Prosen, *Communications in Mathematical Physics* **318**, 809 (2013).
- [51] T. Prosen, *Nuclear Physics B* **886**, 1177 (2014).
- [52] B. Doyon, *Commun. Math. Phys.* **351**, 155 (2017).
- [53] E. H. Lieb and D. W. Robinson, *Commun. Math. Phys.* **28**, 251 (1972).
- [54] S. Bravyi, M. B. Hastings, and F. Verstraete, *Phys. Rev. Lett.* **97**, 050401 (2006).
- [55] D. V. Else, S. D. Bartlett, and A. C. Doherty, *Phys. Rev. B* **88**, 085114 (2013).
- [56] K. Duivenvoorden and T. Quella, *Phys. Rev. B* **88**, 125115 (2013).
- [57] A. J. Beekman, L. Rademaker, and J. van Wezel, *SciPost Phys. Lect. Notes*, 11 (2019).
- [58] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, *Quantum Information Meets Quantum Matter* (Springer, New York, 2019).
- [59] E. T. Jaynes, *Phys. Rev.* **106**, 620 (1957).
- [60] P. Calabrese and J. Cardy, *Journal of Statistical Mechanics: Theory and Experiment* **2004**, P06002 (2004).
- [61] Although a discrete symmetry could in principle be broken, we are only aware of results pointing at its impossibility in specific classes of models.
- [62] There is no standard definition of “relaxation time”, but our considerations are quite independent of the details of the definition.
- [63] P. Calabrese and J. Cardy, *J. Stat. Mech.* **2005**, P04010 (2005).
- [64] P. Bocchieri and A. Loinger, *Phys. Rev.* **107**, 337 (1957).
- [65] A. C. Cassidy, C. W. Clark, and M. Rigol, *Phys. Rev. Lett.* **106**, 140405 (2011).
- [66] J.-S. Caux and F. H. L. Essler, *Phys. Rev. Lett.* **110**, 257203 (2013).
- [67] M. Fagotti and F. H. L. Essler, *Phys. Rev. B* **87**, 245107 (2013).
- [68] B. Wouters, J. De Nardis, M. Brockmann, D. Fioretto, M. Rigol, and J.-S. Caux, *Phys. Rev. Lett.* **113**, 117202 (2014).
- [69] B. Pozsgay, M. Mestyán, M. A. Werner, M. Kormos, G. Zaránd, and G. Takács, *Phys. Rev. Lett.* **113**, 117203 (2014).
- [70] T. Prosen and E. Ilievski, *Phys. Rev. Lett.* **111**, 057203 (2013).
- [71] R. G. Pereira, V. Pasquier, J. Sirker, and I. Affleck, *J. Stat. Mech.* **2014**, P09037 (2014).
- [72] E. Ilievski, M. Medenjak, and T. Prosen, *Phys. Rev. Lett.* **115**, 120601 (2015).
- [73] E. Ilievski, E. Quinn, and J.-S. Caux, *Phys. Rev. B* **95**, 115128 (2017).
- [74] B. Pozsgay, E. Vernier, and M. A. Werner, *Journal of Statistical Mechanics: Theory and Experiment* **2017**, 093103 (2017).
- [75] E. Ilievski and E. Quinn, *SciPost Phys.* **7**, 33 (2019).
- [76] D. Rossini, A. Silva, G. Mussardo, and G. E. Santoro, *Phys. Rev. Lett.* **102**, 127204 (2009).
- [77] Suppose instead that $\langle \mathbf{K}, \mathbf{O} \rangle_\lambda = 0$ for all local operators \mathbf{O} (including $\mathbf{O} = \mathbf{h}_j$). Then condition $\langle -\beta'(\lambda)\mathbf{H} + \mathbf{K}, \mathbf{h}_j \rangle_\lambda = 0$ would imply either $\langle \mathbf{H}, \mathbf{h}_j \rangle = 0$, which is clearly false, or $\beta'(\lambda) = 0$, the latter leading to $\delta \langle \mathbf{O} \rangle_\lambda = 0$ in Eq. (3.4).
- [78] We consider the entropy per site rather than the entropy because the latter is infinite in the thermodynamic limit.
- [79] If $\langle \mathbf{K}, \mathbf{k}_j \rangle$ was infinite $\beta'(\lambda)$ would have to compensate it in order for $\delta \mathfrak{s}[\rho_\infty(\lambda)]$ to remain finite. The entropy density would then not be a smooth function of λ .
- [80] E. Lieb, T. Schultz, and D. Mattis, *Annals of Physics* **16**, 407 (1961).
- [81] We remind the reader that the half-infinite string of Pauli matrices is not well defined in an infinite chain; this is however irrelevant as long as only operators quadratic in Majorana fermions are considered. For a rigorous definition of Jordan-Wigner transformation see instead Ref. [91]. We will use the latter definition in Section IV A where we discuss duality transformation and semilocal operators.
- [82] M. Fagotti, *J. Stat. Mech.* **2014**, P03016 (2014).
- [83] P. Calabrese, F. H. L. Essler, and M. Fagotti, *Journal of Statistical Mechanics: Theory and Experiment* **2012**, P07022 (2012).
- [84] C. Murthy and M. Srednicki, *Phys. Rev. E* **100**, 012146 (2019).
- [85] M. Gluza, J. Eisert, and T. Farrelly, *SciPost Phys.* **7**,

- 38 (2019).
- [86] P. Smacchia, L. Amico, P. Facchi, R. Fazio, G. Florio, S. Pascazio, and V. Vedral, *Phys. Rev. A* **84**, 022304 (2011).
- [87] T. Gombor and B. Pozsgay, *Phys. Rev. E* **104**, 054123 (2021).
- [88] M. Fagotti and F. H. L. Essler, *J. Stat. Mech.* **2013**, P07012 (2013).
- [89] B. Pozsgay, *J. Stat. Mech.* **2013**, P07003 (2013).
- [90] In fact, it will later become clear that the time evolution of $\langle \sigma_\ell^z \rangle$ in the dual XY model is exactly the same as that of $\langle \sigma_\ell^z \sigma_{\ell+1}^z \rangle$ in the XY model shown in Fig. 7.
- [91] H. Araki and T. Matsui, *Communications in Mathematical Physics* **101**, 213 (1985).
- [92] S. Weinberg, *The Quantum Theory of Fields*, Vol. 2 (Cambridge University Press, 1996).
- [93] T. Kennedy and H. Tasaki, *Phys. Rev. B* **45**, 304 (1992).
- [94] T. Kennedy and H. Tasaki, *Communications in Mathematical Physics* **147**, 431 (1992).
- [95] As far as we can see, semilocal terms in $\mathbf{W}(\tau)$, can always be replaced by local terms.
- [96] M. Grady, *Phys. Rev. D* **25**, 1103 (1982).
- [97] Note that the projected ensemble (4.33) gives correct predictions for the stationary expectation values of quasilocal operators, as can be seen by mapping it into the τ representation.
- [98] B. Pozsgay, *Journal of Statistical Mechanics: Theory and Experiment* **2014**, P09026 (2014).
- [99] G. Müller and R. E. Shrock, *Phys. Rev. B* **32**, 5845 (1985).
- [100] A $\mathbb{Z}_2 \times \mathbb{Z}_2$ -semilocal charge would be mapped by $\mathcal{D}_{\mathbb{Z}_2}^{-1}$ into a \mathbb{Z}_2 -semilocal charge in the τ representation, but there are no such charges in the latter — see Section IV C 1.
- [101] J. Shohat, J. Tamarkin, and A. Society, *The Problem of Moments*, *Mathematical Surveys and Monographs* (American Mathematical Society, 1943).
- [102] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States: An Introduction to Quantum Entanglement* (Cambridge University Press, 2006).
- [103] V. Alba, B. Bertini, M. Fagotti, L. Piroli, and P. Ruggiero, *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 114004 (2021).
- [104] J. D. Nardis, B. Doyon, M. Medenjak, and M. Panfil, *Journal of Statistical Mechanics: Theory and Experiment* **2022**, 014002 (2022).
- [105] V. B. Bulchandani, S. Gopalakrishnan, and E. Ilievski, *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 084001 (2021).
- [106] I. Bouchoule and J. Dubail, *Journal of Statistical Mechanics: Theory and Experiment* **2022**, 014003 (2022).
- [107] A. Bastianello, A. D. Luca, and R. Vasseur, *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 114003 (2021).
- [108] M. Borsi, B. Pozsgay, and L. Pristiyák, *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 094001 (2021).
- [109] The existence of odd charges implies that the stationary state after the quench from $|\uparrow\rangle_\tau$ is macroscopically different from that after the quench from $|\downarrow\rangle_\tau$; by symmetry, the nonequilibrium stationary state emerging from $|\cdots \downarrow_{r-1} \downarrow_r \uparrow_{r+1} \uparrow_{r+2} \cdots\rangle_\tau$ has to be macroscopically different from the two of them.
- [110] B. Doyon, J. Dubail, R. Konik, and T. Yoshimura, *Phys. Rev. Lett.* **119**, 195301 (2017).
- [111] J. Dubail, J.-M. Stéphan, J. Viti, and P. Calabrese, *SciPost Phys.* **2**, 002 (2017).
- [112] P. Calabrese and J. Cardy, *Journal of Physics A: Mathematical and Theoretical* **42**, 504005 (2009).
- [113] M. Suzuki, *Physics Letters A* **34**, 338 (1971).
- [114] B.-Q. Jin and V. E. Korepin, *Journal of Statistical Physics* **116**, 79 (2004).
- [115] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, *Phys. Rev. Lett.* **90**, 227902 (2003).
- [116] B. Bertini and M. Fagotti, *Phys. Rev. Lett.* **117**, 130402 (2016).
- [117] B. Bertini, M. Collura, J. De Nardis, and M. Fagotti, *Phys. Rev. Lett.* **117**, 207201 (2016).
- [118] O. A. Castro-Alvaredo, B. Doyon, and T. Yoshimura, *Phys. Rev. X* **6**, 041065 (2016).
- [119] M. Fagotti and P. Calabrese, *Journal of Statistical Mechanics: Theory and Experiment* **2010**, P04016 (2010).
- [120] I. Klich, *Journal of Statistical Mechanics: Theory and Experiment* **2014**, P11006 (2014).
- [121] R. Bhatia, *Variational principles for eigenvalues*, in *Matrix Analysis* (Springer New York, New York, NY, 1997) pp. 57–83.
- [122] M. Fagotti and P. Calabrese, *Phys. Rev. A* **78**, 010306 (2008).
- [123] V. Alba and P. Calabrese, *Proceedings of the National Academy of Sciences* **114**, 7947 (2017), <https://www.pnas.org/doi/pdf/10.1073/pnas.1703516114>.
- [124] V. Alba, B. Bertini, and M. Fagotti, *SciPost Phys.* **7**, 5 (2019).
- [125] B. Bertini, K. Klobas, V. Alba, G. Lagnese, and P. Calabrese, *arXiv:2203.17264* (2022).
- [126] H. Widom, *Advances in Mathematics* **21**, 1 (1976).
- [127] H. Widom, *Proceedings of the American Mathematical Society* **50**, 167 (1975).
- [128] F. Ares, J. G. Esteve, F. Falceto, and A. R. de Queiroz, *Phys. Rev. A* **92**, 042334 (2015).
- [129] F. Ares, J. G. Esteve, F. Falceto, and A. R. de Queiroz, *Phys. Rev. A* **97**, 062301 (2018).
- [130] In case the reader is not comfortable with replacing a density matrix with a traceless operator, we point out that the result does not change if a term proportional to the identity is added to ρ_{eff} .
- [131] R. Haag, *Local Quantum Physics: Fields, Particles, Algebras*, 2nd ed. (Springer Publishing Company, Incorporated, 1996).
- [132] I. Kukuljan, S. Sotiriadis, and G. Takács, *Journal of High Energy Physics* **2020**, 224 (2020).
- [133] H. Alaeian and B. Buča, *Exact bistability and time pseudo-crystallization of driven-dissipative fermionic lattices* (2022).
- [134] V. Marić, S. M. Giampaolo, D. Kuić, and F. Franchini, *New Journal of Physics* **22**, 083024 (2020).
- [135] V. Marić, S. M. Giampaolo, and F. Franchini, *Phys. Rev. B* **105**, 064408 (2022).
- [136] V. Marić, S. M. Giampaolo, and F. Franchini, *Communications Physics* **3**, 220 (2020).
- [137] G. Mussardo, *Statistical Field Theory: An Introduction to Exactly Solved Models in Statistical Physics*, 2nd ed., *Oxford Graduate Texts* (Oxford University Press, Oxford, 2020).
- [138] J. C. Ault, *Journal of the London Mathematical Society* **s1-42**, 497 (1967).
- [139] A. W. Marshall, I. Olkin, and B. C. Arnold, *Matrix*

- theory, in *Inequalities: Theory of Majorization and Its Applications* (Springer New York, New York, NY, 2011) pp. 297–365.
- [140] I. Peschel and V. Eisler, *Journal of Physics A: Mathematical and Theoretical* **42**, 504003 (2009).
- [141] M. Gaudin, *Nuclear Physics* **15**, 89 (1960).