

# BrainViewer: interacting with spatial connectome data at the mesoscale

Seth Daetwiler\*  
Computer Science  
Western Washington University

Angus Read†  
Computer Science  
Western Washington University  
Kameron Decker Harris§  
Computer Science  
Western Washington University

Jessica Stillwell‡  
Computer Science  
Western Washington University

## ABSTRACT

Scientists construct connectomes, comprehensive descriptions of neuronal connections across a brain, in order to better understand and model brain function. Interactive visualizations of these pathways would enable exploratory analysis of such information flows. Current tools can be used to see individual tracing experiments which are used to build mesoscale connectomes of the mouse brain, but not the brain network itself. We present a connectivity visualization program called BrainViewer, which we use with a high-resolution mouse cortical connectome. This has the ability to display connectomes from other datasets when they become available and compare spatial connectivity across multiple brain structures. Our tool, optimized for speed and portability, presents a GUI visualization in 2-D top view and flatmap projections, allowing users to select and explore the connections of every source voxel to everywhere else in the cortex. Anatomists and other neuroscientists will find BrainViewer useful for building understanding beyond the known topography of cortical connectivity.

**Index Terms:** Computing methodologies—Modeling and simulation; Human-centered computing—Interactive systems and tools; Human-centered computing—Visualization systems and tools; Human-centered computing—Visualization application domains; Software and its engineering—Software creation and management;

## 1 INTRODUCTION

Connectivity patterns in the brain reveal important information about brain function. Differences in brain-wide connectivity patterns, or *connectomes*, are at the root of some functional and behavioral differences across animals. Improved connectomic knowledge could lead to new treatments for disease, improved brain-machine interfaces, and influence the development of brain-inspired computing and artificial intelligence.

Topography refers to the functional spatial maps across the brain. For instance, it is known that many sensory systems maintain maps of sensory space which are transformed between brain regions. Retinotopy—the mapping of visual space onto visual regions of cortex—is one example of this and was the original inspiration for deep convolutional neural networks which have revolutionized computer vision [2, 11]. It is now evident that retinotopy arises from topographic patterning of connections between brain areas [22], however such data have been difficult to visualize.

Modern neuronal tracing methods allow researchers to trace the major pathways in the brain. Such tracing experiments have been conducted on the mouse [3, 9, 14, 23], fly [15], marmosets

[6, 12, 20, 21], rat [1] and macaque [13]. Results of these experiments have then been used to build brain models at the mesoscale, i.e. showing the connections between areas containing on the order of thousands of neurons. Oh et al. used the anterograde tracer adeno-associated virus expressing green fluorescent protein to track connections in the mouse brain [14]. After injecting mice with the virus, each brain is then sliced, imaged and processed to reveal a volumetric image of fluorescence. In the dataset we use from the Allen Institute for Brain Science, areas are defined using a common coordinate framework and individual experiments are mapped into this common voxel space [19]. Many of these experiments can be combined into a weighted connectivity matrix which stores projection strength between source and target regions [8, 14], and individual tracing experiments can be visualized in 3D using the BrainExplorer tool [10]. Most connectome work has focused on measuring brain connectivity at the level of regions and various tools exist to display such regional connectivity. New methods have been developed to estimate spatially-resolved connectivity at the resolution of voxels—discrete locations within the brain volume [5, 8].

The closest work to our own is the web-based BrainModules from R. Hira [6] that displays functional and structural connectivity patterns from marmoset and mouse in the browser. However, these are only available at low display resolution and the code is undocumented and unmaintained.

We present a tool *BrainViewer* to display and interact with high resolution spatial brain connectivity. We have built it to work with the mouse voxel connectome from [8], but it is portable and can be used across datasets with a number of capabilities described in Section 2. The user may click on any location in right hemisphere of the mouse cortex and see the outgoing projections of neurons from that location. Our tool provides a 2-D visualization of the connectivity in two projections, flatmap and top view; the relative advantages of each are discussed in the methods, Section 3. Switching between the two projections allows for easy comparison, and the user may also navigate the source voxel using their keyboard. This interactive visualization of a spatial connectome enhances the understanding of the brain’s information processing patterns and allow researchers to generate new and interesting topographic anatomical hypotheses.

## 2 BRAINVIEWER: A CONNECTOME VISUALIZER TOOL

Our tool allows users to navigate and explore the estimated mouse cortical connectivity from Knox et al. [8] in a GUI environment (see Fig. 1). The code is available at <https://github.com/glomerulus-lab/brainviewer>. We include the following features:

- A colormap annotating the strength of connectivity from the source voxel to any given voxel in the plot.
- Click-to-plot allows users to select any voxel on the right hemisphere of either the flatmap or top view, visualizing the connectivity from that source voxel.
- Arrow key navigation which enables users to carefully move the source voxel across the cortex.

\*e-mail: daetwis@wwu.edu

†e-mail: reada2@wwu.edu

‡e-mail: stillwj3@wwu.edu

§e-mail: harri267@wwu.edu, corresponding author





## ACKNOWLEDGMENTS

Thank you to Riichiro Hira and Lydia Ng for inspiration and encouragement. This work was a senior capstone project for the first three authors' bachelor's degrees in Computer Science from Western Washington University.

## REFERENCES

- [1] M. Bota, H.-W. Dong, and L. W. Swanson. From gene networks to brain networks. *Nature Neuroscience*, 6(8):795–799, Aug. 2003. doi: 10.1038/nn1096
- [2] K. Fukushima. Neocognitron: A self-organizing neural network model for a mechanism of pattern recognition unaffected by shift in position. *Biological Cybernetics*, 36(4):193–202, Apr. 1980. doi: 10.1007/BF00344251
- [3] R. Gămănuț, H. Kennedy, Z. Toroczka, M. Ercsey-Ravasz, D. Van Essen, K. Knoblauch, and A. Burkhalter. The Mouse Cortical Connectome Characterized by an Ultra Dense Cortical Graph Maintains Specificity by Distinct Connectivity Profiles. *Neuron*, 97(3):698–715.e10, Feb. 2018. doi: 10.1016/j.neuron.2017.12.037
- [4] C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane, J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, and T. E. Oliphant. Array programming with NumPy. *Nature*, 585(7825):357–362, Sept. 2020. doi: 10.1038/s41586-020-2649-2
- [5] K. D. Harris, S. Mihalas, and E. Shea-Brown. High resolution neural connectivity from incomplete tracing data using nonnegative spline regression. In *Neural Information Processing Systems*, 2016.
- [6] R. Hira. BrainModules. <https://web.ece.ucsb.edu/~riichirohira/TopBM-2.06/>. Accessed: 2022-04-21.
- [7] J. D. Hunter. Matplotlib: A 2d graphics environment. *Computing in Science & Engineering*, 9(3):90–95, 2007. doi: 10.1109/MCSE.2007.55
- [8] J. E. Knox, K. D. Harris, N. Graddis, J. D. Whitesell, H. Zeng, J. A. Harris, E. Shea-Brown, and S. Mihalas. High-resolution data-driven model of the mouse connectome. *Network Neuroscience*, 3(1):217–236, Dec. 2018. doi: 10.1162/netn.a.00066
- [9] L. Kuan, Y. Li, C. Lau, D. Feng, A. Bernard, S. M. Sunkin, H. Zeng, C. Dang, M. Hawrylycz, and L. Ng. Neuroinformatics of the Allen Mouse Brain Connectivity Atlas. *Methods*, 73:4–17, Feb. 2015. doi: 10.1016/j.ymeth.2014.12.013
- [10] C. Lau, L. Ng, C. Thompson, S. Pathak, L. Kuan, A. Jones, and M. Hawrylycz. Exploration and visualization of gene expression with neuroanatomy in the adult mouse brain. *BMC Bioinformatics*, 9(1):153, Mar. 2008. doi: 10.1186/1471-2105-9-153
- [11] Y. LeCun, Y. Bengio, and G. Hinton. Deep learning. *Nature*, 521(7553):436–444, May 2015. doi: 10.1038/nature14539
- [12] P. Majka, T. A. Chaplin, H.-H. Yu, A. Tolpygo, P. P. Mitra, D. K. Wójcik, and M. G. Rosa. Towards a comprehensive atlas of cortical connections in a primate brain: Mapping tracer injection studies of the common marmoset into a reference digital template. *Journal of Comparative Neurology*, 524(11):2161–2181, 2016. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/cne.24023>. doi: 10.1002/cne.24023
- [13] N. T. Markov, M. M. Ercsey-Ravasz, A. R. Ribeiro Gomes, C. Lamy, L. Magrou, J. Vezoli, P. Misery, A. Falchier, R. Quilodran, M. A. Gariel, J. Sallet, R. Gamanut, C. Huissoud, S. Clavagnier, P. Giroud, D. Sappey-Mariniere, P. Barone, C. Dehay, Z. Toroczka, K. Knoblauch, D. C. Van Essen, and H. Kennedy. A Weighted and Directed Interareal Connectivity Matrix for Macaque Cerebral Cortex. *Cerebral Cortex (New York, NY)*, 24(1):17–36, Jan. 2014. doi: 10.1093/cercor/bhs270
- [14] S. W. Oh, J. A. Harris, L. Ng, B. Winslow, N. Cain, S. Mihalas, Q. Wang, C. Lau, L. Kuan, A. M. Henry, M. T. Mortrud, B. Ouellette, T. N. Nguyen, S. A. Sorensen, C. R. Slaughterbeck, W. Wakeman, Y. Li, D. Feng, A. Ho, E. Nicholas, K. E. Hirokawa, P. Bohn, K. M. Joines, H. Peng, M. J. Hawrylycz, J. W. Phillips, J. G. Hohmann, P. Wahnoutka, C. R. Gerfen, C. Koch, A. Bernard, C. Dang, A. R. Jones, and H. Zeng. A mesoscale connectome of the mouse brain. *Nature*, 508(7495):207–214, Apr. 2014. Number: 7495 Publisher: Nature Publishing Group. doi: 10.1038/nature13186
- [15] C.-T. Shih, O. Sporns, S.-L. Yuan, T.-S. Su, Y.-J. Lin, C.-C. Chuang, T.-Y. Wang, C.-C. Lo, R. J. Greenspan, and A.-S. Chiang. Connectomics-based analysis of information flow in the *Drosophila* brain. *Current biology: CB*, 25(10):1249–1258, May 2015. doi: 10.1016/j.cub.2015.03.021
- [16] D. C. Van Essen. Cartography and Connectomes. *Neuron*, 80(3):775–790, Oct. 2013. doi: 10.1016/j.neuron.2013.10.027
- [17] D. C. Van Essen, M. F. Glasser, D. L. Dierker, and J. Harwell. Cortical parcellations of the macaque monkey analyzed on surface-based atlases. *Cerebral Cortex (New York, N.Y.: 1991)*, 22(10):2227–2240, Oct. 2012. doi: 10.1093/cercor/bhr290
- [18] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. J. Carey, Í. Polat, Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E. A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020. doi: 10.1038/s41592-019-0686-2
- [19] Q. Wang, S.-L. Ding, Y. Li, J. Royall, D. Feng, P. Lesnar, N. Graddis, M. Naemi, B. Facer, A. Ho, T. Dolbeare, B. Blanchard, N. Dee, W. Wakeman, K. E. Hirokawa, A. Szafer, S. M. Sunkin, S. W. Oh, A. Bernard, J. W. Phillips, M. Hawrylycz, C. Koch, H. Zeng, J. A. Harris, and L. Ng. The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. *Cell*, 181(4):936–953.e20, May 2020. doi: 10.1016/j.cell.2020.04.007
- [20] A. Watakabe, H. Skibbe, K. Nakae, H. Abe, N. Ichinohe, J. Wang, M. Takaji, H. Mizukami, A. Woodward, R. Gong, J. Hata, H. Okano, S. Ishii, and T. Yamamori. Connectional architecture of the prefrontal cortex in the marmoset brain. Technical report, bioRxiv, Dec. 2021. Section: New Results Type: article. doi: 10.1101/2021.12.26.474213
- [21] A. Woodward, T. Hashikawa, M. Maeda, T. Kaneko, K. Hikishima, A. Iriki, H. Okano, and Y. Yamaguchi. The Brain/MINDS 3D digital marmoset brain atlas. *Scientific Data*, 5(1):180009, Feb. 2018. Number: 1 Publisher: Nature Publishing Group. doi: 10.1038/sdata.2018.9
- [22] J. Zhuang, L. Ng, D. Williams, M. Valley, Y. Li, M. Garrett, and J. Waters. An extended retinotopic map of mouse cortex. *eLife*, 6:e18372, Jan. 2017. doi: 10.7554/eLife.18372
- [23] B. Zingg, H. Hintiryan, L. Gou, M. Y. Song, M. Bay, M. S. Bienkowski, N. N. Foster, S. Yamashita, I. Bowman, A. W. Toga, and H.-W. Dong. Neural Networks of the Mouse Neocortex. *Cell*, 156(5):1096–1111, Feb. 2014. Publisher: Elsevier. doi: 10.1016/j.cell.2014.02.023