General spherical harmonic bra-ket overlap integrals of trigonometric functions

Giuseppe Lingetti[‡]

Dipartimento di Fisica, "Sapienza" Università di Roma & Sezione INFN Roma1, Piazzale Aldo Moro 5, 00185, Roma, Italy

Paolo Pani

Dipartimento di Fisica, "Sapienza" Università di Roma & Sezione INFN Roma1, Piazzale Aldo Moro 5, 00185, Roma, Italy

Abstract. Closed formulas in terms of double sums of Clebsch-Gordan coefficients are computed for the evaluation of bra-ket spherical harmonic overlap integrals of a wide class of trigonometric functions. These analytical expressions can find useful application in problems involving non-separable wave equations, e.g. general-relativistic perturbation theory, electromagnetism, quantum theory, etc, wherein the overlap integrals arise from the coupling among different angular modes.

1. Introduction

The spherical harmonic decomposition is a widely used tool in perturbation theory. Applications are countless and include any linear theory (e.g., electromagnetism [1], quantum mechanics [2], Newtonian gravity [3]), and any perturbative scheme of nonlinear theory (e.g., black-hole perturbation theory within and beyond General Relativity (GR) [4]). The spherical harmonic decomposition finds its main advantage for spherically symmetric systems in order to separate the perturbation equations, but it can also be applied to less symmetric configurations. While the angular modes decouple from each other in spherical symmetry – leading to lower-dimensional differential equations – they typically mix when spherical symmetry is broken. This mode-mixing gives rise to bra-ket integrals that must be explicitly evaluated [4].

A relevant class of these braket integrals involves trigonometric functions:

$$\langle l_1, m_1 | e^{ik\phi} \cos(n\theta) | l_2, m_2 \rangle = \tag{1}$$

$$= \int_{0}^{2\pi} d\phi \int_{0}^{\pi} d\theta \sin \theta Y_{l_1,m_1}^*(\theta,\phi) e^{ik\phi} \cos(n\theta) Y_{l_2,m_2}(\theta,\phi) ,$$

$$\langle l_1, m_1 | e^{ik\phi} \sin(n\theta) | l_2, m_2 \rangle = \tag{2}$$

Due to the fact that $e^{ik\phi} \cos(n\theta)$ and $e^{ik\phi} \sin(n\theta)$ form a complete Fourier basis for $(\theta, \phi) \in [0, \pi] \times [0, 2\pi)$, from the general solutions of Eqs. (1) and (2) in principle one can directly compute the bra-kets of any trigonometric function regular in such interval §. Overlap integrals of this type naturally arise from the spherical harmonic decomposition of wave equations, such as those coming from GR perturbation theory [4]. Important examples are bosonic perturbations of spinning black holes [5,6], or the linear dynamics of "dirty" black holes in GR [7] and in extended theories of gravity [8–10]. In the easiest cases having the analytical results helps in understanding the type of couplings involved: for example from the result of $\langle l_1, m_1 | \cos^2 \theta | l_2, m_2 \rangle$, one can easily deduce that scalar perturbations of a Kerr black hole have couplings between l and $l \pm 2$ modes [5]. In the most cumbersome cases, instead, the number of different couplings can be much higher and/or show involved expressions [4,6], thus closed-form solutions can greatly speed up computations.

To the best of our knowledge, in the literature there is no general, analytical, solution to the bra-ket trigonometric integrals (1) and (2). In this short note we provide a closed-form general solution by exploiting the properties of ultraspherical polynomials and their relationships with trigonometric functions and spherical harmonics.

2. Evaluation of the bra-ket integrals

We recall that spherical harmonics are defined as [11]

$$Y_{l,m}(\theta,\phi) = \sqrt{\frac{(2l+1)(l-m)!}{4\pi(l+m)!}} P_l^m(\cos\theta) e^{im\phi},$$
(3)

where P_l^m are the Legendre associated functions, and consequently their complex conjugate is $Y_{l,m}^* = (-1)^m Y_{l,-m}$. Moreover, we can express their products in the following way [11]:

$$Y_{l_1,m_1}Y_{l_2,m_2} = \sum_{l=|l_1-l_2|}^{l_1+l_2} \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2l+1)}} \langle l_1,0,l_2,0|l,0\rangle$$
(4)
 $\times \langle l_1,m_1,l_2,m_2|l,m_1+m_2\rangle Y_{l,m_1+m_2},$

where $\langle l_1, m_1, l_2, m_2 | l, m \rangle$ are the Clebsch-Gordan coefficients.

By exploiting these properties, bra-kets (1) and (2) can be written as

$$\langle l_1, m_1 | e^{ik\phi} \cos(n\theta) | l_2, m_2 \rangle =$$
(5)

§ In principle, through a Fourier expansion, one could also apply the general solutions to angular functions which are not strictly trigonometric. However, the feasibility of such approach clearly depends on how easily the Fourier expansions can be computed for the problem at hand. General spherical harmonic bra-ket overlap integrals of trigonometric functions

$$=\sum_{l=|l_1-l_2|}^{l_1+l_2} \sqrt{\frac{\left(l_1+\frac{1}{2}\right)\left(l_2+\frac{1}{2}\right)\left(l+k\right)!}{(l-k)!}} (-1)^{m_1} \langle l_1, 0, l_2, 0 | l, 0 \rangle$$

$$\times \langle l_1, -m_1, l_2, m_2 | l, -k \rangle \int_0^{\pi} d\theta \sin \theta \cos(n\theta) P_l^{-k}(\cos \theta) ,$$

$$\langle l_1, m_1 | e^{ik\phi} \sin(n\theta) | l_2, m_2 \rangle = (6)$$

$$=\sum_{l=|l_1-l_2|}^{l_1+l_2} \sqrt{\frac{\left(l_1+\frac{1}{2}\right)\left(l_2+\frac{1}{2}\right)\left(l+k\right)!}{(l-k)!}} (-1)^{m_1} \langle l_1, 0, l_2, 0 | l, 0 \rangle$$

$$\times \langle l_1, -m_1, l_2, m_2 | l, -k \rangle \int_0^\pi d\theta \sin \theta \sin(n\theta) P_l^{-k}(\cos \theta) \, .$$

Functions $\cos(n\theta)$ and $\sin(n\theta)$ can be expressed as finite linear combinations of associated Legendre polynomials:

$$\cos(n\theta) = \sum_{l=0}^{n} a_l^n P_l^0(\cos\theta), \qquad \sin(n\theta) = \sum_{l=0}^{n} b_l^n P_l^1(\cos\theta)$$
(7)

From the definition of the associated Legendre functions [12], we have $P_l^0(\cos\theta) = P_l(\cos\theta)$ and $P_l^1(\cos\theta) = \frac{d}{d\theta} [P_l(\cos\theta)]$, where P_l are the Legendre polynomials. Thus, by taking the derivative of the first equation in (7) with respect to θ and comparing it with the second equation, we get

$$b_l^n = -\frac{1}{n}a_l^n \,. \tag{8}$$

One can therefore focus only on the first equation in (7). Functions $\cos(n\theta)$ are Čebyšëv polynomials $T_n(\cos \theta) = \cos(n\theta)$ [13], hence we can exploit this property for finding the a_l^n coefficients. Functions $T_n(x)$ and $P_l(x)$ are in fact special cases of ultraspherical polynomials $C_n^{\gamma}(x)$ [13]:

$$T_n(x) = \lim_{\gamma \to 0} \frac{n + 2\gamma}{2\gamma} C_n^{\gamma}(x) , \qquad P_l(x) = C_l^{1/2}(x) , \qquad (9)$$

Different types of ultraspherical polynomials can be related through the following expression [13], which is a special case of the connection relation for Jacobi polynomials :

$$C_n^{\gamma}(x) = \sum_{j=0}^{\lfloor n/2 \rfloor} \frac{(\gamma - \beta)_j (\gamma)_{n-j}}{j! (\beta + 1)_{n-j}} \left(\frac{\beta + n - 2j}{\beta}\right) C_{n-2j}^{\beta}(x), \qquad (10)$$

where $(x)_y = \frac{\Gamma(x+y)}{\Gamma(x)}$ is the Pochhammer's symbol for $x, y \in \mathbb{C}$ and $\Gamma(x)$ is Euler's gamma function. From Eqs. (9) and (10) we get the expressions for Čebyšëv polynomials as linear combinations of Legendre polynomials

$$T_n(x) = -\sum_{j=0}^{\lfloor n/2 \rfloor} \frac{n\Gamma(j-1/2)\Gamma(n-j)}{8j!\Gamma(3/2+n-j)} (1+2n-4j)P_{n-2j}(x),$$
(11)

Ultraspherical polynomials are a special case of Jacobi polynomials [13].

and consequently, by using Eq. (8), we can rewrite Eq. (7) as

$$\cos(n\theta) = -\sum_{j=0}^{\lfloor n/2 \rfloor} \frac{n\Gamma(j-1/2)\Gamma(n-j)}{8j!\Gamma(3/2+n-j)} \left(1+2n-4j\right) P_{n-2j}^{0}(\cos\theta), \quad (12)$$

$$\sin(n\theta) = \sum_{j=0}^{\lfloor n/2 \rfloor} \frac{\Gamma(j-1/2) \Gamma(n-j)}{8j! \Gamma(3/2+n-j)} (1+2n-4j) P_{n-2j}^1(\cos\theta).$$
(13)

By using these last results, we can write the general parametric expressions for the bra-kets (5) and (6) in closed form:

$$\langle l_1, m_1 | e^{ik\phi} \cos(n\theta) | l_2, m_2 \rangle =$$

$$= \sum_{l=|l_1-l_2|}^{l_1+l_2} \sum_{j=0}^{\lfloor n/2 \rfloor} \frac{(-1)^{m_1+1}}{2} \frac{n\Gamma(j-1/2)\Gamma(n-j)}{8j!\Gamma(3/2+n-j)} \sqrt{\frac{(2l_1+1)(2l_2+1)(l+k)!}{(l-k)!}} \\ \times (1+2n-4j) \langle l_1, -m_1, l_2, m_2 | l, -k \rangle \langle l_1, 0, l_2, 0 | l, 0 \rangle I(n-2j, 0, l, -k) ,$$

$$\langle l_1, m_1 | e^{ik\phi} \sin(n\theta) | l_2, m_2 \rangle =$$

$$= \sum_{l=|l_1-l_2|}^{l_1+l_2} \sum_{j=0}^{\lfloor n/2 \rfloor} \frac{(-1)^{m_1}}{2} \frac{\Gamma(j-1/2)\Gamma(n-j)}{8j!\Gamma(3/2+n-j)} \sqrt{\frac{(2l_1+1)(2l_2+1)(l+k)!}{(l-k)!}} \\ \times (1+2n-4j) \langle l_1, -m_1, l_2, m_2 | l, -k \rangle \langle l_1, 0, l_2, 0 | l, 0 \rangle I(n-2j, 1, l, -k) ,$$

where we have $I(l, m, l', m') = \int_{-1}^{1} dx P_l^m(x) P_{l'}^{m'}(x)$. This last integral can be explicitly performed [14]:

$$I(l,m,l',m') = \sqrt{\frac{(l+m)!(l'+m')!}{(l-m)!(l'-m')!}} \sum_{j=|l-l'|}^{l+l'} \sqrt{\frac{(j-m-m')!}{(j+m+m')!}} \times \langle l,m,l',m'| \ j,m+m' \rangle \langle l,0,l',0| \ j,0 \rangle I_0(j,m+m') ,$$

where for m > 0 we have

$$I_0(l,m) = \int_{-1}^1 dx P_l^m(x) = \frac{[(-1)^m + (-1)^l] 2^{m-2} m \Gamma(l/2) \Gamma((l+m+1)/2)}{((l-m)/2)! \Gamma((l+3)/2)}$$

Because of $P_0^0(\cos\theta) = 1$, for l, m = 0 we have $I_0(l, m) = 2$, while for m < 0 we can exploit $I_0(l, m) = (-1)^m \frac{(l+m)!}{(l-m)!} I_0(l, -m)$ from the properties of P_l^m . Expression (14) gives an indeterminate result for n = 0, therefore, by taking into account that $\cos(0\theta) = 1 = P_0^0(\cos\theta)$, for this case we need to use the following expression:

$$\langle l_1, m_1 | e^{ik\phi} | l_2, m_2 \rangle = \sum_{l=|l_1-l_2|}^{l_1+l_2} \frac{(-1)^{m_1}}{2} \sqrt{\frac{(2l_1+1)(2l_2+1)(l+k)!}{(l-k)!}}$$

$$\times \langle l_1, -m_1, l_2, m_2 | l, -k \rangle \langle l_1, 0, l_2, 0 | l, 0 \rangle I_0(l, -k) .$$

$$(16)$$

3. Conclusions

We derived the general analytical expressions for the spherical harmonic overlap integrals (1) and (2) of a wide class of trigonometric functions, from which solutions for all continuous trigonometric functions can be easily derived. We expressed the results in closed form in terms of double sums of Clebsch-Gordan coefficients. These results can be useful in the study of wave equations for non-separable systems, e.g. in perturbed quantum mechanical problems or in the context of spherical harmonic decompositions within black-hole perturbation theory.

Acknowledgments

G.L. is grateful to Massimo Vaglio and Enrico Cannizzaro for useful discussions on the topic. We acknowledge the financial support provided under the European Union's H2020 ERC, Starting Grant agreement no. DarkGRA–757480. We also acknowledge support under the MIUR PRIN Grant 2020KR4KN2 "String Theory as a bridge between Gauge Theories and Quantum Gravity" and FARE programmes (GW- NEXT, CUP: B84I20000100001).

Bibliography

- J. D. Jackson, Classical electrodynamics; 2nd ed. Wiley, New York, NY, 1975. https://cds.cern.ch/record/100964.
- J. J. Sakurai, Modern quantum mechanics; rev. ed. Addison-Wesley, Reading, MA, 1994. https://cds.cern.ch/record/1167961.
- [3] E. Poisson and C. M. Will, Gravity: Newtonian, Post-Newtonian, Relativistic. Cambridge University Press, 2014.
- [4] P. Pani, "Advanced Methods in Black-Hole Perturbation Theory," Int. J. Mod. Phys. A 28 (2013) 1340018, arXiv:1305.6759 [gr-qc].
- S. R. Dolan, "Superradiant instabilities of rotating black holes in the time domain," *Phys. Rev. D* 87 (Jun, 2013) 124026.
 https://link.aps.org/doi/10.1103/PhysRevD.87.124026.
- [6] D. Baumann, H. S. Chia, J. Stout, and L. ter Haar, "The spectra of gravitational atoms," Journal of Cosmology and Astroparticle Physics 2019 no. 12, (Dec, 2019) 006–006. https://doi.org/10.1088/1475-7516/2019/12/006.
- [7] A. Dima and E. Barausse, "Numerical investigation of plasma-driven superradiant instabilities," *Classical and Quantum Gravity* 37 no. 17, (Aug, 2020) 175006. https://doi.org/10.1088/1361-6382/ab9ce0.
- [8] V. Cardoso, I. P. Carucci, P. Pani, and T. P. Sotiriou, "Black holes with surrounding matter in scalar-tensor theories," *Phys. Rev. Lett.* 111 (Sep, 2013) 111101. https://link.aps.org/doi/10.1103/PhysRevLett.111.111101.
- [9] V. Cardoso, I. P. Carucci, P. Pani, and T. P. Sotiriou, "Matter around Kerr black holes in scalar-tensor theories: scalarization and superradiant instability," *Phys. Rev. D* 88 (2013) 044056, arXiv:1305.6936 [gr-qc].
- [10] G. Lingetti, E. Cannizzaro, and P. Pani, "Superradiant instabilities by accretion disks in scalar-tensor theories," arXiv:2204.09335 [gr-qc].

- [11] M. Rose, Elementary Theory of Angular Momentum. New York, 1957. https://books.google.it/books?id=MyEFzQEACAAJ.
- [12] Series Expansions of Arbitrary Functions, ch. 2, pp. 48-111. John Wiley and Sons, Ltd, 1989. https://onlinelibrary.wiley.com/doi/abs/10.1002/9783527617210.ch2.
- [13] M. E. H. Ismail, Classical and Quantum Orthogonal Polynomials in One Variable. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2005.
- [14] H. Mavromatis and R. Alassar, "A generalized formula for the integral of three associated legendre polynomials," *Applied Mathematics Letters* 12 no. 3, (1999) 101-105. https://www.sciencedirect.com/science/article/pii/S0893965998001803.