

Fractional quantum Hall effect in semiconductor systems

Zlatko Papić^{a,*}, Ajit C. Balram^{b,c}

^a*School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, United Kingdom*

^b*Institute of Mathematical Sciences, CIT Campus, Chennai 600113, India*

^c*Homi Bhabha National Institute, Training School Complex, Anushaktinagar, Mumbai 400094, India*

ARTICLE INFO

Keywords:

fractional quantum Hall effect, Laughlin wave function, composite fermions, Moore-Read state, non-Abelian statistics, anyons, topological quantum computation

Abstract

The fractional quantum Hall (FQH) effect refers to the strongly-correlated phenomena and the associated quantum phases of matter realized in a two-dimensional gas of electrons placed in a large perpendicular magnetic field. In such systems, topology and quantum mechanics conspire to give rise to exotic physics that manifests via robust quantization of the Hall resistance. In this chapter, we provide an overview of the experimental phenomenology of the FQH effect in GaAs-based semiconductor materials and present its theoretical interpretations in terms of trial wave functions, composite fermion quasiparticles, and enigmatic non-Abelian states. We also highlight some recent developments, including the parton theory and the Dirac composite fermion field theory of FQH states, the role of anisotropy and geometrical degrees of freedom, and quantum entanglement in FQH fluids.

1. Key objectives

- describe the rich phenomenology exhibited by two-dimensional electrons hosted in a semiconductor system placed in a perpendicular magnetic field;
- overview of different theoretical approaches for describing fractional quantum Hall (FQH) phases of matter;
- introduce the concept of non-Abelian FQH states and their potential applications in fault-tolerant topological quantum computation;
- summarize recent developments in the field, including parton theory of the FQH effect, geometric properties of FQH states, entanglement-based and field-theoretic approaches to understanding FQH systems.

2. Notations and acronyms

2DEG: two-dimensional electron gas

IQHE: integer quantum Hall effect

FQHE: fractional quantum Hall effect

LL: Landau level

LLL: lowest Landau level

SLL: second Landau level

PH: particle-hole

CF: composite fermion

MR: Moore-Read

HLR: Halperin-Lee-Read

ν : filling factor

B : magnetic field strength

$\ell = \sqrt{\hbar/eB}$: magnetic length

$z = x - iy$: complex number representation of the electron coordinates (x, y) in the 2D plane

Φ_n : wave function of n filled Landau levels

$\Phi_1 \equiv \prod_{i < j} (z_i - z_j)$: Laughlin-Jastrow factor

P_{LLL} : projection to the lowest Landau level

3. Introduction

The fractional quantum Hall (FQH) effect occurs in a system of electrons confined to two spatial dimensions and a strong magnetic field perpendicular to the two-dimensional (2D) plane. In such systems, the kinetic energy is completely quenched by the magnetic field and Coulomb interaction exerts non-perturbative effects on the behavior of the electrons, typically without analogs in systems of weakly-interacting electrons encountered elsewhere in condensed matter physics. In this chapter, we do not aim to present a historical overview of the FQH effect, but rather highlight the key theoretical concepts for its understanding and to illustrate the connections between a variety of different approaches that have been deployed to tackle the problem. Perhaps most surprisingly, first-quantized wave functions have turned out to be highly valuable tools for understanding FQH phenomena, in contrast to the second-quantized field theories widely used in other areas of many-body physics. To emphasize this unique aspect of the FQH effect, our presentation will focus on introducing physics from the perspective of trial wave functions, motivated by experimental observations.

Before we dive into its fascinating experimental phenomenology, it is instructive to present a high-level description of the core of FQH physics. The following "Theory of Everything" Hamiltonian governing a FQH system should be familiar to most physicists:

$$H = \sum_j \frac{1}{2m_b} \left[p_j + \frac{e}{c} \mathbf{A}(\mathbf{r}_j) \right]^2 + \sum_j U(\mathbf{r}_j) + g\mu_B \mathbf{B} \cdot \mathbf{S} + \frac{e^2}{4\pi\epsilon} \sum_{j < k} \frac{1}{|\mathbf{r}_j - \mathbf{r}_k|}. \quad (1)$$

The first line contains the one-body terms in the problem: the kinetic energy in the presence of a constant external magnetic field $\mathbf{B} = \nabla \times \mathbf{A}$; the potential U incorporating the lattice, disorder and confinement effects; finally, the Zeeman

*Corresponding author

z.papic@leeds.ac.uk (Z. Papić)

ORCID(s): 0000-0002-8451-2235 (Z. Papić); 0000-0002-8087-6015

(A.C. Balram)

energy. The second line is the Coulomb interaction energy between (non-relativistic) electrons.

As a first step, let us simplify the problem down to its bare essentials. While disorder, confinement and lattice effects can be occasionally very important, we will neglect them here and set $U=0$. Most FQH experiments discussed below are performed using GaAs-AlGaAs heterostructures, in which electrons have a band mass of $m_b=0.067m_e$ (where m_e is the electron mass in vacuum), dielectric constant of $\epsilon=12.6\epsilon_0$, and Landé g factor $g=-0.44$. From this, we can estimate the energy required to flip a spin – the Zeeman splitting – to be $E_Z=g\mu_B B \approx 0.3B[\text{T}]\text{K}$, where the magnetic field $B[\text{T}]$ is in units of Tesla. With the exception of Sec. 9, throughout this chapter, we will assume the electron spin is fully polarized due to a large magnetic field.

In order to diagonalize what's left of the Hamiltonian, the usual strategy is to first diagonalize the one-body part of H which gives rise to discrete Landau levels (LLs) of a single electron in a magnetic field. The Landau solution yields two important scales in the problem: the magnetic length

$$\ell = \left(\frac{\hbar}{eB} \right)^{1/2} \approx \frac{25}{\sqrt{B[\text{T}]}} \text{nm}, \quad (2)$$

and the cyclotron energy splitting between LLs

$$\hbar\omega_c = \hbar \frac{eB}{m_b} \approx 20B[\text{T}]\text{K}. \quad (3)$$

On the other hand, the typical Coulomb energy is given by

$$V_C \equiv \frac{e^2}{4\pi\epsilon\ell} \approx 50\sqrt{B[\text{T}]\text{K}}. \quad (4)$$

For the sake of convenience, throughout we shall use natural units where $\ell=1$ and $e^2/(4\pi\epsilon\ell)=1$.

To render the many-body problem as tractable as possible, one often makes a further approximation by choosing $V_C \ll \hbar\omega_c$. In this limit, the Coulomb interaction is too weak to scatter particles across different LLs, i.e., entire dynamics is contained within a single LL and there is no ‘‘LL mixing’’. This assumption is not always warranted and its validity should be tested either numerically or experimentally. Nevertheless, under this approximation, the FQH Hamiltonian takes a remarkably simple form

$$H = \mathcal{P} \left(\frac{e^2}{4\pi\epsilon} \sum_{j < k} \frac{1}{|\mathbf{r}_j - \mathbf{r}_k|} \right) \mathcal{P}, \quad (5)$$

i.e., the Hamiltonian is just the Coulomb interaction which must be diagonalized with the restriction of remaining in a single LL. The latter restriction is formally implemented by the global operator \mathcal{P} which projects out all states that have a component outside the given LL. For most of our discussion below, the LL in question will be the lowest LL (LLL), although we will also mention the second Landau level (SLL) and higher LLs.

The model in Eq. (5) embodies the theoretical essence of the FQH effect: it is believed to capture universal physical properties of broad classes of correlated phases emerging in

the regime of the FQH effect. However, despite its seemingly simple form, it is impossible to analytically diagonalize the Hamiltonian of Eq. (5). The difficulty stems from \mathcal{P} being a global operator defined on a many-electron Hilbert space whose dimension grows exponentially with the number of electrons. Specifically, two main challenges are: (i) Eq. (5), has no parameters whatsoever, including no-small parameters that one might use to attempt a perturbative approach; (ii) If the Coulomb interaction is turned off, one ends up with an exponentially large degeneracy of any LL, again precluding an easy solution.

Broadly speaking, this chapter will focus on the following questions: (i) what are the possible phases of matter that arise from Eq. (5) as one varies the electron density (or, alternatively, the magnetic field)? (ii) what are the universal properties of such phases in the long-wavelength limit, e.g., the types of order and excitations that govern the low-energy physics? (iii) what are the fundamental principles that allow to understand and microscopically model such phases?

4. Phenomenology of the FQH effect

In a typical FQH experiment, one measures the Hall resistance, extracted from the voltage drop perpendicular to the current flowing through a two-dimensional electron gas (2DEG) placed in a perpendicular magnetic field. The 2DEG should have sufficiently low concentration of impurities and high mobility of the electrons. The fabrication of such samples has required major technological advances that we will not touch upon. As found by Tsui, Stormer and Gossard in 1982 Tsui et al. (1982), upon sweeping the magnetic field, the Hall resistance of such a 2DEG exhibits quantization at

$$R_{xy} = \frac{h}{\nu e^2}, \quad (6)$$

where ν was originally found to take the value $1/3$. Subsequently, quantization was observed at other rational values of ν such as $2/3$, $2/5$, $3/7$, etc. On the other hand, ν is related to the electron density ρ and is known as the *filling factor* or *filling fraction*,

$$\nu = \frac{\rho h}{eB}. \quad (7)$$

In an earlier work by von Klitzing, Dorda and Pepper Klitzing et al. (1980), similar resistance quantization had been observed for integral values of ν , i.e., in the regime of integer quantum Hall (IQH) effect. In both cases, the resistance was found to be quantized not only at the special value of ν but also in some neighborhood around the special value, which manifests as a resistance plateau in the experimental trace, as illustrated in the experimental data shown in Fig. 1(a). Furthermore, when R_{xy} is pinned to a plateau, the longitudinal resistance (i.e., the one measured along the direction of the current) exhibits an Arrhenius behavior $R_{xx} \propto \exp(-\Delta/(2k_B T))$, where the energy scale Δ is interpreted as a gap to excitations. R_{xx} vanishes in the limit $T \rightarrow 0$, indicating dissipationless transport. A deep minimum




Figure 1: Experimental measurements of the Hall resistance R_{xy} and the longitudinal resistance R_{xx} as a function of the magnetic field B applied perpendicularly to a 2DEG. (a) Resistance traces in the lowest Landau level from Ref. Pan et al. (2003). Prominent gapped states are shaded in yellow. Other weaker states (without fully developed minimum in R_{xx}) are also indicated. The composite Fermi liquid is shaded purple. (b) In the second Landau level, in addition to gapped states at odd-denominator filling factors, there are also even-denominator states at $\nu=2+1/2$ and $\nu=2+3/8$. Reentrant integer quantum Hall states are shaded in green. Data is reproduced from Ref. Kumar et al. (2010). (c) In higher Landau levels, the FQH plateaus are replaced by states with strongly anisotropic transport, as revealed by the large difference between R_{xx} and R_{yy} . Examples are shaded in red and identified with stripe phases. Reentrant integer quantum Hall states are shaded in green. Data is taken from Ref. Ro et al. (2019).

in R_{xx} is often seen as a precursor of a developing FQH state, however, the definitive observation of a FQH state requires a quantized plateau in R_{xy} .

In both IQH and FQH cases, the R_{xy} quantization is

remarkably accurate to about one part in a billion. This is a manifestation of quantum-mechanical coherence across a semiconductor material on mesoscopic length scales. Moreover, as seen from the appearance of the Planck

constant, the quantization is not described by classical electrodynamics. The latter also predicts that the resistance should vary linearly with the magnetic field. One hint to a topological origin of the phenomenon is that the resistance is quantized, not the resistivity (they both have the same units in two dimensions). Moreover, the Hall quantization was found to be universal: it is independent of the sample type, geometry, and various materials parameters, such as the band mass of the electron or the dielectric constant of the background semiconductor, and is robust to sufficiently weak disorder.

4.1. Incompressible fluids

Performing the transport measurements described above, one obtains remarkable traces of R_{xy} and R_{xx} as a function of the magnetic field B , as we illustrate in Fig. 1 using current state-of-the-art experimental data. At sufficiently low temperatures (typically, in mK range) and very high magnetic fields, the trace is dominated by FQH states which display a quantized plateau in R_{xy} and vanishing R_{xx} . These are signatures of incompressible quantum fluids, which arise from strong Coulomb repulsion between the electrons residing in the partially-filled LLL, as the kinetic energy of the electrons is quenched. FQH quantum fluids are unconventional in that they represent topological phases of matter. While a detailed theory of such topological quantum fluids will be presented in subsequent sections, here we briefly summarize some of their defining properties – see Fig. 2:

- FQH phases are quantum liquids that do not exhibit long-range order for any local order parameter. Thus, FQH phases cannot be described by a Landau-Ginzburg type theory based on spontaneous symmetry breaking Richard E. Prange (1987). For example, the (LL-projected) structure factor of a FQH state, shown in inset of Fig. 2(b), takes the form characteristic of a liquid phase, indicating that the FQH state is uniform in the bulk and distinct from, say, a Wigner crystal.
- The bulk of a FQH fluid exhibits an energy gap Δ , i.e., there is a discontinuity in the chemical potential as one sweeps the magnetic field. Furthermore, there is also a gap for charge-neutral excitations at the fixed magnetic field, as seen in Fig. 2(b). Near the system's boundary, however, there are *gapless* excitations which propagate chirally, in the direction determined by the external magnetic field, see Fig. 2(a). In the semiclassical picture, these excitations can be visualized as “skipping orbits” Halperin (1982), which can also go around weak impurity potentials, avoiding dissipation.
- The ‘topological’ nature of the FQH fluid derives from its sensitivity to the topology of the manifold it lives in. For example, in the infinite plane or when wrapped around a sphere, the fluid has a unique ground state,

but when it is placed on a torus or any higher genus surface, the ground state develops a degeneracy Wen & Niu (1990). At filling factor $\nu=p/q$, with p, q co-prime positive integers, the ground state degeneracy is at least q Haldane (1985). For example, in the $\nu=1/3$ state shown in Fig. 2(b), the ground state is three-fold degenerate.

- By threading magnetic flux or, equivalently, by twisting the torus boundary conditions, the degenerate ground states can mix with one another but remain separated from other excited states at energies $\geq \Delta$. The total Hall conductance carried by the ground state multiplet is proportional to the topological invariant called the Chern number C Thouless et al. (1982). This gives the Hall conductance for each member of the multiplet $\sigma_{xy}=\nu\frac{e^2}{h}$. The topological nature of the Hall conductance explains its robustness at sufficiently low temperatures $k_B T \ll \Delta$.
- Thermal Hall conductance of FQH fluids is given by Kane & Fisher (1997)

$$\kappa_{xy}=c_-[\pi^2 k_B^2/(3h)]T, \quad (8)$$

where the universal coefficient c_- is known as the chiral central charge. This expression holds at temperatures much smaller than the gap and under the assumption that edge modes are fully equilibrated.

- The excitations of incompressible FQH fluids are *anyons* Leinaas & Myrheim (1977), Wilczek (1982): an exotic type of particle which carry a fraction of the electron charge and also have “fractional” exchange statistics Arovas et al. (1984). In particular, for states such as $\nu=1/3$ depicted in Fig. 2(c), the adiabatic exchange of the coordinates of two anyons results in a complex phase, $\Psi_{\text{final}}(z_2, z_1) = e^{i\theta}\Psi_{\text{initial}}(z_1, z_2)$. Bosons and fermions are special cases with $\theta=0$ and $\theta=\pi$, respectively, while Abelian anyons correspond to other real values of θ . Anyons with richer exchange statistics are also possible and will be discussed in Sec. 8. We note that recent experiments have observed statistical properties of the simplest types of Abelian anyons in the $\nu=1/3$ state using mesoscopic interferometry Nakamura et al. (2020) as well as anyon colliders Bartolomei et al. (2020).
- FQH fluids can be described by an emergent Chern-Simons gauge field \mathbf{a} , which attaches magnetic fluxes to particles Zhang et al. (1989). This enables the transmutation of statistics and the existence of anyon excitations. The Chern-Simons field contributes a topological term to the effective action of FQH fluids,

$$\mathcal{L}_{\text{CS}} = \frac{1}{4\pi\nu} \epsilon^{\mu\nu\lambda} a_\mu \partial_\nu a_\lambda, \quad (9)$$

where $\epsilon^{\mu\nu\lambda}$ is the Levi-Civita tensor and we assumed the FQH state is at filling $\nu=1/q$, with Einstein summation convention implied. Thus, the effective low-energy theory of FQH fluids is a variant of topological quantum field theory (see Sec. 10.2 for further discussion).




Figure 2: Incompressible fluids and topological order. (a) Transport experiment probing a FQH state. Robust quantization of the Hall resistance is associated with a current-carrying edge state (yellow), which is insensitive to a low concentration of impurities (depicted in purple). In a semiclassical picture, the edge state forms a narrow one-dimensional channel that propagates ballistically in one direction. (b) On a torus, the $\nu=1/3$ FQH ground state has a three-fold topological degeneracy and carries zero momentum $\mathbf{K}=0$, while the rest of the energy spectrum is separated by a gap. Low energy charge-neutral excitations form a collective mode. Data is obtained using DiagHam open source package DiagHam (n.d.) for exact diagonalization of the Coulomb interaction projected to the lowest LL for $N \leq 12$ electrons. The same plot includes data for both square and hexagonal unit cells, illustrating the robustness of the low-energy physics to geometric details. Inset shows the projected guiding-center structure factor $S_0(\mathbf{k})$, which reveals a liquid ground state. (c) Elementary charged excitations of a FQH fluid behave as anyons: by taking two anyons around each other, the resulting wave function Ψ_{final} is not simply equal to $\pm \Psi_{\text{initial}}$, as it would be for bosons or fermions.

The properties summarized in this section are believed to be shared by all incompressible FQH states, which are

primarily realized in the LLL and SLL, recall Fig. 1(a) and (b). However, as we explain below, the microscopic mechanisms giving rise to these topological fluids can be very different. In particular, the FQH states in the LLL are markedly different from those in the SLL. In contrast to the majority of gapped FQH liquids presented in Fig. 1(a), we note that there also exists a *gapless* liquid state which is similar to the Fermi liquid state in an ordinary 2DEG in zero magnetic field. This state occurs near $\nu=1/2$ where one observes a smooth dependence of both the longitudinal and Hall resistances in Fig. 1(a). Nevertheless, this Fermi-liquid-like state has many exotic properties of its own and it will be discussed separately in Sec. 7.1.

We note here that FQH states extending on either side of $\nu=1/2$ are related by an operation that maps electrons to holes and vice versa, i.e., it relates states at ν with $1-\nu$. This particle-hole (PH) symmetry is exact in the limit $B \rightarrow \infty$. In this limit, any two-body interaction, like the Coulomb one, results in eigenstates that are perfectly PH symmetric. We will return to the consequences of PH symmetry in Secs. 7.3 and 10.2.

4.2. Broken symmetry phases

In addition to incompressible liquids, other types of correlated phases of matter can occur in partially filled LLs, primarily in the regime of low filling factors or in higher LLs, see Fig. 1(b)-(c). While interactions still underpin the formation of such phases, these phases spontaneously break the translation symmetry of the 2D plane by forming crystals or stripe patterns. Such phases are typically weakly-correlated and they have been successfully understood using the more traditional methods, such as Hartree-Fock theory.

4.2.1. Wigner crystals in the lowest Landau level

“Classical” electrons at sufficiently low densities in 2D form a triangular Wigner crystal (WC) to minimize the Coulomb repulsion between them. Strong magnetic field quenches the kinetic energy of the electrons, leaving only interactions to decide the fate of the system. This setting was believed to be ideal to stabilize the WC phase and provided one of the motivations for the Tsui-Stormer-Gossard experiment Tsui et al. (1982).

As it turned out, the FQH liquid has lower energy than WC at high filling factors Yoshioka et al. (1983). However, at very low filling factors, when the density of electrons is sufficiently small, a WC is expected to be stabilized. Recent studies suggest that the FQH fluid can prevail over the WC at filling factors as low as $\nu=1/7$ Pan et al. (2002), Chung et al. (2022). Furthermore, the WC that is stabilized at $\nu < 1/7$ is not that of electrons, but of electron-vortex composites called composite fermions (CFs) Archer et al. (2013), Zuo et al. (2020), that we will introduce in Sec. 6.2.

While direct signatures of the Wigner solid phase have remained elusive, experiments have observed a number of indirect signatures. The presence of a WC has been probed by studying the commensurability oscillations of CFs in a nearby layer Deng et al. (2016). Collective oscillations of the crystalline domains of the WC resulting from disorder

give rise to “pinning resonances” in the frequency-dependent conductivity which have been detected experimentally Williams et al. (1991), Zhu et al. (2010). The melting behavior of the magnetic-field-induced WC suggests that it harbors quantum correlations and is markedly different from the classical WC at zero-field Ma et al. (2020).




Figure 3: Some examples of symmetry-broken phases realized in a two-dimensional electron gas placed in a strong perpendicular magnetic field.

4.2.2. Stripes, nematics and bubble crystals in higher LLs

In higher LLs, interactions can also engender various phases that spontaneously break certain symmetries while exhibiting integral quantization of the Hall resistance, as seen in Fig. 1(b) and (c). In the vicinity of integral fillings, if disorder is sufficiently weak, the additional electrons or holes are expected to form a WC. Using Hartree-Fock theory, Koulakov, Fogler, and Shklovskii Koulakov et al. (1996), Fogler et al. (1996) predicted that the extra electrons or holes in a high LL have a tendency to form a WC, a “bubble” phase – a WC of bubbles of electrons, where each bubble hosts multiple electrons, or stripes of alternating fillings, depending on the filling factor. For example, when a high LL is nearly half full (i.e., the filling factor is $\nu = n + 1/2$ with n being a sufficiently large integer), strong transport anisotropy is observed, as shown in Fig. 1(c), consistent with the formation of stripes Du et al. (1999), Lilly et al. (1999). The direction of the anisotropy aligns with one of the crystal axes, due to the explicit symmetry breaking via the host lattice. Moreover, the anisotropy can flip by 90° as a function of the in-plane magnetic field Lilly et al. (1999) or the density Zhu et al. (2002). Many re-entrant transitions are seen in higher LLs Eisenstein et al. (2002), Xia et al. (2004), indicating competition between numerous approximately degenerate ground states Shibata & Yoshioka (2001) with a common feature of integral Hall resistance quantization.

5. Laughlin states

5.1. Laughlin’s wave function

Soon after the experimental observation of the FQH plateau at $\nu = 1/3$, Laughlin constructed a novel ansatz to describe the electronic ground state Laughlin (1983). Laughlin’s wave function is given by:

lin’s wave function is given by:

$$\Psi_{\nu=1/(2p+1)}^{\text{Laughlin}} = \prod_{i < j} (z_i - z_j)^{2p+1} \exp \left[- \sum_k \frac{|z_k|^2}{4\ell^2} \right] \equiv \Phi_1^{2p+1}, \quad (10)$$

where $z_k = x_k - iy_k$ parametrizes the 2D coordinate of the k^{th} electron as a complex number, and p is a positive integer ($p=1$ for $\nu=1/3$). The $\nu=1$ IQH wave function, $\Phi_1 = \prod_{i < j} (z_i - z_j)$, is referred to as the Laughlin-Jastrow factor. To simplify the notation, below we shall often drop the ubiquitous Gaussian factor and the subscript “ ν ” from the wave functions.

The high power of the Jastrow factor in the Laughlin wave function builds additional correlations in the many-body state (beyond those mandated by the Pauli principle) which keeps the electrons away from each other and thus minimizes their Coulomb repulsion. Because of this, one may expect the wave function in Eq. (10) to provide a good variational description of the exact ground state of the Hamiltonian in Eq. (5). Quite remarkably, it turned out that the overlap between the Laughlin wave function and the exact ground state of Eq. (5) for as many as $N=15$ electrons is consistently above 98%, despite an exponential increase in the size of the Hilbert space with N and the lack of any variational parameters in Eq. (10). This suggests that the Laughlin wave function captures all the essential correlations of the $\nu=1/3$ ground state even in the thermodynamic limit. Indeed, due to the “nice” form of the Laughlin wave function, many of its properties such as energy, density, pair-correlation function etc. have been studied in large systems using the Monte Carlo method Morf & Halperin (1986). The (projected) structure factor of the $\nu=1/3$ state, shown in inset of Fig. 2(b), confirms that the state describes an incompressible liquid.

Laughlin’s theory furthermore predicts that the FQH effect can also occur at $\nu=1/5$ and a quantized Hall plateau at this filling was observed in experiments. Subsequent numerical simulations have shown that the Laughlin wave functions have high overlaps with the exact LLL Coulomb ground small systems at $\nu=1/3$, $1/5$ and $1/7$ Fano et al. (1986), d’Ambrumenil & Reynolds (1988). For $\nu < 1/7$, a Laughlin liquid is expected to give way to a crystalline state Lam & Girvin (1984). By particle-hole conjugation in the LLL, Laughlin’s theory also explains the plateaux observed at fractions $\nu=1-1/(2p+1)$ which can be understood as a $1/(2p+1)$ Laughlin state of “holes”.

5.2. Charged excitations

Laughlin also constructed wave functions for the charged excitations in this system. Laughlin’s wave function for the positively charged excitation, referred to as a *quasihole* (qh), located at a position η , is given by:

$$\Psi_{1/(2p+1)}^{\text{Laughlin,qh}} = \prod_i (z_i - \eta) \Psi_{1/(2p+1)}^{\text{Laughlin}}, \quad (11)$$

while that of the negatively charged *quasiparticle* (qp) is:

$$\Psi_{1/(2p+1)}^{\text{Laughlin,qp}} = \prod_i (2\partial_{z_i} - \bar{\eta}) \Psi_{1/(2p+1)}^{\text{Laughlin}}. \quad (12)$$

In Eq. (12) the derivatives only act on the polynomial part of the wave function and not on the Gaussian factor. To motivate these wave functions, consider the situation where $\eta=0$. The polynomial factor $\prod_i z_i$ pushes electrons away from the origin (the wave function of Eq. (11) vanishes at η) thereby depleting the electron density at the origin resulting in the creation of a positively charged quasihole. The derivatives, on the other hand, pull electrons in towards the origin which leads to a build-up of an excess charge at the origin resulting in the creation of a negatively charged quasiparticle.

The probability density corresponding to Laughlin's ground state and quasihole wave functions can be interpreted as the partition function of a one-component classical plasma in 2D Laughlin (1987). Using this plasma analogy, Laughlin showed that the quasihole excitation carries a fractional charge of $e/(2p+1)$ (we denote the electronic charge by $-e$). Via PH conjugation, one can readily see that the quasiparticle excitation also carries a fractional charge of value $(-e)/(2p+1)$. Shot-noise experiments have verified the presence of $e/3$ charged quasiparticles in the $\nu=1/3$ state Saminadayar et al. (1997), Dolev et al. (2008). Moreover, recent experiments Bartolomei et al. (2020), Nakamura et al. (2020) have confirmed that the quasihole and quasiparticle excitations of the Laughlin state are Abelian anyons with a statistical phase $\theta=2\pi/3$.

5.3. Magnetoroton excitation

Along with charged excitations, any FQH system, including the Laughlin states, also hosts "neutral" excitations which can be viewed as arising from a combination of an equal number of quasiparticles and quasiholes. Girvin, MacDonald, and Platzman (GMP) Girvin et al. (1985) proposed a general ansatz for the lowest-lying neutral excitation as a density wave riding on the ground state. The GMP wave function is written as $\Psi_{\nu,\mathbf{k}}^{\text{GMP}} = \hat{\rho}_{\mathbf{k}} \Psi_{\nu}$, where $\hat{\rho}_{\mathbf{k}}$ is the LLL projected density operator and Ψ_{ν} is the ground state wave function at filling ν . The GMP excitation is termed a *magnetoroton* since its dispersion has a characteristic minimum at a finite wavenumber reminiscent of the roton minimum in the excitation spectrum of liquid Helium. The GMP ansatz gives a reasonable account of the exact low-lying neutral excitation at wavenumbers near and below the roton minimum, $k\ell \lesssim 1$ He et al. (1994). Resonant inelastic light-scattering experiments have mapped out the magnetoroton branch of excitations at $\nu=1/3$ Pinczuk et al. (1993).

5.4. Edge excitations

As we sketched in Fig. 2(a), due to the insulating bulk, the conduction in FQH states occurs in the vicinity of the system's boundary via chiral edge channels. In the FQH effect, electron correlations are strong and one must model the edge channels using the Luttinger liquid description Wen (1990). Theoretical description of FQH edge states is

provided by bosonization, which conveniently expresses the electronic degrees of freedom in terms of bosons. Due to chirality, backscattering in wide samples is minimized and the edge channels are found to propagate ballistically across long distances, with the dispersion $\epsilon_k = \nu k$, where ν is the edge speed and k is the momentum along the edge.

In real samples, however, the edge may undergo a phenomenon of "reconstruction" Chklovskii et al. (1992), Chamon & Wen (1994) where instead of a single channel, the edge consists of compressible and incompressible stripes that form multiple parallel conducting channels, possibly counterpropagating. Unlike the insulating bulk, the physics of the edge is thus far more prone to non-universal effects due to the long-range tail of the Coulomb interaction and details of the confining potential near the boundary of the system.

The edge spectral function can be measured by tunneling an electron laterally into the edge. Wen's theory Wen (1990) predicts a nonlinear current-voltage, $I \propto V^m$, relation at zero temperature, with the edge exponent taking the quantized value $m=2p+1$ for the $\nu=1/(2p+1)$ Laughlin states. Thus, the quantized edge exponent distinguishes a FQH edge from an ordinary Luttinger liquid, whose edge exponent varies continuously with the strength of the interaction. The edge exponent was measured in experiments by Chang et al. Chang et al. (1996) who found its value to be reasonably close to 3 in the $\nu=1/3$ state. However, various non-universal effects have also been observed in these experiments Chang (2003). This makes the study of FQH edges challenging even in the simplest fractions such as $\nu=1/3$.

5.5. Parent Hamiltonian and Haldane pseudopotentials

An important step in validation of Laughlin's theory was Haldane's identification of a parent Hamiltonian for the state in Eq. (10) Haldane (1983). Any homogeneous and isotropic pairwise interaction potential $V(|z_i - z_j|)$, with the particles restricted to a single LL, can be written as Duncan & Haldane (1990)

$$H_{\text{int}} = \sum_{i < j} V(|z_i - z_j|) = \sum_m \sum_{i < j} V_m \hat{P}_{ij}^m, \quad (13)$$

where \hat{P}_{ij}^m projects onto a state in which particles i and j have relative angular momentum m . In the LLL, \hat{P}_{ij}^m projects on the relative state $\psi_{\text{rel}} \sim (z_i - z_j)^m$ of any pair of particles. The numbers V_m are called the Haldane pseudopotentials and they represent an energy penalty for two particles to form a state with relative angular momentum m . For spinless electrons, only odd values of m can appear as ψ_{rel} must be fully antisymmetric in electrons' coordinates.

The parent Hamiltonian of the $\nu=1/3$ Laughlin state is obtained by setting $V_1 > 0$ and all other $V_{m \geq 3} = 0$. This interaction assigns positive energy to any wave function that vanishes as $\psi_{\text{rel}} \sim (z_i - z_j)$, but it allows pairs of particles to form the state $\psi_{\text{rel}} \sim (z_i - z_j)^3$ for free. The latter is just the $p=1$ Laughlin state in Eq. (10), which emerges as the unique and

densest state with exactly zero energy for the V_1 pseudopotential interaction. The quasihole excitations of the Laughlin state, Eq. (11), similarly appear as exact zero-energy states of the same Hamiltonian. Numerics on finite systems show that any other state in the spectrum is separated by a gap on the order $\Delta \propto V_1$ but the persistence of this spectral gap in the thermodynamic limit has not been rigorously proven.

Haldane pseudopotentials greatly facilitate numerical modelling of FQH systems. For example, open-source libraries for exact diagonalization of FQH systems such as DiagHam DiagHam (n.d.) take full advantage of the pseudopotential formalism. On the analytical side, an arbitrary combination of pseudopotentials can be conveniently converted to a real space potential Trugman & Kivelson (1985), which enables systematic studies of parent Hamiltonian of more complex FQH states Simon, Rezayi & Cooper (2007).

6. Hierarchy and composite fermions

Following the discovery of the $\nu=1/3$ state, a zoo of more than a hundred FQH plateaux have been observed, many of which do not fit Laughlin's theory (see Fig. 1). These fractions predominantly belong to the sequence $\nu=n/(2pn\pm1)$ [and their hole-conjugates at $1-n/(2pn\pm1)$], where n and p are positive integers (see Fig. 1). Consequently, new theoretical approaches had to be developed to describe such states. In this chapter we focus on two leading paradigms for understanding the various fractions observed the lowest LL: the Haldane-Halperin hierarchy and Jain's composite fermion theory.

6.1. Haldane-Halperin hierarchy

Haldane Haldane (1983) and Halperin Halperin (1984) proposed a hierarchical construction to obtain FQH states at arbitrary odd-denominator fractions. In the hierarchical construction, one starts with a nearby parent Laughlin state and views the filling factor difference from it as stemming from the presence of additional Laughlin quasiparticles/quasiholes. Assuming that interaction between the additional Laughlin quasiparticles/quasiholes is sufficiently strongly repulsive at short distances and weak in comparison to the parent Laughlin state, one condenses these excitations into Laughlin-like "daughter" FQH states. Iterating such a construction, one can produce candidate FQH states at all odd-denominator fractions – see Fig. 4 for the fractions originating from the $\nu=1/3$ Laughlin state in the first three generations of a hierarchy.

A trial wave function for the first daughter state at filling factor $\nu=1/(2p+1 - q^{-1})$, where q is a positive even integer, with N electrons and N/q Laughlin quasiparticles is constructed as follows:

$$\int \left[\prod_{k=1}^{N/q} d^2\eta_k \right] [\Phi_1^*(\{\bar{\eta}_k\})]^q \Psi_{1/(2p+1)}^{\text{Laughlin,qps}}(\{z\}; \{\eta_k\}), \quad (14)$$

where Φ_1 is the usual Laughlin-Jastrow factor and the term $\Psi_{1/(2p+1)}^{\text{Laughlin,qps}}$ is the Laughlin quasiparticle state of Eq. (12)

generalized to multiple quasiparticles located at positions η_k . Analogous wave functions for the first daughter state at $\nu=1/(2p+1 + q^{-1})$ can be constructed from Laughlin's quasihole wave function of Eq. (11). The same idea can be used to construct wave functions for daughter states further down in the hierarchy.

For small systems at $\nu=2/5$ and $3/7$, the hierarchical wave functions have been constructed numerically and these have high overlaps with the exact LLL Coulomb ground states Greiter (1994). Studies of larger systems and other filling factors are required to substantively test these wave functions. The order of fractions and their stability predicted by the hierarchy theory is not entirely consistent with experiments. For example, the $3/7$ and $5/13$ states appear at the same level in the hierarchy (see Fig. 4) but the FQH effect at $3/7$ is much stronger than that at $5/13$ as is evidenced from their gaps [see Fig. 1(a)]. Nonetheless, universal aspects, such as the charge and braiding statistics of the quasiparticles, predicted by the Haldane-Halperin construction are consistent with experiments.




Figure 4: Daughter fractions obtained from the Haldane-Halperin hierarchy originating from the parent Laughlin state at $\nu=1/3$. Experimentally observed fractions in the lowest Landau level are circumscribed.

6.2. Composite fermions

Jain's theory Jain (1989a) provides a conceptually different microscopic explanation of the $\nu=n/(2pn\pm1)$ FQH states as IQH states of certain emergent topological particles called *composite fermions* (CFs). A composite fermion is a bound state of an electron and an even number ($2p$) of vortices. Due to the vortex-binding, the CFs see a reduced magnetic field $B^*=B-2p\rho\phi_0$ compared to the external magnetic field B seen by the electrons [superscript * is used to denote CF quantities]. The + (parallel) and – (anti-parallel) signs in $\nu=n/(2pn\pm1)$ indicate the direction of the residual magnetic field seen by the CFs with respect to the external magnetic field. In the zeroth-order approximation, the CFs are assumed to be non-interacting, and just like free electrons in a magnetic field, CFs form their own Landau-like levels, called *Lambda levels* (Λ Ls). When an integer number $\nu^*=n$ of these Λ Ls are fully filled, we get an FQH effect of electrons at $\nu=n/(2pn\pm1)$. The existence of Λ Ls has been established in experiment Kang et al. (1993). Recently, in an ultra-high mobility sample, FQH states all the way up to $n=16$ in the $n/(2n\pm1)$ Jain sequence have been observed Chung et al. (2021).

The mapping to an IQH effect leads to the Jain wave

function for the FQH ground state at $\nu=n/(2pn\pm1)$ Jain (1989a)

$$\Psi_{n/(2pn\pm1)}^{\text{Jain}} = \mathcal{P}_{\text{LLL}} \Phi_1^{2p} \Phi_{\pm n}. \quad (15)$$

Here Φ_n is the Slater determinant state of n filled LLs of electrons, while the Jastrow factor Φ_1^{2p} attaches $2p$ vortices to each of the electrons to convert them into CFs, see Fig. 5 for a schematic illustration. For $n=1$ (with the + sign) the Jain wave function is identical to the Laughlin wave function of Eq. (10). For $n>1$, the Jain wave function needs to be explicitly projected to the LLL, as denoted by \mathcal{P}_{LLL} .

Beyond ground state properties, the IQH mapping readily allows the construction of wave functions for the excitations as well, see Fig. 5. The positively charged quasihole called a CF hole is obtained by removing a CF from the topmost filled LL. The negatively charged quasiparticle, referred to as a CF particle, is obtained by adding a CF in the topmost empty LL. The lowest-lying neutral excitation, called the CF-exciton, is obtained by combining a single CF particle and a CF hole. The excitations obtained using CF theory generally exceed the accuracy of other approaches, e.g., the Laughlin and GMP ansatz discussed in Sec. 5, in describing the exact LLL Coulomb excited states.

We note here that although the Haldane-Halperin hierarchical states are topologically equivalent to CF states Read (1990), Wen (1995), the two theories propose different microscopic mechanisms for the origin of FQH effect Jain (2014). Unlike the hierarchy wave functions, the CF wave functions in Eq. (15) have proven amenable to large-scale numerical simulations Jain & Kamilla (1998). These studies have demonstrated that the CF theory provides a highly-accurate description of the majority of FQH states observed in the LLL. For example, neutral excitation spectrum observed using surface acoustic waves Kukushkin et al. (2009) is in good agreement with CF theory Scarola et al. (2000). Nevertheless, some quantitative discrepancy between the experimentally measured Boebinger et al. (1985), Willett et al. (1988), Du et al. (1993), Pan et al. (2020), Villegas Rosales et al. (2021) and theoretically predicted Zhang & Das Sarma (1986), Ortalan et al. (1997), Morf et al. (2002) charge gaps persists to this day. This likely arises from neglecting the effects of the disorder, interface roughening, and screening by gates in theoretical studies. Furthermore, theoretical calculations use simplified models to incorporate the effects of LL mixing and finite-width. More realistic calculations are needed to quantitatively explain the gaps obtained in experimental studies. Finally, certain delicate states, like those observed at $\nu=4/11$ and $5/13$ Samkharadze et al. (2015), Pan et al. (2015), Fig. 1(a), and the states seen in the SLL, Fig. 1(b), lie beyond the paradigm of *weakly-interacting* CFs presented in this section. To understand such states, one needs other approaches, as discussed in subsequent sections below.




Figure 5: Schematic representation of CF ground state, elementary charged and neutral excitations at $\nu=3/7$.

7. Half-filled Landau level

The simplest fraction – half-filling of a LL – has remained in the focus of research since the early days of the FQH effect. The conundrum is that no quantized FQH plateau is observed when it is the LLL that is half-filled, i.e., at $\nu=1/2$ – see Fig. 1(a). Yet, when the *second* LL is half-filled, resulting in a total filling factor $\nu=5/2$, a robust FQH plateau is seen instead [Fig. 1(b)]. In this section, we discuss a novel kind of gapless Fermi-liquid like state that was introduced to describe the $\nu=1/2$, and its pairing instability that is believed to give rise to the exotic gapped state at $\nu=5/2$.

7.1. Gapless composite fermion Fermi liquid in the half-filled Landau level

The $\nu=1/2$ state finds a natural interpretation in the CF theory since at half-filling CFs see no residual magnetic field. Therefore, like electrons at zero field, CFs can form a Fermi liquid state, called the composite Fermi liquid (CFL) Kalmeyer & Zhang (1992), Halperin et al. (1993), which is gapless and thus does not lead to a FQH effect. The CFL is a markedly different state from the ordinary Fermi sea: in the LLL, there is no kinetic energy and interactions alone conspire to create an emergent Fermi-liquid. The wave function for the CFL at even-denominator fillings $\nu=1/(2p)$ is obtained as the $n\rightarrow\infty$ limit of the wave function in Eq. (15):

$$\Psi_{1/(2p)}^{\text{CFL}} = \mathcal{P}_{\text{LLL}} \Phi^{\text{FL}} \Phi_1^{2p}, \quad \Phi^{\text{FL}} \equiv \det [e^{i\vec{k}_i \cdot \vec{r}_m}]. \quad (16)$$

Halperin, Lee, and Read (HLR) Halperin et al. (1993) evaluated many properties of the CFL using a Chern–Simons field theory in which flux quanta are bound to electrons to turn them into CFs. This description is in qualitative agreement with numerous experiments that have verified the presence of the CFL Kang et al. (1993), Willett et al. (1993), Goldman et al. (1994). In Sec. 10.2 we shall discuss one particular experiment, namely the measurement of the CF Fermi wave vector using geometric resonance Willett et al. (1999), Kamurov et al. (2014), Hossain et al. (2020), which has renewed the interest in the CFL state.

7.2. $\nu=5/2$ state

The observation of a FQH state at the even-denominator filling $\nu=5/2$ Willett et al. (1987) indicated that not all FQH

states can be described using the ideas discussed in Sec. 6. Moore and Read Moore & Read (1991) identified conformal field theory (CFT) as an innovative tool for generating trial FQH wave functions (see Ref. Hansson et al. (2017) for an in-depth overview of this technique). Using a particular type of CFT, Moore and Read arrived at the following wave function for a state at $\nu=1/2$:

$$\Psi_{1/2}^{\text{MR-Pf}} = \text{Pf} \left(\frac{1}{z_i - z_j} \right) \prod_{1 \leq i < j \leq N} (z_i - z_j)^2, \quad (17)$$

where the number of particles N is assumed to be even. The key component of this wave function is the Pfaffian – the antisymmetrized sum over pairs:

$$\text{Pf} \left(\frac{1}{z_i - z_j} \right) = \mathbb{A} \left(\frac{1}{z_1 - z_2} \frac{1}{z_3 - z_4} \dots \right), \quad (18)$$

where \mathbb{A} represents the anti-symmetrization operator. The Pfaffian factor makes the overall wave function antisymmetric and therefore a valid trial state for electrons. As first suggested by Greiter, Wen and Wilczek Greiter et al. (1992), this wavefunction can provide a description of the experimental $\nu=5/2$ resistance plateau, provided one assumes that the two lowest LLs of \uparrow and \downarrow spins are completely filled with electrons. Thus, Eq. (17) describes the correlations between fully spin-polarized electrons in the valence (half-filled) SLL.

The Pfaffian factor introduces weak pairing between the CFs Read & Green (2000). Indeed, in the antisymmetrized product over pairs, we recognize the real-space form of the wave function describing a 2D Bardeen-Cooper-Schrieffer (BCS) superconductor. Thus, we can view Eq. (17) as the FQH analog of a $p_x + ip_y$ superconductor. The physical origin of pairing is due to the effective Coulomb interaction being less repulsive at short distances in the SLL compared to LLL. Thus, forming CFs overscreens the Coulomb repulsion resulting in a weak residual attractive interaction between them which leads to a pairing instability of their CFL. One consequence of the electron pairing is an unconventional collective excitation on top of the Moore-Read ground state that was dubbed the “neutral fermion” mode Bonderson et al. (2011), Möller et al. (2011).

The interest in the Moore-Read state was initially fuelled by numerical simulations. Exact diagonalizations performed by Morf Morf (1998) showed that the spin-polarized state at $\nu=5/2$ had lower energy than any partially polarized or spin-singlet state, even in the absence of any Zeeman splitting. Furthermore, it was shown that the state was poised towards a compressible phase Rezayi & Haldane (2000) and can be easily pushed across the phase boundary by slightly varying the Hamiltonian in a way that mimics the effect of an in-plane magnetic field, accounting for experimental observations Eisenstein et al. (1990), Lilly et al. (1999), Pan et al. (1999). The correspondence between numerics and experiment has been made more quantitative by comparisons of the energy gap obtained from exact diagonalization Morf et al. (2002) and DMRG

simulations Feiguin et al. (2008) with the one measured in experiments Kumar et al. (2010).

7.3. Anti-Pfaffian

While PH symmetry is exact for the FQH problem restricted to a single LL, the Moore-Read state in Eq. (17) is not invariant under PH transformation. The state obtained by PH conjugation applied to Eq. (17) is a topologically-distinct phase of matter that was named the “anti-Pfaffian” (aPf) Levin et al. (2007), Lee et al. (2007). The two states have different thermal Hall responses and different edge excitations, which immediately leads to the question: which state describes the $\nu=5/2$ ground state in the thermodynamic limit?

Distinguishing between the aPf and Moore-Read states has proven a considerable challenge. The key to discriminating between the two states is LL mixing, which explicitly breaks PH symmetry and lowers the energy of one of these states with respect to the other. Unfortunately, accurate modeling of LL mixing effects comes with many challenges, in particular its perturbative treatment Bishara & Nayak (2009), Sodemann & MacDonald (2013), Simon & Rezayi (2013), Pakrouski et al. (2015) cannot be rigorously justified. Nevertheless, as recently shown by Rezayi Rezayi (2017), different theoretical approaches converge to the conclusion that LL mixing favors the aPf state.

On the other hand, experimental studies have detected signatures of $e/4$ quasiparticles at $\nu=5/2$ using the interference of edge channels Willett et al. (2009). Shot noise measurements Dolev et al. (2010) and charge-sensing in the bulk using single-electron transistor Venkatachalam et al. (2011) have also reported evidence of $e/4$ quasiparticles. These results are consistent with both Moore-Read and aPf states. The dependence on voltage and temperature of the tunneling current was measured at a point contact in the $\nu=5/2$ state and found to be consistent with the aPf state Radu et al. (2008). Furthermore, upstream neutral modes have been detected at $\nu=5/2$ Bid et al. (2010), Dolev et al. (2011) which is consistent only with the aPf state.

8. Non-Abelian states

Contrasting the case of SLL with that of the LLL in Fig. 1(a)-(b), one notices a few striking differences. One of the most prominent incompressible states in the SLL has an even denominator, $\nu=2+1/2$ Willett et al. (1987) and its strength is roughly comparable with $\nu=2+1/3$, which is expected to be analogous to the $\nu=1/3$ Laughlin state in the LLL Balram et al. (2013). Moreover, FQH effect is also observed at $\nu = 2+3/8$ Xia et al. (2004), Kumar et al. (2010). Even the odd-denominator FQH states in the SLL appear out of order when compared with the LLL sequence: FQH effect has been observed at $1/3$ and $2/5$ in the LLL, but no FQH effect has been confirmed at $3/7$, $4/9$ and $5/11$ but evidence for a plateau at $2+6/13$ exists Kumar et al. (2010). To explain these unconventional fractions, different theoretical approaches have been put forward. A common theme in these approaches are quasiparticles with

non-Abelian braiding statistics, qualitatively different from the Abelian anyons we encountered in Sec. 4.1.

8.1. Non-Abelian anyons and topological quantum computation

Beyond the paired ground state wave function, the quasi-hole excitations of the Moore-Read state in Eq. (17) are particularly interesting: they must be created in pairs and they have non-Abelian braiding statistics. Imagine that $2n$ quasi-holes have been added to the Moore-Read state by increasing the magnetic flux by n flux quanta. In non-Abelian FQH states, fixing the configuration of the quasi-holes does not result in a unique quantum state – instead, there is a degenerate manifold of states. For example, in the Moore-Read state the $2n$ -quasi-hole manifold is 2^{n-1} -fold degenerate and is spanned by some orthogonal basis $\{\psi_1, \psi_2, \dots, \psi_{2^{n-1}}\}$. The braiding of two quasi-holes, depicted in Fig. 2(c), acts as a unitary transformation on this subspace, $\psi_\alpha = U_{\alpha\beta} \psi_\beta$ Nayak & Wilczek (1996). As matrices U do not generally commute, the quasi-holes in the Moore-Read state obey ‘non-Abelian’ braiding statistics.

The degenerate states ψ_α can be interpreted as basis states of a logical qubit, and the quasi-hole braiding operations U can be mapped to the action of quantum gates Freedman et al. (2003). These operations are topologically protected at sufficiently low temperatures $k_B T \lesssim \Delta_{MR}$, where Δ_{MR} is the excitation gap of the Moore-Read state. This is because the qubit states are locally indistinguishable and any decoherence induced by local perturbations will be strongly suppressed. This idea of employing a topologically-protected manifold of non-Abelian FQH quasi-holes to encode and manipulate a quantum bit is known as *topological quantum computation* Nayak et al. (2008).

While the prospect of avoiding the need for error correction in a quantum computer is of immense practical benefit, the particular system of Moore-Read anyons discussed above falls short of this goal as it is unable to perform so-called universal quantum computation. The simplest FQH system capable of universal topological quantum computation is discussed in Sec. 8.2. At this stage, however, topological quantum computation with FQH states remains a theoretical concept, although recent progress in FQH interferometry Willett et al. (2021), Nakamura et al. (2020) paves the way to a potential demonstration of an operating FQH qubit. We note that special types of one-dimensional nanowires can host boundary excitations with properties reminiscent of Moore-Read anyons Kitaev (2001), offering a potentially more practical realization of topological quantum computation. The efforts towards topological quantum computation will rely critically on better understanding and better control of FQH excitations, which may be aided by STM techniques Papić et al. (2018).

8.2. $\nu=12/5$ state

Universal quantum computation can be performed with more complex types of anyons that emerge in the Read-Rezayi (RR) sequence of states Read & Rezayi (1999).

These non-Abelian states generalize the notion of pairing between pairs of particles in the Moore-Read state to a clustering of larger groups (three or more particles). The RR states occur at filling factors $\nu=k/(k+2)$, with $k=1, 2$ being the Laughlin and Moore-Read states, respectively. The $k=3$ RR state is a qualitatively new non-Abelian state hosting the so-called Fibonacci anyons. This state is often referred to as \mathbb{Z}_3 RR state because its wavefunction was originally constructed using a \mathbb{Z}_3 parafermion conformal field theory Read & Rezayi (1999).

Experimental observation of the $\nu=12/5$ FQH plateau by Xia et al. Xia et al. (2004) and subsequently Kumar et al. Kumar et al. (2010) has generated much excitement due to the possibility that it may be described by the PH conjugate of the \mathbb{Z}_3 RR state. Numerical work Read & Rezayi (1999) indicated that the Coulomb ground state at $\nu=12/5$ is close to a phase transition between the Abelian CF state and the PH conjugate of \mathbb{Z}_3 RR state. More recent studies using DMRG Mong et al. (2017) have found that the ground state at $\nu=12/5$ is fully spin-polarized and in the vicinity to a charge-density-wave ordered phase. The latter was found to be much more sensitive to LL mixing at $\nu=13/5$ than at $\nu=12/5$ Pakrouski et al. (2016), accounting for the absence of a quantized plateau at $\nu=13/5$. Due to a much smaller gap in the $\nu=12/5$ state and the scarcity of its experimental realizations, the definitive determination of its nature remains much more challenging compared to $\nu=5/2$.

8.3. Other non-Abelian states

Previously mentioned RR states are a small subset of a family of states associated with the so-called Jack polynomials Feigin et al. (2002). As first realized by Bernevig and Haldane Bernevig & Haldane (2008), the wave function of many FQH states at filling factors $\nu=k/(k+r)$ can be identified with a single Jack polynomial, parametrized by integers k, r and a so-called ‘root’ state. In the case $r=2$ the Jacks coincide with the RR states discussed in Sec. 8.2. An interesting case with $k=2$ and $r=3$ corresponds to the so-called ‘Gaffnian’ state Simon, Rezayi, Cooper & Berdnikov (2007), which has inspired much theoretical work as a potential example of a gapless non-Abelian state Read (2009a). The recursive properties of the Jack polynomials impose a great deal of analytical structure on FQH wave functions, a fact that has been fruitfully employed in numerical computations Bernevig & Regnault (2009).

Bonderson and Slingerland Bonderson & Slingerland (2008) constructed trial wave functions for many of the filling factors where FQH states have been observed in the SLL. The Bonderson-Slingerland approach obtains FQH wave functions as a product of the bosonic version of the Moore-Read state and a composite fermion state. In particular, this approach yields a state that is competitive with the (PH conjugate of) \mathbb{Z}_3 RR state at $\nu=12/5$ Bonderson et al. (2012). On the other hand, the observed filling factor $2+6/13$ Kumar et al. (2010) is not a part of this sequence. Other constructions, based on condensing non-Abelian quasiparticles of the Moore-Read state Hermanns (2010)




Figure 6: Multicomponent FQH effects. (a) Transitions between CF states with different spin polarization (solid dots) as in-plane magnetic field is varied. The lines are fits from the free CF theory, with the g^* factor and the effective mass m^* treated as empirical parameters. B_{eff} is the effective magnetic field. Data from Ref. Du et al. (1995). (b) Phase diagram of GaAs wide quantum wells at $\nu_T=1/2$ as a function of d/ℓ and $\alpha=\Delta_{\text{SAS}}/(e^2/4\pi\epsilon\ell)$. Region corresponding to an incompressible FQH state is shaded in yellow. The interlayer distance d and layer width ℓ are estimated numerically from the self-consistent charge distribution. Figure reproduced from Ref. Shabani et al. (2013). (c) Exciton condensation in $\nu_T=1$ bilayers. Tunneling conductance dI/dV vs interlayer voltage V for a range of total densities N_T , corresponding to $1.6 \lesssim d/\ell \lesssim 2.3$. Trace A, at the highest density, shows a deep suppression of the tunneling near zero bias. By trace D, the lowest density of the four shown, this suppression has been replaced by a tall peak, signalling exciton condensation. The vertical scale is the same for all traces. Figure reproduced from Ref. Spielman et al. (2000).

or by taking products of the bosonic RR states with the Laughlin state Jolicoeur (2007), have also been proposed. The hierarchical construction of Levin and Halperin Levin & Halperin (2009) may account for the FQH effect observed at filling $\nu=2+6/13$. Somewhat surprisingly, the Levin-Halperin states turn out to be Abelian, even though they derive from a non-Abelian Moore-Read parent state. Most of these states, however, result in complicated wave functions that have not been systematically tested in numerical simulations.

9. Multicomponent fractional quantum Hall effect

Up to this point, our discussion was restricted to fully spin-polarized electrons, as is appropriate for the setting when the external magnetic field is large. It turns out that the Zeeman splitting E_Z (see Sec. 3) is strongly suppressed in GaAs: the ratio of E_Z to the Coulomb energy scale for a typical magnetic field $B=9\text{T}$ is $\xi=E_Z/(e^2/(4\pi\epsilon\ell))\approx 0.02$. Experimentally, it is possible to tune ξ by tilting the sample or introducing a parallel magnetic field, suppressing the g -factor using hydrostatic pressure Leadley et al. (1997), Nicholas et al. (1998) or changing the 2DEG's density. Therefore, for certain parameter regimes, it could be favorable for a fraction of the electrons to flip their spin leading to the formation of unpolarized states Chakraborty & Zhang (1984), Chakraborty (2000). We now turn our attention to such “multicomponent” FQH states where the

internal degree of freedom allows the electrons to form new kinds of correlated states without analogs in single-layer FQH systems.

Apart from ordinary spin, the valley degree of freedom could also lead to multi-component FQHE and this plays a particularly important role in materials such as graphene. Here, we just mention that multi-component FQHE has been also observed in multi-valley semiconductor systems such as AlAs Bishop et al. (2007), Padmanabhan et al. (2009). Furthermore, it has also been possible to fabricate two single-layer 2DEGs in close proximity to each other, such that the two layers assume the role of \uparrow, \downarrow spin states. The advantage of studying such bilayer FQH systems is the enhanced tunability of the interactions: one can vary both the tunneling Δ_{SAS} between the two layers, as well as the effective interlayer distance d/ℓ . Note that, conveniently, the physical distance d can be fixed and one varies ℓ by tuning the overall magnetic field (while also adjusting the electron density to remain at the same filling factor). The parameter d/ℓ directly modifies the interlayer Coulomb interaction, which has a strong effect on many-electron states. In experiment, two main architectures were pursued: wide single quantum wells in which electrostatic effects produce a ground subband wave function with a pronounced dumbbell shape Suen et al. (1992) and true double quantum wells consisting of two thin GaAs layers embedded in the alloy $\text{Al}_x\text{Ga}_{1-x}\text{As}$ Eisenstein et al. (1992).

Theoretical understanding of multicomponent FQH states is guided by Halperin's generalization of the Laughlin

wave function in Eq. (10) to FQH systems with a discrete internal quantum number Halperin (1983). Halperin's states are parametrized by non-negative integers m , m' and n and their wave functions are given by

$$\Psi^{(m,m',n)} = \prod_{i<j} (z_i - z_j)^m \prod_{i<j} (w_i - w_j)^{m'} \prod_{i,j} (z_i - w_j)^n, \quad (19)$$

where $\{z_k\}$ and $\{w_k\}$ are the coordinates of the electrons in the "top" and "bottom" layer (or spin state), respectively. The exponents m and m' must be odd integers due to Pauli exclusion principle, while n does not have such a restriction. Moreover, because intra-species correlations are typically stronger than inter-ones we must have $m, m' \geq n$. The state in Eq. (19) corresponds to the filling factor $\nu_T = (m+m'-2n)/(mm'-n^2)$. Most of the time, one is interested in the balanced case where the two layers or spin species are equivalent, i.e., $m = m'$, which can be physically interpreted as each layer forming a $\nu=1/m$ Laughlin state, with interlayer correlations controlled by exponent n .

We note that the Halperin (m, m, n) states can be generalized to bilayer CF states, where the $\nu=1/m$ Laughlin states in each layer are replaced by a Jain state Scarola & Jain (2001). Furthermore, in the case of spin, the CF wave function in Eq. (15) can be directly generalized to accommodate the spin degree of freedom by filling up n_\uparrow LLs with \uparrow -spin electrons (and similarly n_\downarrow LLs filled with electrons of opposite spin) Wu et al. (1993). Depending on n_\uparrow and n_\downarrow , states with full or partial spin polarization can be readily constructed.

9.1. Spinful systems

In single-layer FQH systems, as ξ is varied, transitions between states with different spin polarization can take place and have been extensively studied in experiment, see an illustrative example in Fig. 6(a). For example, at $\nu=4/7$, three CF states can be constructed: a) fully polarized with $n_\uparrow=3$ and $n_\downarrow=0$, b) partially polarized with $n_\uparrow=3$ and $n_\downarrow=1$, and c) spin-singlet with $n_\uparrow=2=n_\downarrow$. At large ξ , the ground state is fully polarized. As ξ is lowered, the fully polarized state transitions to a partially polarized state and eventually at small ξ goes into a spin-singlet state. At the Laughlin fractions, even at zero Zeeman splitting, the ground state is fully spin polarized due to exchange interactions, an effect that has been dubbed quantum Hall ferromagnetism Sondhi et al. (1993).

In the presence of spin, PH symmetry relates states at ν to $2-\nu$. Experimentally in GaAs, spin transitions have predominantly been studied at fractions in the range $1 < \nu < 2$ Clark et al. (1989), Du et al. (1995). The reason for this is at larger fillings, the magnetic field is smaller and thus the starting Zeeman energy is lower which allows access to the least polarized states. Then by tilting the sample, Zeeman energy can be increased and states with higher polarizations can be accessed. The spin phase diagram of the Jain fractions has been worked out in detail both theoretically Park & Jain (1998), Balram, Töke, Wójs & Jain (2015) and experimentally Du et al. (1995), Kukushkin et al. (1999), Liu

et al. (2014) and these are in semi-quantitative agreement with each other.

In the presence of spin, the excitations of fully polarized ground states can also change character. In Secs. 5.3 and 6.2 we discussed excitations which are spin-conserving. However, at low values of ξ , the lowest-lying excitation could involve a spin-flip. At the Laughlin fractions, this lowest-lying spin-flip mode, termed the spin-wave, has energy identical to the Zeeman splitting in the long-wavelength limit Kallin & Halperin (1984). In the fully polarized CF states the spin-wave mode has a roton minimum Mandal & Jain (2001) which lies below the ground state at zero-Zeeman splitting. The presence of these low-lying excitations is indicative of the fact that as ξ is lowered the fully polarized ground state could become unstable giving way to another state with lower polarization.

More exotic excitations involving spin-flip such as skyrmions Sondhi et al. (1993), Fertig et al. (1994), that carry a topological spin texture, can also be stabilized at low values of ξ in the vicinity of a Laughlin fraction. These skyrmions can be viewed as CF particles or CF holes dressed by spin waves. The exchange interaction between CFs is much weaker than that between electrons, so it is a lot harder to realize and detect fractional skyrmions compared to their integer counterparts. Nevertheless, Zeeman-energy dependence of certain excitations does point to the existence of neutral spin-textures in the vicinity of $\nu=1/3$ which have been interpreted as arising from pair(s) of skyrmions and anti-skyrmions Groshaus et al. (2008). More direct evidence for their existence has been obtained from the measurement of the binding energy of fractional skyrmions in the vicinity of $\nu=1/3$ using light-scattering experiments Balram, Wurstbauer, Wójs, Pinczuk & Jain (2015).

9.2. Bilayer systems

The first experimental evidence for unusual states in bilayer systems was the observation of a quantized Hall plateau at $\nu_T=1/2$ in a bilayer system with $d/\ell \lesssim 3$, in stark contrast to a single-layer FQH system at the same filling Suen et al. (1992), Eisenstein et al. (1992). Numerical calculations Chakraborty & Pietiläinen (1987), Yoshioka et al. (1989), He et al. (1993), verified the existence of an incompressible state and identified it with the (3,3,1) Halperin state in Eq. (19). More recently, there has been some renewed interest in the $\nu_T=1/2$ two component systems Shabani et al. (2013, 2009) due to the possible transition into the Moore-Read state as tunneling is increased Ho (1995), Peterson et al. (2010), Zhao et al. (2021). An example of the phase diagram in a wide quantum well at $\nu_T=1/2$ is reproduced in Fig. 6(b).

Perhaps the most remarkable bilayer quantum Hall phenomenon is the exciton condensation observed at total filling $\nu_T=1$ Spielman et al. (2000), Tutuc et al. (2004). The key to demonstrating the existence of excitons was the measurement of interlayer tunneling conductance at zero bias, which revealed a sharp change in the behavior depending on d/ℓ .

When d/ℓ is sufficiently small, one observes a pronounced Josephson-like peak Spielman et al. (2000), see Fig. 6(c). The peak reflects the fact that there is no energy cost associated with the transfer of an electron from one layer to the other. Furthermore, one can also probe the system in a so-called “drag” configuration, where the current is driven through one of the layers and the voltage drop is measured in the other layer. In the exciton phase, the Hall drag resistance becomes accurately quantized Kellogg et al. (2002), Tutuc et al. (2009).

Theoretical explanation of these experimental findings is provided by the Halperin (1,1,1) state of Eq. (19). In this state, each electron is a coherent linear superposition of the two layers Fertig (1989) where the relative phase ϕ is arbitrary and the same for all electrons. Such a state is ferromagnetically ordered in the x - y plane, inclined by the angle ϕ relative to the x -axis and spontaneously chosen by the system. As a consequence of spontaneous U(1) symmetry breaking, the system acquires a gapless Goldstone mode – a linearly dispersing pseudospin wave Moon et al. (1995), which was detected in Ref. Spielman et al. (2001).

While the properties of the $\nu_T=1$ bilayer have been well-understood in the limits of small and large d/ℓ , the nature of the transition at some critical d_c/ℓ and a possibility of intermediate phases remain the subject of on-going work Möller et al. (2008), Milovanović et al. (2015), Zhu et al. (2017), Lian & Zhang (2018), Wagner et al. (2021).

10. Recent developments

In recent years the interest in FQH phases has expanded beyond topology into broader investigations of geometric effects and quantum entanglement. Moreover, there has been a resurgence of interest in the parton description of FQH states and effective field theories that incorporate geometric effects and particle-hole symmetry. This section is an overview of these recent developments in the field.

10.1. Parton theory

Parton theory was initially introduced as a generalization of CF theory to describe a wider class of states Jain (1989b). In the parton theory, one imagines splitting the electron into k (an odd integer) fictitious sub-particles called *partons*. To obtain an incompressible state, each of the partons is placed in an IQH state (or in general, any known incompressible state), resulting in the wave function:

$$\Psi_v^{n_1 n_2 \dots n_k} = \mathcal{P}_{\text{LLL}} \prod_{\beta=1}^k \Phi_{n_\beta}, \quad (20)$$

where β labels the parton species. The parton state described by the wave function given in Eq. (20) is referred to by the shorthand notation “ $n_1 n_2 \dots n_k$ ”. The partons are unphysical entities so they need to be glued back together to recover the physical electrons. At the level of wave functions, this gluing procedure is implemented by setting the different parton species coordinates equal to the electron coordinate (each

Φ_{n_β} in Eq. (20) is built up of all the electronic coordinates.). All the partons are exposed to the same external magnetic field as the underlying electrons and each parton species has the same density as that of the electrons. Therefore, the charge of the β parton $q_\beta = -e\nu/n_\beta$. The constraint that the sum of the parton charges should add to that of the electron relates the parton fillings to the electronic filling, i.e., $\sum_{\beta=1}^k q_\beta = -e \implies \nu = [\sum_{\beta=1}^k n_\beta^{-1}]^{-1}$.

Looking back at the wave functions in Eqs. (10) and (15), it follows that all of the Abelian Laughlin and Jain states are parton states. The parton theory can also describe non-Abelian states. In general, a parton state with a repeated factor of n , with $|n| \geq 2$ hosts non-Abelian excitations Wen (1991). The simplest non-Abelian parton state – the 221 state Jain (1989b) at $\nu=1/2$ – can be interpreted as an f -wave superconductor of CFs Faugno et al. (2019). Recent identification of a parton state that lies in the same universality class as the aPf state but provides a better microscopic representation of the exact Coulomb ground state at $5/2$ Balram et al. (2018) has contributed to the resurgence of interest in partons. Furthermore, it has been shown that certain high-energy excitations of the states in the Jain sequence lie beyond the scope of the CF theory Nguyen et al. (2021) while they lend themselves to a description in terms of partons Balram et al. (2021). At this stage, the parton theory promises to provide a unified framework for describing all of the quantum Hall effects.

10.2. Effective field theory

As we mentioned in Sec. 4.1, many of the exotic properties of FQH states originate from the emergent Chern-Simons gauge field. Based on this idea, a field-theoretic approach to FQH systems was pioneered by Zhang, Hansson and Kivelson Zhang et al. (1989) who showed, using a Chern-Simons field theory, that the FQH effect at Laughlin fractions can be mapped onto a superfluid of composite bosons. Alternatively, the Chern-Simons field theory of CFs was introduced by Lopez and Fradkin Lopez & Fradkin (1991) and further developed by Halperin, Lee and Read Halperin et al. (1993) and extended via a Hamiltonian formalism by Murthy and Shankar Murthy & Shankar (2003). Detailed treatment of these theories is beyond the scope of the current chapter. Below we discuss two important recent developments in formulating an effective field theory of FQH systems: the role of PH symmetry and geometrical degrees of freedom.

The interest in the field-theoretic description of FQH systems, in particular in the role played by PH symmetry, has recently been renewed by geometric resonance experiments, which probe the radius of the cyclotron orbit of CFs. A series of measurements Kamburov et al. (2014), Hossain et al. (2020) have shown that the CF Fermi wave vector k_F^* is determined by the density of minority carriers in the LLL, i.e., the CF Fermi wave vector is set by the electron density for $\nu \leq 1/2$ and by the hole density for $\nu > 1/2$. In other words, the quantity $k_F^* \ell$ is PH-symmetric. The state that underpins the HLR theory does not reside entirely in

the LLL and thus it does not obey PH symmetry manifestly. This puzzle is resolved by going beyond the mean-field level, where it has been shown that the HLR theory produces results that are PH-symmetric and consistent with experiment Wang et al. (2017).

In parallel, Son (2015) proposed a field theory that is valid in the vicinity of half-filling and has a built-in PH symmetry. In Son's theory, CFs are Dirac particles whose density is controlled by the external magnetic field as opposed to by the density of electrons. The PH symmetry of electrons acts as a time-reversal symmetry on Dirac CFs. This results in an absence of $2k_F$ back-scattering that was subsequently observed in DMRG simulations Geraedts et al. (2017), providing evidence in favor of Son's theory. One can ask how different are the predictions of Son's theory from that of the HLR theory? Surprisingly, it turns out that the refined version of HLR theory is consistent with Son's theory on many aspects Wang et al. (2017).

Interestingly, Son's construction suggests a PH-symmetric gapped state at half-filling – dubbed the “PH symmetric Pfaffian” (PH-Pf). The PH-Pf state is topologically distinct from both the Moore-Read state and a Pf state. Intriguingly, recent thermal Hall Banerjee et al. (2018) and heat transport measurements Dutta et al. (n.d.) at $\nu=5/2$ appear to be consistent with only the PH-Pf order. Two candidate wave functions for the PH-Pfaffian phase have been considered in the literature Jolicoeur (2007), Zucker & Feldman (2016), both with a high degree of PH symmetry. The two wave functions also have high overlaps with each other for small systems indicating that they likely describe the same phase. However, numerical studies find no evidence for the existence of the PH-Pf phase in clean systems Balram et al. (2018), Mishmash et al. (2018). Understanding the heat transport measurements at $\nu=5/2$ in light of the numerical results presents one of the foremost challenges in the field.

Apart from PH symmetry, it has recently become apparent that any effective field theory capable of capturing dynamical response of FQH phases must include the *geometric* degrees of freedom FQH systems, which are beyond the remit of purely topological field theories. An important example of a geometric property is the Wen-Zee shift Wen & Zee (1992), related to the response function known as Hall viscosity Avron et al. (1995), Read (2009b). This has highlighted the need for a deeper understanding of “quantum geometry” in FQH fluids Haldane (2011). The quantum geometry is associated with the magnetoroton (GMP) collective excitation of FQH fluids in the long-wavelength limit (cf. Sec. 5.3). In this limit, the GMP mode can be interpreted as a quadrupolar density-wave deformation of the ground state and it has been incorporated into a gravitational field theory in the form of a spin-2 field coupled to an ambient geometry Gromov & Son (2017). Occasionally, this emergent spin-2 degree of freedom is also referred to as “FQH graviton” Yang, Hu, Papić & Haldane (2012), Golkar et al. (2016), Liou et al. (2019). The dynamics of the spin-2 modes Liu et al. (2018) can be induced by mass

anisotropy Yang, Papić, Rezayi, Bhatt & Haldane (2012), Wang et al. (2012), Ghazaryan & Chakraborty (2015) or by tilting the magnetic field Chakraborty & Pietiläinen (1989), Papić (2013). In gapless CFL states, the geometric effects manifest through the shape of the Fermi contour Gokmen et al. (2010), Jo et al. (2017). The investigation of geometric responses of FQH systems to the variations of ambient geometry represents an active research direction You et al. (2014), Bradlyn & Read (2015), Can et al. (2014).

10.3. Entanglement-based approaches

Quantum entanglement has recently emerged as a powerful probe of topological order. One characterizes entanglement by evaluating the system's reduced density matrix, obtained by partitioning the FQH system into two halves and tracing out one of the subsystems. As first shown by Li and Haldane Li & Haldane (2008), the spectrum of the reduced density matrix – the “entanglement spectrum” – contains information about the edge excitations of the system and it can be used to infer the CFT that governs the FQH state.

For FQH wave functions such as the Laughlin or Moore-Read, the number of levels in the entanglement spectrum as a function of momentum in the subsystem becomes universal as the subsystem size increases and it corresponds to the edge spectrum of the corresponding CFT. This “bulk-boundary” correspondence suggests that the bulk of the system and the physical edge are related to each other Dubail et al. (2012). On the other hand, the entanglement spectrum of a physical ground state of the Coulomb interaction is found to contain both a universal CFT part and a non-universal part. The latter is separated by a “entanglement gap” Thomale et al. (2010), which is argued to remain finite in a gapped FQH phase, signaling the robustness of topological order. The entanglement spectrum has found broad use as a tool for topological characterization of phases of matter beyond FQH states discussed in this chapter.

Finally, as first shown by Zaletel and Mong Zaletel & Mong (2012), certain FQH wave functions that can be expressed as correlators of a CFT can also be turned into a matrix-product state (MPS) if one picks the basis of LL orbitals on a cylinder geometry. This leads to a significant advantage in extracting physical properties from FQH wave functions. Exact MPS representations have been found for many states discussed above Estienne et al. (2013). Beyond studies of FQH wave functions, the MPS method has been generalized into a variational density-matrix renormalization group scheme for approximating ground states of realistic systems with Coulomb interactions Zaletel et al. (2013).

11. Conclusions

In this chapter we presented an introduction to the FQH effect observed in semiconductor systems. While it was impossible to give a historically exhaustive account of the entire field, we hope that the multitude of examples illustrates the great richness of emergent many-body phenomena and the diversity of approaches that have been developed over

filling, ν	state	shift, S	chiral central charge, c_-	quasiparticle charge, Q_{qp}	GSD
$1/(2p+1)$	1^{2p+1} , Laughlin	$2p+1$	1	$1/(2p+1)$	$2p+1$
$n/(2pn \pm 1)$	$(\pm n)1^{2p}$, Jain	$(\pm n) + 2p$	$1 \pm (n-1)$	$1/(2pn \pm 1)$	$2pn \pm 1$
$1/(2p)$	Moore-Read	$2p+1$	$3/2$	$1/(4p)$	$6p$
$1/(2p)$	$2^2 1^{2p+1}$, anti-Pfaffian	$2p-3$	$-1/2$	$1/(4p)$	$6p$
$2/5$	$2^3 1^4$, anti- \mathbb{Z}_3 Read-Rezayi	-2	$-4/5$	$1/5$	10

Table 1

Summary of topological properties of main single-component FQH states considered in this chapter. For the states captured by parton theory, we also show their parton label according to Eq. (20). Some of the quantities in this Table can be accessed experimentally: filling factor ν is related to the electrical Hall resistance R_{xy} via Eq. (6), chiral central charge c_- is related to the thermal Hall conductance κ_{xy} via Eq. (8) (any filled LLs provide an additional contribution), Q_{qp} is the charge of fundamental quasiparticle in units of the electron charge which can be accessed via shot-noise experiments or by charge-sensing using a single-electron transistor, the shift S on the sphere is related to the Hall viscosity η_H . The ground state degeneracy (GSD) is quoted for the torus. The composite fermion Fermi liquid (CFL) is not expected to carry a quantized κ_{xy} since its bulk is gapless.

the years in order to understand the novel physics associated with the FQH effect. To emphasize this diversity, our presentation has focused on explaining the experimental observations through model states, which distil the essence of FQH physics into elegant wave functions that can be studied either analytically or in computer simulations. Thus, we conclude by presenting a summary of the properties of the main single-component FQH phases discussed above in Table 1. The table lists some key topological quantum numbers that can be used to characterize an FQH phase, as discussed in Sec. 4.1. Some of these quantities are accessible in experiments and others in numerics.

Throughout this chapter, we have highlighted many open problems and research directions that remain particularly active at the time of writing (early 2022). We note in particular that the material realizations of the FQH effect continue to expand, most notably in the context of graphene and other van der Waals materials, fractional Chern insulators, as well as synthetic quantum simulators based on ultracold atoms and superconducting qubit arrays. We purposefully avoided discussing alternative material realizations of the FQH effect, as some of them are discussed in other chapters of this encyclopedia. These new material platforms may offer additional pathways to testing some of the theoretical ideas discussed above, while they will undoubtedly pose new puzzles to be understood. For a research field that is now four decades old, the FQH effect remains a remarkably active source of exciting new physics.

12. Acknowledgements

We would like to thank Gabor Csathy and Wei Pan for generously sharing their experimental data that we plotted in Fig. 1. The authors express their deep sense of gratitude to all of their collaborators over the years, without whom this chapter would not have been possible. A. C. B. acknowledges the Science and Engineering Research Board (SERB) of the Department of Science and Technology (DST) for financial support through the

Start-up Grant SRG/2020/000154. A. C. B. and Z. P. thank the Royal Society International Exchanges Award IES\R2\202052 for funding support. Some of the numerical calculations reported in this work were carried out on the Nandadevi supercomputer, which is maintained and supported by the Institute of Mathematical Science's High-Performance Computing Center. Z. P. acknowledges support by the Leverhulme Trust Research Leadership Award RL-2019-015.

References

- Archer, A. C., Park, K. & Jain, J. K. (2013), 'Competing crystal phases in the lowest Landau level', *Phys. Rev. Lett.* **111**, 146804.
- Arovas, D., Schrieffer, J. R. & Wilczek, F. (1984), 'Fractional statistics and the quantum Hall effect', *Phys. Rev. Lett.* **53**, 722–723.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.53.722>
- Avron, J. E., Seiler, R. & Zograf, P. G. (1995), 'Viscosity of quantum Hall fluids', *Phys. Rev. Lett.* **75**, 697–700.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.75.697>
- Balram, A. C., Barkeshli, M. & Rudner, M. S. (2018), 'Parton construction of a wave function in the anti-Pfaffian phase', *Phys. Rev. B* **98**, 035127.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.98.035127>
- Balram, A. C., Liu, Z., Gromov, A. & Papić, Z. (2021), 'Very high-energy collective states of partons in fractional quantum Hall liquids', *arXiv e-prints*.
- Balram, A. C., Töke, C., Wójs, A. & Jain, J. K. (2015), 'Phase diagram of fractional quantum Hall effect of composite fermions in multicomponent systems', *Phys. Rev. B* **91**, 045109.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.91.045109>
- Balram, A. C., Wu, Y.-H., Sreejith, G. J., Wójs, A. & Jain, J. K. (2013), 'Role of exciton screening in the $\nu = 7/3$ fractional quantum Hall effect', *Phys. Rev. Lett.* **110**, 186801.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.110.186801>
- Balram, A. C., Wurstbauer, U., Wojs, A., Pinczuk, A. & Jain, J. K. (2015), 'Fractionally charged skyrmions in fractional quantum Hall effect', *Nat Commun* **6**, Article.
URL: <http://dx.doi.org/10.1038/ncomms9981>
- Banerjee, M., Heiblum, M., Umansky, V., Feldman, D. E., Oreg, Y. & Stern, A. (2018), 'Observation of half-integer thermal hall conductance', *Nature* **559**(7713), 205–210.
- Bartolomei, H., Kumar, M., Bisognin, R., Marguerite, A., Berroir, J.-M., Bocquillon, E., Plaçais, B., Cavanna, A., Dong, Q., Gennser, U., Jin, Y. & Fève, G. (2020), 'Fractional statistics in anyon collisions', *Science*

- 368(6487), 173–177.
 URL: <https://science.sciencemag.org/content/368/6487/173>
- Bernevig, B. A. & Haldane, F. D. M. (2008), ‘Model fractional quantum Hall states and jack polynomials’, *Phys. Rev. Lett.* **100**, 246802.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.100.246802>
- Bernevig, B. A. & Regnault, N. (2009), ‘Anatomy of abelian and non-Abelian fractional quantum Hall states’, *Phys. Rev. Lett.* **103**, 206801.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.103.206801>
- Bid, A., Ofek, N., Inoue, H., Heiblum, M., Kane, C. L., Umansky, V. & Mahalu, D. (2010), ‘Observation of neutral modes in the fractional quantum Hall regime’, *Nature* **466**(7306), 585–590.
- Bishara, W. & Nayak, C. (2009), ‘Effect of Landau level mixing on the effective interaction between electrons in the fractional quantum Hall regime’, *Phys. Rev. B* **80**, 121302.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.80.121302>
- Bishop, N. C., Padmanabhan, M., Vakili, K., Shkolnikov, Y. P., De Poortere, E. P. & Shayegan, M. (2007), ‘Valley polarization and susceptibility of composite fermions around a filling factor $\nu = 3/2$ ’, *Phys. Rev. Lett.* **98**, 266404.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.98.266404>
- Boebinger, G. S., Chang, A. M., Stormer, H. L. & Tsui, D. C. (1985), ‘Magnetic field dependence of activation energies in the fractional quantum hall effect’, *Phys. Rev. Lett.* **55**, 1606–1609.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.55.1606>
- Bonderson, P., Feiguin, A. E., Möller, G. & Slingerland, J. K. (2012), ‘Competing topological orders in the $\nu = 12/5$ quantum Hall state’, *Phys. Rev. Lett.* **108**, 036806.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.108.036806>
- Bonderson, P., Feiguin, A. E. & Nayak, C. (2011), ‘Numerical calculation of the neutral fermion gap at the $\nu = 5/2$ fractional quantum hall state’, *Phys. Rev. Lett.* **106**, 186802.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.106.186802>
- Bonderson, P. & Slingerland, J. K. (2008), ‘Fractional quantum Hall hierarchy and the second Landau level’, *Phys. Rev. B* **78**, 125323.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.78.125323>
- Bradlyn, B. & Read, N. (2015), ‘Low-energy effective theory in the bulk for transport in a topological phase’, *Phys. Rev. B* **91**, 125303.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.91.125303>
- Can, T., Laskin, M. & Wiegmann, P. (2014), ‘Fractional quantum hall effect in a curved space: Gravitational anomaly and electromagnetic response’, *Phys. Rev. Lett.* **113**, 046803.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.113.046803>
- Chakraborty, T. (2000), ‘Electron spin transitions in quantum Hall systems’, *Advances in Physics* **49**(8), 959–1014.
 URL: <https://doi.org/10.1080/00018730050198161>
- Chakraborty, T. & Pietiläinen, P. (1987), ‘Fractional quantum Hall effect at half-filled Landau level in a multiple-layer electron system’, *Phys. Rev. Lett.* **59**, 2784–2787.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.59.2784>
- Chakraborty, T. & Pietiläinen, P. (1989), ‘Fractional quantum hall effect in tilted magnetic fields’, *Phys. Rev. B* **39**, 7971–7973.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.39.7971>
- Chakraborty, T. & Zhang, F. C. (1984), ‘Role of reversed spins in the correlated ground state for the fractional quantum Hall effect’, *Phys. Rev. B* **29**, 7032–7033.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.29.7032>
- Chamon, C. d. C. & Wen, X. G. (1994), ‘Sharp and smooth boundaries of quantum Hall liquids’, *Phys. Rev. B* **49**, 8227–8241.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.49.8227>
- Chang, A. M. (2003), ‘Chiral Luttinger liquids at the fractional quantum Hall edge’, *Rev. Mod. Phys.* **75**, 1449–1505.
 URL: <http://link.aps.org/doi/10.1103/RevModPhys.75.1449>
- Chang, A. M., Pfeiffer, L. N. & West, K. W. (1996), ‘Observation of chiral Luttinger behavior in electron tunneling into fractional quantum Hall edges’, *Phys. Rev. Lett.* **77**, 2538–2541.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.77.2538>
- Chklovskii, D. B., Shklovskii, B. I. & Glazman, L. I. (1992), ‘Electrostatics of edge channels’, *Phys. Rev. B* **46**, 4026–4034.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.46.4026>
- Chung, Y. J., Graf, D., Engel, L. W., Rosales, K. A. V., Madathil, P. T., Baldwin, K. W., West, K. W., Pfeiffer, L. N. & Shayegan, M. (2022), ‘Correlated states of 2D electrons near the Landau level filling $\nu = 1/7$ ’, *Phys. Rev. Lett.* **128**, 026802.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.128.026802>
- Chung, Y. J., Villegas Rosales, K. A., Baldwin, K. W., Madathil, P. T., West, K. W., Shayegan, M. & Pfeiffer, L. N. (2021), ‘Ultra-high-quality two-dimensional electron systems’, *Nature Materials* .
 URL: <https://doi.org/10.1038/s41563-021-00942-3>
- Clark, R. G., Haynes, S. R., Suckling, A. M., Mallett, J. R., Wright, P. A., Harris, J. J. & Foxon, C. T. (1989), ‘Spin configurations and quasiparticle fractional charge of fractional-quantum-Hall-effect ground states in the $N=0$ Landau level’, *Phys. Rev. Lett.* **62**, 1536–1539.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.62.1536>
- d’Ambrumenil, N. & Reynolds, A. M. (1988), ‘Fractional quantum Hall states in higher Landau levels’, *Journal of Physics C: Solid State Physics* **21**(1), 119.
 URL: <http://stacks.iop.org/0022-3719/21/i=1/a=010>
- Deng, H., Liu, Y., Jo, I., Pfeiffer, L. N., West, K. W., Baldwin, K. W. & Shayegan, M. (2016), ‘Commensurability oscillations of composite fermions induced by the periodic potential of a Wigner crystal’, *Phys. Rev. Lett.* **117**, 096601.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.117.096601>
- DiagHam (n.d.).
 URL: <https://www.nick-ux.org/diagham>
- Dolev, M., Gross, Y., Chung, Y. C., Heiblum, M., Umansky, V. & Mahalu, D. (2010), ‘Dependence of the tunneling quasiparticle charge determined via shot noise measurements on the tunneling barrier and energetics’, *Phys. Rev. B* **81**, 161303.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.81.161303>
- Dolev, M., Gross, Y., Sabo, R., Gurman, I., Heiblum, M., Umansky, V. & Mahalu, D. (2011), ‘Characterizing neutral modes of fractional states in the second Landau level’, *Phys. Rev. Lett.* **107**, 036805.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.107.036805>
- Dolev, M., Heiblum, M., Stern, A., Umansky, V. & Mahalu, D. (2008), ‘Observation of a quarter of an electron charge at the $\nu = 5/2$ quantum Hall state’, *Nature* **452**, 829 EP –.
 URL: <http://dx.doi.org/10.1038/nature06855>
- Du, R. R., Stormer, H. L., Tsui, D. C., Pfeiffer, L. N. & West, K. W. (1993), ‘Experimental evidence for new particles in the fractional quantum Hall effect’, *Phys. Rev. Lett.* **70**, 2944–2947.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.70.2944>
- Du, R. R., Yeh, A. S., Stormer, H. L., Tsui, D. C., Pfeiffer, L. N. & West, K. W. (1995), ‘Fractional quantum Hall effect around $\nu = 3/2$: Composite fermions with a spin’, *Phys. Rev. Lett.* **75**, 3926–3929.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.75.3926>
- Du, R., Tsui, D., Stormer, H., Pfeiffer, L., Baldwin, K. & West, K. (1999), ‘Strongly anisotropic transport in higher two-dimensional Landau levels’, *Solid State Communications* **109**(6), 389 – 394.
 URL: <http://www.sciencedirect.com/science/article/pii/S003810989800578X>
- Dubail, J., Read, N. & Rezayi, E. H. (2012), ‘Edge-state inner products and real-space entanglement spectrum of trial quantum Hall states’, *Phys. Rev. B* **86**, 245310.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.86.245310>
- Duncan, F. & Haldane, M. (1990), The hierarchy of fractional states and numerical studies, in ‘The quantum Hall effect’, Springer, pp. 303–352.
- Dutta, B., Yang, W., Melcer, R., Kundu, H. K., Heiblum, M., Umansky, V., Oreg, Y., Stern, A. & Mross, D. (n.d.), ‘Distinguishing between non-abelian topological orders in a quantum Hall system’, *Science* **0**(0), eabg6116.
 URL: <https://www.science.org/doi/abs/10.1126/science.abg6116>
- Eisenstein, J. P., Boebinger, G. S., Pfeiffer, L. N., West, K. W. & He, S. (1992), ‘New fractional quantum Hall state in double-layer two-dimensional electron systems’, *Phys. Rev. Lett.* **68**, 1383–1386.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.68.1383>
- Eisenstein, J. P., Cooper, K. B., Pfeiffer, L. N. & West, K. W. (2002), ‘Insulating and fractional quantum Hall states in the first excited Landau

- level', *Phys. Rev. Lett.* **88**, 076801.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.88.076801>
- Eisenstein, J. P., Stormer, H. L., Pfeiffer, L. N. & West, K. W. (1990), 'Evidence for a spin transition in the $\nu = 2/3$ fractional quantum Hall effect', *Phys. Rev. B* **41**, 7910–7913.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.41.7910>
- Estienne, B., Papić, Z., Regnault, N. & Bernevig, B. A. (2013), 'Matrix product states for trial quantum hall states', *Phys. Rev. B* **87**, 161112.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.87.161112>
- Fano, G., Ortolani, F. & Colombo, E. (1986), 'Configuration-interaction calculations on the fractional quantum Hall effect', *Phys. Rev. B* **34**, 2670–2680.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.34.2670>
- Faugno, W. N., Balram, A. C., Barkeshli, M. & Jain, J. K. (2019), 'Prediction of a non-Abelian fractional quantum Hall state with f -wave pairing of composite fermions in wide quantum wells', *Phys. Rev. Lett.* **123**, 016802.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.123.016802>
- Feigin, B., Jimbo, M., Miwa, T. & Mukhin, E. (2002), 'A differential ideal of symmetric polynomials spanned by Jack polynomials at $r\beta = -(r-1)/(k+1)$ ', *International Mathematics Research Notices* **2002**(23), 1223–1237.
 URL: <https://doi.org/10.1155/S1073792802112050>
- Figuin, A. E., Rezayi, E., Nayak, C. & Das Sarma, S. (2008), 'Density matrix renormalization group study of incompressible fractional quantum hall states', *Phys. Rev. Lett.* **100**, 166803.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.100.166803>
- Fertig, H. A. (1989), 'Energy spectrum of a layered system in a strong magnetic field', *Phys. Rev. B* **40**, 1087–1095.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.40.1087>
- Fertig, H. A., Brey, L., Côté, R. & MacDonald, A. H. (1994), 'Charged spin-texture excitations and the hartree-fock approximation in the quantum Hall effect', *Phys. Rev. B* **50**, 11018–11021.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.50.11018>
- Fogler, M. M., Koulakov, A. A. & Shklovskii, B. I. (1996), 'Ground state of a two-dimensional electron liquid in a weak magnetic field', *Phys. Rev. B* **54**, 1853–1871.
- Freedman, M., Kitaev, A., Larsen, M. & Wang, Z. (2003), 'Topological quantum computation', *Bulletin of the American Mathematical Society* **40**(1), 31–38.
 URL: <https://www.ams.org/journals/bull/2003-40-01/S0273-0979-02-00964-3>
- Geraedts, S. D., Repellin, C., Wang, C., Mong, R. S. K., Senthil, T. & Regnault, N. (2017), 'Emergent particle-hole symmetry in spinful bosonic quantum Hall systems', *Phys. Rev. B* **96**, 075148.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.96.075148>
- Ghazaryan, A. & Chakraborty, T. (2015), 'Aspects of anisotropic fractional quantum Hall effect in phosphorene', *Phys. Rev. B* **92**, 165409.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.92.165409>
- Girvin, S. M., MacDonald, A. H. & Platzman, P. M. (1985), 'Collective-excitation gap in the fractional quantum Hall effect', *Phys. Rev. Lett.* **54**, 581–583.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.54.581>
- Gokmen, T., Padmanabhan, M. & Shayegan, M. (2010), 'Transference of transport anisotropy to composite fermions', *Nature Physics* **6**, 621–624.
- Goldman, V. J., Su, B. & Jain, J. K. (1994), 'Detection of composite fermions by magnetic focusing', *Phys. Rev. Lett.* **72**, 2065–2068.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.72.2065>
- Golkar, S., Nguyen, D. X., Roberts, M. M. & Son, D. T. (2016), 'Higher-spin theory of the magnetorotons', *Phys. Rev. Lett.* **117**, 216403.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.117.216403>
- Greiter, M. (1994), 'Microscopic formulation of the hierarchy of quantized Hall states', *Physics Letters B* **336**(1), 48–53.
 URL: <https://www.sciencedirect.com/science/article/pii/0370269394009570>
- Greiter, M., Wen, X. G. & Wilczek, F. (1992), 'Paired Hall states in double-layer electron systems', *Phys. Rev. B* **46**, 9586–9589.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.46.9586>
- Gromov, A. & Son, D. T. (2017), 'Bimetric theory of fractional quantum hall states', *Phys. Rev. X* **7**, 041032.
 URL: <https://link.aps.org/doi/10.1103/PhysRevX.7.041032>
- Groshaus, J. G., Dujovne, I., Gallais, Y., Hirjibehedin, C. F., Pinczuk, A., Tan, Y.-W., Stormer, H., Dennis, B. S., Pfeiffer, L. N. & West, K. W. (2008), 'Spin texture and magnetoroton excitations at $\nu = 1/3$ ', *Phys. Rev. Lett.* **100**, 046804.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.100.046804>
- Haldane, F. D. M. (1983), 'Fractional quantization of the Hall effect: A hierarchy of incompressible quantum fluid states', *Phys. Rev. Lett.* **51**, 605–608.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.51.605>
- Haldane, F. D. M. (1985), 'Many-particle translational symmetries of two-dimensional electrons at rational Landau-level filling', *Phys. Rev. Lett.* **55**, 2095–2098.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.55.2095>
- Haldane, F. D. M. (2011), 'Geometrical description of the fractional quantum Hall effect', *Phys. Rev. Lett.* **107**, 116801.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.107.116801>
- Halperin, B. I. (1982), 'Quantized Hall conductance, current-carrying edge states, and the existence of extended states in a two-dimensional disordered potential', *Phys. Rev. B* **25**, 2185–2190.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.25.2185>
- Halperin, B. I. (1983), 'Theory of the quantized Hall conductance', *Helvetica Physica Acta* **56**(1-3), 75–102.
- Halperin, B. I. (1984), 'Statistics of quasiparticles and the hierarchy of fractional quantized Hall states', *Phys. Rev. Lett.* **52**, 1583–1586.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.52.1583>
- Halperin, B. I., Lee, P. A. & Read, N. (1993), 'Theory of the half-filled Landau level', *Phys. Rev. B* **47**, 7312–7343.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.47.7312>
- Hansson, T. H., Hermanns, M., Simon, S. H. & Viefers, S. F. (2017), 'Quantum hall physics: Hierarchies and conformal field theory techniques', *Rev. Mod. Phys.* **89**, 025005.
 URL: <https://link.aps.org/doi/10.1103/RevModPhys.89.025005>
- He, S., Das Sarma, S. & Xie, X. C. (1993), 'Quantized Hall effect and quantum phase transitions in coupled two-layer electron systems', *Phys. Rev. B* **47**, 4394–4412.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.47.4394>
- He, S., Simon, S. H. & Halperin, B. I. (1994), 'Response function of the fractional quantized Hall state on a sphere. ii. exact diagonalization', *Phys. Rev. B* **50**, 1823–1831.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.50.1823>
- Hermanns, M. (2010), 'Condensing non-abelian quasiparticles', *Phys. Rev. Lett.* **104**, 056803.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.104.056803>
- Ho, T.-L. (1995), 'Broken symmetry of two-component $\nu = 1/2$ quantum Hall states', *Phys. Rev. Lett.* **75**, 1186–1189.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.75.1186>
- Hossain, M. S., Mueed, M. A., Ma, M. K., Villegas Rosales, K. A., Chung, Y. J., Pfeiffer, L. N., West, K. W., Baldwin, K. W. & Shayegan, M. (2020), 'Precise experimental test of the Luttinger theorem and particle-hole symmetry for a strongly correlated fermionic system', *Phys. Rev. Lett.* **125**, 046601.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.125.046601>
- Jain, J. K. (1989a), 'Composite-fermion approach for the fractional quantum Hall effect', *Phys. Rev. Lett.* **63**, 199–202.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.63.199>
- Jain, J. K. (1989b), 'Incompressible quantum Hall states', *Phys. Rev. B* **40**, 8079–8082.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.40.8079>
- Jain, J. K. (2014), 'A note contrasting two microscopic theories of the fractional quantum Hall effect', *Indian Journal of Physics* **88**, 915–929.
 URL: <http://dx.doi.org/10.1007/s12648-014-0491-9>
- Jain, J. K. & Kamilla, R. K. (1998), Composite fermions: Particles of the lowest Landau level, in 'Composite Fermions', World Scientific Pub Co Inc, Singapore, chapter 1, pp. 1–90.
 URL: http://www.worldscientific.com/doi/abs/10.1142/9789812815989_0001
- Jo, I., Rosales, K. A. V., Mueed, M. A., Pfeiffer, L. N., West, K. W., Bald-

- win, K. W., Winkler, R., Padmanabhan, M. & Shayegan, M. (2017), 'Transference of Fermi contour anisotropy to composite fermions', *Phys. Rev. Lett.* **119**, 016402.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.119.016402>
- Jolicoeur, T. (2007), 'Non-abelian states with negative flux: A new series of quantum Hall states', *Phys. Rev. Lett.* **99**, 036805.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.99.036805>
- Kallin, C. & Halperin, B. I. (1984), 'Excitations from a filled Landau level in the two-dimensional electron gas', *Phys. Rev. B* **30**, 5655–5668.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.30.5655>
- Kalmeyer, V. & Zhang, S.-C. (1992), 'Metallic phase of the quantum Hall system at even-denominator filling fractions', *Phys. Rev. B* **46**, 9889–9892.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.46.9889>
- Kamburov, D., Liu, Y., Mueed, M. A., Shayegan, M., Pfeiffer, L. N., West, K. W. & Baldwin, K. W. (2014), 'What determines the Fermi wave vector of composite fermions?', *Phys. Rev. Lett.* **113**, 196801.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.113.196801>
- Kane, C. L. & Fisher, M. P. A. (1997), 'Quantized thermal transport in the fractional quantum Hall effect', *Phys. Rev. B* **55**, 15832–15837.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.55.15832>
- Kang, W., Stormer, H. L., Pfeiffer, L. N., Baldwin, K. W. & West, K. W. (1993), 'How real are composite fermions?', *Phys. Rev. Lett.* **71**, 3850–3853.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.71.3850>
- Kellogg, M., Spielman, I. B., Eisenstein, J. P., Pfeiffer, L. N. & West, K. W. (2002), 'Observation of quantized Hall drag in a strongly correlated bilayer electron system', *Phys. Rev. Lett.* **88**, 126804.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.88.126804>
- Kitaev, A. Y. (2001), 'Unpaired majorana fermions in quantum wires', *Physics-Uspekhi* **44**(10S), 131–136.
URL: <https://doi.org/10.1070%2F1063-7869%2F44%2F10%2F%2Fs29>
- Klitzing, K. v., Dorda, G. & Pepper, M. (1980), 'New method for high-accuracy determination of the fine-structure constant based on quantized Hall resistance', *Phys. Rev. Lett.* **45**, 494–497.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.45.494>
- Koulakov, A. A., Fogler, M. M. & Shklovskii, B. I. (1996), 'Charge density wave in two-dimensional electron liquid in weak magnetic field', *Phys. Rev. Lett.* **76**, 499–502.
- Kukushkin, I. V., Smet, J. H., Scarola, V. W., Umansky, V. & von Klitzing, K. (2009), 'Dispersion of the excitations of fractional quantum Hall states', *Science* **324**(5930), 1044–1047.
URL: <http://www.sciencemag.org/content/324/5930/1044.abstract>
- Kukushkin, I. V., v. Klitzing, K. & Eberl, K. (1999), 'Spin polarization of composite fermions: Measurements of the Fermi energy', *Phys. Rev. Lett.* **82**, 3665–3668.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.82.3665>
- Kumar, A., Csáthy, G. A., Manfra, M. J., Pfeiffer, L. N. & West, K. W. (2010), 'Nonconventional odd-denominator fractional quantum Hall states in the second Landau level', *Phys. Rev. Lett.* **105**, 246808.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.105.246808>
- Lam, P. K. & Girvin, S. M. (1984), 'Liquid-solid transition and the fractional quantum-Hall effect', *Phys. Rev. B* **30**, 473–475.
- Laughlin, R. B. (1983), 'Anomalous quantum Hall effect: An incompressible quantum fluid with fractionally charged excitations', *Phys. Rev. Lett.* **50**, 1395–1398.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.50.1395>
- Laughlin, R. B. (1987), Elementary theory: the incompressible quantum fluid, in 'The quantum Hall effect', Springer, pp. 233–301.
- Leadley, D. R., Nicholas, R. J., Maude, D. K., Utjuzh, A. N., Portal, J. C., Harris, J. J. & Foxon, C. T. (1997), 'Fractional quantum Hall effect measurements at zero $\sim g$ factor', *Phys. Rev. Lett.* **79**, 4246–4249.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.79.4246>
- Lee, S.-S., Ryu, S., Nayak, C. & Fisher, M. P. A. (2007), 'Particle-hole symmetry and the $\nu = 5/2$ quantum Hall state', *Phys. Rev. Lett.* **99**, 236807.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.99.236807>
- Leinaas, J. & Myrheim, J. (1977), 'On the theory of identical particles', *Il Nuovo Cimento B Series 11* **37**(1), 1–23.
URL: <http://dx.doi.org/10.1007/BF02727953>
- Levin, M. & Halperin, B. I. (2009), 'Collective states of non-abelian quasiparticles in a magnetic field', *Phys. Rev. B* **79**, 205301.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.79.205301>
- Levin, M., Halperin, B. I. & Rosenow, B. (2007), 'Particle-hole symmetry and the Pfaffian state', *Phys. Rev. Lett.* **99**, 236806.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.99.236806>
- Li, H. & Haldane, F. D. M. (2008), 'Entanglement spectrum as a generalization of entanglement entropy: Identification of topological order in non-abelian fractional quantum hall effect states', *Phys. Rev. Lett.* **101**, 010504.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.101.010504>
- Lian, B. & Zhang, S.-C. (2018), 'Wave function and emergent su(2) symmetry in the $\nu_T = 1$ quantum hall bilayer', *Phys. Rev. Lett.* **120**, 077601.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.120.077601>
- Lilly, M. P., Cooper, K. B., Eisenstein, J. P., Pfeiffer, L. N. & West, K. W. (1999), 'Evidence for an anisotropic state of two-dimensional electrons in high Landau levels', *Phys. Rev. Lett.* **82**, 394–397.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.82.394>
- Liou, S.-F., Haldane, F. D. M., Yang, K. & Rezayi, E. H. (2019), 'Chiral gravitons in fractional quantum hall liquids', *Phys. Rev. Lett.* **123**, 146801.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.123.146801>
- Liu, Y., Hasdemir, S., Wójs, A., Jain, J. K., Pfeiffer, L. N., West, K. W., Baldwin, K. W. & Shayegan, M. (2014), 'Spin polarization of composite fermions and particle-hole symmetry breaking', *Phys. Rev. B* **90**, 085301.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.90.085301>
- Liu, Z., Gromov, A. & Papić, Z. (2018), 'Geometric quench and nonequilibrium dynamics of fractional quantum hall states', *Phys. Rev. B* **98**, 155140.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.98.155140>
- Lopez, A. & Fradkin, E. (1991), 'Fractional quantum Hall effect and Chern-Simons gauge theories', *Phys. Rev. B* **44**, 5246–5262.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.44.5246>
- Ma, M. K., Villegas Rosales, K. A., Deng, H., Chung, Y. J., Pfeiffer, L. N., West, K. W., Baldwin, K. W., Winkler, R. & Shayegan, M. (2020), 'Thermal and quantum melting phase diagrams for a magnetic-field-induced wigner solid', *Phys. Rev. Lett.* **125**, 036601.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.125.036601>
- Mandal, S. S. & Jain, J. K. (2001), 'Low-energy spin rotons in the fractional quantum Hall effect', *Phys. Rev. B* **63**, 201310.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.63.201310>
- Milovanović, M. V., Dobardžić, E. & Papić, Z. (2015), 'Meron deconfinement in the quantum hall bilayer at intermediate distances', *Phys. Rev. B* **92**, 195311.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.92.195311>
- Mishmash, R. V., Mross, D. F., Alicea, J. & Motrunich, O. I. (2018), 'Numerical exploration of trial wave functions for the particle-hole-symmetric Pfaffian', *Phys. Rev. B* **98**, 081107.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.98.081107>
- Möller, G., Simon, S. H. & Rezayi, E. H. (2008), 'Paired composite fermion phase of quantum hall bilayers at $\nu = \frac{1}{2} + \frac{1}{2}$ ', *Phys. Rev. Lett.* **101**, 176803.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.101.176803>
- Möller, G., Wójs, A. & Cooper, N. R. (2011), 'Neutral fermion excitations in the moore-read state at filling factor $\nu = 5/2$ ', *Phys. Rev. Lett.* **107**, 036803.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.107.036803>
- Mong, R. S. K., Zaletel, M. P., Pollmann, F. & Papić, Z. (2017), 'Fibonacci anyons and charge density order in the 12/5 and 13/5 quantum Hall plateaus', *Phys. Rev. B* **95**, 115136.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.95.115136>
- Moon, K., Mori, H., Yang, K., Girvin, S. M., MacDonald, A. H., Zheng, L., Yoshioka, D. & Zhang, S.-C. (1995), 'Spontaneous interlayer coherence in double-layer quantum Hall systems: Charged vortices and kosterlitz-thouless phase transitions', *Phys. Rev. B* **51**, 5138.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.51.5138>

- Moore, G. & Read, N. (1991), 'Nonabelions in the fractional quantum Hall effect', *Nucl. Phys. B* **360**, 362 – 396.
 URL: <http://www.sciencedirect.com/science/article/pii/0550321391904070>
- Morf, R. H. (1998), 'Transition from quantum Hall to compressible states in the second Landau level: New light on the $\nu = 5/2$ enigma', *Phys. Rev. Lett.* **80**, 1505–1508.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.80.1505>
- Morf, R. H., d'Ambrumenil, N. & Das Sarma, S. (2002), 'Excitation gaps in fractional quantum Hall states: An exact diagonalization study', *Phys. Rev. B* **66**, 075408.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.66.075408>
- Morf, R. & Halperin, B. I. (1986), 'Monte Carlo evaluation of trial wave functions for the fractional quantized Hall effect: Disk geometry', *Phys. Rev. B* **33**, 2221–2246.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.33.2221>
- Murthy, G. & Shankar, R. (2003), 'Hamiltonian theories of the fractional quantum Hall effect', *Rev. Mod. Phys.* **75**, 1101–1158.
 URL: <http://link.aps.org/doi/10.1103/RevModPhys.75.1101>
- Nakamura, J., Liang, S., Gardner, G. C. & Manfra, M. J. (2020), 'Direct observation of anyonic braiding statistics', *Nature Physics* **16**(9), 931–936.
- Nayak, C., Simon, S. H., Stern, A., Freedman, M. & Das Sarma, S. (2008), 'Non-Abelian anyons and topological quantum computation', *Rev. Mod. Phys.* **80**, 1083–1159.
 URL: <http://link.aps.org/doi/10.1103/RevModPhys.80.1083>
- Nayak, C. & Wilczek, F. (1996), '2n-quasihole states realize 2(n-1)-dimensional spinor braiding statistics in paired quantum Hall states', *Nucl. Phys. B* **479**(3), 529–553.
- Nguyen, D. X., Haldane, F. D. M., Rezayi, E. H., Son, D. T. & Yang, K. (2021), 'Multiple magnetorotons and spectral sum rules in fractional quantum Hall systems', *arXiv e-prints*.
- Nicholas, R. J., Leadley, D. R., Maude, D. K., Portal, J. C., Harris, J. J. & Foxon, C. T. (1998), 'Skyrmions and composite fermions in the limit of vanishing zeeman energy', *J. Phys. : Condens. Matter* **10**, 11327–11335.
- Ortalano, M. W., He, S. & Das Sarma, S. (1997), 'Realistic calculations of correlated incompressible electronic states in GaAs-al_xga_{1-x}as heterostructures and quantum wells', *Phys. Rev. B* **55**, 7702–7714.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.55.7702>
- Padmanabhan, M., Gokmen, T. & Shayegan, M. (2009), 'Density dependence of valley polarization energy for composite fermions', *Phys. Rev. B* **80**, 035423.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.80.035423>
- Pakrouski, K., Peterson, M. R., Jolicoeur, T., Scarola, V. W., Nayak, C. & Troyer, M. (2015), 'Phase diagram of the $\nu = 5/2$ fractional quantum Hall effect: Effects of Landau-level mixing and nonzero width', *Phys. Rev. X* **5**, 021004.
 URL: <http://link.aps.org/doi/10.1103/PhysRevX.5.021004>
- Pakrouski, K., Troyer, M., Wu, Y.-L., Das Sarma, S. & Peterson, M. R. (2016), 'Enigmatic 12/5 fractional quantum Hall effect', *Phys. Rev. B* **94**, 075108.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.94.075108>
- Pan, W., Baldwin, K. W., West, K. W., Pfeiffer, L. N. & Tsui, D. C. (2015), 'Fractional quantum Hall effect at Landau level filling $\nu = 4/11$ ', *Phys. Rev. B* **91**, 041301.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.91.041301>
- Pan, W., Du, R. R., Stormer, H. L., Tsui, D. C., Pfeiffer, L. N., Baldwin, K. W. & West, K. W. (1999), 'Strongly anisotropic electronic transport at landau level filling factor $\nu = 9/2$ and $\nu = 5/2$ under a tilted magnetic field', *Phys. Rev. Lett.* **83**, 820–823.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.83.820>
- Pan, W., Kang, W., Lilly, M. P., Reno, J. L., Baldwin, K. W., West, K. W., Pfeiffer, L. N. & Tsui, D. C. (2020), 'Particle-hole symmetry and the fractional quantum hall effect in the lowest landau level', *Phys. Rev. Lett.* **124**, 156801.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.124.156801>
- Pan, W., Stormer, H. L., Tsui, D. C., Pfeiffer, L. N., Baldwin, K. W. & West, K. W. (2002), 'Transition from an electron solid to the sequence of fractional quantum Hall states at very low Landau level filling factor', *Phys. Rev. Lett.* **88**, 176802.
- Pan, W., Stormer, H. L., Tsui, D. C., Pfeiffer, L. N., Baldwin, K. W. & West, K. W. (2003), 'Fractional quantum Hall effect of composite fermions', *Phys. Rev. Lett.* **90**, 016801.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.90.016801>
- Papić, Z. (2013), 'Fractional quantum Hall effect in a tilted magnetic field', *Phys. Rev. B* **87**, 245315.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.87.245315>
- Papić, Z., Mong, R. S. K., Yazdani, A. & Zaletel, M. P. (2018), 'Imaging anyons with scanning tunneling microscopy', *Phys. Rev. X* **8**, 011037.
 URL: <https://link.aps.org/doi/10.1103/PhysRevX.8.011037>
- Park, K. & Jain, J. K. (1998), 'Phase diagram of the spin polarization of composite fermions and a new effective mass', *Phys. Rev. Lett.* **80**, 4237–4240.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.80.4237>
- Peterson, M. R., Papić, Z. & Sarma, S. D. (2010), 'Fractional quantum Hall effects in bilayers in the presence of interlayer tunneling and charge imbalance', *Physical Review B* **82**(23), 235312.
- Pinczuk, A., Dennis, B. S., Pfeiffer, L. N. & West, K. (1993), 'Observation of collective excitations in the fractional quantum Hall effect', *Phys. Rev. Lett.* **70**, 3983–3986.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.70.3983>
- Radu, I. P., Miller, J. B., Marcus, C. M., Kastner, M. A., Pfeiffer, L. N. & West, K. W. (2008), 'Quasi-particle properties from tunneling in the $\nu = 5/2$ fractional quantum Hall state', *Science* **320**(5878), 899–902.
 URL: <https://science.sciencemag.org/content/320/5878/899>
- Read, N. (1990), 'Excitation structure of the hierarchy scheme in the fractional quantum Hall effect', *Phys. Rev. Lett.* **65**, 1502–1505.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.65.1502>
- Read, N. (2009a), 'Conformal invariance of chiral edge theories', *Phys. Rev. B* **79**, 245304.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.79.245304>
- Read, N. (2009b), 'Non-abelian adiabatic statistics and Hall viscosity in quantum Hall states and $p_x + ip_y$ paired superfluids', *Phys. Rev. B* **79**, 045308.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.79.045308>
- Read, N. & Green, D. (2000), 'Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect', *Phys. Rev. B* **61**, 10267–10297.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.61.10267>
- Read, N. & Rezayi, E. (1999), 'Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level', *Phys. Rev. B* **59**, 8084–8092.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.59.8084>
- Rezayi, E. H. (2017), 'Landau level mixing and the ground state of the $\nu = 5/2$ quantum hall effect', *Phys. Rev. Lett.* **119**, 026801.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.119.026801>
- Rezayi, E. H. & Haldane, F. D. M. (2000), 'Incompressible paired Hall state, stripe order, and the composite fermion liquid phase in half-filled Landau levels', *Phys. Rev. Lett.* **84**, 4685–4688.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.84.4685>
- Richard E. Prange, S. M. G., ed. (1987), *The Quantum Hall Effect*, Springer-Verlag New York.
- Ro, D., Deng, N., Watson, J. D., Manfra, M. J., Pfeiffer, L. N., West, K. W. & Csáthy, G. A. (2019), 'Electron bubbles and the structure of the orbital wave function', *Phys. Rev. B* **99**, 201111.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.99.201111>
- Saminadayar, L., Glatli, D. C., Jin, Y. & Etienne, B. (1997), 'Observation of the $e/3$ fractionally charged Laughlin quasiparticle', *Phys. Rev. Lett.* **79**, 2526–2529.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.79.2526>
- Samkharadze, N., Arnold, I., Pfeiffer, L. N., West, K. W. & Csáthy, G. A. (2015), 'Observation of incompressibility at $\nu = 4/11$ and $\nu = 5/13$ ', *Phys. Rev. B* **91**, 081109.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.91.081109>
- Scarola, V. W. & Jain, J. K. (2001), 'Phase diagram of bilayer composite fermion states', *Phys. Rev. B* **64**, 085313.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.64.085313>

- Scarola, V. W., Park, K. & Jain, J. K. (2000), 'Rotons of composite fermions: Comparison between theory and experiment', *Phys. Rev. B* **61**, 13064–13072.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.61.13064>
- Shabani, J., Gokmen, T., Chiu, Y. T. & Shayegan, M. (2009), 'Evidence for developing fractional quantum Hall states at even denominator $1/2$ and $1/4$ fillings in asymmetric wide quantum wells', *Phys. Rev. Lett.* **103**, 256802.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.103.256802>
- Shabani, J., Liu, Y., Shayegan, M., Pfeiffer, L. N., West, K. W. & Baldwin, K. W. (2013), 'Phase diagrams for the stability of the $\nu = \frac{1}{2}$ fractional quantum Hall effect in electron systems confined to symmetric, wide GaAs quantum wells', *Phys. Rev. B* **88**, 245413.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.88.245413>
- Shibata, N. & Yoshioka, D. (2001), 'Ground-state phase diagram of 2d electrons in a high Landau level: A density-matrix renormalization group study', *Phys. Rev. Lett.* **86**, 5755–5758.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.86.5755>
- Simon, S. H. & Rezayi, E. H. (2013), 'Landau level mixing in the perturbative limit', *Phys. Rev. B* **87**, 155426.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.87.155426>
- Simon, S. H., Rezayi, E. H. & Cooper, N. R. (2007), 'Pseudopotentials for multiparticle interactions in the quantum Hall regime', *Phys. Rev. B* **75**, 195306.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.75.195306>
- Simon, S. H., Rezayi, E. H., Cooper, N. R. & Berdnikov, I. (2007), 'Construction of a paired wave function for spinless electrons at filling fraction $\nu = 2/5$ ', *Phys. Rev. B* **75**, 075317.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.75.075317>
- Sodemann, I. & MacDonald, A. H. (2013), 'Landau level mixing and the fractional quantum Hall effect', *Phys. Rev. B* **87**, 245425.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.87.245425>
- Son, D. T. (2015), 'Is the composite fermion a Dirac particle?', *Phys. Rev. X* **5**, 031027.
 URL: <http://link.aps.org/doi/10.1103/PhysRevX.5.031027>
- Sondhi, S. L., Karlhede, A., Kivelson, S. A. & Rezayi, E. H. (1993), 'Skyrmions and the crossover from the integer to fractional quantum Hall effect at small Zeeman energies', *Phys. Rev. B* **47**, 16419–16426.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.47.16419>
- Spielman, I. B., Eisenstein, J. P., Pfeiffer, L. N. & West, K. W. (2000), 'Resonantly enhanced tunneling in a double layer quantum Hall ferromagnet', *Phys. Rev. Lett.* **84**, 5808–5811.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.84.5808>
- Spielman, I. B., Eisenstein, J. P., Pfeiffer, L. N. & West, K. W. (2001), 'Observation of a linearly dispersing collective mode in a quantum Hall ferromagnet', *Phys. Rev. Lett.* **87**, 036803.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.87.036803>
- Suen, Y. W., Engel, L. W., Santos, M. B., Shayegan, M. & Tsui, D. C. (1992), 'Observation of a $\nu = 1/2$ fractional quantum Hall state in a double-layer electron system', *Phys. Rev. Lett.* **68**, 1379–1382.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.68.1379>
- Thomale, R., Sterdyniak, A., Regnault, N. & Bernevig, B. A. (2010), 'Entanglement gap and a new principle of adiabatic continuity', *Phys. Rev. Lett.* **104**, 180502.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.104.180502>
- Thouless, D. J., Kohmoto, M., Nightingale, M. P. & den Nijs, M. (1982), 'Quantized Hall conductance in a two-dimensional periodic potential', *Phys. Rev. Lett.* **49**, 405–408.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.49.405>
- Trugman, S. A. & Kivelson, S. (1985), 'Exact results for the fractional quantum Hall effect with general interactions', *Phys. Rev. B* **31**, 5280–5284.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.31.5280>
- Tsui, D. C., Stormer, H. L. & Gossard, A. C. (1982), 'Two-dimensional magnetotransport in the extreme quantum limit', *Phys. Rev. Lett.* **48**, 1559–1562.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.48.1559>
- Tutuc, E., Pillarisetty, R. & Shayegan, M. (2009), 'Giant frictional drag in strongly interacting bilayers near filling factor one', *Phys. Rev. B* **79**, 041303.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.79.041303>
- Tutuc, E., Shayegan, M. & Huse, D. A. (2004), 'Counterflow measurements in strongly correlated GaAs hole bilayers: Evidence for electron-hole pairing', *Phys. Rev. Lett.* **93**, 036802.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.93.036802>
- Venkatachalam, V., Yacoby, A., Pfeiffer, L. & West, K. (2011), 'Local charge of the $[n\text{gr}] = 5/2$ fractional quantum Hall state', *Nature* **469**(7329), 185–188.
 URL: <http://dx.doi.org/10.1038/nature09680>
- Villegas Rosales, K. A., Madathil, P. T., Chung, Y. J., Pfeiffer, L. N., West, K. W., Baldwin, K. W. & Shayegan, M. (2021), 'Fractional quantum Hall effect energy gaps: Role of electron layer thickness', *Phys. Rev. Lett.* **127**, 056801.
 URL: <https://link.aps.org/doi/10.1103/PhysRevLett.127.056801>
- Wagner, G., Nguyen, D. X., Simon, S. H. & Halperin, B. I. (2021), 's-wave paired composite-fermion electron-hole trial state for quantum Hall bilayers with $\nu = 1$ '.
- Wang, C., Cooper, N. R., Halperin, B. I. & Stern, A. (2017), 'Particle-hole symmetry in the fermion-Chern-Simons and Dirac descriptions of a half-filled Landau level', *Phys. Rev. X* **7**, 031029.
 URL: <http://link.aps.org/doi/10.1103/PhysRevX.7.031029>
- Wang, H., Narayanan, R., Wan, X. & Zhang, F. (2012), 'Fractional quantum Hall states in two-dimensional electron systems with anisotropic interactions', *Phys. Rev. B* **86**, 035122.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.86.035122>
- Wen, X. G. (1990), 'Chiral Luttinger liquid and the edge excitations in the fractional quantum Hall states', *Phys. Rev. B* **41**, 12838–12844.
 URL: <http://link.aps.org/doi/10.1103/PhysRevB.41.12838>
- Wen, X. G. (1991), 'Non-abelian statistics in the fractional quantum Hall states', *Phys. Rev. Lett.* **66**, 802–805.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.66.802>
- Wen, X.-G. (1995), 'Topological orders and edge excitations in fractional quantum Hall states', *Advances in Physics* **44**(5), 405–473.
 URL: <http://www.tandfonline.com/doi/abs/10.1080/00018739500101566>
- Wen, X. G. & Niu, Q. (1990), 'Ground-state degeneracy of the fractional quantum Hall states in the presence of a random potential and on high-genus Riemann surfaces', *Phys. Rev. B* **41**, 9377–9396.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.41.9377>
- Wen, X. G. & Zee, A. (1992), 'Shift and spin vector: New topological quantum numbers for the Hall fluids', *Phys. Rev. Lett.* **69**, 953–956.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.69.953>
- Wilczek, F. (1982), 'Quantum mechanics of fractional-spin particles', *Phys. Rev. Lett.* **49**, 957–959.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.49.957>
- Willett, R., Eisenstein, J. P., Störmer, H. L., Tsui, D. C., Gossard, A. C. & English, J. H. (1987), 'Observation of an even-denominator quantum number in the fractional quantum Hall effect', *Phys. Rev. Lett.* **59**, 1776–1779.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.59.1776>
- Willett, R. L., Pfeiffer, L. N. & West, K. W. (2009), 'Measurement of filling factor $5/2$ quasiparticle interference with observation of charge $e/4$ and $e/2$ period oscillations', *Proceedings of the National Academy of Sciences* **106**(22), 8853–8858.
 URL: <http://www.pnas.org/content/106/22/8853.abstract>
- Willett, R. L., Ruel, R. R., West, K. W. & Pfeiffer, L. N. (1993), 'Experimental demonstration of a Fermi surface at one-half filling of the lowest Landau level', *Phys. Rev. Lett.* **71**, 3846–3849.
 URL: <http://link.aps.org/doi/10.1103/PhysRevLett.71.3846>
- Willett, R. L., Shtengel, K., Nayak, C., Pfeiffer, L. N., Chung, Y. J., Peabody, M. L., Baldwin, K. W., West, K. W. & . (2021), 'Interference measurements of non-abelian $e/4$ and abelian $e/2$ quasiparticle braiding'.
- Willett, R. L., Stormer, H. L., Tsui, D. C., Gossard, A. C. & English, J. H. (1988), 'Quantitative experimental test for the theoretical gap energies in the fractional quantum Hall effect', *Phys. Rev. B* **37**, 8476–8479.
 URL: <https://link.aps.org/doi/10.1103/PhysRevB.37.8476>

- Willett, R. L., West, K. W. & Pfeiffer, L. N. (1999), ‘Geometric resonance of composite fermion cyclotron orbits with a fictitious magnetic field modulation’, *Phys. Rev. Lett.* **83**, 2624–2627.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.83.2624>
- Williams, F. I. B., Wright, P. A., Clark, R. G., Andrei, E. Y., Deville, G., Glattli, D. C., Probst, O., Etienne, B., Dorin, C., Foxon, C. T. & Harris, J. J. (1991), ‘Conduction threshold and pinning frequency of magnetically induced Wigner solid’, *Phys. Rev. Lett.* **66**, 3285–3288.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.66.3285>
- Wu, X. G., Dev, G. & Jain, J. K. (1993), ‘Mixed-spin incompressible states in the fractional quantum Hall effect’, *Phys. Rev. Lett.* **71**, 153–156.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.71.153>
- Xia, J. S., Pan, W., Vicente, C. L., Adams, E. D., Sullivan, N. S., Stormer, H. L., Tsui, D. C., Pfeiffer, L. N., Baldwin, K. W. & West, K. W. (2004), ‘Electron correlation in the second Landau level: A competition between many nearly degenerate quantum phases’, *Phys. Rev. Lett.* **93**, 176809.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.93.176809>
- Yang, B., Hu, Z.-X., Papić, Z. & Haldane, F. D. M. (2012), ‘Model wave functions for the collective modes and the magnetoroton theory of the fractional quantum Hall effect’, *Phys. Rev. Lett.* **108**, 256807.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.108.256807>
- Yang, B., Papić, Z., Rezayi, E. H., Bhatt, R. N. & Haldane, F. D. M. (2012), ‘Band mass anisotropy and the intrinsic metric of fractional quantum Hall systems’, *Phys. Rev. B* **85**, 165318.
URL: <http://link.aps.org/doi/10.1103/PhysRevB.85.165318>
- Yoshioka, D., Halperin, B. I. & Lee, P. A. (1983), ‘Ground state of two-dimensional electrons in strong magnetic fields and $\frac{1}{3}$ quantized Hall effect’, *Phys. Rev. Lett.* **50**, 1219–1222.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.50.1219>
- Yoshioka, D., MacDonald, A. H. & Girvin, S. M. (1989), ‘Fractional quantum Hall effect in two-layered systems’, *Phys. Rev. B* **39**, 1932–1935.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.39.1932>
- You, Y., Cho, G. Y. & Fradkin, E. (2014), ‘Theory of nematic fractional quantum hall states’, *Phys. Rev. X* **4**, 041050.
URL: <http://link.aps.org/doi/10.1103/PhysRevX.4.041050>
- Zaletel, M. P. & Mong, R. S. K. (2012), ‘Exact matrix product states for quantum Hall wave functions’, *Phys. Rev. B* **86**, 245305.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.86.245305>
- Zaletel, M. P., Mong, R. S. K. & Pollmann, F. (2013), ‘Topological characterization of fractional quantum Hall ground states from microscopic Hamiltonians’, *Phys. Rev. Lett.* **110**, 236801.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.110.236801>
- Zhang, F. C. & Das Sarma, S. (1986), ‘Excitation gap in the fractional quantum Hall effect: Finite layer thickness corrections’, *Phys. Rev. B* **33**, 2903–2905.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.33.2903>
- Zhang, S. C., Hansson, T. H. & Kivelson, S. (1989), ‘Effective-field-theory model for the fractional quantum Hall effect’, *Phys. Rev. Lett.* **62**, 82–85.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.62.82>
- Zhao, T., Faugno, W. N., Pu, S., Balram, A. C. & Jain, J. K. (2021), ‘Origin of the $\nu = 1/2$ fractional quantum Hall effect in wide quantum wells’, *Phys. Rev. B* **103**, 155306.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.103.155306>
- Zhu, H., Chen, Y. P., Jiang, P., Engel, L. W., Tsui, D. C., Pfeiffer, L. N. & West, K. W. (2010), ‘Observation of a pinning mode in a Wigner solid with $\nu = 1/3$ fractional quantum Hall excitations’, *Phys. Rev. Lett.* **105**, 126803.
URL: <http://link.aps.org/doi/10.1103/PhysRevLett.105.126803>
- Zhu, J., Pan, W., Stormer, H. L., Pfeiffer, L. N. & West, K. W. (2002), ‘Density-induced interchange of anisotropy axes at half-filled high Landau levels’, *Phys. Rev. Lett.* **88**, 116803.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.88.116803>
- Zhu, Z., Fu, L. & Sheng, D. N. (2017), ‘Numerical study of quantum hall bilayers at total filling $\nu_T = 1$: A new phase at intermediate layer distances’, *Phys. Rev. Lett.* **119**, 177601.
URL: <https://link.aps.org/doi/10.1103/PhysRevLett.119.177601>
- Zucker, P. T. & Feldman, D. E. (2016), ‘Stabilization of the particle-hole Pfaffian order by Landau-level mixing and impurities that break particle-hole symmetry’, *Phys. Rev. Lett.* **117**, 096802.
- Zuo, Z.-W., Balram, A. C., Pu, S., Zhao, J., Jolicœur, T., Wójs, A. & Jain, J. K. (2020), ‘Interplay between fractional quantum Hall liquid and crystal phases at low filling’, *Phys. Rev. B* **102**, 075307.
URL: <https://link.aps.org/doi/10.1103/PhysRevB.102.075307>