

# Inverse Laplace transform based on Widder's method for Tsallis exponential

S.S. Naina Mohammed<sup>‡1</sup>, K. Jeevanandham<sup>1</sup>, A. Basherrudin Mahmud Ahmed<sup>2</sup>, Md. Manirul Ali<sup>3</sup> §, and R. Chandrashekar<sup>4</sup> ||

<sup>1</sup> Department of Physics, Government Arts College, Udumalpet, Tiruppur District - 642126, Tamilnadu, India

<sup>2</sup> School of Physics, Madurai Kamaraj University, Madurai 625021, India

<sup>3</sup> Centre for Quantum Science and Technology, Chennai Institute of Technology, Chennai 600069, India

<sup>4</sup> Centre for Quantum Information, Communication and Computing, Indian Institute of Technology Madras, Chennai 600036, India

**Abstract.** A generalization of the Laplace transform based on the generalized Tsallis  $q$ -exponential is given in the present work for a new type of kernel. We also define the inverse transform for this generalized transform based on the complex integration method. We prove identities corresponding to the Laplace transform and inverse transform like the  $q$ -convolution theorem, the action of generalized derivative and generalized integration on the Laplace transform. We then derive a  $q$ -generalization of the inverse Laplace transform based on the Post-Widder's method which bypasses the necessity for a complex contour integration. We demonstrate the usefulness of this in computing the Laplace and inverse Laplace transform of some elementary functions. Finally we use the Post-Widder's method based inverse Laplace transform to compute the density of states from the partition function for the case of a generalized classical ideal gas and linear harmonic oscillator in  $D$ -dimensions.

‡ naina@gacudpt.in

§ manirul@citchennai.net

|| chandrashekar@physics.iitm.ac.in

## 1. Introduction

A mathematical function ' $f$ ' in the ' $x$ ' domain can be transformed to a function ' $F$ ' in the ' $u$ ' domain using the integral

$$F(u) = \int_a^b f(x) K(u, x) dx. \quad (1)$$

This process is known as an integral transform and the function  $K(u, x)$  is the kernel of the transformation. Here the integral transform maps the problem in a given domain to another domain in which it is simpler to solve. Laplace transform is one of the widely used integral transforms in physics. It is very useful in solving convolution integral equations and differential equations. The inverse Laplace transform is an equally important transform with several applications. In general an inverse Laplace transform is carried out using a Bromwich contour integral in the complex plane [1]. But this practice of using a complex variable technique is done only for the sake of convenience. In fact an inverse Laplace transform based only on real variables was introduced by Post [2] and later it was refined by Widder [3]. This method has been investigated in several works [4, 5, 6] considering different applications.

The kernel of a Laplace transform is an exponential function of the form  $\exp(-st)$ . Tsallis introduced [7] generalizations of the logarithm and the exponential functions as follows:

$$\ln_q(x) \equiv \frac{x^{1-q} - 1}{1 - q}; \quad \exp_q(x) \equiv [1 + (1 - q)x]^{\frac{1}{1-q}}, \quad (2)$$

where  $q \in \mathbb{R}_+$  is the generalization parameter. These functions are generally referred to as Tsallis  $q$ -logarithm and Tsallis  $q$ -exponential and are inversely related. The  $q$ -logarithm and the  $q$ -exponential function reduces to the usual logarithm and exponential functions in the  $q \rightarrow 1$  limit. These generalized functions have been investigated in a wide variety of fields like astrophysics [8, 9], high energy physics [10, 11] neutrino physics [12], mathematical physics [13, 14] and nonequilibrium statistical physics [15].

A generalization of the Laplace transform has been done in Ref. [16, 17, 18, 19] using the Tsallis  $q$ -exponential. Here again the inverse was based on using a contour integral in the complex plane. In our present work we extend the concept of Post and Widder's method to the generalized Laplace transform based on Tsallis  $q$ -exponential.

The work is organized as follows: In Section 2, we give a brief introduction to the Laplace transform, Post-Widders technique and the Tsallis  $q$ -exponential. The Laplace and inverse Laplace based on type - I kernel is defined in the third section. The properties of Laplace transform are derived in Section 4. In Section 5 we give the  $q$ -generalization of the Post-Widder's method for the inverse Laplace transform. The Laplace transform and the inverse Laplace transform based on the Widder's method for different functions are computed in Section 6. As an application of this method we calculate the density of states of a  $D$ -dimensional classical ideal gas and a linear harmonic oscillator from the partition function using the Post-Widder's method based inverse Laplace transform.

## 2. A primer on Laplace transform, Post- Widder's technique and Tsallis $q$ -exponential

In this section we briefly review the concept of Laplace transform and its inverse. Then we describe the Post-Widder's method to calculate the inverse Laplace transform. The Laplace transform of a function  $f(t)$  denoted by  $\mathcal{L}[f(t)] = F(s)$  is defined as

$$F(s) = \int_0^\infty \exp(-st) f(t) dt. \quad (3)$$

Here  $f \in S(\mathbb{R})$  where  $S(\mathbb{R})$  represents the Schwartz space of functions  $f : \mathbb{R} \rightarrow \mathbb{C}$  with  $f \in C^\infty(\mathbb{R})$ , i.e.,  $f$  is infinitely differentiable on  $\mathbb{R}$ . In general  $s = \sigma + i\tau$  with  $\sigma$  and  $\tau$  being real numbers. The integral converges when  $\text{Re}[s] = \sigma > 0$  and for  $\sigma < 0$ ,  $F(s) = 0$ . For the Laplace transform of  $F(s)$ , the inverse transform is defined as

$$f(t) = \frac{1}{2\pi i} \lim_{\tau \rightarrow \infty} \int_{\sigma - i\tau}^{\sigma + i\tau} \exp(st) F(s) ds. \quad (4)$$

The direct Laplace transform Eq. (3) and the inverse Laplace transform Eq. (4) are inverses of each other for the functions  $f \in S(\mathbb{R})$ . In order to calculate the inverse Laplace transform we need to perform a Bromwich contour integration over the complex plane. Below we explain an alternative method introduced by Post and Widder which does a Laplace inverse using only real variables.

Let us consider the  $n^{\text{th}}$  derivative of the Laplace transform  $F(s)$  with respect to the variable 's',

$$\frac{d^n}{ds^n} F(s) \equiv F^{(n)}(s) = (-1)^n \int_0^\infty t^n \exp(-st) f(t) dt. \quad (5)$$

We use the following three steps, (i) First we use the variable transformation  $s = n/x$ , (ii) We follow it by multiplying the numerator and the denominator by  $x^n$  and (iii) Finally we use the variable change  $y = t/x$ , and recast the integral in Eq. (5) to

$$F^{(n)}\left(\frac{n}{x}\right) = (-1)^n x^{n+1} \int_0^\infty (ye^{-y})^n f(xy) dy. \quad (6)$$

Here we would like to point out that the function  $ye^{-y}$  has a single maximum at  $y = 1$  and for the function  $(ye^{-y})^n$  this maximum is sharply peaked at  $y = 1$  and hence Eq. (6) can be rewritten as

$$F^{(n)}\left(\frac{n}{x}\right) \approx (-1)^n x^{n+1} f(x) \int_0^\infty (ye^{-y})^n dy. \quad (7)$$

Evaluating the integral we get

$$F^{(n)}\left(\frac{n}{x}\right) \approx (-1)^n n! \left(\frac{x}{n}\right)^{n+1} f(x). \quad (8)$$

In the limit  $n \rightarrow \infty$ , we can use the transformation  $y = \frac{t}{x}|_{y=1} \Rightarrow x = t$  and so  $s = \frac{n}{x} \Rightarrow s = \frac{n}{t}$  we can rewrite Eq. (8) as

$$f(t) = \lim_{n \rightarrow \infty} \frac{(-1)^n}{n!} s^{n+1} F^{(n)}(s)|_{s=n/t}. \quad (9)$$

Thus we observe that the Laplace transform and the inverse Laplace transform can be expressed as functions of real variables alone. This sequence converges very fast since the rate of convergence is at least  $1/n$ .

The Laplace transform exists for a piecewise continuous function of exponential order. Similarly, the  $q$ -Laplace transform has been defined for a piecewise continuous function of  $q$ -exponential order. The  $q$ -exponential based Laplace transform can be defined using three different types of kernels as noted in Ref. [16] and these kernels are

$$K_I(q; s, t) = \exp_q(-st), \quad (10)$$

$$K_{II}(q; s, t) = [\exp_q(-t)]^s, \quad (11)$$

$$K_{III}(q; s, t) = [\exp_q(t)]^{-s}. \quad (12)$$

Of these three kernels, Laplace transform has been defined using the first and the second kernels in Ref. [16, 18, 19]. In the extensive limit ( $q \rightarrow 1$ ), both these generalized Laplace transforms reduce to the ordinary Laplace transform. For the generalized Laplace transform based on the first kernel [19], the inverse Laplace transform has not been defined. But for the case where the Laplace transform was defined using the second kernel, the inverse transform was defined using the complex integration method.

### 3. Laplace and Inverse transform based on type - I kernel

The Laplace transform due to type-I kernel which was introduced in [19] is examined in this section. For this generalization, the inverse Laplace transform has not been introduced so far and we restrict and present only the  $q < 1$  case here. Below we define the inverse Laplace transform and prove its inverse nature. A function ' $f$ ' is said to be of  $q$ -exponential order ' $c$ ', if there exists  $c, M > 0, T > 0$ , such that  $|f(t)| \leq M \exp_q(ct) \forall t > T$ . If a function is piecewise continuous and is of  $q$ -exponential order  $c$ , then  $F_q(s) = L_q[f(t)]$  exists for  $s > c$  and  $\lim_{s \rightarrow \infty} F(s) = 0$ . Under these conditions, the  $q$ -Laplace transform is defined as

$$L_q[f(t)](s) = F_q(s) = \int_0^\infty f(t) \exp_q(-st) dt. \quad (13)$$

The corresponding inverse Laplace transform is defined as

$$L_q^{-1}[F_q(s)](t) = f(t) = \frac{2-q}{2\pi i} \int_{c-i\infty}^{c+i\infty} F_q(s) [\exp_q(-st)]^{2q-3} ds, \quad (14)$$

where in the limit  $q \rightarrow 1$ , we have  $[\exp_q(-st)]^{2q-3} \rightarrow \exp(st)$ . Here  $c$  is a real constant that exceeds real part of all the singularities of  $F_q(s)$ . To prove the inverse relationship between  $L_q$  and  $L_q^{-1}$ , we verify the following two identities:

$$f(t) = L_q^{-1}[L_q[f(t)]], \quad (15)$$

$$F_q(s) = L_q[L_q^{-1}[F_q(s)]]. \quad (16)$$

Using the inverse Laplace transform we can write (16),

$$\begin{aligned} L_q[L_q^{-1}[F_q(s)]] &= \int_0^\infty \exp_q(-st) L_q^{-1}[F_q(s)] \\ &= \int_0^\infty dt \exp_q(-st) \frac{(2-q)}{2\pi i} \int_{c-i\infty}^{c+i\infty} ds' F_q(s') [\exp_q(-s't)]^{2q-3}. \end{aligned} \quad (17)$$

We can rewrite the above expression as

$$L_q[L_q^{-1}[F_q(s)]] = \frac{(2-q)}{2\pi i} \int_{c-i\infty}^{c+i\infty} ds' F_q(s') \int_0^\infty dt \exp_q(-st) [\exp_q(-s't)]^{2q-3}. \quad (18)$$

The second integral converges when  $\text{Re}[s'] = c < \text{Re}[s]$  and the resulting solution is

$$\mathcal{I}_q(s, s') = \int_0^\infty dt \exp_q(-st) [\exp_q(-s't)]^{2q-3} = \frac{1}{(2-q)} \frac{1}{(s-s')}. \quad (19)$$

Substituting this in Eq. (18) we get

$$L_q[L_q^{-1}[F_q(s)]] = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} ds' \frac{F_q(s')}{(s-s')}. \quad (20)$$

The function  $F_q(s)$  is an entire function and does not have any poles in the complex plane. The integrand  $F_q(s')/(s-s')$  has a simple pole at  $s = s'$ . To evaluate the integral we draw a straight line at  $s'$  and an arc enclosing the pole of the integrand. The solution of the integral is

$$\int_{c-i\infty}^{c+i\infty} ds' \frac{F_q(s')}{(s-s')} = 2\pi i F_q(s). \quad (21)$$

Substituting this in Eq. (20) we can observe that  $L_q[L_q^{-1}[F_q(s)]] = F_q(s)$ . Next we verify the first identity Eq. (15) as follows:

$$L_q^{-1}[L_q[f(t)]] \equiv L_q^{-1}[F_q(s)] = L_q^{-1} \left[ \int_0^\infty dt \exp_q(-st) f(t) \right]. \quad (22)$$

Using the definition of the inverse Laplace transform

$$L_q^{-1}[L_q[f(t)]] = \frac{2-q}{2\pi i} \int_{c-i\infty}^{c+i\infty} [\exp_q(-st)]^{2q-3} \int_0^\infty dt' f(t') \exp_q(-st') ds. \quad (23)$$

Rewriting the above equation (23) we arrive at

$$L_q^{-1}[L_q[f(t)]] = \frac{2-q}{2\pi i} \int_0^\infty dt' f(t') \int_{c-i\infty}^{c+i\infty} [\exp_q(-st)]^{2q-3} \exp_q(-st') ds. \quad (24)$$

Using the definition of the Dirac delta function based on the  $q$ -exponential described in Ref. [20, 21] we get

$$L_q^{-1}[L_q[f(t)]] = \int_0^\infty dt' f(t') \delta(t-t') \equiv f(t). \quad (25)$$

Thus in the present section, we have introduced the inverse for the Laplace transform defined using the type-I kernel using the complex integration technique.

#### 4. Properties of $q$ -Laplace transform

In this section, we list some of the properties of the  $q$ -Laplace transform:

(i)  $I^{st}$  Identity on Limits:

$$\lim_{s \rightarrow \infty} s L_q[f(t)] = \lim_{t \rightarrow 0} \frac{f(t)}{1 + (1-q)}, \quad (26)$$

*Proof:* Let us consider a general convergent function which can be expressed in terms of a power series  $f(t) = \sum_{n=0}^\infty a_n t^n$ . The  $q$ -Laplace transform of this general function is

$$L_q[f(t)] = \int_0^\infty \left( \sum_{n=0}^\infty a_n t^n \right) [1 - (1-q)st]^{\frac{1}{1-q}} dt. \quad (27)$$

Since the function  $f(t)$  is a convergent function we can rewrite the above Equation (27) as

$$L_q[f(t)] = \sum_{n=0}^{\infty} a_n \int_0^{\infty} t^n [1 - (1-q)st]^{\frac{1}{1-q}} dt. \quad (28)$$

To solve Eq. (28) we use the integration by parts method choosing  $u = t^n$  and  $dV = [1 - (1-q)st]^{\frac{1}{1-q}}$ . The resulting solution reads:

$$sL_q[f(t)] = -\frac{1}{1+(1-q)} \left( f(t)[1 - (1-q)st]^{\frac{1}{1-q}+1} \Big|_0^{\infty} - \int_0^{\infty} [1 - (1-q)st] \left[ \frac{d}{dt} f(t) \right] \exp_q(-st) dt \right). \quad (29)$$

On applying the limits corresponding to the integration in Eq. (29) we get

$$sL_q[f(t)] = \frac{f(0)}{1+(1-q)} + \frac{1}{1+(1-q)} \int_0^{\infty} [1 - (1-q)st] \frac{df(t)}{dt} [1 - (1-q)st]^{\frac{1}{1-q}} dt. \quad (30)$$

Under the limiting condition  $s \rightarrow \infty$ , Eq. (30) gives

$$\lim_{s \rightarrow \infty} sL_q[f(t)] = \lim_{s \rightarrow \infty} \frac{f(t)}{1+(1-q)}. \quad (31)$$

We can observe that the RHS is independent of  $s$  and can be expressed as a limiting value of the parameter  $t$  and this gives us

$$\lim_{s \rightarrow \infty} sL_q[f(t)] = \lim_{t \rightarrow 0} \frac{f(t)}{1+(1-q)}. \quad (32)$$

Thus we prove the first identity on the limits of a Laplace transform.

(ii)  $\Pi^{nd}$  Identity on Limits:

$$\lim_{s \rightarrow 0} sL_q[f(t)] = \lim_{t \rightarrow \infty} \frac{f(t)}{1+(1-q)}. \quad (33)$$

*Proof:* To prove this limit let us consider the Eq. (29) and evaluate the limit,  $s \rightarrow 0$  and the resulting expression is

$$\begin{aligned} \lim_{s \rightarrow 0} sL_q[f(t)] &= \lim_{s \rightarrow 0} \frac{f(0)}{1+(1-q)} + \frac{1}{1+(1-q)} \int_0^{\infty} \frac{df}{dt} dt \\ &= \frac{f(0)}{1+(1-q)} + \frac{f(t)}{1+(1-q)} \Big|_0^{\infty}. \end{aligned} \quad (34)$$

Applying the limits we get

$$\lim_{s \rightarrow 0} sL_q[f(t)] = \frac{f(\infty)}{1+(1-q)}. \quad (35)$$

The RHS of the equation can be rewritten as

$$\lim_{s \rightarrow 0} sL_q[f(t)] = \lim_{t \rightarrow \infty} \frac{f(t)}{1+(1-q)}. \quad (36)$$

Thus we prove the second identity on the Laplace transform.

(iii) Scaling

$$L_q[f(at)] = \frac{1}{a} F_q(s/a). \quad (37)$$

*Proof:* Let us consider the Laplace transform of a function  $f(at)$

$$L_q[f(at)] = \int_0^\infty dt [1 - (1-q)st]^{\frac{1}{1-q}} f(at). \quad (38)$$

Substituting  $at = x$ , we get

$$L_q[f(at)] = \frac{1}{a} \int_0^\infty dx \left[ 1 - (1-q)\frac{sx}{a} \right]^{\frac{1}{1-q}} f(x) = \frac{1}{a} F_q(s/a). \quad (39)$$

Hence the scaling relation is proved.

(iv) Shifting

$$F_q(s - s_0) = L_q \left[ f(t) \exp_q \left[ \frac{s_0 t}{1 - (1-q)st} \right] \right]. \quad (40)$$

*Proof:* Let us consider the definition of the Laplace transform

$$F_q(s) = \int_0^\infty \exp_q(-st) f(t) dt, \quad (41)$$

and introduce a shift in 's' as ' $s \rightarrow s - s_0$ ' and this yields

$$F_q(s - s_0) = \int_0^\infty \exp_q(-(s - s_0)t) f(t) dt. \quad (42)$$

This can be rewritten as

$$F_q(s - s_0) = \int_0^\infty \exp_q(-st) \exp_q \left( \frac{s_0 t}{1 - (1-q)st} \right) f(t) dt. \quad (43)$$

Hence we have proved

$$F_q(s - s_0) = L_q \left[ f(t) \exp_q \left( \frac{s_0 t}{1 - (1-q)st} \right) \right]. \quad (44)$$

(v)  $q$ -translation

$$L_q[f(t)] [\exp_q(st_0)]^{1+(1-q)} = L_q \left[ f \left( \frac{t - t_0}{1 - (1-q)st_0} \right) \Theta \left( \frac{t - t_0}{1 - (1-q)st_0} \right) \right]. \quad (45)$$

*Proof:* The RHS of the above identity gives

$$\begin{aligned} I &\equiv L_q \left[ f \left( \frac{t' - t_0}{1 - (1-q)st_0} \right) \Theta \left( \frac{t' - t_0}{1 - (1-q)st_0} \right) \right] \\ &= \int dt' f \left( \frac{t' - t_0}{1 - (1-q)st_0} \right) \Theta \left( \frac{t' - t_0}{1 - (1-q)st_0} \right) \exp_q(-st'), \end{aligned}$$

where  $\Theta(x)$  is the Heaviside step function such that  $\Theta(x) = 0$  when  $x < 0$  and  $\Theta(x) = 1$  for  $x \geq 0$ . Using the scaling  $t = \frac{t' - t_0}{1 - (1-q)st_0}$ , we can rewrite the integral as

$$I = \int_{-\alpha}^\infty dt \exp_q(-st) f(t) [\exp_q(-st_0)]^{2-q} \Theta(t).$$

where  $\alpha = \frac{t_0}{1 - (1-q)st_0}$ . On applying the Heaviside step function we get

$$I = L_q[f(t)] [\exp_q(-st_0)]^{2-q}. \quad (46)$$

Hence the  $q$ -translation identity has been proved.

The properties of linearity and  $q$ -convolution have been established in Ref. [19] and here we are stating them just for the sake of completeness. The two properties of linearity and  $q$ -convolution are

(i) Linearity:

$$L_q[a_1 f_1(t) + a_2 f_2(t)] = a_1 L_q[f_1(t)] + a_2 L_q[f_2(t)]. \quad (47)$$

(ii)  $q$ -convolution:

Let  $f(t)$  and  $g(t)$  be two positive scalar functions of ' $t$ ', and  $F_q(s)$  and  $G_q(s)$  be their  $q$ -Laplace transforms, then

$$L_q[f(t) * g(t)] = F_q(s) * G_q(s), \quad (48)$$

where  $f(t) * g(t) = \int_0^t f(\tau) * g(t - \tau) d\tau$ .

*Laplace transform of  $q$ -derivatives and  $q$ -integrals:* The properties of the  $q$ -Laplace transform based on derivatives and integrals have been derived in Ref. [19]. For the sake of completeness, we give below the expression for the generalized Laplace transform of a derivative function

$$L_q \left[ \frac{d^n}{dt^n} f(t) \right] = - \left( f^{(n-1)}(t) + (1 - \delta_{1n}) \sum_{\ell=1}^{n-1} Q_{\ell-1}(q) s^\ell f^{(n-\ell-1)}(t) \right) \Big|_{t=0} + Q_{n-1}(q) s^n L_{\frac{a_{n+1}}{a_n}}(f(t))(a_n s), \quad (49)$$

where,  $Q_n(q) = \prod_{j=0}^n a_j$  with  $a_j = jq - (j-1)$  and  $\delta_{1j}$  is the Kronecker delta function. The Laplace transform of the integral reads:

$$L_q \left[ \int_0^\infty f(x) dx \right] = \frac{2-q}{s} L_{\frac{1}{2-q}}[f(t)](s(2-q)). \quad (50)$$

In the present work, we derive the  $q$ -Laplace transform of  $q$ -calculus of functions. The concept of  $q$ -calculus was first introduced in [22], where the authors defined the derivative and integral based on  $q$ -deformation. Many such derivatives and integrals were investigated in subsequent works [13, 16, 19]. For the present work we use the  $q$ -derivative defined in [23] and its corresponding  $q$ -deformed integral. The relevant  $q$ -derivative and  $q$ -integral operators are

$$D_q(s) = \frac{1}{1 - (1-q) \left( s \frac{d}{ds} \right)} \frac{d}{ds}; \quad \int d_q x = \int dx \left[ 1 - (1-q)x \frac{d}{dx} \right]. \quad (51)$$

(i) Derivative of Laplace transform:

$$D_q^n(s) \{L_q[f(t)]\}(s) = L_q[(-t)^n f(t)]. \quad (52)$$

*Proof:* Let us consider the first derivative of the  $q$ -Laplace transform

$$\begin{aligned} D_q \{L_q[f(t)]\}(s) &= D_q(s) \left\{ \int_0^\infty dt \exp_q(-st) f(t) \right\} \\ &= \int_0^\infty dt D_q(s) \exp_q(-st) f(t) \\ &= L_q[(-t)f(t)]. \end{aligned} \quad (53)$$

The second derivative of the  $q$ -Laplace transform is

$$D_q^2 \{L_q[f(t)]\}(s) = L_q[(-t)^2 f(t)]. \quad (54)$$

For the  $n^{th}$   $q$ -derivative we get

$$D_q^n \{L_q[f(t)]\}(s) = L_q[(-t)^n f(t)]. \quad (55)$$



(ii) Integral of a Laplace transform:

$$\int_0^\infty d_q s F_q(s) = L_q \left\{ \frac{f(t)}{t} \right\}. \quad (56)$$

Let us consider the definition of the generalized Laplace transform apply the  $q$ -integral operator

$$\int_0^\infty d_q s F_q(s) = \int_s^\infty d_q s \int_0^\infty \exp_q(-st) f(t) dt. \quad (57)$$

Rearranging the order of the integrals and evaluating the  $q$ -integral with respect to 's' gives

$$\int_0^\infty d_q s F_q(s) = \int_0^\infty \frac{1}{t} \exp_q(-st) f(t) dt = L_q \left[ \frac{f(t)}{t} \right]. \quad (58)$$

Hence proved.

## 5. Post-Widders method of inverse Laplace transform

The inverse Laplace transform can be calculated using a Bromwich contour integral over the complex plane. As an alternative we introduce a  $q$ -deformed generalization of the Post-Widder's method to compute the inverse Laplace transform using only real variables. To derive the  $q$ -Widder's formula we will have to rewrite the type I kernel as  $K_q(s, t) = \exp \left( \frac{1}{1-q} \ln(1 - (1-q)st) \right)$ . We then have to take the  $k^{th}$  derivative of the Laplace transform and scale the function  $f(t)$  to  $f(\xi_m t)$  and the resulting expression is

$$F_q^{(k)}(s) = \int_0^\infty dt [-(1-q)t]^k \frac{\Gamma \left( \frac{2-q}{1-q} \right)}{\Gamma \left( \frac{2-q}{1-q} - k \right)} (1 - (1-q)st)^{\frac{1}{1-q}-k} f(t\xi_m). \quad (59)$$

Through a change of variables  $t = xy$  and  $s = k/x$  we get:

$$F_q^{(k)} \left( \frac{k}{x} \right) = [-(1-q)]^k x^{k+1} \frac{\Gamma \left( \frac{2-q}{1-q} \right)}{\Gamma \left( \frac{2-q}{1-q} - k \right)} \int_0^\infty (1 - (1-q)ky)^{\frac{1}{1-q}-k} y^k dy f(xy\xi_m). \quad (60)$$

The function  $y^k (1 - (1-q)ky)^{\frac{1}{1-q}-k}$  has a single maximum which is sharply peaked at  $y = 1$ . Hence we can replace the function  $f(\xi_m xy)$  by  $f(\xi_m x)$  and get

$$F_q^{(k)} \left( \frac{k}{x} \right) = f(\xi_m x) [-(1-q)]^k x^{k+1} \frac{\Gamma \left( \frac{2-q}{1-q} \right)}{\Gamma \left( \frac{2-q}{1-q} - k \right)} \int_0^\infty (1 - (1-q)ky)^{\frac{1}{1-q}-k} y^k dy. \quad (61)$$

The solution of the integral in the above equation is

$$\int_0^\infty (1 - (1-q)ky)^{\frac{1}{1-q}-k} y^k dy = \frac{\Gamma(k+1) \Gamma \left( \frac{2-q}{1-q} - k \right)}{\Gamma \left( \frac{3-2q}{1-q} \right)}. \quad (62)$$

Replacing the value of the integral in Eq. (61) and simplifying we have

$$F_q^{(k)} \left( \frac{k}{x} \right) = (-1)^k f(x\xi_m) x^{k+1} \frac{\Gamma(k+1)}{k^{k+1}(2-q)}. \quad (63)$$

where  $\xi_m = \left[ \frac{1+(1-q)}{Q_m(2-q)} \right]^{\frac{1}{m-1}}$  valid for  $m \geq 2$  and ill defined for  $m = 1$  and the polynomial  $Q_m(q) = \prod_{j=1}^m (1 - (1-q)j)$  exists for only for  $j \geq 1$ . Substituting  $t = \xi_m x$ , the  $q$ -deformed Widder's formula for the inverse Laplace transform is

$$f(t) = \lim_{k \rightarrow \infty} \frac{(-1)^k}{\Gamma(k+1)} F_q^{(k)}(s) (2-q) s^{k+1} \Big|_{s=\frac{k\xi_m}{t}}. \quad (64)$$

## 6. Inverse Laplace transform for some elementary functions

In this section we evaluate the Laplace transform of some elementary functions and then compute their inverse transform using Post-Widders method.

### Algebraic function:

Let us consider an algebraic function  $f(t) = t^{m-1}$ , the  $q$ -Laplace transform of this function is

$$L_q[t^{m-1}] = \int_0^\infty dt \exp_q(-st) t^{m-1}. \quad (65)$$

Evaluating the Laplace transform we get

$$F_q(s) = \frac{\Gamma(m)}{Q_m(2-q)} \frac{1}{s^m}, \quad \text{for } m \geq 2. \quad (66)$$

The inverse of the  $q$ -Laplace transform can be computed using a  $q$ -version of the Widder's formula. For this we calculate the  $k^{th}$ -derivative of  $F_q(s)$

$$F_q^k(s) = \frac{1}{Q_m(2-q)} (-1)^k \frac{\Gamma(m+k)}{s^{m+k}}. \quad (67)$$

Substituting Eq. (67) in the  $q$ -Widder's formula we get

$$\begin{aligned} f(t) &= \lim_{k \rightarrow \infty} \frac{(-1)^k}{\Gamma(k+1)} \left( \frac{1}{Q_m(2-q)} (-1)^k \frac{\Gamma(m+k)}{s^{m+k}} \right) Q_1(2-q) s^{k+1} \Big|_{s=\frac{k\xi_m}{t}}, \\ &= \lim_{k \rightarrow \infty} \frac{(-1)^k}{\Gamma(k+1)} \left( \frac{1}{Q_m(2-q)} (-1)^k \frac{\Gamma(m+k)}{s^{m-1}} \right) Q_1(2-q) \Big|_{s=\frac{k\xi_m}{t}}. \end{aligned} \quad (68)$$

where,  $\xi_m = \left( \frac{Q_1(2-q)}{Q_j(2-q)} \right)^{\frac{1}{j-1}}$  and the subscript ' $j$ ' depends on the exponent of ' $s$ ', since the exponent of ' $s$ ' is ' $m-1$ ' the value of ' $j$ ' is ' $m$ '. Substituting the value of ' $s$ ' we get:

$$f(t) = \frac{Q_1(2-q)}{Q_m(2-q)} \frac{t^{m-1}}{\xi_m^{m-1}} \left( \lim_{k \rightarrow \infty} \frac{k^{1-m} \Gamma(m+k)}{\Gamma(k+1)} \right). \quad (69)$$

On substitution of the limits and simplifying Eq. (69) we get the algebraic limit  $f(t) = t^{m-1}$ . This validates the Post-Widder's method of computing inverse Laplace transform for any Algebraic function.

### Exponential functions:

*Regular exponential:* Let us consider a generalized form of the exponential function  $f(t) = \exp(\pm \sigma_\epsilon t)$  where  $\sigma_\epsilon = i^\epsilon \alpha$  with  $\alpha \in \mathcal{R}$  and  $\epsilon = 0$  or  $1$ . We can write this definition in the following way:

$$f(t) = \exp(\pm \sigma_\epsilon t) = \begin{cases} \exp(\pm \alpha t) & \text{for } \epsilon = 0 \\ \exp(\pm i \alpha t) & \text{for } \epsilon = 1 \end{cases} \quad (70)$$

In terms of hypergeometric functions, the classical exponential given in Eq. (70) can be expressed as:

$$f(t) = \exp(\pm\sigma_\epsilon t) = {}_0F_0(\ ;\ ;\ ;\pm\sigma_\epsilon t). \quad (71)$$

The  $q$ -Laplace transform of  $f(t)$  gives:

$$\begin{aligned} L_q[{}_0F_0(\ ;\ ;\ ;\pm\sigma_\epsilon t)] &= \int_0^\infty dt \exp_q(-st) {}_0F_0(\ ;\ ;\ ;\pm\sigma_\epsilon t) \\ &= \frac{1}{s Q_1(2-q)} {}_1F_1\left(1; \frac{3-2q}{1-q}; \pm \frac{\sigma_\epsilon}{(1-q)s}\right). \end{aligned} \quad (72)$$

To find the  $q$ -inverse Laplace transform using Widder like formula for the Tsallis  $q$ -case, we take the  $k^{th}$ -derivative of  $F_q(s)$  given by Eq. (72) which is

$$F_q^{(k)}(s) = \frac{1}{Q_1(2-q)} \left[ \sum_{n=0}^{\infty} \frac{1}{\left(\frac{3-2q}{1-q}\right)_n} \frac{\left(\pm \frac{\sigma_\epsilon}{1-q}\right)^n}{n!} (-1)^k \frac{\Gamma(n+k+1)}{s^{n+k+1}} \right]. \quad (73)$$

Substituting Eq. (73) in the  $q$ -Widder's formula, Eq. (64) and simplifying it we get:

$$f(t) = \sum_{n=0}^{\infty} \frac{1}{\left(\frac{3-2q}{1-q}\right)_n} \frac{1}{n!} \left(\pm \frac{\sigma_\epsilon t}{(1-q)\xi_m}\right)^n \left[ \lim_{k \rightarrow \infty} k^{1-n} \left(1 + \frac{n}{k}\right) \frac{\Gamma(k+n)}{\Gamma(k+1)} \right]. \quad (74)$$

Applying the limits in Eq. (74) and simplifying the expression we get the initial function  $f(t) = \exp(\pm\sigma_\epsilon t)$  validating the Post-Widder's method.

*Generalized exponential:* In analogy with the exponential, we can write the generalized exponential as  $f(t) = \exp_{q'}(\pm\sigma_\epsilon t)$  where  $\sigma_\epsilon = i^\epsilon \alpha$  with  $\alpha \in \mathcal{R}$  and  $\epsilon = 0$  or  $1$ . This definition of  $f(t)$  can be written as

$$f(t) = \exp_{q'}(\pm\sigma_\epsilon t) = \begin{cases} \exp_{q'}(\pm\alpha t) & \text{for } \epsilon = 0 \\ \exp_{q'}(\pm i\alpha t) & \text{for } \epsilon = 1 \end{cases} \quad (75)$$

The generalized exponential given in equation (75) can be written in terms of hypergeometric functions

$$f(t) = \exp_{q'}(\pm\sigma_\epsilon t) = {}_1F_0\left(-\frac{1}{1-q'}; \ ;\ \mp(1-q')\sigma_\epsilon t\right). \quad (76)$$

Applying the  $q$ -Laplace transform on  $f(t)$  we get:

$$\begin{aligned} L_q[f(t)] &= \int_0^\infty dt \exp_q(-st) {}_1F_0\left(-\frac{1}{1-q'}; \ ;\ \mp(1-q')\sigma_\epsilon t\right) \\ &= \frac{1}{s Q_1(2-q)} {}_2F_1\left(1, -\frac{1}{1-q'}; \frac{3-2q}{1-q}; \mp \frac{(1-q')\sigma_\epsilon}{(1-q)s}\right). \end{aligned}$$

To find the inverse  $q$ -Laplace transform, using the Widder's method we compute the  $k^{th}$  derivative of  $F_q(s)$  given in the previous expression. The derivative so calculated is

$$F_q^{(k)}(s, q') = \sum_{n=0}^{\infty} \frac{\left(-\frac{1}{1-q'}\right)_n (\mp(1-q')\sigma_\epsilon)^n}{Q_{n+1}(2-q) n!} \frac{(-1)^k \Gamma(n+k+1)}{s^{n+k+1}}. \quad (77)$$

Thus, to find the  $q$ -Widders formula we substitute Eq. (77) in Eq. (64) in this case to get:

$$f(t; q') = Q_1(2-q) \left( \sum_{n=0}^{\infty} \frac{\left(-\frac{1}{1-q'}\right)_n (\mp(1-q')\sigma_\epsilon t)^n}{Q_{n+1}(2-q) \xi_m^n n!} \left[ \lim_{k \rightarrow \infty} \frac{(n+k)(k)_n}{k^{2+n}} \right] \right). \quad (78)$$

To get the inverse  $q$ -transform we apply the limits in the above expression and simplify using the identities of Pochhammer and this gives us the generalized exponential function defined in Eq. (78). This result confirms the use of Post-Widder's method.

### Gaussian function:

*Regular Gaussian:* The  $q$ -Laplace transform of a Gaussian function  $f(t) = \exp(-\alpha t^2)$  is

$$L_q [\exp(-\alpha t^2)] = \int_0^\infty dt \exp_q(-st) \exp(-\alpha t^2). \quad (79)$$

and can be written in terms of the hypergeometric function as follows:

$$F_q(s) = \frac{1}{s Q_1(2-q)} {}_2F_2 \left( 1, \frac{1}{2}; \frac{3-2q}{2(1-q)}, \frac{4-3q}{2(1-q)}; -\frac{\alpha}{(1-q)^2 s^2} \right). \quad (80)$$

The  $q$ -inverse Laplace transform can be found using the Widder's method and for this purpose we evaluate the  $k^{th}$ -derivative of  $F_q(s)$  which is

$$F_q^{(k)}(s) = \frac{1}{Q_1(2-q)} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(-\frac{\alpha}{(1-q)^2}\right)^n}{\left(\frac{3-2q}{2(1-q)}\right)_n \left(\frac{4-3q}{2(1-q)}\right)_n} \frac{(-1)^k (2n+k)!}{(2n)!} \frac{1}{s^{2n+k+1}}. \quad (81)$$

Using Eq. (81) in Eq. (64) we can derive the Widder's formula, for the Gaussian function as

$$f(t) = \left[ \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(-\frac{\alpha}{(1-q)^2}\right)^n}{\left(\frac{3-2q}{2(1-q)}\right)_n \left(\frac{4-3q}{2(1-q)}\right)_n} \left( \lim_{k \rightarrow \infty} \frac{(-1)^k}{\Gamma(k+1)} \frac{(-1)^k (2n+k)!}{k^{2n}} \right) \frac{t^{2n}}{(2n)! \xi_m^{2n}} \right]. \quad (82)$$

Applying the limiting value for  $k$  as

$$\lim_{k \rightarrow \infty} \frac{(-1)^k}{\Gamma(k+1)} \frac{(-1)^k (2n+k)!}{k^{2n}} = \lim_{k \rightarrow \infty} \frac{\Gamma(2n+k+1)}{\Gamma(k+1)} k^{1-(2n+1)} \rightarrow 1.$$

Using the identities of Pochhammer symbol  $\left(\frac{3-2q}{2(1-q)}\right)_n \left(\frac{4-3q}{2(1-q)}\right)_n = \frac{1}{4^n} \frac{Q_{2n+1}(2-q)}{(1-q)^{2n} Q_1(2-q)}$ , and the identity of factorials  $(2n)! = 4^n n! \left(\frac{1}{2}\right)_n$  we can simplify Eq. (82). This yields the Gaussian function  $f(t) = \exp(-\alpha t^2)$  confirming the usefulness of Post-Widder's method.

*Generalized Gaussian:* The  $q$ -Laplace transform of a classical Gaussian function  $f(t) = \exp_{q'}(-\alpha t^2)$  is:

$$L_q [\exp_{q'}(-\alpha t^2)] = \int_0^\infty dt \exp_q(-st) \exp_{q'}(-\alpha t^2), \quad (83)$$

which on integration gives the hypergeometric function

$$F_q(s; q') = \frac{1}{s Q_1(2-q)} {}_3F_2 \left( 1, \frac{1}{2}, -\frac{1}{1-q'}; \frac{3-2q}{2(1-q)}, \frac{4-3q}{2(1-q)}; \frac{(1-q')\alpha}{(1-q)^2 s^2} \right). \quad (84)$$

The inverse of the  $q$ -Laplace transform can be computed using the Widder's method and to find it we take the  $k^{th}$ -derivative of the function  $F_q(s)$  as

$$F_q^{(k)}(s; q') = \sum_{n=0}^{\infty} \frac{\left(-\frac{1}{1-q'}\right)_n}{Q_{2n+1}(2-q)} \frac{((1-q')\alpha)^n}{n!} (-1)^k \frac{\Gamma(k+2n+1)}{s^{2n+k+1}}. \quad (85)$$

Substituting Eq. (85) in the  $q$ -Widder's formula in Eq. (64) we get:

$$f(t, q') = \sum_{n=0}^{\infty} \frac{Q_1(2-q) \left(-\frac{1}{1-q'}\right)_n}{Q_{2n+1}(2-q)} \frac{((1-q')\alpha t^2)^n}{n! \xi^{2n}} \frac{1}{s^{2n}} \left[ \lim_{k \rightarrow \infty} \frac{\Gamma(k+2n+1)}{k^{2n} \Gamma(k+1)} \right]. \quad (86)$$

On applying the appropriate limits and using the Pochhammer identities, we will recover the generalized Gaussian function indicating that the Post-Widder's method works.

### Circular functions:

The circular functions (both cosine and sine) can be expressed in terms of hypergeometric function and written in a compact form as follows:

$$f(t) = (\alpha t)^\delta {}_0F_1 \left( ; \frac{1}{2} + \delta; -\frac{1}{4} \alpha^2 t^2 \right) = \begin{cases} \cos \alpha t & \text{for } \delta = 0 \\ \sin \alpha t & \text{for } \delta = 1 \end{cases} \quad (87)$$

The  $q$ -Laplace transform of Eq. (87) is

$$\begin{aligned} F_q(s) &= \frac{1}{s} \left( \frac{\alpha}{s} \right)^\delta \frac{1}{Q_{\delta+1}(2-q)} \\ &\quad \times {}_1F_2 \left( 1; \frac{1}{2} \left( \frac{3-2q}{1-q} + \delta \right), \frac{1}{2} \left( \frac{4-3q}{1-q} + \delta \right); -\frac{\alpha^2}{4(1-q)^2 s^2} \right) \\ &= \begin{cases} \cos \alpha t & \text{for } \delta = 0 \\ \sin \alpha t & \text{for } \delta = 1 \end{cases} \end{aligned} \quad (88)$$

For calculating the inverse  $q$ -Laplace transform, we find the  $k^{th}$  derivative of Eq. (88) as follows:

$$F_q^{(k)}(s) = \frac{1}{Q_{\delta+1}(2-q)} \sum_{n=0}^{\infty} \left( \frac{(-1)^{n+k} \Gamma(2n+\delta+k+1) \alpha^{2n+\delta}}{(1-q)^{2n} \left( \frac{3-2q}{1-q} + \delta \right)_{2n} (2n+\delta)!} \right) \frac{1}{s^{2n+\delta+k+1}}. \quad (89)$$

(i)  $\cos \alpha t$ : For  $\delta = 0$ , the  $k^{th}$  derivative of  $F_q^{(k)}(s)$  can be written from Eq. (89) as

$$F_q^{(k)}(s) = \frac{1}{Q_1(2-q)} \sum_{n=0}^{\infty} \left( \frac{(-1)^{n+k} \Gamma(2n+k+1) \alpha^{2n}}{(1-q)^{2n} \left( \frac{3-2q}{1-q} \right)_{2n} (2n)!} \right) \frac{1}{s^{2n+k+1}}. \quad (90)$$

To calculate the inverse Laplace transform we use the  $q$ -generalization of the Widder's method and for this we substitute Eq. (90) in Eq. (64) to obtain

$$f(t) = \sum_{n=0}^{\infty} \left( \frac{(-1)^n (\alpha t)^{2n}}{(1-q)^{2n} \left( \frac{3-2q}{1-q} \right)_{2n} (2n)! \xi^{2n}} \right) \left[ \lim_{k \rightarrow \infty} \frac{k^{1-(2n+1)} \Gamma(2n+k+1)}{\Gamma(k+1)} \right]. \quad (91)$$

Under the limiting conditions of  $k$  and using the proper Pochhammer identities we get

$$f(t) = \sum_{n=0}^{\infty} \frac{(-\alpha^2 t^2)^n}{(2n)!} = \cos \alpha t. \quad (92)$$

(ii)  $\sin \alpha t$ : Considering  $\delta = 1$ , the  $k^{th}$  derivative of  $F_q^{(k)}(s)$  can be written as

$$F_q^{(k)}(s) = \frac{1}{Q_2(2-q)} \sum_{n=0}^{\infty} \left( \frac{(-1)^{n+k} \Gamma(2n+k+2) \alpha^{2n+1}}{(1-q)^{2n} \left( \frac{4-3q}{1-q} \right)_{2n} (2n+1)!} \right) \frac{1}{s^{2n+k+2}}. \quad (93)$$

Substituting Eq. (93) in  $q$ -Widder's formula in Eq. (64) we get:

$$f(t) = \sum_{n=0}^{\infty} \left( \frac{(-1)^n (\alpha t)^{2n+1}}{(1-q)^{2n} \left( \frac{4-3q}{1-q} \right)_{2n} (2n+1)! \xi^{2n+1}} \right) \left[ \lim_{k \rightarrow \infty} \frac{k^{1-(2n+2)} \Gamma(2n+k+2)}{\Gamma(k+1)} \right]. \quad (94)$$

On application of the limiting value of  $k$  and using the Pochhammer identities we get:

$$f(t) = \sum_{n=0}^{\infty} (-1)^n \frac{(\alpha t)^{2n+1}}{(2n+1)!} = \sin \alpha t. \quad (95)$$

Thus we prove that the inverse of the  $q$ -Laplace transform works.

### Generalized circular functions:

Using the generalized exponential a set of circular functions have been defined in Ref. [24]. In terms of the hypergeometric functions these generalized circular functions can be written in a compact form as

$$\begin{aligned} f(t, q') &= (\alpha t)^\delta {}_0F_1 \left( \frac{1}{2} \left( -\frac{1}{1-q'} + \delta \right), \frac{1}{2} \left( -\frac{q'}{1-q'} + \delta \right); \frac{1}{2} + \delta; -(1-q')^2 \alpha^2 t^2 \right), \\ &= \begin{cases} \cos_{q'} \alpha t & \text{for } \delta = 0 \\ \sin_{q'} \alpha t & \text{for } \delta = 1 \end{cases} \end{aligned} \quad (96)$$

The  $q$ -Laplace transform of the generalized circular function is

$$\begin{aligned} F_q(s; q') &= \frac{1}{s} \left( \frac{\alpha}{s} \right)^\delta \frac{1}{Q_{\delta+1}(2-q)} \times \\ &\quad {}_3F_2 \left( 1, \frac{-a+\delta}{2}, \frac{-a+1+\delta}{2}; \frac{b+\delta}{2}, \frac{b+1+\delta}{2}; -\frac{(1-q')^2 \alpha^2}{(1-q)^2 s^2} \right), \\ &= \begin{cases} L_q [\cos_{q'}(\alpha t)] & \text{for } \delta = 0 \\ L_q [\sin_{q'}(\alpha t)] & \text{for } \delta = 1 \end{cases} \end{aligned} \quad (97)$$

where we have used the relations:

$$a = \frac{1}{1-q'}; \quad b = \frac{1}{1-q} + 2. \quad (98)$$

The  $k^{th}$ -derivative of equation (97) is calculated to find the inverse  $q$ -Laplace transform using the Widder's method. The  $k^{th}$  derivative is

$$\begin{aligned} F_q^{(k)}(s; q') &= \frac{1}{Q_{\delta+1}(2-q)}, \\ &\quad \sum_{n=0}^{\infty} \left( \frac{(-1)^{n+k} \Gamma(2n+\delta+k+1) \left( -\frac{1}{1-q'} + \delta \right)_{2n} \frac{(1-q')^{2n}}{(1-q)^{2n}} \alpha^{2n+\delta}}{Q_{\delta+1}(2-q) \left( \frac{3-2q}{1-q} + \delta \right)_{2n} (2n+\delta)! s^{2n+\delta+k+1}} \right). \end{aligned} \quad (99)$$

(i)  $\cos_{q'} \alpha t$ : For  $\delta = 0$ , the  $k^{th}$ -derivative of  $F_q^k(s; q')$  can be written as

$$F_q^{(k)}(s; q') = \frac{(-1)^k}{Q_1(2-q)} \sum_{n=0}^{\infty} \frac{(-1)^n \left( -\frac{1}{1-q'} \right)_{2n} \Gamma(2n+k+1) \frac{(1-q')^{2n}}{(1-q)^{2n}} \alpha^{2n}}{\left( \frac{3-2q}{1-q} \right)_{2n} (2n)! s^{2n+k+1}}. \quad (100)$$

Substituting in the formula for the Post-Widder's method in Eq. (64) we get:

$$f(t; q') = \sum_{n=0}^{\infty} \left( \frac{\left(-\frac{1}{1-q'}\right)_{2n} \left(\frac{(1-q')\alpha t}{1-q}\right)^{2n}}{(-1)^n \left(\frac{3-2q}{1-q}\right)_{2n} \xi^{2n} (2n)!} \right) \left[ \lim_{k \rightarrow \infty} \frac{k^{1-(2n+1)} \Gamma(2n+k+1)}{\Gamma(k+1)} \right]. \quad (101)$$

Applying the appropriate limits and also using Pochhammer identities we get

$$f(t; q') = \sum_{n=0}^{\infty} (-1)^n \frac{\left(-\frac{1}{1-q'}\right)_{2n} ((1-q')\alpha t)^{2n}}{(2n)!} = \cos_{q'}(\alpha t). \quad (102)$$

(ii)  $\sin_{q'} \alpha t$ : The choice of  $\delta = 1$  gives the generalized sine function. The  $k^{th}$ -derivative of  $F_q^{(k)}(s; q')$  of this function is

$$F_q^{(k)}(s; q') = \frac{(-1)^k}{Q_2(2-q)} \sum_{n=0}^{\infty} \frac{(-1)^n \left(-\frac{q'}{1-q'}\right)_{2n} \Gamma(2n+k+2) \frac{(1-q')^{2n}}{(1-q)^{2n}} \alpha^{2n+1}}{\left(\frac{4-3q}{1-q}\right)_{2n} (2n+1)! s^{2n+k+2}}. \quad (103)$$

Under the appropriate limits and using the proper Pochhammer identities we get

$$f(t, q') = \sum_{n=0}^{\infty} (-1)^{n+1} \left(-\frac{1}{1-q'}\right)_{2n+1} \frac{((1-q')\alpha t)^{2n+1}}{(2n+1)!} = \sin_{q'}(\alpha t). \quad (104)$$

Thus the  $q$ -Laplace transform and the inverse  $q$ -Laplace transform computed for the  $\sin_{q'} \alpha t$  give consistent results proving that the Post-Widder's method works.

### Hyperbolic functions:

The hyperbolic functions  $\cosh \alpha t$  and  $\sinh \alpha t$  can be expressed in terms of the hypergeometric function and can be written in compact form as:

$$f(t) = (\alpha t)^\delta {}_0F_1 \left( ; \frac{1}{2} + \delta; \frac{1}{4} \alpha^2 t^2 \right) = \begin{cases} \cosh \alpha t & \text{for } \delta = 0 \\ \sinh \alpha t & \text{for } \delta = 1 \end{cases} \quad (105)$$

The  $q$ -Laplace transform of Eq. (105) is:

$$\begin{aligned} F_q(s) &= \frac{1}{s} \left(\frac{\alpha}{s}\right)^\delta \frac{1}{Q_{\delta+1}(2-q)} {}_1F_2 \left( 1; \frac{b+\delta}{2}, \frac{b+1+\delta}{2} \frac{\alpha^2}{4(1-q)^2 s^2} \right) \\ &= \begin{cases} L_q[\cosh \alpha t] & \text{for } \delta = 0 \\ L_q[\sinh \alpha t] & \text{for } \delta = 1 \end{cases} \end{aligned} \quad (106)$$

where the factor  $b$  used in Eq. (106) is defined in Eq. (98). The inverse  $q$ -Laplace transform can be calculated using the Post-Widder's method. For this we compute the  $k^{th}$ -derivative of Eq. (106):

$$F_q^{(k)}(s) = \frac{1}{Q_{\delta+1}(2-q)} \sum_{n=0}^{\infty} \left( \frac{(-1)^k \Gamma(2n+\delta+k+1) \alpha^{2n+\delta}}{(1-q)^{2n} \left(\frac{3-2q}{1-q} + \delta\right)_{2n} (2n+\delta)!} \right) \frac{1}{s^{2n+\delta+k+1}}. \quad (107)$$

(i)  $\cosh \alpha t$ : For the  $\cosh \alpha t$  ( $\delta = 0$ ) function, the  $k^{th}$ -derivative is

$$F_q^{(k)}(s) = \frac{1}{Q_1(2-q)} \sum_{n=0}^{\infty} \left( \frac{(-1)^k \Gamma(2n+k+1) \alpha^{2n}}{(1-q)^{2n} \left(\frac{3-2q}{1-q}\right)_{2n} (2n)!} \right) \frac{1}{s^{2n+k+1}}. \quad (108)$$

Substituting in the  $q$ -generalization of Widder's formula, and applying the appropriate limits and Pochhammer identities we get

$$f(t) = \sum_{n=0}^{\infty} \frac{(\alpha^2 t^2)^n}{(2n)!} = \cosh \alpha t. \quad (109)$$

which confirms the  $q$ -Laplace transform as well as the generalized Widder's method defined in our work.

(ii)  $\sinh \alpha t$ : For the choice  $\delta = 1$  the  $k^{th}$ -derivative of  $F_q^{(k)}(s)$  is

$$F_q^{(k)}(s) = \frac{1}{Q_2(2-q)} \sum_{n=0}^{\infty} \left( \frac{(-1)^k \Gamma(2n+k+2) \alpha^{2n+1}}{(1-q)^{2n} \left(\frac{4-3q}{1-q}\right)_{2n} (2n+1)!} \right) \frac{1}{s^{2n+k+2}}. \quad (110)$$

Applying the appropriate limits and using the Pochhammer under the limit  $k \rightarrow \infty$  in the Equation above gives us

$$f(t) = \sum_{n=0}^{\infty} \frac{(\alpha t)^{2n+1}}{(2n+1)!} = \sinh \alpha t. \quad (111)$$

These results confirms that the  $q$ -Laplace transform as well as the inverse transform work as expected.

#### Generalized hyperbolic functions:

The generalized hyperbolic functions *viz* the  $\cosh_{q'}(\alpha t)$  and  $\sinh_{q'}(\alpha t)$  can be expressed in terms of the hypergeometric function and can be written in a compact form as

$$\begin{aligned} f(t, q') &= (\alpha t)^\delta {}_2F_1 \left( \frac{1}{2} \left( -\frac{1}{1-q'} + \delta \right), \frac{1}{2} \left( -\frac{q'}{1-q'} + \delta \right); \frac{1}{2} + \delta; (1-q')^2 \alpha^2 t^2 \right) \\ &= \begin{cases} \cosh_{q'} \alpha t & \text{for } \delta = 0 \\ \sinh_{q'} \alpha t & \text{for } \delta = 1 \end{cases} \end{aligned} \quad (112)$$

The  $q$ -Laplace transform of the  $q$ -hyperbolic function is

$$\begin{aligned} F_q(s; q') &= \frac{1}{s} \left( \frac{\alpha}{s} \right)^\delta \frac{1}{Q_{\delta+1}(2-q)} \times \\ &\quad {}_3F_2 \left( 1, \frac{-a+\delta}{2}, \frac{-(a-1)+\delta}{2}; \frac{b+\delta}{2}, \frac{b+1+\delta}{2}; \frac{(1-q')^2 \alpha^2}{(1-q)^2 s^2} \right) \\ &= \begin{cases} L_q [\cosh_{q'}(\alpha t)] & \text{for } \delta = 0 \\ L_q [\sinh_{q'}(\alpha t)] & \text{for } \delta = 1 \end{cases} \end{aligned} \quad (113)$$

where the factors used in the hypergeometric function are defined in Eq. (98). The  $k^{th}$ -derivative of  $F_q^{(k)}(s)$  is

$$F_q^{(k)}(s; q') = \sum_{n=0}^{\infty} \left( \frac{(-1)^k \Gamma(2n+\delta+k+1) \left( -\frac{1}{1-q'} + \delta \right)_{2n} \frac{(1-q')^{2n}}{(1-q)^{2n}} \alpha^{2n+\delta}}{Q_{\delta+1}(2-q) \left( \frac{3-2q}{1-q} + \delta \right)_{2n} (2n+\delta)! s^{2n+\delta+k+1}} \right). \quad (114)$$

For the case (i)  $\cosh_{q'}(\alpha t)$  we can choose  $\delta = 0$  we can calculate the  $k^{th}$  derivative of  $F_q(s, q')$  by allowing  $\delta = 1$  and substituting this in the  $q$ -Widder's formula and choosing the appropriate limits and using the Pochhammer identities we get:

$$f(t, q') = \sum_{n=0}^{\infty} \frac{\left( -\frac{1}{1-q'} \right)_{2n} ((1-q') \alpha t)^{2n}}{(2n)!} = \cosh_{q'}(\alpha t). \quad (115)$$



By choosing  $\delta = 1$  we can get the results corresponding to the case (ii)  $\sinh_{q'}(\alpha t)$  by repeating the procedure as being done in all the earlier cases. The final result reads:

$$f(t, q') = \sum_{n=0}^{\infty} \left( -\frac{q'}{1-q'} \right)_{2n} (1-q')^{2n} \frac{(\alpha t)^{2n+1}}{(2n+1)!} = \sinh_{q'}(\alpha t). \quad (116)$$

Thus we compute the generalized Laplace transform and its inverse using the Post-Widders method and we observe consistent results.

## 7. Applications in statistical mechanics

The Laplace transform has several applications in physics. Notable among these applications is the use of Laplace transform to interrelate the partition function and the density of states. The partition function is the fundamental quantity using which all the thermodynamic quantities are computed in the canonical ensemble which is used to study a closed system. The thermodynamic quantities are calculated using the density of states in the microcanonical ensemble which is used to describe an isolated system. In both the canonical and the microcanonical ensemble, the number of particles remain fixed. The density of states is the number of states of a given energy in a system and the partition function is the average of the density of states weighted by a Boltzmann distribution. Hence these two quantities *viz* partition function and density of states can be connected by a Laplace transform [25].

To investigate the statistical mechanics of systems involving long-range interactions a generalization of Boltzmann-Gibbs statistics was proposed by Tsallis [7] using the deformed  $q$ -exponential. When the average of the density of states is weighted by the Tsallis statistics i.e., using a deformed  $q$ -exponential, we can connect the partition function to the density matrix using a  $q$ -deformed Laplace transform. While both the partition function and the density of states can be calculated using first principles, the density of states which completely characterizes a isolated system is generally hard to compute. To overcome this one can calculate the partition function in the canonical ensemble and use the inverse Laplace transform to obtain the density of states. In our case we use the partition function of the Tsallis statistics and use the inverse of the  $q$ -Laplace transform and obtain the density of states.

*Classical Ideal gas:*

As an illustration let us consider a system of classical ideal gas in  $D$  dimensions with the Hamiltonian [25, 26, 27, 28]

$$H = \sum_{i=1}^{DN} \frac{p_i^2}{2m}, \quad (117)$$

where  $p$  and  $m$  are the momentum and mass of the gas. The nonextensive statistical mechanics begins with the introduction of a generalization of the entropy. The probability distribution for the microstates can be obtained by optimizing the entropy using physically appropriate constraints. In the course of the development of this field, three different ways of optimizations emerged. Of these three methods, the third constraint method [29, 30] known as the escort probability distribution is the one presently accepted. Using this constraint leads to implicit thermodynamics relations which are hard to solve. To overcome this one can solve the problem in the second constraint approach and the results obtained can be transformed to those in the third constraint using a transformation rule based on the inverse temperature. Since we

are interested in demonstrating the usefulness of Laplace transform, we prefer to avoid the complications arising with the third constraint and instead use the second constraint formulation. But we would like to note that any result we obtain in the second constraint can be transformed to third constraint. The partition function of the system in the second constraint formulation is

$$Z_q(\beta) = \frac{1}{h^{DN} N!} \int_0^\infty d\bar{x} d\bar{p} \exp_q \left( -\beta \sum_{i=1}^{DN} \frac{p_i^2}{2m} \right), \quad (118)$$

where  $\bar{x}$  and  $\bar{p}$  are  $(x_1, x_2, \dots, x_{DN})$ ,  $(p_1, p_2, \dots, p_{DN})$ . On calculating the integral we get:

$$Z_q(\beta) = \frac{V^N (2\pi m)^{\frac{DN}{2}}}{h^{DN} N!} \left( \frac{\Gamma\left(\frac{1}{1-q} + 1\right)}{(1-q)^{\frac{DN}{2}} \Gamma\left(\frac{1}{1-q} + \frac{DN}{2} + 1\right)} \right) \frac{1}{\beta^{\frac{DN}{2}}}. \quad (119)$$

Using the inverse of the  $q$ -Laplace transform one can calculate the density of states. Here we use the  $q$ -generalization of the Widder's method to get:

$$g(E) = \lim_{k \rightarrow \infty} \frac{(-1)^k}{\Gamma(k+1)} Z_q^{(k)}(\beta) \beta^{k+1} Q_1(2-q) \Big|_{\beta = \frac{k \xi_m}{E}}. \quad (120)$$

Substituting for the partition function Eq. (119) in the expression for the density of states Eq. (120), we apply the limiting value for  $\beta$  and simplify the expression to get:

$$g(E) = \frac{V^N (2\pi m)^{\frac{DN}{2}}}{h^{DN} N! \left(\frac{DN}{2} - 1\right)!} E^{\frac{DN}{2} - 1}, \quad (121)$$

which is the density of states of the classical ideal gas [25].

A system is in thermodynamic equilibrium when it is in thermal, mechanical and chemical equilibrium with its surroundings and consequently we need three macroscopic variables to describe the system. The macroscopic variable can be either an extensive variable dependent on the number of particle or an intensive variable which is independent of the particle number. The energy ( $E$ ) of a system is an example for an extensive variable and the temperature ( $T$ ) is an example of an intensive variable. Thus from the thermal, mechanical and chemical equilibrium we can choose from any of the two variables and hence we have in total eight different ensembles. Of the eight ensembles there are four ensembles in which the temperature is fixed and these are referred to as the isothermal ensemble and four in which the heat function is fixed and these ensembles are known as the adiabatic ensembles. The four isothermal ensembles are characterized by the partition function, while the four adiabatic ensembles are characterized by the density of states. The density of states is the number of states with same value for the heat function in a system and the partition function is the average of the density of states weighted by a distribution and in general cases we use a Boltzmann distribution. The eight ensembles can be grouped into four pairs with each pair comprising of an isothermal ensemble and an adiabatic ensemble. The partition of an isothermal ensemble can be related to the density of states of the adiabatic ensemble through a Laplace transform. In the discussion above we saw the correspondence between the microcanonical and the canonical ensemble. Similarly, the constant pressure ensemble and the isoenthalpic-isobaric ensemble are also related through a Laplace transform. A theory of the thermodynamic formulation of all the

eight ensembles are given in Ref. [31] for the generalized statistical mechanics based on Tsallis  $q$ -exponential.

*Linear harmonic oscillator:*

Next let us consider a system of  $N$  number of  $D$ -dimensional non-interacting harmonic oscillators. The Hamiltonian of the oscillator is [25]

$$H = \sum_{i=1}^{DN} \left( \frac{p_i^2}{2m} + \frac{1}{2} m \omega^2 x_i^2 \right), \quad (122)$$

where  $x$  and  $p$  are the position and momentum co-ordinates of the oscillator. There are three different approaches to calculate the probability distribution in the current work we use the second constraint approach. The partition function of the system in the second constraint formulation is

$$Z_q(\beta) = \frac{1}{h^{DN}} \int_0^\infty d\bar{x} d\bar{p} \exp_q \left( -\beta \sum_{i=1}^{DN} \left( \frac{p_i^2}{2m} + \frac{1}{2} m \omega^2 x_i^2 \right) \right), \quad (123)$$

where  $\bar{x}$  and  $\bar{p}$  are  $(x_1, x_2, \dots, x_{DN}), (p_1, p_2, \dots, p_{DN})$ . Evaluating the integral we get:

$$Z_q(\beta) = \frac{1}{(\hbar\omega)^{DN}} \left( \frac{\Gamma\left(\frac{1}{1-q} + 1\right)}{(1-q)^{DN} \Gamma\left(\frac{1}{1-q} + DN + 1\right)} \right) \frac{1}{\beta^{DN}}. \quad (124)$$

To calculate the thermodynamics of a set of isolated harmonic oscillator we use the microcanonical ensemble and so we need to find the density of states. The density of states can be computed from the partition function using the inverse of the  $q$ -Laplace transform. Instead of using the regular complex analysis based approach to calculate the inverse transform we use the  $q$ -generalization of the Widder's method introduced in the present work. Using this method the density of states obtained is

$$g(E) = \lim_{k \rightarrow \infty} \frac{(-1)^k}{\Gamma(k+1)} Z_q^{(k)}(\beta) \beta^{k+1} Q_1(2-q) \Big|_{\beta = \frac{k\epsilon}{E}}. \quad (125)$$

Substituting the partition function Eq. (124) in the expression for the density of states Eq. (125) and substituting the value of  $\beta$  and applying the limiting value of  $k$  we get

$$g(E) = \frac{1}{(\hbar\omega)^{DN} (DN-1)!} E^{DN-1}, \quad (126)$$

which is the expression of the density of states of a system of  $N$  non-interacting  $D$ -dimensional oscillators in the microcanonical ensemble [25]. Thus we use the generalized Post-Widder's method was based inverse Laplace transform to transform the partition function of a non-relativistic classical ideal gas and linear harmonic oscillator to their respective density of states.

## 8. Conclusion

A generalization of the Laplace transform based on the Tsallis  $q$ -exponential is defined in the present work based on the kernel  $K_I(q; s, t) = \exp_q(-st)$ . While this definition was also already introduced in Ref. [19], an inverse transformation was not defined. In our work we define the inverse transform using the complex integration method. We also provide an alternative method based on Post-Widder's method to compute the inverse of the transform using only real variables. Several properties of the Laplace

transform and the inverse transform are verified. Then we compute the Laplace transform of some functions like algebraic functions, Gaussian functions,  $q$ -Gaussian functions, circular functions,  $q$ -circular function, hyperbolic function, and  $q$ -hyperbolic functions. On the resulting expressions we use the inverse Laplace transform based on the Post-Widders method to derive the original functions. Finally we show the applications of these methods in statistical mechanics where we compute the density of states of a  $N$ -Dimensional classical ideal gas and linear harmonic oscillator from their partition function.

## Acknowledgements

Md. Manirul Ali was supported by the Centre for Quantum Science and Technology, Chennai Institute of Technology, India, vide funding number CIT/CQST/2021/RD-007. R. Chandrashekar was supported in part by a seed grant from IIT Madras to the Centre for Quantum Information, Communication and Computing.

## 9. References:

- [1] Schiff J L 1999 *The Laplace transform: theory and applications* (Springer Science & Business Media)
- [2] Post E L 1930 *Transactions of the American Mathematical Society* **32** 723–781
- [3] Widder D V 1934 *Transactions of the American Mathematical Society* **36** 107–200
- [4] Jagerman D 1982 *Bell System Technical Journal* **61** 1995–2002
- [5] Al-Shuaibi A 2001 *Integral Transforms and Special Functions* **11** 225–232
- [6] Soldatov A, Seke J and Bogolubov N 2000 *Condensed Matter Physics* **3** 371–380
- [7] Tsallis C 1988 *Journal of Statistical Physics* **52** 479–487
- [8] Nakamichi A, Joichi I, Iguchi O and Morikawa M 2002 *Chaos, Solitons & Fractals* **13** 595–601
- [9] Sakagami M a and Taruya A 2004 *Continuum Mechanics and Thermodynamics* **16** 279–292
- [10] Tsallis C and Borges E P 2003 Nonextensive statistical mechanics—applications to nuclear and high energy physics *Correlations and Fluctuations in QCD* (World Scientific) pp 326–343
- [11] Tsallis C, Anjos J C and Borges E P 2003 *Physics Letters A* **310** 372–376
- [12] Luciano G G and Blasone M 2021 *The European Physical Journal C* **81** 1–17
- [13] Umarov S and Constantino T 2022 *Mathematical Foundations of Nonextensive Statistical Mechanics* (World Scientific)
- [14] Yamano T 2002 *Physica A: Statistical Mechanics and its Applications* **305** 486–496
- [15] Rajagopal A 1998 *Physica A: Statistical Mechanics and its Applications* **253** 271–289
- [16] Lenzi E, Borges E P and Mendes R 1999 *Journal of Physics A: Mathematical and General* **32** 8551
- [17] Plastino A and Rocca M C 2013 *Physica A: Statistical Mechanics and its Applications* **392** 5581–5591
- [18] Chung W S 2013 *arXiv preprint arXiv:1301.5480*
- [19] Naik S R and Haubold H J 2016 *Axioms* **5** 24
- [20] Mamode M 2010 *Journal of Mathematical Physics* **51** 123509
- [21] Jauregui M and Tsallis C 2010 *Journal of Mathematical Physics* **51** 063304
- [22] Borges E P 2004 *Physica A: Statistical Mechanics and its Applications* **340** 95–101
- [23] Chakrabarti R, Chandrashekar R and Naina Mohammed S 2010 *Physica A: Statistical Mechanics and its Applications* **389** 1571–1584
- [24] Borges E P 1998 *Journal of Physics A: Mathematical and General* **31** 5281
- [25] Greiner W, Neise L and Stöcker H 2012 *Thermodynamics and statistical mechanics* (Springer Science & Business Media)
- [26] Prato D 1995 *Physics Letters A* **203** 165–168
- [27] Chakrabarti R, Chandrashekar R and Naina Mohammed S 2008 *Physica A: Statistical Mechanics and its Applications* **387** 4589–4598
- [28] Abe S 1999 *Physics Letters A* **263** 424–429
- [29] Tsallis C, Mendes R and Plastino A R 1998 *Physica A: Statistical Mechanics and its Applications* **261** 534–554

- [30] Tsallis C, Plastino A R and Alvarez-Estrada R F 2009 *Journal of Mathematical Physics* **50** 043303
- [31] Chandrashekar R and Mohammed Naina Mohammed S 2011 *Journal of Statistical Mechanics: Theory and Experiment* **2011** P05018