

INTERPOLATING FUNCTIONS FOR A FAMILY OF DOMAINS RELATED TO μ -SYNTHESIS

SAMRIDDHO ROY

ABSTRACT. Assuming the existence of an analytic interpolant mapping a two-point data from the unit disc \mathbb{D} to $\tilde{\mathbb{G}}_n$, we describe a class of such interpolating functions where

$$\tilde{\mathbb{G}}_n := \left\{ (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n : q \in \mathbb{D}, y_j = \beta_j + \bar{\beta}_{n-j}q, \beta_j \in \mathbb{C} \text{ and} \right. \\ \left. |\beta_j| + |\beta_{n-j}| < \binom{n}{j} \text{ for } j = 1, \dots, n-1 \right\}.$$

We present the connection of $\tilde{\mathbb{G}}_n$ with the μ -synthesis problem.

1. INTRODUCTION

This article is a sequel of [12] and [13]. In [12] the author and Pal introduced a new family of domains, namely the *extended symmetrized polydisc*, $\tilde{\mathbb{G}}_n$, where

$$\tilde{\mathbb{G}}_n := \left\{ (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n : |q| < 1, y_j = \beta_j + \bar{\beta}_{n-j}q \text{ with} \right. \\ \left. |\beta_j| + |\beta_{n-j}| < \binom{n}{j}, 1 \leq j \leq n-1 \right\},$$

and found a variety of new characterizations of the points of the *symmetrized polydisc*, \mathbb{G}_n , where

$$\mathbb{G}_n = \left\{ \left(\sum_{1 \leq i \leq n} z_i, \sum_{1 \leq i < j \leq n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) : |z_i| < 1, i = 1, \dots, n \right\}.$$

The symmetrized polydisc is directly associated with the spectral interpolation. For a matrix A , the spectral radius $r(A) < 1$ if and only if $\pi_n(\lambda_1, \dots, \lambda_n) \in \mathbb{G}_n$ (see [5]), where $\lambda_1, \dots, \lambda_n$ are eigenvalues of A and π_n is the symmetrization map on \mathbb{C}^n defined by

$$\pi_n(z_1, \dots, z_n) = \left(\sum_{1 \leq i \leq n} z_i, \sum_{1 \leq i < j \leq n} z_i z_j, \dots, \prod_{i=1}^n z_i \right).$$

The Schwarz lemma for the symmetrized bidisc \mathbb{G}_2 was presented in [2] (also see [10]). But there was no Schwarz type lemma for \mathbb{G}_n , $n \geq 3$. Note that $\tilde{\mathbb{G}}_2 = \mathbb{G}_2$ and $\mathbb{G}_n \subsetneq \tilde{\mathbb{G}}_n$ for $n \geq 3$ (see [5], [12]). In [13], the author and Pal produced a Schwarz lemma for $\tilde{\mathbb{G}}_n$ and \mathbb{G}_n and showed that an interpolating function, when exists, may not be unique. The aim of this paper is to describe a class of analytic interpolants when we have such a Schwarz lemma for $\tilde{\mathbb{G}}_n$.

2010 *Mathematics Subject Classification.* 30C80, 32E30, 32F45, 93B50.

Key words and phrases. Interpolation; Invariant distances; Structured singular value; Symmetrized polydisc; extended symmetrized polydisc.

The closure of $\tilde{\mathbb{G}}_n$ is denoted by $\tilde{\Gamma}_n$. to study complex geometry of $\tilde{\mathbb{G}}_n$ and $\tilde{\Gamma}_n$, in [12], we introduced $(n-1)$ fractional linear transformations $\Phi_1, \dots, \Phi_{n-1}$. Recall that, for $z \in \mathbb{C}$ and $y = (y_1, \dots, y_{n-1}, q) \in \mathbb{C}^n$ we define

$$\Phi_j(z, y) = \frac{\binom{n}{j} qz - y_j}{y_{n-j}z - \binom{n}{j}} \quad \text{whenever } y_{n-j}z \neq \binom{n}{j} \text{ and } y_j y_{n-j} \neq \binom{n}{j}^2 q.$$

Theorems 2.5 and 2.7 of [12] characterize the points in $\tilde{\mathbb{G}}_n$ and $\tilde{\Gamma}_n$. The extended symmetrized polydisc is a non-convex but polynomially convex domain for all n . Also $\tilde{\mathbb{G}}_n$ is a starlike domain (see [12]).

Let B_1, \dots, B_k be 2×2 strictly contractive matrices such that $\det B_1 = \det B_2 = \dots = \det B_k$. We define two functions π_{2k+1} and π_{2k} in the following way:

$$\pi_{2k+1}(B_1, \dots, B_k) = \left(\binom{n}{1} [B_1]_{11}, \dots, \binom{n}{k} [B_k]_{11}, \binom{n}{k} [B_k]_{22}, \dots, \binom{n}{1} [B_1]_{22}, \det B_1 \right)$$

and

$$\pi_{2k}(B_1, \dots, B_k) = \left(\binom{n}{1} [B_1]_{11}, \dots, \binom{n}{k-1} [B_{k-1}]_{11}, \binom{n}{k} \frac{([B_k]_{11} + [B_k]_{22})}{2}, \right. \\ \left. \binom{n}{k-1} [B_{k-1}]_{22}, \dots, \binom{n}{1} [B_1]_{22}, \det B_1 \right).$$

Then, by Theorem 2.5 of [12], we have $\pi_{2k}(B_1, \dots, B_k) \in \tilde{\mathbb{G}}_{2k}$ and $\pi_{2k+1}(B_1, \dots, B_k) \in \tilde{\mathbb{G}}_{2k+1}$.

For $0 \neq \lambda \in \mathbb{D}$ and $y \in \tilde{\mathbb{G}}_n$, the Schwarz lemma for $\tilde{\mathbb{G}}_n$ describes the necessary conditions for the existence of an analytic interpolating function from the unit disc to $\tilde{\mathbb{G}}_n$ mapping the origin to the origin and λ to y . In [13], it is shown that unlike the classical Schwarz Lemma there is no uniqueness statement for the interpolating function in the case of $\tilde{\mathbb{G}}_n$. In the next section we describe the class of such analytic interpolating functions for the Schwarz lemma for $\tilde{\mathbb{G}}_n$. Note that, in [11], another version of Schwarz lemma for $\tilde{\mathbb{G}}_n$ was presented. In section 3 we present the relation of $\tilde{\mathbb{G}}_n$ with the μ -synthesis problem. In section 4 we take a step towards the Lempert Theory, we find a class of points for which the Carathéodory pseudo-distance and the Lempert function from the origin coincide. Note that the techniques that are used here are similar to the paper [1].

2. INTERPOLATING FUNCTIONS

The Schwarz Lemma for $\tilde{\mathbb{G}}_n$ tells us that for $0 \neq \lambda_0 \in \mathbb{D}$ and $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \tilde{\mathbb{G}}_n$, one of the necessary condition for the existence of an analytic interpolating functions $\psi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_n$ such that $\psi(0) = (0, \dots, 0)$ and $\psi(\lambda_0) = y^0$ is for each $j = 1, \dots, n-1$, $\|\Phi_j(\cdot, y^0)\|_{H^\infty} \leq |\lambda_0|$ whenever $|y_{n-j}^0| \leq |y_j^0|$. Note that this result generalizes the Schwarz lemma for the symmetrized bidisc \mathbb{G}_2 presented in [2]. In this section we describe the explicit form of such holomorphic interpolating function ψ , assuming its existence, whenever $|y_{n-j}^0| \leq |y_j^0|$ and $\|\Phi_j(\cdot, y^0)\|_{H^\infty} < |\lambda_0|$. Theorem 2.3 is one of the main results of this article, where we portray explicitly how those ψ looks like.

The Schwarz Lemma for \mathbb{G}_n and $\tilde{\mathbb{G}}_n$ only talks about the existential criteria of an interpolating function. But it is important to describe the nature of such interpolating functions. In [2], only an example of an interpolating function with respect to the Schwarz lemma for \mathbb{G}_2 was presented assuming the condition $\|\Phi_1(\cdot, y^0)\|_{H^\infty} = |\lambda_0|$. In

[11], the author and Pal produced an example of an interpolating function related with an other version of Schwarz lemma for $\tilde{\mathbb{G}}_n$ mentioned therein. Here, instead of a particular example, we found the explicit form of all holomorphic interpolating functions arising in both the Schwarz Lemma for \mathbb{G}_n and $\tilde{\mathbb{G}}_n$. Note that, as $\tilde{\mathbb{G}}_2 = \mathbb{G}_2$, Theorem 2.7 provides description of interpolating functions with respect to the Schwarz lemma for \mathbb{G}_2 for the condition $\|\Phi_1(., y^0)\|_{H^\infty} < |\lambda_0|$. We start this section with the following important lemma.

Let $Z \in \mathbb{C}^{2 \times 2}$ be such that $\|Z\| < 1$ and let $0 \leq \rho < 1$. Let

$$\mathcal{K}_Z(\rho) = \begin{bmatrix} [(1 - \rho^2 Z^* Z)(1 - Z^* Z)^{-1}]_{11} & [(1 - \rho^2)(1 - Z Z^*)^{-1} Z]_{21} \\ [(1 - \rho^2) Z^* (1 - Z Z^*)^{-1}]_{12} & [(Z Z^* - \rho^2)(1 - Z Z^*)^{-1}]_{22} \end{bmatrix}. \quad (1)$$

Lemma 2.1. *Let $\lambda_0 \in \mathbb{D} \setminus \{0\}$ and $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \tilde{\mathbb{G}}_n$. For a $j \in \{1, \dots, n-1\}$, suppose $y_j^0 y_{n-j}^0 \neq \binom{n}{j}^2 q^0$, $|y_{n-j}^0| \leq |y_j^0|$ and $\|\Phi_j(., y^0)\| < |\lambda_0|$. For any $\nu > 0$ let*

$$Z_{\nu,j} = \begin{bmatrix} y_j^0 / \binom{n}{j} \lambda_0 & \nu w_j \\ w_j / \nu & y_{n-j}^0 / \binom{n}{j} \end{bmatrix} \quad (2)$$

where $w_j^2 = \frac{y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0}{\binom{n}{j}^2 \lambda_0}$. Let $\mathcal{K}_{Z_{\nu,j}}$ be defined by the equation (1). Also let θ_j, ϑ_j be the roots of the equation

$$z + 1/z = \frac{|\lambda_0|}{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|} \left(\binom{n}{j}^2 - \frac{|y_j^0|^2}{|\lambda_0|^2} - |y_{n-j}^0|^2 + \frac{\binom{n}{j}^2 |q^0|^2}{|\lambda_0|^2} \right).$$

Then, $\|Z_{\nu,j}\| < 1$ if and only if

$$\theta_j < \nu^2 < \vartheta_j. \quad (3)$$

Also $\mathcal{K}_{Z_{\nu,j}}(|\lambda_0|)$ is not positive definite whenever ν satisfies condition (3).

Proof. First note that

$$1 - Z_{\nu,j}^* Z_{\nu,j} = \begin{bmatrix} 1 - \frac{|y_j^0|^2}{\binom{n}{j}^2 |\lambda_0|^2} - \frac{|w|^2}{\nu^2} & -\frac{\bar{y}_j^0 \nu w}{\binom{n}{j} \bar{\lambda}_0} - \frac{y_{n-j}^0 \bar{w}}{\binom{n}{j} \nu} \\ -\frac{y_j^0 \nu \bar{w}}{\binom{n}{j} \lambda_0} - \frac{\bar{y}_{n-j}^0 w}{\binom{n}{j} \nu} & 1 - \nu^2 |w|^2 - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} \end{bmatrix}. \quad (4)$$

Then we have

$$\det(1 - Z_{\nu,j}^* Z_{\nu,j}) = 1 - \frac{|y_j^0|^2}{\binom{n}{j}^2 |\lambda_0|^2} - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} + \frac{|q^0|^2}{|\lambda_0|^2} - \frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2 |\lambda_0|} (\nu + 1/\nu^2). \quad (5)$$

Thus, $\|Z_{\nu,j}\| < 1$ if and only if

$$1 - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} - \nu^2 \frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2 |\lambda_0|} > 0$$

and

$$1 - \frac{|y_j^0|^2}{\binom{n}{j}^2 |\lambda_0|^2} - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} + \frac{|q^0|^2}{|\lambda_0|^2} - \frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2 |\lambda_0|} (\nu^2 + 1/\nu^2) > 0,$$

that is, if and only if

$$\nu^2 < \frac{\binom{n}{j}^2 |\lambda_0| \left(1 - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2}\right)}{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}$$

and $\nu^2 + 1/\nu^2 < \frac{\binom{n}{j}^2 |\lambda_0| \left(1 - \frac{|y_j^0|^2}{\binom{n}{j}^2 |\lambda_0|^2} - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} + \frac{|q^0|^2}{|\lambda_0|^2}\right)}{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}.$

Let

$$R_j \equiv \frac{\binom{n}{j}^2 |\lambda_0| \left(1 - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2}\right)}{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|} = \frac{|\lambda_0| (\binom{n}{j}^2 - |y_{n-j}^0|^2)}{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}. \quad (6)$$

By hypothesis, we have

$$\begin{aligned} & \frac{\binom{n}{j} |y_j^0 - \bar{y}_{n-j}^0 q^0| + |y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2 - |y_{n-j}^0|^2} = \|\Phi_j(\cdot, y^0)\| < |\lambda_0| \\ \Rightarrow & \binom{n}{j} \left| \frac{y_j^0}{\lambda_0} - \bar{y}_{n-j}^0 \frac{q^0}{\lambda_0} \right| + \left| \frac{y_j^0}{\lambda_0} y_{n-j}^0 - \binom{n}{j}^2 \frac{q^0}{\lambda_0} \right| < \binom{n}{j}^2 - |y_{n-j}^0|^2. \end{aligned} \quad (7)$$

Then, using the Theorem 2.5 of [12] (equivalence of conditions (4) and (6)), we have

$$\binom{n}{j}^2 - \frac{|y_j^0|^2}{|\lambda_0|^2} - |y_{n-j}^0|^2 + \binom{n}{j}^2 \frac{|q^0|^2}{|\lambda_0|^2} > 2 \frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{|\lambda_0|}. \quad (8)$$

Since $|y_{n-j}^0| \leq |y_j^0|$ and $\lambda_0 \in \mathbb{D}$, we have $\left(\frac{|y_j^0|^2}{|\lambda_0|^2} + |y_{n-j}^0|^2\right) \geq \left(|y_j^0|^2 + \frac{|y_{n-j}^0|^2}{|\lambda_0|^2}\right)$. Hence

$$\begin{aligned} \binom{n}{j}^2 - |y_j^0|^2 - \frac{|y_{n-j}^0|^2}{|\lambda_0|^2} + \binom{n}{j}^2 \frac{|q^0|^2}{|\lambda_0|^2} & \geq \binom{n}{j}^2 - \frac{|y_j^0|^2}{|\lambda_0|^2} - |y_{n-j}^0|^2 + \binom{n}{j}^2 \frac{|q^0|^2}{|\lambda_0|^2} \\ & > 2 \frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{|\lambda_0|}. \end{aligned} \quad (9)$$

Let, $X_j := \frac{|\lambda_0|}{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|} \left(\binom{n}{j}^2 - |y_j^0|^2 - \frac{|y_{n-j}^0|^2}{|\lambda_0|^2} + \frac{\binom{n}{j}^2 |q^0|^2}{|\lambda_0|^2} \right)$

and $X_{n-j} := \frac{|\lambda_0|}{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|} \left(\binom{n}{j}^2 - \frac{|y_j^0|^2}{|\lambda_0|^2} - |y_{n-j}^0|^2 + \frac{\binom{n}{j}^2 |q^0|^2}{|\lambda_0|^2} \right).$

Then we have, $X_j > 2$ and $X_{n-j} > 2$. Then, $\|Z_{\nu,j}\| < 1$ if and only if

$$\nu^2 < R_j \quad \text{and} \quad \nu^2 + 1/\nu^2 < X_{n-j}.$$

Since $|y_{n-j}^0| \leq |y_j^0|$, we have $y_j^0 \neq \bar{y}_{n-j}^0 q^0$. Otherwise, $y_j^0 = \bar{y}_{n-j}^0 q^0$ and hence $|y_j^0| = |\bar{y}_{n-j}^0| |q^0| < |y_{n-j}^0|$, a contradiction. By hypothesis $\|\Phi_j(\cdot, y^0)\| < |\lambda_0|$. Then

$$\left| y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0 \right| = \left(\binom{n}{j}^2 - |y_{n-j}^0|^2 \right) \|\Phi_j(\cdot, y^0)\| < |\lambda_0| \left(\binom{n}{j}^2 - |y_{n-j}^0|^2 \right).$$


Hence $R_j > 1$. Notice that

$$R_j + 1/R_j - X_{n-j} = \frac{\binom{n}{j}^2 |y_j^0 - \bar{y}_{n-j}^0 q^0|^2}{|\lambda_0|(\binom{n}{j}^2 - |y_{n-j}^0|^2) |y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|} > 0. \quad (10)$$

Hence

$$R_j + 1/R_j > X_{n-j}. \quad (11)$$

The graph of the function $f(x) = x + \frac{1}{x}$ is the following :



The function f is continuous and has a point of minima at $x = 1$ and $f(1) = 2$. Since $X_{n-j} > 2$, the line $x = X_{n-j}$ intercepts the graph of f twice. Note that the equation

$$z + 1/z = \frac{|\lambda_0|}{|y_j y_{n-j} - \binom{n}{j}^2 q|} \left(\binom{n}{j}^2 - \frac{|y_j|^2}{|\lambda_0|^2} - |y_{n-j}|^2 + \frac{\binom{n}{j}^2 |q|^2}{|\lambda_0|^2} \right)$$

is same as the equation $z + 1/z = X_{n-j}$. Since θ_j and ϑ_j are two solutions of the equation $z + 1/z = X_{n-j}$, it is clear from the above figure that $\xi + 1/\xi < X_{n-j}$ if and only if $\theta_j < \xi < \vartheta_j$. From the fact $\theta_j < 1 < \vartheta_j$, $R_j > 1$ and from equation (11), we have $R_j > \vartheta_j$.

If $\theta_j < \nu^2 < \vartheta_j$, then $\nu^2 < R_j$ and $\nu^2 + 1/\nu^2 < X_{n-j}$. Hence $\|Z_{\nu,j}\| < 1$ whenever $\theta_j < \nu^2 < \vartheta_j$. Again $\|Z_{\nu,j}\| < 1$ implies $\nu^2 + 1/\nu^2 < X_{n-j}$, which implies $\theta_j < \nu^2 < \vartheta_j$. Therefore, $\|Z_{\nu,j}\| < 1$ if and only if $\theta_j < \nu^2 < \vartheta_j$.

Next we show that, $\det \mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) < 0$ whenever $\theta_j < \nu^2 < \vartheta_j$. First note that :

$$\begin{aligned} & \mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) \\ &= \begin{bmatrix} [(1 - |\lambda_0|^2 Z_{\nu,j}^* Z_{\nu,j})(1 - Z_{\nu,j}^* Z_{\nu,j})^{-1}]_{11} & [(1 - |\lambda_0|^2)(1 - Z_{\nu,j} Z_{\nu,j}^*)^{-1} Z_{\nu,j}]_{21} \\ [(1 - |\lambda_0|^2) Z_{\nu,j}^* (1 - Z_{\nu,j} Z_{\nu,j}^*)^{-1}]_{12} & [(Z_{\nu,j} Z_{\nu,j}^* - |\lambda_0|^2)(1 - Z_{\nu,j} Z_{\nu,j}^*)^{-1}]_{22} \end{bmatrix}. \end{aligned} \quad (12)$$

Then by simple calculations, we have (see appendix)

$$\mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) \det(1 - Z_{\nu,j}^* Z_{\nu,j}) =$$

$$\begin{bmatrix} 1 - \frac{|y_j^0|^2}{\binom{n}{j}^2} - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} + |q^0|^2 & (1 - |\lambda_0|^2) \left(\frac{w}{\nu} + \frac{q^0}{\lambda_0} \nu \bar{w} \right) \\ -\frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2} \left(\frac{|\lambda_0|}{\nu^2} + \frac{\nu^2}{|\lambda_0|} \right) & -|\lambda_0|^2 + \frac{|y_j^0|^2}{\binom{n}{j}^2} + \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} - \frac{|q^0|^2}{|\lambda_0|^2} \\ (1 - |\lambda_0|^2) \left(\frac{\bar{w}}{\nu} + \frac{\bar{q}^0}{\lambda_0} \nu w \right) & -\frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2} \left(\nu^2 |\lambda_0| + \frac{1}{\nu^2 |\lambda_0|} \right) \end{bmatrix}. \quad (13)$$

Now by a straight forward calculation

$$\det(\mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) \det(1 - Z_{\nu,j}^* Z_{\nu,j})) = -(l_{\nu,j} - k_j)(l_{\nu,j} - k_{n-j}), \quad (14)$$

where

$$l_{\nu,j} = |y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0| \left(\nu^2 + \frac{1}{\nu^2} \right),$$

$$k_j = |y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0| X_j.$$

Since $|y_{n-j}^0| \leq |y_j^0|$,

$$X_j - X_{n-j} = \frac{1 - |\lambda_0|^2}{|(y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0) \lambda_0|} \left(|y_j^0|^2 - |y_{n-j}^0|^2 \right) \geq 0$$

and consequently $k_{n-j} \leq k_j$. If $\theta_j < \nu^2 < \vartheta_j$ then $\nu^2 + \frac{1}{\nu^2} < X_{n-j}$ and hence $l_{\nu,j} < |y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0| X_{n-j} = k_{n-j} \leq k_j$. Thus $\det \mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) < 0$ when $\theta_j < \nu^2 < \vartheta_j$. ■

The next theorem is one of the main results of this article, it describes the class of analytic interpolating functions from the unit disc to $\tilde{\mathbb{G}}_n$. We use two lemmas, Lemma 3.1 and Lemma 3.2, from [1].

Let Z be a strict 2×2 matrix contraction and $\alpha \in \mathbb{C}^2 \setminus \{0\}$. A matricial Möbius transformation \mathcal{M}_Z is defined as

$$\mathcal{M}_Z(X) = (1 - ZZ^*)^{-\frac{1}{2}}(X - Z)(1 - Z^*X)^{-1}(1 - Z^*Z)^{\frac{1}{2}}, \quad X \in \mathbb{C}^{2 \times 2} \text{ and } \|X\| < 1.$$

Then \mathcal{M}_Z is an automorphism of the close unit ball of $\mathbb{C}^{2 \times 2}$ which maps Z to the zero matrix and $\mathcal{M}_Z^{-1} = \mathcal{M}_{-Z}$. Also let

$$u_Z(\alpha) = (1 - ZZ^*)^{-\frac{1}{2}}(\alpha_1 Z e_1 + \alpha_2 e_2), \quad (15)$$

$$v_Z(\alpha) = -(1 - Z^*Z)^{-\frac{1}{2}}(\alpha_1 e_1 + \alpha_2 Z^* e_2)$$

where $\{e_1, e_2\}$ is the standard basis of \mathbb{C}^2 . Then for any 2×2 matrix contraction X , by Lemma 3.1 of [1], $[\mathcal{M}_{-Z}(X)]_{22} = 0$ if and only if there exists $\alpha \in \mathbb{C}^2 \setminus \{0\}$ such that $X^* u_Z(\alpha) = v_Z(\alpha)$.

Remark 2.2. Let $\lambda_0 \in \mathbb{D} \setminus \{0\}$ and $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \tilde{\mathbb{G}}_n$. For a $j \in \{1, \dots, n-1\}$, suppose $y_j^0 y_{n-j}^0 \neq \binom{n}{j}^2 q^0$, $|y_{n-j}^0| \leq |y_j^0|$ and $\|\Phi_j(\cdot, y^0)\| < |\lambda_0|$. Then for any ν satisfying condition (3), using the last lemma and Lemma 3.1 of [1] together, there exists $X \in \mathbb{C}^{2 \times 2}$ such that $\|X\| \leq |\lambda_0|$ and $[\mathcal{M}_{-Z_{\nu,j}}(X)]_{22} = 0$, where $Z_{\nu,j}$ is given by (2).

Recall that the *Schur class* of type 2×2 is the set of analytic functions F on \mathbb{D} with values in the space $\mathbb{C}^{2 \times 2}$ such that $\|F(\lambda)\| \leq 1$ for all $\lambda \in \mathbb{D}$. Also we say that $F \in \mathcal{S}_{2 \times 2}$ if $\|F(\lambda)\| < 1$ for all $\lambda \in \mathbb{D}$.

Theorem 2.3. *Let $\lambda_0 \in \mathbb{D} \setminus \{0\}$ and $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \tilde{\mathbb{G}}_n$ and suppose $y_j^0 y_{n-j}^0 \neq \binom{n}{j}^2 q^0$, $|y_{n-j}^0| \leq |y_j^0|$ and $\|\Phi_j(\cdot, y^0)\| < |\lambda_0|$ for each $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$. Suppose $\psi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_n$ is an analytic function such that $\psi(0) = (0, \dots, 0)$ and $\psi(\lambda_0) = y^0$. Then*

$$\psi(\lambda) = \begin{cases} \pi_{2\lfloor \frac{n}{2} \rfloor + 1} \left(F_1(\lambda), \dots, F_{\lfloor \frac{n}{2} \rfloor}(\lambda) \right) & \text{if } n \text{ is odd} \\ \pi_{2\lfloor \frac{n}{2} \rfloor} \left(F_1(\lambda), \dots, F_{\lfloor \frac{n}{2} \rfloor}(\lambda) \right) & \text{if } n \text{ is even,} \end{cases}$$

where for each $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$,

$$F_j(\lambda) = \mathcal{M}_{-Z_{\nu,j}} \left((BQ_j)(\lambda) \right) \begin{bmatrix} \lambda & 0 \\ 0 & 1 \end{bmatrix}, \quad (16)$$

$Z_{\nu,j}$ is given by (2) and ν satisfies (3), Q_j is a 2×2 Schur function such that

$$Q_j(0)^* \bar{\lambda}_0 u_{Z_{\nu,j}}(\alpha_j) = v_{Z_{\nu,j}}(\alpha_j), \quad (17)$$

where $u_{Z_{\nu,j}}(\alpha_j), v_{Z_{\nu,j}}(\alpha_j)$ are given by equation (15) and $\alpha_j \in \mathbb{C}^2 \setminus \{0\}$ satisfies

$$\langle \mathcal{K}_{Z_{\nu,j}}(|\lambda_0|)\alpha_j, \alpha_j \rangle \leq 0. \quad (18)$$

Proof. Suppose $\psi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_n$ is an analytic function such that $\psi(0) = (0, \dots, 0)$ and $\psi(\lambda_0) = (y_1^0, \dots, y_{n-1}^0, q^0)$. By Fatou's Lemma, ψ has radial limits almost everywhere on \mathbb{T} . We denote the radial limit function of ψ by $\tilde{\psi}$ which maps \mathbb{T} almost everywhere to $\tilde{\Gamma}_n$. Let

$$h_j(\lambda) = \frac{\psi_j(\lambda)\psi_{n-j}(\lambda)}{\binom{n}{j}^2} - \psi_n(\lambda) \quad \text{for } \lambda \in \mathbb{D}.$$

Then h_j is a bounded analytic function on \mathbb{D} (eventually a Schur function by condition (4) of Theorem 2.5 of [12]). By inner-outer factorization, there exist $f_j, g_j \in H^\infty$ such that

$$h_j(\lambda) = f_j(\lambda)g_j(\lambda) \quad \text{for } \lambda \in \mathbb{D},$$

with $|\tilde{f}_j| = |\tilde{g}_j|$ almost everywhere on \mathbb{T} . As $f_j g_j(0) = 0$, without loss of generality assume $g_j(0) = 0$. Consider

$$F_j(\lambda) = \begin{bmatrix} \frac{\psi_j(\lambda)}{\binom{n}{j}} & f_j(\lambda) \\ g_j(\lambda) & \frac{\psi_{n-j}(\lambda)}{\binom{n}{j}} \end{bmatrix} \quad \text{for } \lambda \in \mathbb{D}. \quad (19)$$

Clearly $\pi_n(F_1, \dots, F_{\lfloor \frac{n}{2} \rfloor}) = \psi$. Since $f_j g_j = \frac{\psi_j \psi_{n-j}}{\binom{n}{j}^2} - \psi_n$, we have $\det F_j = \psi_n$. Note that,

$$1 - F_j^* F_j = \begin{bmatrix} 1 - \frac{|\psi_j|^2}{\binom{n}{j}^2} - |g_j|^2 & -\bar{f}_j \frac{\bar{\psi}_j}{\binom{n}{j}} - \bar{g}_j \frac{\psi_{n-j}}{\binom{n}{j}} \\ -\bar{f}_j \frac{\psi_j}{\binom{n}{j}} - g_j \frac{\bar{\psi}_{n-j}}{\binom{n}{j}} & 1 - \frac{|\psi_{n-j}|^2}{\binom{n}{j}^2} - |f_j|^2 \end{bmatrix}$$

and

$$\det(1 - F_j^* F_j) = 1 - \frac{|\psi_j|^2}{\binom{n}{j}^2} - \frac{|\psi_{n-j}|^2}{\binom{n}{j}^2} + |\psi_n|^2 - |f|^2 - |g|^2.$$

Since $|\tilde{f}| = |\tilde{g}|$ almost everywhere on \mathbb{T} , we have

$$[1 - \tilde{F}_j^* \tilde{F}_j]_{11} = 1 - \frac{|\tilde{\psi}_j|^2}{\binom{n}{j}^2} - \left| \frac{\tilde{\psi}_j \tilde{\psi}_{n-j}}{\binom{n}{j}^2} - \tilde{\psi}_n \right|, \quad [1 - \tilde{F}_j^* \tilde{F}_j]_{22} = 1 - \frac{|\tilde{\psi}_{n-j}|^2}{\binom{n}{j}^2} - \left| \frac{\tilde{\psi}_j \tilde{\psi}_{n-j}}{\binom{n}{j}^2} - \tilde{\psi}_n \right|$$

almost everywhere on \mathbb{T} and

$$\det(1 - \tilde{F}_j^* \tilde{F}_j) = 1 - \frac{|\tilde{\psi}_j|^2}{\binom{n}{j}^2} - \frac{|\tilde{\psi}_{n-j}|^2}{\binom{n}{j}^2} + |\tilde{\psi}_n|^2 - \left| \frac{\tilde{\psi}_j \tilde{\psi}_{n-j}}{\binom{n}{j}^2} - \tilde{\psi}_n \right|$$

almost everywhere on \mathbb{T} . Then, using Theorem 2.5 of [12] (by conditions (3), (3') and (5)), we have

$$\left[1 - \tilde{F}_j^* \tilde{F}_j\right]_{11} \geq 0, \quad \left[1 - \tilde{F}_j^* \tilde{F}_j\right]_{22} \geq 0 \quad \text{and} \quad \det(1 - \tilde{F}_j^* \tilde{F}_j) \geq 0 \quad \text{almost everywhere on } \mathbb{T}.$$

Thus $\|\tilde{F}_j\| \leq 1$ almost everywhere on \mathbb{T} . Note that if F_j is constant then the function $(\psi_j, \psi_{n-j}, \psi_n)$ is also constant. Since $\psi(0) = (0, \dots, 0)$, $\psi(\lambda_0) = (y_1^0, \dots, y_{n-1}^0, q^0)$ and $y_j^0 y_{n-j}^0 \neq \binom{n}{j}^2 q^0$, we conclude F_j is non-constant. Therefore by maximum modulus principle, $\|F_j(\lambda)\| < 1$ for all $\lambda \in \mathbb{D}$. Hence $F_j \in S_{2 \times 2}$.

Thus to complete the proof we need to show that F_j can be written in the form (16) for some ν, α_j and Q_j satisfying (3), (18) and (17) respectively. Since $y_j^0 y_{n-j}^0 \neq \binom{n}{j}^2 q^0$, we have $f_j g_j(\lambda_0) = \frac{y_j^0 y_{n-j}^0}{\binom{n}{j}^2} - q^0 \neq 0$ and hence both $f_j(\lambda_0)$ and $g_j(\lambda_0)$ are nonzero. Now

consider $\nu = \frac{f_j(\lambda_0)}{w_j}$ where $w_j^2 = \frac{y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0}{\binom{n}{j}^2 \lambda_0}$. Then $g_j(\lambda_0) = \frac{\lambda_0 w_j^2}{f_j(\lambda_0)} = \frac{\lambda_0 w_j}{\nu}$. Hence

$$F_j(\lambda_0) = \begin{bmatrix} y_j^0 / \binom{n}{j} & \nu w_j \\ \lambda_0 \nu^{-1} w_j & y_{n-j}^0 / \binom{n}{j} \end{bmatrix}.$$

We may assume that $\nu > 0$ (otherwise we can replace F_j by $U^* F_j U$ for a suitable diagonal unitary matrix U). Consider

$$G_j(\lambda) = F_j(\lambda) \begin{bmatrix} \lambda^{-1} & 0 \\ 0 & 1 \end{bmatrix} \quad \text{for all } \lambda \in \mathbb{D}.$$

Then we have $G_j \in S_{2 \times 2}$ and

$$[G_j(0)]_{22} = [F_j(0)]_{22} = 0 \quad \text{and} \quad G_j(\lambda_0) = \begin{bmatrix} y_j^0 / \binom{n}{j} \lambda_0 & \nu w_j \\ \nu^{-1} w_j & y_{n-j}^0 / \binom{n}{j} \end{bmatrix} = Z_{\nu, j}.$$

Then $F_j(\lambda) = G_j(\lambda) \begin{bmatrix} \lambda & 0 \\ 0 & 1 \end{bmatrix}$ for all $\lambda \in \mathbb{D}$. We have already proved that $\|Z_{\nu, j}\| < 1$ if and only if $\theta_j < \nu^2 < \vartheta_j$. Since $G_j \in S_{2 \times 2}$, $\|Z_{\nu, j}\| = \|G_j(\lambda_0)\| < 1$ and hence $\theta_j < \nu^2 < \vartheta_j$. That is, condition (3) holds. Again since $G_j \in S_{2 \times 2}$, $[G_j(0)]_{22} = 0$ and $G_j(\lambda_0) = Z_{\nu, j}$, by part-(2) of Lemma 3.2 of [1], there exists some $\alpha_j \in \mathbb{C}^2 \setminus \{0\}$ such that $\langle \mathcal{K}_{Z_{\nu, j}}(|\lambda_0|) \alpha_j, \alpha_j \rangle \leq 0$ and a Schur function Q_j such that $Q_j(0)^* \bar{\lambda}_0 u_{Z_{\nu, j}}(\alpha_j) = v_{Z_{\nu, j}}(\alpha_j)$ and

$$G_j = \mathcal{M}_{-Z_{\nu, j}} \circ B Q_j,$$

where $u_{Z_{\nu,j}}(\alpha_j), v_{Z_{\nu,j}}(\alpha_j)$ are given by equation (15). Thus the conditions (18) and (17) are satisfied. Hence $F_j(\lambda) = \mathcal{M}_{-Z_{\nu,j}}(BQ_j(\lambda)) \begin{bmatrix} \lambda & 0 \\ 0 & 1 \end{bmatrix}$ that is, F_j can be written in the form (16). Hence the proof is complete. \blacksquare

Remark 2.4. For $0 \neq \lambda \in \mathbb{D}$ and $x \in \mathbb{G}_n$, any analytic function $\phi : \mathbb{D} \rightarrow \mathbb{G}_n$ such that $\phi(0) = (0, \dots, 0)$ and $\phi(\lambda) = x$ then ϕ can be viewed as a map $\phi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_n$ (as $\mathbb{G}_n \subset \tilde{\mathbb{G}}_n$) and by applying Theorem 2.3 we can describe all such function ϕ .

The Schwarz lemma for $\tilde{\mathbb{G}}_n$ says that, for some special case, when the target point y^0 belongs to a particular subset \mathcal{J}_n of $\tilde{\mathbb{G}}_n$, the conditions which are necessary for the existence of the required interpolating function also become sufficient. Recall that when n is odd,

$$\mathcal{J}_n = \left\{ (y_1, \dots, y_{n-1}, q) \in \tilde{\mathbb{G}}_n : y_j = \frac{\binom{n}{j}}{n} y_1, y_{n-j} = \frac{\binom{n}{j}}{n} y_{n-1}, 2 \leq j \leq \left\lfloor \frac{n}{2} \right\rfloor \right\}$$

and for n even,

$$\mathcal{J}_n = \left\{ (y_1, \dots, y_{n-1}, q) \in \tilde{\mathbb{G}}_n : y_{\lfloor \frac{n}{2} \rfloor} = \frac{\binom{n}{\lfloor \frac{n}{2} \rfloor}}{\binom{n}{\lfloor \frac{n}{2} \rfloor}} \frac{y_1 + y_{n-1}}{2n}, y_j = \frac{\binom{n}{j}}{n} y_1, \right. \\ \left. y_{n-j} = \frac{\binom{n}{j}}{n} y_{n-1}, 2 \leq j \leq \left\lfloor \frac{n}{2} \right\rfloor - 1 \right\}.$$

Suppose $\lambda_0 \in \mathbb{D} \setminus \{0\}$, $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \mathcal{J}_n$ such that $\|y_{n-1}^0\| \leq \|y_1^0\|$ and $\|\Phi_1(\cdot, y^0)\| < |\lambda_0|$. Then the Schwarz Lemma for $\tilde{\mathbb{G}}_n$ guarantees the existence of an analytic function $\psi : \mathbb{D} \rightarrow \mathcal{J}_n \subset \tilde{\mathbb{G}}_n$ such that $\psi(0) = (0, \dots, 0)$ and $\psi(\lambda_0) = y^0$ (see [13]).

In Theorem 2.3, assuming the existence of an analytic interpolating function $\psi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_n$ in the Schwarz Lemma for $\tilde{\mathbb{G}}_n$, we explicitly describe the function ψ . The next theorem deals with the converse part when the target point $y^0 \in \mathcal{J}_n$. We show that if ψ is of the form similar to that mentioned in Theorem 2.3, then ψ is an analytic interpolating function from unit disc to $\mathcal{J}_n \subset \tilde{\mathbb{G}}_n$ mapping the origin to the origin and λ_0 to y^0 . The following lemma is just a particular case of Lemma 2.1 and can be proved similarly.

Lemma 2.5. Let $\lambda_0 \in \mathbb{D} \setminus \{0\}$ and $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \mathcal{J}_n$ and suppose that $y_1^0 y_{n-1}^0 \neq n^2 q^0$, $|y_{n-1}^0| \leq |y_1^0|$ and $\|\Phi_1(\cdot, y^0)\| < |\lambda_0|$. For any $\nu > 0$ let

$$Z_\nu = \begin{bmatrix} y_1^0/n\lambda_0 & \nu w \\ \nu^{-1}w & y_{n-1}^0/n \end{bmatrix} \quad (20)$$

where $w^2 = \frac{y_1^0 y_{n-1}^0 - n^2 q^0}{n^2 \lambda_0}$. Let \mathcal{K}_{Z_ν} be defined by the equation (1). Also let θ_1, ϑ_1 be the roots of the equation

$$z + 1/z = \frac{|\lambda_0|}{|y_1^0 y_{n-1}^0 - n^2 q^0|} \left(n^2 - \frac{|y_1^0|^2}{|\lambda_0|^2} - |y_{n-1}^0|^2 + \frac{n^2 |q^0|^2}{|\lambda_0|^2} \right).$$

Then $\|Z_\nu\| < 1$ if and only if

$$\theta_1 < \nu^2 < \vartheta_1. \quad (21)$$

Also $\mathcal{K}_{Z_\nu}(|\lambda_0|)$ is not positive definite for any ν satisfying (21).

For a 2×2 contractive matrix B , let $\hat{\pi}_n$ be the following map

$$\hat{\pi}_n(B) = \begin{cases} \pi_{2[\frac{n}{2}]+1}(B, \dots, B) & \text{if } n \text{ is odd} \\ \pi_{2[\frac{n}{2}]}(B, \dots, B) & \text{if } n \text{ is even.} \end{cases}$$

Then, for $F \in S^{2 \times 2}$, it is clear that $\hat{\pi}_n \circ F$ is a function from \mathbb{D} to \mathcal{J}_n .

Theorem 2.6. *Let $\lambda_0 \in \mathbb{D} \setminus \{0\}$ and $y^0 = (y_1^0, \dots, y_{n-1}^0, q^0) \in \mathcal{J}_n$ and suppose that $y_1^0 y_{n-1}^0 \neq n^2 q^0$, $|y_{n-1}^0| \leq |y_1^0|$ and $\|\Phi_1(\cdot, y^0)\| < |\lambda_0|$. Let \mathcal{R} be the set of analytic functions $\psi : \mathbb{D} \rightarrow \mathcal{J}_n \subset \tilde{\mathbb{G}}_n$ satisfying $\psi(0) = (0, \dots, 0)$ and $\psi(\lambda_0) = y^0$. The function $\hat{\pi}_n \circ F$ belongs to \mathcal{R} where*

$$F(\lambda) = \mathcal{M}_{-Z_\nu}((BQ)(\lambda)) \begin{bmatrix} \lambda & 0 \\ 0 & 1 \end{bmatrix} \quad (22)$$

where Z_ν is given by (20), ν satisfies (21), Q is a 2×2 Schur function such that

$$Q(0)^* \bar{\lambda}_0 u_{Z_\nu}(\alpha) = v_{Z_\nu}(\alpha), \quad (23)$$

where $u_{Z_\nu}(\alpha), v_{Z_\nu}(\alpha)$ are given by equation (15) and $\alpha \in \mathbb{C}^2 \setminus \{0\}$ satisfies

$$\langle \mathcal{K}_{Z_\nu}(|\lambda_0|)\alpha, \alpha \rangle \leq 0. \quad (24)$$

Conversely, every function in \mathcal{R} is of the form $\hat{\pi}_n \circ F$ where F and Z_ν are given by equation (22) and (20), for some choice of ν, α and Q satisfying the conditions (21), (24) and (23), respectively.

Proof. Suppose ν satisfies (21), that is, $\theta_1 < \nu^2 < \vartheta_1$, where θ_1, ϑ_1 are the roots of the equation

$$z + 1/z = \frac{|\lambda_0|}{|y_1^0 y_{n-1}^0 - n^2 q^0|} \left(n^2 - \frac{|y_1^0|^2}{|\lambda_0|^2} - |y_{n-1}^0|^2 + \frac{n^2 |q^0|^2}{|\lambda_0|^2} \right).$$

Then by Lemma 2.5, $\|Z_\nu\| < 1$ and $\mathcal{K}_{Z_\nu}(|\lambda_0|)$ is not positive definite. Further, suppose $\alpha \in \mathbb{C}^2 \setminus \{(0, 0)\}$ such that $\langle \mathcal{K}_{Z_\nu}(|\lambda_0|)\alpha, \alpha \rangle \leq 0$ and also suppose Q is a Schur function such that $Q(0)^* \bar{\lambda}_0 u_{Z_\nu}(\alpha) = v_{Z_\nu}(\alpha)$. Then using part (2) of Lemma 3.2 of [1], there exists a function $G \in S_{2 \times 2}$ such that

$$G = \mathcal{M}_{-Z_\nu} \circ (BQ) \quad \text{and} \quad [G(0)]_{22} = 0, \quad G(\lambda_0) = Z_\nu.$$

Thus the function F , given by equation (22), is

$$F(\lambda) = G(\lambda) \begin{bmatrix} \lambda & 0 \\ 0 & 1 \end{bmatrix} \quad \text{for } \lambda \in \mathbb{D}.$$

It is clear that $F \in S_{2 \times 2}$ and F satisfies the following

$$F(0) = \begin{bmatrix} 0 & * \\ 0 & 0 \end{bmatrix}, \quad F(\lambda_0) = \begin{bmatrix} y_1^0/n & \nu w \\ \lambda_0 \nu^{-1} w & y_{n-1}^0/n \end{bmatrix}.$$

Thus the function $\psi = \hat{\pi}_n \circ F$ is analytic from \mathbb{D} to \mathcal{J}_n such that $\psi(0) = (0, \dots, 0)$ and $\psi(\lambda_0) = (y_1^0, \dots, y_{n-1}^0, q^0) = y^0$. Hence $\psi \in \mathcal{R}$.

The proof of the converse part is similar to the proof of Theorem 2.3. Suppose $\psi \in \mathcal{R}$. Denote the radial limit function of ψ by $\tilde{\psi}$ which, by Fatou's Lemma, maps \mathbb{T} almost everywhere to $\tilde{\Gamma}_n$. Consider the bounded analytic function

$$h(\lambda) = \frac{\psi_1(\lambda) \psi_{n-1}(\lambda)}{n^2} - \psi_n(\lambda) \quad \text{for } \lambda \in \mathbb{D},$$

and write the inner-outer factorization as $h(\lambda) = f(\lambda)g(\lambda)$ for $\lambda \in \mathbb{D}$, where $f, g \in H^\infty$ such that $|\tilde{f}| = |\tilde{g}|$ almost everywhere on \mathbb{T} . Consider $F(\lambda) = \begin{bmatrix} \psi_1(\lambda)/n & f(\lambda) \\ g(\lambda) & \psi_{n-1}(\lambda)/n \end{bmatrix}$, $\lambda \in \mathbb{D}$. Clearly $\hat{\pi}_n \circ F = \psi$. Using similar method, as shown in proof of Theorem 2.3, we can show that $F \in S_{2 \times 2}$.

Again using the method similar to that in proof of Theorem 2.3, we show that F is of the desired form. Consider $\nu = \frac{f(\lambda_0)}{w}$, then $g(\lambda_0) = \frac{\lambda_0 w}{\nu}$. We may assume that $\nu > 0$. Also consider $G(\lambda) = F(\lambda) \begin{bmatrix} \lambda^{-1} & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda \in \mathbb{D}$. We may assume that $\nu > 0$. Then we have $G \in S_{2 \times 2}$, $[G(0)]_{22} = 0$ and $G(\lambda_0) = Z_\nu$. Then $\|Z_\nu\| = \|G(\lambda_0)\| < 1$ and hence $\theta_1 < \nu^2 < \vartheta_1$. By part-(2) of Lemma 3.2 of [1], there exists some $\alpha \in \mathbb{C}^2 \setminus \{0\}$ such that $\langle \mathcal{K}_{Z_\nu}(|\lambda_0|\alpha), \alpha \rangle \leq 0$ and a Schur function Q such that $Q(0)^* \bar{\lambda}_0 u_{Z_\nu}(\alpha) = v_{Z_\nu}(\alpha)$ and $G = \mathcal{M}_{-Z_\nu} \circ BQ$, where $u_{Z_\nu}(\alpha), v_{Z_\nu}(\alpha)$ are given by equation (15). Thus the conditions (21), (24) and (23) are satisfied and F can be written as $F(\lambda) = \mathcal{M}_{-Z_\nu}(BQ(\lambda)) \begin{bmatrix} \lambda & 0 \\ 0 & 1 \end{bmatrix}$. ■

Theorem 2.6 is true for $n \geq 3$, because of the fact that \mathcal{J}_n is defined only for $n \geq 3$. We now present a result similar to Theorem 2.6 for the case $n = 2$. Note that the conditions described in Schwarz lemma for $\tilde{\mathbb{G}}_n$ in [13] are also sufficient for the existence of an analytic interpolating function from the unit disc to the symmetrized bidisc \mathbb{G}_2 (also see [2] and [10]). The next theorem describes and also provide example of the interpolating function related to Schwarz lemma for \mathbb{G}_2 for the condition that is mutually exclusive to that in Theorem 1.4 of [2].

Theorem 2.7. *Let $\lambda_0 \in \mathbb{D} \setminus \{0\}$ and $y^0 = (s_0, p_0) \in \mathbb{G}_2$ be such that $s_0^2 \neq 4p_0$ and*

$$\frac{2|s_0 - \bar{s}_0 p_0| + |s_0^2 - 4p_0|}{4 - |s_0|^2} < |\lambda_0|.$$

Let θ, ϑ be the roots of the equation

$$z + 1/z = \frac{4|\lambda_0|^2 - |s_0|^2 - |s_0|^2|\lambda_0|^2 + 4|p_0|^2}{|\lambda_0||s_0^2 - 4p_0|}$$

and for $\nu > 0$

$$Z_\nu = \begin{bmatrix} s_0/2\lambda_0 & \nu w \\ \nu^{-1}w & s_0/2 \end{bmatrix} \quad (25)$$

where $w^2 = \frac{s_0^2 - 4p_0}{4\lambda_0}$. Suppose a function ψ is given by

$$\psi(\lambda) = \left([F(\lambda)]_{11} + [F(\lambda)]_{22}, \det F(\lambda) \right) \quad \text{for } \lambda \in \mathbb{D},$$

where F is given by equation (22) with Z_ν as in equation (25), for some ν satisfying $\theta < \nu^2 < \vartheta$ and α and Q satisfying the conditions (24) and (23), respectively. Then ψ is an analytic interpolating function from the unit disc to \mathbb{G}_2 satisfying $\psi(0) = (0, 0)$ and $\psi(\lambda_0) = y^0$.

Conversely, any analytic functions $\psi : \mathbb{D} \rightarrow \mathbb{G}_2$ satisfying $\psi(0) = (0, 0)$ and $\psi(\lambda_0) = y^0$ is of the form $\psi(\lambda) = ([F(\lambda)]_{11} + [F(\lambda)]_{22}, \det F(\lambda))$ where F, Z_ν are given by (22), (25), ν satisfies $\theta < \nu^2 < \vartheta$ and α, Q are as in the conditions (24), (23), respectively.

Proof. First note that $\tilde{\mathbb{G}}_2 = \mathbb{G}_2$ and $\hat{\pi}_2 \circ F = ([F(\lambda)]_{11} + [F(\lambda)]_{22}, \det F(\lambda))$. For any $F \in S_{2 \times 2}$ the function $\hat{\pi}_2 \circ F$ is analytic from \mathbb{D} to \mathbb{G}_2 . It is clear, by the definition of Φ_1 for the case $n = 2$, that

$$\|\Phi_1(\cdot, y^0)\| = \frac{2|s_0 - \bar{s}_0 p_0| + |s_0^2 - 4p_0|}{4 - |s_0|^2} < |\lambda_0|.$$

Then the Proof of both the parts are exactly similar to that of Theorem 2.6 upon substituting $q^0 = p_0$, $y_1^0 = s_0$ and $y_{n-1}^0 = s_0$. ■

3. RELATION WITH THE μ -SYNTHESIS PROBLEM

The μ -synthesis problem plays an important role in robust control theory of control engineering. Here μ is used to denote the structured singular value of a matrix relative to a space of linear transformations. For a given linear subspace E of $\mathbb{C}^{n \times m}$ the *structured singular value* of an $m \times n$ matrix B , denoted by $\mu_E(B)$, is defined as

$$\mu_E(B) = \frac{1}{(\inf\{\|X\| : X \in E, I - BX \text{ is singular}\})}.$$

Here $\|X\|$ denotes the operator norm of the matrix X . In the event of $I - BX$ is non-singular for all $X \in E$, we define $\mu_E(B) = 0$. The underlying linear subspace E is considered as a “structure”.

Given a linear subspace $E \subset \mathbb{C}^{n \times m}$, distinct points $\lambda_1, \dots, \lambda_k$ in \mathbb{D} and same number of $m \times n$ target matrices B_1, \dots, B_k , the μ -synthesis problem aims to find an analytic matrix-valued function $f : \mathbb{D} \rightarrow \mathbb{C}^{m \times n}$ such that $f(\lambda_j) = B_j$ for $1 \leq j \leq k$ and $\mu_E(f(\lambda)) < 1$, for all $\lambda \in \mathbb{D}$.

On the other hand, for given distinct points $\lambda_1, \dots, \lambda_k$ in \mathbb{D} and $n \times n$ matrices W_1, \dots, W_k , the spectral Nevanlinna-Pick interpolation problem is to find necessary and sufficient conditions for the existence of an analytic matrix-valued function $F : \mathbb{D} \rightarrow \mathbb{C}^{n \times n}$ such that $F(\lambda_j) = W_j$ for $1 \leq j \leq k$ and $r(F(\lambda)) < 1$, for all $\lambda \in \mathbb{D}$ ($r(F(\lambda))$ denotes the spectral radius of $F(\lambda)$). In [5], it was shown that such interpolation problem into the spectral unit ball of $\mathbb{C}^{n \times n}$ is equivalent to the interpolation problem into the symmetrized polydisc \mathbb{G}_n .

Note that $\mu_E(B) = \|B\|$, when $E = \mathbb{C}^{n \times m}$. For the case of $m = n$ and $E = \{\lambda I_n : \lambda \in \mathbb{C}\}$, I_n denotes the $n \times n$ identity matrix, $\mu_E(B)$ is equal to the spectral radius $r(B)$ and consequently in this case the μ -synthesis problem reduced to the spectral Nevanlinna-Pick interpolation problem. For more details about the μ -synthesis problem and structured singular value a reader is referred to [4, 6].

In this section we describe the relation of the domain $\tilde{\mathbb{G}}_n$ with the μ -synthesis problem. In Theorem 3.1, we show that the points of $\tilde{\mathbb{G}}_n$ can also be characterized using the structured singular value. In Remark 3.3 we show the connection between the μ -synthesis problem with the interpolation problem in $\text{Hol}(\mathbb{D}, \tilde{\mathbb{G}}_n)$.

Consider the linear subspace $E = \left\{ \begin{pmatrix} z & 0 \\ 0 & w \end{pmatrix} : z, w \in \mathbb{C} \right\}$. For a 2×2 matrix $B = [b_{ij}]$ and $X = \begin{pmatrix} z & 0 \\ 0 & w \end{pmatrix}$, we have $\|X\| = \max\{|z|, |w|\}$ and

$$\det(I - BX) = 1 - zb_{11} - wb_{22} - zw \det B.$$

Then for $r > 0$,

$$\begin{aligned} \mu_E(B) &\leq \frac{1}{r} \\ \Leftrightarrow \max\{|z|, |w|\} &\geq r \text{ whenever } 1 - b_{11}z - b_{22}w - \det Bzw = 0 \\ \Leftrightarrow 1 - zb_{11} - wb_{22} - zw \det B &\neq 0 \text{ for any } z, w \in r\mathbb{D}. \end{aligned}$$

Theorem 3.1. $y \in \tilde{\mathbb{G}}_n$ if and only if there exist $\lfloor \frac{n}{2} \rfloor$ number of 2×2 matrices $B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor}$ such that $\det B_1 = \dots = B_{\lfloor \frac{n}{2} \rfloor}$, $y = \pi_n(B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor})$ and $\mu_E(B_j) < 1$ for $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$.

Proof. Suppose $y \in \tilde{\mathbb{G}}_n$. Then for all $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$, by Theorem 2.5 of [12], we have

$$1 - \frac{y_j}{\binom{n}{j}}z - \frac{y_{n-j}}{\binom{n}{j}}w + qzw \neq 0 \text{ whenever } z, w \in r\mathbb{D} \text{ for some } r > 1.$$

For each $1 \leq j \leq \lfloor \frac{n}{2} \rfloor$, consider the matrix

$$\begin{bmatrix} y_j/\binom{n}{j} & w_j \\ w_j & y_{n-j}/\binom{n}{j} \end{bmatrix} = B_j \text{ (say),}$$

where $w_j^2 = \frac{y_j y_{n-j} - \binom{n}{j}^2 q}{\binom{n}{j}^2}$. Then $\mu_E(B_j) \leq \frac{1}{r} < 1$. Clearly $\pi_n(B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor}) = y$.

Conversely, suppose there exist matrices $B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor}$ such that $\det B_1 = \dots = B_{\lfloor \frac{n}{2} \rfloor}$, $y = \pi_n(B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor})$ and $\mu_E(B_j) < 1$ for $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$. Then for each such j there is some $r_j > 1$ so that $\mu_E(B_j) \leq \frac{1}{r_j}$. Therefore

$$1 - z[B_j]_{11} - w[B_j]_{22} - zw \det B_j \neq 0 \text{ for all } z, w \in \overline{\mathbb{D}}.$$

Note that $\pi_n(B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor}) = y$ implies $[B_j]_{11} = \frac{y_j}{\binom{n}{j}}$, $[B_j]_{22} = \frac{y_{n-j}}{\binom{n}{j}}$ and $\det B_j = q$.

Hence we have

$$\binom{n}{j} - y_j z - y_{n-j} w + \binom{n}{j} qzw \neq 0, \text{ for all } z, w \in \overline{\mathbb{D}}.$$

Therefore $y \in \tilde{\mathbb{G}}_n$. ■

Remark 3.2. (1) With the similar line of argument we can show that $y \in \tilde{\Gamma}_n$ if and only if there exist $\lfloor \frac{n}{2} \rfloor$ numbers of 2×2 matrices $B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor}$ such that $\det B_1 = \dots = B_{\lfloor \frac{n}{2} \rfloor}$, $y = \pi_n(B_1, \dots, B_{\lfloor \frac{n}{2} \rfloor})$ and $\mu_E(B_j) \leq 1$ for $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$.

(2) If a point belongs to the special set $\mathcal{J}_n \subset \tilde{\mathbb{G}}_n$, then instead of $\lfloor \frac{n}{2} \rfloor$ number of matrices the existence of only one matrix is sufficient. We can prove, with similar method as in the proof of last theorem, that $y \in \mathcal{J}_n$ if and only if there exist a 2×2 matrix B such that $\hat{\pi}_n(B) = y$ and $\mu_E(B) < 1$. Indeed, for $y \in \mathcal{J}_n$ the matrix

$$B = \begin{bmatrix} y_1/n & w \\ w & y_{n-1}/n \end{bmatrix}, \text{ where } w^2 = \frac{y_1 y_{n-1} - n^2 q}{n^2}, \text{ satisfies required conditions; on}$$

the other hand for any 2×2 matrix B such that $y = \hat{\pi}_n(B)$ and $\mu_E(B) < 1$, we have $n - y_1 z - y_{n-1} w + nqzw \neq 0$, for all $z, w \in \overline{\mathbb{D}}$ which implies $y \in \mathcal{J}_n$.

Let Ω_E be the unit μ_E -ball, that is, $\Omega_E = \{B \in \mathbb{C}^{2 \times 2} : \mu_E(B) < 1\}$. Every analytic map $\phi : \mathbb{D} \rightarrow \tilde{\mathbb{G}}_n$ gives rise to $\lfloor \frac{n}{2} \rfloor$ numbers of analytic functions $F_1, \dots, F_{\lfloor \frac{n}{2} \rfloor}$ from \mathbb{D} to Ω_E such that $\pi_n(F_1, \dots, F_{\lfloor \frac{n}{2} \rfloor}) = \phi$. Indeed, write $\phi = (\phi_1, \dots, \phi_n)$ and consider the matrix valued function

$$F_j = \begin{bmatrix} \phi_j & \phi_j \phi_{n-j} - \phi_n \\ 1 & \phi_{n-j} \end{bmatrix}.$$

It can be seen, with arguments similar to the proof of Theorem 2.3, that $\mu_E(F_j(\lambda)) < 1$ for all $\lambda \in \mathbb{D}$. Clearly each F_j is analytic on \mathbb{D} and $\pi_n(F_1, \dots, F_{\lfloor \frac{n}{2} \rfloor}) = \phi$. In the following remark we find a necessary condition for the μ -synthesis problem where the underlying linear subspace is the space of 2×2 diagonal matrices.

Remark 3.3. Let $\lambda_1, \dots, \lambda_m$ be m distinct points in \mathbb{D} and $B_1, \dots, B_m \in \Omega_E$. If there exists an analytic function $F : \mathbb{D} \rightarrow \Omega_E$ such that $F(\lambda_j) = B_j$ for $1 \leq j \leq m$, then there exists an analytic function $\phi : \mathbb{D} \rightarrow \mathcal{J}_n$ such that $\phi(\lambda_j) = \widehat{\pi_n}(B_j)$ for $1 \leq j \leq m$. In fact, we may take $\phi = \widehat{\pi_n} \circ F$. The problem of finding such F is called the structured Nevanlinna-Pick problem and clearly it has a connection with the finite interpolation problem in $\text{Hol}(\mathbb{D}, \tilde{\mathbb{G}}_n)$.

4. INVARIANT DISTANCES FOR A SUBSET OF $\tilde{\mathbb{G}}_n$

For a domain $\Omega \subset \mathbb{C}^n$ and two points $z, w \in \Omega$, the Carathéodory pseudo-distance between z, w is

$$\mathcal{C}_\Omega(z, w) := \sup \{ \rho(f(z), f(w)) : f \in \mathcal{O}(\Omega, \mathbb{D}) \}$$

and the Lempert function for Ω is defined as

$$\mathcal{L}_\Omega(z, w) := \inf \{ \rho(\alpha, \beta) : \text{there is } f \in \mathcal{O}(\mathbb{D}, \Omega), f(\alpha) = z, f(\beta) = w \},$$

where ρ is the hyperbolic distance. It can be seen that for $z, w \in \mathbb{D}$, $\rho(z, w) = \tanh^{-1}(d(z, w))$ where d is the Möbius distance. The famous Lempert theorem states that $\mathcal{C}_\mathcal{D} = \mathcal{L}_\mathcal{D}$ for a domain $\mathcal{D} \subset \mathbb{C}^n$ if \mathcal{D} is convex (see [8]). In 2004, Agler and Young proved (see [3]) that these two distances coincide for the non-convex domain \mathbb{G}_2 . Later in 2007, Nikolov, Pflug and Zwonek proved in [9] that they do not agree for \mathbb{G}_n if $n \geq 3$. The tetrablock, a domain in \mathbb{C}^3 , was shown to be the second example of a non-convex domain in which those two distances coincide (see [1, 7]). Since \mathbb{E} is linearly isomorphic to $\tilde{\mathbb{G}}_3$, evidently Lempert's theorem holds for $\tilde{\mathbb{G}}_3$. We still do not know whether Lempert theorem holds for $\tilde{\mathbb{G}}_n$, $n > 3$ or not, but we shall prove that $\mathcal{C}_{\tilde{\mathbb{G}}_n} = \mathcal{L}_{\tilde{\mathbb{G}}_n}$ for the points z, w in $\mathcal{J}_n \subset \tilde{\mathbb{G}}_n$, where at least one of z, w is equal to 0. We apply the same techniques as in [1] to establish this.

Theorem 4.1. Let $y = (y_1, \dots, y_{n-1}, q) \in \mathcal{J}_n \subset \tilde{\mathbb{G}}_n$, then

$$\mathcal{C}_{\tilde{\mathbb{G}}_n}(0, y) = \mathcal{L}_{\tilde{\mathbb{G}}_n}(0, y) = \max_{1 \leq j \leq n-1} \left\{ \tanh^{-1} \frac{\binom{n}{j} |y_j - \bar{y}_{n-j} q| + |y_j y_{n-j} - \binom{n}{j}^2 q|}{\binom{n}{j}^2 - |y_{n-j}|^2} \right\}.$$

Proof. It is known that $\mathcal{C}_\Omega \leq \mathcal{L}_\Omega$ for any domain $\Omega \subset \mathbb{C}^n$. We only show $\mathcal{C}_{\tilde{\mathbb{G}}_n}(0, y) \geq \mathcal{L}_{\tilde{\mathbb{G}}_n}(0, y)$ whenever $y \in \mathcal{J}_n$. One can write

$$\begin{aligned} \mathcal{L}_{\tilde{\mathbb{G}}_n}(0, y) &:= \inf \{ \tanh^{-1} |\lambda| : \text{there exists } f \in \mathcal{O}(\mathbb{D}, \tilde{\mathbb{G}}_n) \\ &\quad \text{such that } f(0) = (0, \dots, 0), f(\lambda) = y \}. \end{aligned}$$

For any $f \in \mathcal{O}(\mathbb{D}, \tilde{\mathbb{G}}_n)$ satisfying $f(0) = (0, \dots, 0)$ and $f(\lambda) = (y_1, \dots, y_{n-1}, q)$, using the Schwarz lemma for $\tilde{\mathbb{G}}_n$ in [13], we have $|\lambda| \geq \max_{1 \leq j \leq n-1} \{\|\Phi_j(\cdot, y)\|_{H^\infty}\}$. Thus

$$\mathcal{L}_{\tilde{\mathbb{G}}_n}(0, y) \geq \max_{1 \leq j \leq n-1} \{\tanh^{-1} \|\Phi_j(\cdot, y)\|_{H^\infty}\}.$$

Now consider $\lambda_0 = \max_{1 \leq j \leq n-1} \{\|\Phi_j(\cdot, y)\|_{H^\infty}\}$. Since $y \in \mathcal{J}_n$, again by the Schwarz lemma for $\tilde{\mathbb{G}}_n$ in [13], there exists $f \in \mathcal{O}(\mathbb{D}, \tilde{\mathbb{G}}_n)$ satisfying $f(0) = (0, \dots, 0)$ and $f(\lambda_0) = y$. Therefore

$$\begin{aligned} \mathcal{L}_{\tilde{\mathbb{G}}_n}(0, y) &= \max_{1 \leq j \leq n-1} \{\tanh^{-1} \|\Phi_j(\cdot, y)\|_{H^\infty}\} \\ &= \max_{1 \leq j \leq n-1} \left\{ \tanh^{-1} \frac{\binom{n}{j} |y_j - \bar{y}_{n-j} q| + |y_j y_{n-j} - \binom{n}{j}^2 q|}{\binom{n}{j}^2 - |y_{n-j}|^2} \right\}. \end{aligned}$$

The Carathéodory pseudo-distance can be expressed as

$$\mathcal{C}_{\tilde{\mathbb{G}}_n}(0, y) = \sup \{ \tanh^{-1} |f(y)| : f \in \mathcal{O}(\tilde{\mathbb{G}}_n, \mathbb{D}), f((0, \dots, 0)) = 0 \}.$$

For any $\omega \in \mathbb{T}$ and any $j \in \{1, \dots, n-1\}$, the function $\Phi_j(\omega, \cdot)$ is analytic from $\tilde{\mathbb{G}}_n$ to \mathbb{D} and satisfies $\Phi_j(\omega, (0, \dots, 0)) = 0$. Thus

$$\mathcal{C}_{\tilde{\mathbb{G}}_n}(0, y) \geq \tanh^{-1} |\Phi_j(\omega, y)| \quad \text{for any } \omega \in \mathbb{T}.$$

Hence $\mathcal{C}_{\tilde{\mathbb{G}}_n}(0, y) \geq \tanh^{-1} \|\Phi_j(\cdot, y)\|_{H^\infty}$ for any $1 \leq j \leq n-1$. Therefore

$$\mathcal{C}_{\tilde{\mathbb{G}}_n}(0, y) \geq \max_{1 \leq j \leq n-1} \{\tanh^{-1} \|\Phi_j(\cdot, y)\|_{H^\infty}\} = \mathcal{L}_{\tilde{\mathbb{G}}_n}(0, y).$$

■

Remark 4.2. By the last part of the proof of the last theorem it is clear that, for any point $y \in \tilde{\mathbb{G}}_n$, we have

$$\mathcal{C}_{\tilde{\mathbb{G}}_n}(0, y) \geq \max_{1 \leq j \leq n-1} \{\tanh^{-1} \|\Phi_j(\cdot, y)\|_{H^\infty}\}.$$

5. APPENDIX

Calculation for the implication $12 \Rightarrow 13$:

$$\begin{aligned} (1 - |\lambda_0|^2 Z_{\nu,j}^* Z_{\nu,j}) &= \begin{bmatrix} 1 - \frac{|y_j^0|^2}{\binom{n}{j}^2} - \frac{|\lambda_0|^2 |w|^2}{\nu^2} & -\frac{\lambda_0 \bar{y}_j^0 \nu w}{\binom{n}{j}} - \frac{|\lambda_0|^2 y_{n-j}^0 \bar{w}}{\binom{n}{j} \nu} \\ -\frac{\bar{\lambda}_0 y_j^0 \nu \bar{w}}{\binom{n}{j}} - \frac{|\lambda_0|^2 \bar{y}_{n-j}^0 w}{\binom{n}{j} \nu} & 1 - |\lambda_0|^2 \nu^2 |w|^2 - \frac{|\lambda_0|^2 |y_{n-j}^0|^2}{\binom{n}{j}^2} \end{bmatrix}, \\ (1 - Z_{\nu,j}^* Z_{\nu,j})^{-1} &= \frac{1}{\det(1 - Z_{\nu,j}^* Z_{\nu,j})} \begin{bmatrix} 1 - \nu^2 |w|^2 - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} & \frac{\bar{y}_j^0 \nu w}{\binom{n}{j} \bar{\lambda}_0} + \frac{y_{n-j}^0 \bar{w}}{\binom{n}{j} \nu} \\ \frac{y_j^0 \nu \bar{w}}{\binom{n}{j} \lambda_0} + \frac{\bar{y}_{n-j}^0 w}{\binom{n}{j} \nu} & 1 - \frac{|y_j^0|^2}{\binom{n}{j}^2 |\lambda_0|^2} - \frac{|w|^2}{\nu^2} \end{bmatrix}. \end{aligned}$$

Then,

$$\begin{aligned}
& [\mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) \det(1 - Z_{\nu,j}^* Z_{\nu,j})]_{11} \\
&= \det(1 - Z_{\nu,j}^* Z_{\nu,j}) [(1 - |\lambda_0|^2 Z_{\nu}^* Z_{\nu})(1 - Z_{\nu}^* Z_{\nu})^{-1}]_{11} \\
&= 1 - \frac{|y_j^0|^2}{\binom{n}{j}^2} - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} + |q^0|^2 - \frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2} \left(\frac{|\lambda_0|}{\nu^2} + \frac{\nu^2}{|\lambda_0|} \right). \tag{26}
\end{aligned}$$

Note that, $\det(1 - Z_{\nu,j} Z_{\nu,j}^*) = \det(1 - Z_{\nu,j}^* Z_{\nu,j})$ and

$$(1 - Z_{\nu,j} Z_{\nu,j}^*)^{-1} = \frac{1}{\det(1 - Z_{\nu,j}^* Z_{\nu,j})} \begin{bmatrix} 1 - \frac{|w|^2}{\nu^2} - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} & \frac{y_j^0 \bar{w}}{\binom{n}{j} \nu \lambda_0} + \frac{\bar{y}_{n-j}^0 \nu w}{\binom{n}{j}} \\ \frac{\bar{y}_j^0 w}{\binom{n}{j} \nu \bar{\lambda}_0} + \frac{y_{n-j}^0 \nu \bar{w}}{\binom{n}{j}} & 1 - \frac{|y_j^0|^2}{\binom{n}{j}^2 |\lambda_0|^2} - \nu^2 |w|^2 \end{bmatrix}.$$

It follows that

$$\begin{aligned}
& [\mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) \det(1 - Z_{\nu,j}^* Z_{\nu,j})]_{12} \\
&= \det(1 - Z_{\nu,j}^* Z_{\nu,j}) (1 - |\lambda_0|^2) [(1 - Z_{\nu,j} Z_{\nu,j}^*)^{-1} Z_{\nu,j}]_{21} = (1 - |\lambda_0|^2) \left(\frac{w}{\nu} + \frac{q^0}{\lambda_0} \nu \bar{w} \right).
\end{aligned}$$

Similarly we have

$$[\mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) \det(1 - Z_{\nu,j}^* Z_{\nu,j})]_{21} = (1 - |\lambda_0|^2) \left(\frac{\bar{w}}{\nu} + \frac{\bar{q}^0}{\bar{\lambda}_0} \nu w \right).$$

Clearly

$$(Z_{\nu,j} Z_{\nu,j}^* - |\lambda_0|^2) = \begin{bmatrix} \frac{|y_j^0|^2}{\binom{n}{j}^2 |\lambda_0|^2} + \nu^2 |w|^2 - |\lambda_0|^2 & \frac{y_j^0 \bar{w}}{\binom{n}{j} \nu \lambda_0} + \frac{\bar{y}_{n-j}^0 \nu w}{\binom{n}{j}} \\ \frac{\bar{y}_j^0 w}{\binom{n}{j} \nu \bar{\lambda}_0} + \frac{y_{n-j}^0 \nu \bar{w}}{\binom{n}{j}} & \frac{|w|^2}{\nu^2} + \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} - |\lambda_0|^2 \end{bmatrix}$$

Therefore

$$\begin{aligned}
& [\mathcal{K}_{Z_{\nu,j}}(|\lambda_0|) \det(1 - Z_{\nu,j}^* Z_{\nu,j})]_{22} \\
&= \det(1 - Z_{\nu,j}^* Z_{\nu,j}) [(Z_{\nu} Z_{\nu}^* - |\lambda_0|^2)(1 - Z_{\nu} Z_{\nu}^*)^{-1}]_{22} \\
&= -|\lambda_0|^2 - \frac{|y_j^0|^2}{\binom{n}{j}^2} - \frac{|y_{n-j}^0|^2}{\binom{n}{j}^2} + \frac{|q^0|^2}{|\lambda_0|^2} - \frac{|y_j^0 y_{n-j}^0 - \binom{n}{j}^2 q^0|}{\binom{n}{j}^2} \left(|\lambda_0| \nu^2 + \frac{1}{\nu^2 |\lambda_0|} \right).
\end{aligned}$$

Acknowledgments

The author is supported by the Institute postdoctoral fellowship of TIFR-CAM. The author would like to thank Pof. Sourav Pal for so many helpful discussion and suggestions.

The author would also like to thank the referee for making several valuable suggestions and comments which help in improving the exposition of the paper.

REFERENCES

- [1] A. A. Abouhajar, M. C. White and N. J. Young, A Schwarz lemma for a domain related to μ -synthesis, *J. Geom. Anal.* 17 (2007), 717 – 750.
- [2] J. Agler and N. J. Young, A schwarz lemma for symmetrized bidisc, *Bull. London Math. Soc.*, 33 (2001), 175 – 186.

- [3] J. Agler and N. J. Young, The hyperbolic geometry of the symmetrized bidisc, *J. Geom. Anal.* 14 (2004) 375 – 403.
- [4] Bercovici, H., Foiaş, C. and Tannenbaum, A., Structured interpolation theory, *Operator Theory: Advances and Applications*, 47 (1990), 195-220.
- [5] C. Costara, On the spectral Nevanlinna-Pick problem, *Studia Math.*, 170 (2005), 23 – 55.
- [6] Doyle, J. C., Analysis of feedback systems with structured uncertainties, *IEEE Proceedings D, Control Theory and Applications* 129 (1982), no. 6, 242-250.
- [7] A. Edigarian, L. Kosiński and W. Zwonek, The Lempert theorem and the tetrablock, *J. Geom. Anal.*, 23 (2003), 1818 – 1831.
- [8] L. Lempert, Complex geometry in convex domains, *Proc. Intern. Cong. Math.*, Berkeley, CA, (1986) 759 – 765.
- [9] N. Nikolov, P. Pflug, W. Zwonek, The Lempert function of the symmetrized polydisc in higher dimensions is not a distance, *Proc. Amer. Math. Soc.*, 135 (2007), 2921 – 2928.
- [10] S. Pal and S. Roy, A generalized Schwarz lemma for two domains related to μ -synthesis, *Complex Manifolds*, 5 (2018), 1 – 8.
- [11] S. Pal and S. Roy, A Schwarz lemma for two families of domains and complex geometry, *Complex Var. Elliptic Equ.*, 66 (2021), no. 5, 756 – 782.
- [12] S. Pal and S. Roy, Characterizations of the symmetrized polydisc via another family of domains, *International Journal of Mathematics*, 32 (2021), no. 6, 2150036, 29p.
- [13] S. Pal and S. Roy, A Schwarz lemma for the symmetrized polydisc via another family of domains, *preprint*, available at <https://arxiv.org/abs/2110.04819>.
- [14] S. Pal and S. Roy, The complex geometry and a two point interpolation for two families of domains, <https://arxiv.org/abs/1904.03745>.

(Samriddho Roy) TATA INSTITUTE OF FUNDAMENTAL RESEARCH CENTRE FOR APPLICABLE MATHEMATICS, BANGALORE - 560065, INDIA.

Email address: samriddhoroy@gmail.com