

The Twin Radio galaxy TRG J104454+354055

Gopal-Krishna^{1,*}, Ravi Joshi^{2,6,†}, Dusmanta Patra³, Xiaolong Yang^{4,5}, Luis C. Ho^{6,7}, Paul J. Wiita⁸,
Amitesh Omar⁹

¹UM-DAE Centre for Excellence in Basic Sciences, Mumbai, 400098; India

²Indian Institute of Astrophysics, Koramangla, Bangalore, 560034; India

³S N Bose National Centre for Basic Sciences, Kolkata, 700106; India

⁴Shanghai Astronomical Observatory, Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, Shanghai, 200030, China

⁵Shanghai Key Laboratory of Space Navigation and Positioning Techniques, Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai, 200030, China

⁶Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing, 100871, China

⁷Department of Astronomy, School of Physics, Peking University, Beijing, 100871, China

⁸Department of Physics, The College of New Jersey, 2000 Pennington Rd., Ewing, NJ, 08628-0718, USA

⁹Aryabhata Research Institute of observational sciences, Nainital, 263002; India

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We report observations of a bright dumb-bell system of galaxies ($z = 0.162$) with the upgraded Giant Metrewave Radio Telescope (*u*GMRT), which show that each member of this gravitationally bound pair of galaxies hosts bipolar radio jets extended on 100 kiloparsec scales. Only two cases of such radio morphology have been reported previously, both being dumb-bell systems, as well. The famous first example, 3C 75, was discovered 4 decades ago, and the second case was discovered 3 decades ago. This implies that such ‘Twin-Radio-Galaxies’ (TRGs) are an exceedingly rare phenomenon. As in the case of its two senior cousins, the bi-polar radio jets of the present TRG (J104454+354055) exhibit strong wiggles and are edge-darkened (Fanaroff-Riley class I). However, there are important differences, too. For instance, the jets in the present TRG do not merge and, moreover, show no signs of distortion due to an external crosswind. This makes the present TRG a much neater laboratory for studying the physics of (sideway) colliding jets of relativistic plasma. This TRG has a Wide-Angle-Tail (WAT) neighbour hosted by another bright galaxy belonging to the same group, which appears to be moving towards the TRG.

Key words: galaxies: active, galaxies: jets, quasars: supermassive black holes

1 INTRODUCTION

Galaxy mergers are the bedrock of the evolving cosmic structure leading to hierarchical galaxy formation (e.g., Barnes & Hernquist 1996; Bell et al. 2006). Since practically all large galaxies are believed to harbour a supermassive black hole (SMBH) (Richstone et al. 1998; Kormendy & Ho 2013), the possibility of SMBH coalescence due to galaxy mergers (e.g., Begelman et al. 1980; Gualandris et al. 2017, and references therein) has become a driver for the rapidly advancing new field of gravitational-wave astronomy (e.g., Hughes 2003; Amaro-Seoane et al. 2017; Jenkins 2022). From an observational viewpoint, if one or both members in the merging galaxy pair host a radio-loud Active Galactic Nucleus (AGN), this can serve as a direct tracer of the SMBH merger process, via monitoring by Very Long Baseline Interferometry (VLBI). The prospects become much more exciting when *both* AGN are radio loud (Owen et al. 1985; Simpson & Rawlings 2002). The main techniques used to search for SMBH binaries (i.e., dual AGN) include photometric and spectroscopic measurements and radio interferometric imaging which is

particularly promising because of the high resolution it can provide (e.g., Komossa et al. 2003; Liu et al. 2010; Bogdanović 2015; Shanguan et al. 2016; Rubinur et al. 2021). However, the success rate in finding merging SMBHs with discernible radio jets has been minuscule (e.g. Burke-Spolaor 2011; Tremblay et al. 2016), with just a few reasonably compelling cases reported so far.

On the smallest scale (< 0.1 pc), the BL Lac object OJ 287 remains the only strong case for hosting an SMBH binary, as gleaned from its (paired) optical outbursts, recurring every ~ 12 years (Sillanpää et al. 1988). Its long-term optical light-curve has been modelled in terms of a SMBH binary with an orbital semi-major axis of 0.05 pc, whose secondary crosses the accretion disk of the primary SMBH twice during each orbit (e.g., Lehto & Valtonen 1996; Valtonen et al. 2016; Dey et al. 2021; see, also Britzen et al. 2018). Even on slightly larger (parsec) scale, only a couple of dual radio-AGN candidates seem genuine, the most prominent being the source 0402+379 for which VLBI has yielded a SMBH binary separation of ~ 7 pc (Rodríguez et al. 2006; Burke-Spolaor 2011; Bansal et al. 2017). More recently, a dual radio-AGN, with a SMBH separation of just 0.35 pc has been found in the Seyfert galaxy NGC 7674 at $z = 0.0289$ (Kharb et al. 2017).

It is worth noting that while assessing the genuineness of such radio-loud cases, one needs to guard against the possibility that the

* E-mail: gopaltani@gmail.com

† E-mail: rvjohshirv@gmail.com

candidate dual radio-AGN may turn out to be just (i) a pair of knots in a VLBI jet, or (ii) hot spots in a single compact-symmetric radio source (see, e.g., [Wrobel et al. 2014](#)), or (iii) even a case of (superluminous) gravitational milli-lensing ([Gopal-Krishna, & Subramanian 1996](#)). Naturally, the binarity becomes much less ambiguous when the two radio nuclei are separated on kilo-parsec scale (permitting detailed radio imaging), even though such SMBH pairs are clearly remote from an eventual merger. So far, just two cases of gravitationally bound pairs of radio galaxies with kiloparsec-scale jets have been identified, each hosted by a pair of similarly bright elliptical galaxies within a common envelope ("dumb-bell"). The famous first example of such "Twin Radio Galaxy (TRG)" is the proto-type 3C 75, a Wide-Angle-Tail (WAT) radio source of overall size ~ 0.5 Mpc, located near the centre of the galaxy cluster Abell 400 ($z = 0.023$). It was discovered almost 4 decades ago ([Owen et al. 1985](#)), which testifies to the extreme rarity of this cosmic phenomenon. The second TRG, which was discovered 3 decades ago and has received much less attention, is the source PKS 2149-158 of size ~ 400 kpc, located near the outskirts of the cluster Abell 2382 ($z = 0.062$) ([Parma et al. 1991](#); [Guidetti et al. 2008](#)).

In this paper we report the discovery of the third case of a bound pair of galaxies, each of which has a bi-polar radio jet extended > 100 kpc scale. The dumb-bell galaxy lies at the centre of a galaxy group at $z \sim 0.16$ ([Yuan et al. 2016](#)). This source was initially classified as a X-shaped radio galaxy candidate ([Yang et al. 2019](#); [Joshi et al. 2019](#)), but in a follow-up programme with uGMRT, it was found to be a TRG. Here we report radio-optical properties of this TRG and briefly compare it with the other two TRGs mentioned above. Throughout this paper a cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$ is adopted. This gives a luminosity distance of 780 Mpc for the TRG and a scale of 2.797 kpc per arcsecond at the source redshift of $z = 0.16278$.

2 OBSERVATIONS AND ANALYSIS

The inset in Fig. 1a shows a g, r, z colour composite image from the Dark Energy Camera Legacy survey (DECaLS) ([Dey et al. 2019](#)), of the dumb-bell consisting of the galaxies 'A' & 'B', both with $z \approx 0.162$, and a third, fainter galaxy 'C' of unknown redshift, within the same stellar halo. The colour composite image of another bright galaxy 'D', which has a redshift of 0.160 and is located 2.72 arcmin ($= 456$ kpc) west of the dumb-bell is shown in the inset of Fig. 1c. Particulars of these 4 galaxies are given in Table 1. Low-resolution optical spectroscopic observations of the dumb-bell system were carried out using the HFOSC instrument mounted on the 2m-Himalayan Chandra Telescope ([Prabhu 2014](#)), with grisms Gr#7 (3500–7800 Å) and slit-width of 2.0 arcsec. The data were reduced with the standard IRAF¹ package. The optical spectra were modelled using the penalized PiXel-Fitting method (pPXF, [Cappellari & Emsellem 2004](#); [Cappellari 2017](#)) and the measured stellar velocity dispersions of $\sigma_* = 412.2 \pm 34.8 \text{ km s}^{-1}$ and $254.5 \pm 20.4 \text{ km s}^{-1}$ for the galaxies A and B were used to estimate the black-hole masses ([Greene et al. 2020](#), see their table 5) (see Table 1). The non-detection of [O III]λ5007 nebular emission line for both galaxies implies that they are low-excitation radio galaxies (LERGs) ([Best & Heckman 2012](#)).

The radio observations were made using the upgraded Giant Metrewave Radio Telescope (uGMRT), in the band-5 (900 to 1450

MHz), with 400 MHz bandwidth encompassing 2048 channels ([Swarup et al. 1991](#); [Gupta et al. 2017](#)). The data were reduced following the standard procedures, using the Astronomical Image Processing System (AIPS)². While calibrating the data, bad data were flagged out at various stages. The data for antennas with large errors in antenna-based solutions were checked and flagged over the requisite time ranges. The flux calibration was performed using 3C286 and the flux density scale defined by [Perley & Butler \(2013\)](#). The phase calibration was done using the source 1033+395³. The final image was made after several rounds of phase self-calibration, and one round of amplitude self-calibration. An on-source time of 48 minutes resulted in an rms noise of $34 \mu\text{Jy/beam}$, where the beamsize is $3.15'' \times 1.73''$ (P.A. = 71 deg) (Fig. 1).

3 RESULTS

In the uGMRT map, both members of the dumb-bell system, namely A & B (Table 1) are seen to eject wiggling bi-polar radio jets along roughly the same direction (Fig. 1). Both northern jets are brighter than their southern counterparts and, after maintaining a steady mutual separation over the initial ~ 30 kpc (projected), the two northern jets become diffuse but do not appear to merge (see, Section 4). The (fainter) two jets on the southern side remain separated all along, despite strong wiggling. The observed projected extents of the bi-polar jets associated with A and B are ~ 220 kpc, and ~ 140 kpc, respectively (Fig. 1). The 144 MHz radio maps from the LOFAR Two-meter Sky Survey Data Release 2 (LoTSS-DR2) ([Shimwell et al. 2022](#)), has revealed more extended, bright radio emission connected to the southern jets, whereas a faint, long radio spur of size ~ 60 arcsec is detected on the northern side (Fig. 1). Thus, the source has an overall size of at least 3.0 arcmin (i.e., 0.5 Mpc). However, the TRG morphology is not discernible in the LoTSS map, due to its relatively large beam size of 6×6 arcsec (see, Fig. 1). Consistent with their moderate radio luminosities (Table 1), both radio galaxies exhibit an edge-darkened (Fanaroff-Riley class I) morphology ([Fanaroff & Riley 1974](#); [Ledlow & Owen 1996](#)). Fig. 2 shows the integrated radio spectrum ($S_\nu \propto \nu^\alpha$, where ν and S_ν are the frequency and flux density, respectively), of this dumb-bell system (A + B), which is consistent with a straight line of spectral index $\alpha = -0.74$. For the radio cores of the galaxies A and B, the VLASS maps ([Lacy et al. 2020](#)) give flux densities of $5.6 \pm 0.3 \text{ mJy}$, and $2.8 \pm 0.3 \text{ mJy}$ at 3 GHz, respectively. The galaxy C remained undetected in both VLASS and the present uGMRT observations, setting an upper limit of 1.4 mJy at 1.4 GHz. The uGMRT map of the source D (Fig. 1) shows its radio jets emerging at PA ≈ 70 deg, although after ~ 20 arcsec (≈ 56 kpc) the eastern jet undergoes a sharp bending towards the north-west, giving the source a 'Wide-Angle-Tail' (WAT) morphology (see [Owen & Rudnick 1976](#); [Jones & Owen 1979](#)). From the VLASS, its core is found to have a flux density of $1.1 \pm 0.2 \text{ mJy}$ at 3 GHz. Its integrated radio spectral index is -0.75 , based on the NVSS flux density of $23.4 \pm 1.3 \text{ mJy}$ at 1.4 GHz ([Condon et al. 1998](#)) and LoTSS-DR2 flux density of $127.7 \pm 12.2 \text{ mJy}$ at 144 MHz ([Shimwell et al. 2022](#)).

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

² <http://www.aips.nrao.edu>

³ <http://http://www.vla.nrao.edu/astro/calib/manual/csource.html>




Figure 1. *Left Panel:* The radio-optical overlay shows the u GMRT 1.4 GHz radio contours overlaid on to the r band image from the DECaLS (Dey et al. 2019) of the dumb-bell galaxy pair ‘A’ and ‘B’. The inset shows the g , r , z -band color-composite image from the DECaLS of the dumb-bell galaxy pair together with an apparently close radio-quiet galaxy ‘C’. *Middle panel:* shows the gray-scale u GMRT image of the pair of bi-polar jets (‘Twin-Radio-Galaxy’ TRG) overlaid with the 144 MHz LoTSS-DR2 radio contours. The positions of the respective host galaxies are marked with red dots. *Right Panel:* Radio-optical overlay of the u GMRT 1.4 GHz radio contours of the ‘Wide-Angle-Tail’ (WAT) radio galaxy ‘D’ whose g , r , z -band color-composite DECaLS image is shown within the inset. Contours levels are set at square-root increment of $(2, 4, 8, 16, 32, \dots) \times 14\sigma$ for the TRG and $\times 5\sigma$ for the WAT source, where σ rms noise is $34 \mu\text{Jy beam}^{-1}$ for the our u GMRT maps. The LoTSS counter levels are set at $(1, 16, 81, 256, \dots) \times 3\sigma$, where σ rms noise is $87 \mu\text{Jy beam}^{-1}$.

4 DISCUSSION

Indubitably, a key message from the present study is that Twin-Radio-Galaxies (TRGs), each having a kiloparsec-scale bi-polar radio jet, are an extremely rare phenomenon. The source J104454+354055 reported here becomes the third known TRG, the first one (3C 75) was reported 37 years ago and the second one (PKS 2149-158) 31 years ago. All three TRGs are associated with dumb-bell systems residing in galaxy clusters/groups (Sect. 1). A prognosis of their extreme rarity came long ago from a VLA survey of 88 compact groups of galaxies (Menon & Hickson 1985). These authors concluded that any bright radio emission arising from a galaxy group is usually associated with its brightest member galaxy, while the remaining group members are radio-quiet. They found a strong preference for the radio-loud elliptical galaxy in a group to be the first-ranked; even relatively luminous ellipticals holding second- or third- rank were found to be radio-quiet. A robust theoretical underpinning to this empirical result still remains elusive.

Several parallels can be drawn between the present TRG and its two senior cousins, namely the famous and extensively studied 3C 75 and the TRG PKS 2149-158 (Sect. 1). As mentioned above, the galaxy pairs hosting all these 3 TRGs reside in a group/cluster environment, a circumstance clearly more conducive to the formation of dumb-bell systems. Secondly, all 3 TRGs exhibit FR I morphology, consistent with their moderate radio luminosities. Their radio jets, imaged at 1.4 GHz with a similar resolution of ~ 5 arcsec, too are similarly extended on 100 kpc scales. Also, the apparent separation between their twin host galaxies is around 10 kpc (between 8 and 18 kpc) and the line-of-sight velocity difference between the paired host

galaxies are quite similar: $\sim 430 \text{ km s}^{-1}$ (3C 75) (Lauer et al. 2014), $\sim 546 \text{ km s}^{-1}$ (PKS 2149-158) (Owen et al. 1995; Smith et al. 2004), and $\sim 300 \text{ km s}^{-1}$ (present TRG) (see, Table 1). However, beyond these basic similarities, salient differences are noticeable between the 3 TRGs. Below, we enumerate some of these contrasts, as they might hold vital clues from the perspective of theoretical modelling and numerical simulations, especially once the other two TRGs have been observed in detail comparable to the proto-type 3C 75.

(i) Whilst all 3 TRGs exhibit conspicuously wiggly jets, the two northern jets of 3C 75 are seen to merge into a single entity, fairly early on (a similar situation probably exists in the other TRG, PKS 2149-158, although even its best available (VLA) map lacks sufficient clarity, see (Guidetti et al. 2008). This contrasts with the present TRG in which both bi-polar jets retain their distinct identities much farther out, providing a greater space for their kinematical modelling.

(ii) In 3C 75, situated at the centre of the cluster Abell 400, all four jets undergo a sharp bending towards east/northeast (Owen et al. 1985). This is attributed to dynamic pressure exerted by an intra-cluster crosswind blowing at a speed of $\sim 1200 \text{ km s}^{-1}$ (Hudson et al. 2006), due to the ongoing merger of the host cluster Abell 400 with a subcluster (e.g., Beers et al. 1992; Hudson et al. 2006). This extraneous factor poses a complication to analytical modelling and numerical simulations of the basic phenomenon of twin bi-polar jets (e.g., Molnar et al. 2017; Musoke et al. 2020, see also, Steinborn et al. 2016). A similar structural complexity appears to plague the twin bi-polar jets of the second TRG PKS 2149-158, which exhibit a conspicuous C-shaped bending (Guidetti et al. 2008), again indicating a strong dynamical pressure acting on the jets. In contrast, the present TRG is a much neater system, as the impact of the orbital motion

of the two galaxies A and B on the morphologies of their bi-polar jets is not contaminated by ram-pressure of a cross-wind. This would simplify the dynamical modelling of the twin bi-polar jets, since there is no need to disentangle the effect of the orbital motion of the host galaxy pair from extraneous forces due to cross-wind impinging on the jets. To appreciate this point, we may recall that the orbital motion of the paired host galaxies is expected to impose a mirror-symmetric wiggling on the bi-polar jet pair, as is indeed observed in the uGMRT image of the present system and which also lends further support to the assertion that the two bi-polar jets are physically associated (Fig. 1). In the other two TRGs, the wiggly pattern of jets is massively distorted due to the dynamical impact of the intracluster cross-wind, as mentioned above. The inference then is that the present TRG is a superior laboratory for probing the physics of two nearly parallel wiggly jets of relativistic plasma bouncing sideways against each other, but escaping disruption (Fig. 1). Also, all four jets of the present TRG are bright enough for radio polarimetry, which would enable mapping of their magnetic field and investigating its possible role in preserving the identity of the relativistic plasma jets bouncing against each other.

4.1 The WAT Neighbour

As mentioned in Section 3, the uGMRT map also shows a WAT radio galaxy (D), located 2.72 arcmin (≈ 456 kpc) to the west of the present TRG (Fig. 1c). The twin-jets of the WAT initially advance along PA ~ 70 deg, for ~ 60 kpc, when the eastern jet undergoes a sharp bend towards northwest. Plausibly, the strong dynamic pressure required for the jet bending comes from motion of the host galaxy towards the dumb-bell system and its associated gaseous halo. In order to assess such a possibility, we estimate the zone of gravitational influence of the dumb-bell, i.e., its virial radius, which is roughly the range out to which the gaseous halo of the dumb-bell system might extend.

Using the 2MASS K-band apparent magnitudes of the dumb-bell constituents A and B, of $K = 14.48$, and 14.89 , respectively, we estimate their absolute magnitudes (M_K) taking a distance modulus of 39.4 and K-corrections of -0.27 and -0.28 (Chilingarian et al. 2010; Chilingarian & Zolotukhin 2012). This yields $M_K = -25.1$ and -24.77 for A and B, respectively. These correspond to stellar masses of $\log M_* \sim 11.50 M_\odot$ and $11.35 M_\odot$ for A and B (Cappellari 2013, equation 2), amounting to a total stellar mass of $\log M_* \sim 11.73 M_\odot$ for the galaxy pair. Galaxy stellar-to-halo mass relation (SHMR) implies their dark halo virial masses to be $\log M_{vir} \sim 14.1 M_\odot$ and $\sim 13.8 M_\odot$, respectively (Moster et al. 2010, equation 2; Wechsler & Tinker 2018). Such massive haloes are typically assembled by a redshift of $z_a \sim 0.8$, and have virial radii given by Shull (2014):

$$\frac{R_{vir}}{206 \text{ kpc}} \approx \left(\frac{M_{vir}}{10^{12} M_\odot} \right)^{1/3} (1 + z_a)^{-1} \quad (1)$$

For the total mass of the dumb-bell system ($\log M_{vir} \sim 14.3 M_\odot$), its virial radius is then expected to extend to ~ 670 kpc, i.e., beyond the galaxy D identified with the WAT and is thus available for exerting the dynamic pressure required for the observed bending of its jet (Fig. 1c).

5 CONCLUSIONS

The uGMRT observations reported here have revealed of a pair of bi-polar extended radio jets emanating from a dumb-bell system con-




Figure 2. Integrated radio spectrum of the present TRG J1044+3540, using the flux densities from the 144 MHz LoTSS-DR2 (Shimwell et al. 2022), 325 MHz WENSS (Rengelink et al. 1997) and NVSS 1.4 GHz (Condon et al. 1998). The beamsize for the latter two maps is around 0.75 - 0.9 arcmin. Therefore, the LoTSS-DR2 map was smoothed to a beamwidth of 0.8 arcmin. The measured integrated flux densities at the 3 frequencies are: 590 ± 31 mJy (144 MHz), 333 ± 67 mJy (325 MHz) and 108 ± 22 mJy (1.4 GHz).

taining two gravitationally bound massive elliptical galaxies within a common stellar envelope. This newly discovered ‘Twin-Radio-Galaxy’ (TRG), J104454+354055, is the third TRG found so far. The first (famous) example of this extremely rare class, the TRG 3C 75 was discovered 37 years ago, followed by the discovery of a second (much less known) TRG PKS 2149-158 three decades ago. All four jets of the present TRG are extended on 100-kiloparsec scale and exhibit an edge-darkened radio morphology (FR I), as well as a mirror-symmetric wiggly pattern which is consistent with both host galaxies performing orbital motion about a common gravitational centre (the recent LoTSS/DR2 map shows a total size of ~ 0.5 Mpc). Unlike the other two TRGs, there is no indication that the jets of the present TRG merge with each other early on, or become distorted due to a cross-wind. This goes to make this TRG a much neater exponent of cosmic tango, more amenable to modelling the jet dynamics in TRGs and for studying the impact of sideways collisions of relativistic plasma jets on their survivability (e.g., Achterberg 1988; Musoke et al. 2020). A third bright radio galaxy in this group is found to be a Wide-Angle-Tail (WAT) apparently in motion directed towards the dumb-bell system. Radio spectral and polarimetric imaging of the present TRG, supplemented with X-ray observations yielding gas pressures within the group, and numerical simulations, would be very desirable for a deeper understanding of the evolution of closely interacting synchrotron jets.

ACKNOWLEDGEMENTS

GK acknowledges a Senior Scientist fellowship of the Indian National Science Academy (INSA). DP acknowledges the post-doctoral fellowship of the S. N. Bose National Centre for Basic Sciences, Kolkata, India, funded by the Department of Science and Technology (DST), India. XLY thanks the National Science Foundation of China (12103076), the Shanghai Sailing Program (21YF1455300) and the China Postdoctoral Science Foundation (2021M693267) for support, LCH was supported by the National Science Foundation of China (11721303, 11991052, 12011540375) and the China Manned Space Project (CMS-CSST-2021-A04). This research has made use of the ‘‘K-corrections calculator’’ service available at

Table 1. Basic parameters of the 4 galaxies based on optical and radio μ GMRT observations.

Source	R.A. (J2000)	Dec. (J2000)	redshift z	$\log M_{\text{BH}}$ $M_{\odot}$	Type	Flux density (mJy)			Length (arcsec)			Radio Luminosity $\log(L_R)$ (erg/s)
						Core [†]	North-jet	South-jet	North-jet	South-jet	Total	
A	10 ^h 44 ^m 55.36 ^s	+35°40′53.5″	0.16173	10.2±0.4	LERG	5.6 ± 0.3	21.8	40.8	29.7	51.3	78.9	42.43
B	10 ^h 44 ^m 54.87 ^s	+35°40′55.0″	0.16278	9.3±0.3	LERG	2.8 ± 0.3	36.1	23.5	31.7	20.0	51.3	42.36
C	10 ^h 44 ^m 54.67 ^s	+35°40′57.1″	—	—	—	≤ 1.4 (5 σ)	—	—	—	—	—	—
D	10 ^h 44 ^m 41.85 ^s	+35°40′16.7″	0.16007	9.4±0.2	LERG	1.1 ± 0.2	19.5	20.4	46.6	26.1	73.8	41.87

[†] The core flux densities are estimated using the VLASS radio map at 2.5 GHz (Lacy et al. 2020). The other radio parameters are derived from 1.4 GHz μ GMRT maps (present work).

* The redshift of the galaxies B and D are taken from the SDSS archive (Ahumada et al. 2020).

<http://kcor.sai.msu.ru/>. We thank the staff of the GMRT that made these observations possible. GMRT is run by the National Centre for Radio Astrophysics of the Tata Institute of Fundamental Research. We thank the staff of IAO, Hanle and CREST, Hosakote, that made these observations possible. The facilities at IAO and CREST are operated by the Indian Institute of Astrophysics, Bangalore.

DATA AVAILABILITY

The radio data used in this study will be publicly available at μ GMRT data archive at <https://naps.ncra.tifr.res.in/goa/data/search>

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.

References

- Achterberg A., 1988, *A&A*, **191**, 167
 Ahumada R., et al., 2020, *ApJS*, **249**, 3
 Amaro-Seoane P., et al., 2017, arXiv e-prints, p. arXiv:1702.00786
 Bansal K., Taylor G. B., Peck A. B., Zavala R. T., Romani R. W., 2017, *ApJ*, **843**, 14
 Barnes J. E., Hernquist L., 1996, *ApJ*, **471**, 115
 Beers T. C., Gebhardt K., Huchra J. P., Forman W., Jones C., Bothun G. D., 1992, *ApJ*, **400**, 410
 Begelman M. C., Blandford R. D., Rees M. J., 1980, *Nature*, **287**, 307
 Bell E. F., Phleps S., Somerville R. S., Wolf C., Borch A., Meisenheimer K., 2006, *ApJ*, **652**, 270
 Best P. N., Heckman T. M., 2012, *MNRAS*, **421**, 1569
 Bogdanović T., 2015, in *Gravitational Wave Astrophysics*. p. 103 (arXiv:1406.5193), doi:10.1007/978-3-319-10488-1_9
 Britzen S., et al., 2018, *MNRAS*, **478**, 3199
 Burke-Spolaor S., 2011, *MNRAS*, **410**, 2113
 Cappellari M., 2013, *ApJ*, **778**, L2
 Cappellari M., 2017, *MNRAS*, **466**, 798
 Cappellari M., Emsellem E., 2004, *PASP*, **116**, 138
 Chilingarian I. V., Zolotukhin I. Y., 2012, *MNRAS*, **419**, 1727
 Chilingarian I. V., Melchior A.-L., Zolotukhin I. Y., 2010, *MNRAS*, **405**, 1409
 Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, **115**, 1693
 Dey A., et al., 2019, *AJ*, **157**, 168
 Dey L., Valtonen M. J., Gopakumar A., Lico R., Gómez J. L., Susobhanan A., Komossa S., Pihajoki P., 2021, *MNRAS*, **503**, 4400
 Fanaroff B. L., Riley J. M., 1974, *MNRAS*, **167**, 31P
 Gopal-Krishna, Subramanian K., 1996, *A&A*, **315**, 343
 Greene J. E., Strader J., Ho L. C., 2020, *ARA&A*, **58**, 257
 Gualandris A., Read J. I., Dehnen W., Bortolas E., 2017, *MNRAS*, **464**, 2301
 Guidetti D., Murgia M., Govoni F., Parma P., Gregorini L., de Ruiter H. R., Cameron R. A., Fanti R., 2008, *A&A*, **483**, 699
 Gupta Y., et al., 2017, *Current Science*, **113**, 707
 Hudson D. S., Reiprich T. H., Clarke T. E., Sarazin C. L., 2006, *A&A*, **453**, 433
 Hughes S. A., 2003, *Annals of Physics*, **303**, 142
 Jenkins A. C., 2022, arXiv e-prints, p. arXiv:2202.05105
 Jones T. W., Owen F. N., 1979, *ApJ*, **234**, 818
 Joshi R., et al., 2019, *ApJ*, **887**, 266
 Kharb P., Lal D. V., Merritt D., 2017, *Nature Astronomy*, **1**, 727
 Komossa S., Burwitz V., Hasinger G., Predehl P., Kaastra J. S., Ikebe Y., 2003, *ApJ*, **582**, L15
 Kormendy J., Ho L. C., 2013, *Annual Review of Astronomy and Astrophysics*, **51**, 511
 Lacy M., et al., 2020, *PASP*, **132**, 035001
 Lauer T. R., Postman M., Strauss M. A., Graves G. J., Chisari N. E., 2014, *ApJ*, **797**, 82
 Ledlow M. J., Owen F. N., 1996, *AJ*, **112**, 9
 Lehto H. J., Valtonen M. J., 1996, *ApJ*, **460**, 207
 Liu X., Greene J. E., Shen Y., Strauss M. A., 2010, *ApJ*, **715**, L30
 Menon T. K., Hickson P., 1985, *ApJ*, **296**, 60
 Molnar S. M., Schive H. Y., Birkinshaw M., Chiueh T., Musoke G., Young A. J., 2017, *ApJ*, **835**, 57
 Moster B. P., Somerville R. S., Maubetsch C., van den Bosch F. C., Macciò A. V., Naab T., Oser L., 2010, *ApJ*, **710**, 903
 Musoke G., Young A. J., Molnar S. M., Birkinshaw M., 2020, *MNRAS*, **494**, 5207
 Owen F. N., Rudnick L., 1976, *ApJ*, **205**, L1
 Owen F. N., O’Dea C. P., Inoue M., Eilek J. A., 1985, *ApJ*, **294**, L85
 Owen F. N., Ledlow M. J., Keel W. C., 1995, *AJ*, **109**, 14
 Parma P., de Ruiter H. R., Cameron R. A., 1991, *AJ*, **102**, 1960
 Perley R. A., Butler B. J., 2013, *ApJS*, **204**, 19
 Prabhu T. P., 2014, *Proceedings of the Indian National Science Academy Part A*, **80**, 887
 Rengelink R. B., Tang Y., de Bruyn A. G., Miley G. K., Bremer M. N., Roettgering H. J. A., Bremer M. A. R., 1997, *A&AS*, **124**, 259
 Richstone D., et al., 1998, *Nature*, **395**, A14
 Rodriguez C., Taylor G. B., Zavala R. T., Peck A. B., Pollack L. K., Romani R. W., 2006, *ApJ*, **646**, 49
 Rubinur K., Kharb P., Das M., Rahna P. T., Honey M., Paswan A., Vaddi S., Murthy J., 2021, *MNRAS*, **500**, 3908
 Shangguan J., Liu X., Ho L. C., Shen Y., Peng C. Y., Greene J. E., Strauss M. A., 2016, *ApJ*, **823**, 50
 Shimwell T. W., et al., 2022, *A&A*, **659**, A1
 Shull J. M., 2014, *ApJ*, **784**, 142
 Sillanpää A., Haarala S., Valtonen M. J., Sundelius B., Byrd G. G., 1988, *ApJ*, **325**, 628
 Simpson C., Rawlings S., 2002, *MNRAS*, **334**, 511
 Smith R. J., et al., 2004, *AJ*, **128**, 1558
 Steinborn L. K., Dolag K., Comerford J. M., Hirschmann M., Remus R.-S., Teklu A. F., 2016, *MNRAS*, **458**, 1013
 Swarup G., Ananthakrishnan S., Kapahi V. K., Rao A. P., Subrahmanya C. R., Kulkarni V. K., 1991, *Current Science*, **60**, 95
 Tremblay S. E., Taylor G. B., Ortiz A. A., Tremblay C. D., Helmboldt J. F., Romani R. W., 2016, *Monthly Notices of the Royal Astronomical Society*, **459**, 820
 Valtonen M. J., et al., 2016, *ApJ*, **819**, L37
 Wechsler R. H., Tinker J. L., 2018, *ARA&A*, **56**, 435
 Wrobel J. M., Walker R. C., Fu H., 2014, *ApJ*, **792**, L8
 Yang X., et al., 2019, *ApJS*, **245**, 17
 Yuan Z. S., Han J. L., Wen Z. L., 2016, *MNRAS*, **460**, 3669