

A HYBRIDIZABLE DISCONTINUOUS GALERKIN METHOD FOR THE FULLY COUPLED TIME-DEPENDENT STOKES/DARCY-TRANSPORT PROBLEM

AYCIL CESMELIOGLU¹, DINH DONG PHAM¹ AND SANDER RHEBERGEN²

Abstract. We present a high-order hybridized discontinuous Galerkin (HDG) method for the fully coupled time-dependent Stokes–Darcy-transport problem where the fluid viscosity and source/sink terms depend on the concentration and the dispersion/diffusion tensor depends on the fluid velocity. This HDG method is such that the discrete flow equations are compatible with the discrete transport equation. Furthermore, the HDG method guarantees strong mass conservation in the H^{div} sense and naturally treats the interface conditions between the Stokes and Darcy regions via facet variables. We employ a linearizing decoupling strategy where the Stokes/Darcy and the transport equations are solved sequentially by time-lagging the concentration. We prove well-posedness and optimal a priori error estimates for the velocity and the concentration in the energy norm. We present numerical examples that respect compatibility of the flow and transport discretizations and demonstrate that the discrete solution is robust with respect to the problem parameters.

2020 Mathematics Subject Classification. 65N12, 65N15, 65N30, 76D07, 76S99.

1. INTRODUCTION

Coupled free fluid and porous media flow is encountered in many engineering applications [24, 34] and can be modeled by the Stokes/Darcy equations. Adding a transport equation to this coupled system brings forth a model that can be used to simulate the spread of contaminants towards groundwater resources [4] or biochemical transport in hemodynamics [21].

The accuracy and stability of numerical discretizations of the stationary Stokes/Darcy equations [3, 11–13, 25, 29, 37, 39, 41] and advection-diffusion type transport equations [10, 20, 40, 52] are well studied. However, accuracy and stability are not automatically guaranteed when these discretizations are coupled. In particular, *compatible* discretizations, as defined by [22], are desired to avoid loss of accuracy and loss of conservation properties of the numerical methods used for the transport equation.

The first numerical study on the coupling of the stationary Stokes/Darcy equations with a transport equation was given in [51] where a mixed finite element method (MFEM) is used for the flow problem and the local discontinuous Galerkin method is used for the transport problem. They considered one-way coupling; the concentration is affected by the flow velocity, but the velocity is not affected by a change in concentration. The same problem was studied in [43] by using discontinuous Galerkin (DG) methods for both flow and transport equations. In [28], Ervin et. al considered a fully time-dependent version of the one-way coupled problem where they developed partitioned time-stepping methods by imposing the interface conditions weakly using penalties. One-way coupling was considered also in [16] in which the flow problem was discretized by a strongly mass conservative Embedded-Hybridized DG (EDG-HDG) method while the transport equation was discretized by an EDG method.

Keywords and phrases: Stokes/Darcy flow, coupled flow and transport, advection–diffusion, hybridized methods, discontinuous Galerkin, multiphysics.

¹ Oakland University, Department of Mathematics and Statistics, Rochester Hills, MI, 48307, USA, e-mail: cesmelio@oakland.edu & ddpham@oakland.edu

² University of Waterloo, Department of Applied Mathematics, Waterloo, ON, Canada, e-mail: srheberg@uwaterloo.ca

Less studied is the fully-coupled problem in which, apart from the transport equation depending on the flow velocity, the flow solution is time-dependent and the fluid viscosity and source/sink terms depend on the concentration. To the best of our knowledge, there are only two papers that focus on this fully coupled problem. First, [14] presented an analysis of a weak solution for the case where free flow is governed by the Navier–Stokes equations. The analysis in [14], however, also holds when free flow is governed by the Stokes equations. The only numerical paper on this topic, [45], introduced a stabilized mixed finite element method using nonconforming piece-wise linear Crouzeix–Raviart finite elements for the velocity, a piece-wise constant approximation for the pressure, and a conforming, piece-wise linear, finite element method for the transport equation based on a skew-symmetric formulation.

In this paper, we extend the work in [16] to the fully coupled case. To deal with the non-linearity, we consider a linearizing decoupling strategy, where the Stokes/Darcy and the transport equations are solved sequentially by time-lagging the concentration. We use HDG methods [19] for both the Stokes/Darcy and transport sub-problems at each time step and prove well-posedness and a priori error estimates. These results can easily be extended to the EDG-HDG discretization used for the Stokes/Darcy problem in [16]. Our HDG method for the flow problem provides the transport sub-problem at each time step with an exactly mass conserving and $H(\text{div})$ -conforming velocity field. This renders our scheme robust with respect to the problem parameters. By choosing the polynomial degree in a specific way our flow/transport scheme is also compatible.

Here is an outline for the remainder of this article. In Section 2, we present the fully coupled Stokes/Darcy-transport model and specify the assumptions on the problem parameters. Section 3 sets notation, describes in detail the semi-discrete HDG scheme, and lists the attractive properties of the numerical discretization. Next, Section 4 summarizes standard inequalities and shows continuity, coercivity, and the inf-sup condition for the discretization of the Stokes/Darcy sub-problem. A full discretization of the problem based on a sequential decoupling strategy is introduced in Section 5 while the main results, i.e., a priori error estimates for the velocity, pressure, and concentration, are presented in Section 6. Finally, we present some numerical experiments in Section 7 followed by conclusions in Section 8.

2. THE STOKES/DARCY–TRANSPORT SYSTEM

Let $\Omega \subset \mathbb{R}^{\text{dim}}$, $\text{dim} = 2, 3$, be a bounded polygonal domain. We denote its boundary by $\partial\Omega$ and the outward




FIGURE 1. A depiction of a domain $\Omega \subset \mathbb{R}^2$ with its two sub-domains Ω^s and Ω^d .

unit normal to $\partial\Omega$ by n . Domain Ω consists of two non-overlapping polygonal regions, a free flow region Ω^s and a Darcy flow region Ω^d , such that $\Omega = \Omega^s \cup \Omega^d$. The polygonal interface between Ω^s and Ω^d is denoted by Γ^I and the external boundary of Ω^j is denoted by $\Gamma^j := \partial\Omega \cap \partial\Omega^j$, $j = s, d$. See Figure 1 for a depiction of a domain when $\text{dim} = 2$.

We denote the time interval of interest by $J = [0, T]$. The fully coupled Stokes/Darcy–transport system for the velocity field $u : \Omega \times J \rightarrow \mathbb{R}^{\dim}$, fluid pressure $p : \Omega \times J \rightarrow \mathbb{R}$ and concentration $c : \Omega \times J \rightarrow \mathbb{R}$ is given by

$$\partial_t u - \nabla \cdot (2\mu(c)\varepsilon(u)) + \nabla p = f^s(c) \quad \text{in } \Omega^s \times J, \quad (1a)$$

$$\mathbb{K}^{-1}(c)u + \nabla p = \mathbb{K}^{-1}(c)f^d(c) \quad \text{in } \Omega^d \times J, \quad (1b)$$

$$-\nabla \cdot u = \chi_d(g_p - g_i) \quad \text{in } \Omega \times J, \quad (1c)$$

$$\phi \partial_t c + \nabla \cdot (cu - \tilde{D}(u)\nabla c) = \chi_d(c_I g_i - c g_p) \quad \text{in } \Omega \times J, \quad (1d)$$

$$u = 0 \quad \text{on } \Gamma^s \times J, \quad (1e)$$

$$u \cdot n = 0 \quad \text{on } \Omega^d \times J, \quad (1f)$$

$$\tilde{D}(u)\nabla c \cdot n = 0 \quad \text{on } \partial\Omega \times J, \quad (1g)$$

where $\varepsilon(u) := (\nabla u + (\nabla u)^T)/2$ is the strain rate tensor and χ_d is the characteristic function that takes the value 1 in Ω^d and 0 in Ω^s . Here the fluid viscosity μ , the matrix $\mathbb{K} = \frac{\kappa}{\mu}$, where κ is the permeability matrix of the porous medium, and the body force terms f^s and f^d are concentration dependent functions. The porosity ϕ of the medium in Ω^d is a spatially varying function. In Ω^s we set $\phi = 1$. The functions g_i and g_p denote the source and sink terms related to injection and production wells and c_I is the injected concentration. Furthermore, in the Stokes region $\tilde{D}(u) = d\mathbb{I}$, where \mathbb{I} is the $\dim \times \dim$ identity matrix and d is the diffusion coefficient. In the Darcy region $\tilde{D}(u) = D(u)$, where $D(u)$ denotes the diffusion dispersion tensor in Ω^d .

We will denote the restriction of the velocity u , pressure p , and concentration c to sub-domain Ω^j , $j = s, d$ by, respectively, u^j , p^j , and c^j . Then, on the interface Γ^I , choosing the unit normal vector n to be pointing from Ω^s to Ω^d , we prescribe the following interface conditions that hold for $t \in J$:

$$u^s \cdot n = u^d \cdot n, \quad (2a)$$

$$p^s - (2\mu(c^s)\varepsilon(u^s)n) \cdot n = p^d, \quad (2b)$$

$$-2(\varepsilon(u^s)n) \cdot \tau^\ell = \gamma^\ell u^s \cdot \tau^\ell, \quad \ell = 1, \dots, \dim - 1, \quad (2c)$$

$$c^s = c^d, \quad (2d)$$

$$d\nabla c^s \cdot n = D(u^d)\nabla c^d \cdot n, \quad (2e)$$

where τ^ℓ , $\ell = 1, \dots, \dim - 1$ denote the unit tangent vectors on Γ^I . These conditions enforce the normal continuity of the velocity (2a), the normal continuity of the normal component of the stress (2b), continuity of the concentration (2d), and normal continuity of the concentration flux (2e). Equation (2c), where $\gamma^\ell = \alpha/\sqrt{\tau^\ell \cdot \kappa \tau^\ell}$ with $\alpha > 0$ a constant, is the Beavers–Joseph–Saffman law which enforces a condition on the tangential component of the normal stress [5, 46].

To close the model, we assume the following initial conditions:

$$u^s(x, 0) = u_0^s(x) \quad \text{in } \Omega^s, \quad (3a)$$

$$c(x, 0) = c_0(x) \quad \text{in } \Omega. \quad (3b)$$

We end this section by discussing some assumptions we make on the various functions used in the Stokes/Darcy–transport model. The dispersion-diffusion tensor $D(u)$ in Ω^d satisfies for $u, v \in \mathbb{R}^{\dim}$:

$$D_{\min}|\xi|^2 \leq \xi^T D(u)\xi \quad \forall \xi \in \mathbb{R}^{\dim}, \quad (4a)$$

$$|D(u)| \leq C(1 + |u|), \quad (4b)$$

$$|D(u) - D(v)| \leq C|u - v|, \quad (4c)$$

where D_{\min} and C are positive constants and $|\cdot|$ denotes the Euclidean norm. We assume that μ is Lipschitz continuous in c with Lipschitz constant μ_L and that there exist constants $\phi_*, \phi^*, \mu_*, \mu^* > 0$ such that

$$\phi_* \leq \phi(x) \leq \phi^* \quad \forall x \in \Omega^d, \quad (5a)$$

$$\mu_* \leq \mu(c) \leq \mu^* \quad \forall c \in \mathbb{R}. \quad (5b)$$

The permeability matrix κ is symmetric, uniformly bounded, and elliptic, that is, there exist positive constants $\kappa_* < \kappa^*$ such that

$$\kappa_* |\xi|^2 \leq \xi^T \kappa(x) \xi \leq \kappa^* |\xi|^2 \quad \forall \xi \in \mathbb{R}^{\dim}, \quad \forall x \in \bar{\Omega}^d. \quad (6)$$

From (6) and (5b), we deduce that

$$K_* |\xi|^2 \leq \xi^T \mathbb{K}(c, x) \xi \leq K^* |\xi|^2 \quad \forall \xi \in \mathbb{R}^{\dim}, \quad \forall (c, x) \in \mathbb{R} \times \bar{\Omega}^d, \quad (7)$$

where $K_* = \kappa_*/\mu^*$ and $K^* = \kappa^*/\mu_*$.

The body force functions f^s and f^d are assumed to be Lipschitz continuous in c with Lipschitz constants L_f^s and L_f^d . Note that f^s and f^d depend on x and c , but they do not depend explicitly on t . We will further assume that $0 \leq c_I \leq 1$ a.e. in Ω^d and that $g_i, g_p \geq 0$, $g_i, g_p \in L^\infty(J; L^2(\Omega^d))$ are such that

$$\int_{\Omega^d} (g_i(x, t) - g_p(x, t)) \, dx = 0 \quad \forall t \in J.$$

A weak formulation of the problem defined by eqs. (1) to (3) was presented in [45]. The analysis for a weak solution of a more general version of this problem, in which the free fluid flow is governed by the Navier–Stokes equations, can be found in [14].

3. THE HYBRIDIZED DISCONTINUOUS GALERKIN METHOD

3.1. Preliminaries

We use the same notation that we used previously in [16, 17]. Let $\mathcal{T}^j := \{K\}$ be a shape-regular triangulation of Ω^j , $j = s, d$, into non-overlapping elements (we only consider simplices) such that \mathcal{T}^s and \mathcal{T}^d match at the interface Γ^I . We define the triangulation of the entire domain Ω as $\mathcal{T} := \mathcal{T}^s \cup \mathcal{T}^d$. The maximum diameter over all elements is $h = \max_{K \in \mathcal{T}} h_K$, where h_K stands for the diameter of an element K . The boundary of an element K and its outward unit normal are denoted by ∂K and n , respectively. A facet of an element boundary is an interior facet if it is shared by two neighboring elements and it is a boundary facet if it is a part of $\partial \Omega$. The set of all interior facets and all boundary facets in $\bar{\Omega}^j$ are denoted by \mathcal{F}_i^j and \mathcal{F}_b^j , $j = s, d$, respectively. We also collect the facets that lie on the interface Γ^I in the set \mathcal{F}^I . The set of all facets that lie in $\bar{\Omega}$ and in $\bar{\Omega}^j$ are denoted by \mathcal{F} and \mathcal{F}^j , respectively. We point out that $\mathcal{F}^j = \mathcal{F}_i^j \cup \mathcal{F}_b^j \cup \mathcal{F}^I$, $j = s, d$. Furthermore, we define $\Gamma_0 := \cup_{F \in \mathcal{F}} F$ and $\Gamma_0^j := \cup_{F \in \mathcal{F}^j} F$, $j = s, d$.

The finite element function spaces on Ω for the velocity and pressure are given by

$$\begin{aligned} V_h &:= \{v_h \in [L^2(\Omega)]^{\dim} : v_h \in [P_{k_f}(K)]^{\dim}, \forall K \in \mathcal{T}\}, \\ Q_h &:= \{q_h \in L^2(\Omega) : q_h \in P_{k_f-1}(K), \forall K \in \mathcal{T}\} \cap L_0^2(\Omega), \end{aligned} \quad (8)$$

where $P_k(K)$ denotes the space of polynomials of degree at most k defined on the element K . The finite element spaces for the velocity and pressure traces are given by

$$\begin{aligned} \bar{V}_h &:= \{\bar{v}_h \in [L^2(\Gamma_0^s)]^{\dim} : \bar{v}_h \in [P_{k_f}(F)]^{\dim} \forall F \in \mathcal{F}^s, \bar{v}_h = 0 \text{ on } \Gamma^s\}, \\ \bar{Q}_h^j &:= \{\bar{q}_h^j \in L^2(\Gamma_0^j) : \bar{q}_h^j \in P_{k_f}(F) \forall F \in \mathcal{F}^j\}, \quad j = s, d. \end{aligned} \quad (9)$$

Here $P_k(F)$ denotes the space of polynomials of degree at most k defined on the facet F . Note that functions in \bar{V}_h are not defined on $\Gamma_0^d \setminus \Gamma_I$. The finite element function spaces for the concentration and its trace are defined as

$$C_h = \{c_h \in L^2(\Omega) : c_h \in P_{k_c}(K), \forall K \in \mathcal{T}\}, \quad \bar{C}_h = \{\bar{c}_h \in L^2(\Gamma_0) : \bar{c}_h \in P_{k_c}(F) \forall F \in \mathcal{F}\}. \quad (10)$$

The semi-discrete and fully-discrete HDG methods for the flow and transport equations considered in this article are compatible when $k_c = k_f - 1$ [16]. For this reason we set $k_c = k_f - 1$.

To reduce the notational burden, we define $\mathbf{V}_h := V_h \times \bar{V}_h$, $\mathbf{Q}_h := Q_h \times \bar{Q}_h^s \times \bar{Q}_h^d$, and $\mathbf{Q}_h^j := Q_h^j \times \bar{Q}_h^j$, $j = s, d$. We denote elements in these product spaces by $\mathbf{v}_h := (v_h, \bar{v}_h) \in \mathbf{V}_h$, $\mathbf{q}_h := (q_h, \bar{q}_h^s, \bar{q}_h^d) \in \mathbf{Q}_h$, and $\mathbf{q}_h^j := (q_h^j, \bar{q}_h^j) \in \mathbf{Q}_h^j$, $j = s, d$. In addition, we set $\mathbf{X}_h := \mathbf{V}_h \times \mathbf{Q}_h$. Similarly, we introduce $\mathbf{C}_h = C_h \times \bar{C}_h$ and denote the corresponding elements by $\mathbf{c}_h := (c_h, \bar{c}_h) \in \mathbf{C}_h$.

Next, let us define the function spaces

$$\begin{aligned} V &:= \{v \in [L^2(\Omega)]^{\dim} : v^s \in [H^2(\Omega^s)]^{\dim}, v^d \in [H^1(\Omega^d)]^{\dim}, \\ &\quad v = 0 \text{ on } \Gamma^s, v \cdot n = 0 \text{ on } \Gamma^d, v^s \cdot n = v^d \cdot n \text{ on } \Gamma^I\}, \\ Q &:= \{q \in L_0^2(\Omega) : q^s \in H^1(\Omega^s), q^d \in H^2(\Omega^d)\}, \\ C &:= H^2(\Omega), \end{aligned}$$

and set $X := V \times Q$. As before, we use a superscript j to specify the restriction of these spaces to Ω^j , $j = s, d$. The trace spaces of V restricted to Γ_0^s , Q^j restricted to Γ_0^j , and C restricted to Γ_0 are denoted by, respectively, \bar{V} , \bar{Q}^j , and \bar{C} . The trace operator $\gamma_V : V^s \rightarrow \bar{V}$ restricts functions in V^s to Γ_0^s , and similarly the trace operators $\gamma_{Q^j} : Q^j \rightarrow \bar{Q}^j$ restrict functions in Q^j to Γ_0^j , $j = s, d$. However, when it is clear from the context, we omit the subscript in the trace operator. Analogous to the discrete case, we introduce $\mathbf{V} := V \times \bar{V}$, $\mathbf{Q} := Q \times \bar{Q}^s \times \bar{Q}^d$, and $\mathbf{C} := C \times \bar{C}$. We then define extended function spaces as

$$\mathbf{V}(h) := \mathbf{V}_h + \mathbf{V}, \quad \mathbf{Q}(h) := \mathbf{Q}_h + \mathbf{Q}, \quad \mathbf{C}(h) := \mathbf{C}_h + \mathbf{C},$$

and set $\mathbf{X}(h) := \mathbf{V}(h) \times \mathbf{Q}(h)$.

We close this section by listing various norms on the spaces described above. We refer the reader to [1] for the definitions of the standard Sobolev spaces $W^{m,p}(D)$ and their corresponding norms $\|\cdot\|_{W^{m,p}(D)}$. For ease of notation, we write $\|\cdot\|_{m,p,D}$ instead of $\|\cdot\|_{W^{m,p}(D)}$ with the following simplifications. When $m = 0$, $W^{0,p}(D)$ coincides with $L^p(D)$ and when $p = 2$, $H^m(D) = W^{m,p}(D)$. For $p = 2$, we write $\|\cdot\|_{m,D}$ to denote $\|\cdot\|_{W^{m,2}(D)}$ and for $m = 0$, $p = 2$, we write $\|\cdot\|_D$ instead of $\|\cdot\|_{0,D}$.

On $\mathbf{V}^s(h)$ we define the standard HDG-norm and its strengthened version as follows:

$$\|\mathbf{v}\|_{v,s}^2 := \sum_{K \in \mathcal{T}^s} (\|\nabla v\|_K^2 + h_K^{-1} \|v - \bar{v}\|_{\partial K}^2), \quad \|\mathbf{v}\|_{v',s}^2 := \|\mathbf{v}\|_{v,s}^2 + \sum_{K \in \mathcal{T}^s} h_K^2 |v|_{H^2(K)}^2.$$

On $\mathbf{V}(h)$ we then introduce the norms

$$\begin{aligned} \|\mathbf{v}\|_v^2 &:= \|\mathbf{v}\|_{v,s}^2 + \|v\|_{\Omega^d}^2 + \sum_{j=1}^{\dim} \gamma^j \|\bar{v} \cdot \tau_j\|_{\Gamma^I}^2, \\ \|\mathbf{v}\|_{v'}^2 &:= \|\mathbf{v}\|_v^2 + \sum_{K \in \mathcal{T}^s} h_K^2 |v|_{H^2(K)}^2 = \|\mathbf{v}\|_{v',s}^2 + \|v\|_{\Omega^d}^2 + \sum_{j=1}^{\dim} \gamma^j \|\bar{v} \cdot \tau_j\|_{\Gamma^I}^2, \end{aligned}$$

and note that $\|\mathbf{v}\|_v$ and $\|\mathbf{v}\|_{v'}$ are equivalent on \mathbf{V}_h due to the fact that $\|\cdot\|_{v,s}$ and $\|\cdot\|_{v',s}$ are equivalent on \mathbf{V}_h^s (see, for example, [52, eq. (5.5)]).

On the pressure spaces $\mathbf{Q}^j(h)$, $j = s, d$ and $\mathbf{Q}(h)$, we define, respectively,

$$\|\mathbf{q}^j\|_{p,j}^2 := \|q\|_{\Omega^j}^2 + \sum_{K \in \mathcal{T}^j} h_K \|\bar{q}^j\|_{\partial K}^2, \quad \|\mathbf{q}\|_p^2 := \sum_{j=s,d} \|\mathbf{q}^j\|_{p,j}^2.$$

Finally, for $\mathbf{w}_h \in \mathbf{C}(h)$, we define the following semi-norm:

$$\|\mathbf{w}_h\|_c^2 = \sum_{K \in \mathcal{T}} (\|\nabla w_h\|_K^2 + h_K^{-1} \|w_h - \bar{w}_h\|_{\partial K}^2). \quad (11)$$

3.2. Semi-discrete HDG scheme

The semi-discrete method we propose for the Stokes/Darcy-transport system in eqs. (1) and (2) is as follows: for $t > 0$, find $(\mathbf{u}_h(t), \mathbf{p}_h(t)) \in \mathbf{X}_h$ and $\mathbf{c}_h(t) \in \mathbf{C}_h$ such that

$$\begin{aligned} \sum_{K \in \mathcal{T}^s} \int_K \partial_t u_h \cdot v_h \, dx + B_h^{sd}(\mathbf{c}_h; (\mathbf{u}_h, \mathbf{p}_h), (\mathbf{v}_h, \mathbf{q}_h)) &= \sum_{K \in \mathcal{T}^s} \int_K f^s(c_h) \cdot v_h \, dx \\ &+ \sum_{K \in \mathcal{T}^d} \int_K \mathbb{K}^{-1}(c_h) f^d(c_h) \cdot v_h \, dx + \sum_{K \in \mathcal{T}^d} \int_K (g_p - g_i) q_h \, dx \end{aligned} \quad (12a)$$

and

$$\sum_{K \in \mathcal{T}} \int_K \phi \partial_t c_h w_h \, dx + B_h^{tr}(u_h; \mathbf{c}_h, \mathbf{w}_h) + \sum_{K \in \mathcal{T}^d} \int_K c_h g_p w_h \, dx = \sum_{K \in \mathcal{T}^d} \int_K c_I g_i w_h \, dx, \quad (12b)$$

for all $(\mathbf{v}_h, \mathbf{q}_h) \in \mathbf{X}_h$ and $\mathbf{w}_h \in \mathbf{C}_h$.

The form B_h^{sd} in eq. (12a) collects the discretization terms for the Stokes/Darcy momentum and mass conservation equations as follows:

$$B_h^{sd}(\mathbf{c}; (\mathbf{u}, \mathbf{p}), (\mathbf{v}, \mathbf{q})) := a_h(\mathbf{c}; \mathbf{u}, \mathbf{v}) + \sum_{j=s,d} \left(b_h^j(\mathbf{p}^j, v) + b_h^{I,j}(\bar{p}^j, \bar{v}) \right) + \sum_{j=s,d} \left(b_h^j(\mathbf{q}^j, u) + b_h^{I,j}(\bar{q}^j, \bar{u}) \right). \quad (13)$$

Here $a_h(\cdot, \cdot)$ is defined as

$$a_h(\mathbf{c}; \mathbf{u}, \mathbf{v}) := a_h^s(c; \mathbf{u}, \mathbf{v}) + a_h^d(c; u, v) + a_h^I(\bar{c}; \bar{u}, \bar{v}), \quad (14)$$

where

$$\begin{aligned} a_h^s(c; \mathbf{u}, \mathbf{v}) &:= \sum_{K \in \mathcal{T}^s} \int_K 2\mu(c) \varepsilon(u) : \varepsilon(v) \, dx + \sum_{K \in \mathcal{T}^s} \int_{\partial K} \frac{2\beta_s \mu(c)}{h_K} (u - \bar{u}) \cdot (v - \bar{v}) \, ds \\ &\quad - \sum_{K \in \mathcal{T}^s} \int_{\partial K} 2\mu(c) \varepsilon(u) n^s \cdot (v - \bar{v}) \, ds - \sum_{K \in \mathcal{T}^s} \int_{\partial K} 2\mu(c) \varepsilon(v) n^s \cdot (u - \bar{u}) \, ds, \\ a_h^d(c; u, v) &:= \int_{\Omega^d} \mathbb{K}^{-1}(c) u \cdot v \, dx, \\ a_h^I(\bar{c}; \bar{u}, \bar{v}) &:= \sum_{\ell=1}^{\dim-1} \int_{\Gamma^I} \gamma^\ell \mu(\bar{c}) (\bar{u} \cdot \tau^\ell) (\bar{v} \cdot \tau^\ell) \, ds, \end{aligned}$$

and $\beta_s > 0$ is a penalty parameter. The bilinear forms $b_h^j(\cdot, \cdot)$ and $b_h^{I,j}(\cdot, \cdot)$ in eq. (13), $j = s, d$ are defined as

$$b_h^j(\mathbf{p}^j, v) := - \sum_{K \in \mathcal{T}^j} \int_K p \nabla \cdot v \, dx + \sum_{K \in \mathcal{T}^j} \int_{\partial K} \bar{p}^j v \cdot \mathbf{n}^j \, ds, \quad (15a)$$

$$b_h^{I,j}(\bar{p}^j, \bar{v}) := - \int_{\Gamma^I} \bar{p}^j \bar{v} \cdot \mathbf{n}^j \, ds. \quad (15b)$$

Before defining the terms related to the transport equation, we point out that eqs. (13) to (15) are the same as in [17] when the viscosity and κ are both constants.

The form $B_h^{tr}(u; \mathbf{c}(t), \mathbf{w})$ in eq. (12b) discretizes the advective and diffusive parts of the transport equation:

$$B_h^{tr}(u; \mathbf{c}, \mathbf{w}) = B_h^a(u; \mathbf{c}, \mathbf{w}) + B_h^d(u; \mathbf{c}, \mathbf{w}). \quad (16)$$

The advective part is defined as

$$B_h^a(u; \mathbf{c}, \mathbf{w}) := - \sum_{K \in \mathcal{T}} \int_K c u \cdot \nabla w \, dx + \sum_{K \in \mathcal{T}} \int_{\partial K} c u \cdot \mathbf{n} (w - \bar{w}) \, ds - \sum_{K \in \mathcal{T}} \int_{\partial K^{\text{in}}} u \cdot \mathbf{n} (c - \bar{c}) (w - \bar{w}) \, ds, \quad (17)$$

where ∂K^{in} denotes the inflow portion of the boundary on which $u_h \cdot \mathbf{n} < 0$, and the diffusive part is defined as

$$\begin{aligned} B_h^d(u; \mathbf{c}, \mathbf{w}) := & \sum_{K \in \mathcal{T}} \int_K \tilde{D}(u) \nabla c \cdot \nabla w \, dx + \sum_{K \in \mathcal{T}} \frac{\beta_{tr}}{h_K} \int_{\partial K} [\tilde{D}(u) \mathbf{n}] (c - \bar{c}) \cdot (w - \bar{w}) \mathbf{n} \, ds \\ & - \sum_{K \in \mathcal{T}} \int_{\partial K} [\tilde{D}(u) \nabla c] \cdot \mathbf{n} (w - \bar{w}) \, ds - \sum_{K \in \mathcal{T}} \int_{\partial K} ([\tilde{D}(u) \nabla w] \cdot \mathbf{n}) (c - \bar{c}) \, ds, \end{aligned} \quad (18)$$

where $\beta_{tr} > 0$ is a penalty parameter. Here we pause again to mention that eqs. (16) to (18) are the same as in [16], and a standard extension of the discretization analyzed in [52].

To complete the discretization, we project the initial conditions u_0 and c_0 eq. (3) into \mathbf{V}_h and \mathbf{C}_h , respectively.

3.3. Properties of the numerical scheme

The semi-discrete HDG scheme presented in Section 3.2 has various attractive features. Besides local momentum conservation, a property of all HDG methods, this particular HDG method also conserves mass strongly, according to the definition defined in [35].

To be specific, the discrete velocity enjoys the following properties:

$$-\nabla \cdot u_h = \chi^d \Pi_Q(g_p - g_i) \quad \forall x \in K, \forall K \in \mathcal{T}, \quad (19a)$$

$$\llbracket u_h \cdot \mathbf{n} \rrbracket = 0 \quad \forall x \in F, \forall F \in \mathcal{F} \setminus \mathcal{F}^I, \quad (19b)$$

$$u_h^j \cdot \mathbf{n} = \bar{u}_h \cdot \mathbf{n} \quad \forall x \in F, \forall F \in \mathcal{F}^I, j = s, d, \quad (19c)$$

where $\llbracket \cdot \rrbracket$ is the usual jump operator and \mathbf{n} is the unit normal vector on F . Note that eqs. (19b) and (19c) imply that u_h is $H(\text{div})$ -conforming on the whole domain. More details on eq. (19) can be found in [17, Section 3.3]. Additionally, the scheme is consistent, that is, the solution to eqs. (1) to (3) satisfies eq. (12), as we discuss next.

Lemma 3.1 (Consistency). *Suppose that the solution (u, p, c) to the Stokes/Darcy-transport system eqs. (1) to (3) satisfies $(u, p) \in L^2(0, T; X)$, $\partial_t u \in L^2(0, T; L^2(\Omega^s))$, $c \in L^2(0, T; C)$, and $\partial_t c \in L^2(0, T; L^2(\Omega))$. Then $(\mathbf{u}(t), \mathbf{p}(t), \mathbf{c}(t))$, where $\mathbf{u} := (u, \gamma(u))$, $\mathbf{p} := (p, \gamma(p^s), \gamma(p^d))$, $\mathbf{c} := (c, \gamma(c))$, satisfy the semi-discrete HDG scheme eq. (12) for all $t > 0$.*

Proof. The proof is as in [17, Lemma 1] and [16, Lemma 6] with minor modifications. We do not repeat the proof here, but mention that it is based on integration by parts in $a_h^s(c; \mathbf{u}, \mathbf{v})$, $\sum_{j=s,d} b_h^j(\mathbf{p}^j, v)$, $B_h^a(u, \mathbf{c}, \mathbf{w})$, and $B_h^d(u, \mathbf{c}, \mathbf{w})$, using $\gamma(u) = u$ on Γ_0^s , $\gamma(p^j) = p^j$ on Γ_0^j , $j = s, d$, $\gamma(c) = c$ on Γ_0 , the smoothness of the solution, the continuity of μ and D , and eqs. (1) and (2). \square

4. CONTINUITY, COERCIVITY, AND AN INF-SUP CONDITION

Let us recollect various known inequalities. Throughout this article we denote by $C > 0$ a generic constant that is independent of the mesh size and the time step. From [23, Lemma 1.46, Remark 1.47], for any $K \in \mathcal{T}$, we have

$$\|v\|_{\partial K} \leq Ch_K^{-1/2} \|v\|_K \quad \forall v \in P_k(K). \quad (20)$$

We will also use the following versions of the continuous trace inequality [8, Theorem 1.6.6, (10.3.8)]:

$$\|v\|_{\partial K}^2 \leq C (h_K^{-1} \|v\|_K^2 + h_K \|v\|_{1,K}^2) \quad \forall v \in H^1(K), \quad (21)$$

$$\|v\|_{0,\infty,\partial K} \leq C \quad \forall v \in W^{1,\infty}(K), \quad (22)$$

where C in eq. (22) depends on $\|v\|_{1,\infty,K}$. Regarding the trace on the interface, we have [31, (1.24)], [8, Theorem 1.6.6]:

$$\|v\|_{\Gamma^I} \leq C \|\nabla v\|_{\Omega^s} \quad \forall v \in \{v \in H^1(\Omega^s) : v = 0 \text{ on } \Gamma^s\}, \quad (23)$$

$$\|v\|_{\Gamma^I} \leq C \|v\|_{1,\Omega^s} \quad \forall v \in H^1(\Omega^s). \quad (24)$$

Furthermore, by [31, Theorem 4.4], for any $\mathbf{v}_h \in \mathbf{V}_h$, for $k_f \geq 1$,

$$\|v_h^s\|_{\Gamma^I} \leq C \|\mathbf{v}_h\|_{v,s} \leq C \|\mathbf{v}_h\|_{v'}. \quad (25)$$

Similarly, for any $w_h \in C_h$, for $k_c \geq 1$,

$$\|w_h^s\|_{\Gamma^I} \leq C \|\mathbf{w}_h\|_c. \quad (26)$$

Next, we recall some inverse inequalities from [23, Lemma 1.44, Lemma 1.50]:

$$\|\nabla v\|_K \leq Ch_K^{-1} \|v\|_K \quad \forall v \in P_k(K), \quad (27)$$

$$\|v\|_{0,\infty,K} \leq Ch_K^{-\dim/2} \|v\|_K \quad \forall v \in P_k(K). \quad (28)$$

The following Poincaré-type inequality follows from [31, Proposition 4.5], [6, Remark 1.1]:

$$\|v\|_{\Omega^s} \leq C \|\mathbf{v}\|_{v,s} \quad \forall \mathbf{v} := (v, \mu) \in H^1(\mathcal{T}_h^s) \times \bar{V}_h. \quad (29)$$

The following version of Korn's first inequality is a consequence of [7, (1.19)], [31, Proposition 4.7], [42, p.110]:

$$\|\mathbf{v}_h\|_{v,s} \leq C \sum_{K \in \mathcal{T}^s} (\|\varepsilon(v_h)\|_K^2 + h_K^{-1} \|v_h - \bar{v}_h\|_{\partial K}^2). \quad (30)$$

Continuity and coercivity $a_h(\cdot, \cdot)$, follow from [17, Lemma 2, Lemma 3] keeping in mind that μ satisfies eq. (5b) and \mathbb{K} satisfies eq. (7). They can be stated as follows:

Lemma 4.1 (Continuity and coercivity of a_h). *There exists a constant $C > 0$, independent of h , such that for all $\mathbf{u}, \mathbf{v} \in \mathbf{V}(h)$ and $\mathbf{c} \in \mathbf{C}(h)$,*

$$a_h(\mathbf{c}; \mathbf{u}, \mathbf{v}) \leq C \|\mathbf{u}\|_{v'} \|\mathbf{v}\|_{v'}. \quad (31)$$

In addition, there exists a constant $C_a > 0$, independent of h but dependent on κ^* , μ_* , and μ^* , and a constant $\beta_0^s > 0$ such that if $\beta_s > \beta_0^s$, then

$$a_h(c_h; \mathbf{v}_h, \mathbf{v}_h) \geq C_a \|\mathbf{v}_h\|_v^2 \quad \forall \mathbf{v}_h \in \mathbf{V}_h, \quad \forall c_h \in C_h. \quad (32)$$

The inf-sup condition on the discrete spaces \mathbf{V}_h and \mathbf{Q}_h was proved in [17] in the case of a continuous discrete velocity trace space $\bar{V}_h \cap [C^0(\Gamma_0^s)]^{\dim}$. It is straightforward to show that the inf-sup condition also holds when the discrete velocity trace space is the larger discontinuous \bar{V}_h space.

Theorem 4.2. *There exists a constant $c_{\inf}^* > 0$, independent of h , such that for any $\mathbf{q}_h \in \mathbf{Q}_h$,*

$$c_{\inf}^* \|\mathbf{q}_h\|_p \leq \sup_{\substack{\mathbf{v}_h \in \mathbf{V}_h \\ \mathbf{v}_h \neq \mathbf{0}}} \frac{\sum_{j=s,d} (b_h^j(\mathbf{q}_h^j, v_h) + b_h^{I,j}(\bar{q}_h^j, \bar{v}_h))}{\|\mathbf{v}_h\|_v}. \quad (33)$$

Note that the proof of this theorem as well as the error analysis requires appropriate interpolation operators onto V_h and \bar{V}_h . For V_h we consider the BDM interpolation operator $\Pi_V : [H^1(\Omega)]^{\dim} \rightarrow V_h$ which is such that if $u \in [H^{k_f+1}(K)]^{\dim}$, $K \in \mathcal{T}$, then (see, for example, [33, Lemma 7] and [9, Section III.3]):

$$\int_K q(\nabla \cdot u - \nabla \cdot \Pi_V u) dx = 0, \quad \forall q \in P_{k_f-1}(K), \quad (34a)$$

$$\int_F \bar{q}(u - \Pi_V u) \cdot n ds = 0, \quad \forall \bar{q} \in P_{k_f}(F), \quad F \subset \partial K, \quad (34b)$$

$$\Pi_V u \in H(\text{div}; \Omega), \quad (34c)$$

$$\|u - \Pi_V u\|_{m,K} \leq Ch_K^{\ell-m} \|u\|_{\ell,K}, \quad m = 0, 1, 2 \quad m \leq \ell \leq k_f + 1, \quad (34d)$$

where we remark that F is an edge if $\dim = 2$ and a face if $\dim = 3$. Furthermore, for $u \in W^{1,\infty}(K)$, [32, (2.33)],

$$\|u - \Pi_V u\|_{\infty,K} + h_K \|\nabla(u - \Pi_V u)\|_{\infty,K} \leq Ch_K \|u\|_{1,\infty,K}. \quad (35)$$

The interpolant onto the trace space \bar{V}_h is defined by $\bar{\Pi}_V : [H^1(\Omega^s)]^{\dim} \rightarrow \bar{V}_h$ such that

$$\bar{\Pi}_V u = \begin{cases} (P_{\bar{V}} u)|_F & \text{if } F \in \mathcal{F}^s \setminus \mathcal{F}^I, \\ (\Pi_V u)^s|_F & \text{if } F \in \mathcal{F}^I, \end{cases}$$

where $P_{\bar{V}}$ is the L^2 -projection onto \bar{V}_h . It is straightforward to deduce the following estimates using eq. (21) and the fact that $h = \max_{K \in \mathcal{T}} h_K$: For $v \in [H^\ell(\Omega^s)]^{\dim}$, $1 \leq \ell \leq k_f + 1$,

$$\|v - \bar{\Pi}_V v\|_{\partial K} \leq Ch_K^{\ell-1/2} \|v\|_{\ell,K}, \quad (36)$$

$$\|\Pi_V v - \bar{\Pi}_V v\|_{\partial K} \leq Ch_K^{\ell-1/2} \|v\|_{\ell,K}. \quad (37)$$

We finish this section by noting that by eqs. (34a) to (34c) the solution u of eqs. (1) and (2) under the assumption that $u \in [H^{k+1}(K)]^{\dim}$, for all $K \in \mathcal{T}$, satisfies

$$\sum_{j=s,d} (b_h^j(\mathbf{q}_h^j, u - \Pi_V u) + b_h^{I,j}(q_h^j, \gamma(u^s) - \bar{\Pi}_V u)) = 0 \quad \forall \mathbf{q}_h \in \mathbf{Q}_h. \quad (38)$$

5. FULLY DISCRETE NUMERICAL SCHEME

Let us now describe the fully discrete HDG method and decoupling strategy used to solve the Stokes/Darcy and transport problems sequentially. For the time discretization, we partition the time interval J as: $0 = t^0 < t^1 < \dots < t^N = T$. For simplicity, we assume a uniform partition with $t^{n+1} - t^n = \Delta t$ for $0 \leq n \leq N-1$. We denote a function $h(t)$ at time level t^n by $h^n := h(t^n)$ and for a sequence $\{u^n\}_{n \geq 1}$ we denote by $d_t u^n = (u^n - u^{n-1})/\Delta t$ a first order difference operator.

In the first step of our sequential algorithm, given an initial velocity u_h^0 in the Stokes domain and an initial concentration \mathbf{c}_h^0 , we solve the Stokes/Darcy problem and obtain a velocity in the entire region. This velocity, with properties given by eq. (19), is then substituted into the concentration problem. This approach is repeated for all time steps with the initial velocity and concentration being replaced by the last computed velocity and concentration solutions. We summarize the fully discrete problem in Algorithm 1.

Algorithm 1 Sequential algorithm

Set $u_h^0 = \Pi_V u_0$, $\mathbf{c}_h^0 = (\Pi_C c_0, \bar{\Pi}_C c_0)$.

for $n = 1, \dots, N$ **do**

1. Find $(\mathbf{u}_h^n, \mathbf{p}_h^n) \in \mathbf{X}_h$ such that for all $(\mathbf{v}_h, \mathbf{q}_h) \in \mathbf{X}_h$

$$\begin{aligned} \sum_{K \in \mathcal{T}^s} \int_{\partial K} d_t u_h^n \cdot \mathbf{v}_h \, dx + B_h^{sd}(\mathbf{c}_h^{n-1}; (\mathbf{u}_h^n, \mathbf{p}_h^n), (\mathbf{v}_h, \mathbf{q}_h)) &= \sum_{K \in \mathcal{T}^s} \int_K f^s(\mathbf{c}_h^{n-1}) \cdot \mathbf{v}_h \, dx \\ &+ \sum_{K \in \mathcal{T}^d} \int_K \mathbb{K}^{-1}(\mathbf{c}_h^{n-1}) f^d(\mathbf{c}_h^{n-1}) \cdot \mathbf{v}_h \, dx + \sum_{K \in \mathcal{T}^d} \int_K (g_p^n - g_i^n) q_h \, dx. \end{aligned} \quad (39)$$

2. Find $\mathbf{c}_h^n \in \mathbf{C}_h$ such that for all $\mathbf{w}_h \in \mathbf{C}_h$

$$\sum_{K \in \mathcal{T}} \int_K \phi d_t \mathbf{c}_h^n w_h \, dx + B_h^{tr}(u_h^n; \mathbf{c}_h^n, \mathbf{w}_h) + \sum_{K \in \mathcal{T}^d} \int_K g_p^n \mathbf{c}_h^n w_h \, dx = \sum_{K \in \mathcal{T}^d} \int_K c_I g_i^n w_h \, dx. \quad (40)$$

end for

Remark 5.1. In Algorithm 1, Π_C denotes the L^2 -projection onto C_h and $\Pi_V u_0$ is understood as Π_V applied to the extension of u_0 to Ω by zero assuming $u_0 \in H_0^1(\Omega^s)$. We note that this choice of u_h^0 satisfies normal continuity across the interfaces in Γ_0^s and has zero divergence in Ω^s under the additional assumption that $\nabla \cdot u_0 = 0$ in Ω^s . We further remark that the properties in eq. (19) hold for \mathbf{u}_h^n for each time step $n = 0, \dots, N$.

We conclude this section by stating some preliminary results obtained by Taylor's theorem [15, Lemma 3.2]. For a function z defined on $D \times [0, T]$, assuming enough regularity, we have the following results:

$$\sum_{m=1}^n \|\partial_t z^m - d_t z^m\|_D^2 \leq C \Delta t \|\partial_{tt} z\|_{L^2(0, T; L^2(D))}^2, \quad (41a)$$

$$\Delta t \sum_{m=1}^n \|d_t z^m\|_{\ell, D}^2 \leq \|\partial_t z\|_{L^2(0, T; H^\ell(D))}^2, \quad \ell = 0, 1, \quad (41b)$$

where $\|f\|_{L^2(a, b; X)} := (\int_a^b \|f(t)\|_X^2 dt)^{1/2}$. Note that the inequalities in eq. (41) for $\ell = 0$ were presented in [15, Lemma 3.2] and that it is straightforward to extend eq. (41b) to $\ell = 1$.

6. MAIN RESULTS

In this section we present our main results. For the error estimates we will make use of the following definition of the discrete in time norm:

$$\|f\|_{\ell^2(0,T;X)} = \left(\Delta t \sum_{m=1}^n \|f^m\|_X^2\right)^{1/2}.$$

Before proving a priori error estimates for the discrete velocity, pressure, and concentration, we first state the well-posedness of the discrete Stokes/Darcy problem eq. (39). Well-posedness of the discrete transport problem eq. (40) is proven in Section 6.3 as it depends on results obtained in Section 6.1.

Theorem 6.1. *Let $\beta_s > \beta_s^0$ be as in Lemma 4.1 and $n \geq 1$. Then given $u_h^{n-1} \in V_h$ and $c_h^{n-1} \in C_h$, there exists a unique solution $(u_h^n, p_h^n) \in X_h$ to eq. (39) that satisfies*

$$\|u_h^n\|_v + \|p_h^n\|_p \leq C \left(\frac{1}{\Delta t} \|u_h^{n-1}\|_{\Omega^s} + \|f^s(c_h^{n-1})\|_{\Omega^s} + \frac{1}{K_*} \|f^d(c_h^{n-1})\|_{\Omega^d} + \|g_p^n - g_i^n\|_{\Omega^d} \right). \quad (42)$$

Proof. The result follows by applying the abstract theory for saddle point problems [27, Theorem 2.34] together with Theorem 4.2 and eq. (32). \square

6.1. Error estimates for the discrete velocity

In this section we derive estimates for the error $u_h^n - u^n$, for each $n \geq 0$, given error estimates for the discrete concentration in previous time steps. To do so, we define the following:

$$\begin{aligned} \xi_u^n &:= u^n - \Pi_V u^n, & \zeta_u^n &:= u_h^n - \Pi_V u^n, & \xi_p^n &:= p^n - \Pi_Q p^n, & \zeta_p^n &:= p_h^n - \Pi_Q p^n, \\ \bar{\xi}_u^n &:= \gamma(u^{sn}) - \bar{\Pi}_V u^n, & \bar{\zeta}_u^n &:= \bar{u}_h^n - \bar{\Pi}_V u^n, & \bar{\xi}_p^{jn} &:= \gamma(p^{jn}) - \bar{\Pi}_Q^j p^{jn}, & \bar{\zeta}_p^{jn} &:= \bar{p}_h^j - \bar{\Pi}_Q^j p^{jn}, \end{aligned}$$

where Π_Q is the L^2 -projection onto Q_h and $\bar{\Pi}_Q^j$ is the L^2 -projection onto \bar{Q}_h^j , $j = s, d$. For the case $n = 0$, $\Pi_V u^0$ is understood as Π_V applied to the extension of u_0 to Ω by zero assuming $u_0 \in H_0^1(\Omega^s)$. Therefore, $\zeta_u^0 = 0$. Furthermore, note that the following identities hold:

$$u^n - u_h^n = \xi_u^n - \zeta_u^n, \quad \gamma(u^n) - \bar{u}_h^n = \bar{\xi}_u^n - \bar{\zeta}_u^n, \quad (43)$$

$$p^n - p_h^n = \xi_p^n - \zeta_p^n, \quad \gamma(p^{jn}) - \bar{p}_h^{jn} = \bar{\xi}_p^{jn} - \bar{\zeta}_p^{jn}, \quad j = s, d. \quad (44)$$

To be consistent with the notation used in previous sections, we set $\ell_u^n = (\ell_u^n, \bar{\ell}_u^n)$, $\ell_p^n := (\ell_p^n, \bar{\ell}_p^{sn}, \bar{\ell}_p^{dn})$, and $\ell_p^{jn} := (\ell_p^{jn}, \bar{\ell}_p^{jn})$, for $\ell = \xi, \zeta$ and $j = s, d$.

Here we recall the following results on the interpolation errors [17, Lemma 7, Lemma 8]. Suppose that u is such that $u^s \in [H^\ell(\Omega^s)]^{\dim}$ and $u^d \in [H^{\ell-1}(\Omega^d)]^{\dim}$ for $2 \leq \ell \leq k_f + 1$, and that $p^j \in H^r(\Omega^j)$ for $0 \leq r \leq k_f$ and $j = s, d$. Then

$$\|\xi_u\|_{v',s} \leq Ch^{\ell-1} \|u\|_{H^\ell(\Omega^s)}, \quad (45a)$$

$$\|\xi_u\|_{v'} \leq Ch^{\ell-1} (\|u\|_{H^\ell(\Omega^s)} + \|u\|_{H^{\ell-1}(\Omega^d)}), \quad (45b)$$

$$\|\xi_p^j\|_{p,j} \leq Ch^r \|p\|_{H^r(\Omega^j)}. \quad (45c)$$

Lemma 6.2. *Let (u, p) be the velocity solution of eqs. (1) to (3), $\bar{u} = \gamma(u^s)$. Then for any $n \geq 1$,*

$$\sum_{j=s,d} (b_h^j(\xi_p^{jn}, v_h) + b_h^{I,j}(\bar{\xi}_p^{jn}, \bar{v}_h)) = 0, \quad \forall v_h \in V_h, \quad (46)$$

$$\sum_{j=s,d} (b_h^j(q_h^j, \xi_u^n) + b_h^{I,j}(\bar{q}_h^j, \bar{\xi}_u^n)) = 0, \quad \forall q_h \in Q_h. \quad (47)$$

Proof. The proof is based on the properties of the numerical scheme eq. (19), the properties of the BDM projection Π_V in eq. (34), and the properties of the L^2 -projections Π_Q and $\bar{\Pi}_Q^j$, $j = s, d$. Indeed, eq. (46) follows after noting that $\nabla \cdot v_h \in P_{k_f-1}(K)$, $v_h \cdot n^j, \bar{v}_h \cdot n^j \in P_{k_f}(F)$, and using the definitions of the L^2 -projections Π_Q and $\bar{\Pi}_Q^j$, $j = s, d$,

$$\begin{aligned} \sum_{j=s,d} (b_h^j(\xi_p^{jn}, v_h) + b_h^{I,j}(\bar{\xi}_p^{jn}, \bar{v}_h)) &= \sum_{j=s,d} \left(- \sum_{K \in \mathcal{T}^j} \int_K (p^n - \Pi_Q p^n) \nabla \cdot v_h \, dx \right. \\ &\quad \left. + \sum_{K \in \mathcal{T}^j} \int_{\partial K} (\gamma(p^{jn}) - \bar{\Pi}_Q^j p^{jn}) v_h \cdot n^j \, ds - \int_{\Gamma^I} (\gamma(p^{jn}) - \bar{\Pi}_Q^j p^{jn}) \bar{v}_h \cdot n^j \, ds \right) = 0, \end{aligned}$$

while eq. (47) is exactly the same as eq. (38), evaluated at $t = t_n$, and rewritten by using the definitions of ξ_u^n and $\bar{\xi}_u^n$. \square

Theorem 6.3 (Error equation for eq. (39)). *There holds*

$$\begin{aligned} \sum_{K \in \mathcal{T}^s} \int_K (d_t u_h^n - \partial_t u^n) \cdot v_h \, dx + a_h(c_h^{n-1}; \xi_u^n, v_h) \\ + \sum_{j=s,d} (b_h^j(\xi_p^{jn}, v_h) + b_h^{I,j}(\bar{\xi}_p^{jn}, \bar{v}_h)) + \sum_{j=s,d} (b_h^j(q_h^j, \zeta_u^n) + b_h^{I,j}(\bar{q}_h^j, \bar{\zeta}_u^n)) \\ = a_h(c_h^{n-1}; \xi_u^n, v_h) + a_h(c^n; u^n, v_h) - a_h(c_h^{n-1}; u^n, v_h) \\ + \sum_{K \in \mathcal{T}^s} \int_K [f^s(c_h^{n-1}) - f^s(c^n)] \cdot v_h \, dx + \sum_{K \in \mathcal{T}^d} \int_K [\mathbb{K}^{-1}(c_h^{n-1}) f^d(c_h^{n-1}) - \mathbb{K}^{-1}(c^n) f^d(c^n)] \cdot v_h \, dx. \quad (48) \end{aligned}$$

Proof. By Lemma 3.1 at time $t = t^n$, we have that for all $(v_h, q_h) \in \mathbf{X}_h$,

$$\begin{aligned} \sum_{K \in \mathcal{T}^s} \int_K \partial_t u^n \cdot v_h \, dx + B_h^{sd}(c^n; (u^n, p^n), (v_h, q_h)) &= \sum_{K \in \mathcal{T}^s} \int_K f^s(c^n) \cdot v_h \, dx \\ &\quad + \sum_{K \in \mathcal{T}^d} \int_K \mathbb{K}^{-1}(c^n) f^d(c^n) \cdot v_h \, dx + \sum_{K \in \mathcal{T}^d} \int_K (g_p^n - g_i^n) q_h \, dx. \end{aligned}$$

Subtracting this equation from eq. (39), we obtain

$$\begin{aligned} \sum_{K \in \mathcal{T}^s} \int_K (d_t u_h^n - \partial_t u^n) \cdot v_h \, dx + B_h^{sd}(c_h^{n-1}; (u_h^n, p_h^n), (v_h, q_h)) - B_h^{sd}(c^n; (u^n, p^n), (v_h, q_h)) \\ = \sum_{K \in \mathcal{T}^s} \int_K [f^s(c_h^{n-1}) - f^s(c^n)] \cdot v_h \, dx + \sum_{K \in \mathcal{T}^d} \int_K [\mathbb{K}^{-1}(c_h^{n-1}) f^d(c_h^{n-1}) - \mathbb{K}^{-1}(c^n) f^d(c^n)] \cdot v_h \, dx, \quad (49) \end{aligned}$$

for all $(\mathbf{v}_h, \mathbf{q}_h) \in \mathbf{X}_h$. Then, by eq. (43), the B_h^{sd} terms can be rewritten as

$$\begin{aligned}
& B_h^{sd}(\mathbf{c}_h^{n-1}; (\mathbf{u}_h^n, \mathbf{p}_h^n), (\mathbf{v}_h, \mathbf{q}_h)) - B_h^{sd}(\mathbf{c}^n; (\mathbf{u}^n, \mathbf{p}^n), (\mathbf{v}_h, \mathbf{q}_h)) \\
&= B_h^{sd}(\mathbf{c}_h^{n-1}; (\mathbf{u}_h^n - \mathbf{u}^n, \mathbf{p}_h^n - \mathbf{p}^n), (\mathbf{v}_h, \mathbf{q}_h)) \\
&\quad + B_h^{sd}(\mathbf{c}_h^{n-1}; (\mathbf{u}^n, \mathbf{p}^n), (\mathbf{v}_h, \mathbf{q}_h)) - B_h^{sd}(\mathbf{c}^n; (\mathbf{u}^n, \mathbf{p}^n), (\mathbf{v}_h, \mathbf{q}_h)) \\
&= a_h(\mathbf{c}_h^{n-1}; \boldsymbol{\zeta}_u^n, \mathbf{v}_h) - a_h(\mathbf{c}_h^{n-1}; \boldsymbol{\xi}_u^n, \mathbf{v}_h) + a_h(\mathbf{c}_h^{n-1}; \mathbf{u}^n, \mathbf{v}_h) - a_h(\mathbf{c}^n; \mathbf{u}^n, \mathbf{v}_h) \\
&\quad + \sum_{j=s,d} (b_h^j(\boldsymbol{\zeta}_p^{jn}, \mathbf{v}_h) + b_h^{I,j}(\bar{\boldsymbol{\zeta}}_p^{jn}, \bar{\mathbf{v}}_h)) + \sum_{j=s,d} (b_h^j(\mathbf{q}_h^j, \boldsymbol{\zeta}_u^n) + b_h^{I,j}(\bar{\mathbf{q}}_h^j, \bar{\boldsymbol{\zeta}}_u^n)),
\end{aligned} \tag{50}$$

where we applied eqs. (46) and (47). Combining this with eq. (49) yields the result. \square

The velocity error at each time step depends on the error in concentration from the previous time step. Therefore, for the velocity error estimates, we will need some auxiliary results related to the concentration error. To estimate the error of the concentration, we use the continuous interpolant $\mathcal{I}c \in C_h \cap \mathcal{C}^0(\bar{\Omega})$ of c [8], and we set $\bar{\mathcal{I}}c(t) = \mathcal{I}c|_{\Gamma^0}(t) \in \bar{C}_h$. Denoting the restriction of c to Γ^0 by \bar{c} , we define

$$\begin{aligned}
\xi_c^n &= c^n - \mathcal{I}c^n, & \zeta_c^n &= c_h^n - \mathcal{I}c^n, & \boldsymbol{\xi}_c^n &= (\xi_c^n, \bar{\xi}_c^n), \\
\bar{\xi}_c^n &= \bar{c}^n - \bar{\mathcal{I}}c^n, & \bar{\zeta}_c^n &= \bar{c}_h^n - \bar{\mathcal{I}}c^n, & \boldsymbol{\zeta}_c^n &= (\zeta_c^n, \bar{\zeta}_c^n).
\end{aligned} \tag{51}$$

Note that:

$$c^n - c_h^n = \xi_c^n - \zeta_c^n, \quad \bar{c}^n - \bar{c}_h^n = \bar{\xi}_c^n - \bar{\zeta}_c^n. \tag{52}$$

Furthermore, we have the following interpolation estimate [8, Section 4.4] for $r = 0, 1$ and $1 \leq \ell \leq k_c$:

$$\|\xi_c\|_{r,K} \leq Ch_K^{\ell+1-r} \|c\|_{\ell+1,K}, \tag{53}$$

Theorem 6.4. *Let $c_0^s \in H^{k_c+1}(\Omega^s)$, $c^s \in L^2(0, T; H^{k_c+1}(\Omega^s))$, $c^d \in H^1(0, T; L^2(\Omega^d))$ such that $\partial_t c^s \in L^2(0, T, H^1(\Omega^s))$, and $\bar{c} = \gamma(c)$ on Γ_0 . Then we have the following estimates:*

$$\Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}^s} h_K^2 \|c^m - c_h^{m-1}\|_{1,K}^2 \leq C \left(h^2 (\Delta t)^2 \|\partial_t c\|_{L^2(0,T;H^1(\Omega^s))}^2 + \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}^s} h_K^2 \|c^{m-1} - c_h^{m-1}\|_{1,K}^2 \right), \tag{54a}$$

$$\begin{aligned}
\Delta t \sum_{m=1}^n \|\bar{c}^m - \bar{c}_h^{m-1}\|_{\Gamma^I}^2 &\leq C \left(\Delta t \sum_{m=1}^n \|\boldsymbol{\zeta}_c^{m-1}\|_c^2 \right. \\
&\quad \left. + (\Delta t)^2 \|\partial_t c\|_{L^2(0,T;H^1(\Omega^s))}^2 + h^{2k_c+1} (\Delta t \|c_0\|_{k_c+1,\Omega^s}^2 + \|c\|_{\ell^2(0,T;H^{k_c+1}(\Omega^s))}^2) \right),
\end{aligned} \tag{54b}$$

and

$$\Delta t \sum_{m=1}^n \left(\sum_{K \in \mathcal{T}} (\|c^m - c_h^{m-1}\|_K^2) \right) \leq C \left((\Delta t)^2 \|\partial_t c\|_{L^2(0,T;L^2(\Omega))}^2 + \Delta t \sum_{m=1}^n \left(\sum_{K \in \mathcal{T}} \|c^{m-1} - c_h^{m-1}\|_K^2 \right) \right). \tag{54c}$$

Proof. We first prove eq. (54a). By the triangle inequality and the definition of d_t ,

$$\sum_{K \in \mathcal{T}^s} h_K^2 \|c^m - c_h^{m-1}\|_{1,K}^2 \leq \sum_{K \in \mathcal{T}^s} 2h_K^2 \left((\Delta t)^2 \|d_t c^m\|_{1,K}^2 + \|c^{m-1} - c_h^{m-1}\|_{1,K}^2 \right).$$

Multiplying this by Δt , summing from $m = 1$ to n , and using eq. (41b), we obtain eq. (54a). We now prove eq. (54b). We have

$$\|\bar{c}^m - \bar{c}_h^{m-1}\|_{\Gamma_I}^2 \leq 2(\|\bar{c}^m - \bar{c}^{m-1}\|_{\Gamma_I}^2 + \|\bar{c}^{m-1} - \bar{c}_h^{m-1}\|_{\Gamma_I}^2). \quad (55)$$

Since $\bar{c} = c^s|_{\Gamma_I}$ on Γ_I , and $(c^s)^m - (c^s)^{m-1} \in H^1(\Omega^s)$, by eq. (24), the first term on the right side of eq. (55) is bounded as follows:

$$\|\bar{c}^m - \bar{c}^{m-1}\|_{\Gamma_I}^2 \leq C\|c^m - c^{m-1}\|_{1,\Omega^s}^2 = C(\Delta t)^2 \|d_t c^m\|_{1,\Omega^s}^2. \quad (56)$$

Splitting the second term on the right side of eq. (55) by using $\mathcal{I}c = \bar{\mathcal{I}}c$ for any $F \in \mathcal{F}^I$ gives:

$$\|\bar{c}^{m-1} - \bar{c}_h^{m-1}\|_{\Gamma_I}^2 \leq 2(\|c^{m-1} - \mathcal{I}c^{m-1}\|_{\Gamma_I}^2 + \|\bar{c}_h^{m-1} - \bar{\mathcal{I}}c^{m-1}\|_{\Gamma_I}^2) = 2(\|\xi_c^{m-1}\|_{\Gamma_I}^2 + \|\bar{\zeta}_c^{m-1}\|_{\Gamma_I}^2). \quad (57)$$

The first term on the right hand side of eq. (57) is bounded by eq. (21) and eq. (53) as follows:

$$\|\xi_c^{m-1}\|_{\Gamma_I}^2 \leq \sum_{K \in \mathcal{T}^s} C(h_K^{-1} \|\xi_c^{m-1}\|_K^2 + h_K \|\xi_c^{m-1}\|_{1,K}^2) \leq Ch^{2k_c+1} \|c^{m-1}\|_{k_c+1,\Omega^s}^2. \quad (58)$$

Using eq. (26) and the definition of $\|\cdot\|_c$,

$$\begin{aligned} \|\bar{\zeta}_c^{m-1}\|_{\Gamma_I}^2 &\leq 2(\|\bar{\zeta}_c^{m-1} - (\zeta_c^s)^{m-1}\|_{\Gamma_I}^2 + \|(\zeta_c^s)^{m-1}\|_{\Gamma_I}^2) \\ &\leq 2\left(h \sum_{K \in \mathcal{T}^s} h_K^{-1} \|\bar{\zeta}_c^{m-1} - \zeta_c^{m-1}\|_{\partial K}^2 + \|(\zeta_c^s)^{m-1}\|_{\Gamma_I}^2\right) \leq C\|\zeta_c^{m-1}\|_c^2. \end{aligned} \quad (59)$$

Collecting eqs. (55) to (59), we obtain

$$\|\bar{c}^m - \bar{c}_h^{m-1}\|_{\Gamma_I}^2 \leq C\left((\Delta t)^2 \|d_t c^m\|_{1,\Omega^s}^2 + h^{2k_c+1} \|c^{m-1}\|_{k_c+1,\Omega^s}^2 + \|\zeta_c^{m-1}\|_c^2\right).$$

Equation (54b) now follows after multiplying the above inequality by Δt , summing from 1 to n , and using eq. (41b). We next prove eq. (54c). By the triangle inequality,

$$\sum_{K \in \mathcal{T}} \|c^m - c_h^{m-1}\|_K^2 \leq \sum_{K \in \mathcal{T}} 2((\Delta t)^2 \|d_t c^m\|_K^2 + \|c^{m-1} - c_h^{m-1}\|_K^2). \quad (60)$$

The results follows by multiplying eq. (60) by Δt , summing from $m = 1$ to n , and using eq. (41b) as before. \square

Now that the auxiliary result is established, we proceed with proving error estimates the velocity.

Theorem 6.5. *Let c and u be the solutions of eqs. (1) to (3) such that*

$$\begin{aligned} u^s &\in L^2(0, T; [H^{k_f+1}(\Omega^s)]^{\dim}) \cap L^\infty(0, T; [W^{1,\infty}(\Omega^s)]^{\dim}), \\ \partial_t u^s &\in L^2(0, T; [H^{k_f}(\Omega^s)]^{\dim}), \partial_{tt} u^s \in L^2(0, T; [L^2(\Omega^s)]^{\dim}), \\ u^d &\in L^2(0, T; [H^{k_f}(\Omega^d)]^{\dim}) \cap L^\infty(0, T; [L^\infty(\Omega^d)]^{\dim}), \\ p &\in L^2(0, T; L^2(\Omega)), \nabla p^d \in L^\infty(0, T; [L^\infty(\Omega^d)]^{\dim}), \\ c^s &\in L^2(0, T; H^{k_c+1}(\Omega^s)), \partial_t c^s \in L^2(0, T; H^1(\Omega^s)), \\ c^d &\in L^2(0, T; L^2(\Omega^d)), \partial_t c^d \in L^\infty(0, T; L^2(\Omega^d)), \\ u_0 &\in [H_0^1(\Omega^s)]^{\dim}, \nabla \cdot u^0 = 0, c_0^s \in H^{k_c+1}(\Omega^s), \end{aligned}$$

and let $k_f, k_c \geq \dim - 1$, and $n \geq 1$. Suppose that $u_h^0, \dots, u_h^{n-1} \in V_h$ and $\mathbf{c}_h^0, \dots, \mathbf{c}_h^{n-1} \in \mathbf{C}_h$, the solutions of eq. (39) and eq. (40), respectively, are known and satisfy for $1 \leq i \leq n$,

$$\|\mathbf{c}_h^{i-1} - \mathbf{c}^{i-1}\|_\Omega^2 + \Delta t \sum_{m=1}^i \|\mathbf{c}_h^{m-1} - \mathbf{c}^{m-1}\|_c^2 \leq C((\Delta t)^2 + h^{2k_f} + h^{2k_c}). \quad (61)$$

Then ζ_u^n satisfies:

$$\|\zeta_u^n\|_{\Omega^s}^2 + (\Delta t)^2 \sum_{m=1}^n \|d_t \zeta_u^m\|_{\Omega^s}^2 + \Delta t \sum_{m=1}^n \|\zeta_u^m\|_v^2 \leq C((\Delta t)^2 + h^{2k_f} + h^{2k_c}), \quad (62a)$$

$$\|\zeta_u^n\|_{\Omega^d}^2 \leq C''((\Delta t)^2 + h^{2k_f} + h^{2k_c}), \quad (62b)$$

where the constants depend on $\mu^*, \kappa_*, \mu_L, L_f^s, L_f^d, \sum_{j=1}^{\dim-1} \gamma^j$, and the regularity of u_0, u, c_0 , and c but are independent of the mesh size.

Proof. Proof of eq. (62a):

Setting $(\mathbf{v}_h, \mathbf{q}_h) = (\zeta_u^n, -\zeta_p^n)$ in Theorem 6.3, using $a(a-b) = \frac{1}{2}(a^2 - b^2 + (a-b)^2)$, and the coercivity of a_h eq. (32) yields

$$\begin{aligned} & \frac{1}{2\Delta t} (\|\zeta_u^n\|_{\Omega^s}^2 - \|\zeta_u^{n-1}\|_{\Omega^s}^2) + \frac{\Delta t}{2} \|d_t \zeta_u^n\|_{\Omega^s}^2 + C_a \|\zeta_u^n\|_v^2 \\ &= \sum_{K \in \mathcal{T}^s} \int_K (\partial_t u^n - d_t \Pi_V u^n) \cdot \zeta_u^n \, dx + a_h(\mathbf{c}_h^{n-1}; \xi_u^n, \zeta_u^n) + [a_h(\mathbf{c}^n; \mathbf{u}^n, \zeta_u^n) - a_h(\mathbf{c}_h^{n-1}; \mathbf{u}^n, \zeta_u^n)] \\ &+ \int_{\Omega^s} [f^s(c_h^{n-1}) - f^s(c^n)] \cdot \zeta_u^n \, dx + \int_{\Omega^d} [\mathbb{K}^{-1}(c_h^{n-1}) f^d(c_h^{n-1}) - \mathbb{K}^{-1}(c^n) f^d(c^n)] \cdot \zeta_u^n \, dx \\ &=: I_1 + \dots + I_5. \end{aligned} \quad (63)$$

Using eq. (29) and employing Young's inequality for some $\epsilon > 0$,

$$I_1 \leq (\|\partial_t u^n - d_t u^n\|_{\Omega^s} + \|d_t \xi_u^n\|_{\Omega^s}) \|\zeta_u^n\|_{\Omega^s} \leq C(\|\partial_t u^n - d_t u^n\|_{\Omega^s}^2 + \|d_t \xi_u^n\|_{\Omega^s}^2) + \epsilon \|\zeta_u^n\|_v^2. \quad (64)$$

It follows from eq. (31) and Young's inequality that

$$I_2 \leq C \|\xi_u^n\|_{v'} \|\zeta_u^n\|_v \leq C \|\xi_u^n\|_{v'}^2 + \epsilon \|\zeta_u^n\|_v^2. \quad (65)$$

Consider now I_3 :

$$\begin{aligned} I_3 &= \sum_{K \in \mathcal{T}^s} \int_K 2[\mu(c^n) - \mu(c_h^{n-1})] \varepsilon(u^n) : \varepsilon(\zeta_u^n) \, dx - \sum_{K \in \mathcal{T}^s} \int_{\partial K} 2[\mu(c^n) - \mu(c_h^{n-1})] \varepsilon(u^n) n^s \cdot (\zeta_u^n - \bar{\zeta}_u^n) \, ds \\ &+ \int_{\Omega^d} [\mathbb{K}^{-1}(c^n) - \mathbb{K}^{-1}(c_h^{n-1})] u^n \cdot \zeta_u^n \, dx + \sum_{j=1}^{\dim-1} \int_{\Gamma^j} \gamma^j [\mu(c^n) - \mu(\bar{c}_h^{n-1})] (u^{sn} \cdot \tau^j) (\bar{\zeta}_u^n \cdot \tau^j) \, ds \\ &=: I_{31} + \dots + I_{34}. \end{aligned} \quad (66)$$

The first term on the right side of eq. (66) can be bounded as follows using Lipschitz continuity of μ , the generalized Hölder's inequality for integrals and sums, and Young's inequality:

$$\begin{aligned} I_{31} &\leq 2\mu_L \sum_{K \in \mathcal{T}^s} \|c^n - c_h^{n-1}\|_K \|\varepsilon(u^n)\|_{0,\infty,K} \|\nabla \zeta_u^n\|_K \leq 2\mu_L \|\nabla u^n\|_{0,\infty,\Omega^s} \left(\sum_{K \in \mathcal{T}^s} \|c^n - c_h^{n-1}\|_K^2 \right)^{1/2} \|\zeta_u^n\|_v \\ &\leq C\mu_L^2 \|\nabla u^n\|_{0,\infty,\Omega^s}^2 \sum_{K \in \mathcal{T}^s} \|c^n - c_h^{n-1}\|_K^2 + \epsilon \|\zeta_u^n\|_v^2. \end{aligned} \quad (67)$$

Next we bound I_{32} . By Lipschitz continuity of μ , eq. (21), eq. (34d), generalized Hölder's inequality, and Young's inequality,

$$\begin{aligned} |I_{32}| &\leq 2\mu_L \sum_{K \in \mathcal{T}^s} \|c^n - c_h^{n-1}\|_{\partial K} \|\nabla u^n\|_{0,\infty,K} \|\zeta_u^n - \bar{\zeta}_u^n\|_{\partial K} \\ &\leq 2\mu_L \|\nabla u^n\|_{0,\infty,\Omega^s} \left(\sum_{K \in \mathcal{T}^s} h_K \|c^n - c_h^{n-1}\|_{\partial K}^2 \right)^{1/2} \left(\sum_{K \in \mathcal{T}^s} h_K^{-1} \|\zeta_u^n - \bar{\zeta}_u^n\|_{\partial K}^2 \right)^{1/2} \\ &\leq 2C\mu_L \|\nabla u^n\|_{0,\infty,\Omega^s} \left(\sum_{K \in \mathcal{T}^s} (\|c^n - c_h^{n-1}\|_K^2 + h_K^2 \|c^n - c_h^{n-1}\|_{1,K}^2) \right)^{1/2} \|\zeta_u^n\|_v \\ &\leq C\mu_L^2 \|\nabla u^n\|_{0,\infty,\Omega^s}^2 \sum_{K \in \mathcal{T}^s} (\|c^n - c_h^{n-1}\|_K^2 + h_K^2 \|c^n - c_h^{n-1}\|_{1,K}^2) + \epsilon \|\zeta_u^n\|_v^2. \end{aligned} \quad (68)$$

We bound I_{33} by using the assumption on \mathbb{K} given in eq. (6):

$$I_{33} \leq \frac{\mu_L}{\kappa_*} \|c^n - c_h^{n-1}\|_{\Omega^d} \|u^n\|_{0,\infty,\Omega^d} \|\zeta_u^n\|_v \leq C(\kappa_*^{-1} \mu_L)^2 \|u^n\|_{0,\infty,\Omega^d}^2 \|c^n - c_h^{n-1}\|_{\Omega^d}^2 + \epsilon \|\zeta_u^n\|_v^2. \quad (69)$$

Again by the Lipschitz property of μ and Hölder's inequality,

$$\begin{aligned} I_{34} &\leq \mu_L \|c^n - \bar{c}_h^{n-1}\|_{\Gamma^I} \left(\sum_{j=1}^{\dim-1} \gamma^j \right)^{1/2} \|u^{sn}\|_{0,\infty,\Gamma^I} \left(\sum_{j=1}^{\dim-1} \gamma^j \|\bar{\zeta}_u^n \cdot \tau^j\|_{\Gamma^I}^2 \right)^{1/2} \\ &\leq \mu_L \|c^n - \bar{c}_h^{n-1}\|_{\Gamma^I} \left(\sum_{j=1}^{\dim-1} \gamma^j \right)^{1/2} \|u^n\|_{0,\infty,\Omega^s} \|\zeta_u^n\|_v \\ &\leq C\mu_L^2 \left(\sum_{j=1}^{\dim-1} \gamma^j \right) \|u^n\|_{0,\infty,\Omega^s}^2 \|c^n - \bar{c}_h^{n-1}\|_{\Gamma^I}^2 + \epsilon \|\zeta_u^n\|_v^2. \end{aligned} \quad (70)$$

Combining eqs. (66) to (70),

$$\begin{aligned} I_3 &\leq C\mu_L^2 \|\nabla u^n\|_{0,\infty,\Omega^s}^2 \sum_{K \in \mathcal{T}^s} (\|c^n - c_h^{n-1}\|_K^2 + h_K^2 \|c^n - c_h^{n-1}\|_{1,K}^2) + C(\kappa_*^{-1} \mu_L)^2 \|u^n\|_{0,\infty,\Omega^d}^2 \|c^n - c_h^{n-1}\|_{\Omega^d}^2 \\ &\quad + C\mu_L^2 \left(\sum_{j=1}^{\dim-1} \gamma^j \right) \|u^n\|_{0,\infty,\Omega^s}^2 \|c^n - \bar{c}_h^{n-1}\|_{\Gamma^I}^2 + 4\epsilon \|\zeta_u^n\|_v^2. \end{aligned} \quad (71)$$

Since f^s is Lipschitz continuous in c , with Lipschitz constant L_f^s , and recalling eq. (29),

$$I_4 \leq CL_f^s \|c_h^{n-1} - c^n\|_{\Omega^s} \|\zeta_u^n\|_v \leq C(L_f^s)^2 \|c_h^{n-1} - c^n\|_{\Omega^s}^2 + \epsilon \|\zeta_u^n\|_v^2. \quad (72)$$

Since f^d and μ are Lipschitz continuous in c , with Lipschitz continuity constants L_f^d and μ_L , respectively,

$$\begin{aligned} I_5 &= \int_{\Omega^d} (\mathbb{K}^{-1}(c_h^{n-1})[f^d(c_h^{n-1}) - f^d(c^n)] + [\mu(c_h^{n-1}) - \mu(c^n)]\kappa^{-1}f^d(c^n)) \cdot \zeta_u^n dx \\ &\leq (K_*^{-1}L_f^d + \mu_L\kappa_*^{-1}\|f^d(c^n)\|_{0,\infty,\Omega^d})\|c_h^{n-1} - c^n\|_{\Omega^d}\|\zeta_u^n\|_{\Omega^d} \\ &\leq C(K_*^{-1}L_f^d + \mu_L\kappa_*^{-1}\|f^d(c^n)\|_{0,\infty,\Omega^d})^2\|c_h^{n-1} - c^n\|_{\Omega^d}^2 + \epsilon\|\zeta_u^n\|_v^2. \end{aligned} \quad (73)$$

Combining the above bounds for I_1 to I_5 with eq. (63), letting $\epsilon = \frac{C_a}{16}$ (C_a is the coercivity constant), multiplying by $2\Delta t$, summing from 1 to n , noting that $\zeta_u^0 = 0$, and applying eq. (41), we obtain:

$$\begin{aligned} &\|\zeta_u^n\|_{\Omega^s}^2 + (\Delta t)^2 \sum_{m=1}^n \|d_t \zeta_u^m\|_{\Omega^s}^2 + C_a \Delta t \sum_{m=1}^n \|\zeta_u^m\|_v^2 \\ &\leq C \left[(\Delta t)^2 \|\partial_{tt} u\|_{L^2(0,T;L^2(\Omega^s))}^2 + \|\partial_t \xi_u\|_{L^2(0,T;L^2(\Omega^s))}^2 + 2\Delta t \sum_{m=1}^n \|\xi_u^m\|_{v'}^2 \right. \\ &\quad + \mu_L^2 \Delta t \sum_{m=1}^n \left(\|\nabla u^m\|_{0,\infty,\Omega^s}^2 \sum_{K \in \mathcal{T}^s} h_K^2 \|c^m - c_h^{m-1}\|_{1,K}^2 \right) \\ &\quad + \Delta t \sum_{m=1}^n \left(\kappa_*^{-1} \mu_L \|u^m\|_{0,\infty,\Omega^d} + K_*^{-1} L_f^d + \frac{\mu_L}{\kappa_*} \|f^d(c^m)\|_{0,\infty,\Omega^d} \right)^2 \|c^m - c_h^{m-1}\|_{\Omega^d}^2 \\ &\quad + \mu_L^2 \left(\sum_{j=1}^{\dim-1} \gamma^j \right) \Delta t \sum_{m=1}^n \|u^m\|_{0,\infty,\Omega^s}^2 \|\bar{c}^m - \bar{c}_h^{m-1}\|_{\Gamma_I}^2 \\ &\quad \left. + \Delta t \sum_{m=1}^n ((L_f^s)^2 + \mu_L^2 \|\nabla u^m\|_{0,\infty,\Omega^s}^2) \|c_h^{m-1} - c^m\|_{\Omega^s}^2 \right]. \end{aligned}$$

Next, using eq. (34d) and eq. (45b),

$$\begin{aligned} &\|\zeta_u^n\|_{\Omega^s}^2 + (\Delta t)^2 \sum_{m=1}^n \|d_t \zeta_u^m\|_{\Omega^s}^2 + C_a \Delta t \sum_{m=1}^n \|\zeta_u^m\|_v^2 \\ &\leq C \left[(\Delta t)^2 \|\partial_{tt} u\|_{L^2(0,T;[L^2(\Omega^s)]^{\dim})}^2 + h^{2k_f} (\|\partial_t u\|_{L^2(0,T;[H^{k_f}(\Omega^s)]^{\dim})}^2 + \|u\|_{\ell^2(0,T;[H^{k_f+1}(\Omega^s)]^{\dim})}^2 \right. \\ &\quad + \|u\|_{\ell^2(0,T;[H^{k_f}(\Omega^d)]^{\dim})}^2) + \|\nabla u\|_{L^\infty(0,T;[L^\infty(\Omega^s)]^{\dim})}^2 \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}^s} (h_K^2 \|c^m - c_h^{m-1}\|_{1,K}^2) \\ &\quad + (\|u\|_{L^\infty(0,T;[L^\infty(\Omega^d)]^{\dim})}^2 + \|\nabla p\|_{L^\infty(0,T;[L^\infty(\Omega^d)]^{\dim})}^2 + 1) \Delta t \sum_{m=1}^n \|c^m - c_h^{m-1}\|_{\Omega^d}^2 \\ &\quad + \|u\|_{L^\infty(0,T;[L^\infty(\Omega^s)]^{\dim})}^2 \Delta t \sum_{m=1}^n \|\bar{c}^m - \bar{c}_h^{m-1}\|_{\Gamma_I}^2 \\ &\quad \left. + (1 + \|\nabla u\|_{L^\infty(0,T;[L^\infty(\Omega^s)]^{\dim})}^2) \Delta t \sum_{m=1}^n \|c^m - c_h^{m-1}\|_{\Omega}^2 \right], \end{aligned}$$

where the constant $C > 0$ depends on $\mu_L, \kappa_*, K^*, \gamma^j, L_f^s$ but is independent of h and Δt . Equation (62a) follows by eqs. (54a) to (54c), and assumption eq. (61).

Proof of eq. (62b):

Let $v_h^s = 0$, $\bar{v}_h = 0$, and $\mathbf{q}_h = \mathbf{0}$ in eq. (48). Then

$$\begin{aligned} a_h^d(c_h^{n-1}; \zeta_u^n, v_h) + b_h^d(\zeta_p^{dn}, v_h) &= a_h^d(c_h^{n-1}; \xi_u^n, v_h) + a_h^d(c^n; u^n, v_h) - a_h^d(c_h^{n-1}; u^n, v_h) \\ &\quad + \sum_{K \in \mathcal{T}^d} \int_K [\mathbb{K}^{-1}(c_h^{n-1}) f^d(c_h^{n-1}) - \mathbb{K}^{-1}(c^n) f^d(c^n)] \cdot v_h \, dx. \end{aligned} \quad (74)$$

On the other hand, letting $\mathbf{v}_h = \mathbf{0}$ and $\mathbf{q}_h^s = \mathbf{0}$ in eq. (48), we have

$$b_h^d(\mathbf{q}_h^d, \zeta_u^n) + b_h^{I,d}(\bar{q}_h^d, \bar{\zeta}_u^n) = 0,$$

implying that

$$b_h^d(\mathbf{q}_h^d, \zeta_u^n) = -b_h^{I,d}(\bar{q}_h^d, \bar{\zeta}_u^n) = \int_{\Gamma^I} \bar{q}_h^d \bar{\zeta}_u^n \cdot n^d \, ds. \quad (75)$$

From [30, Lemma 3.2], since $u^{sn} - u_h^{sn} \in H^{\text{div}}(\Omega^s)$, there exists $w \in H^{\text{div}}(\Omega^d)$ such that $\nabla \cdot w = 0$ in Ω^d , $w \cdot n = 0$ on Γ^d and $w \cdot n^d = (\gamma(u^{sn}) - u_h^{sn}) \cdot n^d$ on Γ^I . With this choice of w , and using eqs. (34a) to (34c) we observe that

$$b_h^d(\mathbf{q}_h^d, \Pi_V w) = \int_{\Gamma^I} \bar{q}_h^d (u^{sn} - u_h^{sn}) \cdot n^d \, ds. \quad (76)$$

Adding eqs. (75) and (76), and recalling eqs. (19b) and (19c), the definition of $\bar{\Pi}_V$, and eq. (34b), we obtain

$$b_h^d(\mathbf{q}_h^d, \zeta_u^n + \Pi_V w) = 0 \quad \forall \mathbf{q}_h^d \in \mathbf{Q}_h^d. \quad (77)$$

This leads us to consider $v_h = \zeta_u^n + \Pi_V w$ as test function in eq. (74). Using eqs. (7) and (77) we find that

$$\begin{aligned} \frac{1}{K_*} \|\zeta_u^n\|_{\Omega^d}^2 &\leq -a_h^d(c_h^{n-1}; \zeta_u^n, \Pi_V w) + a_h^d(c_h^{n-1}; \zeta_u^n, \zeta_u^n + \Pi_V w) \\ &\quad + \left(a_h^d(c^n; u^n, \zeta_u^n + \Pi_V w) - a_h^d(c_h^{n-1}; u^n, \zeta_u^n + \Pi_V w) \right) \\ &\quad + \sum_{K \in \mathcal{T}^d} \int_K [\mathbb{K}^{-1}(c_h^{n-1}) f^d(c_h^{n-1}) - \mathbb{K}^{-1}(c^n) f^d(c^n)] \cdot (\zeta_u^n + \Pi_V w) \, dx =: A_1 + A_2 + A_3 + A_4. \end{aligned} \quad (78)$$

We will bound A_1 to A_4 by a series of Cauchy–Schwarz, Hölder’s, triangle, and Young’s inequalities together with the properties of μ and κ . First,

$$A_1 \leq K_*^{-1} \|\zeta_u^n\|_{\Omega^d} \|\Pi_V w\|_{\Omega^d} \leq \epsilon \|\zeta_u^n\|_{\Omega^d}^2 + CK_*^{-2} \|\Pi_V w\|_{\Omega^d}^2,$$

and

$$A_2 \leq K_*^{-1} \|\xi_u^n\|_{\Omega^d} \|\zeta_u^n + \Pi_V w\|_{\Omega^d} \leq CK_*^{-2} \|\xi_u^n\|_{\Omega^d}^2 + \epsilon \|\zeta_u^n\|_{\Omega^d}^2 + K_*^{-1} \|\xi_u^n\|_{\Omega^d} \|\Pi_V w\|_{\Omega^d}.$$

Following the proof of eq. (69),

$$\begin{aligned} A_3 &\leq \mu_L \kappa_*^{-1} \|c^n - c_h^{n-1}\|_{\Omega^d} \|u^n\|_{0,\infty,\Omega^d} (\|\zeta_u^n\|_{\Omega^d} + \|\Pi_V w\|_{\Omega^d}) \\ &\leq C \mu_L^2 \kappa_*^{-2} \|c^n - c_h^{n-1}\|_{\Omega^d}^2 \|u^n\|_{0,\infty,\Omega^d}^2 + \epsilon \|\zeta_u^n\|_{\Omega^d}^2 + \mu_L \kappa_*^{-1} \|c^n - c_h^{n-1}\|_{\Omega^d} \|u^n\|_{0,\infty,\Omega^d} \|\Pi_V w\|_{\Omega^d}, \end{aligned}$$

while as the proof of eq. (73),

$$\begin{aligned} A_4 &\leq (\mu_L \kappa_*^{-1} \|f^d(c^n)\|_{0,\infty,\Omega^d} + K_*^{-1} L_f^d) \|c_h^{n-1} - c^n\|_{\Omega^d} (\|\zeta_u^n\|_{\Omega^d} + \|\Pi_V w\|_{\Omega^d}) \\ &\leq C (\mu_L \kappa_*^{-1} \|f^d(c^n)\|_{0,\infty,\Omega^d} + K_*^{-1} L_f^d)^2 \|c_h^{n-1} - c^n\|_{\Omega^d}^2 + \epsilon \|\zeta_u^n\|_{\Omega^d}^2 \\ &\quad + (\mu_L \kappa_*^{-1} \|f^d(c^n)\|_{0,\infty,\Omega^d} + K_*^{-1} L_f^d) \|c_h^{n-1} - c^n\|_{\Omega^d} \|\Pi_V w\|_{\Omega^d}. \end{aligned}$$

Combining the bounds of A_1 to A_4 with eq. (78), choosing $\epsilon = 1/(8K^*)$, and applying another set of Young's inequalities, we obtain

$$\|\zeta_u^n\|_{\Omega^d}^2 \leq C (\|\Pi_V w\|_{\Omega^d}^2 + \|\zeta_u^n\|_{\Omega^d}^2 + \|c^n - c_h^{n-1}\|_{\Omega^d}^2), \quad (79)$$

where C depends on the problem parameters κ_* , K_* , K^* , μ_L , L_f^d , and the regularity of u , p and c . Noting that from eq. (34d) and [30, Theorem 3.3], [17, Lemma 10],

$$\|\Pi_V w\|_{\Omega^d} \leq C \|w\|_{\Omega^d} \leq Ch^{k_f} (\|u^n\|_{k_f+1,\Omega^s} + \|u\|_{k_f,\Omega^d} + \|g_p - g_i\|_{k_f,\Omega^d}). \quad (80)$$

Then eqs. (34d), (60), (79) and (80) imply

$$\|\zeta_u^n\|_{\Omega^d}^2 \leq Ch^{2k_f} (\|u^n\|_{k_f+1,\Omega^s}^2 + \|u^n\|_{k_f,\Omega^d}^2 + \|g_p - g_i\|_{k_f,\Omega^d}^2 + (\Delta t)^2 \|d_t c^n\|_{\Omega^d}^2 + \|c^{n-1} - c_h^{n-1}\|_{\Omega^d}^2).$$

Equation (62b) is now a consequence of the assumptions on the regularity of the exact solution and eq. (61). \square

The following is a straightforward consequence of Theorem 6.5.

Corollary 6.6. *Let u^n and u_h^n be as defined in Theorem 6.5. Then for all $n \geq 1$,*

$$\|u^n - u_h^n\|_{\Omega} \leq C(\Delta t + h^{k_f} + h^{k_c}), \quad (81a)$$

$$\sum_{K \in \mathcal{T}_h} h_K \|u^n - u_h^n\|_{\partial K}^2 \leq C((\Delta t)^2 + h^{2k_f} + h^{2k_c}). \quad (81b)$$

Before moving on to the next section, we note another consequence of eq. (62) that will prove useful in analysis later on.

Corollary 6.7. *Let u denote the velocity solution to eqs. (1) to (3) satisfying the assumptions in Theorem 6.5 with $k_f, k_c \geq \dim - 1$. Suppose $\Delta t \leq Ch_{k_f}^{\dim/2}$. Then for each $n \geq 1$, the discrete velocity u_h^n that solves eq. (39) satisfies*

$$\|u_h^n\|_{0,\infty,\Omega} \leq C, \quad (82)$$

where C depends on μ^* , κ_* , μ_L , L_f^s , L_f^d , $\sum_{j=1}^{\dim-1} \gamma^j$, and the regularity of u_0 , u , c_0 , and c , but is independent of h , n and Δt .

Proof. The proof is the same as in [16, Lemma 1]. Using eqs. (62a) and (62b), we have

$$\|\zeta_u^n\|_{\Omega}^2 \leq C((\Delta t)^2 + h^{2k_f} + h^{2k_c}). \quad (83)$$

Combining the above estimate with eqs. (28) and (34d), for each $K \in \mathcal{T}$, we have

$$\begin{aligned} \|u_h^n\|_{0,\infty,K} &\leq \|\zeta_u^n\|_{0,\infty,K} + \|\Pi_V u^n\|_{0,\infty,K} \leq Ch_K^{-\dim/2} \|\zeta_u^n\|_K + \|\Pi_V u^n\|_{0,\infty,K} \\ &\leq Ch_K^{-\dim/2} ((\Delta t) + h^{k_f} + h^{k_c}) + \|u^n\|_{0,\infty,K} \\ &\leq Ch_K^{-\dim/2} ((\Delta t) + h^{k_f} + h^{k_c}) + \|u\|_{L^\infty(0,T;[L^\infty(K)]^{\dim})}. \end{aligned}$$

The result eq. (82) follows by using the assumption on Δt and by taking the maximum over $K \in \mathcal{T}$. \square

Remark 6.8. The restrictions on the polynomial degree and the time step are not necessary if we assume that $|D(u)| \leq \bar{D}$ for some positive constant \bar{D} as in [43, 2.12]. It is also possible to avoid these restrictions by using an approach involving a cutoff operator on the velocity solution, as in [49, 50], if one is interested in lower order approximations.

Remark 6.9. Compatibility, as defined in [22], can be achieved by choosing $k_c = k_f - 1$ [16]. However, the requirement that $k_c \geq \dim - 1$ implies $k_f \geq \dim$. Therefore, when $\dim = 2$, our theory supports compatibility only for $k_f \geq 2$ and for $k_f \geq 3$ when $\dim = 3$.

6.2. Error estimate for the pressure

In this section, we briefly discuss the a priori error estimate for the pressure approximation.

Lemma 6.10. *Suppose that the assumptions in Theorem 6.5 hold and that p and \mathbf{p}_h are the pressure solutions to eqs. (1) to (3) and eq. (39), respectively. Then*

$$\begin{aligned} \Delta t \sum_{m=1}^n \|\zeta_p^m\|_p^2 &\leq C \left(\Delta t \sum_{m=1}^n \|\zeta_u^m\|_v^2 + \Delta t \sum_{m=1}^n \|d_t \zeta_u^m\|_{\Omega^s}^2 + \Delta t \sum_{m=1}^n \|\zeta_c^{m-1}\|_c^2 \right. \\ &\quad \left. + \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}} \|c^{m-1} - c_h^{m-1}\|_K^2 + \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}^s} h_K^2 \|c^{m-1} - c_h^{m-1}\|_{1,K}^2 + (\Delta t)^2 + h^{2k_f} + h^{2k_c+1} \right). \end{aligned} \quad (84)$$

Proof. Setting $\mathbf{q}_h = 0$ in the error equation in Theorem 6.3, we obtain:

$$\begin{aligned} \sum_{j=s,d} (b_h^j(\zeta_p^{jn}, v_h) + b_h^{I,j}(\bar{\zeta}_p^{jn}, \bar{v}_h)) \\ = (a_h(c_h^{n-1}; \xi_u^n, v_h) - a_h(c_h^{n-1}; \zeta_u^n, v_h)) + (a_h(c^n; \mathbf{u}^n, v_h) - a_h(c_h^{n-1}; \mathbf{u}^n, v_h)) \\ + \int_{\Omega^s} [f_s(c_h^{n-1}) - f_s(c^n)] \cdot v_h \, dx + \int_{\Omega^d} [\mathbb{K}^{-1}(c_h^{n-1}) f^d(c_h^{n-1}) - \mathbb{K}^{-1}(c^n) f^d(c^n)] \cdot v_h \, dx \\ - \sum_{K \in \mathcal{T}^s} \int_K (d_t u_h^n - \partial_t u^n) \cdot v_h \, dx \\ = : H_1 + \dots + H_5. \end{aligned} \quad (85)$$

By eq. (31) and Young's inequality, and using that $\|\cdot\|_v$ and $\|\cdot\|_{v'}$ are equivalent on \mathbf{V}_h , we have

$$H_1 \leq C(\|\zeta_u^n\|_v + \|\xi_u^n\|_{v'}) \|\mathbf{v}_h\|_v. \quad (86)$$

Following the proof of eq. (71), we can show that

$$\begin{aligned} H_2 &\leq C \left(\mu_L \|\nabla u^n\|_{0,\infty,\Omega^s} \left(\sum_{K \in \mathcal{T}^s} (\|c^n - c_h^{n-1}\|_K^2 + h_K^2 \|c^n - c_h^{n-1}\|_{1,K}^2) \right)^{1/2} \right. \\ &\quad \left. + \kappa_*^{-1} \mu_L \|u^n\|_{0,\infty,\Omega^d} \|c^n - c_h^{n-1}\|_{\Omega^d} + \mu_L \left(\sum_{j=1}^{\dim-1} \gamma^j \right)^{1/2} \|u^n\|_{0,\infty,\Omega^s} \|c^n - \bar{c}_h^{n-1}\|_{\Gamma^I} \right) \|\mathbf{v}_h\|_v. \end{aligned} \quad (87)$$

As in eqs. (72) and (73), we find

$$H_3 \leq L_f^s \|c_h^{n-1} - c^n\|_{\Omega^s} \|\mathbf{v}_h\|_v, \quad (88)$$

$$H_4 \leq (\mu_L \kappa_*^{-1} \|f^d(c^n)\|_{0,\infty,\Omega^d} + K_*^{-1} L_f^d) \|c_h^{n-1} - c^n\|_{\Omega^d} \|\mathbf{v}_h\|_v. \quad (89)$$

Using Cauchy–Schwarz and triangle inequalities, and eq. (34d),

$$\begin{aligned} H_5 &= - \sum_{K \in \mathcal{T}^s} \int_K (d_t \zeta_u^n + (d_t \Pi_V u^n - \partial_t \Pi_V u^n) - \partial_t \xi_u^n) \cdot v_h \, dx \\ &\leq (\|d_t \zeta_u^n\|_{\Omega^s} + \|d_t \Pi_V u^n - \partial_t \Pi_V u^n\|_{\Omega^s} + Ch^{k_f} \|\partial_t u^n\|_{k_f, \Omega^s}) \|\mathbf{v}_h\|_v. \end{aligned} \quad (90)$$

Therefore, combining (85)–(90), dividing both sides by $\|\mathbf{v}_h\|_v$, taking the supremum over $\mathbf{v}_h \in \mathbf{V}_h$, and using Theorem 4.2, we obtain

$$\begin{aligned} c_{\inf}^* \|\zeta_p^n\|_p &\leq C \left(\|\zeta_u^n\|_v + \|\xi_u^n\|_{v'} \right. \\ &\quad + \mu_L \|\nabla u^n\|_{0, \infty, \Omega^s} \left(\sum_{K \in \mathcal{T}^s} (\|c^n - c_h^{n-1}\|_K^2 + h_K^2 \|c^n - c_h^{n-1}\|_{1,K}^2) \right)^{1/2} \\ &\quad + \mu_L \left(\sum_{j=1}^{\dim-1} \gamma^j \right)^{1/2} \|u^n\|_{0, \infty, \Omega^s} \|c^n - \bar{c}_h^{n-1}\|_{\Gamma_I} + L_f^s \|c_h^{n-1} - c^n\|_{\Omega^s} \\ &\quad + (\mu_L \kappa_*^{-1} (\|f^d(c^n)\|_{0, \infty, \Omega^d} + \|u^n\|_{0, \infty, \Omega^d}) + K_*^{-1} L_f^d) \|c_h^{n-1} - c^n\|_{\Omega^d} \\ &\quad \left. + \|d_t \zeta_u^n\|_{\Omega^s} + \|d_t \Pi_V u^n - \partial_t \Pi_V u^n\|_{\Omega^s} + Ch^{k_f} \|\partial_t u^n\|_{k_f, \Omega^s} \right). \end{aligned}$$

Squaring both sides, multiplying by $(c_{\inf}^*)^{-2} \Delta t$, summing from 1 to n , using eqs. (41a) and (45b), stability of Π_V , and the regularity assumptions on u^s , u^d , and ∇p^d yields

$$\begin{aligned} \Delta t \sum_{m=1}^n \|\zeta_p^m\|_p^2 &\leq C \left(\Delta t \sum_{m=1}^n \|\zeta_u^m\|_v^2 + \Delta t \sum_{m=1}^n \|d_t \zeta_u^m\|_{\Omega^s}^2 \right. \\ &\quad + \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}} \|c^m - c_h^{m-1}\|_K^2 + \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}^s} h_K^2 \|c^m - c_h^{m-1}\|_{1,K}^2 \\ &\quad + \Delta t \sum_{m=1}^n \|c^m - \bar{c}_h^{m-1}\|_{\Gamma_I}^2 + (\Delta t)^2 \|\partial_{tt} u\|_{L^2(0, T; [L^2(\Omega^s)]^{\dim})}^2 \\ &\quad \left. + h^{2k_f} (\|u\|_{\ell^2(0, T; [H^{k_f+1}(\Omega^s)]^{\dim})}^2 + \|u\|_{L^2(0, T; [H^{k_f}(\Omega^d)]^{\dim})}^2 + \|\partial_t u\|_{\ell^2(0, T; [H^{k_f}(\Omega^s)]^{\dim})}^2) \right). \end{aligned}$$

Therefore, the result follows by eqs. (54a) to (54c) under the assumptions on the exact solution given in Theorem 6.5. \square

Remark 6.11. An immediate consequence Lemma 6.10 and Theorem 6.5 is

$$\begin{aligned} \Delta t \sum_{m=1}^n \|\zeta_p^m\|_p^2 &\leq C \left(\Delta t + (\Delta t)^{-1} (h^{2k_f} + h^{2k_c}) + \Delta t \sum_{m=1}^n \|\zeta_c^{m-1}\|_c^2 \right. \\ &\quad \left. + \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}} \|c^{m-1} - c_h^{m-1}\|_K^2 + \Delta t \sum_{m=1}^n \sum_{K \in \mathcal{T}^s} h_K^2 \|c^{m-1} - c_h^{m-1}\|_{1,K}^2 \right). \end{aligned}$$

This loss of Δt is due to $\Delta t \sum_{m=1}^n \|d_t \zeta_u^m\|_{\Omega^s}^2$ in eq. (84). However, an improved estimate can be obtained by bounding this term as follows: testing eq. (48) with $\mathbf{v}_h = d_t \zeta_u^m$, multiplying by Δt , summing from $m = 1$ to n , employing a summation-by-parts formula on the terms on the right hand side that are contained in $a_h(\cdot; \cdot, d_t \zeta_u^m)$

to transfer the discrete time derivative on $d_t \zeta_u^m$ to the other terms, and assuming that the exact solution is sufficiently smooth in time, leads to

$$\Delta t \sum_{m=1}^n \|d_t \zeta_u^m\|_{\Omega^s}^2 \leq C(h^{2k_f} + h^{2k_c} + (\Delta t)^2).$$

We do not provide the details of this proof here, but instead refer to [18, p.42].

6.3. Existence and uniqueness of the concentration solution

In this section, we will prove existence and uniqueness of the discrete concentration solution $\mathbf{c}_h^n \in \mathbf{C}_h$ to eq. (40). First observe that assumption eq. (4b) on D implies that for $v \in [L^\infty(\Omega^d)]^{\dim}$,

$$\|D(v)\|_{0,\infty,\Omega^d} \leq C(1 + \|v\|_{0,\infty,\Omega^d}), \quad (91)$$

and together with eq. (22) that

$$\|D(v)\|_{0,\infty,\partial K} \leq C(1 + \|v\|_{0,\infty,\partial K}) \leq C \quad \forall v \in [W^{1,\infty}(K)]^{\dim}, K \in \mathcal{T}^d, \quad (92)$$

where C depends on $\|v\|_{1,\infty,K}$. Therefore by eqs. (82), (91) and (92), there exists a constant $\tilde{D}_{\max} > 0$ that depends on d and the upper bound in eq. (82) such that

$$\|\tilde{D}(u_h^n)\|_{0,\infty,\Omega} \leq \tilde{D}_{\max}, \quad \|\tilde{D}(u_h^n)\|_{0,\infty,\partial K} \leq \tilde{D}_{\max}. \quad (93)$$

With eq. (93), the following coercivity result can be proved following the same steps as the proofs of [16, Lemmas 2 and 3].

Theorem 6.12 (Coercivity of $B_h^{tr}(u_h^n; \mathbf{w}_h, \mathbf{w}_h)$). *There exists a constant $\beta_0 > 0$ such that if $\beta_{tr} > \beta_0^{tr}$, then for all $\mathbf{w}_h \in \mathbf{C}_h$,*

$$B_h^{tr}(u_h^n; \mathbf{w}_h, \mathbf{w}_h) \geq C_{tr} \|\mathbf{w}_h\|_c^2 + \frac{1}{2} \sum_{K \in \mathcal{T}^d} \int_K \nabla \cdot u_h^n w_h^2 \, dx, \quad (94)$$

where $C_{tr} > 0$ is a constant that depends on d, D_{\min} , and the upper bound in eq. (82).

Now that we have coercivity, we proceed with the existence and stability proof for the discrete concentration.

Theorem 6.13. *Let $c_0 \in L^2(\Omega)$ and $g_i - g_p \in L^\infty(0, T; L^\infty(\Omega^d))$. Let $n \geq 1$ and let \mathbf{u}_h^n be the solution to eq. (39) that satisfies eqs. (62a) and (62b). If $d_n \Delta t < 1$, where $d_n =: \frac{1}{\phi_*}(1 + C\|g_i^n - g_p^n\|_{0,\infty,\Omega^d})$, then there exists a unique solution $\mathbf{c}_h^n \in \mathbf{C}_h$ to eq. (40). Furthermore, if $K := \sum_{m=1}^n d_m/(1 - \Delta t d_m)$, then*

$$\phi_* \|c_h^n\|_\Omega^2 + C_{tr} \Delta t \sum_{m=1}^n \|\mathbf{c}_h^m\|_c^2 \leq e^{K\Delta t} (\phi_* \|c_0\|_\Omega^2 + \|g_i\|_{\ell^2(0,T;L^2(\Omega^d))}^2). \quad (95)$$

Proof. Let $\mathbf{w}_h = \mathbf{c}_h^n$ in eq. (40). From the algebraic inequality $(a-b)a \geq \frac{1}{2}(a^2 - b^2)$, eq. (94), and the assumption that $0 \leq c_I \leq 1$ a.e., we have

$$\begin{aligned} \frac{\phi_*}{2\Delta t} (\|c_h^n\|_\Omega^2 - \|c_h^{n-1}\|_\Omega^2) + C_{tr} \|\mathbf{c}_h^n\|_c^2 + \int_{\Omega^d} g_p^n (c_h^n)^2 \, dx &\leq \sum_{K \in \mathcal{T}^d} \int_K c_I g_i^n c_h^n \, dx - \frac{1}{2} \sum_{K \in \mathcal{T}^d} \int_K \nabla \cdot u_h^n (c_h^n)^2 \, dx \\ &\leq \|g_i^n\|_{\Omega^d} \|c_h^n\|_{\Omega^d} + \frac{1}{2} \|\nabla \cdot u_h^n\|_{0,\infty,\Omega^d} \|c_h^n\|_{\Omega^d}^2. \end{aligned}$$

Multiplying this inequality by $2\Delta t$, summing from $m = 1$ to n , noting that $g_p \geq 0$, and recalling eq. (19a) with stability of the L^2 -projections Π_C and Π_Q , we obtain

$$\begin{aligned} & \phi_* \|c_h^n\|_\Omega^2 + 2C_{tr}\Delta t \sum_{m=1}^n \|c_h^m\|_c^2 \\ & \leq \phi_* \|c_h^0\|_\Omega^2 + \Delta t \sum_{m=1}^n (\|g_i^m\|_{\Omega^d}^2 + \|c_h^m\|_{\Omega^d}^2) + \Delta t \sum_{m=1}^n \|\Pi_Q(g_i^m - g_p^m)\|_{0,\infty,\Omega^d} \|c_h^m\|_{\Omega^d}^2 \\ & \leq \phi_* \|c_0\|_\Omega^2 + \|g_i^m\|_{\ell^2(0,T;L^2(\Omega^d))}^2 + \Delta t \sum_{m=1}^n (1 + C\|g_i^m - g_p^m\|_{0,\infty,\Omega^d}) \|c_h^m\|_\Omega^2. \end{aligned}$$

Equation (95) follows after applying Grönwall's inequality [36, Lemma 27]. This stability bound then implies the existence of a unique solution since the system is finite dimensional and linear. \square

6.4. Error estimate for the discrete concentration

This section is devoted to proving an error estimate for the discrete concentration.

Lemma 6.14 (Error equation for eq. (40)).

$$\begin{aligned} & \sum_{K \in \mathcal{T}} \int_K \phi \, d_t \zeta_c^n w_h \, dx + B_h^{tr}(u_h^n; \zeta_c^n, \mathbf{w}_h) + \int_{\Omega^d} g_p^n \zeta_c^n w_h \, dx \\ & = \sum_{K \in \mathcal{T}} \int_K \phi \, d_t \xi_c^n w_h \, dx - \sum_{K \in \mathcal{T}} \int_K \phi (d_t c^n - \partial_t c^n) w_h \, dx + B_h^{tr}(u_h^n; \xi_c^n, \mathbf{w}_h) \\ & \quad + B_h^a(u_h^n - u_h^n; \mathbf{c}^n, \mathbf{w}_h) + B_h^d(u_h^n; \mathbf{c}^n, \mathbf{w}_h) - B_h^d(u_h^n; \mathbf{c}^n, \mathbf{w}_h) + \int_{\Omega^d} g_p^n \xi_c^n w_h \, dx. \end{aligned} \tag{96}$$

Proof. By Lemma 3.1, for $t = t^n$, we have

$$\sum_{K \in \mathcal{T}} \int_K \phi \, \partial_t c^n w_h \, dx + B_h^{tr}(u_h^n; \mathbf{c}^n, \mathbf{w}_h) + \int_{\Omega^d} g_p^n c^n w_h \, dx = \sum_{K \in \mathcal{T}^d} \int_K c_I g_i^n w_h \, dx \quad \forall \mathbf{w}_h \in \mathbf{C}_h, \tag{97}$$

where $\mathbf{c}^n = (c^n, \gamma(c^n))$. Subtracting eq. (97) from eq. (40) yields that for all $\mathbf{w}_h \in \mathbf{C}_h$,

$$\sum_{K \in \mathcal{T}} \int_K \phi (d_t c_h^n - \partial_t c^n) w_h \, dx + B_h^{tr}(u_h^n; \mathbf{c}_h^n, \mathbf{w}_h) - B_h^{tr}(u_h^n; \mathbf{c}^n, \mathbf{w}_h) + \int_{\Omega^d} g_p^n (c_h^n - c^n) w_h \, dx = 0. \tag{98}$$

Next, we rewrite the B_h^{tr} terms in eq. (98) by observing that B_h^{tr} is linear in the second slot and that B_h^a is linear in the first slot:

$$\begin{aligned} & B_h^{tr}(u_h^n; \mathbf{c}_h^n, \mathbf{w}_h) - B_h^{tr}(u_h^n; \mathbf{c}^n, \mathbf{w}_h) \\ & = B_h^{tr}(u_h^n; \mathbf{c}_h^n - \mathbf{c}^n, \mathbf{w}_h) + B_h^a(u_h^n - u_h^n; \mathbf{c}^n, \mathbf{w}_h) + [B_h^d(u_h^n; \mathbf{c}^n, \mathbf{w}_h) - B_h^d(u_h^n; \mathbf{c}^n, \mathbf{w}_h)]. \end{aligned} \tag{99}$$

Using eq. (52), again the linearity of B_h^{tr} in the second slot, and eqs. (98) and (99) completes the proof. \square

Theorem 6.15. *In addition to the assumptions in Theorem 6.5, suppose that*

$$\begin{aligned} c_0 & \in H^{k_c}(\Omega), \quad c \in L^2(0, T; H^{k_c+1}(\Omega)) \cap L^\infty(0, T; W^{1,\infty}(\Omega)), \quad \partial_t c \in L^2(0, T; H^{k_c}(\Omega)), \\ \partial_{tt} c & \in L^2(0, T; L^2(\Omega)), \quad g_i, g_p \in L^\infty(0, T; L^\infty(\Omega^d)). \end{aligned}$$

Then for sufficiently small Δt ,

$$\|\zeta_c^n\|_\Omega^2 + \Delta t \sum_{m=1}^n \|\zeta_c^m\|_c^2 \leq C(h^{2k_f} + h^{2k_c} + (\Delta t)^2), \quad (100)$$

where C depends on ϕ_*, ϕ^*, d, D and the regularity of the solution but is independent of h and Δt .

Proof. Setting $\mathbf{w}_h = \zeta_c^n$ in Lemma 6.14, using the inequality $a(a-b) \geq \frac{a^2-b^2}{2}$, and Theorem 6.12,

$$\begin{aligned} & \frac{\phi_*}{2\Delta t} (\|\zeta_c^n\|_\Omega^2 - \|\zeta_c^{n-1}\|_\Omega^2) + C_{tr} \|\zeta_c^n\|_c^2 + \int_{\Omega^d} g_p^n (\zeta_c^n)^2 dx \\ & \leq \sum_{K \in \mathcal{T}} \int_K \phi d_t \xi_c^n \zeta_c^n dx + \sum_{K \in \mathcal{T}} \int_K \phi (\partial_t c^n - d_t c^n) \zeta_c^n dx + B_h^{tr}(u_h^n; \xi_c^n, \zeta_c^n) \\ & \quad + B_h^a(u^n - u_h^n; c^n, \zeta_c^n) + [B_h^d(u^n; c^n, \zeta_c^n) - B_h^d(u_h^n; c^n, \zeta_c^n)] \\ & \quad - \frac{1}{2} \sum_{K \in \mathcal{T}^d} \int_K \nabla \cdot u_h^n (\zeta_c^n)^2 dx + \int_{\Omega^d} g_p^n \xi_c^n \zeta_c^n dx \\ & := I_1 + \dots + I_7. \end{aligned}$$

Using eq. (5a), the Cauchy-Schwarz inequality, Young's inequality with constant γ , and eq. (53),

$$\begin{aligned} I_1 & \leq \phi^* \left\| \frac{1}{\Delta t} \int_{t^{n-1}}^{t^n} \partial_t \xi_c dt \right\|_\Omega \|\zeta_c^n\|_\Omega \leq \phi^* \frac{1}{\sqrt{\Delta t}} \|\partial_t \xi_c\|_{L^2(t^{n-1}, t^n, L^2(\Omega))} \|\zeta_c^n\|_\Omega \\ & \leq C \frac{(\phi^*)^2 h^{2k_c}}{\phi_* \Delta t} \|\partial_t c\|_{L^2(t^{n-1}, t^n, H^{k_c}(\Omega))}^2 + \gamma(\phi_* \|\zeta_c^n\|_\Omega^2). \end{aligned}$$

Again by eq. (5a), this time using Taylor's theorem in integral form, and applying Young's inequality,

$$\begin{aligned} I_2 & = \sum_{K \in \mathcal{T}} \int_K \phi (\partial_t c^n - d_t c^n) \zeta_c^n dx \leq \phi^* \left\| \frac{1}{\Delta t} \int_{t^{n-1}}^{t^n} (t - t^{n-1}) \partial_{tt} c dt \right\|_\Omega \|\zeta_c^n\|_\Omega \\ & \leq C \frac{(\phi^*)^2 \Delta t}{\phi_*} \|\partial_{tt} c\|_{L^2(t^{n-1}, t^n, L^2(\Omega))}^2 + \gamma(\phi_* \|\zeta_c^n\|_\Omega^2). \end{aligned}$$

The following series of inequalities is dedicated to finding an upper bound for I_3 . By definition of B_h^{tr} ,

$$I_3 = B_h^{tr}(u_h^n; \xi_c^n, \zeta_c^n) = B_h^a(u_h^n; \xi_c^n, \zeta_c^n) + B_h^d(u_h^n; \xi_c^n, \zeta_c^n). \quad (101)$$

We will bound B_h^a and B_h^d separately, starting with B_h^a . Noting that $\xi_c^n - \bar{\xi}_c^n$ vanishes on facets, we have by eq. (17),

$$\begin{aligned} B_h^a(u_h^n; \xi_c^n, \zeta_c^n) & = - \sum_{K \in \mathcal{T}} \int_K \xi_c^n u_h^n \cdot \nabla \zeta_c^n dx + \sum_{K \in \mathcal{T}} \int_{\partial K} \xi_c^n (u_h^n - u^n) \cdot n (\zeta_c^n - \bar{\zeta}_c^n) ds \\ & \quad + \sum_{K \in \mathcal{T}} \int_{\partial K} \xi_c^n u^n \cdot n (\zeta_c^n - \bar{\zeta}_c^n) ds =: I_{311} + I_{312} + I_{313}. \end{aligned} \quad (102)$$

The term I_{311} can be bounded by Hölder's inequality, and eqs. (53) and (82):

$$I_{311} \leq C \|u_h^n\|_{0, \infty, \Omega} \|\xi_c^n\|_\Omega \|\nabla \zeta_c^n\|_\Omega \leq C h^{k_c} \|c^n\|_{k_c, \Omega} \|\zeta_c^n\|_c. \quad (103)$$

By Hölder's inequality and using eq. (22),

$$\begin{aligned} I_{312} &\leq \sum_{K \in \mathcal{T}} \|\xi_c^n\|_{0,\infty,\partial K} \|u_h^n - u^n\|_{\partial K} \|\zeta_c^n - \bar{\zeta}_c^n\|_{\partial K} \\ &\leq C \left(\sum_{K \in \mathcal{T}} h_K \|u_h^n - u^n\|_{\partial K}^2 \right)^{1/2} \|\zeta_c^n\|_c. \end{aligned} \quad (104)$$

Using Hölder's inequality and this time employing eqs. (21), (22) and (53),

$$\begin{aligned} I_{313} &= \sum_{K \in \mathcal{T}} \|\xi_c^n\|_{\partial K} \|u^n\|_{0,\infty,\partial K} \|\zeta_c^n - \bar{\zeta}_c^n\|_{\partial K} \\ &\leq C \left(\sum_{K \in \mathcal{T}} (\|\xi_c^n\|_K^2 + h_K^2 \|\xi_c^n\|_{1,K}^2) \right)^{1/2} \left(\sum_{K \in \mathcal{T}} h_K^{-1} \|\zeta_c^n - \bar{\zeta}_c^n\|_{\partial K}^2 \right)^{1/2} \\ &\leq Ch^{k_c} \|c^n\|_{k_c,\Omega} \|\zeta_c^n\|_c. \end{aligned} \quad (105)$$

Putting eqs. (102) to (105) together and using Young's inequality, we find

$$B_h^a(u_h^n; \xi_c^n, w_h) \leq C \left(\sum_{K \in \mathcal{T}} h_K \|u_h^n - u^n\|_{\partial K}^2 + h^{2k_c} \|c^n\|_{k_c,\Omega}^2 \right) + \epsilon \|\zeta_c^n\|_c^2 \quad (106)$$

We now bound B_h^d in eq. (101). Since $\xi_c^n - \bar{\xi}_c^n = 0$ on ∂K ,

$$B_h^d(u_h^n; \xi_c^n, \zeta_c^n) = \sum_{K \in \mathcal{T}} \int_K \tilde{D}(u_h^n) \nabla \xi_c^n \cdot \nabla \zeta_c^n \, dx - \sum_{K \in \mathcal{T}} \int_{\partial K} [\tilde{D}(u_h^n) \nabla \xi_c^n] \cdot n (\zeta_c^n - \bar{\zeta}_c^n) \, ds =: I_{321} + I_{322}. \quad (107)$$

By Hölder's inequality, eq. (93), and eq. (53),

$$I_{321} \leq \tilde{D}_{\max} \|\nabla \xi_c^n\|_{\Omega} \|\nabla \zeta_c^n\|_{\Omega} \leq Ch^{k_c} \|c\|_{k_c+1,\Omega} \|\zeta_c^n\|_c. \quad (108)$$

Again by Hölder's inequality and this time using eqs. (21), (53) and (93),

$$\begin{aligned} |I_{322}| &\leq \tilde{D}_{\max} \left(\sum_{K \in \mathcal{T}} h_K \|\nabla \xi_c^n\|_{\partial K}^2 \right)^{1/2} \left(\sum_{K \in \mathcal{T}} h_K^{-1} \|\zeta_c^n - \bar{\zeta}_c^n\|_{\partial K}^2 \right)^{1/2} \\ &\leq C \left(\sum_{K \in \mathcal{T}} (\|\nabla \xi_c^n\|_K^2 + h_K^2 \|\nabla \xi_c^n\|_{1,K}^2) \right)^{1/2} \|\zeta_c^n\|_c \leq Ch^{k_c} \|c^n\|_{k_c+1,\Omega} \|\zeta_c^n\|_c. \end{aligned} \quad (109)$$

Hence, the combination of eqs. (107) to (109) and using Young's inequality results in:

$$B_h^d(u_h^n; \xi_c^n, \zeta_c^n) \leq Ch^{k_c} \|c^n\|_{k_c+1,\Omega} \|\zeta_c^n\|_c \leq Ch^{2k_c} \|c^n\|_{k_c+1,\Omega}^2 + \epsilon \|\zeta_c^n\|_c^2. \quad (110)$$

Therefore, from eqs. (106) and (110),

$$I_3 \leq Ch^{2k_c} \|c^n\|_{k_c+1,\Omega}^2 + \sum_{K \in \mathcal{T}} h_K \|u_h^n - u^n\|_{\partial K}^2 + 2\epsilon \|\zeta_c^n\|_c^2.$$

Since $c = \gamma(c)$ on ∂K ,

$$I_4 = - \sum_{K \in \mathcal{T}} \int_K c^n (u^n - u_h^n) \cdot \nabla \zeta_c^n \, dx + \sum_{K \in \mathcal{T}} \int_{\partial K} c^n (u^n - u_h^n) \cdot n (\zeta_c^n - \bar{\zeta}_c^n) \, ds =: I_{41} + I_{42}. \quad (111)$$

Hölder's and Young's inequalities give

$$I_{41} \leq \|u^n - u_h^n\|_{\Omega} \|c^n\|_{0,\infty,\Omega} \|\zeta_c^n\|_c \leq C \|c^n\|_{0,\infty,\Omega}^2 \|u^n - u_h^n\|_{\Omega}^2 + \epsilon \|\zeta_c^n\|_c^2, \quad (112)$$

and

$$I_{42} \leq C \|c^n\|_{0,\infty,\Omega} \left(\sum_{K \in \mathcal{T}_h} h_K \|u^n - u_h^n\|_{\partial K}^2 \right)^{1/2} \|\zeta_c^n\|_c \leq C \|c^n\|_{0,\infty,\Omega}^2 \sum_{K \in \mathcal{T}_h} h_K \|u^n - u_h^n\|_{\partial K}^2 + \epsilon \|\zeta_c^n\|_c^2. \quad (113)$$

Collecting eqs. (111) to (113) leads to

$$I_4 \leq 2\epsilon \|\zeta_c^n\|_c^2 + C \|c^n\|_{0,\infty,\Omega}^2 \left(\|u^n - u_h^n\|_{\Omega}^2 + \sum_{K \in \mathcal{T}_h} h_K \|u^n - u_h^n\|_{\partial K}^2 \right).$$

Since $c = \gamma(c)$ on element boundaries and $\tilde{D}(u^n) - \tilde{D}(u_h^n) = 0$ in Ω^s ,

$$\begin{aligned} I_5 &= \sum_{K \in \mathcal{T}^d} \int_K [D(u^n) - D(u_h^n)] \nabla c^n \cdot \nabla \zeta_c^n \, dx - \sum_{K \in \mathcal{T}^d} \int_{\partial K} [(D(u^n) - D(u_h^n)) \nabla c^n] \cdot n (\zeta_c^n - \bar{\zeta}_c^n) \, ds \\ &=: I_{51} + I_{52}. \end{aligned} \quad (114)$$

Using the Lipschitz property of D eq. (4c), Hölder's and Young's inequalities,

$$I_{51} \leq C \|u^n - u_h^n\|_{\Omega^d} \|\nabla c^n\|_{0,\infty,\Omega^d} \|\nabla \zeta_c^n\|_{\Omega^d} \leq C \|\nabla c^n\|_{0,\infty,\Omega^d}^2 \|u^n - u_h^n\|_{\Omega^d}^2 + \epsilon \|\zeta_c^n\|_c^2. \quad (115)$$

Similarly,

$$\begin{aligned} I_{52} &\leq \sum_{K \in \mathcal{T}^d} C \|u^n - u_h^n\|_{\partial K} \|\nabla c^n\|_{0,\infty,K} \|\zeta_c^n - \bar{\zeta}_c^n\|_{\partial K} \leq C \|c^n\|_{1,\infty,\Omega^d} \left(\sum_{K \in \mathcal{T}^d} h_K \|u^n - u_h^n\|_{\partial K}^2 \right)^{1/2} \|\zeta_c^n\|_c \\ &\leq C \left(\|c^n\|_{1,\infty,\Omega^d}^2 \sum_{K \in \mathcal{T}^d} h_K \|u^n - u_h^n\|_{\partial K}^2 \right) + \epsilon \|\zeta_c^n\|_c^2. \end{aligned} \quad (116)$$

Therefore, substituting eqs. (115) and (116) in eq. (114) yields

$$I_5 \leq 2\epsilon \|\zeta_c^n\|_c^2 + C \|c^n\|_{1,\infty,\Omega^d}^2 \left(\sum_{K \in \mathcal{T}^d} h_K \|u^n - u_h^n\|_{\partial K}^2 + \|u^n - u_h^n\|_{\Omega^d}^2 \right).$$

By eq. (19a), the stability of the L^2 -projection Π_Q and Hölder's inequality,

$$I_6 = -\frac{1}{2} \sum_{K \in \mathcal{T}^d} \int_K \Pi_Q(g_i^n - g_p^n) (\zeta_c^n)^2 \, dx \leq \frac{1}{2\phi_*} \|g_i^n - g_p^n\|_{0,\infty,\Omega^d} (\phi_* \|\zeta_c^n\|_{\Omega}^2).$$

Finally, using Hölder's inequality, eq. (53), and Young's inequality,

$$\begin{aligned} I_7 &\leq \|g_p^n\|_{0,\infty,\Omega^d} \|\xi_c^n\|_{\Omega^d} \|\zeta_c^n\|_{\Omega^d} \leq Ch^{k_c} \|g_p^n\|_{0,\infty,\Omega^d} \|c^n\|_{k_c,\Omega^d} \|\zeta_c^n\|_{\Omega^d} \\ &\leq Ch^{2k_c} \phi_*^{-1} \|g_p^n\|_{0,\infty,\Omega^d}^2 \|c^n\|_{k_c,\Omega^d}^2 + \gamma(\phi_* \|\zeta_c^n\|_{\Omega}^2). \end{aligned}$$

Collecting all bounds, choosing $\epsilon = C_{tr}/12$ (C_{tr} is the coercivity constant), $\gamma = 1/6$, and recalling that $g_p^n \geq 0$, we find:

$$\begin{aligned} & \frac{\phi_*}{2\Delta t} (\|\zeta_c^n\|_\Omega^2 - \|\zeta_c^{n-1}\|_\Omega^2) + \frac{C_{tr}}{2} \|\zeta_c^n\|_c^2 \\ & \leq C \left(\frac{(\phi^*)^2 \phi_*^{-1} h^{2k_c}}{\Delta t} \|\partial_t c\|_{L^2(t^{n-1}, t^n; H^{k_c}(\Omega))}^2 + (\phi^*)^2 \phi_*^{-1} \Delta t \|\partial_{tt} c\|_{L^2(t^{n-1}, t^n; L^2(\Omega))}^2 \right. \\ & \quad + h^{2k_c} \|c^n\|_{k_c+1, \Omega}^2 + \|c^n\|_{1, \infty, \Omega}^2 \|u^n - u_h^n\|_\Omega^2 + (\|c^n\|_{1, \infty, \Omega}^2 + 1) \sum_{K \in \mathcal{T}} h_K \|u^n - u_h^n\|_{\partial K}^2 \\ & \quad \left. + h^{2k_c} \phi_*^{-1} \|g_p^n\|_{0, \infty, \Omega^d}^2 \|c^n\|_{k_c, \Omega^d}^2 \right) + \frac{1}{2} \left(\phi_*^{-1} \|g_i^n - g_p^n\|_{0, \infty, \Omega^d} + 1 \right) (\phi_* \|\zeta_c^n\|_\Omega^2). \end{aligned}$$

Multiplying by $2\Delta t$, summing over m , and using Corollary 6.6,

$$\begin{aligned} & \phi_* \|\zeta_c^n\|_\Omega^2 + C_{tr} \Delta t \sum_{m=1}^n \|\zeta_c^m\|_c^2 \\ & \leq \phi_* \|\zeta_c^0\|_\Omega^2 + C \left[h^{2k_c} \left(\frac{(\phi^*)^2}{\phi_*} \|\partial_t c\|_{L^2(0, T; H^{k_c}(\Omega))}^2 + \|c\|_{\ell^2(0, T; H^{k_c+1}(\Omega))}^2 \right) \right. \\ & \quad + (\phi^*)^2 \phi_*^{-1} (\Delta t)^2 \|\partial_{tt} c\|_{L^2(0, T; L^2(\Omega))}^2 + \Delta t \sum_{m=1}^n \left(\phi_*^{-1} \|g_i^m - g_p^m\|_{0, \infty, \Omega^d} + 1 \right) (\phi_* \|\zeta_c^m\|_\Omega^2) \\ & \quad \left. + \Delta t \sum_{m=1}^n (\|c^m\|_{1, \infty, \Omega}^2 + 1) ((\Delta t)^2 + h^{2k_f} + h^{2k_c}) + h^{2k_c} \Delta t \sum_{m=1}^n \phi_*^{-1} \|g_p^m\|_{0, \infty, \Omega^d}^2 \|c^m\|_{k_c, \Omega^d}^2 \right]. \end{aligned}$$

Using [23, Lemma 1.58] and eq. (53),

$$\|\zeta_c^0\|_\Omega = \|c_h^0 - \mathcal{I}c_0\|_\Omega \leq \|\Pi_C c_0 - c_0\|_\Omega + \|c_0 - \mathcal{I}c_0\|_\Omega \leq Ch^{k_c} \|c_0\|_{k_c, \Omega}.$$

Therefore, the result follows by Grönwall's inequality [36, Lemma 27] assuming that Δt is sufficiently small. \square

By the triangle inequality and eq. (53), we immediately have

$$\|c_h^n - c^n\|_\Omega \leq C(\Delta t + h^{k_f} + h^{k_c}), \quad \forall n \geq 1. \quad (117)$$

7. NUMERICAL EXAMPLES

Algorithm 1 is implemented in the higher-order finite element library Netgen/NGSolve [47, 48]. In all numerical examples we choose $\bar{\Omega} = [0, 1]^2$ with subregions $\bar{\Omega}^d = [0, 1] \times [0, 0.5]$ and $\bar{\Omega}^s = [0, 1] \times [0.5, 1]$. We furthermore choose the penalty parameters as $\beta_s = 6k_f^2$ and $\beta_{tr} = 6k_c^2$ [2, Lemma 1, Section 5].

7.1. Example 1

We first consider the constant coefficient case, i.e., the time-dependent one-way coupled problem in which the numerical solution to the Stokes/Darcy problem is unaffected by the concentration. Let $\alpha = \frac{1}{2}(1 + 4\pi^2)\sqrt{\kappa}$, $\tilde{D} =$

$D = \begin{bmatrix} 0.01 & 0.005 \\ 0.005 & 0.02 \end{bmatrix}$ on Ω , and $T = 0.1$. The source terms and boundary conditions for the Stokes/Darcy-transport problem are chosen such that the exact solution is given by

$$u^s = \left(-\frac{1}{2\pi^2} \sin(\pi x_1 + t) e^{(x_2+t)/2}, \frac{1}{\pi} \cos(\pi x_1 + t) e^{(x_2+t)/2}\right)^T \quad (118a)$$

$$u^d = \left(-2 \sin(\pi x_1 + t) e^{(x_2+t)/2}, \frac{1}{\pi} \cos(\pi x_1 + t) e^{(x_2+t)/2}\right)^T, \quad (118b)$$

$$p^s = \frac{\kappa\mu - 2}{\kappa\pi} \cos(\pi x_1 + t) e^{(x_2+t)/2}, \quad (118c)$$

$$p^d = -\frac{2}{\kappa\pi} \cos(\pi x_1 + t) e^{(x_2+t)/2}, \quad (118d)$$

$$c = \sin(2\pi(x_1 - t)) \cos(2\pi(x_2 - t)). \quad (118e)$$

Note that this solution satisfies all the interface conditions and that $\nabla \cdot u^s = 0$ in Ω^s .

We present our numerical results for a wide range of values for κ and μ : $\kappa = \mu = 1$; $\kappa = 10^3$, $\mu = 10^{-6}$; $\kappa = 1$, $\mu = 10^{-6}$; and $\kappa = 10^{-3}$, $\mu = 10^{-6}$. Since we are primarily interested in the spatial error, to minimize the temporal error as much as possible, we use the third order backward differentiation formulae (BDF3) as time stepping method even though the sequential algorithm 1 is only first order accurate in time. We choose $\Delta t = 0.1h^{k_f}/(k_f + 1)$ and present errors and rates of convergence using $k_f = 2$, $k_c = 1$ in Tables 1 to 3 and using $k_f = 3$, $k_c = 2$ in Tables 4 to 6.

TABLE 1. Errors and rates of convergence at final time $T = 0.1$ for u_h and p_h in the Stokes region Ω^s for the test case in Section 7.1 using $k_f = 2$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^2/3$.

h	dofs	$\ u_h - u\ _{\Omega^s}$	rate	$\ p_h - p\ _{\Omega^s}$	rate	$\ \nabla \cdot u_h\ _{\Omega^s}$
$\kappa = 1, \mu = 1$						
1/4	745	2.6e-04	—	1.2e-02	—	1.4e-16
1/8	3811	2.0e-05	3.7	1.9e-03	2.6	1.8e-16
1/16	14167	2.2e-06	3.2	4.5e-04	2.1	1.6e-16
1/32	57181	2.7e-07	3.1	1.1e-04	2.1	1.7e-16
$\kappa = 10^3, \mu = 10^{-6}$						
1/4	745	2.5e-04	—	1.3e-05	—	8.5e-17
1/8	3811	2.0e-05	3.6	1.9e-06	2.8	9.7e-17
1/16	14167	2.2e-06	3.2	4.5e-07	2.1	9.4e-17
1/32	57181	2.6e-07	3.1	1.1e-07	2.1	9.0e-17
$\kappa = 1, \mu = 10^{-6}$						
1/4	745	6.4e-04	—	2.7e-02	—	8.6e-16
1/8	3811	4.8e-05	3.7	5.3e-03	2.3	1.8e-15
1/16	14167	4.4e-06	3.4	1.2e-03	2.1	3.0e-15
1/32	57181	5.2e-07	3.1	3.0e-04	2.0	6.0e-15
$\kappa = 10^{-3}, \mu = 10^{-6}$						
1/4	745	7.3e-04	—	1.2e+01	—	1.2e-13
1/8	3811	5.0e-05	3.9	1.9e+00	2.6	1.6e-13
1/16	14167	4.1e-06	3.6	4.5e-01	2.1	1.4e-13
1/32	57181	3.8e-07	3.4	1.1e-01	2.1	1.5e-13

Tables 1 and 2 for $k_f = 2$, $k_c = 1$, and Tables 4 and 5 for $k_f = 3$, $k_c = 2$ show that in the Stokes and Darcy regions u_h and p_h both converge optimally in the L^2 -norm with orders $k_f + 1$ and k_f , respectively. This is

TABLE 2. Errors and rates of convergence at final time $T = 0.1$ for u_h and p_h in the Darcy region Ω^d for the test case in Section 7.1 using $k_f = 2$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^2/3$.

h	dofs	$\ u_h - u\ _{\Omega^d}$	rate	$\ p_h - p\ _{\Omega^d}$	rate	$\ \Pi_Q(\nabla \cdot (u_h - u))\ _{\Omega^d}$
$\kappa = 1, \mu = 1$						
1/4	745	3.1e-03	–	9.1e-03	–	5.9e-09
1/8	3811	1.9e-04	4.0	1.4e-03	2.7	9.1e-11
1/16	14167	2.5e-05	2.9	3.7e-04	1.9	1.8e-12
1/32	57181	2.7e-06	3.2	8.4e-05	2.1	3.8e-12
$\kappa = 10^3, \mu = 10^{-6}$						
1/4	745	3.1e-03	–	9.1e-06	–	5.9e-09
1/8	3811	1.9e-04	4.0	1.4e-06	2.7	9.1e-11
1/16	14167	2.5e-05	2.9	3.7e-07	1.9	1.7e-12
1/32	57181	2.7e-06	3.2	8.4e-08	2.1	3.8e-12
$\kappa = 1, \mu = 10^{-6}$						
1/4	745	3.1e-03	–	9.1e-03	–	5.9e-09
1/8	3811	1.9e-04	4.0	1.4e-03	2.7	9.1e-11
1/16	14167	2.5e-05	2.9	3.7e-04	1.9	1.8e-12
1/32	57181	2.7e-06	3.2	8.4e-05	2.1	3.8e-12
$\kappa = 10^{-3}, \mu = 10^{-6}$						
1/4	745	3.1e-03	–	9.1e+00	–	5.9e-09
1/8	3811	1.9e-04	4.0	1.4e+00	2.7	9.1e-11
1/16	14167	2.5e-05	2.9	3.7e-01	1.9	1.8e-12
1/32	57181	2.6e-06	3.2	8.4e-02	2.1	4.7e-12

TABLE 3. Errors and rates of convergence at final time $T = 0.1$ for c_h in Ω , on a mesh with $h = 1/4, 1/8, 1/16, 1/32$, for the test case in Section 7.1 using $k_f = 2$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^2/3$.

	$\kappa = 1, \mu = 1$		$\kappa = 10^3, \mu = 10^{-6}$		$\kappa = 1, \mu = 10^{-6}$		$\kappa = 10^{-3}, \mu = 10^{-6}$	
dofs	$\ c - c_h\ _{\Omega}$	rate	$\ c - c_h\ _{\Omega}$	rate	$\ c - c_h\ _{\Omega}$	rate	$\ c - c_h\ _{\Omega}$	rate
184	9.7e-02	–	9.7e-02	–	9.7e-02	–	9.8e-02	–
944	2.2e-02	2.1	2.2e-02	2.1	2.2e-02	2.1	2.2e-02	2.1
3520	5.4e-03	2.0	5.4e-03	2.0	5.4e-03	2.0	5.0e-03	2.2
14216	1.1e-03	2.3	1.1e-03	2.3	1.1e-03	2.3	1.1e-03	2.2

consistent with our theoretical convergence rate in Corollary 6.6 that predicts at least suboptimal rates for the velocity. The right most columns in these tables demonstrate pointwise mass conservation.

Furthermore, even though the magnitude of the pressure error changes dramatically as we change the values of κ and μ , there is no significant change in the velocity errors. This is more pronounced in the case where both the permeability and the viscosity are small (10^{-3} and 10^{-6}). This confirms that the velocity error bounds in Theorem 6.5 and Corollary 6.6 are independent of the pressure error.

We furthermore observe from Table 3 and Table 6 that c_h converges optimally in the L^2 -norm with order $k_c + 1$. This supports our result eq. (117) that shows at least suboptimal convergence.

7.2. Example 2

We now consider the fully coupled problem by incorporating the influence of the velocity solution on the dispersion/diffusion tensor and the dependence of the viscosity on the concentration solution. The source terms

TABLE 4. Errors and rates of convergence at final time $T = 0.1$ for u_h and p_h in the Stokes region Ω^s for the test case in Section 7.1 using $k_f = 3$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^3/4$.

h	dofs	$\ u_h - u\ _{\Omega^s}$	rate	$\ p_h - p\ _{\Omega^s}$	rate	$\ \nabla \cdot u_h\ _{\Omega^s}$
$\kappa = 1, \mu = 1$						
1/4	1161	5.6e-05	–	4.6e-03	–	2.0e-15
1/8	5993	1.4e-06	5.4	2.2e-04	4.4	3.7e-15
1/16	22081	7.1e-08	4.3	2.1e-05	3.3	4.5e-15
1/32	90241	3.7e-09	4.3	2.3e-06	3.2	6.3e-15
$\kappa = 10^3, \mu = 10^{-6}$						
1/4	1161	2.2e-05	–	7.3e-07	–	1.5e-16
1/8	5993	6.0e-07	5.2	5.7e-08	3.7	1.3e-16
1/16	22081	3.3e-08	4.2	6.4e-09	3.2	1.2e-16
1/32	90241	1.8e-09	4.2	7.7e-10	3.1	1.2e-16
$\kappa = 1, \mu = 10^{-6}$						
1/4	1161	2.2e-05	–	6.9e-04	–	1.5e-16
1/8	5993	6.1e-07	5.2	5.6e-05	3.6	1.5e-16
1/16	22081	3.3e-08	4.2	6.4e-06	3.1	1.2e-16
1/32	90241	1.8e-09	4.2	7.7e-07	3.1	1.2e-16
$\kappa = 10^{-3}, \mu = 10^{-6}$						
1/4	1161	2.6e-05	–	6.9e-01	–	5.5e-14
1/8	5993	9.4e-07	4.8	5.4e-02	3.7	4.2e-14
1/16	22081	4.5e-08	4.4	6.3e-03	3.1	1.9e-14
1/32	90241	2.2e-09	4.3	7.7e-04	3.0	1.0e-14

and boundary conditions for the Stokes/Darcy-transport problem (1) are chosen such that the exact solution is given by eq. (118). We define the diffusion dispersion tensor in Ω and the viscosity according to

$$\tilde{D}(u) = \begin{bmatrix} 1 + u_1^2 & 0 \\ 0 & 1 + u_2^2 \end{bmatrix}, \quad \mu(c) = \mu_0 \left[\left(\frac{\mu_0}{\mu_1} \right)^{1/4} c + (1 - c) \right]^{-4}, \quad (119)$$

where we remark the the viscosity is defined as the quarter-power mixing rule [38] with where $\mu_0 = 0.9$, $\mu_1 = 1.3$.

We use $k_f = 3$, $k_c = 2$, and BDF3 time stepping with $\Delta t = 0.1h^3/4$. We present numerical results for $\kappa = 10^3, 1, 10^{-3}$. We observe from Tables 7 and 8 that when μ is changed from a constant to a concentration dependent function the rate of convergence reduces from $k_f + 1$ to a value between k_f and $k_f + 1$. This is due to our choice $k_c = k_f - 1$ to achieve compatibility and is consistent with our a priori estimates eqs. (62a) and (62b) in the energy norm which imply that the rate of convergence of the velocity approximation is polluted by the concentration approximation. Indeed, from Table 9 we observe that c_h converges in the L^2 -norm with order k_f . Therefore, for the velocity we expect an order of at least $k_f - 1$ in the energy norm and k_f in the L^2 -norm. From the right most columns in Tables 7 and 8 we observe that the discretization is exactly mass conserving.

7.3. Example 3

In this final example, we simulate a more realistic problem similar to [17, Section 6.2] in which the permeability field in the Darcy region is highly heterogeneous. For this, let the boundary of the Stokes region be partitioned as $\Gamma^s = \Gamma_1^s \cup \Gamma_2^s \cup \Gamma_3^s$ where

$$\Gamma_1^s := \{x \in \Gamma^s : x_1 = 0\}, \quad \Gamma_2^s := \{x \in \Gamma^s : x_1 = 1\}, \quad \Gamma_3^s := \{x \in \Gamma^s : x_2 = 1\}.$$

TABLE 5. Errors and rates of convergence at final time $T = 0.1$ for u_h and p_h in the Darcy region Ω^d for the test case in Section 7.1 using $k_f = 3$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^3/4$.

h	dofs	$\ u_h - u\ _{\Omega^d}$	rate	$\ p_h - p\ _{\Omega^d}$	rate	$\ \Pi_Q(\nabla \cdot (u_h - u))\ _{\Omega^d}$
$\kappa = 1, \mu = 1$						
1/4	1161	1.2e-04	–	5.4e-04	–	4.1e-12
1/8	5993	3.6e-06	5.1	4.0e-05	3.8	8.3e-13
1/16	22081	2.2e-07	4.0	5.1e-06	3.0	2.9e-12
1/32	90241	1.3e-08	4.1	6.1e-07	3.1	1.2e-11
$\kappa = 10^3, \mu = 10^{-6}$						
1/4	1161	1.2e-04	–	5.4e-07	–	4.1e-12
1/8	5993	3.6e-06	5.1	4.0e-08	3.8	8.2e-13
1/16	22081	2.2e-07	4.0	5.1e-09	3.0	3.1e-12
1/32	90241	1.3e-08	4.1	6.1e-10	3.1	1.2e-11
$\kappa = 1, \mu = 10^{-6}$						
1/4	1161	1.2e-04	–	5.4e-04	–	4.1e-12
1/8	5993	3.5e-06	5.1	4.0e-05	3.8	7.2e-13
1/16	22081	2.2e-07	4.0	5.1e-06	3.0	3.0e-12
1/32	90241	1.3e-08	4.1	6.1e-07	3.1	1.2e-11
$\kappa = 10^{-3}, \mu = 10^{-6}$						
1/4	1161	1.2e-04	–	5.4e-01	–	4.1e-12
1/8	5993	3.5e-06	5.1	4.0e-02	3.8	8.9e-13
1/16	22081	2.2e-07	4.0	5.1e-03	3.0	3.3e-12
1/32	90241	1.3e-08	4.1	6.1e-04	3.1	1.3e-11

TABLE 6. Errors and rates of convergence at final time $T = 0.1$ for c_h in Ω , on a mesh with $h = 1/4, 1/8, 1/16, 1/32$, for the test case in Section 7.1 using $k_f = 3$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^3/4$.

	$\kappa = 1, \mu = 1$		$\kappa = 10^3, \mu = 10^{-6}$		$\kappa = 1, \mu = 10^{-6}$		$\kappa = 10^{-3}, \mu = 10^{-6}$	
dofs	$\ c - c_h\ _{\Omega}$	rate	$\ c - c_h\ _{\Omega}$	rate	$\ c - c_h\ _{\Omega}$	rate	$\ c - c_h\ _{\Omega}$	rate
318	2.1e-02	–	2.1e-02	–	2.1e-02	–	2.1e-02	–
1644	1.8e-03	3.6	1.8e-03	3.6	1.8e-03	3.6	1.8e-03	3.6
6144	2.2e-04	3.0	2.2e-04	3.0	2.2e-04	3.0	2.2e-04	3.0
24846	2.5e-05	3.2	2.5e-05	3.1	2.5e-05	3.1	2.5e-05	3.2

Similarly, let $\Gamma^d = \Gamma_1^d \cup \Gamma_2^d$ where

$$\Gamma_1^d := \{x \in \Gamma^d \mid x_1 = 0 \text{ or } x_1 = 1\}, \quad \Gamma_2^d := \{x \in \Gamma^d \mid x_2 = 0\}.$$

We impose the following boundary conditions:

$$\begin{aligned}
u &= (x_2(3/2 - x_2)/5, 0) && \text{on } \Gamma_1^s, \\
(-2\mu\varepsilon(u) + p\mathbb{I})n &= 0 && \text{on } \Gamma_2^s, \\
u \cdot n &= 0 \text{ and } (-2\mu\varepsilon(u) + p\mathbb{I})^t = 0 && \text{on } \Gamma_3^s, \\
u \cdot n &= 0 && \text{on } \Gamma_1^d, \\
p &= -0.05 && \text{on } \Gamma_2^d.
\end{aligned}$$

TABLE 7. Errors and rates of convergence at final time $T = 0.1$ for u_h and p_h in the Stokes region Ω^s for the test case in Section 7.2 using $k_f = 3$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^3/4$.

h	dofs	$\ u_h - u\ _{\Omega^s}$	rate	$\ p_h - p\ _{\Omega^s}$	rate	$\ \nabla \cdot u_h\ _{\Omega^s}$
$\kappa = 10^3$						
1/4	1161	9.2e-05	—	8.1e-03	—	2.3e-15
1/8	5993	3.7e-06	4.6	3.9e-04	4.4	5.6e-15
1/16	22081	2.4e-07	4.0	4.1e-05	3.2	6.1e-15
1/32	90241	2.1e-08	3.5	4.7e-06	3.1	8.5e-15
$\kappa = 1$						
1/4	1161	9.6e-05	—	8.1e-03	—	2.6e-15
1/8	5993	3.8e-06	4.7	3.9e-04	4.4	5.5e-15
1/16	22081	2.3e-07	4.1	4.1e-05	3.2	6.1e-15
1/32	90241	1.8e-08	3.6	4.7e-06	3.1	8.6e-15
$\kappa = 10^{-3}$						
1/4	1161	1.8e-03	—	6.9e-01	—	6.3e-15
1/8	5993	5.0e-05	5.1	5.4e-02	3.7	7.1e-15
1/16	22081	3.6e-06	3.8	6.3e-03	3.1	7.4e-15
1/32	90241	4.0e-07	3.2	7.7e-04	3.0	9.2e-15

TABLE 8. Errors and rates of convergence at final time $T = 0.1$ for u_h and p_h in the Darcy region Ω^d for the test case in Section 7.2 using $k_f = 3$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^3/4$.

h	dofs	$\ u_h - u\ _{\Omega^d}$	rate	$\ p_h - p\ _{\Omega^d}$	rate	$\ \Pi_Q(\nabla \cdot (u_h - u))\ _{\Omega^d}$
$\kappa = 10^3$						
1/4	1161	3.5e-03	—	6.6e-07	—	4.2e-12
1/8	5993	3.4e-04	3.4	4.2e-08	4.0	8.4e-13
1/16	22081	3.8e-05	3.1	5.3e-09	3.0	2.9e-12
1/32	90241	4.4e-06	3.1	6.3e-10	3.1	1.2e-11
$\kappa = 1$						
1/4	1161	3.5e-03	—	6.6e-04	—	4.2e-12
1/8	5993	3.4e-04	3.4	4.2e-05	4.0	9.0e-13
1/16	22081	3.8e-05	3.1	5.3e-06	3.0	3.0e-12
1/32	90241	4.4e-06	3.1	6.3e-07	3.1	1.2e-11
$\kappa = 10^{-3}$						
1/4	1161	3.9e-03	—	6.0e-01	—	4.2e-12
1/8	5993	3.4e-04	3.5	4.1e-02	3.9	9.6e-13
1/16	22081	3.8e-05	3.1	5.2e-03	3.0	3.5e-12
1/32	90241	4.4e-06	3.1	6.2e-04	3.1	1.5e-11

The first boundary condition on the left boundary Γ_1^s of Ω^s imposes a parabolic velocity profile. We set the permeability to

$$\kappa = 700(1 + 0.5(\sin(10\pi x_1) \cos(20\pi x_2^2) + \cos^2(6.4\pi x_1) \sin(9.2\pi x_2))) + 100, \quad (120)$$

a plot of which is given in Figure 2. The viscosity is defined by the quarter-power mixing rule as in eq. (119). The other parameters are set as $\mu = 0.1$, $\alpha = 0.5$, $k_f = 3$, $h = 1/80$, $\Delta t = 10^{-3}$, $T = 15$, and the source/sink terms are set to zero. In the Darcy region Ω^d the porosity is set to $\phi = 0.4$. The dispersion/diffusion tensor is

TABLE 9. Errors and rates of convergence at final time $T = 0.1$ for c_h in Ω , on a mesh with $h = 1/4, 1/8, 1/16, 1/32$, for the test case in Section 7.2 using $k_f = 3$, $k_c = k_f - 1$, and BDF3 time stepping with $\Delta t = 0.1h^3/4$.

	$\kappa = 10^3$		$\kappa = 1$		$\kappa = 10^{-3}$	
dofs	$\ c - c_h\ _\Omega$	rate	$\ c - c_h\ _\Omega$	rate	$\ c - c_h\ _\Omega$	rate
318	1.4e-02	—	1.4e-02	—	1.4e-02	—
1644	1.2e-03	3.6	1.2e-03	3.6	1.2e-03	3.6
6144	1.3e-04	3.2	1.3e-04	3.2	1.3e-04	3.2
24846	1.3e-05	3.3	1.3e-05	3.3	1.3e-05	3.3




FIGURE 2. The permeability field in $\Omega^d = [0, 1] \times [0, 0.5]$ defined by eq. (120).

defined as

$$\tilde{D}(u) = \begin{cases} \delta I, & \text{in } \Omega^s, \\ \phi d_m \mathbb{I} + d_l |u| \mathbb{T} + d_t |u| (\mathbb{I} - uu^T / |u|^2), & \text{in } \Omega^d, \end{cases}$$

where d_l , d_t , and d_m represent longitudinal and transverse dispersivities and the molecular diffusivity, respectively, and u^T is the transpose of the vector u . Under the condition $d_l \geq d_t$ (which is usually the case), $D(u)$ satisfies the assumptions eqs. (4a) to (4c) (see, for example, [26], [49, Lemmas 4.3, 4.4], and [44, Lemma 1.3]). In this numerical experiment, we choose $\delta = 10^{-6}$, $d_m = 10^{-5}$, $d_l = 10^{-5}$, and $d_t = 10^{-5}$. The initial velocity is set to zero while the initial concentration of the plume of contaminant is defined as

$$c_0(x) = \begin{cases} 0.95 & \text{if } \sqrt{(x_1 - 0.2)^2 + (x_2 - 0.7)^2} < 0.1, \\ 0.05 & \text{otherwise.} \end{cases}$$

We compute the solution using BDF3 time stepping. Figure 3 shows the computed velocity field after one time step and at the final time. In the Darcy region Ω^d , the flow field avoids areas with low permeability as expected. Figure 4 shows the pressure contours at various times which demonstrates the effect of the concentration on the pressure around the plume of contaminants, especially in the Stokes region. Figure 5 presents the plume of contaminant spreading through the surface water region and infiltrating the porous medium. We plot the solution 6 different instances in time. The contaminant plume stays compact while in the surface water region. Once it reaches the subsurface region it spreads out following a path dictated by the heterogeneous permeability structure of the porous medium.

8. CONCLUSIONS

In this paper, we introduced and analyzed a fully discrete sequential method for the fully coupled Stokes/Darcy-transport problem. The spatial discretization uses the HDG method which is higher-order accurate, strongly mass conservative, and compatible. We remark that the analysis also easily extends to the EDG-HDG method considered in [16, 17]. The sequential method discussed in the article linearizes the problem by time-lagging the concentration and decoupling the Stokes/Darcy and transport subproblems. We proved well-posedness and




FIGURE 3. The velocity field after one time step (left) and at the final time (right) for the example in Section 7.3. The color represents the magnitude of the velocity.

obtained a priori estimates in the energy norm. Finally, we presented numerical results demonstrating mass conservation and robustness with respect to varying permeability and optimal convergence in the L^2 -norm for one-way coupling.

REFERENCES

- [1] R. Adams. *Sobolev spaces*. Academic Press, Amsterdam Boston, 2003.
- [2] M. Ainsworth and G. Fu. Fully computable a posteriori error bounds for hybridizable discontinuous Galerkin finite element approximations. *J. Sci. Comput.*, 77:443–466, 2018.
- [3] S. Badia and R. Codina. Unified stabilized finite element formulations for the Stokes and the Darcy problems. *SIAM J. Numer. Anal.*, 47(3):1971–2000, 2009.
- [4] J. Bear and A. H.-D. Cheng. *Modeling groundwater flow and contaminant transport*, volume 23 of *Theory and Applications of Transport in Porous Media*. Springer, 2010.
- [5] G. S. Beavers and D. D. Joseph. Boundary conditions at a naturally impermeable wall. *J. Fluid. Mech.*, 30(1):197–207, 1967.
- [6] S. C. Brenner. Poincaré-Friedrichs inequalities for piecewise H^1 functions. *SIAM J. Numer. Anal.*, 41(1):306–324, 2003.
- [7] S. C. Brenner. Korn’s inequalities for piecewise H^1 vector fields. *Math. Comp.*, 73(247):1067–1087, 2004.
- [8] S. C. Brenner and L. R. Scott. *The Mathematical Theory of Finite Element Methods*. Springer, 3rd edition, 2010.
- [9] F. Brezzi and M. Fortin. *Mixed and Hybrid Finite Element Methods*, volume 15 of *Springer Series in Computational Mathematics*. Springer-Verlag New York Inc., 1991.
- [10] A. N. Brooks and T. J. R. Hughes. Streamline upwind Petrov–Galerkin formulations for convection dominated flows with particular emphasis on the incompressible Navier–Stokes equation. *Comput. Meth. Appl. Mech. Engrg.*, 32(1):199–259, 1982.
- [11] E. Burman and P. Hansbo. Stabilized Crouzeix–Raviart element for the Darcy–Stokes problem. *Numer. Meth. Part. D. E.*, 21(5):986–97, 2005.
- [12] J. Camaño, G. N. Gatica, R. Oyarzúa, R. Ruiz-Baier, and P. Venegas. New fully-mixed finite element methods for the Stokes–Darcy coupling. *Comput. Method. Appl. M.*, 295:362 – 395, 2015.
- [13] Y. Cao, M. Gunzburger, X. Hu, F. Hua, X. Wang, and W. Zhao. Finite element approximations for Stokes–Darcy flow with Beavers–Joseph interface conditions. *SIAM J. Numer. Anal.*, 47(6):4239–4256, 2010.
- [14] A. Çesmelioglu and B. Rivière. Existence of a weak solution for the fully coupled Navier–Stokes/Darcy–transport problem. *J. Differ. Equations*, 252(7):4138–4175, 2012.
- [15] A. Cesmelioglu and P. Chidyagwai. Numerical analysis of the coupling of free fluid with a poroelastic material. *Numer Methods Partial Differential Eq.*, 36:463–494, 2020.
- [16] A. Cesmelioglu and S. Rhebergen. A compatible embedded-hybridized discontinuous Galerkin method for the Stokes–Darcy–transport problem. *Commun. Appl. Math. Comput.*, 2021.
- [17] A. Cesmelioglu, S. Rhebergen, and G. N. Wells. An embedded-hybridized discontinuous Galerkin method for the coupled Stokes–Darcy system. *J. Comput. Appl. Math.*, 367, 2020.
- [18] N. Chaabane, V. Girault, C. Puelz, and B. Riviere. Convergence of IPDG for coupled time-dependent Navier–Stokes and Darcy equations. *J. Comput. Appl. Math.*, 324:25–48, 2017.




FIGURE 4. Pressure contours at times $t = \Delta t, 3, 6, 9, 12, 15$ for the example in Section 7.3. The color represents the concentration values.




FIGURE 5. The plume of contaminant at times $t = \Delta t, 3, 6, 9, 12, 15$ for the example in Section 7.3. The color represents the concentration values.

- [19] B. Cockburn, J. Gopalakrishnan, and R. Lazarov. Unified hybridization of discontinuous Galerkin, mixed, and continuous Galerkin methods for second order elliptic problems. *SIAM J. Numer. Anal.*, 47(2):1319–1365, 2009.
- [20] B. Cockburn and C.-W. Shu. The local discontinuous Galerkin finite element method for time-dependent convection–diffusion systems. *SIAM J. Numer. Anal.*, 35(6):2440–2463, 1998.
- [21] C. D’Angelo and P. Zunino. Robust numerical approximation of coupled Stokes’ and Darcy’s flows applied to vascular hemodynamics and biochemical transport. *ESAIM: M2AN*, 45(3):447–476, 2011.
- [22] C. Dawson, S. Sun, and M. F. Wheeler. Compatible algorithms for coupled flow and transport. *Comput. Methods Appl. Mech. Engrg.*, 193:2565–2580, 2004.
- [23] D. A. Di Pietro and A. Ern. *Mathematical aspects of discontinuous Galerkin methods*, volume 69 of *Mathématiques et Applications*. Springer–Verlag Berlin Heidelberg, 2012.
- [24] M. Discacciati. *Domain decomposition methods for the coupling of surface and groundwater flows*. PhD thesis, Ecole Polytechnique Federale de Sausanne, Sausanne, Switzerland, 2004.
- [25] M. Discacciati, E. Miglio, and A. Quarteroni. Mathematical and numerical models for coupling surface and groundwater flows. *Appl. Numer. Math.*, 43(1):57 – 74, 2002. 19th Dundee Biennial Conference on Numerical Analysis.
- [26] J. Douglas Jr., R. E. Ewing, and M. F. Wheeler. A time-discretization procedure for a mixed finite element approximation of miscible displacement in porous media. *RAIRO. Anal. numér.*, 17(3):249–265, 1983.
- [27] A. Ern and J.-L. Guermond. *Theory and practice of finite elements*, volume 159 of *Applied Mathematical Sciences*. Springer–Verlag New York, 2004.
- [28] V. Ervin, M. Kubacki, W. Layton, M. Moraiti, Z. Si, and C. Trenchea. Partitioned penalty methods for the transport equation in the evolutionary Stokes–Darcy–transport problem. *Numer. Meth. Part. D. E.*, 35(1):349–374, 2019.
- [29] G. N. Gatica, S. Meddahi, and R. Oyarzúa. A conforming mixed finite-element method for the coupling of fluid flow with porous media flow. *IMA J. Numer. Anal.*, 29:86–108, 2009.
- [30] V. Girault, G. Kanschat, and B. Rivière. Error analysis for a monolithic discretization of coupled Darcy and Stokes problems. *J. Numer. Math.*, 22(2):109–142, 2014.
- [31] V. Girault and B. Rivière. DG approximation of coupled Navier–Stokes and Darcy equations by Beaver–Joseph–Saffman interface condition. *SIAM J. Numer. Anal.*, 47(3):2052–2089, 2009.
- [32] J. Guzmán, C.-W. Shu, and F. Sequeira. H(div) conforming and DG methods for incompressible Euler’s equations. *IMA J. Numer. Anal.*, 37(4):1733–1771, 2016.
- [33] P. Hansbo and M. G. Larson. Discontinuous Galerkin methods for incompressible and nearly incompressible elasticity by Nitsche’s method. *Comput. Methods Appl. Mech. Engrg.*, 191:1895–1908, 2002.
- [34] N. Hanspal, A. Waghode, V. Nassehi, and R. Wakeman. Numerical analysis of coupled stokes/darcy flows in industrial filtrations. *Transport in Porous Media*, 64(1):1573–1634, 2006.
- [35] G. Kanschat and B. Rivière. A strongly conservative finite element method for the coupling of Stokes and Darcy flow. *J. Comput. Phys.*, 229(17):5933–5943, 2010.
- [36] W. Layton. *Introduction to the Numerical Analysis of Incompressible Viscous Flows*. Society for Industrial and Applied Mathematics, Philadelphia, PA, 2008.
- [37] W. Layton, F. Schieweck, and I. Yotov. Coupling fluid flow with porous media flow. *SIAM J. Numer. Anal.*, 40(6):2195–2218, 2002.
- [38] J. Lohrenz, B. G. Bray, and C. R. Clark. Calculating viscosities of reservoir fluids from their compositions. *J. Petrol. Technol.*, 16:1171–1176, 1964.
- [39] A. Márquez, S. Meddahi, and F. J. Sayas. Strong coupling of finite element methods for the Stokes–Darcy problem. *IMA J. Numer. Anal.*, 35(2):969–988, 2015.
- [40] N. C. Nguyen, J. Peraire, and B. Cockburn. An implicit high-order hybridizable discontinuous Galerkin method for linear convection-diffusion equations. *J. Comput. Phys.*, 228(9):3232–3254, 2009.
- [41] B. Rivière. Analysis of a discontinuous finite element method for the coupled Stokes and Darcy problems. *J. Sci. Comput.*, 22(1):479–500, 2005.
- [42] B. Rivière. *Discontinuous Galerkin methods for solving elliptic and parabolic equations*, volume 35 of *Frontiers in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, 2008.
- [43] B. Riviere. Discontinuous finite element methods for coupled surface–subsurface flow and transport problems. In X. Feng, O. Karakashian, and Y. Xing, editors, *Recent Developments in Discontinuous Galerkin Finite Element Methods for Partial Differential Equations: 2012 John H Barrett Memorial Lectures*, pages 259–279. Springer International Publishing, Cham, 2014.
- [44] B. Rivière and N. J. Walkington. Convergence of a discontinuous Galerkin method for the miscible displacement equation under low regularity. *SIAM J. NUMER. ANAL.*, 49(3):1085–1110, 2011.
- [45] H. Rui and J. Zhang. A stabilized mixed finite element method for coupled Stokes and Darcy flows with transport. *Comput. Methods Appl. Mech. Engrg.*, 315:169–189, 2017.
- [46] P. Saffman. On the boundary condition at the surface of a porous media. *Stud. Appl. Math.*, 50:292–315, 1971.
- [47] J. Schöberl. NETGEN an advancing front 2D/3D-mesh generator based on abstract rules. *Computing and Visualization in Science*, 1:41–52, 1997.

- [48] J. Schöberl. C++11 implementation of finite elements in NGSolve. Technical Report ASC Report 30/2014, Institute for Analysis and Scientific Computing, Vienna University of Technology, 2014.
- [49] S. Sun, B. Rivière, and M. F. Wheeler. A combined mixed finite element and discontinuous Galerkin method for miscible displacement problem in porous media. In *Recent Progress in Computational and Applied PDES*, pages 323–351, Boston, MA, 2002. Springer US.
- [50] S. Sun and M. F. Wheeler. Discontinuous Galerkin methods for coupled flow and reactive transport problems. *Applied Numerical Mathematics*, 52(2):273–298, 2005.
- [51] D. Vassilev and I. Yotov. Coupling Stokes–Darcy flow with transport. *SIAM J. Sci. Comput.*, 31(5):3661–3684, 2009.
- [52] G. N. Wells. Analysis of an interface stabilized finite element method: the advection-diffusion-reaction equation. *SIAM J. Numer. Anal.*, 49(1):87–109, 2011.