#### 1 Preliminaries

#### **Operator-norm Trotter product formula on Banach spaces**

#### Valentin A. Zagrebnov

CNRS-Université d'Aix-Marseille Institut de Mathématiques de Marseille CMI - Technopôle Château-Gombert 39, rue F. Joliot Curie 13453 Marseille Cedex 13, France

> In memory of Academician Vasilii Sergeevich Vladimirov on the 100<sup>th</sup> anniversary of his birthday

#### Abstract

In this paper we collect results concerning the *operator-norm* convergent *Trotter* product formula for two semigroups  $\{e^{-tA}\}_{t\geq 0}$ ,  $\{e^{-tB}\}_{t\geq 0}$ , with densely defined generators *A* and *B* in a *Banach* space. Although the *strong* convergence in Banach space for contraction semigroups is known since the seminal paper by Trotter (1959), which after more than three decades was extended to convergence in the *operator-norm* topology in *Hilbert* spaces by Rogava (1993), the operator-norm convergence in a *Banach* space was established only in (2001).

For the first time this result was established under hypothesis that one of the involved into the product formula contraction semigroups, e.g.  $\{e^{-tA}\}_{t\geq 0}$ , is *holomorphic* together with certain conditions of *smallness* on generators *B* and *B*<sup>\*</sup> with respect to generators *A* and *A*<sup>\*</sup>. Note that in spite of a quite strong assumptions on operators *A* and *B* the proof of the operator-norm convergent Trotter product formula on a Banach space is (unexpectedly) involved and technical.

To elucidate the question of how far these conditions are from optimal ones we show an Example of the *operator-norm* convergent Trotter product formula for two semigroups  $\{e^{-tA}\}_{t\geq 0}$  and  $\{e^{-tB}\}_{t\geq 0}$  on a Banach space, where hypothesis on operator *A* is relaxed to condition that *A* is generator of a *contraction* semigroup.

#### **1** Preliminaries

## 1.1 Bounded semigroups on X

For what follows the properties of holomorphic (contraction) semigroups on a Banach space  $\mathfrak{X}$  are essential. Therefore, we start by a suitable for our aim recall of details concerning the bounded, holomorphic semigroups, and fractional powers of their generators. We begin with definitions and properties to introduce certain notations adapted in this section for semigroups on  $\mathfrak{X}$ .

**Definition 1.1.** We would remind that a family  $\{U(t)\}_{t\geq 0}$  of bounded linear operators on a Banach space  $\mathfrak{X}$  is called a one-parameter strongly continuous semigroup if it satisfies the conditions:

(i) U(0) = 1,

(ii) U(s + t) = U(s)U(t) for all  $s, t \ge 0$ , (iii)  $\lim_{t\to+0} U(t)x = x$  for all  $x \in \mathfrak{X}$ .

We recall some immediate consequences of this definition:

- There are constants  $C_A \ge 1$  and  $\gamma_A \in \mathbb{R}$ , depending on the generator of the semigroup, such that  $||U(t)|| \le C_A e^{\gamma_A t}$  for all  $t \ge 0$ .
- t → U(t) is a strongly continuous function from [0, +∞) onto the algebra L(X) of bounded linear operators on X.
- There exists a closed densely defined linear operator A on  $\mathfrak{X}$  with domain dom(A), called the generator of the semigroup, such that  $\lim_{t\to+0} (U(t)x x)/t = -Ax$  for any  $x \in \text{dom}(A)$ , that is, by convention  $U(t) := e^{-tA}$ .
- The resolvent of the generator satisfies the estimate  $||R_A(-\lambda)|| = ||(A + \lambda \mathbb{1})^{-1}|| \le C_A/(\operatorname{Re}(\lambda) \gamma_A)$  for all  $\lambda$  with  $\operatorname{Re}(\lambda) > \gamma_A$ , thus the open half plane with  $\operatorname{Re}(z) < -\gamma_A$  is contained into the resolvent set of *A*, which is defined as  $\rho(A) = \{z \in \mathbb{C} : ||R_A(z)|| < +\infty\}$ .
- If γ<sub>A</sub> ≤ 0, U(t) is called a bounded semigroup (otherwise, U(t) is called a quasibounded semigroup of type γ<sub>A</sub> > 0). For any strongly continuous semigroup, we can construct a bounded semigroup by adding some constant η ≥ γ<sub>A</sub> to its generator. Let à = A + η1, then for the semigroup Q̃(t) generated by Ã, one has ||Q̃(t)|| ≤ C<sub>A</sub>, t ≥ 0, and the open half-plane Re(λ) < η γ<sub>A</sub> is included into the resolvent set ρ(Ã) of Ã. So it is not restrictive to suppose that the considered semigroup U(t) is bounded and that the set {z ∈ C : Re(z) ≤ 0} ⊆ ρ(A).
- If ||U(t)|| ≤ 1, t ≥ 0, the semigroup is called a contraction semigroup. We comment that the method of the preceding remark does not permit to construct a contraction semigroup from a bounded semigroup in general, since it can not change the value of the constant C<sub>A</sub>.

Below we need a characterization of generators of these contraction semigroups. First we recall that the space of linear bounded functionals  $\mathfrak{X}^* = \mathcal{L}(\mathfrak{X}, \mathbb{C})$  is a *dual* of the Banach space  $\mathfrak{X}$  and that  $\mathfrak{X}^*$  is itself a Banach space. Recall that a linear operator A in  $\mathfrak{X}$  is accretive if for all pairs  $(u, \phi) \in \text{dom}(A) \times \mathfrak{X}^*$  with  $||u||_{\mathfrak{X}} = 1$ ,  $||\phi||_{\mathfrak{X}^*} = 1$ ,  $(u, \phi) = 1$ , one has Re $(Au, \phi) \ge 0$ . We also add that a densely defined in  $\mathfrak{X}$  accretive operator A is generator of contraction semigroup if the range of  $\lambda \mathbb{1} + A$  is  $\mathfrak{X}$  for some  $\lambda > 0$ ,

Now we prove a series of estimates indispensable throughout this paper.

**Lemma 1.2.** Let U(t) be a bounded semigroup with boundedly invertible generator *A*, then for all  $t \ge 0$ , and for any  $n \in \mathbb{N}$ , we have:

1 Preliminaries

$$\left(U(t) - \sum_{k=0}^{n} \frac{(-tA)^{k}}{k!}\right) A^{-n-1} = -\int_{0}^{t} d\tau \left(U(\tau) - \sum_{k=0}^{n-1} \frac{(-\tau A)^{k}}{k!}\right) A^{-n}, \quad (1.1)$$

$$\left\| \left( U(t) - \sum_{k=0}^{n} \frac{(-tA)^{k}}{k!} \right) A^{-n-1} \right\| \leq C_{A} \frac{t^{n+1}}{(n+1)!}.$$
 (1.2)

Proof. We proceed by induction, and we first prove that

$$(U(t) - 1)x = -\int_0^t d\tau \, U(\tau) \, A \, x \, , \ x \in \text{dom}(A).$$
(1.3)

Note that for any  $\epsilon > 0$  the semigroup properties yields the representation

$$\int_0^t ds \, U(s) \, U(\epsilon) - \frac{1}{\epsilon} = \int_0^t ds \, U(s+\epsilon) - \frac{U(s)}{\epsilon}$$
$$= \int_t^{t+\epsilon} ds \, \frac{U(s)}{\epsilon} - \int_0^\epsilon ds \, \frac{U(s)}{\epsilon}$$
$$= (U(t) - \mathbb{1}) \frac{1}{\epsilon} \int_0^\epsilon ds \, U(s) \, .$$

Moreover, one also gets:

$$\lim_{\epsilon \to 0} \frac{1}{\epsilon} \int_0^{\epsilon} ds \ U(s) \ x = x \ , \quad x \in \mathfrak{X} \ ,$$
$$\lim_{\epsilon \to 0} \frac{U(\epsilon) - \mathbb{1}}{\epsilon} \ x = -A \ x \ , \quad x \in \operatorname{dom}(A) \ .$$

This proves (1.3), and since A is boundedly invertible, we obtain (1.1) for n = 0. Furthermore, since U(t) is bounded by  $C_A$ , we obtain the estimate (1.2) for n = 0.

Suppose that (1.1) and (1.2) are true for some *n*, then a straightforward calculation leads to (1.1) for n + 1. Hence, using the representation (1.1) and (1.2) for *n* to estimate the integrand, we obtain (1.2) for n + 1, which completes the proof by induction.

Similarly, we obtain a representation for a restricted development of  $(1 + A)^{-1}$ .

**Lemma 1.3.** Let A be as in Lemma 1.2. Then for any  $n \ge 0$ :

$$(\mathbb{1} + A)^{-1}A^{-n-1} = \left(\sum_{k=0}^{n} (-A)^k\right)A^{-n-1} + (-1)^{n+1}(\mathbb{1} + A)^{-1}.$$
 (1.4)

*Proof.* For n = 0, the representation (1.4) follows from the resolvent formula:

$$(\mathbb{1} + A)^{-1} - A^{-1} = -(\mathbb{1} + A)^{-1}A^{-1}.$$
(1.5)

Suppose that (1.4) holds for an integer n > 1, then:

$$(\mathbb{1} + A)^{-1}A^{-n-2} = \left(\sum_{k=0}^{n} (-A)^k\right)A^{-n-2} + (-1)^{n+1}(\mathbb{1} + A)^{-1}A^{-1}.$$
 (1.6)

Applying (1.5) to the last term of (1.6) we get the representation (1.4) for n + 1, and thus for any *n* by induction.

**Lemma 1.4.** If U(t) is a bounded semigroup with boundedly invertible generator A then :

$$\left\|\frac{1}{t^2}\left((\mathbb{1}+tA)^{-1}-U(t)\right)A^{-2}\right\| \le 3C_A/2, \quad t>0.$$
(1.7)

Proof. By Lemma 1.2 one gets

$$\|(U(t) - 1 + tA) \frac{1}{t^2} A^{-2}\| \le \frac{C_A}{2}.$$
(1.8)

On the other hand by Lemma 1.3, we have

$$\left\| \left( (\mathbb{1} + tA)^{-1} - \mathbb{1} + tA \right) \frac{1}{t^2} A^{-2} \right\| = \left\| (\mathbb{1} + tA)^{-1} \right\| \le C_A .$$
 (1.9)

Here the last estimate follows from  $(\mathbb{1} + tA)^{-1} = (1/t)R_A(-1/t)$  and  $||R_A(-\lambda)|| \le C_A/(\lambda + \delta), \delta \ge 0$ , which is valid for bounded semigroups with boundedly invertible generators. Hence (1.7) follows from (1.8) and (1.9).

## 1.2 Holomorphic contraction semigroups on $\mathfrak{X}$

Now let  $U : z \mapsto U(z)$  be a family of operators with z taking their values in the sector of the complex plane:

$$S_{\theta} = \{ z \in \mathbb{C} : z \neq 0 \text{ and } | \arg(z) | < \theta \}$$

$$(1.10)$$

where  $0 < \theta \le \pi/2$ .

**Definition 1.5.** Recall that the family of operators  $\{U(z)\}_{z \in S_{\theta}}$  is a bounded holomorphic semigroup of semi-angle  $\theta \in (0, \pi/2]$  on a Banach space  $\mathfrak{X}$  if it satisfies the following conditions:

- (i) If  $0 < \epsilon < \theta$ , then  $||U(z)|| \le M_{\epsilon}$  for all  $z \in S_{\theta-\epsilon}$  and some  $M_{\epsilon} < \infty$ .
- (ii)  $U(z_1)U(z_2) = U(z_1 + z_2)$  for all  $z_1, z_2 \in S_{\theta}$ .
- (iii)  $U : z \mapsto U(z)$  is analytic function of  $z \in S_{\theta}$ .
- (iv) If  $x \in \mathfrak{X}$  and  $0 < \epsilon < \theta$ , then  $\lim_{z\to 0} U(z) x = x$  provided  $z \in S_{\theta-\epsilon}$ .

Let  $\sigma(A) = \mathbb{C} \setminus \rho(A)$  denote the spectrum of *A*. It can be used for the following characterisation of the holomorphic semigroup generators, [Kato95], Chapter IX:

**Proposition 1.6.** Operator A in a Banach space  $\mathfrak{X}$  is the generator of a bounded holomorphic semigroup of semi-angle  $\theta \in (0, \pi/2]$  if and only if A is a closed operator with a dense domain dom(A) such that:

1 Preliminaries

$$\sigma(A) \subseteq \left\{ z \in \mathbb{C}, |\arg(z)| \le \frac{\pi}{2} - \theta \right\}, \quad 0 < \theta \le \frac{\pi}{2},$$

and

$$\|(z\,\mathbbm{1}+A)^{-1}\| \le \frac{N_{\epsilon}}{|z|} \quad \text{for} \quad N_{\epsilon} > 0 \;, \; z \in S_{\theta+\pi/2-\epsilon}$$

where  $0 < \epsilon < \theta$ .

For applications it is also useful the following property, which is an alternative characterisation of these kind of semigroups.

**Proposition 1.7.** If U(z) is a bounded holomorphic semigroup of semi-angle  $\theta$  with generator A, then for all  $z \in S_{\theta}$  and  $n \in \mathbb{N}$  one has  $U(z)\mathfrak{X} \subseteq \text{dom}(A^n)$ . Moreover, there are positive constants  $C'_A$ ,  $C^{(n)}_A$  such that for t > 0:

$$\left\|\frac{dU(t)}{dt}\right\| = \|AU(t)\| \le \frac{C'_A}{t} \quad \text{and} \quad \left\|\frac{d^n U(t)}{dt^n}\right\| = \|A^n U(t)\| \le \frac{C^{(n)}_A}{t^n} \,. \tag{1.11}$$

Let  $0 < \theta < \pi/2$ . Then estimates (1.11) are valid for complex argument  $z \in S_{\theta}$  with constants depending on  $\theta$ .

**Remark 1.8.** Similarly to strongly continuous semigroups, a family U(z),  $z \in S_{\theta}$  is called a *quasi-bounded* holomorphic semigroup of semi-angle  $\theta$  if there exists a constant  $\beta > 0$  such that restriction of  $\{e^{-\beta z} U(z)\}_{z \in S_{\theta}}$  to  $\mathbb{R}_{0}^{+}$  is a bounded *C*<sub>0</sub>-semigroup. The class of semigroups that we consider here is restricted to *holomorphic contrac-tion* semigroups.

To this aim we recall definition of this notion below, [Kato95], Chapter IX.

**Definition 1.9.** We say that  $\{U_A(z)\}_{z \in S_{\theta}}$ , is a *holomorphic contraction* semigroup with generator  $A \in \mathscr{H}_c(\theta, 0)$ , if its restriction  $\{U_A(t)\}_{t \ge 0}$  to  $\mathbb{R}_0^+$  is a contraction  $C_0$ -semigroup, that is  $\mathscr{H}_c(\theta, 0) := \mathscr{H}(\theta, 0) \cap \mathscr{G}(1, 0)$ .

Note that this class of semigroups is *not* empty and corresponding generators have the following properties:

(i) Let  $\{U_A(t)\}_{t\geq 0}$ , be a *contraction* semigroup with generator  $A \in \mathscr{G}(1,0)$  in a Banach space  $\mathfrak{X}$ , such that  $U_A(t)\mathfrak{X} \subseteq \operatorname{dom}(A)$  for t > 0. If  $||AU(t)|| \leq M_1 t^{-1}$  for some  $M_1 > 0$  and all t > 0, then there exists  $\theta = \arcsin(eM_1)^{-1}(<\pi/2)$  such that  $U_A(t)$  may be analytically continued to contraction holomorphic semigroup of semi-angle  $\theta$ .

(ii) Let *A* be a sectorial operator in a Hilbert space  $\mathfrak{H}$ , i.e. its numerical range  $W = \{(Au, u) : u \in \text{dom}(A) \text{ and } ||u|| = 1\} \subset S_{\pi/2-\theta} \text{ for } 0 < \theta \leq \pi/2.$  If the operator *A* is closed, then it is generator of the holomorphic contraction semigroup of semi-angle  $\theta$ .

(iii) Let A be a generator of holomorphic semigroup on a Banach space  $\mathfrak{X}$ . If A is accretive, then A generates a holomorphic contraction semigroup.

(iv) If *A* is the generator of a strongly continuous group  $\{U_A(t)\}_{t \in \mathbb{R}}$  of contractions  $||U_A(t)|| \le 1$ , then  $\pm A \in \mathcal{G}(1,0)$ , and  $A^2 \in \mathcal{H}_c(\pi/2,0)$ , [EN00], Corollary II.4.9.

## **1.3 Fractional powers of generators**

We scrutinise in this section some properties of fractional powers of the generators for bounded semigroups in a Banach space, see, e.g., [Yos80], Chapter IX.

Recall that fractional power  $A^{\alpha}$ ,  $0 < \alpha < 1$ , of generator of a bounded  $C_0$ -semigroup U(t) ( $||U(t)|| \le C_A$ ) can be expressed by the integral (when it is well-defined):

$$A^{\alpha}x = \frac{1}{\Gamma(-\alpha)} \int_0^{\infty} d\lambda \,\lambda^{-\alpha-1}(U(\lambda) - \mathbb{1}) \, x \,, \quad x \in \operatorname{dom}(A), \tag{1.12}$$

where  $\Gamma(\cdot)$  is the Gamma-function and  $\lambda^{\alpha}$  is chosen to be positive for  $\lambda > 0$ . Since for any  $x \in \text{dom}(A)$  and  $0 < \alpha \le 1$  the integral (1.12) is convergent,  $\text{dom}(A) \subseteq \text{dom}(A^{\alpha})$ . We set  $A^0 := \mathbb{1}$  and define  $A^{\alpha} = A^{\alpha - [\alpha]}A^{[\alpha]}$  for any  $\alpha > 0$ , where  $[\alpha]$  denotes the *integer* part of  $\alpha$ ,

**Proposition 1.10.** For each  $\alpha \in [0, 1]$ , there exists a constant  $C_{A,\alpha}$ , depending only on  $C_A$  and  $\alpha$ , such that for all  $\mu > 0$ ,

$$\left\|A^{\alpha}(A+\mu\mathbb{1})^{-1}\right\| \le \frac{C_{A,\alpha}}{\mu^{1-\alpha}}.$$
 (1.13)

*Proof.* For  $\alpha = 0$  or  $\alpha = 1$ , the result follows directly from the estimate of the resolvent. Let  $0 < \alpha < 1$  and  $x \in \mathfrak{X}$ . Note that  $\operatorname{ran}(A + \mu \mathbb{1})^{-1} = \operatorname{dom}(A) \subseteq \operatorname{dom}(A^{\alpha})$ . Then

$$A^{\alpha}(A+\mu\mathbb{1})^{-1} x = \frac{1}{\Gamma(-\alpha)} \int_0^\infty d\lambda \, \lambda^{-\alpha-1} (U(\lambda)-\mathbb{1}))(A+\mu\mathbb{1})^{-1} x \,. \tag{1.14}$$

We divide the integral (1.14) into two parts:  $0 < \lambda \le \mu^{-1}$  and  $\lambda > \mu^{-1}$ , and we use the representation (1.3):

$$\begin{split} A^{\alpha}(A+\mu\mathbb{1})^{-1}x &= \frac{1}{\Gamma(-\alpha)} \int_{0}^{\mu^{-1}} d\lambda \,\lambda^{-\alpha-1} \int_{0}^{\lambda} dt \,(-U(t))(\mathbb{1}-\mu(A+\mu\mathbb{1})^{-1})x \\ &+ \frac{1}{\Gamma(-\alpha)} \int_{\mu^{-1}}^{\infty} d\lambda \,\lambda^{-\alpha-1} (U(\lambda)-\mathbb{1})(A+\mu\mathbb{1})^{-1} \,x \,. \end{split}$$

Now by the estimate of the resolvent  $||(A + \mu)^{-1}|| \le C_A/\mu$  for all  $\mu > 0$  one obtains:

$$\begin{split} \|A^{\alpha}(A+\mu\mathbb{1})^{-1}x\| &\leq \frac{C_A(1+C_A)\|x\|}{\Gamma(-\alpha)} \left(\int_0^{\mu^{-1}} d\lambda \,\lambda^{-\alpha} + \frac{1}{\mu} \int_{\mu^{-1}}^{\infty} d\lambda \,\lambda^{-\alpha-1}\right) \\ &\leq \frac{C_A(1+C_A)\mu^{\alpha-1}}{\alpha(1-\alpha)\Gamma(-\alpha)} \|x\|. \end{split}$$

Setting  $C_{A,\alpha} := C_A(1 + C_A)/(\alpha(1 - \alpha)\Gamma(-\alpha))$  we obtain the estimate (1.13).

#### 2 Operator-norm Trotter product formula

Next we recall the following well-known property of the semigroup generator A:

**Lemma 1.11.** dom $((A + \delta \mathbb{1})^{\alpha}) = dom(A^{\alpha})$  for all  $\delta > 0$  and  $0 < \alpha < 1$ .

**Theorem 1.12.** Let  $U_A(t)$  be a bounded holomorphic semigroup with generator A, then for any real  $\alpha > 0$ , we have

$$\sup_{t>0} \|t^{\alpha} A^{\alpha} U_A(t)\| = M_{\alpha} < \infty.$$
(1.15)

*Proof.* Let  $0 < \alpha < 1$ . By dom $(A) \subseteq$  dom $(A^{\alpha})$  one gets dom $(A^{\alpha}U_A(t)) = \mathfrak{X}$ . Hence by (1.12) we have

$$A^{\alpha}U_{A}(t) = \frac{1}{\Gamma(-\alpha)} \int_{0}^{\infty} d\lambda \,\lambda^{-\alpha-1} (U_{A}(t+\lambda) - U_{A}(t)) \,. \tag{1.16}$$

Now we split the integral (1.16) in two parts:  $0 < \lambda \le t$  and  $\lambda > t$ , and we use the estimate of the derivative of the holomorphic semigroup (see Proposition 1.7) to obtain

$$\|U_A(t+\lambda) - U_A(t)\| \le \lambda \sup_{t \le \tau \le t+\lambda} \|\partial_\tau U_A(\tau)\| \le \lambda \frac{C_A}{t}.$$
(1.17)

This leads to the estimate

$$\begin{split} \|A^{\alpha}U_{A}(t)\| &\leq \frac{1}{\Gamma(-\alpha)} \left( \int_{0}^{t} d\lambda \,\lambda^{-\alpha} \,\frac{C_{A}'}{t} + \int_{t}^{\infty} d\lambda \,2 \,C_{A} \,\lambda^{-\alpha-1} \right) \\ &\leq \frac{t^{-\alpha}}{\Gamma(-\alpha)} \left( \frac{C_{A}'}{1-\alpha} + \frac{2 \,C_{A}}{\alpha} \right). \end{split}$$

Therefore one obtains (1.15) for  $0 < \alpha < 1$  by setting  $M_{\alpha} := \Gamma(-\alpha)^{-1} (C'_A/(1-\alpha) + 2C_A/\alpha).$ 

For integer powers  $\alpha$ , (1.15) follows directly from Proposition 1.7. Notice that by Proposition 1.7 ran $(U_A(t)) \subseteq \text{dom}(A^n)$  for t > 0. Then result (1.15) follows for any non integer  $\alpha > 1$ , from: the observation that  $\text{dom}(A^{\alpha} = A^{\alpha - [\alpha]}A^{[\alpha]}) \supset \text{dom}(A^{[\alpha]+1})$ , the representation (1.16), and the estimate (1.11) of derivatives of order  $[\alpha] + 1$ .  $\Box$ 

# 2 Operator-norm Trotter product formula

# 2.1 Perturbation of holomorphic contraction semigroups

In this section we prove the operator-norm convergence of the Trotter product formula in Banach space  $\mathfrak{X}$ . Note that convergence of the abstract version of this formula in the strong operator topology is known since [Trot59]. The proof goes via estimate of the rate of convergence in the case of *small* perturbations of the holomorphic contraction semigroups, cf. [CZ01]. We require that operator A generates a holomorphic contraction semigroup, and that perturbation B satisfies the following hypothesis:

(H1) *B* is generator of a contraction semigroup on  $\mathfrak{X}$ .

(H2) There is a real  $\alpha \in [0, 1)$  such that dom $(A^{\alpha}) \subseteq$  dom(B) and that dom $(A^*) \subseteq$  dom $(B^*)$  adjoint operator in the dual space  $\mathfrak{X}^*$ .

Notice that we can suppose the operator *A* boundedly invertible. If it is not the case, one considers  $A + \eta$  for some  $\eta > 0$ . Then by Lemma 1.11 we have dom( $(A + \eta)^{\alpha}$ ) = dom( $A^{\alpha}$ )  $\subseteq$  dom(*B*).

**Remark 2.1.** We note that assumption (H2) implies that *B* is relatively bounded with respect to *A* with the relative bound equals to *zero*. Indeed, for  $\eta > 0$  by dom $(A + \eta \mathbb{1}) \subseteq \text{dom}(A^{\alpha}) \subseteq \text{dom}(B)$  and by Proposition 1.10, (1.13), one gets (here operator *A* is supposed to be boundedly invertible):

$$\|B(A+\eta\mathbb{1})^{-1}\| \le \|BA^{-\alpha}\| \, \|A^{\alpha}(A+\eta\mathbb{1})^{-1}\| \le \frac{C_{\alpha}}{\eta^{1-\alpha}} \, \|BA^{-\alpha}\|.$$
(2.1)

Since the operators  $A^{\alpha}$  and B are closed, the inclusions in (H2) are equivalent to  $A^{\alpha}$ -boundedness of B and the  $A^*$ -boundedness of  $B^*$ . In particular,  $||BA^{-\alpha}|| \le d$  and  $||B^*A^{*-1}|| \le d'$  for some d, d' > 0. Therefore, for any  $x \in \text{dom}(A) \subseteq \text{dom}(B)$ , we have the estimate

$$||Bx|| \le \frac{C_{\alpha} d}{\eta^{1-\alpha}} ||Ax|| + \eta^{\alpha} C_{\alpha} d ||x||$$
(2.2)

and the relative bound in (2.2) can be made infinitesimally small for the large enough shift parameter  $\eta > 0$ .

For this class of (small) perturbations of holomorphic contraction semigroup with generator  $A \in \mathscr{H}_c(\theta, 0)$  (Definition 1.9) one gets the following result:

**Lemma 2.2.** Let  $\{e^{-zA}\}_{z \in S_{\theta}}$  be a holomorphic contraction semigroup of semi-angle  $\theta$  on  $\mathfrak{X}$  and perturbation B satisfy the hypothesis (H1) and (H2). Then the algebraic sum A + B of operators defined on dom(A + B) = dom(A) is also a generator of holomorphic contraction semigroup with the same semi-angle, that is,  $(A + B) \in \mathscr{H}_{c}(\theta, 0)$ .

*Proof.* To this end we verify conditions of Proposition 1.6. Let  $\epsilon \in (0, \theta)$ . Then by (2.2) we obtain inequality

$$\|B(A+z\mathbb{1})^{-1}\| \le \frac{C_{\alpha} \|BA^{-\alpha}\|}{\eta^{1-\alpha}} \|A(A+z\mathbb{1})^{-1}\| + \eta^{\alpha} C_{\alpha} \|BA^{-\alpha}\| \|(A+z\mathbb{1})^{-1}\|,$$

for  $|\arg(z)| < \theta + \pi/2 - \epsilon$ . Seeing that  $||(z \mathbb{1} + A)^{-1}|| \le N_{\epsilon} |z|^{-1}$  for  $N_{\epsilon} > 0$  and  $z \in S_{\theta + \pi/2 - \epsilon}$ , this inequality leads to

$$\|B(A+z\mathbb{1})^{-1}\| \le \frac{C_{\alpha}\|BA^{-\alpha}\|}{\eta^{1-\alpha}} (1+N_{\epsilon}) + \eta^{\alpha}C_{\alpha}\|BA^{-\alpha}\| \frac{N_{\epsilon}}{|z|},$$
(2.3)

#### 2 Operator-norm Trotter product formula

for  $z \in S_{\theta+\pi/2-\epsilon}$ . Therefore, the Neumann series for  $(A + B + z\mathbb{1})^{-1}$  converges if the right hand side of (2.3) is smaller than 1. Since we can choose  $\eta$  and  $z \in S_{\theta+\pi/2-\epsilon}$  such that the right-hand side of the estimate (2.3) becomes smaller than 1, we obtain:

$$||(A + B + z\mathbb{1})^{-1}|| \le \frac{M}{|z - \gamma|}.$$

Here *M* and  $\gamma$  are some positive constants. Then by Proposition 1.6 we conclude that operator  $(A + B) \in \mathcal{H}(\theta, \gamma)$ , that is, it generates a *quasi-bounded* holomorphic semigroup  $\{U_{A+B}(z)\}_{z \in S_{\theta}}$ .

On the other hand, the conditions of lemma imply that *A* and *B* are *accretive*, thus operator A + B is also accretive. Since for  $\lambda < 0$ , and  $|\lambda|$  sufficiently large  $(|\lambda| > \gamma)$ , the point  $\lambda$  is in the resolvent set  $\rho(A + B)$ , we conclude that  $(A + B) \in \mathcal{G}(1, 0)$  generates a contraction semigroup.

Since by  $(A + B) \in \mathcal{H}(\theta, \gamma)$  this semigroup is also holomorphic, one finally obtains the assertion.

The proof of the main theorem of this section involves three technical lemmata. For the two of them we need *only* that  $B(B^*)$  are  $A(A^*)$ -bounded in the Kato sense, i.e., there are positive constants *a* and *b* such that:

$$x \in \operatorname{dom}(A) \subseteq \operatorname{dom}(B), \ ||Bx|| \le a||Ax|| + b||x||, \tag{2.4}$$

$$\phi \in \text{dom}(A^*) \subseteq \text{dom}(B^*), \ ||B^*\phi|| \le a||A^*\phi|| + b||\phi||.$$
(2.5)

If *A* is boundedly invertible, then we can put b = 0 with the relative bound  $a+b||A^{-1}||$  instead of *a*.

**Lemma 2.3.** Let boundedly invertible A and operator B be generators of bounded semigroups. Let B and B<sup>\*</sup> verify (2.4), (2.5) and suppose that operator H = (A + B) with dom(H) = dom(A) is the boundedly invertible generator of a bounded semigroup. Then there exists constant L<sub>1</sub> such that for all  $\tau \ge 0$ :

$$\left\| A^{-1} \left( e^{-\tau B} e^{-\tau A} - e^{-\tau (A+B)} \right) \right\| \le L_1 \tau,$$
 (2.6)

$$\left( e^{-\tau B} e^{-\tau A} - e^{-\tau (A+B)} \right) A^{-1} \right\| \le L_1 \tau.$$
 (2.7)

Proof. By virtue of the identity

$$A^{-1} \left( e^{-\tau B} e^{-\tau A} - e^{-\tau (A+B)} \right) = A^{-1} \left( e^{-\tau B} - \mathbb{1} \right) e^{-\tau A} + A^{-1} \left( e^{-\tau A} - \mathbb{1} \right) + A^{-1} H H^{-1} \left( \mathbb{1} - e^{-\tau H} \right)$$

and by Lemma 1.2 we get (2.6):

$$\begin{aligned} \left\| A^{-1} \left( e^{-\tau B} e^{-\tau A} - e^{-\tau H} \right) \right\| &\leq \left\| \int_{0}^{\tau} ds A^{-1} B e^{-sB} \right\| \\ &+ \left\| A^{-1} \left( e^{-\tau A} - \mathbb{1} \right) \right\| + \left\| A^{-1} H \right\| \left\| H^{-1} \left( \mathbb{1} - e^{-\tau H} \right) \right\| \\ &\leq \left\| A^{-1} B \right\| C_{B} \tau + C_{A} \tau + \left\| A^{-1} H \right\| C_{H} \tau \end{aligned}$$

where coefficients  $C_B$  and  $C_H$  are defined similarly to  $C_A$  in Subsection 1.1.

Finally we remark that (2.5) implies the boundedness of the closed operator  $A^{-1}B$ , and that  $||A^{-1}H|| \leq ||\mathbb{1} + A^{-1}B|| \leq 1 + a + b||A^{-1}||$ . To prove (2.7) one has to use (2.4), and the same line of reasoning as above to put finally  $L_1 = C_Ba' + C_A + C_H(1 + a')$  where  $a' = a + b||A^{-1}||$ .

**Lemma 2.4.** Let A, B and H = A + B be the same as in Lemma 2.3. Then there exists a constant  $L_2$  such that for all  $\tau \ge 0$ :

$$\left\| A^{-1} \left( e^{-\tau B} e^{-\tau A} - e^{-\tau (A+B)} \right) A^{-1} \right\| \le L_2 \tau^2,$$
(2.8)

$$\left| A^{-1} \left( e^{-\tau A} e^{-\tau B} - e^{-\tau (A+B)} \right) A^{-1} \right\| \le L_2 \tau^2.$$
(2.9)

Proof. By virtue of

$$e^{-\tau B}e^{-\tau A} - e^{-\tau H} = (\mathbb{1} - e^{-\tau B})(\mathbb{1} - e^{-\tau A}) + (e^{-\tau A} - (\mathbb{1} + \tau A)^{-1}) + (e^{-\tau B} - (\mathbb{1} + \tau B)^{-1}) + ((\mathbb{1} + \tau H)^{-1} - e^{-\tau H}) + \tau H (\mathbb{1} + \tau H)^{-1} - \tau A (\mathbb{1} + \tau A)^{-1} - \tau B (\mathbb{1} + \tau B)^{-1}$$

and by identity

$$\begin{split} &A^{-1} \left( \tau H (\mathbbm{1} + \tau H)^{-1} - \tau A (\mathbbm{1} + \tau A)^{-1} - \tau B (\mathbbm{1} + \tau B)^{-1} \right) A^{-1} \\ &= \tau^2 \left( (\mathbbm{1} + \tau A)^{-1} + A^{-1} B (\mathbbm{1} + \tau B)^{-1} B A^{-1} - A^{-1} H (\mathbbm{1} + \tau H)^{-1} H A^{-1} \right), \end{split}$$

we obtain the representation

$$\begin{split} A^{-1}(\mathrm{e}^{-\tau B}\mathrm{e}^{-\tau A} - \mathrm{e}^{-\tau H})A^{-1} &= \\ A^{-1}(\mathbbm{1} - \mathrm{e}^{-\tau B})(\mathbbm{1} - \mathrm{e}^{-\tau A})A^{-1} \\ &+ \left(\mathrm{e}^{-\tau A} - (\mathbbm{1} + \tau A)^{-1}\right)A^{-2} + A^{-1}\left(\mathrm{e}^{-\tau B} - (\mathbbm{1} + \tau B)^{-1}\right)A^{-1} \\ &+ A^{-1}H\left((\mathbbm{1} + \tau H)^{-1} - \mathrm{e}^{-\tau H}\right)H^{-2}HA^{-1} + \tau^{2}(\mathbbm{1} + \tau A)^{-1} \\ &+ \tau^{2}A^{-1}B(\mathbbm{1} + \tau B)^{-1}BA^{-1} - \tau^{2}A^{-1}H(\mathbbm{1} + \tau H)^{-1}HA^{-1} \;. \end{split}$$

This presentation yields the following estimate:

#### 2 Operator-norm Trotter product formula

$$\begin{aligned} \frac{1}{\tau^2} \left\| A^{-1} (e^{-\tau B} e^{-\tau A} - e^{-\tau H}) A^{-1} \right\| &\leq \\ & \frac{1}{\tau} \left\| A^{-1} B \right\| \left\| \int_0^{\tau} ds \ e^{-sB} \right\| \frac{1}{\tau} \left\| (\mathbb{1} - e^{-\tau A}) A^{-1} \right\| \\ & + \frac{1}{\tau^2} \left\| \left( e^{-\tau A} - (\mathbb{1} + \tau A)^{-1} \right) A^{-2} \right\| \\ & + \frac{1}{\tau^2} \left\| A^{-1} B \right\| \left\| \int_0^{\tau} ds_1 \int_0^{s_1} ds_2 \ e^{-s_2 B} - \tau^2 (\mathbb{1} + \tau B)^{-1} \right\| \left\| BA^{-1} \right\| \\ & + \frac{1}{\tau^2} \left\| A^{-1} H \right\| \left\| \left( (\mathbb{1} + \tau H)^{-1} - e^{-\tau H} \right) H^{-2} \right\| \left\| HA^{-1} \right\| \\ & + 1 + \left\| A^{-1} B \right\| \left\| BA^{-1} \right\| + \left\| A^{-1} H \right\| \left\| HA^{-1} \right\|. \end{aligned}$$

Now Lemmata 1.2 and 1.4 together with (2.4), (2.5) imply (2.8), where we can take  $L_2 = a'C_AC_B + 3C_A/2 + 3C_Ba'^2/2 + 3C_H(1 + a')^2/2 + 1 + a'^2 + (1 + a')^2$  with  $a' = a + b||A^{-1}||$ . Similarly one obtains (2.9).

Note that for the proof of the third lemma (Lemma 2.4) we do need *conditions* (H2), as well as requirement that semigroups are *contractive*.

**Lemma 2.5.** Let A be a boundedly invertible generator of holomorphic contraction semigroup. If B is generator of a contraction semigroup and there exists  $\alpha \in [0, 1)$  such that dom $(A^{\alpha}) \subseteq \text{dom}(B)$ , then for any  $k \ge 1$  and  $\tau > 0$  one gets the estimates

$$\left\| \left( e^{-\tau B} e^{-\tau A} \right)^k A \right\| \le \frac{L_3}{\tau^{\alpha}} + \frac{C'_A}{k\tau} , \qquad \text{if } \alpha > 0 , \qquad (2.10)$$

$$\left\| \left( e^{-\tau B} e^{-\tau A} \right)^k A \right\| \le \tilde{L}_3 (1 + \ln k) + \frac{C'_A}{k\tau} , \text{ if } \alpha = 0 .$$
 (2.11)

*Proof.* We start with the following chain of estimates:

$$\begin{split} \left\| \left( e^{-\tau B} e^{-\tau A} \right)^{k} A \right\| &\leq \left\| \left( \left( e^{-\tau B} e^{-\tau A} \right)^{k} - e^{-k\tau A} \right) A \right\| + \left\| e^{-k\tau A} A \right\| \\ &\leq \left\| \sum_{j=0}^{k-1} \left( e^{-\tau B} e^{-\tau A} \right)^{k-1-j} \left( e^{-\tau B} - I \right) e^{-\tau A} e^{-j\tau A} A \right\| + \left\| e^{-k\tau A} A \right\| \\ &\leq \sum_{j=0}^{k-1} \left\| \int_{0}^{\tau} ds \ e^{-sB} B A^{-\alpha} \right\| \left\| A^{\alpha} e^{-(j+1)\tau A} A \right\| + \left\| e^{-k\tau A} A \right\| . \end{split}$$

Notice that the second inequality is in particular due to contractions of  $e^{-tA}$  and  $e^{-tB}$ , and to equation (1.3) of Lemma 1.2. From the hypothesis dom( $A^{\alpha}$ )  $\subseteq$  dom(B) we deduce that  $||BA^{-\alpha}|| \leq d$ , see Remark 2.1. By Propositions 1.7 and Theorem 1.12 we get respectively:

$$\left\| e^{-k\tau A} A \right\| \le \frac{C'_A}{k\tau} \quad \text{and} \quad \left\| A^{1+\alpha} e^{-(j+1)\tau A} \right\| \le \frac{M_\alpha}{((j+1)\tau)^{1+\alpha}}.$$

Therefore, we conclude that:

$$\left\| \left( \mathrm{e}^{-\tau B} \mathrm{e}^{-\tau A} \right)^k A \right\| \leq \frac{M_\alpha d}{\tau^\alpha} \sum_{j=0}^{k-1} \frac{1}{(j+1)^{1+\alpha}} + \frac{C'_A}{k \, \tau} \, .$$

Since  $\alpha > 0$ , this gives the announced result (2.10) with

$$L_3 = d M_{\alpha} \sum_{j=1}^{\infty} (1/j)^{1+\alpha}$$

and (2.11) for  $\alpha = 0$  with  $\tilde{L}_3 = ||B|| C'_A$ .

Note that since dom( $A^{\alpha}$ )  $\subseteq$  dom(B) implies dom( $A^{\alpha'}$ )  $\subseteq$  dom(B) for  $\alpha' \ge \alpha$ , the estimate (2.10) is valid in fact for any  $\alpha' \ge \alpha$ .

# 2.2 Convergence rate

**Theorem 2.6.** Let  $\{e^{-zA}\}_{z \in S_{\theta}}$  be a holomorphic contraction semigroup, that is,  $A \in \mathcal{H}_{c}(\theta, 0)$ . Let *B* be generator of a contraction semigroup. If there exists  $\alpha \in [0, 1)$  such that dom $(A^{\alpha}) \subseteq$  dom(B) and dom $(A^{*}) \subseteq$  dom $(B^{*})$ , then there are constants  $M_{1}, M_{2}, \tilde{M}_{2}, \eta > 0$ , such that for any  $t \ge 0$  and n > 2 one gets estimates

$$\left\| \left( e^{-tB/n} e^{-tA/n} \right)^n - e^{-t(A+B)} \right\| \le e^{\eta t} (M_1 + M_2 t^{1-\alpha}) \frac{\ln n}{n^{1-\alpha}}, \ \alpha > 0, \qquad (2.12)$$

$$\left\| \left( e^{-tB/n} e^{-tA/n} \right)^n - e^{-t(A+B)} \right\| \le e^{\eta t} (M_1 + \tilde{M}_2 t) \frac{2 (\ln n)^2}{n}, \ \alpha = 0.$$
 (2.13)

*Proof.* Since *B* satisfies hypothesis (H1) and (H2), by Lemma 2.2 the operator H = (A + B) is generator of a holomorphic contraction semigroup. If operator *A* has no bounded inverse, let  $\tilde{A} := A + \eta$  and  $\tilde{H} := \tilde{A} + B$  for some  $\eta > 0$  (see Remark 2.1). Then both operators are boundedly invertible. As we indicated above, these changes of generators do not modify the domain inclusions. If we want to obtain  $||B\tilde{A}^{-1}|| < 1$  then by the estimate (2.1) we have to choose a sufficiently large shift parameter  $\eta > 0$ . This gives us the estimate  $||\tilde{A}\tilde{H}^{-1}|| = ||(\mathbb{1} + B\tilde{A}^{-1})^{-1}|| \le 1/(1 - a)$  where we set  $a = ||B\tilde{A}^{-1}||$ .

Now we put  $\tau := t/n$ ,  $\tilde{U}(t) := e^{-t\tilde{H}}$ , and  $\tilde{T}(\tau) := e^{-\tau B}e^{-\tau \tilde{A}}$ . To estimate the left-hand side of (2.12) we use

$$\left(e^{-tB/n}e^{-tA/n}\right)^n - e^{-t(A+B)} = (\tilde{T}^n(\tau) - \tilde{U}^n(\tau)) e^{t\eta} , \qquad (2.14)$$

and *telescopic* identity:

2 Operator-norm Trotter product formula

$$\begin{split} \tilde{T}(\tau)^{n} &- \tilde{U}(\tau)^{n} = \sum_{m=0}^{n-1} \tilde{T}(\tau)^{n-m-1} (\tilde{T}(\tau) - \tilde{U}(\tau)) \tilde{U}(\tau)^{m} \\ &= \tilde{T}(\tau)^{n-1} \tilde{A} \tilde{A}^{-1} (\tilde{T}(\tau) - \tilde{U}(\tau)) \\ &+ (\tilde{T}(\tau) - \tilde{U}(\tau)) \tilde{A}^{-1} \tilde{A} \tilde{H}^{-1} \tilde{H} \tilde{U}(\tau)^{n-1} \\ &+ \sum_{m=1}^{n-2} \tilde{T}(\tau)^{n-m-1} \tilde{A} \tilde{A}^{-1} (\tilde{T}(\tau) - \tilde{U}(\tau)) \tilde{A}^{-1} \tilde{A} \tilde{H}^{-1} \tilde{H} \tilde{U}(\tau)^{m} , \end{split}$$

which implies

$$\begin{split} \left\| \tilde{T}(\tau)^{n} - \tilde{U}(\tau)^{n} \right\| &\leq \| \tilde{T}(\tau)^{n-1} \tilde{A} \| \| \tilde{A}^{-1} (\tilde{T}(\tau) - \tilde{U}(\tau)) \| \\ &+ \| (\tilde{T}(\tau) - \tilde{U}(\tau)) \tilde{A}^{-1} \| \| \tilde{A} \tilde{H}^{-1} \| \| \tilde{H} \tilde{U}(\tau)^{n-1} \| \\ &+ \sum_{m=1}^{n-2} \| \tilde{T}(\tau)^{n-m-1} \tilde{A} \| \| \tilde{A}^{-1} (\tilde{T}(\tau) - \tilde{U}(\tau)) \tilde{A}^{-1} \| \| \tilde{A} \tilde{H}^{-1} \| \| \tilde{H} \tilde{U}(\tau)^{m} \| \, . \end{split}$$

Hence by Lemmata 2.3, 2.4, and 2.5 (it is at this point that we use the hypothesis of *contraction*), and by Proposition 1.7 we obtain the estimate :

$$\begin{split} \|\tilde{T}(\tau)^{n} - \tilde{U}(\tau)^{n}\| &\leq \left(\frac{L_{3}}{\tau^{\alpha}} + \frac{C'_{A}}{(n-1)\tau}\right) L_{1}\tau + \frac{L_{1}}{1-a}\frac{C'_{H}}{n-1} \\ &+ \sum_{m=1}^{n-2} \left(L_{3}\tau^{1-\alpha} + \frac{C'_{A}}{n-m-1}\right)\frac{L_{2}}{1-a}\frac{C'_{H}}{m} \\ &\leq L_{3}L_{1}\frac{t^{1-\alpha}}{n^{1-\alpha}} + \frac{L_{1}}{n-1}\left(C'_{A} + \frac{C'_{H}}{1-a}\right) + \frac{L_{3}L_{2}C'_{H}}{1-a}\frac{t^{1-\alpha}}{n^{1-\alpha}}\sum_{m=1}^{n-2}\frac{1}{m} \\ &+ \frac{L_{2}C'_{H}C'_{A}}{1-a}\sum_{m=1}^{n-2}\frac{1}{n-m-1}\cdot\frac{1}{m} \\ &\leq L_{3}L_{1}\frac{t^{1-\alpha}}{n^{1-\alpha}} + \frac{L_{1}}{n-1}\left(C'_{A} + \frac{C'_{H}}{1-a}\right) \\ &+ 2\frac{L_{3}L_{2}C'_{H}}{1-a}t^{1-\alpha}\frac{\ln n}{n^{1-\alpha}} + 4\frac{L_{2}C'_{H}C'_{A}}{1-a}\frac{\ln n}{n}. \end{split}$$
(2.15)

Here we used that:

$$\sum_{m=1}^{n-1} \frac{1}{(n-m)m} = \frac{2}{n} \sum_{m=1}^{n-1} \frac{1}{m} \le \frac{2}{n} (1 + \ln(n-1)) \le 4 \frac{\ln n}{n}.$$

The estimate (2.15) and (2.14) imply the announced result (2.12) for  $\alpha > 0$ , with  $M_1 = 4L_1\left(C'_A + \frac{C'_H}{1-a}\right) + 4\frac{L_2C'_HC'_A}{1-a}$  and  $M_2 = 2L_3L_1 + 2\frac{L_3L_2C'_H}{1-a}$ . In a similar way one gets also estimate for  $\alpha = 0$ :

$$\begin{split} \|\tilde{T}(\tau)^n - \tilde{U}(\tau)^n\| &\leq \tilde{L}_3(1 + \ln(n-1)) L_1 \frac{t}{n} + \frac{L_1 C'_A}{n-1} + \frac{L_1 C'_H}{1-a} \frac{1}{n-1} \\ &+ \sum_{m=1}^{n-2} \left( \tilde{L}_3 \frac{t}{n} (1 + \ln(n-m-1)) + \frac{C'_A}{n-m-1} \right) \frac{L_2}{1-a} \frac{C'_H}{m} \end{split}$$

This estimate together with (2.14) yield (2.13) for  $\tilde{M}_2 = 2 \tilde{L}_3 L_1 + 2 \frac{\tilde{L}_3 L_2 C'_H}{1-a}$ .

**Theorem 2.7.** Let  $\{e^{-zA}\}_{z \in S_{\theta}}$  be a holomorphic contraction semigroup, that is,  $A \in \mathscr{H}_{c}(\theta, 0)$ . Let *B* be generator of a contraction semigroup, and there exists  $\alpha \in [0, 1)$  such that dom $((A^{\alpha})^{*}) \subseteq \text{dom}(B^{*})$  and dom $(A) \subseteq \text{dom}(B)$ . If in addition dom $(A^{*}) \subseteq \text{dom}(B^{*})$  (for the case, when the space  $\mathfrak{X}$  is not reflexive), then there are constants  $M_{3}, M_{4}, \tilde{M}_{4}, \eta > 0$ , such that for any  $t \ge 0$  and n > 2:

$$\left\| \left( e^{-tA/n} e^{-tB/n} \right)^n - e^{-t(A+B)} \right\| \le e^{\eta t} (M_3 + M_4 t^{1-\alpha}) \frac{\ln n}{n^{1-\alpha}}, \ \alpha > 0, \qquad (2.16)$$

$$\left\| \left( e^{-tA/n} e^{-tB/n} \right)^n - e^{-t(A+B)} \right\| \le e^{\eta t} (M_3 + \tilde{M}_4 t) \frac{2 (\ln n)^2}{n}, \ \alpha = 0,$$
 (2.17)

for any  $t \ge 0$  and n > 2.

*Proof.* Let  $\tilde{F}(\tau) := e^{-\tau \tilde{A}} e^{-\tau B}$ . Then by the same arguments as in the proof of Theorem 2.6, one obtains:

$$\begin{split} \tilde{U}(\tau)^{n} &- \tilde{F}(\tau)^{n} = \sum_{m=0}^{n-1} \tilde{U}(\tau)^{n-m-1} (\tilde{U}(\tau) - \tilde{F}(\tau)) \tilde{F}(\tau)^{m} \\ &= \tilde{U}(\tau)^{n-1} \tilde{H} \tilde{H}^{-1} \tilde{A} \tilde{A}^{-1} (\tilde{U}(\tau) - \tilde{F}(\tau)) \\ &+ (\tilde{U}(\tau) - \tilde{F}(\tau)) \tilde{A}^{-1} \tilde{A} \tilde{F}(\tau)^{n-1} \\ &+ \sum_{m=1}^{n-2} \tilde{U}(\tau)^{n-m-1} \tilde{H} \tilde{H}^{-1} \tilde{A} \tilde{A}^{-1} (\tilde{U}(\tau) - \tilde{F}(\tau)) \tilde{A}^{-1} \tilde{A} \tilde{F}(\tau)^{m} \end{split}$$

Notice that the Lemmata 2.3 and 2.4 hold for  $\tilde{F}(\tau)$ . By a simple modification of Lemma 2.5, where one uses  $\|\tilde{A}^{-\alpha}B\| = \|B^*(\tilde{A}^{-\alpha})^*\| < \infty$ , we find that

$$\begin{split} \left\| \tilde{A} \left( e^{-\tau \tilde{A}} e^{-\tau B} \right)^k \right\| &\leq \frac{L_4}{\tau^{\alpha}} + \frac{C'_A}{k \tau} , \qquad \alpha > 0, \\ \left\| \tilde{A} \left( e^{-\tau \tilde{A}} e^{-\tau B} \right)^k \right\| &\leq \tilde{L}_4 (1 + \ln k) + \frac{C'_A}{k \tau} , \quad \alpha = 0. \end{split}$$

These ingredients ensure the estimates (2.16) and (2.17).

**Corollary 2.8.** Under the same conditions as in Theorem 2.6, we have the operatornorm convergence of the symmetrised Trotter formula, i.e., there exists  $M_5$ ,  $M_6$ ,  $\tilde{M}_6$ ,  $\eta > 0$ , such that for any  $t \ge 0$  and n > 2:

$$\left\| \left( e^{-tA/2n} e^{-tB/n} e^{-tA/2n} \right)^n - e^{-t(A+B)} \right\| \le e^{\eta t} \left( M_5 + M_6 t^{1-\alpha} \right) \frac{\ln n}{n^{1-\alpha}},$$
  
for  $0 < \alpha < 1$ , (2.18)  
$$\left\| \left( e^{-tA/2n} e^{-tB/n} e^{-tA/2n} \right)^n - e^{-t(A+B)} \right\| \le e^{\eta t} \left( M_5 + \tilde{M}_6 t \right) \frac{2 \left( \ln n \right)^2}{n},$$

for 
$$\alpha = 0$$
. (2.19)

*Proof.* Since Lemmata 2.3, 2.4, and 2.5 can be easily extended to the symmetrized product  $e^{-\tau A/2}e^{-\tau B}e^{-\tau A/2}$ , the proof of the Theorem 2.6 carries through verbatim to obtain (2.18) and (2.19).

**Remark 2.9.** Seeing that in Theorems 2.6, 2.7 and in Corollary 2.8 the *perturbation B* of *dominating* operator *A* is either *infinitesimally A*-small, or simply *bounded*, the corresponding results in Banach space  $\mathfrak{X}$  are *weaker* than those in Hilbert space  $\mathfrak{H}$ , see [Rog93], [NZ98], [NZ99], and [Zag20]. Recall that in [Rog93], [NZ98] the perturbation *B* in  $\mathfrak{H}$  is *Kato-small* with respect to operator *A* for relative bound *b* < 1. The *fractional* condition (H2) in a Hilbert space  $\mathfrak{H}$  was introduced in [IT97]. Note that in the both cases:  $\mathfrak{X}$  and  $\mathfrak{H}$ , the *dominating* operator *A* is supposed to be generator of a *holomorphic* semigroup.

#### **3 Example**

Resuming Remark 2.9, the question arises: whether the Trotter product formula converges in the *operator-norm* topology if the condition on dominating generator  $A \in \mathscr{H}_c(\theta, 0)$  is relaxed to hypothesis that  $A \in \mathscr{G}(1, 0)$ , i.e., it is generator of a *contraction* (but *not* holomorphic!) semigroup and *B* is a *bounded* generator ?

The aim of this section is to give an answer to this question using *example* of a certain class of generators and semigroups. It turns out that appropriate for this purpose is the class of generators of *evolution semigroups*.

## 3.1 Evolution semigroups and Trotter product formula

To proceed further we need some key notions from the *evolution semigroups* theory and in particular the notion of *solution operator*.

A strongly continuous mapping  $U(\cdot, \cdot) : \Delta \longrightarrow \mathcal{L}(\mathfrak{X})$ , where domain  $\Delta := \{(t, s) : 0 < s \le t \le T\}$  and  $\mathcal{L}(\mathfrak{X})$  is the set of bounded operators on *separable* Banach space  $\mathfrak{X}$ , is called a *solution operator* if the conditions

$$\begin{split} \text{(i)} & \sup_{(t,s)\in\varDelta} \|U(t,s)\|_{\mathcal{L}(\mathfrak{X})} < \infty \;, \\ \text{(ii)} \; U(t,s) &= U(t,r)U(r,s) \;, \;\; 0 < s \leq r \leq t \leq T \;, \end{split}$$

are satisfied. Let us consider the Banach space  $L^p(\mathcal{I}, \mathfrak{X})$  for  $\mathcal{I} := [0, T]$  and  $p \in [1, \infty)$ . The operator  $\mathcal{K}$  is an *evolution generator* of the evolution semigroup  $\{\mathcal{U}(\tau) := e^{-\tau \mathcal{K}}\}_{\tau \ge 0}$  if there is a solution operator such that the *Howland-Evans-Neidhardt* representation, see [How74], [Ev76], [Nei79] and [Nei81]:

$$(e^{-\tau \mathcal{K}} f)(t) = U(t, t - \tau) \chi_{I}(t - \tau) f(t - \tau), \quad f \in L^{p}(I, \mathfrak{X}),$$
(3.1)

holds for a.a.  $t \in I$  and  $\tau \ge 0$ . Seeing that on account of (3.1) the semigroup  $\{e^{-\tau \mathcal{K}}\}_{\tau \ge 0}$  is *nilpotent*:  $e^{-\tau \mathcal{K}}f = 0$  for  $\tau \ge T$ , the evolution generator  $\mathcal{K}$  can never be generator of a holomorphic semigroup.

A simple example of an evolution generator is the differentiation operator, cf. [NSZ20]:

$$(D_0 f)(t) := \partial_t f(t),$$
  

$$f \in \text{dom}(D_0) := \{ f \in W^{1,p}(\mathcal{I}, \mathfrak{X}) : f(0) = 0 \},$$
(3.2)

where  $W^{1,p}(\mathcal{I}, \mathfrak{X})$  is the *Sobolev* space of order (1, p) of *Bochner p*-integrable functions. Then by (3.2) one obviously gets the contraction shift semigroup:

$$(e^{-\tau D_0} f)(t) = \chi_I(t-\tau) f(t-\tau), \quad f \in L^p(I, \mathfrak{X}),$$
(3.3)

for a.a.  $t \in I$  and  $\tau \ge 0$ . Hence, (3.1) implies that the corresponding solution operator of the *non-holomorphic* evolution semigroup  $\{e^{-\tau D_0}\}_{\tau\ge 0}$  is given by  $U_{D_0}(t, s) = \mathbb{1}$ , for all  $(t, s) \in \Delta$ .

Below we consider the operator  $\mathcal{K}_0 := \overline{D_0 + \mathcal{A}}$ , where  $\mathcal{A}$  is the *multiplication* operator induced by generator A of a *holomorphic contraction* semigroup on  $\mathfrak{X}$ . More precisely

$$(\mathcal{A}f)(t) := Af(t), \text{ and } (e^{-\tau \mathcal{A}}f)(t) = e^{-\tau A}f(t),$$
  
$$f \in \operatorname{dom}(\mathcal{A}) := \{f \in L^p(\mathcal{I}, \mathfrak{X}) : Af \in L^p(\mathcal{I}, \mathfrak{X})\}.$$

Then the perturbation of the shift semigroup (3.3) by  $\mathcal{A}$  corresponds to the semigroup with generator  $\mathcal{K}_0$ . One easily checks that  $\mathcal{K}_0$  is an evolution generator of a contraction semigroup on  $L^p(\mathcal{I}, \mathfrak{X})$ , that is never holomorphic [NSZ20]. Indeed, since the generators  $D_0$  and  $\mathcal{A}$  commute, the representation (3.1) for evolution semigroup  $\{e^{-\tau \mathcal{K}_0}\}_{\tau \geq 0}$  takes the form:

$$(\mathrm{e}^{-\tau \mathcal{K}_0} f)(t) = \mathrm{e}^{-\tau A} \chi_{\mathcal{I}}(t-\tau) f(t-\tau), \quad f \in L^p(\mathcal{I}, \mathfrak{X}),$$

for a.a.  $t \in \mathcal{I} = [0, T]$  and  $\tau \ge 0$ . Then by (3.1) the solution operator  $U_0(t, s) = e^{-(t-s)A}$ . Therefore,  $e^{-\tau \mathcal{K}_0} f = 0$  for  $\tau \ge T$ , that is, semigroup  $\{e^{-\tau \mathcal{K}_0}\}_{\tau \ge 0}$  is *nilpotent*.

Furthermore, if  $\{B(t)\}_{t \in I}$  is a *strongly measurable* family of generators of contraction semigroups on  $\mathfrak{X}$ , that is,  $B(\cdot) : I \longrightarrow \mathcal{G}(1,0)$ , then the induced *multiplication* operator  $\mathcal{B}$ :

f

$$(\mathcal{B}f)(t) := B(t)f(t) , \qquad (3.4)$$
  

$$\in \operatorname{dom}(\mathcal{B}) := \left\{ f \in L^p(\mathcal{I}, \mathfrak{X}) : \begin{array}{c} f(t) \in \operatorname{dom}(B(t)) & \text{for a.a. } t \in \mathcal{I} \\ B(t)f(t) \in L^p(\mathcal{I}, \mathfrak{X}) \end{array} \right\} ,$$

is a generator of a contraction semigroup on  $L^p(\mathcal{I}, \mathfrak{X})$ .

In the next Subsection 3.2 we consider perturbation of generator  $\mathcal{K}_0$  by multiplication operator  $\mathcal{B}$  (3.4). Thereupon we construct by means of the *Trotter* product formula approach a corresponding perturbed semigroup.

**Remark 3.1.** We conclude by remarks concerning some *notations* and *definitions* that we use below throughout Section 3.

1. For characterisation the rate of convergence we use, so-called, *Landau's* symbols:

$$g(n) = O(f(n)) \iff \limsup_{n \to \infty} \left| \frac{g(n)}{f(n)} \right| < \infty ,$$
  

$$g(n) = o(f(n)) \iff \limsup_{n \to \infty} \left| \frac{g(n)}{f(n)} \right| = 0 ,$$
  

$$g(n) = \Theta(f(n)) \iff 0 < \liminf_{n \to \infty} \left| \frac{g(n)}{f(n)} \right| \le \limsup_{n \to \infty} \left| \frac{g(n)}{f(n)} \right| < \infty ,$$
  

$$g(n) = \omega(f(n)) \iff \limsup_{n \to \infty} \left| \frac{g(n)}{f(n)} \right| = \infty .$$

2. We also use the notation  $C^{0,\beta}(I) := \{f : I \to \mathbb{C} : \text{there exists some } K > 0 \text{ such that } |f(x) - f(y)| \le K |x - y|^{\beta}, \text{ for any } x, y \in I \text{ and } \beta \in (0, 1] \}.$ 

3. Below we consider the Banach space  $L^p(\mathcal{I}, \mathfrak{X})$  for  $\mathcal{I} := [0, T]$  and  $p \in [1, \infty)$ .

## 3.2 Trotter product formula

We reminisce (cf. Subsection 3.1) that semigroup  $\{\mathcal{U}(\tau)\}_{\tau\geq 0}$ , on the Banach space  $L^p(\mathcal{I}, \mathfrak{X})$  is called the *evolution semigroup* if there is a *solution operator*:  $\{U(t, s)\}_{(t,s)\in\mathcal{A}}$ , such that representation (3.1) holds.

Let  $\mathcal{K}_0$  be the generator of an evolution semigroup  $\{\mathcal{U}_0(\tau)\}_{\tau \geq 0}$  and let  $\mathcal{B}$  be a multiplication operator induced by a measurable family  $\{B(t)\}_{t \in I}$  of generators of contraction semigroups. Note that in this case the multiplication operator  $\mathcal{B}$  (3.4) is a generator of a contraction semigroup  $(e^{-\tau \mathcal{B}} f)(t) = e^{-\tau B(t)} f(t)$ , on the Banach space  $L^p(\mathcal{I}, \mathfrak{X})$ . Since  $\{\mathcal{U}_0(\tau)\}_{\tau \geq 0}$  is an evolution semigroup, then by definition (3.1) there is a propagator  $\{U_0(t, s)\}_{(t,s) \in \mathcal{A}}$  such that the representation

$$(\mathcal{U}_0(\tau)f)(t) = U_0(t, t-\tau)\chi_I(t-\tau)f(t-\tau), \quad f \in L^p(I, \mathfrak{X}),$$

is valid for a.a.  $t \in \mathcal{I}$  and  $\tau \ge 0$ . Then we define  $\tau_n := (t - s)/n$ , for  $n \in \mathbb{N}$ , and

$$G_{j}(t,s;n) := U_{0}(s+j\tau_{n}, s+(j-1)\tau_{n}) e^{-\tau_{n}B(s+(j-1)\tau_{n})}, \quad (t,s) \in \varDelta,$$

where  $j \in \{1, 2, ..., n\}, n \in \mathbb{N}, (t, s) \in \Delta$ , and we set

$$V_n(t,s) := \prod_{j=n}^{1} G_j(t,s;n), \quad n \in \mathbb{N}, \quad (t,s) \in \varDelta.$$

That is, the product is increasingly ordered in j from the right to the left. Then a straightforward computation shows that the representation

$$\left(\left(e^{-\tau\mathcal{K}_0/n}e^{-\tau\mathcal{B}/n}\right)^n f\right)(t) = V_n(t,t-\tau)\chi_I(t-\tau)f(t-\tau), \qquad (3.5)$$

 $f \in L^p(\mathcal{I}, \mathfrak{X})$ , holds for each  $\tau \ge 0$  and a.a.  $t \in \mathcal{I}$ .

**Theorem 3.2.** Let  $\mathcal{K}$  and  $\mathcal{K}_0$  be generators of evolution semigroups on the Banach space  $L^p(\mathcal{I}, \mathfrak{X})$  for some  $p \in [1, \infty)$ . Further, let  $\{B(t) \in \mathcal{G}(1, 0)\}_{t \in \mathcal{I}}$  be a strongly measurable family of generators of contraction semigroups on  $\mathfrak{X}$ . Then

$$\sup_{\tau \ge 0} \left\| e^{-\tau \mathcal{K}} - \left( e^{-\tau \mathcal{K}_0/n} e^{-\tau \mathcal{B}/n} \right)^n \right\|_{\mathcal{L}(L^p(\mathcal{I},\mathfrak{X}))}$$

$$= \operatorname{ess\,sup}_{(t,s) \in \mathcal{A}} \|U(t,s) - V_n(t,s)\|_{\mathcal{L}(\mathfrak{X})}, \quad n \in \mathbb{N}.$$
(3.6)

*Proof.* Let  $\{L(\tau)\}_{\tau \ge 0}$  be the left-shift semigroup on the Banach space  $L^p(\mathcal{I}, \mathfrak{X})$ :

$$(L(\tau)f)(t) = \chi_{I}(t+\tau)f(t+\tau), \quad f \in L^{p}(I,\mathfrak{X}).$$

Using that we get

$$\begin{aligned} & \left( L(\tau) \left( \mathrm{e}^{-\tau \mathcal{K}} - \left( \mathrm{e}^{-\tau/n \mathcal{K}_0} \mathrm{e}^{-\tau \mathcal{B}/n} \right)^n \right) f \right)(t) \\ &= \left\{ U(t+\tau,t) - V_n(t+\tau,t) \right\} \chi_I(t+\tau) f(t) \;, \end{aligned}$$

for  $\tau \ge 0$  and a.a.  $t \in \mathcal{I}$ .

It turns out that for each  $n \in \mathbb{N}$  the operator  $L(\tau) \left( e^{-\tau \mathcal{K}} - \left( e^{-\tau/n\mathcal{K}_0} e^{-\tau \mathcal{B}/n} \right)^n \right)$  is a multiplication operator induced by  $\{ (U(t + \tau, t) - V_n(t + \tau, t)) \chi_I(t + \tau) \}_{t \in I}$ . As a consequence,

$$\begin{aligned} \left\| L(\tau) \left( e^{-\tau \mathcal{K}} - \left( e^{-\tau \mathcal{K}_0/n} e^{-\tau \mathcal{B}/n} \right)^n \right) \right\|_{\mathcal{L}(L^p(\mathcal{I},\mathfrak{X}))} \\ &= \operatorname{ess\,sup}_{t \in \mathcal{I}} \| U(t+\tau,t) - V_n(t+\tau,t) \|_{\mathcal{L}(\mathfrak{X})} \chi_{\mathcal{I}}(t+\tau) , \end{aligned}$$

for each  $\tau \ge 0$ . Note that one has

$$\begin{split} \sup_{\tau \ge 0} & \left\| L(\tau) \left( e^{-\tau \mathcal{K}} - \left( e^{-\tau \mathcal{K}_0/n} e^{-\tau \mathcal{B}/n} \right)^n \right) \right\|_{\mathcal{L}(L^p(\mathcal{I},\mathfrak{X}))} \\ &= \operatorname{ess\,sup}_{\tau \ge 0} \left\| L(\tau) \left( e^{-\tau \mathcal{K}} - \left( e^{-\tau \mathcal{K}_0/n} e^{-\tau \mathcal{B}/n} \right)^n \right) \right\|_{\mathcal{L}(L^p(\mathcal{I},\mathfrak{X}))}. \end{split}$$

This is based on the fact that if  $F(\cdot) : \mathbb{R}^+ \to \mathcal{L}(\mathfrak{X})$  is strongly continuous, then  $\sup_{\tau \ge 0} ||F(\tau)||_{\mathcal{L}(\mathfrak{X})} = \operatorname{ess\,sup}_{\tau \ge 0} ||F(\tau)||_{\mathcal{L}(\mathfrak{X})}$ . Hence, we find

$$\begin{split} \sup_{\tau \ge 0} & \left\| L(\tau) \left( e^{-\tau \mathcal{K}} - \left( e^{-\tau \mathcal{K}_0/n} e^{-\tau \mathcal{B}/n} \right)^n \right) \right\|_{\mathcal{L}(L^p(I,\mathfrak{X}))} \\ & = \operatorname{ess\,sup\,ess\,sup\,}_{t \in I} \| U(t+\tau,t) - V_n(t+\tau,t)) \|_{\mathcal{L}(\mathfrak{X})} \, \chi_I(t+\tau). \end{split}$$

Further, if  $\Phi(\cdot, \cdot) : \mathbb{R}^+ \times \mathcal{I} \to \mathcal{L}(\mathfrak{X})$  is a strongly measurable function, then

$$\operatorname{ess\,sup}_{(\tau,t) \in \mathbb{R}^+ \times I} \| \Phi(\tau,t) \|_{\mathcal{L}(\mathfrak{X})} = \operatorname{ess\,sup}_{\tau \ge 0} \operatorname{ess\,sup}_{t \in I} \| \Phi(\tau,t) \|_{\mathcal{L}(\mathfrak{X})}.$$

Then, taking into account two last equalities, one obtains

$$\begin{split} \sup_{\tau \ge 0} \left\| L(\tau) \left( e^{-\tau \mathcal{K}} - \left( e^{-\tau \mathcal{K}_0/n} e^{-\tau \mathcal{B}/n} \right)^n \right) \right\|_{\mathcal{L}(\mathfrak{X})} \\ &= \operatorname{ess\,sup}_{(\tau,t) \in \mathbb{R}^+ \times \mathcal{I}} \left\| U(t+\tau,t) - V_n(t+\tau,t) \right\|_{\mathcal{L}(\mathfrak{X})} \chi_{\mathcal{I}}(t+\tau) = \\ &= \operatorname{ess\,sup}_{(t,s) \in \mathcal{A}} \left\| U(t,s) - V_n(t,s) \right\|_{\mathcal{L}(\mathfrak{X})}, \end{split}$$

that proves (3.6).

We study bounded perturbations of the evolution generator  $\mathcal{K}_0 = D_0$  (3.2). To this aim we consider  $\mathcal{I} = [0, 1]$ ,  $\mathfrak{X} = \mathbb{C}$  and we denote by  $L^p(\mathcal{I})$  the Banach space  $L^p(\mathcal{I}, \mathbb{C})$ .

For  $t \in I$ , let  $q : t \mapsto q(t) \in L^{\infty}(I)$ . Then, q induces a *bounded* multiplication operator  $\mathcal{B} = Q$  on the Banach space  $L^{p}(I)$ :

$$(Qf)(t) = q(t)f(t), f \in L^p(I).$$

For simplicity we assume that  $q \ge 0$ . Then Q generates on  $L^p(I)$  a contraction semigroup  $\{e^{-\tau Q}\}_{\tau \ge 0}$ . Since generator Q is bounded, the closed operator  $\mathcal{A} := D_0 + Q$ , with domain dom  $(\mathcal{A}) = \text{dom}(D_0)$ , is generator of a  $C_0$ -semigroup on  $L^p(I)$ . By the *Trotter* product formula in the *strong* operator topology it follows immediately that

$$\lim_{n \to \infty} \left( e^{-\tau D_0/n} e^{-\tau Q/n} \right)^n f = e^{-\tau (D_0 + Q)} f, \quad f \in L^p(\mathcal{I}),$$
(3.7)

uniformly in  $\tau \in [0, T]$  on bounded time intervals.

Then we define on Banach space  $\mathfrak{X} = \mathbb{C}$  a family of bounded operators  $\{V(t)\}_{t \in \mathcal{I}}$  by

$$V(t) := e^{-\int_0^t ds \, q(s)}, \quad t \in \mathcal{I}.$$

Note that for almost every  $t \in I$  these operators are positive. Then  $V^{-1}(t)$  exists and it has the form

$$V^{-1}(t) = \mathrm{e}^{\int_0^t ds \, q(s)} \,, \quad t \in \mathcal{I} \,.$$

$$V^{-1}(D_0 + Q) V = D_0 \text{ or } D_0 + Q = V D_0 V^{-1}$$

Hence, the semigroup generated on  $L^p(I)$  by  $D_0 + Q$  gets the explicit form:

$$\left( e^{-\tau (D_0 + Q)} f \right)(t) = \left( \mathcal{V} e^{-\tau D_0} \mathcal{V}^{-1} f \right)(t) =$$

$$= e^{-\int_{t-\tau}^t dy \, q(y)} f(t-\tau) \chi_I(t-\tau) .$$
(3.8)

Since by (3.1) the solution operator U(t, s) that corresponds to evolution semigroup (3.8) is defined by equation

$$\left(\mathrm{e}^{-\tau(D_0+Q)}\right)f(t) = U(t,t-\tau)f(t-\tau)\chi_I(t-\tau)\,,$$

we deduce that it is equal to  $U(t, s) = e^{-\int_{s}^{t} dy q(y)}$ .

Now we study the corresponding Trotter product formula. For a fixed  $\tau \ge 0$  and  $n \in \mathbb{N}$ , we define *approximates*  $\{V_n\}_{n\ge 1}$  by

$$\left(\left(\mathrm{e}^{-\tau D_0/n}\mathrm{e}^{-\tau Q/n}\right)^n f\right)(t) =: V_n(t,t-\tau)\chi_I(t-\tau)f(t-\tau) \ .$$

Then by straightforward calculations, which are similar to (3.5), one finds that approximants have the following explicit form:

$$V_n(t,s) = e^{-\tau_n \sum_{k=0}^{n-1} q(s+k\tau_n)}, \quad (t,s) \in \Delta, \quad \tau_n = (t-s)/n, \quad n \in \mathbb{N}.$$

**Theorem 3.3.** Let  $q \in L^{\infty}(I)$  be non-negative. Then

$$\begin{split} \sup_{\tau \ge 0} \left\| e^{-\tau(D_0 + Q)} - \left( e^{-\tau D_0/n} e^{-\tau Q/n} \right)^n \right\|_{\mathcal{L}(L^p(I))} &= \\ \Theta \left( \operatorname{ess\,sup}_{(t,s) \in \mathcal{A}} \left| \int_s^t dy \, q(y) - \tau_n \, \sum_{k=0}^{n-1} q \, (s+k \, \tau_n) \right| \right), \quad n \in \mathbb{N} \,, \end{split}$$

as  $n \to \infty$ , where  $\Theta$  is the Landau symbol defined in Remark 3.1, see Subsection 3.1.

*Proof.* First, by Theorem 3.2 and by  $U(t, s) = e^{-\int_{s}^{t} dy q(y)}$  we obtain

$$\sup_{\tau \ge 0} \left\| e^{-\tau(D_0 + Q)} - \left( e^{-\tau D_0 / n} e^{-\tau Q / n} \right)^n \right\|_{\mathcal{L}(L^p(I))}$$

$$= \operatorname{ess\,sup}_{(t,s) \in \Delta} \left| e^{-\int_s^t dy \, q(y)} - e^{-\tau_n \sum_{k=0}^{n-1} q(s+k \tau_n)} \right| .$$
(3.9)

Then, using the inequality

$$e^{-\max\{x,y\}} |x-y| \le |e^{-x} - e^{-y}| \le |x-y|, \quad 0 \le x, y,$$

for  $0 \le s < t \le 1$  one finds the estimates

$$e^{-||q||_{L^{\infty}}} R_n(t,s;q) \le \left| e^{-\int_s^t dy q(y)} - e^{-\tau_n \sum_{k=0}^{n-1} q(s+k\tau_n)} \right| \le R_n(t,s;q) ,$$

where

$$R_n(t, s, q) := \left| \int_s^t dy \, q(y) - \tau_n \, \sum_{k=0}^{n-1} q(s+k\,\tau_n) \,\right|, \quad (t,s) \in \Delta \,, \quad n \in \mathbb{N} \,. \tag{3.10}$$

Hence, for the left-hand side of (3.9) we get the estimate

$$e^{-||q||_{L^{\infty}}} R_n(q) \le \sup_{\tau \ge 0} \left\| e^{-\tau(D_0+Q)} - \left( e^{-\tau D_0/n} e^{-\tau Q/n} \right)^n \right\|_{\mathcal{L}(L^p(\mathcal{I}))} \le R_n(q) ,$$

where  $R_n(q) := \operatorname{ess\,sup}_{(t,s)\in\Delta} R_n(t,s;q), n \in \mathbb{N}$ . These estimates together with definition of  $\Theta$  prove the assertion.

Note that by virtue of (3.10) and Theorem 3.3 the operator-norm convergence rate of the Trotter product formula for the pair  $\{D_0, Q\}$  coincides with the convergence rate of the integral *Darboux-Riemann* sum approximation of the *Lebesgue* integral.

# 3.3 Rate of convergence

First we consider the case of a real Hölder-continuous function  $q \in C^{0,\beta}(I)$ .

**Theorem 3.4.** If  $q \in C^{0,\beta}(I)$  is non-negative, then

$$\sup_{\tau \ge 0} \left\| e^{-\tau (D_0 + Q)} - \left( e^{-\tau D_0 / n} e^{-\tau Q / n} \right)^n \right\|_{\mathcal{L}(L^p(I))} = O(1 / n^\beta) ,$$

as  $n \to \infty$ .

*Proof.* One gets for  $\tau_n = (t - s)/n$  and  $n \in \mathbb{N}$ :

$$\int_{s}^{t} dy q(y) - \tau_{n} \sum_{k}^{n-1} q(s + k \tau_{n})$$
  
= 
$$\sum_{k=0}^{n-1} \int_{k \tau_{n}}^{(k+1)\tau_{n}} dy (q(s + y) - q(s + k \tau_{n})) ,$$

which yields the estimate

$$\begin{split} \left| \int_{s}^{t} dy \, q(y) - \tau_{n} \, \sum_{k}^{n-1} q \, (s+k\tau_{n}) \right| \\ & \leq \sum_{k=0}^{n-1} \int_{k\tau_{n}}^{(k+1)\tau_{n}} dy \, | \, q(s+y) - q(s+k\,\tau_{n}) | \, . \end{split}$$

Since  $q \in C^{0,\beta}(I)$ , there is a constant  $L_{\beta} > 0$  such that for  $y \in [k\tau_n, (k+1)\tau_n]$  one has

$$|q(s+y) - q(s+k\tau_n)| \le L_\beta |y-k\tau_n|^\beta \le L_\beta \frac{(t-s)^\beta}{n^\beta}.$$

Hence, we find

$$\left|\int_{s}^{t} dy \, q(y) - \tau_n \, \sum_{k}^{n-1} q(s + k\tau_n)\right| \le L_\beta \, \frac{(t-s)^{1+\beta}}{n^{\beta}} \le L_\beta \, \frac{1}{n^{\beta}} \, ,$$

which proves (cf. Remark 3.1)

$$\operatorname{ess\,sup}_{(t,s)\in\mathcal{A}} \left| \int_{s}^{t} dy \, q(y) - \tau_n \sum_{k}^{n-1} q(s+k\,\tau_n) \right| = O\left(\frac{1}{n^{\beta}}\right).$$

Applying now Theorem 3.3 one completes the proof.

The next natural question is: what happens, when function q is only *continuous*?

**Theorem 3.5.** *If*  $q : \mathcal{I} \to \mathbb{C}$  *is continuous and non-negative, then* 

$$\left\| e^{-\tau(D_0 + Q)} - \left( e^{-\tau D_0/n} e^{-\tau Q/n} \right)^n \right\|_{\mathcal{L}(L^p(I))} = o(1) , \qquad (3.11)$$

as  $n \to \infty$ .

*Proof.* Seeing that q is continuous, for any  $\varepsilon > 0$  there is  $\delta > 0$  such that for  $y, x \in I$  and  $|y - x| < \delta$  we have  $|q(y) - q(x)| < \varepsilon$ . Therefore, if  $1/n < \delta$ , then for  $y \in (k\tau_n, (k+1)\tau_n)$  we have

$$|q(s+y) - q(s+k\tau_n)| < \varepsilon, \quad (t,s) \in \varDelta.$$

Hence,

$$\left|\int_{s}^{t} dy q(y) - \tau_n \sum_{k}^{n-1} q(s+k\tau_n)\right| \leq \varepsilon(t-s) \leq \varepsilon,$$

which yields (cf. Remark 3.1)

$$\operatorname{ess\,sup}_{(t,s)\in\mathcal{A}} \Big| \int_s^t dy \, q(y) - \tau_n \, \sum_k^{n-1} q(s+k\,\tau_n) \Big| = o(1) \, .$$

Now it remains only to apply Theorem 3.3.

Here it is worth to note that for *general* continuous function q one can say *nothing* about the convergence rate. Indeed, it can be shown that in (3.11) the convergence to zero can be *arbitrary slow* (3.12) for a bounded perturbation Q. This is drastically different to the case, when dominating generator corresponds to a holomorphic semigroup and perturbation operator is bounded, cf. (2.17), (2.19) for  $\alpha = 0$ , or to the case of unbounded perturbation, when  $0 < \alpha < 1$ , see (2.16), (2.18).

**Theorem 3.6.** Let  $\delta_n > 0$  be a sequence with  $\delta_n \to 0$  as  $n \to \infty$ . Then there exists a continuous function  $q : \mathcal{I} = [0, 1] \to \mathbb{R}$ , such that

$$\sup_{\tau \ge 0} \left\| e^{-\tau(D_0 + Q)} - \left( e^{-\tau D_0 / n} e^{-\tau Q / n} \right)^n \right\|_{\mathcal{L}(L^p(\mathcal{I}))} = \omega(\delta_n) , \qquad (3.12)$$

as  $n \to \infty$ , where  $\omega$  is the Landau symbol defined in Remark 3.1, see Subsection 3.1.

*Proof.* Taking into account the *Walsh-Sewell* theorem ([WaSe37], Theorem 6), we find that for any sequence  $\{\delta_n\}_{n \in \mathbb{N}}, \delta_n > 0$  satisfying  $\lim_{n \to \infty} \delta_n = 0$  there exists a continuous function  $f : [0, 2\pi] \longrightarrow \mathbb{R}$ , such that

$$\left|\int_0^{2\pi} dx f(x) - \frac{2\pi}{n} \sum_{k=1}^n f(2k\pi/n)\right| = \omega(\delta_n) ,$$

as  $n \to \infty$ . Setting  $q(y) := f(2\pi(1 - y))$  for  $y \in [0, 1]$ , we get a continuous function  $q : [0, 1] \longrightarrow \mathbb{R}$ , such that

$$\left| \int_0^1 dy \, q(y) - \frac{1}{n} \sum_{k=0}^{n-1} q(k/n) \right| = \omega(\delta_n) \, .$$

Given that function q is continuous, we find

$$\underset{(t,s)\in\mathcal{A}}{\operatorname{ess\,sup}} \left| \int_{s}^{t} dy \, q(y) - \tau_{n} \sum_{n=0}^{n-1} q(s+k\,\tau_{n}) \right| \\ \geq \left| \int_{0}^{1} dy \, q(y) - \frac{1}{n} \sum_{k=0}^{n-1} q(k/n) \right|,$$

which yields

$$\operatorname{ess\,sup}_{(t,s)\in\varDelta} \Big| \int_s^t dy \, q(y) - \tau_n \sum_{n=0}^{n-1} q(s+k\,\tau_n) \Big| = \omega(\delta_n) \; .$$

Applying now Theorem 3.3 we prove (3.12).

Our final comment concerns the case, when function  $q : [0,1] \rightarrow \mathbb{R}$  is only *measurable*. Then it can happen that the Trotter product formula for that pair  $\{D_0, Q\}$  *does not* converge in the *operator-norm* topology.

$$\limsup_{n \to \infty} \sup_{\tau \ge 0} \left\| e^{-\tau(D_0 + Q)} - \left( e^{-\tau D_0 / n} e^{-\tau Q / n} \right)^n \right\|_{\mathcal{L}(L^p(I))} > 0.$$
(3.13)

Proof. Let us introduce open intervals:

$$\begin{aligned} \Delta_{0,n} &:= (0, \frac{1}{2^{2n+2}}), \\ \Delta_{k,n} &:= (t_{k,n} - \frac{1}{2^{2n+2}}, t_{k,n} + \frac{1}{2^{2n+2}}), \quad k = 1, 2, \dots, 2^n - 1, \\ \Delta_{2^n,n} &:= (1 - \frac{1}{2^{2n+2}}, 1), \end{aligned}$$

 $n \in \mathbb{N}$ , where

$$t_{k,n}=rac{k}{2^n},\quad k=0,\ldots,n,\quad n\in\mathbb{N}.$$

Notice that  $t_{0,n} = 0$  and  $t_{2^n,n} = 1$ . One easily checks that the intervals  $\Delta_{k,n}$ ,  $k = 0, \ldots, 2^n$ , are mutually disjoint. We introduce the open sets

$$O_n = \bigcup_{k=0}^{2^n} \varDelta_{k,n} \subseteq I, \quad n \in \mathbb{N}.$$

and

$$O=\bigcup_{n\in\mathbb{N}}O_n\subseteq I.$$

Then it is clear that

$$|O_n| = \frac{1}{2^{n+1}}, \quad n \in \mathbb{N}, \text{ and } |O| \le \frac{1}{2}.$$

Therefore, the *Lebesgue* measure of the closed set  $C := I \setminus O \subseteq I$  can be estimated by

$$|C| \ge \frac{1}{2} \ .$$

Using the characteristic function  $\chi_{\mathcal{C}}(\cdot)$  of the set  $\mathcal{C}$  we define

$$q(t) := \chi_C(t), \quad t \in \mathcal{I}$$

The function q is measurable and it satisfies  $0 \le q(t) \le 1$ ,  $t \in I$ . Let  $\varepsilon \in (0, 1)$ . We choose  $s \in (0, \varepsilon)$  and  $t \in (1 - \varepsilon, 1)$  and we set

$$\xi_{k,n}(t,s) := s + k \frac{t-s}{2^n}, \quad k = 0, \dots, 2^n - 1, \quad n \in \mathbb{N}, \quad (t,s) \in \Delta.$$

Note that  $\xi_{k,n}(t,s) \in (0,1), k = 0, \dots, 2^n - 1, n \in \mathbb{N}$ . Moreover, we have

$$t_{k,n} - \xi_{k,n}(t,s) = k \frac{1}{2^n} - s - k \frac{t-s}{2^n} = k \frac{1-t+s}{2^n} - s ,$$

which leads to the estimate

$$|t_{k,n} - \xi_{k,n}(t,s)| \le \varepsilon \ (1 + k/2^{n-1}), \quad k = 0, \dots, 2^n - 1, \quad n \in \mathbb{N}$$

Hence

$$|t_{k,n}-\xi_{k,n}(t,s)|\leq 3\varepsilon, \quad k=0,\ldots,2^n-1, \quad n\in\mathbb{N}.$$

Let  $\varepsilon_n := 1/(3 \cdot 2^{2n+2})$  for  $n \in \mathbb{N}$ . Then we get that  $\xi_{k,n}(t, s) \in \Delta_{k,n}$  for  $k = 0, \ldots, 2^n - 1$ ,  $n \in \mathbb{N}$ ,  $s \in (0, \varepsilon_n)$  and for  $t \in (1 - \varepsilon_n, 1)$ .

Now let

$$S_n(t, s; q) := \tau_n \sum_{k=0}^{n-1} q(s + k\tau_n), \quad (t, s) \in \Delta, \quad \tau_n = (t - s)/n, \quad n \in \mathbb{N}.$$

We consider

$$S_{2^n}(t,s;q) = \frac{t-s}{2^n} \sum_{k=0}^{2^n-1} q\left(s+k \frac{t-s}{2^n}\right) = \frac{t-s}{2^n} \sum_{k=0}^{2^n-1} q\left(\xi_{k,n}(t,s)\right),$$

 $n \in \mathbb{N}, (t, s) \in \Delta$ . If  $s \in (0, \varepsilon_n)$  and  $t \in (1 - \varepsilon_n, 1)$ , then  $S_{2^n}(t, s; q) = 0, n \in \mathbb{N}$  and

$$\left|\int_{s}^{t} dy q(y) - S_{2^{n}}(t,s;q)\right| = \int_{s}^{t} dy q(y), \quad n \in \mathbb{N},$$

for  $s \in (0, \varepsilon_n)$  and  $t \in (1 - \varepsilon_n, 1)$ . In particular, this yields

$$\operatorname{ess\,sup}_{(t,s)\in\mathcal{A}} \left| \int_{s}^{t} dy \, q(y) - S_{2^{n}}(t,s;q) \right| \geq \operatorname{ess\,sup}_{(t,s)\in\mathcal{A}} \int_{s}^{t} dy \, q(y) \geq \int_{\mathcal{I}} dy \, \chi_{C}(y) \geq \frac{1}{2} \, .$$

Hence, we obtain

$$\limsup_{n\to\infty} \operatorname{ess\,sup}_{(t,s)\in\mathcal{A}} \left| \int_s^t dy \, q(y) - S_{2^n}(t,s;q) \right| \ge \frac{1}{2},$$

and applying Theorem 3.3 we finish the prove of (3.13).

**Remark 3.8.** We note that Theorem 3.7 does not exclude the convergence of the Trotter product formula for the pair  $\{D_0, Q\}$  in the *strong* operator topology. We would remind that the same kind of *dichotomy* is known for the Trotter product formula on a Hilbert space, see the *Hiroshi Tamura* example in [Tam00], Theorem B. By virtue of (3.7) and (3.13), Theorem 3.7 yields an example of this dichotomy on a Banach space.

Again, there is a drastic difference between the origin of these conclusions in a Hilbert space ([Tam00], Theorem B) for *unbounded* perturbation of the *holomorphic* semigroup and in a Banach space (Theorem 3.7) for *bounded* perturbation of a (non-holomorphic) *contractive* semigroup.

## 4 Notes

Notes to Section 1. Characterisation of *holomorphic contraction* semigroups at the end of Subsection 1.2 (iv), is due to Corollary II.4.9 [EN00]. For the proof of Lemma 1.11 see, for example [Tan75], Lemma 2.3.5.

Notes to Section 2. Here we extend to the *operator-norm* convergence of the product formula on a Banach space for perturbation *B* with a relative *zero A*-bound for holomorphic semigroup  $\{e^{-tA}\}_{t\geq 0}$  some of the Trotter-Chernoff results, cf. [Trot59], [But20], [Zag20]. This shows that hypothesis of self-adjointness in the case of a Hilbert space [IT97] has only a technical importance.

On the other hand the operator-norm topology is "natural" for holomorphic  $C_0$ semigroups, which may lead one to think that it is an essential hypothesis for the operator-norm convergence of the Trotter product formula. In Section 3 we showed that this hypothesis is also technical, but we have to assume contraction of semigroup {e  ${}^{-tA}$ }<sub>t≥0</sub>. We would like to remark that demand of contraction is not as superfluous as one could suppose. For demonstration we address the reader to instructive example by Trotter [Trot59], where it is called "the norm condition".

This section contains a revision of result [CZ01], Section 3, where the *operator*norm convergence of the Trotter product formula on a Banach space  $\mathfrak{X}$  has been proven (up to our knowledge) for the first time. For a survey of similar results in this direction see [NSZ18b].

Notes to Section 3. In contrast to Section 2, where *operator-norm* convergence holds true if the dominating operator  $A \in \mathscr{H}_c(\theta, 0)$  generates a *holomorphic* contraction semigroup and operator *B* is a *A*-infinitesimally small generator of a *contraction* semigroup (in particular, if *B* is a *bounded* operator), we present *Example* that this is also possible if condition on generator *A* is *relaxed*. The conditions are [NSZ18a]: (1) Operator  $A = \mathcal{K}_0$  generates a *contractive* (not holomorphic!) semigroup. (2)  $B = \mathscr{B}$  is a *bounded* operator.

There it is also demonstrated that the operator-norm convergence generally *fails* (even for bounded operators  $\mathcal{B}$ ) if *unbounded*  $\mathcal{K}_0$  is *not* a *holomorphic* generator and that operator-norm convergence of the Trotter product formula can be *arbitrary slow*, cf. Subsection 3.3. This is again very different to the holomorphic case:  $A \in \mathcal{H}_c(\theta, 0)$  (cf. Subsection 2.2), where the rate of the operator-norm convergence is of the order  $O((\ln n)^2/n)$  for *any* bounded perturbation B ( $\alpha = 0$ ), see Theorems 2.6, 2.7, and Corollary 2.8.

# References

But20.	Ya.A. Butko, The method of Chernoff approximation, pp 19-46. In: J. Banasiak
	et al. (eds.), Semigroups of Operators – Theory and Applications, SOIA 2018, Springer Presedings in Methometics and Statistics vol. 225, Springer Parlin
	Springer Proceedings in Mathematics and Statistics, vol. 525. Springer, Berlin
CZ01.	V. Cachia and V. A. Zagrebnov, <i>Operator-norm convergence of the Trotter product</i>
02011	formula for holomorphic semigroups, J. Oper. Theory <b>46</b> (2001), 199–213.
EN00.	KJ. Engel and R. Nagel, One-parameter Semigroups for Linear Evolution Equa-
	tions, Springer-Verlag, Berlin, 2000.
Ev76.	D. E. Evans. Time dependent perturbations and scattering of strongly continuous
	groups on Banach spaces. Math. Ann., 221(3):275–290, 1976.
How74.	J. S. Howland, Stationary scattering theory for time-dependent Hamiltonians,
	Math. Ann. <b>207</b> (1974) 315–335.
IT97.	T. Ichinose and Hideo Tamura, Error estimate in operator norm for Trotter-Kato
	product formula, Integral Equations Oper. Theory 27 (1997), 195–207.
Kato95.	T. Kato. <i>Perturbation Theory for Linear Operators</i> . (Corrected Printing of the Sec-
N.: 70	ond Edition 1980) Springer-Verlag, Berlin, Heidelberg, 1995.
Nei /9.	H. Neidhardt. Integration of Evolutionsgleichungen mit Hilfe von Evolutionsnalb- gruppen Discortation AdW der DDP Berlin 1070
Nei81	H Neidhardt On abstract linear evolution equations I Math Nachr 103.283-
1001011	298, 1981.
NZ98.	H. Neidhardt and V. A. Zagrebnov, <i>On error estimates for the Trotter-Kato product</i>
	formula, Lett. Math. Phys. 44 (1998), 169-186.
NZ99.	H. Neidhardt and V. A. Zagrebnov, Fractional powers of self-adjoint operators and
	Trotter-Kato product formula, Integral Equations Oper. Theory 35 (1999), 209-
	231.
NSZ18a.	H. Neidhardt, A. Stephan and V. A. Zagrebnov, <i>Remarks on the operator-norm</i>
	<i>convergence of the Trotter product formula</i> , Integral Equations Oper. Theory <b>90:15</b>
NS718b	(2018), pp.1–14.
INSZ100.	Trotter product formula on Hilbert and Banach spaces: a short survey in: Current
	Research in Nonlinear Analysis. Springer Optimization and Its Applications vol.
	135, Springer, Berlin 2018, pp. 229–247.
NSZ20.	H. Neidhardt, A. Stephan, and V. A. Zagrebnov. Convergence rate estimates for
	Trotter product approximations of solution operators for non-autonomous Cauchy
	problems, Publ. RIMS Kyoto Univ. 56 (2020), 83-135.
Rog93.	Dzh. L. Rogava, Error bounds for Trotter-type formulas for self-adjoint operators,
-	Funct. Anal. Appl. <b>27</b> (1993), 217–219.
Tam00.	Hiroshi Tamura, A remark on operator-norm convergence of Trotter-Kato product
Top75	<i>formula</i> , Integral Equations Oper. Theory 37 (2000), 350–356.
Tall/3.	H. Tallade. Equations of evolution, Tokyo, Twallalli, 1975 (III Japanese). Eligitsii translation: Ditman Advanced Publishing Program London, 1979
Trot59	H F Trotter On the products of semigroups of operators Proc. Amer. Math. Soc.
110037.	<b>10</b> (1959), 545–551.
WaSe37.	J. L. Walsh and W. E. Sewell, <i>Note on degree of approximation to an integral by</i>
	Riemann sums, The American Mathematical Monthly 44 (1937), 155–160.
Yos80.	K. Yosida, Functional Analysis, Springer-Verlag, Berlin, 1980.
Zag20.	V. A. Zagrebnov, Notes on the Chernoff product formula, J. Funct. Anal. 279
	(2020), 108696–24pp.