THE SLE LOOP VIA CONFORMAL WELDING OF QUANTUM DISKS

MORRIS ANG, NINA HOLDEN, AND XIN SUN

ABSTRACT. We prove that the SLE_{κ} loop measure arises naturally from the conformal welding of two γ -Liouville quantum gravity (LQG) disks for $\gamma^2 = \kappa \in (0,4)$. The proof relies on our companion work on conformal welding of LQG disks and uses as an essential tool the concept of uniform embedding of LQG surfaces. Combining our result with work of Gwynne and Miller, we get that random quadrangulations decorated by a self-avoiding polygon converge in the scaling limit to the LQG sphere decorated by the $\mathrm{SLE}_{8/3}$ loop. Our result is also a key input to recent work of the first and third coauthors on the integrability of the conformal loop ensemble. Finally, our result can be viewed as the random counterpart of an action functional identity due to Viklund and Wang.

1. Introduction

The Schramm-Loewner evolution (SLE) and Liouville quantum gravity (LQG) are central objects in random conformal geometry. It was shown by Sheffield [She16] and Duplantier-Miller-Sheffield [DMS21] that SLE curves arise as the interfaces of LQG surfaces under conformal welding. This phenomenon is a cornerstone of the mating-of-trees framework [DMS21] for the SLE/LQG coupling and is a fundamental input to the link between LQG and the scaling limits of random planar maps. See e.g. [Law05, GHS19, Gwy20, BP21, She22] for an introduction to SLE, LQG, and their interactions.

Conformal welding results in [She16, DMS21] mainly focus on infinite-volume LQG surfaces. Recently in [AHS20b], we showed that the conformal welding of finite-volume LQG surfaces called two-pointed quantum disks can give rise to some canonical variants of SLE curves with two marked points. In this paper, we show in Theorem 1.1 that when conformally welding two quantum disks without marked points, the interface is another canonical variant of SLE called the *SLE loop*. Moreover, the resulting LQG surface is the so-called quantum sphere (without marked point), which describes the scaling limit of classical random planar map models with spherical topology. For example, in the pure gravity case it corresponds to the Brownian map [Le 13, Mie13, MS15].

As reviewed in Section 1.1, the SLE loop is an important one-parameter family of conformally invariant random Jordan curves whose existence was conjectured by Kontsevitch and Suhov [KS07] and established by Zhan [Zha21]. In particular, the $SLE_{8/3}$ loop introduced by Werner [Wer08] describes the conjectural scaling limit of self-avoiding polygons on planar lattices. Our conformal welding result Theorem 1.1 combined with earlier work of Gwynne and Miller [GM19b, GM16] yields that uniform quadrangulations decorated by a self-avoiding loop converge to the Brownian map decorated by the $SLE_{8/3}$ loop; see Theorem 1.2. For $\kappa \in (8/3, 4]$, the SLE_{κ} loop is closely related to the conformal loop ensemble (CLE) considered in [She09, SW12].

We will state Theorems 1.1 and 1.2 in Section 1.1 and 1.2, respectively, modulo some background material supplied in Section 2. Then we prove Theorems 1.1 and 1.2 in Sections 3 and 4, respectively. Our proof of Theorem 1.1 builds on the conformal welding result in [AHS20b] and the Liouville field description of the quantum surfaces in [AHS22], which will be recalled in Section 2 as well.

1.1. The SLE loop via conformal welding. Kontsevitch and Suhov [KS07], inspired by Malliavin [Mal99], formulated a natural conformal restriction covariance property for loop measures, and they conjectured that there is a unique (up to multiplicative factor) one-parameter family of loop measures

satisfying this property. We call loop measures satisfying this property a Malliavin-Kontsevitch-Suhov (MKS) loop measure and we index the measures by $\kappa \in (0, 4]$.

Werner [Wer08] proved the existence and uniqueness of the MKS loop measure for $\kappa = 8/3$ and constructed the loop measure via the boundaries of Brownian loops. Kemppainen and Werner [KW16] constructed an MKS loop measure for $\kappa \in (8/3, 4]$ using the density measure of a nested simple CLE. For $\kappa = 2$, Benoist and Dubédat [BD16] proved that a measure constructed in [KK17] is an MKS loop measure. Finally, Zhan [Zha21] constructed an MKS loop measure for all $\kappa \in (0, 4]$ via SLE_{κ} equipped with its natural parametrization and also extended the construction to $\kappa \in (4, 8)$. We denote Zhan's MKS loop measure by SLE_{κ}^{loop} . See Section 2.5 for the precise definition. For $\kappa \in (8/3, 4]$, the loop measures of Zhan and Kemppainen-Werner agree; see [AS21, Section 2.3]. We emphasize that SLE_{κ}^{loop} is an infinite measure for each $\kappa \in (0, 4]$.

For each $\gamma \in (0,2)$ there is a natural infinite measure on LQG surfaces with spherical topology called the unmarked quantum sphere. We denote this measure by QS and refer to Section 2 for the precise definition of both this measure and of the other objects we introduce in this and the next paragraph. Let $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be the Riemann sphere. Suppose h is a random field on \mathbb{C} such that the distribution of $(\widehat{\mathbb{C}}, h)$ viewed as a quantum surface is QS. Let η be a sample of $\mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ independent of h for $\kappa = \gamma^2 \in (0, 4]$. It is known that $\mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ is invariant under Möbius transforms [Zha21, Theorem 4.2]. Therefore, although the distribution of h is not uniquely specified, as a curve-decorated quantum surface, the distribution of $(\widehat{\mathbb{C}}, h, \eta)$ is uniquely defined. We denote this loop-decorated quantum surface by $\mathrm{QS} \otimes \mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ and call it the MKS-loop-decorated quantum sphere with parameter γ ; see Section 3 for a precise definition.

For each $\gamma \in (0,2)$ there is also a natural infinite measure on LQG surfaces with disk topology called the unmarked quantum disk. We denote this measure by QD. Let QD(ℓ) be the disintegration of QD over its boundary length, namely QD = $\int_0^\infty \mathrm{QD}(\ell) \, d\ell$. For $\ell > 0$, let $(\mathcal{D}_1, \mathcal{D}_2)$ be a sample from QD(ℓ) × QD(ℓ), so \mathcal{D}_1 and \mathcal{D}_2 have boundary length ℓ . Let Weld($\mathcal{D}_1, \mathcal{D}_2$) be the curve-decorated quantum surface obtained by conformally welding \mathcal{D}_1 and \mathcal{D}_2 along their boundaries such that a uniformly sampled point on the boundary of \mathcal{D}_1 is identified with a uniformly sampled point on the boundary of \mathcal{D}_2 . We denote the distribution of Weld($\mathcal{D}_1, \mathcal{D}_2$) by Weld(QD(ℓ), QD(ℓ)) and define Weld(QD, QD) = $\int_0^\infty \ell \cdot \text{Weld}(\text{QD}(\ell), \text{QD}(\ell)) \, d\ell$; see Section 2.4 for more details.

Theorem 1.1. For $\gamma \in (0,2)$ and $\kappa = \gamma^2$, we have $QS \otimes SLE_{\kappa}^{loop} = CWeld(QD,QD)$ for some constant $C \in (0,\infty)$.

The proof of Theorem 1.1 builds on a welding result from [AHS20b] which says that the conformal welding of two quantum disks, each marked with two points sampled independently from the LQG boundary measure, gives a quantum sphere with two special singularities decorated with a so-called two-sided whole plane SLE_{κ} . Given this result and the construction of Zhan's MKS loop measure, Theorem 1.1 seems plausible. To explain the factor of ℓ in the definition of Weld(QD, QD), we appeal to the following intuition from the discrete: if we have two polygons with ℓ edges, there are ℓ different ways to glue them into a sphere with a self-avoiding loop. The rigorous proof of Theorem 1.1 relies on the idea of uniform embedding introduced in [AHS22] and explained in Section 2.3. The uniform embedding of a quantum surface is a particular random embedding where the fields can be described via Liouville conformal field theory (LCFT) as considered in [DKRV16, HRV18]. Moreover, it is especially convenient to work with when adding or removing marked points on the surface.

Based on the integrability of LQG from mating-of-trees [DMS21] and LCFT (see e.g. [KRV20, ARS22a, RZ22]), it was shown in [AHS22] that conformal welding can be applied to establishing integrability results for the SLE interfaces involved. Theorem 1.1 serves as the starting point of this application to the SLE loop and CLE. In [AS21], this approach was used to obtain the 3-point

 $^{^{1}\}mathrm{We}$ will use the language of probability in the setting of non-probability measures — see Section 2 for precise definitions.

correlation function for the nesting statistics and the electrical thickness of simple CLE. In [ARS22b], it was used to compute the annulus partition function of the ${\rm SLE}_{8/3}$ loop. In a forthcoming work of the first and third coauthors, it will be used to compute the renormalized probability that three given points are close to the same CLE loop on the sphere.

By a limiting argument, Theorem 1.1 can be naturally extended to $\kappa = 4$ using available conformal welding results for SLE₄ [HP21, MMQ19] but since the conformal removability of SLE₄ is not settled, the result would be less definite. Therefore we do not pursue this extension.

Viklund and Wang [VW20] proved a beautiful identity between the Loewner energy of a Jordan curve η on a sphere and the difference between the Dirichlet energy of a function φ on the sphere and the sum of the Dirichlet energies of φ restricted to each component of $\mathbb{S}^2 \setminus \eta$ after applying a uniformizing map. These quantities naturally arise from the large deviation of SLE and LCFT; see [Wan22] and [LRV19]. In particular, the identity can be viewed as a relation between the large deviation rate functions for an SLE loop measure, the LCFT on the sphere, and the LCFT on the disk. See [VW20, Section 1.3] for a discussion of this. Theorem 1.1 can be viewed as a quantum version of this identify.

1.2. The scaling limit of random planar maps decorated by self-avoiding loop. It is believed that the MKS loop with $\kappa=8/3$, namely Werner's loop measure [Wer08], is the scaling limit of the critical self-avoiding loop on a regular planar lattice. We will argue that this result holds in an annealed sense in the setting of planar maps. Namely, the critical Boltzmann measure on quadrangulations decorated with a self-avoiding loop converges to the MKS-loop-decorated quantum sphere with $\gamma=\sqrt{8/3}$ as curve-decorated metric measure spaces. The measure is called critical since the weight assigned to a loop-decorated quadrangulation has been tuned precisely so that the number of vertices of the quadrangulation and the length of the loop have a power-law behavior.

To state this result we will first introduce some notation; see Section 4 for more precise definitions. Let $\mathrm{MS}^n \otimes \mathrm{SAW}^n$ denote the measure on pairs (M,η) where M is a quadrangulation, η is a self-avoiding loop on M, and a pair (M,η) has weight $n^{2.5}12^{-\#\mathcal{F}(M)}54^{-\#\eta}$, where $\#\mathcal{F}(M)$ is the number of faces of M and $\#\eta$ is the number of edges of η . Note that the parameter n does not correspond to any quantity in the quadrangulation beyond the weights we use to define MS^n and as a scaling factor for distances and areas (see Section 4 for the latter). We let $\mathrm{QS} \otimes \mathrm{SLE}_{8/3}^{\mathrm{loop}}$ be as in Theorem 1.1 with $\gamma^2 = \kappa = 8/3$. As we will explain in Section 4, samples from $\mathrm{MS}^n \otimes \mathrm{SAW}^n$ and $\mathrm{QS} \otimes \mathrm{SLE}_{8/3}^{\mathrm{loop}}$ can be viewed as loop-decorated metric measure spaces. In this setting the natural topology for weak convergence is the Gromov-Hausdorff-Prokhorov-uniform (GHPU) topology; see Section 2.6 for a precise definition. For a loop-decorated metric measure space and $c \in (0,1)$ we let A(c) denote the event that the length of the loop is in $[c,c^{-1}]$. We use the notation $M|_{A(c)}$ to stand for the restriction of the measure M to the event A(c), and use the symbol \Rightarrow to indicate weak convergence of finite measures.

Theorem 1.2. There exists $c_0 > 0$ such that for all $c \in (0,1)$,

$$MS^n \otimes SAW^n |_{A(c)} \Rightarrow c_0 \cdot QS \otimes SLE_{8/3}^{loop} |_{A(c)} \quad as \ n \to \infty$$

with respect to the Gromov-Hausdorff-Prokhorov-uniform topology.

Here, we restrict to A(c) to make all measures in Theorem 1.2 finite. Gwynne and Miller proved the counterpart of the theorem in the setting of chordal self-avoiding paths on half-planar quadrangulations [GM16, GM19b], and results from their papers are key inputs to our proof. The other inputs are Theorem 1.1 and an exact discrete counterpart (also observed in [GM19a, CC19]) of Theorem 1.1 given in Observation 4.2.

It is a classical result that the quantum sphere for $\gamma = \sqrt{8/3}$ (also known as the Brownian map) arises as the scaling limit of uniformly sampled planar maps [Le 13, Mie13]. By contrast, our Theorem 1.2 gives the analogous result for a family of *non-uniform* planar maps in the sense that

two planar maps of the same size do not have the same probability of being sampled. Indeed, if (M, η) is sampled from $MS^n \otimes SAW^n$ then the marginal law of M has been reweighted according to the (weighted) number of self-avoiding loops that the map admits. On the other hand, Theorem 1.2 means that this reweighting does not change the scaling limit of the planar map.

For concreteness Theorem 1.2 is stated and proved for quadrangulations, but we remark that the result also holds for random triangulations building on [AHS20a].² By universality we expect that Theorem 1.2 also extends to other families of planar maps decorated by a self-avoiding loop.

Acknowledgements. We are in debt to Yilin Wang for her important insight on SLE loop. In our opinion, her contribution to Theorem 1.1 is as much as ours. We are also grateful to Steffen Rohde, Scott Sheffield, and Dapeng Zhan for helpful discussions. We thank two anonymous referees for helpful comments on an earlier version of this paper. M.A. was supported by the Simons Foundation as a Junior Fellow at the Simons Society of Fellows, and partially supported by NSF grant DMS-1712862. N.H. was supported by Dr. Max Rössler, the Walter Haefner Foundation, and the ETH Zürich Foundation, along with grant 175505 of the Swiss National Science Foundation. X.S. was supported by the Simons Foundation as a Junior Fellow at the Simons Society of Fellows, and by the NSF grant DMS-2027986 and the Career award 2046514.

2. Preliminaries

In this paper we use the language of probability theory in the setting of non-probability measures. If M is a measure on a measurable space (Ω, \mathcal{F}) and X is an \mathcal{F} -measurable function, we call the pushforward measure $M_X = X_*M$ the law of X, and say that X is sampled from M_X . For a finite measure M, we denote its total mass by |M|, and write $M^{\#} = |M|^{-1}M$ for the probability measure proportional to M. We now provide background for the various objects relevant to Theorem 1.1.

2.1. Liouville quantum gravity. We introduce the Gaussian free field (GFF) on various domains. Let $S = \mathbb{R} \times (0, \pi)$ be the infinite strip and let m be the uniform probability measure on $\{0\} \times (0, \pi)$. The Dirichlet inner product is given by $\langle f, g \rangle_{\nabla} = (2\pi)^{-1} \int_{\mathcal{S}} \nabla f \cdot \nabla g$. Consider the space of smooth functions f on S with $\langle f, f \rangle_{\nabla} < \infty$ and $\int_{S} f \, dm = 0$. Let H(S) be its Hilbert space closure with respect to $\langle \cdot, \cdot \rangle_{\nabla}$. Choose an orthonormal basis (f_i) of H(S) and let (ξ_i) be independent standard Gaussian random variables. Then the summation

$$h_{\mathcal{S}} := \sum_{i=1}^{\infty} \xi_i f_i$$

converges in the space of distributions, and we call $h_{\mathcal{S}}$ a GFF on \mathcal{S} normalized so that $\int_{\mathcal{S}} h_{\mathcal{S}} dm = 0$ [DMS21, Section 4.1.4].

Throughout this paper, we fix a choice of LQG parameter $\gamma \in (0,2)$. Suppose $\phi = h_{\mathcal{S}} + g$ where g is a (possibly random) function on $\mathcal{S} \cup \partial \mathcal{S}$ which is continuous at all but finitely many points. For $z \in \mathcal{S} \cup \partial \mathcal{S}$ let $\phi_{\varepsilon}(z)$ be the average of ϕ on $\partial B_{\varepsilon}(z) \cap \mathcal{S}$, and define $\mu_{\phi}^{\varepsilon}(d^2z) := \varepsilon^{\gamma^2/2} e^{\gamma \phi_{\varepsilon}(z)} d^2z$ where d^2z is the Lebesgue measure on \mathcal{S} . The quantum area measure μ_{ϕ} is defined as the almost sure weak limit $\lim_{\varepsilon \to 0} \mu_{\phi}^{\varepsilon}$ [DS11, SW05]. Similarly, we can define the quantum boundary length measure $\nu_{\phi} := \lim_{\varepsilon \to 0} \varepsilon^{\gamma^2/4} e^{\frac{\gamma}{2}\phi_{\varepsilon}(x)} dx$ where dx is the Lebesgue measure on $\partial \mathcal{S}$.

Suppose $f: D \to \widetilde{D}$ is a conformal map between domains D, \widetilde{D} . For a distribution ϕ on D, define

(2.1)
$$f \bullet_{\gamma} \phi = \phi \circ f^{-1} + Q \log |(f^{-1})'|, \qquad Q = \frac{\gamma}{2} + \frac{2}{\gamma}.$$

Consider the set of pairs (D, ϕ) where $D \subset \mathbb{C}$ is open and ϕ is a distribution on D. A quantum surface is an equivalence class of pairs (D, ϕ) where $(D, \phi) \sim_{\gamma} (\widetilde{D}, \widetilde{h})$ if there is a conformal map

²Ewain Gwynne has confirmed in private communication that the techniques in his self-avoiding walk papers with Jason Miller also work for triangulations.

 $f: D \to \widetilde{D}$ such that $\widetilde{\phi} = f \bullet_{\gamma} \phi$, and an *embedding* of the quantum surface is a choice of (D, h) from the equivalence class. This definition is natural because the quantum area and boundary length measures are consistent across elements of an equivalence class: if $(\mathcal{S}, \phi) \sim_{\gamma} (\mathcal{S}, \widetilde{\phi})$ and $f: \mathcal{S} \to \mathcal{S}$ satisfies $\widetilde{\phi} = \phi \circ f^{-1} + Q \log |(f^{-1})'|$, then $\mu_{\widetilde{\phi}} = f_* \mu_{\phi}$ and $\nu_{\widetilde{\phi}} = f_* \nu_{\phi}$ [DS11].

More generally, a loop-decorated quantum surface with m marked points is an equivalence class of tuples $(D, \phi, z_1, \ldots, z_m, \eta)$ with $z_1, \ldots, z_m \in D \cup \partial D$, $\eta : \mathbb{S}^1_{\ell} \to D$ is continuous (i.e., η is a loop on D), and \mathbb{S}^1_{ℓ} is a circle of length $\ell > 0$. We say $(D, \phi, z_1, \ldots, z_m, \eta) \sim_{\gamma} (\widetilde{D}, \widetilde{\phi}, \widetilde{z}_1, \ldots, \widetilde{z}_m, \widetilde{\eta})$ if $\widetilde{\phi} = f \bullet_{\gamma} \phi$, $\widetilde{z}_i = f(z_i)$ for all i, and $\widetilde{\eta}(t) = f(\eta(t+r))$ for some $r \in [0, \ell)$ and all $t \in [0, \ell)$, where we represent \mathbb{S}^1_{ℓ} as the interval $[0, \ell]$ with endpoints identified. We view η as a parametrized and oriented loop with no distinguished starting point. We can similarly define a quantum surface with just m marked points (and no loop).

Now, we recall the radial-lateral decomposition of $h_{\mathcal{S}}$. Let $H_1(\mathcal{S})$ (resp. $H_2(\mathcal{S})$) be the subspace of $H(\mathcal{S})$ comprising functions which are constant (resp. have average zero) on $\{t\} \times (0,\pi)$ for each $t \in \mathbb{R}$. This yields an orthogonal decomposition $H(\mathcal{S}) = H_1(\mathcal{S}) \oplus H_2(\mathcal{S})$.

Definition 2.1. Let $W \ge \frac{\gamma^2}{2}$ and $\beta = Q + \frac{\gamma}{2} - \frac{W}{\gamma^2}$. Let

$$Y_t = \left\{ \begin{array}{ll} B_{2t} - (Q - \beta)t & \text{if } t \ge 0 \\ \widetilde{B}_{-2t} + (Q - \beta)t & \text{if } t < 0 \end{array} \right.,$$

where $(B_s)_{s\geq 0}$ is a standard Brownian motion conditioned on $B_{2s}-(Q-\beta)s<0$ for all s>0, and $(\widetilde{B}_s)_{s\geq 0}$ is an independent copy of $(B_s)_{s\geq 0}$. Let $h^1(z)=Y_{\mathrm{Re}\,z}$ and let $h^2_{\mathcal{S}}$ be the projection of an independent GFF $h_{\mathcal{S}}$ to $H_2(\mathcal{S})$. Set $\widehat{h}=h^1+h^2_{\mathcal{S}}$. Sample an independent real number \mathbf{c} from the measure $[\frac{\gamma}{2}e^{(\beta-Q)c}dc]$ on \mathbb{R} , and let $\phi=\widehat{h}+\mathbf{c}$. Let $\mathcal{M}_2^{\mathrm{disk}}(W)$ be the infinite measure describing the law of $(\mathcal{S},\phi,-\infty,+\infty)/\sim_{\gamma}$. We call a sample from $\mathcal{M}_2^{\mathrm{disk}}(W)$ a quantum disk with two insertions of weight W.

Weight W quantum disks have two marked boundary points. The case W=2 is special since these two points are quantum typical in the following sense.

Proposition 2.2 ([DMS21, Proposition A.8]). Sample $(S, \phi, -\infty, +\infty)/\sim_{\gamma}$ from $\mathcal{M}_2^{\text{disk}}(2)$, then sample independent points $x_1, x_2 \in \partial S$ from the probability measure proportional to ν_{ϕ} . Then the law of $(S, \phi, x_1, x_2)/\sim_{\gamma}$ is $\mathcal{M}_2^{\text{disk}}(2)$.

This motivates the following definition.

Definition 2.3. Sample $(S, \phi, -\infty, +\infty)/\sim_{\gamma}$ from the weighted measure $\nu_{\phi}(\partial S)^{-2}\mathcal{M}_{2}^{\text{disk}}(2)$. Then we call $(S, \phi)/\sim_{\gamma}$ a quantum disk, and denote its law by QD.

Here is a useful perspective on Proposition 2.2 and Definition 2.3. Roughly speaking, given a sample $(S, \phi)/\sim_{\gamma}$ from QD, if we "sample two points from ν_{ϕ} ", then the resulting quantum surface with two marked points has law $\mathcal{M}_2(2)$. Since ν_{ϕ} is a non-probability measure with total mass $\nu_{\phi}(\partial S)$, the sampling operation should induce a weighting by $\nu_{\phi}(\partial S)^2$; this explains the factor $\nu_{\phi}(\partial S)^{-2}$ in Definition 2.3.

Lemma 2.4. The law of the quantum boundary length $\nu_{\phi}(\partial S)$ of a sample $(S, \phi)/\sim_{\gamma}$ from QD is $1_{\ell>0}C\ell^{-\frac{4}{\gamma^2}-2}d\ell$ for some C>0.

Proof. [AHS22, Lemma 3.3] implies that the total boundary length of a sample from $\mathcal{M}_2(2)$ has law $1_{\ell>0}C\ell^{-\frac{4}{\gamma^2}}d\ell$. By Definition 2.3, weighting by ℓ^{-2} gives the corresponding result for QD.

Consequently, we can define a disintegration $\{QD(\ell)\}_{\ell>0}$ of QD on its quantum boundary length, i.e. $QD = \int_0^\infty QD(\ell) d\ell$ for measures $QD(\ell)$ supported on the space of quantum surfaces with

boundary length ℓ . This only specifies QD(ℓ) for a.e. ℓ , but by continuity we can canonically define QD(ℓ) for all ℓ ; see e.g. [DMS21, Section 4.5] or [AHS20b, Section 2.6].

Define the horizontal cylinder $\mathcal{C} := \mathbb{R} \times [0, 2\pi]/\sim$ by identifying $(x, 0) \sim (x, 2\pi)$ for all $x \in \mathbb{R}$. Let m be the uniform measure on $(\{0\} \times [0, 2\pi])/\sim$, and let $H(\mathcal{C})$ be the Hilbert space closure of smooth compactly-supported functions on \mathcal{C} under the Dirichlet inner product. Then, as for \mathcal{S} , we define $h_{\mathcal{C}} = \sum_{i} \alpha_{i} \xi_{i}$ where (ξ_{i}) is an orthonormal basis of $H(\mathcal{C})$ and (α_{i}) are independent standard Gaussians. We call $h_{\mathcal{C}}$ the GFF on \mathcal{C} normalized so that $\int_{\mathcal{C}} h_{\mathcal{C}} dm = 0$.

As before, we can decompose $H(\mathcal{C}) = H_1(\mathcal{C}) \oplus H_2(\mathcal{C})$ where $H_1(\mathcal{C})$ (resp. $H_2(\mathcal{C})$) is the subspace of functions which are constant (resp. have average zero) on $(\{t\} \times [0, 2\pi])/\sim$ for all $t \in \mathbb{R}$.

Definition 2.5. Let W > 0 and $\alpha = Q - \frac{W}{2\alpha}$. Let

$$Y_t = \begin{cases} B_t - (Q - \alpha)t & \text{if } t \ge 0\\ \widetilde{B}_{-t} + (Q - \alpha)t & \text{if } t < 0 \end{cases},$$

where $(B_s)_{s\geq 0}$ is a standard Brownian motion conditioned on $B_s - (Q - \alpha)s < 0$ for all s > 0, and $(\widetilde{B}_s)_{s\geq 0}$ is an independent copy of $(B_s)_{s\geq 0}$. Let $h^1(z) = Y_{\operatorname{Re} z}$ and let $h^2_{\mathcal{C}}$ be the projection of an independent GFF $h_{\mathcal{C}}$ to $H_2(\mathcal{C})$. Set $\widehat{h} = h^1 + h^2_{\mathcal{C}}$. Sample an independent real number \mathbf{c} from the measure $[\frac{\gamma}{2}e^{2(\alpha-Q)c}dc]$ on \mathbb{R} , and let $\phi = \widehat{h} + \mathbf{c}$. Let $\mathcal{M}_2^{\operatorname{sph}}(W)$ be the infinite measure describing the law of $(\mathcal{C}, \phi, -\infty, +\infty)/\sim_{\gamma}$. We call a sample from $\mathcal{M}_2^{\operatorname{sph}}(W)$ a quantum sphere with two insertions of weight W.

The weight $W = 4 - \gamma^2$ is special because the two marked points are independent samples from the quantum area measure [DMS21, Proposition A.13], so the following definition is natural.

Definition 2.6. Sample $(C, \phi, -\infty, +\infty)/\sim_{\gamma}$ from the weighted measure $\mu_{\phi}(C)^{-2}\mathcal{M}_{2}^{\mathrm{sph}}(4-\gamma^{2})$. Then we call $(C, \phi)/\sim_{\gamma}$ a quantum sphere, and denote its law by QS.

Remark 2.7. For $W < \gamma Q$ and compact $I \subset \mathbb{R}$ the $\mathcal{M}_2^{\mathrm{disk}}(W)$ -mass of the event {left boundary length $\in I$ } is finite (so one can condition on boundary length), but for $W = \gamma Q$ this mass is infinite [AHS20b, Lemma 2.16]. The same calculation shows that $\mathcal{M}_2^{\mathrm{sph}}(W)[area \in I] < \infty$ if and only if W < 4.

Finally, we will need an area-weighted variant of $\mathcal{M}_2^{\mathrm{sph}}(W)$.

Definition 2.8. Fix W > 0 and let $(C, \phi, -\infty, +\infty)$ be an embedding of a sample from the quantum-area-weighted measure $\mu_{\phi}(C)\mathcal{M}_{2}^{\mathrm{sph}}(W)$. Given ϕ , sample \mathbf{z} from the probability measure proportional to μ_{ϕ} . We write $\mathcal{M}_{2,\bullet}^{\mathrm{sph}}(W)$ for the law of the marked quantum surface $(C, \phi, -\infty, +\infty, \mathbf{z})/\sim_{\gamma}$.

2.2. The Liouville field. In this section we recall the Liouville field which was constructed in [DKRV16]. Let $\exp: \mathcal{C} \to \widehat{\mathbb{C}}$ be the exponential map $z \mapsto e^z$. Let $h_{\mathcal{C}}$ be the GFF on the cylinder as defined in the previous section, and let $h_{\widehat{\mathbb{C}}} = h_{\mathcal{C}} \circ \exp$. Then $h_{\widehat{\mathbb{C}}}$ is the GFF on $\widehat{\mathbb{C}}$ with average zero on the unit circle. We write $P_{\widehat{\mathbb{C}}}$ for the law of $h_{\widehat{\mathbb{C}}}$. Its covariance kernel is

$$G_{\widehat{\mathbb{C}}}(z,w) = -\log|z-w| + \log|z|_+ + \log|w|_+, \qquad |z|_+ := \max(|z|,1).$$

Definition 2.9. Sample (h, \mathbf{c}) from $P_{\widehat{\mathbb{C}}} \times [e^{-2Qc} dc]$ and let $\phi = h - 2Q \log |\cdot|_+ + \mathbf{c}$. We call ϕ the Liouville field on $\widehat{\mathbb{C}}$ and denote its law by $\mathrm{LF}_{\widehat{\mathbb{C}}}$.

For a finite collection of weights α_i and points $z_i \in \mathbb{C}$, we want to define " $LF_{\widehat{\mathbb{C}}}^{(\alpha_i,z_i)_i} = \prod_i e^{\alpha_i \phi(z_i)} LF_{\widehat{\mathbb{C}}}(d\phi)$ ". This can be understood via regularization and renormalization, see e.g. [AHS22, Lemma 2.6]. We give a direct definition below.

Definition 2.10. Let $(\alpha_i, z_i) \in \mathbb{R} \times \mathbb{C}$ for i = 1, ..., m, where $m \geq 1$ and the z_i are distinct. Sample (h, \mathbf{c}) from $C_{\widehat{\mathbb{C}}}^{(\alpha_i, z_i)_i} P_{\widehat{\mathbb{C}}} \times [e^{(\sum_i \alpha_i - 2Q)c} dc]$ where

$$C_{\widehat{\mathbb{C}}}^{(\alpha_i, z_i)_i} = \prod_{i=1}^m |z_i|_+^{-\alpha_i(2Q - \alpha_i)} e^{\sum_{j=i+1}^m \alpha_i \alpha_j G_{\widehat{\mathbb{C}}}(z_i, z_j)}.$$

Let $\phi = h - 2Q \log |\cdot|_+ + \sum_{i=1}^m \alpha_i G_{\widehat{\mathbb{C}}}(\cdot, z_i) + \mathbf{c}$. We call ϕ the Liouville field on $\widehat{\mathbb{C}}$ with insertions (α_i, z_i) , and denote its law by $LF_{\widehat{\mathbb{C}}}^{(\alpha_i, z_i)_i}$.

For a conformal automorphism $f: \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ and a measure M on the space of distributions on \mathbb{C} , let f_*M be the pushforward of M under the map $\phi \mapsto \phi \circ f^{-1} + Q \log |(f^{-1})'|$. The following change-of-coordinates result is [DKRV16, Theorem 3.5] with different notation. We present the version stated in [AHS22, Proposition 2.29].

Proposition 2.11 ([DKRV16, Theorem 3.5]). For $\alpha \in \mathbb{R}$ let $\Delta_{\alpha} := \frac{\alpha}{2}(Q - \frac{\alpha}{2})$. Let f be a conformal automorphism of $\widehat{\mathbb{C}}$ and let $(\alpha_i, z_i) \in \mathbb{R} \times \mathbb{C}$ satisfy $f(z_i) \neq \infty$ for i = 1, ..., m. Then

$$\mathrm{LF}_{\widehat{\mathbb{C}}} = f_* \mathrm{LF}_{\widehat{\mathbb{C}}}, \quad and \quad \mathrm{LF}_{\widehat{\mathbb{C}}}^{(\alpha_i, f(z_i))_i} = \prod_{i=1}^m |f'(z_i)|^{-2\Delta_{\alpha_i}} f_* \mathrm{LF}_{\widehat{\mathbb{C}}}^{(\alpha_i, z_i)_i}.$$

As the next lemma illustrates, sampling points from quantum measures of the Liouville field corresponds to adding insertions to the Liouville field. We recall the proof for the reader's convenience since a closely related argument will be used later.

Lemma 2.12 ([AHS22, Lemma 2.31]). We have $\mu_{\phi}(d^2u) \operatorname{LF}_{\widehat{\mathbb{C}}}^{(\alpha_i, z_i)_i}(d\phi) = \operatorname{LF}_{\widehat{\mathbb{C}}}^{(\gamma, u), (\alpha_i, z_i)_i}(d\phi) d^2u$.

Proof. Sample $h \sim P_{\widehat{\mathbb{C}}}$. Let $h_{\varepsilon}(u)$ be the average of h on $\partial B_{\varepsilon}(u)$ and write $G_{\widehat{\mathbb{C}},\varepsilon}(z,u) := \mathbb{E}[h(z)h_{\varepsilon}(u)]$ Let f be a non-negative continuous function on the Sobolev space $H^{-1}(\mathbb{C})$, and g a non-negative measurable function on \mathbb{R} . Girsanov's theorem gives

$$\mathbb{E}\Big[f(h)\varepsilon^{\gamma^2/2}e^{\gamma h_\varepsilon(u)}\Big] = \mathbb{E}\Big[f(h+\gamma G_{\widehat{\mathbb{C}},\varepsilon}(\cdot,u))\Big]\mathbb{E}[\varepsilon^{\gamma^2/2}e^{\gamma h_\varepsilon(u)}].$$

With $\mu_h^{\varepsilon}(d^2u) := \varepsilon^{\gamma^2/2} e^{\gamma h_{\varepsilon}(u)} d^2u$ and $\rho_{\varepsilon}(u) := \mathbb{E}[\varepsilon^{\gamma^2/2} e^{\gamma h_{\varepsilon}(u)}]$, integrating against $g(u) d^2u$ gives

$$\mathbb{E}\left[\int_{\mathbb{C}} f(h)g(u)\mu_h^{\varepsilon}(d^2u)\right] = \int_{\mathbb{C}} \mathbb{E}\left[f(h+\gamma G_{\widehat{\mathbb{C}},\varepsilon}(\cdot,u))\right]g(u)\rho_{\varepsilon}(u)\,d^2u.$$

Taking the $\varepsilon \to 0$ limit yields, with $\rho(u)$ defined by $\rho(u) d^2 u = \mathbb{E}[\mu_h(d^2 u)]$,

$$\mathbb{E}\left[\int f(h)g(u)\,\mu_h(d^2u)\right] = \int \mathbb{E}[f(h+\gamma G_{\widehat{\mathbb{C}}}(\cdot,u))]g(u)\rho(u)\,d^2u.$$

See, e.g., $[BP21, Section\ 2.4]$ or $[AHS22, Lemma\ 2.31]$ for details on taking this limit.

Let $c \in \mathbb{R}$ and $q(z) = \sum_i \alpha_i G_{\widehat{\mathbb{C}}}(z, z_i) - 2Q \log |z|_+$. For \widetilde{f} any non-negative continuous function on $H^{-1}(\mathbb{C})$ and \widetilde{g} any non-negative measurable function on \mathbb{R} , choose $f = \widetilde{f}(\cdot + q + c)$ and $g = e^{\gamma(q+c)}\widetilde{g}$. The above equation, together with $\mu_{h+p} = e^{\gamma p}\mu_h$ for any continuous function $p: \mathbb{C} \to \mathbb{R}$, gives

$$\mathbb{E}\left[\int \widetilde{f}(h+q+c)\widetilde{g}(u)\mu_{h+q+c}(d^2u)\right] = \int \mathbb{E}[\widetilde{f}(h+\gamma G_{\widehat{\mathbb{C}}}(\cdot,u)+q+c)]\widetilde{g}(u)e^{\gamma q(u)+\gamma c}\rho(u)\,d^2u.$$

On the other hand, we have $C_{\widehat{\mathbb{C}}}^{(\gamma,u),(\alpha_i,z_i)} = C_{\widehat{\mathbb{C}}}^{(\alpha_i,z_i)_i}C_{\widehat{\mathbb{C}}}^{(\gamma,u)}e^{\gamma q(u)+2\gamma Q\log|u|_+} = C_{\widehat{\mathbb{C}}}^{(\alpha_i,z_i)_i}e^{\gamma q(u)}\rho(u)$, where the first equality holds by definition and the second follows from a direct calculation; see

[AHS22, Lemma 2.12] for a similar calculation. Thus, multiplying the previous identity by $C_{\widehat{\mathbb{C}}}^{(\alpha_i, z_i)_i}$ gives

$$C_{\widehat{\mathbb{C}}}^{(\alpha_i, z_i)_i} \mathbb{E}\left[\int \widetilde{f}(h+q+c)\widetilde{g}(u)\mu_{h+q+c}(d^2u)\right] = \int C_{\widehat{\mathbb{C}}}^{(\gamma, u), (\alpha_i, z_i)_i} \mathbb{E}\left[\widetilde{f}(h+\gamma G_{\widehat{\mathbb{C}}}(\cdot, u)+q+c)\right]\widetilde{g}(u)e^{\gamma c} d^2u.$$

Multiplying by $e^{(\sum_i \alpha_i - 2Q)c}$ and integrating over c gives

$$\mathrm{LF}_{\widehat{\mathbb{C}}}^{(\alpha_{i},z_{i})_{i}}\left[\int_{\mathbb{C}}\widetilde{f}(\phi)\widetilde{g}(u)\mu_{\phi}(d^{2}u)\right] = \int_{\mathbb{C}}\mathrm{LF}_{\widehat{\mathbb{C}}}^{(\gamma,u),(\alpha_{i},z_{i})_{i}}[\widetilde{f}(\phi)]\widetilde{g}(u)\,d^{2}u.$$

The functions \widetilde{f} , \widetilde{g} are arbitrary so the desired result holds.

We need the following Liouville field description of $\mathcal{M}^{\mathrm{sph}}_{2,\bullet}(W)$.

Proposition 2.13. Fix W > 0, let $\alpha = Q - \frac{W}{2\gamma}$ and sample ϕ from $\frac{2\pi\gamma}{(Q-\alpha)^2} LF_{\widehat{\mathbb{C}}}^{(\alpha,0),(\alpha,1),(\gamma,-1)}$. Then the law of $(\widehat{\mathbb{C}}, \phi, 0, 1, -1)/\sim_{\gamma}$ is $\mathcal{M}_{2,\bullet}^{\mathrm{sph}}(W)$.

Proof. [AHS22, Proposition B.7] describes the field of $\mathcal{M}^{\mathrm{sph}}_{2,\bullet}(W)$ in terms of the Liouville field on \mathcal{C} , then [AHS22, Lemma B.4] gives the coordinate change from \mathcal{C} to $\widehat{\mathbb{C}}$.

2.3. Uniform embedding of quantum surfaces. Let $\operatorname{conf}(\widehat{\mathbb{C}})$ denote the space of automorphisms of the Riemann sphere $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. Being a locally compact Lie group, it has a right-invariant Haar measure which is unique modulo multiplicative constant, and since $\operatorname{conf}(\widehat{\mathbb{C}})$ is unimodular the measure is also left-invariant. Let $\mathbf{m}_{\widehat{\mathbb{C}}}$ be such a Haar measure. The following gives an explicit description of $\mathbf{m}_{\widehat{\mathbb{C}}}$; see e.g. [AHS22, Lemma 2.28].

Lemma 2.14. Let \mathfrak{f} be sampled from $\mathbf{m}_{\widehat{\mathbb{C}}}$. Then there is a constant $C \in (0, \infty)$ such that the law of $(\mathfrak{f}(0), \mathfrak{f}(1), \mathfrak{f}(-1))$ is $C|(p-q)(q-r)(r-p)|^{-2} d^2p d^2q d^2r$.

Suppose M is a measure on the space of quantum surfaces which can be embedded in $\widehat{\mathbb{C}}$. Sample a pair (\mathfrak{f}, S) from the product measure $\mathbf{m}_{\widehat{\mathbb{C}}} \times M$, and let ϕ_0 be a distribution on \mathbb{C} chosen in a way measurable with respect to S such that $S = (\widehat{\mathbb{C}}, \phi_0)/\sim_{\gamma}$. We define $\mathbf{m}_{\widehat{\mathbb{C}}} \times M$ to be the law of $\mathfrak{f} \bullet_{\gamma} \phi_0$. We call $\mathbf{m}_{\widehat{\mathbb{C}}} \times M$ the uniform embedding of M. Note that the definition of uniform embedding does not depend on the choice of ϕ_0 . Recall that QS is the law of the quantum sphere from Definition 2.6.

Proposition 2.15 ([AHS22, Theorem 1.2]). There is a constant C such that $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes \mathrm{QS} = C \cdot \mathrm{LF}_{\widehat{\mathbb{C}}}$.

2.4. Conformal welding. Let $\kappa > 0$ and let (D, p, q) be a simply-connected domain with two marked boundary points. SLE_{κ} is a conformally invariant random curve in D from p to q introduced by Schramm [RS05], which describes the scaling limits of many statistical physics models. When $\kappa < 4$, almost surely SLE_{κ} is simple and only intersects ∂D at $\{p,q\}$. We will also need a spherical variant of SLE : for distinct points $p,q \in \mathbb{C}$ and $\rho > -2$, there is a random curve from p to q called whole-plane $\mathrm{SLE}_{\kappa}(\rho)$, see e.g. [MS17, Section 2.1.3] for its definition.

For $\kappa \in (0,8)$ and distinct points $p,q \in \mathbb{C}$, the two-sided whole-plane SLE, which we denote by $SLE_{\kappa}^{p \rightleftharpoons q}$, is the probability measure on pairs of curves (η_1, η_2) on \mathbb{C} connecting p and q where η_1 is a whole-plane $SLE_{\kappa}(2)$ from p to q, and conditioning on η_1 , the curve η_2 is a chordal SLE_{κ} on the complement of η_1 from q to p. This pair of curves (η_1, η_2) satisfies the following resampling property: conditioning on one, the other has the law of chordal SLE_{κ} in the complement, see e.g. [Zha21, Section 2.2].

We need a special case of [AHS20b, Theorem 2.4]. Let $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be the Riemann sphere. Let $\{\mathcal{M}_2^{\text{disk}}(2;\ell_1,\ell_2)\}_{\ell_1,\ell_2}$ be a disintegration of $\mathcal{M}_2^{\text{disk}}(2)$ on its two boundary arc lengths, i.e. $\mathcal{M}_2^{\text{disk}}(2) = \iint \mathcal{M}_2^{\text{disk}}(2;\ell_1,\ell_2) d\ell_1 d\ell_2$, and a sample from $\mathcal{M}_2^{\text{disk}}(2;\ell_1,\ell_2)$ a.s. has boundary lengths (ℓ_1,ℓ_2) .

Proposition 2.16. Fix distinct $p, q \in \mathbb{C}$ and let $(\widehat{\mathbb{C}}, \phi, p, q)$ be an embedding of a sample from $\mathcal{M}_2^{\mathrm{sph}}(4)$. Independently sample (η_1, η_2) from the probability measure $\mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}$, and let D_1 and D_2 be the connected components of $\widehat{\mathbb{C}} \setminus (\eta_1 \cup \eta_2)$ lying to the left and right of η_1 respectively. Then there is a constant C such that the joint law of $(D_1, \phi, p, q)/\sim_{\gamma}$ and $(D_2, \phi, p, q)/\sim_{\gamma}$ is

$$C \int_0^\infty \int_0^\infty \mathcal{M}_2^{\text{disk}}(2;\ell_1,\ell_2) \times \mathcal{M}_2^{\text{disk}}(2;\ell_2,\ell_1) d\ell_1 d\ell_2.$$

The above statement of Proposition 2.16 is in terms of cutting a sphere to get two disks. It can be equivalently expressed in terms of gluing two disks to get a loop-decorated sphere. For fixed $\ell_1, \ell_2 > 0$, a pair of quantum disks sampled from $\mathcal{M}_2^{\text{disk}}(2; \ell_1, \ell_2) \times \mathcal{M}_2^{\text{disk}}(2; \ell_2, \ell_1)$ can be conformally welded along their boundary arcs according to quantum boundary length, to get a quantum surface with the sphere topology decorated by two points and a loop passing through them. For more details on the conformal welding of quantum surfaces, see e.g. [She16, DMS21, GHS19], and see [AHS20b] for more information on the conformal welding of quantum disks.

We now give a more precise definition of the measure Weld(QD, QD) appearing in Theorem 1.1. Let $\ell > 0$ and let $(\mathcal{D}_1, \mathcal{D}_2) \sim \mathrm{QD}(\ell) \times \mathrm{QD}(\ell)$. For i = 1, 2, let $\phi_i : \mathbb{S}^1_\ell \to \mathcal{D}_i$ be a parametrization of the boundary of \mathcal{D}_i according to its quantum boundary length such that ϕ_i traces the boundary in counterclockwise direction when the disk is embedded in \mathbb{D} . Namely, for 0 < s < t < 1, $\phi_i([s,t])$ is an arc on the boundary of \mathcal{D}_i with quantum length t-s, where we represent \mathbb{S}^1_ℓ as the interval $[0,\ell]$ with endpoints identified. Let U be a uniform point on \mathbb{S}^1_ℓ independent of everything else. Let Weld($\mathcal{D}_1, \mathcal{D}_2$) be the curve-decorated quantum surface obtained by conformally welding \mathcal{D}_1 and \mathcal{D}_2 along their boundaries where $\phi_1(t)$ is identified with $\phi_2(U-t)$ for all $t \in \mathbb{S}^1_\ell$. In words, Weld($\mathcal{D}_1, \mathcal{D}_2$) means we conformally weld \mathcal{D}_1 and \mathcal{D}_2 according to their boundary length uniformly at random. Let Weld($\mathrm{QD}(\ell), \mathrm{QD}(\ell)$) be the law of Weld($\mathcal{D}_1, \mathcal{D}_2$), and define Weld($\mathrm{QD}, \mathrm{QD}(\ell) = \int_0^\infty \ell \cdot \mathrm{Weld}(\mathrm{QD}(\ell), \mathrm{QD}(\ell))$.

2.5. **Zhan's construction of the SLE loop measure.** Given a simple loop η and some $d \in [0, 2]$, let $\operatorname{Cont}_{\eta,\varepsilon}$ be ε^{d-2} times Lebesgue area measure restricted to the ε -neighborhood of η . If $\lim_{\varepsilon \to 0} \operatorname{Cont}_{\eta,\varepsilon}$ exists for the weak topology then we denote the limit by $\operatorname{Cont}_{\eta}$ and call it the d-dimensional $\operatorname{Minkowski}$ content of η .

We can view $\mathrm{SLE}_{\kappa}^{p\rightleftharpoons q}$ as a measure on oriented loops by concatenating η_1 and η_2 . Given a loop η sampled from $\mathrm{SLE}_{\kappa}^{p\rightleftharpoons q}$, with probability 1 the dimension of η is $d=1+\frac{\kappa}{8}$ [Bef08] and its d-dimensional Minkowski content Cont_{η} exists [LR15]. The *(unrooted) SLE loop measure* $\mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ on $\mathbb C$ is an infinite measure on oriented loops defined by (see [Zha21, Theorem 4.2])

(2.2)
$$\operatorname{SLE}_{\kappa}^{\operatorname{loop}}(d\eta) = |\operatorname{Cont}_{\eta}|^{-2} \iint_{\mathbb{C} \times \mathbb{C}} |p - q|^{-2(2-d)} \operatorname{SLE}_{\kappa}^{p \rightleftharpoons q}(d\eta) d^{2} p d^{2} q.$$

The operation of forgetting p and q in (2.2) is natural, since given η , the points p, q are conditionally independent points sampled from the Minkowski content measure Cont_{η} on η ; precisely, [Zha21, Theorem 4.2 (i)] states

(2.3)
$$\operatorname{SLE}_{\kappa}^{\operatorname{loop}}(d\eta)\operatorname{Cont}_{\eta}(dp)\operatorname{Cont}_{\eta}(dq) = |p - q|^{\frac{\gamma^2}{4} - 2}\operatorname{SLE}_{\kappa}^{p \rightleftharpoons q}(d\eta) d^2p d^2q.$$

For $\kappa \in (0,4]$, Zhan [Zha21] shows that ${\rm SLE}_{\kappa}^{\rm loop}$ is an example of a Malliavin-Kontsevich-Suhov (MKS) loop measure.

2.6. The Gromov-Hausdorff-Prokhorov-uniform metric. In this subsection we will define precisely the space of compact loop-decorated metric measure spaces and the Gromov-Hausdorff-Prokhorov-uniform metric. This, along with definitions in Section 3, will make precise the statement of Theorem 1.2. We remark that the analogous definitions in the setting of curve-decorated metric measure spaces were first made in [GM17]; see also [Gro99, BBI01, GPW09, Mie09, ADH13].

For a metric space (X,d) let $C_0(X)$ denote the space of parametrized and oriented loops on X with no distinguished starting point. More precisely, identifying the circle \mathbb{S}^1_ℓ of length ℓ with the interval $[0,\ell]$ with endpoints identified, $C_0(X)$ is the space of continuous functions $\eta: \mathbb{S}^1_\ell \to X$, where we identify η and $\tilde{\eta}$ if $\tilde{\eta}(t) = \eta(t+r)$ for some $r \in [0,\ell)$ and all $t \in [0,\ell)$. Let d_d^H denote the d-Hausdorff metric on compact subsets of X and let d_d^P denote the d-Prokhorov metric on finite measures on X. Finally, let d_d^U denote the d-uniform metric on $C_0(X)$, i.e.,

$$\mathbf{d}_d^{\mathrm{U}}(\eta,\widetilde{\eta}) = \inf_{r \in [0,\ell)} \sup_{t \in [0,\ell)} d(\eta(t),\widetilde{\eta}(t+r)).$$

Let \mathbb{M}^{GHPU} be the set of compact loop-decorated metric measure spaces, i.e., \mathbb{M}^{GHPU} is the set of 4-tuples $\mathfrak{X} = (X, d, \mu, \eta)$ where (X, d) is a compact metric space, μ is a finite Borel measure on X, and $\eta \in C_0(X)$. Given elements $\mathfrak{X}_1 = (X_1, d_1, \mu_1, \eta_1)$ and $\mathfrak{X}_2 = (X_2, d_2, \mu_2, \eta_2)$ of \mathbb{M}^{GHPU} , we define their Gromov-Hausdorff-Prokhorov-uniform (GHPU) distance by

$$\mathbf{d}^{\mathrm{GHPU}}(\mathfrak{X}_{1},\mathfrak{X}_{2}) = \inf_{(W,d),\iota_{1},\iota_{2}} \mathbf{d}^{\mathrm{H}}_{D}(\iota_{1}(X_{1}),\iota_{2}(X_{2})) + \mathbf{d}^{\mathrm{P}}_{D}((\iota_{1})_{*}\mu_{1},(\iota_{2})_{*}\mu_{2}) + \mathbf{d}^{\mathrm{U}}_{D}(\iota_{1}\circ\eta_{1},\iota_{2}\circ\eta_{2}),$$

where we take the infimum over all compact metric spaces (W, D) and isometric embeddings $\iota_1: X_1 \to W$ and $\iota_2: X_2 \to W$. It is shown in [GM17] that this defines a complete separable metric in the setting of curve-decorated (rather than loop-decorated) metric measure spaces if we identify two elements of this space which differ by a measure- and curve-preserving isometry. The analogous statement holds in the setting of loop-decorated metric measure spaces since we obtain a loop by considering a curve that forms a loop and identifying two such curves which differ by a time shift.

3. The SLE loop via conformal welding: proof of Theorem 1.1

In Section 3.1 we prove Theorem 1.1 modulo Proposition 3.2, whose proof is given in Section 3.2.

3.1. **Proof of Theorem 1.1.** Fix $\gamma \in (0,2)$ and $\kappa = \gamma^2 \in (0,4)$. Let $M = \mathrm{QS} \otimes \mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ be the law of the loop-decorated quantum surface called the MKS-loop-decorated quantum sphere with parameter γ . Namely, if (S,η) is sampled from $\mathrm{QS} \times \mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ and ϕ is a distribution on $\mathbb C$ chosen in a way measurable with respect to S such that $S = (\widehat{\mathbb C}, \phi)/\sim_{\gamma}$, then M is the law of the loop-decorated quantum surface $(\widehat{\mathbb C}, \phi, \eta)/\sim_{\gamma}$. Theorem 1.1 asserts that $M = C\mathrm{Weld}(\mathrm{QD}, \mathrm{QD})$ for some constant $C \in (0, \infty)$. To prove it, we need a variant of Proposition 2.16 where the marked points on the quantum disks are forgotten.

Lemma 3.1. Let $(\widehat{\mathbb{C}}, h, p, q)$ be an embedding of a sample from $\mathcal{M}_2^{\mathrm{sph}}(4)$. Let (η_1, η_2) be a sample from $\mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}$ independent of h, and let η be the oriented loop obtained by concatenating η_1 and η_2 . Then viewed as a loop-decorated quantum surface the law of $(\widehat{\mathbb{C}}, h, \eta)$ equals $C \int_0^{\infty} \ell^3 \cdot \mathrm{Weld}(\mathrm{QD}(\ell), \mathrm{QD}(\ell)) \, d\ell$ for some constant $C \in (0, \infty)$.

Proof. Let F be the map that forgets the marked points of a quantum surface. By Definition 2.3,

$$\int_0^\infty \ell^2 \mathrm{QD}(\ell) \, d\ell = F_* \mathcal{M}_2^{\mathrm{disk}}(2) = F_* \int_0^\infty \int_0^\infty \mathcal{M}_2^{\mathrm{disk}}(2; \ell_1, \ell_2) \, d\ell_1 \, d\ell_2$$
$$= \int_0^\infty \int_0^\ell F_* \mathcal{M}_2^{\mathrm{disk}}(2; \ell_1, \ell - \ell_1) \, d\ell_1 \, d\ell.$$

In the last equality, we change variables $\ell = \ell_1 + \ell_2$ so $1_{\ell_1,\ell_2>0} d\ell_1 d\ell_2$ corresponds to $1_{\ell>\ell_1>0} d\ell_1 d\ell$. By Proposition 2.2, the measure $F_*\mathcal{M}_2^{\text{disk}}(2;\ell_1,\ell-\ell_1)$ does not depend on the choice of ℓ_1 , and hence must equal $\ell \text{QD}(\ell)$.

Let D_1 and D_2 be the connected components of $\widehat{\mathbb{C}} \setminus \eta$. By Proposition 2.16, the law of the pair of marked quantum surfaces $((D_1, h, p, q)/\sim_{\gamma}, (D_2, h, p, q)/\sim_{\gamma})$ equals

$$C\int_0^\infty \int_0^\ell \mathcal{M}_2^{\text{disk}}(2;\ell_1,\ell-\ell_1) \times \mathcal{M}_2^{\text{disk}}(2;\ell-\ell_1,\ell_1) d\ell_1 d\ell.$$

Applying F to both sides and using $F_*\mathcal{M}_2^{\text{disk}}(2;\ell_1,\ell-\ell_1) = \ell \text{QD}(\ell)$, the law of $((D_1,h)/\sim_{\gamma},(D_2,h)/\sim_{\gamma})$ is $C\int_0^{\infty}\ell^3\text{QD}(\ell)^2\,d\ell$. Finally, since the conformal welding of $(D_1,h,p,q)/\sim_{\gamma}$ and $(D_2,h,p,q)/\sim_{\gamma}$ is determined by the locations of the marked points, and the marked points on each disk are uniformly chosen from quantum length measure (Proposition 2.2), the conformal welding of $(D_1,h)/\sim_{\gamma}$ to $(D_2,h)/\sim_{\gamma}$ is uniform, as desired.

Similarly as for the proof of Proposition 2.15 from [AHS22], we prove Theorem 1.1 by adding three marked points. Suppose $(\widehat{\mathbb{C}}, h, \eta)$ is an embedding of a sample from M weighted by $\mu_h(\mathbb{C})$ times the square of the quantum length of η . Given (h, η) , independently sample p, q from the probability measure proportional to the quantum length measure on η , and r from the probability measure proportional to the quantum area measure, so $p, q \in \eta$ and $r \in \mathbb{C}$. Let M_3 be the law of $(\widehat{\mathbb{C}}, h, \eta, p, q, r)$ viewed as a loop-decorated quantum surface with three marked points. Recall that $\mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}$ is the law of a two-sided whole plane SLE_{κ} from p to q. Moreover, we view a sample (η_1, η_2) from $\mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}$ as an oriented loop by concatenating η_1 with η_2 . Recall $\mathcal{M}_{2,\bullet}^{\mathrm{sph}}(W)$ from Definition 2.8. The following proposition describes M_3 in terms of $\mathcal{M}_{2,\bullet}^{\mathrm{sph}}(W)$ and $\mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}$.

Proposition 3.2. Let $(\widehat{\mathbb{C}}, h, p, q, r)$ be an embedding of a sample from $\mathcal{M}^{\mathrm{sph}}_{2, \bullet}(4)$. Independently sample η from $\mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}$. Let \widetilde{M}_3 be the law of $(\widehat{\mathbb{C}}, h, \eta, p, q, r)$ viewed as a loop-decorated quantum surface with three marked points. Then there exists a constant $C \in (0, \infty)$ such that $M_3 = C\widetilde{M}_3$.

Proof of Theorem 1.1 given Proposition 3.2. See Figure 1. Fix $p,q,r \in \mathbb{C}$. Sample a decorated quantum surface from \widetilde{M}_3 and embed it as $(\widehat{\mathbb{C}},\phi,\eta,p,q,r)$. Let (A,L) be its quantum area and the quantum length of its loop. By Definition 2.8, after weighting by A^{-1} the law of $(\widehat{\mathbb{C}},\phi,\eta,p,q)$ is $\mathcal{M}_2^{\mathrm{sph}}(4) \otimes \mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}$, then by Lemma 3.1 the law of $(\widehat{\mathbb{C}},\phi,\eta)$ is $C \int_0^{\infty} \ell^3 \mathrm{Weld}(\mathrm{QD}(\ell),\mathrm{QD}(\ell)) \, d\ell$ for some C>0. Further weighting by L^{-2} , the law of $(\widehat{\mathbb{C}},\phi,\eta)$ is $C \int_0^{\infty} \ell \mathrm{Weld}(\mathrm{QD}(\ell),\mathrm{QD}(\ell)) \, d\ell = C \mathrm{Weld}(\mathrm{QD},\mathrm{QD})$.

By the definition of M_3 , if we embed a sample from M_3 as $(\widehat{\mathbb{C}}, \phi, \eta, p, q, r)$ and let (A, L) be its quantum area and the quantum length of its loop, then the law of $(\widehat{\mathbb{C}}, \phi, \eta)$ after weighting by $A^{-1}L^{-2}$ is QS \otimes SLE_{κ}^{loop}.

Proposition 3.2 states that M_3 and \widetilde{M}_3 agree up to multiplicative constant, so by the above two paragraphs $QS \otimes SLE_{\kappa}^{loop}$ and Weld(QD,QD) agree up to multiplicative constant.

3.2. **Proof of Proposition 3.2 via the uniform embedding.** We will prove Proposition 3.2 by first establishing Proposition 3.3, which gives its counterpart under the uniform embedding. As for QS in Section 2.3, suppose we sample $(\mathfrak{f},(\widehat{\mathbb{C}},h,\eta,0,1,-1)/\sim_{\gamma})$ from $\mathbf{m}_{\widehat{\mathbb{C}}} \times M_3$. The uniform embedding of M_3 via $\mathbf{m}_{\widehat{\mathbb{C}}}$, which we denote by $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M_3$, is the law of $(\mathfrak{f} \bullet_{\gamma} h, \mathfrak{f} \circ \eta, \mathfrak{f}(0), \mathfrak{f}(1), \mathfrak{f}(-1))$. We can similarly define $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M$ and $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes \widehat{M}_3$.

Proposition 3.3. There exists a constant $C \in (0, \infty)$ such that $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M_3 = C\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes \widetilde{M}_3$.

We first give the uniform embedding of M.

Lemma 3.4. There exists a constant $C \in (0, \infty)$ such that $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M = C \cdot \mathrm{LF}_{\widehat{\mathbb{C}}} \times \mathrm{SLE}_{\kappa}^{\mathrm{loop}}$.

Proof. The measure $\mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ is conformally invariant, namely, for each $f \in \mathrm{conf}(\widehat{\mathbb{C}})$, the law of $f \circ \eta$ is $\mathrm{SLE}_{\kappa}^{\mathrm{loop}}$ if η is sampled from $\mathrm{SLE}_{\kappa}^{\mathrm{loop}}$. Now Lemma 3.4 follows from Proposition 2.15.




FIGURE 1. Illustration for proof of Theorem 1.1. Measures are displayed without multiplicative constants. We denote the quantum area by A and the quantum length of the loop by L.

We now describe the uniform embedding of M_3 .

Lemma 3.5. There exists a constant $C \in (0, \infty)$ such that

$$\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M_3 = C|p-q|^{\frac{\gamma^2}{4}-2} \operatorname{LF}_{\widehat{\mathbb{C}}}^{(\frac{\gamma}{2},p),(\frac{\gamma}{2},q),(\gamma,r)}(d\phi) \operatorname{SLE}_{\kappa}^{p \rightleftharpoons q}(d\eta) d^2p d^2q d^2r.$$

To prove Lemma 3.5 we use an analog of Lemma 2.12 based on the Girsanov theorem. We first review some background on the Minkowski content of SLE and its relation to quantum length. As before we denote the $(1 + \frac{\kappa}{8})$ -dimensional Minkowski content of an SLE_{κ} -type curve η by $Cont_{\eta}$.

Lemma 3.6. Let $d = 1 + \frac{\kappa}{8}$. Let η be sampled from SLE_{κ}^{loop} . Then almost surely

(3.1)
$$\int_{\mathbb{C}^2} \frac{\operatorname{Cont}_{\eta}(dx) \operatorname{Cont}_{\eta}(dy)}{|x-y|^{d-\varepsilon}} < \infty \quad \text{for each } \varepsilon \in (0,d).$$

Proof. By Green's function estimates for chordal SLE (see e.g. [LR15]) (3.1) holds if η is sampled from a chordal SLE_{κ} even after we take expectation over the integral. By local absolutely continuity, (3.1) holds for SLE^{loop}_{κ}.

For each η such that (3.1) holds, the Gaussian multiplicative chaos (GMC) measure (see e.g. [Ber17])

$$\nu_h^{\eta} := \lim_{\varepsilon \to 0} \varepsilon^{\frac{\gamma^2}{8}} e^{\frac{1}{2}\gamma h_{\varepsilon}} \operatorname{Cont}_{\eta}$$

exists, where h is sampled from the Gassian free field measure $P_{\mathbb{C}}$. By [Ben18, Section 3.2], modulo a multiplicative constant, ν_h^{η} is the quantum length of η with respect to h. We now give an analog of Lemma 2.12.

Lemma 3.7. Suppose η is a loop satisfying (3.1). For $\alpha \in \mathbb{R}$ and $z \in \mathbb{C}$, we have

$$\nu_{\phi}^{\eta}(du) \operatorname{LF}_{\widehat{\mathbb{C}}}^{(\alpha,z)}(d\phi) = \operatorname{LF}_{\widehat{\mathbb{C}}}^{(\alpha,z),(\frac{\gamma}{2},u)}(d\phi) \operatorname{Cont}_{\eta}(du).$$

Proof. The proof is identical to that of Lemma 2.12, except we replace the quantum area measure $\mu_{\phi}(d^2u) = \lim_{\varepsilon \to 0} \varepsilon^{\gamma^2/2} e^{\gamma\phi_{\varepsilon}(u)} d^2u$ with the GMC measure $\nu_{\phi}^{\eta}(du) = \lim_{\varepsilon \to 0} \varepsilon^{\frac{\gamma^2}{8}} e^{\frac{1}{2}\gamma\phi_{\varepsilon}} \operatorname{Cont}_{\eta}(du)$. \square

Proof of Lemma 3.5. Since ν_{ϕ}^{η} is the quantum length measure on η modulo a multiplicative constant, by Lemma 3.4, there exists $C \in (0, \infty)$ such that

$$\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M_3 = C\mu_{\phi}(dr) \,\nu_{\phi}^{\eta}(dp) \,\nu_{\phi}^{\eta}(dq) \,\mathrm{LF}_{\widehat{\mathbb{C}}}(d\phi) \,\mathrm{SLE}_{\kappa}^{\mathrm{loop}}(d\eta).$$

By Lemma 3.6 η almost surely satisfies (3.1). Thus, applying Lemma 3.7 twice, we get

$$\nu_{\phi}^{\eta}(dp) \, \nu_{\phi}^{\eta}(dq) \, \mathrm{LF}_{\widehat{\mathbb{C}}}(d\phi) \, \, \mathrm{SLE}_{\kappa}^{\mathrm{loop}}(d\eta)$$

$$= \mathrm{LF}_{\widehat{\mathbb{C}}}^{(\frac{\gamma}{2},p),(\frac{\gamma}{2},q)}(d\phi) \, \mathrm{Cont}_{\eta}(dp) \, \mathrm{Cont}_{\eta}(dq) \, \, \mathrm{SLE}_{\kappa}^{\mathrm{loop}}(d\eta).$$

By Lemma 2.12, we get further that $\mu_{\phi}(dr) \nu_{\phi}^{\eta}(dp) \nu_{\phi}^{\eta}(dq) \operatorname{LF}_{\widehat{\mathbb{C}}}(d\phi) \operatorname{SLE}_{\kappa}^{\operatorname{loop}}(d\eta)$ equals

$$LF_{\widehat{\mathbb{C}}}^{(\frac{\gamma}{2},p),(\frac{\gamma}{2},q),(\gamma,r)}(d\phi)\operatorname{Cont}_{\eta}(dp)\operatorname{Cont}_{\eta}(dq)\operatorname{SLE}_{\kappa}^{\operatorname{loop}}(d\eta)d^{2}r.$$

Comparing against (2.3) completes the proof.

We now switch our attention to $\mathbf{m} \ltimes \widetilde{M}_3$. The following lemma describes the embedding of \widetilde{M}_3 .

Lemma 3.8. Given distinct p,q,r on $\widehat{\mathbb{C}}$, let $\widetilde{M}_3^{p,q,r}$ be the law of (ϕ,η) where $(\widehat{\mathbb{C}},\phi,\eta,p,q,r)$ is an embedding of a sample from \widetilde{M}_3 . Then there exists a constant $C \in (0,\infty)$ such that

$$\widetilde{M}_3^{p,q,r} = C|p-q|^{\frac{\gamma^2}{4}-2}|(p-q)(q-r)(r-p)|^2 \mathrm{LF}_{\widehat{\mathbb{C}}}^{(\frac{\gamma}{2},p),(\frac{\gamma}{2},q),(\gamma,r)} \times \mathrm{SLE}_{\kappa}^{p \rightleftharpoons q}.$$

Proof. By Proposition 2.13, there exists a constant $C \in (0, \infty)$ such that

$$\widetilde{M}_3^{0,1,-1} = C\mathrm{LF}_{\widehat{\mathbb{C}}}^{(\frac{\gamma}{2},0),(\frac{\gamma}{2},1),(\gamma,-1)} \times \mathrm{SLE}_{\kappa}^{0\rightleftharpoons 1} \,.$$

Suppose $f \in \text{conf}(\widehat{\mathbb{C}})$ maps (0,1,-1) to (p,q,r); explicitly, we have $f(z) = \frac{(pq-2qr+rp)z+p(q-r)}{(2p-q-r)z+q-r}$, and

$$f'(0) = \frac{2(p-q)(q-r)(r-p)}{(q-r)^2}, \quad f'(1) = \frac{2(p-q)(q-r)(r-p)}{4(r-p)^2}, \quad f'(-1) = \frac{2(p-q)(q-r)(r-p)}{4(p-q)^2}.$$

We have

$$f'(0)f'(1) = 4(p-q)^2$$
 and $f'(0)f'(1)f'(-1) = 2(p-q)(q-r)(r-p)$.

By Proposition 2.11 the field of $\widetilde{M}_3^{p,q,r}$ is given by

$$f_* LF_{\widehat{\mathbb{C}}}^{(\frac{\gamma}{2},0),(\frac{\gamma}{2},1),(\gamma,-1)} = |f'(0)|^{2\Delta_{\frac{\gamma}{2}}} |f'(1)|^{2\Delta_{\frac{\gamma}{2}}} |f'(-1)|^{2\Delta_{\gamma}} LF_{\widehat{\mathbb{C}}}^{(\frac{\gamma}{2},p),(\frac{\gamma}{2},q),(\gamma,r)},$$

where $\Delta_{\alpha} = \frac{\alpha}{2}(Q - \frac{\alpha}{2})$. Since $\Delta_{\frac{\gamma}{2}} = \frac{1}{2} + \frac{\gamma^2}{16}$ and $\Delta_{\gamma} = 1$, we get the desired result.

Proof of Proposition 3.3. By Lemma 2.14 and the definition of $\widetilde{M}_3^{p,q,r}$ in Lemma 3.8, we see that

(3.2)
$$\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes \widetilde{M}_3 = C \widetilde{M}_3^{p,q,r} |(p-q)(q-r)(r-p)|^{-2} d^2 p \, d^2 q \, d^2 r \text{ for some } C \in (0,\infty).$$

Now Lemmas 3.5 and 3.8 together give $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M_3 = C\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes \widetilde{M}_3$ for a possibly different constant C.

Proof of Proposition 3.2. Given distinct p,q,r on $\widehat{\mathbb{C}}$, let $M_3^{p,q,r}$ be the law of (ϕ,η) where $(\widehat{\mathbb{C}},\phi,\eta,p,q,r)$ is a sample from M_3 . By the definition of uniform embedding, the law of (ϕ,η) sampled from $\mathbf{m}_{\widehat{\mathbb{C}}} \ltimes M_3$ agrees with that of $(\mathfrak{f} \bullet_{\gamma} \phi_0, \mathfrak{f} \circ \eta_0)$ where $(\mathfrak{f},\phi_0,\eta_0) \sim \mathbf{m}_{\widehat{\mathbb{C}}} \times M_3^{0,1,-1}$. The $\mathbf{m}_{\widehat{\mathbb{C}}}$ -law of \mathfrak{f} is described by Lemma 2.14, and by definition, if f is the conformal automorphism of $\widehat{\mathbb{C}}$ sending (0,1,-1) to (p,q,r) and $(\phi_0,\eta_0) \sim M_3^{0,1,-1}$, the law of $(f \bullet_{\gamma} \phi_0, f \circ \eta_0)$ is $M_3^{p,q,r}$. Thus (3.2) holds with M_3 and $M_3^{p,q,r}$ in place of \widehat{M}_3 and $\widehat{M}_3^{p,q,r}$. Consequently

$$M_3^{p,q,r} d^2 p \, d^2 q \, d^2 r = C \widetilde{M}_3^{p,q,r} d^2 p \, d^2 q \, d^2 r \quad \text{for some } C \in (0,\infty).$$

This gives $M_3^{p,q,r} = C\widetilde{M}_3^{p,q,r}$ for almost every p,q,r. Using any such p,q,r, we conclude $M_3 = C\widetilde{M}_3$ as desired.

Remark 3.9 (KPZ relation). As seen in the proof of Lemma 3.5, a crucial fact to our proof is that the exponent $\frac{\gamma^2}{4} - 2$ is equal to $-2(2-d) = \frac{\kappa}{4} - 2$ from (2.2) where $d = 1 + \frac{\kappa}{8}$ is the dimension of SLE_{κ} . As seen in the proof of Lemma 3.7, this comes from $4(\Delta_{\frac{\gamma}{2}} - 1) = -2(2-d)$ where $\Delta_{\alpha} = \frac{\alpha}{2}(Q - \frac{\alpha}{2})$. This is equivalent to $d = 2\Delta_{\frac{\gamma}{2}}$, which is an instance of the Knizhnik-Polyakov-Zamolodchikov (KPZ) relation.

4. The scaling limit on random quandragulation decorated by self-avoiding loop

In this section we prove Theorem 1.2. We start by introducing more precisely the objects appearing in the theorem. Recall that a planar map is a connected graph drawn on the sphere \$\mathbb{S}^2\$ such that no two edges cross, viewed modulo an orientation-preserving homeomorphisms from the sphere to itself. A quadrangulation is a planar map such that all faces have four edges. Le Gall and Miermont [Mie13, Le 13] proved that uniformly sampled quadrangulations converge in the scaling limit to the metric measure space known as the Brownian map for the so-called Gromov-Hausdorff-Prokhorov topology [ADH13].

Define the following constants:

(4.1)
$$\lambda = 12, \quad \theta = 54, \quad a = 5/2 \quad b = 1/2.$$

The constants are chosen such that the number of quadrangulations of a 2p-gon with m faces is of order $\theta^p p^{-b} \lambda^m m^{-a}$ for $m \ge cp^2$ for arbitrary fixed c > 0 [Bro65].³ Let MSⁿ denote the measure on quadrangulations such that a quadrangulation M with m faces has weight $n^a \lambda^{-m}$. For M sampled from MSⁿ, we view M as a metric measure space by considering the graph metric rescaled by $2^{-1/2}n^{-1/4}$ and by giving each vertex mass $2(9n)^{-1}$. With this choice of rescaling, the measure of the set of quadrangulations with mass of order 1 will be of order 1 since the number of quadrangulations with m faces is of order $\lambda^m m^{-a-1}$ [Tut63].

If M is a quadrangulation we say that η is a *self-avoiding loop* on M if η is an ordered set of edges $e_1, \ldots, e_{2k} \in \mathcal{E}(M)$ such e_j and e_i share an end-point if and only if $|i-j| \leq 1$ or $(i,j) \in \{(1,2k),(2k,1)\}$. Let $\#\eta = 2k$ denote the number of edges on η . Let $MS^n \otimes SAW^n$ denote the measure on pairs (M,η) where η is a self-avoiding loop on M and a pair (M,η) has weight

$$n^{2a+b-3}\lambda^{-\#\mathcal{F}(M)}\theta^{-\#\eta}$$
.

For (M, η) sampled from $MS^n \otimes SAW^n$, we view M as a metric measure space as above and view η as a loop on this metric measure space such that the time it takes to trace each edge on the loop is $2^{-1}n^{-1/2}$. Here we include the edges in the metric-measure structure of M so that η can be defined as a continuous curve on M; see e.g. [GM16, Remark 2.4].

It was proved by Miller and Sheffield that quantum surfaces with $\gamma = \sqrt{8/3}$ can be identified with Brownian surfaces [MS15, MS21a, MS21b]. More precisely, a quantum surface sampled from QS with $\gamma = \sqrt{8/3}$ defines a random metric measure space which is equal in law to the Brownian map. In particular, a sample from QS \otimes SLE_{8/3}^{loop} with $\gamma = \sqrt{8/3}$ can be viewed as a loop-decorated metric measure space. We will use this interpretation in this subsection and in the statement of Theorem 1.2; this is a slight abuse of notation since we view QS \otimes SLE_{8/3}^{loop} as a measure on the space of loop-decorated LQG surface in other sections. The loop is parametrized by its quantum length.

The paragraphs above allow for a precise statement of Theorem 1.2. We will now turn to the proof of this theorem, which builds on Theorem 1.1 along with three ingredients given below: Theorem 4.1, Observation 4.2, and (4.2). In order to state these results we first introduce some further notation.

³Our quadrangulated 2p-gons are unrooted. If we consider maps with a root edge on its boundary then the number of maps is of order $\theta^p p^{-b+1} \lambda^m m^{-a}$ instead.

A planar map M is a quadrangulated disk if it is a planar map where all faces have four edges expect for a distinguished face (called the exterior face) which has arbitrary degree and simple boundary. We let ∂M denote the edges on the boundary of the exterior face, and we call $\#\partial M$ the boundary length of M. Let MD^n be the measure on quadrangulated disks such that each quadrangulated disk M has mass $n^{a+b/2-3/2}\lambda^{-\#\mathcal{F}(M)}\theta^{-(\#\partial M)/2}$. We need to choose this mass in order for Observation 4.2 below to be correct; note in particular that the exponents of n and θ have been divided by two as compared to MS^n above since we glue together two disks to form a sphere. If $M \sim \mathrm{MD}^n$ then we view M as a metric measure space by applying the same rescaling as for MS^n above. For $k \in \mathbb{N}$ let $\mathrm{MD}^n(k)$ denote MD^n restricted to quadrangulations with boundary length 2k, and let $\mathrm{MD}^n(k)^\#$ denote $\mathrm{MD}^n(k)$ renormalized to be a probability measure.

If M_1, M_2 are quadrangulated disks with boundary length 2k then we can form a quadrangulation with a self-avoiding loop by choosing uniform boundary edges $e_1 \in \partial M_1, e_2 \in \partial M_2$ and then identifying the boundaries of M_1, M_2 such that e_1 and e_2 are identified. The self-avoiding loop on the sphere represents the boundaries of M_1, M_2 , and we parametrize the loop so that each edge on the loop has length $2^{-1}n^{-1/2}$. Note that the scaling we use of distances along the loop $(2^{-1}n^{-1/2})$ is different from the scaling we use of graph distances in the map $(2^{-1/2}n^{-1/4})$; this choice of exponents (-1/2 and -1/4) cause both distances to be asymptotically non-trivial. If $M_1, M_2 \sim \text{MD}^n(k)^\#$ then we denote the measure on spheres decorated with a self-avoiding loop sampled in this way by $\text{Weld}(\text{MD}^n(k)^\#, \text{MD}^n(k)^\#)$.

Theorem 4.1 ([GM19a, GM19b]). For any $\ell > 0$ the following convergence in law holds for the Gromov-Hausdorff-Prokhorov-uniform topology

$$\operatorname{Weld}(\operatorname{MD}^n(\lceil \ell n^{1/2} \rceil)^{\#}, \operatorname{MD}^n(\lceil \ell n^{1/2} \rceil)^{\#}) \Rightarrow \operatorname{Weld}(\operatorname{QD}(\ell)^{\#}, \operatorname{QD}(\ell)^{\#}).$$

Proof. [GM19a, Theorem 1.5] proves this convergence result when the right side is given by a metric space quotient. By [GM19b] and local absolute continuity we get that this metric space quotient gives the same metric space as the conformal welding of the two disks. \Box

Let $Z_n(k)$ denote the total mass of $MD^n(k)$. It follows from [Bro65] (see his enumeration result cited right below (4.1)) that there is a constant C > 0 such that

(4.2)
$$\frac{Z_n(k)}{n^{a+b/2-3/2}k^{-b-2a+2}} = C(1+o_k(1)),$$

where the $o_k(1)$ is uniform in n. We now define Weld(MDⁿ, MDⁿ) in the same spirit as Weld(QD, QD)

(4.3)
$$\operatorname{Weld}(\operatorname{MD}^n, \operatorname{MD}^n) := \sum_{k=1}^{\infty} 2k Z_n(k)^2 \operatorname{Weld}(\operatorname{MD}^n(k)^{\#}, \operatorname{MD}^n(k)^{\#}),$$

where we recall that samples from $MD^n(k)$ have boundary length 2k and $MD^n(k) = Z_n(k)MD^n(k)^{\#}$. The observation we state next is immediate by combinatorial considerations and was also observed in slightly different forms in e.g. [GM19a, Section 1.3.3] and [CC19]. The key point is that there are 2k ways of welding together two samples from $MD^n(k)$.

Observation 4.2. Weld(MD^n, MD^n) = $MS^n \otimes SAW^n$.

Combining the three ingredients above, we can now conclude the proof of Theorem 1.2.

Proof of Theorem 1.2. For constants $c_1, c_2 > 0$,

$$(4.4) \qquad \text{Weld}(\text{MD}^n, \text{MD}^n)|_{A(c)} = \sum_{k=\lceil c\sqrt{n}\rceil}^{\lceil c^{-1}\sqrt{n}\rceil} 2kZ_n(k)^2 \operatorname{Weld}(\text{MD}^n(k)^\#, \text{MD}^n(k)^\#)$$

$$\Rightarrow c_1 \int_c^{c^{-1}} \operatorname{Weld}(\text{QD}(\ell)^\#, \text{QD}(\ell)^\#) \ell^{-2(b+2a-2)+1} d\ell$$

$$= c_2 \int_c^{c^{-1}} \operatorname{Weld}(\text{QD}(\ell), \text{QD}(\ell)) \ell d\ell,$$

where we use in the last step that the total mass of $QD(\ell)$ is a power law with exponent -7/2 = -(b + 2a - 2), which follows e.g. from Lemma 2.4. The right side of (4.4) is equal to $c_0 \cdot QS \otimes SLE_{8/3}^{loop}|_{A(c)}$ for some $c_0 > 0$ by Theorem 1.1, while it follows from Observation 4.2 that the left side of (4.4) is equal to $MS^n \otimes SAW^n|_{A(c)}$. This concludes the proof.

References

- [ADH13] R. Abraham, J.-F. Delmas, and P. Hoscheit. A note on the Gromov-Hausdorff-Prokhorov distance between (locally) compact metric measure spaces. *Electron. J. Probab.*, 18:no. 14, 21, 2013, arXiv:1202.5464. MR3035742
- [AHS20a] M. Albenque, N. Holden, and X. Sun. Scaling limit of triangulations of polygons. *Electronic Journal of Probability*, 25(none):1 43, 2020.
- [AHS20b] M. Ang, N. Holden, and X. Sun. Conformal welding of quantum disks. arXiv e-prints, September 2020, 2009.08389.
- [AHS22] M. Ang, N. Holden, and X. Sun. Integrability of SLE via conformal welding of random surfaces. *Communications on Pure and Applied Mathematics*, to appear, 2022.
- [ARS22a] M. Ang, G. Remy, and X. Sun. FZZ formula of boundary Liouville CFT via conformal welding. *Journal of the European Mathematical Society*, to appear, 2022.
- [ARS22b] M. Ang, G. Remy, and X. Sun. The moduli of annuli in random conformal geometry. arXiv e-prints, page arXiv:2203.12398, March 2022, 2203.12398.
- [AS21] M. Ang and X. Sun. Integrability of the conformal loop ensemble. arXiv e-print, 2021.
- [BBI01] D. Burago, Y. Burago, and S. Ivanov. A course in metric geometry, volume 33 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2001. MR1835418
- [BD16] S. Benoist and J. Dubédat. An SLE₂ loop measure. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 52(3):1406–1436, 2016.
- [Bef08] V. Beffara. The dimension of the SLE curves. Ann. Probab., 36(4):1421-1452, 2008, math/0211322. MR2435854 (2009e:60026)
- [Ben18] S. Benoist. Natural parametrization of SLE: the Gaussian free field point of view. *Electron. J. Probab.*, 23:Paper No. 103, 16, 2018. MR3870446
- [Ber17] N. Berestycki. An elementary approach to Gaussian multiplicative chaos. Electron. Commun. Probab., 22:Paper No. 27, 12, 2017, 1506.09113. MR3652040
- [BP21] N. Berestycki and E. Powell. Introduction to the Gaussian Free Field and Liouville Quantum Gravity. Available at https://homepage.univie.ac.at/nathanael.berestycki/articles.html, 2021.
- [Bro65] W. G. Brown. Enumeration of quadrangular dissections of the disk. Canadian Journal of Mathematics, 17:302–317, 1965.
- [CC19] A. Caraceni and N. Curien. Self-Avoiding Walks on the UIPQ. In V. Sidoravicius, editor, Sojourns in Probability Theory and Statistical Physics - III, pages 138–165, Singapore, 2019. Springer Singapore.
- [DKRV16] F. David, A. Kupiainen, R. Rhodes, and V. Vargas. Liouville quantum gravity on the Riemann sphere. Comm. Math. Phys., 342(3):869–907, 2016, 1410.7318. MR3465434
- [DMS21] B. Duplantier, J. R. Miller, and S. Sheffield. Liouville quantum gravity as a mating of trees. *Astérisque*, 427, 2021.
- [DS11] B. Duplantier and S. Sheffield. Liouville quantum gravity and KPZ. *Invent. Math.*, 185(2):333–393, 2011, 1206.0212. MR2819163 (2012f:81251)
- [GHS19] E. Gwynne, N. Holden, and X. Sun. Mating of trees for random planar maps and Liouville quantum gravity: a survey. *ArXiv e-prints*, Oct 2019, 1910.04713.
- [GM16] E. Gwynne and J. Miller. Convergence of the self-avoiding walk on random quadrangulations to $SLE_{8/3}$ on $\sqrt{8/3}$ -Liouville quantum gravity. Annales de l'ENS, to appear, 2016, 1608.00956.

- [GM17] E. Gwynne and J. Miller. Scaling limit of the uniform infinite half-plane quadrangulation in the Gromov-Hausdorff-Prokhorov-uniform topology. Electron. J. Probab., 22:1–47, 2017, 1608.00954.
- [GM19a] E. Gwynne and J. Miller. Convergence of the free Boltzmann quadrangulation with simple boundary to the Brownian disk. Ann. Inst. Henri Poincaré Probab. Stat., 55(1):551–589, 2019, 1701.05173. MR3901655
- [GM19b] E. Gwynne and J. Miller. Metric gluing of Brownian and $\sqrt{8/3}$ -Liouville quantum gravity surfaces. Ann. Probab., 47(4):2303–2358, 2019, 1608.00955. MR3980922
- [GPW09] A. Greven, P. Pfaffelhuber, and A. Winter. Convergence in distribution of random metric measure spaces (Λ-coalescent measure trees). Probab. Theory Related Fields, 145(1-2):285–322, 2009, math/0609801. MR2520129
- [Gro99] M. Gromov. Metric structures for Riemannian and non-Riemannian spaces, volume 152 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1999. Based on the 1981 French original [MR0682063 (85e:53051)], With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates. MR1699320
- [Gwy20] E. Gwynne. Random surfaces and Liouville quantum gravity. Notices Amer. Math. Soc., 67(4):484–491, 2020. MR4186266
- [HP21] N. Holden and E. Powell. Conformal welding for critical Liouville quantum gravity. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, 57(3):1229–1254, 2021.
- [HRV18] Y. Huang, R. Rhodes, and V. Vargas. Liouville quantum gravity on the unit disk. Ann. Inst. Henri Poincaré Probab. Stat., 54(3):1694–1730, 2018, 1502.04343. MR3825895
- [KK17] A. Kassel and R. Kenyon. Random curves on surfaces induced from the Laplacian determinant. Ann. Probab., 45(2):932–964, 2017. MR3630290
- [KRV20] A. Kupiainen, R. Rhodes, and V. Vargas. Integrability of Liouville theory: proof of the DOZZ formula. Ann. of Math. (2), 191(1):81–166, 2020. MR4060417
- [KS07] M. Kontsevich and Y. Suhov. On Malliavin measures, SLE, and CFT. Tr. Mat. Inst. Steklova, 258(Anal. i Osob. Ch. 1):107–153, 2007. MR2400527
- [KW16] A. Kemppainen and W. Werner. The nested simple conformal loop ensembles in the Riemann sphere. Probab. Theory Related Fields, 165(3-4):835–866, 2016, 1402.2433. MR3520020
- [Law05] G. F. Lawler. Conformally invariant processes in the plane, volume 114 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2005. MR2129588 (2006i:60003)
- [Le 13] J.-F. Le Gall. Uniqueness and universality of the Brownian map. Ann. Probab., 41(4):2880–2960, 2013, 1105.4842. MR3112934
- [LR15] G. F. Lawler and M. A. Rezaei. Minkowski content and natural parameterization for the Schramm-Loewner evolution. Ann. Probab., 43(3):1082-1120, 2015, 1211.4146. MR3342659
- [LRV19] H. Lacoin, R. Rhodes, and V. Vargas. The semiclassical limit of Liouville conformal field theory. arXiv e-prints, page arXiv:1903.08883, March 2019, 1903.08883.
- [Mal99] P. Malliavin. The canonic diffusion above the diffeomorphism group of the circle. C. R. Acad. Sci. Paris Sér. I Math., 329(4):325–329, 1999. MR1713340
- [Mie09] G. Miermont. Random maps and their scaling limits. In Fractal geometry and stochastics IV, volume 61 of Progr. Probab., pages 197–224. Birkhäuser Verlag, Basel, 2009. MR2762678 (2012a:60017)
- [Mie13] G. Miermont. The Brownian map is the scaling limit of uniform random plane quadrangulations. Acta Math., 210(2):319–401, 2013, 1104.1606. MR3070569
- [MMQ19] O. McEnteggart, J. Miller, and W. Qian. Uniqueness of the welding problem for SLE and Liouville quantum gravity. *Journal of the Institute of Mathematics of Jussieu*, page 1–27, 2019.
- [MS15] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map I: The QLE(8/3,0) metric. Inventiones Mathematicae, to appear, 2015, 1507.00719.
- [MS17] J. Miller and S. Sheffield. Imaginary geometry IV: interior rays, whole-plane reversibility, and space-filling trees. Probab. Theory Related Fields, 169(3-4):729–869, 2017, 1302.4738. MR3719057
- [MS21a] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map II: geodesics and continuity of the embedding. *The Annals of Probability*, 49(6):2732–2829, 2021.
- [MS21b] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map III: the conformal structure is determined. *Probability Theory and Related Fields*, 179:1183–1211, 2021.
- [RS05] S. Rohde and O. Schramm. Basic properties of SLE. Ann. of Math. (2), 161(2):883–924, 2005, math/0106036. MR2153402 (2006f:60093)
- [RZ22] G. Remy and T. Zhu. Integrability of boundary Liouville conformal field theory. Communications in Mathematical Physics, 395(1):179–268, 2022.
- [She09] S. Sheffield. Exploration trees and conformal loop ensembles. Duke Math. J., 147(1):79-129, 2009, math/0609167. MR2494457 (2010g:60184)
- [She16] S. Sheffield. Conformal weldings of random surfaces: SLE and the quantum gravity zipper. Ann. Probab., 44(5):3474–3545, 2016, 1012.4797. MR3551203

- [She22] S. Sheffield. What is a random surface? arXiv e-prints, page arXiv:2203.02470, March 2022, 2203.02470.
- [SW05] O. Schramm and D. B. Wilson. SLE coordinate changes. New York J. Math., 11:659–669 (electronic), 2005, math/0505368. MR2188260 (2007e:82019)
- [SW12] S. Sheffield and W. Werner. Conformal loop ensembles: the Markovian characterization and the loop-soup construction. *Ann. of Math. (2)*, 176(3):1827–1917, 2012, 1006.2374. MR2979861
- [Tut63] W. T. Tutte. A census of planar maps. Canadian Journal of Mathematics, 15:249–271, 1963.
- [VW20] F. Viklund and Y. Wang. Interplay between Loewner and Dirichlet energies via conformal welding and flow-lines. *Geom. Funct. Anal.*, 30(1):289–321, 2020. MR4080509
- [Wan22] Y. Wang. Large deviations of Schramm-Loewner evolutions: A survey. Probability Surveys, 19:351–403, 2022.
- [Wer08] W. Werner. The conformally invariant measure on self-avoiding loops. J. Amer. Math. Soc., 21(1):137–169, 2008. MR2350053
- [Zha21] D. Zhan. SLE loop measures. Probab. Theory Related Fields, 179(1-2):345-406, 2021. MR4221661