arXiv:2205.05766v1 [nlin.SI] 11 May 2022

Weierstrass type projective Riccati equation expansion method and solutions of KdV equation

Sirendaoreji*
Mathematical Science College, Inner Mongolia Normal University,
Huhhot 010022, Inner Mongolia, P. R. China

Abstract

This paper presents two new Weierstrass elliptic function solutions of the projective Riccati equations and four conversion formulas for converting the Weierstrass elliptic functions to the hyperbolic and trigonometric functions. The Weierstrass elliptic function solutions to the projective Riccati equations and the conversion formulas are used to propose the called Weierstrass type projective Riccati equation expansion method. The Weierstrass elliptic function solutions, the solitary wave and the periodic wave solutions of the KdV equation are constructed by using the proposed method. The solitary wave like and the periodic wave solutions of the KdV equation are shown through some figures.

1 Introduction

Exact solutions to nonlinear evolution equations (NLEEs) play an important role in applied mathematics, mathematical physics and nonlinear science. Therefore, the problem of how to find new methods for solving NLEEs has become a hot topic in soliton theory. In the past decades, various direct methods such as the tanh–function method [1], the auxiliary equation method [2], the Riccati equation expansion method [3], the unified Riccati equation expansion method [4], the Jacobi elliptic function expansion method [5], the Weierstrass elliptic function method [6, 7, 8, 9, 10] and the projective Riccati equation expansion method [11, 12, 13, 14, 15], and so on, have been proposed to find exact traveling wave solutions of NLEEs. Among them, the projective Riccati equation expansion method can usually give some new special solutions of NLEEs. Therefore, the researchers give more attention to find new solutions of the projective Riccati equation. So far, two sets of Weierstrass elliptic function solutions of the projective Riccati equation have been given and applied to construct the Weierstrass elliptic function solutions of NLEEs [16, 17, 18]. But most of these Weierstrass elliptic function solutions cannot be transformed into the hyperbolic and trigonometric function solutions because the previously used conversion formulas were determined by the roots of the third order polynomial equation $w^3 - g_2 w - g_3 = 0$. Therefore, this paper aims to give more Weierstrass elliptic function solutions of the projective Riccati equations and establish the so–called conversion formulas which can transform the Weierstrass elliptic function solutions of NLEEs into the hyperbolic and trigonometric function solutions. Further, we shall use the Weierstrass elliptic function solutions of the projective Riccati equations and the

*E-mail: siren@imnu.edu.cn

conversion formulas to propose a direct method which we named as the Weierstrass type projective Riccati equation expansion method. Finally, we shall take the KdV equation as an illustrative example to show the effectiveness of our method.

This paper is organized as follows. In the next section, four sets of Weierstrass elliptic function solutions of the projective Riccati equations and four conversion formulas to transform the Weierstrass elliptic function into the hyperbolic and trigonometric function are constructed. In addition, the Weierstrass type projective Riccati equation expansion method is proposed to find exact traveling wave solutions to NLEEs. In Sec.3, the proposed method is applied to construct the Weierstrass elliptic function solutions, the solitary and periodic wave solutions of the KdV equation. The conclusions are given in Sec.4.

2 Weierstrass type projective Riccati equation expansion method

The Weierstrass elliptic function $w = \wp(\xi, g_2, g_3)$ is defined as the inverse function of the Weierstrass elliptic integral

$$\xi = \int_{\infty}^{w} \frac{dt}{\sqrt{4t^3 - g_2 t - g_3}}, \tag{1}$$

or the solution of the following nonlinear ordinary differential equation

$$\frac{dw}{d\xi} = 4w^3 - g_2 w - g_3, \tag{2}$$

where g_2, g_3 are real parameters and called invariants.

In the following we shall consider the projective Riccati equations of the form

$$\begin{cases} F'(\xi) = pF(\xi)G(\xi), \\ G'(\xi) = q + pG^2(\xi) - rF(\xi), \end{cases} \tag{3}$$

where F,G are unknown functions of the variable ξ and p, q, r are constants. By using the direct assumption approach we can construct the Weierstrass elliptic function solutions of Eqs.(3) as following

$$\begin{cases} F(\xi) = \dfrac{q}{6r} + \dfrac{2}{pr}\wp(\xi, g_2, g_3), \\ G(\xi) = \dfrac{12\wp'(\xi, g_2, g_3)}{p\left[pq + 12\wp(\xi, g_2, g_3)\right]}, \end{cases} \tag{4}$$

$$G^2(\xi) = -\frac{q}{p} + \frac{2r}{p}F(\xi), \tag{5}$$

$$\begin{cases} F(\xi) = \dfrac{5q}{6r} + \dfrac{5q^2 p}{72r\wp(\xi, g_2, g_3)}, \\ G(\xi) = \dfrac{-q\wp'(\xi, g_2, g_3)}{\left[qp + 12\wp(\xi, g_2, g_3)\right]\wp(\xi, g_2, g_3)}, \end{cases} \tag{6}$$

$$G^2(\xi) = -\frac{q}{p} + \frac{2r}{p}F(\xi) - \frac{24r^2}{25pq}F^2(\xi), \tag{7}$$

$$\begin{cases} F(\xi) = \dfrac{(2+q)\left[pq + 12\wp(\xi, g_2, g_3)\right]}{r\left[12p + pq + 12\wp(\xi, g_2, g_3)\right]}, \\ G(\xi) = \dfrac{\wp'(\xi, g_2, g_3)}{\left(\wp(\xi, g_2, g_3) + \frac{p}{2} + \frac{pq}{12}\right)^2 - \frac{p^2}{4}}, \end{cases} \tag{8}$$

$$G^2(\xi) = -\frac{q}{p} + \frac{2r}{p}F(\xi) - \frac{r^2(q+4)}{p(q+2)^2}F^2(\xi), \tag{9}$$

$$\begin{cases} F(\xi) = \dfrac{q(p^2+2)\left[pq + 12\wp(\xi, g_2, g_3)\right]}{pr\left[12q + p^2q + 12p\wp(\xi, g_2, g_3)\right]}, \\ G(\xi) = \dfrac{q\wp'(\xi, g_2, g_3)}{\left(p\wp(\xi, g_2, g_3) + \frac{q}{2} + \frac{p^2q}{12}\right)^2 - \frac{q^2}{4}}, \end{cases} \tag{10}$$

$$G^2(\xi) = -\frac{q}{p} + \frac{2r}{p}F(\xi) - \frac{pr^2(p^2+4)}{q(p+2)^2}F^2(\xi), \tag{11}$$

where the invariants g_2 and g_3 are given by

$$g_2 = \frac{p^2q^2}{12}, g_3 = \frac{p^3q^3}{216}. \tag{12}$$

Here we point out that the solutions (4) and (6) are previously known [16, 17, 18], but the solutions (8) and (10) are new.

We find that the Weierstrass elliptic function can degenerate to the hyperbolic and trigonometric function by using the following conversion formulas

$$\wp(\xi, \frac{\theta^2}{12}, -\frac{\theta^3}{216}) = \frac{\theta}{12} - \frac{\theta}{4}\mathrm{sech}^2(\frac{\sqrt{\theta}}{2}\,\xi), \theta > 0, \tag{13}$$

$$\wp(\xi, \frac{\theta^2}{12}, -\frac{\theta^3}{216}) = \frac{\theta}{12} + \frac{\theta}{4}\mathrm{csch}^2(\frac{\sqrt{\theta}}{2}\,\xi), \theta > 0, \tag{14}$$

$$\wp(\xi, \frac{\theta^2}{12}, -\frac{\theta^3}{216}) = \frac{\theta}{12} - \frac{\theta}{4}\sec^2(\frac{\sqrt{-\theta}}{2}\,\xi), \theta < 0, \tag{15}$$

$$\wp(\xi, \frac{\theta^2}{12}, -\frac{\theta^3}{216}) = \frac{\theta}{12} - \frac{\theta}{4}\csc^2(\frac{\sqrt{-\theta}}{2}\,\xi), \theta < 0, \tag{16}$$

where θ is a real number.

A given NLEE with respect to two variables x and t is of the form

$$P(u, u_x, u_t, u_{xx}, u_{xt}, u_{tt}, u_{xxx}, \cdots) = 0, \tag{17}$$

where the subscripts denote the partial derivatives,P is a polynomial in unknown function $u(x,t)$ and its derivatives.

The Weierstrass type projective Riccati equation expansion method proposed here follows the following five steps.

Step 1: Making the wave transformation

$$u(x,t) = u(\xi), \xi = x - \omega t, \tag{18}$$

we may exchange the Eq. (17) into the following ODE

$$H(u, u', u'', \cdots) = 0, \tag{19}$$

where primes denote the derivatives with respect to ξ.

Step 2: Assume that the Eq. (19) has the truncated formal series solution

$$u(\xi) = a_0 + \sum_{i=1}^{n} F^{i-1}(\xi)\left(a_i F(\xi) + b_i G(\xi)\right), \tag{20}$$

where $F(\xi)$ and $G(\xi)$ are the Weierstrass elliptic function solution of the projective Riccati Eqs. (3), n is an integer number which can be determined by balancing the highest order derivative terms with the highest power nonlinear terms in (19),a_0,a_i, b_i $(i = 1, 2, \cdots, n)$ are undetermined constants and $a_n^2 + b_n^2 \neq 0$.

Step 3: Substituting (20), (3) together with one of the relations (5), (7), (9), (11) into (19) and equating the coefficients of like powers of $F^i(\xi)G^j(\xi)$ to zero yields a set of algebraic equations. Solving this set of algebraic equations with the aid of Maple or Mathematica we can determine the values of ω, a_0 and a_i, b_i $(i = 1, 2, \cdots, n)$.

Step 4: Putting each solutions of the algebraic equations obtained in *Step 3* together with the Weierstrass elliptic function solutions $F(\xi), G(\xi)$ of the projective Riccati equations into (20) and using (18), we obtain the Weierstrass type traveling wave solutions of the Eq. (17).

Step 5: Taking the conversion formulas into the Weierstrass elliptic function type traveling wave solutions obtained in *Step 4*, we get the solitary wave and the periodic wave solutions of the Eq. (17).

3 Solutions of the KdV equation

Now let us consider the famous KdV equation

$$u_t + 6uu_x + u_{xxx} = 0. \tag{21}$$

Taking the wave transformation (18) into (21) we get the following ODE

$$-\omega u'(\xi) + 6u(\xi)u'(\xi) + u'''(\xi) = 0. \tag{22}$$

By using the homogenous balance method we can determine that $n = 2$.Thus the solution of Eq. (22) can be chosen as

$$u(\xi) = a_0 + a_1 F(\xi) + b_1 G(\xi) + a_2 F^2(\xi) + b_2 F(\xi)G(\xi), \tag{23}$$

where $F(\xi), G(\xi)$ are the solution of the projective Riccati equations (3),a_0, a_1, a_2, b_1, b_2 are undermined constants.

(1) Taking (23) with (3),(5) into (22) and setting the coefficients of F^iG^j $(i = 1, 2, 3, 4; j = 0, 1)$ to zero we obtain a set of algebraic equations

$$\begin{cases} 12a_2^2 p = 0, 42a_2 b_2 r = 0, \\ 30a_2 p^2 r + 18a_1 a_2 p + 18b_2^2 r = 0, \\ 30b_2 p r^2 + 30a_1 b_2 r + 30a_2 b_1 r - 18a_2 b_2 q = 0, \\ -a_1 p^2 q + 6a_0 a_1 p - \omega a_1 p + 6b_1^2 r - 6b_1 b_2 q = 0, \\ -b_1 pqr + b_2 pq^2 + 6a_0 b_1 r - 6a_0 b_2 q - 6a_1 b_1 q - \omega b_1 r + \omega b_2 q = 0, \\ 3b_1 p r^2 - 15b_2 pqr + 18a_0 b_2 r + 18a_1 b_1 r - 12a_1 b_2 q - 12a_2 b_1 q - 3\omega b_2 r = 0, \\ 6a_1 p^2 r - 8a_2 p^2 q + 12a_0 a_2 p + 6a_1^2 p - 2\omega a_2 p + 24b_1 b_2 r - 6b_2^2 q = 0. \end{cases}$$

Solving this set of algebraic equations with use of Maple we get

$$a_0 = \frac{1}{6}\left(qp+\omega\right), a_1 = -pr, a_2 = 0, b_1 = 0, b_2 = 0. \tag{24}$$

Substituting (24),(4),(20) into (23) we obtain the Weierstrass type traveling wave solution of the KdV equation

$$u(x,t) = \frac{\omega}{6} - 2\wp(x-\omega t, g_2, g_3), \tag{25}$$

where the invariants g_2, g_3 are determined by (12).

To take the conversion formula (13)–(16) with $\theta = -qp$ into (25) we obtain the solitary wave and periodic wave solutions of the KdV equation

$$\begin{aligned}
u_1^{(1)}(x,t) &= \frac{qp+\omega}{6} - \frac{qp}{2}\,\mathrm{sech}^2\frac{\sqrt{-\mathrm{qp}}}{2}(\mathrm{x}-\omega\mathrm{t}), \mathrm{qp}<0,\\
u_2^{(1)}(x,t) &= \frac{qp+\omega}{6} + \frac{qp}{2}\,\mathrm{csch}^2\frac{\sqrt{-\mathrm{qp}}}{2}(\mathrm{x}-\omega\mathrm{t}), \mathrm{qp}<0,\\
u_3^{(1)}(x,t) &= \frac{qp+\omega}{6} - \frac{qp}{2}\,\sec^2\frac{\sqrt{qp}}{2}(x-\omega t), qp>0,\\
u_4^{(1)}(x,t) &= \frac{qp+\omega}{6} - \frac{qp}{2}\,\csc^2\frac{\sqrt{qp}}{2}(x-\omega t), qp>0,
\end{aligned}$$

where p, q, ω are constants.

(2) Taking (23) with (3), (7) into (22) and setting the coefficients of F^iG^j $(i = 1,2,3,4,5; j = 0,1)$ to zero we obtain the following set of algebraic equations

$$\begin{cases}
-\dfrac{576a_2b_2r^2}{25q} + \dfrac{13824b_2pr^4}{625q^2} = 0,\\
-\dfrac{288b_2^2r^2}{25q} - \dfrac{576a_2p^2r^2}{25q} + 12a_2^2p = 0,\\
18a_2a_1p - \dfrac{144a_1p^2r^2}{25q} - \dfrac{432b_1b_2r^2}{25q} + 18b_2^2r + 30a_2p^2r = 0,\\
-\dfrac{432b_2a_1r^2}{25q} - \dfrac{432b_1a_2r^2}{25q} + \dfrac{3456b_1pr^4}{625q^2} - \dfrac{288b_2pr^3}{5q} + 42a_2b_2r = 0,\\
-2\omega a_2p + 12a_0a_2p + 24b_1b_2r - \dfrac{144b_1^2r^2}{25q} + 6a_1^2p - 6b_2^2q + 6a_1p^2r - 8a_2p^2q = 0,\\
\dfrac{48\omega b_2r^2}{25q} - \dfrac{288b_1a_1r^2}{25q} - \dfrac{288a_0b_2r^2}{25q} - \dfrac{288b_1pr^3}{25q} + \dfrac{246b_2pr^2}{5} - 18a_2b_2q\\
\quad + 30a_2b_1r + 30a_1b_2r = 0,\\
\dfrac{24\omega b_1r^2}{25q} - \dfrac{144a_0b_1r^2}{25q} + \dfrac{171b_1pr^2}{25} - 15b_2rpq - 3\omega b_2r - 12a_2b_1q - 12a_1b_2q\\
\quad + 18a_1b_1r + 18a_0b_2r = 0,\\
-a_1p^2q + 6a_0a_1p - \omega a_1p + 6b_1^2r - 6b_1b_2q = 0,\\
-b_1pqr + b_2pq^2 + 6a_0b_1r - 6a_0b_2q - 6a_1b_1q - \omega b_1r + \omega b_2q = 0.
\end{cases}$$

Using Maple to solve above algebraic equations we obtain

$$a_0 = \frac{pq+\omega}{6}, a_1 = -pr, a_2 = \frac{24pr^2}{25q}, b_1 = 0, b_2 = 2\varepsilon pr\sqrt{-\frac{6p}{25q}}, \tag{26}$$

where $\varepsilon = \pm 1$ and $pq < 0$.

Substituting (26), (6) and (18) into (23) leads the Weierstrass type traveling wave solution of the KdV equation

$$u(x,t) = \frac{9\left[4\omega\wp(\xi) + p^2q^2\right]\wp(\xi) + pq\left[p^2q^2 - 6\varepsilon\sqrt{-6pq}\wp'(\xi)\right]}{216\wp^2(\xi)}, \tag{27}$$

where $pq < 0$,$\wp(\xi) = \wp(x - \omega t, g_2, g_3)$, the invariants g_2,g_3 are determined by (12).

Because $pq < 0$, so it is known from the conversion formulas that the Weierstrass function solution (27) cannot degenerate to the periodic solutions. Thus, substituting the conversion formulas (13) and (14) with $\theta = -qp$ into (27) leads the solitary wave like solutions of the KdV equation as following

$$u_1^{(2)}(x,t) = \frac{(pq+\omega)\cosh^4\eta + 3(3pq-2\omega)\cosh^2\eta - 6\sqrt{6}\varepsilon pq\sinh\eta\cosh\eta + 9\omega}{6\left(\cosh^2\eta - 3\right)^2},$$

$$u_2^{(2)}(x,t) = \frac{(pq+\omega)\sinh^4\eta - 3(3pq-2\omega)\sinh^2\eta + 6\sqrt{6}\varepsilon pq\sinh\eta\cosh\eta + 9\omega}{6\left(\sinh^2\eta + 3\right)^2},$$

where $\eta = \frac{1}{2}\sqrt{-pq}(x - \omega t)$,$pq < 0$.

(3) Inserting (23) with (3), (9) into (22) and setting the coefficients of F^iG^j $(i = 1,2,3,4,5; j = 0,1)$ to zero we obtain the following set of algebraic equations

$$\begin{cases}
12a_2^2p - \dfrac{48b_2^2r^2 + 96a_2p^2r^2 + 12b_2^2r^2q + 24a_2p^2r^2q}{(q+2)^2} = 0, \\
\dfrac{384b_2pr^4 + 24b_2pr^4q^2 + 192b_2pr^4q}{(q+2)^4} - \dfrac{96a_2b_2r^2 + 24a_2b_2r^2q}{(q+2)^2} = 0, \\
18a_2a_1p - \dfrac{24a_1p^2r^2 + 72b_1b_2r^2 + 6a_1p^2r^2q + 18b_1b_2r^2q}{(q+2)^2} + 18b_2^2r + 30a_2p^2r = 0, \\
12a_0a_2p - 2\omega a_2p + 24b_1b_2r + 6a_1^2p - 6b_2^2q + 6a_1p^2r - 8a_2p^2q - \dfrac{6b_1^2r^2q + 24b_1^2r^2}{(q+2)^2} = 0, \\
42a_2b_2r - \dfrac{18b_2a_1r^2q + 18b_1a_2r^2q + 65b_2pr^3q + 260b_2pr^3 + 72b_2a_1r^2 + 72b_1a_2r^2}{(q+2)^2} \\
\quad + \dfrac{6b_1pr^4q^2 + 48b_1pr^4q + 96b_1pr^4}{(q+2)^4} = 0, \\
\dfrac{4b_1pr^2q^2 + 16b_1pr^2q + 4\omega b_1r^2 - 24a_0b_1r^2 + \omega b_1r^2q - 6a_0b_1r^2q}{(q+2)^2} - 19b_2prq + 6b_1pr^2 \\
\quad - 3\omega b_2r - 12a_2b_1q - 12a_1b_2q + 18a_1b_1r + 18a_0b_2r = 0, \\
\dfrac{2\omega b_2r^2q - 14b_1pr^3q - 12b_1a_1r^2q - 12a_0b_2r^2q - 56b_1pr^3}{(q+2)^2} - 18a_2b_2q + 30a_2b_1r \\
\quad + 30a_1b_2r + 39b_2pr^2 + \dfrac{20b_2pr^2q^2 + 80b_2pr^2q + 8\omega b_2r^2 - 48b_1a_1r^2 - 48a_0b_2r^2}{(q+2)^2} = 0, \\
-a_1p^2q + 6a_0a_1p - \omega a_1p + 6b_1^2r - 6b_1b_2q = 0, \\
-2b_1pqr + b_2pq^2 + 6a_0b_1r - 6a_0b_2q - 6a_1b_1q - \omega b_1r + \omega b_2q = 0.
\end{cases}$$

Solving this set of algebraic equations with aid of Maple we obtain

$$a_0 = \frac{\omega}{6} - \frac{2p}{3}, a_1 = -pr, a_2 = 0, b_1 = 0, b_2 = 0, q = -4. \tag{28}$$

From which we calculate that

$$g_2 = \frac{4p^2}{3}, g_3 = -\frac{8p^3}{27}. \tag{29}$$

Substituting (28) with (29),(8) into (23) we get the Weierstrass type traveling wave solution of the KdV equation

$$u(x,t) = \frac{\omega}{6} - \frac{2p}{3} - \frac{2p\left[p - 3\wp(x-\omega t, g_2, g_3)\right]}{2p + 3\wp(x-\omega t, g_2, g_3)}, \tag{30}$$

where the invariants g_2, g_3 are given by (12).

Taking the conversion formulas (13)–(16) with $\theta = 4p$ into (30) leads the solitary and periodic wave solutions of the KdV equation

$$\begin{aligned}
u_1^{(3)}(x,t) &= \frac{\omega - 4p}{6} - 2p\,\mathrm{csch}^2\sqrt{\mathrm{p}}(\mathrm{x}-\omega\mathrm{t}), \mathrm{p} > 0,\\
u_2^{(3)}(x,t) &= \frac{\omega - 4p}{6} + 2p\,\mathrm{sech}^2\sqrt{\mathrm{p}}(\mathrm{x}-\omega\mathrm{t}), \mathrm{p} > 0,\\
u_3^{(3)}(x,t) &= \frac{\omega - 4p}{6} + 2p\,\csc^2\sqrt{-p}(x-\omega t), p < 0,\\
u_4^{(3)}(x,t) &= \frac{\omega - 4p}{6} + 2p\,\sec^2\sqrt{-p}(x-\omega t), p < 0.
\end{aligned}$$

We can see that the above solutions $u_i^{(3)}\,(1,2,3,4)$ are the special case $q = -4$ of the solutions $u_i^{(1)}\,(i = 2,1,4,3)$. Here, we need to pointed out that the case of other NLEEs, the Weierstrass elliptic function solutions (4) and (8) usually can give different types of solutions. Therefore, the solutions (4) and (8) are the different solutions of the projective Riccati equation.




Fig. 1 Plots of the solitary wave like solutions. **(a)** solution $u_1^{(1)}$ with $p = -1, q = 1, \omega = 1.025$. **(b)** solution $u_2^{(2)}$ with $p = 1, q = -1, \omega = 0.995, \varepsilon = 1$. **(c)** solution $u_1^{(4)}$ with $p = -1, q = 1, \omega = 1.025, \varepsilon = -1$.

(4) Taking (23) with (3),(11) into (22) and setting the coefficients of F^iG^j $(i =$

$1,2,3,4,5; j=0,1)$ to zero we obtain the following set of algebraic equations

$$\begin{cases}
-\dfrac{24a_2p^6r^2+96a_2p^4r^2+12b_2^2p^4r^2+48b_2^2p^2r^2}{q(p^2+2)^2}+12a_2^2p=0,\\
-\dfrac{24a_2b_2p^4r^2+96a_2b_2p^2r^2}{q(p^2+2)^2}+\dfrac{24b_2p^9r^4+192b_2p^7r^4+384b_2p^5r^4}{q^2(p^2+2)^4}=0,\\
-\dfrac{6a_1p^6r^2+24a_1p^4r^2+18b_1b_2p^4r^2+72b_1b_2p^2r^2}{q(p^2+2)^2}\\
\quad+30a_2p^2r+18b_2^2r+18a_2a_1p=0,\\
-\dfrac{6b_1^2p^4r^2+24b_1^2p^2r^2}{q(p^2+2)^2}-2\omega a_2p+6a_1^2p+6a_1p^2r-8a_2p^2q\\
\quad+24b_2b1r-6b_2^2q+12a_0a_2p=0,\\
-\dfrac{18b_2a_1p^4r^2+72b_2a_1p^2r^2+18b_1a_2p^4r^2+72b_1a_2p^2r^2+60b_2p^5r^3+240b_2p^3r^3}{q(p^2+2)^2}\\
\quad+42a_2b_2r+\dfrac{6b_1p^9r^4+48b_1p^7r^4+96b_1p^5r^4}{q^2(p^2+2)^4}=0,\\
\dfrac{4b_1p^5r^2+16b_1p^3r^2}{(p^2+2)^2}-15b_2prq+3b_1r^2p-3\omega b_2r-12a_2b_1q-12a_1b_2q+18a_1b_1r\\
\quad+18a_0b_2r+\dfrac{\omega b_1p^4r^2+4\omega b_1p^2r^2-6a_0b_1p^4r^2-24a_0b_1p^2r^2}{q(p^2+2)^2}=0,\\
\dfrac{2\omega b_2p^4r^2+8\omega b_2p^2r^2-12b_1a_1p^4r^2-48b_1a_1p^2r^2-12a_0b_2p^4r^2-48a_0b_2p^2r^2}{q(p^2+2)^2}\\
\quad-\dfrac{12b_1p^5r^3+48b_1p^3r^3}{q(p^2+2)^2}+\dfrac{20b_2p^5r^2+80b_2p^3r^2}{(p^2+2)^2}-18a_2b_2q+30a_2b_1r\\
\quad+30a_1b_2r+30b_2pr^2=0,\\
-a_1p^2q+6a_0a_1p-\omega a_1p+6b_1^2r-6b_1b_2q=0,\\
-b_1pqr+b_2pq^2+6a_0b_1r-6a_0b_2q-6a_1b_1q-\omega b_1r+\omega b_2q=0.
\end{cases}$$

Solution of this set of algebraic equations found by using Maple reads

$$a_0=\frac{qp+\omega}{6}, a_1=-pr, a_2=\frac{p^3r^2(p^2+4)}{q(p^2+2)^2}, b_1=0, b_2=\frac{\varepsilon rp^2}{p^2+2}\sqrt{-\frac{p(p^2+4)}{q}}, \tag{31}$$

where $\varepsilon=\pm1$.

Taking (10),(31),(18) into (23) we obtain the Weierstrass type traveling wave solution of the KdV equation

$$\begin{aligned}
u(x,t)&=\frac{144p\left(p^2q+\omega p+12q\right)\wp^2(\xi)+a\wp(\xi)+864q^2\varepsilon\sqrt{-\frac{p^3+4p}{q}}\wp'(\xi)+a}{6\left(p^2q+12q+12p\wp(\xi)\right)^2},\\
a&=24q\left(p^4q+\omega p^3-12p^2q+12\omega p-72q\right),\\
b&=q^2\left(p^5q+\omega p^4-36p^3q+24\omega p^2+144\omega\right),
\end{aligned} \tag{32}$$

where $\wp(\xi)=\wp(x-\omega t,g_2,g_3)$, the invariants g_2 and g_3 are given by (12).

Note that $pq<0$ in (31) and (32), so (32) can only give the solitary wave solution of the KdV equation. Thus by substituting the conversion formulas (13) and (14) with

$\theta = -qp$ into (32) we obtain the solitary wave like solutions of the KdV equation as follows

$$u_1^{(4)}(x,t) = \frac{24p^2q\varepsilon\sqrt{p^2+4}\,\sinh\eta\cosh\eta + 16(pq+\omega)\cosh^4\eta + r\cosh^2\eta + s}{6\left(4\cosh^2\eta + p^2\right)^2},$$

$$u_2^{(4)}(x,t) = \frac{-24p^2q\varepsilon\sqrt{p^2+4}\,\sinh\eta\cosh\eta + 16(pq+\omega)\sinh^4\eta - r\sinh^2\eta + s}{6\left(4\sinh^2\eta - p^2\right)^2},$$

$$r = 8p(-2p^2q + \omega p - 6q), s = p^3(p^2q + \omega p + 12q),$$

where $\eta = \frac{1}{2}\sqrt{-pq}(x - \omega t), qp < 0$.

It is obvious that the obtained solutions of the KdV equation are related to the coefficients p, q and r of the projective Riccati equations (3). The solitary wave like solutions, the singular solitary wave solutions and the periodic solutions of the KdV equation are plotted in Fig.1,Fig.2 and Fig.3.The graphs of the solutions $u_i^{(3)}\ (i = 1, 2, 3, 4)$ are completely similar to that of the solutions $u_i^{(1)}\ (i = 1, 2, 3, 4)$, so it is omitted.




Fig. 2 Plots of the singular solitary wave solutions. **(a)** solution $u_2^{(1)}$ with $p = 1, q = -1, \omega = 1.025$. **(b)** solution $u_1^{(2)}$ with $p = -1, q = 1, \omega = 1.025, \varepsilon = -1$. **(c)** solution $u_2^{(4)}$ with $p = 1, q = -1, \omega = 1.025, \varepsilon = 1$.

4 Conclusions

In the present paper, the Weierstrass type projective Riccati equation expansion method is proposed to construct Weierstrass elliptic function solutions of NLEEs. At the same time, the conversion formulas are also proposed to transform these Weierstrass elliptic function solutions into the hyperbolic and trigonometric function solutions of NLEEs. Our method is more applicable than the projective Riccati equation expansion method and it can be regarded as an extension of the projective Riccati equation expansion method. In order to explain our method more clear we need to point out the following three points.

(1) The Weierstrass elliptic function solutions (4) and (6) of the projective Riccati equations are known [16, 17, 18], and the solutions (8) and (10) are new.

(2) The known conversion formula such as [16, 17]

$$\wp(\xi, g_2, g_3) = e_2 - (e_2 - e_3)\mathrm{cn}^2\left(\sqrt{e_1 - e_3}\,\xi, m\right), \tag{33}$$




Fig. 3 Plots of the periodic solutions. **(a)** solution $u_3^{(1)}$ with $p = 1, q = 1, \omega = 1.025$. **(b)** solution $u_4^{(1)}$ with $p = 1, q = 1, \omega = 1.025$.

is related to the roots e_1, e_2, e_3 of the third order polynomial equation $w^3 - g_2 w - g_3 = 0$,but our conversion formulas (13)– (16) don't require these roots. Using formula (33), we can transform the Weierstrass elliptic function solutions of NLEEs into the Jacobian elliptic function solutions, and then the hyperbolic and trigonometric function solutions are obtained by taking the limit of modulus $m \to 1$ and $m \to 0$, but they may be incorrect [19, 20]. In contrast, our formulas (13)–(16) can directly convert the Weierstrass elliptic function solutions of NLEEs into the hyperbolic and trigonometric function solutions, and ensure that the obtained solutions are correct. In addition, the conversion formulas (13)–(16) also can be used in other Weierstrass elliptic function methods to transform the Weierstrass elliptic function solutions into the hyperbolic and trigonometric function solutions.

(3) Although we have concerned with the KdV equation, our method can be applied to construct the exact solitary wave and periodic wave solutions of a wide class of NLEEs.

References

[1] Malfliet W. Solitary wave solutions of nonlinear wave equations. Am. J. Phys., **60**(1992): 650–654.

[2] Sirendaoreji, Sun J. Auxiliary equation method for solving nonlinear partial differential equations. Phys. Lett. A, **309**(2003): 387–396.

[3] Peng Y Z. Exact solutions for the generalized variable coefficient KdV equation. Fiz. A, **12**(2003): 9–14.

[4] Sirendaoreji, Unified Riccati equation expansion method and its application to two new classes of Benjamin–Bona–Mahony equations. Nonlinear Dyn., **89**(2017): 333-344.

[5] Liu S K, Fu Z T, Liu S D, Zhao Q. Jacobi elliptic function expansion method and periodic wave solutions of nonlinear wave equations. Phys. Lett. A, **289**(2001): 69–74.

[6] Porubov A V, Velarde M G. Exact periodic solutions of the complex Ginzburg–Landau equation. J. Math. Phys., **40**(1999): 884–896

[7] Huber A. A novel class of solutions for a non-linear third order wave equation generated by the Weierstrass transformation. Chaos Solitons & Fractals, **28**(2006): 972–978.

[8] Nickel J. Elliptic solutions to a generalized BBM equation. Phys. Lett. A, **364**(2007): 221–226.

[9] Yuan W J, Huang Z F, Lai J C, Qi J M. All meromorphic solutions of some algebraic differential equations. Anal. Theory Appl., **30**(2014): 90–107.

[10] Achab A El. Constructing of exact solutions to the nonlinear Schrödinger equation(NLSE) with power–law nonlinearity by the Weierstrass elliptic function method. Optik, **127**(2016): 1229–1232.

[11] Conte R, Musette M. Link between solitary waves and projective Riccati equation. Phys. A: Math. Gen., **25**(1992): 2609–2612.

[12] Zhang G X, Li Z B, Duan Y S. Exact solitary wave solutions of the nonlinear wave equations. Sci. China A, **30**(2000): 1103–1108.

[13] Yan Z Y. Generalized method and its application in the higher–order nonlinear Schrödinger equation in onlinear obtical fibres. Chaos, Solitons & Fractals, **16**(2003): 759–766.

[14] Chen Y, Li B. General projective Riccati equation method and exact solutions for generalized KdV–type and KdV–Burgers–type equations with nonlinear terms of any order. Chaos,Solitons & Fractals,**19**(2004): 977–984.

[15] Shi L M, Zhang S J, Zhu R Y. New solutions to Boussinesq–Burgers equation. Acta Sinica Quant. Opt., **19**(2013): 18–25.

[16] Li D S, Zhang H Q. A new method to construct Weierstrass elliptic function solutions for soliton equations. Acta Physica Sinica, **54**(2005): 5540–5543.

[17] Guo G P. Weierstrass elliptic solutions for MKdV equation. J. Shangqiu Teachers College, **23**(2007): 61–63.

[18] Yu J P, Sun Y L. Weierstrass elliptic solutions to nonlinear evolution equations. Commun. Theor. Phys., **50**(2008): 295–298.

[19] Li Z B. Travelling Wave Solutions of Nonlinear Equations in Mathematical Physics. Science Press, Beijing, (2007).

[20] Sirendaoreji. Travelling Wave Solutions for Nonlinear Wave Equations– Theory and Appliactions of the Auxiliary Equation Method. Science Press, Beijing, (2019).