

Stochastic first-order methods for average-reward Markov decision processes

Tianjiao Li^{*} Feiyang Wu[†] Guanghui Lan^{*}

^{*}School of Industrial and Systems Engineering

[†]School of Computer Science
Georgia Institute of Technology

Abstract

We study the problem of average-reward Markov decision processes (AMDPs) and develop novel first-order methods with strong theoretical guarantees for both policy evaluation and optimization. Existing on-policy evaluation methods suffer from sub-optimal convergence rates as well as failure in handling insufficiently random policies, e.g., deterministic policies, for lack of exploration. To remedy these issues, we develop a novel variance-reduced temporal difference (VRTD) method with linear function approximation for randomized policies along with optimal convergence guarantees, and an exploratory variance-reduced temporal difference (EVRTD) method for insufficiently random policies with comparable convergence guarantees. We further establish linear convergence rate on the bias of policy evaluation, which is essential for improving the overall sample complexity of policy optimization. On the other hand, compared with intensive research interest in finite sample analysis of policy gradient methods for discounted MDPs, existing studies on policy gradient methods for AMDPs mostly focus on regret bounds under restrictive assumptions on the underlying Markov processes (see, e.g., [Abbasi-Yadkori et al., 2019](#)), and they often lack guarantees on the overall sample complexities. Towards this end, we develop an average-reward variant of the stochastic policy mirror descent (SPMD) ([Lan, 2022](#)). We establish the first $\tilde{O}(\epsilon^{-2})$ sample complexity for solving AMDPs with policy gradient method under both the generative model (with unichain assumption) and Markovian noise model (with ergodic assumption). This bound can be further improved to $\tilde{O}(\epsilon^{-1})$ for solving regularized AMDPs. Our theoretical advantages are corroborated by numerical experiments.

1 Introduction

Reinforcement learning (RL) problems are generally formulated as Markov decision processes (MDPs). At each time step, the agent observes the current state and subsequently takes an action, which leads to an instantaneous reward or cost as well as the transition to the next state according to an unknown transition kernel. The eventual goal of the agent is to learn a policy to optimize the reward accrued (or cost paid) over time. MDPs consist of two prominent classes, including the discounted MDPs (DMDPs), which introduce a discount factor to the measure of accumulated reward/cost, and the average-reward MDPs (AMDPs), which measure the success of the system in terms of its steady-state performance. Although DMDP has been studied extensively, it usually leads to poor long-term performance when a system operates for an extended time period. The average-reward criterion used in AMDPs can alleviate this issue, hence has been widely used in many different applications ranging from engineering to natural and social sciences; see, e.g., [Kober et al. \(2013\)](#); [Xu et al. \(2014\)](#) for surveys of applications.

Since the transition kernel is unknown, one important objective in RL studies is to understand its sample complexity, which characterizes how many samples are required to obtain a nearly-optimal policy up to a given accuracy. Finite sample analyses are often carried out under different observation models. The two most well-known observation models considered by the literature are the so-called *generative model* (“i.i.d.” model) and the *Markovian noise model*, where the first one provides an oracle that returns the next state when queried on a state-action pair, while the latter one only returns state transitions of an MDP trajectory. To facilitate the analysis, these observation models are also augmented with various assumptions on the behavior of the underlying MDPs (e.g., mixing unichain and ergodicity). In comparison with significant progress recently made on the finite sample analysis of DMDPs, theoretical studies on the finite sample analysis of AMDPs, especially under the Markovian noise model, are rather limited.

It is well-known that one can formulate AMDPs as a linear programming (LP) problem (see, e.g., Puterman, 2014). As a result, one can utilize the rich theory in LP and specialize some advanced algorithms for solving AMDPs. In particular, to deal with the unknown transition kernels, current research has been focused on utilizing general stochastic primal-dual saddle point optimization methods (see Chapter 4 of Lan, 2020), and some important progresses along this research direction have been made in Wang (2017); Jin and Sidford (2020). In spite of the elegance of the stochastic LP approach, there exist a few significant limitations associated with this approach. Firstly, due to the high dimension of the state space, the size of LP can be huge. One common practice is to use function approximation with linear kernel or neural networks for the value functions. However, the LP approach cannot deal with function approximations. Secondly, in MDP and RL, one usually needs to incorporate certain nonlinear objectives and/or constraints. One prominent example is a regularization given by Kullback–Leibler (KL) divergence, which can be used to model the difference between the desirable policy and a reference policy. The LP approach will fail to handle these nonlinear components. Thirdly, in the current stochastic LP approaches for RL, one needs to have access to a generative model to simulate the probability of moving to the next state for any state-action pair. As a consequence, this type of approach cannot be applied to the Markovian noise setting.

Some classic dynamic programming methods, especially value iteration, have been adapted for AMDPs with unknown transition kernels (Abounadi et al., 2001; Gosavi, 2004; Wan et al., 2021; Zhang et al., 2021b). These methods are often called value-based methods or Q-learning in the RL literature. Even though value-based methods are extensively studied in the DMDP literature and optimal finite sample guarantees are established (Wainwright, 2019; Li et al., 2020), the only known finite sample analysis of Q-learning for AMDPs up to our knowledge is Zhang et al. (2021b). However, Zhang et al. (2021b) only focused on the generative model, and the overall sample complexity is significantly worse than LP methods (Wang, 2017; Jin and Sidford, 2020). It is also noteworthy that a recent work of Jin and Sidford (2021) established a reduction method that solves an AMDP by solving an associated DMDP. Using this reduction technique, while the dependence on the problem parameters, e.g., mixing time, is improved, the dependence on the accuracy measure is sub-optimal, i.e., $\mathcal{O}(1/\epsilon^3)$.

In this paper, we focus on the study of a new class of first-order methods for solving AMDPs that involve both *policy evaluation* and *policy improvement* steps. The framework is also known as the *actor-critic* method in the RL literature, and generalizes the classic policy iteration method. Specifically, the actor aims to perform the policy update while the critic is dedicated to evaluating the value associated with the currently generated policy, which constitutes first-order information

of the nonlinear policy optimization problem. In comparison with the LP method and value-based method, these policy-based first-order methods offer remarkable flexibility for function approximation as well as policy regularization. Meanwhile, the model-free framework of policy-based methods naturally accommodates both the generative and the Markovian noise settings. Recently, there has been considerable interest in the development and analysis of first-order methods for DMDPs. In particular, sublinear convergence guarantees of policy gradient method and its variants are discussed in Agarwal et al. (2021) and Shani et al. (2020). Cen et al. (2021) showed that natural policy gradient (NPG) converges linearly for entropy-regularized DMDPs. Lan (2022) proposed the policy mirror descent (PMD) method which achieved linear convergence for DMDPs, and $\tilde{\mathcal{O}}(1/\epsilon)$ (resp., $\tilde{\mathcal{O}}(1/\epsilon^2)$) sampling complexity for regularized (resp., unregularized) DMDPs. Further studies for these types of methods can be found in Khodadadian et al. (2021); Zhan et al. (2021); Lan et al. (2022); Xiao (2022); Li et al. (2022). Nevertheless, theoretical understanding of policy-based methods for AMDPs remains very limited. It is noteworthy that Abbasi-Yadkori et al. (2019); Lazic et al. (2021) proposed a policy-based method named POLITEX for AMDPs, but their analysis requires restrictive assumptions on the transition kernel and they only established regret bounds without sample complexities (see early related work Bartlett and Tewari, 2012; Auer et al., 2008; Ouyang et al., 2017)).

Under the Markovian noise model, the literature distinguishes two types of actor-critic setup, namely on-policy actor-critic (Konda and Tsitsiklis, 1999), where the samples can be collected based on the current policy, and off-policy actor-critic (Degris et al., 2012), where the samples are collected based on a behavior policy. Both settings have a broad range of applications in practice. The off-policy setting enables offline learning and encourages exploration of the agent (Wan et al., 2021; Zhang et al., 2021a), while in the on-policy setting, the practitioners can continuously improve the current policy and accrue the instantaneous rewards generated by nearly-optimal policies. One significant challenge associated with the on-policy setting exists in the possible lack of exploration, which happens when the policy is not random enough. Consequently, existing on-policy first-order policy optimization methods require restrictive assumptions, e.g., all the iterates of policies are random enough, which is problematic when the optimal policy does not possess this structure (e.g., being deterministic). In spite of intensive research effort of on-policy evaluation (Tsitsiklis and Van Roy, 1999; Yu and Bertsekas, 2009; Zhang et al., 2021b; Mou et al., 2022), one seemingly unresolved problem in RL is whether one can design sampling-efficient on-policy evaluation algorithms for insufficiently random policies and use them for policy optimization (see Remark 1 of Lan, 2022).

1.1 Main contributions

In this work, we make three distinct contributions.

- **Policy evaluation for AMDPs (Critic):** We first propose a simple and novel multiple trajectory method for policy evaluation in the generative model, which achieves $\mathcal{O}(t_{\text{mix}} \log(1/\epsilon))$ sample complexity for ℓ_∞ -bound on the bias of the estimators, as well as $\mathcal{O}(t_{\text{mix}}^2/\epsilon)$ sample complexity for the expected squared ℓ_∞ -error of the estimators. For the on-policy evaluation under Markovian noise, we develop an average-reward variant of the variance-reduced temporal difference (VRTD) algorithm (Khamaru et al., 2021; Li et al., 2021) with linear function approximation, which achieves $\mathcal{O}(t_{\text{mix}}^3 \log(1/\epsilon))$ sample complexity for weighted ℓ_2 -error of the bias of the estimators, as well as an instance-dependent sample complexity for expected weighted ℓ_2 -error of the estimators. The latter complexity improved the one in Zhang et al.

(2021b) by at least a factor of $\mathcal{O}(t_{\text{mix}}^2)$. Meanwhile, we develop a novel exploratory variance-reduced temporal difference (EVRTD) that remedies the exploration issue when the policy is not random enough. The design idea of the EVRTD algorithm naturally extends to DMDPs. We show that the sample complexity of EVRTD is not much worse than VRTD for both AMDP and DMDP.

- **Policy mirror descent for AMDPs (Actor):** Motivated by the policy mirror descent (PMD) algorithm for solving DMDPs (Lan, 2022), we develop the stochastic average-reward PMD method. In the convergence analysis, we handle the expected error and bias error from the critic step separately, leading to sharpened convergence results. For unregularized AMDP, the SPMD method achieves $\mathcal{O}(1/\sqrt{T})$ convergence rate. For regularized AMDP with the ergodicity assumption, we show that all the convergence guarantees for DMDPs in Lan (2022) can be reproduced in the AMDP setting. Furthermore, we establish a unified convergence guarantee on the distance to the optimal policy, with linear rate convergence in terms of deterministic error and $\mathcal{O}(\frac{1}{\omega^2 T})$ convergence (where ω stands for the strongly convex modulus of the regularizer) in terms of stochastic error.
- **Sample complexity:** We establish the overall sample complexity under both the generative model and Markovian noise model. Under the mixing unichain setting with a generative model, by implementing the SPMD algorithm with multiple trajectory method as the critic, we achieve $\tilde{\mathcal{O}}(\frac{t_{\text{mix}}^3 |S| |A|}{\epsilon^2})$ overall sample complexity. As for the ergodic setting with the Markovian observation model, we implement SPMD with VRTD/EVRTD as the critic, which achieves $\tilde{\mathcal{O}}(\epsilon^{-2})$ sample complexity. For regularized AMDPs, this bound is improved to $\tilde{\mathcal{O}}(\frac{1}{\omega^2 \epsilon})$ for obtaining an ϵ -optimal policy in terms of distance to the optimal solution. To the best of our knowledge, these sample complexities are new in the literature on solving average-reward RL problems with policy-based (first-order) methods.

1.2 Notation

For a positive integer n , we define $[n] := \{1, 2, \dots, n\}$. We let $\mathbf{1}_n$ denote the all-ones vector with dimension n , and we will write $\mathbf{1}$ when the dimension is clear in the context. We let e_j denote the j -th standard basis vector in \mathbb{R}^D . Let $\mathbb{I}_S : X \rightarrow \{0, 1\}$ denote the indicator function of the subset $S \subseteq X$. Given a vector $x \in \mathbb{R}^m$, denote its i -th entry by $x_{(i)}$. In situations in which there is no ambiguity, we also use x_i to denote the i -th coordinate of a vector x . Let $\|x\|_1 := \sum_{i=1}^m |x_{(i)}|$, $\|x\|_2 := \sqrt{\sum_{i=1}^m x_{(i)}^2}$ and $\|x\|_\infty := \max_{i \in [m]} |x_{(i)}|$ denote the ℓ_1 , ℓ_2 and ℓ_∞ -norms respectively. For a random variable $X \in \mathbb{R}$, let $\|X\|_{\psi_1} := \inf\{t > 0 : \mathbb{E}[\exp(|X|/t)] \leq 2\}$ and $\|X\|_{\psi_2} := \inf\{t > 0 : \mathbb{E}[\exp(X^2/t^2)] \leq 2\}$ denote the sub-exponential and sub-gaussian norm respectively. Given a matrix A , denote its (i, j) -th entry by $A_{i,j}$. Let $\|A\|_2$ denote the spectral norm of matrix A . Let $\|A\|_\infty := \max_{\|x\|_\infty=1} \|Ax\|_\infty$ denote the ℓ_∞ -operator norm of A . We let $\lambda_{\min}(A)$ denote the smallest eigenvalue of a square matrix A . For a symmetric positive (semi)definite matrix A , define $\langle x, y \rangle_A := x^\top A y$ and the associated (semi)norm $\|x\|_A := \sqrt{x^\top A x}$. We refer to $\|x\|_A$ as the ℓ_A -norm of x . We let Δ_n denote a simplex with dimension n .

2 Background and problem setting

In this section, we introduce some preliminaries for (regularized) AMDPs and the concrete observation models that we study.

2.1 Average-reward Markov decision processes

An AMDP is described by a tuple $\mathcal{M} := (\mathcal{S}, \mathcal{A}, \mathbf{P}, c)$, where \mathcal{S} denotes a finite state space, \mathcal{A} denotes a finite action space, \mathbf{P} is the transition kernel, and c is the cost function. At each time step, the agent takes an action $a \in \mathcal{A}$ at the current state $s \in \mathcal{S}$, then the system moves to some state $s' \in \mathcal{S}$ with probability $\mathbf{P}(s'|s, a)$, while the agent pays the instantaneous cost $c(s, a)$ (or receives the $-c(s, a)$ instantaneous reward). The goal of the agent is to determine a policy which minimizes the long-term cost. A randomized stationary policy of an MDP is a mapping $\pi : \mathcal{S} \rightarrow \Delta_{|\mathcal{A}|}$, i.e., it maps a state $s \in \mathcal{S}$ to a fixed distribution over actions. We denote $P_\pi \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{S}|}$ and $P^\pi \in \mathbb{R}^{(|\mathcal{S}| \times |\mathcal{A}|) \times (|\mathcal{S}| \times |\mathcal{A}|)}$ as the state transition matrix and state-action transition matrix induced by policy π , respectively, where $P_\pi(s, s') = \sum_{a \in \mathcal{A}} \pi(a|s) \mathbf{P}(s'|s, a)$ and $P^\pi((s, a), (s', a')) = \mathbf{P}(s'|s, a) \pi(a'|s')$. Under a given policy π , the (regularized) long-run average cost/reward for state $s \in \mathcal{S}$ is defined as

$$\rho^\pi(s) := \lim_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}_\pi \left[\sum_{t=0}^{T-1} (c(s_t, a_t) + h^\pi(s_t)) \mid s_0 = s \right]. \quad (2.1)$$

Here, h^π is a closed convex function with respect to the policy π , i.e., there exists $\omega \geq 0$ such that

$$h^\pi(s) - [h^{\pi'}(s) + \langle \nabla h^{\pi'}(s, \cdot), \pi(\cdot|s) - \pi'(\cdot|s) \rangle] \geq \omega D_{\pi'}^\pi(s), \quad (2.2)$$

where $\nabla h^{\pi'}$ denotes the subgradient of h at π' and $D_{\pi'}^\pi(s)$ is the Bregman distance between π and π' . In this paper, we fix $D_{\pi'}^\pi(s)$ to be the Kullback-Leibler (KL) divergence, i.e., $D_{\pi'}^\pi(s) = \sum_{a \in \mathcal{A}} \pi(a|s) \log \frac{\pi(a|s)}{\pi'(a|s)}$. It should be noted that our algorithmic framework allows us to use other distance generating functions, e.g., $\|\cdot\|_p^2$ for some $p > 1$. Clearly, if $h^\pi = 0$, then Eq. (2.1) reduces to the classical unregularized average cost function, while if $h^\pi(s) = \omega D_{\pi_0}^\pi(s)$ for $\omega > 0$ then Eq. (2.1) defines the average cost of the entropy-regularized MDP.

In this work, we consider the *unichain* setting, where the induced Markov chain consists of a single recurrent class plus a possibly empty set of transient states for any deterministic stationary policy. We further restrict our attention to *mixing* AMDPs, which satisfy the following assumption.

Assumption 1. *An AMDP instance is mixing if for any feasible policy π , there exists a stationary distribution ν^π such that for any distribution $q \in \Delta_{|\mathcal{S}|}$, the induced Markov chain has mixing time bounded by $t_{\text{mix}} < \infty$, where t_{mix} is defined as*

$$t_{\text{mix}} := \max_{\pi} \left[\arg \min_{t \geq 1} \left\{ \max_{q \in \Delta_{|\mathcal{S}|}} \|((P_\pi)^t)^\top q - \nu^\pi\|_1 \leq \frac{1}{2} \right\} \right].$$

This is a widely used regularity condition for AMDPs (see, e.g., Wang, 2017; Jin and Sidford, 2020), which can be ensured by adding the *aperiodic* assumption to the unichain setting. As a result, for any feasible policy π , the average-reward function does not depend on the initial state (see Section 8 of Puterman, 2014), and we can define a scalar which satisfies for all $s \in \mathcal{S}$

$$\rho^\pi(s) = \rho^\pi := \langle \nu^\pi, c^\pi + h^\pi \rangle,$$

where ν^π is the stationary distribution of the states induced by the policy π , and c^π is the expected cost function induced by policy π , i.e., $c^\pi(s) = \sum_{a \in \mathcal{A}} c(s, a) \pi(a|s)$. Given that one can view ρ^π as a function of π , we will use the notation ρ^π and $\rho(\pi)$ interchangeably. Note that the average-reward function only captures the “steady-state” behavior of the underlining policy, the literature defines the following differential functions to capture the “transient” behavior of the underlying policy. Specifically, the *basic differential value function* (also called bias function; see, e.g., [Puterman \(2014\)](#)) is defined as

$$\bar{V}^\pi(s) := \mathbb{E}_\pi \left[\sum_{t=0}^{\infty} (c(s_t, a_t) + h^\pi(s_t) - \rho^\pi) \mid s_0 = s \right],$$

and the *basic differential action-value function* (or *basic differential Q-function*) is defined as

$$\bar{Q}^\pi(s, a) := \mathbb{E}_\pi \left[\sum_{t=0}^{\infty} (c(s_t, a_t) + h^\pi(s_t) - \rho^\pi) \mid s_0 = s, a_0 = a \right].$$

Moreover, we define the sets of *differential value functions* and *differential action-value functions* (*differential Q-functions*) as the solution sets of the following Bellman equations, respectively,

$$V = c^\pi + h^\pi - \rho^\pi \mathbf{1}_{|\mathcal{S}|} + P_\pi V \quad (2.3)$$

$$Q = c + \tilde{h}^\pi - \rho^\pi \mathbf{1}_{|\mathcal{S}| \times |\mathcal{A}|} + P^\pi Q, \quad (2.4)$$

where $\tilde{h}^\pi(s, a) = h^\pi(s)$. Under Assumption 1, the solution of Eq. (2.3) (resp., Eq. (2.4)) is unique up to an additive constant, which takes the form of $\{\bar{V}^\pi + b \mathbf{1}_{|\mathcal{S}|} \mid b \in \mathbb{R}\}$ (resp., $\{\bar{Q}^\pi + b \mathbf{1}_{|\mathcal{S}| \times |\mathcal{A}|} \mid b \in \mathbb{R}\}$).

Finally, our goal in solving an AMDP is to find an optimal policy π^* that minimizes the long-run average cost. Consequently, the concerned *policy optimization* problem can be formulated as:

$$\min_{\pi} \quad \rho(\pi) \quad \text{s.t.} \quad \pi(\cdot|s) \in \Delta_{|\mathcal{A}|}, \quad \forall s \in \mathcal{S}. \quad (2.5)$$

2.2 Observation models

In RL, one assumes sample access to the stochastic transition kernel and cost function. This work distinguishes two classical observation models, namely the generative model (“i.i.d.” model), and the Markovian noise model.

For the generative model, we have an oracle that, upon query for the state-action pair (s, a) , returns the next state s' and the instantaneous cost $c(s, a)$. If the cost function is noiseless, the setting implicitly indicates the complete knowledge of the cost function.

In the Markovian noise model, we assume that all observed samples come from a single trajectory of a Markov chain induced by the policy π that we want to evaluate. Precisely, we have a sequence of state-action pairs $\{(s_0, a_0), (s_1, a_1), \dots\}$ generated according to the policy π and the transition kernel P . Since solving problem (2.5) requires sample access to all the states in \mathcal{S} , we make the following natural assumption for the Markovian noise model.

Assumption 2. *For any feasible policy π , the Markov chain induced by policy π is irreducible and aperiodic.*

Clearly, this assumption is stronger than Assumption 1 since it indicates that the stationary distribution of any feasible policy is strictly positive.

Given the stochastic nature of the RL setting, a fundamental problem of our interest is *policy evaluation*, for estimating the long-run average reward/cost and the differential value function or

differential action-value function under a fixed policy. Equipped with policy evaluation tools, we can design efficient first-order algorithms for solving the policy optimization problem (2.5).

The rest of this paper is organized as follows. In Section 3, we propose efficient methods to solve the policy evaluation problem under both types of observation models. In Section 4, we develop the stochastic policy mirror descent (SPMD) method for solving AMDPs and establish the overall sample complexity of the actor-critic framework. In Section 5, we provide numerical experiments to corroborate our theoretical guarantees.

3 Critic: Policy evaluation of AMDPs

In this section, we discuss the policy evaluation of AMDPs. To serve as a fundamental building block of policy optimization, we focus on estimating the differential Q-function since it is closely related to the gradient information of the average reward/cost function. Moreover, the results and analyses extend to the estimation of the differential value function naturally. It should be noted that once the policy is fixed, h^π is a fixed function which together with $c(s, a)$ can be viewed as the overall cost function of the policy of interest. Therefore, for sack of simplicity, in this section we use $c(s, a)$ to represent $c(s, a) + h^\pi(s)$. And we let $|c(s, a)| \leq \bar{c}$, for all $s \in \mathcal{S}$ and $a \in \mathcal{A}$.

As illustrated before, we consider both the generative and Markovian noise models. In Section 3.1, we propose a direct multiple trajectory method (Algorithm 1) and establish its convergence guarantees. For the Markovian noise model, we start with policy evaluation of sufficiently random policies and propose the VRTD method (Algorithm 2) along with its convergence results in Section 3.2. Then, in Section 3.3, we design an EVRTD method to remedy the issue of exploration of insufficiently random policies.

3.1 Multiple trajectory method for generative model

Under a generative model, we assume that the AMDP is mixing (Assumption 1). To begin with, we state the following supporting lemma which characterizes the mixing property.

Lemma 1. *Given a policy π , for any integer $k \geq t_{mix}$, we have*

$$\|(P_\pi)^k - \mathbf{1}(\nu^\pi)^\top\|_\infty \leq (\frac{1}{2})^{\lfloor k/t_{mix} \rfloor}.$$

For the sack of completeness, we provide a proof of this lemma in Appendix B.1.

We are now in a position to formally state the multiple trajectory method for estimating both the average reward/cost ρ^π and the basic differential Q-function $\bar{Q}^\pi(s, a)$ for all the state-action pairs $(s, a) \in \mathcal{S} \times \mathcal{A}$.

In view of Algorithm 1, we generate trajectories starting from all state-action pairs to estimate the corresponding basic differential Q-function, which is allowed in the generative model. One should also note that we generate independent trajectories to estimate the average reward/cost and differential Q-function to avoid additional bias. The next proposition provides an ℓ_∞ -convergence guarantee on the bias of the estimators $\hat{\rho}^\pi$ and \hat{Q}^π generated by the multiple-trajectory method, i.e., $|\mathbb{E}[\hat{\rho}^\pi] - \rho^\pi|$ and $\|\mathbb{E}[\hat{Q}^\pi] - \bar{Q}^\pi\|_\infty$, where the expectation is taken with respect to all the samples used in Algorithm 1.

Algorithm 1 Multiple-trajectory method for generative model

1: **Input:** $T, T', M, M' \in \mathbb{Z}_+$ and a feasible policy π .
 2: **for** $i = 1, \dots, M$ **do**
 3: Generate a trajectory with length $T + 1$ following policy π starting from an arbitrary state $s' \in \mathcal{S}$, denoted by $\{(s_0 = s', a_0), (s_1, a_1), \dots, (s_T, a_T)\}$. Let $\rho^i = c(s_T, a_T)$.
 4: **end for**
 5: Calculate $\hat{\rho}^\pi = \frac{1}{M} \sum_{i=1}^M \rho^i$.
 6: **for** $j = 1, \dots, M'$ **do**
 7: **for** $(s, a) \in \mathcal{S} \times \mathcal{A}$ **do**
 8: Generate a trajectory with length $T' + 1$ following policy π starting from (s, a) , denoted by $\{(s_0^j = s, a_0^j = a), (s_1^j, a_1^j), \dots, (s_{T'}^j, a_{T'}^j)\}$. Let $\bar{Q}^j(s, a) = \sum_{t=0}^{T'} (c(s_t^j, a_t^j) - \hat{\rho}^\pi)$
 9: **end for**
 10: **end for**
 11: **Output:** $\hat{\rho}^\pi$ and $\hat{Q}^\pi(s, a) = \frac{1}{M'} \sum_{j=1}^{M'} \bar{Q}^j(s, a)$.

Proposition 1. Fix a feasible policy π . Assume that the epoch length parameters $T, T' > t_{\text{mix}} + 1$. Then the estimators $\hat{\rho}^\pi$ and \hat{Q}^π generated by Algorithm 1 satisfy

$$|\mathbb{E}[\hat{\rho}^\pi] - \rho^\pi| \leq \bar{c} \cdot \left(\frac{1}{2}\right)^{\lfloor T/t_{\text{mix}} \rfloor}, \quad (3.1)$$

and

$$\|\mathbb{E}[\hat{Q}^\pi] - \bar{Q}^\pi\|_\infty \leq \bar{c}(T' + 1) \cdot \left(\frac{1}{2}\right)^{\lfloor T/t_{\text{mix}} \rfloor} + 2\bar{c} \cdot t_{\text{mix}} \cdot \left(\frac{1}{2}\right)^{\lfloor T'/t_{\text{mix}} \rfloor}. \quad (3.2)$$

Proof. For simplicity of notation, we denote $\Delta\rho := \hat{\rho}^\pi - \rho^\pi$, $\Delta\rho^i := \rho^i - \rho^\pi$, thus we have

$$\Delta\rho = \frac{1}{M} \sum_{i=1}^M \Delta\rho^i.$$

We begin by establishing a bias bound for the estimator of the average reward/cost $|\mathbb{E}[\Delta\rho^i]|$,

$$|\mathbb{E}[\Delta\rho^i]| = |(P_\pi)_{s'}^T \cdot c^\pi - \rho^\pi| \leq \|(P_\pi)^T c^\pi - \mathbf{1}(\nu^\pi)^\top c^\pi\|_\infty \leq \|(P_\pi)^T - \mathbf{1}(\nu^\pi)^\top\|_\infty \|c^\pi\|_\infty \leq \bar{c} \cdot \left(\frac{1}{2}\right)^{\lfloor T/t_{\text{mix}} \rfloor}.$$

Consequently, by triangle inequality, we have

$$|\mathbb{E}[\Delta\rho]| \leq \bar{c} \cdot \left(\frac{1}{2}\right)^{\lfloor T/t_{\text{mix}} \rfloor}. \quad (3.3)$$

Now we are ready to bound the bias of the estimator of the basic differential Q-function. For simplicity of notation, we denote $\delta(s, a) := \frac{1}{M'} \sum_{j=1}^{M'} \delta^j(s, a)$, where

$$\delta^j(s, a) := \sum_{t=0}^{T'} \left(c(s_t^j, a_t^j) - \mathbb{E}_\pi[c(s_t, a_t) | s_0 = s, a_0 = a] \right), \quad \forall s \in \mathcal{S}, a \in \mathcal{A}.$$

Let the error induced by truncation be

$$\epsilon_{\text{trunc}}(s, a) := \sum_{t=T'+1}^{\infty} (\rho^\pi - \mathbb{E}_\pi[c(s_t, a_t) | s_0 = s, a_0 = a]), \quad \forall s \in \mathcal{S}, a \in \mathcal{A}.$$

Then by the definition of \hat{Q} and \bar{Q} , we have the following decomposition

$$\hat{Q}^\pi(s, a) - \bar{Q}^\pi(s, a) = \delta(s, a) + (T' + 1)\Delta\rho + \epsilon_{\text{trunc}}(s, a), \quad \forall s \in \mathcal{S}, a \in \mathcal{A}. \quad (3.4)$$

Noticing that $\mathbb{E}[\delta^j(s, a)] = 0$ because the trajectories are sampled following policy π , we have $\mathbb{E}[\delta(s, a)] = 0$. It remains to upper bound $|\mathbb{E}[\epsilon_{\text{trunc}}(s, a)]|$. For any $s \in \mathcal{S}, a \in \mathcal{A}$, we write

$$\begin{aligned} |\epsilon_{\text{trunc}}(s, a)| &= \left| \sum_{s' \in \mathcal{S}} \mathbf{P}(s'|s, a) \sum_{t=T'+1}^{\infty} [(\nu^\pi)^\top c^\pi - (P_\pi)_{s'}^{t-1} c^\pi] \right| \\ &\leq \left\| \sum_{t=T'+1}^{\infty} [\mathbf{1}(\nu^\pi)^\top - (P_\pi)^t] c^\pi \right\|_\infty \leq \sum_{t=T'+1}^{\infty} \|\mathbf{1}(\nu^\pi)^\top - (P_\pi)^t\|_\infty \|c^\pi\|_\infty \\ &\stackrel{(i)}{\leq} \bar{c} \cdot t_{\text{mix}} \cdot \sum_{i=\lfloor T'/t_{\text{mix}} \rfloor}^{\infty} (\frac{1}{2})^i \leq 2\bar{c} \cdot t_{\text{mix}} \cdot (\frac{1}{2})^{\lfloor T'/t_{\text{mix}} \rfloor}, \end{aligned} \quad (3.5)$$

where step (i) follows from Lemma 1. Combining Ineqs. (3.3), (3.4), (3.5), and utilizing triangle inequality, we obtain

$$|\mathbb{E}[\widehat{Q}^\pi(s, a) - \bar{Q}^\pi(s, a)]| \leq \bar{c}(T' + 1) \cdot (\frac{1}{2})^{\lfloor T/t_{\text{mix}} \rfloor} + 2\bar{c} \cdot t_{\text{mix}} \cdot (\frac{1}{2})^{\lfloor T'/t_{\text{mix}} \rfloor}.$$

Noticing that the RHS of the above inequality does not depend on (s, a) , so this is a uniform bound for all the entries of $\mathbb{E}[\widehat{Q}^\pi - \bar{Q}^\pi]$ and we complete the proof. \square

In view of Proposition 1, the number of samples (state-action pairs) required by the multiple trajectory method to find a solution $\widehat{Q}^\pi \in \mathbb{R}^{|\mathcal{S} \times \mathcal{A}|}$ such that $\|\mathbb{E}[\widehat{Q}^\pi] - \bar{Q}^\pi\|_\infty \leq \epsilon$ is bounded on the order $\mathcal{O}\{|\mathcal{S}||\mathcal{A}|t_{\text{mix}} \cdot \log(t_{\text{mix}}/\epsilon)\}$. The next proposition establishes the convergence guarantee of Algorithm 1 in terms of the expected squared ℓ_∞ -error.

Proposition 2. *Fix a feasible policy π . Assume that the parameters $T, T' > t_{\text{mix}} + 1$ and $M \geq M'$. Then the estimators $\widehat{\rho}^\pi$ and \widehat{Q}^π generated by Algorithm 1 satisfy that*

$$\mathbb{E}|\widehat{\rho}^\pi - \rho^\pi|^2 \leq \frac{4\bar{c}^2}{M} + \bar{c}^2 \cdot (\frac{1}{2})^{2\lfloor T/t_{\text{mix}} \rfloor},$$

and

$$\mathbb{E}[\|\widehat{Q}^\pi - \bar{Q}^\pi\|_\infty^2] \leq \frac{v \cdot \bar{c}^2 (T'+1)^2}{M'} (\log(|\mathcal{S}||\mathcal{A}|) + 1) + \bar{c}^2 (3(T' + 1)^2 + 12t_{\text{mix}}^2) \cdot (\frac{1}{2})^{2\lfloor T'/t_{\text{mix}} \rfloor}, \quad (3.6)$$

where $v > 0$ denotes a universal constant.

Proof. First, focus our attention on providing an upper bound on $\mathbb{E}[(\Delta\rho)^2]$. We write

$$\mathbb{E}[(\Delta\rho)^2] = \text{Var}(\Delta\rho) + (\mathbb{E}[\Delta\rho])^2 \stackrel{(i)}{\leq} \frac{4\bar{c}^2}{M} + \bar{c}^2 \cdot (\frac{1}{2})^{2\lfloor T/t_{\text{mix}} \rfloor}, \quad (3.7)$$

where step (i) follows from that each ρ^i is independent of each other and Ineq. (3.1) of Proposition 1. Recalling the decomposition in Eq. (3.4) and utilizing Young's inequality, we have

$$\mathbb{E}[\|\widehat{Q}^\pi - \bar{Q}^\pi\|_\infty^2] \leq 3\mathbb{E}[\|\delta\|_\infty^2] + 3(T' + 1)^2 \mathbb{E}[(\Delta\rho)^2] + 3\|\epsilon_{\text{trunc}}\|_\infty^2, \quad (3.8)$$

where the last term on the RHS does not require an expectation since it does not depend on the samples generated when running Algorithm 1. Invoking an uniform upper bound on ϵ_{trunc} in Ineq. (3.5), it remains to bound the term $\mathbb{E}[\|\delta\|_\infty^2]$. First, we focus on an (s, a) pair, for $j = 1, \dots, M'$,

$$\mathbb{E}[\delta^j(s, a)] = 0, \quad \text{and} \quad |\delta^j(s, a)| \leq 2(T' + 1)\bar{c}.$$

Since each $\delta^j(s, a)$ is bounded and independent with each other, we have that $\delta(s, a) = \frac{1}{M'} \sum_{j=1}^{M'} \delta^j(s, a)$ is a sub-gaussian variable, and there exists an absolute constant $u > 0$ such that $\|\delta(s, a)\|_{\psi_2} \leq \frac{u\bar{c}(T'+1)}{\sqrt{M'}}$. As a result, $\delta(s, a)^2$ is sub-exponential and

$$\|\delta(s, a)^2\|_{\psi_1} = \|\delta(s, a)\|_{\psi_2}^2 \leq \frac{u^2 \bar{c}^2 (T'+1)^2}{M'}.$$

On the other hand, given that $\delta^j(s, a)$ are i.i.d. and zero-mean, we have

$$\mathbb{E}[\delta(s, a)^2] \leq \frac{4\bar{c}^2 (T'+1)^2}{M'}.$$

Then utilizing Lemma 25 of Lan (2022), we have that there exists a universal constant $v > 0$, such that

$$\mathbb{E}[\|\delta\|_\infty^2] = \mathbb{E}[\max_{s \in \mathcal{S}, a \in \mathcal{A}} \delta(s, a)^2] \leq \frac{v\bar{c}^2 (T'+1)^2}{M'} (\log(|\mathcal{S}||\mathcal{A}|) + 1) + \frac{4\bar{c}^2 (T'+1)^2}{M'}. \quad (3.9)$$

Substituting Ineqs. (3.9), (3.7) and (3.5) into Ineq. (3.8) and invoking $M \geq M'$, we obtain

$$\mathbb{E}[\|\hat{Q}^\pi - \bar{Q}^\pi\|_\infty^2] \leq \frac{3v\bar{c}^2 (T'+1)^2}{M'} (\log(|\mathcal{S}||\mathcal{A}|) + 1) + \frac{24\bar{c}^2 (T'+1)^2}{M'} + (3(T'+1)^2 + 12t_{\text{mix}}^2) \bar{c}^2 \cdot \left(\frac{1}{2}\right)^{2\lfloor T'/t_{\text{mix}} \rfloor},$$

which completes the proof. \square

For a fixed policy π , the sample complexity to find an ϵ -accurate estimator $\hat{Q}^\pi \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$ such that $\mathbb{E}[\|\hat{Q}^\pi - \bar{Q}^\pi\|_\infty^2] \leq \epsilon$ requires the parameters M and M' to be bounded by $\tilde{\mathcal{O}}(t_{\text{mix}}^2/\epsilon)$. The total number of samples used is $|\mathcal{S}||\mathcal{A}|M'T' + MT$, which is of the order $\tilde{\mathcal{O}}(|\mathcal{S}||\mathcal{A}|t_{\text{mix}}^3/\epsilon)$.

Observe from Proposition 1 and 2 that the bias of the estimator converges much faster than the expected error. This observation indicates that a careful analysis of “actor”, i.e., splitting the impact of policy evaluation bias and absolute error, may lead to improved overall sample complexity in policy optimization. The technical details of splitting the error terms will be shown in Section 4.

3.2 Evaluation of sufficiently random policies in Markovian noise model

In the Markovian noise model, we assume that Assumption 2 (ergodicity assumption) holds to ensure sample access to all the states. As a consequence, we have the following lemma which characterizes the mixing time property under Assumption 2 (see Bertsekas and Tsitsiklis, 1995).

Lemma 2. *There exist constants $C_P > 0$ and $\varrho \in (0, 1)$ such that for any policy π , we have $\nu^\pi > 0$ and $\max_{s \in \mathcal{S}} \|(P_\pi)_{(s, \cdot)}^t - \nu^\pi\|_1 \leq C_P \cdot \varrho^t$, for all $t \in \mathbb{Z}_+$.*

Remark 1. *Assumption 2 and Lemma 2 guarantee that the mixing time defined in Assumption 1 is bounded as $t_{\text{mix}} \leq \mathcal{O}\left(\frac{\log(C_P)}{\log(1/\varrho)}\right)$.*

For a given policy π , after combining the cost function and the regularization function, we can rewrite the Bellman equation (2.4) as

$$Q = P^\pi Q + c - \rho^\pi \mathbf{1}. \quad (3.10)$$

When analyzing the sampling scheme of the Markovian noise model, we define a natural weighting diagonal matrix $D^\pi := \text{diag}([D_1^\pi; \dots; D_{|S|}^\pi]) \in \mathbb{R}^{(|S| \times |\mathcal{A}|) \times (|S| \times |\mathcal{A}|)}$, where $D_s^\pi := \nu^\pi(s) \cdot \text{diag}(\pi(\cdot|s)) \in$

$\mathbb{R}^{|\mathcal{A}| \times |\mathcal{A}|}$. Clearly, the diagonal elements of D^π are the steady-state distribution of all state-action pairs induced by the policy π . For algorithmic design and analysis, we desire the matrix D^π to be positive definite. However, the matrix D^π can be ill-conditioned or even not positive definite when there exists $s \in \mathcal{S}, a \in \mathcal{A}$ such that $\pi(a|s)$ is close or equal to 0. Towards this end, we first restrict our attention to sufficiently random policy π , and defer the discussion of insufficiently random policy π to Section 3.3. Specifically, we assume there exists some $\underline{\pi} > 0$ such that $\pi(s|a) \geq \underline{\pi}$ for all $s \in \mathcal{S}, a \in \mathcal{A}$, thus $D^\pi \succ 0$. For brevity of notation, we let $P := P^\pi, D := D^\pi, \nu := \nu^\pi$ and $\rho^* := \rho^\pi$.

For large state and action spaces, it is common to seek a low-dimensional approximation to the differential Q-function by implementing linear function approximation. In particular, one chooses $\mathbb{S} := \text{span}\{\psi_1, \dots, \psi_d\}$ for d linearly independent basis vectors $\psi_1, \dots, \psi_d \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$. For each state action pair $s \in \mathcal{S}, a \in \mathcal{A}$, we let $\psi(s, a) := [\psi_1(s, a), \psi_2(s, a), \dots, \psi_d(s, a)]^\top$ denote its feature vector. Additionally, we assume that $\|\psi(s, a)\|_2 \leq 1$ for each state action pair, which can be ensured through feature normalization. Letting $\Pi_{\mathbb{S}, D}$ denote the projection onto the subspace \mathbb{S} with respect to the $\|\cdot\|_D$ -norm, then one can instead solve the following projected Bellman equation:

$$\bar{Q} = \Pi_{\mathbb{S}, D}(P\bar{Q} + c - \rho^* \mathbf{1}).$$

Let $\Psi = [\psi_1, \dots, \psi_d]$, the problem above can be equivalent written as solving for $\theta^* \in \mathbb{R}^d$ satisfying

$$\Psi^\top D \Psi \theta^* = \Psi^\top D P \Psi \theta^* + \Psi^\top D(c - \rho^* \mathbf{1}). \quad (3.11)$$

Next, we define a set of vectors in $\mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$ that have identical value on the support of π , i.e.,

$$\mathcal{I} := \{x \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|} | x(s, a) = 1 \text{ if } \pi(a|s) > 0\}.$$

For the sack of simplicity, we assume that $\mathcal{I} \cap \mathbb{S} = \emptyset$, which is equivalent to $\mathbf{1} \notin \mathbb{S}$ in this section since π is strictly positive. This condition can be easily satisfied by choosing the features appropriately. Then by Theorem 1 of Tsitsiklis and Van Roy (1999), Eq. (3.11) has a unique solution. Furthermore, we define

$$\bar{Q}^* = \bar{Q}^\pi + \hat{b} \mathbf{1} \quad \text{where} \quad \hat{b} := \arg \min_{b \in \mathbb{R}} \{\|\bar{Q}^\pi + b \mathbf{1} - \Psi \theta^*\|_D\}. \quad (3.12)$$

In words, \bar{Q}^* is the projection of $\Psi \theta^*$ onto the set of the differential Q-functions defined in Eq. (2.4).

It is convenient in the analysis to have access to an orthonormal basis spanning the projected space \mathbb{S} . Define the matrix $B \in \mathbb{R}^{d \times d}$ as $B_{i,j} := \langle \psi_i, \psi_j \rangle_D$ for each i, j . Note that $B \succ 0$ due to the linear independence of ψ_i 's and $D \succ 0$. Let $\Phi := [\phi_1, \phi_2, \dots, \phi_d] = \Psi B^{-\frac{1}{2}}$. By construction, the vectors ϕ_1, \dots, ϕ_d satisfy $\langle \phi_i, \phi_j \rangle_D = \mathbb{I}(i = j)$. We further denote

$$\mu := \lambda_{\min}(B) \quad \text{and} \quad M := \Phi D P \Phi^\top. \quad (3.13)$$

Notice that M is a d -dimensional matrix that describes the action of P on the projected space \mathbb{S} .

To solve the projected Bellman equation (3.28), we define the deterministic operator and the corresponding stochastic operator calculated from sample $\xi = \{(s, a), (s', a'), c(s, a)\}$ as

$$g(\theta, \rho) = \Psi^\top D(\Psi \theta - P \Psi \theta - c + \rho \mathbf{1}), \quad (3.14)$$

$$\tilde{g}(\theta, \rho, \xi) = (\langle \psi(s, a) - \psi(s', a'), \theta \rangle - c(s, a) + \rho) \psi(s, a). \quad (3.15)$$

The following lemma characterizes the strong monotonicity property of the operator $g(\theta, \rho)$.

Lemma 3. Under Assumption 2, assume $\pi(a|s) > 0$ for all $s \in \mathcal{S}, a \in \mathcal{A}$, then

$$1 - \beta := \inf_{x \in \mathbb{S}, \|x\|_D=1} x^\top D(I - P)x > 0, \quad (3.16)$$

As a consequence, for any $\theta, \theta' \in \mathbb{R}^d$ and $\rho \in \mathbb{R}$,

$$\langle g(\theta, \rho) - g(\theta', \rho), \theta - \theta' \rangle \geq (1 - \beta) \|\Psi\theta - \Psi\theta'\|_D^2. \quad (3.17)$$

Proof. For any $x \in \mathbb{S}, \|x\|_D = 1$, let the expectation be taken on $s \sim \nu, a \sim \pi(\cdot|s), s' \sim P(\cdot|s, a), a' \sim \pi(\cdot|s')$, then

$$x^\top D(I - P)x = \mathbb{E}[x(s, a)^2] - \mathbb{E}[x(s, a)x(s', a')] \stackrel{(i)}{=} \frac{1}{2} \mathbb{E}[(x(s, a) - x(s', a'))^2] \stackrel{(ii)}{>} 0, \quad (3.18)$$

where step (i) follows from the fact that (s, a) and (s', a') have the same marginal distribution, and step (ii) follows from the fact that $\mathbf{1} \notin \mathbb{S}$ and the Markov chain is irreducible. Meanwhile, given that this feasible region is compact and the function of interest is continuous, by the extreme value theorem, we obtain that $1 - \beta > 0$, which completes the proof. \square

It should be noted that the term $\frac{1}{1-\beta}$, which replaces the role of the effective horizon in the discounted MDPs, is strongly related to the mixing time of the induced Markov chain. It can be shown that if the linear subspace \mathbb{S} is properly chosen, i.e., $\langle \phi_i, \mathbf{1} \rangle_D$ is not extremely close to 1 for all $i \in [d]$, then $\frac{1}{1-\beta}$ can be bounded by $\mathcal{O}(t_{\text{mix}})$ (see Corollary 1.14 of Montenegro et al., 2006).

On the other hand, the data generated by the Markovian noise model is highly correlated, which induces challenges in the algorithmic analysis. To overcome the difficulties, we incorporate the idea of sample skipping from Kotsalis et al. (2022b); Li et al. (2021) to reduce the bias induced by correlated data. The spirit of this sample skipping idea can be traced back to the classical work by Yu (1994). To quantify how the sample skipping works in reducing the bias of the stochastic operator, we establish the following lemmas.

Lemma 4. Let $\xi_t := \{(s_t, a_t), (s_{t+1}, a_{t+1}), c(s_t, a_t)\}$. For every $t, \tau \in \mathbb{Z}_+$, with probability 1,

$$\|\mathbb{E}[\tilde{g}(\theta^*, \rho^*, \xi_{t+\tau+1})|\mathcal{F}_t] - g(\theta^*, \rho^*)\|_2 \leq C_M \cdot \varrho^\tau \|\Psi\theta^* - \bar{Q}^*\|_D. \quad (3.19)$$

where $C_M := \frac{C_P}{\sqrt{\min_{s \in \mathcal{S}, a \in \mathcal{A}} \nu(s) \pi(a|s)}} \|\Psi\|_2 \|I - P\|_2$ and $\mathcal{F}_t := [\xi_1, \dots, \xi_t]$.

Proof. Let $D_t^{\tau+1} := \text{diag}([P(s_{t+\tau+1} = s, a_{t+\tau+1} = a | s_t, a_t)])$ where $s \in \mathcal{S}, a \in \mathcal{A}$. We have

$$\mathbb{E}[\tilde{g}(\theta^*, \rho^*, \xi_{t+\tau+1})|\mathcal{F}_t] = \Psi D_t^{\tau+1} (\Psi\theta^* - P\Psi\theta^* - c + \rho^* \mathbf{1}).$$

Combining the above equality with the definition of $g(\cdot)$ in (3.14) yields

$$\begin{aligned} \|g(\theta^*, \rho^*) - \mathbb{E}[\tilde{g}(\theta^*, \rho^*, \xi_{t+\tau+1})|\mathcal{F}_t]\|_2 &= \|\Psi^\top (D - D_t^{\tau+1}) (\Psi\theta^* - P\Psi\theta^* - c + \rho^* \mathbf{1})\|_2 \\ &= \|\Psi^\top (D - D_t^{\tau+1}) (I - P) (\Psi\theta^* - \bar{Q}^*)\|_2 \\ &\leq C_P \varrho^\tau \|\Psi\|_2 \|I - P\|_2 \|\Psi\theta^* - \bar{Q}^*\|_2 \leq C_M \varrho^\tau \|\Psi\theta^* - \bar{Q}^*\|_D, \end{aligned}$$

as claimed. \square

In view of Lemma 4, the bias of the stochastic operator at the optimal solution depends on the approximation error caused by linear function approximation, and the bound decays geometrically.

Lemma 5. For every $t, \tau \in \mathbb{Z}_+$, $\theta, \theta' \in \mathbb{R}^d$ and $\rho, \rho' \in \mathbb{R}$, with probability 1,

$$\begin{aligned} & \|\mathbb{E}[\tilde{g}(\theta, \rho, \xi_{t+\tau+1})|\mathcal{F}_t] - \mathbb{E}[\tilde{g}(\theta', \rho', \xi_{t+\tau+1})|\mathcal{F}_t] - [g(\theta, \rho) - g(\theta', \rho')]\|_2 \\ & \leq C_M \cdot \varrho^\tau \|\Psi\theta - \Psi\theta'\|_D + C_R \cdot \varrho^\tau |\rho - \rho'|, \end{aligned} \quad (3.20)$$

where $C_R := C_P \|\Psi\|_2$.

Proof. We write

$$\begin{aligned} & \|g(\theta, \rho) - \mathbb{E}[\tilde{g}(\theta, \rho, \xi_{t+\tau+1})|\mathcal{F}_t] - g(\theta', \rho') + \mathbb{E}[\tilde{g}(\theta', \rho', \xi_{t+\tau+1})|\mathcal{F}_t]\|_2 \\ & = \|\Psi^\top (D - D_t^{\tau+1}) ((I - P)\Psi(\theta - \theta') + (\rho - \rho'))\|_2 \\ & \leq C_P \varrho^\tau \|\Psi\|_2 \|I - P\|_2 \|\Psi(\theta - \theta')\|_2 + C_P \varrho^\tau \|\Psi\|_2 |\rho - \rho'| \leq C_M \cdot \varrho^\tau \|\Psi\theta - \Psi\theta'\|_D + C_R \cdot \varrho^\tau |\rho - \rho'|, \end{aligned}$$

as claimed. \square

In contrast to Lemma 4, Lemma 5 captures the bias reduction of the difference between two stochastic operators calculated using the same samples, which also decays in a geometric rate and does not depend on the approximation error. For brevity of notation, we set $C_{\max} := \max\{C_P, C_M, C_R\}$. Now we are ready to formally state the VRTD algorithm for average-reward policy evaluation under Markovian noise in Algorithm 2.

Algorithm 2 Variance-reduced Temporal Difference (VRTD) Method for AMDPs

- 1: **Input:** $\hat{\theta}_0 \in \mathbb{R}^d$, $\eta > 0$, $\tau, \tau' \in \mathbb{Z}_+$, $\{N_k\}_{k=1}^K \subset \mathbb{Z}_+$ and $\{N'_k\}_{k=1}^K \subset \mathbb{Z}_+$.
 - 2: **for** $k = 1, \dots, K$ **do**
 - 3: Set $\theta_1 = \tilde{\theta} = \hat{\theta}_{k-1}$.
 - 4: Collect N'_k state-action samples $\{(\tilde{s}_i, \tilde{a}_i)\}_{i=1}^{N'_k}$ from the single Markov trajectory induced by policy π , where each sample is the last one of $\tau' + 1$ successive state-action samples.
 Calculate $\tilde{\rho} = \frac{1}{N'_k} \sum_{i=1}^{N'_k} c(\tilde{s}_i, \tilde{a}_i)$.
 - 5: Collect N_k samples $\xi_i^k(\tau') = \{(s_i, a_i), (s'_i, a'_i), c(s_i, a_i)\}$ from the single Markov trajectory induced by policy π , where each sample is the last one of $\tau' + 1$ successive samples.
 Calculate $\hat{g}(\tilde{\theta}, \tilde{\rho}) = \frac{1}{N_k} \sum_{i=1}^{N_k} \tilde{g}(\tilde{\theta}, \tilde{\rho}, \xi_i^k(\tau'))$.
 - 6: **for** $t = 1, \dots, T$ **do**
 - 7: Collect a sample $\xi_t(\tau) = \{(s_t, a_t), (s'_t, a'_t), c(s_t, a_t)\}$ from the single Markov trajectory, which is the last one of $\tau + 1$ successive samples.
 - 8:
$$\theta_{t+1} = \theta_t - \eta \left(\tilde{g}(\theta_t, \tilde{\rho}, \xi_t(\tau)) - \tilde{g}(\tilde{\theta}, \tilde{\rho}, \xi_t(\tau)) + \hat{g}(\tilde{\theta}, \tilde{\rho}) \right). \quad (3.21)$$
 - 9: **end for**
 - 10: Output of the epoch:

$$\hat{\theta}_k = \frac{\sum_{t=1}^T \theta_t}{T}, \quad \text{and} \quad \hat{\rho}_k = \tilde{\rho}. \quad (3.22)$$
 - 11: **end for**
-

We introduce the basic idea of Algorithm 2 below. First, it incorporates the idea of the variance-reduced TD learning for solving DMDPs from Li et al. (2021). In particular, this algorithm runs in epochs and utilizes the current best estimator $\tilde{\theta}$ to recenter the updates inside each epoch.

This variance reduction technique allows us to capture the correct stochastic error of the problem. Second, unlike the DMDP setting, policy evaluation of the AMDPs has two objectives, namely ρ^* for average reward/cost and θ^* for parametrized differential Q-function. However, we should expect that estimating ρ^* requires much less sample than estimating θ^* , because the former task is estimating the steady-state behavior of a policy, while the latter one requires solving a linear equation with only sample access to the model. Towards this end, the VRTD algorithm first collects samples and calculates an estimator of ρ^* at the beginning of each epoch and then performs (variance-reduced) stochastic approximation to update the iterates of θ . By utilizing this updating rule, we are able to significantly improve the sample complexity of the TD algorithms that update the two objectives simultaneously introduced by [Tsitsiklis and Van Roy \(1999\)](#) and analyzed by [Zhang et al. \(2021b\)](#). We will provide detailed comparisons of sample complexities after presenting the main convergence results of Algorithm 2.

The following theorem characterizes the convergence of VRTD in terms of expected squared ℓ_D -norm error, which is a natural convergence criterion (see, e.g., [Bhandari et al. \(2018\)](#); [Mou et al. \(2020\)](#); [Li et al. \(2021\)](#) for DMDPs) for policy evaluation in the RL literature.

Theorem 1. *Fix the total number of epochs K and positive integers N and N' . Assume for each epoch $k \in [K]$, the parameters η , T , N_k and N'_k satisfy*

$$\eta \leq \frac{1-\gamma}{845}, \quad T \geq \frac{64}{\mu(1-\beta)\eta}, \quad N_k \geq \max \left\{ \frac{1160}{\mu(1-\beta)^2}, \left(\frac{3}{4}\right)^{K-k} N \right\}, \quad \text{and} \quad N'_k \geq \left(\frac{3}{4}\right)^{K-k} N'. \quad (3.23)$$

Set the output of epoch k to be $\hat{\theta}_k := \frac{\sum_{t=1}^T \theta_t}{T}$. If τ and τ' are chosen to satisfy

$$\varrho^\tau \leq \frac{\sqrt{\mu}\eta}{C_M} \quad \text{and} \quad \varrho^{\tau'} \leq \min \left\{ \frac{\min_{s \in \mathcal{S}, a \in \mathcal{A}} \nu(s) \pi(a|s)}{C_P}, \frac{1}{4C_{\max}+1} \right\}, \quad (3.24)$$

then we have $\mathbb{E}[(\hat{\rho}_K - \rho^*)^2] \leq \frac{5\bar{c}^2}{N'}$, and

$$\mathbb{E}\|\Psi\hat{\theta}_K - \Psi\theta^*\|_D^2 \leq \frac{1}{2K}\|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \frac{3W_1}{N} + \frac{108\bar{c}^2}{N'(1-\beta)^2\mu} + \frac{3\bar{c}^2}{4N'}, \quad (3.25)$$

where $W_1 := 22\text{trace}\{(I_d - M)^{-1}\bar{\Sigma}(I_d - M)^{-\top}\} + \frac{4}{\mu(1-\beta)^2}\mathbb{E}[\|(\bar{Q}^* - \Psi\theta^*)\|_D^2]$ and $\bar{\Sigma} := \text{cov}[B^{-\frac{1}{2}}(\langle \psi(s, a) - \gamma\psi(s', a'), \theta^* \rangle - c(s, a))\psi(s, a)]$, for $s \sim \nu, a \sim \pi(\cdot|s), s' \sim P(\cdot|s, a), a' \sim \pi(\cdot|s')$.

See Appendix B.3 for the detailed proof of this theorem. A few comments on the upper bounds provided in Theorem 1 are in order.

Sample complexity: Noting that the error caused by inexact feature approximation is unavoidable, we set the stopping criterion as finding a solution $\hat{\theta} \in \mathbb{R}^d$ to achieve the accuracy $\mathbb{E}\|\Psi\hat{\theta} - \Psi\theta^*\|_D^2 \leq \mathcal{O}(1)\|\Psi\theta^* - \bar{Q}^*\|_D^2 + \epsilon$. From the statement of Theorem 1, we conclude that the number of required epochs is bounded by $\mathcal{O}\{\log(\|\Psi\theta^0 - \Psi\theta^*\|_D^2/\epsilon)\}$ and the total number of samples is $\sum_{k=1}^K (\tau T + \tau'(N_k + N'_k))$, which is of the order

$$\mathcal{O}\left\{ \frac{t_{\text{mix}}}{(1-\beta)^2\mu} \log\left(\frac{1}{\epsilon}\right) + \frac{t_{\text{mix}}\bar{c}^2}{(1-\beta)^2\mu\epsilon} + \underbrace{\frac{t_{\text{mix}} \cdot \text{trace}\{(I_d - M)^{-1}\bar{\Sigma}(I_d - M)^{-\top}\}}{\epsilon}}_{\text{dominant stochastic error}} \right\}.$$

We add a few comments on the dominant term of the above complexity. It is an instance-dependent characterization of the stochastic error for solving the projected Bellman equation (3.11) with sample access to the transition kernel. A direct extension of Proposition 1 of [Li et al. \(2021\)](#)

demonstrates that this term matches the asymptotic instance-dependent lower bound on stochastic error of solving Eq. (3.11) with i.i.d. samples. We pick up an additional mixing time factor in our complexity result because of the Markovian noise. For fair comparison to the existing literature, we apply a direct upper bound, $\mathcal{O}\{\frac{\|\theta^*\|_2^2}{(1-\beta)^2\mu}\}$, on $\text{trace}\{(I_d - M)^{-1}\bar{\Sigma}(I_d - M)^{-\top}\}$, as well as the upper bound $\mathcal{O}(t_{\text{mix}})$ on $\frac{1}{1-\beta}$, thus arriving at the sample complexity $\mathcal{O}\{\frac{t_{\text{mix}}^3}{\mu} \log(\frac{1}{\epsilon}) + \frac{t_{\text{mix}}^3(\bar{c}^2 + \|\theta^*\|_2^2)}{\mu\epsilon}\}$.

Comparisons with related work: To the best of our knowledge, the only work that established finite sample analysis for solving the AMDP policy evaluation problem under Markovian noise is Zhang et al. (2021b). They analyzed the TD(λ) method proposed by Tsitsiklis and Van Roy (1999), where the algorithm simultaneously updates estimators of both evaluation objectives, i.e., average reward/cost and differential value function, thus worsening their convergence rate. Compared to their guarantees, our sample complexity outperforms by a factor of $\mathcal{O}(\frac{1}{(1-\beta)^2})$ or equivalently $\mathcal{O}(t_{\text{mix}}^2)$. In addition, they establish convergence analysis in ℓ_2 -norm, which is not a natural metric for the underlying problem, thus leading to worse dependence on other problem parameters, i.e., dimension of the transition kernel. It is noteworthy that recently Mou et al. (2022) proposed a variance-reduced stochastic approximation approach that solves the AMDP policy evaluation problem in span semi-norm under the generative model. However, their results do not directly extend to the Markovian noise setting. Meanwhile, it is hard to obtain convergence guarantees of bias by using their metric and analysis techniques.

Next, we switch our attention to the bias bound, which is characterized in the following theorem.

Theorem 2. Fix the total number of epochs K and positive integers N and N' . Assume for each epoch $k \in [K]$, the parameters η , N , N' and T satisfy conditions (3.23) and

$$N_k = N \geq \frac{1160}{\mu(1-\beta)^2}, \quad N'_k = N' \geq \frac{1}{\mu(1-\beta)^2}. \quad (3.26)$$

Set the output of epoch k to be $\hat{\theta}_k := \frac{\sum_{t=1}^T \theta_t}{T}$. Then we have

$$\|\Psi\mathbb{E}[\hat{\theta}_K] - \Psi\theta^*\|_D^2 \leq \frac{1}{2K}\|\Psi\theta^0 - \Psi\theta^*\|_D^2 + (b_1 C_{\max}^2 \varrho^{2\tau'} + \frac{b_2 C_{\max} \varrho^\tau}{(1-\beta)\sqrt{\mu}})(\|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \frac{W_1}{N} + \bar{c}^2), \quad (3.27)$$

where b_1 and b_2 are universal constants.

See Appendix B.4 for the detailed proof of this theorem.

A few comments of the statement of Theorem 2 are in order. First, for finding a solution $\hat{\theta} \in \mathbb{R}^d$ to achieve the accuracy $\|\Psi\mathbb{E}[\hat{\theta}] - \Psi\theta^*\|_D^2 \leq \mathcal{O}(1)\|\Psi\theta^* - \bar{Q}^*\|_D^2 + \epsilon$, we require the parameters τ and τ' to satisfy $\tau, \tau' \geq \mathcal{O}\{t_{\text{mix}} \log(1/\epsilon)\}$, and the total sample complexity to be

$$\mathcal{O}\{\frac{t_{\text{mix}}}{(1-\beta)^2\mu} \log^2(\frac{1}{\epsilon})\}, \quad \text{or equivalently} \quad \mathcal{O}\{\frac{t_{\text{mix}}^3}{\mu} \log^2(\frac{1}{\epsilon})\}.$$

It should be noted that this result naturally extends to policy evaluation of DMDPs under Markovian noise. Towards this end, we are able to improve the $\mathcal{O}(1/T^3)$ convergence rate in Lemma 18 of Lan (2022) to linear convergence, consequently improving the sample complexity of the PMD method under Markovian noise (Proposition 4 and 5 of Lan, 2022). To the best of our knowledge, this geometric convergence rate on the bias of policy evaluation is new to both AMDP and DMDP literature. This result will be used in Section 4 to obtain sharp sample complexities of policy optimization.

3.3 Evaluation of insufficiently random policies in Markovian noise model

In this section, we remove the assumption of sufficiently random policies, i.e., we fix a policy π where $\pi(a|s)$ can be 0 or arbitrarily small for some state-action pairs. This setting induces a possible lack of exploration and makes the on-policy evaluation task more challenging. A classical method to remedy this issue is off-policy learning, where the agent chooses actions according to a behavior policy that encourages exploration. However, in this manner, the agent is not able to collect rewards according to the policy of interest, thus losing the advantages of on-policy reinforcement learning. Towards this end, in this section, we seek an on-policy method to remedies this exploration issue.

The begin with, we first propose an exploratory variance-reduced temporal difference (EVRTD) algorithm that alternates between the given policy π and a slightly perturbed policy, denoted as $\tilde{\pi}$. Again, for brevity of notation, we set $P := P^\pi$, $\nu := \nu^\pi$ and $\rho^* := \rho^\pi$.

The EVRTD algorithm for insufficiently random policy π has similar structure to Algorithm 2, with difference only lying in the collection of samples. Specifically, the EVRTD algorithm replaces steps 5 and 7 in Algorithm 2 with the following ones.

step 5': Run the Markov trajectory induced by policy π for τ' steps and obtain the current state s . Then generate an action $a \sim \tilde{\pi}(\cdot|s)$, the next state $s' \sim P(\cdot|s, a)$, the next action $a' \sim \pi(\cdot|s')$, and collect the tuple $\{(s, a), (s', a'), c(s, a)\}$. Repeat for N_k times, and index the samples as $\xi_i^k(\tau') = \{(s_i, a_i), (s'_i, a'_i), c(s_i, a_i)\}$. Calculate $\hat{g}(\tilde{\theta}, \tilde{\rho}) = \frac{1}{N_k} \sum_{i=1}^{N_k} \tilde{g}(\tilde{\theta}, \tilde{\rho}, \xi_i^k(\tau'))$.

step 7': Run the Markov trajectory induced by policy π for τ steps and obtain the current state s_t . Then generate an action $a_t \sim \tilde{\pi}(\cdot|s_t)$, the next state $s'_t \sim P(\cdot|s_t, a_t)$, the next action $a'_t \sim \pi(\cdot|s'_t)$, and collect the tuple $\{(s_t, a_t), (s'_t, a'_t), c(s_t, a_t)\}$.

Clearly, the EVRTD algorithm is still an on-policy algorithm since in most of the time the agent collects rewards/costs following the policy of interest. We make a few comments on the intrinsic idea of EVRTD algorithm by focusing on one sample tuple $\{(s, a), (s', a'), c(s, a)\}$. Clearly, as τ or τ' grows larger, the distribution of the state s converges to the stationary distribution ν of the original Markov chain induced by policy π . Invoking that $a \sim \tilde{\pi}(\cdot|s)$, $s' \sim P(\cdot|s, a)$ and $a' \sim \pi(\cdot|s')$, the projected fixed point equation we are trying to solve is

$$\Psi^\top \tilde{D} \Psi \theta^* = \Psi^\top \tilde{D} P \Psi \theta^* + \Psi^\top \tilde{D} (c - \rho^* \mathbf{1}), \quad (3.28)$$

where \tilde{D} is defined as $\tilde{D} := \text{diag}([\tilde{D}_1; \dots; \tilde{D}_{|S|}]) \in \mathbb{R}^{(|S| \times |\mathcal{A}|) \times (|S| \times |\mathcal{A}|)}$, and $\tilde{D}_s := \nu^\pi(s) \cdot \text{diag}(\tilde{\pi}(\cdot|s))$. Then Eq. (3.28) can be equivalently written as

$$\bar{Q} = \Pi_{\mathcal{S}, \tilde{D}} (P \bar{Q} + c - \rho^* \mathbf{1}).$$

It is easy to see that instead of solving the Bellman equation induced by a perturbed policy, we are still solving a projected version of the average-reward Bellman equation (3.10) induced by the original policy π . Furthermore, we have $\tilde{D} \succ 0$ when the underlying Markov chain is ergodic and the policy $\tilde{\pi} > 0$. With a slight abuse of notation, in this subsection we define the matrix $B \in \mathbb{R}^{d \times d}$ by letting $B_{i,j} := \langle \psi_i, \psi_j \rangle_{\tilde{D}}$ for each i, j , and let

$$\Phi := [\phi_1, \phi_2, \dots, \phi_d] = \Psi B^{-\frac{1}{2}}, \quad \mu := \lambda_{\min}(B), \quad \tilde{M} := \Phi \tilde{D} P \Phi^\top,$$

and $\tilde{\Sigma} := \text{cov} [B^{-\frac{1}{2}} (\langle \psi(s, a) - \gamma \psi(s', a'), \theta^* \rangle - c(s, a)) \psi(s, a)]$, for $s \sim \nu$, $a \sim \tilde{\pi}(\cdot|s)$, $s' \sim P(\cdot|s, a)$ and $a' \sim \pi(\cdot|s')$. We modify the definition of the deterministic operator correspondingly

$$g(\theta, \rho) = \Psi^\top \tilde{D} (\Psi \theta - P \Psi \theta - c + \rho \mathbf{1}). \quad (3.29)$$

Now switch our attention to the construction of the perturbed policy $\tilde{\pi}$. We first fix a constant $0 < \underline{\pi} < 1$. let $\mathcal{A}_s := \{a \in \mathcal{A} | \pi(a|s) > \underline{\pi}/2\}$, and the policy $\tilde{\pi}$ is constructed as follows,

$$\forall s \in \mathcal{S}, \quad \tilde{\pi}(a|s) := \begin{cases} \underline{\pi}, & \text{if } a \notin \mathcal{A}_s \\ \pi(a|s) \cdot \frac{1 - (|\mathcal{A}| - |\mathcal{A}_s|)\underline{\pi}}{\sum_{a' \in \mathcal{A}_s} \pi(a'|s)}, & \text{if } a \in \mathcal{A}_s \end{cases}. \quad (3.30)$$

We have that $\frac{1 - (|\mathcal{A}| - |\mathcal{A}_s|)\underline{\pi}}{\sum_{a' \in \mathcal{A}_s} \pi(a'|s)} = \frac{1 - (|\mathcal{A}| - |\mathcal{A}_s|)\underline{\pi}}{1 - \sum_{a' \notin \mathcal{A}_s} \pi(a'|s)} \leq \frac{1 - (|\mathcal{A}| - |\mathcal{A}_s|)\underline{\pi}}{1 - (|\mathcal{A}| - |\mathcal{A}_s|)\underline{\pi}/2} \leq 1$. On the other hand, when $\underline{\pi} \leq \frac{1}{|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|}$, we have $\frac{1 - (|\mathcal{A}| - |\mathcal{A}_s|)\underline{\pi}}{\sum_{a' \in \mathcal{A}_s} \pi(a'|s)} \geq 0$. Therefore, we ensure that $\tilde{\pi}(\cdot|s) \in \Delta_{|\mathcal{A}|}$ for all $s \in \mathcal{S}$. It should be noted that the EVRTD method reduces to the VRTD method (Algorithm 2) when $\mathcal{A}_s = \mathcal{A}$ for all $s \in \mathcal{S}$.

Lemma 6. Assume Assumption 2 holds. Recalling the definition of $1 - \beta$ in (3.16), we have

$$1 - \beta > 0. \quad (3.31)$$

Assume $|\mathcal{S}| \geq 2$. When $\min_{s \in \mathcal{S}} |\mathcal{A}_s| \neq |\mathcal{A}|$, if we take

$$\underline{\pi} \leq \frac{(1 - \beta) \min_{s \in \mathcal{S}} \nu(s)}{(|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|)(8 + (1 - \beta) \min_{s \in \mathcal{S}} \nu(s))}, \quad (3.32)$$

and construct the perturbed policy $\tilde{\pi}$ as (3.30), then

$$\inf_{x \in \mathbb{S}, \|x\|_{\tilde{D}}=1} x^\top \tilde{D}(I - P)x \geq \min \left\{ \frac{27\alpha(\underline{\pi})(1 - \beta)}{32}, \frac{1 - \alpha(\underline{\pi})(1 - \beta)}{8} \right\}, \quad (3.33)$$

where $\alpha(\underline{\pi}) := 1 - (|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|)\underline{\pi} \geq \frac{8}{9}$. As a consequence, Eq. (3.28) has a unique solution. Moreover, for any $\theta, \theta' \in \mathbb{R}^d$ and $\rho \in \mathbb{R}$,

$$\langle g(\theta, \rho) - g(\theta', \rho), \theta - \theta' \rangle \geq \min \left\{ \frac{27\alpha(\underline{\pi})(1 - \beta)}{32}, \frac{1 - \alpha(\underline{\pi})(1 - \beta)}{8} \right\} \cdot \|\Psi\theta - \Psi\theta'\|_{\tilde{D}}^2. \quad (3.34)$$

Proof. We first prove Ineq. (3.31). It should be noted that $\|\cdot\|_D$ is a semi-norm when there exists $(s, a) \in \mathcal{S} \times \mathcal{A}$ such that $\pi(a|s) = 0$. However, after removing the state-action pairs where $\pi(a|s) = 0$, we got an irreducible Markov chain with the state-action pairs left. Recalling the fact that $\mathcal{I} \cap \mathbb{S} = \emptyset$, then the proof of Ineq. (3.31) follows from the same argument as the proof of Lemma 3.

Next we prove Ineq. (3.33). The basic idea of this proof is to utilize Ineq. (3.31) and the construction of the perturbed policy $\tilde{\pi}$ to prove the desired lower bound (Ineq. (3.33)) in the weighted ℓ_2 -norm $\|\cdot\|_{\tilde{D}}$. Towards this end, we first define a vector $\hat{v} \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$ as $\hat{v}(s, a) = \nu(s)\underline{\pi}$ if $a \notin \mathcal{A}_s$ and $\hat{v}(s, a) = 0$ otherwise. Let $\hat{D} := \text{diag}(\hat{v})$. Then it is easy to see that $D + \hat{D} \succeq \tilde{D}$. Recalling the definition of $\alpha(\underline{\pi})$, we have for all $s \in \mathcal{S}, a \in \mathcal{A}$,

$$\tilde{\pi}(a|s) \geq \alpha(\underline{\pi})\pi(a|s) + \mathbb{I}_{\{a \notin \mathcal{A}_s\}}(\underline{\pi} - \alpha(\underline{\pi})\pi(a|s)). \quad (3.35)$$

Now we write

$$\begin{aligned} x^\top \tilde{D}(I - P)x &= \mathbb{E} [x(s, a)^2 - x(s, a)x(s', a') | s \sim \nu, a \sim \tilde{\pi}(\cdot|s), s' \sim P(\cdot|s, a), a' \sim \pi(\cdot|s')] \\ &= \underbrace{\frac{1}{2} \mathbb{E} [(x(s, a) - x(s', a'))^2 | s \sim \nu, a \sim \tilde{\pi}(\cdot|s), s' \sim P(\cdot|s, a), a' \sim \pi(\cdot|s')]}_{R_1} \\ &\quad + \underbrace{\frac{1}{2} \|x\|_{\tilde{D}}^2 - \frac{1}{2} \mathbb{E} [x(s', a')^2 | s \sim \nu, a \sim \tilde{\pi}(\cdot|s), s' \sim P(\cdot|s, a), a' \sim \pi(\cdot|s')]}_{R_2}. \end{aligned} \quad (3.36)$$

Next we lower bound R_1 and upper bound R_2 separately. Invoking Ineq. (3.35), we can write

$$\begin{aligned}
R_1 &\geq \alpha(\pi) \mathbb{E}[(x(s, a) - x(s', a'))^2 | s \sim \nu, a \sim \pi(\cdot | s), s' \sim P(\cdot | s, a), a' \sim \pi(\cdot | s')] \\
&\quad + \sum_{s \in \mathcal{S}} \nu(s) \sum_{a \notin \mathcal{A}_s} (\pi - \alpha(\pi) \pi(a | s)) \sum_{s' \in \mathcal{S}} P(s' | s, a) \sum_{a' \in \mathcal{A}} \pi(a' | s') (x(s, a) - x(s', a'))^2 \\
&\stackrel{(i)}{\geq} 2(1 - \beta) \alpha(\pi) \|x\|_D^2 + \frac{\pi}{2} \sum_{s \in \mathcal{S}} \nu(s) \sum_{a \notin \mathcal{A}_s} \sum_{s' \in \mathcal{S}} P(s' | s, a) \sum_{a' \in \mathcal{A}} \pi(a' | s') (x(s, a) - x(s', a'))^2 \\
&\stackrel{(ii)}{\geq} 2(1 - \beta) \alpha(\pi) \|x\|_D^2 + \frac{\pi}{2} \sum_{s \in \mathcal{S}} \nu(s) \sum_{a \notin \mathcal{A}_s} \sum_{s' \in \mathcal{S}} P(s' | s, a) \sum_{a' \in \mathcal{A}} \pi(a' | s') (\frac{1}{2} x(s, a)^2 - x(s', a')^2) \\
&\stackrel{(iii)}{\geq} 2(1 - \beta) \alpha(\pi) \|x\|_D^2 + \frac{1}{4} \|x\|_D^2 - \frac{\pi}{2} (|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|) \sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \pi(a' | s') x(s', a')^2 \\
&\stackrel{(iv)}{\geq} \frac{31(1-\beta)\alpha(\pi)}{16} \|x\|_D^2 + \frac{1}{4} \|x\|_D^2 \stackrel{(iv)}{\geq} \min \left\{ \frac{31(1-\beta)\alpha(\pi)}{16}, \frac{1}{4} \right\} \|x\|_D^2,
\end{aligned} \tag{3.37}$$

where step (i) follows from Eq. (3.18) and (3.31), step (ii) follows from the fact that $(x - y)^2 = x^2 - 2xy + y^2 \geq \frac{x^2}{2} - y^2$, step (iii) follows from $P(s' | s, a) \leq 1$ and $\sum_{s \in \mathcal{S}} \nu(s) = 1$, and step (iv) follows from that the definition of π ensures $\pi \leq \frac{\alpha(\pi)(1-\beta) \min_{s \in \mathcal{S}} \nu(s)}{8(|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|)}$, and step (iv) follows from $D + \hat{D} \succeq \tilde{D}$.

Next we upper bound R_2 . Write

$$\begin{aligned}
R_2 &= \sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \pi(a' | s') x(s', a')^2 \left(\sum_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}} \nu(s) \pi(a | s) P(s' | s, a) \right) \\
&\quad + \sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \pi(a' | s') x(s', a')^2 \left(\sum_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}} \nu(s) (\tilde{\pi}(a | s) - \pi(a | s)) P(s' | s, a) \right) \\
&\stackrel{(i)}{=} \sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \nu(s') \pi(a' | s') x(s', a')^2 \\
&\quad + \underbrace{\sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \pi(a' | s') x(s', a')^2 \left(\sum_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}} \nu(s) (\tilde{\pi}(a | s) - \pi(a | s)) P(s' | s, a) \right)}_{R_3},
\end{aligned} \tag{3.38}$$

where step (i) follows from the fact that ν is the stationary distribution under policy π . Next we bound R_3 on the RHS of the above equality.

$$\begin{aligned}
R_3 &\stackrel{(i)}{\leq} \sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \pi(a' | s') x(s', a')^2 \left(\sum_{s \in \mathcal{S}} \sum_{a \notin \mathcal{A}_s} \nu(s) (\tilde{\pi}(a | s) - \pi(a | s)) P(s' | s, a) \right) \\
&\stackrel{(ii)}{\leq} \sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \pi(a' | s') x(s', a')^2 \left(\sum_{s \in \mathcal{S}} \nu(s) (|\mathcal{A}| - |\mathcal{A}_s|) \pi \right) \\
&\stackrel{(iii)}{\leq} \frac{\alpha(\pi)(1-\beta)}{8} \sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} \nu(s') \pi(a' | s') x(s', a')^2,
\end{aligned}$$

where step (i) follows from that $\tilde{\pi}(a | s) - \pi(a | s) > 0$ only when $a \notin \mathcal{A}_s$, step (ii) follows from $P(s' | s, a) \leq 1$, and step (iii) follows from $\pi \leq \frac{\alpha(\pi)(1-\beta) \min_{s \in \mathcal{S}} \nu(s)}{8(|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|)}$. Combining the bound of R_3 with Eq. (3.38), we obtain

$$\begin{aligned}
R_2 &\leq \left(1 + \frac{\alpha(\pi)(1-\beta)}{8} \right) \sum_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}} \nu(s) \pi(a | s) x(s, a)^2 \\
&\stackrel{(i)}{\leq} \left(1 + \frac{\alpha(\pi)(1-\beta)}{8} \right) \sum_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}} \nu(s) \tilde{\pi}(a | s) x(s, a)^2 \\
&\quad + \left(1 + \frac{\alpha(\pi)(1-\beta)}{8} \right) \sum_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}_s} \nu(s) (\pi(a | s) - \tilde{\pi}(a | s)) x(s, a)^2 \\
&\stackrel{(ii)}{\leq} \left(1 + \frac{\alpha(\pi)(1-\beta)}{8} \right) \|x\|_D^2 + \left(1 + \frac{\alpha(\pi)(1-\beta)}{8} \right) \left(\frac{1}{\alpha(\pi)} - 1 \right) \|x\|_D^2 \\
&\stackrel{(iii)}{\leq} \left(1 + \frac{\alpha(\pi)(1-\beta)}{4} \right) \|x\|_D^2.
\end{aligned} \tag{3.39}$$

where step (i) follows from $\pi(a|s) - \tilde{\pi}(a|s) \geq 0$ only when $a \in \mathcal{A}_s$, step (ii) follows from Ineq. (3.35), and step (iii) follows from that the definition of $\underline{\pi}$ and $\alpha(\underline{\pi})$ ensures $(1 + \frac{\alpha(\underline{\pi})(1-\beta)}{8})(\frac{1}{\alpha(\underline{\pi})} - 1) \leq \frac{\alpha(\underline{\pi})(1-\beta)}{8}$. Combining the Ineqs. (3.36), (3.37) and (3.39) yields

$$x^\top \tilde{D}(I - P)x \geq (\min\{\frac{31(1-\beta)\alpha(\underline{\pi})}{32}, \frac{1}{8}\} + \frac{1}{2} - (\frac{1}{2} + \frac{\alpha(\underline{\pi})(1-\beta)}{8}))\|x\|_{\tilde{D}}^2 = \min\{\frac{27(1-\beta)\alpha(\underline{\pi})}{32}, \frac{1-\alpha(\underline{\pi})(1-\beta)}{8}\}\|x\|_{\tilde{D}}^2,$$

which completes the proof. \square

Given Lemma 6 above, we prove that Eq. (3.28) has a unique solution and establish the strong monotonicity of the deterministic operator; consequently, we are able to establish the convergence guarantees of EVRTD by proving the analogs of Theorem 1 and 2.

Proposition 3. *We note in passing that after replacing $1 - \beta$ with $\min\{\frac{27\alpha(\underline{\pi})(1-\beta)}{32}, \frac{1-\alpha(\underline{\pi})(1-\beta)}{8}\}$, ℓ_D -norm with $\ell_{\tilde{D}}$ -norm, M with \tilde{M} , $\tilde{\Sigma}$ with $\tilde{\Sigma}$, and slightly modifying some universal constants, the conclusions of Theorem 1 and 2 also hold for the EVRTD algorithm.*

In conclusion, we develop the EVRTD algorithm, where the agent follows the original policy π in most of the time and follows the perturbed policy $\tilde{\pi}$ only when collecting samples for stochastic approximation updates. The EVRTD algorithm recovers the convergence guarantees of VRTD in a weighted ℓ_2 -metric induced by the positive definite matrix $\tilde{D} \succ \frac{\alpha(\underline{\pi}) \cdot \underline{\pi}}{2} I$. Notice that when our desired guarantee is in ℓ_∞ -norm or ℓ_2 -norm, we need to pay for additional dependence on $1/\lambda_{\min}(\tilde{D})$.

Remark 2. *The design idea of the EVRTD algorithm naturally extends to the DMDP setting, by replacing the $1 - \beta$ with $1 - \gamma$, where γ is the discount factor. We provide proof of the analog of Lemma 6 for DMDPs in Appendix A.*

4 Actor: Policy mirror descent for AMDPs

Now we switch our attention to the problem of policy optimization. Our goal in this section is to develop a computationally efficient first-order method to solve the AMDP problem (2.5) in the stochastic setting. To start with, we first explore the first-order information of the average reward/cost function and establish a variational inequality (VI) formulation of the problem.

4.1 A variational inequality formulation

We first establish the following lemma for the gradient of the long-run average-reward function.

Lemma 7 (Gradient of average reward/cost). *For any $s \in \mathcal{S}$ and $a \in \mathcal{A}$,*

$$\frac{\partial \rho(\pi)}{\partial \pi(a|s)} = \nu^\pi(s) (\bar{Q}^\pi(s, a) + \nabla h^\pi(s, a)).$$

where $\nabla h^\pi(s, \cdot)$ denotes the gradient of $h^\pi(s)$ w.r.t. policy π .

Proof. By the relationship $\bar{V}^\pi(s_0) = \sum_{a' \in \mathcal{A}} \pi(a'|s_0) \bar{Q}^\pi(s_0, a')$, we have

$$\frac{\partial \bar{V}^\pi(s_0)}{\partial \pi(a|s)} = \sum_{a' \in \mathcal{A}} \left[\frac{\partial \pi(a'|s_0)}{\partial \pi(a|s)} \bar{Q}^\pi(s_0, a') + \pi(a'|s_0) \frac{\partial \bar{Q}^\pi(s_0, a')}{\partial \pi(a|s)} \right].$$

Using the relationship $\bar{Q}^\pi(s_0, a') = c(s_0, a') + h^\pi(s_0) - \rho(\pi) + \sum_{s'} P(s'|s_0, a') \bar{V}^\pi(s')$ and rearranging the terms, we obtain

$$\frac{\partial \rho(\pi)}{\partial \pi(a|s)} = \sum_{a' \in A} \left[\frac{\partial \pi(a|s_0)}{\partial \pi(a|s)} \bar{Q}^\pi(s_0, a') + \pi(a'|s_0) (\nabla h^\pi(s_0) + \sum_{s'} P(s'|s_0, a') \frac{\partial \bar{V}^\pi(s')}{\partial \pi(a|s)}) \right] - \frac{\partial \bar{V}^\pi(s_0)}{\partial \pi(a|s)} \quad (4.1)$$

Now taking a weighted sum of $\frac{\partial \rho(\pi)}{\partial \pi(a|s)}$ over ν^π , we arrive at

$$\begin{aligned} \frac{\partial \rho(\pi)}{\partial \pi(a|s)} &= \sum_{s_0 \in S} \nu^\pi(s_0) \left\{ \sum_{a' \in A} \left[\frac{\partial \pi(a'|s_0)}{\partial \pi(a|s)} \bar{Q}^\pi(s_0, a') + \pi(a'|s_0) (\nabla h^\pi(s_0) + \sum_{s'} P(s'|s_0, a') \frac{\partial \bar{V}^\pi(s')}{\partial \pi(a|s)}) \right] - \frac{\partial \bar{V}^\pi(s_0)}{\partial \pi(a|s)} \right\} \\ &= \sum_{s_0 \in S} \nu^\pi(s_0) \sum_{a' \in A} \left(\frac{\partial \pi(a'|s_0)}{\partial \pi(a|s)} \bar{Q}^\pi(s_0, a') + \pi(a|s_0) \nabla h^\pi(s_0) \right) \\ &\quad + \sum_{s' \in S} \sum_{s_0 \in S} \nu^\pi(s_0) \sum_{a' \in A} \pi(a'|s_0) P(s'|s_0, a') \frac{\partial \bar{V}^\pi(s')}{\partial \pi(a|s)} - \sum_{s_0 \in S} \nu^\pi(s_0) \frac{\partial \bar{V}^\pi(s_0)}{\partial \pi(a|s)} \\ &= \sum_{s_0 \in S} \nu^\pi(s_0) \sum_{a' \in A} \left(\frac{\partial \pi(a'|s_0)}{\partial \pi(a|s)} \bar{Q}^\pi(s_0, a') + \pi(a|s_0) \nabla h^\pi(s_0) \right) \end{aligned}$$

Notice that $\frac{\partial \pi(a'|s')}{\partial \pi(a|s)} = 1$ if $a' = a, s' = s$, otherwise being 0. Hence,

$$\frac{\partial \rho(\pi)}{\partial \pi(a|s)} = \nu^\pi(s) (\bar{Q}^\pi(s, a) + \nabla h^\pi(s)), \quad (4.2)$$

which completes the proof. \square

In view of lemma 7, the optimality condition of (2.5) suggests us to solve the following VI

$$\mathbb{E}_{s \in \nu^*} [\langle \bar{Q}^{\pi^*}, \pi(\cdot|s) - \pi^*(\cdot|s) \rangle + h^\pi(s) - h^{\pi^*}(s)] \geq 0. \quad (4.3)$$

This VI can be solved efficiently since it satisfies certain generalized monotonicity condition (Facchinei and Pang, 2003; Dang and Lan, 2015; Kotsalis et al., 2022a), thanks to the following performance difference lemma (generalized from Zhang and Ross, 2021).

Lemma 8 (Performance difference lemma). *Given any two policies π and π' , we have*

$$\rho(\pi') - \rho(\pi) = \mathbb{E}_{s \sim \nu^{\pi'}} [\langle \bar{Q}^\pi(s, \cdot), \pi'(\cdot|s) - \pi(\cdot|s) \rangle + h^{\pi'}(s) - h^\pi(s)]. \quad (4.4)$$

Proof. To prove the performance difference lemma, we expand the right hand side of Eq. (4.4).

$$\begin{aligned} &\mathbb{E}_{s \sim \nu^{\pi'}} [\langle \bar{Q}^\pi(s, \cdot), \pi'(\cdot|s) - \pi(\cdot|s) \rangle + h^{\pi'}(s) - h^\pi(s)] \\ &= \mathbb{E}_{s \sim \nu^{\pi'}, a \sim \pi'(\cdot|s)} [\bar{Q}^\pi(s, a) - \bar{V}^\pi(s) + h^{\pi'}(s) - h^\pi(s)] \\ &= \mathbb{E}_{s \sim \nu^{\pi'}, a \sim \pi'(\cdot|s)} [c(s, a) + h^{\pi'}(s) - \rho(\pi) + \mathbb{E}_{s' \sim P(\cdot|s, a)} [\bar{V}^\pi(s')] - \bar{V}^\pi(s)] \\ &\stackrel{(i)}{=} \rho(\pi') - \rho(\pi) + \mathbb{E}_{s' \sim \nu^{\pi'}} [\bar{V}^\pi(s')] - \mathbb{E}_{s \sim \nu^{\pi'}} [\bar{V}^\pi(s)] = \rho(\pi') - \rho(\pi), \end{aligned}$$

where step (i) follows from the fact that if $s \sim \nu^{\pi'}, a \sim \pi'(\cdot|s)$ and $s' \sim P(\cdot|s, a)$, then $s' \sim \nu^{\pi'}$. \square

As a consequence of Lemma 8, we arrive at

$$\rho(\pi^*) - \rho(\pi) = \mathbb{E}_{s \sim \nu^*} [\langle \bar{Q}^\pi(s, \cdot), \pi^*(\cdot|s) - \pi(\cdot|s) \rangle + h^{\pi^*}(s) - h^\pi(s)].$$

Since $\rho(\pi) - \rho(\pi^*) \geq 0$ for any feasible policy π , we conclude that

$$\mathbb{E}_{s \sim \nu^*} [\langle \bar{Q}^\pi(s, \cdot), \pi(\cdot|s) - \pi^*(\cdot|s) \rangle + h^\pi(s) - h^{\pi^*}(s)] \geq 0, \quad (4.5)$$

which implies the VI in Ineq. (4.3) satisfies the generalized monotonicity condition. To solve the given VI efficiently in the stochastic setting, we will design efficient stochastic iterative algorithms and provide theoretical guarantees in the following subsections.

4.2 Stochastic policy mirror descent for AMDPs

We start with introducing the following prox-mapping step that solves the average-reward policy optimization problem (2.5) iteratively:

$$\pi_{k+1}(\cdot|s) = \arg \min_{p(\cdot|s) \in \Delta_{|\mathcal{A}|}} \left\{ \lambda_k \left[\langle \bar{Q}^{\pi_k}(s, \cdot), p(\cdot|s) \rangle + h^p(s) \right] + D_{\pi_k}^p(s) \right\}, \forall s \in \mathcal{S}. \quad (4.6)$$

However, in the RL setting the exact basic differential Q-function \bar{Q}^{π_k} is unavailable, thus we replace with its stochastic estimator calculated with samples ζ_k , denoted as \bar{Q}^{π_k, ζ_k} , which leads to the following actor-critic type stochastic policy mirror descent (SPMD) algorithm. To make the

Algorithm 3 Stochastic policy mirror descent (SPMD) for AMDPs

- 1: **Input:** initial points $\pi_0(a|s) = 1/|\mathcal{A}|$, $\forall a \in \mathcal{A}, s \in \mathcal{S}$ and stepsize parameters λ_k ;
- 2: **for** $k = 0, 1, \dots, K$ **do**
- 3: **Critic step:** Implement a policy evaluation algorithm to evaluation \bar{Q}^{π_k} ,

$$\bar{Q}^{\pi_k, \zeta_k} = \text{critic}(\pi_k, \zeta_k). \quad (4.7)$$

- 4: **Actor step:** Implement the stochastic policy mirror descent update,

$$\pi_{k+1}(\cdot|s) = \arg \min_{p(\cdot|s) \in \Delta_{|\mathcal{A}|}} \left\{ \lambda_k \left[\langle \bar{Q}^{\pi_k, \zeta_k}(s, \cdot), p(\cdot|s) \rangle + h^p(s) \right] + D_{\pi_k}^p(s) \right\}, \forall s \in \mathcal{S}. \quad (4.8)$$

- 5: **end for**
-

policy evaluation error tractable, we define the following error measurements,

$$\mathbb{E}_{\zeta_k} \left[\bar{Q}^{\pi_k, \zeta_k} \right] = \tilde{Q}^{\pi_k}, \quad \min_{b \in \mathbb{R}} \left\| \tilde{Q}^{\pi_k} + b\mathbf{1} - \bar{Q}^{\pi_k} \right\|_{\infty} \leq \varsigma_k \quad (4.9)$$

$$\mathbb{E}_{\zeta_k} \left[\min_{b \in \mathbb{R}} \left\| \bar{Q}^{\pi_k, \zeta_k} + b\mathbf{1} - \bar{Q}^{\pi_k} \right\|_{\infty}^2 \right] \leq \sigma_k^2, \quad \mathbb{E}_{\zeta_k} \left[\min_{b \in \mathbb{R}} \left\| \bar{Q}^{\pi_k, \zeta_k} + b\mathbf{1} \right\|_{\infty}^2 \right] \leq \kappa_k^2. \quad (4.10)$$

for some $\sigma_k, \varsigma_k, \kappa_k \geq 0$. Note that we utilize these error measures because the update rule of Algorithm 3 is invariant with adding $b\mathbf{1}$ to \bar{Q}^{π_k, ζ_k} . As shown in Section 3, the bias term ς_k can be reduced much faster than the expected error term σ_k^2 , which enables us to sharpen the analysis. For convenience, we denote the σ -field $\mathcal{F}_s := \sigma(\zeta_0, \dots, \zeta_s)$ and $\delta_k := \bar{Q}^{\pi_k, \zeta_k} - \bar{Q}^{\pi_k}$.

To start with the convergence analysis, the following lemma (i.e., three-point lemma) characterizes the optimality condition of (4.8) in Algorithm 3 (see, e.g., Lemma 3.5 of Lan, 2020). Recall the definition of the convexity modulus $\omega \geq 0$ in Ineq. (2.2).

Lemma 9. *For any $p(\cdot|s) \in \Delta_{|\mathcal{A}|}$, we have*

$$\begin{aligned} & \lambda_k \left[\langle \bar{Q}^{\pi_k, \zeta_k}(s, \cdot), \pi_{k+1}(\cdot|s) - p(\cdot|s) \rangle + h^{\pi_{k+1}}(s) - h^p(s) \right] + D_{\pi_k}^{\pi_{k+1}}(s) \\ & \leq D_{\pi_k}^p(s) - (1 + \lambda_k \omega) D_{\pi_{k+1}}^p(s). \end{aligned} \quad (4.11)$$

As illustrated before, we distinguish two settings, namely the mixing unichain setting and the uniform ergodic setting as in Section 3. In the former setting, we only consider solving the

unregularized AMDPs ($h^\pi = 0$) under the generative model. In the latter setting, we solve both regularized and unregularized AMDPs under the Markovian noise model. Note that the regularized AMDPs can also be solved in the former setting when the subgradient of the regularizer is bounded, e.g., $\|\cdot\|_p^2$ for some $p > 1$. Moreover, for strongly convex regularizers, the complexity in Theorem 3 can be improved from $\mathcal{O}(\epsilon^{-2})$ to $\mathcal{O}(\epsilon^{-1})$ by exploiting the convex optimization literature (see Lan, 2020).

4.3 SPMD for unregularized AMDPs in the mixing unichain setting

The following theorem establishes the convergence guarantees of the SPMD algorithm for unregularized AMDPs.

Theorem 3. Suppose that $h^\pi = 0$. Set $\lambda_k = \lambda = \frac{\sqrt{2\log|\mathcal{A}|}}{\kappa\sqrt{K}}$ in Algorithm 3. Assume $\kappa_k \leq \kappa$. Then

$$\mathbb{E}[\sum_{t=0}^{K-1} (\rho(\pi_k) - \rho(\pi^*))] \leq \kappa\sqrt{2K\log|\mathcal{A}|} + 2K\varsigma. \quad (4.12)$$

Assume that r is uniformly chosen from $\{0, 1, \dots, K-1\}$, we have

$$\mathbb{E}[\rho(\pi_r) - \rho(\pi^*)] \leq \frac{\kappa\sqrt{2\log|\mathcal{A}|}}{\sqrt{K}} + 2\varsigma.$$

Proof. Given $h^\pi = 0$, we have $\omega = 0$. By Lemma 9 applied with $\lambda_k = \lambda$ and $p = \pi^*$, we have

$$\lambda\langle \bar{Q}^{\pi_k, \zeta_k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi^*(\cdot|s) \rangle + D_{\pi_k}^{\pi_{k+1}}(s) \leq D_{\pi_k}^{\pi^*}(s) - D_{\pi_{k+1}}^{\pi^*}(s).$$

Rearranging the terms gives us

$$\begin{aligned} & \lambda\langle \bar{Q}^{\pi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle \\ & \leq D_{\pi_k}^{\pi^*}(s) + \lambda\langle \bar{Q}^{\pi_k, \zeta_k}(\cdot|s), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle - D_{\pi_{k+1}}^{\pi^*}(s) - D_{\pi_k}^{\pi_{k+1}}(s) + \lambda\langle \delta_k(\cdot|s), \pi^*(\cdot|s) - \pi_k(\cdot|s) \rangle \\ & \stackrel{(i)}{\leq} D_{\pi_k}^{\pi^*}(s) + \lambda(\min_{b \in \mathbb{R}} \|\bar{Q}^{\pi_k, \zeta_k}(\cdot|s) + b\mathbf{1}\|_\infty) \|\pi_{k+1}(\cdot|s) - \pi_k(\cdot|s)\|_1 - \frac{1}{2} \|\pi_{k+1}(\cdot|s) - \pi_k(\cdot|s)\|_1^2 \\ & \quad - D_{\pi_{k+1}}^{\pi^*}(s) + \lambda\langle \delta_k(\cdot|s), \pi^*(\cdot|s) - \pi_k(\cdot|s) \rangle \\ & \stackrel{(ii)}{\leq} D_{\pi_k}^{\pi^*}(s) + \frac{\lambda^2}{2} \min_{b \in \mathbb{R}} \|\bar{Q}^{\pi_k, \zeta_k}(\cdot|s) + b\mathbf{1}\|_\infty^2 - D_{\pi_{k+1}}^{\pi^*}(s) + \lambda\langle \delta_k(\cdot|s), \pi^*(\cdot|s) - \pi_k(\cdot|s) \rangle, \end{aligned}$$

where step (i) follows from $\langle \mathbf{1}, \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle = 0$, Cauchy-Schwarz inequality and $D_{\pi_k}^{\pi_{k+1}}(s) \geq \frac{1}{2} \|\pi_{k+1}(\cdot|s) - \pi_k(\cdot|s)\|_1^2$, step (ii) follows from Young's inequality. By taking expectation with respect to \mathcal{F}_{k-1} , we obtain that

$$\begin{aligned} \lambda\langle \bar{Q}^{\pi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle & \leq D_{\pi_k}^{\pi^*}(s) + \frac{\lambda^2\kappa^2}{2} - \mathbb{E}[D_{\pi_{k+1}}^{\pi^*}(s)|\mathcal{F}_{k-1}] + \lambda\langle \mathbb{E}[\delta_k(\cdot|s)|\mathcal{F}_{k-1}], \pi^*(\cdot|s) - \pi_k(\cdot|s) \rangle \\ & \leq D_{\pi_k}^{\pi^*}(s) + \frac{\lambda^2\kappa^2}{2} - \mathbb{E}[D_{\pi_{k+1}}^{\pi^*}(s)|\mathcal{F}_{k-1}] + 2\lambda\varsigma. \end{aligned}$$

Taking expectation with respect to ν^* and full expectation with respect to \mathcal{F}_{-1} , and taking telescope sum from 0 to $K-1$, we have

$$\mathbb{E}[\sum_{k=0}^{K-1} \mathbb{E}_{s \sim \nu^*} [\lambda\langle \bar{Q}^{\pi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle]] \leq D(\pi_0, \pi^*) + \frac{K\lambda^2\kappa^2}{2} + 2K\lambda\varsigma.$$

where $D(\pi, \pi') := \mathbb{E}_{s \sim \nu^*}[D_{\pi'}^{\pi}(s)]$. Using Lemma 8 we arrive at

$$\mathbb{E}[\sum_{k=0}^{K-1} \lambda(\rho(\pi_k) - \rho(\pi^*))] \leq D(\pi_0, \pi^*) + \frac{K\lambda^2\kappa^2}{2} + 2K\lambda\varsigma.$$

Invoking that r is uniformly chosen from $\{0, \dots, K-1\}$ gives us

$$\mathbb{E}[\rho(\pi_r) - \rho(\pi^*)] \leq \frac{D(\pi_0, \pi^*)}{\lambda K} + \frac{\lambda\kappa^2}{2} + 2\varsigma.$$

By applying the upper bound $D(\pi_0, \pi^*) \leq \log |\mathcal{A}|$ and the choice of λ , we complete the proof. \square

Below we discuss the sample complexity of finding an ϵ -optimal solution, i.e., a solution $\hat{\pi}$, such that $\mathbb{E}[\rho(\hat{\pi}) - \rho(\pi^*)] \leq \epsilon$, under the generative model. We implement the multiple trajectory method (Algorithm 1) as the critic solver in (4.7). Let T_k, T'_k, M_k, M'_k denote the parameters T, T', M, M' in Algorithm 1 for estimating \bar{Q}^{π_k, ζ_k} respectively.

Corollary 1 (Sample complexity under generative model). *Consider the mixing AMDPs which satisfy Assumption 1. Suppose the number of iterations K is given a priori and $\lambda = \frac{\sqrt{2 \log |\mathcal{A}|}}{\kappa \sqrt{K}}$. If $T'_k \geq T' = t_{\text{mix}} \cdot \log_2 \left(\frac{4t_{\text{mix}}}{\epsilon} \right)$, $T_k \geq T = t_{\text{mix}} \cdot \log_2 \left(\frac{2(T'+1)}{\epsilon} \right) + 1$ and $M_k = M = M'_k = M' = 1$, then an ϵ -optimal solution can be found in at most $\mathcal{O}(\kappa^2 \log |\mathcal{A}| / \epsilon^2)$ SPMD iterations and the total number of state-action transition samples can be bounded by $\tilde{\mathcal{O}}\left(\frac{|\mathcal{S}||\mathcal{A}|t_{\text{mix}}^3}{\epsilon^2}\right)$.*

Proof. Notice that we can upper bound κ by $\kappa \leq (\bar{c} + \bar{h})T' = \tilde{\mathcal{O}}(t_{\text{mix}})$, where $c(s, a) \leq \bar{c}$ and $h(s) \leq \bar{h}$. Then the total number of iterations K is bounded by $\tilde{\mathcal{O}}(t_{\text{mix}}^2 / \epsilon^2)$. Moreover, for each SPMD iteration, we require $M_k T_k + |\mathcal{S}||\mathcal{A}|M'_k T'_k = \tilde{\mathcal{O}}(|\mathcal{S}||\mathcal{A}|t_{\text{mix}})$ samples for policy evaluation. Therefore, the total number of samples can be bounded by $\tilde{\mathcal{O}}\left(\frac{|\mathcal{S}||\mathcal{A}|t_{\text{mix}}^3}{\epsilon^2}\right)$, as desired. \square

To the best of our knowledge, the result in Corollary 1 is the first sample complexity result of policy gradient/actor-critic methods for solving AMDPs. This complexity result matches the lower bound $\Omega\left(\frac{t_{\text{mix}}|\mathcal{S}||\mathcal{A}|}{\epsilon^2}\right)$ proved in Jin and Sidford (2021) in terms of dependence on ϵ as well as on $|\mathcal{S}||\mathcal{A}|$ up to logarithmic factors. The dependence of t_{mix} appears to be worse than the state-of-the-art LP approaches (Wang, 2017; Jin and Sidford, 2020) where the complexity is $\tilde{\mathcal{O}}\left(\frac{t_{\text{mix}}^2|\mathcal{S}||\mathcal{A}|}{\epsilon^2}\right)$. We notice that this performance difference between policy gradient type methods and model-based or value-based methods also exists in the DMDP setting; With policy gradient type methods, the best-known dependence on the effective horizon is $\tilde{\mathcal{O}}(1/(1-\gamma)^5)$, failing to match the $\tilde{\mathcal{O}}(1/(1-\gamma)^3)$ results achieved by model-based or value-based methods. Nevertheless, the policy gradient type methods still play an important role in the RL literature, partially because it naturally accommodates the Markovian noise setting, especially in an on-policy manner.

4.4 SPMD for unregularized/regularized AMDPs in the ergodic setting

In this section, we take the Markovian noise model as the observation model and assume the ergodicity assumption (Assumption 2) holds correspondingly.

We utilize the EVRTD method with linear function approximation introduced in Section 3.2 as the “critic” solver. Specifically, when estimating \bar{Q}^{π_k} , the first task is to find a π_k satisfying Ineq. (3.32). As such, some rough estimation of the mixing time and stationary distribution is required, which has been the topic of active research (e.g., Hsu et al., 2019; Wolfer and Kontorovich,

2019). Note that this task is also required for obtaining the parameters τ and τ' . After obtaining π_k , we implement the EVRTD algorithm introduced in Section 3.3 to evaluate \bar{Q}^{π_k} . Note that if $\pi_k(a|s) > \frac{\pi_k}{2}$ for all $s \in \mathcal{S}$, $a \in \mathcal{A}$, EVRTD reduces to the original VRTD method (Algorithm 2).

The SPMD algorithm naturally benefit from the linear function approximation of the policy evaluation steps. If we use the same feature matrix Ψ for the policy evaluation steps of all the policy iterates, the updating rule of Algorithm 3 reduces to the updates of a low dimensional vector in \mathbb{R}^d , which will be illustrated as follows. Let $\theta_k \in \mathbb{R}^d$ denote the low-dimensional output of evaluating policy π_k . If we consider the case where $h^\pi(s) = \omega \cdot \sum_{a \in \mathcal{A}} \pi(a|s) \log \pi(a|s)$ for $\omega \geq 0$, then by utilizing the Karush-Kuhn-Tucker conditions and recursive arguments, we can write the solution of step (4.8) as

$$\begin{aligned} \pi_{k+1}(\cdot|s) &\propto \exp\left(-\frac{\lambda_k}{1+\lambda_k\omega}(\Phi\theta_k)(s, \cdot) + \frac{1}{1+\lambda_k\omega} \log \pi_k(\cdot|s)\right) \\ &\propto \exp\left((\Phi\tilde{\theta}_{k+1})(s, \cdot)\right) \propto \exp\left((\Phi\left[-\frac{\lambda_k}{1+\lambda_k\omega}\theta_k + \frac{1}{1+\lambda_k\omega}\tilde{\theta}_k\right])(s, \cdot)\right), \end{aligned}$$

where $\tilde{\theta}_k := -\sum_{t=0}^{k-1} \left[\left(\prod_{l=t}^{k-1} \frac{1}{1+\lambda_l\omega}\right) \lambda_t \theta_t\right]$. It indicates that π_{k+1} is a function of $\tilde{\theta}_{k+1}$, which can be calculated with $\tilde{\theta}_k$ and θ_k . Towards this end, for $k \geq 1$, the critic step (4.7) can be equivalently written as $\theta_k = \text{critic}(\tilde{\theta}_k, \zeta_k)$. In words, we only need to calculate the policy π_k for generating the Markov trajectory in the critic step, and we only maintain the low dimensional vectors $\tilde{\theta}_k$ when implementing Algorithm 3. Note that SPMD method can also benefit from the linear function approximation when solving AMDPs with other regularization.

To establish the convergence guarantees, we require some additional notation. For a fixed policy π , we let θ_π^* denote the solution of the projected Bellman equation (3.28) induced by policy π . Since the approximation error introduced by linear function approximation cannot be eliminated, we can only establish the convergence to a neighborhood of the optimal solution whose diameter is the approximation error defined as $\epsilon_{\text{approx}} := \max_{\pi} \min_{b \in \mathbb{R}} \|\Psi\theta_\pi^* + b\mathbf{1} - \bar{Q}^\pi\|_\infty$. Moreover, we let $\bar{\mu}$ be a lower bound of μ defined in (3.13) across all feasible policies.

4.4.1 Unregularized AMDPs

For unregularized AMDPs, notice that Theorem 3 still holds given that Assumption 2 is stronger than Assumption 1. We first implement the policy mirror descent in Theorem 3 with EVRTD critic solver and establish the following sample complexity.

Corollary 2 (Sample complexity of unregularized AMDPs). *Consider the mixing AMDPs which satisfy Assumption 2. Suppose the number of iterations K of Algorithm 3 is given a priori and $\lambda = \frac{\sqrt{2 \log |\mathcal{A}|}}{\kappa \sqrt{K}}$. For each policy π_k , implement the EVRTD algorithm with parameter selection in Theorem 2 for $\log(1/\epsilon)$ epochs. Then a solution $\hat{\pi}$, which satisfies that $\mathbb{E}[\rho(\hat{\pi}) - \rho(\pi^*)] \leq \mathcal{O}(1)\epsilon_{\text{approx}} + \epsilon$, can be found in at most $\mathcal{O}(\kappa^2 \log |\mathcal{A}|/\epsilon^2)$ SPMD iterations and the total number of state-action transition samples can be bounded by $\tilde{\mathcal{O}}\left(\frac{t_{\text{mix}}^3 \kappa^2}{\bar{\mu} \epsilon^2}\right)$. Moreover, we have $\kappa^2 \leq 2 \max_{\pi} \|\Psi\theta_\pi^*\|_\infty^2 + \mathcal{O}\left(\max_{\pi} \frac{\|\theta_\pi^*\|_2^2}{\lambda_{\min}(D^\pi)}\right)$.*

Proof. For each SPMD iteration, to ensure policy evaluation bias to satisfy $\varsigma_k \leq \epsilon + \mathcal{O}(1)\epsilon_{\text{approx}}$, we require $\tilde{\mathcal{O}}\left(\frac{t_{\text{mix}}^3}{\bar{\mu}}\right)$ samples for policy evaluation by Theorem 2. Therefore, the total number of samples can be bounded by $\tilde{\mathcal{O}}\left(\frac{t_{\text{mix}}^3 \kappa^2}{\bar{\mu} \epsilon^2}\right)$. Next, we proof an upper bound for κ^2 . For $k = 0, \dots, K-1$,

by Theorem 1 we can upper bound κ_k^2 by

$$\begin{aligned}\kappa_k^2 &\leq 2\|\Psi\theta_{\pi_k}^*\|_\infty^2 + 2\mathbb{E}\|\bar{\mathcal{Q}}^{\pi_k, \zeta_k} - \Psi\theta_{\pi_k}^*\|_\infty^2 \leq 2\|\Psi\theta_{\pi_k}^*\|_\infty^2 + 2\mathbb{E}[\|\bar{\mathcal{Q}}^{\pi_k, \zeta_k} - \Psi\theta_{\pi_k}^*\|_\infty^2 / \lambda_{\min}(\tilde{D}^{\pi_k})] \\ &\leq 2\|\Psi\theta_{\pi_k}^*\|_\infty^2 + \mathcal{O}\left(\frac{\|\theta_{\pi_k}^*\|_2^2}{\lambda_{\min}(\tilde{D}^{\pi_k})}\right) \leq 2\max_{\pi}\|\Psi\theta_{\pi}^*\|_\infty^2 + \mathcal{O}\left(\max_{\pi}\frac{\|\theta_{\pi}^*\|_2^2}{\lambda_{\min}(\tilde{D}^{\pi})}\right),\end{aligned}$$

as desired. \square

4.4.2 Regularized AMDPs

Next, we consider the regularized AMDPs. By Assumption 2, there exists $\Gamma \in (0, 1)$, such that for all feasible policy π ,

$$\min_{s \in \mathcal{S}} \nu^\pi(s) \geq 1 - \Gamma.$$

Under this assumption, the convergence guarantees of the SPMD method for solving AMDPs are mostly similar to the the SPMD method for DMDPs presented in Lan (2022). To demonstrate this claim, we first establish two supporting lemmas whose analogs are Lemma 12 and 13 in Lan (2022).

Lemma 10. *For any $s \in \mathcal{S}$, we have*

$$\begin{aligned}\frac{1}{1-\Gamma}(\rho(\pi_{k+1}) - \rho(\pi_k)) &\leq \langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s, \cdot), \pi_{k+1}(\cdot | s) - \pi_k(\cdot | s) \rangle + h^{\pi_{k+1}}(s) - h^{\pi_k}(s) \\ &\quad + \frac{1}{\lambda_k} D_{\pi_k}^{\pi_{k+1}}(s) + \frac{\lambda_k \min_{b \in \mathbb{R}} \|\delta_k + b\mathbf{1}\|_\infty^2}{2(1-\Gamma)}\end{aligned}$$

Proof. Using Lemma 8, we obtain

$$\begin{aligned}\rho(\pi_{k+1}) - \rho(\pi_k) &= \mathbb{E}_{s' \sim \nu^{\pi_{k+1}}} [\langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s', \cdot), \pi_{k+1}(\cdot | s') - \pi_k(\cdot | s') \rangle + h^{\pi_{k+1}}(s') - h^{\pi_k}(s') \\ &\quad - \langle \delta_k(s', \cdot), \pi_{k+1}(\cdot | s') - \pi_k(\cdot | s') \rangle] \\ &\stackrel{(i)}{\leq} \mathbb{E}_{s' \sim \nu^{\pi_{k+1}}} [\langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s', \cdot), \pi_{k+1}(\cdot | s') - \pi_k(\cdot | s') \rangle + h^{\pi_{k+1}}(s') - h^{\pi_k}(s') \\ &\quad + \frac{1}{2\lambda_k} \|\pi_{k+1}(\cdot | s') - \pi_k(\cdot | s')\|_1^2 + \frac{\lambda_k \min_{b \in \mathbb{R}} \|\delta_k + b\mathbf{1}\|_\infty^2}{2}] \\ &\stackrel{(ii)}{\leq} \mathbb{E}_{s' \sim \nu^{\pi_{k+1}}} [\langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s', \cdot), \pi_{k+1}(\cdot | s') - \pi_k(\cdot | s') \rangle + h^{\pi_{k+1}}(s') - h^{\pi_k}(s') \\ &\quad + \frac{1}{\lambda_k} D_{\pi_k}^{\pi_{k+1}}(s') + \frac{\lambda_k \min_{b \in \mathbb{R}} \|\delta_k + b\mathbf{1}\|_\infty^2}{2}],\end{aligned}\tag{4.13}$$

where step (i) follows from Young's inequality and the fact that $\langle \mathbf{1}, \pi_{k+1}(\cdot | s') - \pi_k(\cdot | s') \rangle = 0$, step (ii) follows from the strong convexity of $D_{\pi_k}^{\pi_{k+1}}$ with respect to ℓ_1 -norm. Recalling the three-point lemma (Ineq. (4.11)) and taking $p = \pi_k$, we obtain that for all $s' \in \mathcal{S}$,

$$\begin{aligned}\langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s', \cdot), \pi_{k+1}(\cdot | s') - \pi_k(\cdot | s') \rangle + h^{\pi_{k+1}}(s') - h^{\pi_k}(s') + \frac{1}{\lambda_k} D_{\pi_k}^{\pi_{k+1}}(s') \\ \leq -\frac{1}{\lambda_k} [(1 + \lambda_k \omega) D_{\pi_{k+1}}^{\pi_k}(s')] \leq 0,\end{aligned}$$

which implies that

$$\begin{aligned}\mathbb{E}_{s' \sim \nu^{\pi_{k+1}}} [\langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s', \cdot), \pi_{k+1}(\cdot | s') - \pi_k(\cdot | s') \rangle + h^{\pi_{k+1}}(s') - h^{\pi_k}(s') + \frac{1}{\lambda_k} D_{\pi_k}^{\pi_{k+1}}(s')] \\ \leq \nu^{\pi_{k+1}}(s) [\langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s, \cdot), \pi_{k+1}(\cdot | s) - \pi_k(\cdot | s) \rangle + h^{\pi_{k+1}}(s) - h^{\pi_k}(s) + \frac{1}{\lambda_k} D_{\pi_k}^{\pi_{k+1}}(s)] \\ \leq (1 - \Gamma) [\langle \bar{\mathcal{Q}}^{\pi_k, \zeta_k}(s, \cdot), \pi_{k+1}(\cdot | s) - \pi_k(\cdot | s) \rangle + h^{\pi_{k+1}}(s) - h^{\pi_k}(s) + \frac{1}{\lambda_k} D_{\pi_k}^{\pi_{k+1}}(s)],\end{aligned}$$

and the desired result follows from combining the above inequality with Ineq. (4.13). \square

Lemma 11. For any $k \geq 0$, we have

$$\begin{aligned} & \mathbb{E}\left[\frac{1}{1-\Gamma}(\rho(\pi_{k+1}) - \rho(\pi^*)) + \left(\frac{1}{\lambda_k} + \omega\right)D(\pi_{k+1}, \pi^*)\right] \\ & \leq \mathbb{E}\left[\frac{\Gamma}{1-\Gamma}(\rho(\pi_k) - \rho(\pi^*)) + \frac{1}{\lambda_k}D(\pi_k, \pi^*)\right] + 2\varsigma_k + \frac{\lambda_k \sigma_k^2}{2(1-\Gamma)}, \end{aligned}$$

where the expectation is taken with respect to \mathcal{F}_{-1} , and $D(\pi, \pi') := \mathbb{E}_{s \sim \nu^*}[D_\pi^{\pi'}(s)]$.

Proof. By applying Lemma 9 with $p = \pi^*$ we obtain

$$\begin{aligned} & \lambda_k \left[\langle \bar{Q}^{\pi_k, \zeta_k}(s, \cdot), \pi_{k+1}(\cdot | s) - \pi^*(\cdot | s) \rangle + h^{\pi_{k+1}}(s) - h^{\pi^*}(s) \right] + D_{\pi_k}^{\pi_{k+1}}(s) \\ & \leq D_{\pi_k}^{\pi^*}(s) - (1 + \lambda_k \omega) D_{\pi_{k+1}}^{\pi^*}(s), \end{aligned}$$

Combining the above inequality with Lemma 10 implies that

$$\begin{aligned} & \langle \bar{Q}^{\pi_k, \zeta_k}(s, \cdot), \pi_k(\cdot | s) - \pi^*(\cdot | s) \rangle + h^{\pi_k}(s) - h^{\pi^*}(s) + \frac{1}{1-\Gamma}(\rho(\pi_{k+1}) - \rho(\pi_k)) \\ & \leq \frac{1}{\lambda_k} D_{\pi_k}^{\pi^*}(s) - \left(\frac{1}{\lambda_k} + \omega\right) D_{\pi_{k+1}}^{\pi^*}(s) + \frac{\lambda_k \min_{b \in \mathbb{R}} \|\delta_k + b\mathbf{1}\|_\infty^2}{2(1-\Gamma)}. \end{aligned}$$

Taking full expectation with \mathcal{F}_{-1} and ν^* , and using Lemma 8, we arrive at

$$\mathbb{E}\left[(\rho(\pi_k) - \rho(\pi^*)) + \frac{1}{1-\Gamma}(\rho(\pi_{k+1}) - \rho(\pi_k))\right] \leq \mathbb{E}\left[\frac{1}{\lambda_k} D_{\pi_k}^{\pi^*}(s) - \left(\frac{1}{\lambda_k} + \omega\right) D_{\pi_{k+1}}^{\pi^*}(s)\right] + 2\varsigma_k + \frac{\lambda_k \sigma_k^2}{2(1-\Gamma)}.$$

Rearranging terms yields the desired result. \square

Remark 3. In view of Lemma 11 and its analog (Lemma 13) in Lan (2022), it can be easily proved that Theorem 5 and 6 in Lan (2022) still holds by replacing $f(\cdot)$ with $\rho(\cdot)/(1-\Gamma)$ and γ with Γ . Similarly, the convergence guarantees for deterministic PMD in Lan (2022) can also be extended to AMDPs.

Nevertheless, all the convergence guarantees in Lan (2022) failed to decompose the convergence rate into a deterministic error part with linear convergence rate and a stochastic error part with sublinear convergence rate, which motivates us to derive the following theorem.

Theorem 4. Assume that $\omega > 0$. If K is fixed and $\varsigma_k \leq \varsigma$, $\sigma_t \leq \sigma$ for any $0 \leq k \leq K$, set

$$\lambda_k = \lambda = \min\left\{\frac{1-\Gamma}{\omega\Gamma}, \frac{q \log K}{\omega K}\right\}, \quad \text{with} \quad q = 2\left(1 + \log\left(\frac{2\Gamma(1-\Gamma)\omega^2 D(\pi_0, \pi^*) + 2\Gamma(1-\Gamma)\omega(\rho(\pi_0) - \rho(\pi^*))}{\sigma^2}\right)\right), \quad (4.14)$$

then

$$\mathbb{E}[D(\pi_{K+1}, \pi^*)] \leq \frac{\Gamma^K}{\omega}(\rho(\pi_0) - \rho(\pi^*)) + \Gamma^K D(\pi_0, \pi^*) + \frac{\sigma^2}{\omega^2 \Gamma(1-\Gamma)K}(1 + q \log K) + \frac{4\varsigma}{\omega\Gamma}.$$

Proof. By Lemma 11, $\lambda_k = \lambda$, $\varsigma_k \leq \varsigma$, $\sigma_k \leq \sigma$, we have that with $\theta_k > 0$

$$\begin{aligned} & \theta_k \mathbb{E}\left[\frac{1}{1-\Gamma}(\rho(\pi_{k+1}) - \rho(\pi^*)) + \left(\frac{1}{\lambda} + \omega\right)D(\pi_{k+1}, \pi^*)\right] \\ & \leq \theta_k \mathbb{E}\left[\frac{\Gamma}{1-\Gamma}(\rho(\pi_k) - \rho(\pi^*)) + \frac{1}{\lambda}D(\pi_k, \pi^*)\right] + 2\theta_k \varsigma + \frac{\theta_k \lambda \sigma^2}{2(1-\Gamma)}. \end{aligned}$$

Set $\theta_k = \theta^k$ and $\theta \leq \min\{\frac{1}{\Gamma}, 1 + \omega\lambda\}$, then by taking telescope sum from $k = 0$ to K , we obtain

$$\begin{aligned} & \theta^K \cdot \mathbb{E}\left[\frac{1}{1-\Gamma} (\rho(\pi_{K+1}) - \rho(\pi^*)) + \left(\frac{1}{\lambda} + \omega\right) D(\pi_{K+1}, \pi^*)\right] \\ & \leq \frac{\Gamma}{1-\Gamma} (\rho(\pi_0) - \rho(\pi^*)) + \frac{1}{\lambda} D(\pi_0, \pi^*) + \frac{2\theta^K \zeta}{1-\theta^{-1}} + \frac{\theta^K \lambda \sigma^2}{2(1-\Gamma)(1-\theta^{-1})}. \end{aligned}$$

From $\theta \leq \min\{\frac{1}{\Gamma}, 1 + \omega\lambda\}$, we have $(1 - \theta^{-1})^{-1} \geq \max\{\frac{1}{1-\Gamma}, \frac{1+\omega\lambda}{\omega\lambda}\}$. Using this relationship and multiplying $\theta^{-K}\lambda$ on both sides of the above inequality, we obtain

$$\begin{aligned} & \lambda \mathbb{E}\left[\frac{1}{1-\Gamma} (\rho(\pi_{K+1}) - \rho(\pi^*))\right] + (1 + \omega\lambda) \mathbb{E}[D(\pi_{K+1}, \pi^*)] \\ & \leq \theta^{-K} \frac{\lambda \Gamma}{1-\Gamma} (\rho(\pi_0) - \rho(\pi^*)) + \theta^{-K} D(\pi_0, \pi^*) + \frac{2\lambda \zeta}{1-\Gamma} + \frac{2(1+\omega\lambda)\zeta}{\omega} + \frac{\lambda^2 \sigma^2}{2(1-\Gamma)^2} + \frac{(1+\omega\lambda)\lambda \sigma^2}{2(1-\Gamma)\omega}. \end{aligned} \quad (4.15)$$

Take $\lambda = \min\{\frac{1-\Gamma}{\omega\Gamma}, \frac{q \log K}{\omega K}\}$ and $\theta = \min\{\frac{1}{\Gamma}, 1 + \frac{q \log K}{K}\}$, then we have

$$\theta^{-K} \leq \Gamma^K + (1 + \frac{q \log K}{K})^{-K} \leq \Gamma^K + \exp(-\frac{1}{2} q \log K) \leq \Gamma^K + \frac{1}{K^{q/2}}. \quad (4.16)$$

Invoking the definition that $q = 2(1 + \log(\frac{2\Gamma(1-\Gamma)\omega^2 D(\pi_0, \pi^*) + 2\Gamma(1-\Gamma)\omega(\rho(\pi_0) - \rho(\pi^*))}{\sigma^2}))$, we have

$$\frac{\lambda \Gamma}{(1-\Gamma)K^{q/2}} \leq \frac{\sigma^2}{2\Gamma(1-\Gamma)\omega^2 K} \quad \text{and} \quad \frac{D(\pi_0, \pi^*)}{K^{q/2}} \leq \frac{\sigma^2}{2\Gamma(1-\Gamma)\omega^2 K}. \quad (4.17)$$

Combining Ineqs. (4.15), (4.16) and (4.17) yields

$$\begin{aligned} & \lambda \mathbb{E}\left[\frac{1}{1-\Gamma} (\rho(\pi_{K+1}) - \rho(\pi^*))\right] + \mathbb{E}[D(\pi_{K+1}, \pi^*)] \\ & \leq \frac{\Gamma^K}{\omega} (\rho(\pi_0) - \rho(\pi^*)) + \Gamma^K D(\pi_0, \pi^*) + \frac{\lambda \Gamma}{(1-\Gamma)K^{q/2}} + \frac{D(\pi_0, \pi^*)}{K^{q/2}} + \frac{4\zeta}{\omega\Gamma} + \frac{\lambda \sigma^2}{\omega\Gamma(1-\Gamma)} \\ & \leq \frac{\Gamma^K}{\omega} (\rho(\pi_0) - \rho(\pi^*)) + \Gamma^K D(\pi_0, \pi^*) + \frac{\sigma^2}{\omega^2 \Gamma(1-\Gamma)K} (1 + q \log K) + \frac{4\zeta}{\omega\Gamma}, \end{aligned}$$

which completes the proof. \square

Theorem 4 establishes the convergence guarantee for the distance between the output policy and the optimal policy. Meanwhile, we are allowed to output the policy generated by the last iteration instead of randomly output a policy as in Theorem 3. The next corollary provides the sample complexity for obtaining a convergence guarantee on the output policy.

Corollary 3 (Sample complexity for regularized AMDPs). *Consider the mixing AMDPs which satisfy Assumption 2. Suppose the number of iterations K is given a priori and λ satisfies (4.14). For each policy π_k , implement the EVRTD algorithm with parameter selection in Theorem 2 for $\log(1/\epsilon)$ epochs, then a solution $\hat{\pi}$, which satisfies that $\mathbb{E}[D(\hat{\pi}, \pi^*)] \leq \mathcal{O}(1) \frac{\epsilon_{\text{approx}}}{\omega\Gamma} + \epsilon$, can be found in at most $\tilde{\mathcal{O}}(\frac{\sigma^2}{\omega^2(1-\Gamma)\epsilon})$ SPMD iterations and the total number of state-action transition samples can be bounded by $\tilde{\mathcal{O}}(\frac{\sigma^2 t_{\text{mix}}^3}{\omega^2(1-\Gamma)\bar{\mu}\epsilon})$. Moreover, we have $\sigma^2 \leq 2\epsilon_{\text{approx}}^2 + \mathcal{O}(\max_{\pi} \frac{\|\theta_{\pi}^*\|_2^2}{\lambda_{\min}(\tilde{D}^{\pi})})$.*

Proof. For each SPMD iteration, to ensure policy evaluation bias to satisfy $\varsigma_k \leq \epsilon + \mathcal{O}(1)\epsilon_{\text{approx}}/(\omega\Gamma)$, we require $\tilde{\mathcal{O}}(\frac{t_{\text{mix}}^3}{\bar{\mu}})$ samples for policy evaluation by Theorem 2. Therefore, the total number of samples can be bounded by $\tilde{\mathcal{O}}(\frac{\sigma^2 t_{\text{mix}}^3}{\omega^2(1-\Gamma)\bar{\mu}\epsilon})$. Next, we proof an upper bound for σ^2 . For $k = 0, \dots, K-1$, by Theorem 1 we can upper bound σ_k^2 by

$$\begin{aligned} \sigma_k^2 & \leq 2\epsilon_{\text{approx}}^2 + 2\mathbb{E}\|\bar{\mathcal{Q}}^{\pi_k, \zeta_k} - \Psi\theta_{\pi_k}^*\|^2 \leq 2\epsilon_{\text{approx}}^2 + 2\mathbb{E}\|\bar{\mathcal{Q}}^{\pi_k, \zeta_k} - \Psi\theta_{\pi_k}^*\|_{\infty}^2 / \lambda_{\min}(\tilde{D}^{\pi_k}) \\ & \leq 2\epsilon_{\text{approx}}^2 + \mathcal{O}\left(\frac{\|\theta_{\pi_k}^*\|_2^2}{\lambda_{\min}(\tilde{D}^{\pi_k})}\right) \leq 2\epsilon_{\text{approx}}^2 + \mathcal{O}\left(\max_{\pi} \frac{\|\theta_{\pi}^*\|_2^2}{\lambda_{\min}(\tilde{D}^{\pi})}\right), \end{aligned}$$

as desired. \square

To the best of our knowledge, both sample complexities in Corollary 2 and 3 appear to be new in the AMDP literature. The former result extends the result of the generative model to the Markovian noise model, while the latter result further exploits the ergodicity structure as well as the impact of the strongly convex regularizer, thus providing an improved $\tilde{O}(\epsilon^{-1})$ convergence guarantee in terms of the KL divergence between the output solution and the optimal solution.

5 Numerical experiments

In this section, we report numerical experiments for our proposed algorithms, i.e., VRTD, EVRTD and SPMD methods. To generate a comprehensive performance profile, we conduct experiments on both randomly generated AMDPs and the popular evaluation benchmark – OpenAI Gym (Brockman et al., 2016).

5.1 Randomly generated MDPs

We consider randomly generated MDPs with state space $|\mathcal{S}| = 400, 600, 800$ and action space $|\mathcal{A}| = 15$. The transition kernel \mathbf{P} is generated by first constructing a tensor of size $\mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}| \times |\mathcal{S}|}$, where each entry is a random number, then taking softmax normalization to ensure $\sum_{s' \in \mathcal{S}} \mathbf{P}(s'|s, a) = 1$ and $\mathbf{P}(s'|s, a) > 0$. For a fixed state s , the cost function $c(s, \cdot)$ is a constant across all the actions. We randomly pick 95 percent of the states to have zero cost, while the cost of the other 5 percent of the states are generated randomly. In both cases, random numbers are uniformly taken in $[0, 10]$. Clearly, under this construction, the Markov chain induced by any feasible policy is irreducible and aperiodic. Moreover, our random construction leads to a unique and deterministic optimal policy almost surely, which is desirable for corroborating the advantages of the EVRTD critic solver.

Using the constructed MDP instances, we test the performance of the SPMD method with EVRTD as the critic (namely SPMD-EVRTD), and compare its performance against the SPMD method with VRTD as the critic (namely SPMD-VRTD). Figure 1 plots the average cost function of the current policies against the number of iterations of the SPMD method.




Figure 1. Comparing the performance of SPMD method with different critic solvers. For the red curves, SPMD takes VRTD method as the critic solver. For the blue curves, SPMD takes EVRTD as the critic solver. In the iterations marked by crosses, EVRTD implements the perturbation rule (3.30); in the other iterations, EVRTD reduces to VRTD.

In all three experiments, we can see that for the blue curves, the average cost function in the later iterations decreases continuously. In comparison, it is hard for the SPMD method without the EVRTD solver to make progress in the later iterations. As a result, this set of experiments showcases that the VRTD method fails to efficiently evaluate the insufficiently random policies due

to lack of exploration, and the EVRTD method can remedy this issue to enable fast and robust convergence of the SPMD method.

5.2 OpenAI Gym

For the experiments conducted on OpenAI Gym, we choose two environments (LunarLander and MountainCar), and compare the performance of SPMD against TRPO (Schulman et al., 2015), PPO (Schulman et al., 2017), and Q-learning. Specifically, TRPO and PPO utilize neural networks for value function approximation as well as policy parametrization, and the networks are updated using stochastic gradient descent. We use state-of-the-art implementations for both TRPO and PPO in `stable_baseline3` (Raffin et al., 2021). Additionally, we incorporate undiscounted Q-learning with linear function approximation, which utilizes least-square updates for linear function estimators described in Bertsekas and Ioffe (1996).

For linear function approximation in SPMD and Q-learning, we implement radial basis function (RBF) kernels to construct the feature space. All the RBF centers are computed prior to learning, using randomly sampled state-action pairs from the environment. The policy-based algorithms (SPMD, PPO, TRPO) perform policy updates after each episode with a specified length, and evaluate its performance by sampling multiple trajectories. Note that even though TRPO and PPO are trained using discounted objectives, their performance is evaluated without discounting, thus leading to fair comparisons with our proposed SPMD method.




Figure 2. Compare SPMD method with various methods. Policies are evaluated using 10 independent trajectories in each evaluation point. We highlight running average over 50 evaluation points.

From Figure 2, it is clear that our proposed SPMD method exhibits efficient and robust performance in both experiments. On the other hand, although TRPO and PPO perform well in MountainCar, they suffer from unstable performance in LunarLander. The Q-learning approach achieves comparable policy improvement with SPMD in MountainCar, but significantly underperforms in LunarLander.

6 Conclusions

In this paper, we investigated the problem of AMDPs, making contributions to both policy evaluation and optimization. For policy evaluation, we develop the multiple trajectory method and the VRTD method for the generative model and Markovian noise model, respectively. Moreover, we develop the EVRTD method to handle insufficiently random policies in the on-policy evaluation setting. We further extend the design idea of EVRTD to DMDPs. For policy optimization, we

develop an average-reward stochastic policy mirror descent (SPMD) method and establish $\tilde{O}(\epsilon^{-2})$ overall sample complexity under both generative model (with unichain assumption) and Markovian noise model (with ergodic assumption). Moreover, we improve the sampling complexity to $\tilde{O}(\epsilon^{-1})$ for solving regularized AMDPs.

Our work leaves a few interesting directions for future studies. First, we restrict our attention to the unichain problem with mixing property (see Assumption 1). It would be interesting to see if efficient methods with finite sample guarantees can be developed to solve AMDPs with (weakly) communicating or multi-chain structures. Second, note that under the generative model, our proposed $\tilde{O}\left(\frac{t_{\text{mix}}^3 |S||\mathcal{A}|}{\epsilon^2}\right)$ sample complexity for using first-order method to solve AMDPs is $\mathcal{O}(t_{\text{mix}})$ worse compared with state-of-the-art LP methods. A similar performance gap also exists in the DMDP literature. It is an important open question to understand whether this performance gap is fundamental, if not, how to fill this performance gap.

Acknowledgments

TL and GL were supported in part by the National Science Foundation grant CCF-1909298 and Office of Naval Research grant N00014-20-1-2089.

References

- Y. Abbasi-Yadkori, P. Bartlett, K. Bhatia, N. Lazic, C. Szepesvari, and G. Weisz. Politex: Regret bounds for policy iteration using expert prediction. In *International Conference on Machine Learning*, pages 3692–3702. PMLR, 2019.
- J. Abounadi, D. P. Bertsekas, and V. S. Borkar. Learning algorithms for Markov decision processes with average cost. *SIAM Journal on Control and Optimization*, 40(3):681–698, 2001.
- A. Agarwal, S. M. Kakade, J. D. Lee, and G. Mahajan. On the theory of policy gradient methods: Optimality, approximation, and distribution shift. *Journal of Machine Learning Research*, 22(98):1–76, 2021.
- P. Auer, T. Jaksch, and R. Ortner. Near-optimal regret bounds for reinforcement learning. *Advances in neural information processing systems*, 21, 2008.
- P. L. Bartlett and A. Tewari. Regal: A regularization based algorithm for reinforcement learning in weakly communicating MDPs. *arXiv preprint arXiv:1205.2661*, 2012.
- D. P. Bertsekas and S. Ioffe. Temporal differences-based policy iteration and applications in neuro-dynamic programming. *Lab. for Info. and Decision Systems Report LIDS-P-2349*, MIT, Cambridge, MA, 14, 1996.
- D. P. Bertsekas and J. N. Tsitsiklis. Neuro-dynamic programming: an overview. In *Proceedings of 1995 34th IEEE conference on decision and control*, volume 1, pages 560–564. IEEE, 1995.
- J. Bhandari, D. Russo, and R. Singal. A finite time analysis of temporal difference learning with linear function approximation. In *Conference on learning theory*, pages 1691–1692. PMLR, 2018.

- G. Brockman, V. Cheung, L. Pettersson, J. Schneider, J. Schulman, J. Tang, and W. Zaremba. Openai gym. *arXiv preprint arXiv:1606.01540*, 2016.
- S. Cen, C. Cheng, Y. Chen, Y. Wei, and Y. Chi. Fast global convergence of natural policy gradient methods with entropy regularization. *Operations Research*, 2021.
- C. D. Dang and G. Lan. On the convergence properties of non-euclidean extragradient methods for variational inequalities with generalized monotone operators. *Computational Optimization and applications*, 60(2):277–310, 2015.
- T. Degris, M. White, and R. S. Sutton. Off-policy actor-critic. *arXiv preprint arXiv:1205.4839*, 2012.
- F. Facchinei and J. Pang. *Finite-Dimensional Variational Inequalities and Complementarity Problems, Volumes I and II*. Comprehensive Study in Mathematics. Springer-Verlag, New York, 2003.
- A. Gosavi. Reinforcement learning for long-run average cost. *European journal of operational research*, 155(3):654–674, 2004.
- D. Hsu, A. Kontorovich, D. A. Levin, Y. Peres, C. Szepesvári, and G. Wolfer. Mixing time estimation in reversible Markov chains from a single sample path. *The Annals of Applied Probability*, 29(4):2439–2480, 2019.
- Y. Jin and A. Sidford. Efficiently solving MDPs with stochastic mirror descent. In *International Conference on Machine Learning*, pages 4890–4900. PMLR, 2020.
- Y. Jin and A. Sidford. Towards tight bounds on the sample complexity of average-reward MDPs. In *International Conference on Machine Learning*, pages 5055–5064. PMLR, 2021.
- K. Khamaru, A. Pananjady, F. Ruan, M. J. Wainwright, and M. I. Jordan. Is temporal difference learning optimal? an instance-dependent analysis. *SIAM Journal on Mathematics of Data Science*, 3(4):1013–1040, 2021.
- S. Khodadadian, P. R. Jhunjhunwala, S. M. Varma, and S. T. Maguluri. On the linear convergence of natural policy gradient algorithm. *arXiv preprint arXiv:2105.01424*, 2021.
- J. Kober, J. A. Bagnell, and J. Peters. Reinforcement learning in robotics: A survey. *The International Journal of Robotics Research*, 32(11):1238–1274, 2013.
- V. Konda and J. Tsitsiklis. Actor-critic algorithms. *Advances in neural information processing systems*, 12, 1999.
- G. Kotsalis, G. Lan, and T. Li. Simple and optimal methods for stochastic variational inequalities, I: operator extrapolation. *SIAM Journal on Optimization*, 32(3):2041–2073, 2022a.
- G. Kotsalis, G. Lan, and T. Li. Simple and optimal methods for stochastic variational inequalities, II: Markovian noise and policy evaluation in reinforcement learning. *SIAM Journal on Optimization*, 32(2):1120–1155, 2022b.
- G. Lan. *First-order and stochastic optimization methods for machine learning*. Springer, 2020.

- G. Lan. Policy mirror descent for reinforcement learning: Linear convergence, new sampling complexity, and generalized problem classes. *Mathematical programming*, pages 1–48, 2022.
- G. Lan, Y. Li, and T. Zhao. Block policy mirror descent. *arXiv preprint arXiv:2201.05756*, 2022.
- N. Lazic, D. Yin, Y. Abbasi-Yadkori, and C. Szepesvari. Improved regret bound and experience replay in regularized policy iteration. In *International Conference on Machine Learning*, pages 6032–6042. PMLR, 2021.
- G. Li, Y. Wei, Y. Chi, Y. Gu, and Y. Chen. Sample complexity of asynchronous Q-learning: Sharper analysis and variance reduction. *Advances in neural information processing systems*, 33: 7031–7043, 2020.
- T. Li, G. Lan, and A. Pananjady. Accelerated and instance-optimal policy evaluation with linear function approximation. *arXiv preprint arXiv:2112.13109*, 2021.
- Y. Li, T. Zhao, and G. Lan. Homotopic policy mirror descent: Policy convergence, implicit regularization, and improved sample complexity. *arXiv preprint arXiv:2201.09457*, 2022.
- R. Montenegro, P. Tetali, et al. Mathematical aspects of mixing times in Markov chains. *Foundations and Trends in Theoretical Computer Science*, 1(3):237–354, 2006.
- W. Mou, A. Pananjady, and M. J. Wainwright. Optimal oracle inequalities for solving projected fixed-point equations. *arXiv preprint arXiv:2012.05299*, 2020.
- W. Mou, K. Khamaru, M. J. Wainwright, P. L. Bartlett, and M. I. Jordan. Optimal variance-reduced stochastic approximation in banach spaces. *arXiv preprint arXiv:2201.08518*, 2022.
- Y. Ouyang, M. Gagrani, A. Nayyar, and R. Jain. Learning unknown Markov decision processes: A thompson sampling approach. *Advances in neural information processing systems*, 30, 2017.
- M. L. Puterman. *Markov decision processes: Discrete stochastic dynamic programming*. John Wiley & Sons, 2014.
- A. Raffin, A. Hill, A. Gleave, A. Kanervisto, M. Ernestus, and N. Dormann. Stable-baselines3: Reliable reinforcement learning implementations. *Journal of Machine Learning Research*, 22 (268):1–8, 2021.
- J. Schulman, S. Levine, P. Abbeel, M. Jordan, and P. Moritz. Trust region policy optimization. In *International conference on machine learning*, pages 1889–1897. PMLR, 2015.
- J. Schulman, F. Wolski, P. Dhariwal, A. Radford, and O. Klimov. Proximal policy optimization algorithms. *arXiv preprint arXiv:1707.06347*, 2017.
- L. Shani, Y. Efroni, and S. Mannor. Adaptive trust region policy optimization: Global convergence and faster rates for regularized MDPs. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 34, pages 5668–5675, 2020.
- J. N. Tsitsiklis and B. Van Roy. Average cost temporal-difference learning. *Automatica*, 35(11): 1799–1808, 1999.

- M. J. Wainwright. Variance-reduced Q-learning is minimax optimal. *arXiv preprint arXiv:1906.04697*, 2019.
- Y. Wan, A. Naik, and R. S. Sutton. Learning and planning in average-reward Markov decision processes. In *International Conference on Machine Learning*, pages 10653–10662. PMLR, 2021.
- M. Wang. Primal-dual π learning: Sample complexity and sublinear run time for ergodic Markov decision problems. *arXiv preprint arXiv:1710.06100*, 2017.
- G. Wolfer and A. Kontorovich. Estimating the mixing time of ergodic Markov chains. In *Conference on Learning Theory*, pages 3120–3159. PMLR, 2019.
- L. Xiao. On the convergence rates of policy gradient methods. *arXiv preprint arXiv:2201.07443*, 2022.
- X. Xu, L. Zuo, and Z. Huang. Reinforcement learning algorithms with function approximation: Recent advances and applications. *Information Sciences*, 261:1–31, 2014.
- B. Yu. Rates of convergence for empirical processes of stationary mixing sequences. *The Annals of Probability*, pages 94–116, 1994.
- H. Yu and D. P. Bertsekas. Convergence results for some temporal difference methods based on least squares. *IEEE Transactions on Automatic Control*, 54(7):1515–1531, 2009.
- W. Zhan, S. Cen, B. Huang, Y. Chen, J. D. Lee, and Y. Chi. Policy mirror descent for regularized reinforcement learning: A generalized framework with linear convergence. *arXiv preprint arXiv:2105.11066*, 2021.
- S. Zhang, Y. Wan, R. S. Sutton, and S. Whiteson. Average-reward off-policy policy evaluation with function approximation. In *International Conference on Machine Learning*, pages 12578–12588. PMLR, 2021a.
- S. Zhang, Z. Zhang, and S. T. Maguluri. Finite sample analysis of average-reward TD learning and Q-learning. *Advances in Neural Information Processing Systems*, 34, 2021b.
- Y. Zhang and K. W. Ross. On-policy deep reinforcement learning for the average-reward criterion. *arXiv preprint arXiv:2106.07329*, 2021.

A Extension of Lemma 6 to DMDPs

As mentioned in Section 3.3, the EVRTD method naturally extends to the DMDP setting. To illustrate the extension, we proof the analog of Lemma 6 in the DMDP setting, then the convergence guarantees of VRTD for DMDPs in Li et al. (2021) follow.

Lemma 12. *Let $\gamma \in (0, 1)$ denote the discount factor for the underlying discounted Markov decision process. Assume $|\mathcal{S}| \geq 2$. For $s \in \mathcal{S}$, if we take*

$$\underline{\pi} \leq \frac{(1-\gamma) \min_{s \in \mathcal{S}} \nu(s)}{(|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|)(8 + (1-\gamma) \min_{s \in \mathcal{S}} \nu(s))}, \quad (\text{A.1})$$

and construct the perturbed policy $\tilde{\pi}$ as (3.30), then

$$\inf_{x \in \mathbb{S}, \|x\|_{\tilde{D}}=1} x^\top \tilde{D}(I - \gamma P)x \geq \frac{3}{4}(1 - \gamma). \quad (\text{A.2})$$

Proof. First, we utilize the Cauchy-Schwarz inequality to obtain

$$x^\top \tilde{D}(I - \gamma P)x = \|x\|_{\tilde{D}}^2 - \gamma x^\top \tilde{D}Px \geq \|x\|_{\tilde{D}}^2 - \gamma \|Px\|_{\tilde{D}} \|x\|_{\tilde{D}}. \quad (\text{A.3})$$

Towards this end, the rest of this proof focus on establishing an upper bound on $\|Px\|_{\tilde{D}}$. We write

$$\|Px\|_{\tilde{D}}^2 \leq \sum_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}} \nu(s) \tilde{\pi}(a|s) \left(\sum_{s' \in \mathcal{S}} \sum_{a' \in \mathcal{A}} P(s'|s, a) \pi(a'|s') x(s', a')^2 \right),$$

which follows from Jensen's inequality. Notice that the RHS of the above inequality is R_2 in the proof of Lemma 6. Applying the upper bound of R_2 (with β replaced by γ) in (3.39) gives us

$$\|Px\|_{\tilde{D}}^2 \leq \left(1 + \frac{\alpha(\underline{\pi})(1-\gamma)}{4}\right) \|x\|_{\tilde{D}}^2,$$

where $\alpha(\underline{\pi}) := 1 - (|\mathcal{A}| - \min_{s \in \mathcal{S}} |\mathcal{A}_s|) \underline{\pi} \leq 1$. Therefore, we have $\|Px\|_{\tilde{D}} \leq \left(1 + \frac{\alpha(\underline{\pi})(1-\gamma)}{4}\right) \|x\|_{\tilde{D}}$. Substituting this inequality into Ineq. (A.3) yields

$$x^\top \tilde{D}(I - \gamma P)x \geq \|x\|_{\tilde{D}}^2 - \gamma \left(1 + \frac{\alpha(\underline{\pi})(1-\gamma)}{4}\right) \|x\|_{\tilde{D}}^2 = \left(1 - \frac{\gamma \alpha(\underline{\pi})}{4}\right) (1 - \gamma) \|x\|_{\tilde{D}}^2 \geq \frac{3}{4}(1 - \gamma) \|x\|_{\tilde{D}}^2,$$

which completes the proof. \square

B Proofs of Section 3

We now turn to proofs of the results in Section 3.

B.1 Proof of Lemma 1

Fix a feasible policy π . By the fact of $(\nu^\pi)^\top P_\pi = (\nu^\pi)^\top$, we have that for any positive integer z ,

$$\left((P_\pi)^z - \mathbf{1}(\nu^\pi)^\top\right) \left(P_\pi - \mathbf{1}(\nu^\pi)^\top\right) = (P_\pi)^{z+1} - \mathbf{1}(\nu^\pi)^\top. \quad (\text{B.1})$$

On the other hand, for any positive integer z ,

$$\|(P_\pi)^{z+1} - \mathbf{1}(\nu^\pi)^\top\|_\infty = \left\| \left((P_\pi)^z - \mathbf{1}(\nu^\pi)^\top\right) P_\pi \right\|_\infty \leq \|(P_\pi)^z - \mathbf{1}(\nu^\pi)^\top\|_\infty. \quad (\text{B.2})$$

Noting that $k \geq t_{\text{mix}}$ and utilizing the two relationships above, we obtain that

$$\begin{aligned} \|(P_\pi)^k - \mathbf{1}(\nu^\pi)^\top\|_\infty &\stackrel{(i)}{\leq} \|(P_\pi)^{t_{\text{mix}} \cdot \lfloor k/t_{\text{mix}} \rfloor} - \mathbf{1}(\nu^\pi)^\top\|_\infty \stackrel{(ii)}{\leq} \left\| \left((P_\pi)^{t_{\text{mix}}} - \mathbf{1}(\nu^\pi)^\top \right)^{\lfloor k/t_{\text{mix}} \rfloor} \right\|_\infty \\ &\leq \left(\|(P_\pi)^{t_{\text{mix}}} - \mathbf{1}(\nu^\pi)^\top\|_\infty \right)^{\lfloor k/t_{\text{mix}} \rfloor}, \end{aligned}$$

where step (i) follows from Ineq. (B.2), and step (ii) follows from Ineq. (B.1). Next, we bound the term $\|(P_\pi)^{t_{\text{mix}}} - \mathbf{1}(\nu^\pi)^\top\|_\infty$. We have

$$\|(P_\pi)^{t_{\text{mix}}} - \mathbf{1}(\nu^\pi)^\top\|_\infty = \max_{s \in \mathcal{S}} \sum_{s' \in \mathcal{S}} |(P_\pi)^{t_{\text{mix}}}(s, s') - \nu^\pi(s')| \leq \max_{q \in \Delta_{|\mathcal{S}|}} \|((P_\pi)^{t_{\text{mix}}})^\top q - \nu^\pi\|_1 \stackrel{(i)}{\leq} \frac{1}{2},$$

where (i) follows from Assumption 1. Combining two relationships above yields the desired result.

B.2 Supporting lemmas for Theorem 1 and 2

We begin by establishing a few useful lemmas which are frequently used in the proof of the main theorems in Section 3. As a consequence of Lemma 3, we establish the following supporting lemma.

Lemma 13. *For any $\theta \in \mathbb{R}^d$, we have that the matrix $I_d - M$ is invertible and*

$$\|(I_d - M)^{-1}\theta\|_2 \leq \frac{1}{1-\beta}\|\theta\|_2. \quad (\text{B.3})$$

Proof. By Cauchy-Schwarz inequality, we have

$$\|(I_d - M)\theta\|_2 \|\theta\|_2 \geq \theta^\top \Phi^\top D(I - P)\Phi\theta \stackrel{(i)}{\geq} (1 - \beta)\|\Phi\theta\|_D^2 \stackrel{(ii)}{=} (1 - \beta)\|\theta\|_2^2,$$

where step (i) follows from Lemma 3, step (ii) follows from $\Phi^\top D\Phi = I_d$. As a result, we obtain that $\|\theta\|_2 \leq \frac{1}{1-\beta}\|(I_d - M)\theta\|_2$. Meanwhile, by Lemma 3, it is clear that $I_d - \frac{1}{2}(M + M^\top) = \frac{1}{2}(\Phi^\top D(I - P)\Phi + \Phi^\top (I - P)^\top D\Phi)$ is positive definite, thus $I_d - M$ is invertible. \square

The following lemma characterizes the Lipschitz continuity of the operator g .

Lemma 14. *For any $\theta, \theta' \in \mathbb{R}^d$, $\rho \in \mathbb{R}$, we have*

$$\|g(\theta, \rho) - g(\theta', \rho)\|_2 \leq 2\|\Psi\theta - \Psi\theta'\|_D \leq 2\|\theta - \theta'\|_2.$$

Proof. Assume $s \sim \nu, a \sim \pi(\cdot|s), s' \sim P(\cdot|s, a), a' \sim \pi(\cdot|s')$. Let $\Upsilon := \langle \psi(s, a), \theta - \theta' \rangle$ and $\Upsilon' := \langle \psi(s', a'), \theta - \theta' \rangle$. We have

$$\begin{aligned} \|g(\theta, \rho) - g(\theta', \rho)\|_2 &= \|\Psi^\top D(I - P)\Psi(\theta - \theta')\|_2 = \|\mathbb{E}[\psi(s, a)(\Upsilon - \Upsilon')]\|_2 \\ &\stackrel{(i)}{\leq} \mathbb{E}[|\Upsilon - \Upsilon'|] \leq \sqrt{\mathbb{E}[\Upsilon^2]} + \sqrt{\mathbb{E}[(\Upsilon')^2]} \stackrel{(ii)}{=} 2\|\Psi\theta - \Psi\theta'\|_D \stackrel{(iii)}{\leq} 2\|\theta - \theta'\|_2, \end{aligned}$$

where step (i) follows from Young's inequality and $\|\psi(s, a)\|_2 \leq 1$, and (ii) follows from the fact that Υ and Υ' have the same distribution and $\mathbb{E}[\Upsilon^2] = \|\Psi\theta - \Psi\theta'\|_D^2$, step (iii) follows from Cauchy-Schwarz inequality and $\|\psi(s, a)\|_2 \leq 1$. \square

Next we establish a few supporting lemmas that characterize the properties of the stochastic operator defined above. Lemma 15 and 16 upper bound the variance of the stochastic operator.

Lemma 15. For every $\theta, \theta' \in \mathbb{R}^d$, $\rho, \rho' \in \mathbb{R}$, and $\xi := \{(s, a), (s', a'), c^\pi(s, a)\}$ where $s \sim \nu^\pi, a \sim \pi(\cdot|s), s' \sim P(\cdot|s, a)$, we have

$$\mathbb{E} \|\tilde{g}(\theta, \rho, \xi) - \tilde{g}(\theta', \rho', \xi) - (g(\theta, \rho) - g(\theta', \rho'))\|_2^2 \leq 8\|\Psi\theta - \Psi\theta'\|_D^2 + 2|\rho - \rho'|^2. \quad (\text{B.4})$$

Proof. First, it is easy to see that given $s \sim \nu^\pi$, we have $\mathbb{E}[\tilde{g}(\theta, \rho, \xi) - \tilde{g}(\theta', \rho', \xi)] = g(\theta, \rho) - g(\theta', \rho')$. For notation brevity, we let $\Upsilon := \langle \psi(s, a), \theta - \theta' \rangle$ and $\Upsilon' := \langle \psi(s', a'), \theta - \theta' \rangle$. Since $s \in \nu^\pi$, Υ and Υ' have the same distribution, we can write

$$\begin{aligned} & \mathbb{E} \|\tilde{g}(\theta, \rho, \xi) - \tilde{g}(\theta', \rho', \xi) - (g(\theta, \rho) - g(\theta', \rho'))\|_2^2 \\ & \leq \mathbb{E} \|\tilde{g}(\theta, \rho, \xi) - \tilde{g}(\theta', \rho', \xi)\|_2^2 = \mathbb{E} \|(\Upsilon - \Upsilon' + \rho - \rho')\psi(s, a)\|_2^2 \\ & \stackrel{(i)}{\leq} 2\mathbb{E} \|(\Upsilon - \Upsilon')\psi(s, a)\|_2^2 + 2\mathbb{E} \|(\rho - \rho')\psi(s, a)\|_2^2 \\ & \stackrel{(ii)}{\leq} 4\mathbb{E}[\Upsilon^2] + 4\mathbb{E}[(\Upsilon')^2] + 2(\rho - \rho')^2 \stackrel{(iii)}{\leq} 8\|\Psi\theta - \Psi\theta'\|_D^2 + 2(\rho - \rho')^2, \end{aligned}$$

where step (i) follows from Young's inequality, step (ii) follows from the fact that $\|\psi(s, a)\|_2 \leq 1$ and Young's inequality, and step (iii) follows from the fact that Υ and Υ' have the same distribution and $\mathbb{E}[\Upsilon^2] = \|\Psi(\theta - \theta')\|_D^2$. \square

Lemma 16 (see Lemma 12 of Li et al. (2021)). For $t \in \mathbb{Z}_+$, $\theta, \theta' \in \mathbb{R}^d$, $\rho, \rho' \in \mathbb{R}$, and $\tau \in \mathbb{Z}_+$ such that

$$C_P \cdot \varrho^\tau \leq \min_{s \in \mathcal{S}, a \in \mathcal{A}} \nu(s)\pi(a|s), \quad (\text{B.5})$$

we have with probability 1,

$$\mathbb{E} \|\tilde{g}(\theta, \rho, \xi_{t+\tau}) - \tilde{g}(\theta', \rho', \xi_{t+\tau}) - (g(\theta, \rho) - g(\theta', \rho'))\|_2^2 | \mathcal{F}_t \leq 16\|\Psi\theta - \Psi\theta'\|_D^2 + 4|\rho - \rho'|^2. \quad (\text{B.6})$$

Recall the definition of \hat{g} , N_k , N'_k in Algorithm 2 and $C_{\max} := \max\{C_P, C_M, C_R\}$.

Lemma 17. For every $\theta, \theta' \in \mathbb{R}^d$, $\rho, \rho' \in \mathbb{R}$, $\tau' \in \mathbb{Z}_+$ satisfied Ineq. (B.5) and $\varrho^{\tau'} \leq \frac{1}{4C_{\max}+1}$, then

$$\mathbb{E} \|\hat{g}(\theta, \rho) - \hat{g}(\theta', \rho') - g(\theta, \rho) + g(\theta', \rho')\|_2^2 \leq \frac{20}{N_k} \|\Psi\theta - \Psi\theta'\|_\Pi^2 + \frac{8}{N_k} |\rho - \rho'|^2. \quad (\text{B.7})$$

Proof. We write

$$\mathbb{E} \|\hat{g}(\theta, \rho) - \hat{g}(\theta', \rho') - g(\theta, \rho) + g(\theta', \rho')\|_2^2 = \frac{1}{N_k^2} \sum_{i=1}^{N_k} \mathbb{E} \|\Lambda_i\|_2^2 + \frac{2}{N_k^2} \sum_{i=1}^{N_k-1} \sum_{j>i} \mathbb{E} \langle \Lambda_i, \Lambda_j \rangle,$$

where $\Lambda_i = \tilde{g}(\theta, \rho, \xi_i^k(\tau')) - \tilde{g}(\theta', \rho', \xi_i^k(\tau')) - g(\theta, \rho) + g(\theta', \rho')$. Invoking Lemma 17 we can bound $\frac{1}{N_k^2} \sum_{i=1}^{N_k} \mathbb{E} \|\Lambda_i\|_2^2 \leq \frac{16}{N_k} \|\Psi\theta - \Psi\theta'\|_\Pi^2 + \frac{4}{N_k} |\rho - \rho'|^2$. On the other hand, we have

$$\begin{aligned} \frac{2}{N_k^2} \sum_{i=1}^{N_k-1} \sum_{j>i} \mathbb{E} \langle \Lambda_i, \Lambda_j \rangle & \stackrel{(i)}{\leq} \frac{2}{N_k^2} \sum_{i=1}^{N_k-1} \sum_{j>i} \mathbb{E} [C_{\max} \rho^{(j-i)\tau'} \|\Lambda_i\|_2 (\|\Psi\theta - \Psi\theta'\|_D + |\rho - \rho'|)] \\ & \stackrel{(ii)}{\leq} \frac{16}{N_k^2} \sum_{i=1}^{N_k-1} \sum_{j>i} C_{\max} \rho^{(j-i)\tau'} (\|\Psi\theta - \Psi\theta'\|_D^2 + |\rho - \rho'|^2) \\ & \leq \frac{16C_{\max}\rho^{\tau'}}{N_k(1-\rho^{\tau'})} (\|\Psi\theta - \Psi\theta'\|_D^2 + |\rho - \rho'|^2) \stackrel{(iii)}{\leq} \frac{4}{N_k} (\|\Psi\theta - \Psi\theta'\|_D^2 + |\rho - \rho'|^2), \end{aligned}$$

where (i) follows from Cauchy-Schwarz inequality and Lemma 5, (ii) follows from Young's inequality and Lemma 17, (iii) follows from $\varrho^{\tau'} \leq \frac{1}{4C_{\max}+1}$. Combining the upper bounds yields the result. \square

Lemma 18. For $\tilde{\rho}$ defined in step 4 of Algorithm 2, if τ' satisfies $\varrho^{\tau'} \leq \frac{1}{4C_{\max}+1}$, then with probability 1,

$$\mathbb{E}[(\tilde{\rho} - \rho^*)^2] \leq \frac{5\bar{c}^2}{N'_k}. \quad (\text{B.8})$$

Proof. Recall the definition of $(\tilde{s}_i, \tilde{a}_i)$ in Algorithm 2. Define the vectors $\mathcal{D}, \mathcal{D}_j^i \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$ as $\mathcal{D} := [\nu^\pi(s) \cdot \pi(a|s)]$ and $\mathcal{D}_j^i := [\mathbf{P}(\tilde{s}_j = s, \tilde{a}_j = a | \tilde{s}_i, \tilde{a}_i)]$ where $s \in \mathcal{S}, a \in \mathcal{A}$. We have

$$\begin{aligned} \mathbb{E}[(\tilde{\rho} - \rho^*)^2] &= \frac{1}{N'^2_k} \sum_{i=1}^{N'_k} \mathbb{E}|c(\tilde{s}_i, \tilde{a}_i) - \rho^*|^2 + \frac{2}{N'^2_k} \sum_{i=1}^{N'_k-1} \sum_{j=i+1}^{N'_k} \mathbb{E}[(c(\tilde{s}_i, \tilde{a}_i) - \rho^*) \mathbb{E}[c(\tilde{s}_j, \tilde{a}_j) - \rho^* | \mathcal{F}_i]] \\ &\stackrel{(i)}{\leq} \frac{1}{N'^2_k} \sum_{i=1}^{N'_k} \mathbb{E}|c(\tilde{s}_i, \tilde{a}_i) - \rho^*|^2 + \frac{2}{N'^2_k} \sum_{i=1}^{N'_k-1} \sum_{j=i+1}^{N'_k} \bar{c} \cdot \mathbb{E}[(c(\tilde{s}_i, \tilde{a}_i) - \rho^*) \|\mathcal{D}_j^i - \mathcal{D}\|_1] \\ &\stackrel{(ii)}{\leq} \frac{4\bar{c}^2}{N'_k} + \frac{2}{N'^2_k} \sum_{i=1}^{N'_k-1} \sum_{j=i+1}^{N'_k} 2C_P \cdot \varrho^{\tau'(j-i)} \bar{c}^2 \leq \frac{4\bar{c}^2}{N'_k} + \frac{4C_P \bar{c}^2 \varrho^{\tau'}}{N'_k(1-\varrho^{\tau'})} \leq \frac{5\bar{c}^2}{N'_k}, \end{aligned}$$

as claimed, where step (i) follows from Cauchy-Schwarz inequality, and step (ii) follows from Lemma 2 and $\|c\|_\infty \leq \bar{c}$. \square

B.3 Proof of Theorem 1

We first focus on the progress of a single epoch $k \in [K]$. Let $\underline{\theta}$ satisfy $g(\underline{\theta}, \tilde{\rho}) - g(\tilde{\theta}, \tilde{\rho}) + \hat{g}(\tilde{\theta}, \tilde{\rho}) = 0$. The vector $\underline{\theta}$ can be interpreted as the fixed point that the VRTD algorithm is trying to solve inside the given epoch. The following lemma establish an upper bound on $\|\Psi\underline{\theta} - \Psi\theta^*\|_D^2$.

Lemma 19. Consider a single epoch with index $k \in [K]$. We have

$$\begin{aligned} \mathbb{E}[\|\Psi\underline{\theta} - \Psi\theta^*\|_D^2] &\leq \frac{9}{N_k} \text{trace} \left((I_d - M)^{-1} \bar{\Sigma} (I_d - M)^{-\top} \right) + \frac{3}{2N_k(1-\beta)^2\mu} \mathbb{E}\|\Psi\theta^* - \bar{Q}^\pi\|_D^2 \\ &\quad + \frac{60}{N_k(1-\beta)^2\mu} \mathbb{E}\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2 + \frac{15\bar{c}^2}{N'_k(1-\beta)^2\mu} + \frac{120\bar{c}^2}{N_k N'_k(1-\beta)^2\mu}. \end{aligned} \quad (\text{B.9})$$

Proof of the lemma is postponed to Section B.3.1. Given the upper bound on $\mathbb{E}[\|\Psi\underline{\theta} - \Psi\theta^*\|_D^2]$, we are able to derive the progress in a single epoch $k \in [K]$, which is characterized in the following proposition.

Proposition 4. Consider a single epoch with index $k \in [K]$. Suppose that the parameters η , N_k and T satisfy

$$\eta \leq \frac{1-\gamma}{845}, \quad T \geq \frac{64}{\mu(1-\gamma)\eta}, \quad \text{and} \quad N_k \geq \frac{1160}{\mu(1-\gamma)^2}. \quad (\text{B.10})$$

Set the output of this epoch to be $\hat{\theta}_k := \frac{\sum_{t=1}^T \theta_t}{T}$. If τ and τ' are chosen to satisfy

$$\varrho^\tau \leq \frac{\sqrt{\mu}\eta}{C_M} \quad \text{and} \quad \varrho^{\tau'} \leq \min\left\{\frac{\min_{s \in \mathcal{S}, a \in \mathcal{A}} \nu(s) \pi(a|s)}{C_P}, \frac{1}{4C_{\max}+1}\right\}, \quad (\text{B.11})$$

then we have

$$\begin{aligned} \mathbb{E}[\|\Psi\hat{\theta}_k - \Psi\theta^*\|_D^2] &\leq \frac{\mathbb{E}[\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2]}{2} + \frac{22\text{trace}\{(I_d - M)^{-1} \bar{\Sigma} (I_d - M)^{-\top}\}}{N_k} \\ &\quad + \frac{4\mathbb{E}[\|(\bar{Q}^* - \Psi\theta^*)\|_D^2]}{\mu(1-\beta)^2 N_k} + \frac{36\bar{c}^2}{N'_k(1-\beta)^2\mu} + \frac{285\bar{c}^2}{N_k N'_k(1-\beta)^2\mu}. \end{aligned} \quad (\text{B.12})$$

Proof of the proposition is postponed to Section B.3.2. Now we take Proposition 4 as given and finish the proof of Theorem 1. Recall the definition $W_1 := 22\text{trace}\{(I_d - M)^{-1}\bar{\Sigma}(I_d - M)^{-\top}\} + \frac{4}{\mu(1-\beta)^2}\mathbb{E}[\|(\bar{Q}^* - \Psi\theta^*)\|_D^2]$. Recursively using Ineq. (B.12) yields

$$\begin{aligned}\mathbb{E}\|\Psi\hat{\theta}_K - \Psi\theta^*\|_D^2 &\leq \frac{1}{2^K}\|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \sum_{k=1}^K \frac{1}{2^{K-k}} \left(\frac{W_1}{N_k} + \frac{36\bar{c}^2}{N'_k(1-\beta)^2\mu} + \frac{285\bar{c}^2}{N_k N'_k(1-\beta)^2\mu} \right) \\ &\stackrel{(i)}{\leq} \frac{1}{2^K}\|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \sum_{k=1}^K \left(\frac{2}{3} \right)^{K-k} \left(\frac{W_1}{N} + \frac{36\bar{c}^2}{N'(1-\beta)^2\mu} + \frac{\bar{c}^2}{4N'} \right) \\ &\leq \frac{1}{2^K}\|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \frac{3W_1}{N} + \frac{108\bar{c}^2}{N'(1-\beta)^2\mu} + \frac{3\bar{c}^2}{4N'},\end{aligned}\tag{B.13}$$

as desired, where step (i) follows from $N_k \geq (\frac{3}{4})^{K-k}N$, $N'_k \geq (\frac{3}{4})^{K-k}N'$ and $N_k \geq \frac{1160}{\mu(1-\gamma)^2}$.

B.3.1 Proof of Lemma 19

For notational simplicity, we define $\hat{\delta}_g(\theta, \rho) := \hat{g}(\theta, \rho) - g(\theta, \rho)$ for $\theta \in \mathbb{R}^d, \rho \in \mathbb{R}$. By the definition of $g(\cdot)$ and $\underline{\theta}$, we obtain

$$\begin{aligned}\Psi^\top D\Psi(\theta^* - \underline{\theta}) &= \Psi^\top D(c - \rho^*\mathbf{1}) + \Psi^\top DP\Psi\theta^* - (\Psi^\top D(c - \tilde{\rho}\mathbf{1}) + \Psi^\top DP\Psi\underline{\theta} - \hat{\delta}_g(\tilde{\theta}, \tilde{\rho})) \\ &= \Psi^\top DP\Psi(\theta^* - \underline{\theta}) + \Psi^\top D\mathbf{1}(\tilde{\rho} - \rho^*) + \hat{\delta}_g(\tilde{\theta}, \tilde{\rho}).\end{aligned}$$

Invoking the fact that $\Phi = \Psi B^{-\frac{1}{2}}$ and $I_d - M$ is invertible, we have that

$$B^{\frac{1}{2}}(\theta^* - \underline{\theta}) = (I_d - M)^{-1}B^{-\frac{1}{2}}(\Psi^\top D\mathbf{1}(\tilde{\rho} - \rho^*) + \hat{\delta}_g(\tilde{\theta}, \tilde{\rho})).\tag{B.14}$$

Therefore, further utilizing the relationship $\Phi^\top D\Phi = I_d$ and $\Phi = \Psi B^{-\frac{1}{2}}$ yields

$$\mathbb{E}\|\Psi\theta^* - \Psi\underline{\theta}\|_D^2 = \mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}(\Psi^\top D\mathbf{1}(\tilde{\rho} - \rho^*) + \hat{\delta}_g(\tilde{\theta}, \tilde{\rho}))\|_2^2.\tag{B.15}$$

By applying Young's inequality we obtain that

$$\begin{aligned}\mathbb{E}\|\Psi\theta^* - \Psi\underline{\theta}\|_D^2 &\leq 3\mathbb{E}|\tilde{\rho} - \rho^*|^2 \cdot \|(I_d - M)^{-1}B^{-\frac{1}{2}}\Psi^\top D\mathbf{1}\|_2^2 \\ &\quad + 3\mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}(\hat{\delta}_g(\tilde{\theta}, \tilde{\rho}) - \hat{\delta}_g(\theta^*, \rho^*))\|_2^2 + 3\mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}\hat{\delta}_g(\theta^*, \rho^*)\|_2^2 \\ &\stackrel{(i)}{\leq} \frac{15\bar{c}^2}{N'_k(1-\beta)^2\mu} + 3\mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}(\hat{\delta}_g(\tilde{\theta}, \tilde{\rho}) - \hat{\delta}_g(\theta^*, \rho^*))\|_2^2 \\ &\quad + 3\mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}\hat{\delta}_g(\theta^*, \rho^*)\|_2^2 \\ &\stackrel{(ii)}{\leq} \frac{15\bar{c}^2}{N'_k(1-\beta)^2\mu} + \frac{120\bar{c}^2}{N_k N'_k(1-\beta)^2\mu} + \frac{60\mathbb{E}\|\Psi\theta^* - \Psi\underline{\theta}\|_D^2}{N_k(1-\beta)^2\mu} + 3\mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}\hat{\delta}_g(\theta^*, \rho^*)\|_2^2.\end{aligned}\tag{B.16}$$

where (i) follows from Jensen's inequality $\|\Psi^\top D\mathbf{1}\|_2^2 \leq \sum_{s \in \mathcal{S}, a \in \mathcal{A}} \nu(s)\pi(a|s)\|\psi(s, a)\|_2^2 \leq 1$, Lemma 13 and 18, step (ii) follows from Lemma 13, 17 and 18.

By Lemma 15 of Li et al. (2021), we have that

$$\begin{aligned}\mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}\hat{\delta}_g(\theta^*, \rho^*)\|_2^2 \\ \leq \left(\frac{2}{N_k} + \frac{2C_M\varrho^{\tau'}}{(1-\varrho^{\tau'})N_k} \right) \text{trace}\{(I_d - M)^{-1}\bar{\Sigma}(I_d - M)^{-\top}\} + \frac{C_M\varrho^{\tau'}}{\alpha(1-\varrho^{\tau'})\mu(1-\beta)^2N_k} \|\Psi\theta^* - \bar{Q}^*\|_D^2.\end{aligned}$$

Invoking the condition that $\varrho^{\tau'} \leq \frac{1}{2C_M+1}$, we obtain

$$\mathbb{E}\|(I_d - M)^{-1}B^{-\frac{1}{2}}\widehat{\delta}_g(\theta^*, \rho^*)\|_2^2 \leq \frac{3}{N_k}\text{trace}\{(I_d - M)^{-1}\bar{\Sigma}(I_d - M)^{-\top}\} + \frac{1}{2\mu(1-\beta)^2N_k}\|\Psi\theta^* - \bar{Q}^*\|_D^2.$$

Putting together the bounds proved above and Ineq. (B.16) yields the desired result.

B.3.2 Proof of Proposition 4

We first provide an upper bound on the term $\mathbb{E}\|\Psi\widehat{\theta}_k - \Psi\theta\|_D^2$. For simplicity, we define $\widetilde{\delta}_g(\theta, \rho, \xi_t) := \widetilde{g}(\theta, \rho, \xi_t) - g(\theta, \rho)$ for $\theta \in \mathbb{R}^d$ and $\rho \in \mathbb{R}$. By the updates of the VRTD algorithm, we have

$$\theta_{t+1} - \underline{\theta} = \theta_t - \eta(\widetilde{g}(\theta_t, \widetilde{\rho}, \xi_t) - \widetilde{g}(\widetilde{\theta}, \widetilde{\rho}, \xi_t) + \widehat{g}(\widetilde{\theta}, \widetilde{\rho})) - \underline{\theta}.$$

Invoking the fact that $\widehat{g}(\widetilde{\theta}, \widetilde{\rho}) = g(\widetilde{\theta}, \widetilde{\rho}) - g(\underline{\theta}, \widetilde{\rho})$, we obtain

$$\theta_{t+1} - \underline{\theta} = \theta_t - \underline{\theta} - \eta(g(\theta_t, \widetilde{\rho}) - g(\underline{\theta}, \widetilde{\rho})) + \eta(\widetilde{\delta}_g(\widetilde{\theta}, \widetilde{\rho}, \xi_t) - \widetilde{\delta}_g(\theta_t, \widetilde{\rho}, \xi_t)). \quad (\text{B.17})$$

By taking $\|\cdot\|_2^2$ and expectation on both sides of the above equation, we have

$$\begin{aligned} & \mathbb{E}\|\theta_{t+1} - \underline{\theta}\|_2^2 \\ & \leq \mathbb{E}\|\theta_t - \underline{\theta}\|_2^2 - 2\eta\langle g(\theta_t, \widetilde{\rho}) - g(\underline{\theta}, \widetilde{\rho}), \theta_t - \underline{\theta} \rangle + 2\eta\mathbb{E}\langle \widetilde{\delta}_g(\widetilde{\theta}, \widetilde{\rho}, \xi_t) - \widetilde{\delta}_g(\theta_t, \widetilde{\rho}, \xi_t), \theta_t - \underline{\theta} \rangle \\ & \quad + 2\eta^2\mathbb{E}\|g(\theta_t, \widetilde{\rho}) - g(\underline{\theta}, \widetilde{\rho})\|_2^2 + 2\eta^2\mathbb{E}\|\widetilde{\delta}_g(\widetilde{\theta}, \widetilde{\rho}, \xi_t) - \widetilde{\delta}_g(\theta_t, \widetilde{\rho}, \xi_t)\|_2^2 \\ & \stackrel{(ii)}{\leq} \mathbb{E}\|\theta_t - \underline{\theta}\|_2^2 + (8\eta^2 - 2\eta(1 - \beta))\mathbb{E}\|\Psi\theta_t - \Psi\underline{\theta}\|_D^2 + 32\eta^2\mathbb{E}\|\Psi\widetilde{\theta} - \Psi\theta_t\|_D^2 \\ & \quad + 2\eta\langle \widetilde{\delta}_g(\widetilde{\theta}, \widetilde{\rho}, \xi_t) - \widetilde{\delta}_g(\theta_t, \widetilde{\rho}, \xi_t), \theta_t - \underline{\theta} \rangle \\ & \stackrel{(iii)}{\leq} \mathbb{E}\|\theta_t - \underline{\theta}\|_2^2 + (8\eta^2 - 2\eta(1 - \beta) + \eta C_M \varrho^\tau / \sqrt{\mu})\mathbb{E}\|\Psi\theta_t - \Psi\underline{\theta}\|_D^2 \\ & \quad + (32\eta^2 + \eta C_M \varrho^\tau / \sqrt{\mu})\mathbb{E}\|\Psi\widetilde{\theta} - \Psi\theta_t\|_D^2, \end{aligned} \quad (\text{B.18})$$

where step (ii) follows from using Lemma 3 to bound the second term, Lemma 14 to bound the forth term and Lemma 15 to bound the last term. Step (iii) follows from

$$\begin{aligned} & \langle \widetilde{\delta}_g(\widetilde{\theta}, \widetilde{\rho}, \xi_t) - \widetilde{\delta}_g(\theta_t, \widetilde{\rho}, \xi_t), \theta_t - \underline{\theta} \rangle \\ & \leq \|\widetilde{\delta}_g(\widetilde{\theta}, \widetilde{\rho}, \xi_t) - \widetilde{\delta}_g(\theta_t, \widetilde{\rho}, \xi_t)\|_2 \|\theta_t - \underline{\theta}\|_2 \leq C_M \varrho^\tau \|\Psi\theta_t - \Psi\widetilde{\theta}\|_D \cdot \frac{1}{\sqrt{\mu}} \|\Psi\theta_t - \Psi\underline{\theta}\|_D \\ & \leq \frac{C_M \varrho^\tau}{2\sqrt{\mu}} (\|\Psi\theta_t - \Psi\widetilde{\theta}\|_D^2 + \|\Psi\theta_t - \Psi\underline{\theta}\|_D^2), \end{aligned}$$

where the first inequality follows from Cauchy-Schwarz inequality, the second inequality follows from Lemma 5 and $\|\theta\|_2 \leq (1/\sqrt{\mu})\|\Psi\theta\|_D$, and the third inequality follows from Young's inequality. Further using Young's inequality $\|\Psi\widetilde{\theta} - \Psi\theta_t\|_D^2 \leq 2\|\Psi\theta_t - \Psi\underline{\theta}\|_D^2 + 2\|\Psi\widetilde{\theta} - \Psi\underline{\theta}\|_D^2$ to Ineq. (B.18) yields

$$\begin{aligned} \mathbb{E}\|\theta_{t+1} - \underline{\theta}\|_2^2 & \leq \mathbb{E}\|\theta_t - \underline{\theta}\|_2^2 + (8\eta^2 + 64\eta^2 - 2\eta(1 - \beta) + 3\eta C_M \varrho^\tau / \sqrt{\mu})\mathbb{E}\|\Psi\theta_t - \Psi\underline{\theta}\|_D^2 \\ & \quad + (64\eta^2 + 2\eta C_M \varrho^\tau / \sqrt{\mu})\mathbb{E}\|\Psi\underline{\theta} - \Psi\widetilde{\theta}\|_D^2 \\ & \stackrel{(i)}{\leq} \mathbb{E}\|\theta_t - \underline{\theta}\|_2^2 - \eta(1 - \beta)\mathbb{E}\|\Psi\theta_t - \Psi\underline{\theta}\|_D^2 + 66\eta^2\mathbb{E}\|\Psi\underline{\theta} - \Psi\widetilde{\theta}\|_D^2, \end{aligned} \quad (\text{B.19})$$

where step (i) follows from the assumption $\eta \leq \frac{1-\beta}{75}$ and $\varrho^\tau \leq \frac{\sqrt{\mu}\eta}{C_M}$. By taking a telescopic sum of Ineq. (B.18) from $t = 1$ to T , we obtain

$$\sum_{t=1}^T \eta(1-\beta) \mathbb{E}[\|\Psi\theta_t - \Psi\theta\|_D^2] + \mathbb{E}[\|\theta_{T+1} - \theta\|_2^2] \leq \mathbb{E}[\|\theta_1 - \theta\|_2^2] + 66T\eta^2 \|\Psi\theta - \Psi\tilde{\theta}\|_D^2.$$

Utilizing the inequality $\|\theta\|_2^2 \leq \frac{1}{\mu} \|B^{\frac{1}{2}}\theta\|_2^2 = \frac{1}{\mu} \|\Psi\theta\|_D^2$, we then have

$$\sum_{t=1}^T \eta(1-\beta) \mathbb{E}[\|\Psi\theta_t - \Psi\theta\|_D^2] \leq \frac{1}{\mu} \mathbb{E}[\|\Psi\theta_1 - \Psi\theta\|_D^2] + 66T\eta^2 \|\Psi\theta - \Psi\tilde{\theta}\|_D^2.$$

Noting in addition that $\theta_1 = \tilde{\theta}$ then yields

$$\begin{aligned} \sum_{t=1}^T \mathbb{E}[\|\Psi\theta_t - \Psi\theta\|_D^2]/T &\leq \frac{1}{T\eta(1-\beta)} \left[\frac{1}{\mu} \mathbb{E}[\|\Psi\theta_1 - \Psi\theta\|_D^2] + 66T\eta^2 \|\Psi\theta - \Psi\tilde{\theta}\|_D^2 \right] \\ &\leq \frac{2/\mu + 132T\eta^2}{T\eta(1-\beta)} (\mathbb{E}[\|\Psi\theta - \Psi\theta^*\|_D^2] + \mathbb{E}[\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2]). \end{aligned} \quad (\text{B.20})$$

Combining the bound on $\|\Psi\theta - \Psi\theta^*\|_D^2$ shown in Lemma 19 and the bound on $\frac{\mathbb{E}[\|\Psi\theta_t - \Psi\theta\|_D^2]}{T}$ from Ineq. (B.20), we obtain

$$\begin{aligned} \sum_{t=1}^T \frac{\mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2]}{T} &\leq \sum_{t=1}^T 2\mathbb{E}[\|\Psi\theta_t - \Psi\theta\|_D^2]/T + 2\mathbb{E}[\|\Psi\theta^* - \Psi\theta\|_D^2] \\ &\leq \frac{4/\mu + 264T\eta^2}{T\eta(1-\beta)} \mathbb{E}[\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2] + \left(\frac{4/\mu + 264T\eta^2}{T\eta(1-\beta)} + 2 \right) \mathbb{E}[\|\Psi\theta - \Psi\theta^*\|_D^2] \\ &\leq \frac{4/\mu + 264T\eta^2}{T\eta(1-\beta)} \mathbb{E}[\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2] + \left(\frac{4/\mu + 264T\eta^2}{T\eta(1-\beta)} + 2 \right) \left(\frac{120\mathbb{E}[\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2] + 3\mathbb{E}[\|\bar{Q}^* - \Psi\theta^*\|_D^2]}{2\mu(1-\beta)^2 N_k} \right. \\ &\quad \left. + \frac{15\bar{c}^2}{N'_k(1-\beta)^2\mu} + \frac{120\bar{c}^2}{N_k N'_k(1-\beta)^2\mu} + \frac{9}{N_k} \text{trace}\{(I_d - M)^{-1} \bar{\Sigma} (I_d - M)^{-\top}\} \right). \end{aligned} \quad (\text{B.21})$$

Then the final term takes the desired form; we are left with bounding the first two terms in the display above. First, recall by assumption that $\eta \leq \frac{1-\beta}{845}$, and $T \geq \frac{64}{\mu(1-\beta)\eta}$, so that

$$\frac{4/\mu + 264T\eta^2}{T\eta(1-\beta)} \leq \frac{4/\mu + 64 \cdot 264\eta/\mu(1-\beta)}{64/\mu} \leq \frac{3}{8}. \quad (\text{B.22})$$

Second, utilizing the condition $N_k \geq \frac{1160}{\mu(1-\beta)^2}$, we have

$$\left(\frac{4/\mu + 264T\eta^2}{T\eta(1-\beta)} + 2 \right) \frac{60}{\mu(1-\beta)^2 N_k} \leq \frac{1}{8}. \quad (\text{B.23})$$

Substituting the bounds (B.22) and (B.23) into inequality (B.21), we obtain

$$\frac{\sum_{t=1}^T \mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2]}{T} \leq \frac{\mathbb{E}[\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2]}{2} + \frac{22\text{trace}\{(I_d - M)^{-1} \bar{\Sigma} (I_d - M)^{-\top}\}}{N_k} + \frac{4\mathbb{E}[\|\bar{Q}^\pi - \Psi\theta^*\|_D^2]}{\mu(1-\beta)^2 N_k} + \frac{36\bar{c}^2}{N'_k(1-\beta)^2\mu} + \frac{\bar{c}^2}{4N'_k}. \quad (\text{B.24})$$

By definition of $\hat{\theta}_k = \frac{\sum_{t=1}^T \theta_t}{T}$ and Jensen's inequality, we have $\mathbb{E}[\|\Psi\hat{\theta}_k - \Psi\theta^*\|_D^2] \leq \frac{\sum_{t=1}^T \mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2]}{T}$, which completes the proof of Proposition 4.

B.4 Proof of Theorem 2

Similar as the proof of Theorem 1, we first focus on a single epoch. For simplicity, we let $\bar{\theta}_t$ denote $\mathbb{E}[\theta_t]$, let $\bar{\theta}$ denote $\mathbb{E}[\theta]$. We first establish a lemma to bound the term $\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2$.

Lemma 20. *We have*

$$\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2 \leq \frac{3e^{2\tau'}}{\mu(1-\beta)^2(1-e^{\tau'})^2} \left(\frac{C_P^2 \cdot \bar{c}^2}{N^{\tau'}} + \frac{2C_M^2 \mathbb{E}[\|\Psi\bar{\theta} - \Psi\theta^*\|_D^2] + C_M^2 \|\Psi\theta^* - \bar{Q}^*\|_D^2 + 8C_R^2 \cdot \bar{c}^2}{N^2} \right).$$

The proof of this lemma is postponed to Section B.4.1.

Now we switch our attention to bound the iterates. Recall the definition $\tilde{g}_g(\theta, \rho, \xi_t) := \tilde{g}(\theta, \rho, \xi_t) - g(\theta, \rho)$. By taking expectation on both sides of Eq. (B.17) and using the linearity of g , we obtain

$$\bar{\theta}_{t+1} - \bar{\theta} = \bar{\theta}_t - \bar{\theta} - \eta(g(\bar{\theta}_t, \mathbb{E}[\tilde{\rho}]) - g(\bar{\theta}, \mathbb{E}[\tilde{\rho}])) + \eta \mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t)].$$

Taking squared ℓ_2 -norm on both sides of the above equality yields

$$\begin{aligned} \|\bar{\theta}_{t+1} - \bar{\theta}\|_2^2 &= \|\bar{\theta}_t - \bar{\theta}\|_2^2 - 2\eta \langle g(\bar{\theta}_t, \mathbb{E}[\tilde{\rho}]) - g(\bar{\theta}, \mathbb{E}[\tilde{\rho}]), \bar{\theta}_t - \bar{\theta} \rangle + 2\eta \langle \mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t)], \bar{\theta}_t - \bar{\theta} \rangle \\ &\quad + \eta^2 \| (g(\bar{\theta}_t, \mathbb{E}[\tilde{\rho}]) - g(\bar{\theta}, \mathbb{E}[\tilde{\rho}])) - \mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t)] \|_2^2 \\ &\leq \|\bar{\theta}_t - \bar{\theta}\|_2^2 - 2\eta \langle g(\bar{\theta}_t, \mathbb{E}[\tilde{\rho}]) - g(\bar{\theta}, \mathbb{E}[\tilde{\rho}]), \bar{\theta}_t - \bar{\theta} \rangle + 2\eta \langle \mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t)], \bar{\theta}_t - \bar{\theta} \rangle \\ &\quad + 2\eta^2 \|g(\bar{\theta}_t, \mathbb{E}[\tilde{\rho}]) - g(\bar{\theta}, \mathbb{E}[\tilde{\rho}])\|_2^2 + 2\eta^2 \|\mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t)]\|_2^2, \end{aligned} \quad (\text{B.25})$$

where the last inequality follows from Young's inequality. By Jensen's inequality, we have that

$$\|\mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t)]\|_2^2 \leq \mathbb{E} \|\mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t) | \mathcal{F}_{t-1}]\|_2^2 \leq C_M^2 e^{2\tau} \mathbb{E}[\|\Psi\theta_t - \Psi\bar{\theta}\|_D^2]. \quad (\text{B.26})$$

Using similar idea, we obtain another upper bound

$$\begin{aligned} &\langle \mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t)], \bar{\theta}_t - \bar{\theta} \rangle \\ &\stackrel{(i)}{\leq} \mathbb{E} \|\mathbb{E}[\tilde{\delta}_g(\bar{\theta}, \tilde{\rho}, \xi_t) - \hat{\delta}_g(\bar{\theta}_t, \tilde{\rho}, \xi_t) | \mathcal{F}_{t-1}]\|_2 \|\bar{\theta}_t - \bar{\theta}\|_2 \\ &\stackrel{(ii)}{\leq} \frac{C_M e^{\tau}}{\sqrt{\mu}} \mathbb{E}[\|\Psi\theta_t - \Psi\bar{\theta}\|_D] \|\Psi\bar{\theta}_t - \Psi\bar{\theta}\|_D \stackrel{(iii)}{\leq} \frac{C_M e^{\tau}}{2\sqrt{\mu}} (\mathbb{E}[\|\Psi\theta_t - \Psi\bar{\theta}\|_D^2] + \|\Psi\bar{\theta}_t - \Psi\bar{\theta}\|_D^2), \end{aligned} \quad (\text{B.27})$$

where step (i) follows from Cauchy-Schwarz inequality and Jensen's inequality, step (ii) follows from $\|\theta\|_2 \leq \frac{1}{\sqrt{\mu}} \|\Psi\theta\|_D$ and Lemma 4, step (iii) follows from Jensen's inequality and Young's inequality. Combining Ineqs. (B.25), (B.26), (B.27) we obtain

$$\begin{aligned} &\|\bar{\theta}_{t+1} - \bar{\theta}\|_2^2 \\ &\leq \|\bar{\theta}_t - \bar{\theta}\|_2^2 - 2\eta \langle g(\bar{\theta}_t, \mathbb{E}[\tilde{\rho}]) - g(\bar{\theta}, \mathbb{E}[\tilde{\rho}]), \bar{\theta}_t - \bar{\theta} \rangle + \frac{\eta C_M e^{\tau}}{\sqrt{\mu}} (\mathbb{E}[\|\Psi\theta_t - \Psi\bar{\theta}\|_D^2] + \|\Psi\bar{\theta}_t - \Psi\bar{\theta}\|_D^2) \\ &\quad + 2\eta^2 \|g(\bar{\theta}_t, \mathbb{E}[\tilde{\rho}]) - g(\bar{\theta}, \mathbb{E}[\tilde{\rho}])\|_2^2 + 2\eta^2 C_M^2 e^{2\tau} \mathbb{E}[\|\Psi\theta_t - \Psi\bar{\theta}\|_D^2] \\ &\stackrel{(i)}{\leq} \|\bar{\theta}_t - \bar{\theta}\|_2^2 + (8\eta^2 - 2\eta(1-\beta) + \frac{\eta C_M e^{\tau}}{\sqrt{\mu}}) \|\Psi\bar{\theta}_t - \Psi\bar{\theta}\|_D^2 + (\frac{\eta C_M e^{\tau}}{\sqrt{\mu}} + 2\eta^2 C_M^2 e^{2\tau}) \mathbb{E}[\|\Psi\theta_t - \Psi\bar{\theta}\|_D^2] \\ &\stackrel{(ii)}{\leq} \|\bar{\theta}_t - \bar{\theta}\|_2^2 - \eta(1-\beta) \|\Psi\bar{\theta}_t - \Psi\bar{\theta}\|_D^2 + (\frac{2\eta C_M e^{\tau}}{\sqrt{\mu}} + 4\eta^2 C_M^2 e^{2\tau}) (\mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2] + \mathbb{E}[\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2]). \end{aligned}$$

where step (i) follows from Lemma 3 and 14, step (ii) follows from Young's inequality, $\eta \leq \frac{1-\beta}{9}$ and $\varrho^\tau \leq \frac{\sqrt{\mu}\eta}{C_M}$. By taking a telescope sum of the above inequality from $t = 1$ to T , we have

$$\begin{aligned} & \sum_{t=1}^T \eta(1-\beta) \|\Psi\bar{\theta}_t - \Psi\bar{\theta}\|_D^2 \\ & \leq \|\bar{\theta}_1 - \bar{\theta}\|_D^2 + \left(\frac{2\eta C_M \varrho^\tau}{\sqrt{\mu}} + 4\eta^2 C_M^2 \varrho^{2\tau}\right) (T\mathbb{E}[\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2] + \sum_{t=1}^T \mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2]). \end{aligned}$$

By definition, we have $\hat{\theta}_k = \frac{\sum_{t=1}^T \theta_t}{T}$; noting in addition that $\theta_1 = \bar{\theta}$, we have

$$\begin{aligned} & T\eta(1-\beta) \|\Psi\mathbb{E}[\hat{\theta}_k] - \Psi\bar{\theta}\|_D^2 \\ & \leq \frac{1}{\mu} \|\Psi\mathbb{E}[\bar{\theta}] - \Psi\bar{\theta}\|_D^2 + \left(\frac{2\eta C_M \varrho^\tau}{\sqrt{\mu}} + 4\eta^2 C_M^2 \varrho^{2\tau}\right) (T\mathbb{E}[\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2] + \sum_{t=1}^T \mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2]) \\ & \leq \frac{2}{\mu} (\|\Psi\mathbb{E}[\bar{\theta}] - \Psi\theta^*\|_D^2 + \|\Psi\theta^* - \Psi\bar{\theta}\|_D^2) \\ & \quad + \left(\frac{2\eta C_M \varrho^\tau}{\sqrt{\mu}} + 4\eta^2 C_M^2 \varrho^{2\tau}\right) (T\mathbb{E}[\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2] + \sum_{t=1}^T \mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2]). \end{aligned} \quad (\text{B.28})$$

Recalling the upper bound of $\sum_{t=1}^T \mathbb{E}[\|\Psi\theta_t - \Psi\theta^*\|_D^2]/T$ in Ineq. (B.24) and Young's inequality $\|\Psi\mathbb{E}[\hat{\theta}_k] - \Psi\theta^*\|_D^2 \leq 2\|\Psi\mathbb{E}[\bar{\theta}] - \Psi\bar{\theta}\|_D^2 + 2\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2$, we have

$$\begin{aligned} \|\Psi\mathbb{E}[\hat{\theta}_k] - \Psi\theta^*\|_D^2 & \leq \frac{4}{\mu T \eta(1-\beta)} \|\Psi\mathbb{E}[\bar{\theta}] - \Psi\theta^*\|_D^2 + \left(2 + \frac{4}{\mu T \eta(1-\beta)}\right) \|\Psi\theta^* - \Psi\bar{\theta}\|_D^2 \\ & \quad + \frac{4\eta C_M \varrho^\tau / \sqrt{\mu} + 8\eta^2 C_M^2 \varrho^{2\tau}}{\eta(1-\beta)} \left(\frac{3}{2} \mathbb{E}[\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2] + \frac{W_1}{N_k} + \frac{36\bar{c}^2}{N'_k(1-\beta)^2\mu} + \frac{\bar{c}^2}{4N'_k}\right), \end{aligned}$$

where $W_1 := 22\text{trace}\{(I_d - M)^{-1}\bar{\Sigma}(I_d - M)^{-\top}\} + \frac{4}{\mu(1-\beta)^2} \mathbb{E}[\|(\bar{Q}^* - \Psi\theta^*)\|_D^2]$. Noticing that if the conditions (3.23) and (3.26) are satisfied, we have that all assumptions of Theorem 1 are satisfied. Therefore, by utilizing Theorem 1 and $\bar{\theta} = \hat{\theta}_{k-1}$, $N_k = N$ and $N'_k = N'$, we obtain

$$\begin{aligned} \|\Psi\mathbb{E}[\hat{\theta}_k] - \Psi\theta^*\|_D^2 & \leq \frac{4}{\mu T \eta(1-\beta)} \|\Psi\mathbb{E}[\bar{\theta}] - \Psi\theta^*\|_D^2 + \left(2 + \frac{4}{\mu T \eta(1-\beta)}\right) \|\Psi\theta^* - \Psi\bar{\theta}\|_D^2 \\ & \quad + \frac{4\eta C_M \varrho^\tau / \sqrt{\mu} + 8\eta^2 C_M^2 \varrho^{2\tau}}{\eta(1-\beta)} \left(\frac{3}{2^k} \|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \frac{11W_1}{2N} + \frac{198\bar{c}^2}{N'(1-\beta)^2\mu} + \frac{11\bar{c}^2}{8N'}\right), \end{aligned}$$

Finally, invoking the bound of $\|\Psi\theta^* - \Psi\bar{\theta}\|_D^2$ in Lemma 20 and the fact that $\frac{4}{\mu T \eta(1-\beta)} \leq \frac{1}{2}$, we obtain

$$\begin{aligned} & \|\Psi\mathbb{E}[\hat{\theta}_k] - \Psi\theta^*\|_D^2 \\ & \leq \frac{1}{2} \|\Psi\mathbb{E}[\bar{\theta}] - \Psi\theta^*\|_D^2 + \frac{15\varrho^{2\tau'}}{2\mu(1-\beta)^2(1-\varrho^{\tau'})^2} \left(\frac{C_P^2 \cdot \bar{c}^2}{N'^2} + \frac{2C_M^2 \mathbb{E}[\|\Psi\bar{\theta} - \Psi\theta^*\|_D^2] + C_M^2 \|\Psi\theta^* - \bar{Q}^\pi\|_D^2 + 8C_R^2 \cdot \bar{c}^2}{N^2}\right) \\ & \quad + \frac{4\eta C_M \varrho^\tau / \sqrt{\mu} + 8\eta^2 C_M^2 \varrho^{2\tau}}{\eta(1-\beta)} \left(\frac{3}{2^k} \|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \frac{11W_1}{2N} + \frac{198\bar{c}^2}{N'(1-\beta)^2\mu} + \frac{11\bar{c}^2}{8N'}\right). \end{aligned}$$

Invoking that $N \geq \frac{1}{\mu(1-\beta)^2}$ and $N' \geq \frac{1}{\mu(1-\beta)^2}$, we have with constants b_1 and b_2 ,

$$\|\Psi\mathbb{E}[\hat{\theta}_k] - \Psi\theta^*\|_D^2 \leq \frac{1}{2} \|\Psi\mathbb{E}[\hat{\theta}_{k-1}] - \Psi\theta^*\|_D^2 + \left(b_1 C_{\max}^2 \varrho^{2\tau'} + \frac{b_2 C_{\max} \varrho^\tau}{(1-\beta)\sqrt{\mu}}\right) (\|\Psi\theta^0 - \Psi\theta^*\|_D^2 + \frac{W_1}{N} + \bar{c}^2).$$

By recursively using this inequality for epochs, we achieve the desired result.

B.4.1 Proof of Lemma 20

Taking expectation on both sides of Eq. (B.14), we obtain

$$B^{\frac{1}{2}}(\theta^* - \bar{\theta}) = (I_d - M)^{-1} B^{-\frac{1}{2}} (\Psi^\top D \mathbf{1}(\mathbb{E}[\tilde{\rho}] - \rho^*) + \mathbb{E}[\hat{g}(\tilde{\theta}, \tilde{\rho}) - g(\tilde{\theta}, \tilde{\rho})]). \quad (\text{B.29})$$

Invoking the fact that $\Phi^\top D \Phi = I_d$ and $\Phi^\top = B^{-\frac{1}{2}} \Psi^\top$ yields

$$\begin{aligned} \|\Psi\theta^* - \Psi\bar{\theta}\|_D^2 &= \|(I_d - M)^{-1} B^{-\frac{1}{2}} (\Psi^\top D \mathbf{1}(\mathbb{E}[\tilde{\rho}] - \rho^*) + \mathbb{E}[\hat{g}(\tilde{\theta}, \tilde{\rho}) - g(\tilde{\theta}, \tilde{\rho})])\|_2^2 \\ &\stackrel{(i)}{\leq} 3(\mathbb{E}[\tilde{\rho}] - \rho^*)^2 \cdot \|(I_d - M)^{-1} B^{-\frac{1}{2}} \Psi^\top D \mathbf{1}\|_2^2 + 3\|(I_d - M)^{-1} B^{-\frac{1}{2}} \mathbb{E}[\hat{g}(\theta^*, \rho^*) - g(\theta^*, \rho^*)]\|_2^2 \\ &\quad + 3\|(I_d - M)^{-1} B^{-\frac{1}{2}} \mathbb{E}[\hat{g}(\tilde{\theta}, \tilde{\rho}) - \hat{g}(\theta^*, \rho^*) - g(\tilde{\theta}, \tilde{\rho}) + g(\theta^*, \rho^*)]\|_2^2 \\ &\stackrel{(ii)}{\leq} 3\left(\frac{C_P \varrho^{\tau'}}{N'(1-\varrho^{\tau'})}\right)^2 \cdot \frac{\bar{c}^2}{(1-\beta)^2 \mu} + \frac{3}{(1-\beta)^2 \mu} \left(\frac{\varrho^{\tau'}}{N(1-\varrho^{\tau'})}\right)^2 C_M^2 \|\Psi\theta^* - \bar{Q}^\pi\|_D^2 \\ &\quad + \frac{3}{(1-\beta)^2 \mu} \cdot \left(\frac{\varrho^{\tau'}}{N(1-\varrho^{\tau'})}\right)^2 \cdot (2C_M^2 \mathbb{E}[\|\Psi\tilde{\theta} - \Psi\theta^*\|_D^2] + 8C_R^2 \cdot \bar{c}^2) \end{aligned}$$

where step (i) follows from Young's inequality, step (ii) follows from Assumption 2, Lemma 4 and Lemma 5. Then the desired result immediately follows from the above inequality.